An injective martingale coupling

David Hobson[∗] Dominykas Norgilas†

March 6, 2023

Abstract

We give an injective martingale coupling; in particular, given measures μ and ν in convex order on R such that ν is continuous, we construct a martingale coupling π of the two measures with disintegration $\pi(dx, dy) = \pi_x(dy)\mu(dx)$ such that $\#\{x : y \in \text{supp}(\pi_x)\} = 1$ for ν almost all y . Equivalently we find a martingale coupling π such that π has disintegration $\pi(dx, dy) = \nu(dy)\delta_{\theta(y)}(dx)$ for some function θ .

Keywords: martingale couplings, Strassen's theorem, convex order. 2010 Mathematics Subject Classification: 60G42.

1 Introduction

Suppose μ and ν are measures on R and in convex order. Define $\mathcal{M}(\mu, \nu)$ to be the set of martingale couplings of μ and ν , i.e., the set of probability measures π on $\mathbb{R}\times\mathbb{R}$ such that π has first marginal μ , second marginal ν and satisfies the martingale property $\int_{x \in A} (y - x) \pi(dx, dy) = 0$ for all Borel sets $A \subseteq \mathbb{R}$. Equivalently, π is the joint law of a pair of random variables (X, Y) such that $\mathcal{L}(X) = \mu$, $\mathcal{L}(Y) = \nu$ and $\mathbb{E}^{\pi}[Y|X] = X$. For $\pi \in \mathcal{M}(\mu, \nu)$ we can write π in terms of its disintegration $\pi(dx, dy) = \mu(dx)\pi_x(dy)$.

Ruodu Wang asks: if ν is continuous then is there a martingale coupling $\pi \in \mathcal{M}(\mu, \nu)$ such that $\#\{x : y \in \text{supp}(\pi_x)\} \leq 1$, or equivalently such that $\mathcal{L}(X|Y)$ is a point mass? We show the answer is yes by constructing such a π . This means that given Y we can recover X, and thus $X = \theta(Y)$. We call such a martingale coupling an *injective* martingale coupling and write $\mathcal{IM}(\mu, \nu)$ for the set of injective martingale couplings.

[∗]Department of Statistics, University of Warwick d.hobson@warwick.ac.uk

[†]Department of Mathematics, University of Michigan dnorgila@umich.edu

In the forward direction the martingale property means that we cannot have $Y =$ $\phi(X)$ except in the trivial situation when $\mu = \nu$. However, one can look for π such that supp (π_x) is small. Many constructions have this property. For example, at least when μ is continuous, the construction of Hobson and Neuberger [\[7\]](#page-38-0) and the left-curtain coupling of Beiglböck and Juillet [\[2\]](#page-38-1) are such that $\#\{y : y \in \text{supp}(\pi_x)\} \leq 2$, for all $x \in \mathbb{R}$, and the construction of Hobson and Klimmek [\[6\]](#page-38-2) is such that $\#\{y : y \in \mathbb{R}\}$ $\text{supp}(\pi_x) \leq 3$, for all $x \in \mathbb{R}$. We are interested in the reverse direction, can we make the pre-image of y under the martingale coupling a singleton for $(\nu \text{ almost})$ all y.

It is clear that there is no possibility of an injective martingale coupling (except in a few special cases) if ν has atoms^{[1](#page-1-0)}. For example, consider the case where $\mu \sim U[-1,1]$ and $\nu = \frac{1}{2}$ $rac{1}{2}\delta_{-1}+\frac{1}{2}$ $\frac{1}{2}\delta_1$. Then $\mathcal{M}(\mu, \nu)$ is a singleton for which $\pi_x(dy) = \frac{(x+1)}{2}\delta_1(dy) +$ $(1-x)$ $\frac{-x}{2}$ $\delta_{-1}(dy)$. Then, for $y = 1$, $\{x : y \in \text{supp}(\pi_x)\} = (-1, 1]$, and for $y = -1$, ${x : y \in \text{supp}(\pi_x)} = [-1, 1)$, and in either case (i.e., *v*-almost surely), modulo a μ -null set, $\{x : y \in \text{supp}(\pi_x)\}\$ is equal to supp (μ) .

Further, if there exists an injective coupling then there is no expectation of uniqueness, as the following example shows.

Example 1.1. This example is taken from Hobson and Neuberger [\[7,](#page-38-0) Section 6.3]. Suppose $\mu \sim U[-1,1]$ and $\nu \sim U[-2,2]$. For each $a \in [-1,1]$ there exists a pair of monotonic increasing surjective functions $f_{HN}^a: [-1,1] \mapsto [-2,a]$ and $h_{HN}^a: [-1,1] \mapsto$ $[a, 2]$ such that for $i = 0, 1$,

$$
\int_{-1}^{x} z^{i} \frac{dz}{2} = \int_{-2}^{f_{HN}^{a}(x)} z^{i} \frac{dz}{4} + \int_{a}^{h_{HN}^{a}(x)} z^{i} \frac{dz}{4}, \quad x \in \mathbb{R}.
$$
 (1)

If we set $\pi^{f_{HN}^a,h_{HN}^a}(dx,dy) = \mu(dx)\pi_x^{f_{HN}^a,h_{HN}^a}(dy)$ where for $f(x) \leq x \leq h(x)$

$$
\pi_x^{f,h}(dy) = \frac{h(x) - x}{h(x) - f(x)} \delta_{f(x)}(dy) + \frac{x - f(x)}{h(x) - f(x)} \delta_{h(x)}(dy),
$$

then $\pi^{f_{HN}^a, h_{HN}^a} \in \mathcal{M}(U[-1,1], U[-2,2])$. (The pair of conditions in [\(1\)](#page-1-1) are exactly the conditions required to ensure that the initial law mass in $[-1, x)$ maps onto the target law mass in $[-2, f_{HN}^a(x)) \cup [a, h_{HN}^a(x)]$ in a way which preserves both mass and mean.) Moreover it is clear that $\pi^{f_{HN}^a, h_{HN}^a}$ is an injective martingale coupling; for $y \in (a, 2]$ we have $\{x : y \in \text{supp}(\pi_x)\} = \{x : y = h_{HN}^a(x)\} = \{(h_{HN}^a)^{-1}(y)\}$ and for $y \in [-2, a)$, ${x : y \in supp(\pi_x)} = {(f_{HN}^a)^{-1}(y)}$. On the other hand, ${x : a \in supp(\pi_x)}$ $\{(h_{HN}^a)^{-1}(a), (f_{HN}^a)^{-1}(a)\} = \{-1, 1\}$, but since $\nu(\{a\}) = 0$ the injectiveness holds. By solving [\(1\)](#page-1-1) explicitly, for each $a \in [-1,1]$ we obtain

$$
h_{HN}^a(x) = \frac{2x + a + \sqrt{4 + a^2 - 4ax}}{2} = 2x + a - f_{HN}^a(x), \quad x \in [-1, 1].
$$

¹One such special case is when μ is a point mass.

Note that when $a = 0$ we find that $f_{HN}^a(x) = x - 1$ and $h_{HN}^a(x) = x + 1$.

The martingale coupling in Hobson and Neuberger [\[7\]](#page-38-0) was designed to have other properties and not to be an injective coupling, and it seems difficult to extend the argument to the general case. (Indeed, the general construction in [\[7,](#page-38-0) Section 6.3.] fails to be injective.) Instead, including all cases where ν is atom free, we shall design a new martingale coupling which is injective for a wide class of pairs of measures (μ, ν) in convex order. By so doing we prove the following theorem:

Theorem 1.2. Suppose $\mu \leq_{cx} \nu$ where μ is an arbitrary measure and ν is continuous. Then $\mathcal{IM}(\mu, \nu) \neq \emptyset$.

Remark 1.3. Note that the main result extends very nicely to multiple marginals. Suppose $(\nu_k)_{0 \leq k \leq n}$ are increasing in convex order and ν_k is continuous for $k \geq 1$. Then there is a martingale coupling such that if π denotes the joint law of a martingale $Y = (Y_k)_{0 \le k \le n}$ with the given marginals then $\pi(dy_0, dy_1, \ldots, dy_n) = \nu_n(dy) \prod_{i=0}^{n-1} \delta_{\theta_i}(dy_i)$ for some functions $(\theta_i)_{0\leq i\leq n-1}$. In particular we can recover the whole history of Y from its terminal value.

Whilst finishing the current paper, we became aware of Nutz et al. [\[10\]](#page-38-3), where (among other things) the authors established Theorem [1.2.](#page-2-0) (In the language of [\[10\]](#page-38-3), $\mathcal{IM}(\mu, \nu)$ is the set of *backward Monge* martingale couplings, but we think that the term injective is more suggestive.) Both Nutz et al. and this paper make extensive use of the left-curtain martingale coupling π^{lc} , introduced by Beiglböck and Juillet [\[2\]](#page-38-1). However, π^{lc} is not injective - typically (when ν is continuous) for the disintegration π_x^{lc} of π^{lc} we have $\#\{x : y \in \text{supp}(\pi_x)\} \leq 2$ but there is a large set of y such that $y \in \{x : y \in \text{supp}(\pi_x)\}\$ along with another point.

The main insight of [\[10\]](#page-38-3) is that by restricting μ to a set $A_{\text{NWZ}} = \{d\mu/(d\mu + d\nu) \geq$ $d\nu/(d\mu + d\nu)$, and then by embedding this restriction to ν via the shadow measure $S^{\nu}(\mu|_{A_{\text{NWZ}}})$ (see Beiglböck and Juillet [\[3\]](#page-38-4)), one can find $\pi^{A_{\text{NWZ}}} \in \mathcal{IM}(\mu|_{A_{\text{NWZ}}}, S^{\nu}(\mu|_{A_{\text{NWZ}}}))$. In particular, the left-curtain coupling of $\mu|_{A_{\text{NWZ}}}$ and $S^{\nu}(\mu|_{A_{\text{NWZ}}})$ is injective. Then one (inductively) repeats this process for the remaining masses $(\mu - \mu|_{A_{\text{NWZ}}})$ and $(\nu - S^{\nu}(\mu|_{A_{\text{NWZ}}}))$. The corresponding coupling is baptized the *barcode* backward martingale coupling.

Our proof of Theorem [1.2](#page-2-0) also relies on the properties of the left-curtain martingale coupling π^{lc} . It uses the left-curtain coupling in two ways. First it uses the left-curtain coupling to divide the problem into a countable family of sub-problems, such that for each sub-problem the corresponding pair (μ, ν) is still in convex order, but has additional structure. Second, for each sub-problem we use a modified version of the leftcurtain coupling to construct an injective coupling. The key idea is to choose a special

starting point $x_0 \in \mathbb{R}$ and a (maximal) interval $[x_0, x_1]$ to the right of x_0 and couple $\mu|_{[x_0,x_1)}$ with $S^{\nu}(\mu|_{[x_0,x_1)})$ using the left-curtain coupling. For a well-chosen interval $[x_0, x_1]$ this coupling is injective, and can be described in semi-explicit terms (given the results of [\[1\]](#page-38-5)) using potentials and tangents. Then in the next step we consider an interval (x_2, x_0) to the left of x_0 and embed $\mu|_{(x_2, x_0)}$ in $\nu - S^{\nu}(\mu|_{[x_0, x_1)})$ via the right-curtain coupling. Again, for an appropriate choice of x_2 the construction remains injective, and the resulting disintegration $\pi_x(dy)$ can be calculated semi-explicitly. Inductively we will obtain a sequence of points $\ldots < x_{2k} < \ldots < x_2 < x_0 < x_1 < \ldots < x_{2k+1} < \ldots$ such that $\mu|_{[x_{2k},x_{2(k-1)]}}$ and $\mu|_{[x_{2k+1},x_{2k+3]}]}$ are embedded using inductive right and left-curtain couplings. We will explicitly construct two locally strictly monotonic functions such that the coupling concentrates on the graph of these two functions. What distinguishes this paper from Nutz et al. [\[10\]](#page-38-3), apart from the different constructions, is our emphasis on describing the injective coupling as fully as possible, whereas [\[10\]](#page-38-3) focusses on existence and general properties.

The outline of the paper is as follows. In the next section we introduce some notation, including the notion of a lifted martingale coupling. In Section [3](#page-9-0) we construct a pair of functions which ultimately will form the main part of the construction of an injective martingale coupling. In Section [4](#page-18-0) we show that an injective martingale coupling exists in a case with certain regularity properties by constructing a lifted martingale coupling. Then in Section [5](#page-30-0) we show how the left curtain coupling of Beiglböck and Juillet $|2|$ can be used to divide the problem for general measures μ and (continuous) ν into a countable family of sub-problems, each of which almost satisfies the regularity conditions of Section [3.](#page-9-0) Finally, in Section [6](#page-34-0) we show how each sub-problem arising from the left-curtain decomposition can be further sub-divided into a countable family of problems which do satisfy the regularity conditions. Putting it all together we deduce the existence of an injective martingale coupling.

2 Notation and preliminaries

2.1 Convex Hulls

For a continuous function $\mathcal{H}: \mathbb{R} \to \mathbb{R}_+$ let \mathcal{H}^c denote the convex hull of \mathcal{H} . For $z \in \mathbb{R}$ let $X^+_{\mathcal{H}}(z) = \sup\{w : w \leq z, \mathcal{H}(w) = \mathcal{H}^c(w)\}\$ and $Z^-_{\mathcal{H}}(z) = \inf\{w : w \geq z : w \leq z\}$ $\mathcal{H}(w) = \mathcal{H}^c(w)$ with the convention that $\sup \emptyset = -\infty$ and $\inf \emptyset = \infty$. Note that $X_{\mathcal{H}}^{+}(z) = z$ if and only if $Z_{\mathcal{H}}^{-}(z) = z$ (and also if and only if $\mathcal{H}(z) = \mathcal{H}^{c}(z)$). Now, for $z \in \mathbb{R}$, let $\psi_-(z) := (\mathcal{H}^c)'(z-)$, where $f'(z-)$ denotes the left derivative (if it exists) of a measurable function $f : \mathbb{R} \to \mathbb{R}$, and similarly, let $\psi_{+}(z) := (\mathcal{H}^{c})'(z+)$, where

 $f'(z+)$ denotes the right derivative $f'(z+)$. Define $X_{\mathcal{H}}^{-}(z) := \inf\{w : w \leq z, \mathcal{H}^c(w) =$ $L_{\mathcal{H}c}^{z,\psi_{-}(z)}(w)$ } and $Z_{\mathcal{H}}^{+}(z) := \sup\{w : w \geq z, \mathcal{H}^{c}(w) = L_{\mathcal{H}c}^{z,\psi_{+}(z)}(w)\}\,$, where, for $f : \mathbb{R} \to \mathbb{R}$, $\phi \in \mathbb{R}$ and $z \in \mathbb{R}$ we define $L_f^{z,\phi}$ $t^{z,\phi}_f$ to be the straight line $L_f^{z,\phi}$ $f_f^{z,\varphi}(k) = f(z) + \phi(k - z).$

Note that $X_{\mathcal{H}}^{-}(z) \leq X_{\mathcal{H}}^{+}(z) \leq z \leq Z_{\mathcal{H}}^{-}(z) \leq Z_{\mathcal{H}}^{+}(z)$, for all $z \in \mathbb{R}$. Indeed, $X_{\mathcal{H}}^{-}(z) \leq$ $z \leq Z^+_{\mathcal{H}}(z)$ for all $z \in \mathbb{R}$, and thus if $X^+_{\mathcal{H}}(z) = z = Z^-_{\mathcal{H}}(z)$ the claim is immediate. On the other hand, if $X^+_{\mathcal{H}}(z) < Z^-_{\mathcal{H}}(z)$, then $\mathcal{H}^c(z) < \mathcal{H}(z)$, \mathcal{H}^c is linear on $(X^+_{\mathcal{H}}(z), Z^-_{\mathcal{H}}(z)) \ni z$, and it follows that $X_{\mathcal{H}}^-(z) \leq X_{\mathcal{H}}^+(z) < z < Z_{\mathcal{H}}^-(z) \leq Z_{\mathcal{H}}^+(z)$. Also, in the latter case $(\mathcal{H}^c)'(z)$ is well-defined and $(\mathcal{H}^c)'(z) = \psi_-(z) = \psi_+(z)$.

Suppose \mathcal{H} is such that its left and right derivatives exist everywhere and that $\mathcal{H}'(z-) \geq \mathcal{H}'(z+)$ everywhere. Then $\psi_- = \psi_+$. This is clear at any z for which $\mathcal{H}(z) > \mathcal{H}^c(z)$. If $\mathcal{H}(z) = \mathcal{H}^c(z)$ then

$$
\psi_{-}(z) \leq \psi_{+}(z) = (\mathcal{H}^{c})'(z+) \leq \mathcal{H}'(z+) \leq \mathcal{H}'(z-) \leq (\mathcal{H}^{c})'(z-) = \psi_{-}(z).
$$

where the first inequality is true for any convex function, the two equalities are by definition, the second and fourth inequalities hold since $\mathcal{H}(z) = \mathcal{H}^c(z)$ and the third inequality holds by our hypothesis on H .

2.2 Measures and convex order

Let χ be a measure on R, and denote its support by supp (χ) ; supp (χ) is the smallest closed set $E \subseteq \mathbb{R}$ with $\chi(E) = \chi(\mathbb{R})$. Let $\alpha_{\chi} := \inf\{k : k \in \text{supp}(\chi)\}\$ and $\beta_{\chi} := \sup\{k : k \in \text{supp}(\chi)\}\$ $k \in \text{supp}(\chi)$. Let $I_{\chi} = \{z : \chi((-\infty, z]) \chi([z, \infty)) > 0\}$. Then $(\alpha_{\chi}, \beta_{\chi}) \subseteq I_{\chi} \subseteq [\alpha_{\chi}, \beta_{\chi}],$ and, provided χ has an atom at α_{χ} (resp. β_{χ}), we have $\alpha_{\chi} \in I_{\chi}$ (resp. $\beta_{\chi} \in I_{\chi}$). Let $\mathcal{P} = \{ \chi : \int_{\mathbb{R}} |z|^i \chi(dz) < \infty \text{ for } i = 0, 1 \}$. For $p \in (0, \infty)$ define $\mathcal{P}_p = \{ \chi \in \mathcal{P} : \chi(\mathbb{R}) = 0 \}$ p } so that \mathcal{P}_1 is the set of integrable probability measures.

For $\eta, \chi \in \mathcal{P}$ we write $\eta \leq \chi$ if $\eta(A) \leq \chi(A)$ for all Borel $A \subseteq \mathbb{R}$.

For $\chi \in \mathcal{P}$ we denote by $G_{\chi} : [0, \chi(\mathbb{R})] \to \mathbb{R}$ a quantile of function of χ , i.e., a generalized inverse of $x \mapsto \chi((-\infty, x])$. There are two canonical versions of G_{χ} : the left-continuous and right-continuous versions correspond to $\overrightarrow{G}_x(u) = \sup\{k \in$ $\mathbb{R} : \chi((-\infty, k]) < u$ (with convention sup $\emptyset = -\infty$) and $\overleftarrow{G}_\chi(u) = \inf\{k \in \mathbb{R} :$ $\chi((-\infty, k]) > u$ } (with convention inf $\emptyset = \infty$), for $u \in [0, \chi(\mathbb{R})]$, respectively. In particular, an arbitrary version of the quantile function G_χ satisfies $\overrightarrow{G}_\chi \leq G_\chi \leq \overleftarrow{G}_\chi$ on $[0, \chi(\mathbb{R})]$. Note that G_χ may take values $-\infty$ and ∞ at the left and right endpoints of [0, $\chi(\mathbb{R})$], respectively. For any quantile function we set $G_{\chi}(0-) = -\infty$ and $G_{\gamma}(\chi(\mathbb{R})+)=\infty.$

For $\chi \in \mathcal{P}$ let $\bar{\chi} = \int_{\mathbb{R}} z \chi(dz)$ and define $P_{\chi}: \mathbb{R} \mapsto \mathbb{R}_+$ by $P_{\chi}(k) = \int_{\mathbb{R}} (k-z)^+ \chi(dz)$ and $C_\chi : \mathbb{R} \to \mathbb{R}_+$ by $C_\chi(k) = \int_{\mathbb{R}} (z - k)^+ \chi(dz)$. Then P_χ (respectively C_χ) is an increasing (respectively decreasing) convex function. Note that put-call parity gives that $P_{\chi}(k) - C_{\chi}(k) = k\chi(\mathbb{R}) - \bar{\chi}.$

For $\chi \in \mathcal{P}$ and $u \in [0, \chi(\mathbb{R})]$, define $\chi_u \in \mathcal{P}_u$ by

$$
\chi_u = \chi|_{(-\infty, G_\chi(u))} + (u - \chi((-\infty, G_\chi(u)))\delta_{G_\chi(u)}, \quad u \in [0, \chi(\mathbb{R})].
$$

Note that χ_u is independent of the choice of G_χ . Then for $u \in [0, \chi(\mathbb{R})], \chi_u \leq \chi$, $\chi_u(\mathbb{R}) = u$ and $P_{\chi_u}(k) = P_{\chi}(k)$ for $k \leq G_{\chi}(u+)$. The measure χ_u consists of the leftmost part of χ of total mass u. We treat χ_0 as the zero measure, whereas $\chi_{\chi(\mathbb{R})} = \chi$.

For a pair of measures $\mu, \nu \in \mathcal{P}$ such that for $i = 0, 1, \int_{\mathbb{R}} z^i \mu(dz) = \int_{\mathbb{R}} z^i \nu(dz)$ define $D = D_{\mu,\nu} : \mathbb{R} \mapsto \mathbb{R}$ by $D_{\mu,\nu}(z) = P_{\nu}(z) - P_{\mu}(z) = C_{\nu}(z) - C_{\mu}(z)$. (The equality of these two alternative expressions follows from put-call parity.) Then, $\lim_{z\to\pm\infty}D_{\mu,\nu}(z)=0$. Moreover, $D_{\mu,\nu} \geq 0$ if and only if μ and ν are in convex order, which is denoted by $\mu \leq_{cx} \nu$ (i.e., $\int_{\mathbb{R}} f d\mu \leq \int_{\mathbb{R}} f d\nu$ for all convex $f : \mathbb{R} \to \mathbb{R}$). By a classical result of Strassen [\[11\]](#page-39-0), $\mathcal{M}(\mu, \nu)$ is non-empty if and only if $\mu \leq_{cx} \nu$.

For a pair of measures $\mu, \nu \in \mathcal{P}$ we write $\mu \leq_{pcx} \nu$ if $\int_{\mathbb{R}} f d\mu \leq \int_{\mathbb{R}} f d\nu$ for all nonnegative and convex $f : \mathbb{R} \to \mathbb{R}$. Given $\mu \leq_{pcx} \nu$, the set $\{\theta : \mu \leq_{cx} \theta \leq \nu\}$ is non-empty and admits the minimal element w.r.t. \leq_{cx} . In particular, the so-called shadow measure of μ in ν , denoted by $S^{\nu}(\mu)$, is an element of $\{\theta : \mu \leq_{cx} \theta \leq \nu\}$ such that $S^{\nu}(\mu) \leq_{cx} \chi$ for all $\chi \in \{\theta : \mu \leq_{cx} \theta \leq \nu\}$. See Beiglböck and Juillet [\[2,](#page-38-1) Lemma 4.6].

2.3 Reduction of the problem to irreducible components

If $\mu \leq_{cx} \nu$ and $x \in \mathbb{R}$ is such that $D_{\mu,\nu}(x) = 0$ then, following Hobson [\[5,](#page-38-6) page 254], for any martingale coupling $\pi \in \mathcal{M}(\mu, \nu)$ we must have that no mass can cross x, i.e., $\pi((-\infty, x] \times (x, \infty)) = 0 = \pi([x, \infty) \times (-\infty, x])$. In this case, the task of constructing π can be separated into a pair of subproblems involving the mass to the left and right of x. More generally, the set $\{x \in \mathbb{R} : D_{\mu,\nu}(x) = 0\}$ gives rise to the following 'irreducible' decomposition of μ and ν .

Lemma 2.1 (Beiglböck and Juillet [\[2,](#page-38-1) Theorem A.4]). Let $\mu, \nu \in \mathcal{P}$ with $\mu \leq_{cx} \nu$. Represent $\{k \in \mathbb{R} : D_{\mu,\nu}(k) > 0\}$ as a union of disjoint open intervals, $\bigcup_{k \geq 1} I_k$. Let $I_0 = \mathbb{R} \setminus \bigcup_{k \geq 1} I_k$. Set $\mu_k = \mu|_{I_k}$ so that $\mu = \sum_{k \geq 0} \mu_k$.

There exists the unique decomposition $\nu = \sum_{k\geq 0} \nu_k$ such that $\mu_0 = \nu_0$, $\mu_k \leq_{cx} \nu_k$ and $\{x \in \mathbb{R} : D_{\mu_k, \nu_k}(x) > 0\} = I_k$ for each $k \ge 1$. Moreover, any martingale coupling $\pi \in \mathcal{M}(\mu, \nu)$ admits the unique decomposition $\pi = \sum_{k \geq 0} \pi_k$ where $\pi_k \in \mathcal{M}(\mu_k, \nu_k)$ for all $k \ge 0$, and $supp(\pi_0) \subseteq \{(x, y) \in \mathbb{R}^2 : x = y, D_{\mu,\nu}(x) = 0\}.$

Corollary 2.2. If $\mu = \nu$ then $\mathcal{IM}(\mu, \nu) = \mathcal{M}(\mu, \nu) = {\pi_0} \neq \emptyset$.

Proof. If $\mu = \nu$ then $D_{\mu,\nu} \equiv 0$, and by Lemma [2.1](#page-5-0) we have that supp $(\pi) \subseteq \{(x, y) \in$ \mathbb{R}^2 : $x = y$ for all $\pi \in \mathcal{M}(\mu, \nu)$. In particular, $\pi(dx, dy) = \mu(dx)\delta_x(dy)$ is the unique element of $\mathcal{IM}(\mu, \nu)$. \Box

Without loss of generality we make the following standing assumption.

Standing Assumption 2.3. $\mu \leq_{cx} \nu$ and $\mu \neq \nu$.

Proposition 2.4. Let $\mu, \nu \in \mathcal{P}$ with $\mu \leq_{cx} \nu$ and suppose that ν is continuous. Let $(\mu_k, \nu_k)_{k>0}$ be as in Lemma [2.1](#page-5-0) and let $\pi_0(dx, dy) = \mu_0(dx)\delta_x(dy)$. If $\pi_k \in \mathcal{IM}(\mu_k, \nu_k)$ for all $k \geq 1$, then $\pi := \sum_{k \geq 0} \pi_k$ is such that $\pi \in \mathcal{IM}(\mu, \nu)$.

Proof. All that remains to show is the injective property. For each $k \geq 1$ let I_k be an open interval as in Lemma [2.1,](#page-5-0) and let α_k and β_k be the left and right end points of I_k , respectively. Without the continuity property on ν we may have that ν_k places mass on $[\alpha_k, \beta_k]$ and then if $\beta_k = \alpha_{k'}$ for some $k' \neq k$ (or $\beta_k \in \text{supp}(\nu_0)$) we may have that $\beta_k \in \text{supp}(\nu_k) \cap \text{supp}(\nu_{k'})$ (or $\beta_k \in \text{supp}(\nu_k) \cap \text{supp}(\nu_0)$) and $\#\{x : \beta_k \in \text{supp}(\pi_x)\} > 1$. But, by our assumption on ν , ν_k places mass on $(\alpha_k, \beta_k) = I_k$. Then, given y, either $y \in I_k$ for a unique $k \geq 1$ and since $\pi_k \in \mathcal{IM}(\mu_k, \nu_k)$ there exists a unique $x \in I_k$ with $y \in \text{supp}(\pi_x)$, or $y \in I_0 = \mathbb{R} \setminus \bigcup_{k \geq 1} I_k$ and the only x for which it is possible to have $y \in \text{supp}(\pi_x)$ is $x = y$. \Box

2.4 Lifted martingale couplings

To deal with atoms in the initial law we consider the notion of a lifted martingale coupling introduced by Beiglböck and Juillet $[3]$ (see also Hobson and Norgilas $[9]$). This allows us to reinterpret the problem as one of finding a pair of functions, so that instead of mapping masses to sets, single points are mapped to at most two points in a way which respects the martingale property.

Fix $\chi \in \mathcal{P}$ and, for each $u \in [0, \chi(\mathbb{R})]$, recall the definition of $\chi_u \in \mathcal{P}_u$ (see Sec-tion [2.2\)](#page-4-0). The *quantile lift* $\hat{\chi}^Q$ of χ is a measure on $(0, \chi(\mathbb{R})) \times I_{\chi}$ with support $\{(v, G_\chi(v)); 0 < v < \chi(\mathbb{R})\}$ such that $\hat{\chi}^Q(du, dx) = du \delta_{G_\chi(u)}(dx)$. Then, for each $u \in (0, \chi(\mathbb{R})), \int_{v \in (0, u)} dv \delta_{G_{\chi}(v)}(dx) = \chi_u(dx).$

Given $\mu \leq_{cx} \nu$ a quantile-lifted martingale coupling of μ and ν is a measure $\hat{\pi}^Q$ on $(0, \mu(\mathbb{R})) \times I_{\mu} \times I_{\nu}$ given by $\hat{\pi}^Q(du, dx, dy) = du \delta_{G_{\mu}(u)}(dx) \hat{\pi}_{u,x}^Q(dy)$ such that $\hat{\pi}_{u,x}^Q(dy) =$ $\hat{\pi}^Q_n$ $\int_{u,G_\mu(u)}^Q(dy), \int_y \hat{\pi}_{u,x}^Q(dy) = 1, \int_y y \hat{\pi}_{u,x}^Q(dy) = x \text{ for } \hat{\mu}^Q \text{-a.e. } (u,x), \text{ and } \int_{u \in (0,\mu(\mathbb{R}))} \int_{x \in I_\mu} \hat{\pi}^Q(du, dx, dy) = 0$ $\nu(dy)$. Then if $\pi(dx, dy) := \int_{u \in (0,\mu(\mathbb{R}))} \hat{\pi}^Q(du, dx, dy)$ it follows that $\pi \in \mathcal{M}(\mu, \nu)$. Let $\mathcal{M}_{QL}(\mu, \nu)$ be the set of quantile-lifted martingale couplings of μ and ν .

Definition 2.5. A quantile-lifted martingale coupling $\hat{\pi}^Q \in \mathcal{M}_{QL}(\mu, \nu)$ is injective if for v-a.e. $y \in \mathbb{R}$ there exists the unique $u \in (0, \mu(\mathbb{R}))$ such that $y \in supp(\hat{\pi}_{u}^Q)$ $\big\{ \begin{matrix} Q_u, & \ldots, & Q_{\mu}(u) \end{matrix} \big\}.$

Let $\mathcal{IM}_{QL}(\mu, \nu)$ be the set of injective quantile-lifted martingale couplings of μ to ν . It follows from the definitions that if $\hat{\pi}^Q \in \mathcal{IM}_{QL}(\mu, \nu)$ then π defined by $\pi(dx, dy) :=$ $\int_{u\in(0,\mu(\mathbb{R}))} \hat{\pi}^Q(du, dx, dy)$ is such that $\pi \in \mathcal{IM}(\mu, \nu)$. Hence, the goal in future sections will be to show that $\mathcal{IM}_{QL}(\mu, \nu)$ is non-empty from which the existence of an injective martingale coupling follows immediately.

Sometimes it is convenient to define a quantile lift for sub-measures in a more general way which allows for superposition. In what follows sets J and J_k are assumed to be Borel.

Definition 2.6. Suppose $\eta \in \mathcal{P}$ and $J \subseteq (0, \eta(\mathbb{R}))$. Let $\eta^J \in \mathcal{P}$ be given by $\eta^J(A) =$ $Leb(\lbrace u \in J : G_{\eta}(u) \in A \rbrace)$, for all $(Borel)$ $A \subseteq \mathbb{R}$ so that $\eta^{J}(dx) = \int_{u \in (0, \eta(\mathbb{R}))} du I_{\lbrace u \in J \rbrace} \delta_{G_{\eta}(du)}(dx)$.

The J-quantile lift of η^J in η , denoted by $\hat{\eta}^{Q,J}$, is a measure on $(0, \eta(\mathbb{R})) \times I_{\eta}$ with support $(u, G_{\eta}(u))_{u \in J}$ such that $\hat{\eta}^{Q,J}(du, dx) = I_{\{u \in J\}} du \delta_{G_{\eta}(u)}(dx)$.

Suppose $\chi \in \mathcal{P}$ is such that $\eta^J \leq_{cx} \chi$. Then a J-quantile-lifted martingale coupling of η^J and χ , denoted by $\hat{\pi}^{Q,J}$, is a measure on $J \times I_{\eta^J} \times I_{\chi}$ such that $\hat{\pi}^{Q,J}$ is of the form

$$
\hat{\pi}^{Q,J}(du, dx, dy) = I_{\{u \in J\}} du \delta_{G_{\eta}(u)}(dx) \hat{\pi}^{Q}_{u,x}(dy),
$$

where $\hat{\pi}_{u,x}^Q(dy) = \hat{\pi}_{u,x}^Q$ $\int_{u,G_\eta(u)}^Q(dy)$, $\int_y \hat{\pi}_{u,x}^Q(dy) = 1$, $\int_y y\hat{\pi}_{u,x}^Q(dy) = x$ $\hat{\eta}^{Q,J}$ almost everywhere, and

$$
\int_{u \in (0,\eta(\mathbb{R}))} \int_{x \in I_{\eta}} \hat{\pi}^{Q,J}(du, dx, dy) = \chi(dy).
$$

Note that if $J = (0, \eta(\mathbb{R}))$ then $\hat{\eta}^{Q,J} = \hat{\eta}^Q$.

Given a quantile-lifted martingale coupling, it is straight-forward to construct Jquantile-lifted martingale couplings. Suppose $\eta, \chi \in \mathcal{P}$ are such that $\eta \leq_{cx} \chi$ and suppose $\hat{\pi}^Q \in \mathcal{M}_{QL}(\eta, \chi)$. Define $\hat{\pi}^{Q,J}$ by

$$
\hat{\pi}^{Q,J}(du, dx, dy) = I_{\{u \in J\}} \hat{\pi}^Q(du, dx, dy)
$$

Then it is easy to see that $\hat{\pi}^{Q,J}$ is a J-quantile-lifted martingale coupling of η^J and $\chi^{\hat{\pi}^{Q,J}}$ where $\chi^{\hat{\pi}^{Q},J}(dy) = \int_{u} \int_{x} \hat{\pi}^{Q,J}(du, dx, dy)$. Note that $\chi^{\hat{\pi}^{Q},J}$ is such that $\eta^{J} \leq_{cx} \chi^{\hat{\pi}^{Q},J} \leq \chi$.

The following lemma allows us to translate between J-quantile lifts of sub-measures and quantile lifts of measures. For $J \subseteq (0,\infty)$ let G_J be the left-continuous inverse of the map $u \mapsto \int_0^u I_{\{v \in J\}} dv$.

Lemma 2.7. 1. Suppose $\chi \in \mathcal{P}$ and $J \subseteq (0, \chi(\mathbb{R}))$. Then $^J \hat{\chi}^Q$ given by $^J \hat{\chi}^Q(du, dx) =$ $du\delta_{G_\chi(G_J(u))}(dx)$ is a quantile lift of χ^J .

2. For $\eta, \zeta \in \mathcal{P}$ with $\eta \leq_{cx} \zeta$ let $\hat{\pi}^Q \in \mathcal{M}_{QL}(\eta, \zeta)$. Suppose $J \subseteq (0, \eta(\mathbb{R}))$. Then $J\hat{\pi}^Q$ given by

$$
J_{\hat{\pi}}^Q(du, dx, dy) = du \delta_{G_{\eta}(G_J(u))}(dx) \hat{\pi}^Q_{G_J(u), G_{\eta}(G_J(u))}(dy)
$$

is a quantile-lifted martingale coupling of η^J and $\zeta^{\hat{\pi}^{Q},J}$ where $\zeta^{\hat{\pi}^{Q},J}(dy) := \int_u \int_x \hat{\pi}^{Q,J}(du, dx, dy)$ is such that $\eta^J \leq_{cx} \zeta^{\hat{\pi}^Q, J} \leq \zeta$.

Proof. For 1., it is easy to check that $G_{\chi}(u) = G_{\chi}(G_{J}(u))$ and hence that ${}^{J}\hat{\chi}^{Q}$ has support $(v, G_{\chi^J}(v))_{0 \le v \le Leb(J)}$. Further, $\int_u^J \hat{\chi}^Q(du, dx) = \int_u \delta_{G\chi(G_J(u))}(dx)$ which after the change of variable $v = G_J(u)$ (note that $du = I_{\{v \in J\}} dv$) becomes $\int_u^J \hat{\chi}^Q(du, dx) =$ $\int_v I_{\{v \in J\}} \delta_{G\chi(v)}(dx) = \chi^J(dx).$

For 2., we use the same transformation: $\int_u \int_y^J \hat{\pi}^Q(du, dx, dy) = \int_u du \delta_{G_\eta(G_J(u))}(dx)$ $\int_{\nu} I_{\{v \in J\}} \delta_{G_{\eta}(v)}(dx) = \eta^{J}(dx)$ and $\int_{u} \int_{x}^{J} \hat{\pi}^{Q}(du, dx, dy) = \int_{u} \int_{x} du \delta_{G_{\eta}(G_{J}(u))}(dx) \hat{\pi}^{Q}_{G_{J}(u), G_{\eta}(G_{J}(u))}(dy) =$ $\int_v \int_x dv I_{\{v \in J\}} \delta_{G\chi(v)}(dx) \hat{\pi}_{v,\mathbf{v}}^Q$ $\mathcal{L}_{v,G_\eta(v)}^Q(dy) = \int_v \int_x \hat{\pi}_{v,x}^{Q,J}(dv,dx,dy) = \zeta^{\hat{\pi}^{Q},J}(dy).$

The idea behind introducing J-quantile-lifted martingale couplings is that it allows for superposition.

Lemma 2.8. Suppose $\mu \in \mathcal{P}$. Suppose $(J_k)_{k>1}$ is a countable partition of J where $J\subseteq (0,\mu(\mathbb{R})).$

- (i) Suppose $\hat{\mu}^{Q,J_k}$ is a J_k -quantile lift of $\mu_k := \mu^{J_k}$. Then $\sum_k \hat{\mu}^{Q,J_k}$ is a J-quantile lift of μ^J .
- (ii) Further, for $(\mu_k)_{k>1}$ as above and for $(\nu_k)_{k>1}$ such that $\mu_k \leq_{cx} \nu_k$, suppose that for each $k \geq 1$ there exists a J_k -quantile-lifted martingale coupling of μ_k and ν_k . Then there exists a J-quantile-lifted martingale coupling of μ^{J} and $\sum_{k\geq 1}\nu_{k}$.

Proof. For (i): clearly, $\sum_k \hat{\mu}^{Q,J_k}$ is a measure on $(0, \mu(\mathbb{R})) \times I_\mu$ with support $\cup_k (u, G_\mu(u))_{u \in J_k} =$ $(u, G_\mu(u))_{u \in \cup_k J_k}$ and

$$
\sum_{k} \hat{\mu}^{Q,J_k}(du, dx) = \sum_{k} I_{\{u \in J_k\}} du \delta_{G_{\mu}(u)}(dx) = I_{\{u \in J\}} du \delta_{G_{\mu}(u)}(dx).
$$

For (ii): if $\hat{\pi}^{Q,J_k}$ given by $\hat{\pi}^{Q,J_k}(du, dx, dy) = I_{\{u \in J_k\}} du \delta_{G_{\mu_k}(u)}(dx) \hat{\pi}^{Q,J_k}_{u,x}(dy)$ is a J_k quantile-lifted martingale coupling of μ_k and ν_k then

$$
\sum_{k} \hat{\pi}^{Q,J_k}(du, dx, dy) = \sum_{k} I_{\{u \in J_k\}} du \delta_{G_{\mu_k}(u)}(dx) \hat{\pi}^{Q,J_k}_{u,x}(dy) = I_{\{u \in J\}} du \delta_{G_{\mu}(u)}(dx) \hat{\pi}^{Q}_{u,x}(dy)
$$

where $\hat{\pi}_{u,x}^Q(dy) = \hat{\pi}_{u,x}^{Q,J_k}$ for $u \in J_k$. It follows that $\sum_k \hat{\pi}^{Q,J_k}$ is a J-quantile-lifted martingale coupling of μ^J and $\sum_{k\geq 1} \nu_k$. \Box

Proposition 2.9. Suppose μ , $(J_k)_{k\geq 1}$, $(\mu_k)_{k\geq 1}$ and $(\nu_k)_{k\geq 1}$ are as in Lemma [2.8](#page-8-0) and suppose $J = (0, \mu(\mathbb{R}))$ and $\nu_k \wedge \nu_{k'} = 0$ for all $k \neq k'$. Suppose that for each k there exists an injective J_k -quantile-lifted martingale coupling of μ_k and ν_k . Then there exist an injective quantile-lifted martingale coupling of μ and $\nu = \sum_{k\geq 1} \nu_k$.

Proof. Suppose $\hat{\pi}_k^Q$ κ_k^Q is an injective J_k -quantile-lifted martingale coupling of μ_k and ν_k . Then, by the arguments in Lemma [2.8,](#page-8-0) $\sum_{k\geq 1} \hat{\pi}_k^Q \in \mathcal{M}_{QL}(\mu, \nu)$.

Further, by the fact that $\nu_k \wedge \nu_{k'} = 0$, v-almost every y lies in the support of at most one set supp (ν_k) . Since each $\hat{\pi}_k^Q \in \mathcal{IM}_{QL}(\mu_k, \nu_k)$, such a y can have at most one pre-image under $\sum_{k\geq 1} \hat{\pi}^Q_k$ $\frac{Q}{k}$. \Box

3 A building block for injective martingale couplings

In the light of Proposition [2.4,](#page-6-0) from now on we assume that $\mu \leq_{cx} \nu$ are such that ${x : D_{\mu,\nu}(x) > 0}$ is an interval and is equal to $I_{\nu} = (\alpha_{\nu}, \beta_{\nu})$ where $\alpha_{\nu} < \beta_{\nu}$. We will also assume that ν is continuous.

Standing Assumption 3.1. ν is atom-free: $\nu({x}) = 0$ for all $x \in \mathbb{R}$.

It follows from Standing Assumption [3.1](#page-9-1) that $D'_{\mu,\nu}(\alpha_\nu) = 0 = D'_{\mu,\nu}(\beta_\nu)$ whenever $-\infty < \alpha_{\nu} < \beta_{\nu} < \infty$, or more generally, $\lim_{k \downarrow \alpha_{\nu}} D'_{\mu,\nu}(k+) = 0 = \lim_{k \uparrow \beta_{\nu}} D'_{\mu,\nu}(k-)$.

Let G_{μ} be an arbitrary quantile function of μ and, for $u \in [0, \mu(\mathbb{R})]$, recall the definition of μ_u (see Section [2.2\)](#page-4-0). For each $0 \le v \le u \le \mu(\mathbb{R})$ define $\mathcal{E}_{v,u}^{\mu,\nu} = \mathcal{E}_{u,v} : \mathbb{R} \mapsto$ \mathbb{R}_+ by

$$
\mathcal{E}_{v,u}(k) = \begin{cases}\nD_{\mu,\nu}(k) + C_{\mu_v}(k), & k \le G_{\mu}(v) \\
D_{\mu,\nu}(k), & G_{\mu}(v) < k < G_{\mu}(u) \\
P_{\nu}(k) - P_{\mu_u}(k), & k \ge G_{\mu}(u).\n\end{cases}
$$
\n(2)

Note that the definition of $\mathcal{E}_{v,u}$ is independent of the choice of G_{μ} . By design $\mathcal{E}_{v,u}$ is continuous, including at $G_{\mu}(v)$ and $G_{\mu}(u)$. Note that on $k \leq G_{\mu}(v)$ we have that $\mathcal{E}_{v,u}$ does not depend on u; similarly, on $k \geq G_{\mu}(u)$, $\mathcal{E}_{v,u}$ does not depend on v.

Recall that, for measures $\eta \leq_{pcx} \nu$, $S^{\nu}(\eta)$ denotes the shadow of η in ν . Note that if $\eta \leq \mu \leq_{cx} \nu$ then $\eta \leq_{pcx} \nu$. The next lemma describes the shadow of $\mu_u - \mu_v$ in terms of $\mathcal{E}_{u,v}$ and its convex hull.

Lemma 3.2. Suppose $\mu \leq_{cx} \nu$ and let $0 \leq v \leq u \leq \mu(\mathbb{R})$. Then

$$
P_{\nu-S^{\nu}(\mu_u-\mu_v)}(k)=\mathcal{E}_{v,u}^c(k)-(\overline{\mu_v}-vk), \quad k \in \mathbb{R}.
$$

In particular, the second (distributional) derivative of $\mathcal{E}_{v,u}^c$ corresponds to the measure $\nu-S^{\nu}(\mu_u-\mu_v).$

Proof. Let $l_v : \mathbb{R} \to \mathbb{R}$ be given by $l_v(k) = \overline{\mu_v} - vk, k \in \mathbb{R}$. By Beiglböck et al. [\[1,](#page-38-5) Theorem 4.7],

$$
P_{\nu-S^{\nu}(\mu_u-\mu_v)}=P_{\nu}-P_{S^{\nu}(\mu_u-\mu_v)}=(P_{\nu}-P_{\mu_u-\mu_v})^c,
$$

and therefore it is enough to show that $\mathcal{E}_{v,u}^c - l_v = (P_\nu - P_{\mu_u - \mu_v})^c$. On the other hand, by Beiglböck et al. [\[1,](#page-38-5) Lemma 2.4] and linearity of l_v we have that

$$
(\mathcal{E}_{v,u} - l_v)^c = (\mathcal{E}_{v,u}^c - l_v)^c = \mathcal{E}_{v,u}^c - l_v.
$$

We now show that $(\mathcal{E}_{v,u} - l_v) = (P_\nu - (P_{\mu_u} - P_{\mu_v})) = (P_\nu - P_{\mu_u - \mu_v})$. Suppose $k \leq G_{\mu}(v)$. Then

$$
\mathcal{E}_{v,u}(k) = D_{\mu,\nu}(k) + C_{\mu_v}(k) = D_{\mu,\nu}(k) + P_{\mu_v}(k) + l_v(k)
$$

= $D_{\mu,\nu}(k) + P_{\mu}(k) + l_v(k)$
= $P_{\nu}(k) + l_v(k)$
= $P_{\nu}(k) - (P_{\mu_u}(k) - P_{\mu_v}(k)) + l_v(k),$

where we used that $(P_{\mu_u} - P_{\mu_v}) = 0$ on $(-\infty, G_{\mu}(v))$. Now suppose $G_{\mu}(v) < k <$ $G_{\mu}(u)$. Then

$$
(P_{\nu}(k) - (P_{\mu_u}(k) - P_{\mu_v}(k)) = (P_{\nu}(k) - (P_{\mu}(k) - P_{\mu_v}(k))
$$

= $D_{\mu,\nu}(k) + P_{\mu_v}(k) = D_{\mu,\nu}(k) - l_v(k) = \mathcal{E}_{v,u} - l_v(k),$

as required. Finally, if $k \geq G_{\mu}(u)$, then

$$
\mathcal{E}_{v,u}(k) = P_{\nu}(k) - P_{\mu_u}(k)
$$

= $(P_{\nu}(k) - (P_{\mu_u}(k) - P_{\mu_v}(k)) - P_{\mu_v}(k) = (P_{\nu}(k) - (P_{\mu_u}(k) - P_{\mu_v}(k)) + l_v(k)).$

Corollary 3.3. Fix $0 \le v \le u \le \mu(\mathbb{R})$. Then $\mathcal{E}_{v,u}^c$ is continuously differentiable. Moreover, for any $k \in \mathbb{R}$ with $\mathcal{E}_{v,u}(k) = \mathcal{E}_{v,u}^c(k)$, we have that $\mathcal{E}_{v,u}'(k) = (\mathcal{E}_{v,u}^c)'(k)$.

Proof. Since $(\nu - S^{\nu}(\mu_u - \mu_v)) \leq \nu$, $(\nu - S^{\nu}(\mu_u - \mu_v))$ is also continuous, and therefore $P_{\nu-S^{\nu}(\mu_u-\mu_v)}$ is continuously differentiable. Then Lemma [3.2](#page-9-2) implies the desired differentiability of $\mathcal{E}_{v,u}^c$.

Now suppose that $k \in \mathbb{R}$ is such that $\mathcal{E}_{v,u}(k) = \mathcal{E}_{v,u}^c(k)$. Since $\mathcal{E}_{v,u} \geq \mathcal{E}_{v,u}^c$ on \mathbb{R} , $\mathcal{E}'_{v,u}(k-) \leq (\mathcal{E}_{v,u}^c)'(k) \leq \mathcal{E}'_{v,u}(k+)$. But if one (or more) of the inequalities is strict, then the definition of $\mathcal{E}_{v,u}$ implies that $\nu({k}) > 0$, a contradiction. □

Corollary 3.4. $\mathcal{E}_{v,u}^c$ is linear on $[X_{\mathcal{E}_v}^-]$ $[\varepsilon_{v,u}(z), Z^+_{\mathcal{E}_{v,u}}(z)]$

Proof. Since, by Corollary [3.3,](#page-10-0) $(\mathcal{E}_{v,u}^c)'(z-) = (\mathcal{E}_{v,u}^c)'(z+)$ for each $z \in \mathbb{R}$, the claim follows immediately from the definitions of $X_{\mathcal{E}_{\alpha}}^ \bar{\varepsilon}_{v,u}(z)$ and $Z_{\mathcal{E}_u}^+$ $\vec{\varepsilon}_{v,u}(z).$ \Box

Our ultimate goal is to define a pair of functions $(M, N) = (M(u), N(u))_{0 \le u \le u(\mathbb{R})}$ such that (M, N) define an injective lifted martingale coupling. See Figure [1.](#page-11-0) We will do this by defining (M, N) on a sequence of domains, whose union is $(0, \mu(\mathbb{R}))$. In this section we will give some preliminary results which are applicable to a single domain.

We begin by defining a family of functions $\overrightarrow{m}_{.,\cdot,\cdot}$, which will later be used to define (M, N) . See Figure [1.](#page-11-0) Recall that \overrightarrow{G}_{μ} is the left-continuous quantile function of μ .

Figure 1: The construction of $\overrightarrow{m}_{v,u}$ and $\overrightarrow{n}_{v,u}$. For $v < u < u_1 < u_2$, the dotted curve represents \mathcal{E}_{v,u_2} , the dashed curve corresponds to \mathcal{E}_{v,u_1} , while the dash-dotted curve represents $\mathcal{E}_{v,u}$. The solid curve corresponds to $D_{\mu,\nu}$. Note that $D_{\mu,\nu} \leq \mathcal{E}_{v,u} \leq$ $\mathcal{E}_{v,u_1} \leq \mathcal{E}_{v,u_2}$ everywhere, $\mathcal{E}_{v,w} = D_{\mu,\nu}$ on $[\overrightarrow{G}_{\mu}(v), \overrightarrow{G}_{\mu}(w)]$ for $w \in \{u, u_1, u_2\}$, and $\mathcal{E}_{v,u_i} = \mathcal{E}_{v,u}$ on $(-\infty, \overrightarrow{G}_{\mu}(v))$ for $i = 1, 2$. Furthermore, for $i = 1, 2$, the straight line going through $(\overrightarrow{m}_{v,u}(u_i), \mathcal{E}_{v,u_i}(\overrightarrow{m}_{v,u}(u_i)))$ and $(\overrightarrow{n}_{v,u}(u_i), \mathcal{E}_{v,u_i}(\overrightarrow{n}_{v,u}(u_i)))$ corresponds to the linear section on $[\overrightarrow{m}_{v,u}(u_i), \overrightarrow{n}_{v,u}(u_i)]$ of the convex hull \mathcal{E}_{v,u_i}^c of \mathcal{E}_{v,u_i} . In particular, $\vec{m}_{v,u}(u_2) \leq \vec{m}_{v,u}(u_1) \leq \vec{n}_{v,u}(u_1) \leq \vec{n}_{v,u}(u_2).$

Definition 3.5. Fix $v, u \in [0, \mu(\mathbb{R})]$ with $v \leq u$. Define $\overrightarrow{m}_{v,u}, \overrightarrow{n}_{v,u} : [u, \mu(\mathbb{R})] \to \mathbb{R}$ by

$$
\overrightarrow{m}_{v,u}(l) = X_{\mathcal{E}_{v,l}}(\overrightarrow{G}_{\mu}(l)) \quad \text{and} \quad \overrightarrow{n}_{v,u}(l) = Z_{\mathcal{E}_{v,l}}^+(\overrightarrow{G}_{\mu}(l)), \quad l \in [u, \mu(\mathbb{R})].
$$

Lemma 3.6. Fix $0 \le v \le u \le l \le \mu(\mathbb{R})$. Then, for all $w \in [l, \mu(\mathbb{R})]$, $S^{\nu}(\mu_w - \mu_v) = \nu$ on $(\vec{m}_{v,u}(l), \vec{n}_{v,u}(l))$ and $\mathcal{E}_{v,w}^c$ is linear on $[\vec{m}_{v,u}(l), \vec{n}_{v,u}(l)].$

Proof. If $\overrightarrow{m}_{v,u}(l) = \overrightarrow{n}_{v,u}(l)$ then there is nothing to prove so we may assume $\overrightarrow{m}_{v,u}(l)$ < $\overrightarrow{n}_{v,u}(l)$. By the associativity of the shadow measure (see Beiglböck and Juillet [\[2,](#page-38-1) Theorem 4.8]), for $v \le u \le l \le w$, $S^{\nu}(\mu_w - \mu_v) = S^{\nu}(\mu_l - \mu_v) + S^{\nu - S^{\nu}(\mu_l - \mu_v)}(\mu_w - \mu_l)$ and therefore $S^{\nu}(\mu_l - \mu_v) \leq S^{\nu}(\mu_w - \mu_v) \leq \nu$. Hence,

$$
\nu - S^{\nu}(\mu_w - \mu_v) \leq \nu - S^{\nu}(\mu_l - \mu_v).
$$

By Corollary [3.4,](#page-10-1) $\mathcal{E}_{v,l}^c$ is linear on $[\vec{m}_{v,u}(l), \vec{n}_{v,u}(l)]$, and then, see Lemma [3.2,](#page-9-2) ν – $S^{\nu}(\mu_l - \mu_v)$ (and thus also $\nu - S^{\nu}(\mu_w - \mu_v)$) does not charge $(\vec{m}_{v,u}(l), \vec{n}_{v,u}(l))$. Hence, by Lemma [3.2](#page-9-2) again, $\mathcal{E}_{v,w}^c$ is linear on $[\overrightarrow{m}_{v,u}(l), \overrightarrow{n}_{v,u}(l)]$. Moreover, since $(\nu - S^{\nu}(\mu_w - \mu_w))$ $(\mu_v)(\overrightarrow{(m}_{v,u}(l), \overrightarrow{n}_{v,u}(l))) = 0$ it follows that $S^{\nu}(\mu_w - \mu_v)(\overrightarrow{(m}_{v,u}(l), \overrightarrow{n}_{v,u}(l))) = \nu((\overrightarrow{m}_{v,u}(l), \overrightarrow{n}_{v,u}(l))).$ Then, since $S^{\nu}(\mu_w - \mu_v) \leq \nu$ we conclude that $S^{\nu}(\mu_w - \mu_v) = \nu$ on $(\overrightarrow{m}_{v,u}(l), \overrightarrow{n}_{v,u}(l))$.

Corollary 3.7. Fix $0 \le v \le u \le l \le \mu(\mathbb{R})$. If $l \in (0, \mu(\mathbb{R}))$ then $\alpha_{\nu} < \overrightarrow{m}_{v,u}(l) \le$ $\overrightarrow{n}_{v,u}(l) < \beta_{v}$.

Proof. Note that $\overrightarrow{m}_{v,u}(l) \leq \overrightarrow{G}_{\mu}(l) \leq \overrightarrow{n}_{v,u}(l)$ and $\alpha_{\nu} \leq \alpha_{\mu} \leq \overrightarrow{G}_{\mu}(l) \leq \underline{\beta}_{\nu} \leq \beta_{\nu}$. Further, by the continuity of ν , if $\alpha_{\mu} = \alpha_{\nu}$ then $\overrightarrow{G}_{\mu}(l) > \alpha_{\mu} = \alpha_{\nu}$; otherwise $\overrightarrow{G}_{\mu}(l) \geq \alpha_{\mu} > \alpha_{\nu}$. Similarly, $\overrightarrow{G}_{\mu}(l) < \beta_{\nu}$. Hence, $\overrightarrow{m}_{v,u}(l) < \beta_{\nu}$ and $\alpha_{\nu} < \overrightarrow{n}_{v,u}(l)$.

Observe that if $\vec{m}_{v,u}(l) \leq \alpha_{\nu}$ and $\beta_{\nu} \leq \vec{n}_{v,u}(l)$ then by Lemma [3.6,](#page-11-1) $0 = \nu(\mathbb{R})$ – $S^{\nu}(\mu_l - \mu_v)(\mathbb{R}) = \mu(\mathbb{R}) - (\mu_l(\mathbb{R}) - \mu_v(\mathbb{R})) = v + \mu(\mathbb{R}) - l > v \geq 0$, a contradiction. Hence, in order to prove that $\alpha_{\nu} < \overrightarrow{m}_{v,u}(l) \leq \overrightarrow{n}_{v,u}(l) < \beta_{\nu}$ it is sufficient to show that neither $\alpha_{\nu} < \overrightarrow{m}_{v,u}(l) < \beta_{\nu} \leq \overrightarrow{n}_{v,u}(l)$ nor $\overrightarrow{m}_{v,u}(l) \leq \alpha_{\nu} < \overrightarrow{n}_{v,u}(l) < \beta_{\nu}$.

Suppose $\alpha_{\nu} < \overrightarrow{m}_{v,u}(l) < \beta_{\nu} \le \overrightarrow{n}_{v,u}(l)$. Then by Lemma [3.6](#page-11-1) and the continuity of ν we have that $\nu - S^{\nu}(\mu_l - \mu_v)$ is concentrated on $(\alpha_{\nu}, \overrightarrow{m}_{v,u}(l)) \subseteq (\alpha_{\nu}, \overrightarrow{G}_{\mu}(l))$. On the other hand, by the associativity of the shadow measure $\mu_v + (\mu - \mu_l) = \mu - (\mu_l - \mu_v) \leq_{cx}$ $\nu - S^{\nu}(\mu_l - \mu_v)$. But, since $l \in (0, \mu(\mathbb{R}))$, $(\mu - \mu_l)(\mathbb{R}) > 0$ and $\mu - \mu_l$ is concentrated on $[\vec{G}_{\mu}(l), \infty)$. It follows that $\mu - \mu_l$ cannot be embedded in $\nu - S^{\nu}(\mu_l - \mu_v)$ in a way which respects the martingale property. A fortiori, $\mu_v + (\mu - \mu_l)$ cannot be embedded in $\nu - S^{\nu}(\mu_l - \mu_v)$ in a way which respects the martingale property, a contradiction to the fact that $\mu - (\mu_l - \mu_v) \leq_{cx} \nu - S^{\nu}(\mu_l - \mu_v)$. We conclude that $\overrightarrow{n}_{v,u}(l) < \beta_{\nu}$ whenever $\alpha_{\nu} < \overrightarrow{m}_{v,u}(l).$

Now suppose that $\vec{m}_{v,u}(l) \le \alpha_v < \vec{n}_{v,u}(l) < \beta_v$. Then, by Lemma [3.6,](#page-11-1) $\nu S^{\nu}(\mu_l - \mu_v)$ is concentrated on $(\vec{n}_{v,u}(l), \beta_{\nu})$. Suppose that $v > 0$. Then $\mu_v(\mathbb{R}) = v > 0$ and μ_v is concentrated on $(-\infty, \overrightarrow{G}_{\mu}(v)] \subseteq (-\infty, \overrightarrow{n}_{v,u}(l)]$. In this case μ_v cannot be embedded in $\nu-S^{\nu}(\mu_l-\mu_v)$ in a martingale way, a contradiction. Finally suppose that $v=0$, so that $\mu - (\mu_l - \mu_v) = \mu - \mu_l$ and $\mathcal{E}_{v,l} = \mathcal{E}_{0,l} = D$ on $(-\infty, \overrightarrow{G}_{\mu}(l+)]$. Then, since $\overrightarrow{m}_{v,u} \leq \alpha_{\nu}$, we have that $\mathcal{E}_{0,l}^c(\overrightarrow{m}_{v,u}) = D(\overrightarrow{m}_{v,u}) = 0$ (or $\lim_{k \downarrow \overrightarrow{m}_{v,u}} \mathcal{E}_{0,l}^c(k) = \lim_{k \downarrow \overrightarrow{m}_{v,u}} D(k) = 0$ in the case $\vec{m}_{v,u} = -\infty$) and $(\mathcal{E}_{0,l}^c)'(\vec{m}_{v,u}) = D'(\vec{m}_{v,u}) = 0$ (or $(\mathcal{E}_{0,l}^c)'(\vec{m}_{v,u}+) = 0$ in the case

 $\overrightarrow{m}_{v,u} = -\infty$). But then $D(\overrightarrow{n}_{v,u}(l)) = \mathcal{E}_{0,l}(\overrightarrow{n}_{v,u}(l)) = \mathcal{E}_{0,l}(\overrightarrow{n}_{v,u}(l)) = 0$, a contradiction since $\overrightarrow{n}_{v,u}(l) < \beta_{\nu}$ and $D_{\mu,\nu} > 0$ on $(\alpha_{\nu}, \beta_{\nu})$.

 \Box

Corollary 3.8. Fix $v, u \in [0, \mu(\mathbb{R}))$ with $v \leq u$. $\overrightarrow{n}_{v,u}(\cdot)$ is non-decreasing on $[u, \mu(\mathbb{R})]$.

Proof. Note that since $\overrightarrow{n}_{v,u}$ only depends on u via the domain on which it is defined, it is enough to prove the claim for $u = v$. Let $v = u \leq w \leq \mu(\mathbb{R})$. Either $\overrightarrow{G}_{\mu}(w) \geq \overrightarrow{n}_{v,v}(v)$ and then $\overrightarrow{n}_{v,v}(v) \leq \overrightarrow{G}_{\mu}(w) \leq \overrightarrow{n}_{v,v}(w)$ or $\overrightarrow{G}_{\mu}(w) < \overrightarrow{n}_{v,v}(v)$ and then $\overrightarrow{m}_{v,v}(v) \leq$ $\overrightarrow{G}_{\mu}(v) \leq \overrightarrow{G}_{\mu}(w) < \overrightarrow{\mathcal{D}}_{v,v}(v)$. In the latter case, since $[\overrightarrow{m}_{v,v}(w), \overrightarrow{n}_{v,v}(w)]$ is the largest interval containing $\vec{G}_{\mu}(w)$ on which $\mathcal{E}_{v,w}^c$ is linear, and since (by Lemma [3.6\)](#page-11-1) $\mathcal{E}_{v,w}^c$ is linear on $[\overrightarrow{m}_{v,v}(v), \overrightarrow{n}_{v,v}(v)]$ we must have $\overrightarrow{m}_{v,v}(w) \leq \overrightarrow{m}_{v,v}(v) \leq \overrightarrow{n}_{v,v}(v) \leq \overrightarrow{n}_{v,v}(w)$. Hence in both cases we have $\overrightarrow{n}_{v,v}(v) \leq \overrightarrow{n}_{v,v}(w)$ so that $\overrightarrow{n}_{v,v}$ is non-decreasing. \Box

Lemma 3.9. Let $v, u, r_1, r_2 \in [0, \mu(\mathbb{R})]$ be such that $v \leq u \leq r_1 \leq r_2$. If $\overrightarrow{m}_{v,u}(r_2) \leq$ $\overrightarrow{n}_{v,u}(r_1)$ then $\overrightarrow{m}_{v,u}(r_2) \leq \overrightarrow{m}_{v,u}(r_1)$.

Proof. If $r_1 = r_2$ then there is nothing to prove. So suppose $r_1 < r_2$.

If $\vec{m}_{v,u}(r_1) = \vec{n}_{v,u}(r_1)$ then, by hypothesis, $\vec{m}_{v,u}(r_2) \leq \vec{n}_{v,u}(r_1) = \vec{m}_{v,u}(r_1)$ and we are done. Hence in the rest of the proof we suppose that $\vec{m}_{v,u}(r_1) < \vec{n}_{v,u}(r_1)$.

If $\overrightarrow{m}_{v,u}(r_2) = \overrightarrow{G}_{\mu}(r_2)$ then either $\overrightarrow{m}_{v,u}(r_2) = \overrightarrow{G}_{\mu}(r_2) < \overrightarrow{m}_{v,u}(r_1)$ (and there is nothing to prove) or $\vec{m}_{v,u}(r_1) \leq \vec{G}_{\mu}(r_2) = \vec{m}_{v,u}(r_2) \leq \vec{n}_{v,u}(r_1)$. In the latter case, since $[\vec{m}_{v,u}(r_2), \vec{n}_{v,u}(r_2)]$ is the largest interval (containing $\vec{G}_{\mu}(r_2)$) on which \mathcal{E}_{v,r_2}^c is linear (note that, by Lemma [3.6,](#page-11-1) \mathcal{E}_{v,r_2}^c is linear on $[\vec{m}_{v,u}(r_1), \vec{n}_{v,u}(r_1)]$), we conclude that $\overrightarrow{m}_{v,u}(r_2) \leq \overrightarrow{m}_{v,u}(r_1)$. Hence in the rest of the proof we suppose that $\overrightarrow{m}_{v,u}(r_2) < \overrightarrow{G}_{\mu}(r_2)$, so that $\overrightarrow{m}_{v,u}(r_2) < \overrightarrow{n}_{v,u}(r_2)$.

We now have that \mathcal{E}_{v,r_2}^c is linear on both $[\overrightarrow{m}_{v,u}(r_1), \overrightarrow{n}_{v,u}(r_1)]$ and $[\overrightarrow{m}_{v,u}(r_2), \overrightarrow{n}_{v,u}(r_2)]$ where (by Corollary [3.8\)](#page-13-0) $\overrightarrow{n}_{v,u}(r_1) \leq \overrightarrow{n}_{v,u}(r_2)$. Since $\overrightarrow{m}_{v,u}(r_2) \leq \overrightarrow{n}_{v,u}(r_1)$ these intervals either overlap or meet at a point. In the latter case, since ν is atom-less, the slopes of both linear sections of \mathcal{E}_{v,r_2}^c must be the same at $\vec{m}_{v,u}(r_2) \equiv \vec{n}_{v,u}(r_1)$ (i.e., \mathcal{E}_{v,r_2}^c cannot have kink at $\vec{m}_{v,u}(r_2)$). Therefore in both cases \mathcal{E}_{v,r_2}^c is linear on $[\vec{m}_{v,u}(r_1), \vec{n}_{v,u}(r_2)]$. But $G_{\mu}(r_2) \in [\vec{m}_{v,u}(r_1), \vec{n}_{v,u}(r_2)]$, and therefore we must have that $\overrightarrow{m}_{v,u}(r_2) \leq \overrightarrow{m}_{v,u}(r_1)$. \Box

For $v, u \in [0, \mu(\mathbb{R}))$ with $v \leq u$ define

$$
\overline{w}_{v,u} := \inf\{w \in (u, \mu(\mathbb{R})): \overrightarrow{n}_{v,u}(w) \leq \overrightarrow{G}_{\mu}(w+) \}
$$

with convention inf $\emptyset = \mu(\mathbb{R})$.

Lemma 3.10. 1. Suppose $0 \le v \le u < \mu(\mathbb{R})$. If $\overrightarrow{n}_{v,u}(u) > \overrightarrow{G}_{\mu}(u+)$ then $\overline{w}_{v,u} > u$.

2. Suppose $0 \le v \le u < \overline{w}_{v,u} \le \mu(\mathbb{R})$. Then $\overrightarrow{G}_{\mu}(\overline{w}_{v,u}) \le \overrightarrow{n}_{v,u}(\overline{w}_{v,u}) \le \overrightarrow{G}_{\mu}(\overline{w}_{v,u}+)$. and $D(\overrightarrow{n}_{v,u}(\overline{w}_{v,u})) = \mathcal{E}_{v,u}(\overrightarrow{n}_{v,u}(\overline{w}_{v,u})) = \mathcal{E}_{v,u}^c(\overrightarrow{n}_{v,u}(\overline{w}_{v,u})).$

Proof. Since $\overrightarrow{G}_{\mu}(u+) < \overrightarrow{n}_{v,u}(u)$, by the right-continuity of $\overrightarrow{G}_{\mu}(\cdot+)$ there exists $\overline{\epsilon} > 0$ such that for all $\epsilon \in (0, \overline{\epsilon}]$

$$
\overrightarrow{G}_{\mu}(u+) \leq \overrightarrow{G}_{\mu}(u+\epsilon) \leq \overrightarrow{G}_{\mu}((u+\epsilon)+) < \overrightarrow{n}_{v,u}(u).
$$

Then for $\epsilon \in [0,\bar{\epsilon}]$ we have $\overrightarrow{G}_{\mu}((u+\epsilon)+)<\overrightarrow{n}_{v,u}(u)<\overrightarrow{n}_{v,u}(u+\epsilon)$ and therefore $\overline{w}_{v,u} \geq u + \epsilon > u.$

For the second part, suppose $0 \le v \le u < \overline{w}_{v,u} \le \mu(\mathbb{R})$. That $\overrightarrow{G}_{\mu}(\overline{w}_{v,u}) \le$ $\overrightarrow{n}_{v,u}(\overline{w}_{v,u})$ is clear from the definition of $\overrightarrow{n}_{v,u}$, and thus we now prove that $\overrightarrow{n}_{v,u}(\overline{w}_{v,u}) \leq$ $\vec{G}_{\mu}(\overline{w}_{v,u}+)$. This is immediate if $\overline{w}_{v,u} = \mu(\mathbb{R})$ since we have defined $G_{\mu}(\mu(\mathbb{R})+) = \infty$ for any quantile function G_{μ} of μ . Suppose $\overline{w}_{v,u} < \mu(\mathbb{R})$, so that $\{w \in (u, \mu(\mathbb{R})]:$ $\overrightarrow{n}_{v,u}(w) \leq \overrightarrow{G}_{\mu}(w+)\}\neq \emptyset$. Either $\overrightarrow{n}_{v,u}(\overline{w}_{v,u}) \leq \overrightarrow{G}_{\mu}(\overline{w}_{v,u}+)$ and we are done, or there exists $(w_k)_{k\geq 1}$ in $(u, \mu(\mathbb{R})]$ such that $w_k \downarrow \overline{w}_{v,u}$ (as $k \uparrow \infty$) and $\overrightarrow{n}_{v,u}(w_k) \leq \overrightarrow{G}_{\mu}(w_k)$ for each $k \ge 1$. Then, by Corollary [3.8,](#page-13-0) $\overrightarrow{n}_{v,u}(\cdot)$ is monotonic increasing on $[u, \mu(\mathbb{R})]$ and we have that

$$
\overrightarrow{n}_{v,u}(\overline{w}_{v,u}) \leq \lim_{k \uparrow \infty} \overrightarrow{n}_{v,u}(w_k) \leq \lim_{k \uparrow \infty} \overrightarrow{G}_{\mu}(w_k+) = \overrightarrow{G}_{\mu}(\overline{w}_{v,u}+).
$$

 \Box

We have shown that $\overrightarrow{n}_{v,u}$ is increasing on $[u, \mu(\mathbb{R})]$ and now we would like to show that $\vec{m}_{v,u}$ is decreasing. This is not true on $[u, \mu(\mathbb{R})]$; however, we will show that it is true on $[u,\overline{w}_{v,u}].$

Lemma 3.11. Fix $v, u \in [0, \mu(\mathbb{R}))$ with $v \leq u$. Then $\overrightarrow{m}_{v,u}(\cdot)$ is non-increasing on $(u,\overline{w}_{v,u}).$

Proof. If $u = \overline{w}_{v,u}$ there is nothing to prove. So, suppose $u < \overline{w}_{v,u}$.

Fix $l, w \in (u, \overline{w}_{v,u}]$ with $l < w$. We show that $\overrightarrow{m}_{v,u}(l) \leq \overrightarrow{m}_{v,u}(w)$.

Suppose that $\overrightarrow{m}_{v,u}(w) \leq \overrightarrow{n}_{v,u}(l)$. Then by Lemma [3.9](#page-13-1) with $r_1 = l$ and $r_2 = w$, we have $\overrightarrow{m}_{v,u}(w) \leq \overrightarrow{m}_{v,u}(l)$ as required.

The alternative is that $\overrightarrow{n}_{v,u}(l) < \overrightarrow{m}_{v,u}(w)$. We show that this case cannot happen by finding a contradiction. Define $I_{v,u,w} = \{k \in (l, w] : \overrightarrow{n}_{v,u}(l) < \overrightarrow{m}_{v,u}(k)\}\.$ Clearly, since $\overrightarrow{n}_{v,u}(l) < \overrightarrow{m}_{v,u}(w)$ we have $w \in I_{v,u,w}$. Define $\tilde{l} := \inf\{k : k \in I_{v,u,w}\}$. We show first that $\tilde{l} > l$, second that $\tilde{l} \notin I_{v,u,w}$ (so that $\tilde{l} < w$) and third that if $\tilde{l} < w$ then there exists $\tilde{\epsilon} > 0$ such that $[\tilde{l}, \tilde{l} + \tilde{\epsilon}) \cap I_{v,u,w} = \emptyset$. But, this contradicts the definition of \tilde{l} as the infimum of elements of $I_{v,u,w}$.

So, suppose $\overrightarrow{n}_{v,u}(l) < \overrightarrow{m}_{v,u}(w)$. Since $l < \overline{w}_{v,u}$ we have that $\overrightarrow{G}_{\mu}(l) \leq \overrightarrow{G}_{\mu}(l+) <$ $\overrightarrow{n}_{v,u}(l)$, and there exists $\overline{\epsilon} > 0$ with $l + \overline{\epsilon} < w$ such that $\overrightarrow{m}_{v,u}(l+\epsilon) \leq \overrightarrow{G}_{\mu}(l+\epsilon) \leq \overrightarrow{n}_{v,u}(l)$ for all $\epsilon \in [0, \bar{\epsilon}]$. In particular, $\overrightarrow{m}_{v,u}(l+\epsilon) \leq \overrightarrow{n}_{v,u}(l)$ for all $\epsilon \in [0, \bar{\epsilon}]$, and it follows that $l > l + \bar{\epsilon} > l.$

Now we show that $\overrightarrow{m}_{v,u}(\cdot)$ is non-increasing on $[l, \tilde{l})$. Let $l_1, l_2 \in [l, \tilde{l})$ with $l_1 < l_2$. Since $l_1 < \overline{w}_{v,u}$ we have that $\overrightarrow{m}_{v,u}(l_1) \leq \overrightarrow{G}_{\mu}(l_1) \leq \overrightarrow{G}_{\mu}(l_1+) < \overrightarrow{n}_{v,u}(l_1)$ and, since $l_1 < \tilde{l}, \; \overrightarrow{n}_{v,u}(l) \geq \overrightarrow{m}_{v,u}(l_1)$. Then, applying Lemma [3.9](#page-13-1) with $r_1 = l$ and $r_2 = l_1$ we have that $\vec{m}_{v,u}(l_1) \leq \vec{m}_{v,u}(l)$. Since $l_2 < \tilde{l}$, similarly as for l_1 , we have that $\vec{m}_{v,u}(l_2) \leq$ $\vec{m}_{v,u}(l) \leq \vec{n}_{v,u}(l)$, and therefore $\vec{m}_{v,u}(l_2) \leq \vec{n}_{v,u}(l_1)$. Then, by Lemma [3.9](#page-13-1) with $r_1 = l_1$ and $r_2 = l_2$, $\overrightarrow{m}_{v,u}(l_2) \leq \overrightarrow{m}_{v,u}(l_1)$ as required.

We now show that $\tilde{l} \notin I_{v,u,w}$. Note that $\mathcal{E}_{v,\tilde{l}}^c$ is linear on $[\overrightarrow{m}_{v,u}(k), \overrightarrow{n}_{v,u}(k)]$ for all $k \in [l, \tilde{l})$. By the monotonicity of $\overrightarrow{m}_{v,u}(\cdot)$ and $\overrightarrow{n}_{v,u}(\cdot)$ on $[l, \tilde{l})$, by Lemma [3.6](#page-11-1) and the fact that for each $k \in [l, \tilde{l}), \nu - S^{\nu}(\mu_k - \mu_v)$ does not charge $(\overrightarrow{m}_{\underline{v},u}(k), \overrightarrow{n}_{\underline{v},\underline{u}}(k)))$ we have that $\mathcal{E}_{v,\tilde{l}}^c$ is linear on $[\overrightarrow{m}_{v,u}(\tilde{l}-), \overrightarrow{n}_{v,u}(\tilde{l}-)]$. Since $\overrightarrow{m}_{v,u}(k) \leq \overrightarrow{G}_{\mu}(k) \leq \overrightarrow{G}_{\mu}(k+) < \overrightarrow{n}_{v,u}(k)$ for all $k \in [l, \tilde{l})$, $\overrightarrow{m}_{v,u}(\tilde{l}-) \leq \overrightarrow{G}_{\mu}(\tilde{l}) \leq \overrightarrow{n}_{v,u}(\tilde{l}-)$. But $[\overrightarrow{m}_{v,u}(\tilde{l}), \overrightarrow{n}_{v,u}(\tilde{l})]$ is the largest interval (containing $\overrightarrow{G}_{\mu}(\tilde{l})$) on which $\mathcal{E}^c_{v,\tilde{l}}$ is linear. It follows that $\overrightarrow{m}_{v,u}(\tilde{l}) \leq \overrightarrow{m}_{v,u}(\tilde{l}) \leq \overrightarrow{m}_{v,u}(\tilde{l})$ $\overrightarrow{m}_{v,u}(l) < \overrightarrow{n}_{v,u}(l)$, and therefore $\tilde{l} \notin I_{v,u,w}$ and $\tilde{l} < w$.

Finally, we have that

$$
\overrightarrow{m}_{v,u}(\tilde{l}) \leq \overrightarrow{G}_{\mu}(\tilde{l}) \leq \overrightarrow{G}_{\mu}(\tilde{l}+) < \overrightarrow{n}_{v,u}(\tilde{l}).
$$

Since $\tilde{l} < w$, there exists $0 < \tilde{\epsilon} < (w - \tilde{l})$ such that, for all $\epsilon \in [0, \tilde{\epsilon}], \overrightarrow{G}_{\mu}(\tilde{l} + \epsilon) \leq \overrightarrow{n}_{v,u}(\tilde{l}).$ But, since $[\overrightarrow{m}_{v,u}(\tilde{l}+\epsilon), \overrightarrow{n}_{v,u}(\tilde{l}+\epsilon)]$ is the largest interval (containing $\overrightarrow{G}_{\mu}(\tilde{l}+\epsilon)$) on which $\mathcal{E}^c_{v,\tilde{l}+\epsilon}$ is linear, we have that $\overrightarrow{m}_{v,u}(\tilde{l}+\epsilon) \leq \overrightarrow{m}_{v,u}(\tilde{l}) < \overrightarrow{n}_{v,u}(l)$. Hence $[\tilde{l}, \tilde{l}+\tilde{\epsilon}) \cap I_{v,u,w} = \emptyset$ and l is not the infimum of elements of $I_{v,u,w}$. \Box

Corollary 3.12. Fix $v, u \in [0, \mu(\mathbb{R}))$ with $v \leq u$. If $\overrightarrow{G}_{\mu}(u+) < \overrightarrow{n}_{v,u}(u)$ then $\overline{w}_{v,u} > u$ and $\overrightarrow{m}_{v,u}(\cdot)$ is non-increasing on $[u,\overline{w}_{v,u}].$

Proof. By Lemma [3.10,](#page-13-2) if $\overrightarrow{G}_{\mu}(u+) < \overrightarrow{n}_{v,u}(u)$ then $\overline{w}_{v,u} > u$. Then, from Lemma [3.11](#page-14-0) we have that $\overrightarrow{m}_{v,u}(\cdot)$ is non-increasing on $(u,\overline{w}_{v,u}]$ so it only remains to consider the behaviour at u.

By hypothesis we have that $\overrightarrow{m}_{v,u}(u) \leq \overrightarrow{G}_{\mu}(u) \leq \overrightarrow{G}_{\mu}(u+\) \leq \overrightarrow{n}_{v,u}(u)$, and therefore $\mathcal{E}_{v,u}^c$ is linear on $[\overrightarrow{m}_{v,u}(u), \overrightarrow{n}_{v,u}(u)]$. By the left-continuity of $\overrightarrow{G}_{\mu}(\cdot)$, and since $u < \mu(\mathbb{R})$, there exists $\hat{\epsilon} \in (0, \underline{\mu}(\mathbb{R}) \wedge \overline{w}_{v,u} - u]$ such that for all $\epsilon \in [0, \hat{\epsilon}], \ \overrightarrow{G}_{\mu}(u + \epsilon) \leq \overrightarrow{n}_{v,u}(u)$. Then $\overrightarrow{m}_{v,u}(u+\epsilon) \leq \overrightarrow{G}_{\mu}(u+\epsilon) \leq \overrightarrow{n}_{v,u}(u)$ and, by Lemma [3.9](#page-13-1) with $r_1 = u$ and $r_2 = u+\epsilon$, $\overrightarrow{m}_{v,u}(u+\epsilon) \leq \overrightarrow{m}_{v,u}(u)$ for all $\epsilon \in [0,\hat{\epsilon}].$ □

Lemma 3.13. Let $v, u \in [0, \mu(\mathbb{R}))$ with $v \leq u < \overline{w}_{v,u}$. Suppose that either $v = u$, or $v < u$ and $\overrightarrow{m}_{v,u}(u+) \leq \overrightarrow{G}_{\mu}(v+)$. Then

- (i) $S^{\nu}(\mu_l \mu_v) = \nu|_{(\overrightarrow{m}_{v,u}(l), \overrightarrow{n}_{v,u}(l))}$, for all $l \in (u, \overline{w}_{v,u}]$.
- (ii) $\overrightarrow{n}_{v,u}$ is strictly increasing and $\overrightarrow{m}_{v,u}$ is strictly decreasing on $(u,\overline{w}_{v,u})$.

Proof. (i) Fix $l \in (u, \overline{w}_{v,u})$. By Lemma [3.6,](#page-11-1) $S^{\nu}(\mu_l - \mu_v) = \nu$ on $(\overrightarrow{m}_{v,u}(l), \overrightarrow{n}_{v,u}(l))$. In order to conclude that $S^{\nu}(\mu_l - \mu_v) = \nu|_{(\vec{m}_{v,u}(l),\vec{\pi}_{v,u}(l))}$ it remains to show that $S^{\nu}(\mu_l - \mu_v)$ $(\mu_v)(\mathbb{R}\setminus(\overrightarrow{m}_{v,u}(l),\overrightarrow{n}_{v,u}(l)))=0.$

Suppose that $\overrightarrow{m}_{v,u}(l) \leq \overrightarrow{G}_{\mu}(v)$. Note that the second derivative of $\mathcal{E}_{v,l}$ on $(-\infty, \overrightarrow{G}_{\mu}(v)) \cup$ $(\overrightarrow{G}_{\mu}(l), \infty)$ corresponds to ν . Then since ν is continuous and $\mathcal{E}_{v,l}^c = \mathcal{E}_{v,l}$ on $(-\infty, \overrightarrow{m}_{v,u}(l)] \cup$ $[\vec{n}_{v,u}(l),\infty)$ we have that $(\nu - S^{\nu}(\mu_l - \mu_v)) = \nu$ on $(-\infty, \vec{m}_{v,u}(l)) \cup [\vec{n}_{v,u}(l),\infty)$, and therefore $S^{\nu}(\mu_l - \mu_v)((\overrightarrow{m}_{v,u}(\underline{l}), \overrightarrow{n}_{v,u}(l))) = S^{\nu}(\mu_l - \mu_v)(\mathbb{R})$, and we are done.

Now suppose $\overrightarrow{m}_{v,u}(l) > \overrightarrow{G}_{\mu}(v)$. Since $\overrightarrow{m}_{v,u}(\cdot)$ is non-increasing on $(u, \overline{w}_{v,u})$ (see Lemma [3.11\)](#page-14-0), $\vec{m}_{v,u}(l) \leq \vec{m}_{v,u}(u+) \leq \vec{G}_{\mu}(v+)$ (the second inequality follows directly if $v = u$, or by assumption in the case $v < u$). It follows that μ does not charge $(\overrightarrow{G}_{\mu}(v), \overrightarrow{m}_{v,u}(l)) \subseteq (\overrightarrow{G}_{\mu}(v), \overrightarrow{G}_{\mu}(v+))$. Therefore, $\mathcal{E}_{v,l}$ is convex on $(-\infty, \overrightarrow{m}_{v,u}(l))$ and its second (distributional) derivative on $(-\infty, \overrightarrow{m}_{v,u}(l))$ corresponds to ν . Since $\mathcal{E}_{v,l}^c =$ $\mathcal{E}_{v,l}$ on $[\vec{n}_{v,u}(l),\infty)$ and hence $\nu-S^{\nu}(\mu_l-\mu_v)=\nu$ on $[\vec{n}_{v,u}(l),\infty)$, we again conclude that $S^{\nu}(\mu_l - \mu_v)((\vec{m}_{v,u}(l), \vec{n}_{v,u}(l))) = S^{\nu}(\mu_l - \mu_v)(\mathbb{R}).$

We now show that the assertion holds for $l = \overline{w}_{v,u}$ as well. By the monotonicity of $\overrightarrow{m}_{v,u}$ and $\overrightarrow{n}_{v,u}$ on $(u,\overline{w}_{v,u})$ we have that $\overrightarrow{m}_{v,u}(\overline{w}_{v,u}) \leq \overrightarrow{m}_{v,u}(\overline{w}_{v,u}-)$ and $\overrightarrow{n}_{v,u}(\overline{w}_{v,u}-) \leq$ $\overrightarrow{n}_{v,u}(\overline{w}_{v,u})$. On the other hand, $S^{\nu}(\mu_{\overline{w}_{v,u}} - \mu_v) = \nu$ on $((\overrightarrow{m}_{v,u}(\overline{w}_{v,u}-), \overrightarrow{n}_{v,u}(\overline{w}_{v,u}-))$ and $S^{\nu}(\mu_{\overline{w}_{v,u}} - \mu_v)(\mathbb{R}) = \overline{w}_{v,u} - v = \lim_{l \uparrow \overline{w}_{v,u}} l - v = \lim_{l \uparrow \overline{w}_{v,u}} S^{\nu}(\mu_l - \mu_v)(\mathbb{R}) =$ $\lim_{l\uparrow \overline{w}_{v,u}} \nu((\overrightarrow{m}_{v,u}(l),\overrightarrow{n}_{v,u}(l))) = \nu((\overrightarrow{m}_{v,u}(\overline{w}_{v,u}-),\overrightarrow{n}_{v,u}(\overline{w}_{v,u}-)))$. It follows that $S^{\nu}(\mu_{\overline{w}_{v,u}}-\mu)$ μ_v) = $\nu|_{(\overrightarrow{m}_{v,u}(\overline{w}_{v,u}-),\overrightarrow{n}_{v,u}(\overline{w}_{v,u}-))} = \nu|_{(\overrightarrow{m}_{v,u}(\overline{w}_{v,u}),\overrightarrow{n}_{v,u}(\overline{w}_{v,u}))}$, where the last equality follows from the fact that $\mathcal{E}_{v,\overline{w}_{v,u}}^c$ is linear on $[\overrightarrow{m}_{v,u}(\overline{w}_{v,u}), \overrightarrow{n}_{v,u}(\overline{w}_{v,u})]$ and therefore, in the case $\overrightarrow{m}_{v,u}(\overline{w}_{v,u}) < \overrightarrow{m}_{v,u}(\overline{w}_{v,u-})$ (resp. $\overrightarrow{n}_{v,u}(\overline{w}_{v,u-}) < \overrightarrow{n}_{v,u}(\overline{w}_{v,u}))$, $\nu((\overrightarrow{m}_{v,u}(\overline{w}_{v,u}), \overrightarrow{m}_{v,u}(\overline{w}_{v,u-})))$ 0 (resp. $\nu((\overrightarrow{n}_{v,u}(\overline{w}_{v,u}-), \overrightarrow{n}_{v,u}(\overline{w}_{v,u})]) = 0).$

(ii) Let $u < l < w \leq \overline{w}_{u,v}$. We cannot have both $\overrightarrow{m}_{v,u}(l) = \overrightarrow{m}_{v,u}(w)$ and $\overrightarrow{n}_{v,u}(l) =$ $\overrightarrow{n}_{v,u}(w)$ else $w-l=S^{\nu-S^{\nu}(\mu_l-\mu_v)}(\mu_w-\mu_l)(\mathbb{R})=0$, a contradiction. Suppose $\overrightarrow{m}_{v,u}(w)$ < $\overrightarrow{m}_{v,u}(l)$ and $\overrightarrow{n}_{v,u}(l) = \overrightarrow{n}_{v,u}(w)$. Then $S^{\nu-S^{\nu}(\mu_l-\mu_v)}(\mu_w-\mu_l) = \nu|_{(\overrightarrow{m}_{v,u}(w),\overrightarrow{m}_{v,u}(l))}$. But this cannot hold since $\mu_w - \mu_l$ places all its mass at or to the right of $\overline{G}_{\mu}(l)$ and $\overrightarrow{G}_{\mu}(l) \geq \overrightarrow{m}_{v,u}(l)$ and so the shadow measure $S^{\nu-S^{\nu}(\mu_l-\mu_v)}(\mu_w-\mu_l)$, being a measure in convex order with respect to $\mu_w - \mu_l$, must place some mass to the right of $\vec{m}_{v,u}(l)$.

Now we show that $\vec{m}_{v,u}(w) = \vec{m}_{v,u}(l)$ and $\vec{n}_{v,u}(l) < \vec{n}_{v,u}(w)$ cannot happen either. Since $l < \overline{w}_{v,u}$, $\overrightarrow{G}_{\mu}(l) \leq \overrightarrow{G}_{\mu}(l+)< \overrightarrow{n}_{v,u}(l)$. Then there exists $\tilde{\epsilon} > 0$ such that $l + \tilde{\epsilon} < w$ and for all $\epsilon \in [l, l+\tilde{\epsilon}]$ we have that $\overrightarrow{m}_{v,u}(w) = \overrightarrow{m}_{v,u}(l) = \overrightarrow{m}_{v,u}(l+\epsilon) \leq \overrightarrow{G}_{\mu}(l+\epsilon) \leq \overrightarrow{n}_{v,u}(l)$. But then $S^{\nu-S^{\nu}(\mu_l-\mu_v)}(\mu_{l+\epsilon}-\mu_l)$ concentrates on $(\overrightarrow{n}_{v,u}(l), \overrightarrow{n}_{v,u}(l+\epsilon))$ and thus to the right of $\vec{G}_{\mu}(l + \epsilon)$, while $\mu_{l+\epsilon} - \mu_l$ places all its mass at or to the left of $\vec{G}_{\mu}(l + \epsilon)$, a contradiction.

Corollary 3.14. Suppose that $v, u \in [0, \mu(\mathbb{R}))$ with $v \leq u$. Suppose that $\overrightarrow{G}_{\mu}(u+)$ $\overrightarrow{n}_{v,u}(u)$ and either $v = u$, or $v < u$ and $\overrightarrow{m}_{v,u}(u+) \leq \overrightarrow{G}_{\mu}(v+)$. Then $\overrightarrow{n}_{v,u}$ is strictly increasing and $\overrightarrow{m}_{v,u}$ is strictly decreasing on $[u,\overline{w}_{v,u}].$

 \Box

Proof. Monotonicity follows from Corollaries [3.8](#page-13-0) and [3.12;](#page-15-0) strict monotonicity on $[u,\overline{w}_{v,u}]$ then follows from strict monotonicity on $(u,\overline{w}_{v,u}]$. П

Let
$$
u, v \in [0, \mu(\mathbb{R})]
$$
 with $u \le v$. Define $\overleftarrow{m}_{u,v}, \overleftarrow{n}_{u,v} : [0, u] \to \mathbb{R}$ by
\n
$$
\overleftarrow{m}_{u,v}(l) = X_{\mathcal{E}_{l,v}}(\overleftarrow{G}_{\mu}(l)) \text{ and } \overleftarrow{n}_{u,v}(l) = Z_{\mathcal{E}_{l,v}}^+(\overleftarrow{G}_{\mu}(l)), \quad l \in [0, u].
$$

The following results follow by symmetry, working right-to-left instead of left-to-right.

Lemma 3.15. $Fix\ 0 \le l \le u \le v \le \mu(\mathbb{R})$. Then, for all $w \in [0, l]$, $S^{\nu}(\mu_v - \mu_w) = \nu$ on $(\overleftarrow{m}_{u,v}(l), \overleftarrow{n}_{u,v}(l))$ and $\mathcal{E}_{w,v}^c$ is linear on $[\overleftarrow{m}_{u,v}(l), \overleftarrow{n}_{u,v}(l)].$

Corollary 3.16. Fix $0 \leq l \leq u \leq v \leq \mu(\mathbb{R})$. If $l \in (0, \mu(\mathbb{R}))$ then $\alpha_{\nu} < \overleftarrow{m}_{u,v}(l) \leq$ $\overleftarrow{n}_{u,v}(l) < \beta_{\nu}.$

Corollary 3.17. Fix $u, v \in (0, \mu(\mathbb{R})]$ with $u \leq v$. $\overleftarrow{m}_{u,v}(\cdot)$ is non-decreasing on [0, u].

For $u, v \in (0, \mu(\mathbb{R})]$ with $u \leq v$ define $\underline{w}_{u,v} := \sup\{w \in [0, u) : \overleftarrow{m}_{u,v}(w) \geq \overleftarrow{G}_{\mu}(w-) \}$ with convention $\sup \emptyset = 0$.

Lemma 3.18. 1. Suppose $0 < u \le v \le \mu(\mathbb{R})$. If $\overleftarrow{m}_{u,v}(u) < \overleftarrow{G}_{\mu}(u-)$ then $\underline{w}_{u,v} < u$.

2. If $0 \leq \underline{w}_{u,v} < u \leq v \leq \mu(\mathbb{R})$. Then $\overleftarrow{G}_{\mu}(\underline{w}_{u,v}) \leq \overleftarrow{m}_{u,v}(\underline{w}_{u,v}) \leq \overleftarrow{G}_{\mu}(\underline{w}_{u,v})$ and $D(\overleftarrow{m}_{u,v}(\underline{w}_{u,v})) = \mathcal{E}_{u,v}(\overleftarrow{m}_{u,v}(\underline{w}_{u,v})) = \mathcal{E}_{u,v}^c(\overleftarrow{m}_{u,v}(\underline{w}_{u,v})).$

Lemma 3.19. Fix $u, v \in (0, \mu(\mathbb{R}))$ with $u \leq v$. Then $\overleftarrow{n}_{u,v}(\cdot)$ is non-increasing on $\lfloor \underline{w}_{u,v}, u \rfloor$.

Corollary 3.20. Fix $u, v \in (0, \mu(\mathbb{R})]$ with $u \leq v$. If $\underline{w}_{u,v} < u$ and $\overleftarrow{m}_{u,v}(u) < \overleftarrow{G}_{\mu}(u-)$ then $\underline{w}_{u,v} < u$ and $\overleftarrow{n}_{u,v}(\cdot)$ is non-increasing on $[\underline{w}_{u,v}, u]$.

Lemma 3.21. Let $v, u \in (0, \mu(\mathbb{R}))$ with $\underline{w}_{u,v} < u \leq v$. Suppose that either $u = v$, or $u < v$ and $\overleftarrow{G}_u(v-) \leq \overleftarrow{n}_{u,v}(u-)$. Then

(i) $S^{\nu}(\mu_v - \mu_l) = \nu|_{(\overleftarrow{m}_{u,v}(l), \overleftarrow{n}_{u,v}(l))},$ for all $l \in [\underline{w}_{u,v}, u)$.

(ii) $\overleftarrow{n}_{u,v}$ is strictly decreasing and $\overleftarrow{m}_{u,v}$ is strictly increasing on $[\underline{w}_{u,v}, u)$.

Corollary 3.22. Suppose that $v, u \in (0, \mu(\mathbb{R})]$ with $u \leq v$. Suppose that $\overline{m}_{v,u}(u)$ $\overleftarrow{G}_{\mu}(u-)$ and either $v = u$ or $u < v$ and $\overleftarrow{G}_{\mu}(v-) \leq \overleftarrow{n}_{v,u}(u-)$. Then $\overrightarrow{n}_{u,v}$ is strictly decreasing and $\vec{m}_{v,u}$ is strictly increasing on $[\underline{w}_{u,v}, u]$.

4 The construction in the regular case

The goal of this section is to construct an injective lifted martingale coupling of μ and ν in a fairly general, but still regular case. We may restrict attention to the irreducible case, so our general setting is pairs of measures in the following set.

Definition 4.1. Suppose $\mu, \nu \in \mathcal{P}$. Then $(\mu, \nu) \in \mathcal{K}$ if ν is continuous, if $\mu \leq_{cx} \nu$ and if $\{x : D_{\mu,\nu}(x) > 0\} = I_{\nu}$.

We begin by outlining the principles which govern our approach.

Given $(\mu, \nu) \in \mathcal{K}$ the first step is to choose a suitable starting point $u_0 \in [0, \mu(\mathbb{R}))$ and then to consider $(\mathcal{E}_{u_0,u})_{u_0\leq u\leq \mu(\mathbb{R})}$ and to define $u_1=\overline{w}_{u_0,u_0}$. Assuming that $u_1>u_0$, we define M and N on $[u_0, u_1]$ via $M(u) = \overrightarrow{m}_{u_0, u_0}(u)$ and $N(u) = \overrightarrow{m}_{u_0, u_0}(u)$. These functions are strictly monotonic (on the domain where they have been defined) and moreover, for each $u \in [u_0, u_1]$ we have that $S^{\nu}(\mu_u - \mu_{u_0}) = \nu|_{(M(u), N(u))}$. In particular, the functions M and N can be used to define a lifted martingale coupling of $\mu_{u_1} - \mu_{u_0}$ and $\nu|_{(M(u_1),N(u_1))}$. Moreover, by the strict monotonicity of M and N this coupling is injective.

If $u_0 = 0$ it will then follow by Lemma [5.10](#page-34-1) below that $u_1 = \mu(\mathbb{R})$ and then the construction is complete, and we have described an injective martingale coupling. More generally, if $u_0 > 0$ then this need not be the case. Then we proceed by induction. If $u_0 > 0$ (and then $u_1 \leq \mu(\mathbb{R})$) we consider $(\mathcal{E}_{u,u_1})_{0 \leq u \leq u_0}$ and set $u_2 = \underline{w}_{u_0,u_1}$. It will turn out that $u_2 < u_0$ and we can extend the definitions of M and N to $[u_2, u_1]$ such that M and N are strictly monotonic on $[u_2, u_0]$ and such that $S^{\nu}(\mu_{u_1} - \mu_u) = \nu|_{(M(u), N(u))}$. In particular, $S^{\nu}(\mu_{u_1} - \mu_{u_2}) = \nu|_{(M(u_2), N(u_2))}$ and M and N now defined on $[u_2, u_0]$ can be used to define a lifted martingale coupling of $\mu_{u_1} - \mu_{u_2}$ and $\nu|_{(M(u_2),N(u_2))}$. It follows from the properties of M and N that the coupling is injective.

If $u_1 = \mu(\mathbb{R})$ then $\mu_2 = 0$ and the construction terminates. Otherwise we consider $(\mathcal{E}_{u_2,u})_{u\geq u_1}$ and proceed by induction working alternately from left-to-right and then right-to-left.

Suppose we have $0 \leq u_{2k} < u_{2k-2} < \ldots u_0 < u_1 < \ldots u_{2k-1} < \mu(\mathbb{R})$ and M and N defined on $[u_{2k}, u_{2k-1}]$. Then we define $u_{2k+1} \in (u_{2k-1}, \mu(\mathbb{R})]$ by $u_{2k+1} = \overline{w}_{u_{2k}, u_{2k-1}}$, and extend the domain of definition of M and N to $[u_{2k}, u_{2k+1}]$ such that for $u \in$ $(u_{2k-1}, u_{2k+1}], S^{\nu}(\mu_u - \mu_{u_{2k}}) = \nu|_{(M(u), N(u))}.$ Either $u_{2k} = 0$ and then $\mu_{2k+1} = \mu(\mathbb{R})$ and the construction terminates, or $0 < u_{2k} < u_{2k-2} < \ldots < u_0 < u_1 < \ldots < u_{2k-1} <$ $u_{2k+1} \leq \mu(\mathbb{R})$. Then we define $u_{2k+2} \in [0, u_{2k})$ by $u_{2k+2} = \underline{w}_{u_{2k}, u_{2k+1}}$ and extend the definitions of M and N to $[u_{2k+2}, u_{2k+1}].$

Either $u_j = 0$, $\mu_{j+1} = \mu(\mathbb{R})$ for some (even) j, or $u_j = \mu(\mathbb{R})$, $\mu_{j+1} = 0$ for some (odd) j, and we have a pair of functions $\{M, N\}$ is defined on $[0, \mu(\mathbb{R}),$ or we have a pair of decreasing and increasing sequences such that $0 < \ldots < u_{2k} < u_{2k-2} < \ldots < u_0 <$ $u_1 < \ldots < u_{2k-1} < u_{2k+1} < \ldots < \mu(\mathbb{R})$ and a pair of functions $\{M, N\}$ is defined on (u_{∞}, u^{∞}) where $u_{\infty} := \lim_{k} u_{2k}$ and $u^{\infty} := \lim_{k} u_{2k+1}$.

The main issues are: first, to argue that $u_1 > u_0$ and thereafter $u_{2k} < u_{2k-2}$ and $u_{2k+1} > u_{2k-1}$ at least until u_{2k} reaches zero or u_{2k+1} reaches $\mu(\mathbb{R})$ and the construction terminates; second, to show that if the construction does not terminate then $u_{\infty} = 0$ and $u^{\infty} = \mu(\mathbb{R})$; third to prove that $S^{\nu}(\mu_{u_{2k+1}} - \mu_{u_{2k}}) = \nu|_{(M(u_{2k+1}), N(u_{2k+1}))}$ and that $(M(u), N(u))_{0\leq u\leq u(\mathbb{R})}$ define an lifted martingale coupling; and fourth to justify that M and N have appropriate monotonicity properties so that this coupling is injective.

In fact, it is not the case that the construction will work as described in the general case. For this reason we introduce a pair of subspaces $\mathcal{K}_{*} \subseteq \mathcal{K}_{R} \subseteq \mathcal{K}$. We will show that the construction outlined above defines an injective coupling on \mathcal{K}_{*} . Later we show that the general case can be reduced to this case.

Let G_{μ} be any quantile function of μ .

Definition 4.2. Suppose $(\mu, \nu) \in \mathcal{K}$. Then $(\mu, \nu) \in \mathcal{K}_R$ if (see Figure [2\)](#page-20-0)

- 1. $\exists a \in \mathbb{R}$ such that $\mu \leq \nu$ on $(-\infty, a)$; let a_0 be the largest such value and let $u_0 := \mu((-\infty, a_0))$ so that $G_{\mu}(u_0+) = a_0$; note that if $a \in (G_{\mu}(u_0), G_{\mu}(u_0+))$ is such that $\mu \leq \nu$ on $(-\infty, a)$ then $\mu \leq \nu$ on $(-\infty, G_{\mu}(u_0+))$; let $a_1 = \inf\{a :$ $\mu = \nu$ on (a, a_0) , with the convention inf $\emptyset = a_0$.
- 2. The tangent line L_{a_0} to $D_{\mu,\nu}$ at a_0 is such that $D_{\mu,\nu} < L_{a_0}$ on (a_0, β_{ν}) .
- 3. for all $b \in (\alpha_{\nu}, a_1)$ the tangent line L_b to D at b, given by $L_b(k) = D_{\mu,\nu}(b) + (k$ b) $D'_{\mu,\nu}(b)$, is such that there exists c with $c > a_0$ such that $D_{\mu,\nu} > L_b$ on (a_0, c) , $D_{\mu,\nu} < L_b$ on (c, ∞) and $D'_{\mu,\nu}(b) > D'_{\mu,\nu}(c-)$.

Note that since $\mu \leq \nu$ on $(-\infty, a_0)$ and ν is continuous, μ is also continuous on this set. Then $D_{\mu,\nu}$ is continuously differentiable and convex on $(-\infty, a_0)$ and has a unique tangent at each $b < a_0$.

Lemma 4.3. Suppose $(\mu, \nu) \in \mathcal{K}_R$. Then $u_0 < \mu(\mathbb{R})$.

Proof. If $u_0 = \mu(\mathbb{R})$ then $\mu \leq \nu$ on \mathbb{R} and therefore $\mu = \nu$, since $\mu \leq_{cx} \nu$. Then $D_{\mu,\nu} \equiv 0$, but this is a case which we have ruled out by Standing Assumption [2.3\)](#page-6-1). \Box

Figure 2: Plot of $D_{\mu,\nu}$ for $(\mu,\nu) \in \mathcal{K}_R$. The line L_{a_0} satisfies $D_{\mu,\nu} > L_{a_0}$ on $(-\infty, a_1)$, $D_{\mu,\nu} = L_{a_0}$ on $[a_1, a_0]$ and $D_{\mu,\nu} < L_{a_0}$ on (a_0, ∞) . On the other hand, the line L_b , that is tangent to $D_{\mu,\nu}$ at $b < a_1$, crosses $D_{\mu,\nu}$ at $c > a_0$ and satisfies $D_{\mu,\nu} \geq L_b$ on $(-\infty, c]$ and $D_{\mu,\nu} < L_b$ on (c,∞) .

Lemma 4.4. Suppose $(\mu, \nu) \in \mathcal{K}_R$.

- (i) Suppose $0 < v \leq u_0 \leq u < \overline{w}_{v,u} \leq \mu(\mathbb{R})$ and $\overrightarrow{m}_{v,u}(\overline{w}_{v,u}) < \overrightarrow{G}_{\mu}(v+\) \wedge a_1$. Then $\overrightarrow{m}_{v,u}(\overline{w}_{v,u}) < \overrightarrow{G}_{\mu}(v).$
- (ii) Suppose $0 \leq \underline{w}_{u,v} < u \leq u_0 \leq v < \mu(\mathbb{R})$, $\overleftarrow{m}_{u,v}(\underline{w}_{u,v}) < a_1$ and $\overleftarrow{n}_{u,v}(\underline{w}_{u,v}) >$ $\overleftarrow{G}_{\mu}(v-)$. Then $\overleftarrow{n}_{u,v}(\underline{w}_{u,v}) > \overleftarrow{G}_{\mu}(v)$.

Proof. We first prove (i). Suppose the conclusion does not hold, so that $\vec{G}_{\mu}(v) \leq$ $\overrightarrow{m}_{v,u}(\overline{w}_{v,u}) < \overrightarrow{G}_{\mu}(v+)$. Then

$$
\mathcal{E}_{v,\overline{w}_{v,u}}^c(k) = \mathcal{E}_{v,\overline{w}_{v,u}}(k) = D_{\mu,\nu}(k)
$$
\n(3)

for $k = \overrightarrow{m}_{v,u}(\overline{w}_{v,u})$. Suppose that $\overrightarrow{n}_{v,u}(\overline{w}_{v,u}) < r_{\nu} \leq \infty$. Then, by the results of Lemma [3.10,](#page-13-2) [\(3\)](#page-20-1) also holds at $k = \overrightarrow{n}_{v,u}(\overline{w}_{v,u})$. Then if $b := \overrightarrow{m}_{v,u}(\overline{w}_{v,u})$ (where $b < a_1$) by hypothesis) we find that $\overrightarrow{n}_{v,u}(\overline{w}_{v,u})$ plays the role of c in Definition [4.2](#page-19-0) in the sense that $D_{\mu,\nu} > L_b$ on (a_0, c) and $D_{\mu,\nu} < L_b$ on (c, ∞) . However, $D'_{\nu,\mu}(b) = D'_{\nu,\mu}(c)$ and so the condition $D'_{\nu,\mu}(b) > D'_{\nu,\mu}(c-)$ is not satisfied. Hence $(\nu,\mu) \notin \mathcal{K}_R$, a contradiction. We conclude that $\overrightarrow{m}_{v,u}(\overline{w}_{v,u}) < \overrightarrow{G}_{\mu}(v)$.

It is left to argue that $\overrightarrow{n}_{v,u}(\overline{w}_{v,u}) < r_{\nu}$. Suppose not; so that $\overrightarrow{n}_{v,u}(\overline{w}_{v,u}) \geq r_{\nu}$. Then $\lim_{k\to \vec{n}_{v,u}(\overline{w}_{v,u})} D(k) = 0$ and the line joining $\mathcal{E}_{v,\overline{w}_{v,u}}^c(\overrightarrow{m}_{v,u}(\overline{w}_{v,u}))$ and $\mathcal{E}_{v,\overline{w}_{v,u}}^c(\overrightarrow{n}_{v,u}(\overline{w}_{v,u}))$

must have slope equal to zero, from which it follows that $\mathcal{E}_{v,u}(\overrightarrow{m}_{v,u}(\overrightarrow{w}_{v,u})) = 0$. But this contradicts the fact that $\mathcal{E}_{v,u} > 0$ on $(-\infty, r_{\nu})$, since $v > 0$.

Part (ii) follows similarly, where again we have $b = \overline{m}_{v,u}(\overline{w}_{v,u}) < a_1$ by hypothesis. \Box

Lemma 4.5. Suppose $(\mu, \nu) \in \mathcal{K}_R$.

- i) Suppose $0 \le u < \mu(\mathbb{R})$ and $\overline{w}_{0,u} > u$. Then $\overline{w}_{0,u} = \mu(\mathbb{R})$.
- ii) Suppose $0 < u \leq u_0$ and $\underline{w}_{u,\mu(\mathbb{R})} < u$. Then $\underline{w}_{u,\mu(\mathbb{R})} = 0$.

Proof. i) Suppose $\overline{w}_{0,u} < \mu(\mathbb{R})$. It follows that $\mathcal{E}_{0,\overline{w}_{0,u}} = D$ on $(-\infty, \overrightarrow{G}_{\mu}(\overline{w}_{0,u}+))$. Note that $\vec{G}_{\mu}(0+) = \alpha_{\mu} \leq a_1$. Then, since $\overline{w}_{0,\mu} \in (0, \mu(\mathbb{R}))$, by Corollary [3.7](#page-12-0) we have that $b := \overrightarrow{m}_{0,u}(\overline{w}_{0,u}) > \alpha_{\nu}$, so that $\mathcal{E}_{0,\overline{w}_{0,\mu}}^c(b) = D(b) > 0$. On the other hand, by Lemma [3.10,](#page-13-2) $c := \overrightarrow{n}_{0,u}(\overline{w}_{0,u}) \in [\overrightarrow{G}_{\mu}(\overline{w}_{0,u}), \overrightarrow{G}_{\mu}(\overline{w}_{0,u}+)]$, and therefore $\mathcal{E}_{0,\overline{w}_{0,u}}^c(c) = D(c)$. Then, using the same arguments as in the proof of Lemma [4.4,](#page-20-2) we have that b and c violate Definition [4.2,](#page-19-0) and hence we must have that $\overline{w}_{0,u} = \mu(\mathbb{R})$.

ii) Suppose $\underline{w}_{u,\mu(\mathbb{R})} > 0$. Since $\underline{w}_{u,\mu(\mathbb{R})} \in (0, u)$, we have by Corollary [3.16](#page-17-0) that $c := \overleftarrow{n}_{u,\mu(\mathbb{R})}(\underline{w}_{u,\mu(\mathbb{R})}) < \beta_{\nu}$, and therefore $\mathcal{E}^c_{\underline{w}_{u,\mu(\mathbb{R})},\mu(\mathbb{R})}(c) = D(c) > 0$. On the other hand, by Lemma [3.18,](#page-17-1) $b := \overleftarrow{m}_{u,\mu(\mathbb{R})}(\underline{w}_{u,\mu(\mathbb{R})}) \in [\overleftarrow{G}(\underline{w}_{u,\mu(\mathbb{R})}-), \overleftarrow{G}(\underline{w}_{u,\mu(\mathbb{R})})]$ and $\mathcal{E}^c_{\underline{w}_{u,\mu(\mathbb{R})},\mu(\mathbb{R})}(b) =$ $D(b)$. Moreover, the derivatives of $\mathcal{E}^c_{\underline{w}_{u,\mu(\mathbb{R})},\mu(\mathbb{R})}$ at b and c must match, and be equal to the derivatives of $D = D_{\mu,\nu}$ at these points. But, $D'(c) < 0 < D'(b)$, a contradiction. \Box

Now we turn to the inductive step. We begin with some definitions.

Definition 4.6. For j even with $j \geq 2$ let $\mathcal{U}_j = \{(u_0, \ldots, u_j) \in [0, \mu(\mathbb{R})]^{j+1} : 0 \leq u_j < \infty \}$ $u_{j-2} < \ldots u_0 < u_1 < \ldots u_{j-3} < u_{j-1} \leq \mu(\mathbb{R})$. For j odd with $j \ge 1$ let $\mathcal{U}_j = \{(u_0, \ldots u_j) \in [0, \mu(\mathbb{R})]^{j+1} : 0 \le u_{j-1} < u_{j-3} < \ldots u_0 < \ell \}$ u_1 < ... u_{j-2} < u_j ≤ $\mu(\mathbb{R})$. For $j = \infty$ let $\mathcal{U}_{\infty} = \{(u_0, \ldots u_{2k-1}, u_{2k}, u_{2k+1}, \ldots) \in [0, \mu(\mathbb{R})]^{\infty} : 0 < \ldots < u_{2k} < \infty \}$ $u_{2k-2} < \ldots u_0 < u_1 < \ldots < u_{2k-1} < u_{2k+1} < \ldots < u(\mathbb{R})$.

Fix $j \in \mathbb{N} \cup \{\infty\}$ and $(u_0, u_1, \ldots, u_j) \in \mathcal{U}_j$ (or $(u_0, u_1, \ldots) \in \mathcal{U}_\infty$ if $j = \infty$, but in a slight abuse of notation we still write $(u_0, u_1, \ldots, u_j) \in \mathcal{U}_j$.

Set $A_j = [u_j, u_{j-1}]$ if $j \ge 2$ is even, $A_j = [u_{j-1}, u_j]$ if $j \ge 1$ is odd, and $A_j =$ $(\lim_{k \uparrow \infty} u_{2k}, \lim_{k \uparrow \infty} u_{2k+1})$ if $j = \infty$.

Let $J = \inf\{j \in \mathbb{N} : u_j = 0 \text{ or } u_j = \mu(\mathbb{R})\}$. For $j \geq 0$ set $v_j = |u_{j+1} - u_j|$. Then $(v_j)_{0 \le j \le J}$ is an increasing sequence and if $J < \infty$ then $v_J = \mu(\mathbb{R})$. Set $\mathbb{V} = \bigcup_{j=0}^J \{v_j\}$. For $u \in [0, \mu(\mathbb{R})]$ let $j(u) = \inf\{j \geq 0 : v_j \geq u\}$ with $\inf \emptyset = 0$. Define $\phi : [0, \mu(\mathbb{R})] \mapsto$ $[0, \mu(\mathbb{R})]$ by

Figure 3: Plots of ϕ (see the top figure) and alternately increasing and decreasing functions N and M (see the bottom figure). In the bottom figure the dotted curve represents $u \mapsto G_{\mu}(u)$. The solid curve above (resp. below) G_{μ} corresponds to the graph of $u \mapsto N(u)$ (resp. $u \mapsto M(u)$). Note that N (resp. M) is non-decreasing (resp. non-increasing) on $[u_0, u_1]$, non-increasing (resp. non-decreasing) on $[u_2, u_0]$ and again non-decreasing (resp. non-increasing) on $[u_1, u_3]$.

Definition 4.7.

$$
\phi(v) = \begin{cases} u_{j(v)} + v, & \text{if } j(v) \text{ is even,} \\ u_{j(v)} - v, & \text{if } j(v) \text{ is odd,} \end{cases} v \in [0, \mu(\mathbb{R})].
$$

Then ϕ is a right-continuous bijection, and moreover ϕ is continuous with $|\phi'| = 1$ on $[0, \mu(\mathbb{R})] \setminus \mathbb{V}$ and $\phi(\mathbb{V}) = \{u_j : j \ge 0\}$. See Figure [3.](#page-22-0)

Definition 4.8. Suppose $j \leq J+1$.

 $L: A_j \to \mathbb{R}$ is alternately increasing of order $j \in \mathbb{N} \cup \{\infty\}$ with respect to $(u_0, \ldots, u_j) \in$ \mathcal{U}_j if $L \circ \phi$ is increasing on $[0, v_{j-1}]$.

 $L: A_j \to \mathbb{R}$ is alternately decreasing of order $j \in \mathbb{N} \cup \{\infty\}$ with respect to $(u_0, \ldots, u_j) \in$ \mathcal{U}_j if $L \circ \phi$ is decreasing on $[0, v_{j-1}]$.

Remark 4.9. If L_1 is alternately decreasing and L_2 is alternately increasing, both of order $j \in \mathbb{N} \cup \infty$ and with respect to $(u_0, u_1, \ldots, u_j) \in \mathcal{U}_j$, and if $L_1(u_0) \leq L_2(u_0)$ then $L_1(w) \leq L_2(v)$ for all $(v, w) \in A \times A \setminus \{(u_0, u_0)\}.$

Suppose $(\mu, \nu) \in \mathcal{K}_{*}$ so that $\overline{w}_{u_0, u_0} > u_0$. Now set $u_1 = \overline{w}_{u_0, u_0}$. We define M, N on $[u_0, u_1 = \overline{w}_{u_0, u_0}]$ via $M(u) = \overline{m}_{u_0, u_0}(u)$ and $N(u) = \overline{m}_{u_0, u_0}(u)$. Assuming that $[u_0, u_1] \subsetneq$ $[0, \mu(\mathbb{R})]$ we then define $u_2 = \underline{w}_{u_0, u_1}$, and $M(u) = \overleftarrow{m}_{u_0, u_1}(u)$ and $N(u) = \overleftarrow{m}_{u_0, u_1}(u)$ on $[u_2, u_0]$. We proceed inductively: given k even we define $u_{k+1} > u_{k-1}$ via $u_{k+1} =$ $\overline{w}_{u_k, u_{k-1}}$ and M, N on $(u_{k-1}, u_{k+1}]$ by $M(u) = \overrightarrow{m}_{u_k, u_{k-1}}(u)$ and $N(u) = \overrightarrow{n}_{u_k, u_{k-1}}(u);$ given k odd we define $u_{k+1} < u_{k-1}$ via $u_{k+1} = \underline{w}_{u_{k-1},u_k}$ and M, N on $[u_{k+1}, u_{k-1}]$ by $M(u) = \overleftarrow{m}_{u_{k-1}, u_k}(u)$ and $N(u) = \overleftarrow{m}_{u_{k-1}, u_k}(u)$.

Note that if J is finite then the construction terminates at $J + 1$. Otherwise the construction continues indefinitely. For j even with $j \geq 2$ let $\mathcal{P}_E(j,(u_0,u_1,\ldots,u_j))$ be the statement

- 1. $(u_0, \ldots u_j) \in \mathcal{U}_j;$
- 2. $\overleftarrow{m}_{u_{j-2},u_{j-1}}(u_j) < a_1;$
- 3. N (respectively, M) is alternately increasing (respectively, decreasing) of order j with respect to $(u_0, \ldots u_i);$

4.
$$
S^{\nu}(\mu_{u_{j-1}} - \mu_{u_j}) = \nu|_{(M(u_j), N(u_j))};
$$

5. if
$$
j \leq J
$$
, $N(u_j) := \overleftarrow{n}_{u_{j-2},u_{j-1}}(u_j) > \overleftarrow{G}_{\mu}(u_{j-1})$ and $\overleftarrow{G}_{\mu}(u_j-) \leq M(u_j) = \overleftarrow{m}_{u_{j-2},u_{j-1}}(u_j) \leq \overleftarrow{G}_{\mu}(u_j)$.

For j odd let $\mathcal{P}_O(j,(u_0,u_1,\ldots,u_i))$ be the statement (to cover $j=1$ we define $u_{-1} := u_0$

- 1. $(u_0, \ldots u_j) \in \mathcal{U}_j;$
- 2. if $j \geq 3$, $\overleftarrow{m}_{u_{j-3},u_{j-2}}(u_{j-1}) < a_1$;
- 3. N (respectively, M) is alternately increasing (respectively, decreasing) of order j with respect to $(u_0, \ldots u_i);$

4.
$$
S^{\nu}(\mu_{u_j} - \mu_{u_{j-1}}) = \nu|_{(M(u_j), N(u_j))};
$$

5. if
$$
j \leq J
$$
, $M(u_j) := \overrightarrow{m}_{u_{j-1}, u_{j-2}}(u_j) < \overrightarrow{G}_{\mu}(u_{j-1})$ and $\overrightarrow{G}_{\mu}(u_j) \leq N(u_j) = \overrightarrow{n}_{u_{j-1}, u_{j-2}}(u_j) \leq \overrightarrow{G}_{\mu}(u_j+),$

Lemma 4.10. Suppose $j \geq 2$ is even, $j \leq J$, and $\mathcal{P}_E(j, (u_0, u_1, \ldots, u_j))$ holds. Then $\overrightarrow{n}_{u_j, u_{j-1}}(u_{j-1}) = \overleftarrow{n}_{u_{j-2}, u_{j-1}}(u_j)$ and $\overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1}) = \overleftarrow{m}_{u_{j-2}, u_{j-1}}(u_j)$.

Suppose $j \ge 1$ is odd, $j \le J$, and $\mathcal{P}_O(j,(u_0,u_1,\ldots,u_j))$ holds. Then for $j \ge 3$, $\overleftarrow{m}_{u_{j-1},u_j}(u_{j-1}) = \overrightarrow{m}_{u_{j-1},u_{j-2}}(u_j)$ and $\overleftarrow{n}_{u_{j-1},u_j}(u_{j-1}) = \overrightarrow{n}_{u_{j-1},u_{j-2}}(u_j)$. For $j = 1$, $\overleftarrow{m}_{u_0,u_1}(u_0) = \overrightarrow{m}_{u_0,u_0}(u_1)$ and $\overleftarrow{h}_{u_0,u_1}(u_0) = \overrightarrow{n}_{u_0,u_0}(u_1)$.

Proof. Suppose $j \geq 2$ is even. By hypothesis,

$$
\overleftarrow{m}_{u_{j-2},u_{j-1}}(u_j) \leq \overleftarrow{G}_{\mu}(u_j) \leq \overrightarrow{G}_{\mu}(u_{j-1}) < \overleftarrow{n}_{u_{j-2},u_{j-1}}(u_j).
$$

Note that $\mathcal{E}^c_{u_j, u_{j-1}}$ is linear on $[\overline{m}_{u_{j-2}, u_{j-1}}(u_j), \overleftarrow{n}_{u_{j-2}, u_{j-1}}(u_j)]$. But $[\overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1}), \overrightarrow{n}_{u_j, u_{j-1}}(u_{j-1})]$ is the largest interval containing $\overrightarrow{G}_{\mu}(u_{j-1})$ on which $\mathcal{E}^c_{u_j, u_{j-1}}$ is linear, and therefore

$$
\overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1}) \leq \overleftarrow{m}_{u_{j-2}, u_{j-1}}(u_j) < \overleftarrow{n}_{u_{j-2}, u_{j-1}}(u_j) \leq \overrightarrow{n}_{u_j, u_{j-1}}(u_{j-1}).
$$

Since $(\mathcal{E}_{u_j,u_{j-1}}^c)'$ is continuous everywhere and constant on $[\overrightarrow{m}_{u_j,u_{j-1}}(u_{j-1}), \overrightarrow{n}_{u_j,u_{j-1}}(u_{j-1})],$ and since $\overleftrightarrow{G}_{\mu}(u_j)$ and $\overrightarrow{G}_{\mu}(u_{j-1})$ lie in this interval, we have that $\overrightarrow{L}_{\mathcal{E}_{\mathcal{E}_{\mu}}^{c}}(u_{j-1}),(\mathcal{E}_{u_j,u_{j-1}}^{c})'(\overrightarrow{G}_{\mu}(u_{j-1})-)$ $\varepsilon_{u_j,u_{j-1}}^c$ = $\varepsilon_{u_j,u_{j-1}}^c$ = $\varepsilon_{u_j,u_{j-1}}^c$ = L $\overleftarrow{G}_{\mu}(u_j),({\mathcal E}^c_{u_j,u_{j-1}})'(\overleftarrow{G}_{\mu}(u_j)-)$ $\varepsilon_{u_j, u_{j-1}}^c$. Then

$$
\overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1}) = X_{\mathcal{E}_{u_j, u_{j-1}}}(\overrightarrow{G}_{\mu}(u_{j-1})) = X_{\mathcal{E}_{u_j, u_{j-1}}}(\overleftarrow{G}_{\mu}(u_j)) = \overleftarrow{m}_{u_{j-2}, u_{j-1}}(u_j)
$$

and

$$
\overleftarrow{n}_{u_{j-2},u_{j-1}}(u_j) = Z^+_{\mathcal{E}_{u_j,u_{j-1}}}(\overleftarrow{G}_{\mu}(u_j)) = Z^+_{\mathcal{E}_{u_j,u_{j-1}}}(\overrightarrow{G}_{\mu}(u_{j-1})) = \overrightarrow{n}_{u_j,u_{j-1}}(u_{j-1}).
$$

 \Box

The result for j odd follows symmetrically.

Proposition 4.11. Suppose $(\mu, \nu) \in \mathcal{K}_R$. Suppose j is even, $2 \leq j \leq J$ and $(u_0, u_1, \ldots, u_j) \in$ \mathcal{U}_j . Suppose $\mathcal{P}_E(j,(u_0,u_1,\ldots,u_j))$ holds. Then, $(u_0,u_1,\ldots,u_j,\overline{w}_{u_j,u_{j-1}}) \in \mathcal{U}_{j+1}$ and $\mathcal{P}_O(j+1, (u_0, u_1, \ldots, u_j, \overline{w}_{u_j, u_{j-1}}))$ holds.

Proof. Since $j \leq J$, if $u_j = 0$ then $u_{j-1} < \mu(\mathbb{R})$.

Set $u_{j+1} = \overline{w}_{u_j, u_{j-1}}$. Since $\mathcal{P}_E(j, (u_0, u_1, \ldots, u_j))$ holds $N(u_j) = \overleftarrow{n}_{\frac{u_{j-2}, u_{j-1}}{u_j}}(u_j)$ $\vec{G}_{\mu}(u_{j-1}+)$ and then by Lemma [4.10,](#page-24-0) $\vec{n}_{u_j, u_{j-1}}(u_{j-1}) = \vec{n}_{u_{j-2}, u_{j-1}}(u_j) > \vec{G}_{\mu}(u_{j-1}+)$. It follows from Lemma [3.10](#page-13-2) that $u_{j+1} = \overline{w}_{u_j, u_{j-1}} > u_{j-1}$ and hence that $(u_0, u_1, \ldots, u_j, \overline{w}_{u_j, u_{j-1}}) \in$ U_{i+1} .

The condition that $\overleftarrow{m}_{u_{j-2},u_{j-1}}(u_j) < a_1$ is inherited directly from the inductive hypothesis.

Since $\vec{G}_{\mu}(u_{j-1}+) < \vec{n}_{u_j,u_{j-1}}(u_{j-1}),$ by Corollary [3.12](#page-15-0) we have $\vec{m}_{u_j,u_{j-1}}(u_{j-1}+) \leq$ $\overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1})$ and then by Lemma [4.10](#page-24-0) and the inductive hypothesis,

$$
\overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1}+) \leq \overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1}) = \overleftarrow{m}_{u_{j-2}, u_{j-1}}(u_j) \leq \overleftarrow{G}_{\mu}(u_j) = \overrightarrow{G}_{\mu}(u_j+). \tag{4}
$$

Then by Corollary [3.14,](#page-17-2) $\overrightarrow{n}_{u_j, u_{j-1}}$ is strictly increasing on $[u_{j-1}, u_{j+1}]$. Moreover, $\overrightarrow{n}_{u_j, u_{j-1}}(u_{j-1}) =$ $\overleftarrow{n}_{u_{j-2},u_{j-1}}(u_j)$ and N is alternately increasing of order j with respect to $(u_0, u_1, \ldots u_j)$. Using these facts we now show that N is alternately increasing of order $j + 1$ with respect to $(u_0, u_1, \ldots u_{i+1})$.

Since N is alternately increasing of order j it is sufficient to show that for $u_{i-1} \leq$ $v < w \leq u_{j+1}$ and for $u_j \leq v \leq u_{j-1} < w \leq u_{j+1}$ we have that $N(v) < N(w)$. But, for $u_{j-1} \leq v \leq w \leq u_{j+1}$ the result follows from the strict monotonicity of $\overrightarrow{n}_{u_i,u_{i-1}}$ whence $N(v) = \overrightarrow{n}_{u_j, u_{j-1}}(v) < \overrightarrow{n}_{u_j, u_{j-1}}(w) = N(w)$. Similarly, for $u_j \le v \le u_{j-1}$ $w \le u_{j+1}$ we have $N(v) \le N(u_j) = \overleftarrow{n}_{u_{j-2}, u_{j-1}}(u_j) = \overrightarrow{n}_{u_j, u_{j-1}}(u_{j-1}) < N(w)$, where the first inequality follows from the fact that N is alternately increasing of order j and the second inequality follows from the monotonicity of $\overrightarrow{n}_{u_j, u_{j-1}}$ on $[u_{j-1}, u_{j+1}]$.

The proof that M is alternately decreasing is similar.

The fact that $S^{\nu}(\mu_{u_{j+1}} - \mu_{u_j}) = \nu_{(M(u_{j+1}),N(u_{j+1}))}$ follows from Lemma [3.13](#page-15-1) where we use the result $\vec{m}_{u_j, u_{j-1}}(u_{j-1}+) \leq \vec{G}_{\mu}(u_j+)$ derived in [\(4\)](#page-25-0) to verify the hypotheses of the lemma.

Suppose $u_j = 0$. By the second paragraph of the proof, $u_{j+1} = \overline{w}_{u_i, u_{i-1}} > u_{j-1}$. Moreover,

$$
\overrightarrow{m}_{0=u_j,u_{j-1}}(u_{j+1}) < \overrightarrow{m}_{0=u_j,u_{j-1}}(u_{j-1}) = \overleftarrow{m}_{u_{j-2},u_{j-1}}(0) \leq \overleftarrow{G}_{\mu}(0) = \overrightarrow{G}_{\mu}(0+).
$$

Then by Lemma [4.5,](#page-21-0) $u_{j+1} = \mu(\mathbb{R})$. Then $\mathcal{P}_O(j+1, (u_0, u_1, \ldots, u_j = 0, u_{j+1} = \mu(\mathbb{R}))$ holds. Since $j = J$ we do not have to check the final statement of $\mathcal{P}_O(j,(u_0, u_1, \ldots, u_{j+1})).$

Now suppose $u_j > 0$. In order to show that $\mathcal{P}_O(j+1, (u_0, u_1, \ldots, u_j, \overline{\underline{w}}_{u_j, u_{j-1}}))$ holds it only remains to show that $\overrightarrow{G}_{\mu}(u_{j+1}) \leq N(u_{j+1}) = \overrightarrow{n}_{u_j, u_{j-1}}(u_{j+1}) \leq \overrightarrow{G}_{\mu}(u_{j+1}+)$ and

that $M(u_{j+1}) = \overrightarrow{m}_{u_j, u_{j-1}}(u_{j+1}) < \overrightarrow{G}_{\mu}(u_j)$. The inequalities for N follow immediately from Lemma [3.10.](#page-13-2) For M we have using strict monotonicity of $\vec{m}_{u_j, u_{j-1}}$, Lemma [4.10,](#page-24-0) the inductive statement and Lemma [4.4](#page-20-2)

$$
\overrightarrow{m}_{u_j, u_{j-1}}(u_{j+1}) < \overrightarrow{m}_{u_j, u_{j-1}}(u_{j-1}) = \overleftarrow{m}_{u_{j-2}, u_{j-1}}(u_j) \leq \overleftarrow{G}_{\mu}(u_j) = \overrightarrow{G}_{\mu}(u_j+).
$$

Then, by Lemma [4.4,](#page-20-2) $M(u_{j+1}) = \overrightarrow{m}_{u_j, u_{j-1}}(u_{j+1}) < \overrightarrow{G}_{\mu}(u_j)$ where we use the fact that $\overrightarrow{m}_{u_j, u_{j-1}}(u_{j+1}) < a_1,$

The following lemma, is the parallel result for odd i .

Proposition 4.12. Suppose $(\mu, \nu) \in \mathcal{K}_R$. Suppose j is odd, $1 \leq j \leq J$ and $(u_0, u_1, \ldots, u_j) \in$ \mathcal{U}_j . Suppose $\mathcal{P}_O(j,(u_0,u_1,\ldots,u_j))$ holds, and if $j\geq 3$, $\mathcal{P}_E(j-1,(u_0,u_1,\ldots,u_{j-1})$ holds. Then, $(u_0, u_1, \ldots, u_j, \underline{w}_{u_{j-1},u_j}) \in \mathcal{U}_{j+1}$ and $\mathcal{P}_E(j+1, (u_0, u_1, \ldots, u_j, \underline{w}_{u_{j-1},u_j}))$ holds.

 \Box

Proof. Mutatis mutandis, the majority of the proof is identical. The only exception is that it is now additionally necessary to show that $\overleftarrow{m}_{u_{j-1}u_j}(u_{j+1}) < a_1$.

Suppose $j \ge 3$. Then, by strict monotonicity of $\overleftarrow{m}_{u_{j-1},u_j}$ and $\overrightarrow{m}_{u_{j-1}u_{j-2}}$, Lemma [4.10](#page-24-0) (twice) and the fact that we are assuming that $\mathcal{P}_E(j-1,(u_0,u_1,\ldots,u_{j-1})$ holds,

$$
\overleftarrow{m}_{u_{j-1},u_j}(u_{j+1}) < \overleftarrow{m}_{u_{j-1},u_j}(u_{j-1}) = \overrightarrow{m}_{u_{j-1},u_{j-2}}(u_j) < \overrightarrow{m}_{u_{j-1},u_{j-2}}(u_{j-2}) = \overrightarrow{m}_{u_{j-3},u_{j-2}}(u_{j-1}) < a_1.
$$

The remaining case is when $j = 1$. Note that if $u_0 = 0$ then $J = 0$, a contradiction to $j \leq J$ and hence we must have $u_0 > 0$. We are required to show that whenever $u_0 > 0$ we have $\overleftarrow{m}_{u_0, u_1}(u_2) < a_1$. Since $(\mu, \nu) \in \mathcal{K}_R$, \mathcal{E}_{u_0, u_1} lies below the tangent L_{a_1} (to D at a_1) to the right of a_0 . Now suppose $u_1 > u_0 > 0$ is fixed and consider $(\mathcal{E}_{u,u_1})_{u \leq \underline{u_0}}$ and D. Note that $\mathcal{E}_{u_2,u_1} = D$ on $\left[\overleftarrow{G}_{\mu}(u_2 -) = \overrightarrow{G}_{\mu}(u_2), \overleftarrow{G}_{\mu}(u_2)\right]$. We have that $D'(\overleftarrow{G}_{\mu}(u_2-)) \leq D'(\overleftarrow{H}_{u_0, u_1}(u_2))) \leq D'(\overleftarrow{G}_{\mu}(u_2))$. But then $L_{\overleftarrow{H}_{u_0, u_1}(u_2)}$ has slope less than $D'(a_0) = D'(a_1)$: if not then $L_{\overleftarrow{m}_{\nu_0, u_1}(u_2)}$ cannot touch \mathcal{E}_{u_2, u_1} to the right of a_0 . In particular, $\overleftarrow{m}_{u_0, u_1}(u_2) < a_1$ and then $\overrightarrow{G}_{\mu}(u_2) \leq \overleftarrow{m}_{u_0, u_1}(u_2) < a_1$. □

It remains to show that the statement $\mathcal{P}_O(1,(u_0,u_1))$ holds. However, there is no guarantee that $u_1 = \overline{w}_{u_0, u_0} > u_0$. Hence we introduce:

Definition 4.13. $\mathcal{K}_{*} = \{(\mu, \nu) \in \mathcal{K}_{R} : \overline{w}_{u_0, u_0} > u_0\}.$

Proposition 4.14. Suppose $(\mu, \nu) \in \mathcal{K}_{*}$. Then $\mathcal{P}_O(1, (u_0, u_1 = \overline{w}_{u_0, u_0}))$ holds.

Proof. The proof is a simplified version of the proof of Proposition [4.11.](#page-24-1)

Since $\overline{w}_{u_0,u_0} > u_0$ by hypothesis it follows that $(u_0, u_1) \in \mathcal{U}_1$. Also, by Lemma [3.13](#page-15-1) in the case $v = u$, $S^{\mu}(\mu_{u_1} - \mu_{u_0}) = \nu|_{(M(u_1), N(u_1))}$ and M (respectively N) is strictly decreasing (increasing) on $[u_0, u_1]$. Finally, by Lemma [3.10,](#page-13-2) $\overrightarrow{G}_{\mu}(\overline{w}_{u_0, u_0}) \leq \overrightarrow{n}_{u_0, u_0}(\overline{w}_{u_0, u_0}) \leq$ $\overrightarrow{G}_{\mu}(\overline{w}_{u_0, u_0}+)$, whereas by definition $M(u_1) = \overrightarrow{m}_{u_0, u_0}(u_1) < \overrightarrow{m}_{u_0, u_0}(u_0) \le \overrightarrow{G}_{\mu}(u_0)$. \Box **Theorem 4.15.** Suppose $(\mu, \nu) \in \mathcal{K}_*$. Then the pair (M, N) defines an injective martingale coupling.

Proof. By Proposition [4.14,](#page-26-0) $\mathcal{P}_O(1,(u_0,u_1))$ holds. Then, alternating between Proposi-tions [4.12](#page-26-1) and [4.11](#page-24-1) it follows that for each $j \leq J+1$, the statement $\mathcal{P}_E(j,(u_0,u_1,\ldots u_j))$ holds if j is even and $\mathcal{P}_O(j,(u_0, u_1, \ldots u_i))$ holds if j is odd.

It remains to deduce that the pair (M, N) defines an injective lifted martingale coupling of μ and ν . If $J < \infty$ then the construction terminates: if J is even then $u_J = 0$ and $u_{J+1} = \mu(\mathbb{R})$; if J is odd then $u_J = \mu(\mathbb{R})$ and $u_{J+1} = 0$. If $J = \infty$, then define $u_{\infty} = \lim_{k \to \infty} u_{2k}$ and $u^{\infty} = \lim_{k \to \infty} u_{2k+1}$. It follows that $0 \le u_{\infty} < u_0 < u^{\infty} \le \mu(\mathbb{R})$. It remains to show that if $J = \infty$ then $u_{\infty} = 0$, $u^{\infty} = \mu(\mathbb{R})$ and that (M, N) define a coupling of μ to ν . If this is the case then (M, N) will define an injective coupling by the fact that M and N are alternately decreasing and increasing respectively and Remark [4.9.](#page-23-0)

We first show that if $J = \infty$ then $(u_{\infty}, u^{\infty}) = (0, \mu(\mathbb{R}))$. The idea is that if the assertion does not hold, then $b = \lim_{k \to \infty} M(u_{2k})$ and $c = \lim_{k \to \infty} N(u_{2k+1})$ violate the assumption that $(\mu, \nu) \in \mathcal{K}_R$.

Define $m_{\infty} = \lim_{k \uparrow \infty} M(u_{2k}), n_{\infty} = \lim_{k \uparrow \infty} N(u_{2k}), m^{\infty} = \lim_{k \uparrow \infty} M(u_{2k+1}), n^{\infty} =$ $\lim_{k \uparrow \infty} N(u_{2k+1}),$ and note that $M(u_{2k}) \geq M(u_{2k+1}) \geq M(u_{2k+2})$ and $N(u_{2k}) \leq N(u_{2k+1}) \leq N(u_{2k+2})$ so that $m_{\infty} = m^{\infty}$ and $n_{\infty} = n^{\infty}$. Furthermore, $M(u_{2k+1}) < \overrightarrow{G}_{\mu}(u_{2k}) = \overleftarrow{G}_{\mu}(u_{2k-}) \le$ $M(u_{2k}) \leq \overleftarrow{G}_{\mu}(u_{2k})$ so that $m_{\infty} = m^{\infty} = \overleftarrow{G}_{\mu}(u_{\infty})$. Similarly we obtain that $n_{\infty} =$ $n^{\infty} = \overrightarrow{G}_{\mu}(u^{\infty}).$

We have both $\lim_{k\to\infty} \mathcal{E}_{u_{2k},u_{2k-1}}(M(u_{2k})) = \lim_{k\to\infty} D_{\mu,\nu}(M(u_{2k})) = D_{\mu,\nu}(m_\infty)$ and $\lim_{k\to\infty} \mathcal{E}_{u_{2k},u_{2k-1}}(N(u_{2k})) = \mathcal{E}_{u_{\infty},u_{\infty}}(n_{\infty}) = D_{\mu,\nu}(n_{\infty}).$ Moreover, $\lim_{k\to\infty} D'_{\mu,\nu}(M(u_{2k})) =$ $\lim_{k \to \infty} {\mathcal{E}}'_{u_{2k}, u_{2k-1}}(M(u_{2k})) = \lim_{k \to \infty} P'_{\nu}(M(u_{2k})) - u_{2k} = P'_{\nu}(m_{\infty}) - u_{\infty} = {\mathcal{E}}'_{u_{\infty}, u_{\infty}}(m_{\infty}) =$ $D'_{\mu,\nu}(m_{\infty})$. Similarly, $\lim_{k\to\infty} (\mathcal{E}_{u_{2k},u_{2k-1}})'(N(u_{2k})) = \lim_{k\to\infty} {P'_{\nu}(N(u_{2k})) - u_{2k-1}} =$ $P'_\nu(n_\infty) - u^\infty = D'_{\mu,\nu}(n_\infty)$. On the other hand, by construction of N, M, and using Lemma [3.21,](#page-17-3) Corollary [3.3](#page-10-0) and the properties of the convex hull, we have that

$$
\mathcal{E}'_{u_{2k},u_{2k-1}}(M(u_{2k})) = \mathcal{E}'_{u_{2k},u_{2k-1}}(N(u_{2k})) = \frac{\mathcal{E}_{u_{2k},u_{2k-1}}(N(u_{2k})) - \mathcal{E}_{u_{2k},u_{2k-1}}(M(u_{2k}))}{N(u_{2k}) - M(u_{2k})}.
$$

Since $\mathcal{E}_{u_{2k}, u_{2k-1}}(k) = D(k)$ for $k \in \{M(u_{2k}), N(u_{2k})\}$, we find

$$
D'_{\mu,\nu}(m_{\infty}) = \frac{D_{\mu,\nu}(n_{\infty}) - D_{\mu,\nu}(m_{\infty})}{n_{\infty} - m_{\infty}} = D'_{\mu,\nu}(n_{\infty} -),
$$

so that $L_{m_{\infty}}$ given by $L_{m_{\infty}}(x) = D_{\mu,\nu}(x) + (x - m_{\infty})D'_{\mu,\nu}(m_{\infty})$ agrees with $D_{\mu,\nu}$ at n_{∞} . Moreover, $\mathcal{E}_{u_{2k},u_{2k-1}} > L_{\mathcal{E}_{u_{2k},u_{2k-1}}}^{M(u_{2k}),D'_{\mu,\nu}(M(u_{2k}))}$ $\epsilon_{u_{2k},u_{2k-1}}^{u_{1}(u_{2k}),\mu_{\mu,\nu}(u_{1}(u_{2k}))}$ on $(M(u_{2k}),N(u_{2k}))$. Since, for any version G_μ of the quantile function of μ , we have that $\mathcal{E}_{u_{2k},u_{2k-1}} = D_{\mu,\nu}$ on

 $[G_\mu(u_{2k}), G_\mu(u_{2k-1}+)]$, it follows that $D_{\mu,\nu} > L_{\mathcal{E}_{\text{max}}(u_{2k-1})}^{M(u_{2k}),D'_{\mu,\nu}(M(u_{2k}))}$ $\epsilon_{\mu_2\mu_3,\mu_2\mu_{k-1}}^{u_1(u_2k),\nu_{\mu,\nu}(m_1(u_2k))}$ on $(G_\mu(u_{2k}), G\mu(u_{2k-1}+)).$ Letting $k \uparrow \infty$ we conclude $D_{\mu,\nu} > L_{\mathcal{E}_{u_{\infty},u_{\infty}}}^{m_{\infty},D'_{\mu,\nu}(m_{\infty})} \equiv L_{D_{\mu,\nu}}^{m_{\infty},D'_{\mu,\nu}(m_{\infty})}$ $\sum_{\mu,\nu}^{m_{\infty},\nu}$ $\mu,\nu}^{m_{\infty},\nu}$ on (m_{∞},n_{∞}) .

It follows that with $b = m_{\infty}$ (and supposing $m_{\infty} > \alpha_{\nu}$) the corresponding c in Definition [4.2](#page-19-0) is n_{∞} . But then $D'_{\mu,\nu}(m_{\infty}) = D'_{\mu,\nu}(n_{\infty})$ contradicting the fact that $(\mu, \nu) \in \mathcal{K}_R$. Thus it follows that $m_{\infty} = \alpha_{\nu}$. Then also $u_{\infty} = 0$, $D(n_{\infty}) = \beta_{\nu}$ and $u^{\infty} = \mu(\mathbb{R})$, as claimed.

It is left to show that (M, N) define a lifted martingale coupling of μ and ν . Recall the definition of ϕ in Definition [4.7.](#page-21-1) Define $G : [0, \mu(\mathbb{R})] \to \mathbb{R}$ by

$$
\tilde{G}(v) = \begin{cases} \overrightarrow{G}_{\mu}(\phi(v)), & \text{if } j(v) \text{ is even}, \\ \overleftarrow{G}_{\mu}(\phi(v)), & \text{if } j(v) \text{ is odd}, \end{cases} v \in [0, \mu(\mathbb{R})].
$$

Let $\tilde{\mu}$ be a measure on $[0,1] \times \mathbb{R}$ defined by $\tilde{\mu}(du, dx) = du \delta_{\tilde{G}(u)}(dx)$. It is easy to see that, by construction, the first and second marginals of $\tilde{\mu}$ are λ and μ , respectively.

Now, by Beiglböck and Juillet $[3,$ Theorem 2.1, there exists the unique lifted martingale coupling $\tilde{\pi}$ (a measure on $[0, \mu(\mathbb{R})] \times \mathbb{R} \times \mathbb{R}$) of μ and ν , that, for each $u \in [0, \mu(\mathbb{R})]$, embeds $\tilde{\mu}_{[0,u]} := \int_0^u \delta_{\tilde{G}(v)} dv$ into $S^{\nu}(\tilde{\mu}_{[0,u]})$. More precisely,

$$
\tilde{\pi}(du, dx, dy) = du \delta_{\tilde{G}(u)}(dx) \chi_{\tilde{R}(u), \tilde{G}(u), \tilde{S}(u)},
$$

where

$$
\chi_{\tilde{R}(u),\tilde{G}(u),\tilde{S}(u)} = \begin{cases} \delta_{\tilde{G}(u)}, & \text{if } \tilde{G}(u) \in \text{supp}(\nu - S^{\nu}(\tilde{\mu}_{[0,u]})),\\ \frac{\tilde{S}(u)-\tilde{G}(u)}{\tilde{S}(u)-\tilde{R}(u)}\delta_{\tilde{R}(u)} + \frac{\tilde{G}(u)-\tilde{R}(u)}{\tilde{S}(u)-\tilde{R}(u)}\delta_{\tilde{S}(u)}, & \text{otherwise}, \end{cases}
$$

and $\tilde{R}(u) = \sup\{k \in \text{supp}(\nu - S^{\nu}(\tilde{\mu}_{[0,u]})) \cap (-\infty, \tilde{G}(u)]\}, \ \tilde{S}(u) = \inf\{k \in \text{supp}(\nu S^{\nu}(\tilde{\mu}_{[0,u]})) \cap [\tilde{G}(u),\infty)\}.$

Fix $\tilde{u} \in [0, \mu(\mathbb{R})] \setminus V$. Suppose that $j(\tilde{u})$ is even (the case when $j(\tilde{u})$ is odd follows by symmetry). Then

$$
u_{j(\tilde{u})} < u_{j(\tilde{u})} + \tilde{u} < u_{j(\tilde{u})+1}.
$$

Then $\tilde{G}(\tilde{u}) = \overrightarrow{G}_{\mu}(u_{j(\tilde{u})} + \tilde{u})$ and $\tilde{\mu}_{[0,\tilde{u}]} = \mu_{u_{j(\tilde{u})} + \tilde{u}} - \mu_{u_{j(\tilde{u})}}$. It follows from Lemma [3.13\(](#page-15-1)i) that

$$
S^{\nu}(\tilde{\mu}_{[0,\tilde{u}]}) = S^{\nu}(\mu_{u_{j(\tilde{u})} + \tilde{u}} - \mu_{u_{j(\tilde{u})}}) = \nu|_{(M(u_{j(\tilde{u})} + \tilde{u}), N(u_{j(\tilde{u})} + \tilde{u}))},
$$

and therefore $R(\tilde{u}) = M(u_{j(\tilde{u})} + \tilde{u})$ and $S(\tilde{u}) = N(u_{j(\tilde{u})} + \tilde{u})$.

We conclude that, for each even $j \geq 0$, and $u \in [0, \mu(\mathbb{R})]$ with $u_{j-1} < u_j + u < u_j+1$ (or $u_0 < u_0 + u < u_1$ in the case $j = 0$), we have that

$$
\chi_{\tilde{R}(u),\tilde{G}(u),\tilde{S}(u)} = \frac{N(u_j+u) - \overrightarrow{G}_{\mu}(u_j+u)}{N(u_j+u) - M(u_j+u)} \delta_{M(u_j+u)} + \frac{\overrightarrow{G}_{\mu}(u_j+u) - M(u_j+u)}{N(u_j+u) - M(u_j+u)} \delta_{N(u_j+u)}.
$$

Similarly, for each odd $j \geq 1$, and $u \in [0, \mu(\mathbb{R})] \setminus \mathbb{V}$ with $u_{j+1} < u_j - u < u_{j-1}$, we have that

$$
\chi_{\tilde{R}(u),\tilde{G}(u),\tilde{S}(u)} = \frac{N(u_j - u) - \overleftarrow{G}_{\mu}(u_j - u)}{N(u_j - u) - M(u_j - u)} \delta_{M(u_j - u)} + \frac{\overleftarrow{G}_{\mu}(u_j - u) - M(u_j - u)}{N(u_j - u) - M(u_j - u)} \delta_{N(u_j - u)}.
$$

 \Box

Since V is countable, it follows that (M, N) induces the coupling $\tilde{\pi}$.

Example 4.16. Suppose $\mu \sim U[-1, 1]$ and $\nu \sim U[-2, 2]$. Then $G_{\mu} : [0, 1] \mapsto [-1, 1]$ is given by $G_{\mu}(u) = 2u - 1$ and $u_0 = 0$. Simple calculations show that

$$
\mathcal{E}_{0,u}(x) = \begin{cases} \frac{(2+x)^2}{8} & -2 < x \le -1\\ \frac{2-x^2}{8} & -1 < x \le 2u - 1\\ \frac{(2+x)^2}{8} - u(1+x-u) & 2u - 1 < x < 2. \end{cases}
$$

Solving $\mathcal{E}'_{0,u}(m) = \mathcal{E}'_{0,u}(n)$ for $-2 < m < -1 < 2u - 1 < n$ we find $M(u) = N(u) - 4u$. Then solving $\mathcal{E}'_{0,u}(n) = \frac{\mathcal{E}_{0,u}(n) - \mathcal{E}_{0,u}(m)}{n-m}$ together with $n-m=4u$ we find $N(u) = 3u - 1$ and $M(u) = -u - 1$.

Define $\hat{\pi}_{u.G.(u)}^{Q,M,N}$ $u_{u,G_\mu(u)}^{Q,M,N}(dy) = \frac{1}{4}\delta_{-u-1}(dy) + \frac{3}{4}\delta_{3u-1}(dy)$. We have that $\int_{u \in [0,1]} du \delta_{2u-1}(dx) =$ 1 $\frac{1}{2} I_{\{|x| \leq 1\}} dx = \mu(dx)$ and considering $y < -1$ and $y > -1$ separately, $\int_{u \in [0,1]} du \delta_{2u-1}(dx) \hat{\pi}_{u,G_\mu(u)}^{Q,M,N}$ $\omega_{u,G_\mu(u)}^{Q,M,N}(dy)=$ 1 $\frac{1}{4}I_{\{|y|\leq2\}}dy=\nu(dy)$. Hence $\hat{\pi}^{Q,M,N}$ defined by $\hat{\pi}^{Q,M,N}(du,dx,dy)=du\delta_{G_{\mu}(u)}(dx)\hat{\pi}^{Q,M,N}_{u,G_{\mu}(u)}$ $\int_{u,G_\mu(u)}^{Q,M,N}(dy)$ is a quantile-lifted, martingale coupling of μ and ν .

Observe that for this example $N(u) > G_{\mu}(u+)$ on $(0,1)$ and so $u_1 = 1$. Also $\lim_{u \uparrow 1} M(u) = -2 = \alpha_{\nu}$ and $\lim_{u \uparrow 1} N(u) = 2 = \beta_{\nu}$.

Example 4.17. Consider the case where $\mu = \frac{1}{2}$ $\frac{1}{2}U(-1,1)+\frac{1}{2}U(-2,2)$ and $\nu=U(-2,2)$. Then $a_0 = -1$. Also $G_{\mu}(u) = -2 + 8u$ for $0 < u \le \frac{1}{8}$ $\frac{1}{8}$; $G_{\mu}(u) = \frac{8}{3} \left(u - \frac{1}{2}\right)$ $(\frac{1}{2})$ for $\frac{1}{8} < u \leq \frac{7}{8}$ $\frac{7}{8}$; $G_{\mu}(u) = 8u - 6$ for $\frac{7}{8} < u < 1$. Hence $u_0 = \frac{1}{8}$ $\frac{1}{8}$. Also $D_{\mu,\nu}(x) = \frac{(x+2)^2}{16}$ for $-2 < x \le -1$; $D_{\mu,\nu}(x) = \frac{2-x^2}{16}$ for $-1 < x \le 1$; $D_{\mu,\nu}(x) = \frac{(\tilde{2}-x)^2}{16}$ for $1 < x < 2$.

Consider $\mathcal{E}_{u_0=\frac{1}{8},u}$. For any $u \geq u_0$, we find for $-2 \leq x \leq -1$, $\mathcal{E}_{\frac{1}{8},u}(x) = \frac{1}{16}(5+6x+$ 2x²), and for $-1 < x \le G_{\mu}(u)$, $\mathcal{E}_{\frac{1}{8},u}(x) = D_{\mu,\nu}(x)$.

For $\frac{1}{8} < u < \frac{7}{8}$ and $G_{\mu}(u) < x \leq 2$ we have $\mathcal{E}_{\frac{1}{8},u}(x) = \frac{x^2}{8} - (u - \frac{1}{2})$ $(\frac{1}{2})x + \frac{4}{3}$ $rac{4}{3}(u-\frac{1}{2})$ $(\frac{1}{2})^2 + \frac{1}{8}$ $\frac{1}{8}$. Then for $\frac{1}{8} < u < \frac{7}{8}$ we can solve for $N(u) \geq G_{\mu}(u)$ and $M(u) < -1$ using

$$
\mathcal{E}'_{\frac{1}{8},u}(M(u)) = \mathcal{E}'_{\frac{1}{8},u}(N(u)) = \frac{\mathcal{E}_{\frac{1}{8},u}(N(u)) - \mathcal{E}_{\frac{1}{8},u}(M(u))}{N(u) - M(u)}.
$$
(5)

These equations have solution $M(u) = -\frac{11}{12} - \frac{2u}{3}$ $\frac{2u}{3}$, $N(u) = \frac{10u}{3} - \frac{17}{12}$. In particular $M(\frac{7}{8})$ $\frac{7}{8}$) = $-\frac{3}{2}$ $rac{3}{2}$ and $N(\frac{7}{8})$ $\frac{7}{8}$) = $\frac{3}{2}$ and in the very first part of the construction we have that the mass in $[-1, 1)$ is mapped to $(-3/2, 3/2)$.

Since $N > G_{\mu}$ on $\left(\frac{1}{8}\right)$ $\frac{1}{8}, \frac{7}{8}$ $(\frac{7}{8})$ we have that $u_1 > \frac{7}{8}$ $\frac{7}{8}$.

For $u > \frac{7}{8}$ and $G_{\mu}(u) < x \leq 2$, $\mathcal{E}_{\frac{1}{8},u}(x) = \frac{x^2}{8} - (u - \frac{1}{2})$ the same equations as in we find that $N(u) = \frac{48u^2 - 52u + \frac{71}{4}}{8u - 1}$ and $M(u) = N(u) - 4u + \frac{1}{2}$ $(\frac{1}{2})x + 4u^2 - 6u + \frac{5}{2}$ $\frac{5}{2}$. Then, solving $\frac{1}{2}$, at least for $u \leq u_1$. In particular, solving $N(u) = G_{\mu}(u)$ we find that u_1 is the solution $in\(\frac{7}{8})$ $\frac{7}{8}$, 1) to $64u^2 - 16u - 47 = 0$.

Now we solve for $M(u)$ and $N(u)$ on the interval (u_2, u_0) where $u_2 \in (0, u_0)$ is to be found. For $v \le u_0 < u_1 \le u$ and $x \ge G_\mu(u)$ we find $\mathcal{E}_{v,u}(x) = \frac{x^2}{8} + x(\frac{1}{2} - u) +$ $4u^2 - 6u + \frac{5}{2}$ $\frac{5}{2}$. In order to find M and N on (u_2, u_0) we need to find \mathcal{E}_{v,u_1} for v such that $G(v) \in [M(u_1), a_0]$ and $G(v) \leq M(u_1)$, but, if the goal is purely to calculate u_2 , then it is sufficient to only consider $v \leq G_{\mu}^{-1}(M(u_1)) = \frac{1}{16}$. There we find $\mathcal{E}_{v,u_1} =$ $\frac{x^2}{8} + x\left(\frac{1}{2} - v\right) + 4v^2 - 2v + \frac{1}{2}$ $\frac{1}{2}$. Solving for M and N we find, on $v \leq \frac{1}{16}$, $N(v) =$ $M(v) + 4(u - v)$ where $M(v) = \frac{2u_1^2 + 4u_1v - 6u_1 - 6v^2 + 2v + 2}{(u_1 - v)}$ $\frac{-\alpha u_1 - \alpha v + 2v + 2}{(u_1 - v)}$. Solving for $G_\mu(v) = M(v)$ we find that $u_2 = u_1 - \sqrt{2u_1 - 1}$.

These arguments can be iterated. We find that $u_{2k+1} = u_{2k} + \sqrt{1 - u_{2k}}$ and $u_{2k+2} =$ $u_{2k+1} - \sqrt{2u_{2k+1} - 1}$. Clearly $u_{2k} \downarrow 0$ and $u_{2k+1} \uparrow 1$.

5 Reducing the problem to countably many intervals

The goal of this section is to explain how to divide the general problem with $(\mu, \nu) \in \mathcal{K}$ into countably many intervals, in such a way that if we can construct an injective mapping on each interval then we can construct an injective map overall.

To this end we use the left-curtain coupling introduced by Beiglböck and Juillet $[2]$, and studied further by Henry-Labordaire and Touzi [\[4\]](#page-38-8) and Hobson and Norgilas [\[8,](#page-38-9) [9\]](#page-38-7), although having defined the intervals using the left-curtain coupling we use a completely different construction to define the injective coupling, namely the construction of the previous section. Beiglböck and Juillet $[2]$ studied existence and uniqueness of the leftcurtain coupling and showed the construction was optimal for a class of Martingale Optimal Transport problems; Henry-Labordaire and Touzi [\[4\]](#page-38-8) extended the optimality to a wider class of problems and gave a constructive proof under certain regularity conditions on the measures and Hobson and Norgilas [\[8\]](#page-38-9) extended the construction to the case where μ and ν are general measures. Most relevantly for this work Hobson and Norgilas [\[9\]](#page-38-7) give a graphical representation of the construction in the general case.

Recall the definition of $\mathcal{E}_{0,u}$ which can be written as $\mathcal{E}_{0,u} = P_{\nu} - P_{\mu_u}$

Set $S(u) = Z_{\mathcal{E}_0}^{-1}$ $\overline{\mathcal{E}}_{\mathcal{E}_{0,u}}(\overrightarrow{G}_{\mu}(u))$ and $R(u) = X_{\mathcal{E}_{0,u}}^{-1}$ $\overline{\vec{c}_{0,u}}(\overrightarrow{G}_{\mu}(u))$. Note that the definition of the lower function $R(u)$ used by Hobson and Norgilas [\[9\]](#page-38-7) is slightly different from $X_{\mathcal{E}_0}^ \overline{\mathcal{E}}_{0,u}(\overrightarrow{G}_{\mu}(u))$, and is given by $\inf\{w : w \leq \overrightarrow{G}_{\mu}(u), D(w) = L^{z,(\mathcal{E}_{0,u}^c)/2}_{\mathcal{E}_{0,u}^c}\}$ $\varepsilon_{0,u}^{c}$ (*w*). However,

given that $\mathcal{E}_{0,u} = D$ on $(-\infty, \overrightarrow{G}_{\mu}(u+)]$, it is easy to see that two definitions coincide.

Theorem 5.1 (Hobson and Norgilas [\[9,](#page-38-7) Theorem 3.8]). Suppose $\mu \leq_{cx} \nu$. Define $\hat{\pi}^{Q,LC}_{\overrightarrow{C}}$ $\frac{Q, LC}{u, \overrightarrow{G}_{\mu}(u)}$ by

$$
\hat{\pi}_{u,\overrightarrow{G}_{\mu}(u)}^{Q,LC}(dy) = \frac{S(u) - \overrightarrow{G}_{\mu}(u)}{S(u) - R(u)} \delta_{R(u)}(dy) + \frac{\overrightarrow{G}_{\mu}(u) - R(u)}{S(u) - R(u)} \delta_{R(u)}(dy)
$$
(6)

on $R(u) < S(u)$ and $\hat{\pi}^{Q, LC}_{\vec{\sigma}}$ $\frac{Q,LC}{w,\overrightarrow{G}_{\mu}(u)}(dy) = \delta_{\overrightarrow{G}_{\mu}(u)}(dy)$ otherwise. Then $\hat{\pi}^{Q,LC}$ defined by $\hat{\pi}^{Q,LC}(du,dx,dy) = dy \delta_{\overrightarrow{G}_{\mu}(u)}(dx) \hat{\pi}^{Q,LC}_{u,\overrightarrow{G}_{u}}$ $u_{u,\overrightarrow{G}_{\mu}(u)}^{Q,LC} \in \mathcal{M}_{QL}(\mu,\nu)$. $\hat{\pi}^{Q,LC}$ is the left-curtain quantilelifted martingale coupling of μ and ν

Definition 5.2 (Hobson and Norgilas [\[9,](#page-38-7) Definition 3.5]). Given a left-continuous increasing function $g : (0, c) \mapsto \mathbb{R}$, a pair of functions $r, s : (0, c) \mapsto \mathbb{R}$ is said to be left-monotone with respect to g on $I = (0, c)$ if $r \leq g \leq s$ and if for $u < u'$ we have $s(u) \leq s(u')$ and $r(u') \notin (r(u), s(u))$.

Proposition 5.3 (Hobson and Norgilas [\[9,](#page-38-7) Theorem 4.8]). Let $R, S : (0, \mu(\mathbb{R})) \mapsto \mathbb{R}$ be defined as above by $R(u) = X_{\mathcal{E}_0}^{-1}$ $\overline{\vec{\varepsilon}_{0,u}}(\overrightarrow{G}_{\mu}(u))$ and $S(u)=Z_{\overline{\varepsilon}_{0}}^{-1}$ $\overline{\mathcal{E}}_{0,u}(\overrightarrow{G}_{\mu}(u))$. Then (R,S) is left monotone with respect to \overrightarrow{G}_{μ} on $\mathcal{I} = (0, \mu(\mathbb{R}))$.

- **Lemma 5.4** (Hobson and Norgilas [\[9,](#page-38-7) Lemma 4.1, Proposition 6.1]). 1. If $u \in (0, \mu(\mathbb{R}))$ is such that $S(u) > \overrightarrow{G}_{\mu}(u+)$ then $R(u) < \overrightarrow{G}_{\mu}(u)$.
	- 2. S is left-continuous and R satisfies $R(v) \leq \liminf_{u \uparrow v} R(u)$.

Under our assumption that ν is continuous we have a bit more structure.

Lemma 5.5. Suppose Standing Assumption [3.1](#page-9-1) holds. Then S is strictly increasing.

Proof. If S takes the value $\{y\}$ on an interval $(\underline{u}, \overline{u}] \subseteq (0, \mu(\mathbb{R}))$ then $\nu(\{y\}) \geq \int_{\underline{u}}^{\overline{u}}$ $\overrightarrow{G}_{\mu}(u)-R(u)$ $\frac{\sum_{\mu}(u)-R(u)}{S(u)-R(u)}du >$ 0, but this contradicts our standing assumption.

Since S is strictly increasing it has a continuous inverse S^{-1} .

Now want to partition into $[0, \mu(\mathbb{R})]$ into disjoint intervals. Let $\mathcal{A}_{\leq} = \{u : u \in$ $(0, \mu(\mathbb{R}))$, $\overrightarrow{G}_{\mu}(u+) < S(u)$. Then, see the end of Section 5 in [\[9\]](#page-38-7) for details (and especially Equation (2.7) in [\[9\]](#page-38-7)), A_{\leq} is a countable union of disjoint open intervals, and R is strictly decreasing on each of these intervals. Let \tilde{A}_k with endpoints $u_k < v_k$ be such an interval. Let $\beta_k = S(v_k)$ and $\alpha_k = \lim_{u \uparrow v_k} R(u)$. Let $A_k = (t_k := S^{-1}(\alpha_k), v_k =$ $S^{-1}(\beta_k)$). Note that $A_k = (t_k, u_k] \cup \tilde{A}_k$ and $\inf_{u \in \tilde{A}_k} R(u) = \inf_{u \in A_k} R(u) = \alpha_k$.

Lemma 5.6. Given $k \neq k'$ either $A_k \subsetneq A_{k'}$ or $A_{k'} \subsetneq A_k$ or $A_k \cap A_{k'} = \emptyset$.

Proof. Consider A_k , $A_{k'}$ for some $k \neq k'$. Note that we cannot have $A_k = A_{k'}$. Without loss of generality we may assume $v_{k'} < v_k$. If $v_{k'} \leq t_k$ then $A_k \cap A_{k'} = \emptyset$. If $t_k \leq t_{k'}$ then $A_{k'} \subseteq A_k$. Finally we show that the case $t_{k'} < t_k < v_{k'}$ cannot happen. Note that if $t_{k'} < t_k < v_{k'}$ then from the strict monotonicity of S we must have that $S(t_k) < S(v_{k'})$ and so $S(S^{-1}(\alpha_k)) < S(S^{-1}(\beta_{k'}))$ and hence $\alpha_k < \beta_{k'}$. We conclude that $R(\hat{u}) < \beta_{k'}$ for some $\hat{u} \in \tilde{A}_k$. Furthermore, since $t_{k'} < t_k$ and S^{-1} is non-decreasing, $\alpha_{k'} \leq \alpha_k$. But then, for all $u \in (\hat{u}, v_k)$, and using Part (ii) of Lemma [5.4](#page-31-0) for the first inequality,

$$
R(v_{k'}) \leq \alpha_{k'} < \alpha_k < R(u) < S(v_{k'}),
$$

which contradicts the left-monotonicity of (R, S) .

In the following definition the LC in the subscript refers to 'left-curtain coupling' and the S to 'simple' in the sense that the set $\{u : R(u) < S(u)\}$ takes the form of a single interval.

Definition 5.7. $(\mu, \nu) \in \mathcal{K}_{SLC}$ if $(\mu, \nu) \in \mathcal{K}$ and, in the construction of the lifted leftcurtain martingale coupling of μ and ν , $\exists u_0 \in (0, \mu(\mathbb{R}))$ such that $S = \overrightarrow{G}_{\mu}$ on $(0, u_0]$ and $S(u) > \overrightarrow{G}_{\mu}(u+)$ on $(u_0, \mu(\mathbb{R}))$, whence also R is decreasing on $(u_0, \mu(\mathbb{R}))$.

Define $\tilde{J}_k = A_k \setminus (\cup_{k' \neq k: A_{k'} \subsetneq A_k} A_{k'})$ and let $J_k = \text{interior}(\tilde{J}_k)$.

Proposition 5.8. Suppose $\mu \leq_{cx} \nu$. There exists a partition of $(0, \mu(\mathbb{R}))$ into countably many disjoint sets $(J_k)_{k\geq 1}$ and a quantile-lifted martingale coupling $\hat{\pi}^Q \in \mathcal{M}_{QL}(\mu, \nu)$ such that if $\mu_k = \mu^{J_k}$ and $\nu_k = \nu^{J_k} = \int_{u \in J_k} \int_{x \in I_\mu} \hat{\pi}^Q(du, dx, dy)$, then $(\mu - \sum_{k \ge 1} \mu_k) =$ $(\nu - \sum_{k\geq 1} \nu_k)$ and $\mu_k \leq_{cx} \nu_k$. Moreover, this partition and coupling can be chosen such that for each $k \geq 1$ we have $(\mu_k, \nu_k) \in \mathcal{K}_{SLC}$ and such that $\nu_k \wedge \nu_{k'} = 0$ for $k \neq k'$.

Proof. Consider the left-curtain mapping of μ to ν with associated lift specified by (R, S) . By Lemma [5.6](#page-31-1) for $k \neq k'$ we have $J_k \cap J_{k'} = \emptyset$ and then $\nu^{J_k} \wedge \nu^{J_{k'}} = 0$.

Proposition 5.9. $\mathcal{K}_{SLC} \subseteq \mathcal{K}_R$.

Proof. Suppose $(\mu, \nu) \in \mathcal{K}_{SLC}$. Then $S = \overrightarrow{G}_{\mu}$ on $(0, u_0]$ and $\mathcal{A}_{\leq} = (u_0, \mu(\mathbb{R}))$ for some $u_0 \in (0, \mu(\mathbb{R})).$

Now we argue that (μ, ν) satisfy the first numbered property of Definition [4.2.](#page-19-0) Consider a random variable $Y(U, V)$, defined as in Hobson and Norgilas [\[9,](#page-38-7) Theorem 3.7], where U and V are independent $U[0, 1]$ random variables. In particular, $Y(u, v) = \overrightarrow{G}_u(u)$ on $R(u) = S(u)$ and

$$
Y(u,v) = R(u)I_{\{v \le \frac{S(u) - \vec{G}_{\mu}(u)}{S(u) - R(u)}\}} + S(u)I_{\{v > \frac{S(u) - \vec{G}_{\mu}(u)}{S(u) - R(u)}\}}\tag{7}
$$

 \Box

 \Box

otherwise. Then $Y = Y(U, V) \sim \nu$ and $Y(u, V) = \overrightarrow{G}_{\mu}(u)$ almost surely whenever $u \in \{v \in (0,1) : \overrightarrow{G}_{\mu}(v) = S(v)\}\$. Since, by assumption, $S(u) = \overrightarrow{G}_{\mu}(u)$ for all $u \in (0, u_0]$, we have that $Y(U, V) = \overrightarrow{G}_{\mu}(U)I_{\{U \le u_0\}} + Y(U, V)I_{\{U > u_0\}}$ almost surely. But $(\overrightarrow{G}_{\mu}(U)I_{\{U\leq u_0\}})$ is distributed (up to normalisation of total mass) according to $\mu|_{(-\infty,\vec{G}_{\mu}(u_0))} + \delta_{\vec{G}_{\mu}(u_0)}(u_0 - \mu((-\infty,\vec{G}_{\mu}(u_0))))$, from which we have that $\nu \geq \mu|_{(-\infty,\overrightarrow{G}_{\mu}(u_0))} + \delta_{\overrightarrow{G}_{\mu}(u_0)}(u_0 - \mu((-\infty,\overrightarrow{G}_{\mu}(u_0))))$. Since ν is atom-less we must have that $u_0 = \mu((-\infty, \overrightarrow{G}_{\mu}(u_0)))$ and therefore $\nu \geq \mu|_{(-\infty, \overrightarrow{G}_{\mu}(u_0)))} = \mu|_{(-\infty, \overrightarrow{G}_{\mu}(u_0+))}$. It follows that $\nu \geq \mu$ on $(-\infty, \overrightarrow{G}_{\mu}(u_0+))$. Then set $a_0 := G(u_0+)$ and note that $u_0 = \mu((-\infty, a_0)).$

If $\nu \geq \mu$ on $(-\infty, a_2)$ for some $a_2 > a_0$ then $\nu([a_0, a_2)) \geq \mu([a_0, a_2)) > 0$ and D is convex on $(-\infty, a_2)$. Then we can pick $u_1 > u_0$ with $\overrightarrow{G}_{\mu}(u_0+) = a_0 \leq \overrightarrow{G}_{\mu}(u_1) < a_2$ and then convexity of D on $(-\infty, a_2)$ ensures that \mathcal{E}_{u_1} is convex everywhere, since $\mathcal{E}_{u_1} \geq D$ and $\vec{G}_{\mu}(u_1) < a_2$. It follows that $S(u_1) = \vec{G}_{\mu}(u_1)$, contradicting the fact that $S > \vec{G}_{\mu}$ on $(u_0, \mu(\mathbb{R}))$. This proves that our candidate quantities u_0 and a_0 satisfy the first property of Definition [4.2.](#page-19-0)

Now we argue that the middle listed property of Definition [4.2](#page-19-0) holds, namely that the tangent L_{a_0} to D at a_0 lies above $D_{\mu,\nu}$ on (a_0,∞) .

There are four possible cases: either $L_{a_0} > D_{\mu,\nu}$ on (a_0,∞) ; or there exists $c_2 >$ $c_1 > a_0$ with $D_{\mu,\nu}(c_1) < L_{a_0}(c_1)$ and $D_{\mu,\nu}(c_2) \ge L_{a_0}(c_2)$; or there exists $d_1 > a_0$ such that $L_{a_0} \leq D_{\mu,\nu}$ on (a_0, d_1) and $L_{a_0}(d_1) < D_{\mu,\nu}(d_1)$; or there exists $e > a_0$ such that $L_{a0} = D_{\mu,\nu}$ on [a₀, e]. The last case cannot happen else a₀ is not maximal. Therefore it is sufficient to show that the second and third cases also lead to a contradiction.

For the second case, suppose there exists $c_2 > c_1 > a_0$ with $D_{\mu,\nu}(c_1) < L_{a_0}(c_1)$ and $D_{\mu,\nu}(c_2) \geq L_{a_0}(c_2)$. Without loss of generality we may assume $D_{\mu,\nu}(c_2) = L_{a_0}(c_2)$. Choose w_2 such that $\overrightarrow{G}_{\mu}(w_2) \leq c_2 \leq \overrightarrow{G}_{\mu}(w_2+)$ and consider \mathcal{E}_{0,w_2} . Note $w_2 < \mu(\mathbb{R})$. Given the existence of c_1 , there exists $c_3 \in (a_0, c_2)$ such that $\mathcal{E}_{0,w_2}^c(c_3) = \mathcal{E}_{0,w_2}(c_3)$ $L_{a_0}(c_3)$. Choose w_3 such that $\overrightarrow{G}_{\mu}(w_3) \leq c_3 \leq \overrightarrow{G}_{\mu}(w_3+)$. Then $S(w_3) \leq \overrightarrow{G}_{\mu}(w_3+)$. But this contradicts the fact that $\mathcal{A} = (u_0, \mu(\mathbb{R}))$.

For the third case, suppose there exists $d_1 > a_0$ such that $L_{a_0} \leq D_{\mu,\nu}$ on (a_0, d_1) and $L_{a_0}(d_1) < D_{\mu,\nu}(d_1)$. There exists $d_2 \in (a_0, d_1)$ such that $L_{a_0} \le D_{\mu,\nu}$ on (a_0, d_2) , $L_{a_0}(d_2) < D_{\mu,\nu}(d_2)$ and $D'_{\mu,\nu}(d_2-) > L'_{a_0}(d_2)$. Choose v_2 such that $\overrightarrow{G}_{\mu}(v_2) \leq d_2 \leq$ $\vec{G}_{\mu}(v_2+)$. Then $\mathcal{E}_{0,v_2}^c > L_{a_0}$ on $[d_2,\infty)$. It follows that there exists $d_3 \in (a_0, d_2]$ such that $\mathcal{E}_{0,v_2}^c(d_3) = \mathcal{E}_{0,v_2}(d_3)$. Choose v_3 such that $\overrightarrow{G}_{\mu}(v_3) \leq d_3 \leq \overrightarrow{G}_{\mu}(v_3+)$. Then $S(v_3) \leq \overrightarrow{G}_{\mu}(v_3+)$ which again gives us a contradiction.

Now consider the final part of Definition [4.2.](#page-19-0) Note that for $b \in \left(\inf\{k : D_{\mu,\nu}(k)\right)$ 0}, a1) we have that D′ µ,ν(b) exists and is positive. Let L^b := L b,D′ µ,ν (b−) $D_{\mu,\nu}^{(0,-)}$. We must have that $\{k > b : D_{\mu,\nu}(k) > L_b(k)\}\neq \emptyset$. Define $\overline{c} := \sup\{k > b : D_{\mu,\nu}(k) > L_b(k)\}.$ Continuity of $D_{\mu,\nu}$ (and L_b) implies that $D_{\mu,\nu}(\overline{c})=L_b(\overline{c})$. Moreover, since $D_{\mu,\nu}$ is convex and non-decreasing on $(-\infty, a_0)$, and $\lim_{k\to\infty} D_{\mu,\nu}(k) = 0$, we have that $a_0 < \overline{c} < \infty$. There are three cases: either $L_b < D_{\mu,\nu}$ on $[a_0,\overline{c})$; or $L_b \le D_{\mu,\nu}$ on $[a_0,\overline{c})$ and there exists $\tilde{c} \in (a_0, \overline{c})$ for which $D_{\mu,\nu}(\tilde{c}) = L_b(\tilde{c});$ or $\{k \in (b, \overline{c}) : D_{\mu,\nu}(k) < L_b(k)\}\neq \emptyset$. We show the first case leads to $D'_{\mu,\nu}(b) = L'_{b} > D'_{\mu,\nu}(\overline{c}-)$, and that the second and third cases cannot happen.

Case 1: $L_b < D_{\mu,\nu}$ on $[a_0,\overline{c})$. Suppose $D_{\mu,\nu} < L_b$ on (\overline{c},∞) and $D'_{\mu,\nu}(b) = D'_{\mu,\nu}(\overline{c})$. Let $\overline{u} := \inf \{ u \in (u_0, \mu(\mathbb{R})) : \overrightarrow{G}_{\mu}(u) \leq \overline{c} \leq \overrightarrow{G}_{\mu}(u+)\}.$ Note that $\overrightarrow{G}_{\mu}(\overline{u}) \leq \overline{c} \leq \overrightarrow{G}_{\mu}(\overline{u}+)$ and $\overline{u} > u_0$. Then $S(\overline{u}) \leq \overrightarrow{G}_{\mu}(\overline{u}_{+}),$ a contradiction (since $\overline{u} \in (u_0, \mu(\mathbb{R})) = \mathcal{A}_{<}$). Hence $D'_{\mu,\nu}(b) = L'_b > D'_{\mu,\nu}(\overline{c})$.

Case 2: $L_b < D_{\mu,\nu}$ on $[a_0,\overline{c})$ and there exists $\tilde{c} \in (a_0,\overline{c})$ with $D_{\mu,\nu}(\tilde{c}) = L_b(\tilde{c})$. In this case we can take $\tilde{u} \in (u_0, \mu(\mathbb{R})) = A_{\leq \varepsilon}$ such that $\overrightarrow{G}_{\mu}(\tilde{u}) \leq \tilde{c} \leq \overrightarrow{G}_{\mu}(\tilde{u}^+),$ and then $S(\tilde{u}) \leq \overrightarrow{G}_{\mu}(\tilde{u}+)$, a contradiction.

Case 3: ${k \in (b,\overline{c}) : D_{\mu,\nu}(k) < L_b(k)} \neq \emptyset$. In this case we can find $b' < b$ (with $D'_{\mu,\nu}(b') > 0$ for which there exist $c_1, c_2 \in \mathbb{R}$ with $a_0 < c_1 < \overline{c} < c_2$ and such that $L_{b'} \leq D_{\mu,\nu}$ on $(-\infty,c_2]$ and $L_{b'}(k) = D_{\mu,\nu}(k)$ for $k \in \{c_1,c_2\}$. Then, similarly as in Case 2, we can take $\hat{u} \in (u_0, \mu(\mathbb{R}))$ such that $\overrightarrow{G}_{\mu}(\hat{u}) \leq c_1 \leq \overrightarrow{G}_{\mu}(\hat{u}+)$ and then $S(\hat{u}) \leq \overrightarrow{G}_{\mu}(\hat{u}^{\perp}),$ a contradiction. \Box

Lemma 5.10. Suppose $(\mu, \nu) \in \mathcal{K}_R$. If $u_0 = 0$ then $u_1 = \mu(\mathbb{R})$ and $(\mu, \nu) \in \mathcal{K}_*$.

Proof. In this case, for each $u \in (0, \mu(\mathbb{R}))$, $\mathcal{E}_{u_0,u} = \mathcal{E}_{0,u}$ and $N(u) = S(u)$. But, by Proposition [5.9,](#page-32-0) $S(u) > G(u+)$ for $u \in (u_0, \mu(\mathbb{R})) = (0, \mu(\mathbb{R}))$. Hence $u_1 = \mu(\mathbb{R}) > 0 =$ \square u_0 .

6 Extending the existence result to \mathcal{K}_{SLC} and then to all cases with ν continuous.

Theorem 6.1. Suppose $(\mu, \nu) \in \mathcal{K}_{SLC}$. Then $\mathcal{IM}(\mu, \nu) \neq \emptyset$.

Proof. If $(\mu, \nu) \in \mathcal{K}_{*}$ then the result is covered by Theorem [4.15.](#page-26-2) So we assume $(\mu, \nu) \in$ $\mathcal{K}_{SLC} \setminus \mathcal{K}_*.$

The idea of the proof is to use the left-curtain martingale coupling to construct a decreasing sequence of nested intervals $[0, \mu(\mathbb{R})] = J_0 \supseteq J_1 \supseteq J_2 \supseteq \cdots$ such that $\lim_j Leb(J_j) = 0$ and such that we can find $\hat{\pi}_k \in \mathcal{IM}_{QL}(\mu^{J_k \setminus J_{k+1}}, \nu^{J_k \setminus J_{k+1}})$ for each $k \geq 1$. Then we use Proposition [2.9](#page-8-1) to combine these martingale couplings to give an injective martingale coupling of μ and ν . Note that $\cup_{j>0}(J_k \setminus J_{k+1}) = [0, \mu(\mathbb{R})]$.

Let (R, S) be the (left-monotone with respect to \overrightarrow{G}_μ) functions which define the quantile-lifted left-curtain martingale coupling of μ and ν as in Theorem [5.1.](#page-31-2) R and S remain defined relative to μ and ν throughout this proof. Note that since $(\mu, \nu) \in \mathcal{K}_{SLC}$ we have that $S = \overrightarrow{G}_{\mu}$ on $(0, u_0]$, and since $\mu \leq \nu$ there, and ν is continuous, we have that S is strictly increasing on $(0, u_0]$. Hence S^{-1} exists and is continuous below $S(u_0)$.

Let $J_0 = [0, \mu(\mathbb{R})], \mu_0 = \mu^{J_0} = \mu, \nu_0 = \nu^{J_0} = \nu$. By Lemma [5.10,](#page-34-1) since $(\mu, \nu) \notin \mathcal{K}_*$ we must have $u_0 > 0$. Further, if $u_0 = \mu(\mathbb{R})$ then $\mu = \nu$, a case excluded by Standing Assumption [2.3.](#page-6-1) Hence $0 < u_0 < \mu(\mathbb{R})$.

Since $(\mu, \nu) \in \mathcal{K}_{SLC}$, by Proposition [5.9](#page-32-0) we must have that $D_{\mu,\nu}$ has the properties listed in Definition [4.2.](#page-19-0) There are two cases: either $a_1 = a_0$ or $a_1 < a_0$. We begin by arguing that these cases correspond to $S(u_0) = R(u_0+)$ and $S(u_0) > R(u_0+)$ respectively.

If $R(u_0+) < S(u_0)$ then for $u \in (v_0 := S^{-1}(R(u_0+)), u_0)$ we have $S(u) = \overrightarrow{G}_{\mu}(u)$. Then $\nu \geq \mu$ on $(R(u_0+), S(u_0))$. But, by the left-monotonicity property of (R, S) with respect to \vec{G}_{μ} we have $R(v) \le R(u_0 + s) \le S(u_0) \le S(v)$ for all $v > u$ and hence ${u: R(u) \text{ or } S(u) \in (R(u_0+), S(u_0))} = (v_0, u_0).$ Hence $\nu = \mu$ on $(R(u_0+), S(u_0))$ and $a_1 \leq R(u_0+) \leq S(u_0) = a_0$. Further we must have $a_1 = R(u_0+)$ since for each $a < a_1$, ${u: R(u) < \overrightarrow{G}_{\mu}(u+)< S(u) \text{ and } R(u) \in (a, a_0)} \neq \emptyset$. Conversely, suppose $R(u_0+)$ = $S(u_0)$. Then for each $a < S(u_0)$, there exists $v > u_0$ such that $R(v) \in (a, R(u_0))$ and then $\nu(a, a_0) > \mu(a, a_0)$. Hence $a_1 = a_0$.

Suppose $a_1 < a_0$, or equivalently $R(u_0+) < S(u_0)$. Let $\mu = \mu|_{(a_1,a_0)}$ and $\mu = \nu|_{(a_1,a_0)}$. Then $\mu = \nu$. Let $\mu = \mu - \mu$ and $\dot{\nu} = \nu - \dot{\nu}$. Since $\mu = \dot{\nu}$ we have $\dot{\mu} \leq_{cx} \dot{\nu}$. Moreover, $D_{\mu,\nu} = D_{\mu,\nu}$ so that $(\mu, \nu) \in \mathcal{K}_R$. Also, $(\overrightarrow{G}_{\mu}, \overrightarrow{R}, \overrightarrow{S})$ define the left-curtain coupling of (μ, ν) if and only if $\hat{S} = \overrightarrow{G}_{\mu} = \overrightarrow{G}_{\mu}$ on $(0, v_0]$ and for $v_0 < v < \mu(\mathbb{R}),$

$$
\dot{R}(v) = R(v - (u_0 - v_0)), \quad \dot{S}(v) = S(v - (u_0 - v_0)).
$$

It follows that $(\hat{\mu}, \hat{\nu}) \in \mathcal{K}_{SLC}$. Note that $\hat{R}(v_0+) = R(u_0+) = S(v_0) = \hat{S}(v_0)$.

Suppose that there exists an injective quantile-lifted martingale coupling π^Q of μ and *ν*. Since $\mu = \nu$ there is only one quantile-lifted martingale coupling π^Q of μ and ν and for this coupling mass stays in the same place. This is clearly injective. Then, since both supp $(\nu) \cap \text{supp}(\nu) \subseteq \{a_1, a_0\}$ (a set of zero ν -measure) we can combine π^Q and π^Q to give an injective quantile-lifted coupling of μ and ν .

The above arguments allow us to reduce the general case to one in which $R(u_0+)$ $S(u_0)$. From now on we assume that we are in this case. Define $f : (u_0, \mu(\mathbb{R})) \mapsto$ $(0, \mu(\mathbb{R})]$ by $f(v) = v - S^{-1} \circ R(v)$. Then since S is increasing and R is decreasing on $(u_0, \mu(\mathbb{R}))$ we have that f is increasing. Moreover, since S is strictly increasing on $(0, u_0)$ and $R(u_0+) = S(u_0)$ we have that $\lim_{v \downarrow u_0} f(v) = 0$.

Fix $\epsilon \in (0,1)$. Suppose inductively that there exists $J_k = [\underline{u}_k = (S^{-1} \circ R)(\overline{u}_k), \overline{u}_k]$ with $u_0 \in J_k$ and $\overline{u}_k - \underline{u}_k \leq \epsilon^k$ such that $(\mu_k := \mu^{J_k}, \nu_k := \nu^{J_k}) \in \mathcal{K}_{SLC}$ where ν^{J_k} is calculated using the left-curtain martingale coupling of μ and ν , i.e. $\nu^{J_k}(dy) =$ $\int_{u\in J_k}\int_{x\in I_\mu}\pi^{Q,LC}(du, dx, dy)$. Clearly the inductive hypothesis is true for $k=0$, and if it holds true for all k then $\lim_k Leb(J_k) = \lim_k (\overline{u}_k - \underline{u}_k) = 0$.

By the definition of u_0 it must be the case that $S(u) > \overrightarrow{G}_{\mu}(u+)$ on (u_0, \overline{u}_k) . Then, by the arguments of the previous paragraph we can choose $\overline{u}_{k+1} \in [u_0, \overline{u}_k)$ such that $S(\overline{u}_{k+1}) > \overrightarrow{G}_{\mu}(\overline{u}_{k+1}+)$ and $f(\overline{u}_{k+1}) \leq \epsilon(\overline{u}_{k} - \underline{u}_{k})$. Set $\underline{u}_{k+1} = S^{-1}(R(\overline{u}_{k+1}))$. Note that $u_{k+1} \leq u_0 < \overline{u}_{k+1}.$

Set $J_{k+1} = [\underline{u}_{k+1}, \overline{u}_{k+1}]$ and $\tilde{J}_{k+1} = J_k \setminus J_{k+1}$. Let $\mu_{k+1} = \mu^{J_{k+1}}, \nu_{k+1} = \nu^{J_{k+1}}, \tilde{\mu}_{k+1} =$ $\mu^{J_{k+1}}$ and $\tilde{\nu}_{k+1} = \nu^{J_{k+1}}$. We argue that $(\tilde{\mu}_{k+1}, \tilde{\nu}_{k+1}) \in \mathcal{K}_{*}$ and hence $\mathcal{IM}_{QL}(\tilde{\mu}_{k+1}, \tilde{\nu}_{k+1})$ is non-empty, and $(\mu_{k+1}, \nu_{k+1}) \in \mathcal{K}_{SLC}$, which will complete the inductive hypothesis.

In this way we get a family $(\tilde{\mu}_k, \tilde{\nu}_k)_{k\geq 1}$ such that for each $k, (\tilde{\mu}_k, \tilde{\nu}_k) \in \mathcal{K}_*$, and such that $(\tilde{J}_k)_{k\geq 1}$ is a partition of $(0,\mu(\mathbb{R}))$. Combining Theorem [4.15](#page-26-2) and Proposition [2.9](#page-8-1) we conclude $\mathcal{IM}(\mu, \nu)$ is non-empty.

It only remains to argue that for each k we have $(\mu_{k+1}, \nu_{k+1}) \in \mathcal{K}_{SLC}$ and $(\tilde{\mu}_{k+1}, \tilde{\nu}_{k+1}) \in$ \mathcal{K}_{*} . Note that if $(\mu_{k+1}, \nu_{k+1}) \in \mathcal{K}_{*} \subseteq \mathcal{K}_{SLC}$ for any k then the construction can terminate at this point.

For the first conclusion, define $\{G_{k+1}, S_{k+1}, R_{k+1}\} : (0, \overline{u}_{k+1} - \underline{u}_{k+1} = \mu_{k+1}(\mathbb{R})) \mapsto \mathbb{R}$ by $G_{k+1}(u) = G_{\mu}(\underline{u}_{k+1} + u), S_{k+1}(u) = S(\underline{u}_{k+1} + u)$ and $R_{k+1}(u) = R(\underline{u}_{k+1} + u)$. See Figure [4.](#page-37-0) Then the triple $(G_{k+1}, S_{k+1}, R_{k+1})$ define a lifted left-curtain coupling of μ_{k+1} and ν_{k+1} . Moreover, $S_{k+1} = G_{k+1}$ on $[0, u_0 - \underline{u}_{k+1}]$ and $S_{k+1}(u) > G_{k+1}(u+)$ on $(u_0 - \underline{u}_{k+1}, \mu_{k+1}(\mathbb{R})].$ Hence $(\mu_{k+1}, \nu_{k+1}) \in \mathcal{K}_{SLC}$.

The second conclusion is similar, but needs an extra argument to show that $(\tilde{\mu}_{k+1}, \tilde{\nu}_{k+1}) \in$ \mathcal{K}_* . Define $\{\tilde{G}_{k+1}, \tilde{S}_{k+1}, \tilde{R}_{k+1}\}$: $[0, \overline{u}_k - \overline{u}_{k+1} + \underline{u}_{k+1} - \underline{u}_k = \tilde{\mu}_{k+1}(\mathbb{R})] \mapsto \mathbb{R}$ by, for $H \in \{G, S, R\}$

$$
\tilde{H}_{k+1}(u) = \begin{cases} H(\underline{u}_k + u) & 0 < u \le \underline{u}_{k+1} - \underline{u}_k \\ H(\overline{u}_{k+1} - \underline{u}_{k+1} + \underline{u}_k + u) & \underline{u}_{k+1} - \underline{u}_k < u < \tilde{\mu}_{k+1}(\mathbb{R}) \end{cases}
$$

Then the triple $(\tilde{G}_{k+1}, \tilde{S}_{k+1}, \tilde{R}_{k+1})$ define a lifted left-curtain coupling of $\tilde{\mu}_{k+1}$ and $\tilde{\nu}_{k+1}$. Moreover, $\tilde{S}_{k+1} = \tilde{G}_{k+1}$ on $[0, \tilde{u}_0^{k+1} := \underline{u}_{k+1} - \underline{u}_k]$ and $\tilde{S}_{k+1}(u) > \tilde{G}_{k+1}(u+)$ on $(\underline{u}_{k+1}-\underline{u}_k,\tilde{\mu}_{k+1}(\mathbb{R}))$. Hence $(\tilde{\mu}_{k+1},\tilde{\nu}_{k+1})\in \mathcal{K}_{SLC}$. Further, by construction $\tilde{S}_{k+1}((\underline{u}_{k+1}-\underline{u}_k,\tilde{\mu}_{k+1}(\mathbb{R}))$ u_k \rightarrow $\tilde{G}_{k+1}((u_{k+1} - u_k) + \text{)}$ and $\tilde{\nu}_{k+1}$ places no mass on the interval $[\tilde{G}_{k+1}((u_{k+1} - u_k) + \text{)}]$ (\underline{u}_k) +), $\tilde{S}_{k+1}((\underline{u}_{k+1} - \underline{u}_k)$ +). Then, recall the definition of $\vec{\tau}$ in Definition [3.5,](#page-11-2) for $l \in [\tilde{u}_0^{k+1}, \tilde{\mu}_{k+1}(\mathbb{R})],$ we can define $Z^+_{\tilde{\mathcal{E}}_{\tilde{\mathcal{E}}}}$ $\tilde{\varepsilon}_{\tilde{u}_{\alpha}^{k+1},\tilde{u}_{\alpha}^{k+1}}(\tilde{G}_{k+1}(l))$ where $\tilde{\mathcal{E}}_{v,u} = \mathcal{E}_{v,u}^{\tilde{\mu}_{k+1},\tilde{\nu}_{k+1}}$. It is clear that $Z_{\tilde{\varepsilon}}^+$ $(\tilde{G}_{k+1}(\tilde{u}_0^{k+1} + \delta)) \geq \tilde{S}_k$ $\tilde{\mathcal{E}}_{\tilde{u}_0^{k+1},\tilde{u}_0^{k+1}}$ $(\tilde{G}_{k+1}(\tilde{u}_0^{k+1} + \delta)) \ge \tilde{S}_{k+1}((\underline{u}_{k+1} - \underline{u}_k) +) > \tilde{G}_{k+1}((\tilde{u}_0^{k+1} + \delta) +)$ for sufficiently small δ . In particular, $(\tilde{\mu}_{k+1}, \tilde{\nu}_{k+1}) \in \mathcal{K}_{*}$.

Figure 4: The decomposition of J_k into J_{k+1} and \tilde{J}_{k+1} . We can recreate the top figure by inserting the lower-left figure into the lower right figure at the x-coordinate $\underline{u}_{k+1} - \underline{u}_k$.

It remains to put it all together.

Proof of Theorem [1.2.](#page-2-0) The quantile-lifted left curtain martingale coupling and Proposition [5.8](#page-32-1) can be used to divide the problem into countably many intervals on each of which $(\mu_k, \nu_k) \in \mathcal{K}_{SLC}$. On each such interval Theorem [6.1](#page-34-2) implies there exists an injective quantile-lifted martingale coupling. Finally, Proposition [2.9](#page-8-1) gives that there is an injective quantile-lifted martingale coupling of μ and ν . \Box

References

- [1] BEILGLBÖCK, M., HOBSON D.G., NORGILAS D.: The potential of the shadow measure. Electronic Communications in Probability, 27:1-–12, (2022).
- [2] BEIGLBÖCK M., JUILLET N.: On a problem of optimal transport under marginal martingale constraints. The Annals of Probability, 44(1):42–106, (2016).
- [3] BEIGLBÖCK M., JUILLET N.: Shadow couplings. Transactions of American Mathematical Society, 374:4973–5002, (2021).
- [4] HENRY-LABORDÈRE P., TOUZI N.: An explicit martingale version of the onedimensional Brenier theorem. Finance and Stochastics, 20(3):635–668, (2016).
- [5] HOBSON D.G.: The maximum maximum of a martingale. In: Azéma, J., Yor, M., Emery, M., Ledoux, M. (eds) Séminaire de Probabilités XXXII. Lecture Notes in Mathematics, vol 1686. Springer, Berlin, Heidelberg, (1998).
- [6] Hobson D.G., Klimmek M.: Robust price bounds for the forward starting straddle. Finance and Stochastics, $19(1):189-214$, (2015) .
- [7] HOBSON D.G., NEUBERGER A.: Robust bounds for forward start options. Mathematical Finance, 22(1):31–56, (2012).
- [8] HOBSON D.G., NORGILAS D.: The left-curtain martingale coupling in the presence of atoms. The Annals of Applied Probability, 29(3):1904–1928, (2019).
- [9] Hobson D.G., Norgilas D.: A construction of the left-curtain coupling. Electronic Journal of Probability, 27:1–46, (2022)
- [10] NUTZ M., WANG R., ZHANG Z.: Martingale Transports and Monge Maps. Preprint, https://arxiv.org/abs/2209.14432, (2022)

[11] Strassen V.: The existence of probability measures with given marginals. The Annals of Mathematical Statistics, 36(2):423–439, (1965).