# **Compact Binary Formation in Open Star Clusters I: High Formation Efficiency of Gaia BHs and Their Multiplicities**

Ataru Tanikawa<sup>1\*</sup>, Savannah Cary<sup>2</sup>, Minori Shikauchi<sup>3,4,5</sup>, Long Wang<sup>6,7</sup>, Michiko S. Fujii<sup>2</sup>

<sup>1</sup>Department of Earth Science and Astronomy, College of Arts and Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8902, Japan <sup>2</sup>Department of Astronomy, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>3</sup>Department of Physics, the University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

<sup>4</sup>Research Center for the Early Universe (RESCEU), the University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

<sup>5</sup>Department of Physics and Astronomy, the University of British Columbia, 6224 Agricultural Road, Vancouver, BC, V6T 1Z1, Canada

<sup>6</sup>School of Physics and Astronomy, Sun Yat-sen University, Daxue Road, Zhuhai, 519082, China

<sup>7</sup>CSST Science Center for the Guangdong-Hong Kong-Macau Greater Bay Area, Zhuhai, 519082, China

Accepted XXX. Received YYY; in original form ZZZ

#### ABSTRACT

Gaia BHs, black hole (BH) binaries discovered from database of an astrometric telescope *Gaia*, pose a question to the standard binary evolution model. We have assessed if Gaia BHs can be formed through dynamical capture in open clusters rather than through isolated binary evolution. We have performed gravitational *N*-body simulations of 100 open clusters with  $10^5 M_{\odot}$  in total for each metallicity Z = 0.02, 0.01, and 0.005. We have discovered one Gaia BH-like binary escaping from an open cluster, and found that the formation efficiency of Gaia BHs in open clusters (~  $10^{-5}M_{\odot}^{-1}$ ) is larger than in isolated binaries (~  $10^{-8}M_{\odot}^{-1}$ ) by 3 orders of magnitude. The Gaia BH-like binary is the inner binary of a triple star system. Gaia BHs can have tertiary stars frequently, if they are formed in open clusters. Combining additional *N*-body simulations with 8000 open clusters with  $8 \times 10^6 M_{\odot}$ , we have estimated the number of Gaia BHs in the Milky Way disk to  $10^4 - 10^5$  (depending on the definitions of Gaia BHs), large enough for the number of Gaia BHs discovered so far. Our results indicate that the discoveries of Gaia BHs do not request the reconstruction of the standard binary evolution model, and that Gaia BHs are a probe for the dynamics of open clusters already evaporated.

Key words: stars: black holes – galaxies: star clusters: general – Astrometry and celestial mechanics – (stars:) binaries (including multiple): close

#### **1 INTRODUCTION**

Black holes (BHs) are the final state of massive stars. Albeit of their darkness, they have been discovered in binary systems, such as X-ray binaries (Casares et al. 2017), spectroscopic binaries (Shenar et al. 2022), astrometric binaries (El-Badry et al. 2023a,b; Tanikawa et al. 2023; Chakrabarti et al. 2023), and gravitational wave sources (The LIGO Scientific Collaboration et al. 2021). The discoveries of astrometric BH binaries, also known as Gaia BH1 and BH2 (El-Badry et al. 2023a,b, respectively) in Gaia DR3 (Gaia Collaboration et al. 2022), challenge the standard binary evolution model. Gaia BH1 and BH2 (hereafter, Gaia BHs) have ~  $1M_{\odot}$  visible stars as companions, orbital periods of ~  $10^2$ - $10^3$  days, and moderately high eccentricities (~ 0.5). Isolated binaries can form such configurations only if common envelope ejection is ~ 10 times more efficient than previously expected (El-Badry et al. 2023a).

Another possible formation channel of Gaia BHs is that a BH dynamically captures a ~  $1M_{\odot}$  visible star in open clusters. This is similar to double BH formation in open clusters (Ziosi et al. 2014;

Kumamoto et al. 2019, 2020; Di Carlo et al. 2019; Banerjee 2021; Rastello et al. 2021). As for double BHs, such dynamical capture can happen in globular clusters (Portegies Zwart & McMillan 2000; Tanikawa 2013; Rodriguez et al. 2016; Askar et al. 2017; Kremer et al. 2022), first star clusters (Wang et al. 2022), and galactic centers (Tagawa et al. 2020). However, Gaia BHs should not originate from globular clusters, first star clusters, and galactic centers, since they are Milky Way disk components, and have near solar metallicities. They are likely to be formed in open clusters if their formation channel is dynamical capture.

Gaia BH1 and BH2 have different orbital periods by an order of magnitude, and different types of visible stars: a main-sequence (MS) star for Gaia BH1 and a red giant star for Gaia BH2. They may have different origins. What can be said with certainty is that Gaia BHs have common features in which they are difficult to be formed through isolated binary evolution, if common envelope is ejected efficiently (El-Badry et al. 2023a,b). Thus, we explore possibility that Gaia BHs are formed through dynamical capture in open clusters.

In this paper, we report an extremely high formation efficiency of Gaia BHs in open clusters. Moreover, our simulations show that they frequently host tertiary stars. The structure of this paper is as follows.

<sup>\*</sup> E-mail: tanikawa@ea.c.u-tokyo.ac.jp

In section 2, we describe our numerical method to follow Gaia BH formation in open clusters and isolated binaries. In section 3, we show our simulation results. In section 4, we conclude our results, and discuss the implication of our results for Gaia BHs discovered recently. We focus on a Gaia BH-like binary formed in 100 open clusters with  $10^5 M_{\odot}$  and metallicity of Z = 0.005, while we discuss the dependence of the formation efficiency of Gaia BHs on initial conditions of open clusters in Appendix.

#### 2 METHOD

We follow the dynamical evolution of open clusters by means of a gravitational *N*-body code PeTar (Wang et al. 2020b). The PeTar code is based on the particle-tree and particle-particle algorithm (Oshino et al. 2011; Iwasawa et al. 2017) with the aid of FDPS (Iwasawa et al. 2016, 2020). Binary orbital evolution and close encounters are treated by the slow-down algorithmic regularization method (SDAR: Wang et al. 2020a). The external potential is calculated by GALPY (Bovy 2015). We can see comparison between PeTar and NBODY6++GPU (Nitadori & Aarseth 2012; Wang et al. 2015; Kamlah et al. 2022) for mass scale of small globular clusters in Wang et al. (2020b), and open clusters in Appendix A.

The PeTar code is coupled with the BSE code (Hurley et al. 2002; Banerjee et al. 2020) to solve single and binary star evolutions. We overview the model of single and binary star evolution. Single stars evolve on a track developed by Hurley et al. (2000) with stellar winds formulated by Belczynski et al. (2010). Massive stars either cause supernovae or collapse to BHs at the end of their lives, and leave behind NSs and BHs with masses modeled by the rapid model (Fryer et al. 2012) with modification of pair instability supernovae modeled by Belczynski et al. (2016b). Binary evolution model is the same as Banerjee et al. (2020). It contains various binary evolution processes, such as tidal interaction, wind accretion, stable mass transfer, and common envelope. Here, we only describe parameters of common envelope which Gaia BHs formation is the most sensitive to (El-Badry et al. 2023a). A binary can experience common envelope evolution modeled as the  $\alpha$  formalism (Webbink 1984), where we adopt  $\alpha = 3$  and  $\lambda$  of Claeves et al. (2014).

We generate 100 open clusters for each metallicity of Z = 0.02, 0.01, and 0.005. The total mass of each open cluster is  $10^3 M_{\odot}$ , and then the total mass of open clusters is  $10^5 M_{\odot}$  for each metallicity. We adopt the Plummer model for the initial phase space distribution of stars. We set the half-mass radius to 0.8 pc. Each cluster orbits around Milky Way at 8 kpc with 220 km s<sup>-1</sup>. The initial binary fraction is 100 %. Although the binary fraction appears too high, it is not sensitive to the formation efficiency and multiplicity of Gaia BHs as seen in Appendix B. The primary stars of binaries have Kroupa's initial mass function (Kroupa 2001) with the minimum and maximum masses of 0.08 and  $150M_{\odot}$ , respectively. The mass ratio, orbital period, and eccentricity distribution of binaries follow initial conditions from Sana et al. (2012). The initial conditions are generated using MCLUSTER (Küpper et al. 2011). We follow their evolutions over 500 Myr. Although the initial central number density seems to be high (~  $10^3 \text{ pc}^{-3}$ ), the central number density of visible stars (defined as stars except BHs, neutron stars, and white dwarfs) drop to  $\sim 3 \text{ pc}^{-3}$  at 500 Myr.

For comparison, we also calculated isolated binary evolution. We prepare two types of initial conditions for calculations of isolated binary evolution. Like the binaries in our open clusters, the 1st type of initial conditions for isolated binary evolution are also adopted from Sana et al. (2012). We find that the 1st type of initial conditions

Table 1. Summary of the Gaia BH-like binaryat 500 Myr.

Parameters	Values	Remarks
Metallicity (Z)	0.005	
BH mass Secondary mass Period Eccentricity	$21.4M_{\odot}$ $0.82M_{\odot}$ $8.3 \times 10^2$ days 0.3-0.8	MS star Oscillating
Tertiary mass Outer period Outer eccentricity Mutual inclination	$1.59 M_{\odot}$ $1.2 \times 10^{6}$ days 0.689 34-59 deg	MS star Oscillating

never form Gaia BHs, because the minimum mass ratio is 0.1. In our single star evolution model,  $\geq 20M_{\odot}$  stars can leave behind BHs. BHs cannot have  $\leq 1M_{\odot}$  stars as companions. Thus, we prepare the 2nd type of initial conditions in which the minimum mass ratio is reduced to 0.0005, such that even  $150M_{\odot}$  stars can have  $0.08M_{\odot}$  stars as companions. We would like to remark that the 2nd type of initial conditions is much more likely to form Gaia BHs than the initial conditions of open clusters unless anything prohibits Gaia BHs from forming. We also note that we do not adopt such a small mass ratio for initial binaries in open clusters. We follow their evolution over 30 Myr. Until that, all BH progenitors evolve to BHs.

Throughout this paper, we compare properties of BH binaries for open clusters at 500 Myr, and for isolated binaries at 30 Myr. We underestimate the formation efficiency of BH binaries with secondary masses of  $\geq 2.5 M_{\odot}$ , because  $\geq 2.5 M_{\odot}$  stars evolve to remnant objects until 500 Myr. However, this does not affect our purpose. Gaia BHs discovered so far have secondary stars of  $\sim 1 M_{\odot}$ .

Our simulations generate 59 BH binaries. Such BH binaries have various secondary masses, periods, and eccentricities. In this paper, we define Gaia BHs as BH binaries with secondary masses of  $\leq 1.1 M_{\odot}$ , periods of 100-2000 days, and eccentricities of 0.3-0.9 unless stated.

#### **3 RESULTS**

We find that one open cluster with Z = 0.005 contains a BH binary similar to Gaia BHs, hereafter called "Gaia BH-like binary". We summarize the properties of the Gaia BH-like binary in Table 1. Its BH mass is  $21.4M_{\odot}$ . Its companion (hereafter, secondary) star is a MS star and has a mass of  $0.82M_{\odot}$ . The binary period is  $8.3 \times 10^2$ days, and its eccentricity oscillates from 0.3 to 0.8 due to von Zeipel-Lidov-Kozai (ZLK) mechanism (von Zeipel 1910; Lidov 1962; Kozai 1962). We find that the Gaia BH-like binary is the inner binary of a triple star system, and it is because of a tertiary star that we see ZLK oscillation of the Gaia BH-like binary. The tertiary star is a MS star and has  $1.59M_{\odot}$  at 500 Myr. The outer period and eccentricity are  $1.2 \times 10^6$  days and 0.689, respectively.

Figure 1 shows the position and velocity of the Gaia BH-like binary at 500 Myr. We can see that the Gaia BH-like binary has escaped from the open cluster located at the coordinate origin. The Gaia BH-like binary will be long-lived after 500 Myr as long as it is not perturbed by the tertiary star (described later). This Gaia BHlike binary escapes from the open cluster through either of two-body



**Figure 1.** Positions (top) and velocities (bottom) of the Gaia BH-like binary (star points) and other stars (dots) at 500 Myr. All the stars including BHs, neutron stars, and white dwarfs, are plotted. The central number density of visible stars is  $\sim 3.4 \text{ pc}^{-3}$ .

relaxation or close binary interaction<sup>1</sup>. Looking at their position and velocity in Figure 1, the Gaia BH-like binary belongs to a tidal tail of the open cluster. This means that it will be identified as a Galactic disk component long after the open cluster has evaporated.

We show the time evolution of the Gaia BH-like binary and its progenitor in Figure 2, and illustrate the formation process of the Gaia BH-like binary in Figure 3. At the initial time, its BH progenitor has  $22.6M_{\odot}$  and a companion with  $9.32M_{\odot}$ . The binary period and eccentricity are  $2.8 \times 10^3$  days and 0.167, respectively. At 9.49 Myr, its BH progenitor evolves to a  $16.7M_{\odot}$  BH. At 21.5 Myr, this binary is perturbed by an open cluster star, excites its eccentricity to nearly 1, and merges. This merger leaves behind a  $21.4M_{\odot}$  BH. Until 188 Myr, this BH does not form any hard binary. Although soft binaries are occasionally formed, their binary periods are at least  $\sim 10^7$  days. Such soft binaries are quickly disrupted by perturbations of other stars. At 188 Myr, the BH captures an open cluster star, and form a relatively hard binary with a period of  $5.3 \times 10^5$  days. After this, the BH companion is replaced with another star several times (although omitted in Figure 3). At 244 Myr the BH companion becomes a  $1.59M_{\odot}$  MS star to be finally the tertiary star. At 260 Myr, the BH companion is exchanged with a  $0.82M_{\odot}$  MS star to be finally the secondary star. The superseded companion is not fully ejected from the Gaia BH-like binary, and stays as the tertiary star. Soon after this interaction, this Gaia BH-like binary with the tertiary star escapes from the open cluster.

We can see in Figure 2 that the Gaia BH-like binary eccentricity and mutual inclination between the orbital planes of the inner and outer binaries (hereafter, mutual inclination) are oscillating after the Gaia BH-like binary escapes due to ZLK mechanism. The ZLK oscillation is modulated, and the modulation amplitude appears inconstant. This is because the cadence of the snapshot (4 Myr) is longer than the timescale of the ZLK oscillation. In order to catch the ZLK oscillation by sufficiently high cadence of the snapshot, we recalculate the orbital evolution of the triple system. For this purpose, we extract the masses, positions, and velocities of the triple-system components at 400 Myr, and follow their evolution over 2 Gyr by means of the SDAR code (Wang et al. 2020a), where the SDAR code



**Figure 2.** Time evolution of the Gaia BH-like binary and its progenitor. Secondary and tertiary stars are defined as temporary companions of the BH and its progenitor at each time. We indicate the distance from the center, component masses, binary periods, binary eccentricities, and mutual inclination from top to bottom. These quantities are plotted for each 4 Myr. The shaded regions show binary periods of 100 - 2000 days, which tends to be detected by *Gaia*.

is incorporated into the PeTar code to treat close binary and multiple star systems as described above. We approximate that the triple system is completely isolated from other stars, since the triple system escapes from the open cluster at 400 Myr. We do not take into account the tertiary star evolution to a red giant and white dwarf phases.

Figure 4 shows the eccentricity and mutual inclination of the Gaia BH-like binary. The cadence of the snapshot is 0.1 Myr, which is sufficiently high to show the ZLK oscillation. The oscillation on several Myr and modulation on several 100 Myr are due to the quadrupole-level and octupole-level interactions, respectively (Antognini 2015). The modulation amplitude is constant, although it appears not to be constant due to the low cadence of the snapshot (4 Myr) in Figure 2. We can see that the Gaia BH-like binary eccentricity rises to at most 0.8, and its pericenter distance is reduced down to at least 0.97 au. Since this pericenter distance is much larger than the secondary radius, the Gaia BH-like binary will not merge. Note that the secondary star, a 0.82 MS star, will not evolve to a giant star within the Hubble time. The tertiary star evolves to a red giant star and then a white dwarf over ~ 2 Gyr. The red giant star will not fill the Roche lobe of

<sup>&</sup>lt;sup>1</sup> Such a binary with massive objects can be ejected from an open cluster (Fujii & Portegies Zwart 2011)



**Figure 3.** Illustration of the Gaia BH-like binary formation. Objects indicated by "BH", "2" and "3" are the BH, secondary and tertiary stars in the Gaia BH-like binary, respectively. Object "A" is the initial companion of the BH progenitor. Objects "B", "C", "D" are perturbers for the BH and its companions. Although the Gaia BH-like binary experience many interactions during 188-244 Myr, we omit them. Thus, an ejected object at 244 Myr is not actually object "C".

the outer binary, because the pericenter distance of the outer binary is  $2.0 \times 10^2$  au much larger than the asymptotic giant branch radius ~ 1 au. The tertiary star gradually decreases its mass and influence on the Gaia BH-like binary. The tertiary star evolution is expected not to trigger merger or disruption of the Gaia BH-like binary.

In Figure 5, we compare the formation efficiency among BH binaries in open clusters to the 1st and 2nd types of isolated binary evolution. We focus on only binary periods of 100 - 2000 days, which tends to be detected by *Gaia*<sup>2</sup>. The formation efficiency of all BH binaries in open clusters is much smaller than both the types



**Figure 4.** Time evolution of the Gaia BH-like binary eccentricity (top) and mutual inclination (bottom) over 2 Gyr from 400 Myr. The quantities are plotted for each 0.1 Myr.



**Figure 5.** Formation efficiency of BH binaries per  $M_{\odot}$  for each bin in open clusters (top), the 1st type of isolated binaries (middle), and the 2nd type of isolated binaries (bottom). The stellar type of the companions are limited to MS stars. The grey, blue, and red histograms indicate All BH binaries, BH binaries with  $\leq 1.1 M_{\odot}$  companions, and those with  $\leq 1.1 M_{\odot}$  companions and eccentricities from 0.3 to 0.9 (i.e. Gaia BHs). The shaded regions show binary periods of 100 – 2000 days, which tends to be detected by *Gaia*.

<sup>&</sup>lt;sup>2</sup> The binary periods of Gaia BH1 and BH2 are  $\sim$  180 day (El-Badry et al. 2023a) and  $\sim$  1300 day (El-Badry et al. 2023b; Tanikawa et al. 2023).



**Figure 6.** Formation efficiency of BH binaries per  $M_{\odot}$  for each bin in open clusters, and the 1st and 2nd types of isolated binaries as a function of secondary masses. The top, middle, and bottom panels indicate Z = 0.005, 0.01, and 0.02, respectively. All the secondary stars are MS stars, and all the binary periods and eccentricities are 100-2000 days and 0.3-0.9, respectively.

of isolated binaries. However, when we limit BH binaries to Gaia BHs, i.e. those with light companions  $(1.1M_{\odot})$  and moderately high eccentricities (0.3 - 0.9), the formation efficiency is reversed. It is  $\sim 10^{-5}M_{\odot}^{-1}$  in open clusters, while it is  $\sim 10^{-8}M_{\odot}^{-1}$  even in the 2nd type of isolated binaries. Thus, the formation efficiency is larger by 3 orders of magnitudes.

We again emphasize that the formation of Gaia BHs in open clusters is much higher than in isolated binaries with the respect of initial conditions. The numbers of binaries with >  $20M_{\odot}$  and <  $1.1M_{\odot}$  stars are 0, 0, and  $5.4 \times 10^{-5}$  per  $1M_{\odot}$  in open clusters and the 1st and 2nd types of isolated binaries, respectively. If we limit their orbital periods to 100 - 2000 days, the number is  $9.2 \times 10^{-6}$  per  $1M_{\odot}$  in the 2nd types of isolated binaries. Note that >  $20M_{\odot}$  stars evolve to BHs long before 500 Myr, and <  $1.1M_{\odot}$  stars are still MS stars at 500 Myr if they evolve as single stars; such binaries can be potentially progenitors of Gaia BHs. On the other hand, the formation efficiencies of Gaia BHs are ~  $10^{-5}$ , 0, and ~  $10^{-8}M_{\odot}^{-1}$  in open clusters are more disadvantageous to form Gaia BHs than the 2nd type of isolated binaries. However, open clusters can form many more Gaia BHs.

Figure 6 shows the formation efficiency of BH binaries as a function of secondary masses. From the top panel, we can confirm that, for the secondary mass of less than  $1.1M_{\odot}$ , the formation efficiency in open clusters is larger than in isolated binaries by 3 orders of magnitude for Z = 0.005. If BH binaries with the secondary mass of  $\sim 2M_{\odot}$  can be regarded as Gaia BHs, there is another Gaia BH in a Z = 0.01 open cluster (see the middle panel in Figure 6). Such a BH binary is also formed in open clusters more easily than in isolated binaries by 3 orders of magnitude. Note that we find no tertiary star for this BH binary.

The mass distribution in Figure 6 may have the sharp peak at ~  $1M_{\odot}$  for the open clusters with Z = 0.005 due to the small sampling. If we increase the number of open cluster simulations to much more than 100, the sharp peak is likely to be smeared out around ~  $1M_{\odot}$ . Thus, some of them may exceed  $1M_{\odot}$ , and may not be categorized as Gaia BHs. Nevertheless, it should have a minor effect on the formation efficiency of Gaia BHs in open clusters.

The low formation efficiency in isolated binaries is consistent with the argument of El-Badry et al. (2023a). Here, we review its reason. Let's consider a binary with a BH progenitor ( $\gtrsim 20 M_{\odot}$ ) and a ~  $1 M_{\odot}$ star. Its period is ~  $10^3$  days, and then its separation is ~  $10^3 R_{\odot}$ . When the BH progenitor evolves to a giant star, it fills its Roche lobe, and starts Roche lobe overflow. The Roche lobe overflow is typically unstable, since the donor star (here, the BH progenitor) is much more massive than the accretor (here, the  $1M_{\odot}$  star). Thus, the binary experiences common envelope evolution. The common envelope evolution ends with a merger of the BH progenitor core and ~  $1M_{\odot}$  star. This is because the gravitational binding energy between the core and star is too small to fully eject the common envelope. Even if the binary survives against the common envelope evolution, its orbital period is much shorter than the orbital periods of Gaia BHs, and its eccentricity is reduced to  $\leq 0.2$  (e.g. Kruckow et al. 2021; Trani et al. 2022). The BH natal kick could excite its eccentricity if its velocity is sufficiently high. However, the barycenter velocities of the Gaia BHs are too small to suppose such high BH natal kick velocities.

In contrast to isolated binaries, the formation of Gaia BHs in open clusters avoids all the difficulties described above. As illustrated in Figure 3, Gaia BHs can be formed through dynamical capture after BH progenitors evolve to BHs in open clusters. If we dare to mention the difficulty of Gaia BH formation in open clusters, BHs may rarely capture  $\sim 1M_{\odot}$  stars. Generally, BHs capture more massive stars more easily, and there are many  $\gtrsim 1M_{\odot}$  MS and white dwarfs in open clusters.

Taking into account only the Gaia BH-like binary, we estimate the number of Gaia BHs formed in open clusters over Milky Way as follows:

$$N_{\text{GaiaBH,MW}} \sim 1.6 \times 10^4 \left(\frac{\eta}{10^{-5} M_{\odot}}\right) \left(\frac{M_{\text{MW}}}{6.1 \times 10^{10} M_{\odot}}\right) \\ \times \left(\frac{f_{Z=0.005}}{0.26}\right) \left(\frac{f_{\text{cluster}}}{0.1}\right), \tag{1}$$

where  $\eta$  is the formation efficiency obtained from our simulation results,  $M_{\text{MW}}$  is Milky Way stellar mass (e.g. Licquia & Newman 2015), and  $f_{Z=0.005}$  is the mass fraction of Z = 0.005 in the Milky Way, which is calculated by integrating stellar mass between Z =0.003 and 0.007 in a Milky Way model of Wagg et al. (2022) and Shikauchi et al. (2023). The fraction  $f_{\text{cluster}}$  is the mass fraction of stars formed in open clusters (Misiriotis et al. 2006; Piskunov et al. 2007). We assume that the lifetime of Gaia BHs is larger than the Hubble time because of the small mass of the Gaia BH-like binary. Since the size of Milky Way disk is ~ 10 kpc, the number of Gaia BHs can be ~ 160 within ~ 1 kpc. We expect that the formation efficiency in open clusters will explain the number of Gaia BHs, even if the number of Gaia BHs (currently, 2) will grow in the future as seen in the lists of BH binary candidates (Andrews et al. 2022; Shahaf et al. 2023).

#### **4 CONCLUSION AND DISCUSSION**

Our main conclusions in this paper are as follows.

(i) Formation efficiency of Gaia BHs in open clusters is higher than in isolated binaries by 3 orders of magnitude.

(ii) Gaia BHs are in multiple star systems with a high probability.

We find a Gaia BH-like binary in our simulations albeit of the small total mass of open clusters. The Gaia BH-like binary has many points similar to Gaia BHs. The visible star has ~  $1M_{\odot}$ , and the binary period and eccentricity are  $10^2$ - $10^3$  days and ~ 0.5, respectively. Its barycenter orbit indicates that it is a Galactic disk component. Since the visible star is dynamically captured by the BH after the BH formation, it should not be polluted by supernova ejecta from the BH, which is consistent with the chemical abundance pattern of Gaia BH2 (El-Badry et al. 2023b). The visible star has subsolar metallicity similar to Gaia BH1, whose visible star has  $[Fe/H] = -0.2 \pm 0.05$ (El-Badry et al. 2023a). The number of such BH binaries is large enough to explain the presence of Gaia BHs. The Gaia BH-like binary contains different points from Gaia BHs, especially Gaia BH2. The visible star of Gaia BH-like binary has subsolar metallicity and does not evolve to a red giant star within the Hubble time, while Gaia BH2's visible star has solar metallicity, and already evolves to a red giant star. However, if we simulate the dynamical evolutions of many more open clusters, we are likely to generate BH binaries similar to Gaia BH2.

We expect that Gaia BHs frequently have tertiary stars if they are formed in open clusters. The frequency of the presence of tertiary stars is highly uncertain. If we estimate the frequency from the small samples, the frequency is 100 %, because the Gaia BH-like binary, which is just one Gaia BH in our simulation, has a tertiary star. Even if we include the BH binary with the secondary mass of ~  $2M_{\odot}$ , this frequency is not changed much because of its short lifetime. Even if we take into account more many runs in Appendix B, the frequency is still high  $\sim 50$  %. Nevertheless, we recognize that this is a rough estimate. In order to determine the frequency accurately, we need to perform many more cluster simulations, and acquire a larger number of samples than now. On the other hand, there are no reports for such tertiary stars in Gaia BHs. Its reason may be just that no one searches for such tertiary stars (but see El-Badry et al. 2021). Moreover, the tertiary star of the Gaia BH-like binary becomes a white dwarf after  $\sim 2$  Gyr, and may be too faint to be detected by *Gaia* at the present day. Thus, we are not always able to discover such tertiary stars if anv

As for multiplicity, our results show that a visible binary can orbit a BH. In other words, the BH itself is a tertiary star of the multiple star system. We observe several binaries orbiting BHs in our simulations. However, their outer binary periods are  $> 10^5$  days. They cannot be discovered by *Gaia*, since their periods are much larger than *Gaia*'s operation duration, a few  $10^3$  days. We may generate such systems detectable by *Gaia* if we increase the number of open cluster simulations again.

It appears that there are large uncertainties in our results, in other words, the formation efficiency of Gaia BHs and multiplicity frequency. This is derived from only one Gaia BHs formed from open clusters with  $3 \times 10^5 M_{\odot}$  in total. However, we expect that our results would be robust. This is because our results are not different from those in Appendix B by an order of magnitude. In Appendix B, we

investigate open clusters with  $8 \times 10^6 M_{\odot}$  in total and the number of Gaia BH-like binaries is 11.

At the moment when we posted this paper to arXiv, only Shikauchi et al. (2020) have previously investigated Gaia BHs formed in open clusters, although many studies have studied Gaia BHs formed in isolated binaries (Mashian & Loeb 2017; Breivik et al. 2017; Yamaguchi et al. 2018; Kinugawa & Yamaguchi 2018; Yalinewich et al. 2018; Shao & Li 2019; Andrews et al. 2019; Wiktorowicz et al. 2020; Chawla et al. 2022; Shikauchi et al. 2022, 2023). They have not found BH binaries with  $\sim 1M_{\odot}$  MS stars despite that they treat open clusters with  $9 \times 10^5 M_{\odot}$  in total, 3 times larger than those in this paper. No Gaia BH-like binary in Shikauchi et al. (2020) can be explained by small statistics. No Gaia BH-like binary can be formed even in some of model grids in which open clusters have  $10^6 M_{\odot}$  in total according to Appendix B regardless of binary fractions.

We should make a caveat that the presence of tertiary stars of Gaia BHs does not mean the evidence of Gaia BH formation in open clusters. A few 10 % of galactic field stars are in triple and quadruple systems (Tokovinin 2014; Moe & Di Stefano 2017). Such multiple systems may form Gaia BHs with tertiary stars. We expect that this possibility will be investigated owing to rapid progress for studying evolution of multiple systems (Hamers et al. 2021; Toonen et al. 2022; Stegmann et al. 2022; Dorozsmai et al. 2023), although it has not yet.<sup>3</sup> Comparison between tertiary stars of Gaia BHs formed in open clusters and multiple systems will helpful for identifying the origin of Gaia BHs. In any case, what we can say now is that the open cluster scenario will be ruled out if no Gaia BHs have tertiary stars at all.

In the near future, we will perform a larger number of open cluster simulations than in this paper. These open clusters will have different mass and density, and will be in different tidal fields. It will reveal the formation efficiency of Gaia BHs in open clusters more accurately. We will analyze the distribution of BH and visible star masses, binary periods, eccentricities, and frequency of the presence of tertiary stars. It is also important to obtain the distribution of the tertiary star masses, and outer binary periods and eccentricities. If an outer binary has a short enough period to cause the ZLK oscillation of its inner BH binary, we may observe an eccentricity variability of the BH binary (Hayashi & Suto 2020). We may also see visible binaries orbiting BHs with periods of 100-2000 days. The presence of such binaries may be indicated by differences between spectroscopic and astrometric mass functions for each binary (Tanikawa et al. 2023). These distributions should be a clue to elucidate the formation process of Gaia BHs.

This paper indicates two important possibilities. First, we do not need to rebuild the theory of common envelope evolution, which is carefully constructed to explain several types of compact binaries, such as double BHs (e.g. Belczynski et al. 2016a). Second, if Gaia BHs are dominantly formed in open clusters, they can be a useful probe to study open cluster dynamics.

#### ACKNOWLEDGMENTS

This research could not be accomplished without the support by Grants-in-Aid for Scientific Research (17H06360, 19K03907) from the Japan Society for the Promotion of Science. M.F. is supported by The University of Tokyo Excellent Young Researcher Program. S.C.

<sup>&</sup>lt;sup>3</sup> El-Badry et al. (2023a,b) form Gaia BHs from triple systems. However these Gaia BHs do not have tertiary stars, since the inner binaries merge in the halfway of their evolutions.

is supported by the Fulbright U.S. Student Program, which is sponsored by the U.S. Department of State and Japan-U.S. Educational Commission. Its contents are solely the responsibility of the author and do not necessarily represent the official views of the Fulbright Program, the Government of the United States, or the Japan-U.S. Educational Commission. M.S. thanks a support by Research Fellowships of Japan Society for the Promotion of Science for Young Scientists, by Forefront Physics and Mathematics Program to Drive Transformation (FoPM), a World-leading Innovative Graduate Study (WINGS) Program, the University of Tokyo, and by JSPS Overseas Challenge Program for Young Researchers. L.W. thanks the support from the one-hundred-talent project of Sun Yat-sen University, the Fundamental Research Funds for the Central Universities, Sun Yatsen University (22hytd09) and the National Natural Science Foundation of China through grant 12073090 and 12233013. Numerical simulations are carried out on Small Parallel Computers at Center for Computational Astrophysics, National Astronomical Observatory of Japan, the Yukawa Institute Computer Facility, and Cygnus/Pegasus at the CCS, University of Tsukuba.

#### DATA AVAILABILITY

Results will be shared on reasonable request to authors. The data is generated by the software PeTar and SDAR, which are available in GitHub at https://github.com/lwang-astro/PeTar and https://github.com/lwang-astro/SDAR, respectively. The initial conditions of star cluster models are generated by the software MCLUSTER (Küpper et al. 2011), which is available in GitHub at https://github.com/lwang-astro/mcluster.

### APPENDIX A: COMPARISON BETWEEN PETAR AND NBODY6++GPU

We make comparison between simulation results of PeTar and NB0DY6++GPU (Nitadori & Aarseth 2012; Wang et al. 2015; Kamlah et al. 2022). Indeed, Wang et al. (2020b) have already done in detail. However, they have treated small globular clusters with ~  $10^5 M_{\odot}$ , while we treat open clusters with ~  $10^3 M_{\odot}$ . Thus, it might be worth comparing simulation results between PeTar and NB0DY6++GPU with respect of cluster mass scale of ~  $10^3 M_{\odot}$ .

We generate 10 open clusters for each code. The metallicity is Z = 0.02, and initial binary fractions are 20 %. We do not take into account galactic tides, since NB0DY6++GPU does not support combination with GALPY. The other setup is the same as section 2. Note that we do not remove any stars from simulations even when stars go away from open clusters. This is true for simulations in section 2.

Figures A1 and A2 shows the cluster evolution and binary properties at 500 Myr. We do not find any different features between PeTar and NBODY6++GPU results. This indicates that PeTar works well even for mass scale of open clusters.

## APPENDIX B: EFFECTS OF INITIAL BINARY FRACTIONS

We investigate how the formation efficiency of Gaia BHs depend on initial binary fractions. We adopt the fractions of 0, 20, 50, and 100 % for open clusters with Z = 0.02 and 0.005. We generate 1000 open clusters for each fraction and each metallicity in order to reduce statistical uncertainties.



Figure A1. (Top) Time evolution of the total masses of single and binary stars. (Bottom) Time evolution of 10, 50, and 90 % Lagrangian radii.

The top panel of Figure B2 shows that the formation efficiency of Gaia BHs in open clusters is not sensitive to initial binary fractions in open clusters. Although the error bars are large, the average values are not dependent on the initial binary fractions. The bottom panel of Figure B2 indicates little dependence of multiplicity on initial binary fractions. Although the results suffer from small statistics, the multiplicity does not monotonically increase with initial binary fractions does not affect our results in section 3.

The top panel of Figure B2 shows  $\eta \sim 2 \times 10^{-6} M_{\odot}^{-1}$  for all binary



Figure A2. (Top) Distribution of binary semi-major axes and eccentricity at 500 Myr. The point sizes show binary masses on linear scale. (Bottom) Distribution of binary mass and mass ratio at 500 Myr. The point size indicates binary semi-major axes on logarithmic scale.

fractions and metallicities. Then, we can rewrite Equation (1) as

$$N_{\text{GaiaBH,MW}} \sim 1.2 \times 10^4 \left(\frac{\eta}{2 \times 10^{-6} M_{\odot}}\right) \left(\frac{M_{\text{MW}}}{6.1 \times 10^{10} M_{\odot}}\right) \\ \times \left(\frac{f_{\text{cluster}}}{0.1}\right). \tag{B1}$$

This is similar to the estimate in Equation (1).

We show the formation efficiency and multiplicity fraction of Gaia BHs. Here, we change the definition of Gaia BHs to mimic the definition in Di Carlo et al. (2023), such that period is 0 - 4000 days, eccentricity is 0.0 - 1.0, and the companion mass is  $< 0.8M_{\odot}$ . Compared to the definition in section 2, the formation efficiency is



**Figure B1.** (Top) Formation efficiency of Gaia BHs in open clusters (solid curves) as a function of initial binary fractions. Error bars indicate standard errors under assumption that Gaia BH formation is a Poisson process. Dashed lines show the formation efficiency in isolated binaries. In order to be easy to see, the initial binary fractions are shifted by 0.125 leftward and rightward for Z = 0.02 and 0.05, respectively. The definition of Gaia BHs is the same as section 2. The black dotted curve indicates the formation efficiency proportional to the initial binary fraction. (Bottom) Fraction of Gaia BHs in multiple star systems as a function of initial binary fractions. We calculate standard errors, assuming that Gaia BHs obtain tertiary stars in a Poisson process. The initial binary fractions are shifted in the same way as the top panel.

increased to ~  $10^{-5}M_{\odot}^{-1}$  by 10 times. This is in good agreement with the results of Di Carlo et al. (2023). The multiplicity of Gaia BHs in open clusters is ~ 0.5. As seen in the top panel, the formation efficiency may increase with metallicity decreasing. However, metallicity dependence is beyond the scope of this paper.

In summary, the formation efficiency of Gaia BHs is quite consistent with the results of Rastello et al. (2023) and Di Carlo et al. (2023). Our formation efficiency looks different from the formation efficiency of Di Carlo et al. (2023). However, the difference comes from the difference of the definitions of Gaia BHs.

#### REFERENCES

Andrews J. J., Breivik K., Chatterjee S., 2019, ApJ, 886, 68

- Andrews J. J., Taggart K., Foley R., 2022, arXiv e-prints, p. arXiv:2207.00680 Antognini J. M. O., 2015, MNRAS, 452, 3610
- Askar A., Szkudlarek M., Gondek-Rosińska D., Giersz M., Bulik T., 2017, MNRAS, 464, L36

Banerjee S., 2021, MNRAS, 503, 3371



**Figure B2.** The same as Figure B2, except for the definition of Gaia BHs. Period and eccentricity criteria are relaxed, such that the former and latter are 0 - 4000 days, and 0.0 - 1.0, respectively. The companion criteria become more strict, such that  $< 0.8M_{\odot}$ . Note that  $< 0.8M_{\odot}$  stars do not evolve to giant stars within the Hubble time.

- Banerjee S., Belczynski K., Fryer C. L., Berczik P., Hurley J. R., Spurzem R., Wang L., 2020, A&A, 639, A41
- Belczynski K., Bulik T., Fryer C. L., Ruiter A., Valsecchi F., Vink J. S., Hurley J. R., 2010, ApJ, 714, 1217
- Belczynski K., Holz D. E., Bulik T., O'Shaughnessy R., 2016a, Nature, 534, 512
- Belczynski K., et al., 2016b, A&A, 594, A97
- Bovy J., 2015, ApJS, 216, 29
- Breivik K., Chatterjee S., Larson S. L., 2017, ApJ, 850, L13
- Casares J., Jonker P. G., Israelian G., 2017, X-Ray Binaries. p. 1499, doi:10.1007/978-3-319-21846-5\_111
- Chakrabarti S., et al., 2023, AJ, 166, 6
- Chawla C., Chatterjee S., Breivik K., Moorthy C. K., Andrews J. J., Sanderson R. E., 2022, ApJ, 931, 107
- Claeys J. S. W., Pols O. R., Izzard R. G., Vink J., Verbunt F. W. M., 2014, A&A, 563, A83
- Di Carlo U. N., Giacobbo N., Mapelli M., Pasquato M., Spera M., Wang L., Haardt F., 2019, MNRAS, 487, 2947
- Di Carlo U. N., Agrawal P., Rodriguez C. L., Breivik K., 2023, arXiv e-prints, p. arXiv:2306.13121
- Dorozsmai A., Toonen S., Vigna-Gómez A., de Mink S. E., Kummer F., 2023, arXiv e-prints, p. arXiv:2307.04793
- El-Badry K., Rix H.-W., Heintz T. M., 2021, MNRAS, 506, 2269
- El-Badry K., et al., 2023a, MNRAS, 518, 1057
- El-Badry K., et al., 2023b, MNRAS, 521, 4323
- Fryer C. L., Belczynski K., Wiktorowicz G., Dominik M., Kalogera V., Holz D. E., 2012, ApJ, 749, 91
- Fujii M. S., Portegies Zwart S., 2011, Science, 334, 1380
- Gaia Collaboration et al., 2022, arXiv e-prints, p. arXiv:2206.05595
- Hamers A. S., Rantala A., Neunteufel P., Preece H., Vynatheya P., 2021, MNRAS, 502, 4479
- Hayashi T., Suto Y., 2020, ApJ, 897, 29
- Hurley J. R., Pols O. R., Tout C. A., 2000, MNRAS, 315, 543

- Hurley J. R., Tout C. A., Pols O. R., 2002, MNRAS, 329, 897
- Iwasawa M., Tanikawa A., Hosono N., Nitadori K., Muranushi T., Makino J., 2016, PASJ, 68, 54
- Iwasawa M., Oshino S., Fujii M. S., Hori Y., 2017, PASJ, 69, 81
- Iwasawa M., Namekata D., Nitadori K., Nomura K., Wang L., Tsubouchi M., Makino J., 2020, PASJ, 72, 13
- Kamlah A. W. H., et al., 2022, MNRAS, 511, 4060
- Kinugawa T., Yamaguchi M. S., 2018, arXiv e-prints,
- Kozai Y., 1962, AJ, 67, 591
- Kremer K., Ye C. S., Kıroğlu F., Lombardi J. C., Ransom S. M., Rasio F. A., 2022, ApJ, 934, L1
- Kroupa P., 2001, MNRAS, 322, 231
- Kruckow M. U., Neunteufel P. G., Di Stefano R., Gao Y., Kobayashi C., 2021, ApJ, 920, 86
- Kumamoto J., Fujii M. S., Tanikawa A., 2019, MNRAS, 486, 3942
- Kumamoto J., Fujii M. S., Tanikawa A., 2020, MNRAS, 495, 4268
- Küpper A. H. W., Maschberger T., Kroupa P., Baumgardt H., 2011, MNRAS, 417, 2300
- Licquia T. C., Newman J. A., 2015, ApJ, 806, 96
- Lidov M. L., 1962, Planet. Space Sci., 9, 719
- Mashian N., Loeb A., 2017, MNRAS, 470, 2611
- Misiriotis A., Xilouris E. M., Papamastorakis J., Boumis P., Goudis C. D., 2006, A&A, 459, 113
- Moe M., Di Stefano R., 2017, ApJS, 230, 15
- Nitadori K., Aarseth S. J., 2012, MNRAS, 424, 545
- Oshino S., Funato Y., Makino J., 2011, PASJ, 63, 881
- Piskunov A. E., Schilbach E., Kharchenko N. V., Röser S., Scholz R. D., 2007, A&A, 468, 151
- Portegies Zwart S. F., McMillan S. L. W., 2000, ApJ, 528, L17
- Rastello S., Mapelli M., Di Carlo U. N., Iorio G., Ballone A., Giacobbo N., Santoliquido F., Torniamenti S., 2021, MNRAS, 507, 3612
- Rastello S., Iorio G., Mapelli M., Arca-Sedda M., Di Carlo U. N., Escobar G. J., Torniamenti S., Shenar T., 2023, arXiv e-prints, p. arXiv:2306.14679
- Rodriguez C. L., Chatterjee S., Rasio F. A., 2016, Phys. Rev. D, 93, 084029
- Sana H., et al., 2012, Science, 337, 444
- Shahaf S., Bashi D., Mazeh T., Faigler S., Arenou F., El-Badry K., Rix H. W., 2023, MNRAS, 518, 2991
- Shao Y., Li X.-D., 2019, ApJ, 885, 151
- Shenar T., et al., 2022, Nature Astronomy, 6, 1085
- Shikauchi M., Kumamoto J., Tanikawa A., Fujii M. S., 2020, PASJ, 72, 45
- Shikauchi M., Tanikawa A., Kawanaka N., 2022, ApJ, 928, 13
- Shikauchi M., Tsuna D., Tanikawa A., Kawanaka N., 2023, arXiv e-prints, p. arXiv:2301.07207
- Stegmann J., Antonini F., Moe M., 2022, MNRAS, 516, 1406
- Tagawa H., Haiman Z., Kocsis B., 2020, ApJ, 898, 25
- Tanikawa A., 2013, MNRAS, 435, 1358
- Tanikawa A., Hattori K., Kawanaka N., Kinugawa T., Shikauchi M., Tsuna D., 2023, ApJ, 946, 79
- The LIGO Scientific Collaboration et al., 2021, arXiv e-prints, p. arXiv:2111.03606
- Tokovinin A., 2014, AJ, 147, 87
- Toonen S., Boekholt T. C. N., Portegies Zwart S., 2022, A&A, 661, A61
- Trani A. A., Rieder S., Tanikawa A., Iorio G., Martini R., Karelin G., Glanz H., Portegies Zwart S., 2022, Phys. Rev. D, 106, 043014
- Wagg T., Broekgaarden F. S., de Mink S. E., Frankel N., van Son L. A. C., Justham S., 2022, ApJ, 937, 118
- Wang L., Spurzem R., Aarseth S., Nitadori K., Berczik P., Kouwenhoven M. B. N., Naab T., 2015, MNRAS, 450, 4070
- Wang L., Nitadori K., Makino J., 2020a, MNRAS, 493, 3398
- Wang L., Iwasawa M., Nitadori K., Makino J., 2020b, MNRAS, 497, 536
- Wang L., Tanikawa A., Fujii M., 2022, MNRAS, 515, 5106
- Webbink R. F., 1984, ApJ, 277, 355
- Wiktorowicz G., Lu Y., Wyrzykowski Ł., Zhang H., Liu J., Justham S., Belczynski K., 2020, ApJ, 905, 134
- Yalinewich A., Beniamini P., Hotokezaka K., Zhu W., 2018, MNRAS, 481, 930
- Yamaguchi M. S., Kawanaka N., Bulik T., Piran T., 2018, ApJ, 861, 21

### 10 Ataru Tanikawa

- Ziosi B. M., Mapelli M., Branchesi M., Tormen G., 2014, MNRAS, 441, 3703
- von Zeipel H., 1910, Astronomische Nachrichten, 183, 345
- This paper has been typeset from a  $T_{\ensuremath{E}} X/I \!\! \ensuremath{\Delta} T_{\ensuremath{E}} X$  file prepared by the author.