Evaluating Sensitivity to the Stick-Breaking Prior in Bayesian Nonparametrics (Rejoinder)

Ryan Giordano^{*,‡}, Runjing Liu^{†,‡}, Michael I. Jordan[†], and Tamara Broderick^{*}

1 Introduction

We feel very grateful to have our work carefully read and commented on by so many insightful respondents. We would like to thank Professor Steel and the editorial board of Bayesian Analysis for selecting our work and making this discussion possible. Statistical robustness is a venerable topic of conversation and we have no doubt that the present discussion will continue far into the future.

We might roughly categorize points made in the responses as follows:¹

- 1. Would a different model or summary statistic be more or less robust than the Dirichlet process (DP) and number of clusters?
- 2. Can (or should) one form variational Bayes (VB) approximations to different models from the BNP literature?
- 3. Can one form a local sensitivity metric to different quantities of interest, modeling or fitting parameters, different posterior approximation procedures, or some combination of all of these?

Questions in category (1) and (2) are natural and important, since our work is based on arguably the most canonical Bayesian nonparametric prior (the DP), and a fairly vanilla VB approximation (a mean field approximation to a truncated stick-breaking representation). The evaluation of our robustness ideas with respect to a wider range of priors and approximations is certainly warranted. Nevertheless, in the present rejoinder we will focus on questions in category (3), largely because we feel that our use of local sensitivity metrics constitutes our work's most distinctive contribution.

Can one form a local robustness metric for a particular problem? In section 2 of the present rejoinder, we will argue: typically, yes, quite directly, in Markov chain Monte Carlo (MCMC) applications as well as VB. In section 2.1 we derive local robustness metrics for a select few settings that were described in the responses. After reading

^{*}Department of EECS, MIT, 77 Massachusetts Ave., 38-401, Cambridge, MA 02139

[†]Department of Statistics, 367 Evans Hall, UC Berkeley, Berkeley, CA 94720

[‡]Equal contribution.

¹Please think of these categories as atom locations from an Indian buffet process, not a Chinese restaurant process; respondents engaged at times with multiple categories simultaneously. For example, **Gil-Leyva and Mena** ask how to form a VB approximation to an exchangeable stick breaking (ESB) prior (item 2) in order to form a local sensitivity metric (item 3) to assess whether the ESB prior is robust for pointwise density estimation (item 1).

section 2, we hope that all readers of this rejoinder feel able and empowered to form and investigate local robustness metrics for their own particular problems.

However, in the subsequent section 3, we will argue that simply forming a local robustness metric is not enough: the hard work is showing that it is useful. Computability, interpretability, and the ability of a local robustness metric to *extrapolate* well, are more important — and more difficult to establish — than mere computation of derivatives. It is this work of establishing usefulness that we have endeavored to undertake in the present paper, and to which we wish to call attention as a foundation for further work.

As might be expected in a topic as established as robustness, the points made in this rejoinder are not new, and have in fact been argued in the past by many of our own respondents. Nevertheless, we hope that by emphasizing the relative ease of computing derivatives and the relative difficulty of showing their utility in particular contexts, we can help advance the research agenda in this important and challenging area.

2 What does it take to do local robustness?

A great deal of statistical inference—including Bayesian statistics—fixes some hyperparameter ω and then performs posterior inference using some combination of two types of estimators:

- The solution to a system of estimating equations: $\hat{\theta}_{opt} := \theta$ such that $G(\theta, \omega) = 0$.
- A posterior moment from a density known up to a constant: $\hat{\theta}_{samp} := \underset{\mathcal{P}(\zeta|\omega)}{\mathbb{E}} [H(\zeta)].$

An example of $\hat{\theta}_{opt}$ could be the parameter that sets the gradient of a VB loss function to zero (as in our paper), and an example of $\hat{\theta}_{samp}$ could be a posterior mean. In practice, we may not be able to compute either exactly: $\hat{\theta}_{opt}$ might be approximated using numerical optimization, and $\hat{\theta}_{samp}$ may be approximated using Markov chain Monte Carlo (MCMC). Below, we will briefly discuss the consequences of using approximations, but assume for the moment that we can compute $\hat{\theta}_{opt}$ and $\hat{\theta}_{samp}$ to a desired accuracy. A practitioner might then ask, "what would happen if ω had taken on some different value?" The techniques of local robustness approximately answer this question by forming a series approximation using derivatives of the maps $\omega \mapsto \hat{\theta}_{opt}(\omega)$ and $\omega \mapsto \hat{\theta}_{samp}(\omega)$.

There are simple, general formulas for both these derivatives, under certain common (but not universal) regularity conditions. For notational simplicity, take ω to be a scalar for the moment. Furthermore, let us take $\hat{\theta}_{opt}(\omega)$ to be finite-dimensional, $\mathcal{P}(\zeta|\omega)$ to be defined as a Radon-Nikodym derivative with respect to a common dominating measure for all ω , assume that we can exchange integration and differentiation as needed, and assume all needed partial derivatives exist. Then

$$\frac{d\hat{\theta}_{\rm opt}(\omega)}{d\omega}\bigg|_{\omega_0} = \left(\frac{\partial G(\theta,\omega)}{\partial \theta}\bigg|_{\theta=\hat{\theta}_{\rm opt}(\omega_0),\omega=\omega_0}\right)^{-1} \frac{\partial G(\theta,\omega)}{\partial \omega}\bigg|_{\theta=\hat{\theta}_{\rm opt}(\omega_0),\omega=\omega_0} \tag{1}$$

Giordano, Liu, Jordan, and Broderick

$$\frac{d\hat{\theta}_{\mathrm{samp}}(\omega)}{d\omega}\bigg|_{\omega_0} = \operatorname{Cov}_{\mathcal{P}(\zeta|\omega_0)}\left(H(\zeta), \left.\frac{\partial\log\mathcal{P}(\zeta|\omega)}{\partial\omega}\right|_{\omega=\omega_0}\right).$$
(2)

These two formulas have been noted many times in the literature, though we feel it is worth calling attention to the simplicity of their form. The estimating equation derivative in eq. (1) is formed using the implicit function theorem (Krantz and Parks, 2012) and is used, explicitly or implicitly, in many local robustness works (Hampel, 1974; Thomas and Cook, 1989; Hattori and Kato, 2009; Shi et al., 2016), as well as our own present paper. The sampling derivative in eq. (2) is formed by differentiating under the integral using a dominated convergence theorem (Billingsley, 2008, Theorem 5.4) and appears widely, in various forms, in the local Bayesian robustness literature and beyond (Diaconis and Freedman, 1986; Ruggeri and Wasserman, 1993; Gustafson, 1996; Mohamed et al., 2020).

Importantly, eqs. (1) and (2) can be computed nearly automatically using automatic differentiation, using only the original solution $\hat{\theta}_{opt}(\omega_0)$ or the ability to compute moments of $\mathcal{P}(\zeta|\omega_0)$. Furthermore, higher-order derivatives can be computed mechanically by repeatedly applying eqs. (1) and (2) to themselves. For example, higher-order versions of eq. (1) can be found in Giordano et al. (2019a). In practice, one can use a corresponding numerical approximation to $\hat{\theta}_{opt}(\omega_0)$ or draws from $\mathcal{P}(\zeta|\omega_0)$ to approximate the derivatives. As observed by **Griffin and Kalli**, the key practical difficulty with eq. (1) is the solution of a linear system, and the key practical difficulty with eq. (2) is Monte Carlo error.

One might contrast eqs. (1) and (2) with approaches that differentiate the optimization procedure or the sampling procedure directly, as in chapter 6 of Maclaurin (2016) and Jacobi et al. (2018), respectively, both of which require considerable bespoke computational effort, even with automatic differentiation. The simplicity of eqs. (1) and (2) comes at a cost, however, of assuming (respectively) that $\hat{\theta}_{opt}$ actually solves the estimating equation, or that we are actually able to approximate draws from $\mathcal{P}(\zeta|\omega_0)$. Studying eqs. (1) and (2) in the presence of violations of these assumptions is exciting and ongoing work, most notably in the setting of optimization (Bae et al., 2022).

Although eqs. (1) and (2) apply to scalar ω , they extend readily to multivariate and even functional derivatives, since one can use scalar derivatives to differentiate along a path in a multivariate space. Different notions of "derivative" differ only in how they conceptually bundle these path derivatives together—as a basis for a gradient in finite dimensional vector spaces (Fleming, 2012), as a basis for a tangent plane in a geometric perspective (McInerney, 2013; Murray and Rice, 1993), as Hadamard or Fréchet derivatives in infinite-dimensional spaces according to the smoothness of the underlying function (Averbukh and Smolyanov, 1967; Zeidler, 1986). In each case, however, for a particular path, the derivative is always formally the same, and computable by eqs. (1) and (2).

Given estimators of the form $\hat{\theta}_{opt}$, $\hat{\theta}_{samp}$, or some smooth combination of them, the capacity to imagine a parameterized set of perturbations of interest (and a class of

univariate paths through it), one can form local robustness measures—even for infinitedimensional perturbations—using little more than eqs. (1) and (2), univariate calculus, and the chain rule. In the following section, we will demonstrate this point in a few of the settings described by the respondents.

2.1 Some requested derivatives

We now demonstrate our claim that derivatives are typically straightforward to compute by doing so for several of the settings requested by our respondents: case influence measures (Cook, 1977), exchangeable stick breaking (ESB) processes (Gil-Leyva and Mena, 2021), empirical Bayes (EB) procedures (McAuliffe et al., 2006), and robustness to the loss function. For ESB processes and EB we will discuss some interesting challenges that arise from working with infinite-dimensional priors in BNP models.

Many respondents noted that our (classical) approach to deriving local robustness measures extends readily to other settings. Readers who are similarly convinced that it is not difficult to compute local robustness derivatives for a wide range of applications, for both MCMC and optimization-based statistical procedures, can safely skip to section 3.

For this short rejoinder we have selected only a few settings from the responses to address in detail, and we have chosen to prioritize the settings that are most unlike the results in our paper. Unfortunately, doing so means forgoing discussion of some ideas which seem particularly promising to us, such as the proposal of **Miscouridou and Panero** to apply local robustness techniques to the VB approximations for generalized gamma processes given in Lee et al. (2016).

Throughout this section, we will take ζ to denote all parameters of a model and X to denote observed data, so that the posterior is $\mathcal{P}(\zeta|X)$. Let $\phi(\zeta)$ be some quantity of interest. Note that, in eq. (2), we need only differentiate $\log \mathcal{P}(\zeta, X|\omega)$ rather than $\log \mathcal{P}(\zeta|X, \omega)$, because the normalizing constant $\mathcal{P}(X|\omega)$ does not depend on ζ and does not contribute to the covariance.

Case influence. MacEachern and Lee connect our work to a long history of frequentist and Bayesian "case influence" literature. This literature attempts to quantify the importance of individual datapoints or groups of datapoints on a particular inferential procedure. MacEachern and Lee point to a set of works, beginning with Cook (1977), which is particularly concerned with "outliers" or "gross errors" as popularized by Huber (1964).² Indeed, the idea of using local approximations to robustness under generic data perturbations goes back even farther—at least as far as von Mises (1947)—and has been employed for asymptotic theory (Serfling, 1980; Shao and Tu, 2012; van der Vaart and Wellner, 1996), design and analysis of robust estimators (Hampel, 1974), approximation of cross-validation in machine learning (Koh and Liang, 2017; Giordano et al., 2019b) and more.

To connect this broad literature to our work, we can augment each datapoint with a scalar-valued weight, w_n , in such a way that $w_n = 1$ represents no change, and $w_n = 0$

 $^{^{2}}$ A short historical account of this branch of robust statistics is given by Stigler (2010).

represents omitting the datapoint from the model. Specifically, letting $w = (w_1, \ldots, w_N)$ and $X = (X_1, \ldots, X_N)$, we can write the log likelihood in a Bayesian model as

$$\log \mathcal{P}(X,\zeta|w) = \sum_{n=1}^{N} w_n \log \mathcal{P}(X_n|\zeta) + \log \mathcal{P}(\zeta)$$

with $\mathcal{P}(\zeta|X, w)$ representing the corresponding posterior. With unit weights, we recover the original posterior: $\mathcal{P}(\zeta|X, w = (1, \ldots, 1)) = \mathcal{P}(\zeta|X)$. When $w_n = 0$ but all other weights are one, data point n is left out. Similarly, one can drop or replicate any set of data points using the appropriate configuration of zeros, ones, or other integers.

The advantage of writing $\log \mathcal{P}(\zeta | X, w)$ in this way is that we can take a particular w_n to be our hyperparameter ω and apply eqs. (1) and (2) to form a local approximation to leaving out (or replicating) sets of datapoints. The form of the derivative for estimating equations resulting from eq. (1) is the well-known empirical influence function for Mestimators (see, e.g., eq. 2.3.5 of Hampel et al. (1986)). Perhaps less widely known is the corresponding result for MCMC estimators, which is

$$\frac{\partial \underset{\mathcal{P}(\zeta|X,w)}{\mathbb{E}} [\phi(\zeta)]}{\partial w_n} \bigg|_{w=(1,\dots,1)} = \underset{\mathcal{P}(\zeta|X)}{\operatorname{Cov}} (\phi(\zeta), \log \mathcal{P}(X_n|\zeta)).$$
(3)

The right-hand side of eq. (3) can be estimated from MCMC samples. Then the quantity given in eq. (3) is precisely the "Bayesian empirical influence function," evaluated at X_n , for the statistic $\underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} [\phi(\zeta)]$. As with the frequentist influence function, eq. (3) may be used to approximate all sorts of case deletion schemes from both the frequentist and Bayesian literature—as long as one can show that it provides a good approximation to the effect of actually removing the points.³

Dirichlet-driven ESB models. Gil-Leyva and Mena ask about local sensitivity analvsis for Dirichlet-driven exchangeable stick breaking (ESB) models (Gil-Levva and Mena, 2021). The joint stick distribution in an ESB model is controlled by a parameter $\rho \in [0, 1]$ that smoothly transitions between stick-breaking priors with independent and identically distributed sticks and stick-breaking priors for which all the sticks take a common value. Gil-Leyva and Mena take the quantity of interest to be the posterior estimate of the density of the data generating distribution evaluated at a point—a quantity which we can call $\phi(\zeta)$ —and ask how $\mathbb{E}_{\mathcal{P}(\zeta|X,\rho)}[\phi(\zeta)]$ depends on ρ . Gil-Leyva and Mena run an MCMC chain, but then, in order to compute local robustness measures, attempt to construct a VB approximation to this posterior, observing that one would need either

 $^{^{3}}$ We should note that a first-order approximation is inadequate when taking some form of KL divergence from the original posterior as the quantity of interest, as is done in much of the literature cited by MacEachern and Lee (e.g., Johnson and Geisser, 1983; McCulloch, 1989; Carlin and Polson, 1991; Thomas et al., 2018). This KL divergence is minimized at $w = (1, \ldots, 1)$, so the first derivative with respect to the weights is zero, and one must form a local approximation using a second-order derivative. However, all our comments in the present rejoinder, particularly section 3, apply to local second-order approximations as well as to first-order approximations.

to make a (potentially limiting) mean field assumption on the sticks or deal with a computationally intractable normalizing constant.

We will avoid the question of how to construct a VB approximation in their setting, and derive instead a local sensitivity measure that can be used with an MCMC chain as long as the stick-breaking distribution can be effectively truncated at K sticks for some finite K. Let the truncated stick lengths be denoted by v_1, \ldots, v_K . We can imagine several ways to truncate an ESB model, but for the present purposes, one would need to be able to sample from the truncated model, and the prior $\mathcal{P}(v_1, \ldots, v_K | \rho)$ would need to be tractable and smooth for any draw from the MCMC chain.⁴ By eq. (2) we then have

$$\frac{\partial \mathop{\mathbb{E}}_{\mathcal{P}(\zeta|X,\rho)} \left[\phi(\zeta)\right]}{\partial \rho} \bigg|_{\rho=\rho_0} = \mathop{\mathrm{Cov}}_{\mathcal{P}(\zeta|X,\rho_0)} \left(\phi(\zeta), \left.\frac{\partial \log \mathcal{P}(v_1,\dots,v_K|\rho)}{\partial \rho}\right|_{\rho=\rho_0}\right).$$
(4)

The preceding sampling covariance can in principle be approximated from MCMC samples, with no need to form a VB approximation.

Carefully considering the implications of truncating ESB models is beyond the scope of this rejoinder, but it is worth noting the challenges for local robustness if one does not truncate, especially since Gil-Leyva and Mena (2021) use a slice sampler and do not truncate the stick-breaking distribution. If the log prior contained an infinite number of terms, it is not obvious that one could apply the dominated convergence to derive eq. (2). There exist sampling schemes that in fact sample only a finite number of sticks without truncation; see, e.g., Gil-Leyva and Mena (2021) and Walker (2007). Similarly, one might hope that one could apply eq. (2) without truncation by conditioning on auxiliary random variables in the log $\mathcal{P}(\zeta|\omega)$ term in eq. (2). But this term cannot be conditional on quantities that are random in $\mathcal{P}(\zeta|\omega)$. Additionally, the truncation will, in general, have an effect; here, the quantity of interest (the posterior estimate of the data density at a point) has nonzero correlation with all sticks. Although the posterior density at a point plausibly has diminishing correlation with sticks that come later in the process, one could design adversarial quantities of interest—e.g., the value of the 10,000-th stick—for which truncation would be quite inaccurate. Developing tractable sensitivity measures for infinite-dimensional posteriors is an interesting problem, though we suspect that eq. (4) will still be informative using straightforward finite truncation, especially given that any error in the linear approximation may well be larger than the error induced by truncation.

Empirical Bayes. Rebaudo, Fasano, Franzolini, and Müller ask whether Empirical Bayes (EB) methods for setting a DP prior might be more robust. One might answer such a question empirically by forming local robustness measures for EB procedures, which we will now undertake.

⁴Note that for the un-truncated Dirichlet-driven ESB model, the density of any finite number of sticks is tractable and smooth as a function of ρ : see Appendix E, Section 5 of the supplementary material to Gil-Leyva and Mena (2021) where the needed density is derived as part of a Gibbs sampler for ρ .

Giordano, Liu, Jordan, and Broderick

Concretely, McAuliffe et al. (2006) rely on an EB procedure to choose the DP concentration parameter. Their EB procedure takes the following general form. Fix some hyperparameter ω , which might be a case weight (see above), a perturbation of the base measure, etc. The EB procedure then finds a prior parameter $\hat{\alpha}$ that satisfies, for some F and G,

$$G(\hat{\alpha}, m(\hat{\alpha}, \omega)) = 0 \quad \text{where} \quad m(\alpha, \omega) := \mathop{\mathbb{E}}_{\mathcal{P}(\zeta | X, \alpha, \omega)} [F(\zeta)].$$
(5)

Specifically, to set the concentration parameter of a DP prior using EB, McAuliffe et al. (2006) take α to be the DP concentration parameter, $F(\zeta)$ to denote the number of clusters observed for the dataset X of size N, and $G(\alpha, m) = \sum_{n=1}^{N} \frac{\alpha}{\alpha+n-1} - m$ (see their eq. 8).

EB procedures such as this one take the form of an estimating equation that depends on a posterior moment, which can be differentiated by eqs. (1) and (2) and the chain rule. Note that $\hat{\alpha}$ depends on ω , which we write as $\hat{\alpha}(\omega)$. Additionally, write $\hat{\alpha}_0 := \hat{\alpha}(\omega_0)$ and $m_0 := m(\hat{\alpha}_0, \omega_0)$. To compute a local robustness measure for the posterior expectation of $\phi(\zeta)$, we must compute

$$\frac{\partial \mathop{\mathbb{E}}_{\mathcal{P}(\zeta|X,\hat{\alpha}(\omega),\omega)}[\phi(\zeta)]}{\partial \omega}\bigg|_{\omega_{0}},$$

accounting for the ω dependence in both the empirical Bayes procedure and the final posterior expectation. This derivative can be readily computed by applying the chain rule and eqs. (1) and (2). The result, given below in eq. (6), is a bit tedious, but its computation is entirely mechanical (and automatable) and applies to any EB procedure of the form in eq. (5).⁵

$$\frac{\partial \sum_{\mathcal{P}(\zeta|X,\hat{\alpha}(\omega),\omega)} [\phi(\zeta)]}{\partial \omega} \bigg|_{\omega_{0}} = \frac{\operatorname{Cov}}{\mathcal{P}(\zeta|X,\hat{\alpha}_{0},\omega_{0})} \left(\phi(\zeta), \frac{\partial \log \mathcal{P}(\zeta,X|\hat{\alpha}_{0},\omega)}{\partial \omega} \bigg|_{\omega_{0}} + \frac{\partial \log \mathcal{P}(\zeta,X|\alpha,\omega_{0})}{\partial \alpha} \bigg|_{\hat{\alpha}_{0}} \frac{d\hat{\alpha}(\omega)}{d\omega} \bigg|_{\omega_{0}} \right)^{1} \left(\frac{\partial G(\hat{\alpha}_{0},m)}{\partial \alpha} \bigg|_{\omega_{0}} + \frac{\partial G(\hat{\alpha}(m))}{\partial \omega} \bigg|_{\omega_{0}} \right)^{1} \left(\frac{\partial G(\hat{\alpha}_{0},m)}{\partial m} \bigg|_{m_{0}} \frac{\partial m(\hat{\alpha}_{0},\omega)}{\partial \omega} \bigg|_{\omega_{0}} \right),$$

$$\frac{\partial G(\alpha,m(\alpha,\omega_{0}))}{\partial \alpha} \bigg|_{\hat{\alpha}_{0}} = \frac{\partial G(\alpha,m_{0})}{\partial \alpha} \bigg|_{\hat{\alpha}_{0}} + \frac{\partial G(\hat{\alpha}_{0},m)}{\partial m} \bigg|_{m_{0}} \frac{\partial m(\alpha,\omega_{0})}{\partial \alpha} \bigg|_{\hat{\alpha}_{0}},$$

$$\frac{\partial m(\alpha,\omega_{0})}{\partial \alpha} \bigg|_{\hat{\alpha}_{0}} = \frac{\operatorname{Cov}}{\mathcal{P}(\zeta|X,\hat{\alpha}_{0},\omega_{0})} \left(F(\zeta), \frac{\partial \log \mathcal{P}(\zeta,X|\alpha,\omega_{0})}{\partial \omega} \bigg|_{\hat{\alpha}_{0}} \right),$$
and
$$\frac{\partial m(\alpha_{0},\omega)}{\partial \omega} \bigg|_{\omega_{0}} = \operatorname{Cov}_{\mathcal{P}(\zeta|X,\hat{\alpha}_{0},\omega_{0})} \left(F(\zeta), \frac{\partial \log \mathcal{P}(\zeta,X|\hat{\alpha}_{0},\omega)}{\partial \omega} \bigg|_{\omega_{0}} \right).$$
(6)

⁵For compactness, we have suppressed some evaluation notation in this display; for example, we write $\hat{\alpha}_0$ in place of $\alpha = \hat{\alpha}_0$. The evaluation is always done for the parameter with respect to which we are differentiating.

Evaluating Sensitivity to the Stick-Breaking Prior in BNP

One might ask whether eq. (6) can be applied to the EB procedure used by McAuliffe et al. (2006) for the base measure. Unfortunately, since McAuliffe et al. (2006) estimate the base measure nonparametrically, the space of possible base measures is infinite-dimensional, and so one cannot apply eq. (1) directly. Versions of eq. (1) for infinite-dimensional parameters exist (see, e.g., Chapter 4 of Zeidler, 1986), though applying them in practice seems to be challenging and beyond the scope of this short rejoinder. Alternatively, one could represent the base measure using a large but finite basis and apply eq. (6).

Loss functions. MacEachern and Lee ask whether we can compute sensitivity to the loss function in a Bayesian analysis. Formally, for a posterior $\mathcal{P}(\zeta|X)$ and loss function L, under common regularity conditions we have

$$\hat{\theta}_{\text{loss}} := \underset{\theta}{\operatorname{argmin}} \underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} \left[L(\zeta, \theta) \right] \quad \Leftrightarrow \underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} \left[\left. \frac{\partial L(\zeta, \theta)}{\partial \theta} \right|_{\theta = \hat{\theta}_{\text{loss}}} \right] = 0.$$
(7)

We will consider the common situation described in eq. (7). However, we note that, by exchanging the order of local robustness derivatives and posterior expectations, this approach could be naturally extended to estimators of the form given in Lee and MacEachern (2014), i.e., $\underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} \left[\operatorname{argmin}_{\theta} \int L(y,\theta)\zeta(dy) \right]$ for a distribution-valued ζ .

Equation (7) defines an estimating equation for $\hat{\theta}_{loss}$. We can parameterize a path to a different loss function using $L(\zeta, \theta, \omega) = L(\zeta, \theta) + \omega \Delta(\zeta, \theta)$ for some $\Delta(\zeta, \theta)$. Then, apply eq. (1) to eq. (7), and interchange differentiation and integration to get

$$\frac{d\hat{\theta}_{\text{loss}}(\omega)}{d\omega}\bigg|_{\omega=0} = -\left(\mathbb{E}_{\mathcal{P}(\zeta|X)}\left[\frac{\partial^2 L(\zeta,\theta)}{\partial\theta\partial\theta}\bigg|_{\theta=\hat{\theta}_{\text{loss}}}\right]\right)^{-1}\mathbb{E}_{\mathcal{P}(\zeta|X)}\left[\frac{\partial\Delta(\zeta,\theta)}{\partial\theta}\bigg|_{\theta=\hat{\theta}_{\text{loss}}}\right].$$
 (8)

For example, to estimate the effect of replacing the mean with the median, we could take $L(\zeta, \theta) = \frac{1}{2}(\zeta - \theta)^2$, $\Delta = |\zeta - \theta| - L(\zeta, \theta)$, and

$$\operatorname{Median}(\mathcal{P}(\zeta|X)) - \underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} [\zeta] = \hat{\theta}_{\operatorname{loss}}(1) - \hat{\theta}_{\operatorname{loss}}(0) \approx \left. \frac{d\hat{\theta}_{\operatorname{loss}}(\omega)}{d\omega} \right|_{\omega=0} (1-0)$$
$$= \underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} \left[\mathbb{I} \left(\zeta > \underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} [\zeta] \right) \right] - \underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} \left[\mathbb{I} \left(\zeta < \underset{\mathcal{P}(\zeta|X)}{\mathbb{E}} [\zeta] \right) \right]. \tag{9}$$

For example, this approximation reasonably asserts that the median will exceed the mean when the posterior is asymmetric, with a greater mass to the right of the mean than to the left. (But we will discuss some of its limitations in section 3 below.)

Since Δ could be any function satisfying basic regularity conditions, one could in principle use eq. (8) to explore the space of loss functions—if one can believe that the approximation provided by $\Delta \mapsto \frac{d\hat{\theta}_{loss}(\omega)}{d\omega}\Big|_{\omega=0}$ is a good one uniformly over the candidate set of perturbations Δ . However, again, we do not necessarily recommend this for this particular path through the space of loss functions. On the contrary, we will use this example in section 3 below as an example of a derivative that may not serve its intended purpose very well.

3 What makes a derivative useful?

In section 2.1 above, we derived local robustness measures for several settings requested by our respondents, and we expect that readers can readily derive most of the rest for themselves. Have we solved all their problems? Certainly not! To the contrary, we will argue that the computation of derivatives is straightforward, but showing their utility is harder.

In our view, a useful derivative should (at least) satisfy a few related "usefulness desiderata": (1) be readily computable to the desired accuracy, (2) be easily interpretable, and, most importantly, (3) extrapolate well so as to provide a reasonable approximation to the "global robustness" problem. We have endeavored to show that certain derivatives are at least plausibly useful, according to these criteria, for DP priors in VB approximations. In addition to considering Fréchet differentiability—which is, arguably, a rather low bar for a derivative to pass—we primarily demonstrated our local robustness measure's ability to extrapolate through careful experiments and comparison with refitting. In certain situations, such as case influence in large datasets, one can sometimes prove good extrapolation by bounding the second derivative under readily interpretable conditions (as in Giordano et al., 2019b).

Evaluating the usefulness desiderata for the derivatives given in section 2.1 above is worth doing, and it is the work which constitutes most of the effort. We do not believe that all the results of section 2.1 will pass the test. Let us focus on the loss function example, though many of these potential problems apply to the other settings as well.

Computability. The expected Hessian inside the inverse in eq. (8) will have Monte Carlo error if estimated with MCMC, which will bias the inverse. Furthermore, if the difference between L and $L + \Delta$ occurs mostly in the tails, the expectation of the derivative of Δ may suffer from high MCMC noise.

Interpretability. The loss function derivative in eq. (8) will behave pathologically as an approximation to losses that are pointwise close to the original loss function, but have very large derivatives near $\hat{\theta}_{loss}$. Without a judiciously constrained search space to exclude such alternative loss functions, eq. (8) will provide poor guidance when exploring the space of alternative loss functions. Unfortunately, a more complex search space comes at a cost, which is the computational difficulty of optimizing a linear form (i.e., the derivative in eq. (8), viewed as a functional of Δ) over this space.

Extrapolation. The example of the mean and median shows that the derivative eq. (8) may not always extrapolate well in common use cases. Though we know that the mean and median can be arbitrarily different in general, the approximation to the difference in eq. (9) cannot be larger in magnitude than one.

Attempting to investigate and repair these deficiencies—e.g., by improved MCMC sampling, alternative paths through the space of loss functions, and the selection of search sets in the space of loss functions—is an interesting and valuable project, and one that may require considerably more effort than derivation of the local robustness approximation.

The usefulness desiderata sufficed for our objective, which was primarily to provide

Evaluating Sensitivity to the Stick-Breaking Prior in BNP

a tool for quickly exploring the space of stick-breaking priors in a way that is not too computationally or technically burdensome, for a particular quantity of interest. On the other hand, we do not attempt to detect sensitivity of the entire model (e.g., with a whole-distribution divergence measure), we do not assert that a large worst-case derivative implied non-robustness (e.g., the corresponding prior may have looked subjectively unreasonable), and we do not assert that our good results on extrapolation will necessarily hold in very different settings (we primarily showed good extrapolation via experiment). In this sense, our objectives are somewhat different than much of the classical robustness literature, which we see as attempting to provide more universal notions of "robustness." For example, the foundational works of Ruggeri and Wasserman (1993), Basu et al. (1996), Gustafson (1996), to which we are quite indebted, appear to take as their task the *definition* of a single number which can be interrogated, relatively free of context, to ascertain whether a posterior is "robust" or not. This goal is reflected in how their techniques are used, for example, in Basu (2000). A similar goal of finding universal metrics of "data importance" motivates much of the case deletion literature; it is perhaps for this reason that many authors focus on various forms of whole-model KL divergence or likelihood ratios (see, e.g., Johnson and Geisser, 1983; Cook, 1986; Carlin and Polson, 1991). The production of a universally valid local robustness metric requires even stricter conditions on the derivative than our usefulness desiderata; e.g., it must extrapolate well in all directions, its worst-case perturbation must lead to a subjectively reasonable prior, the researcher must actually care about whole-model sensitivity and not just a particular posterior quantity, and so on.

Our context-specific approach and a more universalist approach need not be at odds. On the contrary, the intuition and best practices arising from routine and systematic assessment of prior assumptions might lead, in the end, to better and more universally applicable metrics of robustness. Similarly, asymptotic analysis of the sort advocated by **Ascolani, Catalano, and Prünster** can inform and be informed by the robustness of particular finite-data settings.

Where local approximations fail to satisfy the usefulness desiderata, there is ample room for creativity. The analysis of **Griffin and Kalli**, which both clearly demonstrates a failure of a linear approximation to extrapolate and suggests a solution, seems exemplary to us. Their idea of performing sensitivity analysis separately for a number of local modes seems promising; to their suggestion we might add forming a (second order) approximation to the value of the ELBO at these modes as well, in order to assess when the relative ordering of the modes changes. Local approximations might also augment other schemes, such as suggesting quadratic transforms for the importance sampling techniques of MacEachern and Peruggia (2000).

We hope that researchers will feel empowered by this work to creatively explore the space of model perturbations, relatively unencumbered by the difficulty of deriving local robustness measures, but attentive to their ability to answer useful questions in their own modeling contexts.

10

References

- Averbukh, V. and Smolyanov, O. (1967). "The theory of differentiation in linear topological spaces." Russian Mathematical Surveys, 22(6): 201–258. 3
- Bae, J., Ng, N., Lo, A., Ghassemi, M., and Grosse, R. (2022). "If Influence Functions are the Answer, Then What is the Question?" In Advances in Neural Information Processing Systems. 3
- Basu, S. (2000). Bayesian Robustness and Bayesian Nonparametrics, 223–240. New York, NY: Springer New York. 10
- Basu, S., Jammalamadaka, S. R., and Liu, W. (1996). "Local posterior robustness with parametric priors: Maximum and average sensitivity." In *Maximum Entropy and Bayesian Methods*, 97–106. Springer. 10
- Billingsley, P. (2008). Probability and Measure. John Wiley & Sons. 3
- Carlin, B. and Polson, N. (1991). "An expected utility approach to influence diagnostics." Journal of the American Statistical Association, 86(416): 1013–1021. 5, 10
- Cook, D. (1977). "Detection of influential observation in linear regression." *Technometrics*, 19(1): 15–18. 4
- Cook, R. (1986). "Assessment of local influence." Journal of the Royal Statistical Society: Series B (Methodological), 48(2): 133–155. 10
- Diaconis, P. and Freedman, D. (1986). "On the consistency of Bayes estimates." The Annals of Statistics, 1–26. 3
- Fleming, W. (2012). Functions of Several Variables. Springer Science & Business Media.
 3
- Gil-Leyva, M. and Mena, R. (2021). "Stick-breaking processes with exchangeable length variables." Journal of the American Statistical Association, 1–14. 4, 5, 6
- Giordano, R., Jordan, M. I., and Broderick, T. (2019a). "A higher-order swiss army infinitesimal jackknife." arXiv preprint arXiv:1907.12116. 3
- Giordano, R., Stephenson, W., Liu, R., Jordan, M. I., and Broderick, T. (2019b). "A Swiss army infinitesimal jackknife." In *The 22nd International Conference on Artifi*cial Intelligence and Statistics, 1139–1147. PMLR. 4, 9
- Gustafson, P. (1996). "Local sensitivity of posterior expectations." The Annals of Statistics, 24(1): 174–195. 3, 10
- Hampel, F. (1974). "The influence curve and its role in robust estimation." Journal of the American Statistical Association, 69(346): 383–393. 3, 4
- Hampel, F., Ronchetti, E., Rousseeuw, P., and Stahel, W. (1986). Robust Statistics: The Approach Based on Influence Functions. Wiley-Interscience; New York. 5
- Hattori, S. and Kato, M. (2009). "Approximate subject-deletion influence diagnostics for Inverse Probability of Censoring Weighted (IPCW) method." Statistics and Probability Letters, 79(17): 1833–1838. 3

- Huber, P. J. (1964). "Robust estimation of a location parameter." The Annals of Mathematical Statistics, 35(1): 73-101. URL http://www.jstor.org/stable/2238020 4
- Jacobi, L., Joshi, M., and Zhu, D. (2018). "Automated sensitivity analysis for Bayesian inference via Markov chain Monte Carlo: Applications to Gibbs sampling." Available at SSRN 2984054. 3
- Johnson, W. and Geisser, S. (1983). "A predictive view of the detection and characterization of influential observations in regression analysis." Journal of the American Statistical Association, 78(381): 137–144. 5, 10
- Koh, P. and Liang, P. (2017). "Understanding black-box predictions via influence functions." In International Conference on Machine Learning (ICML). 4
- Krantz, S. and Parks, H. (2012). The Implicit Function Theorem: History, Theory, and Applications. Springer Science & Business Media. 3
- Lee, J., James, L., and Choi, S. (2016). "Finite-dimensional BFRY priors and variational Bayesian inference for power law models." Advances in Neural Information Processing Systems. 4
- Lee, J. and MacEachern, S. (2014). "Inference functions in high dimensional Bayesian inference." Statistics and its Interface, 7(4): 477–486.
- MacEachern, S. and Peruggia, M. (2000). "Importance link function estimation for Markov chain Monte Carlo methods." Journal of Computational and Graphical Statistics, 9(1): 99–121. 10
- Maclaurin, D. (2016). "Modeling, Inference and Optimization With Composable Differentiable Procedures." 3
- McAuliffe, J., Blei, D., and Jordan, M. I. (2006). "Nonparametric empirical Bayes for the Dirichlet process mixture model." *Statistics and Computing*, 16: 5–14. 4, 7, 8
- McCulloch, R. (1989). "Local model influence." Journal of the American Statistical Association, 84(406): 473–478. 5
- McInerney, A. (2013). First Steps in Differential Geometry. Springer. 3
- Mohamed, S., Rosca, M., Figurnov, M., and Mnih, A. (2020). "Monte Carlo gradient estimation in machine learning." Journal of Machine Learning Research, 21(132): 1–62. 3
- Murray, M. and Rice, J. (1993). Differential geometry and statistics, volume 48. CRC Press. 3
- Ruggeri, F. and Wasserman, L. (1993). "Infinitesimal sensitivity of posterior distributions." Canadian Journal of Statistics, 21(2): 195–203. 3, 10
- Serfling, R. (1980). Approximation Theorems of Mathematical Statistics, volume 162. John Wiley & Sons. 4

- Shao, J. and Tu, D. (2012). The Jackknife and Bootstrap. Springer Science & Business Media. 4
- Shi, L., Lu, J., Zhao, J., and Chen, G. (2016). "Case deletion diagnostics for GMM estimation." Computational Statistics & Data Analysis, 95: 176–191. 3
- Stigler, S. (2010). "The changing history of robustness." The American Statistician, 64(4): 277–281. 4
- Thomas, W. and Cook, D. (1989). "Assessing influence on regression coefficients in generalized linear models." *Biometrika*, 76(4): 741–749. 3
- Thomas, Z., MacEachern, S., and Peruggia, M. (2018). "Reconciling curvature and importance sampling based procedures for summarizing case influence in Bayesian models." *Journal of the American Statistical Association*, 113(524): 1669–1683. 5
- van der Vaart, A. and Wellner, J. (1996). Empirical Processes and Weak Convergence. Springer, New York. 4
- von Mises, R. (1947). "On the asymptotic distribution of differentiable statistical functions." The Annals of Mathematical Statistics, 18(3): 309–348. 4
- Walker, S. (2007). "Sampling the Dirichlet mixture model with slices." Communications in Statistics Simulation and Computation, 36(1): 45–54. 6
- Zeidler, E. (1986). Nonlinear Functional Analysis and its Applications I: Fixed-point Theorems. Springer-Verlag. 3, 8