Topological densities in Einstein-scalar-Gauss-Bonnet gravity

M. Bousder¹*and Z Sakhi²

¹Faculty of Science, Mohammed V University in Rabat, Rabat, Morocco

²Quantum Physics and Applications Team, LPMC, Faculty of Science Ben M'sik,

Casablanca Hassan II University, Morocco

April 4, 2023

Abstract

The present work is devoted to studying the background dynamical evolution of a scalar field in Einstein-Gauss-Bonnet gravity in maximally symmetric spacetime. This study is useful for giving meaning to the presence of two Gauss-Bonnet vacua, instead of using the spherically symmetric bubbles of the "true" vacuum expand in the "false" vacuum. The theory admits two possible effective cosmological constants, which lead to two maximally symmetric vacuum solutions. The first solution corresponds to the dynamics of dark energy. When there is matter, the second solution describes dark matter. In Einstein-Gauss-Bonnet gravity, we establish the expression of the topological mass spectrum which depends on the golden ratio and its inverse. In the Schwarzschild limit, the topological density corresponds to the standard model radiation energy density. We find the mass loss rate which gives the evolution of mass over time.

1 Introduction

The observations of the rotation of galaxies and gravitational lenses indicate the presence of a new mass, called dark matter (DM) hiding in galaxies, which does not interact with radiation and matter, but can be detected by its gravitational effect. The Λ CDM model is a cosmological model, parametrized by a cosmological constant Λ associated with cold dark matter. It is often called the standard Big Bang model because it is the

^{*}mostafa.bousder@gmail.com

simplest model that accounts for the properties of the cosmos: the large-scale structure of the observable universe and the distribution of galaxies, the abundance of light elements (hydrogen, helium and lithium) and the expansion of the universe. ACDM model assumes that general relativity theory correctly describes gravity on a cosmological scale. However, the ΛCDM model presents several problems, such as the cosmological constant, fine-tuning problem, and the problem of cosmic coincidence [1]. Recent developments of non-trivial extension of Lovelock theory, namely Einstein-Gauss-Bonnet (EGB) theory have been proposed [2], providing new insights into the 4-dimensional theory of gravity [3]. Their idea is that before taking the limit $D \rightarrow 4$, they multiplied the Gauss-Bonnet (GB) term by the factor 1/(D-4). The divergent factor 1/(D-4) is canceled by the vanishing GB contributions, which leads to a theory of gravity with only two dynamical degrees of freedom, which is in contradiction with Lovelock theorem [4] which describes the gravity at $D \geq 5$. However, it was shown in several papers that the idea of the limit $D \longrightarrow 4$ is not clearly defined, as well as the absence of proper action [5, 6, 7, 8]. It was explicitly confirmed by a direct product D-dimensional spacetime or by adding a counter term, before taking the limit $D \longrightarrow 4$, which can be seen as a class of Horndeski theory [9] but with 2 + 1-dofs. Although the EGB gravity is currently debatable, the spherically symmetric black hole solution is still meaningful and worthy of study [10]. The 4D symmetrical static and spherical black hole solution in EGB gravity were obtained [11], also, solutions of static and spherically symmetric compact stars [12, 13, 14]. Many researchers have studied the mass-radius profile and the maximum mass in EGB gravity [15, 16]. Consequently, it is possible to describe the matter inside compact objects (COs) and the dynamical evolution of the matter at high density and the behavior of violent events. The Einstein-scalar-Gauss-Bonnet (EsGB) gravity [17], are the class of classical scalar-tensor theories that have second-order EOMs, as a special case the Horndeski gravity [18]. The EGB gravity admits two maximally symmetric vacuum solutions as the Einstein vacuum in $\alpha \to 0$, and the Gauss-Bonnet vacuum in $\alpha \neq 0$ [19, 20]. Previous studies [21] including the effect of varying the cosmological constant, showed that the correspondence between ordinary thermodynamic systems and black hole mechanics would be completed to include a variable cosmological constant. The bubble nucleation probability depends on the curvature coupling of the Higgs fields, which is a renormalizable parameter of the Standard Model (SM) in curved spacetime [22]. In the effective field theory, the thermal and quantum fluctuations to overcome the barrier are characterized by the decrease of the vacuum. The bubble nucleation in thermal fluctuations can be described in terms of Euclidean time coordinate by instantons [23]. The action of the Coleman-de Luccia instanton determines the rate of vacuum decay [24]. The classic solutions for switching from a false vacuum to a true vacuum are called instantons. We suppose that the Boltzmann constant k_B , the reduced Planck constant \hbar and light speed c, are such that: $k_B = \hbar = c = 1$.

We begin in section 2 with a discussion of the equation of motion for a coupling between a scalar field and the model, the chameleon mechanism and the scalaron mass are investigated in section 3. Section 4 is devoted to the two branches of solutions for the effective cosmological constant in a maximally symmetric vacuum. In Section 5, we study the scalar dark matter in the exterior region of CO by the EsGB and the regions of validity of the GB functional coupling. In Section 6, we explore the applications for the 4D Einstein-Gauss-Bonnet black hole. Finally, the paper ends in Section 7 by summarizing our main results.

2 Einstein-scalar-Gauss-Bonnet gravity

We start by the action of Einstein-scalar-Gauss-Bonnet (EsGB) gravity [25, 26] in 4dimensions as

$$S = \int d^4x \sqrt{-g} \left(\frac{M_p^2}{2} R + f(\phi) \mathcal{G} + \mathcal{L}_{DM} \right) + S_m, \qquad (2.1)$$

where S_m is the matter action, $M_p = \frac{c^4}{8\pi G} \approx 2 \times 10^{18} [GeV]$, R is the Ricci scalar, $f(\phi)$ is a Gauss-Bonnet coupling function, which is ultraviolet (UV) corrections to Einstein theory. We define the Gauss-Bonnet invariant [27] as

$$\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}.$$
(2.2)

The variation with respect to the field ϕ gives us the equation of motion for the scalaron field

$$\Box \phi - \partial_{\phi} V(\phi) + \mathcal{G} \partial_{\phi} f(\phi) = 0.$$
(2.3)

The variation of the action over the metric $g_{\mu\nu}$ simplified by the Bianchi identity gives the equations of motion in [25]. In the Jordan frame, the scalar DM Lagrangian reads [28]

$$\mathcal{L}_{DM} = -\frac{1}{2}g^{\mu\nu}\partial_{\mu}\phi\partial_{\nu}\phi - V(\phi)$$
(2.4)

the kinetic term is invariant under the transformation $\phi \to -\phi$ of the Z_2 symmetry. The metric of a spatially flat homogeneous and isotropic universe in FLRW model is given by:

$$ds^{2} = -dt^{2} + a^{2}(t) \sum_{i=1}^{3} (dx^{i})^{2}, \qquad (2.5)$$

where a(t) is a dimensionless scale factor, from which we define the Ricci scalar R and the GB invariant \mathcal{G} in FLRW geometry as

$$R = 6\left(2H^2 + \dot{H}\right) \qquad \mathcal{G} = 24H^2\left(\dot{H} + H^2\right).$$
(2.6)

We start by considering $\phi = \phi(t)$. where $\dot{\phi} = \partial_t \phi$, $f'(\phi) = \partial_{\phi} f(\phi)$ and $H = \dot{a}/a$ is the Hubble parameter. Eq.(2.3) can be written as the Klein Gordon equation in a simple form

$$\ddot{\phi} + 3H\left[\dot{\phi} - 8f'(\phi)H\left(\dot{H} + H^2\right)\right] + V'(\phi) = 0.$$
(2.7)

3 Scalaron mass in EsGB gravity

Recently, there has been a renewed interest in the relationship between dark matter and the scalaron mass (i.e. the mass of fields ϕ) [31]. We will study later the scalaron mass m_{ϕ} which will describe the dark matter. To describe the mass of any scalar field, we use the Klein-Gordon equation $\Box \phi = \partial_{\phi} V_{eff}(\phi)$. Note that Eq.(2.3) gives the same form as in the last equation. Since \mathcal{G} in Eq.(2.3), does not depend on ϕ , in this case, the effective potential is evaluated as follows

$$V_{eff}(\phi) = V(\phi) - f(\phi)\mathcal{G}.$$
(3.1)

We notice that the effective potential of the scalaron includes the Gauss-Bonnet coupling and the Gauss-Bonnet invariant. In other words, the Gauss-Bonnet term affects the potential structure of the scalaron; thus, the scalaron mass depends on the matter's contribution. The particles of the field ϕ come from the fluctuation around the minimum of the effective potential $V_{eff}(\phi)$. The mass of small fluctuations around ϕ_{\min} Eq.(??) give the scalaron mass [55] is determined as

$$m_{\phi}^2 = \frac{\partial^2}{\partial \phi^2} V_{eff}(\phi_{\min}), \qquad (3.2)$$

where $V_{eff}(\phi_{\min})$ as a minimum value of the scalaron effective potential V_{eff} . Also we have $\mathcal{G} = \frac{\partial_{\phi}^2 V(\phi)}{\partial_{\phi}^2 f(\phi)} - \frac{\partial_{\phi}^2 V_{eff}(\phi)}{\partial_{\phi}^2 f(\phi)}$. To make progress, let us express the \mathcal{G} in a more convenient form

$$\mathcal{G} = \frac{\partial_{\phi}^2 V\left(\phi_{\min}\right)}{\partial_{\phi}^2 f\left(\phi_{\min}\right)} - \frac{m_{\phi}^2}{\partial_{\phi}^2 f\left(\phi_{\min}\right)}.$$
(3.3)

For $f(\phi) = f_0 e^{k\phi}$ and $V(\phi) = V_0 e^{-k\phi} [25, 3]$ we obtain

$$f(\phi_{\min})\mathcal{G} = V(\phi_{\min}) - \frac{m_{\phi}^2}{k^2}, \qquad (3.4)$$

From Eqs. (3.1, 3.3), we get

$$V_{eff}(\phi_{\min}) = \frac{m_{\phi}^2}{k^2}.$$
 (3.5)

where $\partial_{\phi}^2 = \partial^2/\partial \phi^2$ and $V_{eff,\min}$ as a minimum value of the scalaron effective potential V_{eff} . The minimum of the potential at $\phi = \phi_{\min}$ should satisfy $\partial_{\phi}V_{eff}(\phi_{\min}) = 0$. It is shown that scalar fields can explain the abundance of dark matter. The scalaron mass change according to the trace of the energy-momentum tensor [33]. The minimum of the potential at $\phi = \phi_{\min}$ should satisfy $\partial_{\phi}V_{eff}(\phi_{\min}) = 0$, which give a special value of the GB invariant as

$$\mathcal{G} = -\frac{V_0}{f_0} e^{-4R_0\phi_{\min}}$$
(3.6)

also we have

$$V_{eff}(\phi) = V(\phi) \left[1 + e^{4R_0(\phi - \phi_{\min})} \right]$$
(3.7)

If $\phi_{\min} = 0$, we get $V_{eff}(\phi) = 2V_0 \cosh(2R_0\phi)$. Substituting Eqs.(??,3.6) into Eq.(3.1) then into Eq.(3.2), the scalaron mass can then be expressed as

$$m_{\phi} = 2\sqrt{2V_0}R_0 e^{-R_0\phi_{\min}}.$$
(3.8)

It is shown that scalar fields can explain the abundance of dark matter. The scalaron mass change according to the trace of the energy-momentum tensor [33]. In the large curvature limit [34] we have $R_0\phi_{\min} = 1$, one finds

$$R \longrightarrow R_0 = 1/\phi_{\min} = M_p^{-2}\rho, \qquad (3.9)$$

where ρ is the matter matter-energy. We can then express the scalaron mass m_{ϕ} as a function of the matter-energy density

$$m_{\phi} = \frac{2\sqrt{2V_0}}{eM_p^2}\rho.$$
 (3.10)

The scalaron field ϕ becomes dynamical in the low energy density environment on the cosmological scale. Since the mass m_{ϕ} depends on the matter density ρ , thus, the scalaron becomes heavy in the high-density region of matter. This feature is called the chameleon mechanism which is one of the screening mechanism in the modified gravity [35]. The chameleon mechanisms is defined when the scalaron mass depends on the environment surrounding the scalaron field. The scalaron ϕ is regarded as a dynamical dark matter and can be a dark matter candidate. Similar topics had been researched in many literatures [35, 36].

4 Maximally symmetric vacuum solutions

Recently, there are several papers which study the Particle production induced by vacuum decay [45]. Following these concepts, here, we will interpret the Gauss-Bonnet vacua by the production of dark matter particles. When gravity is taken into consideration, the vacua are those with maximally symmetric spaces [22]. In the maximally symmetric space, the scalar curvature of de Sitter space is given by

$$R = \frac{2D}{D-2}\Lambda,\tag{4.1}$$

where Λ is the cosmological constant. In the case of the positive Λ , we have the de Sitter solution. In the large limit, we obtain $1/\phi_{\min} = \frac{2D}{D-2}\Lambda_0$. In 4-dimensional vacuum, the equation of motion has a solution that $R_{\mu\nu} = \Lambda g_{\mu\nu}$, implies that $R = 4\Lambda$. The Minkowski spacetime as the vacuum of the $\Lambda_0 = 0$. The singularity problem [37] corresponds to $R_0 \to \infty$ (the curvature singularity) or $\phi_{\min} = 0$. From Eq.(3.8), Eq.(3.9) and Eq.(4.1), for $e^{-2R_0\phi_{\min}} \approx 1 - 2R_0\phi_{\min}$ and $R_0 \longrightarrow R$ one obtain

$$2\phi_{\min}R^2 - R + \frac{m_{\phi}}{2\sqrt{2V_0}} = 0, \qquad (4.2)$$

which can reduce to

$$4\phi_{\min}\left(\frac{D}{D-2}\right)^{2}\Lambda^{2} - \frac{D}{D-2}\Lambda + \frac{m_{\phi}}{4\sqrt{2V_{0}}} = 0.$$
 (4.3)

In maximally symmetric vacuum solutions, there are two branches of solutions for the effective cosmological constant,

$$\Lambda_{\pm} = \frac{D-2}{8D\phi_{\min}} \left(1 \pm \sqrt{1 - \frac{4m_{\phi}}{\sqrt{2V_0}}\phi_{\min}} \right), \qquad (4.4)$$

In the limit where $2m_{\phi}\phi_{\min} \ll \sqrt{2V_0}$, the two branches are given by

$$\Lambda_{+} = \frac{D-2}{4D\phi_{\min}} \left(1 - \frac{m_{\phi}\phi_{\min}}{\sqrt{2V_0}} \right), \qquad (4.5)$$

$$\Lambda_{-} = \frac{D-2}{4D} \frac{m_{\phi}}{\sqrt{2V_0}}.$$
(4.6)

Using Eq. (3.10) and $1/\phi_{\min} = \frac{2D}{D-2}\Lambda_0$, the above equations can be further rewritten as

$$\Lambda_{+} = \frac{\Lambda_{0}}{2} \left(1 - \frac{2\phi_{\min}\rho}{eM_{p}^{2}} \right); \quad \Lambda_{-} = \Lambda_{0} \frac{\phi_{\min}\rho}{eM_{p}^{2}}. \tag{4.7}$$

We notice that $\Lambda_{+} = \frac{\Lambda_{0}}{2} - \Lambda_{-}$. In particular, the large curvature limit $1/4 = \Lambda_{0}\phi_{\min}$, we have

$$\frac{\Lambda_0}{2} = \Lambda_+ + \Lambda_-; \ \Lambda_- = \frac{\rho}{4eM_p^2}; \qquad \Lambda_0 = \frac{1}{4\phi_{\min}} = \frac{\rho}{4M_p^2},$$
(4.8)

or equivalently

$$\Lambda_{+} = \frac{(e-2)}{2e} \Lambda_{0} ; \Lambda_{-} = \frac{\Lambda_{0}}{e}; \quad \Lambda_{0} = \frac{\rho}{4M_{p}^{2}}$$

$$(4.9)$$

Let us now comment on the two solutions above. The cosmological constant Λ_+ is equivalent to that found by [26] which is proportional to the mass of the scalar field (chameleonic dark matter [33]). If ρ decreases over time, the value of Λ_+ increases to reach Λ_0 . On the other hand, the second branch Λ_- depends on the matter density. We notice that $\Lambda_+ > 0$ and $\Lambda_- > 0$, instead of studying the false vacuum forming inside the true vacuum in the bubble geometry [39]. The two roots $(\Lambda_{-}, \Lambda_{+})$ can represent two faces of the same true vacuum. In a vacuum (without matter), the chameleon mechanism will be zero, which corresponds to dark energy. The chameleon mechanism appears when there is the matter Eq.(3.10), the vacua are those with Λ_{-} . The vacuum Λ_{-} will be spontaneously produced with ordinary matter fields. The Gauss-Bonnet vacuum becomes a chameleon if there is matter, which solves the problem of the Gauss-Bonnet vacuum suffering from perturbative ghost instability [19]. This justifies that the matter has an impact on the vacuum, is that the two vacua are separated by a domain wall, composed of ordinary matter in the thin wall approximation [39]. In particular, the large curvature limit $1/\phi_{\min} = R_0 = 4\Lambda_0$ Eq.(4.9), must satisfy $\Lambda_0 = \rho/4M_p^2$ and $\Lambda_{-} = \rho/4eM_{p}^{2}$, we can show that $\Lambda_{+} + \Lambda_{-} = \Lambda_{0}$, which corresponds to $\Lambda_{-}/\Lambda_{0} \approx$ $0,368 \equiv 36,8\%$ and $\Lambda_+/\Lambda_0 \approx 0,132 \equiv 13,2\%$. The percentage $\Lambda_-/\Lambda_0 \equiv 36,8\%$ is close to the density of matter and dark matter in the universe [43]. It is interesting to note that 50% = (36, 8% + 13, 2%) of space-time gets a mass in a region occupied by $(\Lambda_{-}, \Lambda_{+})$. In this case, we introduce the parameter X which explains the lack of 50% of $\Lambda_0: \Lambda_0 = \Lambda_+ + \Lambda_- + X$, i.e. $X/\Lambda_0 = 0, 5 \equiv 50\%$. So, we can associate the parameter X to dark energy.

Energy content	parameter	pourcentage
Matter+Dark matter	Λ_{-}	$\Lambda_{-}/\Lambda_{0} \equiv 36,8\%$
Dark energy	$\Lambda_+ + X$	$\left(\Lambda_{+}+X\right)/\Lambda_{0}=63,2\%$
Universe	Λ_0	100%

Table 1: Numerical estimate of the percentages of ordinary matter, dark matter and dark energy in the universe [43].

We notice that the dark energy is described by the parameter Λ_+ and an unknown parameter X table (2).

5 Functional coupling and Barrow entropy

In this section, we explain in detail how to construct the equation of motion of the Einstein-Gauss-Bonnet gravity. We begin by reviewing the the 4D EsGB action

$$\mathcal{S} = \frac{M_p^2}{2} \int d^4x \sqrt{-g} \left(R + f(\phi) \mathcal{G} - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right) + \mathcal{S}_m$$

where $M_p = 1/(8\pi G_N) = 1.221 \times 10^{19} GeV$ is the reduced Planck mass, R is the Ricci scalar, S_m is the matter action and $f(\phi)$ is a functional coupling of the scalar field ϕ . In the above equation $(\mu, \nu) = (0, 1, 2, 3)$. We define the GB term as

$$\mathcal{G} \equiv R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}.$$
(5.1)

The Kretschmann scalar is $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$. The variation with respect to the field ϕ gives us the equation of motion for the scalar field

$$\Box \phi = \partial_{\phi} V_{eff}(\phi) \,, \tag{5.2}$$

where $\Box \equiv \nabla_{\mu} \nabla^{\mu}$ and the effective potential is

$$V_{eff}(\phi) = V(\phi) - f(\phi)\mathcal{G}.$$
(5.3)

Varying the action (??) over the metric $g_{\mu\nu}$, we obtain the following equations of motion:

$$G^{\mu\nu} + \mathcal{K}^{\mu\nu} + f\left(\phi\right)\mathcal{H}^{\mu\nu} + \frac{1}{2}\left[\mathcal{T}^{\mu\nu}_{\phi} - g^{\mu\nu}V_{eff}\left(\phi\right)\right] = \frac{1}{2}\kappa^{2}T^{\mu\nu},$$

where the Einstein tensor is $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2}g^{\mu\nu}R$, the matter stress tensor is $T^{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta S_m}{\delta g_{\mu\nu}}$. On the other hand the $\mathcal{K}^{\mu\nu}$ and $\mathcal{H}^{\mu\nu}$ are given by

$$\mathcal{K}^{\mu\nu} = 4[G^{\mu\nu}\Box + \frac{1}{2}R\nabla^{\mu}\nabla^{\nu} + (g^{\mu\nu}R^{\rho\sigma} - R^{\mu\rho\nu\sigma})\nabla_{\rho}\nabla_{\sigma}, -R^{\nu\rho}\nabla_{\rho}\nabla^{\mu} + R^{\mu\rho}\nabla_{\rho}\nabla^{\nu}]f(\phi)$$
(5.4)

$$\mathcal{H}^{\mu\nu} = 2R^{\mu\rho\sigma\tau}R^{\nu}{}_{\rho\sigma\tau} - RR^{\mu\nu} + \frac{1}{2}R^{\mu}{}_{\rho}R^{\nu\rho} - R^{\mu\rho\sigma\tau}R^{\nu}{}_{\rho\sigma\tau}.$$
(5.5)

The tensor $\mathcal{K}^{\mu\nu}$ represents an operator which acts on $f(\phi)$. The energy-momentum tensor for the scalar field is

$$\mathcal{T}^{\mu\nu}_{\phi} = \nabla^{\mu}\phi\nabla^{\nu}\phi - \frac{1}{2}g^{\mu\nu}\nabla_{\rho}\phi\nabla^{\rho}\phi.$$
(5.6)

The stress tensor for anisotropic compact object is given as

$$T^{\mu\nu} = (\rho + P_t) u^{\mu} u^{\nu} + P_t g^{\mu\nu} + (P - P_t) \chi^{\mu} \chi^{\nu}, \qquad (5.7)$$

with energy density $\rho_{CO} = \rho(r) \equiv c^2 \rho(r)$, transverse pressure $P_t(r)$ and radial pressure P(r) of the homogeneously distributed matter in the compact object (CO), where u^{μ} is the four-velocity of the fluid and χ^{μ} is the unit space-like vector in the radial direction. In limit $\alpha \longrightarrow 0$, one can see that this is equivalent to the original form given by the Bekenstein-Hawking entropy. One can check that for D = 2, the topological term characterized by a vanishes Lovelock coupling α . The Barrow entropy [59] is a new black hole entropy which is given by

$$S = \pi \left(\frac{A}{A_0}\right)^{1+\delta/2},\tag{5.8}$$

where $0 \le \delta \le 1$, A is the black hole horizon area and A_0 is the Planck area [56]. When $\delta = 0$, the area law is restored, i.e. $S = \frac{A}{4G}$ (where $A_0 = 4G$) while $\delta = 1$ represents the most intricate and fractal structure of the horizon.

The form the functional $f(\phi)$ can take $f(\phi) = e^{-\gamma\phi}$ [60], where γ is a constant, which corresponds to EGB gravity coupled with dilaton that arises as a low-energy limit of the string theory [61]. Using the same principle as this entropy, we assume that for the compact object (or black hole) exterior, there is a presence of scalar fields ϕ , while inside the compact object is replaced by the Gauss-Bonnet (GB) coupling α as

coupling constant=
$$f(\phi)$$
, CO exterior $(D = 4)$
coupling constant= $\frac{\alpha}{D-4}$, CO interior $(D \to 4)$. (5.9)

The GB coupling α is measured in km^2 . In the CO exterior interior, we have rescaled the coupling constant $\alpha \to \alpha/(D-4)$. The negative (positive) α leads to a decrease (increase) of the CO radius and the maximum mass [63]. If $\alpha < 0$ the solution is still the anti-de Sitter (AdS) space, if $\alpha > 0$ the solution is the de Sitter (dS) space [62]. We investigate in detail the impact of the Gauss-Bonnet coupling on the properties of an anisotropic compact object, such as mass, radius and the factor of compactness. Considering the limit $D \to 4$, and it has an effect on gravitational dynamics in 4D. Additionally, at the CO boundary (r = R), the GB coupling must be continuous, i.e. $f(\phi) \equiv \frac{\alpha}{D-4}$. On the other hand, the function $f(\phi)$ describes the star's exterior region. To study the equations of motion inside and outside CO, we differentiate between two cases:

In CO interior $(D \to 4)$ we have:

$$G^{\mu\nu} + \alpha \left(\mathcal{H}^{\mu\nu} + \frac{1}{2} g^{\mu\nu} \mathcal{G} \right) = \frac{\kappa^2}{2} T^{\mu\nu}.$$
 (5.10)

In CO exterior (D = 4) we have:

$$G^{\mu\nu} + \mathcal{K}^{\mu\nu} + f\left(\phi\right) \left[\mathcal{H}^{\mu\nu} - \frac{1}{2f\left(\phi\right)}g^{\mu\nu}\left(\nabla_{\lambda}\phi\nabla^{\lambda}\phi + 4V_{eff}\left(\phi\right)\right)\right] = \frac{1}{2}\kappa^{2}T^{\mu\nu},\qquad(5.11)$$

with $g_{\mu\nu}\mathcal{T}^{\mu\nu}_{\phi} = -\nabla_{\lambda}\phi\nabla^{\lambda}\phi$. In CO interior $(D \to 4)$ we have: The GB invariant can be greatly simplified to the matter density [64] and using Eq.(5.3) we obtain

$$\rho_{f(\phi)} = 4\mathcal{G} - \frac{4}{f(\phi)} \left(\frac{1}{4} \nabla_{\lambda} \phi \nabla^{\lambda} \phi + V(\phi) \right), \quad \text{CO exterior } (D = 4) \\
\rho_{CO} = \mathcal{G}, \quad \text{CO interior } (D \to 4).$$
(5.12)

In this case, the term ρ_{CO} represents the density of matter in compact objects, and $\rho_{f(\phi)}$ is the density of dark matter surrounding these objects. Note that the relation between $\rho_{f(\phi)}$ and ρ_{CO} highlight the chameleon dark matter [65, 66]. Formally, at the points where the dark matter density ρ_{DM} equal to $\rho_{f(\phi)}$. For $V(\phi) \approx -\frac{1}{4} \nabla_{\lambda} \phi \nabla^{\lambda} \phi$, we obtain $\rho_{DM} \approx 4\rho_{CO}$, which is in good agreement with the observation data of the percentages of dark matter and the matter in the Universe [69, 57]: $\rho_{DM} \equiv 80\%$ and $\rho_{matter}(\rho_{CO}) \equiv 20\%$.

Next, we assume that $\phi = \phi(t)$. In cosmological and quintessence behavior [67, 68], the energy density ρ_{ϕ} and pressure P_{ϕ} of the scalar field are given by

$$\rho_{\phi} = \frac{1}{2}\dot{\phi}^{2} + V(\phi), \qquad (5.13)$$

$$P_{\phi} = \frac{1}{2}\dot{\phi}^{2} - V(\phi).$$

The quintessence models describe dark energy with a scalar field ϕ . In this case, ρ_{ϕ} and P_{ϕ} are respectively, the density and the pressure of the dark energy (DE). The Planck Collaboration [69] provides constraints on the equation of state $\omega_{\phi} = P_{\phi}/\rho_{\phi} \approx -1.028 \pm 0.032$. Starting from Eqs. (5.13), we obtain

$$\rho_{DM} = 4\rho_{CO} + \frac{1}{f(\phi)} \left(P_{\phi} - 3\rho_{\phi} \right).$$
 (5.14)

For small $f(\phi)$, the DM chameleon effect vanishes and the DM density depends only on $(P_{\phi} - 3\rho_{\phi})/f(\phi)$.

The scalar field always sits at the minimum of its effective potential. We assume that a massive scalar field begins oscillating about a minimum. The mass of small fluctuations around ϕ_{\min} gives a new scalar field mass as effective mass by $m_{eff}^2 = \frac{\partial^2}{\partial \phi^2} V_{eff}(\phi) \Big|_{\phi = \phi_{\min}}$. From Eqs. (5.3,5.13,5.14) we obtain

$$m_{eff}^{2} = \frac{1}{2} \frac{\partial^{2}}{\partial \phi^{2}} \left(\rho_{\phi} - P_{\phi} + \frac{\rho_{CO}}{\rho_{DM} - 4\rho_{CO}} \left(3\rho_{\phi} - P_{\phi} \right) \right)_{\phi = \phi_{\min}}.$$
 (5.15)

From Eq. (5.14), the functional coupling is given by

$$f(\phi) = \frac{P_{\phi} - 3\rho_{\phi}}{\rho_{DM} - 4\rho_{CO}}.$$
(5.16)

In the void ($\rho_{CO} = 0, \omega_{\phi} \sim -1$), we have $f(\phi) = -4\rho_{\phi}/\rho_{DM}$. In the CO surface, we assume that $\rho_{DM} \approx 0$ and $P_{\phi} \approx 0$, so we get $f_{surface}(\phi) \approx 3\rho_{\phi}/4\rho_{CO}$. Since ρ_{ϕ} represents

the density of DE according to quintessence, the effect of DE is weak on the CO surface, which shows that $f_{surface}(\phi) (\propto \rho_{\phi}) \rightarrow 0$. Inside CO, we have $(P_{\phi} = \rho_{\phi} = 0)$, i.e. $f(\phi) = 0$, which exactly corresponds with the assumption Eq. (5.9). For this reason, we exclude $f(\phi)$ inside matter, and we replace it with the GB coupling α (see the next section).

6 EGB primordial black holes

We start by the action of Einstein-Gauss-Bonnet (EGB) gravity [48] in 4-dimensions as

$$\mathcal{I} = \int d^D x \sqrt{-g} \left(\frac{M_p^2}{2} R - 2\Lambda + f(\phi) \left(R^2 - 4R_{\mu\nu} R^{\mu\nu} + R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right) \right), \qquad (6.1)$$

where S_m is the matter action, $M_p^2 = \frac{c^4}{8\pi G} \approx 2 \times 10^{18} \, [GeV]$, R is the Ricci scalar. We define the Gauss-Bonnet invariant as $\mathcal{G} = \mathcal{L}_2 = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$, where $\alpha_2 = \alpha/(D-4)$ is the Gauss-Bonnet coupling have dimensions of $[length]^2$, that represent ultraviolet (UV) corrections to Einstein theory. To solving the field equation we obtain the black hole solution $ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)$. Taking the limit $D \longrightarrow 4$, we obtain the exact solution in closed form

$$f(r) \approx 1 + \frac{r^2}{2\alpha} \left(1 - \sqrt{1 + 4\alpha \left(\frac{2M}{r^3} \pm \frac{1}{l^2}\right)} \right). \tag{6.2}$$

This last solution could be obtained directly from the derivation done in [49]. In the limit $r \to \infty$ with vanishing black hole charge, we asymptotically obtain the GR Schwarzschild solution. In the limit $\alpha \to 0$, we can recover the Reissner-Nordström-AdS solution. If $\alpha < 0$ the solution is still an AdS space, if $\alpha > 0$ the solution is a de Sitter (dS) space, [50]. The solutions show that the event horizon is located at $R_{\pm} = M \pm \sqrt{M^2 - \alpha}$, where $R_H = R_+$ and R_- are the event horizon and the Cauchy horizon radius of the EGB black hole [47]. We can express the ADM mass M of the black hole in terms of R_H by solving f(r) = 0 for $r = R_H$ resulting in

$$M = \frac{l^{-2}R_H^4 + R_H^2 + \alpha}{2R_H}.$$
(6.3)

The Hawking temperature of the EBG black hole can be calculated as

$$T = \frac{3l^{-2}R_H^4 + R_H^2 - \alpha}{8\pi\alpha R_H + 4\pi R_H^3}.$$
(6.4)

The thermodynamic volume $V = \frac{4\pi R_H^3}{3}$ is the conjugate variable to the pressure. The parameters V and A are the conjugate quantities of the pressure P and GB coupling

parameter α . The event horizon in spacetime can be located by solving the metric equation: f(r) = 0, and from Eq.(6.2) we obtain

$$\rho_{BH} + P = \frac{3}{8\pi R_H^2} + \frac{3\alpha}{8\pi R_H^4},\tag{6.5}$$

where $M = \rho_{BH}V$, $V = 4\pi r^3/3$ and $P = 3/8\pi l^2$. For the limit $\alpha \longrightarrow 0$, we can recover the density $\rho_{\Lambda} = 3/16\pi R_H^2$ of the holographic dark energy (DE) [70]. In the spatially flat homogeneous and isotropic universe in the FLRW, the (modified) Friedmann equations can be obtained [48] $H^2 + \alpha H^4 = \frac{8\pi G}{3}\rho + \frac{\Lambda}{3}$ and $(H^2 + \alpha H^4) \dot{H} = -4\pi G (\rho + P)$. This equation looks like the Eq.(6.5), i.e. can describe the dynamics in space-time associated to a black hole. For Gauss–Bonnet branch, we introduce the topological density as

$$\rho_{\alpha} = \rho_{\alpha}^{(2)} = \frac{\alpha}{16\pi R_{H}^{4}}.$$
(6.6)

From Eq. (6.5) we obtain the Van der Waals equation:

$$\left(P + \rho_{BH}\right) \left(V - \frac{4\pi\alpha R_H}{3}\right) = \frac{R_H}{2}.$$
(6.7)

In the limit $\alpha \longrightarrow 0$ and $P \gg \rho$, we can recover the ideal gas law. The critical point occurs when P(V). We note that the Gauss-Bonnet coupling α represente measure of the average attraction between particles. Before proceeding further, we note that the mass M can be interpreted as a chemical enthalpy [51], which is the total energy of the black hole [52] including both the energy PV and the internal energy E and required to displace the vacuum energy of its environment. The Hubble horizon mass is connected with the Smarr formula as

$$M_H + PV = \frac{R_H}{2} + \frac{\alpha}{2R_H},\tag{6.8}$$

where $M_H = \frac{4\pi R_H^3}{3}\rho$ is the Hubble horizon mass. We have also

$$R_H \approx 2M - \frac{\alpha}{2M}.\tag{6.9}$$

We can recover the Schwarzschild radius for $\alpha \approx 0$. This expression gives an interpretation of the difference between Schwarzschild and EGB black holes. From Eqs. (6.8,6.9), we obtain the the Smarr formula:

$$M = M_H + PV + \frac{\alpha}{4M} \left(1 - \frac{1}{1 - \frac{\alpha}{4M^2}} \right).$$
(6.10)

In the limit $\alpha \longrightarrow 0$, we can recover the standard Smarr formula. Let us mention that for the AdS-Schwarzschild limit and using $M \approx M_H$, we get $PV \propto = -\alpha/4M \propto T$. This expression gives an interpretation to the term that depends on α , as a solution of AdS-GB gravity in the presence of a perfect fluid [53]. This is not valid for the dS-GB black holes.

$$PV = (M - M_H) - \frac{\alpha}{4M} \left(1 - \frac{1}{1 - \frac{\alpha}{4M^2}} \right).$$
(6.11)

For Gauss-Bonnet branch we obtain $\left(1 - \frac{4M^2}{\alpha}\right)\left(1 - \frac{\alpha}{4M^2}\right) = 1$. One of two solutions to this equation is the Golden ratio $(\alpha > 0)$ in de Sitter (dS) space: $\frac{\alpha}{4M^2} = \frac{1+\sqrt{5}}{2}$ or $\frac{4M^2}{\alpha} = \frac{1+\sqrt{5}}{2}$. If $\alpha < 0$ the solution is still an AdS space: $\frac{\alpha}{4M^2} = \frac{1-\sqrt{5}}{2}$ or $\frac{4M^2}{\alpha} = \frac{1-\sqrt{5}}{2}$. We obtain the spectrum of mass in the Gauss-Bonnet branch is

$$M_{\alpha} = \frac{\sqrt{\gamma \alpha}}{2},\tag{6.12}$$

with $\gamma = \left\{\frac{2}{1-\sqrt{5}}, \frac{1-\sqrt{5}}{2}, \frac{2}{1+\sqrt{5}}, \frac{1+\sqrt{5}}{2}\right\}$. It can be written as $\gamma \approx \{-1.618, -0.618, 0.618, 1.618\}$. This shows that the Gauss-Bonnet coupling represents a topological black hole mass.

α	γ	M_{α}
-0.5	-1.618	0.449
-0.5	-0.618	0.277
0.5	0.618	0.277
0.5	1.618	0.449

Table 2: Numerical estimate of the values of the PBH mass, according to the frequency f and the interval of N between 10^{-47} and 60. Such as $f = 2.561 \times 10^{33} \frac{N}{M}$.

Taking the derivative with respect to time t in both sides of Eq. (6.9), thus the mass loss rate of a black hole is obtained as

$$\frac{dM}{dt} \approx \frac{R_H}{2} \left(1 - \frac{\alpha}{2M^2} \right), \tag{6.13}$$

with $\dot{R}_H = \frac{dR_H}{dt}$. Using the 4D Stefan-Boltzmann law [54, 58] $\frac{dM}{dt} = -\pi^2 A T^4/60$, with T and A are temperature and area of black hole, respectively. Evidently, we have $\alpha/M^2 \propto T^4$. For the Gauss-Bonnet branch and from Eq. (6.12), we have $\frac{dM}{dt} \approx \frac{\dot{R}_H}{2} (1-2\gamma)$. Taking the Friedmann equation $HR_H = 1$ [70] and from Eq. (6.9) we have

$$M_{\pm}(t) = \frac{1}{4H(t)} \pm \frac{1}{4} \left(\frac{1}{H(t)} + 4\alpha\right)^{1/2}.$$
(6.14)

Focusing on $M_{-}(t)$, one can find the condition $\frac{1}{H}\left(\frac{1}{H}-1\right) \geq \frac{4\alpha}{\theta^2}$. For $\alpha = 0.5$ we plot M_{\pm} vs H Fig. (1, left). Using H = q/t in the radiation-dominated era (q = 1/2) for $\alpha = -0.5$, we plot M_{\pm} vs t Fig. (1, right). In Fig. (1, right), the domain in red, represents the interval where the evolution of the mass with respect to time begins to increase. While,



Figure 1: Left, curves of M_{\pm} vs H from Eq. (6.14)with $\alpha = 0.5$. Right, M_{+} vs t (dashed line) and M_{-} vs t (solid line) for the values q = 1/2 and $\alpha = -0.5$.

before this interval, the mass decreases over time which is due to Hawking evaporation. According to this figure, two types of PBHs have opposite properties such as the evolution of their masses over time.

7 Conclusion

The coupling between the scalar field ϕ and the 4D Einstein-Gauss-Bonnet gravity action is studied. We have studied the model describing the compact stars in 4D Einstein-Gauss-Bonnet gravity surrounded by scalar dark matter. We have made a comparison between the star interior and the exterior region. In this case, the Gauss-Bonnet coupling describes the interior structure of the black hole, while the coupling function $f(\phi)$ describes the black hole exterior region. We have shown that $f(\phi) = 0$ inside the black hole. By solving the equations of motion based on the chameleon mechanism. We refer to ϕ as a chameleon field, since its physical properties, such as its mass, depend sensitively on the environment. It can be used to demonstrate the relation between the cosmological constant and the matter density. The Gauss-Bonnet vacuum have a chameleon structure, and the mass m_{ϕ} appears with a small fluctuations. Moreover, we have studied the vacua in maximally symmetric solutions, and we obtain two branches of solutions, i.e. there are two such vacua. The first solution corresponds to the dark energy and the second represents the chameleonic vacua or dark matter. We have found strong mixing between the two vacua, which is presented by the chameleon mechanism. We also discussed the equation of state parameter in the model of the scalar field ϕ minimally coupled to EGB gravity, also in the maximally symmetric space. The percentages of the effective cosmological constants are in good agreement with those of dark matter and dark energy in the Universe.

We have considered 4D Einstein-Gauss-Bonnet gravity. We establish the relationship between a topological mass spectrum and the golden ratio. We have obtained the mass loss rate, which gives by taking the Friedmann equation of type $HR_H = 1$ [70], the evolution of the mass over time, and the evaporation Hawking time. For such a choice and fixed values of Lovelock coupling, we have plotted the mass of PBH versus the Hubble parameter and time.

References

- Del Popolo, Antonino, and Morgan Le Delliou. "Small scale problems of the ΛCDM model: a short review." Galaxies 5.1 (2017): 17.
- [2] D. Glavan, & C. Lin, Einstein-Gauss-Bonnet Gravity in Four-Dimensional Spacetime, Phys. Rev. Lett. 124(8) (2020), 081301.
- [3] Easson, D. A., Manton, T., & Svesko, A. (2020). D→4 Einstein-Gauss-Bonnet gravity and beyond. Journal of Cosmology and Astroparticle Physics, 2020(10), 026.
- [4] D. Lovelock, The four-dimensionality of space and the Einstein tensor, J. Math. Phys. 13(6) (1972), 874-876.
- Bonifacio, J., Hinterbichler, K., & Johnson, L. A. (2020), Amplitudes and 4D Gauss-Bonnet Theory, Phys. Rev. D, 102(2), 024029.
- [6] Gürses, M., Şişman, T. Ç., & Tekin, B. (2020). Comment on "Einstein-Gauss-Bonnet Gravity in Four-Dimensional Spacetime", Phys. Rev. Lett., 125(14), 149001.
- [7] Wang, D., & Mota, D. (2021). 4D Gauss–Bonnet gravity: Cosmological constraints, H0 tension and large scale structure, Phys. Dark Universe, 100813.
- [8] Wu, C. H., Hu, Y. P., & Xu, H. (2021). Hawking evaporation of Einstein–Gauss– Bonnet AdS black holes in D≥ 4 dimensions, Eur. Phys. J. C, 81(4), 1-9.
- [9] Horndeski, G. W. (1974). Second-order scalar-tensor field equations in a fourdimensional space. Int. J. Theor. Phys., 10(6), 363-384.
- [10] Guo, M., & Li, P. C. (2020), Innermost stable circular orbit and shadow of the 4 D Einstein–Gauss–Bonnet black hole, Eur. Phys. J. C, 80(6), 1-8.

- [11] Konoplya, R. A., & Zinhailo, A. F., 2020, Eur. Phys. J. C, 80(11), 1-13.
- [12] Banerjee, A., Tangphati, T., & Channuie, P., 2021, Astrophys. J., 909(1), 14.
- [13] Hansraj, S., Banerjee, A., Moodly, L., & Jasim, M. K., 2020, Class. Quantum Gravity, 38(3), 035002.
- [14] Banerjee, A., Tangphati, T., Samart, D., & Channuie, P., 2021, Astrophys. J., 906(2), 114.
- [15] Banerjee, A., & Singh, K. N., 2021, Phys. Dark Universe, 31, 100792.
- [16] Tangphati, T., Pradhan, A., Errehymy, A., & Banerjee, A., 2021, Phys. Lett. B, 136423.
- [17] East, W. E., & Ripley, J. L. (2021). Evolution of Einstein-scalar-Gauss-Bonnet gravity using a modified harmonic formulation. Physical Review D, 103(4), 044040.
- [18] Horndeski, G. W. (1974). Second-order scalar-tensor field equations in a fourdimensional space. International Journal of Theoretical Physics, 10(6), 363-384.
- [19] Charmousis, C., & Padilla, A. (2008). The instability of vacua in Gauss-Bonnet gravity. Journal of High Energy Physics, 2008(12), 038.
- [20] Ma, L., & Lü, H. (2020). Vacua and exact solutions in lower-D limits of EGB. The European Physical Journal C, 80(12), 1-10.
- [21] Panah, B. E., Jafarzade, K., & Hendi, S. H. (2020). Charged 4D Einstein-Gauss-Bonnet-AdS black holes: shadow, energy emission, deflection angle and heat engine. Nuclear Physics B, 961, 115269.
- [22] Markkanen, T., Nurmi, S., Rajantie, A., & Stopyra, S. (2018). The 1-loop effective potential for the Standard Model in curved spacetime. Journal of High Energy Physics, 2018(6), 1-40.
- [23] Linde, A. D. (1983). Decay of the false vacuum at finite temperature. Nuclear Physics B, 216(2), 421-445.
- [24] Coleman, S., & Luccia, F. D. (1980). Gravitational effects on and of vacuum decay. In EUCLIDEAN QUANTUM GRAVITY (pp. 295-305).
- [25] Nojiri, S. I., Odintsov, S. D., & Sasaki, M. (2005). Gauss-Bonnet dark energy. Physical Review D, 71(12), 123509.

- [26] Glavan, D., & Lin, C. (2020). Einstein-Gauss-Bonnet Gravity in Four-Dimensional Spacetime. Physical Review Letters, 124(8), 081301.
- [27] Bousder, M., Salmani, E., El Fatimy, A., & Ez-Zahraouy, H. (2023). Holographic dark energy satisfying the energy conditions in Lovelock gravity. Annals of Physics, 169282.
- [28] Catà, O., Ibarra, A., & Ingenhütt, S. (2016). Dark matter decays from nonminimal coupling to gravity. Physical review letters, 117(2), 021302.
- [29] Aoki, K., Gorji, M. A., & Mukohyama, S. (2020). A consistent theory of $D \longrightarrow 4$ Einstein-Gauss-Bonnet gravity. Physics Letters B, 135843.
- [30] Caldwell, R. R., Kamionkowski, M., & Weinberg, N. N. (2003). Phantom energy: dark energy with w<-1 causes a cosmic doomsday. Physical Review Letters, 91(7), 071301.
- [31] Pi, Shi, et al. "Scalaron from R2-gravity as a heavy field." Journal of Cosmology and Astroparticle Physics 2018.05 (2018): 042.
- [32] Khoury, Justin, and Amanda Weltman. "Chameleon cosmology." Physical Review D 69.4 (2004): 044026.
- [33] Katsuragawa, Taishi, and Shinya Matsuzaki. "Cosmic history of chameleonic dark matter in F (R) gravity." Physical Review D 97.6 (2018): 064037.
- [34] Gannouji, Radouane, M. Sami, and I. Thongkool. "Generic f (R) theories and classicality of their scalarons." Physics Letters B 716.2 (2012): 255-259.
- [35] Katsuragawa, T., & Matsuzaki, S. (2018). Cosmic history of chameleonic dark matter in F (R) gravity. Physical Review D, 97(6), 064037.
- [36] Choudhury, S., Sen, M., & Sadhukhan, S. (2016). Can dark matter be an artifact of extended theories of gravity?. The European Physical Journal C, 76(9), 1-24.
- [37] Frolov, A. V. (2008). Singularity problem with f (R) models for dark energy. Physical review letters, 101(6), 061103.
- [38] Crisostomo, J., Troncoso, R., & Zanelli, J. (2000). Black hole scan. Physical Review D, 62(8), 084013.
- [39] Shu, F. W. (2020). Vacua in novel 4D Einstein-Gauss-Bonnet gravity: Pathology and instability?. Physics Letters B, 811, 135907.

- [40] Ema, Y., Jinno, R., & Nakayama, K. (2020). High-frequency graviton from inflaton oscillation. Journal of Cosmology and Astroparticle Physics, 2020(09), 015.
- [41] Elder, B., Khoury, J., Haslinger, P., Jaffe, M., Müller, H., & Hamilton, P. (2016). Chameleon dark energy and atom interferometry. Physical Review D, 94(4), 044051.
- [42] Brax, P., van de Bruck, C., Davis, A. C., Khoury, J., & Weltman, A. (2004). Detecting dark energy in orbit: The cosmological chameleon. Physical Review D, 70(12), 123518.
- [43] Aghanim, N., Akrami, Y., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., ... & Roudier, G. (2020). Planck 2018 results-VI. Cosmological parameters. Astronomy & Astrophysics, 641, A6.
- [44] García-Aspeitia, M. A., & Hernández-Almada, A. (2021). Einstein–Gauss–Bonnet gravity: Is it compatible with modern cosmology?. Physics of the Dark Universe, 100799.
- [45] Hashiba, S., Yamada, Y., & Yokoyama, J. I. (2021). Particle production induced by vacuum decay in real time dynamics. Physical Review D, 103(4), 045006.
- [46] Krishnan, C., Mohayaee, R., Colgáin, E. Ó., Sheikh-Jabbari, M. M., & Yin, L. (2021). Does Hubble tension signal a breakdown in FLRW cosmology?. Classical and Quantum Gravity, 38(18), 184001.
- [47] P. G. Fernandes, Charged black holes in AdS spaces in 4D Einstein Gauss-Bonnet gravity, Phys. Lett. B, 135468 (2020).
- [48] Fernandes, P. G., Carrilho, P., Clifton, T., & Mulryne, D. J. (2022). The 4D Einstein-Gauss-Bonnet theory of gravity: a review. Class. Quant. Grav.
- [49] M. Cvetič, S. I. Nojiri and S. D. Odintsov, Black hole thermodynamics and negative entropy in de Sitter and anti-de Sitter Einstein–Gauss–Bonnet gravity, Nucl. Phys. B, 628(1-2), 295-330, (2002).
- [50] P. G. Fernandes, Charged black holes in AdS spaces in 4D Einstein Gauss-Bonnet gravity, Phys. Lett. B, 135468 (2020).
- [51] D. Kastor, S. Ray and J. Traschen, Enthalpy and the Mechanics of AdS Black Holes, Class. Quant. Grav., 26 195011, (2009).
- [52] D. Kubizňák, R. B. Mann and M. Teo, Black hole chemistry: thermodynamics with Lambda, Class. Quant. Grav., 34(6), 063001 (2017).

- [53] N. Farhangkhah, New black hole solutions of Gauss-Bonnet gravity in the presence of a perfect fluid, Iran. J. Phys. Res., 17(5), 729-735 (2019).
- [54] C. H. Wu, Y. P. Hu and H. Xu. Hawking evaporation of Einstein–Gauss–Bonnet AdS black holes in $D \ge 4$ dimensions, Eur. Phys. J. C, **81(4)**, 1-9 (2021).
- [55] Bousder, M. (2022). A new constant behind the rotational velocity of galaxies. Journal of Cosmology and Astroparticle Physics, 2022(01), 015.
- [56] Bousder, M. (2021). Quantum f (R) gravity and AdS/CFT. arXiv preprint arXiv:2104.14434.
- [57] Bousder, M., Sakhi, Z., & Bennai, M. (2020). A new unified model of dark matter and dark energy in 5-dimensional f (R, φ) gravity. International Journal of Geometric Methods in Modern Physics, 17(13), 2050183.
- [58] S. B. Giddings, Phys. Lett. B, **754**, 39-42 (2016).
- [59] Barrow, J. D. (2020). The area of a rough black hole. Phys. Lett. B, 808, 135643.
- [60] Antoniou, G., Bakopoulos, A., & Kanti, P., 2018, Phys. Rev. D, 97(8), 084037.
- [61] Maeda, K. I., Ohta, N., & Sasagawa, Y. 2009, Phys. Rev. D, 80(10), 104032.
- [62] Fernandes, P. G. (2020), Phys. Lett. B, 135468.
- [63] Doneva, D. D., & Yazadjiev, S. S., 2021, J. Cosmol. Astropart. Phys., 2021(05), 024.
- [64] Bean, R., Flanagan, E. E., & Trodden, M. 2008, Phys. Rev. D, 78(2), 023009.
- [65] Khoury, J., & Weltman, A., 2004, Phys. Rev. D, 69(4), 044026.
- [66] Folomeev, V., Aringazin, A., & Dzhunushaliev, V., 2013, Phys. Rev. D, 88(6), 063005.
- [67] Matos, T., & Urena-Lopez, L. A., 2000, Class. Quantum Gravity, 17(13), L75.
- [68] 4Arbey, A., & Coupechoux, J. F., 2021, J. Cosmol. Astropart. Phys., 2021(01), 033.
- [69] Aghanim, N., Akrami, Y., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., ... & Roudier, G., 2020, Astron. Astrophys., 641, A6.
- [70] M. Li, A model of holographic dark energy. Phys. Lett. B, **603(1-2)**, 1-5 (2004).