

Hyper Suprime-Cam Year 3 Results: Cosmology from Galaxy Clustering and Weak Lensing with HSC and SDSS using the Emulator Based Halo Model

Hironao Miyatake^{1,2,3,*} Sunao Sugiyama^{3,4,†} Masahiro Takada^{3,‡} Takahiro Nishimichi^{5,3,6} Xiangchong Li^{7,3}
 Masato Shirasaki^{8,9} Surhud More^{10,3} Yosuke Kobayashi^{11,3} Atsushi J. Nishizawa^{12,13} Markus M. Rau¹⁴
 Tianqing Zhang⁷ Ryuichi Takahashi¹⁵ Roohi Dalal¹⁶ Rachel Mandelbaum⁷ Michael A. Strauss¹⁶
 Takashi Hamana⁸ Masamune Oguri^{17,18,4,3} Ken Osato^{17,19} Robert Armstrong²⁰ Arun Kannawadi¹⁶
 Yutaka Komiyama²¹ Robert H. Lupton¹⁶ Nate B. Lust¹⁶ Satoshi Miyazaki²² Hitoshi Murayama^{23,24,3}
 Yuki Okura⁸ Paul A. Price¹⁶ Tomomi Sunayama^{25,1} Philip J. Tait²² Masayuki Tanaka⁸ and Shiang-Yu Wang²⁶

¹*Kobayashi-Maskawa Institute for the Origin of Particles and the Universe (KMI), Nagoya University, Nagoya, 464-8602, Japan*

²*Institute for Advanced Research, Nagoya University, Nagoya 464-8601, Japan*

³*Kavli Institute for the Physics and Mathematics of the Universe (WPI),*

The University of Tokyo Institutes for Advanced Study (UTIAS), The University of Tokyo, Chiba 277-8583, Japan

⁴*Department of Physics, The University of Tokyo, Bunkyo, Tokyo 113-0031, Japan*

⁵*Center for Gravitational Physics and Quantum Information,*

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

⁶*Department of Astrophysics and Atmospheric Sciences, Faculty of Science,*

Kyoto Sangyo University, Motoyama, Kamigamo, Kita-ku, Kyoto 603-8555, Japan

⁷*McWilliams Center for Cosmology, Department of Physics,*

Carnegie Mellon University, Pittsburgh, PA 15213, USA

⁸*National Astronomical Observatory of Japan, National Institutes of Natural Sciences, Mitaka, Tokyo 181-8588, Japan*

⁹*The Institute of Statistical Mathematics, Tachikawa, Tokyo 190-8562, Japan*

¹⁰*The Inter-University Centre for Astronomy and Astrophysics, Post bag 4, Ganeshkhind, Pune 411007, India*

¹¹*Department of Astronomy/Steward Observatory, University of Arizona,*

933 North Cherry Avenue, Tucson, AZ 85721-0065, USA

¹²*Gifu Shotoku Gakuen University, Gifu 501-6194, Japan*

¹³*Institute for Advanced Research/Kobayashi Maskawa Institute, Nagoya University, Nagoya 464-8602, Japan*

¹⁴*High Energy Physics Division, Argonne National Laboratory, Lemont, IL 60439, USA*

¹⁵*Faculty of Science and Technology, Hirosaki University, 3 Bunkyo-cho, Hirosaki, Aomori 036-8561, Japan*

¹⁶*Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA*

¹⁷*Center for Frontier Science, Chiba University, Chiba 263-8522, Japan*

¹⁸*Research Center for the Early Universe, The University of Tokyo, Bunkyo, Tokyo 113-0031, Japan*

¹⁹*Department of Physics, Graduate School of Science, Chiba University, Chiba 263-8522, Japan*

²⁰*Lawrence Livermore National Laboratory, Livermore, CA 94551, USA*

²¹*Department of Advanced Sciences, Faculty of Science and Engineering,*

Hosei University, 3-7-2 Kajino-cho, Koganei-shi, Tokyo 184-8584, Japan

²²*Subaru Telescope, National Astronomical Observatory of Japan, 650 N Aohoku Place Hilo HI 96720 USA*

²³*Berkeley Center for Theoretical Physics, University of California, Berkeley, CA 94720, USA*

²⁴*Theory Group, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*

²⁵*Department of Astronomy and Steward Observatory, University of Arizona, 933 N Cherry Ave, Tucson, AZ 85719, USA*

²⁶*Institute of Astronomy and Astrophysics, Academia Sinica, Taipei 10617, Taiwan*

(Dated: April 4, 2023)

We present cosmology results from a blinded joint analysis of cosmic shear, $\xi_{\pm}(\vartheta)$, galaxy-galaxy weak lensing, $\Delta\Sigma(R)$, and projected galaxy clustering, $w_p(R)$, measured from the Hyper Suprime-Cam three-year (HSC-Y3) shape catalog and the Sloan Digital Sky Survey (SDSS) DR11 spectroscopic galaxy catalog – a 3×2 pt cosmology analysis. We define luminosity-cut, and therefore nearly volume-limited, samples of SDSS galaxies to serve as the tracers of w_p and as the lens samples for $\Delta\Sigma$ in three spectroscopic redshift bins spanning the range $0.15 < z < 0.7$. For the ξ_{\pm} and $\Delta\Sigma$ measurements, we use a single sample of about seven million source galaxies over 416 deg^2 , selected from HSC-Y3 based on having photometric redshifts (photo- z) greater than 0.75. The deep, high-quality HSC-Y3 data enable significant detections of the $\Delta\Sigma$ signals, with integrated signal-to-noise ratio $S/N \sim 24$ in the range $3 \leq R/[h^{-1}\text{Mpc}] \leq 30$ over the three lens samples. ξ_{\pm} has $S/N \sim 19$ in the range $8' \leq \vartheta \leq 50'$ and $30' \leq \vartheta \leq 150'$ for ξ_+ and ξ_- , respectively. For cosmological parameter inference, we use the `Dark Emulator` package, combined with a halo occupation distribution prescription for the relation between galaxies and halos, to model w_p and $\Delta\Sigma$ down to quasi-nonlinear scales, and we estimate cosmological parameters after marginalizing over nuisance parameters. In our baseline analysis we employ an *uninformative* flat prior of the residual photo- z error, given by $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$, to model a residual bias in the mean redshift of HSC source galaxies. Comparing the relative lensing amplitudes for $\Delta\Sigma$ in the three redshift bins and for ξ_{\pm} with the single HSC source galaxy sample allows us to calibrate the photo- z parameter Δz_{ph} to the precision of $\sigma(\Delta z_{\text{ph}}) \simeq 0.09$. With these methods, we obtain a robust constraint on the cosmological parameters for the flat Λ CDM model: $S_8 = \sigma_8(\Omega_m/0.3)^{0.5} = 0.763_{-0.036}^{+0.040}$ (68% C.I.), or the best-constrained parameter given by $S'_8 = \sigma_8(\Omega_m/0.3)^{0.22} = 0.721 \pm 0.028$, determined with about 4%

fractional precision. Our HSC-Y3 data exhibits about 2.5σ tension with the *Planck* inferred S_8 value for the Λ CDM model, and hints at a non-zero residual photo- z bias implying that the true mean redshift of the HSC galaxies at $z \gtrsim 0.75$ is higher than that implied by the original photo- z estimates.

I. INTRODUCTION

The cosmological standard model assuming the initial conditions predicted by an inflationary scenario, Λ Cold Dark Matter (Λ CDM) model, has been successful in explaining a variety of observations [e.g. 1]. Wide-area imaging galaxy surveys in optical and near-infrared wavelengths offer exciting opportunities to address fundamental questions in cosmology, such as the nature of dark matter and the origin of cosmic acceleration [e.g. 2]. Current-generation imaging surveys such as the Subaru Hyper Suprime-Cam¹ [HSC; 3–7], the Dark Energy Survey² [DES; 8], and the Kilo-Degree Survey³ [KiDS; 9] have used precise measurements of weak gravitational lensing effects to obtain tight constraints on cosmological parameters. Intriguingly, the cosmological models inferred from these large-scale structure probes consistently exhibit a lower value of σ_8 or S_8 , which characterizes the clustering amplitude in the late-time universe [Ref. 10, for a recent review], than do cosmological models inferred from the *Planck* cosmic microwave background (CMB) measurements [11]. This discrepancy might hint at the possibility of new physics beyond the standard Λ CDM model.

Challenges of large-scale structure probes lie in systematic effects/errors inherent both in observations and theory. One of the important observational systematic effects, relevant to weak lensing cosmology, arises from imperfect photometric redshift estimates (hereafter referred to as photo- z). Due to the limited information carried by broad-band photometry and/or difficulties in uniform and accurate characterization of individual galaxy photometry, photometric redshift estimates are not perfect. Hence, photo- z estimates need to be calibrated using a representative calibration sample of galaxies that have accurate redshift estimates; ideally we need a representative spectroscopic sample but the COSMOS catalog which provides 30-band photo- z 's is currently a main calibration sample for photo- z 's of faint galaxies [12].

The main systematic effects on the theory side lie in the difficulties in accurately modeling nonlinear structure formation, and the unknown relation between the distributions of matter and galaxies, where the latter is referred to as the galaxy bias uncertainty. The physical processes inherent in the formation and evolution of galaxies cannot yet be accurately and fully modeled from first principles. Nevertheless, on large scales, i.e., beyond a few 10 Mpc where gravity is a driving force of structure formation, the linear theory of structure formation is quite accurate, and predicts that the galaxy distribution for

any type of galaxy is related to the underlying matter distribution by a scale-independent factor, i.e the linear bias parameter [13]. On smaller scales, the bias function is scale-dependent, due to the mode coupling in nonlinear structure formation.

Combining multiple cosmological probes provides a promising way to mitigate the aforementioned systematic effects in cosmological inference. In this paper, we combine the projected correlation function of galaxies (w_p), galaxy-galaxy weak lensing ($\Delta\Sigma$), and cosmic shear correlations (ξ_{\pm}) – so-called 3×2 pt cosmology analysis, measured from the photometric HSC three-year (hereafter HSC-Y3) data covering about 416 deg^2 of the sky and the spectroscopic SDSS galaxy catalogs. By cross-correlating the positions of SDSS galaxies with shapes of the background HSC galaxies, we can measure the $\Delta\Sigma$ signal, which in turn allows us to infer the average matter distribution around the SDSS galaxies. We then combine the $\Delta\Sigma(R)$ measurement with the auto-correlation function of galaxies in the same sample, $w_p(R)$, as a function of projected separation R to observationally infer the galaxy bias function of the SDSS galaxies, including its scale-dependence. We will use a *single* sample of the HSC source galaxies to perform weak lensing measurements for $\Delta\Sigma$ for each of the SDSS galaxy subsamples that are subdivided into three spectroscopic redshift bins spanning the range $z = [0.15, 0.7]$. Comparing the relative $\Delta\Sigma$ amplitudes in the three redshift bins and the cosmic shear signal ξ_{\pm} , for a given sample of HSC source galaxies, enables us to *calibrate* any residual error in the mean photometric redshifts of HSC source galaxies, as proposed in Oguri and Takada [14] [also see 6, 15].

To resolve the modeling difficulties of clustering observables on small scales, we use the halo model approach [16–19]. Dark matter halos are self-gravitating systems where galaxies form. Clustering statistics of halos such as the halo mass function and the halo-matter and halo-halo correlation functions can be accurately modeled down to small scales using N -body simulations for a given cosmological model. Based on this motivation, Nishimichi *et al.* [20] used an ensemble of N -body simulations for different cosmologies to build an emulation package, dubbed as `Dark Emulator`, that enables fast and accurate computations of the halo clustering quantities as a function of halo masses, redshift and separations for an input cosmological model. As shown in Nishimichi *et al.* [20] [also see 6, 15, 21], the “scale-dependent” halo bias in the halo-matter and halo-halo correlation functions, relative to the matter correlation function, carries useful cosmological information beyond the linear theory.

The purpose of this paper is to use the combined 3×2 pt measurements from the photometric HSC-Y3 galaxies and the spectroscopic SDSS galaxies to estimate cosmological parameters while mitigating the impact of the systematic photo- z error and the galaxy bias uncertainty. We carried out a similar analysis with the HSC-Y1 data [6], but there are some important differences. First, while the HSC-Y1 analysis used only

* miyatake@kmi.nagoya-u.ac.jp

† sunao.sugiyama@ipmu.jp

‡ masahiro.takada@ipmu.jp

¹ <https://hsc.mtk.nao.ac.jp/ssp/>

² <https://www.darkenergysurvey.org>

³ <http://kids.strw.leidenuniv.nl>

measurements of $\Delta\Sigma$ and w_p , in this paper we include the cosmic shear measurements, ξ_{\pm} , to improve the precision of cosmological parameter inference and the calibration of the residual photo- z error parameter (hereafter Δz_{ph}). Second, we employ a completely uninformative flat prior of Δz_{ph} , $\mathcal{U}(-1, 1)$, in our baseline analysis method. Much narrower priors have been used in other weak lensing analyses; e.g., the HSC-Y1 2×2 pt analysis [6] used a Gaussian prior with width $\sigma(\Delta z_{\text{ph}}) = 0.1$ and many other weak lensing cosmology analyses use a prior with width $\mathcal{O}(10^{-2})$ [e.g. 8, 22] for the mean redshift of source galaxies. We will show that the statistical power of the HSC-Y3 data enables us to calibrate the Δz_{ph} parameter to a precision of $\sigma(\Delta z_{\text{ph}}) \sim 0.1$. For theoretical templates, we combine `Dark Emulator` and the halo occupation distribution, which gives a phenomenological description of the galaxy-halo connection, to model the $\Delta\Sigma$ and w_p observables down to quasi-nonlinear scales. We will validate our model and method using a synthetic data vector of the clustering observables, taking into account the covariance matrix for the HSC-Y3 and SDSS observables. In this paper, we will pay particular attention to a stringent test of the flat Λ CDM model, especially whether the HSC-Y3 data exhibits a tension in the S_8 constraint with the *Planck* result.

We perform a *blinded* cosmology analysis at the catalog and analysis levels to avoid confirmation bias. We carry out various tests for systematic errors in the measurements and do extensive validation tests of the method and model. During the blinded analysis stage, we determine the analysis setup, including the uninformative uniform prior of Δz_{ph} , without access to the values of cosmological parameters, and we agree not to make any changes in our analysis methodology after we unblind. We will explicitly mention any results that were found “post-unblinding”. This paper is one of a series of the HSC-Y3 cosmology papers: More, Sugiyama *et al.* [23] give detailed descriptions of the measurements used in the 3×2 pt analysis, Sugiyama *et al.* [24] use exactly the same 3×2 pt observables as those in this paper to perform a cosmology analysis using a perturbation theory based model, Li *et al.* [25] show cosmology results using the real-space cosmic shear tomography, and Dalal *et al.* [26] show cosmology results using the Fourier-space cosmic shear tomography. The two 3×2 pt papers (this paper and Sugiyama *et al.* [24]) use the same blinded shape catalog of the HSC data. Li *et al.* [25] and Dalal *et al.* [26] use different blinded catalogs. Thus we use three different blinded catalogs for our cosmology analyses. We compared the cosmological parameters from the 3×2 pt analyses and the real- and Fourier-space cosmic shear analyses only after unblinding. We believe that our analysis strategy and method allow us to obtain a robust, convincing result for both the cosmological parameters and the residual photo- z error, without being subject to confirmation bias.

This paper is organized as follows. In Section II we describe the HSC three-year shape catalog and the spectroscopic SDSS galaxy catalog that are used in this paper. In Section III we describe our analysis method: the theoretical templates based on the halo model and the likelihood analysis. In Section IV we describe our blinding strategy for the cosmology analysis. In Section V we show the main results of this pa-

per: our cosmological constraints, the robustness to different systematics, and the degree of tension of our results with the *Planck* inferred cosmology. In Section VI we give a detailed discussion of our cosmology results: the impact of the residual photo- z error and the assembly bias, and the cosmological results when combined with external constraints on Ω_m . Finally we give our conclusions in Section VII. We give technical details of our method and tests of systematic effects in several Appendices.

Throughout this paper we use the natural unit $c = 1$ for the speed of light. Unless stated otherwise, we quote the central value of a parameter from the mode value of the posterior parameter that has the highest probability in the marginalized 1D posterior distribution in the chain: $\mathcal{P}(p_{\text{mode}}) = \text{maximum}$. The justification of the use of mode as the central value is described in Dalal *et al.* [26]. We quote the 68% credible interval for the parameter(s) from the highest density interval of parameter(s) satisfying

$$\int_{\mathbf{p} \in \mathcal{P} > \mathcal{P}_{68}} d\mathbf{p} \mathcal{P}(\mathbf{p}) = 0.68, \quad (1)$$

where $\mathcal{P}(\mathbf{p})$ is the 1D or 2D marginalized posterior distribution. The 95% credible interval is similarly defined.

II. DATA

A. HSC-Y3 Data: Source Galaxies for Galaxy-galaxy Weak Lensing

HSC is a wide-field imaging camera on the prime focus of the 8.2m Subaru Telescope [27–30]. The HSC Subaru Strategic Program (HSC SSP) survey started in 2014, and used 330 Subaru nights to conduct a five-band (*grizy*) wide-area imaging survey [3]. The combination of HSC’s wide field-of-view (1.77 deg^2), superb image quality (typically $0.6''$ seeing FWHM in *i* band), and large photon-collecting power makes it one of the most powerful instruments for weak lensing measurements. The HSC SSP survey consists of three layers; Wide, Deep, and Ultradeep. The Wide layer, which is designed for weak lensing cosmology, covers about $1,100 \text{ deg}^2$ of the sky with a 5σ depth of $i \sim 26$ ($2''$ aperture for a point source). Since the *i*-band images are used for galaxy shape measurements in weak lensing analyses, they are preferentially taken under good seeing conditions.

In this paper, we use the HSC three-year (hereafter HSC-Y3) galaxy shape [31] and photo- z catalogs [32], constructed from the S19A internal data release (released in September 2019) of data acquired from March 2014 to April 2019. In the following subsections, we describe details of the shape and photo- z catalogs.

1. HSC-Y3 galaxy shape catalog

In this paper, we use the HSC-Y3 shape catalog [31] from the S19A images that were processed with `hscPipe v7` [33].

In `hscPipe v7`, there were a number of improvements to the PSF modelling, image warping kernel, background subtraction and bright star masks, which have improved the quality of the shape catalog in HSC-Y3 compared to the HSC Year 1 shape catalog [34, 35]. The detailed selection of galaxies that form the shape catalog is presented in Li *et al.* [31]. Briefly, the shape catalog consists of galaxies selected from the “full-depth full-color region” in all five filters. Apart from some basic quality cuts related to pixel level information, we select extended objects with an extinction corrected `cmodel` magnitude $i < 24.5$, i -band $\text{SNR} \geq 10$, resolution > 0.3 , $> 5\sigma$ detection in at least two bands other than i , a 1 arcsec diameter aperture magnitude cut of $i < 25.5$, and a blendedness cut in the i -band of $10^{-3.8}$.

The shape catalog consists of 35.7 million galaxies spanning an area of about 430 deg^2 , with an effective number density of 19.9 arcmin^{-2} . It is divided into six disjoint regions: XMM, VVDS, GAMA09H, WIDE12H, GAMA15H and HECTOMAP [see Fig. 2 in Ref. 31]. The shape measurements in the catalog were calibrated using detailed image simulations, such that the galaxy-property-dependent multiplicative shear bias uncertainty is less than $\sim 10^{-2}$. Li *et al.* [31] also presented a number of systematics tests and null tests, and quantify the level of residual systematics in the shape catalog that could affect the cosmological science analyses carried out using the data. Li *et al.* [31] flag residual additive biases due to PSF model shape residual correlations and star galaxy shape correlations as systematics requiring special attention and marginalization, so we will also investigate the effect of these systematics on the cosmic shear measurements.

As described in detail in companion papers, More *et al.* [23], Li *et al.* [25] and Dalal *et al.* [26], we find a significant source of B -mode systematics in the cosmic shear correlation functions for a $\sim 20 \text{ deg}^2$ patch in the GAMA09H region, and we remove this problematic region from the following analysis. The resultant total area of the HSC data is about 416 deg^2 .

2. Source galaxy catalog for galaxy-galaxy weak lensing

Given the depth of the HSC-Y3 data, we can define a secure sample of source galaxies behind lens galaxies, for galaxy-galaxy weak lensing measurements. In this paper we use three distinct samples of lens galaxies selected from the Data Release 11 (DR11) of spectroscopic SDSS galaxies up to $z = 0.7$, as described below. To select background galaxies behind the SDSS galaxies, we use photo- z estimates of each HSC galaxy. The HSC-Y3 shape catalog is accompanied by a photo- z catalog of galaxies based on three different methods [12]. MIZUKI [36] is a template fitting based photo- z estimation code. DEMPZ [37] and DNNZ [32] on the other hand provide machine-learning-based estimates of the galaxy photo- z 's. Each of these methods provides an estimate of the posterior distribution of the redshift for individual galaxies, denoted as $P(z_s)$. In this paper we employ the DEMPZ photo- z catalog as our fiducial choice to define a sample of background galaxies by requiring that the posterior that the galaxy

has redshift less than 0.75 be less than 1% [38–40]:

$$\int_{z_{1,\text{max}}+0.05}^7 dz_s P_i(z_s) \geq 0.99, \quad (2)$$

where $z_{1,\text{max}}=0.70$ is the maximum redshift of the lens samples. Such cuts significantly reduce the contamination of source galaxies which are physically associated with the lens galaxies and which would dilute the weak lensing signal. The total number of galaxies in our source sample is $\sim 24\%$ of the original HSC-Y3 shape catalog, with an effective number density of 4.9 galaxies per square arcmin. The mean redshift of the sample, estimated from the stacked photo- z posterior, is $\langle z_s \rangle \simeq 1.3$.

Photo- z uncertainties are one of the most important systematic effects in weak lensing cosmology, and could cause significant biases in the cosmological parameters if unknown residual systematic errors in photo- z exist. To minimize the impact of possible systematic photo- z error, we will employ the method in Oguri and Takada [14] that enables a self-calibration of such residual photo- z errors, using a *single* sample of photometric source galaxies for the weak lensing measurements as we will later describe in detail.

B. Lens Galaxy Sample

We use the large-scale structure sample compiled as part of DR11⁴ [41] of the SDSS-III Baryon Oscillation Spectroscopic Survey (BOSS) project [42] for measurements of the clustering of galaxies and as lens galaxies for the weak lensing signal measurements. The lens galaxy sample used in this paper is the same as that used in the first year analysis of HSC data (Sugiyama *et al.* [7], Miyatake *et al.* [15]) [also see 43]. We use a luminosity-limited catalog of SDSS galaxies in order for it to be approximately volume limited (see More *et al.* [23] for the details). We describe the resultant catalog here briefly.

The BOSS is a spectroscopic survey of galaxies and quasars selected from the imaging data obtained by the SDSS-I/II and covers an area of approximately $11,000 \text{ deg}^2$ [44] using the dedicated 2.5m SDSS Telescope [45]. Imaging data obtained in five photometric bands (*ugriz*) as part of the SDSS I/II surveys [46–48] were augmented with an additional $3,000 \text{ deg}^2$ in SDSS DR9 to cover a larger portion of the sky in the southern region [42, 49–51]. These data were processed by a series of photometric processing pipelines [52–54], and corrected for Galactic extinction [55] to obtain a reliable photometric catalog which serves as an input to select targets for spectroscopy [42]. The resulting spectra were processed by an automated pipeline to perform redshift determination and spectral classification [56]. The BOSS large-scale structure (LSS) samples are selected using algorithms focused on galaxies at different redshifts: $0.15 < z < 0.35$ (LOWZ) and $0.43 < z < 0.7$ (CMASS). In addition to the galaxies targeted by the BOSS project, we also use galaxies which pass

⁴ <https://www.sdss.org/dr11/>

the target selection but had already been observed as part of the SDSS-I/II project (legacy galaxies). These legacy galaxies are subsampled in each sector so that they obey the same completeness as that of the LOWZ/CMASS samples in their respective redshift ranges [57].

We use three galaxy subsamples divided into three redshift bins: “LOWZ” galaxies in the redshift range $z = [0.15, 0.35]$ and two subsamples of “CMASS” galaxies, hereafter called “CMASS1” and “CMASS2”, which are obtained from the division of CMASS galaxies into two redshift bins, $z = [0.43, 0.55]$ and $z = [0.55, 0.70]$, respectively. As shown in Fig. 1 of Miyatake *et al.* [6], we define each of the subsamples by selecting galaxies having absolute magnitudes $M_i - 5 \log h < -21.5$, -21.9 and -22.2 for the LOWZ, CMASS1 and CMASS2 subsamples, respectively. The comoving number densities for the *Planck* cosmology are $\bar{n}_g/[10^{-4} (h^{-1} \text{Mpc})^{-3}] \simeq 1.8, 0.74$ and 0.45 , respectively, which are a few times smaller than those of the entire parent (color-cut and flux-limited) LOWZ and CMASS samples.

III. MODELING AND ANALYSIS METHOD

In this paper, we use three clustering observables to perform the cosmological parameter inference – the so-called $3 \times 2\text{pt}$ analysis. To be more precise, we use (i) the average excess surface mass density profile, denoted as $\Delta\Sigma(R)$, that is measured from the galaxy-galaxy weak lensing combining the photometric HSC source galaxy sample and each of the three spectroscopic SDSS lens subsamples over the overlapping 416 deg^2 area of HSC-Y3 and BOSS, (ii) the projected correlation function, denoted as $w_p(R)$, for each of the spectroscopic SDSS subsamples used as lens samples in the $\Delta\Sigma$ analysis measured from the entire BOSS regions of about $8,300 \text{ deg}^2$ area, and (iii) the cosmic shear correlation functions, denoted as $\xi_{\pm}(\vartheta)$, for the HSC source sample measured from the HSC-Y3 416 deg^2 area. The details of the measurements, null and systematics tests and covariance matrix are described in the companion paper, More *et al.* [23]. In this section we describe our model of these clustering observables within the ΛCDM framework and our method of Bayesian based parameter inference.

A. Model

1. Dark Emulator

To model $\Delta\Sigma$ and w_p , we use the publicly-available code, Dark Emulator⁵, developed in Nishimichi *et al.* [20]. Dark Emulator is a software package enabling fast, accurate computations of halo clustering quantities for an input flat $w\text{CDM}$ cosmological model. Dark Emulator is based

TABLE I. The set of six cosmological parameters used in our analysis, which specify a model within the flat-geometry ΛCDM framework. For an input ΛCDM model, Dark Emulator outputs the halo clustering quantities (see text for details). The column labeled “parameters” lists the six cosmological parameters. The column labeled “supported range” denotes the range of parameters that is supported by Dark Emulator.

parameters	supported range [min,max]
Ω_{de}	[0.54752, 0.82128]
$\ln(10^{10} A_s)$	[2.4752, 3.7128]
$\omega_b \equiv \Omega_b h^2$	[0.0211375, 0.0233625]
$\omega_c \equiv \Omega_c h^2$	[0.10782, 0.13178]
n_s	[0.916275, 1.012725]

on an ensemble set of cosmological N -body simulations, each of which was performed with 2048^3 particles for a box with length 1 or 2 $h^{-1}\text{Gpc}$ on a side, for 101 flat $w\text{CDM}$ cosmological models. The $w\text{CDM}$ cosmology is parametrized by six parameters, $\mathbf{p} = \{\omega_b, \omega_c, \Omega_{\text{de}}, \ln(10^{10} A_s), n_s, w_{\text{de}}\}$, where $\omega_b (\equiv \Omega_b h^2)$ and $\omega_c (\equiv \Omega_c h^2)$ are the physical density parameters of baryons and CDM, respectively, h is the Hubble parameter, $\Omega_{\text{de}} \equiv 1 - (\omega_b + \omega_c + \omega_\nu)/h^2$ is the density parameter of dark energy for a flat-geometry universe, A_s and n_s are the amplitude and tilt parameters of the primordial curvature power spectrum normalized at $k_{\text{pivot}} = 0.05 \text{ Mpc}^{-1}$, and w_{de} is the equation of state parameter for dark energy. In the following we focus on flat ΛCDM cosmological models with $w_{\text{de}} = -1$.

For the N -body simulations, the effect of finite neutrino mass was included by fixing the neutrino density parameter $\omega_\nu \equiv \Omega_\nu h^2$ to 0.00064. This value corresponds to a total mass of three neutrino species of 0.06 eV, the lower bound of the normal mass hierarchy [58]. The presence of massive neutrinos affects the linear transfer function, where the total matter fluctuation was computed including massive neutrinos by CAMB [59] and was scaled back to the initial redshift of the simulations using the linear growth factor with the neutrino density included in the matter content. The subsequent non-linear growth was followed consistently in an N -body simulation, including the neutrino density as a part of matter density [see 20, for details]. Since we focus on the σ_8 parameter⁶, i.e., the present-day normalization of the linear matter power spectrum instead of the amplitude of the primordial fluctuations, this approximate treatment has little impact on our primary constraints from the HSC-Y3 and SDSS data.

The particle mass for the fiducial *Planck* cosmology is $m = 1.02 \times 10^{10} h^{-1} M_\odot$ for the higher resolution simulations used as the basis for Dark Emulator. The emulator uses halos with mass greater than $10^{12} h^{-1} M_\odot$, corresponding to about 100 simulation particles.

⁵ https://github.com/DarkQuestCosmology/dark_emulator_public

⁶ σ_8 is the parameter often used in the literature for the normalization of the linear matter power spectrum, corresponding to the rms linear mass density fluctuations within a top-hat sphere of radius $8 h^{-1} \text{Mpc}$.

For each N -body simulation realization (each redshift output) for a given cosmological model, Nishimichi *et al.* [20] constructed a catalog of halos using `Rockstar` [60], which identifies halos and subhalos based on clustering of N -body particles in position and velocity space. Then they constructed the catalog of central halos at each output. In this step, halo mass is defined using the spherical overdensity with respect to the halo center (defined as the position with the maximum mass density): $M \equiv M_{200m} = (4\pi/3)R_{200m}^3 \times (200\bar{\rho}_{m0})$, where R_{200m} is the spherical halo boundary radius within which the mean mass density is 200 times $\bar{\rho}_{m0}$, where $\bar{\rho}_{m0}$ is the present-day mean matter density. By combining the outputs of N -body simulations and the halo catalogs at multiple redshifts in the range $z = [0, 1.48]$, they built an emulator, dubbed `Dark Emulator`, which enables fast and accurate computations of the following quantities:

- $\frac{dn_h}{dM}(M; z, \mathbf{p})$: the halo mass function for halos in the mass range $[M, M + dM]$,
- $\xi_{hm}(r; M, z, \mathbf{p})$: the halo-matter cross-correlation function for a sample of halos in the mass range $[M, M + dM]$, and
- $\xi_{hh}(r; M, M', z, \mathbf{p})$: the halo-halo auto-correlation function for two samples of halos with masses $[M, M + dM]$ and $[M', M' + dM']$, respectively,

for an input set of parameters, halo mass M (and M' for the cross-correlation function between two halo samples), redshift z , and cosmological parameters \mathbf{p} .

Fig. 2 of More *et al.* [23] showed that the LOWZ, CMASS1 and CMASS2 galaxies in our samples likely reside on host halos with typical masses greater than $10^{13}h^{-1}M_\odot$, assuming a concordance flat Λ CDM model consistent with the CMB and current large-scale structure data. Hence `Dark Emulator` can be safely used to compute the model predictions of $\Delta\Sigma$ and w_p for these SDSS galaxies.

In addition, `Dark Emulator` outputs ancillary quantities, such as the linear halo bias (the large-scale limit of the halo bias), the Tinker model of the linear halo bias [61] (see below), the linear matter power spectrum, the linear rms mass fluctuations of halo mass scale M ($\sigma_m^L(M)$), and σ_8 .

The supported range of each cosmological parameter for `Dark Emulator` is given in Table I. These ranges are sufficiently broad that they cover the range of cosmological constraints from current state-of-the-art large-scale structure probes such as the Subaru HSC cosmic shear results [4, 5]. Since σ_8 and Ω_m are the primary parameters to which large-scale structure probes are sensitive, we also quote the supported ranges of these *derived* parameters: $0.55 \lesssim \sigma_8 \lesssim 1.2$ and $0.17 \lesssim \Omega_m \lesssim 0.45$, as shown in Fig. 2 of Nishimichi *et al.* [20]. In this paper we use `Dark Emulator` to perform cosmological parameter inference in a multi-dimensional parameter space by comparing the model templates of $\Delta\Sigma$ and w_p with the signals measured from the SDSS and HSC-Y3 data.

The Bayesian parameter inference method we use might occasionally sample models that are outside the supported range of Λ CDM models in `Dark Emulator`. In this case, we

make the following simple extrapolation of the model predictions:

$$\begin{aligned} \xi_{hm}(r; \mathbf{p}_\notin) &\rightarrow \frac{b^{\text{Tinker}}(\mathbf{p}_\notin)}{b^{\text{Tinker}}(\mathbf{p}_{\text{edge}})} \frac{\xi_{\text{mm}}^L(r; \mathbf{p}_\notin)}{\xi_{\text{mm}}^L(r; \mathbf{p}_{\text{edge}})} \xi_{\text{hm}}^{\text{DE}}(r; \mathbf{p}_{\text{edge}}), \\ \xi_{hh}(r; \mathbf{p}_\notin) &\rightarrow \left(\frac{b^{\text{Tinker}}(\mathbf{p}_\notin)}{b^{\text{Tinker}}(\mathbf{p}_{\text{edge}})} \right)^2 \\ &\quad \times \frac{\xi_{\text{mm}}^L(r; \mathbf{p}_\notin)}{\xi_{\text{mm}}^L(r; \mathbf{p}_{\text{edge}})} \xi_{\text{hh}}^{\text{DE}}(r; \mathbf{p}_{\text{edge}}), \end{aligned} \quad (3)$$

where \mathbf{p}_\notin is a set of six cosmological parameters at a Λ CDM model that is outside the supported range (Table I), \mathbf{p}_{edge} is a set of parameters at the edge Λ CDM model inside the supported range, $b^{\text{Tinker}}(\mathbf{p}_\notin)$ and $b^{\text{Tinker}}(\mathbf{p}_{\text{edge}})$ are the linear bias parameters at the two Λ CDM models with \mathbf{p}_\notin and \mathbf{p}_{edge} that are computed based on the fitting formula of Tinker *et al.* [61], ξ_{mm}^L is the linear-theory prediction for the matter two-point correlation function at the respective model, and $\xi_{\text{hh}}^{\text{DE}}$ and $\xi_{\text{hm}}^{\text{DE}}$ are the `Dark Emulator` outputs at the edge model. Here we use an emulator built on `CLASS` [62, 63] to compute the linear-theory matter correlation, $\xi_{\text{mm}}^L(r)$, for models outside the supported range [see Appendix A in Ref. 20, for details]. We define \mathbf{p}_{edge} by replacing only the parameter(s) outside the supported range with their value(s) at the edge of the supported range, while keeping the other parameter(s) at their input value(s). In the above extrapolation, we simply assume that the halo-matter cross-correlation and the halo auto-correlation follow the linear theory predictions ($\xi_{hm} \simeq b\xi_{\text{mm}}$ and $\xi_{hh} \simeq b^2\xi_{\text{mm}}$), and that the ratio of $\xi_{\text{hh}}^L(\mathbf{p}_\notin)$ and $\xi_{\text{hh}}(\mathbf{p}_\notin)$ can be accurately captured by a similar ratio between $\xi_{\text{hh}}^L(\mathbf{p}_{\text{edge}})$ and $\xi_{\text{hh}}(\mathbf{p}_{\text{edge}})$. For the extrapolation we can adopt any input value for A_s , but need to adopt values in the specific ranges for ω_c and Ω_{de} , as we will explain in Table II in Section III E.

Our code outputs the model predictions regardless of whether the cosmological parameters are inside or outside the supported range. This treatment is important, because we perform a blinded cosmological analysis of the HSC and SDSS data. If `Dark Emulator` were to return an error message indicating that an outside model has been sampled, we could unintentionally and prematurely unblind our analysis.

After unblinding our cosmology analysis, we confirmed that all models within the 95% credible interval of S_8 in the chains for our baseline analysis are within the emulator supported range for $\ln(10^{10}A_s)$ and Ω_{de} ⁷, the most important parameters that are sensitive to S_8 ⁸.

⁷ This means that models outside of the 95% credible interval of S_8 are occasionally computed with the extrapolation.

⁸ For this discussion, we used the chains for the models that have ω_c within a $\pm 5\sigma$ range of the *Planck* constraint, because ω_c is not well-constrained by the observables used in this paper.

2. Galaxy-galaxy weak lensing: $\Delta\Sigma(R)$

The details of the galaxy-galaxy weak lensing measurements are presented in More *et al.* [23], but in this section we briefly review the measurement method that we will later use to introduce a residual photo- z error parameter. In particular, we will give the concept of our method to calibrate a residual systematic error in the mean source redshift that affects the weak lensing observables.

Cross-correlating the positions of *spectroscopic* SDSS galaxies with shapes of background *photometric* HSC galaxies enables us to probe the average mass distribution around the lens SDSS galaxies – galaxy-galaxy weak lensing [64]. Throughout this paper we use the average excess surface mass density profile, $\Delta\Sigma(R)$, as the galaxy-galaxy weak lensing observable, where $\Delta\Sigma$ has units of [$hM_\odot \text{pc}^{-2}$] and is given as a function of the projected comoving separation R with units of [$h^{-1}\text{Mpc}$]. An estimator of $\Delta\Sigma(R_i)$ for the i -th radial bin R_i is given [e.g. see Ref. 43], roughly by the following form:

$$\widehat{\Delta\Sigma}(R_i) \simeq \frac{1}{2\mathcal{R} \sum_{\text{ls}} w_{\text{ls}}} \sum_{\text{ls} \in R_i} w_{\text{ls}} \langle \Sigma_{\text{cr}}^{-1} \rangle_{\text{ls}}^{-1} \epsilon_{t,\text{ls}} \Big|_{R_i = \chi(z_1) \Delta\theta_{\text{ls}}}, \quad (4)$$

where the summation “ls” runs over all lens-source pairs that lie in the i -th radial bin $R_i \equiv \chi(z_1) \Delta\theta_{\text{ls}}$, $\chi(z_1)$ is the comoving angular diameter distance to the l -th SDSS lens galaxy at the spectroscopic redshift z_1 , and $\Delta\theta_{\text{ls}}$ is the angular separation between the lens and source in each pair; $\epsilon_{t,\text{ls}}$ is the tangential component of ellipticity of the s -th HSC source galaxy⁹; \mathcal{R} is the shear responsivity [64, 65] which accounts for conversion of “distortion” ($[a^2 - b^2]/[a^2 + b^2]$) to “shear” ($[a - b]/[a + b]$); w_{ls} is the weight, for which we employ an inverse variance weighting that is nearly optimal in the shape-noise dominated regime, following [66] [also see 67]. Additionally, we need to subtract the lensing signal around random points, correct for the additive and multiplicative shear calibration factors [31], and correct for the multiplicative and additive selection bias. Details of the estimator which we used for actual measurements can be found in More *et al.* [23].

The measured weak lensing signal in Eq. (4) depends on the *true* redshift distribution of source galaxies. Hence, to obtain an *unbiased* estimate of $\Delta\Sigma$ for the lens sample, we need the average surface mass density to convert shear to $\Delta\Sigma$, in the *ensemble average sense*:

$$\begin{aligned} \langle \Sigma_{\text{cr}}^{-1} \rangle_{\text{ls}} &= \frac{\int_0^\infty dz_s p_s(z_s) \Sigma_{\text{cr}}^{-1}(z_l, z_s)}{\int_0^\infty dz_s p_s(z_s)} \\ &= 4\pi G(1 + z_1) \chi(z_1) \left[1 - \chi(z_1) \left\langle \frac{1}{\chi(z_s)} \right\rangle_{p_s(z_s)} \right], \quad (5) \end{aligned}$$

⁹ Here we denote the dependence of each lens-source pair, i.e. “ls” in the subscript, because the tangential shear component of the s -th HSC source galaxy shape is defined with respect to the line connecting the source and lens galaxies on the sky.

for a flat-geometry universe, where $p_s(z_s)$ is the *true* redshift distribution of source galaxies in the sample. The factor $(1 + z_1)$ arises from our use of comoving coordinates in the projected separation and we set $\Sigma_{\text{cr}}^{-1} = 0$ when $z_s < z_1$ in Eq. (5). In the second equality on the r.h.s. of Eq. (5), we explicitly show that the dependence of source galaxy redshifts enters only into the average of the inverse of the comoving angular diameter distances to source galaxies over the true redshift distribution of source galaxies: $\langle 1/\chi(z_s) \rangle_{p_s(z_s)}$. On the other hand, there is no uncertainty in the dependence of lens redshifts on an evaluation of $\langle \Sigma_{\text{cr}}^{-1} \rangle$, because we use the spectroscopic galaxy subsamples (LOWZ, CMASS1 and CMASS2) as the lens sample. Hence we stress that, as long as a correct value of $\langle 1/\chi(z_s) \rangle_{p_s(z_s)}$ is evaluated, the shape of the redshift distribution of source galaxies, such as a high-redshift tail or an outlier redshift population of source galaxies, does not cause a bias in $\Delta\Sigma$.

However, the true redshift of individual sources is not available, and we have to use photo- z estimates. To estimate $\widehat{\Delta\Sigma}(R)$ in Eq. (4), we use, in our baseline method, the posterior distribution of photo- z 's for source galaxies to compute $\langle \Sigma_{\text{cr}}^{-1} \rangle$, where the photo- z distribution is generally different from the true redshift distribution $p_s(z_s)$, even in the average sense. We will later introduce a nuisance parameter to model the effect of residual systematic error in the mean source redshift, or equivalently a residual error in the estimate of $\langle \Sigma_{\text{cr}}^{-1} \rangle$. The nuisance parameter can be calibrated from the relative amplitudes between $\Delta\Sigma$ signals for the three lens subsamples and the cosmic shear signals, because the average $\langle \Sigma_{\text{cr}}^{-1} \rangle$ has characteristic dependences on the lens redshifts, as proposed in Oguri and Takada [14]. More exactly speaking, we will implement the self-calibration method along with the estimator used in the measurement, properly taking into account the weight (w_{ls}) for each lens-source pair (see around Eq. 15 in More *et al.* [23]), as we will explain below in detail.

As seen in Eq. (4), the estimation of $\Delta\Sigma(R)$ involves conversion of the observed angular separation between source and lens, $\Delta\theta$, to the comoving separation R and the multiplicative factor of $\langle \Sigma_{\text{cr}}^{-1} \rangle_{\text{ls}}$. To do this, we need to assume a “reference” cosmology, which generally differs from the underlying true cosmology. In Section III B 2 we will describe how to include the effect of varying cosmological models on parameter inference.

Given an unbiased estimate of $\Delta\Sigma$ for a lens sample, we need the theoretical template in cosmology inference. We employ the following two-component model for $\Delta\Sigma(R)$:

$$\Delta\Sigma(R; z_1) = \Delta\Sigma_{\text{gG}}(R; z_1) + \Delta\Sigma_{\text{mag}}(R; z_1). \quad (6)$$

The first term on the right-hand side is the standard contribution to the galaxy-galaxy weak lensing signal, which we refer to as the cross-correlation of the lens galaxies (“g”) and gravitational-lens (“G”) inferred mass in the large scale structure containing the lens sample. Note that $\Delta\Sigma_{\text{gG}}$ is the *standard* excess surface mass density profile of lens galaxies, used in galaxy-galaxy weak lensing. The second term is the contribution caused by the lensing magnification effect, which arises from correlations between shapes of source galaxies and the mass distribution in the foreground structures of lens

galaxies along the line-of-sight to source galaxies due to the fact that lens galaxies can preferentially reside in overdensity regions [68]. Below we describe our models for each contribution within the Λ CDM model framework. Throughout this paper, we model the clustering observables of each SDSS galaxy sample using the theoretical model prediction at a representative redshift, denoted as z_1 : $z_1 \simeq 0.26, 0.51$ and 0.63 for the LOWZ, CMASS1 and CMASS2 samples, respectively. That is, we ignore the possible redshift evolution of the clustering observables within each redshift bin for simplicity. In More *et al.* [23], we confirm that this is a reasonable approximation by looking at variations in the measured clustering and lensing signals within each redshift bin for each of the three subsamples.

The excess surface mass density profile $\Delta\Sigma$ for a given sample of lens galaxies is expressed as [e.g. 66, 69]:

$$\Delta\Sigma_{\text{gG}}(R; z_1) = \bar{\rho}_{\text{m}0} \int \frac{k dk}{2\pi} P_{\text{gm}}(k; z_1) J_2(kR), \quad (7)$$

where $J_2(x)$ is the second-order Bessel function and $P_{\text{gm}}(k; z_1)$ is the cross-power spectrum between galaxies and matter at redshift z_1 . Hereafter we omit z_1 in the argument for notational simplicity.

As described above, `Dark Emulator` outputs halo clustering properties for an input cosmology. To obtain the model predictions for the observable quantities for SDSS galaxies, we need a model for the galaxy-halo connection. For this, we use the halo occupation distribution [HOD 70, 71] [also see 6, 15]. Under this setup we can compute $P_{\text{gm}}(k)$ for a given model:

$$P_{\text{gm}}(k) = \frac{1}{\bar{n}_{\text{g}}} \int dM \frac{dn_{\text{h}}}{dM} \langle N_{\text{c}} \rangle(M) [1 + \lambda_{\text{s}}(M) \tilde{u}_{\text{s}}(k; M, z)] \times P_{\text{hm}}(k; M), \quad (8)$$

where the mean number density of galaxies is given by

$$\bar{n}_{\text{g}} = \int dM \frac{dn_{\text{h}}}{dM} \langle N_{\text{c}} \rangle(M) [1 + \lambda_{\text{s}}(M)], \quad (9)$$

$\langle N_{\text{c}} \rangle(M)$ is the HOD of central galaxies, $\langle N_{\text{c}} \rangle(M) \lambda_{\text{s}}(M)$ is the HOD of satellite galaxies, and $\tilde{u}_{\text{s}}(k; M)$ is the Fourier transform of the average radial profile of satellite galaxies in a host halo with mass M . All the quantities are evaluated at a representative redshift z_1 of the lens galaxies in the LOWZ, CMASS1 or CMASS2 subsample. The impact of using representative redshifts instead of integrating over the lens redshift range is less than $\sim 6\%$ of the square root of the diagonal element of covariance, according to the discussion in Sugiyama *et al.* [7]. Here we use `Dark Emulator` to compute the halo mass function dn_{h}/dM and the halo-matter cross power spectrum, $P_{\text{hm}}(k; M)$, for an input cosmological model, where $P_{\text{hm}}(k; M)$ is obtained from the Fourier transform of the `Dark Emulator` output, $\xi_{\text{hm}}(r; M)$.

We employ the following models for the central and satel-

lite HODs in our baseline analysis:

$$\begin{aligned} \langle N_{\text{c}} \rangle(M) &= \frac{1}{2} \left[1 + \text{erf} \left(\frac{\log M - \log M_{\text{min}}}{\sigma_{\log M}} \right) \right], \\ \langle N_{\text{s}} \rangle(M) &\equiv \langle N_{\text{c}} \rangle(M) \lambda_{\text{s}}(M) \\ &= \langle N_{\text{c}} \rangle(M) \left(\frac{M - \kappa M_{\text{min}}}{M_1} \right)^\alpha, \end{aligned} \quad (10)$$

where $\text{erf}(x)$ is the error function. For our fiducial prescription, we assume that satellite galaxies reside only in a halo that already hosts a central galaxy. Our fiducial HOD model is specified by the five parameters $\{M_{\text{min}}, \sigma_{\log M}, \kappa, M_1, \alpha\}$.

For $\tilde{u}_{\text{s}}(k; M)$ in Eq. (8), throughout this paper, we assume that satellite galaxies follow a Navarro-Frenk-White (NFW) profile [72]. To compute the NFW profile as a function of halo mass and redshift for a given cosmological model, we use the halo mass-concentration relation computed using the publicly-available code `Colossus`¹⁰ [73].

For an extended cosmological analysis, we include parameters to model the effect of off-centered ‘‘central’’ galaxies or the ‘‘incompleteness’’ of central galaxies [69, 74], where the incompleteness effect models the possibility that some massive halos might not host a central galaxy in the sample due to color and magnitude cuts. We use the model in Miyatake *et al.* [15] to model the effects.

We model the second term in Eq. (6), following the method in Ref. [68] [also see Eq. 4 in Ref. 7], as

$$\begin{aligned} \Delta\Sigma_{\text{mag}}(R) &= 2(\alpha_{\text{mag}} - 1) \int_0^{z_{\text{max}}} dz_1 p_1(z_1) \int_0^{z_{\text{max}}} dz_s p_s(z_s) \\ &\times \int \frac{\ell d\ell}{2\pi} \Sigma_{\text{cr}}(z_1, z_s) C_\kappa(\ell; z_1, z_s) J_2\left(\frac{\ell R}{\chi}\right), \end{aligned} \quad (11)$$

where $C_\kappa(\ell)$ is the cosmic shear convergence power spectrum for source galaxies at redshifts z_1 and z_s , defined as

$$\begin{aligned} C_\kappa(\ell; z_1, z_s) &\equiv \int_0^{\chi_{\text{H}}} d\chi \frac{W(\chi, \chi_1) W(\chi, \chi_s)}{\chi^2} \\ &\times P_{\text{mm}}^{\text{NL}}\left(k = \frac{\ell + 1/2}{\chi}; z\right), \end{aligned} \quad (12)$$

with the lensing efficiency function, $W(\chi, \chi_s)$, for lens and source at distances χ and χ_s :

$$W(\chi, \chi_s) \equiv \frac{3\Omega_{\text{m}}}{2} H_0^2 (1+z) \chi \left(1 - \frac{\chi}{\chi_s} \right). \quad (13)$$

Here we used the relation between redshift and comoving distance, via relations $\chi = \chi(z)$, for a given cosmological model; $p_1(z_1)$ in Eq. (11) denotes the redshift distribution of lens galaxies (LOWZ, CMASS1 or CMASS2), normalized as $\int_0^{z_{\text{max}}} dz_1 p_1(z_1) = 1$; α_{mag} is the power-law slope of number counts of the lens galaxies around a magnitude cut in each

¹⁰ <http://www.benediktdiemer.com/code/colossus/>

sample [see Eq. 10 and Fig. 2 in Ref. 6, for the estimated value and error]; $P_{\text{mm}}^{\text{NL}}(k)$ is the nonlinear matter power spectrum for which we use `halofit` [75] for a given cosmological model. Note that $\Delta\Sigma_{\text{mag}}$ does not depend on the models for galaxy bias or galaxy-halo connection. In Eq. (11) we take into account the redshift distribution of both the lens (SDSS) and source (HSC) galaxies, which is different from our treatment in the HSC-Y1 cosmology analyses [6, 7]. As shown in Miyatake *et al.* [6], $\Delta\Sigma_{\text{mag}}$ leads to about 1%, 7% and 10% contributions to the total power of $\Delta\Sigma$ for the LOWZ, CMASS1 and CMASS2 subsamples, respectively, for the *Planck* cosmology [76]. Including the $\Delta\Sigma_{\text{mag}}$ contribution in the theoretical template adds some cosmological information. In our analysis we treat the magnitude slope α_{mag} as a nuisance parameter, with a Gaussian prior with width $\sigma(\alpha_{\text{mag}}) = 0.5$ around the central value taken from the measurement value (see Fig. 2 of Ref. [6]). Note that α_{mag} is different from α , which is a parameter of the satellite HOD.

3. Projected auto-correlation function: $w_{\text{p}}(R)$

As a second clustering observable of the LOWZ, CMASS1 and CMASS2 galaxy subsamples used in the galaxy-galaxy weak lensing measurements, we use the projected spatial correlation function, denoted as $w_{\text{p}}(R)$. We model $w_{\text{p}}(R)$ as

$$w_{\text{p}}(R; z_1) \equiv 2f_{\text{corr}}^{\text{RSD}}(R; z_1) \int_0^{\Pi_{\text{max}}} d\Pi \xi_{\text{gg}}(\sqrt{R^2 + \Pi^2}; z_1), \quad (14)$$

where we take $\Pi_{\text{max}} = 100 h^{-1} \text{Mpc}$ as our fiducial choice and $\xi_{\text{gg}}(r)$ is the real-space, three-dimensional correlation function of galaxies. To compute the radial and projected separations, Π and R , between galaxies in each pair from their observed redshifts and angular positions, we assume the reference cosmological model as done in our $\Delta\Sigma$ analysis above; the flat-geometry model with $\Omega_{\text{m}}^{\text{ref}} = 0.279$. The prefactor $f_{\text{corr}}^{\text{RSD}}(R)$ is a correction factor that accounts for the effect of redshift-space distortion (RSD); we assume the linear Kaiser RSD [13] to compute $f_{\text{corr}}^{\text{RSD}}$ following the method in van den Bosch *et al.* [77] (see Eq. 48 in the paper) [also see Ref. 15].

To use Eq. (14), we must first compute the three-dimensional correlation function of galaxies for a given set of model parameters. The three-dimensional correlation function ξ_{gg} is given as

$$\xi_{\text{gg}}(r; z_1) = \int_0^\infty \frac{k^2 dk}{2\pi^2} P_{\text{gg}}(k; z_1) j_0(kr), \quad (15)$$

where $j_0(x)$ is the zero-th order spherical Bessel function, and $P_{\text{gg}}(k)$ is the auto-power spectrum of galaxies. Once the power spectrum $P_{\text{gg}}(k)$ is given for an input of model parameters, we can compute the model prediction of $w_{\text{p}}(R)$ according to Eq. (14).

In the halo model, P_{gg} can be divided into two contributions, i.e., the 1- and 2-halo terms, as

$$P_{\text{gg}}(k) = P_{\text{gg}}^{\text{1h}}(k) + P_{\text{gg}}^{\text{2h}}(k), \quad (16)$$

where the 1-halo term describes correlations between galaxies within the same host halo, and the 2-halo term describes correlations between galaxies residing in different halos. In our method, we compute the auto-power spectrum as

$$\begin{aligned} P_{\text{gg}}^{\text{1h}}(k) &= \frac{1}{\bar{n}_{\text{g}}^2} \int dM \frac{dn_{\text{h}}}{dM} \langle N_{\text{c}} \rangle (M) [2\lambda_{\text{s}}(M) \tilde{u}_{\text{s}}(k; M) + \lambda_{\text{s}}(M)^2 \tilde{u}_{\text{s}}(k; M)^2], \\ P_{\text{gg}}^{\text{2h}}(k) &= \frac{1}{\bar{n}_{\text{g}}^2} \left[\int dM \frac{dn_{\text{h}}}{dM} \langle N_{\text{c}} \rangle (M) \{1 + \lambda_{\text{s}}(M) \tilde{u}_{\text{s}}(k; M)\} \right] \\ &\quad \times \left[\int dM' \frac{dn_{\text{h}}}{dM'} \langle N_{\text{c}} \rangle (M') \{1 + \lambda_{\text{s}}(M') \tilde{u}_{\text{s}}(k; M')\} \right] P_{\text{hh}}(k; M, M'). \end{aligned} \quad (17)$$

Here we use `Dark Emulator` to compute dn_{h}/dM and $P_{\text{hh}}(k; M, M')$, the power spectrum between halos with masses M and M' for an input cosmological model. Note that in our fiducial model we assume that satellite galaxies reside in halos that host a central galaxy in our sample. Miyatake *et al.* [15] confirmed that fitting the model to mock observables computed for the case that satellite galaxies are populated in halos irrespective of whether the halos host central galaxies in the sample resulted in a negligible shift in S_8 , for our baseline analysis setup.

4. Cosmic shear correlation functions: $\xi_{\pm}(\vartheta)$

As the third clustering observable in our cosmology analysis, we use the measured two-point correlation functions of galaxy shapes in the HSC source sample used in the $\Delta\Sigma$ measurement, denoted as $\xi_{\pm}(\vartheta)$. We model $\xi_{\pm}(\vartheta)$ as a sum of the following three contributions, taking into account contamination of intrinsic alignments (IA):

$$\xi_{\pm}(\vartheta) = \xi_{\text{GG},\pm}(\vartheta) + \xi_{\text{GI},\pm}(\vartheta) + \xi_{\text{II},\pm}(\vartheta). \quad (18)$$

The “+” and “−” correlation functions are measured from different combinations of the correlations of the two ellipticity components of source galaxy shapes in each pair, $\xi_{\pm} \leftarrow$

$(\epsilon_+ \epsilon_+ \pm \epsilon_\times \epsilon_\times)$, where ϵ_+ is the ellipticity component along the R.A. or Dec coordinate direction, and ϵ_\times is its 45-degree rotated component. The first term is the ‘‘gravitational-gravitational’’ term (i.e. cosmic shear, ‘‘GG’’), the third term is the ‘‘intrinsic-intrinsic’’ (‘‘II’’) IA contribution [78–81], and the second term is the ‘‘gravitational-intrinsic’’ correlation (‘‘GI’’) [82] that arises in pairs of galaxies for which common large-scale structure in the line of sight affects the intrinsic shapes of one of the galaxies and the gravitational lensing shear on the other.

The GG term in Eq. (18) is given in terms of the cosmic shear convergence power spectrum, $C_\kappa(\ell)$, as

$$\xi_{\text{GG},\pm}(\vartheta) \equiv \int \frac{\ell d\ell}{2\pi} C_\kappa(\ell) J_{0,4}(\ell\vartheta), \quad (19)$$

where $J_{0,4}(x)$ is the zero-th order (for ξ_+) or fourth-th order (for ξ_-) Bessel function. Using the flat-sky approximation and Limber’s approximation [83], $C_\kappa(\ell)$ is computed from the line-of-sight integral of the nonlinear matter power spectrum as

$$C_\kappa(\ell) = \int_0^{\chi_H} d\chi \frac{q(\chi)^2}{\chi} P_{\text{mm}}^{\text{NL}}\left(k = \frac{\ell + 1/2}{\chi}, z\right), \quad (20)$$

where χ_H is the comoving horizon radius, and z is given by the inverse of $\chi = \chi(z)$. To model $P_{\text{mm}}^{\text{NL}}$ for a given cosmological model, we employ `halofit` [75] in the same way as used in $\Delta\Sigma_{\text{mag}}$ in Eq. (6). The lensing efficiency function $q(\chi)$ [also see 84] is defined as

$$q(\chi) \equiv \int_{z=z(\chi)}^{z_{\text{max}}} dz_s p_s(z_s) W(\chi, \chi_s). \quad (21)$$

where $W(\chi, \chi_s)$ is defined by Eq. (13). We note that we use the same redshift distribution of source galaxies, $p_s(z_s)$, as used in the $\Delta\Sigma$ measurement. Adding the cosmic shear information in parameter inference further helps the self-calibration of the residual photo- z errors of the HSC source galaxies, as we will show later.

To model the IA correlation functions, in this paper we adopt the NLA model [85] in our baseline model. In this model, the II and GI correlation functions are given by

$$\xi_{\text{II/GI},\pm}(\vartheta) = \int \frac{\ell d\ell}{2\pi} C_{\text{II/GI}}(\ell) J_{0,4}(\ell\vartheta), \quad (22)$$

with

$$\begin{aligned} C_{\text{II}}(\ell) &= \int_0^{\chi_H} d\chi F^2(\chi) \frac{p_s(\chi)p_s(\chi)}{\chi^2} P_{\text{mm}}^{\text{NL}}\left(k = \frac{\ell + 1/2}{\chi}, z\right), \\ C_{\text{GI}}(\ell) &= 2 \int_0^{\chi_H} d\chi F(\chi) \frac{q(\chi)p_s(\chi)}{\chi^2} P_{\text{mm}}^{\text{NL}}\left(k = \frac{\ell + 1/2}{\chi}, z\right). \end{aligned} \quad (23)$$

Here the redshift- and cosmology-dependent factor, $F(\chi)$, relating the galaxy ellipticity and the gravitational tidal field is parametrized as

$$F(z) = -A_{\text{IA}} C_1 \rho_{\text{cr},0} \frac{\Omega_m}{D(z)} \left(\frac{1+z}{1+z_0}\right)^{\eta_{\text{IA}}}, \quad (24)$$

where A_{IA} is a dimensionless amplitude parameter, $\rho_{\text{cr},0}$ is the critical density of the Universe at $z = 0$, and $D(z)$ is the linear growth factor normalized to unity at $z = 0$. The additional redshift dependence is assumed to have a power-law form, with power-law index parameter η_{IA} . We use a single parameter model of IA by fixing $\eta_{\text{IA}} = 0$ as our cosmic shear data is only for a single redshift bin, that is, does not contain the redshift information of the IA effect. The normalization constant factor C_1 is set to $5 \times 10^{-14} h^{-2} M_\odot^{-1} \text{Mpc}^3$ at $z_0 = 0.62$ by convention, which is motivated by the observed ellipticity variance in SuperCOSMOS [86]. This model has previously been used in cosmic shear cosmology analyses [4, 87]. While this is merely a phenomenological model of the IA effect, our cosmological constraints are from the joint information of w_p , $\Delta\Sigma$ and ξ_\pm , we expect its effect is small. Indeed, we will find that the cosmological constraints are changed very little even if we ignore the IA contamination in the model template.

We note that galaxy-galaxy weak lensing is not affected by IA contamination, as long as the redshifts of source galaxies do not overlap with those of lens galaxies, which we believe is the case for our source galaxy selection.

B. Modeling Residual Systematic Errors

In this section, we present a method to account for the effects of residual systematic errors on our cosmology analysis. In what follows, we include the systematic effects in the theoretical templates rather than in the signals to keep the data vector and the covariance matrix invariant.

1. Residual systematic photo- z uncertainty: Δz_{ph}

Photo- z uncertainty is one of the most important systematic effects in weak lensing measurements, i.e. $\Delta\Sigma$ and $\xi_\pm(\vartheta)$ in our data vector. As detailed in More *et al.* [23], the redshift distribution of HSC source galaxies was inferred by combining the individual photo- z posteriors with the cross-correlation clustering measurement of HSC galaxies with the CAMIRA sample of Luminous Red Galaxies (LRGs) that have accurate photo- z estimates (typically a few per cent in $\sigma(z)/(1+z)$), based on the method in Rau *et al.* [88]. However, we were not able to fully calibrate the redshift distribution due to the lack of a calibration sample of CAMIRA LRGs at $z \gtrsim 1$ (more exactly speaking, the photo- z accuracies of LRGs at $1 \lesssim z \lesssim 1.2$ are degraded, and there are no LRGs available at $z \gtrsim 1.2$). Hence we take into account the possibility that there is an unknown residual systematic error in the mean redshift of source galaxies. To study the impact of such residual photo- z calibrated uncertainty, we introduce a nuisance parameter, denoted as Δz_{ph} , to model a systematic shift in the mean source redshift by shifting the posterior distribution of source redshifts, given as $z^{\text{est}} = z^{\text{true}} + \Delta z_{\text{ph}}$ [6, 14, 89]. That is, we use the shifted photo- z distribution to model the true distribution as

$$p^{\text{true}}(z) = p^{\text{est}}(z + \Delta z_{\text{ph}}). \quad (25)$$

A positive Δz_{ph} corresponds to the true mean redshift being lower than what is inferred from the photo- z posterior, and vice-versa. Note that the discussion around Eq. (5) gives a justification of this shifted model for the galaxy-galaxy weak lensing ($\Delta\Sigma$) and Zhang *et al.* [90] gave a quantitative justification for the HSC-Y3 analyses on the cosmic shear signals.

For $\Delta\Sigma$ (Eq. 6), we first need to recompute the averaged lensing efficiency $\langle \Sigma_{\text{cr}}^{-1} \rangle$ and the weight w_{ls} using the shifted redshift distribution (Eqs. 4 and 5): we define the correction factor as

$$f_{\Delta\Sigma}(\Delta z_{\text{ph}}) \equiv \frac{\sum_{\text{ls}} w_{\text{ls}} \langle \Sigma_{\text{c}}^{-1} \rangle_{\text{ls}}^{\text{true}} / \langle \Sigma_{\text{c}}^{-1} \rangle_{\text{ls}}^{\text{est}}}{\sum_{\text{ls}} w_{\text{ls}}}. \quad (26)$$

We compute the correction factor for each of the three lens subsamples, LOWZ, CMASS1 and CMASS2. In our method, we multiply the correction factor by the model template of $\Delta\Sigma$, rather than varying the signal, as

$$\Delta\Sigma^{\text{corr}}(R, \Delta z_{\text{ph}}; z_1) = f_{\Delta\Sigma}(\Delta z_{\text{ph}}; z_1) \Delta\Sigma(R; z_1). \quad (27)$$

Note that $\Delta\Sigma$ includes both the galaxy-galaxy weak lensing and the magnification term in Eq. (6): $\Delta\Sigma = \Delta\Sigma_{\text{gG}} + \Delta\Sigma_{\text{mag}}$, since the correction factor is an overall factor that is applied to the estimator of $\widehat{\Delta\Sigma}$ (Eq. 4). In the theoretical template, in addition to the overall factor, we properly use the shifted redshift distribution of source galaxies to re-compute the magnification bias term, $\Delta\Sigma_{\text{mag}}$. Also note that the definition of $f_{\Delta\Sigma}$ is the inverse of the similar correction factor f_{ph} used in the HSC-Y1 papers [6, 7].

Similarly, we recompute the model prediction for the cosmic shear correlation functions $\xi_{\pm}(\vartheta)$ using the shifted redshift distribution of source galaxies.

2. Correction for the reference cosmology used in our measurement

In the measurements of w_{p} and $\Delta\Sigma$, we need to assume a ‘‘reference’’ cosmology to convert the angular separation between galaxies in each pair to the projected separation R , and the redshift difference to the radial separation, Π . For $\Delta\Sigma$, we also need the reference cosmology to convert the shear to $\Delta\Sigma$. Throughout our series of papers, the reference cosmology is a flat Λ CDM model with $\Omega_{\text{m}}^{\text{ref}} = 0.279$. However, the reference cosmology generally differs from the true underlying cosmology, and we need to correct for this discrepancy in our cosmology analysis. We denote a cosmology taken in the parameter inference as \mathbb{C} and the reference cosmology as \mathbb{C}^{ref} . The corrections for R and Π are obtained as

$$\begin{aligned} R &= \frac{\chi(z_1; \mathbb{C})}{\chi(z_1; \mathbb{C}^{\text{ref}})} R^{\text{ref}}, \\ \Pi &= \frac{E(z_1; \mathbb{C}^{\text{ref}})}{E(z_1; \mathbb{C})} \Pi^{\text{ref}}, \end{aligned} \quad (28)$$

where $E(z)$ is the normalized, dimension-less Hubble rate at redshift z , defined as $E(z) \equiv H(z)/H_0$. Thus we include the

measurement corrections in the theoretical templates of $\Delta\Sigma$ and w_{p} as

$$\Delta\Sigma^{\text{ref}}(R^{\text{ref}}; \Delta z_{\text{ph}}, \mathbb{C}, z_1) = f_{\Delta\Sigma}(\Delta z_{\text{ph}}; \mathbb{C}, z_1) \Delta\Sigma(R; \mathbb{C}, z_1), \quad (29)$$

$$\begin{aligned} w_{\text{p}}^{\text{ref}}(R^{\text{ref}}; \mathbb{C}, z_1) &= 2f_{\text{corr}}^{\text{RSD}}(R; \mathbb{C}, z_1) \frac{E(z_1; \mathbb{C})}{E(z_1; \mathbb{C}^{\text{ref}})} \\ &\times \int_0^{\Pi_{\text{max}}} d\Pi \xi_{\text{gg}}(\sqrt{R^2 + \Pi^2}; \mathbb{C}, z_1), \end{aligned} \quad (30)$$

where R and Π are given by R^{ref} or Π^{ref} and the cosmological parameters ($\Omega_{\text{m}}^{\text{ref}}$ and Ω_{m} for a flat model) in the \mathbb{C}^{ref} and \mathbb{C} models (Eq. 28). Note that we adopt $\Pi_{\text{max}} = [E(\mathbb{C}^{\text{ref}})/E(\mathbb{C})] \Pi_{\text{max}}^{\text{ref}} = [E(\mathbb{C}^{\text{ref}})/E(\mathbb{C})] \times 100 h^{-1} \text{Mpc}$, as we use the fixed $\Pi_{\text{max}}^{\text{ref}} = 100 h^{-1} \text{Mpc}$ in the measurement. Also note that $\Delta\Sigma(R)$ and $\xi_{\text{gg}}(r)$ on the r.h.s. of the above equations are computed from theory (Dark Emulator in our method) for a given cosmological model (\mathbb{C}). The overall correction factor for $\Delta\Sigma$ is defined as

$$f_{\Delta\Sigma}(\Delta z_{\text{ph}}; \mathbb{C}, z_1) \equiv \frac{\sum_{\text{ls}} w_{\text{ls}} \langle \Sigma_{\text{c}}^{-1} \rangle_{\text{ls}}^{\text{true}, \mathbb{C}} / \langle \Sigma_{\text{c}}^{-1} \rangle_{\text{ls}}^{\text{est}, \mathbb{C}^{\text{ref}}}}{\sum_{\text{ls}} w_{\text{ls}}}. \quad (31)$$

Thus this correction factor accounts for both the effects of residual photo- z errors (Δz_{ph}) and the measurement correction (\mathbb{C}). We evaluate the model templates, $\Delta\Sigma^{\text{ref}}$ and $w_{\text{p}}^{\text{ref}}$, at the discrete sampling points of R^{ref} as used in the data vector in More *et al.* [23].

3. Residual multiplicative shear error

In order to account for possible residual errors in the shape calibration, we introduce a nuisance parameter which quantifies the residual multiplicative bias Δm and shifts the theoretical templates of the lensing observables:

$$\Delta\Sigma^{\text{corr}}(R, \Delta m; z_1) = (1 + \Delta m) \Delta\Sigma(R; z_1), \quad (32)$$

$$\xi_{\pm}^{\text{corr}}(\vartheta, \Delta m) = (1 + \Delta m)^2 \xi_{\pm}(\vartheta). \quad (33)$$

Since we use a single source sample for both the galaxy-galaxy lensing and cosmic shear, we use the same residual multiplicative bias parameter for the theoretical templates of $\Delta\Sigma$ for the three lens subsamples and for ξ_{\pm} . Hence comparing these data vectors allows us to calibrate the Δm parameter, simultaneously with the calibration of the photo- z error parameter Δz_{ph} .

4. PSF systematics

As discussed in the HSC-Y3 shape catalog paper [31] [also see 91], PSF leakage and residual PSF modeling error contaminate the measured cosmic shear correlation functions. Such residual PSF systematic errors could produce artificial two-point correlations and hence bias the cosmic shear measurements. Here we examine the impact of these systematics in

our cosmic shear measurements, following the method used for the Year 1 analyses Hikage *et al.* [4], Hamana *et al.* [5] [also see 92], where we assume that the measured galaxy shapes have an additional additive bias given by

$$\epsilon^{(\text{sys})} = \alpha_{\text{psf}} \epsilon^{\text{P}} + \beta_{\text{psf}} \epsilon^{\text{Q}}. \quad (34)$$

The first term, referred to as PSF leakage, represents a systematic error proportional to the PSF model ellipticity ϵ^{P} due to the imperfect PSF correction. The second term represents the systematic error associated with the difference between the model PSF ellipticity, ϵ^{P} , and the true PSF ellipticity estimated from individual “reserved” stars ϵ^{star} , i.e. $\epsilon^{\text{Q}} \equiv \epsilon^{\text{P}} - \epsilon^{\text{star}}$ [92]. Non-zero residual PSF ellipticities ϵ^{Q} indicate an imperfect PSF estimate, which will propagate to estimates of galaxy shears. Note that the above PSF systematics causes additive shear bias, so does not affect the galaxy-galaxy weak lensing.

When the observed galaxy ellipticity is contaminated by $\epsilon^{(\text{sys})}$, these systematic terms cause an additional contamination to the measured cosmic shear correlation functions as

$$\xi_{\text{psf},\pm}(\vartheta) = \alpha_{\text{psf}}^2 \hat{\xi}_{\pm}^{\text{PP}}(\vartheta) + 2\alpha_{\text{psf}}\beta_{\text{psf}} \hat{\xi}_{\pm}^{\text{PQ}}(\vartheta) + \beta_{\text{psf}}^2 \hat{\xi}_{\pm}^{\text{QQ}}(\vartheta), \quad (35)$$

where $\hat{\xi}_{\pm}^{\text{PP}}$, $\hat{\xi}_{\pm}^{\text{QQ}}$ and $\hat{\xi}_{\pm}^{\text{PQ}}$ represent the auto-correlation of the model PSF ellipticity $\epsilon_{\pm}^{\text{P}}$, the auto-correlation of the residual PSF ellipticity $\epsilon_{\pm}^{\text{Q}}$, and the cross-correlation of $\epsilon_{\pm}^{\text{P}}$ and $\epsilon_{\pm}^{\text{Q}}$, respectively. The hat notation, “ $\hat{}$ ”, denotes the correlation function measured from the HSC data using the model PSF and the reserved stars (see [23]). The coefficients α_{psf} and β_{psf} are estimated by cross-correlating $\epsilon_{\pm}^{\text{P}}$ and $\epsilon_{\pm}^{\text{Q}}$ with the observed galaxy ellipticities, as

$$\begin{aligned} \hat{\xi}_{\pm}^{\text{PQ}}(\vartheta) &= \alpha_{\text{psf}} \hat{\xi}_{\pm}^{\text{PP}}(\vartheta) + \beta_{\text{psf}} \hat{\xi}_{\pm}^{\text{PQ}}(\vartheta), \\ \hat{\xi}_{\pm}^{\text{QQ}}(\vartheta) &= \alpha_{\text{psf}} \hat{\xi}_{\pm}^{\text{PQ}}(\vartheta) + \beta_{\text{psf}} \hat{\xi}_{\pm}^{\text{QQ}}(\vartheta), \end{aligned} \quad (36)$$

where $\hat{\xi}_{\pm}^{\text{PQ}}$ and $\hat{\xi}_{\pm}^{\text{QQ}}$ are the measured cross-correlations between the galaxy ellipticities, used for the cosmic shear data

vector, and $\epsilon_{\pm}^{\text{P}}$ and $\epsilon_{\pm}^{\text{Q}}$. As discussed in More *et al.* [23] (also see Li *et al.* [25], Zhang *et al.* [91]), we used the measurements of mock galaxy shape catalogs and the real star catalog to estimate the statistical errors of the $\hat{\xi}_{\pm}^{\text{GP}}$ and $\hat{\xi}_{\pm}^{\text{GQ}}$ measurements, where the errors take into account the cosmic variance. By comparing the measured $\hat{\xi}_{\pm}^{\text{GP}}$ and $\hat{\xi}_{\pm}^{\text{GQ}}$ with Eq. (36) using the measured $\hat{\xi}_{\pm}^{\text{PP}}$, $\hat{\xi}_{\pm}^{\text{PQ}}$ and $\hat{\xi}_{\pm}^{\text{QQ}}$, we found $\alpha_{\text{psf}} = -0.0292 \pm 0.0129$ and $\beta_{\text{psf}} = -2.59 \pm 1.65$ for our fiducial source catalog (see Fig. 16 of More *et al.* [23]).

To take into account the impact of the residual PSF modeling errors on parameter inference, we add the PSF error contamination $\xi_{\text{psf},\pm}$ (Eq. 35) to the model cosmic correlation function ξ_{\pm} in Eq. (18) and then estimate parameters by varying the parameters α_{psf} and β_{psf} using Gaussian priors with widths inferred from the above errors.

The above method of PSF systematics takes into account the PSF systematics up to the second-order moment of PSF. The HSC-Y3 cosmic shear cosmology papers, Li *et al.* [25] and Dalal *et al.* [26], used the more sophisticated, accurate method developed in Zhang *et al.* [91], which accounts for the effects up to fourth-moment PSF leakage and fourth-moment PSF modeling error on cosmic shear correlations. Using the same method, we also measured up to the α and β coefficients of the fourth-moments of PSF for the HSC source galaxy sample used in this paper. We then generated synthetic cosmic shear data vectors including the measured PSF systematic effects up to the fourth-order moment and checked that the estimated S_8 remains almost unchanged compared to our baseline analysis method using the α and β coefficients of the PSF second-moment with the priors described above. The main reason for this is that most of the constraining power is from the galaxy clustering information of SDSS galaxies. The details are given in Appendix A [also see 24].

C. Summary: Model Templates

For convenience, here we write down the model templates used in cosmology inference where we explicitly show which parameters are used in the templates of each observable:

$$\begin{aligned} \Delta\Sigma^{\text{t}}(R^{\text{ref}}, z_1 | \mathbb{C}, \boldsymbol{\theta}_{\text{g}}, \Delta z_{\text{ph}}, \Delta m, \alpha_{\text{mag}}(z_1)) &= (1 + \Delta m) \Delta\Sigma^{\text{ref}}(R^{\text{ref}}, z_1 | \mathbb{C}, \boldsymbol{\theta}_{\text{g}}, \Delta z_{\text{ph}}, \alpha_{\text{mag}}(z_1)), \\ w_{\text{p}}^{\text{t}}(R^{\text{ref}}, z_1 | \mathbb{C}, \boldsymbol{\theta}_{\text{g}}) &: \text{Eqs. (15)–(17), (30)}, \\ \xi_{\pm}^{\text{t}}(\vartheta | \mathbb{C}, \Delta z_{\text{ph}}, A_{\text{IA}}, \eta_{\text{IA}}, \alpha_{\text{psf}}, \beta_{\text{psf}}) &= (1 + \Delta m)^2 \xi_{\pm}(\vartheta | \mathbb{C}, \Delta z_{\text{ph}}, A_{\text{IA}}, \eta_{\text{IA}}) + \xi_{\text{psf},\pm}(\vartheta | \alpha_{\text{psf}}, \beta_{\text{psf}}), \end{aligned} \quad (37)$$

where \mathbb{C} denotes a cosmological model sampled in parameter inference within the flat-geometry Λ CDM model characterized by five cosmological parameters, $\boldsymbol{\theta}_{\text{g}}$ is a set of parameters to model the galaxy-halo connection (five parameters for each of LOWZ, CMASS1 and CMASS2 in our baseline model), $\Delta\Sigma^{\text{ref}}$ is given by Eq. (29), and others are nuisance parameters to model the residual systematic errors. For our baseline

model, we have 28 parameters in total: 5 (\mathbb{C}) + 3 \times 5 ($\boldsymbol{\theta}_{\text{g}}$) + 8 ($\Delta z_{\text{ph}}, \Delta m, \alpha_{\text{mag}}(z_1), A_{\text{IA}}, \alpha_{\text{psf}}, \beta_{\text{psf}}$).

D. Computation Time

We use `Dark Emulator` to compute the model predictions, $\Delta\Sigma^t(R)$ and $w_p^t(R)$, for an input model. We use the publicly available `FFTLLog` code [93] to perform the Hankel transforms in Eqs. (7), (11), 15, and (19); for our analysis we use the updated code in Fang *et al.* [94]. Since our data vector is given by discrete bins of R^{ref} or ϑ , we properly take into account the weighted average of the model predictions within the bin width, more precisely $\Delta \ln R^{\text{fid}} = 0.246$ for $\Delta\Sigma^t$, $\Delta \ln R^{\text{fid}} = 0.169$ for w_p^t , and $\Delta \ln \vartheta = 0.242$ for $\xi_{\pm}(\vartheta)$, respectively. With our current analysis pipeline, we can compute the model predictions of $\Delta\Sigma^t$ for all three lens samples (LOWZ, CMASS1 and CMASS2) in about 2 CPU seconds in total, those of w_p^t in about 2 seconds total, and those of ξ_{\pm} in about 0.15 seconds for a given model. This is fast enough to enable cosmological parameter inference in a high dimensional parameter space (28 parameters in our baseline setup).

E. Parameter Estimation Method

We assume that the likelihood of data for a given model follows a multivariate Gaussian distribution:

$$\ln \mathcal{L}(\mathbf{d}|\boldsymbol{\theta}) = -\frac{1}{2} \sum_{i,j} [d_i - t_i(\boldsymbol{\theta})] \mathbf{C}_{ij}^{-1} [d_j - t_j(\boldsymbol{\theta})], \quad (38)$$

where \mathbf{d} is the data vector, \mathbf{t} is the model prediction for the data vector given the model parameters ($\boldsymbol{\theta}$), \mathbf{C}^{-1} is the inverse of the covariance matrix, and the summation runs over indices corresponding to the dimension of the data vector. Note that non-Gaussianity in the likelihood might affect our results as indicated by Lin *et al.* [95], although they showed that the non-Gaussianity does not cause a significant bias in the parameter value, and rather changes the size of the confidence region. We will leave this question to future studies. Please see More *et al.* [23] for our method to construct the covariance matrix using the mock catalogs of SDSS and HSC galaxies. In our baseline analysis, the data vector consists of $\Delta\Sigma(R)$ in 9 logarithmically-spaced radial bins within $3 \leq R/[h^{-1}\text{Mpc}] \leq 30$, and $w_p(R)$ in 16 logarithmically-spaced radial bins within $2 \leq R/[h^{-1}\text{Mpc}] \leq 30$, for each galaxy subsample (LOWZ, CMASS1 and CMASS2), and 8 and 7 logarithmically-spaced angular bins within $8 \lesssim \vartheta/[\text{arcmin}] \lesssim 50$ and $30 \lesssim \vartheta/[\text{arcmin}] \lesssim 150$ for ξ_+ and ξ_- , respectively. We therefore use $90(= 3 \times (9 + 16) + 8 + 7)$ data points in total. When we use the data vector in a more limited range of separations, we take the submatrix of the full covariance matrix computed in More *et al.* [23], corresponding to that range of separations, and then invert the matrix to obtain the inverse of the covariance matrix, $[\mathbf{C}_{\text{sub}}]^{-1}$. We also note that our analysis does not include the observed galaxy abundance, \bar{n}_g , as an observable in the likelihood.

Table II summarizes the parameters and the priors used in our analysis. We include five cosmological parameters given by $\mathbb{C} = \{\Omega_{\text{de}}, \ln(10^{10} A_s), \omega_b, \omega_c, n_s\}$ for the flat Λ CDM model and five HOD parameters for each of the LOWZ,

TABLE II. Model parameters and priors used in our cosmological inference. The label $\mathcal{U}(a, b)$ denotes a uniform (or equivalently flat) prior with minimum a and maximum b , while $\mathcal{N}(\mu, \sigma)$ denotes a normal (or Gaussian) prior with mean μ and width σ . The parameters above the horizontal double lines are the parameters used in our baseline analysis: five cosmological parameters, five HOD parameters for each of the LOWZ, CMASS1 and CMASS2 subsamples, two nuisance parameters to model residual photo- z and multiplicative shear biases, three parameters (α_{mag}) to model the magnitude slope of the galaxy number counts that characterizes the magnification bias effect on $\Delta\Sigma$ for each of the LOWZ, CMASS1 and CMASS2 subsamples, two parameters to model residual PSF modeling errors in the cosmic shear 2pt functions, and one parameter to model the IA contamination to cosmic shear: $28 = 5 + 3 \times 5 + 2 + 3 + 2 + 1$ in total. The parameters below the double lines are used in the extended models.

Parameter	Prior
Cosmological parameters	
Ω_{de}	$\mathcal{U}(0.4594, 0.9094)$
$\ln(10^{10} A_s)$	$\mathcal{U}(1.0, 5.0)$
ω_c	$\mathcal{U}(0.0998, 0.1398)$
ω_b	$\mathcal{N}(0.02268, 0.00038)$
n_s	$\mathcal{N}(0.9649, 3 \times 0.0042)$
HOD parameters	
$\log M_{\text{min}}(z_i)$	$\mathcal{U}(12.0, 14.5)$
$\sigma_{\log M}^2(z_i)$	$\mathcal{U}(0.01, 1.0)$
$\log M_1(z_i)$	$\mathcal{U}(12.0, 16.0)$
$\kappa(z_i)$	$\mathcal{U}(0.01, 3.0)$
$\alpha(z_i)$	$\mathcal{U}(0.5, 3.0)$
Magnification bias	
α_{mag} (LOWZ)	$\mathcal{N}(2.26, 0.5)$
α_{mag} (CMASS1)	$\mathcal{N}(3.56, 0.5)$
α_{mag} (CMASS2)	$\mathcal{N}(3.73, 0.5)$
Residual photo-z/Shear errors	
Δz_{ph}	$\mathcal{U}(-1.0, 1.0)$
Δm	$\mathcal{N}(0.0, 0.01)$
Residual PSF modeling errors	
α_{psf}	$\mathcal{N}(-0.03, 0.01)$
β_{psf}	$\mathcal{N}(-1.66, 1.33)$
IA contamination to cosmic shear	
A_{IA}	$\mathcal{U}(-5.0, 5.0)$
Add. galaxy-halo connection paras	
Off-centering parameters	
$p_{\text{off}}(z_i)$	$\mathcal{U}(0, 1)$
$R_{\text{off}}(z_i)$	$\mathcal{U}(0.01, 1)$
Incompleteness parameters	
$\alpha_{\text{incomp}}(z_i)$	$\mathcal{U}(0, 5)$
$\log M_{\text{incomp}}(z_i)$	$\mathcal{U}(12, 15.3)$

CMASS1, and CMASS2 samples. For ω_b , we employ a Gaussian prior with a mean and width inferred from Big Bang nucleosynthesis (BBN) constraints [11, 96–98]. For n_s , we employ a Gaussian prior inferred from the *Planck* 2018 “TT,EE,TE+lowE” constraints [11]: $n_s = 0.9649 \pm (3 \times 0.0042)$, where we employ a Gaussian width three times wider than the 1σ uncertainty (0.0042) of the *Planck* constraint. We employ these priors since the clustering observables $\Delta\Sigma$ and w_p are not sensitive to ω_b and n_s . For Ω_{de} and ω_c , we adopt broad, flat priors in the ranges that correspond to

about $\pm 30\sigma$ and $\pm 15\sigma$, respectively, compared to the 1σ error of the *Planck* constraints for the flat Λ CDM model. These ranges correspond to the supported range of the extrapolation of Dark Emulator (for details see Section III A 1). Since there is no limitation on $\ln(10^{10}A_s)$ in the extrapolation, we employ a broad and uninformative flat prior.

In addition we include $\alpha_{\text{mag}}(z_i)$ to model possible uncertainty in the magnitude slope of the number counts in modeling the magnification for each lens sample: we use the measured value of α_{mag} for the central value (see Section III A and Fig. 2 in Miyatake *et al.* [6]) and employ a Gaussian prior with a width of $\sigma(\alpha_{\text{mag}}) = 0.5$. This is a conservative choice, since the Gaussian width is much wider than the measurement error on α_{mag} , but we show that the results remain almost unchanged even when fixing α_{mag} to the central (measured) value.

We also include nuisance parameters, Δz_{ph} and Δm , to model residual uncertainties in the source photo- z error and the multiplicative shear bias. Since we use a single population of source galaxies, we need to adopt just one Δz_{ph} and one Δm parameter to model the impact on the galaxy-galaxy weak lensing signals for all three lens galaxy samples and the cosmic shear correlation functions. As we described in Section III B 1, Δz_{ph} contains uncertainty, so we adopt the most conservative choice, i.e., a completely uninformative flat, wide prior of Δz_{ph} , given by $\mathcal{U}(-1.0, 1.0)$ in our baseline setup.

We will demonstrate that our method enables a self-calibration of Δz_{ph} . We made this choice while the analysis was still blinded, i.e. before the unblinding. We also consider a Gaussian prior with width $\sigma(\Delta z_{\text{ph}}) = 0.1$ and mean $\Delta z_{\text{ph}} = 0$, as done in the Year 1 analysis [6]. This allows us to study how the cosmological parameter inference is altered by this informative prior. This prior is still wider than the width of a few $O(10^{-2})$ that is inferred from the photo- z method in Rau *et al.* [88]. If we use only a subset of the observables, either only the 2×2 pt or the cosmic shear correlations, the analysis cannot constrain Δz_{ph} . Hence, for analyses aimed at internal consistency tests of the data, we used a Gaussian prior whose width and mean are given by the posterior of Δz_{ph} obtained from the baseline 3×2 pt analysis. This prior is denoted by the superscript mark “*” in Table II and hereafter.

For Δm , we employ a prior range that corresponds to about 1σ statistical uncertainties in the shape measurement calibration [35] [also see Table 6 in Ref. 4]. We will discuss the case where the prior range of Δm is broadened in Section V A.

For α_{psf} and β_{psf} , which model the residual PSF modeling errors, we use the measured values for the central values and employ a Gaussian prior with width given by the 1σ measurement uncertainty for each of α_{psf} and β_{psf} . The details of estimation of these parameters are given in More *et al.* [23].

The parameters we described above are the model parameters used for our baseline analysis. We also employ the extended halo model to check how the cosmological parameters obtained from the baseline analysis are robust against possible variations in the model template. For the extended model, we consider the effects of off-centered central galaxies and the incompleteness of central galaxies [also see 6, 15, for details].

Table II gives the parameters to model these effects.

We then obtain the posterior distribution $\mathcal{P}(\boldsymbol{\theta}|\mathbf{d})$ of our parameters $\boldsymbol{\theta}$ given the data \mathbf{d} , by performing Bayesian inference:

$$\mathcal{P}(\boldsymbol{\theta}|\mathbf{d}) \propto \mathcal{L}(\mathbf{d}|\boldsymbol{\theta})\Pi(\boldsymbol{\theta}), \quad (39)$$

where $\Pi(\boldsymbol{\theta})$ is the prior distribution of the parameters. Throughout this paper we focus on the marginalized posterior distributions of the derived parameters, Ω_m , σ_8 , and $S_8 \equiv \sigma_8(\Omega_m/0.3)^{0.5}$, where $\Omega_m = 1 - \Omega_{\text{de}}$ for a flat cosmological model. While $\ln(10^{10}A_s)$ is sampled in logarithmic space with a flat prior, we account for the Jacobian or weight to effectively produce a flat prior in linear space of σ_8 when obtaining the posterior distribution of σ_8 as a derived parameter (see Section IV A in Sugiyama *et al.* [99] and Section V G in Dalal *et al.* [26] for a detailed discussion). However, the effect is negligible because the Jacobian is nearly constant in the range of our credible interval on σ_8 .

To estimate the posterior distribution of parameters in our multi-dimensional parameter space, we use the importance nested sampling algorithm implemented in the publicly-available software package `MultiNest` [100–102] and its python wrapper, `PyMultiNest` [103]. We use `nlive = 600`, `tol = 0.1` for the hyper parameters of `MultiNest`. However, we found that `MultiNest` tends to underestimate the credible interval, e.g., that of S_8 by $\sim 10\%$. This is because `MultiNest` samples a parameter with a Gaussian prior in Table II, in a limited volume that is specified by another hyper parameter `efr`. We checked that we can avoid this inaccuracy by treating the Gaussian prior as an additive term to the likelihood of our observables, rather than injecting the prior to the prior interface in `MultiNest`. We use this implementation for our baseline 3×2 pt analysis. On the other hand, we use the standard `MultiNest` implementation with `nlive = 600`, `tol = 0.1`, and `efr = 0.5` for model/method validations and internal consistency tests described in Section V B, since we need to save computing time to run chains for each setup listed in Table III. Still, we note that the central value of cosmological parameter is stable for both the implementations (typically only a few percent difference). We describe detailed investigations, such as a convergence of `MultiNest` chains and comparison with the Metropolis algorithm, in Appendix D.

F. Analysis Setups

To perform the cosmological parameter inference, we must specify the analysis setups, including the range of scales and combinations of observables to use. Table III summarizes the setups used in this paper.

The setup labeled “ 3×2 pt” is our baseline setup that serves as a reference; if we identify any internal consistency test that significantly deviates from the cosmological parameters obtained from this reference setup, we should consider changing the baseline setup.

If we use either $\Delta\Sigma$ or w_p alone, the parameter inference suffers from severe degeneracies, especially between the galaxy bias (and therefore the HOD model parameters) and

TABLE III. A summary of the analysis setups. The first column identifies each analysis setup. The scale cuts “ (X, Y) ” denote the lower scale cuts applied to $w_p(R)$ and $\Delta\Sigma(R)$, meaning that we use w_p and $\Delta\Sigma$ for $X \leq R/[h^{-1}\text{Mpc}] \leq 30$ and $Y \leq R/[h^{-1}\text{Mpc}] \leq 30$, respectively, in the cosmology analysis. The symbol * at the end of the setup label indicates an analysis using a prior of photo- z shift parameter derived from the baseline $3 \times 2\text{pt}$ analysis, given by $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(-0.06, 0.08)$, while \dagger denotes the analysis with a Gaussian prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$. $\mathcal{D}(\boldsymbol{\theta})$ and $\mathcal{D}(\mathbf{d})$ denote the dimension of parameters and data vector, respectively, in each analysis.

setup label	description	$\mathcal{D}(\boldsymbol{\theta}), \mathcal{D}(\mathbf{d})$
$3 \times 2\text{pt}$	<i>baseline analysis</i> $\Delta\Sigma + w_p + \xi_{\pm}$, with $(2,3)h^{-1}\text{Mpc}$ scale cuts for w_p and $\Delta\Sigma$	28, 90
$2 \times 2\text{pt}^*$	$2 \times 2\text{pt}$ ($\Delta\Sigma + w_p$), w/o ξ_{\pm} , using Δz_{ph} posterior from $3 \times 2\text{pt}$ analysis as a prior	25, 75
cosmic shear*	ξ_{\pm} alone, using Δz_{ph} posterior from $3 \times 2\text{pt}$ analysis as a prior	10, 15
$3 \times 2\text{pt}, R_{\text{min}} = (4, 6) h^{-1}\text{Mpc}^*$	$3 \times 2\text{pt}$, with the minimum scale cuts $R_{\text{min}} = (4, 6) h^{-1}\text{Mpc}$ for w_p and $\Delta\Sigma$	28, 72
$3 \times 2\text{pt}, R_{\text{min}} = (8, 12) h^{-1}\text{Mpc}^*$	$3 \times 2\text{pt}$, with the minimum scale cuts $R_{\text{min}} = (8, 12) h^{-1}\text{Mpc}$ for w_p and $\Delta\Sigma$	28, 51
$3 \times 2\text{pt}$, w/o LOWZ	$3 \times 2\text{pt}$ w/o LOWZ	22, 65
$3 \times 2\text{pt}$, w/o CMASS1	$3 \times 2\text{pt}$ w/o CMASS1	22, 65
$3 \times 2\text{pt}$, w/o CMASS2	$3 \times 2\text{pt}$ w/o CMASS2	22, 65
$2 \times 2\text{pt}$, w/o LOWZ*	$2 \times 2\text{pt}$ w/o LOWZ	18, 51
$2 \times 2\text{pt}$, w/o CMASS1*	$2 \times 2\text{pt}$ w/o CMASS1	18, 51
$2 \times 2\text{pt}$, w/o CMASS2*	$2 \times 2\text{pt}$ w/o CMASS2	18, 51
no photo- z error	$3 \times 2\text{pt}$, but fixing $\Delta z_{\text{ph}} = 0$	27, 90
no shear error	$3 \times 2\text{pt}$, but fixing $\Delta m = 0$	27, 90
fix mag. bias	$3 \times 2\text{pt}$, but fixing $\alpha_{\text{mag}} = \mu$	25, 90
no PSF error	$3 \times 2\text{pt}$, but fixing $\alpha_{\text{psf}} = \beta_{\text{psf}} = 0$	26, 90
no IA	$3 \times 2\text{pt}$, but fixing $A_{\text{IA}} = 0$	27, 90
extreme IA	$3 \times 2\text{pt}$, but fixing $A_{\text{IA}} = 5$	27, 90
$3 \times 2\text{pt}^\dagger$	$3 \times 2\text{pt}$ with a prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$	28, 90
$2 \times 2\text{pt}^\dagger$	$2 \times 2\text{pt}$ with a prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$	24, 75
cosmic shear †	cosmic shear with a prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$	11, 15
$2 \times 2\text{pt}$	$2 \times 2\text{pt}$ with a prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$	25, 75
cosmic shear	ξ_{\pm} with a prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$	10, 15
XMM ($\sim 33 \text{ deg}^2$)*	$3 \times 2\text{pt}$, but using the signals of the XMM field alone	28, 90
GAMA15H ($\sim 41 \text{ deg}^2$)*	$3 \times 2\text{pt}$, but using the signals of the GAMA15H field alone	28, 90
HECTOMAP ($\sim 43 \text{ deg}^2$)*	$3 \times 2\text{pt}$, but using the signals of the HECTOMAP field alone	28, 90
GAMA09H ($\sim 78 \text{ deg}^2$)*	$3 \times 2\text{pt}$, but using the signals of the GAMA09H field alone	28, 90
VVDS ($\sim 96 \text{ deg}^2$)*	$3 \times 2\text{pt}$, but using the signals of the VVDS field alone	28, 90
WIDE12H ($\sim 121 \text{ deg}^2$)*	$3 \times 2\text{pt}$, but using the signals of the WIDE12H field alone	28, 90
DEMPz & WX	DEMPz & WX is used to infer the source redshift distribution and for $\Delta\Sigma$ measurement.	28, 90
MIZUKI	MIZUKI is used for source sample selection and $\Delta\Sigma$ measurement	28, 90
DNNz	DNNz is used for source sample selection and $\Delta\Sigma$ measurement	28, 90
w/o star weight	$3 \times 2\text{pt}$, but without using star weight when computing w_p	28, 90
offcentering	$3 \times 2\text{pt}$, but including the off-centering effect in galaxy-halo connection	34, 90
incompleteness	$3 \times 2\text{pt}$, but including the incompleteness effect in galaxy-halo connection	34, 90
Additional tests after unblinding		
$\sigma(\Delta m) = 0.1$ prior	$3 \times 2\text{pt}$, but using prior $\Pi(\Delta m) = \mathcal{N}(0, 0.1)$	28, 90
$\sigma(\Delta z_{\text{ph}}) = 0.2$ prior	$3 \times 2\text{pt}$, but using prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.2)$	28, 90
2 cosmo paras	$3 \times 2\text{pt}$, but varying only $(\Omega_{\text{de}}, \ln(10^{10} A_s))$ and fixing (ω_c, ω, n_s) to their <i>Planck</i> values	25, 90

the cosmological parameters that encode information about the power spectrum amplitude, as shown in our validation paper [15] (see Fig. 9). Hence, in the following we show only the results of the joint analysis of $\Delta\Sigma$ and w_p , even when not combined with ξ_{\pm} . The “ $2 \times 2\text{pt}^*$ ” setup is such a case, but we set the prior on the residual photo- z parameter Δz_{ph} to the

one obtained from the baseline $3 \times 2\text{pt}$ analysis because without the prior cosmological constraints are quite weak. We also run the cosmic shear-only analysis with the same prior which

is labeled as “cosmic shear*¹¹”.

One important choice in the analysis is the “scale cuts”, i.e., the range of scales (R) in $\Delta\Sigma(R)$ and $w_p(R)$ used in the cosmological analysis. There are two competing considerations. To increase the statistical constraining power on cosmological parameters, we want to include information from $\Delta\Sigma(R)$ and $w_p(R)$ down to as small separations as possible. However, the observables at such small scales may be more strongly affected by physical systematic effects inherent in galaxy formation/physics, which are difficult to accurately model. As we described in our validation paper [15], the scale cuts of $2 h^{-1}\text{Mpc}$ for w_p and $3 h^{-1}\text{Mpc}$ for $\Delta\Sigma$ (hereafter $(2, 3) h^{-1}\text{Mpc}$ scale cuts) are reasonable choices to obtain unbiased estimates of the cosmological parameters, with reasonably small credible intervals given the statistical power of HSC-Y3 and SDSS. The scale cuts of $(2, 3) h^{-1}\text{Mpc}$ are larger than the virial radii of massive halos, so we do not include information from scales that are deeply in the 1-halo term regime in our cosmology analysis. Nevertheless we note that the galaxy-galaxy weak lensing signal around the scale cut is sensitive to the *interior* mass inside that radius. Thus, we can extract the average mass of halos hosting the SDSS galaxies which in turn helps constrain the large-scale bias of SDSS galaxies via the scaling relation of halo bias with halo mass, encoded in `DARK EMULATOR`, when combined with the measurement of w_p . To study the impact of the scale cut choice, we also study the results for scale cuts of $(4, 6)$ and $(8, 12) h^{-1}\text{Mpc}$, respectively.

For the scale cuts of ξ_{\pm} we follow the same scale cuts in $\xi_{\pm}(\vartheta)$ as those in Li *et al.* [25], because we did not find any significant residual B -mode signal in the range. For this paper, we use the cosmic shear signals in the range of $8 \lesssim \vartheta/\text{arcmin} \lesssim 50$ and $30 \lesssim \vartheta/\text{arcmin} \lesssim 150$ for ξ_+ and ξ_- , respectively.

As an internal consistency test, we perform various analyses to assess the robustness of our results to various splits of our data and the sensitivity of our results to the different analysis choices. We perform the analyses excluding one of the LOWZ, CMASS1 or CMASS2 subsample for both “ $3\times 2\text{pt}$ ” and “ $2\times 2\text{pt}$ ” analysis. To study the impact of the nuisance parameters on our results, we perform the analysis by fixing either one of the nuisance parameters, rather than varying it, in the parameter inference: the residual photo- z error $\Delta z_{\text{ph}} = 0$, the multiplicative error $\Delta m = 0$, the magnification bias parameters α_{mag} to their measured values (see Table II), the PSF systematics parameters $\alpha_{\text{psf}} = \beta_{\text{psf}} = 0$, intrinsic alignment $A_{\text{IA}} = 0$, or $A_{\text{IA}} = 5$, respectively.

To check the impact of the prior choice of the residual photo- z error parameter, which is a key parameter in our analysis, we perform the analysis using a Gaussian prior given by

¹¹ Note that the Gaussian prior employed for internal consistency tests, i.e., $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(-0.06, 0.08)$ is slightly different from what we show as the baseline result, $\Delta z_{\text{ph}} = -0.05 \pm 0.09$, in Section V A. This is because we use the Δz_{ph} constraint from the standard `MULTINEST` implementation for this Gaussian prior rather than the setup used for the baseline $3\times 2\text{pt}$ analysis described in Section III E. This is the case for the Gaussian prior on Δz_{ph} used in Sugiyama *et al.* [24].

$\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$, for the $3\times 2\text{pt}$ analysis or the $2\times 2\text{pt}$ - or the cosmic shear-only analysis. We also perform the analysis using an uninformative flat prior, $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$, for the $2\times 2\text{pt}$ - or the cosmic shear-only analysis.

We also perform field-by-field analyses which are labeled by the field name, e.g. “XMM” in Table III, which is the result when using the HSC-Y3 data only in the XMM region [31]. Note that the signals measured in different fields are almost independent.

To check for possible systematic biases arising from the different photo- z estimate methods, we perform the analyses using the lensing signals that are measured using the different photo- z method instead of our fiducial method (DEMPZ). More exactly, we use the different method to *select* source galaxies based on the same selection cut (Eq. 2) and then use the inferred source redshift distribution in the weak lensing observables (the signal of $\Delta\Sigma$ and the theory of ξ_{\pm}). For “DEMPZ&WX”, we use the source redshift distribution inferred by the “DEMPZ&WX” method that is obtained by combining the DempZ photo- z estimates and the clustering with the CAMIRA LRGs in Rau *et al.* [88]. We also use the MIZUKI photo- z method or the DNNZ photo- z method, respectively. We use the uninformative flat prior, $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$ for all these tests as in our baseline analysis. Note that the DempZ&WX analysis uses the same source sample as that in the baseline analysis, but the other methods use the different source samples.

The analysis labeled “w/o star weight” is for testing the impact of the star weight used in the clustering measurement of w_p (see More *et al.* [23] for details) on our results, where the star weight is designed to account for the systematic effect related to seeing and stellar density in the target selection of SDSS galaxies. We use the w_p data measured without the star weight in [23], instead of our default w_p , for the baseline $3\times 2\text{pt}$ analysis.

We also perform the analyses using the extended models that are more flexible models for the galaxy-halo connection; we use the extended models including the effects of off-centered central galaxies or the incompleteness effect of central galaxies [104]. For both extended models, we introduce two additional model parameters, as indicated in the rows “off-centering” or “incompleteness”.

In addition, after unblinding our cosmology results (see next section), we further decided to perform the analyses labeled “ $\sigma(\Delta m) = 0.1$ prior” and “ $\sigma(\Delta z_{\text{ph}}) = 0.2$ prior”. For these, we employ Gaussian priors, $\Pi(\Delta m) = \mathcal{N}(0, 0.1)$ or $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.2)$, to study the impact of the prior choices on the cosmological parameters. Here $\sigma(\Delta m) = 0.1$ is much wider than the fiducial prior, $\sigma(\Delta m) = 0.01$ (see Table II), and this test is for assessing the self-calibration power of the residual shear calibration factor with our $3\times 2\text{pt}$ method (therefore, we consider the overwhelmingly pessimistic shear calibration error). We also perform an analysis with the cosmological parameters other than $(\Omega_m, \ln(10^{10} A_s))$ fixed to the *Planck* 2015 “TT,TE,EE+lowP” constraints [76] to check how the parameters that are not well constrained by our data vector affect our cosmological constraints. This setup is labeled “2 cosmo” in Table III.

G. Validation of Modeling and Analysis Choices

To check the robustness of our results to our modeling and analysis choices, we perform various validation tests. This includes tests of different samplers, different models of galaxy-halo connection, different models for baryonic feedback, as well as different models of observational systematics including residual systematic error in the mean source redshift and PSF systematics. Please also see Miyatake *et al.* [15] for the detailed validation tests, where the scale cuts of w_p and $\Delta\Sigma$, $R = 2$ and $3 h^{-1}\text{Mpc}$, were validated in the sense that estimated parameters, such as S_8 , do not have significant biases compared to their statistical errors. For the tests in this paper, we also include the synthetic data of cosmic shear correlation functions ξ_{\pm} and perform the validation tests using the covariance matrix for the HSC-Y3 data. The details of the validation tests are given in Appendix A, and here we give a summary of the important points.

For w_p and $\Delta\Sigma$ we have to study the impact of uncertainties in the galaxy-halo connection on the cosmological constraints. Since HOD is an empirical prescription for the galaxy-halo connection, our philosophy is that we should include a sufficient number of the HOD parameters and then extract the cosmological information from the halo clustering quantities, accurately modeled by `Dark Emulator`, after marginalizing over the galaxy-halo connection parameters. For the validation tests of our analysis method, we generate various types of mock SDSS galaxies, where we employ different ways to populate galaxies into halos in N -body simulations and then generate the synthetic data vectors of w_p and $\Delta\Sigma$ from the mock catalogs [also see 15]. We then apply our baseline analysis pipeline using the fiducial HOD model to the synthetic data vector to assess whether our method can recover the input cosmological parameters used in the mock catalogs. As shown in Appendix A, our method can recover the S_8 value with an accuracy better than $\sim 0.5\sigma$ for most of the mock SDSS galaxy catalogs.

Nevertheless, reflecting on the fact that there is no established theory of the halo-galaxy connection, we also consider the “worst-case” scenario in order for us to be ready for surprises due to unknown systematic effects. We prepare extreme mock catalogs of SDSS galaxies, where we implement a non-standard prescription of the galaxy-halo connection, e.g. the mock catalogs including the overwhelmingly large assembly bias effect and the off-centering effect of central galaxies. These worst-case scenario mocks change w_p and $\Delta\Sigma$ at scales around the scale cuts, $R = 2$ and $3 h^{-1}\text{Mpc}$, and could cause a significant bias in S_8 (more than 0.5σ). Even for these worst-case scenarios, we have a useful diagnostic to flag such an extreme systematic effect on the cosmological parameters. Since the halo model based theory includes the linear-theory prediction at large scales, which can be applied to any types of galaxies [105], the method can recover the underlying cosmological parameters *if* employing sufficiently large scale cuts, such as $R = 8$ and $12 h^{-1}\text{Mpc}$ [15, 99]. Hence, if a significant systematic effect exists in the actual SDSS galaxies, the estimated central value of S_8 would have a systematic shift with changing the scale cuts. Based on these findings, we

can employ the different scale cuts for the actual cosmology analysis to monitor a change in cosmological parameters as an indicator of potential systematic effects.

For the systematic effects on the cosmic shear signal ξ_{\pm} , our modeling method is very similar to that of Li *et al.* [25]. We employ exactly the same scale cuts in ξ_{\pm} as those in Li *et al.* [25]. All the validation tests are passed in the sense that the S_8 value is recovered using synthetic data vectors of ξ_{\pm} where different models of systematic effects (baryonic feedbacks and PSF systematics) are implemented. We did not find any flag in our analysis method arising from modeling inaccuracy and systematic effects in cosmic shear.

One of the most important validation tests is to assess the sensitivity of our method to a residual systematic error in the mean source redshift of HSC galaxies, Δz_{ph} . As described in Sugiyama *et al.* [24] in detail, we implement a non-zero shift (non-zero Δz_{ph}) in the source redshift distribution to generate synthetic data vectors of $\Delta\Sigma$ and ξ_{\pm} and then assess whether our analysis method can recover the input Δz_{ph} as well as the S_8 parameter. As shown in Appendix A, we find that, if a non-zero shift of Δz_{ph} by more than $|\Delta z_{\text{ph}}| \sim 0.1$ exists in the synthetic data, the use of the uninformative flat prior of Δz_{ph} allows us to recover the underlying true S_8 value to within the credible interval. In other words, if we employ an informative prior on Δz_{ph} , such as $\mathcal{N}(0, 0.1)$, it could yield a significant bias in S_8 larger than the 1σ statistical error. Thus, the HSC-Y3 data has the statistical power to calibrate Δz_{ph} to the precision of $\sigma(\Delta z_{\text{ph}}) \simeq 0.1$ and then recover S_8 . This gives a validation of our analysis method using an uninformative prior of Δz_{ph} , $\mathcal{U}(-1, 1)$, even if the method gives a large credible interval in exchange. In this sense, our cosmology analysis can be considered robust and conservative.

IV. BLINDING SCHEME AND UNBLINDING CRITERIA

To avoid confirmation bias we perform our cosmological analysis in a blind fashion. We employ a two-tier blinding strategy to avoid unintentional unblinding during the cosmological analysis. The two tiers are as follows:

- *Catalog level:* The analysis team performs the cosmological analysis using three different weak lensing shape catalogs. Only one is the true catalog and the other two are fake catalogs (see below for details). The analysis team members do not know which is the true catalog.
- *Analysis level:* When the analysis team makes plots comparing the measurements with theoretical models, the y -axis values (e.g., the amplitudes of $\Delta\Sigma$) are hidden and the analysis team is not allowed to see the values of cosmological parameters used in the theoretical models. When the analysis team makes plots showing the credible intervals of cosmological parameters (i.e. the posterior distribution), the central value(s) of parameter(s) are shifted by the mode of the baseline 3×2 pt analysis (without looking at the mode value), and only the range of the credible interval(s) can be seen. With this

blinding method, we can evaluate how the cosmological parameters change by each analysis setup compared to those of the baseline analysis. Finally, the analysis team does not compare the posterior of cosmological parameter(s) or the best-fit model predictions with external results such as the *Planck* CMB cosmology, prior to unblinding.

See Section IIB of More *et al.* [23] for details of how the fake catalogs were constructed in a manner that prevents accidental unblinding by the analysis team. The use of these catalogs means that the analysis team must perform the same analyses three times, but this method avoids the need for re-analysis once the catalogs are unblinded.

The set of the three shape catalogs used in this paper is shared with the two companion papers, More *et al.* [23] and Sugiyama *et al.* [24]. The first of these presents the details of the measurements of clustering observables used in this paper. The companion paper Sugiyama *et al.* [24] presents the cosmological parameter estimation from the same signals as that of this paper, but using the perturbation theory based model template that is compared to the signals on scales with the larger scale cuts of $(8, 12)h^{-1}\text{Mpc}$ for w_p and $\Delta\Sigma$, respectively.

We imposed the following criteria for deciding to unblind our results:

- Analysis pipeline codes are made available to collaboration members and some specific members are assigned to review each part of the code.
- Various analysis setups, described in Table III, are tested to ensure that the cosmological constraints are robust to the different setups. We check whether a significant shift in S_8 , $> 0.5\sigma$, (where σ is from the baseline $3\times 2\text{pt}$ analysis), is found from any of the analysis setups. If such a shift is found, we check whether it is consistent with the statistical scatter by comparing the shift with the distribution of S_8 values estimated from a set of hypothetical analyses using 100 realizations of the synthetic noisy data vector. In particular we check how the S_8 value changes for different scale cuts as a flag for the assembly bias effect (see Section VIB for detailed discussion).
- Internal consistency tests to check whether an estimation of the key cosmological parameter changes, compared to that from the baseline analysis method, using subsets of data vector and/or different analysis methods, where the different methods are based on more flexible models of galaxy-halo connection including the incompleteness and off-centering effects of central galaxies (see Table II).
- Quantify the goodness of fit of the best-fit model predictions to the data vector in each of the three blind catalogs.

Once the collaboration agrees to unblind the analysis, the analysis-level unblinding is first removed by the analysis team. The final catalog-level unblinding happens a few hours

later. The analysis team resolved that the results would be published regardless of the outcome, once the results are unblinded. In addition, the analysis method could not be changed or modified after unblinding. Following these agreed rules, we unblinded the results at our regular telecon on Dec 3, 2022 in JST. The figures shown in this paper, unless otherwise noted, were made prior to unblinding, with only the axes changed after unblinding to show true values.

V. RESULTS

In this section we show the main results of this paper, the cosmological parameters estimated from the joint measurements of $\Delta\Sigma(R)$, $w_p(R)$ and $\xi_{\pm}(\vartheta)$ for the HSC-Y3 and SDSS catalogs. All the analyses in this section are done before unblinding, and the results are presented without any change after unblinding.

A. ΛCDM Constraints

The shaded contours in Fig. 1 are the 1d and 2d posterior distributions of the key parameters, S_8 , σ_8 and Ω_m for flat ΛCDM model, obtained from the baseline $3\times 2\text{pt}$ analysis setup of the HSC-Y3 data as given in Table III. The central value and credible interval for each parameter are given as

$$\begin{aligned}\Omega_m &= 0.382_{-0.047}^{+0.031}(0.401), \\ \sigma_8 &= 0.685_{-0.026}^{+0.035}(0.696), \\ S_8 &= 0.763_{-0.036}^{+0.040}(0.805), \\ \Delta z_{\text{ph}} &= -0.05 \pm 0.09,\end{aligned}\tag{40}$$

where the number in parentheses denotes the value for the MAP (*maximum a posteriori*) model in the sampled chains. The HSC-Y3 $3\times 2\text{pt}$ analysis achieves about 5% fractional accuracy in the S_8 parameter.

Fig. 2 shows that the best-fit (MAP) model fits all the measured quantities simultaneously over the range of radial or angular separations that are used in the cosmology analysis. We note that our cosmology analysis does not include the information in the deeply nonlinear regime such as the 1-halo term regime, e.g. $R \lesssim$ a few $h^{-1}\text{Mpc}$ corresponding to the virial radii of massive halos. Nevertheless we stress that the use of `Dark Emulator` is critical to accurately model the clustering observables on scales in the mildly nonlinear regime. If we use the simpler perturbation-theory based model (Sugiyama *et al.* [24]), it cannot describe the signals well in the range of scales we use in this paper [also see 7].

In Fig 3 we evaluate the goodness-of-fit of the best-fit model to the measured signal. To do this evaluation, we generate 100 realizations of noisy mock data vectors using the “full” covariance matrix; the full covariance includes the elements in radial or angular bins outside those used in our cosmology analysis and the cross-covariance terms that describe correlated scatter between the observables, i.e., the galaxy-galaxy weak lensing signals for the different lens subsamples




FIG. 1. The 1d and 2d posterior distributions in the sub-space of S_8 , σ_8 and Ω_m for the flat Λ CDM cosmology. The blue dark (light) shaded regions denote the 68% (95%) credible interval, respectively, for our HSC-Y3 3×2 pt baseline analysis in Table III. For comparison we show the results for other recent cosmological analyses. The red contours are from the DES-Y3 3×2 pt analysis [8]. The blue contours are from the KiDS-1000 analysis [9] with cosmic shear (“CS”) and galaxy-galaxy weak lensing (“GGL”) (see text for details). The green contours are the *Planck* 2018 results using the primary CMB anisotropy information (“TT,TE,EE+lowE”) [11]. Note that the degeneracy direction of the HSC-Y3 result in each 2d subspace of the parameters are different from those of DES-Y3 and KiDS-1000, since the relative constraining powers of the cosmological parameters for different observables are different.

and cosmic shear as shown in Fig. 6 in More *et al.* [23]. The histogram in Fig. 3 shows the distribution of the χ^2 -value of the MAP model prediction for each of the 100 realizations. We find that the χ^2 values tend to exceed that expected from the degrees of freedom, $\nu = 90 - 28 = 62$ (see Table III). We ascribe this excess to severe parameter degeneracies; some of the model parameters, especially the HOD parameters, are not well constrained by the data vector. The histogram can be compared to the χ^2 value of the actual HSC-Y3 and SDSS analysis (solid black line), showing that the observed χ^2 value

is near the middle of the distribution. Hence, we conclude that the best-fit model is quite acceptable.

B. Internal Consistency Tests

As self-consistency tests, we performed the cosmological parameter estimation for each of the different setups listed in Table III, before unblinding. The cosmological constraints for each setup are shown in Fig. 4 and the mode, credible inter-




FIG. 2. The green solid line in each panel denotes the model prediction at the *maximum a posteriori* (MAP) for the baseline analysis in Fig. 1, while the data points with error bars are the measured signals. The upper-row three panels are for the projected correlation functions of galaxies, $w_p(R)$, for the LOWZ, CMASS1 and CMASS2 samples in the redshift ranges $z = [0.15, 0.35]$, $[0.43, 0.55]$ and $[0.55, 0.70]$, respectively. The middle-row three panels are for the galaxy-galaxy weak lensing using the HSC galaxies as source sample, $\Delta\Sigma(R)$, for the same LOWZ, CMASS1, and CMASS2 samples as lens samples, respectively. The bottom-row two panels are for the cosmic shear correlation functions, $\xi_{\pm}(\vartheta)$. For illustration purpose, we show $R \times w_p(R)$, $R \times \Delta\Sigma(R)$ and $\vartheta \times \xi_{\pm}(\vartheta)$. The red shaded regions around the green line denote the 68% and 95% credible intervals of the model predictions in each separation bin, which are computed from the posterior distributions in the Bayesian cosmology inference. Note that the errors are computed from the diagonal components of the covariance matrix. The blue-color shaded region in each panel denotes the range of projected or angular separation bins that is used for the cosmology analysis.

val, and MAP value of each parameter are summarized in Table V of Appendix B. We find that the S_8 parameter is robust to these different tests, changing by $< 1\sigma$ in each case. The exception is the test in which the residual photo- z error parameter is fixed to $\Delta z_{\text{ph}} = 0$, i.e. the case in which the mean redshift of HSC source galaxies is assumed to be perfectly estimated based on their photo- z 's. The test using the fixed $\Delta z_{\text{ph}} = 0$ gives a smaller error bar in S_8 and gives a sizable shift in the central value of S_8 (1.5σ where σ is taken

from the test with fixed Δz_{ph}). This indicates the existence of a non-zero residual photo- z error as discussed below. Some tests show a scattered shift in the values of Ω_m and σ_8 , but such shifts are also seen in the validation tests using the mock catalogs. We did not find any significant shift of S_8 compared to the tests using the mock data, or did not find any evidence of unknown systematic effects in our results. We also note that the shifts in these parameters are likely due to projection effects of the non-Gaussian posterior distribution in the




FIG. 3. An evaluation of the goodness-of-fit of the best-fit (MAP) model in Fig. 1. The histogram shows the distribution of the χ^2 values at the MAP model, obtained by applying the same baseline analysis to 100 noisy mock datasets (see text for details). The vertical black line denotes the measured χ^2 -value ($\chi^2 = 85.1$) at MAP for the actual analysis of HSC-Y3 and SDSS data. The probability of finding the χ^2 value larger than the observed value (p value) is about 41%.

full-dimensional parameter space. In Appendix B we give a detailed discussion of the internal consistency tests.

In Fig. 5 we compare the cosmological constraints from the different cosmology analyses using the HSC-Y3 data. Although the result “ 3×2 pt large-scale” uses the same data vector as that in this paper, the analysis uses the perturbation theory based theoretical template to compare with the measurements at $R > 8$ and $12 h^{-1} \text{Mpc}$ for w_p and $\Delta\Sigma$, respectively, where the perturbation theory model is valid (Sugiyama *et al.* [24]) The other two results are from the cosmic shear tomography analyses using the cosmic shear two-point correlations [25] and the power spectra [26]. The three sets of analyses (both 3×2 pt analyses, the real-space cosmic shear analysis and the Fourier-space cosmic shear analysis), performed blinded cosmology analyses using different blinded catalogs, and we made this comparison plot after the unblinding. The cosmological results from all the four analyses, especially the S_8 results, are in good agreement with each other. This is quite encouraging, because the constraining power of the 3×2 pt analyses is mainly from the clustering (w_p) information of SDSS galaxies and the two cosmic shear constraints are sensitive to different scales in the cosmic shear information of HSC-Y3 data due to the different scale cuts in the real- and Fourier-space. We also emphasize that the two cosmic shear analyses adopted the uninformative prior on the residual photo- z error parameters for the two high-redshift tomographic bins, z_3 and z_4 , that correspond to redshifts at $z \gtrsim 0.9$. If the cosmic shear analyses employ informative priors on the photo- z error parameters, the S_8 parameter is shifted and the agreement in Fig. 5 cannot be realized. Hence all the cosmological analyses indicate a non-zero residual systematic error in the photo- z estimates of such high-redshift

HSC galaxies.

C. Comparison with External Data and S_8 Tension

In this section, we discuss the comparison of our HSC-Y3 result with external cosmology results (see Fig. 1). Note that, as we described in Section IV, we never compared the HSC-Y3 results with external cosmology results during the blind analysis stage, and made plots like Fig. 1 only after unblinding. For the CMB constraints, we consider the “Planck2018” results [11], from the analysis where the primary CMB temperature and E -mode polarization anisotropy information (“TT, EE, TE+lowE”) are used and the neutrino mass is fixed to 0.06 eV as we did in our setup. To obtain the posterior distribution we used the public chain “base/plikHM_TTTEEE_lowl_lowE” available from the website¹². For the posterior distribution of the DES Year 3 (“DES-Y3”) result, we used the public chain¹³, which is the result obtained from the 3×2 pt cosmological analysis using the photometric “MagLim” samples for both lens and source galaxies [8]. For the “KiDS-1000” result, we used the public chain¹⁴ to exhibit the result from a joint analysis of cosmic shear (“CS” in the legend) and galaxy-galaxy (“GGL”) weak lensing in Heymans *et al.* [9], where the *spectroscopic* samples in BOSS and the 2-degree Field Lensing Survey (2dFLenS) were used as the lens samples in the galaxy-galaxy weak lensing measurements. Note that the 3×2 pt results from KiDS-1000 include BAO information in the BOSS galaxies that can give a tighter constraint on Ω_m , so we instead refer the above 2×2 pt CS \times GGL result as KiDS-1000. For both the DES-Y3 and KiDS-1000 analyses, the weak lensing observables are angular correlation functions – $\gamma_T(\theta)$ and/or $w(\theta)$ – rather than $\Delta\Sigma(R)$ and $w_p(R)$.

Fig. 1 shows that the HSC-Y3 result is generally consistent with the DES-Y3 and KiDS-1000 results within the credible intervals. However, the degeneracy direction of the HSC-Y3 result in each 2d subspace of the parameters are different from those of DES-Y3 and KiDS-1000, as the relative constraining powers of different observables for the cosmological parameters (after marginalizing over other parameters) are different. For our case, the galaxy-galaxy clustering of BOSS galaxies has the most constraining power, after lifting the parameter degeneracies between the galaxy bias and the cosmological parameters with the weak lensing information as we will discuss below.

Extending the definition of S_8 to $S'_8 \equiv \sigma_8(\Omega_m/0.3)^\alpha$ with α being a free parameter, we find that the best-constrained parameter is $\alpha \simeq 0.22$: with this value, we find $S'_8 \simeq$

¹² https://pla.esac.esa.int/pla/aio/product-action?COSMOLOGY.FILE_ID=COM_CosmoParams_fullGrid_R3.01.zip

¹³ <https://des.ncsa.illinois.edu/releases/y3a2/Y3key-products>

¹⁴ https://kids.strw.leidenuniv.nl/DR4/KiDS-1000_3x2pt_Cosmology.php




FIG. 4. A summary of the cosmological parameters and the residual photo- z error parameter Δz_{ph} , estimated from each of the different analysis setups in Table III. The vertical dashed line in the panel of Δz_{ph} denotes $\Delta z_{\text{ph}} = 0$, i.e. the case of no residual photo- z error or equivalently the case that the mean redshift estimate of HSC source galaxies inferred from the photo- z estimates is perfect. As our default choice, we employ the uninformative flat prior on the residual photo- z error, $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$. As explained in Table III, the analysis with superscript “*” denotes an analysis using the informative Gaussian prior on the photo- z error, given by $\mathcal{N} = (-0.06, 0.08)$, which is inferred from the posterior of the baseline $3 \times 2\text{pt}$ analysis. The analysis with superscript “†” denotes an analysis using the Gaussian prior on Δz_{ph} with mean around $\Delta z_{\text{ph}} = 0$, given by $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$. The analysis with label “ $\sigma(\Delta z_{\text{ph}}) = 0.2$ prior” denotes the result using the Gaussian prior, $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.2)$.

0.721 ± 0.0279 , the fractional precision is 4%. This precision is 30% smaller than the $\sigma(S_8) \simeq 0.040$ for the standard S_8 in Eq. (40). If we compare the width of the 2d contour along the narrowest direction in the (Ω_m, S_8) sub-space in Fig. 1, the HSC-Y3 result is comparable with the KiDS-1000 result, but is somewhat larger than the DES-Y3 result. This is partly due to our use of an uninformative prior on the resid-

ual photo- z error parameter (Δz_{ph}). If we employ a tighter prior on Δz_{ph} such as the prior width inferred from the original photo- z estimate, $\sigma(\Delta z_{\text{ph}}) \simeq O(10^{-2})$, we can obtain a tighter credible interval, however, the central value of S_8 shows a non-negligible shift. Hence Δz_{ph} is a key parameter in our analysis to obtain a robust estimate of the cosmological parameters, and we decided to adopt the uninformative prior




FIG. 5. Comparison of the cosmological constraints from the different cosmological analyses of the HSC-Y3 data. The result labeled as “ 3×2 pt large scale” uses the same data vector as that in this paper, but uses the perturbation theory based model template when comparing the predictions to the measurements on large scales (Sugiyama *et al.* [24]). “Cosmic shear tomography: Real” is the result from the cosmic shear tomography analysis using the real-space cosmic shear correlation functions with 4 tomographic redshift bins (Li *et al.* [25]). “Cosmic shear tomography: Fourier” is from the cosmic shear tomography using the cosmic shear power spectra (Dalal *et al.* [26]). The two cosmic shear results use the different scale cuts.

of Δz_{ph} during the blind analysis stage before revealing the central value of S_8 . In Section VI A, we will give a more detailed discussion of how different treatments of the residual photo- z error, e.g. informative vs. uninformative prior, alter our cosmological constraints.

Fig. 1 displays a 2σ -level tension between the HSC-Y3 3×2 pt result and the *Planck* 2018 result. To quantify the possible tension, we use the methods developed in Park and Rozo [106] and Raveri *et al.* [107], which are called *eigentension* and *tensiometer*, respectively [also see 108].

For the *eigentension* method, we start by diagonalizing the covariance matrix of cosmological parameters to find the eigenvectors and eigenvalues. Among the 5 cosmological parameters in our 3×2 pt analysis n_s and ω_b are prior dominated, so we focus on the parameters, σ_8 , Ω_m , and ω_c . When we diagonalize the covariance matrix of these parameters, obtained from the chains in our baseline analysis, we find the two eigenvectors, $(e_0, e_1) = (\sigma_8 \omega_c^{0.19} \Omega_m^{0.38}, \omega_c \Omega_m^{0.54} \sigma_8^{-0.40})$, are well constrained by the HSC-Y3 3×2 pt observables compared to the prior widths, while the third eigenvector is prior-dominated. If we compute the posterior distribution of the eigenvector differences, defined as $(\Delta e_0, \Delta e_1) \equiv (e_0, e_1)_{\text{HSC-Y3}} - (e_0, e_1)_{\text{Planck}}$, from the two chains of the

HSC-Y3 baseline analysis and *Planck* 2018, we find that the point where the two data sets are consistent with each other, i.e., $(\Delta e_0, \Delta e_1) = (0, 0)$, is located at $\sim 2.5\sigma$ in the posterior. Note that this method allows us to compute the posterior distributions of these parameter differences from the existing chains of HSC-Y3 and *Planck*, as long as the two datasets are independent [109]. Thus, we conclude that the HSC-Y3 result displays a 2.5σ tension with the *Planck* 2018 result.

To implement the *tensiometer* method, we use the publicly-released code¹⁵. This code allows us to generate the posterior distribution of the three parameter differences, $(\Delta\sigma_8, \Delta\Omega_m, \Delta\omega_c)$, from the two chains of the HSC-Y3 baseline analysis and *Planck* 2018 using machine learning modeling of the posterior distribution with normalizing flows. It then quantifies a disagreement significance in the full 3d parameter space: we find a 2.7σ tension between the HSC-Y3 and the *Planck* 2018 results, in close agreement with the estimate from *eigentension* above.

Hence, we conclude that the HSC-Y3 3×2 pt result has about 2.5σ tension with the *Planck* 2018 constraints within the flat-geometry Λ CDM framework.

VI. DISCUSSION

A. An Implication of Residual Systematic Photo- z Error

A notable aspect of this study, compared to other weak lensing cosmology analyses, is that we estimate the cosmological parameters employing an *uninformative* prior on the residual photo- z error parameter of source galaxies; $\Delta z_{\text{ph}} : \mathcal{U}(-1, 1)$ (see Table II). This is a conservative setup which is equivalent to the case in which we do not adopt any prior knowledge about the mean redshift of HSC source galaxies. In this section, we show how the self-calibration of the photo- z error parameter is achieved by our method, and also study how the cosmological parameters change when using different priors on Δz_{ph} .

The left panel of Fig. 6 shows how the residual photo- z error parameter Δz_{ph} is calibrated by combining the different observables. For the 2×2 pt ($\Delta\Sigma \times w_p$) analysis and the cosmic shear, we also employ the flat prior $\mathcal{U}(-1, 1)$ for Δz_{ph} . The figure shows that the different observables are complementary to each other and Δz_{ph} is accurately estimated by combining the three observables. The self-calibration of Δz_{ph} is achieved by comparing the galaxy-galaxy weak lensing signals for the three lens subsamples (LOWZ, CMASS1 and CMASS2) at the three different spectroscopic redshifts and the cosmic shear signals for the same source galaxies. The baseline 3×2 pt analysis suggests a non-zero value of $\Delta z_{\text{ph}} = -0.05 \pm 0.09$, i.e. a 1σ -level hint of a non-zero residual systematic error in the mean source redshift. This indicates that the true mean redshift of HSC source galaxies is higher than the photo- z estimate by $|\Delta z_{\text{ph}}| = 0.05$. Furthermore, if we

¹⁵ <https://github.com/mraveri/tensiometer>




FIG. 6. The importance of the uninformative prior on the residual photo- z error parameter (Δz_{ph}) in the cosmology analysis. *Left panel:* The 1d and 2d posterior distributions obtained using the different observables: the baseline ($3 \times 2\text{pt}$), the $2 \times 2\text{pt}$ ($\Delta\Sigma \times w_p$), and the cosmic shear correlations. For all the analyses, we employ the flat prior of Δz_{ph} : $\mathcal{U}(-1, 1)$ as our baseline analysis. *Right:* The posterior distributions for the $3 \times 2\text{pt}$ analyses when using the different priors of Δz_{ph} : the baseline analysis (the flat prior), the Gaussian prior of $\mathcal{N}(0, 0.1)$ and the case fixing $\Delta z_{\text{ph}} = 0$, respectively. The result of $\Delta z_{\text{ph}} = 0$ approximately corresponds to the case where the redshift distribution of HSC source galaxies is as inferred from the photo- z estimate, because the prior is as informative as $\sigma(\Delta z_{\text{ph}}) \sim 10^{-2}$. The different treatments of Δz_{ph} affect the mode value and the size of the credible interval of the cosmological parameters.

focus on the $\Omega_m = 0.3$ cross section of the posterior, where $\Omega_m \simeq 0.3$ is indicated by measurements of BAO or the galaxy clustering [e.g. 21], one fits an even larger bias, $\Delta z_{\text{ph}} \sim -0.2$, as we will below study in more detail. This case also implies a lower value of S_8 , since Δz_{ph} and S_8 are positively correlated. Hence, employing an uninformative flat prior of Δz_{ph} is important to obtain an unbiased estimate of S_8 , if the non-zero Δz_{ph} is genuine. Encouragingly, a similar residual photo- z error for HSC source galaxies at high redshifts is also implied by the real- and Fourier-space cosmology analyses of HSC-Y3 cosmic shear tomography in Li *et al.* [25] and Dalal *et al.* [26], respectively. For these cosmology analyses, the photo- z error parameters for the two high- z bins (corresponding roughly to our source galaxy sample) are calibrated by the cosmic shear signals relative to those in the lower redshift bins that are more reliably estimated by the photo- z and the cross-correlation method [88]. Thus the photo- z error calibration by the cosmic shear methods is somewhat independent from the calibration of the $3 \times 2\text{pt}$ method in this paper.

B. Assembly Bias

One concern in the halo model based analysis is the effect of possible assembly bias of the SDSS galaxies on the cosmological parameters [15, 110]. Even if we use the w_p and $\Delta\Sigma$ information on scales greater than the size of most massive

halos, $R \gtrsim 2$ and $3 h^{-1}\text{Mpc}$, respectively, the galaxy-galaxy lensing ($\Delta\Sigma$) contains information on the interior mass of halos hosting the SDSS galaxies, which in turn lifts degeneracies in the galaxy-halo connection in the clustering amplitudes of the 2-halo term regime. If the SDSS galaxies are affected by assembly bias, it could cause a bias in the cosmological parameters, because of a breakdown in the *simple* galaxy-halo connection as a function of halo mass. To test the impact of possible assembly bias, we perform the parameter estimation using the different scale cuts, $R = [4, 6]$ or $[8, 12] h^{-1}\text{Mpc}$, respectively. On sufficiently large scales, galaxy clustering properties are governed by gravity, and the correlation coefficient function of galaxy clustering approaches to the simple relation irrespective of galaxy types including a galaxy sample with assembly bias, given by $\xi_{\text{gm}}/[\xi_{\text{mm}}\xi_{\text{gg}}]^{1/2} \simeq 1$ [see Fig. 6 in Ref. 111, for the results using the Illustris hydrodynamical simulations] [also see 15, 20]. As demonstrated in Miyatake *et al.* [15], if we adopt the large scale cuts of $[8, 12] h^{-1}\text{Mpc}$, the clustering signals are safely in the 2-halo term regime, and the cosmology analysis can recover the cosmological parameters even if the assembly bias effect exists.

Fig. 7 shows the posterior distribution when using the different scale cuts, $R = [4, 6]$ or $[8, 12] h^{-1}\text{Mpc}$ for the w_p and $\Delta\Sigma$ signals, respectively, in the cosmology analysis. Even if we use the largest scale cut $R = [8, 12] h^{-1}\text{Mpc}$, the cosmological parameters are almost unchanged. In fact, the cosmological parameters for the baseline analysis are also consistent




FIG. 7. The posterior distributions of the cosmological parameters when using different scale cuts of $R = [4, 6]$ or $[8, 12] h^{-1}\text{Mpc}$ for the w_p and $\Delta\Sigma$ signals, respectively, in the cosmology analysis. The gray contours are the same as those in Fig. 1. Note that we used the Gaussian prior of the residual photo- z error parameter, $\mathcal{N}(-0.06, 0.08)$, that is inferred from the baseline $3\times 2\text{pt}$ analysis and the results include the cosmic shear information too, where we used the same range of the angular separations as in the baseline analysis.

with the results obtained using the perturbation theory based method in Sugiyama *et al.* [24] as shown in Fig. 5. Thus, the clustering signals do not exhibit any signature of the assembly bias effect. While the results for $R = [4, 6] h^{-1}\text{Mpc}$ show a shift in the posterior distribution, we checked that the shift is caused primarily by upward scatter in the data points of $\Delta\Sigma$ around the scale cut for the LOWZ and CMASS1 subsamples (see Fig. 2), which causes the code to prefer unphysical regions of the HOD parameters and then leads to a shift in cosmological parameters. We found that, if we remove the scattered data points, the shift in cosmological parameters does not occur.

As a further sanity check, we run the cosmology analyses for 100 realizations of the mock data vector that do not include the assembly bias effect. Note that for this test, we employ the photo- z prior indicated from the $(2,3)h^{-1}\text{Mpc}$ scale cut analysis for the $(8, 12) h^{-1}\text{Mpc}$ scale-cut analysis in each realization as we did in the actual analysis. Fig. 8 shows in the difference in the S_8 values with different scale cuts we found from the real HSC-Y3 data occur with a reasonable chance. The two arrows in the figure show a shift in the S_8 values found from the two assembly bias mocks that we use in our validation tests (see Appendix A), and the S_8 difference is located at the tail of the 100 realizations, significantly displaced from the measured difference value. Thus we conclude that our cosmology results are unlikely to be affected by assembly




FIG. 8. The shaded histogram shows the expected distribution of the differences between the S_8 values obtained from the $3\times 2\text{pt}$ analyses using the different scale cuts of $(2,3)$ and $(8, 12)h^{-1}\text{Mpc}$ for w_p and $\Delta\Sigma$, respectively, assuming that w_p and $\Delta\Sigma$ are not contaminated by the assembly bias effect. To obtain the distribution, we perform the same cosmology analysis on each of the 100 realizations of the noisy mock data vector. Note that in this inference simulation, we employ the prior on the residual photo- z error, Δz_{ph} obtained from the fiducial analysis of $(2,3)h^{-1}\text{Mpc}$ scale cut to each realization for the $(8,12)$ -analyses, as we did for the actual analysis. When there is no assembly bias effect, the S_8 values from the $(2,3)$ - and $(8,12)$ -scale cuts should be consistent with each other, and the actual observed difference of S_8 , as denoted by the vertical solid line, is consistent with the distribution from the synthetic data vector. The probability of finding ΔS_8 larger than the observed value (p -value) is about 50%. The two arrows indicated by “assembly- b ” and “assembly- b -ext” denote the expected difference values of S_8 obtained from the simulated synthetic data, where assembly bias effects with different amplitudes are included.

bias.

C. Post-unblinding Analysis: The Impact of Ω_m Prior

During the blind analysis stage (see Section IV), we did not compare the posterior distribution of the cosmological parameters in our analysis with any external datasets. Our main result of Fig. 1 indicates a higher Ω_m than the *Planck* constraint. Ω_m is well constrained by the baryon acoustic oscillation information in galaxy clustering [112] for flat-geometry ΛCDM model. Since we did not intentionally include the BAO information of SDSS galaxy clustering in our analysis, here we study how a BAO-motivated prior of Ω_m affects our results, as part of our post-unblinding analysis. To do this, we employ the Gaussian prior given by $\Omega_m : \mathcal{N}(0.3, 0.01)$ in the baseline $3\times 2\text{pt}$ analysis, where the central value of Ω_{m0} and the width of $\sigma(\Omega_{m0}) = 0.01$ are roughly consistent with the constraints obtained from the BAO analyses [113, 114] [also see 115, for the CMB-independent constraint].




FIG. 9. The posterior distribution of the parameters when the Gaussian prior of $\Pi(\Omega_m) = \mathcal{N}(0.3, 0.01)$ as motivated by BAO constraints is added. This analysis was done as a part of the post-unblinding analysis.

Fig. 9 shows the 2d posterior distributions of the parameters. We find

$$\begin{aligned} S_8 &= 0.732^{+0.027}_{-0.029}(0.738), \\ \Delta z_{\text{ph}} &= -0.133^{+0.077}_{-0.084}(-0.132). \end{aligned} \quad (41)$$

The prior of Ω_m slightly lowers the central value of S_8 , and also indicates a larger central value of $|\Delta z_{\text{ph}}|$ than we found for the fiducial analysis (Eq. 40). If we apply the eigentension method to quantify a tension between the HSC $3 \times 2\text{pt}$ analysis and the *Planck* result, we find the tension is at about 2.4σ , almost no change from the result in Section V C.

VII. CONCLUSION

In this paper we have carried out a cosmology analysis combining three clustering observables, the projected correlation function (w_p), galaxy-galaxy weak lensing ($\Delta\Sigma$), and cosmic shear correlation functions (ξ_{\pm}). These quantities are measured from the spectroscopic SDSS galaxy samples as lens samples of $\Delta\Sigma$ and tracers of the w_p , and the HSC-Y3 photometric galaxy samples for source galaxies in the $\Delta\Sigma$ and ξ_{\pm} measurements. One of the most important aspects of our analysis is that we use a *single* source sample in the weak lensing measurements, allowing us to self-calibrate the residual error in the mean redshift of source galaxies, which is one of the most important systematic effects in weak lensing cosmology. We do so by comparing the relative $\Delta\Sigma$ -amplitudes for the

three spectroscopic lens subsamples and the cosmic shear signal, as suggested in Oguri and Takada [14]. We employ a completely *uninformative* flat prior, $\mathcal{U}(-1, 1)$, for the residual photo- z error parameter Δz_{ph} in our cosmology analysis. We showed that, with the statistical power of the HSC-Y3 data, we can estimate the cosmological parameters and the photo- z parameter Δz_{ph} simultaneously. We decided on this analysis setup during the blind analysis stage, without looking at the estimated values of cosmological parameters, and we froze the analysis method including the flat prior of Δz_{ph} before unblinding. This allowed us to obtain a *robust* estimate of the cosmological parameters, which minimizes the impact of the possible photo- z errors, even at the cost of larger error bars of the cosmological parameters.

The parameters we obtained for the flat ΛCDM model are: $S_8 = 0.763^{+0.040}_{-0.036}$ and $\Delta z_{\text{ph}} = -0.05 \pm 0.09$ after marginalizing over a number of other parameters. Thus we have estimated S_8 with a fractional precision of 5%. Extending the S_8 definition to $S'_8 = \sigma_8(\Omega_m/0.3)^\alpha$, we showed that our method gives the best constraint with $\alpha = 0.22$; we find $S'_8 = 0.721 \pm 0.028$ with $\alpha = 0.22$, about 4% fractional precision. These S_8 values are lower than indicated by the *Planck* CMB result. Using the tension metric in Refs. [106] and [107], we quantified the tension to be about 2.5σ . We plan to use extended models such as dark energy models, i.e., $w \neq -1$, or model templates with varying neutrino masses for the HSC-Y3 $3 \times 2\text{pt}$ analysis to study whether the S_8 tension is relaxed. This requires a joint likelihood analysis of the HSC-Y3 $3 \times 2\text{pt}$ and the *Planck* data using such extended models.

We also showed that when our HSC-Y3 analysis is combined with the external BAO constraints on Ω_m with $\mathcal{N}(0.3, 0.01)$, the parameters are changed to $S_8 = 0.732^{+0.027}_{-0.029}$ and $\Delta z_{\text{ph}} = -0.133^{+0.077}_{-0.084}$. This result indicates a 2σ -level residual photo- z error, implying that the mean redshift of the HSC galaxies at $z \gtrsim 0.7$ is *higher* by $|\Delta z| = 0.133$ than implied by the photo- z estimates. Interestingly, such a large photo- z bias for the high-redshift HSC galaxies is also indicated in the companion works of the HSC cosmic shear tomography analyses (Li *et al.* [25] and Dalal *et al.* [26]). For the cosmic shear analyses, the photo- z 's at the high redshifts are calibrated by the cosmic shear amplitudes at different redshift bins, while the photo- z error in our method is calibrated by the combination of the galaxy-galaxy lensing signal of the spectroscopic SDSS galaxies and the cosmic shear.

We stress that we carried out all our cosmology analyses, the $3 \times 2\text{pt}$ analyses and the real- and Fourier-space cosmic shear analyses, using different blinded catalogs. We did not compare the cosmological constraints from the different methods during the blind analysis stage. After unblinding we found that all cosmological constraints are in agreement with each other, and also indicate a non-zero residual photo- z error for the high-redshift HSC galaxies. The significance of the non-zero residual photo- z error and the consistency tests of these HSC cosmology results are studied and presented in the upcoming paper, Sugiyama *et al.* (in prep.) using a mock analysis of these HSC cosmology analyses taking into account the cross covariances between the different observables. Thus the

HSC results might suggest an unknown systematic error in the photo- z estimates for high-redshift galaxies that are not calibrated out by the COSMOS data. The upcoming spectroscopic samples to be delivered from the DESI¹⁶ and PFS surveys [116] will be very powerful samples for calibrating these high-redshift photo- z 's using the clustering redshift method to higher redshifts, $z \gtrsim 1$ [88]. If we can constrain the photo- z systematics to the precision of $\sigma(\Delta z_{\text{ph}}) \sim \mathcal{O}(10^{-2})$, we can significantly improve the precision of our S_8 constraint even with the current HSC-Y3 data.

There are various directions to improve the cosmological constraints in this paper. First of all the cosmological analysis in this paper is based on the HSC Year 3 dataset of 416 deg^2 , which is about one-third of the full HSC dataset covering about $1,100 \text{ deg}^2$. Obviously, it is worth pursuing this possible S_8 -tension with the full HSC dataset. In addition, this paper uses the projected correlation function of the SDSS galaxies for the joint analysis. In this paper we intentionally did not include the BAO information or the redshift-space distortion (RSD) effect that can be measured from the redshift-space three-dimensional correlation function or power spectrum. Since the BAO and RSD information are very powerful probes of cosmological parameters, it would be very interesting to explore a full joint analysis of the galaxy-galaxy weak lensing, the cosmic shear and the redshift-space power spectrum. In our future work, we will do this, using an emulator-based method similar to that in this paper to model the redshift-space power spectrum of galaxies based on the redshift-space halo power spectrum and the halo occupation method [21]. Lastly, our method can be applied for the Stage-IV surveys, i.e., ground-based survey: Vera C. Rubin Observatory Legacy Survey of Space and Time [LSST; 117]), and space-based surveys: Euclid [118] and the Nancy Grace Roman space telescope [Roman; 119], by which statistical uncertainties will be significantly improved.

ACKNOWLEDGMENTS

This work was supported in part by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, and JSPS KAKENHI Grant Numbers JP18H04350, JP18H04358, JP19H00677, JP19K14767, JP20H00181, JP20H01932, JP20H04723, JP20H05850, JP20H05855, JP20H05856, JP20H05861, JP21J00011, JP21H05456, JP21J10314, JP21H01081, JP21H05456, JP22K03634, JP22K03655 and JP22K21349 by Japan Science and Technology Agency (JST) CREST JPMHCR1414, by JST AIP Acceleration Research Grant Number JP20317829, Japan, and by Basic Research Grant (Super AI) of Institute for AI and Beyond of the University of Tokyo. SS was supported in part by International Graduate Program for Excellence in Earth-Space Science (IGPEES), WINGS Program, the University of Tokyo. YK is supported in part by the David

and Lucile Packard foundation. RM is supported by a grant from the Simons Foundation (Simons Investigator in Astrophysics, Award ID 620789). RD acknowledges support from the NSF Graduate Research Fellowship Program under Grant No. DGE-2039656.

The Hyper Suprime-Cam (HSC) collaboration includes the astronomical communities of Japan and Taiwan, and Princeton University. The HSC instrumentation and software were developed by the National Astronomical Observatory of Japan (NAOJ), the Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), the University of Tokyo, the High Energy Accelerator Research Organization (KEK), the Academia Sinica Institute for Astronomy and Astrophysics in Taiwan (ASIAA), and Princeton University. Funding was contributed by the FIRST program from Japanese Cabinet Office, the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the Japan Society for the Promotion of Science (JSPS), Japan Science and Technology Agency (JST), the Toray Science Foundation, NAOJ, Kavli IPMU, KEK, ASIAA, and Princeton University. This paper makes use of software developed for the Large Synoptic Survey Telescope. We thank the LSST Project for making their code available as free software at <http://dm.lsst.org>

The Pan-STARRS1 Surveys (PS1) have been made possible through contributions of the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max-Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, Queen's University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under Grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation under Grant No. AST-1238877, the University of Maryland, and Eotvos Lorand University (ELTE) and the Los Alamos National Laboratory.

Based in part on data collected at the Subaru Telescope and retrieved from the HSC data archive system, which is operated by Subaru Telescope and Astronomy Data Center at National Astronomical Observatory of Japan.

Appendix A: Model Validation with Mock Galaxy Catalogs

In this section we describe the validation tests of our modeling and analysis methods that we perform as one of the blinding criterion. Table IV summarizes the synthetic data vectors used for validation tests. For $\Delta\Sigma$ and w_p we use the same data vector as described in Miyatake *et al.* [15] except for the synthetic data labelled as “ $\Delta z_{\text{ph}} = -0.2$ ” and “ $\Delta z_{\text{ph}} = -0.2^\dagger$ ”. For ξ_{\pm} , we use the data vector described in Sugiyama *et al.* [24]. Note that we take extreme cases for the baryonic effect on ξ_{\pm} by setting $A_{\text{bary}} = 1.6$ and $T_{\text{AGN}} = 7.3$. The detailed procedures to generate the syn-

¹⁶ <https://www.desi.lbl.gov>

thetic data vectors for “ $\Delta z_{\text{ph}}^{\text{in}} = -0.2$ ” and “ $\Delta z_{\text{ph}}^{\text{in}} = -0.2^\dagger$,” assuming that the estimated redshift distribution of source galaxies is systematically lower than the true distribution by $|\Delta z_{\text{ph}}^{\text{in}}| = 0.2$ are described in Appendix A of Sugiyama *et al.* [24]. Here $\Delta z_{\text{ph}}^{\text{in}} = -0.2$ is about 2σ away from the central value of Δz_{ph} in our fiducial analysis, i.e. $\Delta z_{\text{ph}} = -0.05 \pm 0.09$. Hence this validation test gives the worst case scenario for the impact of residual photo- z error. Note that the validation test for “ $\Delta z_{\text{ph}}^{\text{in}} = -0.2^\dagger$ ” assumes the informative prior of Δz_{ph} given by $\mathcal{N}(0, 0.1)$, aimed at studying how the informative prior gives a biased estimate in the cosmological parameters in the presence of the photo- z bias given by $\Delta z_{\text{ph}}^{\text{in}} = -0.2$. We apply the baseline analysis pipeline to each of the synthetic data vector to estimate the cosmological parameters using the covariance matrix for the HSC-Y3 data.

Fig. 10 shows the summary of the validation tests. As described in the main text, we do not find any significant deviation from the input cosmological parameters, except for “assembly- b -ext,” “assembly- b ,” and “ $\Delta z_{\text{ph}} = -0.2^\dagger$.” In “assembly- b -ext” and “assembly- b ,” we assume the large assembly bias amplitudes, so these tests give the worst case scenario in our cosmological constraints if the SDSS galaxies are affected by such large assembly bias effects, although there has been no detection of assembly bias for actual SDSS galaxies. As described in Section VIB, a possible assembly bias signature can be identified from actual data analysis: if the assembly bias effect exists, we expect that using the different scale cuts of w_p and $\Delta\Sigma$ in the cosmology analysis would lead to a systematic shift in S_8 . For example, if we employ the sufficiently large scale cuts such as $R = (8, 12)h^{-1}\text{Mpc}$ for w_p and $\Delta\Sigma$, where the linear theory or perturbation theory model is valid, the cosmological parameters are safely recovered. For the actual SDSS data, we did not observe such a systematic shift in S_8 , so we concluded that the SDSS galaxies do not display any evidence of the assembly bias effect.

The row of “ $\Delta z_{\text{ph}}^{\text{in}} = -0.2^\dagger$ ” in Fig. 10 shows a significant bias in S_8 by $> 1\sigma$ if we employ an informative Gaussian prior on Δz_{ph} given by $\mathcal{N}(0, 0.1)$ even when the photo- z bias given by $\Delta z_{\text{ph}} = -0.2$ exists. On the other hand, the row of “ $\Delta z_{\text{ph}}^{\text{in}} = -0.2^*$ ” shows that the input cosmological parameters are safely recovered if we employ an uninformative, flat prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$ as our baseline analysis. Note that the photo- z bias introduced in the data vector is also recovered in this case. This means that the HSC-Y3 data has a calibration power of Δz_{ph} to the precision of $\sigma(\Delta z_{\text{ph}}) \sim 0.1$. Based on these findings, we decided to implement an analysis setup that uses the uninformative, flat prior of Δz_{ph} .

The row of “4th-order PSF” shows the results when including the fourth-moment PSF leakage and fourth-moment PSF modeling error in the synthetic data of ξ_\pm . As described in Sugiyama *et al.* [24], we used the method in Zhang *et al.* [91] to measure the fourth-moment PSF leakage and fourth-moment PSF modeling errors from the HSC galaxy sample used in this paper. Then we include the measured PSF systematics contamination in the synthetic data of ξ_\pm and then

apply the baseline analysis pipeline to the synthetic data vector including w_p and $\Delta\Sigma$. The result shows that the cosmological constraints are not affected by the PSF systematics. The impact is smaller than that found from the cosmic shear analyses of HSC-Y3 data [25, 26], because the constraining power in our 3×2 pt analysis is mainly from the clustering information of SDSS galaxies, not from the cosmic shear signal.

Appendix B: Details of Internal Consistency Tests

In this section we show the results for various consistency tests for the different analysis setups and/or different subsets of the data vector, as listed in Table III. Table V shows the mean and 68% upper and lower credible intervals with the MAP in parenthesis, i.e., the numbers plotted in Fig. 4. Figs. 11–20 show the 1- or 2-dimensional posterior distributions of the different analysis setups in which the same kind of consistency tests are grouped and compared with the fiducial 3×2 pt analysis.

Appendix C: Posterior Distributions of All Parameters

Fig. 23 shows the posterior distribution of all parameters sampled in our baseline analysis.

Appendix D: Robustness of Parameter Sampling

1. Nestcheck

In this section, we present the results of the convergence test of MultiNest sampling for the baseline 3×2 pt analysis. We use the nestcheck diagnostic to test the convergence of the MultiNest chain, implemented as nestcheck[121]. Fig. 24 shows the result of the convergence test by nestcheck for the main cosmological parameters, Ω_m , σ_8 , and S_8 . In the top right panel, we can see that the chain covers sufficient posterior volume. The left panels show the uncertainty of the posterior distributions, estimated by bootstrapping the original MultiNest chain, and indicating that our estimate of the posterior distributions is robust.

2. Sampler difference

As an additional test of convergence of our parameter estimates, we compare the result of the nested sampling by MultiNest to the result with the standard Metropolis algorithm in Fig. 25. The difference between the posterior estimates is almost negligible (the difference in mode and 68% credible interval is $\sim 2\%$ for S_8), and thus we conclude that our parameter inference by MultiNest is robust.

TABLE IV. A summary of mock signals used for the validation tests (see Miyatake *et al.* [104] for the details). All the mock catalogs, except for “cent-incomp.” and “FoF-halo” catalogs, have the same HOD in the average sense, but use the different ways to populate galaxies into halos in N -body simulations. The column “satellite gals.” denotes a model of the spatial distribution of satellite galaxies in the host halo. In the columns of $\Delta\Sigma$, w_p , and ξ_{\pm} , “ \checkmark ” or “ $-$ ” denote whether they are modified from the fiducial mock or not, respectively. Note that the “assembly- b ” and “assembly- b -ext” are the worst-case scenarios, where we implemented the overwhelmingly large assembly bias in the sense that the catalogs give the larger clustering amplitudes in w_p than those of the fiducial mocks (where the halo bias is simply given by the host halo mass) by a factor of 1.3 and 1.5 for the mock LOWZ, CMASS1 and CMASS2 galaxies (Fig. 5 of Miyatake *et al.* [104]).

setup label	HOD	satellite gals.	$\Delta\Sigma$	w_p	ξ_{\pm}	description
3×2 pt	fid.	NFW	-	-	-	fiducial model
2×2 pt*	fid.	NFW	-	-	N/A	without ξ_{\pm} , using Δz_{ph} posterior from 3×2 pt analysis as a prior
cosmic shear*	N/A	N/A	N/A	N/A	-	without $\Delta\Sigma$ and w_p , using Δz_{ph} posterior from 3×2 pt analysis as a prior
nonfidNsat	fid.	NFW	\checkmark	\checkmark	-	populate satellites irrespectively of centrals
sat-dm-dist	fid.	DM part.	\checkmark	\checkmark	-	populate satellites according to N -body particles
sat-sub	fid.	subhalos	\checkmark	\checkmark	-	populate satellites according to subhalos
off-cent1	fid.	NFW	\checkmark	\checkmark	-	all centrals off-centered, with Gaussian profile
off-cent2	fid.	NFW	\checkmark	\checkmark	-	a fraction (0.34) of “off-centered” centrals, assuming Gaussian profile
off-cent3	fid.	NFW	\checkmark	\checkmark	-	similar to “off-cent1”, but with NFW profile
off-cent4	fid.	NFW	\checkmark	\checkmark	-	similar to “off-cent2”, but with NFW profile
baryon	fid.	NFW	\checkmark	-	-	mimic the baryonic effect of <i>Illustris</i> on the halo mass profile
assembly- b -ext	fid.	NFW	\checkmark	\checkmark	-	populate galaxies according to concentrations of host halos
assembly- b	fid.	NFW	\checkmark	\checkmark	-	similar to “assembly- b -ext”, but introduce scatters
cent-incomp.	$\langle N_c \rangle$ mod.	NFW	\checkmark	\checkmark	-	include an “incomplete” selection of centrals
FoF-halo	mod.	FoF halos	-	-	\checkmark	use FoF halos to populate galaxies
HMCode v2015	fid.	NFW	-	-	\checkmark	ξ_{\pm} is generated by HMCode v2015 with $A_{\text{bary}} = 1.6$ or 2.8
HMCode v2020	fid.	NFW	-	-	\checkmark	ξ_{\pm} is generated by HMCode v2020 with $T_{\text{AGN}} = 7.3$ or 8.3
$\Delta z_{\text{ph}}^{\text{in}} = -0.2$	fid.	NFW	\checkmark	-	\checkmark	$\Delta\Sigma$ and ξ_{\pm} with $\Delta z_{\text{ph}}^{\text{in}} = -0.2$, analyzed with a prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$
$\Delta z_{\text{ph}}^{\text{in}} = -0.2^{\dagger}$	fid.	NFW	\checkmark	-	\checkmark	$\Delta\Sigma$ and ξ_{\pm} with $\Delta z_{\text{ph}}^{\text{in}} = -0.2$, analyzed with a prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$
4th-order PSF	fid.	NFW	-	-	\checkmark	Include the 4th-order moment PSF systematics into ξ_{\pm}

- [1] S. Dodelson and F. Schmidt, *Modern Cosmology* (2020).
- [2] D. H. Weinberg, M. J. Mortonson, D. J. Eisenstein, C. Hirata, A. G. Riess, and E. Rozo, Observational probes of cosmic acceleration, *Phys. Rep.* **530**, 87 (2013), arXiv:1201.2434.
- [3] H. Aihara, N. Arimoto, R. Armstrong, S. Arnouts, N. A. Bahcall, S. Bickerton, J. Bosch, K. Bundy, P. L. Capak, J. H. H. Chan, *et al.*, The Hyper Suprime-Cam SSP Survey: Overview and survey design, *Publ. Astron. Soc. Japan* **70**, S4 (2018), arXiv:1704.05858 [astro-ph.IM].
- [4] C. Hikage, M. Oguri, T. Hamana, S. More, R. Mandelbaum, M. Takada, F. Köhlinger, H. Miyatake, A. J. Nishizawa, H. Aihara, R. Armstrong, J. Bosch, J. Coupon, A. Ducout, *et al.*, Cosmology from cosmic shear power spectra with Subaru Hyper Suprime-Cam first-year data, *Publ. Astron. Soc. Japan* **71**, 43 (2019), arXiv:1809.09148 [astro-ph.CO].
- [5] T. Hamana, M. Shirasaki, S. Miyazaki, C. Hikage, M. Oguri, S. More, R. Armstrong, A. Leauthaud, R. Mandelbaum, H. Miyatake, A. J. Nishizawa, M. Simet, M. Takada, H. Aihara, *et al.*, Cosmological constraints from cosmic shear two-point correlation functions with HSC survey first-year data, *Publ. Astron. Soc. Japan* **72**, 16 (2020), arXiv:1906.06041 [astro-ph.CO].
- [6] H. Miyatake, S. Sugiyama, M. Takada, T. Nishimichi, M. Shirasaki, Y. Kobayashi, R. Mandelbaum, S. More, M. Oguri, K. Osato, *et al.*, Cosmological inference from an emulator based halo model. II. Joint analysis of galaxy-galaxy weak lensing and galaxy clustering from HSC-Y1 and SDSS, *Phys. Rev. D* **106**, 083520 (2022), arXiv:2111.02419 [astro-ph.CO].
- [7] S. Sugiyama, M. Takada, H. Miyatake, T. Nishimichi, M. Shirasaki, Y. Kobayashi, R. Mandelbaum, S. More, R. Takahashi, K. Osato, *et al.*, HSC Year 1 cosmology results with the minimal bias method: HSC xBOSS galaxy-galaxy weak lensing and BOSS galaxy clustering, *Phys. Rev. D* **105**, 123537 (2022), arXiv:2111.10966 [astro-ph.CO].
- [8] T. M. C. Abbott, M. Aguena, A. Alarcon, S. Allam, O. Alves, A. Amon, F. Andrade-Oliveira, J. Annis, S. Avila, D. Bacon, *et al.*, Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and weak lensing, *Phys. Rev. D* **105**, 023520 (2022), arXiv:2105.13549 [astro-ph.CO].
- [9] C. Heymans, T. Tröster, M. Asgari, C. Blake, H. Hildebrandt, B. Joachimi, K. Kuijken, C.-A. Lin, A. G. Sánchez, J. L. van den Busch, *et al.*, KiDS-1000 Cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering constraints, *Astronomy & Astrophysics* **646**, A140 (2021), arXiv:2007.15632 [astro-ph.CO].
- [10] E. Abdalla, G. F. Abellán, A. Aboubrahim, A. Agnello, Ö. Akarsu, Y. Akrami, G. Alestas, D. Aloni, L. Amendola, L. A. Anchordoqui, *et al.*, Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies, *Journal of High Energy Astrophysics* **34**, 49 (2022), arXiv:2203.06142 [astro-ph.CO].
- [11] Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday,




FIG. 10. Summary of model validation tests for the 3×2 pt analysis. Constraints on cosmological parameters Ω_m , σ_8 , and S_8 and the photo- z shift parameter Δz_{ph} are shown in each panel from left to right. The input cosmological parameters used for making synthetic data vectors are indicated by the vertical dashed lines in the panels. The shaded regions are the 68% confidence intervals from the constraints in the fiducial “ 3×2 pt” case.

- R. B. Barreiro, N. Bartolo, S. Basak, R. Battye, K. Benabed, J. P. Bernard, *et al.*, Planck 2018 results. VI. Cosmological parameters, *Astronomy & Astrophysics* **641**, A6 (2020), arXiv:1807.06209 [astro-ph.CO].
- [12] A. J. Nishizawa, B.-C. Hsieh, M. Tanaka, and T. Takata, Photometric Redshifts for the Hyper Suprime-Cam Subaru Strategic Program Data Release 2, arXiv e-prints, arXiv:2003.01511 (2020), arXiv:2003.01511 [astro-ph.GA].
- [13] N. Kaiser, On the spatial correlations of Abell clusters., *Astrophys. J. Lett.* **284**, L9 (1984).
- [14] M. Oguri and M. Takada, Combining cluster observables and stacked weak lensing to probe dark energy: Self-calibration of systematic uncertainties, *Phys. Rev. D* **83**, 023008 (2011), arXiv:1010.0744 [astro-ph.CO].
- [15] H. Miyatake, Y. Kobayashi, M. Takada, T. Nishimichi, M. Shirasaki, S. Sugiyama, R. Takahashi, K. Osato, S. More, and Y. o. Park, Cosmological inference from an emulator based halo model. I. Validation tests with HSC and SDSS mock catalogs, *Phys. Rev. D* **106**, 083519 (2022), arXiv:2101.00113 [astro-ph.CO].
- [16] U. Seljak, Analytic model for galaxy and dark matter clustering, *Mon. Not. Roy. Astron. Soc.* **318**, 203 (2000), arXiv:astro-ph/0001493.
- [17] C. Ma and J. N. Fry, Deriving the Nonlinear Cosmological Power Spectrum and Bispectrum from Analytic Dark Matter Halo Profiles and Mass Functions, *Astrophys. J.* **543**, 503 (2000), arXiv:astro-ph/0003343.
- [18] J. A. Peacock and R. E. Smith, Halo occupation numbers and galaxy bias, *Mon. Not. Roy. Astron. Soc.* **318**, 1144 (2000), arXiv:astro-ph/0005010.
- [19] R. Scoccimarro, R. K. Sheth, L. Hui, and B. Jain, How Many Galaxies Fit in a Halo? Constraints on Galaxy Formation Efficiency from Spatial Clustering, *Astrophys. J.* **546**, 20 (2001), arXiv:astro-ph/0006319 [astro-ph].
- [20] T. Nishimichi, M. Takada, R. Takahashi, K. Osato, M. Shirasaki, T. Oogi, H. Miyatake, M. Oguri, R. Murata, Y. Kobayashi, *et al.*, Dark Quest. I. Fast and Accurate Emulation of Halo Clustering Statistics and Its Application to Galaxy Clustering, *Astrophys. J.* **884**, 29 (2019), arXiv:1811.09504 [astro-ph.CO].
- [21] Y. Kobayashi, T. Nishimichi, M. Takada, and H. Miyatake, Full-shape cosmology analysis of the SDSS-III BOSS galaxy power spectrum using an emulator-based halo model: A 5% determination of σ_8 , *Phys. Rev. D* **105**, 083517 (2022), arXiv:2110.06969 [astro-ph.CO].
- [22] C. Heymans, T. Tröster, M. Asgari, C. Blake, H. Hildebrandt, B. Joachimi, K. Kuijken, C.-A. Lin, A. G. Sánchez, J. L. van den Busch, *et al.*, KiDS-1000 Cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering constraints, *Astronomy & Astrophysics* **646**, A140 (2021), arXiv:2007.15632 [astro-ph.CO].
- [23] S. More *et al.*, (in prep.).
- [24] S. Sugiyama *et al.*, (in prep.).
- [25] X. Li *et al.*, (in prep.).

TABLE V. Summary of the main cosmological parameters constrained in this work, Ω_m , σ_8 , and S_8 . The estimates are presented in the format of mode $^{+68\% \text{ upper}}_{-68\% \text{ lower}}$ (MAP, mean). The analysis setup for each row is summarized in Table III.

	Ω_m	σ_8	S_8
$3 \times 2\text{pt}$	$0.382^{+0.031}_{-0.047}$ (0.401, 0.367)	$0.685^{+0.035}_{-0.026}$ (0.696, 0.696)	$0.763^{+0.040}_{-0.036}$ (0.805, 0.768)
$2 \times 2\text{pt}^*$	$0.397^{+0.025}_{-0.040}$ (0.413, 0.382)	$0.683^{+0.030}_{-0.024}$ (0.678, 0.693)	$0.776^{+0.032}_{-0.027}$ (0.796, 0.780)
cosmic shear *	$0.380^{+0.095}_{-0.089}$ (0.454, 0.375)	$0.632^{+0.103}_{-0.066}$ (0.623, 0.674)	$0.735^{+0.039}_{-0.040}$ (0.767, 0.737)
$3 \times 2\text{pt}$, $R_{\min} = (4, 6) h^{-1}\text{Mpc}^*$	$0.312^{+0.044}_{-0.040}$ (0.347, 0.318)	$0.759^{+0.056}_{-0.050}$ (0.753, 0.767)	$0.785^{+0.028}_{-0.028}$ (0.809, 0.785)
$3 \times 2\text{pt}$, $R_{\min} = (8, 12) h^{-1}\text{Mpc}^*$	$0.355^{+0.037}_{-0.053}$ (0.405, 0.341)	$0.690^{+0.070}_{-0.035}$ (0.660, 0.721)	$0.760^{+0.035}_{-0.029}$ (0.767, 0.763)
$3 \times 2\text{pt}$, w/o LOWZ	$0.374^{+0.031}_{-0.054}$ (0.398, 0.357)	$0.708^{+0.040}_{-0.032}$ (0.684, 0.718)	$0.785^{+0.043}_{-0.053}$ (0.788, 0.781)
$3 \times 2\text{pt}$, w/o CMASS1	$0.380^{+0.030}_{-0.048}$ (0.415, 0.367)	$0.670^{+0.031}_{-0.029}$ (0.658, 0.675)	$0.742^{+0.041}_{-0.038}$ (0.774, 0.745)
$3 \times 2\text{pt}$, w/o CMASS2	$0.364^{+0.041}_{-0.039}$ (0.447, 0.364)	$0.690^{+0.036}_{-0.030}$ (0.682, 0.699)	$0.764^{+0.036}_{-0.036}$ (0.833, 0.767)
$2 \times 2\text{pt}$, w/o LOWZ *	$0.389^{+0.028}_{-0.049}$ (0.400, 0.374)	$0.700^{+0.039}_{-0.030}$ (0.718, 0.713)	$0.792^{+0.034}_{-0.035}$ (0.830, 0.793)
$2 \times 2\text{pt}$, w/o CMASS1 *	$0.381^{+0.032}_{-0.047}$ (0.402, 0.371)	$0.677^{+0.036}_{-0.028}$ (0.653, 0.688)	$0.760^{+0.032}_{-0.031}$ (0.756, 0.762)
$2 \times 2\text{pt}$, w/o CMASS2 *	$0.358^{+0.055}_{-0.035}$ (0.423, 0.369)	$0.694^{+0.039}_{-0.029}$ (0.673, 0.705)	$0.774^{+0.033}_{-0.032}$ (0.800, 0.779)
no photo- z error	$0.394^{+0.023}_{-0.031}$ (0.394, 0.383)	$0.690^{+0.026}_{-0.021}$ (0.693, 0.699)	$0.791^{+0.018}_{-0.020}$ (0.794, 0.788)
no shear error	$0.383^{+0.027}_{-0.044}$ (0.366, 0.369)	$0.687^{+0.028}_{-0.025}$ (0.693, 0.694)	$0.765^{+0.038}_{-0.035}$ (0.765, 0.768)
fix mag. bias	$0.381^{+0.029}_{-0.045}$ (0.407, 0.367)	$0.686^{+0.032}_{-0.025}$ (0.688, 0.696)	$0.766^{+0.037}_{-0.035}$ (0.801, 0.769)
no PSF error	$0.393^{+0.024}_{-0.043}$ (0.408, 0.375)	$0.684^{+0.029}_{-0.025}$ (0.681, 0.690)	$0.768^{+0.037}_{-0.035}$ (0.794, 0.770)
no IA	$0.374^{+0.032}_{-0.040}$ (0.358, 0.366)	$0.688^{+0.031}_{-0.024}$ (0.692, 0.696)	$0.766^{+0.035}_{-0.036}$ (0.756, 0.767)
extreme IA	$0.385^{+0.027}_{-0.046}$ (0.373, 0.369)	$0.687^{+0.029}_{-0.024}$ (0.684, 0.694)	$0.766^{+0.037}_{-0.037}$ (0.763, 0.768)
$3 \times 2\text{pt}^\dagger$	$0.389^{+0.025}_{-0.036}$ (0.394, 0.378)	$0.689^{+0.025}_{-0.025}$ (0.692, 0.694)	$0.773^{+0.031}_{-0.027}$ (0.794, 0.777)
$2 \times 2\text{pt}^\dagger$	$0.397^{+0.024}_{-0.042}$ (0.355, 0.381)	$0.690^{+0.030}_{-0.025}$ (0.728, 0.699)	$0.779^{+0.035}_{-0.028}$ (0.792, 0.786)
cosmic shear †	$0.451^{+0.089}_{-0.108}$ (0.509, 0.387)	$0.624^{+0.113}_{-0.063}$ (0.611, 0.688)	$0.756^{+0.044}_{-0.043}$ (0.795, 0.760)
$2 \times 2\text{pt}$	$0.375^{+0.029}_{-0.039}$ (0.371, 0.364)	$0.655^{+0.028}_{-0.030}$ (0.652, 0.661)	$0.719^{+0.039}_{-0.034}$ (0.725, 0.727)
cosmic shear	$0.228^{+0.164}_{-0.070}$ (0.280, 0.299)	$0.655^{+0.135}_{-0.109}$ (0.710, 0.691)	$0.624^{+0.094}_{-0.070}$ (0.686, 0.660)
XMM ($\sim 33 \text{ deg}^2$) *	$0.356^{+0.031}_{-0.044}$ (0.361, 0.345)	$0.641^{+0.037}_{-0.033}$ (0.647, 0.642)	$0.693^{+0.050}_{-0.055}$ (0.710, 0.688)
GAMA15H ($\sim 41 \text{ deg}^2$) *	$0.344^{+0.042}_{-0.035}$ (0.368, 0.347)	$0.708^{+0.046}_{-0.040}$ (0.707, 0.714)	$0.768^{+0.046}_{-0.047}$ (0.782, 0.766)
HECTOMAP ($\sim 43 \text{ deg}^2$) *	$0.382^{+0.029}_{-0.043}$ (0.398, 0.370)	$0.708^{+0.043}_{-0.035}$ (0.725, 0.717)	$0.794^{+0.045}_{-0.042}$ (0.835, 0.795)
GAMA09H ($\sim 78 \text{ deg}^2$) *	$0.387^{+0.029}_{-0.042}$ (0.411, 0.375)	$0.681^{+0.042}_{-0.029}$ (0.663, 0.693)	$0.773^{+0.044}_{-0.041}$ (0.776, 0.774)
VVDS ($\sim 96 \text{ deg}^2$) *	$0.356^{+0.030}_{-0.043}$ (0.373, 0.346)	$0.650^{+0.034}_{-0.030}$ (0.643, 0.659)	$0.704^{+0.038}_{-0.037}$ (0.717, 0.705)
WIDE12H ($\sim 121 \text{ deg}^2$) *	$0.333^{+0.045}_{-0.030}$ (0.326, 0.340)	$0.698^{+0.039}_{-0.034}$ (0.735, 0.706)	$0.744^{+0.038}_{-0.029}$ (0.765, 0.749)
DEMPZ & WX	$0.374^{+0.030}_{-0.040}$ (0.403, 0.364)	$0.682^{+0.030}_{-0.024}$ (0.673, 0.690)	$0.758^{+0.034}_{-0.038}$ (0.780, 0.757)
MIZUKI	$0.383^{+0.028}_{-0.039}$ (0.371, 0.370)	$0.678^{+0.024}_{-0.023}$ (0.681, 0.682)	$0.757^{+0.031}_{-0.033}$ (0.757, 0.756)
DNNZ	$0.381^{+0.031}_{-0.044}$ (0.390, 0.368)	$0.696^{+0.043}_{-0.026}$ (0.710, 0.711)	$0.784^{+0.042}_{-0.040}$ (0.810, 0.786)
w/o star weight	$0.371^{+0.029}_{-0.041}$ (0.382, 0.360)	$0.690^{+0.031}_{-0.026}$ (0.679, 0.698)	$0.765^{+0.033}_{-0.039}$ (0.766, 0.763)
offcentering	$0.386^{+0.026}_{-0.039}$ (0.378, 0.373)	$0.685^{+0.028}_{-0.022}$ (0.682, 0.691)	$0.767^{+0.037}_{-0.032}$ (0.766, 0.770)
incompleteness	$0.381^{+0.027}_{-0.041}$ (0.406, 0.368)	$0.694^{+0.030}_{-0.024}$ (0.679, 0.701)	$0.776^{+0.032}_{-0.037}$ (0.790, 0.775)
$\sigma(\Delta m) = 0.1$ prior	$0.381^{+0.028}_{-0.044}$ (0.401, 0.368)	$0.685^{+0.032}_{-0.024}$ (0.701, 0.694)	$0.765^{+0.036}_{-0.037}$ (0.811, 0.767)
$\sigma(\Delta z_{\text{ph}}) = 0.2$ prior	$0.387^{+0.025}_{-0.045}$ (0.395, 0.372)	$0.683^{+0.031}_{-0.020}$ (0.697, 0.693)	$0.768^{+0.032}_{-0.034}$ (0.800, 0.769)
2 cosmo paras	$0.359^{+0.018}_{-0.019}$ (0.367, 0.356)	$0.691^{+0.023}_{-0.021}$ (0.703, 0.695)	$0.756^{+0.031}_{-0.028}$ (0.778, 0.757)

[26] R. Dalal *et al.*, (in prep.).

[27] S. Miyazaki, Y. Komiyama, S. Kawanomoto, Y. Doi, H. Furusawa, T. Hamana, Y. Hayashi, H. Ikeda, Y. Kamata, H. Karoji, M. Koike, T. Kurakami, S. Miyama, T. Morokuma, *et al.*, Hyper Suprime-Cam: System design and verification of image quality, Publ. Astron. Soc. Japan **70**, S1 (2018).

[28] Y. Komiyama, Y. Obuchi, H. Nakaya, Y. Kamata, S. Kawanomoto, Y. Utsumi, S. Miyazaki, F. Uraguchi, H. Furusawa, T. Morokuma, T. Uchida, H. Miyatake, S. Mineo, H. Fujimori, *et al.*, Hyper Suprime-Cam: Camera dewar design, Publ. Astron. Soc. Japan **70**, S2 (2018).

[29] H. Furusawa, M. Koike, T. Takata, Y. Okura, H. Miyatake, R. H. Lupton, S. Bickerton, P. A. Price, J. Bosch, N. Yasuda,




FIG. 11. The posterior distributions for baseline “3×2pt,” “2×2pt*,” and “cosmic shear*” analysis setups in Table III. The contours show the 68% and 96% credible intervals. The constraints from “2×2pt*” and “cosmic shear*” are consistent with the “3×2pt” analysis. Note that since we use the prior on the residual photo- z errors (Δz_{ph}) for “2×2pt*” and “cosmic shear*” derived from the “3×2pt” analysis, the constraining power of “2×2pt*” is similar to “3×2pt,” but that of “cosmic shear*” is still weaker than “3×2pt.” For the constraints from the 2×2pt and cosmic shear analysis with the uniform prior on Δz_{ph} , see the left panel in Fig. 6.




FIG. 12. Similar to Fig. 11, but the results from “3×2pt, w/o LOWZ,” “3×2pt, w/o CMASS1,” and “3×2pt, w/o CMASS2” analysis setups in Table III. We do not see any significant changes in the cosmological constraints from the baseline “3×2pt” analysis when excluding one of the lens samples in our analysis.




FIG. 13. Similar to Fig. 11, but the results from “2×2pt, w/o LOWZ*,” “2×2pt, w/o CMASS1*,” and “2×2pt, w/o CMASS2*” analysis setups in Table III. We do not see any significant changes in the cosmological constraints from the baseline “3×2pt” analysis when excluding one of the lens samples in our analysis.

S. Mineo, Y. Yamada, S. Miyazaki, F. Nakata, *et al.*, The on-site quality-assurance system for Hyper Suprime-Cam: OS-QAH, Publ. Astron. Soc. Japan **70**, S3 (2018).

- [30] S. Kawanomoto, F. Uruguchi, Y. Komiyama, S. Miyazaki, H. Furusawa, F. Finet, T. Hattori, S.-Y. Wang, N. Yasuda, and N. Suzuki, Hyper Suprime-Cam: Filters, Publ. Astron. Soc. Japan **70**, 66 (2018).
- [31] X. Li, H. Miyatake, W. Luo, S. More, M. Oguri, T. Hamana, R. Mandelbaum, M. Shirasaki, M. Takada, R. Armstrong, *et al.*, The three-year shear catalog of the Subaru Hyper Suprime-Cam SSP Survey, Publ. Astron. Soc. Japan **74**, 421 (2022), arXiv:2107.00136 [astro-ph.CO].
- [32] A. Nishizawa *et al.*, (in prep.).
- [33] J. Bosch, R. Armstrong, S. Bickerton, H. Furusawa, H. Ikeda, M. Koike, R. Lupton, S. Mineo, P. Price, T. Takata, *et al.*, The Hyper Suprime-Cam software pipeline, Publ. Astron. Soc. Japan **70**, S5 (2018), arXiv:1705.06766 [astro-ph.IM].
- [34] R. Mandelbaum, H. Miyatake, T. Hamana, M. Oguri, M. Simet, R. Armstrong, J. Bosch, R. Murata, F. Lanusse, A. Leauthaud, *et al.*, The first-year shear catalog of the Subaru Hyper Suprime-Cam Subaru Strategic Program Survey, Publ. Astron. Soc. Japan **70**, S25 (2018), arXiv:1705.06745 [astro-ph.CO].
- [35] R. Mandelbaum, F. Lanusse, A. Leauthaud, R. Armstrong, M. Simet, H. Miyatake, J. E. Meyers, J. Bosch, R. Murata, S. Miyazaki, *et al.*, Weak lensing shear calibration with simu-




FIG. 14. Similar to Fig. 11, but the results from “no photo- z error,” “no shear error,” and “fix mag. bias” analysis setups in Table III. When the source redshift distribution is fixed, corresponding to the case “no photo error”, the credible interval of the cosmological parameters are significantly tightened, but the central values are shifted. In particular, fixing the source redshift distribution as indicated from the photo- z estimates rather than using the flat prior $\Pi(\Delta z_{\text{ph}}) = \mathcal{U}(-1, 1)$ in our baseline analysis leads to a higher value of S_8 than that of the baseline analysis. We do not see any significant changes in the cosmological constraints for the other cases.




FIG. 15. Similar to Fig. 11, but the results from “no PSF error,” “no IA error,” and “extreme IA” analysis setups in Table III. For these setups, we do not see any significant changes in the cosmological constraints from the baseline analysis.

lations of the HSC survey, *Mon. Not. Roy. Astron. Soc.* **481**, 3170 (2018), arXiv:1710.00885 [astro-ph.CO].

- [36] M. Tanaka, Photometric Redshift with Bayesian Priors on Physical Properties of Galaxies, *Astrophys. J.* **801**, 20 (2015), arXiv:1501.02047.
- [37] B. C. Hsieh and H. K. C. Yee, Estimating Luminosities and Stellar Masses of Galaxies Photometrically without Determining Redshifts, *Astrophys. J.* **792**, 102 (2014), arXiv:1407.5151.
- [38] M. Oguri, A cluster finding algorithm based on the multiband identification of red sequence galaxies, *Mon. Not. Roy. Astron. Soc.* **444**, 147 (2014), arXiv:1407.4693 [astro-ph.CO].
- [39] E. Medezinski, M. Oguri, A. J. Nishizawa, J. S. Speagle, H. Miyatake, K. Umetsu, A. Leauthaud, R. Murata, R. Mandelbaum, C. Sifón, *et al.*, Source selection for cluster weak lensing measurements in the Hyper Suprime-Cam survey, *Publ. Astron. Soc. Japan* **70**, 30 (2018), arXiv:1706.00427 [astro-ph.CO].
- [40] H. Miyatake, N. Battaglia, M. Hilton, E. Medezinski, A. J. Nishizawa, S. More, S. Aiola, N. Bahcall, J. R. Bond, E. Calabrese, S. K. Choi, M. J. Devlin, J. Dunkley, R. Dunner, *et al.*, Weak-lensing Mass Calibration of ACTPol Sunyaev-Zel’dovich Clusters with the Hyper Suprime-Cam Survey, *Astrophys. J.* **875**, 63 (2019), arXiv:1804.05873 [astro-ph.CO].
- [41] S. Alam, F. D. Albareti, C. Allende Prieto, F. Anders, S. F. Anderson, T. Anderton, B. H. Andrews, E. Armengaud, É. Aubourg, S. Bailey, *et al.*, The Eleventh and




FIG. 16. Similar to Fig. 11, but the results from “ $3 \times 2\text{pt}^\dagger$,” “ $2 \times 2\text{pt}^\dagger$,” and “cosmic shear † ” analysis setups in Table III. Here the analyses with superscript † uses the Gaussian prior on the photo- z error prior parameter, $\Pi(\Delta z_{\text{ph}}) = \mathcal{N}(0, 0.1)$. The use of the Gaussian prior still affects the S_8 constraints for the “ $3 \times 2\text{pt}^\dagger$,” “ $2 \times 2\text{pt}^\dagger$ ” analyses.




FIG. 17. Similar to Fig. 11, but the results from “XMM ($\sim 33 \text{ deg}^2$)*,” “GAMA15H ($\sim 41 \text{ deg}^2$)*,” “HECTOMAP ($\sim 43 \text{ deg}^2$)*,” “GAMA09H ($\sim 78 \text{ deg}^2$)*,” “VVDS ($\sim 96 \text{ deg}^2$)*,” and “WIDE12H ($\sim 121 \text{ deg}^2$)*” analysis setups in Table III. Given that the analyses of each field are almost uncorrelated, the cosmological constraints are consistent with each other.




FIG. 18. Similar to Fig. 11, but the results from “DEMPZ & WX,” “MIZUKI,” and “DNNz” analysis setups in Table III. “The DNNz” analysis exhibits the largest deviation in S_8 , but it is still $\sim 0.5\sigma$.




FIG. 19. Similar to Fig. 11, but the results from “w/o star weight” analysis setup in Table III. We do not see any significant changes in the cosmological constraints in the “w/o star weight” analysis.

Twelfth Data Releases of the Sloan Digital Sky Survey: Final Data from SDSS-III, *Astrophys. J. Suppl.* **219**, 12 (2015), arXiv:1501.00963 [astro-ph.IM].

- [42] K. S. Dawson, D. J. Schlegel, C. P. Ahn, S. F. Anderson, É. Aubourg, S. Bailey, R. H. Barkhouser, J. E. Bautista, A. r. Beifiori, A. A. Berlind, V. Bhardwaj, D. Bizyaev, C. H. Blake, *et al.*, The Baryon Oscillation Spectroscopic Survey of SDSS-III, *The Astronomical J.* **145**, 10 (2013), arXiv:1208.0022 [astro-ph.CO].
- [43] H. Miyatake, S. More, R. Mandelbaum, M. Takada, D. N. Spergel, J.-P. Kneib, D. P. Schneider, J. Brinkmann, and J. R. Brownstein, The Weak Lensing Signal and the Clustering of BOSS Galaxies. I. Measurements, *Astrophys. J.* **806**, 1 (2015), arXiv:1311.1480.
- [44] K. N. Abazajian, J. K. Adelman-McCarthy, M. A. Agüeros, S. S. Allam, C. Allende Prieto, D. An, K. S. J. Anderson, S. F. Anderson, J. Annis, N. A. Bahcall, *et al.*, The Seventh Data Release of the Sloan Digital Sky Survey, *Astrophys. J. Suppl.* **182**, 543 (2009), arXiv:0812.0649 [astro-ph].
- [45] J. E. Gunn, W. A. Siegmund, E. J. Mannery, R. E. Owen, C. L. Hull, R. F. Leger, L. N. Carey, G. R. Knapp, D. G. York, W. N. Boroski, *et al.*, The 2.5 m Telescope of the Sloan Digital Sky Survey, *The Astronomical J.* **131**, 2332 (2006), arXiv:astro-ph/0602326 [astro-ph].
- [46] M. Fukugita, T. Ichikawa, J. E. Gunn, M. Doi, K. Shimasaku, and D. P. Schneider, The Sloan Digital Sky Survey Photometric System, *The Astronomical J.* **111**, 1748 (1996).
- [47] J. A. Smith, D. L. Tucker, S. Kent, M. W. Richmond, M. Fukugita, T. Ichikawa, S.-i. Ichikawa, A. M. Jorgensen, A. Uomoto, J. E. Gunn, *et al.*, The u’g’r’i’z’ Standard-Star System, *The Astronomical J.* **123**, 2121 (2002), arXiv:astro-ph/0201143 [astro-ph].
- [48] M. Doi, M. Tanaka, M. Fukugita, J. E. Gunn, N. Yasuda, Ž. Ivezić, J. Brinkmann, E. de Haars, S. J. Kleinman,




FIG. 20. Similar to Fig. 11, but the results from “off-centering” and “incompleteness” analysis setups in Table III. We do not see any significant changes in the cosmological constraints in these analysis.




FIG. 21. Similar to Fig. 11, but the results from “ $\sigma(\Delta m) = 0.1$ prior” and “ $\sigma(\Delta z_{\text{ph}}) = 0.2$ prior” analysis setups in Table III. Note that these are the post-unblinding analyses. We do not see a significant change in the “ $\sigma(\Delta m) = 0.1$ prior” analysis, meaning that the degradation of constraining power is mostly due to the use of flat prior on the residual photo- z errors. We see only a slight change in the “ $\sigma(\Delta z_{\text{ph}}) = 0.2$ ” analysis without improvement in the constraining power. This means that with this prior width, we are reaching to the limit of the flat prior.




FIG. 22. Similar to Fig. 11, but the results from “2 cosmo paras” analysis setup in Table III. Note that this is the post-unblinding analyses. When fixing cosmological parameters other than Ω_{de} and $\ln(10^{10} A_s)$ to the Planck *Planck* CMB constraints [11], we obtain better cosmological constraints, but the shift in Δz_{ph} is unchanged.

J. Krzesinski, *et al.*, Photometric Response Functions of the Sloan Digital Sky Survey Imager, *The Astronomical J.* **139**, 1628 (2010), arXiv:1002.3701 [astro-ph.IM].

- [49] D. J. Eisenstein, D. H. Weinberg, E. Agol, H. Aihara, C. Allende Prieto, S. F. Anderson, J. A. Arns, É. Aubourg, S. Bailey, E. Balbinot, *et al.*, SDSS-III: Massive Spectroscopic Surveys of the Distant Universe, the Milky Way, and Extra-Solar Planetary Systems, *The Astronomical J.* **142**, 72 (2011), arXiv:1101.1529 [astro-ph.IM].
- [50] C. P. Ahn, R. Alexandroff, C. Allende Prieto, S. F. Anderson, T. Anderton, B. H. Andrews, É. Aubourg, S. Bailey, E. Balbinot, R. Barnes, *et al.*, The Ninth Data Release of the Sloan Digital Sky Survey: First Spectroscopic Data from the SDSS-III Baryon Oscillation Spectroscopic Survey, *Astrophys. J. Suppl.* **203**, 21 (2012), arXiv:1207.7137 [astro-ph.IM].
- [51] H. Aihara, C. Allende Prieto, D. An, S. F. Anderson, É. Aubourg, E. Balbinot, T. C. Beers, A. A. Berlind, S. J. Bickerton, D. Bizyaev, *et al.*, The Eighth Data Release of the Sloan Digital Sky Survey: First Data from SDSS-III, *Astrophys. J. Suppl.* **193**, 29 (2011), arXiv:1101.1559 [astro-ph.IM].
- [52] R. Lupton, J. E. Gunn, Z. Ivezić, G. R. Knapp, and S. Kent, The SDSS Imaging Pipelines, in *Astronomical Data Analysis Software and Systems X*, *Astronomical Society of the Pacific Conference Series*, Vol. 238, edited by J. Harnden, F. R., F. A. Primini, and H. E. Payne (2001) p. 269, arXiv:astro-ph/0101420 [astro-ph].
- [53] J. R. Pier, J. A. Munn, R. B. Hindsley, G. S. Hennessy, S. M. Kent, R. H. Lupton, and Ž. Ivezić, Astrometric Calibration of the Sloan Digital Sky Survey, *The Astronomical J.* **125**, 1559 (2003), arXiv:astro-ph/0211375 [astro-ph].




FIG. 23. The posterior distributions of all parameters sampled in the baseline 3×2 pt analysis.

- [54] N. Padmanabhan, D. J. Schlegel, D. P. Finkbeiner, J. C. Barntine, M. R. Blanton, H. J. Brewington, J. E. Gunn, M. Harvanek, D. W. Hogg, Ž. Ivezić, *et al.*, An Improved Photometric Calibration of the Sloan Digital Sky Survey Imaging Data, *Astrophys. J.* **674**, 1217 (2008), arXiv:astro-ph/0703454 [astro-ph].
- [55] D. J. Schlegel, D. P. Finkbeiner, and M. Davis, Maps of Dust Infrared Emission for Use in Estimation of Reddening and Cosmic Microwave Background Radiation Foregrounds, *Astrophys. J.* **500**, 525 (1998), arXiv:astro-ph/9710327 [astro-ph].
- [56] A. S. Bolton, D. J. Schlegel, É. Aubourg, S. Bailey, V. Bhardwaj, J. R. Brownstein, S. Burles, Y.-M. Chen, K. Dawson, D. J. Eisenstein, *et al.*, Spectral Classification and Redshift Measurement for the SDSS-III Baryon Oscillation Spectroscopic Survey, *The Astronomical J.* **144**, 144 (2012), arXiv:1207.7326 [astro-ph.CO].
- [57] L. Anderson, É. Aubourg, S. Bailey, F. Beutler, V. Bhardwaj, M. Blanton, A. S. Bolton, J. Brinkmann, J. R. Brownstein, A. Burden, *et al.*, The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: baryon acoustic oscillations in the Data Releases 10 and 11 Galaxy samples, *Mon. Not. Roy. Astron. Soc.* **441**, 24 (2014), arXiv:1312.4877




FIG. 24. The result of `nestcheck`[120] for the baseline 3×2 pt analysis, sampled in the real data analysis. The top panel shows the posterior volume as a function of the prior volume, X . The left panels show the uncertainty of the posterior distribution from an input nested sampling chain, where the uncertainty is estimated by bootstrapping the chain.

[astro-ph.CO].

- [58] I. Esteban, M. C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni, and T. Schwetz, Global analysis of three-flavour neutrino oscillations: synergies and tensions in the determination of θ_{23} , δ_{CP} , and the mass ordering, *Journal of High Energy Physics* **2019**, 106 (2019), arXiv:1811.05487 [hep-ph].
- [59] A. Lewis, A. Challinor, and A. Lasenby, Efficient computation of CMB anisotropies in closed FRW models, *Astrophys. J.* **538**, 473 (2000), arXiv:astro-ph/9911177 [astro-ph].
- [60] P. S. Behroozi, R. H. Wechsler, and H.-Y. Wu, The ROCKSTAR Phase-space Temporal Halo Finder and the Velocity Offsets of Cluster Cores, *Astrophys. J.* **762**, 109 (2013), arXiv:1110.4372 [astro-ph.CO].
- [61] J. L. Tinker, B. E. Robertson, A. V. Kravtsov, A. Klypin, M. S. Warren, G. Yepes, and S. Gottlöber, The Large-scale Bias of Dark Matter Halos: Numerical Calibration and Model Tests, *Astrophys. J.* **724**, 878 (2010), arXiv:1001.3162 [astro-ph.CO].
- [62] J. Lesgourgues, The Cosmic Linear Anisotropy Solving System (CLASS) I: Overview, arXiv e-prints, arXiv:1104.2932 (2011), arXiv:1104.2932 [astro-ph.IM].
- [63] D. Blas, J. Lesgourgues, and T. Tram, The Cosmic Linear Anisotropy Solving System (CLASS). Part II: Approximation schemes, *JCAP* **2011**, 034 (2011), arXiv:1104.2933 [astro-ph.CO].




FIG. 25. The comparison of posterior estimates of our baseline analysis from the nested sampling by `MultiNest` and the Markov-Chain Monte Carlo sampling by the standard Metropolis algorithm. We do not see significant difference in the cosmological constraints from `MultiNest` and the Metropolis algorithm, meaning `MultiNest` provides sufficiently accurate estimate of credible levels.

- [64] R. Mandelbaum, A. Tasitsiomi, U. Seljak, A. V. Kravtsov, and R. H. Wechsler, Galaxy-galaxy lensing: dissipationless simulations versus the halo model, *Mon. Not. Roy. Astron. Soc.* **362**, 1451 (2005), arXiv:astro-ph/0410711.
- [65] G. M. Bernstein and M. Jarvis, Shapes and Shears, Stars and Smears: Optimal Measurements for Weak Lensing, *The Astronomical J.* **123**, 583 (2002), astro-ph/0107431.
- [66] R. Mandelbaum, A. Slosar, T. Baldauf, U. Seljak, C. M. Hirata, R. Nakajima, R. Reyes, and R. E. Smith, Cosmological parameter constraints from galaxy-galaxy lensing and galaxy clustering with the SDSS DR7, *Mon. Not. Roy. Astron. Soc.* **432**, 1544 (2013), arXiv:1207.1120.
- [67] M. Shirasaki, M. Takada, H. Miyatake, R. Takahashi, T. Hamana, T. Nishimichi, and R. Murata, Robust covariance estimation of galaxy-galaxy weak lensing: validation and limitation of jackknife covariance, *Mon. Not. Roy. Astron. Soc.* **470**, 3476 (2017), arXiv:1607.08679 [astro-ph.CO].
- [68] S. Unruh, P. Schneider, S. Hilbert, P. Simon, S. Martin, and J. C. Puertas, The importance of magnification effects in galaxy-galaxy lensing, *Astronomy & Astrophysics* **638**, A96 (2020), arXiv:1910.06400 [astro-ph.CO].
- [69] C. Hikage, R. Mandelbaum, M. Takada, and D. N. Spergel, Where are the Luminous Red Galaxies (LRGs)? Using correlation measurements and lensing to relate LRGs to dark matter haloes, *Mon. Not. Roy. Astron. Soc.* **435**, 2345 (2013), arXiv:1211.1009 [astro-ph.CO].
- [70] Y. P. Jing, H. J. Mo, and G. Börner, Spatial Correlation Function and Pairwise Velocity Dispersion of Galaxies: Cold Dark Matter Models versus the Las Campanas Survey, *Astrophys. J.* **494**, 1 (1998), arXiv:astro-ph/9707106 [astro-ph].

- [71] Z. Zheng *et al.*, Theoretical Models of the Halo Occupation Distribution: Separating Central and Satellite Galaxies, *Astrophys. J.* **633**, 791 (2005), arXiv:astro-ph/0408564.
- [72] J. F. Navarro, C. S. Frenk, and S. D. M. White, A Universal Density Profile from Hierarchical Clustering, *Astrophys. J.* **490**, 493 (1997), arXiv:astro-ph/9611107.
- [73] B. Diemer, COLOSSUS: A Python Toolkit for Cosmology, Large-scale Structure, and Dark Matter Halos, *Astrophys. J. Suppl.* **239**, 35 (2018), arXiv:1712.04512 [astro-ph.CO].
- [74] C. Hikage, M. Takada, and D. N. Spergel, Using galaxy-galaxy weak lensing measurements to correct the finger of God, *Mon. Not. Roy. Astron. Soc.* **419**, 3457 (2012), arXiv:1106.1640 [astro-ph.CO].
- [75] R. Takahashi, M. Sato, T. Nishimichi, A. Taruya, and M. Oguri, Revising the halofit model for the nonlinear matter power spectrum, *The Astrophysical Journal* **761**, 152 (2012).
- [76] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, *et al.*, Planck 2015 results. XIII. Cosmological parameters, *Astronomy & Astrophysics* **594**, A13 (2016), arXiv:1502.01589.
- [77] F. C. van den Bosch, S. More, M. Cacciato, H. Mo, and X. Yang, Cosmological constraints from a combination of galaxy clustering and lensing - I. Theoretical framework, *Mon. Not. Roy. Astron. Soc.* **430**, 725 (2013), arXiv:1206.6890 [astro-ph.CO].
- [78] A. Heavens, A. Refregier, and C. Heymans, Intrinsic correlation of galaxy shapes: implications for weak lensing measurements, *Mon. Not. Roy. Astron. Soc.* **319**, 649 (2000), astro-ph/0005269.
- [79] R. A. C. Croft and C. A. Metzler, Weak-Lensing Surveys and the Intrinsic Correlation of Galaxy Ellipticities, *Astrophys. J.* **545**, 561 (2000), arXiv:astro-ph/0005384 [astro-ph].
- [80] J. Lee and U.-L. Pen, Cosmic Shear from Galaxy Spins, *Astrophys. J. Lett.* **532**, L5 (2000), arXiv:astro-ph/9911328 [astro-ph].
- [81] P. Catelan, M. Kamionkowski, and R. D. Blandford, Intrinsic and extrinsic galaxy alignment, *Mon. Not. Roy. Astron. Soc.* **320**, L7 (2001), arXiv:astro-ph/0005470 [astro-ph].
- [82] C. M. Hirata, R. Mandelbaum, U. Seljak, J. Guzik, N. Padmanabhan, C. Blake, J. Brinkmann, T. Budavari, A. Connolly, I. Csabai, *et al.*, Galaxy-galaxy weak lensing in the Sloan Digital Sky Survey: intrinsic alignments and shear calibration errors, *Mon. Not. Roy. Astron. Soc.* **353**, 529 (2004), arXiv:astro-ph/0403255 [astro-ph].
- [83] D. N. Limber, The Analysis of Counts of the Extragalactic Nebulae in Terms of a Fluctuating Density Field. II., *Astrophys. J.* **119**, 655 (1954).
- [84] M. Takada and B. Jain, Cosmological parameters from lensing power spectrum and bispectrum tomography, *Mon. Not. Roy. Astron. Soc.* **348**, 897 (2004), arXiv:astro-ph/0310125.
- [85] S. Bridle and L. King, Dark energy constraints from cosmic shear power spectra: impact of intrinsic alignments on photometric redshift requirements, *New Journal of Physics* **9**, 444 (2007), arXiv:0705.0166 [astro-ph].
- [86] M. L. Brown, A. N. Taylor, N. C. Hambly, and S. Dye, Measurement of intrinsic alignments in galaxy ellipticities, *Mon. Not. Roy. Astron. Soc.* **333**, 501 (2002), arXiv:astro-ph/0009499 [astro-ph].
- [87] H. Hildebrandt, M. Viola, C. Heymans, S. Joudaki, K. Kuijken, C. Blake, T. Erben, B. Joachimi, D. Klaes, L. Miller, *et al.*, KiDS-450: cosmological parameter constraints from tomographic weak gravitational lensing, *Mon. Not. Roy. Astron. Soc.* **465**, 1454 (2017), arXiv:1606.05338 [astro-ph.CO].
- [88] M. M. Rau, R. Dalal, T. Zhang, X. Li, A. J. Nishizawa, S. More, R. Mandelbaum, M. A. Strauss, and M. Takada, Weak Lensing Tomographic Redshift Distribution Inference for the Hyper Suprime-Cam Subaru Strategic Program three-year shape catalogue, (2022), arXiv:2211.16516 [astro-ph.CO].
- [89] D. Huterer, M. Takada, G. Bernstein, and B. Jain, Systematic errors in future weak-lensing surveys: requirements and prospects for self-calibration, *Mon. Not. Roy. Astron. Soc.* **366**, 101 (2006), arXiv:astro-ph/0506030.
- [90] T. Zhang, M. M. Rau, R. Mandelbaum, X. Li, and B. Moews, Photometric redshift uncertainties in weak gravitational lensing shear analysis: models and marginalization, *Mon. Not. Roy. Astron. Soc.* **518**, 709 (2023), arXiv:2206.10169 [astro-ph.CO].
- [91] T. Zhang, X. Li, R. Dalal, R. Mandelbaum, M. A. Strauss, A. Kannawadi, H. Miyatake, A. Nicola, A. A. Plazas Malagón, M. Shirasaki, *et al.*, A General Framework for Removing Point Spread Function Additive Systematics in Cosmological Weak Lensing Analysis, arXiv e-prints, arXiv:2212.03257 (2022), arXiv:2212.03257 [astro-ph.CO].
- [92] M. A. Troxel, N. MacCrann, J. Zuntz, T. F. Eifler, E. Krause, S. Dodelson, D. Gruen, J. Blazek, O. Friedrich, S. Samuroff, *et al.*, Dark Energy Survey Year 1 results: Cosmological constraints from cosmic shear, *Phys. Rev. D* **98**, 043528 (2018), arXiv:1708.01538 [astro-ph.CO].
- [93] A. J. S. Hamilton, Uncorrelated modes of the non-linear power spectrum, *Mon. Not. Roy. Astron. Soc.* **312**, 257 (2000), astro-ph/9905191.
- [94] X. Fang, T. Eifler, and E. Krause, 2D-FFTLLog: efficient computation of real-space covariance matrices for galaxy clustering and weak lensing, *Mon. Not. Roy. Astron. Soc.* **497**, 2699 (2020), arXiv:2004.04833 [astro-ph.CO].
- [95] C.-H. Lin, J. Harnois-Déraps, T. Eifler, T. Pospisil, R. Mandelbaum, A. B. Lee, S. Singh, and LSST Dark Energy Science Collaboration, Non-Gaussianity in the weak lensing correlation function likelihood - implications for cosmological parameter biases, *Mon. Not. Roy. Astron. Soc.* **499**, 2977 (2020), arXiv:1905.03779 [astro-ph.CO].
- [96] E. Aver, K. A. Olive, and E. D. Skillman, The effects of He I $\lambda 10830$ on helium abundance determinations, *JCAP* **07**, 011, arXiv:1503.08146 [astro-ph.CO].
- [97] R. J. Cooke, M. Pettini, and C. C. Steidel, One Percent Determination of the Primordial Deuterium Abundance, *Astrophys. J.* **855**, 102 (2018), arXiv:1710.11129 [astro-ph.CO].
- [98] N. Schöneberg, J. Lesgourgues, and D. C. Hooper, The BAO+BBN take on the Hubble tension, *JCAP* **10**, 029, arXiv:1907.11594 [astro-ph.CO].
- [99] S. Sugiyama, M. Takada, Y. Kobayashi, H. Miyatake, M. Shirasaki, T. Nishimichi, and Y. Park, Validating a minimal galaxy bias method for cosmological parameter inference using HSC-SDSS mock catalogs, *Phys. Rev. D* **102**, 083520 (2020), arXiv:2008.06873 [astro-ph.CO].
- [100] F. Feroz and M. P. Hobson, Multimodal nested sampling: an efficient and robust alternative to Markov Chain Monte Carlo methods for astronomical data analyses, *Mon. Not. Roy. Astron. Soc.* **384**, 449 (2008), arXiv:0704.3704 [astro-ph].
- [101] F. Feroz, M. P. Hobson, and M. Bridges, MULTINEST: an efficient and robust Bayesian inference tool for cosmology and particle physics, *Mon. Not. Roy. Astron. Soc.* **398**, 1601 (2009), arXiv:0809.3437 [astro-ph].
- [102] F. Feroz, M. P. Hobson, E. Cameron, and A. N. Pettitt, Importance Nested Sampling and the MultiNest Algorithm, *The*

- Open Journal of Astrophysics **2**, 10 (2019), arXiv:1306.2144 [astro-ph.IM].
- [103] J. Buchner, A. Georgakakis, K. Nandra, L. Hsu, C. Rangel, M. Brightman, A. Merloni, M. Salvato, J. Donley, and D. Kocevski, X-ray spectral modelling of the AGN obscuring region in the CDFS: Bayesian model selection and catalogue, *Astronomy & Astrophysics* **564**, A125 (2014), arXiv:1402.0004 [astro-ph.HE].
- [104] H. Miyatake, Y. Kobayashi, M. Takada, T. Nishimichi, M. Shirasaki, S. Sugiyama, R. Takahashi, K. Osato, S. More, and Y. o. Park, Cosmological inference from emulator based halo model I: Validation tests with HSC and SDSS mock catalogs, arXiv e-prints, arXiv:2101.00113 (2020), arXiv:2101.00113 [astro-ph.CO].
- [105] T. Nishimichi, G. D'Amico, M. M. Ivanov, L. Senatore, M. Simonović, M. Takada, M. Zaldarriaga, and P. Zhang, Blinded challenge for precision cosmology with large-scale structure: Results from effective field theory for the redshift-space galaxy power spectrum, *Phys. Rev. D* **102**, 123541 (2020), arXiv:2003.08277 [astro-ph.CO].
- [106] Y. Park and E. Rozo, Concordance cosmology?, *Mon. Not. Roy. Astron. Soc.* **499**, 4638 (2020), arXiv:1907.05798 [astro-ph.CO].
- [107] M. Raveri, G. Zacharegkas, and W. Hu, Quantifying concordance of correlated cosmological data sets, *Phys. Rev. D* **101**, 103527 (2020), arXiv:1912.04880 [astro-ph.CO].
- [108] P. Lemos, M. Raveri, A. Campos, Y. Park, C. Chang, N. Weaverdyck, D. Huterer, A. R. Liddle, J. Blazek, R. Cawthon, *et al.*, Assessing tension metrics with dark energy survey and Planck data, *Mon. Not. Roy. Astron. Soc.* **505**, 6179 (2021), arXiv:2012.09554 [astro-ph.CO].
- [109] T. Charnock, R. A. Battye, and A. Moss, Planck data versus large scale structure: Methods to quantify discordance, *Phys. Rev. D* **95**, 123535 (2017), arXiv:1703.05959 [astro-ph.CO].
- [110] H.-R. Yu, J. Harnois-Déraps, T.-J. Zhang, and U.-L. Pen, Information content in the angular power spectrum of weak lensing: wavelet method, *Mon. Not. Roy. Astron. Soc.* **421**, 832 (2012), arXiv:1012.0444 [astro-ph.CO].
- [111] B. Hadzhiyska, S. Bose, D. Eisenstein, and L. Hernquist, Extensions to models of the galaxy-halo connection, *Mon. Not. Roy. Astron. Soc.* **501**, 1603 (2021), arXiv:2008.04913 [astro-ph.CO].
- [112] D. J. Eisenstein *et al.*, Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies, *Astrophys. J.* **633**, 560 (2005), arXiv:astro-ph/0501171.
- [113] S. Alam, M. Aubert, S. Avila, C. Balland, J. E. Bautista, M. A. Bershad, D. Bizyaev, M. R. Blanton, A. S. Bolton, J. Bovy, *et al.*, Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory, *Phys. Rev. D* **103**, 083533 (2021), arXiv:2007.08991 [astro-ph.CO].
- [114] Y. Kobayashi, T. Nishimichi, M. Takada, and R. Takahashi, Cosmological information content in redshift-space power spectrum of SDSS-like galaxies in the quasilinear regime up to $k = 0.3 \text{ h Mpc}^{-1}$, *Phys. Rev. D* **101**, 023510 (2020), arXiv:1907.08515 [astro-ph.CO].
- [115] O. H. E. Philcox, M. M. Ivanov, M. Simonović, and M. Zaldarriaga, Combining full-shape and BAO analyses of galaxy power spectra: a 1.6% CMB-independent constraint on H_0 , *JCAP* **2020**, 032 (2020), arXiv:2002.04035 [astro-ph.CO].
- [116] M. Takada, R. S. Ellis, M. Chiba, J. E. Greene, H. Aihara, N. Arimoto, K. Bundy, J. Cohen, O. Doré, G. Graves, *et al.*, Extragalactic science, cosmology, and Galactic archaeology with the Subaru Prime Focus Spectrograph, *Publ. Astron. Soc. Japan* **66**, R1 (2014), arXiv:1206.0737.
- [117] Ž. Ivezić, S. M. Kahn, J. A. Tyson, B. Abel, E. Acosta, R. Allsman, D. Alonso, Y. AlSayyad, S. F. Anderson, J. Andrew, J. R. P. Angel, *et al.*, LSST: From Science Drivers to Reference Design and Anticipated Data Products, *Astrophys. J.* **873**, 111 (2019), arXiv:0805.2366 [astro-ph].
- [118] R. Laureijs, J. Amiaux, S. Arduini, J. . Auguères, J. Brinchmann, R. Cole, M. Cropper, C. Dabin, L. Duvet, A. Ealet, *et al.*, *et al.*, Euclid Definition Study Report, ArXiv e-prints (2011), arXiv:1110.3193 [astro-ph.CO].
- [119] D. Spergel, N. Gehrels, C. Baltay, D. Bennett, J. Breckinridge, M. Donahue, A. Dressler, B. S. Gaudi, T. Greene, O. Guyon, C. Hirata, *et al.*, Wide-Field Infrared Survey Telescope-Astrophysics Focused Telescope Assets WFIRST-AFTA 2015 Report, ArXiv e-prints (2015), arXiv:1503.03757 [astro-ph.IM].
- [120] E. Higson, nestcheck: error analysis, diagnostic tests and plots for nested sampling calculations, *Journal of Open Source Software* **3**, 916 (2018).
- [121] E. Higson, W. Handley, M. Hobson, and A. Lasenby, nestcheck: diagnostic tests for nested sampling calculations, *Monthly Notices of the Royal Astronomical Society* **483**, 2044 (2019).