INVOLUTIONS, LINKS, AND FLOER COHOMOLOGIES

HOKUTO KONNO, JIN MIYAZAWA, AND MASAKI TANIGUCHI

ABSTRACT. We develop a version of Seiberg–Witten Floer cohomology/homotopy type for a spin^c 4-manifold with boundary and with an involution which reverses the spin^c structure, as well as a version of Floer cohomology/homotopy type for oriented links with non-zero determinant. This framework generalizes the previous work of the authors regarding Floer homotopy type for spin 3manifolds with involutions and for knots. Based on this Floer cohomological setting, we prove Frøyshov-type inequalities which relate topological quantities of 4-manifolds with certain equivariant homology cobordism invariants. The inequalities and homology cobordism invariants have applications to the topology of unoriented surfaces, the Nielsen realization problem for non-spin 4-manifolds, and non-smoothable unoriented surfaces in 4-manifolds.

Contents

1.	Introduction	2
1.1.	Background	2
1.2.	A constraint on involutions on closed 4-manifolds	3
1.3.	New link concordance invariants	4
1.4.	Applications to non-orientable surfaces in D^4 bounded by torus knots	7
1.5.	Nielsen realization for non-spin 4-manifolds	8
1.6.	Non-smoothable and non-orientable surfaces	9
1.7.	Structural conjecture	10
1.8.	Structure of the paper	10
1.9.	Acknowledgement	11
2.	Involution I	11
3.	Floer homotopy type of rational homology 3-spheres with involution	
	and Frøyshov type invariants	15
3.1.	Representations	15
3.2.	Numerical invariants $d, \underline{d}, \overline{d}$	16
3.3.	Duality	21
3.4.	Spectrum classes	22
3.5.	Seiberg–Witten Floer stable homotopy type for non-spin case	24
3.6.	Seiberg–Witten Floer cohomology for involutions	25

3.7. Cobordisms	27	
3.8. Proof of Frøyshov type inequalities	28	
3.9. Connected sum formula	29	
3.10. SWF-spherical	31	
4. Floer homotopy type of links and Frøyshov type invariants	31	
4.1. Floer homotopy type of links	32	
4.2. Frøyshov type invariants for links	32	
4.3. κ invariants for links	35	
5. Applications	36	
5.1. Applications to non-smoothable actions	36	
5.2. Applications to non-smoothable and non-orientable surfaces	37	
5.3. Applications to the Nielsen realization problem	37	
5.4. Several inequalities for knots	40	
5.5. Applications to non-orientable genera and normal Euler numbers	41	
References		

1. INTRODUCTION

1.1. **Background.** We introduce a version of Seiberg–Witten Floer cohomology theory for 3-manifolds with involutions, as well as a version of Seiberg–Witten Floer cohomology theory for oriented links with non-zero determinant, which can be seen as a generalization of a theory developed by the authors in [21].

In [21], for a given oriented 3-manifold Y with a spin structure \mathfrak{s} and an odd involution ι , the authors constructed Seiberg–Witten Floer Floer homotopy theory/K-theory and proved a 10/8-type inequality in the setting, based on Manolescu's Seiberg–Witten Floer homotopy type [32].

In this paper, we extend such a theory for (possibly non-spin) spin^c structures called *real spin^c* structures defined as follows. A real spin^c structure on a manifold is a pair of a spin^c structure \mathfrak{s} and an orientation-preserving smooth involution ι on the manifold such that

 $\iota^*\mathfrak{s}\cong\overline{\mathfrak{s}},$

where $\overline{\mathfrak{s}}$ is the conjugate spin^c structure of \mathfrak{s} . We call a manifold equipped with a real spin^c structure a *real spin^c manifold*. In this paper, we mainly consider involutions that have codimension-2 non-empty fixed sets.

For a real spin^c 3- or 4-manifold, we shall define an involutive symmetry I of the Seiberg–Witten equations acting anti-linearly on the domain and the codomain of the map corresponding to the equations. When a spin^c structure is spin, the involution I is the same as the involution $I = j \circ \tilde{\iota}$ used in previous paper [21], where $\tilde{\iota}$ is a lift of ι to a spin structure and j denotes the third basis vector of the

 $\mathbf{2}$

quaternion. We shall use information of Seiberg–Witten theory corresponding to the fixed point set of I.

There are several preceding studies of Seiberg–Witten theory for real spin^c 3or 4-manifolds, which may be called *real Seiberg–Witten theory*. In [52], Tian and Wang introduced real Seiberg–Witten invariants for hermitian almost complex 4-manifolds by counting the moduli spaces corresponding to the *I*-fixed point parts. Recently, Li [28] introduced real monopole Floer homology for real spin^c 3-manifolds, which can be seen as the associated Floer homology to consider the relative version of real Seiberg–Witten invariants for real 4-manifolds in [52], by developing Floer theory for the *I*-fixed point parts.

It is natural to conjecture that our framework gives a Floer homotopy refinement of Li's real monopole Floer homology for real spin^c 3-manifolds under the assumption that $b_1 = 0$, as Manolescu's Seiberg–Witten Floer homotopy type gives a spacial refinement of monopole Floer homology [29].

While we shall mainly focus on involutions with codimension-2 non-empty fixed point sets, the corresponding theory for free involution has been known as Pin⁻(2) Seiberg–Witten theory developed by Nakamura [41, 42]. Also, when we consider a real spin structure, the corresponding Floer homotopy type or 4-dimensional invariant can be seen as the fixed point parts of Montague's [38] $\mathbb{Z}_4 \times_{\mathbb{Z}_2}$ Pin(2)equivariant homotopy spectra $SWF(\Sigma(L), \mathfrak{s})$ and $\mathbb{Z}_4 \times_{\mathbb{Z}_2}$ Pin(2)-equivariant Bauer– Furuta invariant with respect to the element [(i, j)], where *i* is a generator of \mathbb{Z}_4 .

Our main results are Frøyshov-type inequalities for real spin^c 4-manifolds with boundary, and for surfaces bounded by links. We first exhibit results for closed 4-manifolds below.

1.2. A constraint on involutions on closed 4-manifolds. Let (W, \mathfrak{s}, ι) be a real spin^c 4-manifold, i.e. W is an oriented smooth 4-manifold, \mathfrak{s} is a spin^c structure on W, and $\iota : W \to W$ is an orientation-preserving smooth involution such that $\iota^*\mathfrak{s} \cong \overline{\mathfrak{s}}$. We shall define the notion that ι is of odd type, and see that ι is of odd type if the fixed-point set W^{ι} is non-empty and of codimension-2. Let $\sigma(W)$ denote the signature of W and $b^+(W)$ denote the maximal dimension of positive-definite subspaces of $H^2(W;\mathbb{R})$. We denote by $b_{\iota}^+(W)$ the maximal dimension of the ι -invariant part of positive-definite subspaces of $H^2(W;\mathbb{R})$. One of our main theorems is stated as follows:

Theorem 1.1. Let (W, \mathfrak{s}, ι) be a closed oriented smooth real spin^c 4-manifold with $b_1(W) = 0$. Suppose that ι is of odd type. If $b_{\iota}^+(W) = b^+(W)$, then we have

(1)
$$c_1(\mathfrak{s})^2 - \sigma(W) \le 0.$$

For negative-definite 4-manifolds, the inequality (1) is well-known to be held for every spin^c structure. The theorem states that we have the inequality (1) also for non-negative-definite 4-manifolds, as far as $b_{\iota}^+(W) = b^+(W)$ and \mathfrak{s} is real.

Remark 1.2. One can apply Theorem 1.1 to 4-manifolds with involution obtained from double branched covering spaces along surfaces in 4-manifolds. By describing (1) in terms of the base 4-manifolds, we obtain the following from Theorem 1.1: for a closed 4-manifold X and a smoothly embedded (possibly non-orientable) surface S satisfying

$$[S] \equiv 0 \mod 2 \text{ and } b^+(X) + \frac{1}{2}b_1(S) - \frac{1}{4}S \circ S = 0,$$

one has

(2)
$$c_1(\mathfrak{s})^2 - 2\sigma(X) + \frac{1}{2}S \circ S \le 0$$

for any spin^c structure on the double branched covering space $\Sigma(S)$ which satisfies $\iota^*\mathfrak{s} \cong \overline{\mathfrak{s}}$, where ι is the covering involution on $\Sigma(S)$ and $S \circ S$ denotes the evaluation of the normal Euler class of S.

1.3. New link concordance invariants. Theorem 1.1 is generalized to the case of 4-manifolds with boundary and it will be translated as constraints of surfaces embedded into 4-manifolds via the double branched cover construction. In order to write Frøyshov type inequality for 4-manifolds with boundary, we need to introduce Frøyshov type invariants. As such a Frøyshov invariant, we shall define a linkinvariant

$$\delta_R(L) \in \frac{1}{16}\mathbb{Z}$$

for an oriented link with non-zero determinant, where $\Sigma(L)$ is the double branched covering space along L and t_L is the spin structure corresponding to the orientation of L. Here R stands for real Seiberg-Witten Frøyshov invariant. In order to summarize basic properties of $\delta_R(L)$, it is convenient to use notion of χ -concordance introduced in [8]. Let us first review the definition of χ -concordance. A marked link is a link in S^3 equipped with a marked component. For given oriented marked links L_0 and L_1 , we call L_0 and L_1 are χ -concordant if $-L_0^* \# L_1$ bounds a smoothly properly embedded surface F in D^4 such that

- (i) F is a disjoint union of one disk together with annuli and Möbius bands;
- (ii) the boundary of the disk component of F is the marked component of $-L_0^* \# L_1;$
- (iii) we require orientable components of F to be oriented compatible with the orientation of $-L_0^* \# L_1$.

where L^* means the mirror image of L and the connected sum $-L_0^* \# L_1$ is taken along marked components. In [8], it is proven that the set $\tilde{\mathcal{L}}$ of all χ -concordant classes of oriented marked links forms an abelian group with respect to the connected sum along marked components. The group $\hat{\mathcal{L}}$ is called the *link concordance* group. In this paper, we focus on the subgroup $\tilde{\mathcal{F}}$ of $\tilde{\mathcal{L}}$ generated by oriented marked links whose determinants are *non-zero*. For an embedded surface S in a given 4-manifold X, we denote by $\Sigma(S)$ the double branched cover of X along Σ if it exists.

Theorem 1.3. The invariant $\delta_R(L)$ associated to an oriented link with non-zero determinant satisfies the following properties:

- (i) The quantity $\delta_R(L)$ is a χ -concordance invariant. Thus, we have a welldefined map $\delta_R : \tilde{\mathcal{F}} \to \frac{1}{16}\mathbb{Z}.$
- (ii) The map $\delta_R : \tilde{\mathcal{F}} \to \frac{1}{16}\mathbb{Z}$ is a homomorphism. (iii) Let L and L' be oriented links in S³ with non-zero determinants, X be an oriented smooth compact connected cobordism from S^3 to S^3 , and S be a

compact connected properly and smoothly (possibly non-orientable) embedded cobordism in X from L to L' such that the homology class $[S] \equiv 0 \mod 2$. Suppose that $b_1(X) = 0$,

(3)
$$b^+(X) + \frac{1}{2}b_1(S) - \frac{1}{4}S \circ S - \frac{1}{2}\sigma(L) + \frac{1}{2}\sigma(L') = 0,$$

and there is a spin^c structure \mathfrak{s} on $\Sigma(S)$ such that $\iota^*\mathfrak{s} \cong \overline{\mathfrak{s}}$ and whose restrictions are compatible with orientations of L and L'.

Then, we have

(4)
$$\delta_R(L) + \frac{1}{16} \left(\langle c_1(\mathfrak{s})^2, [\Sigma(S)] \rangle - 2\sigma(X) - \frac{1}{2}S \circ S - \sigma(L) + \sigma(L') \right) \le \delta_R(L')$$

where $\sigma(L)$ denotes the signature of L (with the sign convention $\sigma(T(2,3)) = -2$), and $S \circ S$ means the self-intersection number (or normal Euler number) of S in X.

As an advantage of the inequality (34), which we may call *real Frøyshov inequality*, we can directly treat non-orientable surfaces.

Remark 1.4. The inequality (34) implies (2) if we put L and L' are the unknot. Also note that the quantity $b^+(X) + \frac{1}{2}b_1(S) - \frac{1}{4}S \circ S - \frac{1}{2}\sigma(L) + \frac{1}{2}\sigma(L')$ is non-negative since it can be written as $b^+(\Sigma(S)) - b_{\iota}^+(\Sigma(S))$.

Remark 1.5. When we restrict δ_R as a function on the knot concordance group, we obtain a homomorphism

$$\delta_R: \mathcal{C} \to \frac{1}{16}\mathbb{Z}.$$

For two bridge knots, we have $\delta_R = -\frac{1}{16}\sigma$. It is natural to ask δ_R is a slice torus invariant ([27, 30]) up to constant multiplication. However, (34) does not imply slice genus bounds and, for torus knots, it is *not* equal to their slice genera up to constant multiplication.

As analog of invariants introduced by Stoffregen [50] $\bar{\delta}(Y, \mathfrak{t})$, $\underline{\delta}(Y, \mathfrak{t})$ corresponding to $\bar{d}(Y, \mathfrak{t})$, $\underline{d}(Y, \mathfrak{t})$ in involutive Heegaard Floer homology [16], we also define invariants

$$\bar{\delta}_R(L)$$
 and $\underline{\delta}_R(L) \in \frac{1}{16}\mathbb{Z}$

for any link L with non-zero determinant.

When \mathfrak{s} is spin and (3) is satisfied, we also have inequalities similar to (34). See Theorem 4.6 for the details. Moreover, we also have the following Theorem B type inequality:

Theorem 1.6. The invariants $\underline{\delta}_R(L)$ and $\overline{\delta}_R(L)$ associated to an oriented link with non-zero determinant satisfies the following property: Let L, L', X, S be as in (iii) of Theorem 1.3, except for that we suppose that \mathfrak{s} is a spin structure and that

(5)
$$b^+(X) + \frac{1}{2}b_1(S) - \frac{1}{4}S \circ S - \frac{1}{2}\sigma(L) + \frac{1}{2}\sigma(L') = 1,$$

instead of (3). Then, we have

(6)
$$\underline{\delta}_R(L) + \frac{1}{16} \left(-2\sigma(X) + \frac{1}{2}S \circ S + \sigma(L) - \sigma(L') \right) \le \overline{\delta}_R(L').$$

Remark 1.7. If we take the unknot as both L and L' and the trivial null-homologous 2-disk embedded in a spin 4-manifold X as S, then the inequality (31) implies $-\frac{1}{8}\sigma(X) \leq 0$ if $b^+(X) = 1$. This recovers Theorem B proven by Donaldson [10].

By specifying Theorem 1.3 to $X = S^3 \times [0, 1]$, L' = U and L is a knot, we have the following constraint on topology of non-orientable surfaces in D^4 :

Theorem 1.8. Let S be a (possibly non-orientable) smoothly and properly embedded connected surface in D^4 bounded by a given knot K. Suppose that

- (i) the determinant of K is 1;
- (ii) the Manolescu–Owen invariant [35] $\delta(K) = \delta(\Sigma(K))$ is zero;
- (iii) $\delta_R(K) < 0.$

Then, we have

$$-\sigma(K) + \frac{1}{2}S \circ S + 1 \le b_1(S).$$

We shall use Theorem 1.8 to obstruct certain types of non-orientable surfaces bounded by torus knots in the next subsection.

Remark 1.9. As a corollary of Theorem 1.3, one can provide crossing change formula for $\delta_R(L)$, $\bar{\delta}_R(L)$ and $\underline{\delta}_R(L)$. Let us summarize the results for the case of knots: Let K and K' be knots in S^3 . Suppose that K' is obtained from K by a positive crossing change, where our convention on the positive crossing change is the same as in [21]. Then, we have the following:

(i) Suppose $\sigma(K) - \sigma(K') = 2$. Then the inequalities

$$\delta_R(K) + \frac{1}{8} \le \delta_R(K'), \quad \underline{\delta}_R(K) + \frac{1}{8} \le \underline{\delta}_R(K'), \quad \overline{\delta}_R(K) + \frac{1}{8} \le \overline{\delta}_R(K')$$

hold

(ii) Suppose $\sigma(K) - \sigma(K') = 0$. Then the inequalities

$$\delta_R(K') - \frac{1}{4} \le \delta_R(K), \quad \underline{\delta}_R(K') - \frac{1}{4} \le \underline{\delta}_R(K), \quad \overline{\delta}_R(K') - \frac{1}{4} \le \overline{\delta}_R(K)$$

hold.

We compute $\delta_R(L), \bar{\delta}_R(L), \underline{\delta}_R(L)$ for some class of links:

Proposition 1.10. We have the following computations:

(i) For a two bridge link K(p,q), we have

$$\delta_R(K(p,q)) = \underline{\delta}_R(K(p,q)) = \overline{\delta}_R(K(p,q)) = -\frac{\sigma(K(p,q))}{16}.$$

(ii) Let p, q be coprime odd integers and T(p, q) be the torus knot of type (p, q). Then we have

$$\delta_R(T(p,q)) = \underline{\delta}_R(T(p,q)) = \overline{\delta}_R(T(p,q)) = -\frac{\overline{\mu}(\Sigma(2,p,q))}{2},$$

where $\bar{\mu}$ denote the Neumann–Siebenmann invariant [43].

Also, the following computations follow from Montague's computations of equivariant Seiberg–Witten Floer stable homotopy types [38].

Example 1.11. For Montesinos knots of type $M(2, 3, 6n \pm 1)$, we have the following computations (see Example 3.21):

Montesinos knot	$\bar{\delta}_R$	δ_R	δ_R
M(2, 3, 12k - 1)	1/2	1/2	0
M(2, 3, 12k - 5)	0	0	-1/2
M(2, 3, 12k + 1)	0	0	0
M(2, 3, 12k + 5)	1/2	1/2	1/2

Now, we focus on applications of Theorem 1.1 and Theorem 1.3.

1.4. Applications to non-orientable surfaces in D^4 bounded by torus knots. Recently, using several types of Floer theories, the *non-orientable 4-genus* has been studied, for example see [1,5,6,31,48]. For a given knot K in S^3 , we focus on topological types of smoothly and properly embedded possibly non-orientable surfaces in D^4 bounded by K. For such a surface S, the *non-orientable genus* h(S) of Sis defined by $h(S) := b_1(S)$. The non-orientable 4-genus is the minimum number of h(S) for all such S. Moreover, there is also another topological invariant of the embedding S, called the *normal Euler number* $S \circ S$. The number $S \circ S$ is defined as the evaluation of the twisted Euler class of a normal bundle of S by the fundamental class of S (in the orientation local system of S). The following question is natural to ask regarding topology of non-orientable surfaces:

Question 1.12 ([1]). For a given knot K in S^3 , what is the set of realizable pairs

$$(S \circ S, h(S)) \in \mathbb{Z} \times \mathbb{Z}_{>0},$$

where S are properly and smoothly embedded (possibly non-orientable) surfaces in D^4 bounded by K.

Even for torus knots, Question 1.12 has not been solved completely. We focus on torus knots of type T(3, q). By using the Heegaard Floer *d*-invariant and the upsilon invariant Υ [44], Allen [1] detected the set all such possible pairs $(S \circ S, h(S))$ for positive torus knots T(3, 6n + 4) and T(3, 6n + 5). By a similar argument, she [1] detected the realizable pairs $(S \circ S, h(S))$ for T(3, 6n + 1) and T(3, 6n + 2) except for the following cases:

- (i) for T(3, 6n + 1), (e, h) = (8/3(1 n) + 2 + 2m, 1 + m);
- (ii) for T(3, 6n + 2), (e, h) = (8/3(2 n) + 2 + 2m, 3 + m),

where m is a non-negative integer. Related the remaining parts, Allen conjectured the following.

Conjecture 1.13 ([1]). Both of (i) and (ii) are not realizable.

Theorem 1.8 enables us to prove the half of the conjecture.

Theorem 1.14. The case (i) is not realizable.

Combined with Allen's result, we can detect all realizable pairs.

Corollary 1.15. The set of realizable pairs $(S \circ S, h(S))$ for T(3, 6n + 1) is given by

$$(-2 - 16n \pm 2m, 1 + m + 2l), m, l \ge 0.$$

1.5. Nielsen realization for non-spin 4-manifolds. Next application is about the Nielsen realization problem. Given a smooth manifold W, a subgroup G of $\pi_0(\text{Diff}(W))$ is said to be *realizable in* Diff(W) when there is a section $s: G \to$ Diff(W) of the natural map $\text{Diff}(W) \to \pi_0(\text{Diff}(W))$ over G. Recall that, for a (-1)-sphere S in an oriented 4-manifold W, we obtain an orientation-preserving diffeomorphism $\rho_S: W \to W$ called the *reflection*, which is locally modeled on the complex conjugation on $-\mathbb{CP}^2$. It is easy to see that the diffeomorphism ρ_S generates an order 2 subgroup of $\pi_0(\text{Diff}(W))$. (See Lemma 5.4.)

Theorem 1.16. Let W be a closed oriented smooth 4-manifold with $\sigma(W) \neq -2$. Let \mathfrak{s} be a spin^c structure on W and let S be a smoothly embedded (-1)-sphere in W. If $c_1(\mathfrak{s})^2 - \sigma(W) > 0$ and $\rho_S^* \mathfrak{s} \cong \overline{\mathfrak{s}}$, then ρ_S is not homotopic to any smooth involution.

Example 1.17. Let W' be a closed oriented spin smooth 4-manifold with $\sigma(W') < 0$. Set $W = W' \# (-\mathbb{CP}^2)$. Let S be the canonical (-1)-sphere of the $(-\mathbb{CP}^2)$ component. Then $\rho_S : W \to W$ is not homotopic to any smooth involution. Indeed, let \mathfrak{s} be a spin^c structure on W such that $c_1(\mathfrak{s})|_{W'} = 0$ and $c_1(\mathfrak{s})|_{-\mathbb{CP}^2}$ is a generator of $H^2(-\mathbb{CP}^2)$. Then the reflection $\rho_S : W \to W$ satisfies the assumption of Theorem 1.16 for \mathfrak{s} .

Example 1.17 immediately implies:

Corollary 1.18. Let W' be a closed oriented spin smooth 4-manifold with $\sigma(W') < 0$. Set $W = W' \# (-\mathbb{CP}^2)$ and let S be the exceptional sphere in the $(-\mathbb{CP}^2)$ component. Then the order 2 subgroup G of $\pi_0(\text{Diff}(W))$ generated by the mapping
class $[\rho_S]$ of the reflection about S is not realizable in Diff(W).

To the best of the author's knowledge, Corollary 1.18 gives the first example of a non-spin 4-manifold that admits non-realizable finite subgroup of the mapping class group. The first example of a 4-manifold that is shown to admit a nonrealizable finite subgroup is some nilmanifold [47] due to Raymond and Scott. (Note that every nilmanifold is parallelizable and hence spin.) Recently, many simplyconnected spin 4-manifolds are shown to admit non-realizable finite subgroups: for K3 by Baraglia and the first author [4] and by Farb and Looijenga [11], and later for more general spin 4-manifolds by the first author [20].

Remark 1.19. After the first version of this paper appeared on arXiv, the authors were informed that, in their upcoming work [2], Arabadji and Baykur give other examples of non-liftable subgroups of mapping class groups of non-spin 4-manifolds, including irreducible 4-manifolds and definite 4-manifolds.

Remark 1.20. Generalizing Corollary 1.18, one may also obtain a non-realizable order 2 element of $\pi_0(\text{Diff}(W))$ for $W = W' \# n(-\mathbb{CP}^2)$ for a spin W and for a general $n \geq 1$ by considering connected sums of ρ_S along fixed points.

Remark 1.21. The assumption that $c_1(\mathfrak{s})^2 - \sigma(W) > 0$ in Theorem 1.16 cannot be dropped in general. To see this, let us consider $-\mathbb{CP}^2$. The model reflection $\rho_S : -\mathbb{CP}^2 \to -\mathbb{CP}^2$ about the exceptional sphere S is a smooth involution, while any spin^c structure \mathfrak{s} on $-\mathbb{CP}^2$ is reversed by ρ_S . This is consistent in Theorem 1.16 as $c_1(\mathfrak{s})^2 - \sigma(-\mathbb{CP}^2) \leq 0$. More generally, let f be an orientation-preserving diffeomorphism of $W = \mathbb{CP}^2 \# n(-\mathbb{CP}^2)$ with $n \leq 8$ for which $f_* : H_2(W) \to H_2(W)$ is an involution. Then it follows from a result by Lee [25, Corollary 1.5, Remark 1.7] that f is topologically isotopic (hence homotopic) to some smooth involution. This is consistent in Corollary 1.18, as $\mathbb{CP}^2 \# n(-\mathbb{CP}^2)$ is not diffeomorphic to a manifold of the form $W' \# (-\mathbb{CP}^2)$ with spin W' with $\sigma(W') < 0$.

We also give a comparison result on the smooth and topologival Nielsen realization problems. As a topological version of the above realization problem, given a subgroup G of $\pi_0(\text{Homeo}(W))$, we say that G is realized in Homeo(W) if there is a section of the natural map $\text{Homeo}(W) \to \pi_0(\text{Homeo}(W))$ over G. One may ask whether there is a discrepancy between the realization problems in the smooth and topological categories. Such a comparison result was obtained first by Baraglia and the first author [4, Theorem 1.2], and it was generalized in [20, Theorem 1.3] by the first author. These results treated only spin 4-manifolds. The following theorem is the first comparison result for a non-spin 4-manifold:

Theorem 1.22. For p > 0 and $q, r \ge 0$, set $W = pK3 \# qS^2 \times S^2 \# r(-\mathbb{CP}^2)$. Then there exists a subgroup G of $\pi_0(\text{Diff}(W))$ of order 2 that satisfies the following properties:

- The group G is not realized in Diff(W).
- Define a subgroup $G' \subset \pi_0(\operatorname{Homeo}(W))$ to be the image of G under the map $\pi_0(\operatorname{Diff}(W)) \to \pi_0(\operatorname{Homeo}(W))$. Then G' is non-trivial and not realized in $\operatorname{Homeo}(W)$.

By the use of Theorem 1.1, we provide the two applications: regarding nonsmoothable and non-orientable surfaces and the Nielsen realization problem.

1.6. Non-smoothable and non-orientable surfaces. While non-smoothable orientable surfaces have been well studied by the use of Theorem A, 10/8 inequality and adjunction inequality combined with Freedman's theory (for example, see [14, Proof of Lemma 9.4.2 and Addendum 9.4.4]), to the best of our knowledge, construction of non-smoothable and non-orientable surfaces have not been developed well. By combining Rochlin's theorem, Theorem A, or 10/8 inequality with surgery technique, several constraints have been proven [23, 26, 36]. In [26], Levine–Ruberman–Strle gave genus bounds when the ambient 4-manifold is definite or the embedded surface is characteristic. Below we give non-orientable and non-smoothable surfaces, based on a constraint on surfaces that is not characteristic and the ambient 4-manifold is not definite:

Theorem 1.23. Let m, k > 0 and $n \ge 0$ and set

$$X = (m+n)\mathbb{CP}^2 \# (-n-8)\mathbb{CP}^2.$$

Then there exists a locally flat embedding of $k\mathbb{RP}^2$ into X that is not topologically isotopic to any smoothly embedded surface. More precisely:

- The homology class of the image of the embedding of kℝP² is zero in H₂(X; ℤ/2). In particular, the image of the embedding is not a characteristic surface, i.e. [kℝP²] ≠ w₂(X) in H²(X; ℤ/2).
- The normal Euler number is given by 4m + 2k.

Let us remark that Theorem 1.23 is almost a rephrase of the following application to detect non-smoothable involutions:

Theorem 1.24. Let m, k > 0 and $n \ge 0$ and set

 $W = (m+2n)\mathbb{CP}^2 \# (-m-2n-k-16)\mathbb{CP}^2.$

Then there exists an orientation-preserving non-smoothable locally linear involution $\iota: W \to W$ with $b_{\iota}^+(W) = m$, $b_{\iota}^-(W) = n + 8$.

Remark 1.25. Non-smoothable involutions on spin 4-manifolds have been extensively studied by Nakamura [40], Kato [17] and Baraglia [3]. Baraglia [3, Proposition 7.3] gave also a homological constraint on smooth involution on connected sums of copies of \mathbb{CP}^2 and $-\mathbb{CP}^2$. However, this constraint works for an involution ι on W with $b_{\iota}^+(W) = 0$, which is complementary to Theorem 1.24.

1.7. Structural conjecture. The invariants $\underline{\delta}_R$ and $\overline{\delta}_R$ are analogs of the invariants of homology 3-spheres $\underline{\delta}$ and $\overline{\delta}$ in Seiberg–Witten theory [50], which are conjectured to be equal to \underline{d} and \overline{d} in involutive Heegaard Floer homology [16]. One can ask if several analogous phenomena also hold for our invariants $\underline{\delta}_R$ and $\overline{\delta}_R$. We conjecture the following which can be seen as an analog of [49, Theorem 1.3] and [15, Theorem 1.2] for our invariants.

Conjecture 1.26. For any knot K, we have

$$\lim_{n \to \infty} \frac{\underline{\delta}_R(\#_n K)}{n} = \delta(K) \text{ and } \lim_{n \to \infty} \frac{\overline{\delta}_R(\#_n K)}{n} = \delta(K).$$

Moreover, as a 'real' version of Manolescu–Lidman's isomorphism [29], we conjecture the following:

Conjecture 1.27. For any oriented link L with non-zero determinant, we have

$$\begin{split} &\widehat{HMR}_*(L,\mathfrak{s}_L;\mathbb{Z}_2)\cong H^{\mathbb{Z}_2}_*(SWF(L);\mathbb{Z}_2),\\ &\widetilde{HMR}_*(L,\mathfrak{s}_L;\mathbb{Z}_2)\cong cH^{\mathbb{Z}_2}_*(SWF(L);\mathbb{Z}_2)\\ &\overline{HMR}_*(L,\mathfrak{s}_L;\mathbb{Z}_2)\cong tH^{\mathbb{Z}_2}_*(SWF(L);\mathbb{Z}_2), \end{split}$$

where $H_*^{\mathbb{Z}_2}$, $cH^{\mathbb{Z}_2}$ and $tH^{\mathbb{Z}_2}$ are \mathbb{Z}_2 -equivariant Borel, coBorel and Tate homologies respectively and HMR° are real monopole Floer homologies introduced in [28] for the spin structure \mathfrak{s}_L determined by orientation of L.

1.8. Structure of the paper. We finish off this introduction with an outline of the contents of this paper. In Section 2, we define a symmetry I of Seiberg–Witten equation for a given real spin^c structure. In Section 3, using the I-invariant part of Manolescu's Floer homotopy type, we introduce three invariants $d(Y, \mathfrak{s}, \iota)$, $\overline{d}(Y, \mathfrak{s}, \iota)$, and $\underline{d}(Y, \mathfrak{s}, \iota)$ for 3-manifolds with real spin^c structures (Y, \mathfrak{s}, ι) . Moreover, we prove several fundamental properties of these invariants including Frøyshov type inequality, connected sum formula, and duality formula. In Section 4, by setting Y to be the double branched covering space of an oriented link L in S^3 with non-zero determinant, we obtain invariants for L: d(L), $\overline{d}(L)$, and $\underline{d}(L)$. Moreover, we prove these invariants are χ -concordant invariants, and d(L) defines a homomorphism on the subgroup of the link concordance group generated by oriented based links with non-zero determinant. In Section 5, we prove all applications in the introduction

such as non-smoothable and non-orientable surfaces in 4-manifolds (Theorem 1.23), obstruction to Nielsen realization problem (Theorem 1.16) and non-orientable surfaces in D^4 bounded by torus knots (Theorem 1.14).

1.9. Acknowledgement. We would like to thank Joshua Sabloff for answering our question about his paper [48]. The authors also thank Kouki Sato, Tye Lidman, Peter Feller, Mike Miller for their helpful discussions. The authors would like to thank Jiakai Li and Ian Montague for enlightening discussions. The first author was partially supported by JSPS KAKENHI Grant Numbers 19K23412 and 21K13785, and Overseas Research Fellowships. The second author was supported by JSPS KAKENHI Grant Number 21J22979 and WINGS-FMSP program at the Graduate school of Mathematical Science, the University of Tokyo. The third author was supported by JSPS KAKENHI Grant Number 20K22319 and RIKEN iTHEMS Program.

2. Involution I

In this section, we explain the definition of the anti-linear involution I which covers the involution ι . Let W be an oriented smooth 4-manifold and \mathfrak{s} be a spin^c structure on W. Let $\iota : W \to W$ be a smooth involution that preserves the orientation of W and satisfies that $\iota^*\mathfrak{s} \cong \overline{\mathfrak{s}}$. We also assume that $W^{\iota} \neq \emptyset$ and $H^1(W,\mathbb{Z})$ is 0. In this section, we fix an ι -invariant metric on W.

Note that a spin^c structure \mathfrak{s} gives rise to a $\mathbb{Z}/2\mathbb{Z}$ -graded Clifford module $S = S^+ \oplus S^-$ with a hermitian metric, and \mathfrak{s} is determined uniquely by S [22, Section 1.1]. In this section, we identify \mathfrak{s} with S.

Firstly, we give an anti-linear involution on \mathfrak{s} if the involution ι satisfies a condition which is called odd type. We will prove later that such an involution is unique up to gauge transformations.

Definition 2.1. Let \mathfrak{s} be a spin^c structure on W and we will denote by $S = S^+ \oplus S^$ the spinor bundle of \mathfrak{s} and will denote by ρ its Clifford multiplication. Let $\overline{\mathfrak{s}}$ be a complex conjugate spin^c structure whose spinor bundle \overline{S} is the conjugate of S, while Clifford multiplication is unchanged as real-linear map. We define the pull-back of the spin^c structure \mathfrak{s} , say $\iota^*\mathfrak{s}$, as follows: The spinor bundle of $\iota^*\mathfrak{s}$ is ι^*S as a complex vector bundle. We define its Clifford multiplication by $(\xi, \phi) \mapsto$ $\rho((\iota^{-1})^*(\xi))\phi$ where $\xi \in T_x^*W$ and $\phi \in \iota^*S_x = S_{\iota(x)}$. To simplify the notation, we write ι when it should be ι^{-1} since ι is an involution.

We now give an anti-linear map on S which covers ι which may not be an involution.

Definition 2.2. Let $\iota^* \colon \iota^* \mathfrak{s} \to \mathfrak{s}$ be a natural map that covers ι . Note that this is a bijection. Let us take an isomorphism of the spin^c structures $\varphi \colon \overline{\mathfrak{s}} \to \iota^* \mathfrak{s}$ and complex conjugate $c \colon \mathfrak{s} \to \overline{\mathfrak{s}}$. Let us define $I_{\varphi} = \iota^* \circ \varphi \circ c$. This is an anti-linear map on S which covers the involution ι .

Note that the definition of the isomorphism between the spin^c structures \mathfrak{s} and \mathfrak{s}' is an isomorphism of complex vector bundle φ from the spinor bundle S of \mathfrak{s} to that of \mathfrak{s}' which satisfies $\varphi(\rho(\xi)\phi) = \rho(\xi)\varphi(\phi)$ for $\xi \in T_x^*W$ and $\phi \in S_x$. From the definition of I_{φ} , we can easily prove the lemma below.

Lemma 2.3. The anti-linear map I_{φ} satisfies

$$I_{\varphi}(\rho(\xi)\phi) = \rho(\iota^*(\xi))I_{\varphi}(\phi)$$

for all $\xi \in T_x W$ and $\phi \in S_x$. Moreover, I_{φ} preserves the hermitian metric on S.

We will prove that there is a gauge transformation u such that $(uI_{\varphi})^2 = \pm 1$. We prove two lemmas for preparation.

Lemma 2.4. There exists a gauge transformation $u_{\varphi} \colon W \to U(1)$ such that $I_{\varphi}^2 = u_{\varphi}$.

Proof. We see that I_{φ}^2 is an automorphism of the spin^c structure \mathfrak{s} . Thus I_{φ}^2 is a gauge transformation.

Lemma 2.5. The gauge transformation u_{φ} in Lemma 2.4 satisfies that $u_{\varphi}(\iota^{-1}(x)) = u_{\varphi}(x)$.

Proof. Note that for $\psi \in \Gamma(\iota^*\mathfrak{s})$, $\iota^*(\psi)(x) = \psi(\iota^{-1}(x))$. We see that for all $\phi \in \Gamma(\mathfrak{s})$, $I^3(\phi(x)) = \psi_{-1}(x)I_{-1}(\phi(x))$

$$I_{\varphi}^{3}(\phi(x)) = u_{\varphi}(x)I_{\varphi}(\phi(x))$$

= $I_{\varphi}(u_{\varphi}(x)\phi(x))$
= $\iota^{*} \circ \varphi(\overline{u_{\varphi}(x)}c(\phi(x)))$
= $\iota^{*}(\overline{u_{\varphi}(x)}\varphi \circ c(\phi(x)))$
= $\overline{u_{\varphi}(\iota^{-1}(x))}I_{\varphi}(\phi(x)).$

Thus we have $u_{\varphi}(\iota^{-1}(x)) = \overline{u_{\varphi}(x)}$.

Now we prove the following proposition.

Lemma 2.6. There exists a gauge transformation u such that $(uI_{\varphi})^2 = \pm 1$.

Proof. From Lemma 2.5 and the assumption that $H^1(W,\mathbb{Z})^{-\iota^*} = 0$, we have that there exists a real valued smooth function f such that $f(\iota^{-1}(x)) = -f(x)$ and $u_{\varphi}(x) = \pm \exp(if(x))$. We set $u(x) = \exp(-if(x)/2)$. Then we have

$$\begin{split} uI_{\varphi} \circ uI_{\varphi}(\phi(x)) &= u(x)I_{\varphi}(u(x)I_{\varphi}(\phi(x))) \\ &= u(x)\iota^*(\overline{u(x)}\varphi \circ c(\phi(x))) \\ &= u(x)\overline{u(\iota^{-1}(x))}I_{\varphi}^2(\phi(x)) \\ &= u(x)\overline{u(\iota^{-1}(x))}u_{\varphi}(x). \end{split}$$

From the definition of u, we see $\overline{u(\iota^{-1}(x))} = u(x)$ and $u(x)^2 = \pm \overline{u_{\varphi}(x)}$. Thus we have $(uI_{\varphi})^2 = \pm 1$.

From the following lemma, we see that an anti-linear involution on S which satisfies some conditions is unique up to ι invariant gauge transformations.

Lemma 2.7. Let $I_1, I_2: \mathfrak{s} \to \mathfrak{s}$ be an anti-linear map which satisfies that I_1, I_2 covers ι and I_1, I_2 is compatible with Clifford multiplication ρ in the following sense:

$$I_i(\rho(X)\phi(x)) = \rho(d\iota(X))I_i(\phi(x)). \ (i = 1, 2)$$

Then there exists a gauge transformation u_0 such that $I_2 = u_0I_1$. If I_1 and the gauge transformation u_0 satisfies that $I_1^2 = \pm 1$ and $(u_0I_1)^2 = \pm I_1^2$, then u_0 is an ι invariant gauge transformation and $(u_0I_1)^2 = I_1^2$.

Proof. From the assumption of the compatibility of the Clifford multiplication of I_1, I_2 , we see that $I_1 \circ I_2^{-1}$ is an automorphism of the spin^c structure \mathfrak{s} . Thus this is a gauge transformation u_0 .

Let us show the second half of the lemma. Note that $I_1(\phi)(\iota^{-1}(x)) = I_1(\phi(x))$ for all $\phi \in \Gamma(\mathfrak{s})$. We have

$$u_0(x)I_1(u_0(\iota^{-1}(x))I_1(\phi(x))) = u_0(x)\overline{u_0(\iota^{-1}(x))}I_1^2(\phi(x)) = \pm I_1^2(\phi(x)).$$

Thus u_0 satisfies that $u_0(x)\overline{u_0(\iota^{-1}(x))} = \pm 1$. From the assumption that $W^{\iota} \neq \emptyset$, we have $u_0(x)\overline{u_0(\iota^{-1}(x))} = 1$.

From Lemma 2.7, we see that the sign $(uI_{\varphi})^2 = \pm 1$ does not depend on the choice of φ and u. Moreover, If we have $(u'I_{\varphi'})^2 = \pm 1$ for another choice of a gauge transformation u' and an isomorphism $\varphi' : \bar{\mathfrak{s}} \to \iota^* \mathfrak{s}$, we have that uI_{φ} coincides with $u'I_{\varphi'}$ up to some ι -invariant gauge transformation. For abbreviation, we write uI_{φ} for I.

Definition 2.8. We say that an involution ι is of odd type for \mathfrak{s} when $I^2 = 1$.

Lemma 2.9. Suppose that $W^{\iota} \neq \emptyset$. Then W^{ι} is of codimension-2 if and only if ι is an odd type.

Proof. Let $x_0 \in W$ be an ι fixed point. Let $U(x_0)$ an ι invariant normal coordinate chart centered at a point x_0 . In the open set $U(x_0)$, there is the unique spin structure \mathfrak{s}_0 . We fix the trivialization $S|_{U(x_0)} \cong U(x_0) \times (\mathbb{H} \oplus \mathbb{H})$ of the spinor bundle of \mathfrak{s}_0 which satisfies that the Clifford action ρ is represented by 4×4 matrices as follows:

$$\rho(e_0) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \ \rho(e_1) = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \ \rho(e_1) = \begin{pmatrix} 0 & j \\ j & 0 \end{pmatrix}, \ \rho(e_1) = \begin{pmatrix} 0 & k \\ k & 0 \end{pmatrix}.$$

(In our convantion, ijk = 1.) In this chart, the restrictions of the spin^c structures $\mathfrak{s}, \overline{\mathfrak{s}}$ and $\iota^*\mathfrak{s}$ are isomorphic to the spin^c structure \mathfrak{s}_0 . Using this identification, we fix an isomorphism of the spin^c structure $\varphi: \overline{\mathfrak{s}} \to \iota^*\mathfrak{s}$. Let $\cdot j: \mathfrak{s}_0 \to \mathfrak{s}_0$ is an anti-linear map given by the right multiplication of j. Then we see that $\tilde{\iota} = \iota^* \circ \varphi \circ c \circ (\cdot j): \mathfrak{s}_0 \to \mathfrak{s}_0$ is a lift of the involution ι to the spin^c structure \mathfrak{s}_0 . Therefore $\tilde{\iota}_{x_0}$ can be written by $[g, u] \in \mathrm{Spin}^c(4) \cong (\mathrm{Spin}(4) \times U(1))/\{(1, 1), (-1, -1)\}$ where g is a lift of $d\iota_{x_0}$ to $\mathrm{Spin}(4)$. Thus we see that for $\phi \in S_{x_0} \cong \mathbb{H} \oplus \mathbb{H}$,

$$I_{x_0}(\phi) = g\phi j u^{-1}$$

and we have $I^2(\phi) = -g^2 \phi$. Hence ι is odd if and only if $g^2 = -1$. One can easily check that $g^2 = -1$ if and only if fixed point sets of ι is codimension 2.

From Lemma 2.9, we may obtain a convenient sufficient condition that ι is an odd type:

Lemma 2.10. Let W be a closed, oriented smooth 4-manifold. Let $\iota : W \to W$ be an orientation-preserving smooth involution. Let \mathfrak{s} be a spin^c structure on W and suppose that $\iota^*\mathfrak{s} = \overline{\mathfrak{s}}$. If $\sigma_\iota(W) \neq \sigma(W)/2$, then ι is an odd type. *Proof.* Assume that ι is an even type. By the *G*-signature theorem, $\sigma_{\iota}(W)$ can be obtained by adding $\sigma(W)/2$ to contributions from W^{ι} . However, by Lemma 2.9, W^{ι} consists only of isolated points, and the contribution from isolated fixed points are zero for a general involution.

As well as for involutions on 4-manifolds discussed until here, we can repeat similar arguments for involutions on 3-manifolds. We summarize it below:

Theorem 2.11. Let Y be a closed, oriented three-manifold and t be a spin^c structure on Y. Let $\iota: Y \to Y$ be an involution such that $Y^{\iota} \neq \emptyset$, $\iota^* \mathfrak{t} \cong \overline{\mathfrak{t}}$, and $H^1(Y,\mathbb{Z})^{-\iota^*} = 0$. Then we have an anti-linear map $I: \mathfrak{t} \to \mathfrak{t}$ such that I covers ι and I is compatible with Clifford multiplication ρ in the following sense:

$$I(\rho(X)\phi(x)) = \rho(d\iota(X))I(\phi(x)).$$

The choice of I is unique up to ι invariant gauge transformations. Moreover, I satisfies that $I^2 = \pm 1$ and $I^2 = 1$ if and only if the fixed point set of ι is codimension 2.

From now, we assume that ι is of odd type.

Next, we show that the Seiberg–Witten equation is equivariant with the involution $-\iota^* \oplus I$.

Proposition 2.12. Let us define an involution on $\Omega^1(W) \oplus \Gamma(S^+)$ and $\Omega^+(W) \oplus \Gamma(S^-)$ by $-\iota^* \oplus I$. We have the Seiberg–Witten equations with ι -invariant Riemannian metric on W is equivariant with this involution.

Proof. Let A'_0 be a spin^c connection on \mathfrak{s} . We define the covariant derivative $d_{A_0}: \Omega(W) \otimes \Gamma(S) \to \Omega(W) \otimes \Gamma(S)$ to be

$$d_{A_0} := \frac{1}{2} (d_{A'_0} + (\iota^* \otimes I) \circ d_{A'_0} \circ (\iota^* \otimes I)).$$

It is easy to check that this is an $\iota^* \otimes I$ invariant spin^c connection on \mathfrak{s} . Moreover, we have that the curvature form $F_{A_0^{\tau}}$ of det S^+ induced by the connection A_0 satisfies that $\iota^* F_{A_0^{\tau}} = -F_{A_0^{\tau}}$ since F_{A_0} is an imaginary-valued 2-form and I is an anti-linear. We set A_0 to be a reference connection. Let $A = A_0 + \sqrt{-1a}$. We only need to show that the non-linear terms of the Seiberg–Witten equations are equivariant with the involution. From Lemma 2.3, we have

(7)
$$I(\sqrt{-1}\rho(a)\phi) = -\sqrt{-1}I(\rho(a)\phi) = -\sqrt{-1}\rho(\iota^*a)I(\phi)$$

for any $a \in \Omega^*(W)$ and $\phi \in \Gamma(S^+)$. Thus we have that the Dirac equation is equivariant to the involution.

Next, we show that the equation of curvature

$$F_{A^{\tau}}^{+} = \sqrt{-1}d^{+}a + F_{A_{0}^{\tau}}^{+} = -\sqrt{-1}\tau(\phi,\phi)$$

is equivariant under the involution. The quadratic form $\tau: S^+ \otimes S^+ \to \Lambda^+$ is characterized by the following relation:

$$\langle b, \tau(\phi_1, \phi_2) \rangle_{\Lambda^+} = -\langle \sqrt{-1}\rho(b)\phi_1, \phi_2 \rangle_{S^+}$$

where $b \in \Omega^+(W)$ and $\phi_1, \phi_2 \in \Gamma(S^+)$. The inner product $\langle, \rangle_{\Lambda^+}$ and \langle, \rangle_{S^+} are invariant under the involution ι^* and I respectively. From (7), we have

$$\begin{split} \langle b, \iota^* \tau(\phi_1, \phi_2) \rangle_{\Lambda^+} &= \langle \iota^* b, \tau(\phi_1, \phi_2) \rangle_{\Lambda^+} \\ &= -\langle \sqrt{-1} \rho(\iota^* b) \phi_1, \phi_2 \rangle_{S^+} \\ &= -\langle I(\sqrt{-1} \rho(\iota^* b) \phi_1), I(\phi_2) \rangle_{S^+} \\ &= \langle \sqrt{-1} \rho(b) I(\phi_1), I(\phi_2) \rangle_{S^+} \\ &= -\langle b, \tau(I(\phi_1), I(\phi_2)) \rangle_{\Lambda^+}. \end{split}$$

This completes the proof.

We can prove that the Seiberg–Witten equation on 3-manifolds with involution ι is invariant under the involution $-\iota^* \oplus I$ in a similar way.

3. FLOER HOMOTOPY TYPE OF RATIONAL HOMOLOGY 3-SPHERES WITH INVOLUTION AND FRØYSHOV TYPE INVARIANTS

In [21], we defined a (\mathbb{Z}_4 -equivariant) "doubled" Seiberg–Witten Floer stable homotopy type of spin rational homology 3-spheres with involution $DSWF_{\mathbb{Z}_4}(Y, \mathfrak{t}, \iota)$. The doubling construction was considered mainly to define a K-theoretic Frøyshovtype invariant easily. In this section, we define Seiberg–Witten Floer stable homotopy type of real spin^c rational homology 3-spheres $SWF_{\mathbb{Z}_2}(Y, \mathfrak{t}, \iota)$, without taking double, and define a Frøyshov type invariant

$$\delta_R(Y,\mathfrak{t},\iota)\in\frac{1}{16}\mathbb{Z}$$

applying the \mathbb{Z}_2 -equivariant ordinary cohomology to $SWF_{\mathbb{Z}_2}(Y, \mathfrak{t}, \iota)$. Moreover, when \mathfrak{t} is spin, we also define two Frøyshov type invariants

$$\underline{\delta}_{R}(Y,\mathfrak{t},\iota), \bar{\delta}_{R}(Y,\mathfrak{t},\iota) \in \frac{1}{16}\mathbb{Z}$$

applying \mathbb{Z}_4 -equivariant ordinary cohomology to $SWF_{\mathbb{Z}_4}(Y, \mathfrak{t}, \iota)$. These invariants are analogues of Stoffregen's invariants $\underline{\delta}_R$ and $\overline{\delta}_R$. Throughout this section, we consider cohomologies with coefficient $\mathbb{F} = \mathbb{Z}_2$.

3.1. **Representations.** First let us consider \mathbb{Z}_2 -representations. Let \mathbb{R} be the trivial real 1-dimensional representation of \mathbb{Z}_2 , and \mathbb{C} denote the complex 1-dimensional representation of \mathbb{Z}_2 defined as the scalar multiplication of $\mathbb{Z}_2 = \{1, -1\}$. For a finite-dimensional vector space V, let V^+ denote the one-point compactification of V.

Recall that we defined the group G to be the cyclic group of order 4 generated by $j \in Pin(2)$, i.e.

$$G = \{1, j, -1, -j\}.$$

Define a subgroup H of G by

$$H = \{1, -1\} \subset G.$$

Let \mathbb{R} denote the trivial 1-dimensional real representation of G. Let \mathbb{R} be the 1-dimensional real representation space of G defined by the surjection $G \to \mathbb{Z}_2 = \{1, -1\}$ and the scalar multiplication of \mathbb{Z}_2 on \mathbb{R} .

Let \mathbb{C} be a 1-dimensional complex representation of G defined via the surjection $G \to \mathbb{Z}_2$ and the scalar multiplication of \mathbb{Z}_2 on \mathbb{C} . Note that, for an even natural number s, say 2t, there is an isomorphism of real representations $\mathbb{R}^{2s} \cong \mathbb{C}^t$. We introduce also a G-representation \mathbb{C} (the same notation of the complex number) which is the complex 1-dimensional representation defined by assigning $j \in G$ to i in \mathbb{C} .

3.2. Numerical invariants d, \underline{d} , \overline{d} . We first define the main ingredient of the Frøyshov invariant using \mathbb{Z}_2 -equivariant cohomology, following Stoffregen's formulation [50].

Definition 3.1. Let G be a group and H be a subgroup of G. Let \mathcal{V} be a coutable direct sum of a fixed 1-dimensional real representation G. Let X be a pointed finite G-CW complex. We call X a space of type (G, H)-SWF if

- X^H is G-homotopy equivalent to V^+ , where V is a finite dimensional subspace of \mathcal{V} .
- *H* acts freely on $X \setminus X^H$.

The dimension dim V is called the *level* of X. We put $\mu(X) \in \mathbb{Q}/2\mathbb{Z}$ by $\mu(X) = \dim V/2 \mod 2$.

The situation we have in mind is either

$$(G,H) = (\mathbb{Z}_2,\mathbb{Z}_2)$$
 or $(\mathbb{Z}_4,\mathbb{Z}_2)$.

We often drop H in the notation in (G, H).

First, let us consider $(G, H) = (\mathbb{Z}_2, \mathbb{Z}_2)$. Note that

$$\tilde{H}^*_{\mathbb{Z}_2}(S^0) \cong \mathbb{Z}_2[W]_{\mathbb{Z}_2}$$

where W is of degree 1. For a space X of type \mathbb{Z}_2 -SWF, define

$$d(X) = \min\left\{ \begin{array}{l} m \ge 0 \end{array} \middle| \exists x \in \tilde{H}^m_{\mathbb{Z}_2}(X), W^l x \ne 0 \; (\forall l \ge 0) \end{array} \right\}$$
$$= \min\left\{ \begin{array}{l} m \ge 0 \end{array} \middle| \exists x \in \tilde{H}^m_{\mathbb{Z}_2}(X), 0 \ne \iota^* x \in \widetilde{H}^*_G(X^H) \end{array} \right\}.$$

(See [50, Equation (19)].) By an equivariant localization theorem (see, e.g., [50, Theorem 2.3]), we have $d(X) < +\infty$.

Next, let us consider $(G, H) = (\mathbb{Z}_4, \mathbb{Z}_2)$. In this case, we have

$$\widetilde{H}^*_G(S^0) \cong \mathbb{Z}_2[U,Q]/(Q^2=0),$$

where deg U = 2 and deg Q = 1. For a space X of type \mathbb{Z}_4 -SWF, define

$$\overline{d}(X) = \min\left\{ m \equiv 2\mu(X) \ge 0 \pmod{2} \mid \exists x \in \tilde{H}^m_{\mathbb{Z}_4}(X), \ U^l x \neq 0 \ (\forall l \ge 0) \right\}$$

and

$$\underline{d}(X) = \min\left\{ m \equiv 2\mu(X) + 1 \ge 0 \pmod{2} \mid \exists x \in \tilde{H}^m_{\mathbb{Z}_4}(X), \ U^l x \neq 0 \ (\forall l \ge 0) \right\} - 1.$$

Again, by an equivariant localization theorem (see, e.g., [50, Theorem 2.3]), we have that $\overline{d}(X), \underline{d}(X) < \infty$.

We also use alternative descriptions of invariants \underline{d} and \overline{d} by using *infinity version* of equivariant cohomology. For a space X of type G-SWF, we define

$$\infty \widetilde{H}_G^*(X) := \begin{cases} \operatorname{Im}(\widetilde{H}_G(X) \to U^{-1}\widetilde{H}_G(X)) \text{ if } G = \mathbb{Z}_4\\ \operatorname{Im}(\widetilde{H}_G(X) \to W^{-1}\widetilde{H}_G(X)) \text{ if } G = \mathbb{Z}_2 \end{cases}$$

We have the following classification result of ideals of $U^{-1}\mathbb{Z}_2[U,Q]/(Q^2=0)$.

Lemma 3.2. Any graded ideal \mathcal{J} of $U^{-1}\mathbb{Z}_2[U,Q]/(Q^2=0)$ such that $U^{-1}\mathcal{J} = U^{-1}\mathbb{Z}_2[U,Q]/(Q^2=0)$ have the following form:

$$\mathcal{J} = (U^i, QU^j)$$

for some $i \ge j \ge 0$.

Proof. The proof is essentially the same as the proof of [34, Lemma 2.8].

If
$$\infty \widetilde{H}_{G}^{n+*}(X) = (U^{i}, QU^{j})$$
, then we have
(8) $\underline{d}(X) = i + n \text{ and } \overline{d}(X) = j + n.$

In particular, this expression (8) enables us to check

$$\underline{d}(X) \le \overline{d}(X).$$

Moreover, we have the following properties with respect to stabilizations:

Lemma 3.3. Let X be a space of type $G = \mathbb{Z}_4$ SWF, and V a representation of $G = \mathbb{Z}_4$. Then, we have

(9)
$$\underline{d}(\Sigma^V X) = \underline{d}(X) + \dim V \text{ and } \overline{d}(\Sigma^V X) = \overline{d}(X) + \dim V.$$

Similarly, for a space X of type $G = \mathbb{Z}_2$ SWF and a \mathbb{Z}_2 -representation V, we have (10) $d(\Sigma^V X) = d(X) + \dim V.$

Proof. As it is proven in [34, Proposition 2.2], for any finite-dimensional representation V of $G = \mathbb{Z}_4$ or \mathbb{Z}_2 , we have the suspension isomorphism of $\widetilde{H}_G(S^0)$ -modules:

(11)
$$\widetilde{H}^*_G(\Sigma^V X; \mathbb{Z}_2) \cong \widetilde{H}^{*-\dim V}_G(X; \mathbb{Z}_2)$$

The equations (9) and (10) follow from (11).

Later we shall use the following elementary lemmas. Recall that, given a G-vector bundle $E \to B$, the (mod 2) G-equivariant Euler class $e_G(E) \in H^*_G(B)$ is defined by

$$e_G(E) = e(EG \times_G E \to EG \times_G B) \in H^*_G(B).$$

Lemma 3.4. The cohomology class $W \in \tilde{H}^*_{\mathbb{Z}_2}(S^0)$ coincides with $e_{\mathbb{Z}_2}(\tilde{\mathbb{R}})$, where $\tilde{\mathbb{R}}$ is regarded as a \mathbb{Z}_2 -vector bundle over {pt}.

Proof. By definition, we have

$$e_{\mathbb{Z}_2}(\tilde{\mathbb{R}}) = e(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} \tilde{\mathbb{R}} \to B\mathbb{Z}_2) \in \tilde{H}^1_{\mathbb{Z}_2}(S^0).$$

The right-hand side coincides with $w_1(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} \tilde{\mathbb{R}})$ and it is the generator of $H^1(B\mathbb{Z}_2) = \tilde{H}^1_{\mathbb{Z}_2}(S^0)$. This completes the proof. \Box

Lemma 3.5. The cohomology class $Q \in \tilde{H}^*_{\mathbb{Z}_4}(S^0)$ is the image of $W \in \tilde{H}^*_{\mathbb{Z}_2}(S^0)$ under the natural map $\tilde{H}^*_{\mathbb{Z}_2}(S^0) \to \tilde{H}^*_{\mathbb{Z}_4}(S^0)$ induced from the surjection $\mathbb{Z}_4 \to \mathbb{Z}_2$.

Proof. The surjection $\mathbb{Z}_4 \to \mathbb{Z}_2$ induces a surjection $\tilde{H}_1(B\mathbb{Z}_4) \to \tilde{H}_1(B\mathbb{Z}_2)$. Passing to the dual, this induces the isomorphism of the cohomologies $\tilde{H}^1(B\mathbb{Z}_2) \to \tilde{H}^1(B\mathbb{Z}_4)$, thus the generator W of $\tilde{H}^1(B\mathbb{Z}_2)$ maps to the generator Q of $\tilde{H}^1(B\mathbb{Z}_4)$.

For S^0 , the simplest example of the space of type SWF, it is easy to see that the above invariants coincide,

$$d(S^0) = \underline{d}(S^0) = \overline{d}(S^0) = 0.$$

Below we exhibit an example for which the invariants d and \underline{d} , \overline{d} are distinct, following [34, Example 2.10].

Example 3.6. Set $G = \mathbb{Z}_4$, and let \tilde{G} denote the unreducible suspension of G. We regard \tilde{G} as a based space by choosing one of the cone point of \tilde{G} as the base point. This space \tilde{G} is obviously a space of type \mathbb{Z}_4 -SWF. We claim that

(12)
$$\underline{d}(\tilde{G}) = 0, \quad d(\tilde{G}) = 1, \quad \overline{d}(\tilde{G}) = 2.$$

To see this, first note that the \mathbb{Z}_2 -invariant part of \tilde{G} is given by $\tilde{G}^{\mathbb{Z}_2} = S^0$. The cone of the inclusion map $S^0 \hookrightarrow \tilde{G}$ is given by the reduced suspension $\Sigma^{\mathbb{R}}G_+$ of $G_+ = G \sqcup \text{pt}$. Thus we obtain the long exact sequence

(13)
$$\cdots \to H^*_G(\Sigma^{\mathbb{R}}G_+) \to \tilde{H}^*_G(\tilde{G}) \to H^*_G(S^0) \to H^{*+1}_G(\Sigma^{\mathbb{R}}G_+) \to \cdots$$

It follows from this long exact sequence combined with

$$\tilde{H}_G^{*+1}(\Sigma^{\mathbb{R}}G_+) \cong \tilde{H}_G^*(G_+) \cong H^*(\mathrm{pt})$$

that the restriction map $\tilde{H}_{G}^{m}(\tilde{G}) \to \tilde{H}_{G}^{m}(S^{0})$ is isomorphic if $m \geq 2$. On the other hand, it is clear that $\tilde{H}_{G}^{0}(\tilde{G}) = 0$. This combined with the above long exact sequence (13) implies that the restriction $\tilde{H}_{G}^{1}(\tilde{G}) \to \tilde{H}_{G}^{1}(S^{0})$ is isomorphic. The claims that $\underline{d}(\tilde{G}) = 0$ and that $\overline{d}(\tilde{G}) = 2$ immediately follow from these computations.

Next, we prove the claim $d(\tilde{G}) = 1$. First, we obviously have $\tilde{H}^0_{\mathbb{Z}_2}(\tilde{G}) = 0$. Second, by combining a long exact sequence for $\tilde{H}^*_{\mathbb{Z}_2}$ analogous to (13) with that

$$\tilde{H}^2_{\mathbb{Z}_2}(\Sigma^{\mathbb{R}}G_+) \cong \tilde{H}^1_{\mathbb{Z}_2}(G_+) \cong H^1(\mathrm{pt} \sqcup \mathrm{pt}) = 0,$$

we can see that the restriction map $\tilde{H}^1_{\mathbb{Z}_2}(\tilde{G}) \to \tilde{H}^1_{\mathbb{Z}_2}(S^0)$ is surjective. The claim $d(\tilde{G}) = 1$ directly follows from these observations.

Next we see a certain monotonicity for the quantities $d, \underline{d}, \overline{d}$:

Lemma 3.7. Let X and X' be spaces of type \mathbb{Z}_2 -SWF at the same level. Suppose that there exists a pointed \mathbb{Z}_2 -equivariant map $f : X \to X'$ whose \mathbb{Z}_2 -fixed-point set map is a homotopy equivalence. Then we have

$$d(X) \le d(X').$$

Similarly, for spaces X, X' of type \mathbb{Z}_4 -SWF at the same level, if there exists a pointed \mathbb{Z}_4 -equivariant map $f: X \to X'$ whose \mathbb{Z}_2 -fixed-point set map is a \mathbb{Z}_4 -homotopy equivalence, then we have

$$\overline{d}(X) \le \overline{d}(X'), \quad \underline{d}(X) \le \underline{d}(X').$$

Proof. This is a standard argument, but we give a sketch of the proof for the reader's convenience. We first show the claim for the invariant d. Set $(G, H) = (\mathbb{Z}_2, \mathbb{Z}_2)$. Consider the following commutative diagram:

$$\begin{array}{cccc} X^H & \stackrel{f^H}{\longrightarrow} & (X')^H \\ \downarrow & & \downarrow \\ X & \stackrel{f}{\longrightarrow} & X'. \end{array}$$

Here the vertical maps are inclusions. By applying $\widetilde{H}_G^*,$ one has a commutative diagram

$$\widetilde{H}^*_G(X^H) \xleftarrow{(f^H)^*} \widetilde{H}^*_G((X')^H)
\uparrow \qquad \uparrow \qquad \uparrow
\widetilde{H}^*_G(X) \xleftarrow{f^*} \widetilde{H}^*_G(X').$$

It follows from the assumption that $(f^H)^*$ is a homotopy equivalence combined with the equivariant localization theorem that f^* is an isomorphism in large enough degree. From this combined with that f^* commutes with the W-action, the commutative diagram above implies that

(14)
$$f^*\left(\left\{ x' \in \tilde{H}^*_G(X') \mid W^l x' \neq 0 \; (\forall l \ge 0) \right\} \right) \subset \left\{ x \in \tilde{H}^*_G(X) \mid W^l x \neq 0 \; (\forall l \ge 0) \right\}.$$

The desired inequality $d(X) \leq d(X')$ follows from this.

For the invariants $\overline{d}, \underline{d}$, we may repeat the above argument for $(G, H) = (\mathbb{Z}_4, \mathbb{Z}_2)$, with replacing W with U in (14).

We also note a smash formula of d:

Lemma 3.8. Let X and X' be spaces of type \mathbb{Z}_2 -SWF at the level l and l'. Then, $X \wedge X'$ becomes a space of type \mathbb{Z}_2 -SWF at the level l + l' and

$$d(X \wedge X') \ge d(X) + d(X')$$

holds.

Proof. Set $(G, H) = (\mathbb{Z}_2, \mathbb{Z}_2)$. The following commutative diagram is a key ingredient in the proof:

$$\widetilde{H}^*_G((X \wedge X')^H) \xrightarrow{=_{\#}H} \widetilde{H}^*_G(X^H) \otimes \widetilde{H}^*_G((X')^H) \\
\xrightarrow{(\iota \wedge \iota')^*} \downarrow \qquad \qquad \iota^* \otimes \iota'^* \uparrow \\
\widetilde{H}^*_G(X \wedge X') \xrightarrow{\#} \widetilde{H}^*_G(X) \otimes \widetilde{H}^*_G(X'),$$

where the horizontal maps are induced from the inclusion maps $X \to X \wedge X'$ and $X' \to X \wedge X'$ and the vertical maps are induced from inclusions $X^H \to X$ and $(X')^H \to X'$. Take an element $x \in \tilde{H}^n_G(X \wedge X')$ so that $\iota^{\otimes} x \neq 0$ and $d(X \wedge X') = n$. Then, for an expression $(\iota \otimes \iota')^* \circ \#(x) = \sum_i a_i \otimes b_i$ with $a_i \in \tilde{H}^*_G(X^H) \setminus \{0\}$ and $b_i \in \tilde{H}^*_G((X')^H) \setminus \{0\}$, each of a_1 and b_1 are images of elements of $\tilde{H}^*_G(X)$ and $\tilde{H}^*_G(X')$ under the restriction maps, respectively. Then, by the definition of d, one has $d(X) \leq \deg(a_1)$ and $d(X') \leq \deg(b_1)$. On the other hand, we have $\deg(a_1) + \deg(b_1) = \deg(x) = n$. This completes the proof. \Box

At the end of this subsection, we provide a key ingredient of a Theorem B type theorem in our theory.

Lemma 3.9. For spaces X, X' of type \mathbb{Z}_4 -SWF, if there exists a pointed \mathbb{Z}_4 equivariant map $f : X \to X'$ whose \mathbb{Z}_2 -fixed-point set map is induced from a \mathbb{Z}_4 -injective linear map whose image is of codimension-1, then we have

$$\underline{d}(X) \le \overline{d}(X')$$

Proof. Set $(G, H) = (\mathbb{Z}_4, \mathbb{Z}_2)$. Let s, s' denote the levels of X and X' respectively. By the assumption on f^H , we have that s + 1 = s'. Recall that the \mathbb{Z}_4 -equivariant Thom isomorphism implies that $\widetilde{H}^*_G(X^H) \cong \widetilde{H}^*_G(S^0)_{[s]}$, where [s] denotes the degree s-shift. Let $[s] : \widetilde{H}^*_G(S^0) \to \widetilde{H}^*_G(S^0)_{[s]}$ denote the degree s-map defined by just the degree shift.

As in the proof of Lemma 3.7, consider the commutative diagram

(15)

$$\begin{aligned}
\widetilde{H}_{G}^{*}(S^{0}) & \widetilde{H}_{G}^{*}(S^{0}) \\
[s] \downarrow & [s'] \downarrow \\
\widetilde{H}_{G}^{*}(X^{H}) \stackrel{(f^{H})^{*}}{\longleftarrow} \widetilde{H}_{G}^{*}((X')^{H}) \\
[s] i^{*} \uparrow & (i')^{*} \uparrow \\
\widetilde{H}_{G}^{*}(X) \stackrel{(i')^{*}}{\longleftarrow} \widetilde{H}_{G}^{*}(X'),
\end{aligned}$$

where $i: X^H \to X$ and $i': (X')^H \to X'$ are inclusions.

We claim that

$$[-s] \circ (f^H)^* \circ [s'] : \widetilde{H}^*_G(S^0) \to \widetilde{H}^*_G(S^0)$$

is given by multiplication by Q. To see this, note that $\widetilde{H}^*_G(X^H)$ and $\widetilde{H}^*_G(X^H)$ are rank 1 free $\widetilde{H}^*_G(S^0)$ -modules generated by the *G*-equivariant mod 2 Thom classes τ_G and τ'_G of $\mathbb{R}^s \to \{\text{pt}\}$ and $\mathbb{R}^{s'} \to \{\text{pt}\}$, respectively. By the assumption on f^H , we have

$$(f^H)^*\tau'_G = e_G(\tilde{\mathbb{R}})\tau_G$$

where $e_G(\tilde{\mathbb{R}})$ is the *G*-equivariant mod 2 Euler class of $\tilde{\mathbb{R}} \to \{\text{pt}\}$. Since *G* acts on $\tilde{\mathbb{R}}$ as $\{\pm 1\}$ -multiplication, we have that $e_G(\tilde{\mathbb{R}}) = Q$ from Lemmas 3.4 and 3.5. This proves the claim.

In each of $\widetilde{H}^*_G(X^H)$ and $\widetilde{H}^*_G((X')^H)$, there are exactly two U-towers that are degree shifts of $\{U^l\}_{l\geq 0}$ and $\{QU^l\}_{l\geq 0}$ in $\widetilde{H}^*_G(S^0)$. For i = 0, 1, let $\mathcal{T}_i \subset \widetilde{H}^*_G(X^H)$

and $\mathcal{T}'_i \subset \widetilde{H}^*_G((X')^H)$ denote these towers such that every $x \in \mathcal{T}_i$ and $x' \in \mathcal{T}'_i$ satisfy that

$$\deg x \equiv 2\mu(X) + i$$
 and $\deg x' \equiv 2\mu(X') + i \mod 2$.

It follows from the above claim that the restriction of $(f^H)^*$ on \mathcal{T}'_0 gives a bijection onto \mathcal{T}_1 . This combined with the commutative diagram (15) implies that $\underline{d}(X) \leq \overline{d}(X')$.

3.3. **Duality.** We prove a duality theorem for d, \overline{d} , and \underline{d} , which can be seen as an analog of [34, Proposition 2.13]. In order to prove the duality, we use the following duality relations among equivariant cohomologies. See [37, Section XVI. 8], [34, Subsection 2.2] for the definition of (equivariant) V-dual.

If X and X' are V-dual with $m = \dim V$, the coBorel cohomology of X' can be viewed realized as follow:

$$c\tilde{H}^G_*(X;\mathbb{Z}_2) \cong \tilde{H}^{m-*}_G(X';\mathbb{Z}_2).$$

Moreover, there is a long exact sequence relating Borel homology with co-Borel and Tate homologies:

$$\cdots \to c\tilde{H}^G_*(X;\mathbb{Z}_2) \to t\tilde{H}^G_*(X;\mathbb{Z}_2) \to \tilde{H}^G_{*-1}(X;\mathbb{Z}_2) \to \cdots$$

For our purpose, it is also convenient to describe d, \overline{d} , and \underline{d} in terms of homologies as in [34, (17), (18), and (19)]. We first define

$$\infty \widetilde{H}^G_*(X) := \bigcap_{m \ge 0} \operatorname{Im} U^m : \widetilde{H}^G_*(X) \to \widetilde{H}^G_{*-2m}(X).$$

Then, alternatively, for a space X of type $G = \mathbb{Z}_4$ -SWF of level s, we can write

$$\begin{split} \underline{d}(X) &= \min \left\{ \left. r \equiv s \mod 3 \right| \, \exists x, \ 0 \neq x \in \infty \widetilde{H}_r^G(X) \right\}, \\ \overline{d}(X) &= \min \left\{ \left. r \equiv s + 1 \mod 3 \right| \, \exists x, \ 0 \neq x \in \infty \widetilde{H}_r^G(X) \right\}, \\ d(X) &= \min \left\{ \left. r \equiv s \mod 2 \right| \, \exists x, \ 0 \neq x \in \infty \widetilde{H}_r^{\mathbb{Z}_2}(X) \right\}. \end{split}$$

Lemma 3.10. Let G be \mathbb{Z}_2 . If X and X' are equivariantly V-dual, then

$$d(X) = \dim V - d(X')$$

hold. Moreover, if $G = \mathbb{Z}_4$, then we have

$$\underline{d}(X) = \dim V - \overline{d}(X').$$

Proof. Suppose the level of X is s and $m = \dim V$. Then the level of X' is m - s by duality. The proof for d is an easier version of the proof for <u>d</u> and <u>d</u>. Thus, we only write a proof of the second equality. The main strategy is the same as the proof of [34, Proposition 2.13]. We have the exact sequence connecting Borel, coBorel and Tate cohomologies:

$$\cdots \to c\tilde{H}^G_*(X;\mathbb{Z}_2) \to t\tilde{H}^G_*(X;\mathbb{Z}_2) \to \tilde{H}^G_{*-1}(X;\mathbb{Z}_2) \to \cdots$$

Since X and X' are V-dual, from the associated isomorphism

$$cH^G_*(X;\mathbb{Z}_2) \cong H^{m-*}_G(X';\mathbb{Z}_2),$$

we have

$$\cdots \to \tilde{H}_G^{m-*}(X';\mathbb{Z}_2) \to \mathrm{t}\tilde{H}_*^G(X;\mathbb{Z}_2) \to \tilde{H}_{*-1}^G(X;\mathbb{Z}_2) \to \cdots$$

Moreover, the localization theorem implies

$$t\tilde{H}^G_*(X;\mathbb{Z}_2) \cong (U^{-1}\mathbb{Z}_2[U,Q]/Q^2 = 0)_{s+2}$$

Then by the same arguemnt given in [34, Proof of Propsotion 2.13], after shifting degrees by s + 2, the exact sequence above induces

$$\cdots \to \infty \widetilde{H}_G^{m-s-2+*}(X') \to U^{-1}\mathbb{Z}_2[U,Q]/(Q^2=0) \to \widetilde{H}_{*+s}^G(X) \to \cdots$$

which connects the infinity versions of homology and cohomology directly. Since we have a concrete formula of infinity versions using \underline{d} and \overline{d} . This enables us to write

$$\infty \widetilde{H}_{G}^{m-s-2+*}(X') = \begin{cases} \mathbb{Z}_{2} \text{ if } * = m - s - 2 + \underline{d}(X') + 2j \ j \ge 0, \\ \mathbb{Z}_{2} \text{ if } * = m - s - 3 + \overline{d}(X') + 2j \ j \ge 0, \\ 0 \text{ if otherwise,} \end{cases}$$
$$\infty \widetilde{H}_{*+s}^{G}(X) = \begin{cases} \mathbb{Z}_{2} \text{ if } * = -s + \underline{d}(X) + 2j \ j \ge 0, \\ \mathbb{Z}_{2} \text{ if } * = -s + 1 + \overline{d}(X) + 2j \ j \ge 0, \\ 0 \text{ if otherwise.} \end{cases}$$

This completes the proof.

3.4. Spectrum classes. Let (G, H) be either $(\mathbb{Z}_2, \mathbb{Z}_2)$ or $(\mathbb{Z}_4, \mathbb{Z}_2)$, where, in the latter case, \mathbb{Z}_2 is regarded as a subgroup of \mathbb{Z}_4 in a natural way. When $(G, H) = (\mathbb{Z}_2, \mathbb{Z}_2)$, we set

$$\mathcal{V} = \oplus_{\mathbb{N}} \mathbb{R}, \quad \mathcal{W} = \oplus_{\mathbb{N}} \mathbb{R},$$

and When $(G, H) = (\mathbb{Z}_4, \mathbb{Z}_2)$, we set

$$\mathcal{V}=\oplus_{\mathbb{N}} ilde{\mathbb{R}},\quad \mathcal{W}=\oplus_{\mathbb{N}}\mathbb{C}.$$

Following [33, Section 4], consider a triple (X, m, n), where X is a space of type G-SWF, and $m \in \mathbb{Z}$ and $n \in \mathbb{Q}$. For a finite dimensional representation V of G and a pointed G-space X, we denote by $\Sigma^V X$ the suspension $V^+ \wedge X$. Set

$$\mathbb{K}_G = \begin{cases} \mathbb{\bar{R}} & \text{if} \quad G = \mathbb{Z}_2 \\ \mathbb{C} & \text{if} \quad G = \mathbb{Z}_4, \end{cases} \qquad k_G = \dim_{\mathbb{R}} \mathbb{K}_G = \begin{cases} 1 & \text{if} \quad G = \mathbb{Z}_2 \\ 2 & \text{if} \quad G = \mathbb{Z}_4 \end{cases}$$

where $\mathbb{R}, \tilde{\mathbb{R}}, \mathbb{C}, \tilde{\mathbb{C}}$ are the representations we introduced in Subsection 3.1.

Definition 3.11. For such triples (X, m, n), (X', m', n'), we say that they are *G*-stably equivalent to each other if $n - n' \in \mathbb{Z}$ and there exist finite dimensional subspaces V, V' of \mathcal{V} and W, W' of \mathcal{W} and a pointed *G*-homotopy equivalence

$$\Sigma^V \Sigma^W X \to \Sigma^{V'} \Sigma^{W'} X',$$

where $\dim_{\mathbb{R}} V - \dim_{\mathbb{R}} V' = m' - m$ and $\dim_{\mathbb{K}_G} W - \dim_{\mathbb{K}_G} W' = n' - n$. Define \mathfrak{C}_G as the set of *G*-stable equivalence classes of triples (X, m, n). An element of \mathfrak{C}_G is called a *spectrum class*.

22

Informally, we may think of the triple (X, m, n) as the formal desuspension of X by V and by W, where $V \subset \mathcal{V}, W \subset \mathcal{W}$ with $\dim_{\mathbb{R}} V = m, \dim_{\mathbb{K}_G} W = n$, so symbolically one may write

$$(X,m,n) = \begin{cases} \Sigma^{-m\mathbb{R}}\Sigma^{-n\mathbb{R}}X & \text{if} \quad G = \mathbb{Z}_2, \\ \Sigma^{-m\mathbb{R}}\Sigma^{-n\mathbb{C}}X & \text{if} \quad G = \mathbb{Z}_4. \end{cases}$$

For $G = \mathbb{Z}_2$, as in [32], we have *canonical* stable equivalences

$$((V \oplus V)^+ \land X, 2 \dim_{\mathbb{R}} V + m, n) \simeq (X, m, n),$$
$$((W \oplus W)^+ \land X, m, 2 \dim_{\mathbb{R}} W + n) \simeq (X, m, n).$$

The reason why we take twice of $\dim_{\mathbb{R}} V$ and $\dim_{\mathbb{R}} W$ and put a copy of V and Wat the first factor is that $GL(N, \mathbb{R})$ for N > 0 is not connected and thus there is no canonical choice of trivialization of a given real representation. When $G = \mathbb{Z}_4$, we actually have a simpler formula for W: there is a canonical stable equivalence

$$(W^+ \wedge X, m, \dim_{\mathbb{R}} W + n) \simeq (X, m, n),$$

since \mathcal{W} is a complex representation of G.

As well as the non-equivariant case, we can define the notion of local equivalence, which was introduced by Stoffregen [51], in our G-equivariant setting:

Definition 3.12. Let (X, m, n), (X', m', n') be triples as above and let $l \in \mathbb{Q}$. A *G-stable map* $(X, m, n) \to (X', m', n')$ of height *l* is a pointed *G*-map

$$\Sigma^V \Sigma^W X \to \Sigma^{V'} \Sigma^{W'} X'$$

for some subspaces $V, V' \subset \mathcal{V}$ and $W, W' \subset \mathcal{W}$ with $\dim_{\mathbb{R}} V - \dim_{\mathbb{R}} V' = m' - m$ and and $\dim_{\mathbb{K}_G} W - \dim_{\mathbb{K}_G} W' = n' - n + l$. A *G*-stable map $(X, m, n) \to (X', m', n')$ is called a *G*-local map if $\dim_{\mathbb{R}} V - \dim_{\mathbb{R}} V' = m' - m$ and it induces a *G*-homotopy equivalence on the *H*-fixed-point sets. We say that (X, m, n) and (X', m', n')are *G*-locally equivalent if there exist *G*-local maps $(X, m, n) \to (X', m', n')$ and $(X', m', n') \to (X, m, n)$.

The G-local equivalence is evidently an equivalence relation, and we call an equivalence class for this relation a G-local equivalence class. The set of G-local equivalence classes is denoted by \mathcal{LE}_G . We write an element of \mathcal{LE}_G by $[(X, m, n)]_{\text{loc}}$. Evidently the G-stable equivalence implies the G-local equivalence, and thus we have a natural surjection $\mathfrak{C}_G \to \mathcal{LE}_G$.

Fix $G = \mathbb{Z}_2$ or \mathbb{Z}_4 . For a triple (X, m, n) above, we define

$$\tilde{H}^*_G(X, m, n) := \tilde{H}^{*+m+k_G \cdot n}_G(X).$$

Then we may assign each element $\mathcal{X} = [(X, m, n)] \in \mathfrak{C}_G$ to the isomorphism class of equivariant singular cohomology,

$$\tilde{H}^*_G(\mathcal{X}) = [\tilde{H}^*_G(X, m, n)],$$

as a graded $\widetilde{H}^*_G(S^0)$ -module. If $G = \mathbb{Z}_2$, we set

$$d(X, m, n) := d(X) - m - n \in \mathbb{Q}.$$

If $G = \mathbb{Z}_4$, we set

$$\overline{d}(X,m,n) := \overline{d}(X) - m - 2n \in \mathbb{Q},$$

$$\underline{d}(X,m,n) := \underline{d}(X) - m - 2n \in \mathbb{Q}.$$

Obviously, the invariants $d, \overline{d}, \underline{d}$ descend to invariants

 $d(\mathcal{X}), \quad \overline{d}(\mathcal{X}), \quad d(\mathcal{X})$

for $\mathcal{X} \in \mathfrak{C}_G$.

The invariants $d(\mathcal{X})$, $\overline{d}(\mathcal{X})$ and $\underline{d}(\mathcal{X})$ satisfy the following inequalities:

Lemma 3.13. First we suppose $(G, H) = (\mathbb{Z}_2, \mathbb{Z}_2)$. Let (X, m, n), (X', m', n') be triples as above. If there is a stable local G-map of height l, then

$$d(X, m, n) + l \le d(X', m', n').$$

Next, we suppose $(G, H) = (\mathbb{Z}_4, \mathbb{Z}_2)$. If there is a stable local G-map of height l, then we have

$$\overline{d}(X,m,n) + 2l \le \overline{d}(X',m',n'), \quad \underline{d}(X,m,n) + 2l \le \underline{d}(X',m',n').$$

Proof. This follows from Lemma 3.7.

As a corollary of Lemma 3.13, we have the following.

Corollary 3.14. The rational numbers $d(\mathcal{X})$ is a \mathbb{Z}_2 -local equivalence invariant and $\overline{d}(\mathcal{X})$ and $d(\mathcal{X})$ are \mathbb{Z}_4 -local equivalence invariants.

Lemma 3.15. Let $(G, H) = (\mathbb{Z}_4, \mathbb{Z}_2)$ and let (X, m, n), (X', m', n') be triples as above. Suppose that there is a G-stable map

$$f: \Sigma^V \Sigma^W X \to \Sigma^{V'} \Sigma^{W'} X'$$

from (X,m,n) to (X',m',n') of height l as in Definition 3.12 such that f^H : $\Sigma^V X^H \to \Sigma^{V'} (X')^H$ is induced from a linear injection whose image is of codimension-1. Then we have

$$\underline{d}(X, m, n) + 2l \le d(X', m', n').$$

Proof. This follows from Lemma 3.9.

3.5. Seiberg–Witten Floer stable homotopy type for non-spin case. In previous paper, for a spin rational homology 3-sphere with odd involution, we defined Seiberg–Witten Floer stable homotopy type. In this section, we extend our theory to a theory of non-spin case.

Let (Y, \mathfrak{t}) be a spin^c rational homology 3-sphere and ι be a smooth orientationpreserving involution on Y. Suppose that ι also preserves the given spin^c structure t on Y and is of odd type. If t comes from a spin structure, we set $(G, H) = (\mathbb{Z}_4, \mathbb{Z}_2)$, and otherwise set $(G, H) = (\mathbb{Z}_2, \mathbb{Z}_2)$.

Fix an ι -invariant metric g on Y. Once we fix $\lambda \ll 0 \ll \mu$, by Manolescu's construction [32], we obtain a G-equivariant Conley index $I^{\mu}_{\lambda}(Y, \mathfrak{t}, g)$ for a finitedimensional approximation of the Seiberg–Witten flow. Recall that, the finitedimensional approximation of the configuration space is decomposed into

$$V^{\mu}_{\lambda}(Y,\mathfrak{t},g)\oplus W^{\mu}_{\lambda}(Y,\mathfrak{t},g).$$

Here each direct summand is isomorphic to some finite dimensional subspaces of \mathcal{V} and \mathcal{W} as *G*-representations. The *H*-invariant part of $I^{\mu}_{\lambda}(Y, \mathfrak{t}, g)$ is given by $V^{\mu}_{\lambda}(Y, \mathfrak{t}, g)$.

As explained in Section 2, the involution I on the configuration space is determined only by ι up to ι -invariant gauge transformation. Henceforth, we pick a choice of I. Let us set

$$I^{\mu}_{\lambda}(Y,\mathfrak{t},\iota,g) = I^{\mu}_{\lambda}(Y,\mathfrak{t},g)^{I}, \quad V^{\mu}_{\lambda}(Y,\mathfrak{t},\iota,g) = V^{\mu}_{\lambda}(Y,\mathfrak{t},g)^{I}, \quad W^{\mu}_{\lambda}(Y,\mathfrak{t},\iota,g) = W^{\mu}_{\lambda}(Y,\mathfrak{t},g)^{I}$$

As in [32], set

(16)
$$n(Y,\mathfrak{t},g) = \operatorname{ind}_{\mathbb{K}_G} D_W + \frac{2}{k_G} \frac{\sigma(W)}{8} \in \frac{1}{8}\mathbb{Z},$$

where W is a compact spin Riemann 4-manifold bounded by (Y, g) and $\operatorname{ind}_{\mathbb{C}} D_W$ is the complex spin Dirac index on W with spectral boundary condition.

Definition 3.16. Given $Y, \mathfrak{t}, \iota, g$ as above, define an element $SWF_G(Y, \mathfrak{t}, \iota) \in \mathfrak{C}_G$ by

$$SWF_G(Y,\mathfrak{t},\iota) := [(I^{\mu}_{\lambda}(Y,\mathfrak{t},\iota,g), \dim_{\mathbb{R}} V^0_{\lambda}, \dim_{\mathbb{K}_G} W^0_{\lambda} + n(Y,\mathfrak{t},g)/2)].$$

We call $SWF_G(Y, \mathfrak{t}, \iota)$ the *G*-equivariant Seiberg-Witten Floer stable homotopy type or Seiberg-Witten Floer *G*-spectrum class for the involution ι .

As in [21], $SWF_G(Y, \mathfrak{t}, \iota)$ is an invariant of (Y, \mathfrak{t}, ι) :

Proposition 3.17. The spectrum class $SWF_G(Y, \mathfrak{t}, \iota) \in \mathfrak{C}_G$ is an invariant of (Y, \mathfrak{t}, ι) , independent of λ, μ , and g.

Definition 3.18. A *G*-stable map $(X, m, n) \to (X', m', n')$ is called a *G*-local map if it induces a *G*-homotopy equivalence on the *H*-fixed-point sets. We say that (X, m, n) and (X', m', n') are *G*-locally equivalent if there exist *G*-local maps $(X, m, n) \to (X', m', n')$ and $(X', m', n') \to (X, m, n)$.

The *G*-local equivalence is evidently an equivalence relation, and we call an equivalence class for this relation a *G*-local equivalence class. The set of *G*-local equivalence classes is denoted by \mathcal{LE}_G . We write an element of \mathcal{LE}_G by $[(X, m, n)]_{\text{loc}}$. Evidently the *G*-stable equivalence implies the *G*-local equivalence, and we have a natural surjection $\mathfrak{C}_G \to \mathcal{LE}_G$.

3.6. Seiberg–Witten Floer cohomology for involutions. As in the last subsection, let (Y, \mathfrak{t}) be a spin rational homology 3-sphere and ι be a smooth orientationpreserving involution ι . Suppose that ι also preserves the given spin structure \mathfrak{t} and is of odd type.

Definition 3.19. For $G = \mathbb{Z}_2$ or \mathbb{Z}_4 , define the *G*-Seiberg-Witten Floer cohomology for the involution ι by

$$SWFH_G(Y, \mathfrak{t}, \iota) := H^*_G(SWF_G(Y, \mathfrak{t}, \iota)),$$

defined as the isomorphism class of an $\tilde{H}_G(S^0)$ -module. We define also the *Frøyshov*type invariants by

$$\begin{split} \delta_R(Y, \mathfrak{t}, \iota) &:= \frac{1}{2} d(SWF_{\mathbb{Z}_2}(Y, \mathfrak{t}, \iota)) \\ \bar{\delta}_R(Y, \mathfrak{t}, \iota) &:= \frac{1}{2} \overline{d}(SWF_{\mathbb{Z}_4}(Y, \mathfrak{t}, \iota)) \text{ and} \\ \underline{\delta}_R(Y, \mathfrak{t}, \iota) &:= \frac{1}{2} \underline{d}(SWF_{\mathbb{Z}_4}(Y, \mathfrak{t}, \iota)). \end{split}$$

Note that

$$\delta_R(Y,\mathfrak{t},\iota), \underline{\delta}_R(Y,\mathfrak{t},\iota), \overline{\delta}_R(Y,\mathfrak{t},\iota) \in \frac{1}{16}\mathbb{Z}$$

since we have that $n(Y, \mathfrak{t}, g) \in \frac{1}{8}\mathbb{Z}$.

Lemma 3.20. The isomorphism class $SWF_G(Y, \mathfrak{t}, \iota)$ and the rational numbers $\delta_R(Y, \mathfrak{t}, \iota), \ \bar{\delta}_R(Y, \mathfrak{t}, \iota)$ and $\underline{\delta}_R(Y, \mathfrak{t}, \iota)$ are invariants of (Y, \mathfrak{t}, ι) .

Proof. This is a direct consequence of Proposition 3.17.

When Y is an integral homology 3-sphere, we often drop the unique spin structure \mathfrak{t} on Y from our notation.

Example 3.21. For a Brieskorn sphere

$$Y = \Sigma(p,q,r) = \{ z_1^p + z_2^q + z_3^r = 0 \},\$$

let $\iota: Y \to Y$ be the involution defined as the complex conjugation. Namely, define $\iota(z_1, z_2, z_3) = (\bar{z}_1, \bar{z}_2, \bar{z}_3)$. The fixed-point set Y^{ι} is of codimension-2, and hence ι is of odd type. In fact, Y^{ι} is known as the Montesinos knot M(p, q, r).

Recall that an odd involution gives rise to a $\mathbb{Z}_4 \times_{\mathbb{Z}_2} \operatorname{Pin}(2)$ -symmetry on the Seiberg–Witten equations ([7, 38]). For several series of Brieskorn spheres, Montague [38, Subsubsection 8.3.1] computed the $\mathbb{Z}_4 \times_{\mathbb{Z}_2} \operatorname{Pin}(2)$ -equivariant Seiberg– Witten Floer spectra and their *I*-invariant parts (which correspond to the $\langle j\mu \rangle$ invariant parts in the notation of [38]). In particular, in our notation, the local equivalence class of $SWF_{\mathbb{Z}_4}(Y,\iota)$ for $Y = \Sigma(2,3,6n \pm 1)$ is given by

$$\begin{split} [SWF_{\mathbb{Z}_4}(\Sigma(2,3,12k-1),\iota)]_{\rm loc} &= [(\tilde{G},0,0)]_{\rm loc},\\ [SWF_{\mathbb{Z}_4}(\Sigma(2,3,12k-5),\iota)]_{\rm loc} &= [(\tilde{G},0,1/4)]_{\rm loc},\\ [SWF_{\mathbb{Z}_4}(\Sigma(2,3,12k+1),\iota)]_{\rm loc} &= [(S^0,0,0)]_{\rm loc},\\ [SWF_{\mathbb{Z}_4}(\Sigma(2,3,12k+5),\iota)]_{\rm loc} &= [(S^0,0,-1/4)]_{\rm loc}. \end{split}$$

for $k \geq 1$. Using this combined with Example 3.6, we can immediately calculate δ , $\underline{\delta}_R$, $\overline{\delta}_R$ for $(\Sigma(2,3,6n\pm 1),\iota)$, which is summarized in the following table:

Brieskorn sphere equipped with ι	$\overline{\delta}_R$	δ_R	δ_R
$(\Sigma(2,3,12k-1),\iota)$	1/2	1/2	0
$(\Sigma(2,3,12k-5),\iota)$	0	0	-1/2
$(\Sigma(2,3,12k+1),\iota)$	0	0	0
$(\Sigma(2,3,12k+5),\iota)$	1/2	1/2	1/2

3.7. Cobordisms. Let (Y_0, \mathfrak{t}_0) and (Y_1, \mathfrak{t}_1) be spin^c closed 3-manifolds with $b_1(Y_i) = 0$. We do not assume that Y_0 and Y_1 are connected. Suppose that we have an involution ι_i on each of Y_i is an odd type. Let (W, \mathfrak{s}) be a smooth spin 4-dimensional oriented cobordism with $b_1(W) = 0$. We assume that there is an odd involution ι on W such that $\iota|_{Y_i} = \iota_i$ for i = 0, 1. Let \mathbb{S}^{\pm} be positive and negative spinor bundles on W and let \mathbb{S}_i be the spinor bundles on Y_i i = 0, 1.

Here we consider the Sobolev norms L_k^2 for the spaces $\Omega^*(W)$ and $\Gamma(\mathbb{S}^{\pm})$ obtained from ι -invariant metrics and ι -invariant connections for a fixed integer $k \geq 3$. The relative Bauer–Furuta invariant of W introduced by Manolescu [32] gives a map between the Seiberg–Witten Floer stable homotopy types of Y_0 and Y_1 . This is obtained from the Seiberg–Witten map on W, which is given as a finite-dimensional approximation of a map

(17)
$$SW: \Omega^{1}_{CC}(W) \times \Gamma(\mathbb{S}^{+}) \to \Omega^{+}(W) \times \Gamma(\mathbb{S}^{-}) \times \hat{V}(-Y_{0})^{\mu}_{-\infty} \times \hat{V}(Y_{1})^{\mu}_{-\infty}$$

for large μ . Here

$$\Omega^1_{CC}(W) = \left\{ \left. a \in \Omega^1(W) \right| \, d^*a = 0, d^*\mathbf{t}_i a = 0, \int_{Y_i} \mathbf{t}_i * a = 0 \right\}$$

is the space of 1-forms satisfying the double Coulomb condition. This is introduced by Khandhawit in [19]. Here for a general rational homology sphere Y equipped with the spin^c structure, $\hat{V}(Y, \mathfrak{t})_{-\infty}^{\mu}$ is a subspace of $\hat{V}(Y, \mathfrak{t}) = \text{Ker } d^* \times \Gamma(\mathbb{S})$ which is defined as the direct sum of eigenspaces whose eigenvalues are less than μ . The $\Omega^+(W) \times \Gamma(\mathbb{S}^-)$ -factor of the map SW is given as the Seiberg–Witten equations, and the $\hat{V}(-Y_0)_{-\infty}^{\mu} \times \hat{V}(Y_1)_{-\infty}^{\mu}$ -factor is given, roughly, as the restriction of 4dimensional configurations to 3-dimensional ones. Taking the *I*-invariant part of (17), we obtain a *G*-equivariant map and a finite-dimensional approximation of this gives us a *G*-equivariant map of the form

(18)
$$f: \Sigma^{m_0 \mathbb{\bar{R}}} \Sigma^{n_0 \mathbb{K}_G} I_{-\mu}^{-\lambda}(Y_0) \to \Sigma^{m_1 \mathbb{\bar{R}}} \Sigma^{n_1 \mathbb{K}_G} I_{\lambda}^{\mu}(Y_1),$$

where $I_{\lambda}^{\mu}(Y_i) = I_{\lambda}^{\mu}(Y_i, \mathfrak{t}_i, \iota_i, g_i)$, and $m_i, n_i^{\pm} \geq 0$ and $-\lambda, \mu$ are sufficiently large. The representation spaces $\mathbb{\tilde{R}}$ and \mathbb{K}_G are the representations of G which we introduced in Subsection 3.4.

Let us denote by $V_i(\tilde{\mathbb{R}})^{\mu}_{\lambda}$ the vector space $V(\tilde{\mathbb{R}})^{\mu}_{\lambda}$, a finite-dimensional approximation of the *I*-invariant part, for Y_i , and let us use similar notations also for other representations.

Lemma 3.22. We have

(19)
$$m_{0} - m_{1} = \dim_{\mathbb{R}}(V_{1}(\mathbb{R})_{\lambda}^{0}) - \dim_{\mathbb{R}}(V_{0}(\mathbb{R})_{-\mu}^{0}) - b^{+}(W) + b_{\iota}^{+}(W),$$

$$n_{0} - n_{1} = \dim_{\mathbb{K}_{G}}(V_{1}(\mathbb{K}_{G})_{\lambda}^{0}) - \dim_{\mathbb{K}_{G}}(V_{0}(\mathbb{K}_{G})_{-\mu}^{0})$$

(20)
$$-\frac{1}{k_{G}}\left(\frac{c_{1}(\mathfrak{s})^{2} - \sigma(W)}{8} + n(Y_{1},\mathfrak{t}_{1},g_{1}) - n(Y_{0},\mathfrak{t}_{0},g_{0})\right).$$

Proof. From [19, Theorem 1.], we can calculate the index of the linearization of the map (17). Taking the $(-\iota^*, I)$ -invariant part of the finite dimensional approximation of f, we have that $m_0 - m_1$ and $n_0 - n_1$ coincides with the index of $-\iota^*$ and I fixed part of the form part and spinor part respectively. Since I is an anti-linear involution, $n_0 - n_1$ is half of the Dirac index.

3.8. **Proof of Frøyshov type inequalities.** Let Y be an oriented closed rational homology 3-sphere. Let (\mathfrak{t}, ι) be a real spin^c structure on Y, i.e. \mathfrak{t} is a spin^c structure on Y and $\iota : Y \to Y$ is an orientation-preserving smooth involution on Y such that $\iota^*\mathfrak{t} \cong \overline{\mathfrak{t}}$. As in dimension 4, we shall define the notion that ι is of *odd type*, which is satisfied if the fixed-point set Y^{ι} is non-empty and of codimension-2. For this triple (Y, \mathfrak{t}, ι) , we have the Frøyshov-type invariant

$$\delta_R(Y,\mathfrak{t},\iota) \in \frac{1}{16}\mathbb{Z}.$$

First, we state and prove the main inequality of this paper in the most general form, which relates $\delta_R(Y, \mathfrak{t}, \iota)$ with topological quantities:

Theorem 3.23. Let $(Y_0, \mathfrak{t}_0), (Y_1, \mathfrak{t}_1)$ be spin^c rational homology 3-spheres. Let ι_0, ι_1 be smooth involutions on Y_0, Y_1 . Suppose that ι_0, ι_1 preserve the given orientations and spin^c structures $\mathfrak{t}_0, \mathfrak{t}_1$ on Y_0, Y_1 respectively, and suppose that ι_0, ι_1 are of odd type. Let (W, \mathfrak{s}) be a smooth compact oriented spin^c cobordism with $b_1(W) = 0$ from (Y_0, \mathfrak{t}_0) to (Y_1, \mathfrak{t}_1) . Suppose that there exists a smooth involution ι on W such that ι preserves the given orientation and spin^c structure \mathfrak{s} on $W, b^+(W) - b_{\iota}^+(W) = 0$ and that the restriction of ι to the boundary is given by ι_0, ι_1 . Then we have

(21)
$$\delta_R(Y_0, \mathfrak{t}_0, \iota_0) + \frac{c_1(\mathfrak{s})^2 - \sigma(W)}{16} \le \delta_R(Y_1, \mathfrak{t}_1, \iota_1).$$

Proof. The relative Bauer–Furuta invariant for (W, \mathfrak{s}, ι) gives a map

$$f: \Sigma^V \Sigma^W SWF(Y_0, \mathfrak{t}_0, \iota_0) \to \Sigma^{V'} \Sigma^{W'} SWF(Y_1, \mathfrak{t}_1, \iota_1)$$

and we have $\dim_{\mathbb{R}}(V) - \dim_{\mathbb{R}}(V') = b^+(W) - b^+_{\iota}(W)$ and $\dim_{\mathbb{R}}(W) - \dim_{\mathbb{R}}(W') = (c_1(\mathfrak{s})^2 - \sigma(W))/8$. The claim of theorem follows from Lemma 3.7 applied to this f.

If we focus on the case that \mathfrak{t} comes from a spin structure, for which one has $\mathfrak{t} \cong \overline{\mathfrak{t}}$, we may define invariants analogous to invariants introduced by Stoffregen [50] $\overline{\delta}_R(Y,\mathfrak{t})$, $\underline{\delta}_R(Y,\mathfrak{t})$, corresponding to $\overline{d}(Y,\mathfrak{t})$, $\underline{d}(Y,\mathfrak{t})$ in involutive Heegaard Floer homology [16]. For a spin rational homology 3-sphere (Y,\mathfrak{t}) and an orientaion-preserving smooth involution $\iota: Y \to Y$ such that $\iota^*\mathfrak{t} \cong \mathfrak{t}$, when ι is of odd type, we have two invariants

$$\bar{\delta}_R(Y,\mathfrak{t},\iota), \ \underline{\delta}_R(Y,\mathfrak{t},\iota) \in \frac{1}{16}\mathbb{Z},$$

which satisfy inequalities

$$\underline{\delta}_R(Y,\mathfrak{t},\iota) \leq \delta_R(Y,\mathfrak{t},\iota) \leq \bar{\delta}_R(Y,\mathfrak{t},\iota)$$

On these invariants, we have the following:

Theorem 3.24. Let (Y_0, \mathfrak{t}_0) , (Y_1, \mathfrak{t}_1) be spin rational homology 3-spheres. Let ι_0, ι_1 be smooth involutions on Y_0, Y_1 . Suppose that ι_0, ι_1 preserve the given orientations and spin structures $\mathfrak{t}_0, \mathfrak{t}_1$ on Y_0, Y_1 respectively, and suppose that ι_0, ι_1 are of odd type. Let (W, \mathfrak{s}) be a smooth compact oriented spin cobordism with $b_1(W) = 0$ from (Y_0, \mathfrak{t}_0) to (Y_1, \mathfrak{t}_1) . Suppose that there exists a smooth involution ι on W such that ι preserves the given orientation and spin structure \mathfrak{s} on W, $b^+(W) - b^+_{\iota}(W) = 0$ and that the restriction of ι to the boundary is given by ι_0, ι_1 . Then we have

(22)
$$\bar{\delta}_R(Y_0, \mathfrak{t}_0, \iota_0) - \frac{\sigma(W)}{16} \le \bar{\delta}_R(Y_1, \mathfrak{t}_1, \iota_1),$$

(23)
$$\underline{\delta}_R(Y_0, \mathfrak{t}_0, \iota_0) - \frac{\sigma(W)}{16} \le \underline{\delta}_R(Y_1, \mathfrak{t}_1, \iota_1).$$

Moreover, if $b^+(W) - b^+_{\iota}(W) = 1$, then we have

(24)
$$\underline{\delta}_R(Y_0, \mathfrak{t}_0, \iota_0) - \frac{\sigma(W)}{16} \le \overline{\delta}_R(Y_1, \mathfrak{t}_1, \iota_1).$$

Proof. When \mathfrak{s} is spin and $b^+(W) - b^+_\iota(W) = 0$, the Bauer–Furuta invariant for (W, \mathfrak{s}, ι) ,

$$f: \Sigma^{V} \Sigma^{W} SWF(Y_{0}, \mathfrak{t}_{0}, \iota_{0}) \to \Sigma^{V'} \Sigma^{W'} SWF(Y_{1}, \mathfrak{t}_{1}, \iota_{1}),$$

is a \mathbb{Z}_4 -equivariant local map of height $-\sigma(W)/8$ since $\dim_{\mathbb{R}}(V) - \dim_{\mathbb{R}}(V') = b^+(W) - b^+_{\iota}(W)$ and $\dim_{\mathbb{R}}(W) - \dim_{\mathbb{R}}(W') = -\sigma(W)/8$. So, the first two inequalities follow from Lemma 3.13 applied to this f. The third inequality follows from Lemma 3.9 applied to \mathbb{Z}_4 -equivariant relative Bauer–Furuta invariants of (W, \mathfrak{s}) . \Box

The inequality (24) can be seen as a real and 4-manifold with boundary version of Donaldson's Theorem B [9].

3.9. Connected sum formula. In this section, we prove the following connected sum formula of $[SWF(Y, \mathfrak{t}, \iota)]_{\text{loc}}$ and δ . We first write a statement of connected sum for $[SWF(Y, \mathfrak{t}, \iota)]_{\text{loc}}$.

First, we recall the operation of equivariant connected sum. For more details, see [21, Definition 3.43]. Let $(Y_0, \mathfrak{t}_0, y_0)$ and $(Y_1, \mathfrak{t}_1, y_1)$ be spin^c rational homology 3-spheres with base points. Let ι_0 and ι_1 are involutions on Y_0 and Y_1 respectively. Suppose that the fixed-point set of ι_i are codimension-2 and $\iota_i(y_i) = y_i$, and that ι_i preserves the spin structure \mathfrak{t}_i for i = 0, 1. We give an orientation $o(\iota_i)$ of the set of fixed points of the involution ι_i . For these data, the equivariant connected sum

$$(Y_0, \mathfrak{t}_0, y_0, \iota_0, o(\iota_0)) \# (Y_1, \mathfrak{t}_1, y_1, \iota_1, o(\iota_1))$$

is defined. We sometimes drop y_0, y_1 and $o(\iota_1), o(\iota_2)$ from our notation.

The standard 3-handle cobordism between the connected sum and disjoint union of two manifolds can be considered equivariantly. Namely, we have an oriented homology cobordism

 $W_{01}: (Y_0, \mathfrak{t}_0, \iota_0, o(\iota_0)) \# (Y_1, \mathfrak{t}_1, \iota_1, o(\iota_1)) \to (Y_0, \mathfrak{t}_0, \iota_0, o(\iota_0)) \sqcup (Y_1, \mathfrak{t}_1, \iota_1, o(\iota_1))$

that is equipped with an involution ι whose restriction to the boundary is given by ι_0, ι_1 .

Lemma 3.25. The cobordism W_{01} admits a spin^c structure \mathfrak{s} whose restriction to the boundary is given by $\mathfrak{t}_0, \mathfrak{t}_1$ such that $\iota^* \mathfrak{s} \cong \overline{\mathfrak{s}}$.

Proof. From homological computations, one can verify $H^2(W_{01}, \partial W_{01}; \mathbb{Z}) = 0$. Thus, we know an extension of spin^c structure of $\mathfrak{t}_0 \cup \mathfrak{t}_1 \cup \mathfrak{t}_0 \# \mathfrak{t}_1$ exists and is unique up to isomorphism. The connected sum theorem can be formulated as follows.

Theorem 3.26. Let $(Y_0, \mathfrak{t}_0, \iota_0)$ and $(Y_1, \mathfrak{t}_1, \iota_1)$ are spin^c \mathbb{Z}_2 -homology 3-spheres with odd involutions ι_0 and ι_1 . Suppose that the fixed-point set of ι_i is non-empty and connected for each i = 0, 1. Then, we have

$$[SWF_G(Y_0,\mathfrak{t}_0,\iota_0)\wedge SWF_G(Y_1,\mathfrak{t}_1,\iota_1)]_{\text{loc}}=[SWF_G(Y^{\#},\mathfrak{t}^{\#},\iota^{\#})]_{\text{loc}},$$

where $(Y^{\#}, \mathfrak{t}^{\#}, \iota^{\#}) := (Y_0, \mathfrak{t}_0, \iota_0) \# (Y_1, \mathfrak{t}_1, \iota_1).$

Proof. This follows from the same argument in the proof of [21, Theorem 3.48] although the doubled Floer homotopy types are used in [21]. The local maps of both directions are given as relative Bauer–Furuta invariants for spin^c cobordism $\pm W_{01}$ with involution.

Theorem 3.27. Let $(Y_0, \mathfrak{t}_0, \iota_0)$ and $(Y_1, \mathfrak{t}_1, \iota_1)$ are spin^c \mathbb{Z}_2 -homology 3-spheres with odd involutions ι_0 and ι_1 .

$$\delta_R(Y,\iota,\mathfrak{t}) + \delta_R(Y',\iota',\mathfrak{t}') = \delta_R(Y\#Y',\iota\#\iota',\mathfrak{t}\#\mathfrak{t}').$$

In order to prove Theorem 3.27, we first prove the following duality result:

Lemma 3.28. Let (Y, \mathfrak{t}, ι) be an oriented spin rational homology sphere with odd involution. Then we have

$$\delta_R(Y, \mathfrak{t}, \iota) = -\delta_R(Y, \mathfrak{t}, \iota).$$

Moreover, suppose \mathfrak{t} is spin, we have

$$\underline{\delta}_R(Y,\mathfrak{t},\iota) = -\overline{\delta}_R(-Y,\mathfrak{t},\iota).$$

Proof of Lemma 3.28. We first claim that $I^{\lambda}_{-\lambda}(Y,\iota,g)$ and $I^{\lambda}_{-\lambda}(-Y,\iota,g)$ are equivariantly $V^{\lambda}_{-\lambda}(Y,\iota,g) \oplus W^{\lambda}_{-\lambda}(Y,\iota,g)$ -dual. This can be obtained as the restriction of the V-dual map [38, Proposition 6.23]. Then the inequalities of the conclusion directly follow from Lemma 3.10.

Proof of Theorem 3.27. From Lemma 3.8 and Theorem 3.26, the following inequality has already been proven:

$$\delta_R(Y,\iota,\mathfrak{t}) + \delta_R(Y',\iota',\mathfrak{t}') \le \delta_R(Y\#Y',\iota\#\iota',\mathfrak{t}\#\mathfrak{t}').$$

The remaining part is to see the opposite inequality. To see this, using the above inequality to orientation reversals, we obtain

$$\delta_R(-Y,\iota,\mathfrak{t}) + \delta_R(-Y',\iota',\mathfrak{t}') \leq \delta_R(-(Y\#Y'),\iota\#\iota',\mathfrak{t}\#\mathfrak{t}').$$

Then, we use Lemma 3.28 and obtain

$$\delta_R(Y,\iota,\mathfrak{t}) + \delta_R(Y',\iota',\mathfrak{t}') \ge \delta_R(Y\#Y',\iota\#\iota',\mathfrak{t}\#\mathfrak{t}').$$

This completes the proof.

3.10. **SWF-spherical.** We introduce the easiest case to calculate our invariants d, \underline{d} and \overline{d} .

Definition 3.29. We say that a triple (Y, t, ι) of a rational homology 3-sphere, spin^c structure, and an odd involution is *SWF-spherical* if

$$[SWF_G(Y,\mathfrak{t},\iota)]_{\mathrm{loc}} = [(S^0,m,n)]_{\mathrm{loc}}$$

for some $m \in \mathbb{Z}$ and $n \in \mathbb{Q}$.

The following computations are obvious from the definitions of invariants δ_R , $\underline{\delta}_R$ and $\overline{\delta}_R$.

Proposition 3.30. If a triple (Y, t, ι) is SWF-spherical, then

$$\delta_R(Y,\iota,\mathfrak{t}) = \underline{\delta}_R(Y,\iota,\mathfrak{t}) = \overline{\delta}_R(Y,\iota,\mathfrak{t}) = -\frac{m}{2} - \frac{n}{2}$$

for m, n satisfying $[SWF_G(Y, \mathfrak{t}, \iota)]_{loc} = [(S^0, m, n)].$

Now, we prove Proposition 1.10 using Proposition 3.30.

Proof of Proposition 1.10. From [21], for lens spaces, we have the following computation: Let p, q be coprime natural numbers. Regard the lens space $Y = \Sigma(K(p,q)) = L(p,q)$, and equip Y with the standard metric g, which has positive scalar curvature. The complex conjugation on Y defines an involution ι on Y that preserves g. The fixed-point set of ι is non-empty and of codimension-2, which is called two bridge knot/link. Through the same discussion given in [21, Example 3.56], we have

$$SWF_G(L(p,q), t, \iota) = [(S^0, 0, n(L(p,q), \mathfrak{t}, g)/2].$$

This combined with Proposition 3.30 completes the computations for δ , $\bar{\delta}_R$ and $\underline{\delta}_R$. For torus knots, the proof is similar and we use the argument of the proof of [21, Theorem 3.58] instead of [21, Example 3.56].

4. Floer homotopy type of links and Frøyshov type invariants

In [21], we defined a Floer homotopy type of a knot

$$K \mapsto DSWF(K)$$

which we called the doubled Floer homotopy type for a knot K. For some reason related to K-theory, we made the doubling operation. In this paper, we give a Floer homotopy type SWF(K) without taking double. Moreover, we generalize the invariant SWF(K) to that of oriented links L in S^3 with non-zero determinant:

$$L \mapsto SWF(L)$$
.

Using them, for links with non-zero determinant, we introduce Frøyshov type invariants

$$L \mapsto \delta_R(L), \underline{\delta}_R(L), \overline{\delta}_R(L) \in \frac{1}{16}\mathbb{Z}.$$

4.1. Floer homotopy type of links. Let L be a link in S^3 with non-zero determiant. Then, we can associate the double branched cover

$$\Sigma(L) \to S^3$$

uniquely and a covering involution $\iota : \Sigma(L) \to \Sigma(L)$. Since our theory uses real spin^c structures, we need to confirm compatibility between the covering involutions and spin(^c) structures on the double branched covering spaces. The following correspondence between orientations of L and spin structures on $\Sigma(L)$ is proven by Turaev [53]:

Lemma 4.1. There is a canonical one-to-one correspondence between the set of isomorphism classes of spin structures on $\Sigma(L)$ and {orientations on L}/{±1}.

We need to clarify the correspondence in Lemma 4.1 to see if the spin structures are preserved by the involution.

For a given link L, we take an open tubular neighborhood N_L of L in $\Sigma(L)$, where \tilde{L} is the lift of L in $\Sigma(L)$. Let \mathfrak{s} be the unique spin structure on $\Sigma(L) \setminus N_L$ coming from the pull-back of the spin structure on $S^3 \setminus L$, which is the restriction of the spin structure on S^3 . We pick an orientation o of L. Corresponding to o, one can associate an element h(0) of $H^1(\Sigma(L) \setminus N_L; \mathbb{Z}_2)$ as follows:

$$h(o)(l) := \frac{1}{2} \operatorname{lk}(p(l), L) : H_1(\Sigma(L) \setminus N_L; \mathbb{Z}_2) \to \mathbb{Z}_2,$$

where p is the covering projection of $\Sigma(L)$ and lk is the linking number with respect to the orientation o. Thus, we have a spin structure $\mathfrak{s} + h(o)$ on $\Sigma(L) \setminus N_L$, which has the unique extension to the spin structure on $\Sigma(L)$ denote by $\mathfrak{s}(o)$. Then, the map

(25) {orientations on
$$L\}/\{\pm 1\} \to \operatorname{Spin}(\Sigma(L)); o \mapsto \mathfrak{s}(o)$$

gives a bijection, where $\text{Spin}(\Sigma(L))$ denotes the set of ismorphism classes of spin structures on $\Sigma(L)$. From the constructions of the map (25), we can ensure that the spin structure $\mathfrak{s} + h(o)$ is preserved by the covering involution ι . Thus we have:

Lemma 4.2. Any spin structure on $\Sigma(L)$ is preserved by the involution ι .

For a given orientation of L, we denote by \mathfrak{t}_L the spin structure on $\Sigma(L)$ corresponding to the orientation. Note that $\mathfrak{t}_L \cong \mathfrak{t}_{-L}$.

Definition 4.3. For a given oriented link L with non-zero determinant, we define

$$SWF(L) := SWF(\Sigma(L), \iota, \mathfrak{t}_L).$$

Obviously, the \mathbb{Z}_4 -equivariant stable homotopy type SWF(L) is invariant under isotopy for links.

4.2. Frøyshov type invariants for links. Now, we define the three invariants: **Definition 4.4.** For a given oriented link *L*, we define

$$\delta_R(L) := \delta_R(\Sigma(L), \iota, \mathfrak{t}_L), \quad \bar{\delta}_R(L) := \bar{\delta}_R(\Sigma(L), \iota, \mathfrak{t}_L), \quad \underline{\delta}_R(L) := \underline{\delta}_R(\Sigma(L), \iota, \mathfrak{t}_L),$$

where \mathfrak{t}_L is the spin structure corresponding to the given orientation of L. For the correspondence between orientations of L and spin structures on $\Sigma(L)$, see (25). Similarly, we define

$$\kappa(L) := \kappa(\Sigma(L), \iota, \mathfrak{t}_L).$$

Here $\kappa(-)$ is the invariant defined in [21, Theorem 1.1].

Before proving Theorem 1.3, we need to calculate several fundamental quantities of double branched covering spaces.

Let (X, S) be a connected (possibly non-orientable) link cobordism from (S^3, L) to (S^3, L') with $H_1(X; \mathbb{Z}_2) = 0$. Suppose the determinants L and L' are nonzero. As in the closed case given in [39, Corollary 2.10], we see that the surface S in X determines a unique double branched covering space if and only if $0 \equiv$ $[S] \in H_2(X, \partial X; \mathbb{Z}_2)$. The followings are fundamental computations regarding the homologies of double-branched covering spaces.

Lemma 4.5. The following equalities hold:

(26)
$$\sigma(\Sigma(S)) = 2\sigma(X) - \frac{1}{2}S \circ S - \sigma(L) + \sigma(L'),$$

(27) $b_2(\Sigma(S)) = 2b_2(X) + b_1(S),$

(28)
$$b^+(\Sigma(S)) = 2b^+(X) + \frac{1}{2}b_1(S) - \frac{1}{4}S \circ S - \frac{1}{2}\sigma(L) + \frac{1}{2}\sigma(L'),$$

(29) $b_1(\Sigma(S)) = b_3(\Sigma(S)) = 0.$

Moreover, if L and L' are knots, there is no 2-torsion in the cohomology $H^2(\Sigma(S);\mathbb{Z})$.

Proof. The equality (29) follows from the exact sequence given in [24]. The Mayer– Vietoris argument given in [21, Lemma 4.2] enables us to confirm (26) by the following discussion. It is confirmed in [18, Lemma 1.1] that for a Seifert surface S_L in S^3 bounded by L giving $\sigma(L)$, the double branched covering N_L for its push into D^4 satisfies

(30)
$$\sigma(L) = \sigma(N_L).$$

From G-signature theorem for closed 4-manifolds and (30), we obtain (26). Next, we see (29). Since $\Sigma(S) \to X$ is a degree 2-branched cover, we have $\chi(\Sigma(S)) = 2\chi(X) - \chi(S)$. From (29), we have $b_2(\Sigma(S)) = 2b_2(X) + b_1(S)$. We note (28) follows from (27) and (26). We finally show there is no 2-torsion in $H^*(\Sigma(S);\mathbb{Z})$ when L and L' are knots. From [24], the sequence

$$H_2(X;\mathbb{Z}_2) \to H_1(X,S;\mathbb{Z}_2) \to H_1(\Sigma(S);\mathbb{Z}_2) \to 0.$$

is exact. The pair exact sequence for (X, S) combined with connectivity of S implies $H_1(X, S; \mathbb{Z}_2) = 0$. Thus we have $H_1(\Sigma(S); \mathbb{Z}_2) = 0$, and there is no 2-torsion in $H_1(\Sigma(S); \mathbb{Z}) = 0$. This implies there is no 2-torsion on $H^2(\Sigma(S))$. This completes the proof.

We now prove Theorem 1.3(iii).

Proof of Theorem 1.3(iii). This follows from Theorem 3.23 combined with the computation of homological invariants in Lemma 4.5. \Box

We also have similar inequalities for $\bar{\delta}_R$ and $\underline{\delta}_R$. Note that Theorem 1.6 follows from (33).

Theorem 4.6. Let L and L' be links in S^3 with non-zero determinants, X be an oriented smooth compact connected cobordism from S^3 to S^3 , and S be a compact connected properly and smoothly (possibly non-orientable) embedded cobordism in X from L to L' such that the homology class $[S] \equiv 0 \mod 2$. Suppose $H_1(X; \mathbb{Z}) = 0$, $PD[S] \equiv w_2(X) \in H^2(X; \mathbb{Z}_2)$,

$$b^{+}(X) + g(S) - \frac{1}{4}S \circ S - \frac{1}{2}\sigma(L) + \frac{1}{2}\sigma(L') = 0$$

and there is a spin structure \mathfrak{s} on $\Sigma(S)$ such that $\iota^*\mathfrak{s} \cong \mathfrak{s}$ and whose restrictions are compatible with orientations of L and L'.

Then, we have

(31)
$$\bar{\delta}_R(L) + \frac{1}{16} \left(-2\sigma(X) + \frac{1}{2}S \circ S + \sigma(L) - \sigma(L') \right) \leq \bar{\delta}_R(L')$$

(32)
$$\underline{\delta}_R(L) + \frac{1}{16} \left(-2\sigma(X) + \frac{1}{2}S \circ S + \sigma(L) - \sigma(L') \right) \leq \underline{\delta}_R(L')$$

where $S \circ S$ means the self intersection number of S in X.

Moreover, when

$$b^+(X) + g(S) - \frac{1}{4}S \circ S - \frac{1}{2}\sigma(L) + \frac{1}{2}\sigma(L') = 1,$$

we have

(33)
$$\underline{\delta}_R(L) + \frac{1}{16} \left(-2\sigma(X) + \frac{1}{2}S \circ S + \sigma(L) - \sigma(L') \right) \leq \bar{\delta}_R(L')$$

Proof of Theorem 4.12. These inequalities follow from Theorem 3.23 and Theorem 3.24 combined with Lemma 4.5. $\hfill \Box$

As we discussed in the introduction, our invariant will be χ -concordance invariant. Thus, we introduce the notion of χ -concordant. First, we introduce a notion of χ -slice:

Definition 4.7. A link L in S^3 is χ -slice if L bounds a smoothly properly embedded surface F in D^4 without closed components, and with $\chi(F) = 1$. If L is oriented we require F to be compatibly oriented for orientable components of F.

Note that χ -sliceness gives a generalization of sliceness of knots. In order to take connected sum, we also need a base point. A *marked link* is a link in S^3 equipped with a marked component.

Definition 4.8. For given oriented marked links L_0 and L_1 , we say L_0 and L_1 are χ -concordant if $-L_0^* \# L_1$ bounds a smoothly properly embedded surface F in D^4 such that

- (i) F is a disjoint union of one disk together with annuli and Mobius bands;
- (ii) the boundary of the disk component of F is the marked component of $-L_0^* \# L_1$;
- (iii) we require orientable components of F to be oriented compatible with the orientation of $-L_0^* \# L_1$.

Here the conncted sum $-L_0^* \# L_1$ is taken along marked points.

If L_0 and L_1 are χ -concordant, then $-L_0^* \# L_1$ is χ -slice. On the other hand, the converse is not true. In [8], it is proven that the set \mathcal{L} of all χ -concordant classes of oriented mariked links forms an abelian group with respect to the connected sum along marked components. The group $\tilde{\mathcal{L}}$ is called the *link concordance group*. In this paper, we focus on the subgroup $\tilde{\mathcal{F}}$ of $\tilde{\mathcal{L}}$ generated by oriented mariked links whose determinants are non-zero.

We first provide a homotopy theoretic homomorphism:

Proposition 4.9. The map $[L] \mapsto [SWF(L)]_{loc}$ gives a homomorphism from the link concordance group $\widetilde{\mathcal{F}}$ to \mathcal{LE}_G , where $G = \mathbb{Z}_4$.

Proof. As \mathbb{Z}_2 -equivariant spin manifolds, we have

 $(\Sigma(L\#L'),\iota_{L\#L'},\mathfrak{t}_{L\#L'})\cong(\Sigma(L),\iota_L,\mathfrak{t}_L)\#(\Sigma(L'),\iota_{L'},\mathfrak{t}_{L'}).$

where ι_L denotes the covering involution of double branched covering space. Also, taking local equivalence class $[-]_{loc}$ is a homomorphism from Theorem 3.26. This completes the proof.

We next prove a relation between our invariants and χ -sliceness.

Lemma 4.10. If L is χ -slice, then $\underline{\delta}_R(L)$, $\overline{\delta}_R(L)$ and $\kappa(L)$ are zero. Moreover, the quantity $\delta_R(L)$ is a χ -concordace invariant.

Proof. Suppose L is χ -slice, i.e. $-L_0^* \# L_1$ bounds a smoothly properly embedded surface S in D^4 without closed components, and with $\chi(S) = 1$. Then [8, Proposition 2.6] implies the double branched cover gives a rational spin homology D^4 bounded by $\Sigma(L)$ with a spin structure corresponding to the orientation of L. Then Theorem 4.6 and [21, Theorem 1.1 (iv)] implies

$$0 \leq \delta_R(L), 0 \leq \underline{\delta}_R(L), 0 \leq \delta_R(L), 0 \leq \kappa(L).$$

Also, if L is χ -slice then, $-L^*$ is also χ -slice. It implies that $\underline{\delta}_R(L)$, $\overline{\delta}_R(L)$ and $\kappa(L)$ are zero.

Suppose L and L' are χ -concordant. Then $-L_0^* \# L_1$ is χ -slice. This implies $\delta_R(-L_0^* \# L_1) = 0$. On the other hand, since δ is homomorphism, we see $\delta_R(L_0) =$ $\delta_R(L_1).$ \square

As a corollary of Theorem 3.27, we have;

Corollary 4.11. The invariant $\delta_R(L)$ gives a homomorphism from $\tilde{\mathcal{F}}$ to $\frac{1}{16}\mathbb{Z}$.

4.3. κ invariants for links. We gave a generalization of the kappa invariant for knots defined in [21] to that of links with non-zero determinant in Definition 4.4:

$$L \mapsto \kappa(L) \in \frac{1}{16}\mathbb{Z}$$

We also note general properties of the invariant for links. In particular, we provide 10/8-inequality for surfaces bounded by links as (iv) in the the following theorem.

Theorem 4.12. The invariant κ satisfies the following properties:

- (i) $2\kappa(L) \equiv -\frac{\sigma(L)}{8} \mod 2$, (ii) $\kappa(L) + \kappa(-L^*) \ge 0$,

(iii) Let L and L' be oriented links in S³, X be an oriented smooth compact connected cobordism from S³ to S³ with H₁(X; Z) = 0, and S be an oriented compact connected properly and smoothly embedded cobordism in X from L to L' such that the homology class [S] of S is divisible by 2 and PD(w₂(X)) = [S]/2 mod 2. Then, we have

(34)
$$-\frac{\sigma(X)}{8} + \frac{9}{32}S \circ S - \frac{9}{16}\sigma(L') + \frac{9}{16}\sigma(L) \le b^+(X) + g(S) + \kappa(L') - \kappa(L).$$

Proof. For a proof of (i), we take a properly and smoothly embedded and connected oriented surface S in D^4 bounded by L. For such a surface, the signature of L is given by $\sigma(\Sigma(S))$ from Lemma 4.5. Thus, one can describe the Rochlin invariant of $\Sigma(L)$ by $\frac{1}{8}\sigma(\Sigma(S))$. Then Theorem 1.1 (i) in [21] implies

$$-2\kappa(L) = \mu(\Sigma(L), \mathfrak{t}_L) = \frac{1}{8}\sigma(\Sigma(S)).$$

This completes the proof of (i). The inequality (ii) followed from [21, Theorem 1.1 (iii)]. The inequality (iii) follows by combining Lemma 4.5 with [21, Theorem 1.1(iv)].

5. Applications

In this section, we prove applications in the introduction such as non-smoothable and non-orientable surfaces in 4-manifolds Theorem 1.23, obstruction to the Nielsen realization problem Theorem 1.16 and non-orientable surfaces in D^4 bounded by torus knots Theorem 1.14.

5.1. Applications to non-smoothable actions. We start with proving an application to non-smoothable actions, Theorem 1.24.

Proof of Theorem 1.24. First, we shall construct a locally linear involution $\iota: W \to W$. Define a topological 4-manifold W' by

$$W' = mS^2 \times S^2 \# k(-\mathbb{CP}^2) \# 2(-E_8) \# 2nS^2 \times S^2.$$

By Freedman theory [12], we have a homeomorphism $h: W \to W'$. Define orientation-preserving diffeomorphisms $f_0: S^2 \times S^2 \to S^2 \times S^2$ and $f_1: -\mathbb{CP}^2 \to -\mathbb{CP}^2$ by $f_0(x,y) = (y,x)$ and $f_1([z_0:z_1:z_2]) = [\bar{z}_0:\bar{z}_1:\bar{z}_2]$. Both of f_0 and f_1 have codimension-2 fixed-point set, diffeomorphic to S^2 and \mathbb{RP}^2 respectively, and we may form an equivariant connected sum

$$f_{\#} = \#_m f_0 \#_k f_1 : mS^2 \times S^2 \# k(-\mathbb{CP}^2) \to mS^2 \times S^2 \# k(-\mathbb{CP}^2).$$

Choose a point $x_0 \in mS^2 \times S^2 \# k(-\mathbb{CP}^2)$ outside the fixed-point set of $f_{\#}$, and attach two copies of $(-E_8)\# nS^2 \times S^2$ along x_0 and $f_{\#}(x_0)$. Now we obtain a locally linear involution f' of W', and define $\iota = h^{-1} \circ f' \circ h$.

It is straightforward to check that

$$b_{f_0}^+(S^2 \times S^2) = 1, \quad b_{f_0}^-(S^2 \times S^2) = 0, \quad b_{f_1}^+(-\mathbb{CP}^2) = 0, \quad b_{f_1}^-(-\mathbb{CP}^2) = 0,$$
 and thus we have

(35)
$$b_{\iota}^{+}(W) = m, \quad b_{\iota}^{-}(W) = n+8.$$

Let $c' \in H^2(W'; \mathbb{Z})$ be a characteristic element defined by

$$c' = (0_S, E_1, \dots, E_k, 0_{ES}),$$

where 0_S and 0_{ES} denote the zero elements of $H^2(mS^2 \times S^2)$ and $H^2(2(-E_8) \# 2nS^2 \times S^2)$ respectively, and E_i are copies of a generator of $H^2(-\mathbb{CP}^2)$. Let $c = h^*c'$ and let \mathfrak{s} be a spin^c structure on W with $c_1(\mathfrak{s}) = c$. Since $f_1^*E = -E$, we have that $\iota^*\mathfrak{s} = \overline{\mathfrak{s}}$.

Since ι has codimension-2 fixed-point set, by Lemma 2.9 ι is of odd type. Hence it follows from (35) and and Theorem 1.1 that $c_1(\mathfrak{s})^2 - \sigma(W) \leq 0$. However, since we have $c_1(\mathfrak{s})^2 - \sigma(W) = 16$ by a direct calculation, this is a contradiction.

5.2. Applications to non-smoothable and non-orientable surfaces. The above argument on non-smoothable actions can be translated to a result on smoothable and unorientable surfaces, Theorem 1.23:

Proof of Theorem 1.23. Let us use the notation of the statement and the proof of Theorem 1.24. Then it follows that X is homeomorphic to W/ι . Let S be the fixed-point set of $\iota: W \to W$. Then S is homeomorphic to $k\mathbb{RP}^2$ and is locally-flatly embedded in W, and hence also in X. If S is topologically isotopic to a smoothly embedded surface, it induces a smooth odd involution that acts on homology just as ι does, but it is a contradiction by the proof of Theorem 1.24.

We shall see what the homology class of S is. As the fixed-point set of f_0 : $S^2 \times S^2 \to S^2 \times S^2$ is the diagonal, which descends to a 2-sphere in \mathbb{CP}^2 that represent $2[\mathbb{CP}^1] \in H_2(\mathbb{CP}^2)$. On the other hand, $-\mathbb{CP}^2/f_1$ is diffeomorphic to S^4 . Thus we have that the homology class of S is zero in $H_2(X;\mathbb{Z}/2)$.

The normal Euler number $[k\mathbb{RP}^2]^2$ can be immediately deduced from Lemma 4.5.

5.3. Applications to the Nielsen realization problem. In this subsection, we shall prove a result on the Nielsen realization problem for non-spin 4-manifolds, Theorem 1.16. First, we recall the definition of a diffeomorphism called the *reflection* about a (-1)-sphere in a 4-manifold:

Definition 5.1. Let W be a smooth oriented 4-manifold W and S a (-1)-sphere S in W. Then one may form an orientation-preserving diffeomorphism

$$\rho_S: W \to W$$

called the *reflection*, constructed as follows: first, let $\rho : \mathbb{CP}^2 \to \mathbb{CP}^2$ be the complex conjugation, $[z_0 : z_1 : z_2] \mapsto [\bar{z}_0 : \bar{z}_1 : \bar{z}_2]$. Take an embedded 4-disk D^4 in \mathbb{CP}^2 . Let ρ' be a diffeomorphism that fixes D^4 pointwise obtained by deforming ρ near D^4 by isotopy. By reversing orientation, it induces a diffeomorphism $\rho' : -\mathbb{CP}^2 \to -\mathbb{CP}^2$. In general, for a (-1)-sphere S in W, a tubular neighborhood of S is diffeomorphic to a tubular neighborhood of $-\mathbb{CP}^1$ in $-\mathbb{CP}^2$, which is the fixed point set of ρ . Since a tubular neighborhood of $-\mathbb{CP}^1$ in $-\mathbb{CP}^2$ is diffeomorphic also to $\mathbb{CP}^2 \setminus \operatorname{int} D^4$, we may implant ρ' into W by extending it by the identity, which we denote by $\rho_S : W \to W$. To prove our results on the Nielsen realization problem, we need to take care of Dehn twitsts. First, let us recall the definition of the Dehn twist along an embedded annulus:

Definition 5.2. Let W be a 4-manifold and let $A \cong S^3 \times [0, 1]$ be an embedded annulus in W. Then one may define an orientation-preserving diffeomorphism

$$\tau: W \to W$$

called the *Dehn twist along* A as follows. Define the *model Dehn twist* $\tau_0 : S^3 \times [0,1] \to S^3 \times [0,1]$ by $\tau_0(y,t) = (l(t) \cdot y,t)$, where $l : [0,1] \to SO(4)$ is the homotopically non-trivial loop in SO(4). As τ_0 is supported inside $S^3 \times (0,1)$, one may extend τ_0 by the identity and get a diffeomorphism $\tau : W \to W$.

Note that, by $\pi_1(SO(4)) = \mathbb{Z}/2$, the square of the Dehn twist τ along an annulus is smoothly isotopic to the identity through an isotopy supported inside the annulus.

Given a closed smooth 4-manifold W, let $\text{Diff}(W, D^4)$ denote the group of diffeomorphisms of W that are identity on a fixed embedded 4-disk $D^4 \subset W$. Let

$$p: \pi_0(\operatorname{Diff}(W, D^4)) \to \pi_0(\operatorname{Diff}(W))$$

be the map induced from the natural inclusion $\text{Diff}(W, D^4) \hookrightarrow \text{Diff}(W)$. Note that p is surjective since any diffeomorphism can be isotoped to a diffeomorphism that fixes D^4 pointwise.

Lemma 5.3 (Giansiracusa [13, Corollary 2.5, Proposition 3.1]). Let W be a simplyconnected closed smooth 4-manifold W. Then we have the following:

- (1) ker p is isomorphic to either $\mathbb{Z}/2$ or 0, and it is generated by the Dehn twist around the boudnary $\partial(W \setminus D^4)$.
- (2) If W is diffeomorphic to $W' # \mathbb{CP}^2$ or $W' # (-\mathbb{CP}^2)$ for some 4-manifold W', then ker p = 0.

In summary, one has an exact sequence

 $0 \to \ker p \to \pi_0(\operatorname{Diff}(W, D^4)) \xrightarrow{p} \pi_0(\operatorname{Diff}(W)) \to 0$

such that ker p is either $\mathbb{Z}/2$ or 0 generated by the Dehn twist, and ker p = 0 if W contains either \mathbb{CP}^2 or $-\mathbb{CP}^2$ as a connected sum factor.

Using Lemma 5.3, we see that the reflection about a (-1)-sphere gives an order 2 mapping class:

Lemma 5.4. For an oriented 4-manifold W and a (-1)-sphere S in W, the mapping class $[\rho_S]$ of the reflection is of order 2 in $\pi_0(\text{Diff}(W))$.

Proof. First, clearly ρ is an order 2 element in $\text{Diff}(\mathbb{CP}^2)$, and so is the mapping class $[\rho]$ in $\pi_0(\text{Diff}(\mathbb{CP}^2))$. By Lemma 5.3, the natural surjection

$$p: \pi_0(\operatorname{Diff}(\mathbb{CP}^2, D^4)) \to \pi_0(\operatorname{Diff}(\mathbb{CP}^2)))$$

is isomorphic. This implies that the relative mapping class $[\rho']$ is of order 2 in $\pi_0(\text{Diff}(\mathbb{CP}^2, D^4))$. The claim of the lemma now follows immediately by the construction of ρ_S .

We shall also use the following lemma about simultaneous Dehn twists on punctured S^4 :

Lemma 5.5 ([20, Lemma 4.3]). Let N > 0. Define S_0^4 to be a N-punctured 4sphere, $S_0^4 = S^4 \setminus (\bigsqcup_{i=1}^N D_i^4)$. Let $\tau_{S_0^4} : S_0^4 \to S_0^4$ be a diffeomorphism defined by performing the Dehn twists near all ∂D_i^4 simultaneously. Then $\tau_{S_0^4}$ is smoothly isotopic to the identity in the group $\text{Diff}(\partial S_0^4, \partial S_0^4)$ of diffeomorphisms which fix the boundary pointwise.

Now we are ready to prove the main result in this subsection, Theorem 1.16:

Proof of Theorem 1.16. Suppose that ρ_S is homotopic to a smooth involution ι : $W \to W$. Then we have $\iota^* \mathfrak{s} \cong \overline{\mathfrak{s}}$.

Recall that the induced action of ρ_S on homology $H_2(W)$ is given by $(\rho_S)_*(x) = x + 2(x \cdot [S])[S]$. Set $\varphi = (\rho_S)_* : H_2(W) \to H_2(W)$. Then we have

$$b_{\varphi}^{+}(W) = b^{+}(W), \quad b_{\varphi}^{-}(W) = b^{-}(W) - 1, \quad \sigma_{\varphi}(W) = \sigma(W) + 1.$$

Here $b_{\varphi}^+(W)$ and $b_{\varphi}^-(W)$ denote the maximal dimension of the φ -invariant part of $H^+(W)$ and of $H^-(W)$ respectively, and we set $\sigma_{\varphi}(W) = b_{\varphi}^+(W) - b_{\varphi}^-(W)$. By the assumption that $\sigma(W) \neq -2$, we have $\sigma_{\varphi}(W) \neq \sigma(W)/2$. It follows from Lemma 2.10 that ι is of odd type. Combined this with $b_{\varphi}^+(W) = b^+(W)$, we may apply Theorem 1.1 to ι , but this contradicts the assumption that $c_1(\mathfrak{s})^2 - \sigma(W) > 0$.

Proof of Theorem 1.22. As the case that r = 0 has been treated in [20, Theorem 1.3], we suppose that r > 0. Set $W' = pK3\#qS^2 \times S^2$. In [20, Proof of Theorem 1.3], a locally linear topological involution $f: W' \to W'$ with the following properties was constructed:

- $b_f^+(W') = 3p + q, \ \sigma_f(W') = -5p + q,$
- the fixed point set of f is non-empty and of codimension-2.
- there is a diffeomorphism $g: W' \to W'$ such that g^2 is smoothly isotopic to the identity and $g_* = f_*$ on $H_2(W'; \mathbb{Z})$.

Let $\rho : -\mathbb{CP}^2 \to -\mathbb{CP}^2$ be the complex conjugation. The fixed point set of ρ is of codimension-2: it is given by $\mathbb{RP}^2 \subset -\mathbb{CP}^2$. Choose points in \mathbb{RP}^2 , and form an equivariant connected sum of *r*-copies of ρ along those points to get an involution on $r(-\mathbb{CP}^2)$, denoted by $\#_r\rho$.

On the other hand, recall that the fixed point set of f is also codimension-2. Also, since f is locally linear, we have the notion of tangential representation of f at a fixed point of f. Since f is of order 2 and has codimension-2 fixed point set, the representation is diag(-1, -1, 1, 1) in some coordinate, which coincides with the tangential representation of $\#_r \rho$ at a fixed point. Thus we may form an equivariant connected sum of f with $\#_r \rho$, which we denote by $\tilde{f}: W \to W$.

Let $g': W' \to W'$ be a diffeomorphism obtained by deforming g by isotopy so that g' has a pointwise fixed 4-disk $D_1^4 \subset W'$. Similarly, by deforming $\#_r \rho$ by isotopy, we may get a diffeomorphism $g'': r(-\mathbb{CP}^2) \to r(-\mathbb{CP}^2)$ that has a fixed 4-disk D_2^4 . Let S_0^4 be the 2-punctured 4-sphere, $S_0^4 = S^4 \setminus (\sqcup_{i=1}^2 D_i^4)$. Regard W as the connected sum of W' with $r(-\mathbb{CP}^2)$ along D_1^4 and D_2^4 ,

$$W = (W' \setminus D_1^4) \cup_{\partial D_1^4} S_0^4 \cup_{\partial D_2^4} (r(-\mathbb{CP}^2) \setminus D_2^4).$$

Form a diffeomorphism $\tilde{g} : W \to W$ by gluing g' with g'' along D_1^4 and D_2^4 . Evidently, \tilde{g} is supported outside S_0^4 . By construction, the induced action of \tilde{g} on homology coincides with that of \tilde{f} . This combined with a result by Quinn [46] and Perron [45] implies that \tilde{g} is topologically isotopic to \tilde{f} . Also, it is easy to see that

(36)
$$b_{\tilde{a}}^+(W) = 3p + q, \quad \sigma_{\tilde{a}}(W) = -5p + q$$

using the action of f on homology.

We claim that \tilde{g}^2 is smoothly isotopic to the identity. First, let τ_1 and τ_2 denote the Dehn twists on W along ∂D_1^4 and ∂D_2^4 respectively. Recall that g^2 is smoothly isotopic to the identity and $(\#_r \rho)^2 = 1$. It follow from this together with (1) of Lemma 5.3 that \tilde{g}^2 is smoothly isotopic to one of the following: $\tau_1 \circ \tau_2$, τ_1 , τ_2 and id. Also, by (2) of Lemma 5.3, τ_2 is smoothly isotopic to the identity in Diff $(r(-\mathbb{CP}^2) \setminus \text{Int} D_2^4, \partial D_2^4)$. So it suffices to prove that τ_1 is smoothly isotopic to the identity. However, Lemma 5.5 implies that τ_1 is smoothly isotopic to τ_2 , which is smoothly isotopic to the identity again by (2) of Lemma 5.3. This proves the claim.

Let G be the subgroup of $\pi_0(\text{Diff}(W))$ generated by the mapping class of \tilde{g} . By the above claim, G is of order 2. As \tilde{g} is topologically isotopic to \tilde{f} , the group G' defined as the image of G under the map $\pi_0(\text{Diff}(W)) \to \pi_0(\text{Homeo}(W))$ is realized in Homeo(W). Also, the group G' is non-trivial since the action of \tilde{f} on homology is non-trivial. This proves the statement on G' in the theorem.

What remains to show is that G cannot be realized in Diff(W). This is equivalent to prove that \tilde{g} is not smoothly isotopic to a smooth involution. Suppose that \tilde{g} is smoothly isotopic to a smooth involution $\iota : W \to W$. Let \mathfrak{s} be the spin^c structure on W obtained as the connected sum of the unique spin structure on W' with *r*-copies of the spin^c structure on $-\mathbb{CP}^2$ whose first Chern class generates $H^2(-\mathbb{CP}^2)$. Then it is obvious that $\iota^*\mathfrak{s} \cong \bar{\mathfrak{s}}$ and $c_1(\mathfrak{s}) - \sigma(W) > 0$. It follows from this combined with Lemma 2.10 and (36) that ι is of odd type. Again by (36), we have $b^+(W) = b_{\iota}^+(W)$. Thus we have a contradiction by Theorem 1.1. This proves the claim and completes the proof of the theorem. \Box

5.4. Several inequalities for knots. In this subsection, we give a proof of Theorem 1.8 which describes a bound for the first Betti numbers of non-orientable surfaces embedded into D^4 .

Proof of Theorem 1.8. Let S be a (possibly non-orientable) smoothly and properly embedded connected surface in D^4 bounded by a given knot K. Suppose the determinant of K is 1, the Manolescu–Owen invariant $\delta(\Sigma(K))$ [35] is zero and $\delta_R(K) < 0$. As a known topological obstruction (which can be regarded as $b^+(S) \ge$ 0), we have

$$\left|\sigma(K) - \frac{1}{2}S \circ S\right| \le b_1(S).$$

Thus it is sufficient to suppose

$$-\sigma(K) + \frac{1}{2}S \circ S = b_1(S)$$

and obtain a contradiction. In order to use Theorem 1.3(iii), we verify the assumptions. We first note that (3) follows from $-\sigma(K) + \frac{1}{2}S \circ S = b_1(S)$. Since K is determinant-1, $\Sigma(K)$ is a homology 3-sphere. So, the intersection form of $\Sigma(S)$ is unimodular. From Lemma 4.5, we know

$$-\sigma(\Sigma(S)) = \frac{1}{2}S \circ S - \sigma(K) = b_1(S) = b_2(\Sigma(S)).$$

This implies the intersection form of $\Sigma(S)$ is negative definite. Now, since we supposed the Manolescu–Owen invariant $\delta(\Sigma(K))$ is zero, from Frøyshov inequality and Elikies's theorem, we know that the intersection form of $\Sigma(S)$ is the direct sum of copies of (-1). From Lemma 4.5, we know that the double branched covering space $\Sigma(S)$ has no 2-torsions in $H^2(\Sigma(S);\mathbb{Z})$. Thus, the set of isomorphism classes of spin^c structures on $\Sigma(S)$ is determined by their first Chern classes in $H^2(\Sigma(S);\mathbb{Z})$. Note that the action of ι^* on $H^2(\Sigma(S);\mathbb{Z})$ gives a decomposition

$$H^{2}(\Sigma(S);\mathbb{Z}) = H^{2}(\Sigma(S);\mathbb{Z})^{\iota^{*}} \oplus H^{2}(\Sigma(S);\mathbb{Z})^{-\iota^{*}}$$

as the eigenvalue decomposition of ι corresponding to +1 and -1 respectively. Note that the dimension dim $H^2(\Sigma(S);\mathbb{Z})^{\iota^*} = b_2(\Sigma(S)/\iota) = b_2(D^4) = 0$. This implies $H^2(\Sigma(S);\mathbb{Z}) = H^2(\Sigma(S);\mathbb{Z})^{-\iota^*}$. So, by taking a spin^c structure corresponding to the sum of basis of $\bigoplus (-1)$ and by applying Theorem 1.3(iii), we have

$$0 \leq \delta(K)$$

This gives a contradiction to $\delta(K) < 0$.

We next prove crossing change formulae for δ_R , $\overline{\delta}_R$ and $\underline{\delta}_R$.

Proof of Remark 1.9. Let K and K' be knots in S^3 . Suppose K' is obtained from K by a positive crossing change. By blown up, we obtain a smooth annulus cobordism S in $I \times S^3 \# - \mathbb{CP}^2$ from (S^3, K) to (S^3, K') such that

$$[S] = 2 \in \mathbb{Z} = H_2(-\mathbb{CP}^2) = H_2(I \times S^3 \# - \mathbb{CP}^2).$$

Suppose $2 = \sigma(K) - \sigma(K)$. Then, for the unique spin structure on $\Sigma(S)$, the assumptions of Theorem 1.3(iii) are satisfied. Then Theorem 1.3(iii) implies the desired inequality. In the case of $\sigma(K) - \sigma(K') = 0$, we use $\overline{I \times S^3 \# (-\mathbb{CP}^2)}$ instead of $I \times S^3 \# (-\mathbb{CP}^2)$ and Theorem 1.3(iii) again.

5.5. Applications to non-orientable genera and normal Euler numbers.

Proof of Theorem 1.14. Note that K = T(3, 6n+1) is determinant 1, the Manolescu– Owen's invariant is known to be zero and $\delta_R(T(3, 6n+1)) = -\frac{1}{2} < 0$. Thus all assumptions of Theorem 1.8 are satisfied. For a smooth embedded connected surfaces in D^4 bounded by K satisfying

$$(e,h) = (8/3(1-n) + 2 + 2m, 1+m),$$

we can check $-\sigma(K) + \frac{1}{2}S \circ S = b_1(S)$. However, from Theorem 1.8, we have the stronger inequality $-\sigma(K) + \frac{1}{2}S \circ S + 1 \leq b_1(S)$ which gives a contradiction. This completes the proof.

References

- Samantha Allen, Nonorientable surfaces bounded by knots: a geography problem (2020), available at arXiv:2007.14332.
- [2] M. Arabadji and I. Baykur, The Nielsen realization for non-spin 4-manifolds, in preparation.
- [3] David Baraglia, Obstructions to smooth group actions on 4-manifolds from families Seiberg-Witten theory, Adv. Math. 354 (2019), 106730, 32. MR3981995
- [4] David Baraglia and Hokuto Konno, A note on the Nielsen realization problem for K3 surfaces, arXiv:1908.03970 (2019). to appear in Proc. Amer. Math. Soc.
- [5] Joshua Batson, Nonorientable slice genus can be arbitrarily large, Math. Res. Lett. 21 (2014), no. 3, 423–436. MR3272020
- [6] Fraser Binns, Sungkyung Kang, Jonathan Simone, and Paula Truöl, On the nonorientable four-ball genus of torus knots (2021), available at arXiv:2109.09187.
- [7] Jim Bryan, Seiberg-Witten theory and Z/2^p actions on spin 4-manifolds, Math. Res. Lett. 5 (1998), no. 1-2, 165–183. MR1617929
- [8] Andrew Donald and Brendan Owens, Concordance groups of links, Algebr. Geom. Topol. 12 (2012), no. 4, 2069–2093. MR3020201
- S. K. Donaldson, An application of gauge theory to four-dimensional topology, J. Differential Geom. 18 (1983), no. 2, 279–315. MR710056
- [10] _____, Floer homology groups in Yang-Mills theory, Cambridge Tracts in Mathematics, vol. 147, Cambridge University Press, Cambridge, 2002. With the assistance of M. Furuta and D. Kotschick. MR1883043
- [11] Benson Farb and Eduard Looijenga, The Nielsen realization problem for K3 surfaces, arXiv:2104.08187 (2021). to appear in J. Differential Geom.
- [12] Michael Hartley Freedman, The topology of four-dimensional manifolds, J. Differential Geometry 17 (1982), no. 3, 357–453. MR679066
- [13] Jeffrey Giansiracusa, The stable mapping class group of simply connected 4-manifolds, J. Reine Angew. Math. 617 (2008), 215–235. MR2400996
- [14] Robert E. Gompf and András I. Stipsicz, 4-manifolds and Kirby calculus, Graduate Studies in Mathematics, vol. 20, American Mathematical Society, Providence, RI, 1999. MR1707327
- [15] Kristen Hendricks, Jennifer Hom, and Tye Lidman, Applications of involutive Heegaard Floer homology, J. Inst. Math. Jussieu 20 (2021), no. 1, 187–224. MR4205781
- [16] Kristen Hendricks and Ciprian Manolescu, Involutive Heegaard Floer homology, Duke Math. J. 166 (2017), no. 7, 1211–1299. MR3649355
- [17] Yuya Kato, Nonsmoothable actions of Z₂ × Z₂ on spin four-manifolds, Topology Appl. 307 (2022), Paper No. 107868, 13. MR4365044
- [18] Louis H. Kauffman and Laurence R. Taylor, Signature of links, Trans. Amer. Math. Soc. 216 (1976), 351–365. MR388373
- [19] Tirasan Khandhawit, A new gauge slice for the relative Bauer-Furuta invariants, Geom. Topol. 19 (2015), no. 3, 1631–1655. MR3352245
- [20] Hokuto Konno, Dehn twists and the Nielsen realization problem for spin 4-manifolds, arXiv:2203.11631 (2022). to appear in Algebr. Geom. Topol.
- [21] Hokuto Konno, Jin Miyazawa, and Masaki Taniguchi, Involutions, knots, and Floer K-theory (2021), available at arXiv:2110.09258.
- [22] Peter Kronheimer and Tomasz Mrowka, Monopoles and three-manifolds, New Mathematical Monographs, vol. 10, Cambridge University Press, Cambridge, 2007. MR2388043
- [23] Terry Lawson, Normal bundles for an embedded rp2 in a positive definite 4-manifold, Journal of Differential Geometry 22 (1985), no. 2, 215–231.
- [24] Ronnie Lee and Steven H. Weintraub, On the homology of double branched covers, Proc. Amer. Math. Soc. 123 (1995), no. 4, 1263–1266. MR1224618
- [25] Seraphina Eun Bi Lee, Isotopy classes of involutions of del Pezzo surfaces, arXiv:2202.10616 (2022).
- [26] Adam Levine, Daniel Ruberman, and Sašo Strle, Nonorientable surfaces in homology cobordisms, Geometry & Topology 19 (2015), no. 1, 439–494.
- [27] Lukas Lewark, Rasmussen's spectral sequences and the sln-concordance invariants, Advances in mathematics 260 (2014), 59–83.
- [28] Jiakai Li, Monopole Floer Homology and Real Structures (2022), available at arXiv:2211.10768.

- [29] Tye Lidman and Ciprian Manolescu, The equivalence of two Seiberg-Witten Floer homologies, Astérisque 399 (2018), vii+220. MR3818611
- [30] Charles Livingston, Computations of the ozsváth-szabó knot concordance invariant, Geometry & Topology 8 (2004), no. 2, 735–742.
- [31] Andrew Lobb, A counterexample to Batson's conjecture, Math. Res. Lett. 26 (2019), no. 6, 1789. MR4078695
- [32] Ciprian Manolescu, Seiberg-Witten-Floer stable homotopy type of three-manifolds with $b_1 = 0$, Geom. Topol. 7 (2003), 889–932. MR2026550
- [33] _____, On the intersection forms of spin four-manifolds with boundary, Math. Ann. 359 (2014), no. 3-4, 695–728. MR3231012
- [34] _____, Pin(2)-equivariant Seiberg-Witten Floer homology and the triangulation conjecture,
 J. Amer. Math. Soc. 29 (2016), no. 1, 147–176. MR3402697
- [35] Ciprian Manolescu and Brendan Owens, A concordance invariant from the Floer homology of double branched covers, Int. Math. Res. Not. IMRN 20 (2007), Art. ID rnm077, 21. MR2363303
- [36] William Massey, Proof of a conjecture of whitney, Pacific Journal of Mathematics 31 (1969), no. 1, 143–156.
- [37] J. P. May, Equivariant homotopy and cohomology theory, CBMS Regional Conference Series in Mathematics, vol. 91, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1996. With contributions by M. Cole, G. Comezaña, S. Costenoble, A. D. Elmendorf, J. P. C. Greenlees, L. G. Lewis, Jr., R. J. Piacenza, G. Triantafillou, and S. Waner. MR1413302
- [38] Ian Montague, Seiberg-Witten Floer K-theory and cyclic group actions on spin four-manifolds with boundary (2022), available at arXiv:2210.08565.
- [39] Seiji Nagami, Existence of Spin structures on double branched covering spaces over fourmanifolds, Osaka J. Math. 37 (2000), no. 2, 425–440. MR1772842
- [40] Nobuhiro Nakamura, Bauer-Furuta invariants under Z₂-actions, Math. Z. 262 (2009), no. 1, 219–233. MR2491607
- [41] _____, Pin⁻(2)-monopole equations and intersection forms with local coefficients of fourmanifolds, Math. Ann. 357 (2013), no. 3, 915–939. MR3118618
- [42] _____, Pin⁻(2)-monopole invariants, J. Differential Geom. 101 (2015), no. 3, 507–549. MR3415770
- [43] Walter D. Neumann, An invariant of plumbed homology spheres, Topology Symposium, Siegen 1979 (Proc. Sympos., Univ. Siegen, Siegen, 1979), 1980, pp. 125–144. MR585657
- [44] Peter S. Ozsváth, András I. Stipsicz, and Zoltán Szabó, Concordance homomorphisms from knot Floer homology, Adv. Math. 315 (2017), 366–426. MR3667589
- [45] B. Perron, Pseudo-isotopies et isotopies en dimension quatre dans la catégorie topologique, Topology 25 (1986), no. 4, 381–397. MR862426
- [46] Frank Quinn, Isotopy of 4-manifolds, J. Differential Geom. 24 (1986), no. 3, 343–372. MR868975
- [47] Frank Raymond and Leonard L. Scott, Failure of Nielsen's theorem in higher dimensions, Arch. Math. (Basel) 29 (1977), no. 6, 643–654. MR467773
- [48] Joshua M. Sabloff, On a refinement of the non-orientable 4-genus of torus knots (2022), available at arXiv:2206.11200.
- [49] Matthew Stoffregen, Manolescu invariants of connected sums, Proceedings of the London Mathematical Society 115 (2017), no. 5, 1072–1117.
- [50] _____, A remark on Pin(2)-equivariant Floer homology, Michigan Math. J. 66 (2017), no. 4, 867–884. MR3720328
- [51] _____, Pin(2)-equivariant Seiberg-Witten Floer homology of Seifert fibrations, Compos. Math. 156 (2020), no. 2, 199–250. MR4044465
- [52] Gang Tian and Shuguang Wang, Orientability and real Seiberg-Witten invariants, Internat. J. Math. 20 (2009), no. 5, 573–604. MR2526308
- [53] V. G. Turaev, Classification of oriented Montesinos links by means of invariants of spin structures, 1985, pp. 130–146, 178. Studies in topology, V. MR806564

Graduate School of Mathematical Sciences, the University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8914, Japan

Email address: konno@ms.u-tokyo.ac.jp

Graduate School of Mathematical Sciences, the University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8914, Japan

Email address: miyazawa@ms.u-tokyo.ac.jp

Department of Mathematics, Graduate School of Science, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

 $Email \ address: \verb+taniguchi.masaki.7m@kyoto-u.ac.jp$