Minimal models, modular linear differential equations and modular forms of fractional weights

Kiyokazu Nagatomo^{[1](#page-0-0)} and Yuichi Sakai^{[2](#page-0-1)}

¹ Department of Pure and Applied Mathematics Graduate School of Information Science and Technology Osaka University, Suita, Osaka 565-0871, Japan

² Fundamental Educational Center Kurume Institute of Technology 2228-66, Kamitsu, Kurume, Fukuoka 830-0052, Japan e-mail: dynamixaxs@gmail.com

Abstract

We show that modular forms of fractional weights on principal congruence subgroups of odd levels, which are found by T. Ibukiyama, naturally appear as characters being multiplied η^{ceff} of the so-called minimal models of type $(2, p)$, where c_{eff} is the effective central charge of a minimal model. Using this fact and modular invariance property of the space of characters, we give a different proof of the result showned by Ibukiyama that is the explicit formula representing $SL_2(\mathbf{Z})$ on the space of the ibukiyama modular forms. We also find several pairs of spaces of the Ibukiyama modular forms on $\Gamma(m)$ and $\Gamma(n)$ with $m|n$ having the property that the former are included in the latter. Finally, we construct vector-valued modular forms of weight $k \in \frac{1}{5}\mathbb{Z}_{>0}$ starting from the Ibukiyama modular forms of weight $k \in \frac{1}{5}\mathbb{Z}_{>0}$ with some multiplier system by a symmetric tensor product of this representation and show that the components functions coincides with the solution space of some monic modular linear differential equation of weight k and the order $1 + 5k$.

Introduction

In this paper we study modular forms of fractional weights found by T. Ibukiyama [\[10\]](#page-16-0) from the conformal field theoretic point of view. The principal goal of this paper is to relate these modular forms to the characters of the so-called *minimal model* of type $(2, p)$ with an odd integer $p > 3$, which is a well-known example of 2D conformal field theories. It was proved in [\[17\]](#page-16-1) that the space of characters of any minimal model of type (P,Q) is equal to the solution space of some monic linear differential equation of order $(P-1)(Q-1)/2$. We shows, in this paper, that the Ibukiyama modular forms on $\Gamma(p)$ satisfy some monic linear *modular* differential equation (MLDE) of the same order $(p-1)/2$ but a different weight. The space of the Ibukiyama modular forms, where v_p is a certain multiplier system, is equal to the space $M_{\frac{p-3}{2p}}(\Gamma(p), v_p)$ of modular forms of weights $\frac{p-3}{2p}$ on $\Gamma(p)$ for any odd integer $p < 15$. (See Lemma 1 and Lemma 2 in section 3 in [\[9\]](#page-16-2).) However, for $p = 15$ and 23, it was shown in section 4 of [\[9\]](#page-16-2) that $M_{\frac{p-3}{2p}}(\Gamma(p), v_p)$ do not coincide with the space of Ibukiyama modular forms. The dimension is known [\[10\]](#page-16-0) Lemma 1.7.

It is known that the character ch_M of an arbitrary module M over a rational simple self-dual vertex operator algebra (VOA) V of CFT and of finite type $(C_2$ -cofinite) is a modular function on

¹The first author is supported in part by International Center of Theoretical Physics, Italy, and Max-Planck institute for Mathematics, Germany.

²The second author is supported in part by JSPS KAKENHI Grant numbers JP21K03183, JP18K03215 and JP16H06336.

the congruence subgroup $\Gamma(n)$ of $SL_2(\mathbf{Z})$, where n is the minimal positive integer^{[1](#page-1-0)} such that $n(h-\mathbf{Z})$ $c/24$) is integral for any conformal weight h of all (inequivalent) simple V-modules, where c is the central charge of V. (See Theorem I and Theorem 3.10 in $[4]$.) The central charges and conformal weights of a simple VOA of finite type are rational numbers, and number of their modules is finite.) Let c_{eff} be the *effective central charge* defined by $c_{\text{eff}} = c - 24h_{\text{min}}$, where h_{min} is the minimal element of the (finite) set of conformal weights. Then $\eta^{c_{\text{eff}}} \ch_M$ is a modular function, where ch_M is a character of the minimal model of type $(2, p)$ with the multiplier system that of η^{c} on $\Gamma(n)$. For instance, for the minimal model type $(2,5)$ we have $n_5 = 60$ and $c_{\text{eff}} = 2/5$. Therefore $\eta^{2/5}$ ch_M is a modular function on $\Gamma(60)$ with the multiplier of $\eta^{2/5}$. We will show that the Ibukiyama modular form of weight $\frac{p-3}{2p}$ on the congruence group $\Gamma(p)$ with some multiplier system v_p equals $\eta^{c_{\text{eff}}}$ ch_M. It implies that ch_M is a modular form with some non-trivial multiplier system on $\Gamma(p)$ since the multiplier system of the Ibukiyama modular forms on $\Gamma(p)$ is not equal to that of η^{c} on $\Gamma(p)$.

There are at least two interesting problems of modular forms of fractional weight. (a) Any $f \in M_*(\Gamma(p), \nu)$ is standard, where ν is a multiplier, i.e., whether there exists a positive integer m such that $f^m \in M_*(\Gamma(p), 1)$. An arbitrary Ibukiyama modular form is proved to be standard (Theorem 1.1 (3), Section 1.1) in [\[10\]](#page-16-0). However, it is not known whether modular functions $\eta^{\text{c}_{\text{eff}}}$ ch_M for any minimal models always satisfy this important property. Of course they are standard on $\Gamma(24) \cap \Gamma(n)$. (b) let $M_k(\Gamma(m), \mu)$ and $M_k(\Gamma(n), \nu)$ be spaces of modular forms of weight k on $\Gamma(m)$ and $\Gamma(n)$ with multiplier systems μ and ν satisfying $\nu|_{\Gamma(m)} = \mu$ for m|n. The question is that if m|n we have $M_k(\Gamma(m), \mu) \subset M_k(\Gamma(n), \nu)$. Note that any spaces of modular forms of (half-)*integral weight* of congruence groups of $SL_2(\mathbf{R})$ have the property (b). In this paper we show that the pair of the spaces $M_*(\Gamma(5), v_5) := \bigoplus_{m=0}^{\infty} M_{\frac{m}{5}}(\Gamma(5), v_5^m)$ and $M_*(\Gamma(15), v_{15}) := \bigoplus_{m=0}^{\infty} M_{\frac{2m}{5}}(\Gamma(15), v_{15}^m)$ and $M_*(\Gamma(7), v_7) := \bigoplus_{m=0}^{\infty} M_{2m/7}(\Gamma(7), v_7^m)$ and $M_*(\Gamma(21), v_{21}) := \bigoplus_{m=0}^{\infty} M_{3m/7}(\tilde{\Gamma(21)}, v_{21}^m)$ have the property (a).

Let f be an arbitrary (holomorphic or meromorphic) modular form of weight k on a Fuchsian group $\Gamma \subset PSL_2(\mathbf{R})$ such that $\mathfrak{H}\backslash\Gamma$ is genus 0. Then f satisfies a certain ordinary equation of order $k + 1$ with a local coordinate defined by a modular invariant function (Hauptmoduln) (e.g. p. 61, Propposition 21 [\[20\]](#page-16-3)) Let $k \in \frac{1}{5}\mathbb{Z}_{\geq 0}$. We construct vector-valued modular forms by using the symmetric representation of an action $SL_2(\mathbf{Z})$ on $M_k(\Gamma(5), v_5^{5k})$, and prove that the space spanned by components function of this VVMF is equal to the solution space of some MLDE is of weight k and order $1 + 5k$. Namely, vector-valued modular forms, which are constructed by fractional modular forms, provide examples of differential equations whose orders are higher than the weight $+1$.

In §[1](#page-2-0) we review the notion of multipliers systems, and introduce the multiplier systems which are used to define the Ibukiyama modular forms. The definition of the Ibukiyama modular forms is also given, and the invariance of the space of the Ibukiyama modular forms under some action π_p of $SL_2(\mathbf{Z})$ is mentioned in the same section. We briefly explain the notion of vertex operator algebras in §[2.](#page-3-0) The minimal models are introduced in §[3.](#page-4-0) Here we collect several facts of characters of minimal models which are used in the following sections. We further prove the formula that gives an explicit formula of $n_p (= n)$ of the minimal model of type $(2, p)$. We clarify in §[4](#page-6-0) the relation between the Ibukiyama modular forms and characters of the minimal models of type $(2, p)$ in Theorem [6.](#page-6-1) We then give a different proof of modular invariance of the Ibikuyama modular forms under the action of π_p by using the modular invariance property of the space of characters of minimal models. In §[5](#page-8-0) we derive MLDEs whose solution spaces are equal to the space of characters of the minimal models of type $(2, p)$ and then the space of the Ibukiyama modular forms, respectively. However, it is shown that they are immediate consequence of differential equations of Milas, and we *cannot* obtain any new way to find: using the fractional modular forms supersingular j-invariant as they did. Here we give several examples which satisfy the property (a) . In the final section [7](#page-14-0) we study the case $p = 5$. We compute an action of $SL_2(\mathbf{Z})$ on a basis of $M_{\frac{1}{5}}(\Gamma(5), v_5)$

¹See equation [\(8\)](#page-7-0) in Thereom [8](#page-7-1) for the general formula of $n = n_p$ of the minimal model of type $(2, p)$.

and show that vector-valued modular forms the space of the symmetric tensor representation of degree 5k associated with this representation above is equal to the solution space of some MLDE of the order $1 + 5k$.

1 Ibukiyama's modular forms

In this section we review the Ibukiyama modular forms which are main interest of this paper. Let denote $e(z) = e^{2\pi i z}$ with any complex number z. For any element $\vec{m} = (m_1, m_2) \in \mathbb{Q}^2$ we define the theta function in the complex upper half-plain \mathfrak{H} of *characteristic* \vec{m} by

$$
\Theta_{\vec{m}}(\tau, z) = \sum_{n \in \mathbf{Z}} e\left(\frac{1}{2}\tau(n + m_1)^2 + (n + m_1)(z + m_2)\right)
$$

for any $\tau \in \mathfrak{H}$ and any $z \in \mathbb{C}$. We denote the value of $\Theta_{\vec{m}}(\tau, z)$ at $z = 0$ by $\Theta_{\vec{m}}(\tau)$, which is called the theta constant. Let $\ell > 0$ be an odd prime number and ε_1 , ε_2 be odd integers. Let $m_1 = \varepsilon_1/\ell$ and $m_2 = \varepsilon_2/\ell$. Then and $\Theta_{\vec{m}}(\tau, z)$ is a modular form of weight $1/2$ on $\Gamma(\ell)$ (cf. [\[5\]](#page-15-1)).

Before giving definition of the multilinear systems we need the notion of "automorphy factor." Let $\Gamma \subset PSL_2(\mathbf{Z})$ be a Fuchsian group. An *automorphy factor* $j(\gamma, \tau)$ is a complex-valued function in $\Gamma \times \mathfrak{H}$, which is holomorphic in the complex upper half-plane with $j(\gamma_1 \gamma_2, \tau) = j(\gamma_1, \gamma_2 \tau)j(\gamma_2, \tau)$ for any $\gamma_1, \gamma_2 \in \Gamma$.

Let $v : \Gamma \to \mathbf{C}^\times$ $(\gamma \mapsto v(\gamma))$ be a map and let r be a rational number. For any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, let denote $j(\gamma, \tau) = v(\gamma)(c\tau + d)^r$, where we take the principal value for the branch of the rational power. We say that $v(\gamma)$ is a multiplier system if $j(\gamma_1\gamma_2, \tau) = j(\gamma_1, \gamma_2\tau)j(\gamma_2, \tau)$ for any $\gamma_1, \gamma_2 \in \Gamma$, and $j(\gamma, \tau)$ is an automorphy factor of weight r. We say that a holomorphic function $f(\tau)$ on \mathfrak{H} is a holomorphic modular form of weight r with a multiplier system $v(\gamma)$ if $f(\gamma \tau) = f(\tau)j(\gamma, \tau)$ for each $\gamma \in \Gamma$ and f is holomorphic at each cusp of Γ .

We denote by $v_0: SL_2(\mathbf{Z}) \to \mathbf{C}^\times$ the multiplier system of $\eta(\tau)^{3/p}$ for any odd integer $p > 3$ and for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbf{Z})$, i.e. $\eta(\gamma \tau)^{3/p} = v_0(\gamma) \eta(\tau)^{3/p} (c\tau + d)^{3/2p}$, and set $v_p = v_0^{p^2 - 1}$. Then v_p is a multiplier system.

The next theorem is proved in [\[10\]](#page-16-0).

Theorem 1 (Theorem 1.2 in [\[10\]](#page-16-0)). Let $p > 3$ be an odd integer. Then $v_p = v_0^{p^2-1}$ is a multiplier system on $\Gamma(p)$ and $j_p(\gamma, \tau) = v_p(\gamma)(c\tau + d)^{\frac{p-3}{2p}}$ is an automorphy factor of degree $\frac{p-3}{2p}$.

Definition. Let f be a modular form of non-integral weight k on $\Gamma(n)$. Then f is called *standard* if there is a positive integer m such that f^m is a modular form of integral weight with a trivial multiplier on some group including Γ_n .

We can now give the definition of the Ibukiyama modular forms.

Definition. Let $p > 2$ be an odd integer. For any odd integer $1 \leq r \leq p-2$, the *Ibukiyama modular* forms $f_r(\tau)$, which are holomorphic function in the complex upper-half plane \mathfrak{H} , are defined by

$$
f_r(\tau) = \frac{\Theta_{\vec{m}}(p\tau)}{\eta(\tau)^{3/p}} e\left(\frac{(p-1)(r-1)}{4p}\right), \quad \vec{m}_r = \left(\frac{r}{2p}, \frac{1}{2}\right).
$$

Theorem 2 ([\[10\]](#page-16-0)). Let $p > 3$ be an odd integer.

(a) For any odd integer r with $1 \leq r \leq p-2$, the Ibukiyama modular form f_r is a holomorphic modular form of weight $\frac{p-3}{2p}$ on $\Gamma(p)$ with the multiplier system v_p .

(b) The set of the functions $\{f_r\}_{r=1}^{p-2}$ is linearly independent.

(c) We have $v_p(\gamma)^p = 1$ for any $\gamma \in \Gamma(p)$. In particular, $(f_r)^p$ is a holomorphic modular form of integral weight $\frac{p-3}{2}$ on $\Gamma(p)$ for any $1 \le r \le p-2$, and then any modular form f_r is standard.

In §[4](#page-6-0) we relate the Ibukiyama modular forms to characters of the minimal model of type $(2, p)$ with an arbitrary odd integer $p > 3$. We denote the space of modular forms with multiplier systems v_p on $\Gamma(p)$ by $M_*(\Gamma(p), v_p) = \bigoplus_{m=0}^{\infty} M_{\frac{m(p-3)}{2p}}(\Gamma(p), v_p^m)$. However, at the moment, we cannot give a different proof, by using the relation to the minimal model, that f_r is standard on $\Gamma(p)$.

Definition. Let $p > 3$ be an odd integer. We define a left action π_p of $SL_2(\mathbf{Z})$ on $M_{\frac{p-3}{2p}}(\Gamma(p), v_p)$ by $\pi_p(\gamma) f(\tau) = j_p(\gamma, \tau)^{-1} f(\gamma(\tau))$ for any $f \in M_{\frac{p-3}{2p}}(\Gamma(p), v_p)$ and any $\gamma \in SL_2(\mathbf{Z})$.

Theorem 3 (Theorem 1.3 of [\[10\]](#page-16-0)). Let $p > 3$ be an odd integer.

(a) The function j_p is a factor of automorphy of $SL_2(\mathbb{Z})$, and is a prolongation of the factor of the automorphy of $\Gamma(p)$ of weight $(p-3)/2$ satisfying $j_p(-I_2, \tau) = (-1)^{(p+1)/2}$.

(b) The space of the Ibukiyama modular forms is spanned by f_r with all odd integers $1 \le r \le p-2$ is invariant under the action π_p of $SL_2(\mathbf{Z})$ as

$$
\pi_p(S)f_r = \sum_{\substack{1 \le s \le p-2, \\ s \equiv 1 \bmod 2}} \frac{1}{\sqrt{p}} \Big[e\Big(\frac{r-s}{4} + \frac{rs}{4p} + \frac{3p-1}{8}\Big) + (-1)^{\frac{p+1}{2}} e\Big(-\frac{r-s}{4} - \frac{rs}{4p} - \frac{3p-1}{8}\Big) \Big] f_s,
$$
\n
$$
\pi_p(T)f_r = e\Big(\frac{r^2 - p^2}{8p}\Big) f_r, \quad \pi_p(-I_2)f_r = (-1)^{(p+1)/2} f_r,
$$
\n(1)

where $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

We will show later that equation (1) is nothing but the modular invariance of the space of characters of the minimal model of type $(2, p)$ in §[4](#page-6-0)

2 Vertex operator algebras

The goal of this paper is to show that any Ibukiyama modular form of a fractional weight is equal to some character of a minimal model model of type $(2, p)$ multiplied by some power of the eta function. (See Theorem [6](#page-6-1) of §[4.](#page-6-0)) We will therefore explain the minimal models and their characters. To this end, we first collect definitions and basic properties related to vertex operator algebras.

A vertex operator algebra (VOA) is a quadruple $(V, Y, 1, \omega)$ with the following properties. The vector space V over the complex number field C is graded as $V = \bigoplus_{n=0}^{\infty} V_n$ with dim $V_n < \infty$ for all n. The element $1 \in V_0$ and $\omega \in V_2$ are called the vacuum vector and the Virasoro vector, respectively, and a linear map $Y: V \to \text{End}_{\mathbf{C}}(V)[[z, z^{-1}]]$ $(a \mapsto Y(a, z) = \sum_{n \in \mathbf{Z}} a_n z^{-n-1})$ satisfies a number of axioms. (For further information about axioms, see [\[14\]](#page-16-4) and [\[16\]](#page-16-5).) We define $L_n \in$ End(V) for any integer n by $Y(\omega, z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$. Then the set $\{L_n, id_V \mid n \in \mathbb{Z}\}\)$ forms a module over the Virasoro algebra on V. (The definition of the Virasoro algebra is given in §[3.](#page-4-0)) The action of the central element C is required to be a complex number, which is denoted by $c_V \in \mathbf{C}$ called the *central charge* of V in the literature. For each nonnegative integer n any element of a homogeneous subspace V_n has weight n with respect to L_0 , i.e., $L_0v = nv$ for all $v \in V_n$. In this paper we always suppose that $V_0 = \mathbf{C} \cdot \mathbf{1}$, and then V is called of CFT type. Let V be a VOA with central charge c_V . A weak V-module is a pair (M, Y_M) of a vector space M and a linear map $Y_M : V \to \text{End}_{\mathbf{C}}(M)[[z, z^{-1}]]$ with required natural conditions as a module on V. A weak V-module M is also a module over the Virasoro algebra with the same central charge $c_V \in \mathbb{C}$ by letting $Y_M(\omega, z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$. (For more details, see §3 in [\[14\]](#page-16-4) and §4 in [\[16\]](#page-16-5).)

A weak V-module is said to be a V-module if there are finitely many complex numbers h_i such that $M = \bigoplus_i \bigoplus_{n=0}^{\infty} M_{h_i+n}$, where M_h is a *finite-dimensional* eigenspace of L_0 with an eigenvalue h . If M is a simple V -module (in the obvious meaning), then there exists a complex number h such that $M = \bigoplus_{n=0}^{\infty} M_{h+n}$ with $M_h \neq 0$. This complex number h is called the *conformal*

weight of the simple V-module M . Then the (formal) *character* of the simple V-module M is defined by

$$
\operatorname{ch}_M(\tau) = \operatorname{tr}_M q^{L_0 - \frac{c_V}{24}} = q^{h - \frac{c_V}{24}} \sum_{n=0}^{\infty} (\dim M_{h+n}) q^n \quad (q = e^{2\pi i \tau}).
$$

Definition. (a) We say that a vertex operator algebra V is *rational* if any V-module is completely reducible, and is of finite type if the codimension of the subspace of V , which is spanned by elements $a_{(-2)} b$ for all $a, b \in V$, is finite. It is known that conformal weights and central charge of such a V are rational numbers. (cf. [\[2\]](#page-15-2)).

(b) We denote by \mathfrak{ch}_V the space spanned by characters of all (inequivalent) simple V-modules.

It was shown in Theorem 4.4.1 of [\[21\]](#page-16-6) that if V is of finite type, then the number of simple V modules is finite (up to isomorphism), and the character ch_M of any simple V-module M converges in the complex upper half-plane \mathfrak{H} . Moreover, if rational VOA V is of CFT and of finite type, then the space \mathfrak{ch}_V is invariant under the slash 0 action |0 of $SL_2(\mathbf{Z})$. This fact is now called the modular invariance property of the space \mathfrak{ch}_V .

3 Minimal models

The minimal models is one of the important examples of rational simple VOAs of CFT and finite type. The Virasoro algebra denoted here by Vir is a Lie algebra spanned by L_n ($n \in \mathbb{Z}$) and a central element C with commutation relations

$$
[L_m, L_n] = (m - n)L_{m+n} + \frac{m^3 - m}{12} \delta_{m, -n} C \quad (m, n \in \mathbf{Z}).
$$

Let $\text{Vir}_{\geq 0} = \bigoplus_{n=0}^{\infty} \mathbf{C} L_n \oplus \mathbf{C} C$, and let c and h be complex numbers. We denote by $\mathbf{C} v_{c,h}$ the one-dimensional $U(\text{Vir}_{\geq 0})$ -module that is defined by $L_n v_{c,h} = h \, \delta_{n,0} v_{c,h}$ and $C v_{c,h} = c v_{c,h}$, where $U(Vir)$ is the universal enveloping algebra of the Lie algebra Vir. Then the induced module $M(c, h) = U(\text{Vir}) \otimes_{U(\text{Vir}_{\geq 0})} \mathbf{C} v_{c, h}$ is called the *Verma module* of the Virasoro algebra with central charge c and the highest weight h . If Verma module has the *unique* maximal proper submodule $J(c, h)$, then we denote its simple quotient by $L(c, h) = M(c, h)/J(c, h)$. Let $\langle L_{-1} v_c, 0 \rangle$ be a left Vir-submodule of $M(c, 0)$ which is generated by $L_{-1}v_{c, 0}$. One can verify that the space $\langle L_{-1} v_{c, 0} \rangle$ is a proper submodule of $M(c, 0)$. We denote by $V(c, 0)$ the quotient Virmodule $M(c, 0)/\langle L_{-1} v_{c, 0} \rangle$, and by 1 the image of $v_{c, 0}$ under the natural surjection. It was proved in Theorem 4.3 (p. 162) and Remark (p. 163) of [\[8\]](#page-16-7) that $V(c, 0)$ and $L(c, 0)$ are VOAs with the vacuum vector 1 and the Virasoro vector $\omega = L_{-2}$, respectively. Then by applying the Poincaré-Birkhoff-Witt theorem to $U(Vir)$ one can find that the space $V(c, 0)$ has a basis which consists of 1 and $L_{-n_1} \cdots L_{-n_r}$ 1 $(r > 0, n_i \in \mathbb{Z}, n_1 \geq \cdots \geq n_r > 1)$, whose weights induced by L_0 are 0 and $\sum_{i=1}^{r} n_i$, respectively. Therefore the character of Vir-module $V(c, 0)$ is given by

$$
\mathrm{ch}_{V(c,h)}(\tau) = q^{-c/24} \prod_{n>1} (1-q^n)^{-1} = q^{-c/24} \left(1+q^2+q^3+2q^4+\cdots\right).
$$

A nonzero vector v of a (highest weight) Vir-module is called *singular* if $L_n v = 0$ for all positive integers n. If $V(c, 0)$ contains a singular vector v with a *positive weight*, then $V(c, 0)$ is not simple since $U(\text{Vir})v$ is a proper vertex operator subalgebra of V. Using the Kac determinant formula (cf. [\[13\]](#page-16-8)), it was shown in [\[7\]](#page-15-3) and [\[6\]](#page-15-4) that $V(c, 0)$ contains a (positive weight) singular vector if and only if its central charge is given by

$$
c = c_{P,Q} = 1 - \frac{6(P - Q)^2}{PQ} \tag{2}
$$

with coprime positive integers P and Q. It was moreover proved in [\[6\]](#page-15-4), [\[7\]](#page-15-3) and in Theorem 6.5 in [\[11\]](#page-16-9) that the maximal proper submodule $J(c_{P,Q}, 0)$ of $M(c_{P,Q}, 0)$ is generated by two singular vectors, $L_{-1}v_{c_{P,Q}}$ and $w_{P,Q}$, where the latter has weight $(P-1)(Q-1)$. We denote by $L(c_{P,Q}, 0)$ the simple Virasoro VOA $M(c_{P,Q}, 0)/J(c_{P,Q}, 0)$ which is called the *minimal model* of type (P,Q) . Let P and Q be coprime integers and define rational numbers by

$$
h_{P,Q\,;\,r,\,s} = \frac{(rQ - sP)^2 - (P - Q)^2}{4PQ}.
$$
\n⁽³⁾

Notations. We denote $L(c_{P,Q}, 0)$ and $L(c_{P,Q}, h_{P,Q}; r, s)$ by $L_{P,Q}$ and $L_{P,Q}; r, s$, respectively. In particular, one can see $L_{P,Q} = L_{P,Q+1,1}$.

The following facts are well known (cf. Theorem 6.13 in [\[11\]](#page-16-9)).

Theorem 4. Let P and Q be coprime integers with $0 < P < Q$.

(a) The vertex operator algebra $L_{P,Q}$ is rational, simple and is of CFT and finite type.

(b) For any simple $L_{P,Q}$ -module M there exist integers $0 < r < P$ and $0 < s < Q$ such that M is isomorphic to $L_{P,Q;\,r,s}.$

(c) The $L_{P,Q}$ -modules $L(c_{P,Q}, h_1)$ and $L(c_{P,Q}, h_2)$ are isomorphic if and only if $h_1 = h_2$.

(d) The rational number $h_{P,Q; r,s}$ is the conformal weight of the $L_{P,Q}$ -module $L_{P,Q; r,s}$.

Furthermore, it is known that the character ch $P, Q; r, s$ is written in the form

$$
\text{ch}_{P,Q\,;\,r,\,s}(\tau) \,=\, \frac{1}{\eta(\tau)} \big(\theta_{PQ,\,Qr-Ps}(\tau) - \theta_{PQ,\,Qr+Ps}(\tau)\big)\,,\quad \theta_{a,\,b}(\tau) \,=\, \sum_{n\in\mathbf{Z}} q^{a(n+b/2a)^2} \tag{4}
$$

for any half-integers a and b . (See Theorem 6.13 of [\[11\]](#page-16-9).) This set of characters coincides with the solution space of some monic linear differential equation and of order $\frac{(P-1)(Q-1)}{2}$. (See Theorem 6.1) of [\[17\]](#page-16-1), Theorem 6.1 of [\[18\]](#page-16-10) and Theorem 2.1 of [\[19\]](#page-16-11))

Let P and Q are coprime positive integers greater than one. Let r and s be integers with $1 \leq$ $r \leq P-1$ and $1 \leq s \leq Q-1$. Then central charge and conformal weights of the minimal model of type (P,Q) can be expressed as

$$
c = c_{P,Q} = 1 - \frac{6(P-Q)^2}{PQ},
$$
 $h_{r,s} = \frac{(rQ-Ps)^2 - (P-Q)^2}{4PQ}.$

We now compute the *effective central charge* and the *the minimal conformal weight* of the minimal model of type (P,Q) .

Lemma 5. Let P and Q be coprime positive integers greater than one, and let r and s be integers with $1 \leq r \leq P-1$ and $1 \leq s \leq Q-1$. Then the minimum $h_{\min}^{P,Q}$ of the set of all conformal weights $\{h_{r,s}^{P,Q}\}$ and effective central charge $c_{\text{eff}}^{P,Q}$ are given by

$$
h_{\min}^{P,Q} = \frac{1 - (P - Q)^2}{4PQ} \quad \text{and} \quad c_{\text{eff}}^{P,Q} = \frac{PQ - 6}{PQ}.
$$
 (5)

Proof. Since P and Q are coprime, we have $r_1P - Qs_1 = 1$ for some integers r_1 and s_1 . Then any solution $(r, s) \in \mathbb{Z}^2$ of the Diophantine equation $Pr - Qs = 1$ is written as $r = r_1 + Qk$, $s = s_1 + Pk$ with an integer k. Obviously $(r_1, P) = 1$ and $(s_1, Q) = 1$, and then we can take r with $1 \le r \le P-1$ for some integer k since $(r, P) = 1$ and $P > 1$. Hence we have $Q - 1 \leq rQ - Q = Ps \leq Q(P - 1) - 1$ and $0 < (Q-1)/P \le s \le Q-(Q+1)/Q \le Q-1$. Thus we have found r and s so that $Qr-Ps=1$ and $1 \leq r \leq P-1$ and $1 \leq s \leq Q-1$. Therefore the minimum value of $h_{r,s}^{P,Q}$ is given by the first of equation [\(5\)](#page-5-0). Then the formula of the effective central charge immediately follows by the definition. □ We now obtain

Corollary. Let c_{eff} and h_{min} be the effective central charge and the minimal conformal weight of the minimal model of type $(2, p)$. Then we have

$$
h_{\min} = \frac{1 - (2 - p)^2}{8p} \quad and \quad c_{\text{eff}} = 1 - \frac{3}{p},
$$

where $h_{\min} = h_{\min}^{2,p}$ and $c_{\text{eff}} = c_{\text{eff}}^{2,p}$, respectively.

4 Minimal models and Ibukiyama modular forms

Now, we can relate the Ibukiyama modular forms to characters of the minimal models of type $(2, p)$. We show that any character of the minimal model of type $(2, p)$ multiplied by η^{c} is equal to an Ibukiyama modular form up to a multiple constant.

Theorem 6. Let $L_{2,p}$ be the minimal model of type $(2, p)$ with an odd integer $p (> 3)$. Then we have

$$
\eta(\tau)^{c_{\rm eff}} \ch_{2,\,p;\,1,\,s}(\tau) \,=\, e\left(\frac{2ps-1+p-p^2}{8p}\right) f_{p-2s}(\tau) \,,\quad c_{\rm eff} \,=\, 1-\frac{3}{p} \,. \tag{6}
$$

 \Box

The set of ${p-2s \mid 0 < s < p/2}$ equals the set ${1 \le r \le p-2 \mid r \text{ is odd}}$, and where f_r (the index r is any element in latter set) is the Ibukiyama modular form given in Theorem [3](#page-3-2).

Proof. Using equation [\(4\)](#page-5-1) one can compute the left-hand side of the expected identity [\(6\)](#page-6-2) as

$$
\eta(\tau)^{\text{c}}^{\text{ref}} \text{ ch}_{2, p; 1, s}(\tau) = \eta(\tau)^{-3/p} \sum_{n \in \mathbb{Z}} \left[e(\tau)^{\frac{p}{2} \left(\frac{2n}{2} + \frac{p-2s}{2p} \right)^2} - e(\tau)^{\frac{p}{2} \left(\frac{2n-1}{2} + \frac{p-2s}{2p} \right)^2} \right]
$$
\n
$$
= \eta(\tau)^{-3/p} \sum_{n \in \mathbb{Z}} (-1)^n e(\tau)^{\frac{p}{2} \left(n + \frac{p-2s}{2p} \right)^2} = e\left(\frac{2s-p}{4p} \right) \eta(\tau)^{-3/p} \Theta_{\left(\frac{p-2s}{2p}, \frac{1}{2} \right)}(p\tau, 0)
$$
\n
$$
= e\left(\frac{2ps - 1 + p - p^2}{4p} \right) f_{p-2s}(\tau),
$$

and hence equation [\(6\)](#page-6-2) holds.

We now give a different proof of Theorem [3](#page-3-2) (b), i.e. Theorem 1.2 in [\[10\]](#page-16-0). Using Proposition 6.3 in [\[11\]](#page-16-9) one can see that the action of $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ on the character of $L_{2, p; 1, s}$ is given by

$$
\operatorname{ch}_{2, p; 1, s}(-1/\tau) = \frac{2}{\sqrt{p}} \sum_{r=1}^{\frac{p-1}{2}} (-1)^{(1+r)(1+s)} \sin\left(\frac{\pi}{2}(2-p)\right) \sin\left(\frac{\pi s r}{p}(2-p)\right) \operatorname{ch}_{2, p; 1, r}(\tau).
$$

Therefore we have

$$
\pi_p(S)f_{p-2s} = e\left(\frac{3p-1}{8}\right)\frac{2}{\sqrt{p}}\sum_{r=1}^{\frac{p-1}{2}}\cos\left(2\pi\left(\frac{p}{4}+\frac{rs}{p}-s\right)\right)f_{p-2r}
$$

\n
$$
= e\left(\frac{3p-1}{8}\right)\frac{1}{\sqrt{p}}\sum_{r=1}^{\frac{p-1}{2}}\left[e\left(-s+\frac{p}{4}+\frac{rs}{p}\right)+e\left(s-\frac{p}{4}-\frac{rs}{p}\right)\right]f_{p-2r}
$$

\n
$$
= e\left(\frac{3p-1}{8}\right)\frac{1}{\sqrt{p}}\sum_{r=1}^{\frac{p-1}{2}}\left[e\left(\frac{(p-2s)-(p-2r)}{4}+\frac{(p-2s)(p-2r)}{4p}\right)\right]
$$

\n
$$
+ e\left(-\frac{(p-2s)-(p-2r)}{4}-\frac{(p-2s)(p-2r)}{4p}\right)f_{p-2r}.
$$

Hence it follows for any odd integer s with $1 \leq s \leq p-2$ that

$$
\pi_p(S)f_s = e\left(\frac{3p-1}{8}\right) \frac{1}{\sqrt{p}} \sum_{\substack{1 \le r \le p-2, \\ r \equiv 1 \bmod 2}} \left[e\left(\frac{s-r}{4} + \frac{sr}{4p}\right) + e\left(-\frac{s-r}{4} - \frac{sr}{4p}\right) \right] f_r
$$

=
$$
\frac{1}{\sqrt{p}} \sum_{\substack{1 \le r \le p-2, \\ r \equiv 1 \bmod 2}} \left[e\left(\frac{s-r}{4} + \frac{sr}{4p} + \frac{3p-1}{8}\right) + (-1)^{\frac{p+1}{2}} e\left(-\frac{s-r}{4} - \frac{sr}{4p} - \frac{3p-1}{8}\right) \right] f_r.
$$

From the definition of π_p we obtain $\pi_p(T) f_s = e \left(\frac{1-p^2}{8n} \right)$ 8p $\left(e^{\frac{s^2-1}{8p}}\right) f_s(\tau) = e^{\frac{s^2-p^2}{8p}}$ $\frac{-p^2}{8p}$) $f_s(\tau)$. Therefore we have proved Theorem [3](#page-3-2) (b).

The characters of a VOA are not always modular functions on $SL_2(\mathbf{Z})$. However, it was proved by Dong, Lin and Ng that they are modular functions on some principal congruence subgroups of $SL_2(\mathbf{Z})$.

Theorem 7 (Theorem I and Theorem 3.10 in [\[4\]](#page-15-0)). Let V be a rational simple self-dual vertex operator algebra of CFT and of finite type, and let $\{M^i\}$ be a complete list of simple V-modules up to isomorphism. Then each character χ_{M^i} is a modular function on the principal congruence subgroup of $SL_2(\mathbf{Z})$ of level n that is the smallest positive integer such that $n(\lambda_i - c/24)$ is an integer for all i, where λ_i is the conformal weight of M^i .

Let m be a positive integer with m|n. The character χ_{M} is a modular function on $\Gamma(n)$ but is not always modular on the group $\Gamma(m)$. We denote n in Theorem [7](#page-7-2) by n_p for the minimal models of type $(2, p)$. For instance, we have $n_5 = 60$, however, the Ibukiyama modular form (given by [\(6\)](#page-6-2)) is defined on $\Gamma(5)$. Therefore the $\chi_{M,i} (= \mathrm{ch}_{M,i})$ must be a modular function with a certain multiplier system on $\Gamma(5)$.

We now give a formula of n_p of the minimal model of type $(2, p)$. Any character of minimal models of type $(2, p)$ with an odd integer $p > 3$ has an q-expansion

$$
\frac{1}{\eta(q)} \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{p}{2}\left(n + \frac{p-2s}{2p}\right)^2} = q^{\frac{(p-2s)^2}{8p} - \frac{1}{24}} + O\left(q^{\frac{(p-2s)^2}{8p} - \frac{1}{24} + 1}\right)
$$
(7)

for $s = 1, 2, \ldots, (p-1)/2$. We will find the smallest positive integer n_p such that $n\left(\frac{(p-2s)^2}{8p}-\frac{1}{24}\right)$ $n\left(\frac{3(p-2s)^2-p}{24n}\right)$ $\frac{(2s)^2-p}{24p}$ is an integer for all s. We now let $p=2r+3$ with $r>0$. Then we have

Theorem 8. Let $p = 2r + 3$ with a positive integer r. Then the minimal denominator of the irreducible fraction of $\frac{3(p-2s)^2-p}{24p}$ $\frac{(-2s)^2 - p}{24p}$ for any $0 \le s \le (p-1)/2$ is equal to

$$
\frac{12(2r+3)}{\gcd(4,r)\gcd(3,r)}.
$$
\n
$$
(8)
$$

Proof. Let $r = 12m + \ell$, where $0 \leq \ell \leq 11$ and m is a nonnegative integer. We will give the proof in the case $\ell = 0$. The cases $1 \leq \ell \leq 11$ are described in the appendix and they are very similar to the case of $\ell = 0$.

Now, on the one hand $\frac{12(2r+3)}{\gcd(4,r)\gcd(3,r)} = 24m+3$, on the other hand,

$$
\frac{3(p-2s)^2-p}{24p}\,=\,\frac{3(24m+3-2s)^2-24m-3}{24(24m+3)}\,=\,\frac{(24m+3-2s)^2-8m-1}{8(24m+3)}\,.
$$

Since $(24m + 3 - 2s)^2 - 8m - 1 \equiv 0 \pmod{8}$, we see that the denominator of irreducible fraction $\frac{3(p-2s)^2-p}{24p}$ $\frac{22S(1-p)}{24p}$ is equal to $24m+3$. Other cases for $\ell=1,\ldots,11$ are proved in the same way as mentioned in the appendix.

 \Box

ℓ	0	1	2	3	4	5	6	7	8	9	10	11
n	p	12p	6p	4p	12p	12p	4p	12p	12p	4p	6p	12p

Table 1: Table of n_p

Corollary. Let n_p be the integer in Theorem [7](#page-7-2) of the minimal model of type $(2, p)$. Then it is equal to $\frac{12(2r+3)}{\gcd(4,r)\gcd(3,r)}$.

Let $p = 2r + 3$ with a positive integer r. The following Table [4](#page-7-0) is a part of the list of the denominator of irreducible fraction of $\frac{12(2r+3)}{\gcd(4,r)\gcd(3,r)}$ for $r \equiv \ell \pmod{12}$.

5 Modular linear differential equations and vector-valued modular forms

In this section we recall the basic fact that the space spanned by characters of all simple modules over a minimal model equals the solution space of some monic MLDE whose definition is given below. Let k be a positive integer and let E_{2k} be the normalized Eisenstein series of weight $2k$, which is defined by $E_{2k}(\tau) = 1 - (4k/B_{2k}) \sum_{n=1}^{\infty} \sigma_{2k-1}(n) q^n$ and $q = e^{2\pi i \tau}$ ($\tau \in \mathfrak{H}$), where $\sigma_j(n)$ for $n \in \mathbb{N}$ is the sum of the jth powers of the positive divisors of n, and where B_n is the nth Bernoulli number.

Let \mathfrak{d} : Hol(5) \to Hol(5) be an *operator* that is defined by $\mathfrak{d}_k(f) = D(f) - \frac{k}{12}E_2(f)$ for each rational number k, where $D = (2\pi i)^{-1}d/d\tau$ for any $f \in M_k(SL_2(\mathbf{Z}))$, which is called the *Serre* operator. It is easy to show that $\mathfrak{d}: M_*(SL_2(\mathbf{Z})) \to M_{*+2}(SL_2(\mathbf{Z}))$. The *i*th *iterated Serre operator* is defined by $\mathfrak{d}_k^i = \mathfrak{d}_{k+2(i-1)} \circ \cdots \circ \mathfrak{d}_{k+2} \circ \mathfrak{d}_k$ with $\mathfrak{d}_k^0 = 1$, any differential equation of the *n*th order is written as

$$
\mathfrak{d}_k^n(f) + \sum_{i=0}^{n-1} P_i \mathfrak{d}_k^i(f) = 0
$$
\n(9)

is called a *monic modular linear differential equation* (monic MLDE) of weight k if P_i is a holomorphic modular form of weight $2(n-i)$ for any i. Since any holomorphic modular form of weight two in $\mathfrak H$ is 0, the coefficient P_{n-1} of [\(9\)](#page-8-1) is always zero. It was shown by Mason (Theorem 4.1) in [\[15\]](#page-16-12)) that the solution space of any monic MLDE of weight k is closed under the usual slash k action $|_k$ of $SL_2(\mathbf{Z})$.

Definition. (a) Let k be a rational number. A vector-valued modular form (VVMF) of weight k on $SL(2, \mathbb{Z})$ with a multiplier system $v : SL(2, \mathbb{Z}) \to \mathbb{C}$ with $v(-I) = (-1)^k$ is a column vectorvalued holomorphic function $F = {}^t (f_1, f_2, \ldots, f_n)$ in the upper half-plane $\mathfrak H$ equipped with an ndimensional representation $\pi : SL(2, \mathbf{Z}) \to GL_n(\mathbf{C})$ that is given by $j(\gamma, \tau)^{-1} F(\gamma(\tau)) = \pi(\gamma)F(\tau)$ for all $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ with $j(\gamma, \tau) = v(\gamma)(c\tau + d)^k$, and each component function f_j has a Fourier expansion $f_j = q^{\lambda_j} \left(\sum_{i=0}^{\infty} a_i^j q^i \right)$ in 5, with $a_0^j \neq 0$ and $\lambda_j \in \mathbf{Q}$.

(b) A vector-valued modular form F is called *normalized* if $\lambda_1 > \lambda_2 > \cdots > \lambda_r$ and $f_j = 0$ $(r+1 \leq$ $j \leq n$). If the component functions of F are linearly independent, then there is an invertible matrix A such that AF is normalized. In particular, if $\lambda_1, \ldots, \lambda_n$ are mutually distinct, then they are automatically normalized after consecutive interchanges of indices. The VVMF F is called strictly normalized when $r = n$.

We have the following fact which was also proved by G. Mason (Theorem 4.3 in [\[15\]](#page-16-12)). Very recently we have proved a similar result for the theta gamma group in [\[1\]](#page-15-5).

Proposition 9. The space spanned by all component functions $\{f_j\}_{j=1}^n$ of a strictly normalized vector-valued modular form of weight k is equal to the solution space of some monic modular linear differential equation of the nth order if and only if $n(n+k-1) - 12\sum_{i=1}^{n} \lambda_i = 0$, where each λ_j $(1 \leq j \leq n)$ is the leading exponent of f_j .

Since any monic MLDE has a unique regular singular point at $q = 0$ one can use the Frobenius method described in Chapter XVI in [\[12\]](#page-16-13) to obtain a basis of the solution space of a monic MLDE. Suppose that $f = q^{\lambda} (1 + \sum_{i=0}^{\infty} a_i q^i)$ is a solution of equation [\(9\)](#page-8-1). Then, substituting f into (9) and taking the coefficient of the lowest power term, one can see that λ is a root of an algebraic equation (called an *indicial equation*) $\Psi(t) = 0$, where $\Psi(t) = \prod_{\ell=0}^{n-1} (t - \frac{\ell}{6}) + \sum_{i=0}^{n-2} P_{i,0} \prod_{\ell=0}^{i-1} (t - \frac{\ell}{6})$ and $P_i = \sum_{n=0}^{\infty} P_{i,n} q^n$. This polynomial is called the *indicial polynomial* of the monic MLDE [\(9\)](#page-8-1) and the roots of the indicial equation are called indicial roots.

Lemma 10. If any pair of two different indicial roots have a non-integral difference, then there exists a unique solution of the form $f = q^{\lambda} (1 + \sum_{n=1}^{\infty} a_n q^n)$, where λ is any indicial root and the coefficients a_n ($n \in \mathbb{Z}_{\geq 0}$) are uniquely determined from a_0 by some recursive relations and indicial roots. If there is a double incidicial root, then there always exists a logarithmic solution that is a polynomial of τ with coefficients in $q^{\lambda} \mathbf{C}((q))$ with a complex number λ .

It is important to know that the modular linear differential equation associated to a rational simple vertex operator algebra has no logarithmic solutions even if two different indicial roots have an integral difference.

6 Some properties of the Ibukiyama modular forms

We have established some relations between the Imukiyama modular forms and the characters of the minimal models of type $(2, p)$. It is therefore natural to ask whether we can obtain modular forms of fractional weights from the characters of any minimal model of type (P, Q) . The characters of the minimal model of type (P,Q) is a modular function on $\Gamma(n)$, where n is defined in Theorem [7.](#page-7-2) Of course, we can define fractional modular forms $\eta^{\text{c}_{\text{eff}}}$ ch_M on $\Gamma(n)$ with the multiplier system which is the same as that of η^{c} and then η^{c} ch_M is standard on $\Gamma(n)$. However, it is not clear to us whether η^{c} ^{eff} ch_M is standard on a smaller congruence group $\Gamma(m)$. Moreover, we want to know whether the following property holds for the Ibukiyama modular forms

(A) Let m and n be positive integers with m|n. Then $M_k(\Gamma(m), \nu_m) \subset M_k(\Gamma(n), \mu_n)$ for $k \in \mathbf{Q}$ with $\mu_n|\Gamma(m)=\nu_m$, where ν and μ are multiplier systems of $\Gamma(m)$ and $\Gamma(n)$.

We will give four examples of spaces of the Ibukiyama modular forms of fractional weights, and show that the set of pairs $(m, n) = (5, 15)$ and $(7, 21)$ satisfies (A). Since the space of Ibukiyama modular forms $M_{\frac{p-3}{2}}(\Gamma(p), v_p)$ is invariant by the $SL_2(\mathbf{Z})$ -action, it may be equal to the solution space of of some MLDE. In fact, we will give the corresponding monic MLDE. For any positive odd integer $p > 3$ we set

$$
f_s(\tau) := e\left(\frac{2ps - 1 + p - p^2}{4p}\right) f_{p-2s}
$$
 for $1 \le s \le \frac{p-1}{2}$.

First, it was shown in Lemma 1.7 of [\[10\]](#page-16-0) that $\dim M_{\frac{p-3}{2p}}(\Gamma(p), v_p) > 1$ with any odd integer $p > 3$. In particular, $\dim M_{\frac{p-3}{2p}}(\Gamma(p), v_p) = 2$ and 3 for $p = 5$ and 7, respectively. (See p. 325, lemma1.7) of Section 1.3 in [\[10\]](#page-16-0)) To show (A) for the pairs $(m, n) = (5, 15)$ and $(7, 21)$ it is, in principle, necessary to know basis of $M_{\frac{1}{5}}(\Gamma(5), v_5)$ and $M_{2/7}(\Gamma(7), v_7)$. Each basis of the space $M_{\frac{1}{5}}(\Gamma(5), v_5)$ and $M_{2/7}(\Gamma(7), v_7)$ is known (pp. 326–332, Section 2.1–2.3 in [\[10\]](#page-16-0)) (however, in the cases $p = 15, 21$ they are not known); the former basis consists of

$$
f_1^{[5]}(\tau) := \eta(\tau)^{2/5} ch_{2, 5; 1, 1}(\tau) = \eta(\tau)^{-3/5} \sum_{n \in \mathbb{Z}} (-1)^n e(\tau)^{\frac{5}{2} \left(n + \frac{3}{10}\right)^2}, \tag{10}
$$

$$
f_2^{[5]}(\tau) := \eta(\tau)^{2/5} ch_{2, 5; 1, 2}(\tau) = \eta(\tau)^{-3/5} \sum_{n \in \mathbb{Z}} (-1)^n e(\tau)^{\frac{5}{2}(n + \frac{1}{10})^2}, \tag{11}
$$

and the associated monic MLDE is $\mathfrak{d}^2_{1/5}(f) - \frac{11}{3600} E_4(f) = 0$. The latter has the effective central charge of 4/7. Then the Ibukiyama modular forms are $f_1^{[7]}(\tau) := \eta(\tau)^{4/7}$ ch_{2,7;1,1}(τ), $f_2^{[7]} :=$ $\eta(\tau)^{4/7}$ ch_{2,7;1,2}(τ) and $f_3^{[7]} := \eta(\tau)^{4/7}$ ch_{2,7;1,3}(τ). More precisely, they are written as $f_j^{[7]}$ $j^{[1]}(\tau) =$ $\eta(\tau)^{-3/7} \sum_{n \in \mathbf{Z}} (-1)^n e(\tau)^{\frac{7}{2}\left(n+\frac{7-2s}{14}\right)^2}$ for $1 \leq j \leq 3$, which form a basis of the space $M_{2/7}(\Gamma(7), v_7)$, and then the q-expansion of $f_j^{[7]}$ for $1 \le j \le 3$ is given by

$$
f_1^{[7]}(\tau) = q^{3/7} \left(1 - \frac{4}{7}q + \frac{15}{49}q^2 + \frac{43}{343}q^3 + \frac{965}{2401}q^4 + \frac{1524}{16807}q^5 - \frac{39026}{117649}q^6 - \frac{1874288}{5764801}q^7 + \cdots \right),
$$

\n
$$
f_2^{[7]}(\tau) = q^{1/7} \left(1 + \frac{3}{7}q - \frac{13}{49}q^2 + \frac{148}{343}q^3 + \frac{1266}{2401}q^4 - \frac{8528}{16807}q^5 + \frac{38870}{117649}q^6 + \frac{213504}{5764801}q^7 + \cdots \right),
$$

\n
$$
f_3^{[7]}(\tau) = 1 + \frac{3}{7}q + \frac{36}{49}q^2 - \frac{48}{343}q^3 - \frac{400}{2401}q^4 + \frac{3183}{16807}q^5 + \frac{50140}{117649}q^6 + \frac{13535}{5764801}q^7 + \cdots
$$

\n(12)

The associated monic MLDE is $\mathfrak{d}^3_{2/7}(f) - \frac{5}{252} E_4 \mathfrak{d}_{2/7}(f) + \frac{85}{74088} E_6(f) = 0.$

We next treat fractional modular forms associated to the minimal model of type $(2, 15)$. Since the effective central charge of this case is 4/5 the Ibukiyama modular forms, which do not form a basis of $M(\Gamma(15), v_{15})$, are written as

$$
f_1^{[15]}(\tau) = \eta(\tau)^{4/5} ch_{2,15;1,1}(\tau), \quad f_2^{[15]}(\tau) = \eta(\tau)^{4/5} ch_{2,15;1,2}(\tau), \quad f_3^{[15]}(\tau) = \eta(\tau)^{4/5} ch_{2,15;1,3}(\tau),
$$

\n
$$
f_4^{[15]}(\tau) = \eta(\tau)^{4/5} ch_{2,15;1,4}(\tau), \quad f_5^{[15]}(\tau) = \eta(\tau)^{4/5} ch_{2,15;1,5}(\tau), \quad f_6^{[15]}(\tau) = \eta(\tau)^{4/5} ch_{2,15;1,6}(\tau),
$$

\n
$$
f_7^{[15]}(\tau) = \eta(\tau)^{4/5} ch_{2,15;1,7}(\tau).
$$

It is not difficult to see that $f_j^{[15]}(\tau) = \eta(\tau)^{-1/5} \sum_{n \in \mathbb{Z}} (-1)^n e(\tau)^{\frac{15}{2}(n+\frac{2j-1}{30})^2}$ for any $1 \le j \le 7$ and their Fourier expansions are given by

$$
f_1^{[15]}(\tau) = q^{7/5} \left(1 - \frac{4}{5}q + \frac{3}{25}q^2 + \frac{1}{125}q^3 + \frac{79}{625}q^4 - \frac{776}{15625}q^5 + \frac{16609}{78125}q^6 + \cdots \right),
$$

\n
$$
f_2^{[15]}(\tau) = q \left(1 + \frac{1}{5}q - \frac{17}{25}q^2 + \frac{16}{125}q^3 + \frac{84}{625}q^4 + \frac{1199}{15625}q^5 + \cdots \right),
$$

\n
$$
f_3^{[15]}(\tau) = q^{2/3} \left(1 + \frac{1}{5}q + \frac{8}{25}q^2 - \frac{84}{125}q^3 + \frac{159}{625}q^4 + \frac{1324}{15625}q^5 + \frac{22604}{78125}q^6 + \cdots \right),
$$

\n
$$
f_4^{[15]}(\tau) = q^{2/5} \left(1 + \frac{1}{5}q + \frac{8}{25}q^2 + \frac{41}{125}q^3 - \frac{341}{625}q^4 + \frac{3199}{15625}q^5 + \frac{23229}{78125}q^6 + \cdots \right),
$$

\n
$$
f_5^{[15]}(\tau) = q^{1/5} \left(1 + \frac{1}{5}q + \frac{8}{25}q^2 + \frac{41}{125}q^3 + \frac{284}{625}q^4 - \frac{9301}{15625}q^5 + \frac{32604}{78125}q^6 + \cdots \right),
$$

\n
$$
f_6^{[15]}(\tau) = q^{1/15} \left(1 + \frac{1}{5}q + \frac{8}{25}q^2 + \frac{41}{125}q^3 + \frac{284}{625}q^4 + \frac{6324}{15625}q^5 - \frac{29896}{78125}q^6 + \cdots \right),
$$

\n
$$
f_7^{[15]}(\tau)
$$

The set $\{f_j^{[15]}\}\$ spans a *subspace* of $M_{2/5}(\Gamma(15), v_{15})$, and it is not difficult by using [\(12\)](#page-10-0) and [\(13\)](#page-10-1)

and the fact that the Strum bound of $\Gamma(15)$ is 1440 to find that

$$
f_2^{[5]}(\tau)^2 = f_4^{[15]}(\tau) - f_7^{[15]}(\tau) \,, \quad f_2^{[5]}(\tau) f_1^{[5]}(\tau) = f_3^{[15]}(\tau) \,, \quad f_1^{[5]}(\tau)^2 = f_1^{[15]}(\tau) + f_6^{[15]}(\tau) \,.
$$

(cf. p. 8, Lemma 3 in [\[9\]](#page-16-2).), where the left-hand sides of the above equations are basis of $M_{2/5}(\Gamma(5), v_5^2)$. We now prove $v_5^2 = v_{15}$ since both hand sides of [\(6\)](#page-10-1) must have the same weights. Since for any integer p (> 3) there are multiplier systems v_0 corresponding $p = 5, 15$ which we denote it by $v_{0,5}$ and $v_{0,15}$ by $v_{0,5}(\gamma) = \frac{\eta^{3/5}(\gamma \tau)}{(c\tau+d)^{3/10}n^{3/5}}$ $\frac{\eta^{3/5}(\gamma\tau)}{(c\tau+d)^{3/10}\eta^{3/5}(\tau)}$ for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(5)$ and $v_{0,15}(\gamma) = \frac{\eta^{1/5}(\gamma\tau)}{(c\tau+d)^{1/10}\eta^{1/5}}$ $(c\tau+d)^{1/10}\eta^{1/5}(\tau)$ for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(15)$, respectively. Then we immediately see $v_{0,5}(\gamma) = v_{0,15}(\gamma)^3$ for any $\gamma \in \Gamma(15)$. It also holds that $v_{0,15}^{40} = 1$ for any $\gamma \in \Gamma(15)$ since $\eta(\tau)^8 = (\eta(\tau)^{1/5})^{40}$ is a modular form of weight 4 on $\Gamma(3)$. Thus for any $\gamma \in \Gamma(15)$ we can see that $v_p = v_0^{p^2-1}$ weight of v_{15} is $2/5$.

$$
\frac{v_{15}(\gamma)}{v_5(\gamma)^2} = \frac{v_{0,15}(\gamma)^{15^2 - 1}}{v_{0,5}(\gamma)^{2(5^2 - 1)}} = \frac{v_{0,15}(\gamma)^{15^2 - 1}}{v_{0,15}(\gamma)^{6(5^2 - 1)}} = v_{0,15}(\gamma)^{80} = (v_{0,15}(\gamma)^{40})^2 = 1.
$$

Therefore one can conclude $M_{2m/5}(\Gamma(5), v_5^{2m}) \subset M_{2m/5}(\Gamma(15), v_{15}^m)$ for any positive integer m which implies (A). The monic MLDE of the 7th order whose solution space consists of the Ibukiyama modular forms of weight $2/5$ on $\Gamma(15)$; any function of [\(13\)](#page-10-1) is a solution of the MLDE given by

$$
\mathfrak{d}_{2/5}^7(\mathsf{f})-\frac{21}{50}E_4\mathfrak{d}_{2/5}^5(\mathsf{f})+\frac{529}{2700}E_6\mathfrak{d}_{2/5}^4(\mathsf{f})-\frac{3283}{90000}E_8\mathfrak{d}_{2/5}^3(\mathsf{f})+\frac{7733}{2430000}E_{10}\mathfrak{d}_{2/5}^2(\mathsf{f})\\-\frac{248501}{729000000}\Big(E_4^3-\frac{31233600}{22591}\Delta\Big)\mathfrak{d}_{2/5}(\mathsf{f})+\frac{248501}{4374000000}E_{14}(\mathsf{f})=0\,.
$$

Finally, we discuss modular forms of weight $3/7$ on $\Gamma(21)$, i.e. fractional modular forms associated to the minimal model of type $(2, 21)$. Since the effective central charge is $6/7$ the Ibukiyama modular forms are given by

$$
f_1^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,1}(\tau), \quad f_2^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,2}(\tau), \quad f_3^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,3}(\tau),
$$

\n
$$
f_4^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,4}(\tau), \quad f_5^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,5}(\tau), \quad f_6^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,6}(\tau),
$$

\n
$$
f_7^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,7}(\tau), \quad f_8^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,8}(\tau), \quad f_9^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,9}(\tau),
$$

\n
$$
f_{10}^{[21]}(\tau) = \eta(\tau)^{6/7} ch_{2,21;1,10}(\tau).
$$

It is not difficult to see that $f_i^{[21]}$ $j_j^{[21]}(\tau) = \eta(\tau)^{-3/21} \sum_{n \in \mathbf{Z}} (-1)^n e(\tau)^{\frac{21}{2}(n+\frac{2j-1}{42})^2}$ for any $1 \le j \le 10$, and then their Fourier expansions are given by

$$
f_{1}^{[21]}(\tau) = q^{15/7} \left((1 - \frac{6}{7}q + \frac{4}{49}q^{2} - \frac{1}{343}q^{3} + \frac{194}{2401}q^{4} - \frac{825}{16807}q^{5} + \frac{16351}{117649}q^{6} + \cdots \right),
$$

\n
$$
f_{2}^{[21]}(\tau) = q^{12/7} \left(1 + \frac{1}{7}q - \frac{38}{49}q^{2} + \frac{27}{343}q^{3} + \frac{187}{2401}q^{4} + \frac{533}{16807}q^{5} + \frac{10576}{117649}q^{6} + \cdots \right),
$$

\n
$$
f_{3}^{[21]}(\tau) = q^{4/3} \left(1 + \frac{1}{7}q + \frac{11}{49}q^{2} - \frac{267}{343}q^{3} + \frac{383}{2401}q^{4} + \frac{484}{16807}q^{5} + \frac{20082}{117649}q^{6} + \cdots \right),
$$

\n
$$
f_{4}^{[21]}(\tau) = q \left(1 + \frac{1}{7}q + \frac{11}{49}q^{2} + \frac{76}{343}q^{3} - \frac{1675}{2401}q^{4} + \frac{1856}{16807}q^{5} + \frac{19739}{117649}q^{6} + \cdots \right),
$$

\n
$$
f_{5}^{[21]}(\tau) = q^{5/7} \left(1 + \frac{1}{7}q + \frac{11}{49}q^{2} + \frac{76}{343}q^{3} + \frac{726}{2401}q^{4} - \frac{12550}{16807}q^{5} + \frac{29343}{117649}q^{6} + \cdots \right),
$$

\n
$$
f_{6}^{[21]}(\tau) = q^{10/21} \left(1 + \frac{1}{7}q + \frac{11}{49}q^{2} + \frac{76}{343}q^{3} + \frac{726}{24
$$

The monic MLDE such that each function of [\(14\)](#page-12-0) is a solution is given by

$$
\begin{aligned} \mathfrak{d}_{3/7}^{10}(\mathsf{f})-\frac{451}{336}E_{4}\mathfrak{d}_{3/7}^{8}(\mathsf{f})+\frac{39545}{37044}E_{6}\mathfrak{d}_{3/7}^{7}(\mathsf{f})-\frac{9757}{43904}E_{8}\mathfrak{d}_{3/7}^{6}(\mathsf{f})-\frac{4330919}{37340352}E_{10}\mathfrak{d}_{3/7}^{5}(\mathsf{f})\\ &+\frac{5130680923}{58549671936}\Big(E_{4}^{3}-\frac{3204764098560}{5130680923}\Delta\Big)\mathfrak{d}_{3/7}^{4}(\mathsf{f})-\frac{6729854203}{263473523712}E_{14}\mathfrak{d}_{3/7}^{3}(\mathsf{f})\\ &+\frac{1403994215923}{354108415868928}E_{4}\Big(E_{4}^{3}-\frac{1664226941276160}{1403994215923}\Delta\Big)\mathfrak{d}_{3/7}^{2}(\mathsf{f})\\ &-\frac{3323559970259}{13013484283183104}E_{6}\Big(E_{4}^{3}-\frac{1706576837437440}{3323559970259}\Delta\Big)\mathfrak{d}_{3/7}^{[1]}(\mathsf{f})\\ &-\frac{637460013}{42313823813632}E_{4}^{2}\Big(E_{4}^{3}-\frac{16388096}{12099}\Delta\Big)\mathsf{f}=0\,. \end{aligned}
$$

The set $\{f_j^{[21]}\}\$ spans a *subspace* of $M_{3/7}(\Gamma(21), v_{21})$, and it is not difficult to see that

$$
\begin{array}{ll} f_{3}^{[7]}(\tau)^{3} = f_{10}^{[21]}(\tau)^{2} + 2f_{10}^{[21]}(\tau)f_{4}^{[21]}(\tau) - f_{7}^{[21]}(\tau)f_{5}^{[21]}(\tau) + f_{4}^{[21]}(\tau)^{2} + f_{7}^{[21]}(\tau)f_{2}^{[21]}(\tau) \,,\\ f_{3}^{[7]}(\tau)^{2}f_{2}^{[7]}(\tau) = f_{10}^{[21]}(\tau)f_{8}^{[21]}(\tau) + f_{10}^{[21]}(\tau)f_{1}^{[21]}(\tau) + f_{8}^{[21]}(\tau)f_{4}^{[21]}(\tau) + f_{4}^{[21]}(\tau)f_{1}^{[21]}(\tau) \,,\\ f_{3}^{[7]}(\tau)^{2}f_{1}^{[7]}(\tau) = f_{8}^{[21]}(\tau)f_{7}^{[21]}(\tau) + f_{7}^{[21]}(\tau)f_{1}^{[21]}(\tau) \,,\\ f_{2}^{[7]}(\tau)^{3} = f_{8}^{[21]}(\tau)f_{7}^{[21]}(\tau) + f_{7}^{[21]}(\tau)f_{1}^{[21]}(\tau) + f_{5}^{[21]}(\tau)^{2} - 2f_{5}^{[21]}(\tau)f_{2}^{[21]}(\tau) + f_{2}^{[21]}(\tau)^{2} \,,\\ f_{2}^{[7]}(\tau)^{2}f_{3}^{[7]}(\tau) = f_{10}^{[21]}(\tau)^{2} + f_{7}^{[21]}(\tau)f_{4}^{[21]}(\tau) \,,\\ f_{2}^{[7]}(\tau)^{2}f_{1}^{[7]}(\tau) = f_{10}^{[21]}(\tau)f_{5}^{[21]}(\tau) - f_{10}^{[21]}(\tau)f_{2}^{[21]}(\tau) + f_{5}^{[21]}(\tau)f_{4}^{[21]}(\tau) - f_{4}^{[21]}(\tau)f_{2}^{[21]}(\tau) \,,\\ f_{1}^{[7]}(\tau)^{3} = f_{10}^{[21]}(\tau)f_{7}^{[21]}(\tau) - f_{8}^{[21]}(\tau)f_{2}^{[21]}(\tau) + f_{7}^{[21]}(\tau)f_{4}^{[21]}(\tau) - f_{1}^{[21]}(\tau)^{2} \,,\\ f_{
$$

by using the Strum bound of $\Gamma(21)$. We still need to prove that $v_7^3 = v_{21}^2$. As before, we define two maps $v_{0,5} : \Gamma(7) \to \mathbf{C}$ and $v_{0,15} : \Gamma(21) \to \mathbf{C}$ by $v_{0,7}(\gamma) = \frac{\eta^{3/7}(\gamma \tau)}{(c\tau+d)^{3/14}\eta^{3/2}}$ $\frac{\eta^{3/7}(\gamma\tau)}{(c\tau+d)^{3/14}\eta^{3/7}(\tau)}$ for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(7)$ and $v_{0,21}(\gamma) = \frac{\eta^{1/7}(\gamma\tau)}{(c\tau+d)^{1/14}\eta^{1/4}}$ $\frac{\eta^{1/7}(\gamma\tau)}{(c\tau+d)^{1/14}\eta^{1/7}(\tau)}$ for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(21)$, respectively. Then it is not difficult to see $v_{0,7}(\gamma)^2 = v_{0,21}(\gamma)^3$ for $\gamma \in \Gamma(21)$. We also obtain $(v_{0,21})^{56} = 1$ for any $\gamma \in \Gamma(21)$ since $\eta(\tau)^8 = (\eta^{1/7}(\tau))^{56}$ is a modular form of weight 4 on $\Gamma(3)$. Therefore we have

$$
\frac{v_{21}(\gamma)^2}{v_7(\gamma)^3} = \frac{v_{0,21}(\gamma)^{2(21^2-1)}}{v_{0,7}(\gamma)^{3(7^2-1)}} = \frac{v_{0,21}(\gamma)^{2(21^2-1)}}{v_{0,21}(\gamma)^{9(7^2-1)}} = v_{0,21}(\gamma)^{448} = (v_{0,15}(\gamma)^{56})^8 = 1.
$$

Hence one can see $M_{6m/7}(\Gamma(7), v_7^{3m}) \subseteq M_{6m/7}(\Gamma(21), v_{21}^{2m})$ for any positive integer m.

For odd integer $p > 13$ the set of the Ibukiyama modular forms is not always a basis of the space of modular forms of fractional weight $(p-3)/2p$ on $\Gamma(p)$ with multiplier systems v_p (cf. [\[9\]](#page-16-2)), and we do not know if (A) holds between $M_*(\Gamma(m), v_m)$ and $M_*(\Gamma(n), v_n)$ when $m, n > 13$. However, it was already seen that the set of pairs $p = 5, 15$ and $p = 7, 21$ satisfy (A). This suggests;

Conjecture. Let $\{\eta^{c_{\text{eff}}} \ch_{2, p; 1, s}\}$ and $\{\eta^{c'_{\text{eff}}} \ch_{2, p'; 1, s'}\}$ be the sets of modular forms on $\Gamma(p)$ and $\Gamma(p')$ with multiplier systems of the minimal models of type $(2, p)$ and of type $(2, p')$, respectively. Then the former and the latter have the property (A) if and only if $p|p'$.

We now show that each set of the Ibykiyama modular forms spans the solution space of some monic modular linear differential equation. We start with an easy lemma.

Lemma 11. Let $\Gamma \subset PSL_2(\mathbf{R})$ be a Fuchsian group. Suppose that $f \in M_k(\Gamma)$ is a solution of the monic modular linear differential equation $\mathfrak{d}_k^n(f) + \sum_{j=2}^n P_{2j} \mathfrak{d}_k^{n-j}(f) = 0$. Then $g = \eta^{\ell} f$ with an $\ell \in \mathbf{Z}$ is a solution of a monic linear differential equation $\mathfrak{d}^n_{k+\frac{\ell}{2}}(g) + \sum_{j=2}^n P_{2j} \mathfrak{d}^{n-j}_{k+\frac{\ell}{2}}$ $lim_{k+\frac{\ell}{2}}(g) = 0$ of weight $k + \frac{\ell}{2}$.

Proof. It follows from the definition of the iterated Serre operators and the equality $\mathfrak{d}_k(\eta)$ = $(1/24 - k/12)E_2\eta$ that $\mathfrak{d}_{k+\frac{\ell}{2}}^{n+1}(\eta^{\ell}f) = \mathfrak{d}_{k+\frac{\ell}{2}+2}^{n}(\mathfrak{d}_{k+\frac{\ell}{2}}(\eta^{\ell}f)) = \mathfrak{d}_{k+\frac{\ell}{2}+2}^{n}(\eta^{\ell}\mathfrak{d}_{k}(f))$ since $\mathfrak{d}_{k+\frac{\ell}{2}}^{n}(\eta^{\ell}f) =$ $\eta^{\ell} \mathfrak{d}^n_k(f)$. П

Theorem 12. Let $L_{2,p}$ be the minimal model of type $(2, p)$ with an odd integer $p > 3$, and let $\text{ch}_{2, p; 1, s}$ for $0 < s \leq (p-1)/2$ be the characters of the simple $L_{2, p}$ -module $L_{2, p; 1, s}$. (a) There exists a monic modular linear differential equation

$$
\mathfrak{d}_0^{\frac{p-1}{2}}(f) + \sum_{j=2}^{(p-1)/2} P_{2j} \mathfrak{d}_0^{\frac{p-1}{2}-j}(f) = 0 \tag{15}
$$

whose solution space is equal to the space spanned by the characters $\text{ch}_{2, p; 1, (s+1)/2}$ for all $0 < s \leq$ $(p-1)/2$, where P_{2j} is a holomorphic modular form of weight 2j on the full modular group. (b) The space of the Ibukiyama modular forms of weight $\frac{p-3}{2p}$ on $\Gamma(p)$ is equal to the solution space of some monic modular linear differential equation of weight $\frac{p-3}{2p}$ of the order $\frac{p-1}{2}$ on the full modular group, which is written is the form

$$
\mathfrak{d}\frac{\frac{p-1}{2}}{\frac{p-3}{2p}}(f) + \sum_{j=2}^{\frac{p-1}{2}} P_{2j} \mathfrak{d}\frac{\frac{p-3}{2p} - j}{\frac{p-3}{2p}}(f) = 0, \qquad (16)
$$

where each P_{2j} is the same modular form appearing in equation [\(15\)](#page-13-0).

Proof. The statement (a) follows from and Theorem [9.](#page-9-0) Lemma [11](#page-13-1) with $n = (p-1)/2$ and $k = 0$ shows the assertion (b). shows the assertion (b).

It must be mentioned that equation [\(15\)](#page-13-0) was first found in the form which uses $D = (2\pi i)^{-1}d/d\tau$ in Theorem 6.1 of [\[18\]](#page-16-10). Since equation [\(15\)](#page-13-0) and Milas' linear differential equation are monic and have the same solution space, they must coincide and then the latter is automatically monic MLDEs. We expected, as an application of the Theorem [12](#page-13-2) (b), that we can obtain supersingular j-invariant from the 0th order term of equation (16) as in [\[19\]](#page-16-11). However, this fails since 0th order terms of (16) and the Milas' one are equal.

7 Vector-valued modular forms associated to modular forms of fractional weights on $\Gamma(5)$

Let $f(\tau)$ be a (holomorphic or meromorphic) modular form of positive weight k on a Fuchsian group $\Gamma \subset PSL_2(\mathbf{R})$ and $t(\tau)$ a modular function with respect to Γ . Then the function $f(t)$ $f(t(\tau))$ satisfies a linear differential equation of order $k + 1$ with algebraic coefficients, or with polynomial coefficients if Γ\ \mathfrak{H} has genus 0 and $t(\tau)$ generates the field of modular functions on Γ.

Here we give examples of vector-valued modular forms of weight $k = m/5$ with $m \in \mathbb{Z}_{\geq 0}$ but of the minimal order of ordinary linear differential equation which has these modular forms as solutions with order $>$ weight $+1$. (See Proposition 21 p.61) of [\[20\]](#page-16-3)). The space of characters of the minimal model of type $(2, 5)$ multiplied by $\eta^{2/5}$ and the space of modular forms of weight $1/5$ with level five equal as we have seen in Theorem [6.](#page-6-1) We will take a symmetric tensor product representation of $SL_2(\mathbf{Z})$ -actin $\pi_5 : M_{1/5}(\Gamma(5), v_5) \to GL_2(\mathbf{C})$, which give a VVMF, and the set spanned by components function of the VVMF is equal to the solution space of some monic MLDE of order $1+5k$.

Let ${}^t(g_1, g_2) \in Hol(5) \times Hol(5)$. We denote $\pi_5(g_1, g_2)(\gamma) : SL_2(\mathbf{Z}) \to GL_2(\mathbf{C})$ a representation of $SL_2(\mathbf{Z})$ on $GL_2(\mathbf{C})$ defined by for any $\gamma \in SL_2(\mathbf{Z})$. Then one can show easily that (17)

$$
\pi_5^{\ t}\left(f_1^{[5]}, f_2^{[5]}\right)(T) = \begin{pmatrix} e\left(-\frac{2}{5}\right) & 0\\ 0 & e\left(-\frac{3}{5}\right) \end{pmatrix}, \ \pi_5^{\ t}\left(f_1^{[5]}, f_2^{[5]}\right)(S) = \frac{e\left(\frac{3}{4}\right)}{\sqrt{10}} \begin{pmatrix} -\sqrt{5} + \sqrt{5} & \sqrt{5} - \sqrt{5} \\ \sqrt{5} - \sqrt{5} & \sqrt{5} + \sqrt{5} \end{pmatrix}.
$$
\n(17)

It is proved in Lemma 1.7 in [\[10\]](#page-16-0) that dim $M_k(\Gamma(5)) = 1 + 5k$ for any $k \in \frac{1}{5}\mathbb{Z}_{>0}$. Let $m + 1 =$ $5k + 1$ (= dim $M_k(\Gamma(5))$), and define a VVMF $\mathsf{F} = {}^t(f_0, \ldots, f_m)$ with $f_\ell = (\mathsf{f}_2^{[5]})^{m-\ell}(\mathsf{f}_1^{[5]})^{\ell}$ for any $0 \leq \ell \leq m$. Then it is shown in [\[3\]](#page-15-6) that representation [\(17\)](#page-14-1) gives

$$
(\mathsf{F}|_{k}\gamma)(\tau) = A_{\gamma} \mathsf{F}(\tau) = \begin{pmatrix} a_0^{m+1} & (m+1)a_0^m b_0 & \cdots & b_0^{m+1} \\ a_0^m c_0 & \cdots & \cdots & b_0^m d_0 \\ \vdots & \vdots & \ddots & \vdots \\ c_0^{m+1} & (m+1)c_0^m d_0 & \cdots & d_0^{m+1} \end{pmatrix} \mathsf{F}(\tau) = \text{Sym}^{m+1} \, {}^t(\pi_5(\mathsf{f}_2^{[5]}, \mathsf{f}_1^{[5]})(\gamma)) \, \mathsf{F}(\tau)
$$

for any $\gamma \in SL_2(\mathbf{Z})$, where $\pi_5^{\{t\}}(f_1^{[5]}, f_2^{[5]})(\gamma) = \begin{pmatrix} a_0 & b_0 \\ c_0 & d_0 \end{pmatrix}$. More explicitly, the (m, n) entry of the matrix A_{γ} is given by $\sum_{i+j=m+1-n} {m+1-n \choose j} {n \choose i} a_0^j b_0^i c_0^i d_0^{n-i}$. Then the representation A_{γ} of $\gamma \in \Gamma(5)$ associated to F of $\Gamma(5)$ is the symmetric tensor representation $\text{Sym}^{m+1}(\pi_5(f_1^{[5]}, f_2^{[5]}))$ of degree $1+m$.

We now study the monic MLDE which is associated to the vector-valued modular form F. A short calculation shows $(f_2^{[5]})^{5k-\ell} (f_1^{[5]})^{\ell} = e(\tau)^{\ell/5} + O(e(\tau)^{(\ell+1)/5})$ for any $0 \leq \ell \leq 5k$. Since $\dim M_k(\Gamma(5), v_5^{1+5k}) = 1 + 5k$ for any $k \in \frac{1}{5}\mathbb{Z}_{\geq 0}$ and t is proved in Lemma 1.7 in [\[10\]](#page-16-0) that $\dim M_k(\Gamma(5)) = 1 + 5k$ for any $k \in \frac{1}{5}\mathbb{Z}_{\geq 0}$. (31) the leading exponents of $(f_2^{[5]})^{5k-\ell}(f_1^{[5]})^{\ell}$ are $\ell/5$, the sum of $\ell/5$ multiplied by 12 is equal to

$$
12\sum_{\ell=0}^{5k} \frac{\ell}{5} = 6k(5k+1) = d(d+k-1), \quad d = 1+5k.
$$

Therefore there exists some monic MLDE whose solution space is equal to the space $M_k(\Gamma(5), v_5^{1+5k})$. Hence we have

Theorem 13. Let m is $k = \frac{1}{5}\mathbf{Z}_{>0}$. There exists a monic modular linear differential equation of weight k and order $1 + 5k$ on $SL_2(\mathbb{Z})$, whose solution space coincides with the space of modular forms of weight k with multipliers v_5^{5k} on $\Gamma(5)$ for any k.

For the case of the characters of the minimal model of type $(2, 5)$, the symmetric representation and the MLDE are known (cf. [\[18\]](#page-16-10)).

8 Appendix

We now complete the proof of Theorem [8.](#page-7-1) The case $\ell = 0$ was done already, and then we show the theorem for $0 < \ell < 12$.

$\frac{12(2r+3)}{2r+3}$ gcd(4,r) gcd(3,r)			$12(24m+5)$ $6(24m+7)$ $4(24m+9)$ $12(24m+11)$	
$12(2r+3)$ gcd(4,r)gcd(3,r)	$12(24m+13)$ $4(24m+15)$ $2(24m+17)$ $12(24m+19)$			
		10		$12 (\ell = 0)$
$12(2r+3)$ $gcd(4,r) gcd(3,\overline{r})$	$4(24m + 21)$		$6(24m+23 \mid 12(24m+25))$	12p

Table 2: The values of Theorem [8](#page-7-1)

References

- [1] Y. Arike, K. Nagatomo and Y. Sakai, Vertex operator supper algebras and the theta group, preprint.
- [2] G. Anderson, G. Moore, Rationality in conformal field theory, Commmu. Math. Phys., 117, 441-450 (1988)
- [3] N. O. Azaiez, Formes quasi-modulaires sur des groupes modulaires co-compacts et restrictions des formes modulaires de Hilbert aux courbes modulaires (2005) <https://tel.archives-ouvertes.fr/tel-00011122/document>
- [4] C. Dong, X. Lin and S. H. Ng, Congruence property in conformal field theory, Algebra and Number Theory, 9, no. 9, 2121–2166 (2015)
- [5] H. Farkas and I. Kra, Riemann surfaces, First edition, Springer Verlag, New York, 1981.
- [6] B. Feigin and D. Fuchs, Verma modules over the Virasoro algebra, Lect. Notes in Math., 1060, 230–245 (1982)
- [7] B. Feigin and D. Fuchs, Representation theory of the Virasoro algebra, In Representation theory of Lie groups and related topics, Adv. Stud. Contemp. Math., 7, New York, Gordon and Breach, 465–554, 1990.
- [8] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, Duke Math. J., 66, No. 1, 123–168 (1992)
- [9] T. Ibukiyama, Graded rings of modular forms of rational weight, Research in Number Theory 6.1, 1–13, (2020)
- [10] T. Ibukiyama, Modular forms of rational weights and modular varieties, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, 70, no. 1, 315–339 (2000)
- [11] K. Iohara and Y. Koga, Representation theory of the Virasoro algebra, Springer-Verlag London, 2011.
- [12] E. L. Ince, Ordinary differential equations, Dover Publications, New York, 1944.
- [13] V. G. Kac, A. K. Raina and N. Rozhkovskaya, Bombay Lectures on highest weight representations of infinite dimensional Lie algebras, Second Edition, World Scientific Publishing Co. Pte. Ltd. 2013.
- [14] J. Lepowsky and H.-S. Li, Introduction to vertex operator algebras and their representations, Prog. Math., Birkhäuser, Boston, 2004.
- [15] G. Mason, Vector-valued modular forms and linear differential operators, Int. J. Number Theory, 3, no.3, 377–390 (2007)
- [16] A. Matsuo and K. Nagatomo, Axioms for a vertex algebra and the locality of quantum fields, MSJ Memoirs, 4, Mathematical Society of Japan, 1999.
- [17] A. Milas, Ramanujan's "Lost Notebook" and the Virasoro algebra, Commun. Math. Phys., 251, 567–588 (2004)
- [18] A. Milas, Virasoro algebra, Dedekind η -function and specialized MacDonald identities, Transformation Groups, 9, No. 3, 273–288 (2004)
- [19] A. Milas, E. Mortenson and K. Ono, Number theoretic properties of Wronskians of Andrews-Gordon series, Int. J. Number Theory, 4, no. 2, 323–337 (2008)
- [20] D. Zagier, Elliptic modular forms and their applications. In J. Bruinier, G. Harder, G. van der Geer and D. Zagier, The 1-2-3 of Modular Forms : Lectures at a Summer School in Nordfjordeid, Norway (ed. K. Ranestad). Universitext, Springer-Verlag, Berlin–Heidelberg–New York (2008), 1–103.
- [21] Y. Zhu, Modular invariance of characters of vertex operator algebras, J. Amer. Math. Soc., 9, 237–302 (1996)