SPECTRAL ANALYSIS OF A CLASS OF LÉVY-TYPE PROCESSES AND CONNECTION WITH SOME SPIN SYSTEMS

GRÉGOIRE VÉCHAMBRE¹

ABSTRACT. We consider a class of Lévy-type processes on which spectral analysis technics can be made to produce optimal results, in particular for the decay rate of their survival probability and for the spectral gap of their ground state transform. This class is defined by killed symmetric Lévy processes under general random time-changes satisfying some integrability assumptions. Our results reveal a connection between those processes and a family of spin systems. This connection, where the free energy and correlations play an essential role, is, up to our knowledge, new, and relates some key properties of objects from the two families. When the underlying Lévy process is a Brownian motion, the associated spin system turns out to have interactions of a rather nice form that are a natural alternative to the quadratic interactions from the lattice Gaussian Free Field. More general Lévy processes give rise to more general interactions in the associated spin systems.

MSC 2020. Primary: 60G53, 60G51, 60J25, 60J35, 82B20 Secondary: 60J55, 82B31

Keywords. Lévy-type processes, Lévy processes with random time changes, spectral analysis, ground state, ground state transform, spectral gap, survival probability, spin systems, interface models, partition function, free energy, infinite-volume Gibbs states, correlations

1. INTRODUCTION

A fundamental problem about Markov processes is to determine the asymptotic behavior of their survival probability (when their life-time is finite) or the speed of convergence toward their stationary distribution (when they are ergodic). Obtaining such estimates for Markov processes is a well-studied issue on which spectral analysis methods offer great insight [8], [52], [53], [16], [15]. However, even though those methods are extremely powerful, they unfortunately do not often provide explicit optimal estimates. This is especially true for Markov processes with jumps such as Lévy-type processes (see for example Chapter 6.1 of [6]). In particular, when the survival probability, or distance to the stationary distribution, of a Markov process decays exponentially fast, one can hope to establish the positivity of the decay rate, and sometimes to produce an interval that contains it, but it is unusual to obtain exact expressions of the decay rate. Unfortunately, there seem to be no hope to develop a methodology that would produce optimal estimates and exact expressions in full generality. It seems that a more reasonable approach is to isolate subclasses of sufficiently nice Markov processes and to develop optimal methodologies that are tailored for each class. For classes of Markov processes with jumps, this is already a challenging issue.

The goal of the present paper is to isolate a non-trivial and fairly large class of Lévy-type processes for which we can set up an adapted spectral analysis methodology that gets through, allowing to provide optimal estimates and useful exact expressions. The most natural and convenient way to represent this class is via randomly time-changed Lévy processes (but see also (1.4) below for an SDE representation). Several important classes of Markov processes admit representations via randomly time-changed Lévy processes, which are natural and interesting objects. Let us mention in particular Positive self-similar Markov processes (pssMp's) that include Bessel processes [27], [24], [35], Continuous-state Branching processes (CSBP's) [26], [25], some generalizations of CSBP's [29], [30], diffusions in random potentials (which can be represented as functions of randomly time-changed Brownian motions) [7], [20], [47], [50], or also the skew-product representation for planar Brownian motion [28], and many others. In our case, we consider a class of processes that are represented by killed symmetric Lévy processes under rather general random time-changes. Let us mention that some of the questions we study on these processes may be reformulated in terms of integral functionals of Lévy processes (see Section 1.1 below for more details and a brief account on that topic). The main specificity of the class we are interested in is the integrability assumptions on the characteristic exponent of the underlying Lévy process and on a function appearing in the random time-change. Those assumptions seem to be what make nice the class of processes and allow our methodology to work efficiently. We show that several key properties of those processes are related to properties of a class of spin systems. In particular, we prove optimal estimates for the asymptotic of the survival probability of a process in our class and show that the decay rate can be expressed in terms of a normalized version of the free energy of a spin system whose interactions are determined by the potential density of the underlying Lévy process. When the killing rate r goes to 0, the study of the spin system allows, under some conditions, to determine the exact asymptotic of the decay rate of the survival probability in terms of r. We also study the ground state transform of a process in our class and show that its spectral gap can be expressed in terms of the free energy and of the decay rate of correlations of the associated spin system. Further properties of the processes and of their ground state transforms are related to aspects of the spin systems such as infinite-volume Gibbs states. This surprising connection between our class of processes and equilibrium statistical mechanics is, up to our knowledge, new. While this connection turns out to be useful for understanding the processes we are interested in, it also opens a way for a deeper study of the class of spin systems, which is interesting on its own. In particular, when the underlying Lévy process is a Brownian motion, the associated spin system turns out to have interactions of a rather nice form that are a natural alternative to the quadratic interactions from the lattice Gaussian Free Field (GFF), and more general Lévy processes give rise to more general interactions in the associated spin system.

An interesting aspect of our methodology and results is that they are available for rather general random time-changes while, as we can see in the end of Section 1.1, several classical classes of Markov processes consist of functions of Lévy processes with a specific random time-change. The ideas introduced in the present paper can be applied to more general settings. In particular we aim to apply, in a subsequent study, those ideas to a fairly large class of Lévy-type processes on Lie groups.

1.1. A family of Lévy-type processes. Let ξ be a real symmetric Lévy process. Let $\psi_{\xi}(\cdot)$ be the characteristic exponent of ξ , i.e. for any $t \ge 0$ and $y \in \mathbb{R}$ we have $\mathbb{E}[e^{iy\xi(t)}] = e^{t\psi_{\xi}(y)}$. According to the Lévy-Khintchine formula and the symmetry of ξ , $\psi_{\xi}(\cdot)$ can be expressed by

$$\psi_{\xi}(y) = -\frac{A_{\xi}}{2}y^2 + \int_{\mathbb{R}} (e^{iyu} - 1 - yu\mathbf{1}_{[-1,1]}(u))\Pi_{\xi}(du),$$

where A_{ξ} is a non-negative number and Π_{ξ} is a symmetric measure on \mathbb{R} satisfying $\int_{\mathbb{R}} (1 \wedge u^2) \Pi_{\xi}(du) < \infty$. A_{ξ} and Π_{ξ} are called respectively the Gaussian component and the Lévy measure of ξ .

We now prepare for the definition of a process $X_x^{m,\xi,r}$ as a time changed version of $x + \xi(\cdot)$, where $x \in \mathbb{R}$ is the starting position of the process. As for many classical processes, we consider a time change given by an integral functional of ξ , but we do not impose a specific form for the random time-change. More precisely, we consider a time change defined as the inverse of the function $(t \mapsto \int_0^t m(x + \xi(s))ds)$, where m is a general positive function satisfying some assumptions. We assume the function m to be square-integrable on \mathbb{R} . This integrability assumption is not met by some classical families of Markov processes that can be represented via randomly time-changed Lévy processes (see the end of this subsection). However, in our case, it will be essential to define and study key quantities and operators, and to draw a connection with a spin system. We also assume m to be continuous on \mathbb{R} and vanishing at ∞ and $-\infty$ (we denote this by $m \in C_0(\mathbb{R})$). The assumptions on mare gathered as follows:

$$m \in \mathcal{C}_0(\mathbb{R}) \cap L^2(\mathbb{R}), \ \forall x \in \mathbb{R}, m(x) > 0.$$
 (Condition 1)

We will sometimes make the stronger assumption

$$n \in \mathcal{C}_0(\mathbb{R}) \cap L^1(\mathbb{R}), \ \forall x \in \mathbb{R}, m(x) > 0.$$
 (Condition 1')

Note that (Condition 1') implies (Condition 1).

Let r > 0. The killed symmetric Lévy process ξ^r is defined by $\xi^r(s) := \xi(s)$ for $s < e_r$ and $\xi^r(s) := \dagger$ for $s \ge e_r$, where e_r is an exponential random variable with parameter r, independent of ξ , and \dagger is a cemetery state. For any $x \in \mathbb{R}$ we define a random time-change as follows:

$$A_x(s) := \int_0^s m(x + \xi(u)) du.$$
 (1.1)

Since *m* is bounded, continuous, and positive on \mathbb{R} , $A_x(\cdot)$ is almost surely continuous and increasing on $[0, \infty)$. In particular it has an inverse $A_x^{-1}(\cdot)$ that is almost surely continuous and increasing on $[0, A_x(\infty))$. For $t \ge A_x(\infty)$ we set $A_x^{-1}(t) := \infty$ by convention. We refer to Lemma 41 for a condition for $A_x(\infty)$ to be infinite. For any

 $x \in \mathbb{R}$, we define the càd-làg process $X_x^{m,\xi,r}$ by $X_x^{m,\xi,r}(t) := x + \xi^r(A_x^{-1}(t))$, or more precisely,

$$X_x^{m,\xi,r}(t) := \begin{cases} x + \xi(A_x^{-1}(t)) \text{ if } A_x^{-1}(t) < e_r, \\ \dagger \text{ if } A_x^{-1}(t) \ge e_r. \end{cases}$$
(1.2)

Note that $X_x^{m,\xi,r}$ is almost surely killed in finite time. Indeed, let $\zeta_x := A_x(e_r) \in (0,\infty)$, then note from (1.2) that $X_x^{m,\xi,r}(t) \in \mathbb{R}$ for $t < \zeta_x$ and that $X_x^{m,\xi,r}(t) = \dagger$ for $t \ge \zeta_x$. We write $X^{m,\xi,r}$ to refer to the process defined in (1.2) without specification of a starting point. To $X^{m,\xi,r}$ we naturally associate a family $(P_t)_{t\ge 0}$ of linear operators, defined on the space $\mathcal{B}_b(\mathbb{R})$ of bounded measurable functions $f: \mathbb{R} \to \mathbb{C}$, by

$$P_t f(x) := \mathbb{E}\left[f(x + \xi(A_x^{-1}(t)))\mathbf{1}_{A_x^{-1}(t) < e_r}\right].$$
(1.3)

It is proved in Lemma 43 that $X^{m,\xi,r}$ is an homogeneous Markovian process, so that the family $(P_t)_{t\geq 0}$ is a semigroup. The following proposition establishes that $X^{m,\xi,r}$ is even a Feller process (or sub-Feller process, in the terminology of some authors).

Proposition 1. Assume that r > 0 and (Condition 1) holds, then the semigroup $(P_t)_{t\geq 0}$ defined by (1.3) is a Feller semigroup in the sense of Definition 1.2 of [6].

Proposition 1 is proved in Section 6.2, together with some useful facts about $X^{m,\xi,r}$. Proposition 1 requires that r > 0 and that (Condition 1) holds. In the rest of this paper, we will always make those assumptions when dealing with $(P_t)_{t\geq 0}$ so that, in particular, $(P_t)_{t\geq 0}$ enjoys all the properties of Feller semigroups.

Our primary goal is to characterize the decay rate of the survival probability $\mathbb{P}(\zeta_x > t)$ of $X_x^{m,\xi,r}$ via spectral analysis. In general, estimating the survival probability of a Markov process is not an easy issue. For pssMp's without positive jumps, several properties of their life-time are investigated in [38] via elaborated tools from complex analysis, yielding in particular the behavior of the survival probability of those processes. In our case, since $\zeta_x = A_x(e_r) \leq e_r ||m||_{\infty}$, we have the trivial bound $\mathbb{P}(\zeta_x > t) \leq e^{-rt/||m||_{\infty}}$, but what can be said about the exact decay rate of $\mathbb{P}(\zeta_x > t)$? Roughly, the further away the process $X_x^{m,\xi,r}$ is from 0, the faster time runs, so the faster killing occurs. Therefore, the behavior of $\mathbb{P}(\zeta_x > t)$ strongly depends on the behaviors of both the underlying Lévy process ξ and the function m (that determines the random time-change). Note also that, since $\zeta_x = A_x(e_r) = \int_0^{e_r} m(x+\xi(s))ds$, studying $\mathbb{P}(\zeta_x > t)$ amounts to studying the right distribution tail of the integral functional $\int_0^{e_r} m(x+\xi(s))ds$, which is also not an easy issue. Integral functionals of Lévy processes, or more generally of Markov processes, are well studied objects with a particular focus on determining conditions for finiteness/infiniteness of those functionals [21], [22]; this has applications ranging in several domains such as asymptotic properties of Schrödinger semigroups [3] or properties of SDEs driven by stable Lévy processes [2]. The moments of those functionals are studied in [13]. Our results show that $\mathbb{P}(\zeta_x > t)$ has order $K_{m,\xi,r}(x)e^{-t\gamma_{m,\xi,r}}$, where $\gamma_{m,\xi,r}$ and $K_{m,\xi,r}(x)$ are characterized uniquely in terms of a spin system. Under some conditions, we moreover determine the asymptotic of the coefficient $\gamma_{m,\xi,r}$ when r is small, that is, when the killing occurs at a low rate, so that the behavior of ξ has a strong influence. In this last case, it appears from our results that the trivial bound $\mathbb{P}(\zeta_x > t) \leq e^{-rt/\|m\|_{\infty}}$ is particularly bad. We also establish further spectral properties of the process $X^{m,\xi,r}$ and of its ground state transform, which draws more connections with the associated spin system. In particular, we express the spectral gap of the ground state transform of $X^{m,\xi,r}$ in terms of the associated spin system.

The survival probabilities of other classes of Markov processes have already been studied. For example, the integral functionals expressing the life-time of pssMp's are the so-called *exponential functionals of Lévy processes*, which have been intensively studied [5], [32], [19], [36], [34], [39], [41], [48]. Determining the asymptotic of the survival probability of pssMp's requires knowledge on the right distribution tail of exponential functionals, which have been studied in particular in [43], [31], [44], [45]. In the other way, non-trivial properties of exponential functionals of Lévy processes are sometimes derived from the study of pssMp's [37], [38]. Many nice properties of exponential functionals come from the exponential function that allows to exploit the independence of increments of Lévy processes. This includes the computation of moments of exponential functionals (see for example [5]), and more generally a useful functional equation satisfied by their Mellin transforms [32], [31], [23], [40]. In our case, the exponential function is replaced by a general function m satisfying (Condition 1), so we are deprived from those nice properties of exponential functionals (note also that our case excludes exponential functionals, since (Condition 1) is not satisfied by the exponential function).

In expressing $\gamma_{m,\xi,r}$, a key object that we use is the potential density of the killed Lévy process ξ^r . It is defined as the density of the potential measure $V_{\xi}^r(dx) := \int_0^\infty e^{-rt} \mathbb{P}(\xi(t) \in dx) dt$. For the potential density to be well-defined we need to assume

$$\int_{|y|>1} \frac{1}{|\psi_{\xi}(y)|} dy < \infty.$$
 (Condition 2)

Since ξ is symmetric, (Condition 2) coincides with conditions (43.5) and (43.6) from [46]. The later imply absolute continuity of potential measures (see Remark 43.6 in [46], see also Lemma 50 in the Appendix), which has important consequences in potential theory. In particular, it implies that any one point set is reached by ξ with positive probability (see Theorems 43.3 and 43.5 in [46]). Note also that (Condition 2) implies that ξ is of type C in the sense of Definition 11.9 of [46]. Indeed, if ξ was of type A or B, then from Lemma 43.11 in [46] we would have $|\psi_{\xi}(y)| = o(|y|)$ as |y| is large, so (Condition 2) would not be satisfied. Recall from Theorem 21.9 of [46] that a Lévy process is of type C in the sense of Definition 11.9 of [46] if and only if its sample paths on finite intervals have infinite variation almost surely. Under (Condition 2), for any r > 0, we denote by $v_{\xi}^{r}(\cdot)$ the potential density of ξ^{r} (see Lemma 50 for an expression and some properties). One can see that, under (Condition 2), $\mathbb{E}[\zeta_x] = \mathbb{E}[A_x(e_r)] = \int_{\mathbb{R}} v_{\xi}^{r}(y)m(x+y)dy$. This suggests that, if we assume (Condition 2), we may hope for an expression of $\gamma_{m,\xi,r}$ in term of m and $v_{\xi}^{r}(\cdot)$ as well. In our case, the potential density plays an important role in our results and in our methodology, so we assume (Condition 2). According to Theorem 43.9 of [46], (Condition 2) is never satisfied by Lévy processes in \mathbb{R}^d for d > 1. This is why we restrict our study to the case of real valued processes. Finally, if ξ is a symmetric α -stable Lévy process on \mathbb{R} with $\alpha > 1$, then $-\psi_{\xi}(y) = c|y|^{\alpha}$ for some c > 0 (see for example Theorem 14.15 in [46]) so (Condition 2) is clearly satisfied.

Remark 2 (A parallel with CSBP's). (Condition 2) may seem familiar as it resembles Grey's condition for a CSBP, which involves the Laplace exponent of the underlying Lévy process and is necessary and sufficient for extinction of the CSBP in finite time with positive probability [17]. Moreover, under the Grey condition, it is possible to derive exact expressions related to a CSBP, for example for the probability of non-extinction until time t [17].

In the literature CSBP's are sometimes represented via SDEs. It allows to define their extensions and to study them via stochastic analysis technics [10], [30], [33], [18]. In our case, we can also give an alternative definition of $X_x^{m,\xi,r}$ as follows: Let $(X(t))_{t\geq 0}$ be the solution of the SDE

$$X(t) = x + \int_0^t \sqrt{\frac{A_{\xi}}{m(X(s))}} dW(s) + \int_0^t \int_{[-1,1]} \int_0^{\frac{1}{m(X(s-))}} z \tilde{M}_1(ds, dz, du)$$

$$+ \int_0^t \int_{\mathbb{R} \setminus [-1,1]} \int_0^{\frac{1}{m(X(s-))}} z M_2(ds, dz, du),$$
(1.4)

where W, $\tilde{M}_1(ds, dz, du)$ and $M_2(ds, dz, du)$ are independent. W is a standard Brownian motion. $M_1(ds, dz, du)$ (resp. $M_2(ds, dz, du)$) is a Poisson random measure on $[0, \infty) \times [-1, 1] \times [0, \infty)$ (resp. $[0, \infty) \times (\mathbb{R} \setminus [-1, 1]) \times [0, \infty)$) with intensity measure $\mathbf{1}_{z \in [-1,1]} ds \times \Pi_{\xi}(dz) \times du$ (resp. $\mathbf{1}_{z \notin [-1,1]} ds \times \Pi_{\xi}(dz) \times du$), and $\tilde{M}_1(ds, dz, du) := M_1(ds, dz, du) - \mathbf{1}_{z \in [-1,1]} ds \times \Pi_{\xi}(dz) \times du$. Let e_r be an independent exponential random variable with parameter r. Provided that the SDE (1.4) is well-posed we set $X_x^{m,\xi,r}(t) := X(t)$ when $\int_0^t (1/m(X(s))) ds < e_r$ and $X_x^{m,\xi,r}(t) := \frac{1}{2}$ when $\int_0^t (1/m(X(s))) ds \ge e_r$. The $X_x^{m,\xi,r}$ obtained by this procedure has the same law as the one defined by (1.2). We will not justify the well-posedness of the SDE (1.4), nor the equivalence of the two definitions. We only work with the definition (1.2) of $X^{m,\xi,r}$ (via (1.3)), which allows us to conveniently exploit properties of the Lévy process ξ .

Remark 3. If ξ is a symmetric α -stable Lévy process on \mathbb{R} with $\alpha > 1$, then we see by the time-change method that the solution of (1.4) is equal in law to the solution of the more compact SDE $dX(t) = m(X(t))^{-1/\alpha} d\xi(t)$.

 $X^{m,\xi,r}$ is defined via (1.2) and a choice of ξ , m and r satisfying " ξ is symmetric", r > 0, (Condition 1) and (Condition 2). Some classical families of Markov processes can be represented via randomly time-changed Lévy processes, as in (1.2), but with different assumptions on ξ , m and r. Let us recall, for some of these classical families, which choices of ξ , m and r they correspond to.

- pssMp's: There is no restriction on the Lévy process ξ , one takes $m(x) := e^{\alpha x}$, where $\alpha > 0$ is the selfsimilarity index, and $r \ge 0$. The pssMp is then defined as the exponential of the resulting time-changed Lévy process [27], [24], [35]. See also [49] for generalizations of pssMp's where the Lévy process ξ is replaced by more general processes involving exponential functionals of bivariate Lévy processes.
- CSBP's: ξ has to be chosen as a Lévy process with no negative jumps killed upon hitting {0}, one takes $m(x) = x^{-1}$ and r = 0, see [26], [25].
- Generalizations of CSBP's: One can see that the process considered in [29] corresponds to taking ξ and r as for CSBP's but $m(x) = x^{-\theta}$ for some $\theta > 0$. In [30], their process is parametrized by three functions $\gamma_0, \gamma_1, \gamma_2$. The particular case $\gamma_0 = \gamma_1 = \gamma_2$ corresponds to taking ξ and r as for CSBP's and $m(x) = 1/\gamma_0(x)$.

1.2. Some spin systems. We now step in a framework of equilibrium statistical mechanics and define the spin system that will be related to the process $X^{m,\xi,r}$. All the spin system-related objects mentioned in this subsection will later be seen to be related to properties of the process $X^{m,\xi,r}$ (or its ground state transform), and thus allow to draw a connection between the later and the spin system. To introduce the model we adopt (and recall) the terminology of equilibrium statistical mechanics that is used for example in [14]. We fix $n \ge 2$ and consider a spin system on the one dimensional box $\Lambda_n := \{1, \ldots, n\} \subset \mathbb{Z}$. Each site $j \in \Lambda_n$ is viewed as a particle and has a spin $\omega_j \in \mathbb{R}$. A configuration of the system is determined by the spins of all particles, that is, by a vector $(\omega_1, \ldots, \omega_n) \in \mathbb{R}^{\Lambda_n}$. \mathbb{R}^{Λ_n} is thus called the *configuration space* of the system. To each configuration $(\omega_1,\ldots,\omega_n) \in \mathbb{R}^{\Lambda_n}$ we associated an energy $\mathcal{H}_{\Lambda_n}(\omega_1,\ldots,\omega_n)$. We assume nearest-neighbors interactions in the system (with periodic boundary condition) that favors agreement of the spins of neighboring sites. For this, we more precisely assume that the contribution to the energy of the pair of neighboring sites $\{j, j+1\} \subset \Lambda_n$ is given by $V_{int}(\omega_{j+1}-\omega_j)$ where V_{int} is assumed to be a continuous even function that converges to ∞ at ∞ . Spin systems with this form of interactions fall into the category of gradient models. Additionally, we penalize large values of the spins by assuming that each site $j \in \Lambda_n$ also adds a contribution $U_{mass}(\omega_j)$ to the energy, where U_{mass} is assumed to be a continuous function that converges to ∞ at ∞ . The energy of the configuration $(\omega_1,\ldots,\omega_n)\in\mathbb{R}^{\Lambda_n}$ is thus given by

$$\mathcal{H}_{\Lambda_n}(\omega_1,\ldots,\omega_n) = V_{int}(\omega_2-\omega_1) + \cdots + V_{int}(\omega_n-\omega_{n-1}) + V_{int}(\omega_1-\omega_n) + U_{mass}(\omega_1) + \cdots + U_{mass}(\omega_n).$$
(1.5)

The function $\mathcal{H}_{\Lambda_n} : \mathbb{R}^{\Lambda_n} \to \mathbb{R}$ is classically called the *Hamiltonian*. That function allows to define a probability measure called the *Gibbs distribution* on the configuration space \mathbb{R}^{Λ_n} via

$$\mu_{\Lambda_n}(A) = \frac{1}{Z_n} \int_A e^{-\mathcal{H}_{\Lambda_n}(\omega_1,\dots,\omega_n)} d\omega_1 \dots d\omega_n, \qquad (1.6)$$

for any Borel set $A \subset \mathbb{R}^{\Lambda_n}$. In the above expression, Z_n denotes the *partition function* of the system of n particles and is defined by

$$Z_n := \int_{\mathbb{R}^{\Lambda_n}} e^{-\mathcal{H}_{\Lambda_n}(\omega_1, \dots, \omega_n)} d\omega_1 \dots d\omega_n.$$
(1.7)

Of course, the Gibbs measure $\mu_{\Lambda_n}(\cdot)$ from (1.6) is well-defined only if $Z_n < \infty$. We will work with particular choices of V_{int} and U_{mass} that ensure $Z_n < \infty$ for all $n \ge 2$. The normalized free energy of the system is defined by

$$\mathcal{E} := \lim_{n \to \infty} -\frac{1}{n} \log(Z_n), \tag{1.8}$$

when the latter limit exists. That quantity is often related to some physical properties of spin systems. The model presented above ranges in the category of *effective interface models*. Such models often play the role of approximations for interfaces occurring in more realistic models. A particularly famous example of effective interface model is the *lattice Gaussian Free Field* (GFF). In particular, the lattice GFF on Λ_n corresponds to the above model with the choice $V_{int}(x) = \beta x^2$ and $U_{mass}(x) = \lambda x^2$, for some $\beta, \lambda \ge 0$ (see Chapter 8 of [14]).

Among objects of interest are the limit distributions of spins in fixed finite regions, as the size n of the global system goes to infinity. The distribution of the spins in the region $\{1, \ldots, k\}$ within the system of size n is determined by the quantities

$$L_n^k(f_1,\ldots,f_k) := \int_{\mathbb{R}^{\Lambda_n}} f_1(\omega_1)\ldots f_k(\omega_k)\mu_{\Lambda_n}(d\omega_1\ldots d\omega_n) = \frac{1}{Z_n} \int_{\mathbb{R}^{\Lambda_n}} f_1(\omega_1)\ldots f_k(\omega_k) e^{-\mathcal{H}_{\Lambda_n}(\omega_1,\ldots,\omega_n)} d\omega_1\ldots d\omega_n,$$
(1.9)

for $f_1, \ldots, f_k \in \mathcal{C}_b(\mathbb{R})$ (where $\mathcal{C}_b(\mathbb{R})$ denotes the space of bounded continuous functions $f : \mathbb{R} \to \mathbb{C}$). In equilibrium statistical mechanics, a function of the spins of a finite set of particles, such as $\prod_{j=1}^k f_j(\omega_j)$, is classically called a *local observable* and quantities such as $L_n^k(f_1, \ldots, f_k)$ are called *finite volume thermal averages of local observables* (see for example [51]). For any $k \ge 1$, $(L_n^k)_{n\ge 2\vee k}$ defines a sequence of probability measures on \mathbb{R}^k . For any $n\ge 2$ we see that $L_n^1(d\omega_1) = l_1^n(\omega_1)d\omega_1$ where

$$l_1^n(\omega_1) := \frac{1}{Z_n} \int_{\mathbb{R}^{n-1}} e^{-\mathcal{H}_{\Lambda_n}(\omega_1,\dots,\omega_n)} d\omega_2\dots d\omega_n.$$
(1.10)

Since we are interested in the infinite volume limit, we consider

$$\mathcal{L}_k(d\omega_1\dots d\omega_k) := \lim_{n \to \infty} L_n^k(d\omega_1\dots d\omega_k), \tag{1.11}$$

when the latter limit exists for the convergence in distributions. The limit laws \mathcal{L}_k are often called *infinite-volume* Gibbs states. Determining such limits of marginals is closely related to the problem of identifying *infinite volume*

Gibbs measures (see Chapters 6 and 8 in [14]). However, the study of infinite volume Gibbs measures is beyond the scope of this article so we will not develop that aspect.

In equilibrium statistical mechanics, it is also of interest to study the decay rate of correlations of local observables, since it provides information on correlation lengths in the global system. Let us consider, in the infinite volume limit system, observables of the spins of two particles that are at distance k from each other (for example the first and the $(k+1)^{th}$ particle). We assume that those observables are given by functions $f, g \in C_b(\mathbb{R})$ of the spins of the two particles. Then their correlation is given by

$$C_k(f,g) := \int_{\mathbb{R}^{k+1}} f(\omega_1)g(\omega_{k+1})\mathcal{L}_{k+1}(d\omega_1\dots d\omega_{k+1}) - \left(\int_{\mathbb{R}} f(\omega)\mathcal{L}_1(d\omega)\right) \left(\int_{\mathbb{R}} g(\omega)\mathcal{L}_1(d\omega)\right).$$
(1.12)

An important question is to determine the exact decay rate of those correlations as k goes to infinity.

Let us notice a convenient duality satisfied by our spin systems.

Remark 4 (Fourier duality for the partition function). Let m be a function satisfying (Condition 1') and let \hat{m} denote the characteristic function of the measure m(x)dx, ie $\hat{m}(z) := \int_{\mathbb{R}} e^{-izx} m(x)dx$. Fix r > 0, and let ξ be a symmetric Lévy process satisfying (Condition 2). Then under the choice $V_{int} := -\log(v_{\xi}^{r}(\cdot))$ and $U_{mass} := -\log(m(\cdot))$ we have that, for all $n \geq 2$, Z_n (defined by (1.7)) is well-defined and satisfies $Z_n = \hat{Z}_n/(2\pi)^n$ where

$$\hat{Z}_n := \int_{\mathbb{R}^n} \frac{\hat{m}(z_2 - z_1) \times \dots \times \hat{m}(z_n - z_{n-1}) \times \hat{m}(z_1 - z_n)}{(r - \psi_{\xi}(z_1)) \times \dots \times (r - \psi_{\xi}(z_n))} dz_1 \dots dz_n.$$
(1.13)

This is justified in Appendix B. In particular we have

$$Z_2 = \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} \frac{|\hat{m}(z_1 - z_2)|^2}{(r - \psi_{\xi}(z_1)) \times (r - \psi_{\xi}(z_2))} dz_1 dz_2.$$
(1.14)

Remark 4 can be interpreted as a duality between a system and a dual system, obtained by taking the Fourier transforms of $e^{-U_{mass}}$ and $e^{-V_{int}}$ and reversing their roles. Under some conditions of symmetry and positivity, the dual system can also satisfy the requirements of our definitions. In Section 4.2, this duality will facilitate the study of the behavior of the normalized free energy \mathcal{E} as r goes to 0, in order to prove Corollary 7 below.

In the spin system we introduced above, a rather natural choice for the interaction function V_{int} would be $V_{int}(x) = \beta |x|$, where the parameter β is usually interpreted physically as the inverse of the temperature. A key observation is that this can be related to the potential density of a killed Brownian motion. Indeed, when ξ^r is a standard Brownian motion killed at rate r then $v_{\xi}^r(x) = e^{-\sqrt{2r}|x|}/\sqrt{2r}$ (see e.g. Example 30.11 of [46]). This leads us to naturally wonder how the spin system with interaction function $V_{int}(x) = -\beta |x|$ can be related to a killed Brownian motion. More generally, this suggests that negative logarithms of potential densities of general symmetric Lévy processes are good candidates to generalize classical interaction functions in spin systems, and motivates investigating the relation between 1) the systems with those general interaction functions and 2) the corresponding killed Lévy processes. In the following subsection we state our main results that establish the connection between the two objects. To the best of our knowledge, such a connection between the two problems is new (both in the Brownian case and in the general case). Let us mention that a limited parallel can be made between the connection we just mentioned and the so-called random walk representations for spin systems. Those representations connect spin systems to random walks on their lattices and are fundamental tools for the study of the lattice GFF in particular and of other spin systems. Good references on such representations are [12], [11], [9], and Chapter 8 of [14] for the particular case of the lattice GFF. The connection between a spin system and its associated random walk allows to get a hand on correlations in the system and usually involves the potential of the random walk (through its Green function). Those aspects bear some similarities with the connection we present in the following subsection. However, one of the limits to the analogy is that, in our case, the spin system is connected to a process on \mathbb{R} instead of a process living on the same lattice.

1.3. Main results. The following result establishes the exponential decay of the survival probability $\mathbb{P}(\zeta_x > t)$ and relates in particular the decay rate with the normalized free energy of the spin system defined in Section 1.2.

Theorem 5. Assume that r > 0 and (Condition 1) and (Condition 2) hold. There is a positive constant $\gamma_{m,\xi,r}$ and, for any $x \in \mathbb{R}$, a positive constant $K_{m,\xi,r}(x)$, such that

$$\mathbb{P}(\zeta_x > t) \underset{t \to \infty}{\sim} K_{m,\xi,r}(x) e^{-t\gamma_{m,\xi,r}}.$$
(1.15)

Under the choice $V_{int} := -\log(v_{\xi}^r(\cdot))$ and $U_{mass} := -\log(m(\cdot))$, the coefficients $(Z_n)_{n\geq 2}$ and the normalized free energy \mathcal{E} (defined in respectively (1.7) and (1.8)) are well-defined and we have

$$\gamma_{m,\xi,r} = e^{\mathcal{E}},\tag{1.16}$$

and the infinite-volume Gibbs states \mathcal{L}_k (defined in (1.10)) are well-defined and absolutely continuous for all $k \geq 1$. Let us denote their densities by ℓ_k . Then $\ell_1/m \in \mathcal{C}_0(\mathbb{R}) \cap L^1(\mathbb{R})$, ℓ_1 is positive on \mathbb{R} , ℓ_1 is the point-wise limit of the density l_1^n (defined in (1.10)) as n goes to infinity, and

$$K_{m,\xi,r}(x) = \sqrt{\frac{\ell_1(x)}{m(x)}} \int_{\mathbb{R}} \sqrt{m(z)\ell_1(z)} dz.$$

$$(1.17)$$

If moreover we assume that (Condition 1') holds then we additionally have the bound

$$\frac{1}{\|m\|_{L^1(\mathbb{R})}} \times \frac{1}{v_{\xi}^r(0)} \le \gamma_{m,\xi,r} \le \frac{\|m\|_{L^1(\mathbb{R})} \times v_{\xi}^r(0)}{Z_2}.$$
(1.18)

Remark 6. The periodic boundary condition in the spin system appears naturally from our study of $X^{m,\xi,r}$, however that boundary condition can be removed. Indeed, if for $n \ge 2$ we define

$$\mathcal{H}^{f}_{\Lambda_{n}}(\omega_{1},\ldots,\omega_{n}) = V_{int}(\omega_{2}-\omega_{1}) + \cdots + V_{int}(\omega_{n}-\omega_{n-1}) + U_{mass}(\omega_{1}) + \cdots + U_{mass}(\omega_{n}), \tag{1.19}$$

 $Z_n^f := \int_{\mathbb{R}^{\Lambda_n}} e^{-\mathcal{H}_{\Lambda_n}^f(\omega_1, \dots, \omega_n)} d\omega_1 \dots d\omega_n \text{ and } \mathcal{E}_f \text{ as the limit of } -\log(Z_n^f)/n, \text{ then under the assumptions of Theorem 5 and the choice } V_{int} := -\log(v_{\xi}^r(\cdot)) \text{ and } U_{mass} := -\log(m(\cdot)), \mathcal{E}_f \text{ is also well-defined and we have } \mathcal{E}_f = \mathcal{E}. \text{ This is justified in Appendix C.}$

In the following result, we characterize the asymptotic behavior of $\gamma_{m,\xi,r}$ as r goes to 0 when the characteristic exponent $\psi_{\xi}(\cdot)$ is regular enough near zero.

Corollary 7 (Behavior for small r). Assume that (Condition 1') and (Condition 2) hold. Assume moreover that there are $\alpha \geq 1$ and c > 0 such that $-\psi_{\xi}(y) \sim c|y|^{\alpha}$ as y goes to 0. Then, for the coefficient $\gamma_{m,\xi,r}$ from Theorem 5 we have

$$\gamma_{m,\xi,r} \underset{r \to 0}{\sim} \frac{1}{\|m\|_{L^1(\mathbb{R})}} \times \frac{c^{\frac{1}{\alpha}}\pi}{\int_0^\infty \frac{1}{1+u^\alpha} du} r^{\frac{\alpha-1}{\alpha}} \quad if \, \alpha > 1,$$

$$(1.20)$$

$$\gamma_{m,\xi,r} \underset{r \to 0}{\sim} \frac{1}{\|m\|_{L^1(\mathbb{R})}} \times c\pi \left(\log\left(\frac{1}{r}\right) \right)^{-1} \quad if \ \alpha = 1.$$

$$(1.21)$$

Corollary 7 shows in particular that the bound $\gamma_{m,\xi,r} \geq r/||m||_{\infty}$ (resulting from the trivial inequality $\mathbb{P}(\zeta_x > t) \leq e^{-rt/||m||_{\infty}}$) is bad for small values of r. We give, in Section 4.2, a proof of Corollary 7 that mainly relies on spin system-related objects. More precisely, it relies on the inequality (1.18) that bounds the free energy in terms of the partition function, and on the duality from Remark 4 that allows to determine the asymptotic of the partition function as r goes to 0. A byproduct of this is that (1.18) is optimal for small values of r.

Example 8. Assume that (Condition 1') and (Condition 2) hold and that $\mathbb{E}[\xi(1)^2] < \infty$. In this case, since $\mathbb{E}[\xi(1)] = 0$ by the symmetry of ξ , we have $-\psi_{\xi}(y) \sim_{y \to 0} \mathbb{E}[\xi(1)^2]y^2/2$. By Corollary 7 applied with $c = \mathbb{E}[\xi(1)^2]/2$ and $\alpha = 2$ we get

$$\gamma_{m,\xi,r} \underset{r \to 0}{\sim} \frac{\sqrt{2r\mathbb{E}[\xi(1)^2]}}{\|m\|_{L^1(\mathbb{R})}}$$

Example 9. Assume that (Condition 1') holds and that ξ is a symmetric α -stable Lévy process on \mathbb{R} with $\alpha > 1$, then $-\psi_{\xi}(y) = c|y|^{\alpha}$ for some c > 0 (see for example Theorem 14.15 in [46]). In particular all the assumptions from Corollary 7 are satisfied (with this α and c) so $\gamma_{m,\xi,r}$ satisfies the estimate (1.20).

Remark 10. Under more general assumptions on the Lévy process ξ , an equivalent for $\gamma_{m,\xi,r}$ as r goes to 0 will be provided in Remark 40 from Section 4.2.

The following theorem establishes further spectral properties of the semigroup of the process $X^{m,\xi,r}$. $L^2(m)$ denotes the space of measurable functions $f : \mathbb{R} \to \mathbb{C}$ such that $\int_{\mathbb{R}} |f(y)|^2 m(y) dy < \infty$.

Theorem 11. Assume that r > 0 and (Condition 1) and (Condition 2) hold. The following claims hold:

- $(P_t)_{t\geq 0}$ extends uniquely to a strongly continuous contraction semigroup on $L^2(m)$ (we still denote this extension by $(P_t)_{t\geq 0}$) and, for t > 0, $P_t(L^2(m)) \subset C_0(\mathbb{R})$.
- There is an orthonormal Hilbert basis $(h_n)_{n\geq 1}$ of $L^2(m)$ such that for any $n\geq 1$, $h_n\in \mathcal{C}_0(\mathbb{R})\cap L^2(\mathbb{R})\subset L^2(m)$ and $P_t.h_n=e^{-t\lambda_n^X}h_n$ where $0<\lambda_1^X\leq \lambda_2^X\leq \ldots$ For any $f\in L^2(m)$ and t>0 we have

$$\forall x \in \mathbb{R}, \ P_t.f(x) = \sum_{n \ge 1} e^{-t\lambda_n^X} \langle f, h_n \rangle_{L^2(m)} h_n(x),$$
(1.22)

where the series of functions in the right-hand side converges absolutely in $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$.

Spectral analysis of a class of Lévy-type processes and connection with Interacting Particles Systems

• We have $\lambda_1^X < \lambda_2^X$ and

$$\lambda_1^X = \gamma_{m,\xi,r}, \qquad h_1(x) = \sqrt{\frac{\ell_1(x)}{m(x)}},$$
(1.23)

where $\gamma_{m,\xi,r}$ and ℓ_1 are as in Theorem 5.

• If moreover we assume that (Condition 1') holds then $(P_t)_{t\geq 0}$ has the strong Feller property in the sense that, for any t > 0, we have $P_t(\mathcal{B}_b(\mathbb{R})) \subset \mathcal{C}_0(\mathbb{R})$.

The function h_1 is sometimes referred to as the ground state and the quantity λ_1^X as the ground state energy. Note from (1.23) and Theorem 5 that h_1 is positive on \mathbb{R} so, under the assumptions of that theorem, we can define a family of measures $(\tilde{p}_t(x, dy))_{t \geq 0, x \in \mathbb{R}}$ on \mathbb{R} by

$$\tilde{p}_t(x, dy) := e^{t\lambda_1^X} \frac{h_1(y)}{h_1(x)} \mathbb{P}(X_x^{m,\xi,r}(t) \in dy).$$
(1.24)

The following proposition justifies that a Markov process can be associated to $(\tilde{p}_t(x, dy))_{t \ge 0, x \in \mathbb{R}}$. Such a process is classically called a *ground state transform*.

Proposition 12. Assume that r > 0 and (Condition 1) and (Condition 2) hold. There exists an homogeneous Markov process $\tilde{X}^{m,\xi,r}$ on \mathbb{R} such that $\mathbb{P}(\tilde{X}^{m,\xi,r}_x(t) \in dy) = \tilde{p}_t(x,dy)$ for any $t \ge 0$ and $x \in \mathbb{R}$. This process $\tilde{X}^{m,\xi,r}$ is unique, up to equality in law.

We now state results that show how the spin system from Section 1.2 is related to the Markov process $\tilde{X}^{m,\xi,r}$. In particular, the following proposition shows that the infinite-volume Gibbs states \mathcal{L}_k can be represented from the process $\tilde{X}^{m,\xi,r}$ taken at arrival times of a Poisson process. As mentioned in the end of Section 1.2, that representation can be paralleled with the random walk representations for spin systems, or also with the fact that the lattice GFF in dimension one can be represented by a random walk with Gaussian increments (see Exercise 8.6 in [14]).

Proposition 13. Assume that r > 0 and (Condition 1) and (Condition 2) hold. Let the distributions $(\mathcal{L}_k)_{k\geq 1}$ be as in Theorem 5. Let $(T_j)_{j\geq 1}$ be the sequence of arrival times of a standard Poisson process with parameter λ_1^X , independent of $\tilde{X}^{m,\xi,r}$. Set $T_0 := 0$ for convenience and denote $\tilde{X}_{\mathcal{L}_1}^{m,\xi,r}$ for the process $\tilde{X}^{m,\xi,r}$ with initial distribution $\mathcal{L}_1(dy)$. Then for any $k \geq 1$ the random vector $(\tilde{X}_{\mathcal{L}_1}^{m,\xi,r}(T_0), \ldots, \tilde{X}_{\mathcal{L}_1}^{m,\xi,r}(T_k))$ has law \mathcal{L}_{k+1} .

The following corollary shows that the correlations (defined in (1.12)) in the spin system from Section 1.2 decay exponentially in k.

Corollary 14 (Correlation decay). Assume that r > 0 and (Condition 1) and (Condition 2) hold. Under the choice $V_{int} := -\log(v_{\xi}^{r}(\cdot))$ and $U_{mass} := -\log(m(\cdot))$ in the spin system from Section 1.2, there exist C > 0 and a continuous non-zero bilinear form $B(\cdot, \cdot) : C_{b}(\mathbb{R}) \times C_{b}(\mathbb{R}) \to \mathbb{C}$ such that for any $f, g \in C_{b}(\mathbb{R})$ we have

$$e^{k\mathcal{C}}C_k(f,g) \xrightarrow[k \to \infty]{} B(f,g).$$
 (1.25)

The following theorem shows that the rate of correlation decay in the spin system from Section 1.2 is related to the spectral gap of the process $\tilde{X}^{m,\xi,r}$.

Theorem 15. Assume that r > 0 and (Condition 1) and (Condition 2) hold. $\tilde{X}^{m,\xi,r}$ is ergodic and its unique stationary distribution is $\mathcal{L}_1(dy)$ (where $\mathcal{L}_1(dy)$ is as in Theorem 5). For any $f \in \mathcal{C}_b(\mathbb{R})$ and $t \ge 0$ we have the following spectral gap inequality

$$\int_{\mathbb{R}} \left| \mathbb{E} \left[f\left(\tilde{X}_x^{m,\xi,r}(t) \right) \right] - \int_{\mathbb{R}} f(y) \mathcal{L}_1(dy) \right|^2 \mathcal{L}_1(dx) \le e^{-2tc} \int_{\mathbb{R}} \left| f(x) - \int_{\mathbb{R}} f(y) \mathcal{L}_1(dy) \right|^2 \mathcal{L}_1(dx),$$
(1.26)

where $c := e^{\mathcal{E}}(e^{\mathcal{C}}-1)$. Here \mathcal{E} is the normalized free energy, as in Theorem 5, and \mathcal{C} is the decay rate of correlations, as in Corollary 14. Moreover, the choice $c = e^{\mathcal{E}}(e^{\mathcal{C}}-1)$ is the largest choice of c such that (1.26) holds true for all $f \in \mathcal{C}_b(\mathbb{R})$.

Let us finish this section with two particular examples where several quantities and functions from the above theorems can be obtained explicitly.

Example 16. Let m(x) := 1/(1+|x|), $\xi := B$, where B denotes the standard Brownian motion, and r = 1/2 (this corresponds to choosing $V_{int}(x) = |x|$ and $U_{mass}(x) = \log(1+|x|)$ in the spin system from Section 1.2). Then we can show that the ground state h_1 from Theorem 11 has the simple expression $h_1(x) = \sqrt{2/3}(1+|x|)e^{-|x|}$ and that $\lambda_1^X = 1$. This is justified in Section 4.4. As a consequence, we get from (1.23), (1.16) and (1.17) that the

quantities from Theorem 5 have the following expressions: $\gamma_{m,B,1/2} = 1$, $\mathcal{E} = 0$, $\ell_1(x) = 2(1 + |x|)e^{-2|x|}/3$ and $K_{m,B,1/2}(x) = 4(1 + |x|)e^{-|x|}/3$.

Example 17. Let $m(x) := (2x^2 + 6|x| + 3)/(1 + |x|)^4$, $\xi := B$, where B denotes the standard Brownian motion, and r = 1/2 (this corresponds to choosing $V_{int}(x) = |x|$ and $U_{mass}(x) = 4\log(1 + |x|) - \log(2x^2 + 6|x| + 3)$ in the spin system from Section 1.2). Then we can show that the ground state h_1 from Theorem 11 has the simple expression $h_1(x) = ce^{-x^2/(1+|x|)}$ where c is a normalizing constant, and that $\lambda_1^X = 1/2$. The justification of this is similar to the justification of Example 16 in Section 4.4. As a consequence, we get from (1.23), (1.16) and (1.17) that the quantities from Theorem 5 have the following expressions: $\gamma_{m,B,1/2} = 1/2$, $\mathcal{E} = -\log(2)$, $\ell_1(x) = c^2(2x^2 + 6|x| + 3)e^{-2x^2/(1+|x|)}/(1+|x|)^4$ and $K_{m,B,1/2}(x) = \tilde{c}e^{-x^2/(1+|x|)}$, for some constant $\tilde{c} > 0$.

The above examples illustrate in particular cases the fact that, whenever we are able to get information on the eigenfunctions and eigenvalues of the generator of $X^{m,\xi,r}$, this yields information on the associated spin system. More generally, the connections built in the above results between, on one hand the processes $X^{m,\xi,r}$ and $\tilde{X}^{m,\xi,r}$, and on the other hand the associated spin system, allow to transform information obtained on one of these two objects into information on the other.

1.4. Sketch of proofs and organization of the paper. In Section 2 we study some linear operators that provide some understanding on the generator of $X^{m,\xi,r}$. More precisely, in Subsection 2.1 we introduce those operators and study their basic properties. In Subsection 2.2 we study the traces of the successive compositions of one of these operators. In Subsection 2.3 we prove that a duality holds between that operator and the generator of $X^{m,\xi,r}$.

In Section 3 we basically diagonalize the semigroup $(P_t)_{t\geq 0}$ of $X^{m,\xi,r}$ and deduce the asymptotic behavior of the survival probability $\mathbb{P}(\zeta_x > t)$. More precisely, in Subsection 3.1 we use the duality from Section 2 to construct and study a basis of eigenfunctions of the generator of $X^{m,\xi,r}$ on some natural Hilbert space, and to study the sequence of eigenvalues of that generator. In Subsection 3.2 we deduce a decomposition of the semigroup $(P_t)_{t\geq 0}$ and we study the special properties of the first eigenvalue and its associated eigenfunction. In Subsection 3.3 we apply the previous results to obtain a representation of the survival probability $\mathbb{P}(\zeta_x > t)$ and determine its asymptotic behavior. In Subsection 3.4 we study how the first eigenvalue of the generator of $X^{m,\xi,r}$ is related to the spin system from Section 1.2. In Subsection 3.5 we study the infinite-volume Gibbs states \mathcal{L}_k defined in (1.11) and relate them to the first eigenfunction of the generator of $X^{m,\xi,r}$.

In Section 4 we prove the main results on $X^{m,\xi,r}$ from Section 1.3. More precisely, in Subsection 4.1 we gather the pieces collected in Section 3 to prove Theorem 5. In Subsection 4.2 we prove Corollary 7 by studying the potential density of the killed Lévy process ξ at 0 and using the duality from Remark 4 and the inequality (1.18). In Subsection 4.3 we prove Theorem 11 from the results of Section 3. In Subsection 4.4 we justify Example 16.

In Section 5 we prove the rest of the main results from Section 1.3 that are about the ground state transform of $X^{m,\xi,r}$ and its relation with the spin system from Section 1.2. More precisely, in Subsection 5.1 we prove Proposition 12, in Subsection 5.2 we prove Proposition 13, in Subsection 5.3 we prove Theorem 15, and in Subsection 5.4 we prove Corollary 14.

In Section 6 we establish properties of the process $X^{m,\xi,r}$ that are used all along the paper. More precisely, in Subsection 6.1 we study properties of the random time change $A_x^{-1}(\cdot)$. In Subsection 6.2 we proves the Markov property for $X^{m,\xi,r}$ and the Feller property (Proposition 1). In Subsection 6.3 we study the support of $X^{m,\xi,r}$. In Subsection 6.4 we study the generator of $X^{m,\xi,r}$. In Subsection 6.5 we study some technical properties of the semigroup $(P_t)_{t\geq 0}$ of $X^{m,\xi,r}$.

Appendix A contains some technical facts about the real symmetric Lévy process ξ , Appendices B, and C contain the proofs of Remarks 4 and 6 respectively.

1.5. Another route for the survival probability. The spectral properties of the semigroup of $X^{m,\xi,r}$ are a main focus of this paper and the direct object of several of our results. This is why our approach is mainly spectral, in the sense that it relies substantially on establishing and using spectral properties of the semigroup of $X^{m,\xi,r}$. However, for the asymptotic of the survival probability of $X^{m,\xi,r}$ studied in Theorem 5, other approaches are also possible. One can start by noticing that the moments of the survival time ζ_x are related to the spin system from Section 1.2. Indeed, recall from Section 1.1 that $\zeta_x = A_x(e_r)$ so, since $A_x(\cdot)$ is an additive functional of a Markov process, we can use formula (4) from [13] and get

$$\frac{\mathbb{E}[(\zeta_x)^n]}{n!} = \int_{\mathbb{R}^n} v_{\xi}^r(\omega_1 - x) v_{\xi}^r(\omega_2 - \omega_1) \dots v_{\xi}^r(\omega_n - \omega_{n-1}) m(\omega_1) \dots m(\omega_n) d\omega_1 \dots d\omega_n.$$
(1.27)

The right-hand side is quite similar to the expression (1.7) of the partition function Z_n with the choice $V_{int} := -\log(v_{\xi}^r(\cdot))$ and $U_{mass} := -\log(m(\cdot))$, but with a different boundary condition. Let us denote by $\tilde{Z}_n(x)$ the right-hand side of (1.27). This suggests that the following alternative route may be used to prove (1.15)-(1.16) (and maybe (1.17)) or results in the same spirit:

- (1) Prove directly that $-\log(\tilde{Z}_n(x))/n$ converges to a limit that does not depend on x and that the free energy \mathcal{E} (defined in (1.8)) is well-defined, and identify the limit of $-\log(\tilde{Z}_n(x))/n$ with \mathcal{E} .
- (2) Using $\mathbb{E}[(\zeta_x)^n] = n! \tilde{Z}_n(x)$, the previous point, and the Cauchy, ÄiHadamard theorem, show that the Laplace transform of ζ_x has a pole at $e^{\mathcal{E}}$ and study relevant properties of this pole.
- (3) Prove or use an appropriate Tauberian theorem to deduce an estimate similar to (1.15) for the right distribution tail of ζ_x . (1.16) would then follow from the pole being located at $e^{\mathcal{E}}$.
- (4) Prove directly the existence of the infinite-volume Gibbs state \mathcal{L}_1 (defined in (1.10)) and of its positive density ℓ_1 , and identify the constant, in the equivalent of $\mathbb{P}(\zeta_x > t)$ found in the previous point, with $\sqrt{\ell_1(x)/m(x)} \int_{\mathbb{R}} \sqrt{m(z)\ell_1(z)} dz$. This would yield (1.17).

It is certainly not too difficult to carry out at least some steps of the above route by (or along with) using spectral properties of the semigroup of $X^{m,\xi,r}$, but in this case the approach that we use in this paper seems to be more direct. See Remark 37 for a moment-based proof of the lower bound in (1.18), provided that (1.15) has been proved. We are not sure how easy or difficult it would be to carry out each point of the route outlined above without making any use of spectral properties, and if further results like Corollary 7 can be obtained in this way.

1.6. Facts and notations. Let $\mathcal{F}^{\xi} := (\mathcal{F}^{\xi}_t)_{t\geq 0}$ and $\mathcal{F}^X := (\mathcal{F}^X_t)_{t\geq 0}$ denote the right continuous filtrations generated by respectively the Lévy process ξ and the process $X^{m,\xi,r}_x$.

Let $C_0(\mathbb{R})$ denote the space of continuous functions $f : \mathbb{R} \to \mathbb{C}$ that converge to 0 at ∞ and $-\infty$. $C_0(\mathbb{R})$ is equipped with the norm $\|\cdot\|_{\infty}$, defined by $\|f\|_{\infty} := \sup_{x \in \mathbb{R}} |f(x)|$, which make it a Banach space. We sometimes denote $(C_0(\mathbb{R}), \|\cdot\|_{\infty})$ to emphasis that we consider $C_0(\mathbb{R})$ equipped with this norm. For $k \geq 1$, let $C_0^k(\mathbb{R})$ denote the space of functions $f \in C_0(\mathbb{R})$ that are k times differentiable and whose successive derivatives of order 1, 2, ..., k belong to $C_0(\mathbb{R})$. $L^1(\mathbb{R})$ denotes the space of measurable functions $f : \mathbb{R} \to \mathbb{C}$ such that $\|f\|_{L^1(\mathbb{R})} := \int_{\mathbb{R}} |f(y)| dy < \infty$. $\mathcal{B}_b(\mathbb{R})$ and $\mathcal{C}_b(\mathbb{R})$ denote respectively the spaces of bounded measurable functions and bounded continuous functions $f : \mathbb{R} \to \mathbb{C}$.

 $L^2(\mathbb{R})$ and $L^2(m)$ denote the Hilbert spaces of measurable functions $f : \mathbb{R} \to \mathbb{C}$ such that, respectively, $\int_{\mathbb{R}} |f(y)|^2 dy < \infty$ and $\int_{\mathbb{R}} |f(y)|^2 m(y) dy < \infty$. We denote by $\langle \cdot, \cdot \rangle_{L^2(\mathbb{R})}$ and $\langle \cdot, \cdot \rangle_{L^2(m)}$ respectively the natural inner products on these spaces. We also denote $||f||_{L^2(\mathbb{R})} := (\langle f, f \rangle_{L^2(\mathbb{R})})^{1/2}$. $L^2((-1,1))$ denotes the Hilbert space of measurable functions $g : (-1,1) \to \mathbb{C}$ such that $\int_{(-1,1)} |g(y)|^2 dy < \infty$. We denote by $\langle \cdot, \cdot \rangle_{L^2((-1,1))}$ the natural inner product on $L^2((-1,1))$ and $||g||_{L^2((-1,1))} := (\langle g, g \rangle_{L^2((-1,1))})^{1/2}$.

For the Fourier transform we use the convention

$$(\mathcal{F}f)(x) := \int_{\mathbb{R}} f(y)e^{-2i\pi yx}dy, \quad (\mathcal{F}^{-1}f)(x) := \int_{\mathbb{R}} f(y)e^{2i\pi yx}dy, \quad \text{for } f \in L^1(\mathbb{R}).$$
(1.28)

According to Riemann-Lebesgue lemma, \mathcal{F} and \mathcal{F}^{-1} are contractions from $L^1(\mathbb{R})$ to $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$. The Fourier inversion theorem says that, if $f \in L^1(\mathbb{R})$ is such that $\mathcal{F}f \in L^1(\mathbb{R})$ (equivalently, $\mathcal{F}^{-1}f \in L^1(\mathbb{R})$) then $\mathcal{F}^{-1}(\mathcal{F}f) = \mathcal{F}(\mathcal{F}^{-1}f) = f$. With the convention (1.28), the extensions of \mathcal{F} and \mathcal{F}^{-1} to $L^2(\mathbb{R})$ (that we also denote by \mathcal{F} and \mathcal{F}^{-1}) are isometric and satisfy Plancherel's identity, i.e. for any $f, g \in L^2(\mathbb{R})$ we have $\langle f, g \rangle_{L^2(\mathbb{R})} = \langle \mathcal{F}f, \mathcal{F}g \rangle_{L^2(\mathbb{R})} = \langle \mathcal{F}^{-1}f, \mathcal{F}^{-1}g \rangle_{L^2(\mathbb{R})}$.

We denote the null function by **0**. For any functional Banach space \mathcal{U} equipped with a norm $\|\cdot\|_{\mathcal{U}}$, we denote the operator norm by $\||\cdot\||$, i.e. for any continuous operator $T: \mathcal{U} \to \mathcal{U}$, $\||T\|| := \sup_{f \in \mathcal{U} \setminus \{\mathbf{0}\}} \|T.f\|_{\mathcal{U}} / \|f\|_{\mathcal{U}}$. We denote by $\mathcal{L}(\mathcal{U})$ the space of continuous operators of \mathcal{U} equipped with the operator norm, which makes it a Banach space. For an operator T and $n \geq 1$, T^n denotes the operator obtained by n successive compositions of T.

According to Proposition 1, $(P_t)_{t\geq 0}$ is a Feller semigroup under (Condition 1) and r > 0. As mentioned after Proposition 1, we will always make those assumptions when dealing with $(P_t)_{t\geq 0}$ so that, in particular, $(P_t)_{t\geq 0}$ enjoys all the properties of Feller semigroups. We refer to [6] for background on the theory of Feller semigroups and their generators. The domain $\mathcal{D}(\mathcal{A}_{X^m,\xi,r})$ is the subspace of $\mathcal{C}_0(\mathbb{R})$ defined by $\{f \in \mathcal{C}_0(\mathbb{R}) \text{ s.t. } \lim_{t\to 0} t^{-1}(P_t.f-f) \text{ exists in } (\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})\}$. The generator $\mathcal{A}_{X^m,\xi,r} : \mathcal{D}(\mathcal{A}_{X^m,\xi,r}) \to \mathcal{C}_0(\mathbb{R})$ is defined by $\mathcal{A}_{X^m,\xi,r}f := \lim_{t\to 0} t^{-1}(P_t.f-f)$ for $f \in \mathcal{D}(\mathcal{A}_{X^m,\xi,r})$. $\mathcal{D}(\mathcal{A}_{X^m,\xi,r})$ is dense in $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$ (see for example Theorem 3.2.6 in [1]). It is known (see for example Theorem 3.2.6 in [1]) that, if $g \in \mathcal{D}(\mathcal{A}_{X^m,\xi,r})$, then $P_t.g \in \mathcal{D}(\mathcal{A}_{X^{m,\xi,r}})$ for any $t \ge 0$ and $\frac{d}{dt}P_t.g = \mathcal{A}_{X^{m,\xi,r}}P_t.g = P_t.\mathcal{A}_{X^{m,\xi,r}}g$. We refer to this as the differentiation rule for the semigroup. It is important to note that the limit involved in the definitions of this derivative is a limit in $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$.

It is classical that the un-killed (resp. killed) Lévy process ξ (resp. ξ^r) is also a Feller process, that is, its semigroup is Feller in the sense of Definition 1.2 of [6]. We denote its generator by \mathcal{A}_{ξ} (resp. \mathcal{A}_{ξ^r}) and the domain of its generator by $\mathcal{D}(\mathcal{A}_{\xi})$ (resp. $\mathcal{D}(\mathcal{A}_{\xi^r})$). Note that $\mathcal{D}(\mathcal{A}_{\xi}) = \mathcal{D}(\mathcal{A}_{\xi^r})$ and that $\mathcal{A}_{\xi^r}f = \mathcal{A}_{\xi}f - rf$. For any $\alpha > 0$, the α -potential measure of ξ is defined by $V_{\xi}^{\alpha}(dx) := \int_{0}^{\infty} e^{-\alpha t} \mathbb{P}(\xi(t) \in dx) dt$. The resolvent operator at α associated with ξ is defined by $V_{\xi}^{\alpha}f(x) := \int_{\mathbb{R}} f(x+y)V_{\xi}^{\alpha}(dy)$ for $f \in \mathcal{C}_{0}(\mathbb{R})$.

2. Some operators, their traces, and a duality

2.1. Definition of the operators and basic properties. We now introduce some definitions. Assume that r > 0 and (Condition 2) holds and let

$$\varphi_r(x) := \frac{2}{v_\xi^r(0)} \int_0^x \frac{1}{-\psi_\xi(2\pi y) + r} dy.$$

We see from Lemma 50 and the symmetry of ψ_{ξ} that we have $\varphi_r(\infty) = 1$ and $\varphi_r(-\infty) = -1$. Let φ_r^{-1} : $(-1,1) \to \mathbb{R}$ be the inverse function of φ_r . For any $y \in \mathbb{R}$, let $\phi_y^r : (-1,1) \to \mathbb{C}$ be the function defined by $\phi_y^r(x) := e^{-2i\pi y \varphi_r^{-1}(x)}/\sqrt{2}$. It can be seen using dominated convergence that $(y \mapsto \phi_y^r)$ is continuous from \mathbb{R} to $L^2((-1,1))$.

Let $Q_r : L^2(\mathbb{R}) \to L^2(\mathbb{R})$ be the operator defined via $Q_r \cdot f := \mathcal{F}^{-1}((-\psi_{\xi}(2\pi \cdot) + r)^{-1/2} \times f)$. Note that $(-\psi_{\xi}(2\pi \cdot) + r)^{-1/2}$ is bounded by $r^{-1/2}$. Therefore, if $f \in L^2(\mathbb{R}), (-\psi_{\xi}(2\pi \cdot) + r)^{-1/2} \times f \in L^2(\mathbb{R})$ so $\mathcal{F}^{-1}((-\psi_{\xi}(2\pi \cdot) + r)^{-1/2} \times f) \in L^2(\mathbb{R})$, i.e. $Q_r \cdot f \in L^2(\mathbb{R})$. Q_r is thus well defined. Also, it is easy to check that Ker $Q_r = \{\mathbf{0}\}$.

The following lemma states some properties of Q_r and relates it with ϕ_u^r .

Lemma 18. Assume that r > 0 and (Condition 2) holds.

- (1) Q_r is a bounded operator and $||Q_r||^2 \leq 1/r$.
- (2) $Q_r(L^2(\mathbb{R})) \subset C_0(\mathbb{R})$. In particular, for any $f \in L^2(\mathbb{R})$ and $y \in \mathbb{R}$, $Q_r(f(y))$ is defined for every $y \in \mathbb{R}$ and is continuous.
- (3) There is a bijective isometry $J: L^2(\mathbb{R}) \to L^2((-1,1))$ such that

$$\forall f \in L^2(\mathbb{R}), \forall y \in \mathbb{R}, \ Q_r.f(y) = \sqrt{v_{\xi}^r(0)} \langle J.f, \phi_y^r \rangle_{L^2((-1,1))}.$$
(2.29)

An important point of the proof of the third point is to choose a suitable Hilbert basis of $L^2(\mathbb{R})$ on which Q_r acts nicely. The action of Q_r on the elements of the basis is then determined by noticing a duality (2.33) between the Fourier transforms of some functions built from the basis, and Fourier coefficients of the functions ϕ_y^r .

Proof of Lemma 18. Using the definition of Q_r and Plancherel's identity we get

 ϵ

$$\|Q_r \cdot f\|_{L^2(\mathbb{R})}^2 = \|\mathcal{F}^{-1}((-\psi_{\xi}(2\pi\cdot) + r)^{-1/2} \times f)\|_{L^2(\mathbb{R})}^2 = \|(-\psi_{\xi}(2\pi\cdot) + r)^{-1/2} \times f\|_{L^2(\mathbb{R})}^2$$

$$\leq \|(-\psi_{\xi}(2\pi\cdot) + r)^{-1/2}\|_{\infty}^2 \cdot \|f\|_{L^2(\mathbb{R})}^2 = \|f\|_{L^2(\mathbb{R})}^2/r.$$
(2.30)

This proves the first claim.

Note from (Condition 2) that $(-\psi_{\xi}(2\pi\cdot)+r)^{-1/2} \in L^2(\mathbb{R})$. Therefore, if $f \in L^2(\mathbb{R}), (-\psi_{\xi}(2\pi\cdot)+r)^{-1/2} \times f \in L^1(\mathbb{R})$ by Cauchy-Schwartz inequality so $Q_r \cdot f = \mathcal{F}^{-1}((-\psi_{\xi}(2\pi\cdot)+r)^{-1/2} \times f) \in \mathcal{C}_0(\mathbb{R})$. This proves the second claim.

Let us define

$$e_n(x) := \sqrt{\frac{1}{v_{\xi}^r(0)}} \times \frac{e^{in\pi\varphi_r(x)}}{\sqrt{-\psi_{\xi}(2\pi x) + r}}, \ n \in \mathbb{Z}.$$
(2.31)

It is not difficult to check that $(e_n)_{n \in \mathbb{Z}}$ is an Hilbert basis of $L^2(\mathbb{R})$. Let $(g_n)_{n \in \mathbb{Z}}$ be the classical Hilbert basis of $L^2((-1,1))$, namely, $g_n(x) := e^{in\pi x}/\sqrt{2}$ for $n \in \mathbb{Z}$. We define an isometry $J : L^2(\mathbb{R}) \to L^2((-1,1))$ via $J.e_n := g_n$ for $n \in \mathbb{Z}$ and linear isometrical extension. Since Q_r is continuous on $L^2(\mathbb{R})$ we have for any $f \in L^2(\mathbb{R})$ that

$$Q_r.f = \sum_{k \in \mathbb{Z}} \langle f, e_k \rangle_{L^2(\mathbb{R})} Q_r.e_k.$$
(2.32)

We have $(-\psi_{\xi}(2\pi \cdot) + r)^{-1/2} \times e_k \in L^2(\mathbb{R}) \cap L^1(\mathbb{R})$ so, for $k \in \mathbb{Z}$ and $y \in \mathbb{R}$ we have

$$Q_{r}.e_{k}(y) = \left(\mathcal{F}^{-1}\frac{e_{k}}{\sqrt{-\psi_{\xi}(2\pi\cdot)+r}}\right)(y) = \sqrt{\frac{1}{v_{\xi}^{r}(0)}} \int_{\mathbb{R}} \frac{e^{ik\pi\varphi_{r}(x)}}{-\psi_{\xi}(2\pi x)+r} e^{2i\pi xy} dx$$
$$= \sqrt{v_{\xi}^{r}(0)} \int_{-1}^{1} \frac{e^{2i\pi y\varphi_{r}^{-1}(u)}}{\sqrt{2}} \frac{e^{ik\pi u}}{\sqrt{2}} du$$
$$= \sqrt{v_{\xi}^{r}(0)} \times \overline{\langle \phi_{y}^{r}, g_{k} \rangle_{L^{2}((-1,1))}},$$
(2.33)

where we have made the change of variable $u = \varphi_r(x)$. Since $\langle f, e_k \rangle_{L^2(\mathbb{R})} = \langle J.f, J.e_k \rangle_{L^2((-1,1))} = \langle J.f, g_k \rangle_{L^2((-1,1))}$, we get for any $y \in \mathbb{R}$,

$$\sum_{k\in\mathbb{Z}}\langle f,e_k\rangle_{L^2(\mathbb{R})}Q_r.e_k(y) = \sqrt{v_{\xi}^r(0)}\sum_{k\in\mathbb{Z}}\langle J.f,g_k\rangle_{L^2((-1,1))}\overline{\langle \phi_y^r,g_k\rangle_{L^2((-1,1))}} = \sqrt{v_{\xi}^r(0)}\langle J.f,\phi_y^r\rangle_{L^2((-1,1))},$$

where the last equality comes from Parseval's identity. Combining the above with (2.32) we get that (2.29) holds for a.e. $y \in \mathbb{R}$. Since $Q_r f \in C_0(\mathbb{R})$ and $(y \mapsto \phi_y^r)$ is continuous, both sides of the equality are continuous so (2.29) holds for every $y \in \mathbb{R}$. This proves the third claim.

Lemma 19. Assume that r > 0 and (Condition 2) holds. There is a constant C > 0 such that for any $h \in L^2((-1,1))$ we have $\int_{\mathbb{R}} |\langle \phi_y^r, h \rangle_{L^2((-1,1))}|^2 dy \leq C ||h||_{L^2((-1,1))}^2$.

Proof. Lemma 18 shows that the claim holds with $C = 1/rv_{\xi}^{r}(0)$.

For any $y \in \mathbb{R}$, let $P_y^r : L^2((-1,1)) \to L^2((-1,1))$ be defined by $P_y^r \cdot f := \langle f, \phi_y^r \rangle_{L^2((-1,1))} \phi_y^r$. Since $\|\phi_y^r\|_{L^2((-1,1))} = 1$, P_y^r is the orthogonal projection on $Span\{\phi_y^r\}$ and $\||P_y^r\|| = 1$. From the continuity of $y \mapsto \phi_y^r$ we can see that $y \mapsto P_y^r$ is continuous from \mathbb{R} to $\mathcal{L}(L^2((-1,1)))$.

The following lemma defines a linear operator R_r and shows some of its useful properties.

Lemma 20. Assume that r > 0 and (Condition 1) and (Condition 2) hold.

- (1) For any $h \in L^2((-1,1))$ we have $\int_{\mathbb{R}} m(y) \|P_y^r \cdot h\|_{L^2((-1,1))} dy < \infty$. We can thus define an operator $R_r : L^2((-1,1)) \to L^2((-1,1))$ by $R_r \cdot h := \int_{\mathbb{R}} m(y)(P_y^r \cdot h) dy$, where the integral is in the sense of Bochner.
- (2) For any $f, g \in L^2((-1,1))$ we have $\langle R_r.f, g \rangle_{L^2((-1,1))} = \int_{\mathbb{R}} m(y) \langle P_y^r.f, g \rangle_{L^2((-1,1))} dy$.
- (3) R_r is a bounded operator, Ker $R_r = \{0\}$, and R_r is compact and self-adjoint in $L^2((-1,1))$. In particular there is a Hilbert basis of $L^2((-1,1))$ consisting of eigenfunctions of R_r .
- (4) The eigenvalues of R_r are real, positive, and form a sequence converging to 0. Let $\lambda_1^R \ge \lambda_2^R \ge \ldots$ be the ordered sequence of eigenvalues of R_r , repeated with their respective multiplicities.

Proof. Using the definition of P_y^r we get

$$\forall h \in L^{2}((-1,1)), \ |\langle \phi_{y}^{r}, h \rangle_{L^{2}((-1,1))}|^{2} = \langle P_{y}^{r}.h, h \rangle_{L^{2}((-1,1))} = \|P_{y}^{r}.h\|_{L^{2}((-1,1))}^{2}.$$
(2.34)

Therefore Lemma 19 can be re-written as $\int_{\mathbb{R}} \|P_y^r \cdot h\|_{L^2((-1,1))}^2 dy \leq C \|h\|_{L^2((-1,1))}^2$. Combining this with $m \in L^2(\mathbb{R})$ (from (Condition 1)) and Cauchy-Schwartz inequality we get

$$\int_{\mathbb{R}} m(y) \|P_y^r . h\|_{L^2((-1,1))} dy \le \sqrt{C} \|m\|_{L^2(\mathbb{R})} \|h\|_{L^2((-1,1))} < \infty,$$
(2.35)

which proves the first claim of the lemma. By properties of Bochner integrals, if T is a continuous linear operator from $L^2((-1,1))$ to another Banach space, then the Bochner integral $\int_{\mathbb{R}} m(y)(T.P_y^r.h)dy$ is well-defined and we have $T.R_r.h = \int_{\mathbb{R}} m(y)(T.P_y^r.h)dy$. For any $g \in L^2((-1,1))$, we can apply this to the continuous linear operator $\langle \cdot, g \rangle_{L^2((-1,1))}$ from $L^2((-1,1))$ to \mathbb{R} , which yields the second claim of the lemma.

By properties of Bochner integrals we have

$$\|R_r.h\|_{L^2((-1,1))} \le \int_{\mathbb{R}} m(y) \|P_y^r.h\|_{L^2((-1,1))} dy.$$
(2.36)

Combining with (2.35) we get that R_r is a bounded operator. Using the second claim of the lemma, (2.34) and (2.29) from Lemma 18 we get for any $h \in L^2((-1, 1))$,

$$\langle R_r.h,h\rangle_{L^2((-1,1))} = \int_{\mathbb{R}} m(y) |\langle \phi_y^r,h\rangle_{L^2((-1,1))}|^2 dy = \frac{1}{v_{\xi}^r(0)} \int_{\mathbb{R}} m(y) |Q_r.J^{-1}.h(y)|^2 dy,$$
(2.37)

where J^{-1} denotes the inverse of the bijective isometry J from Lemma 18. Since m is positive on \mathbb{R} (by (Condition 1)), Ker $Q_r = \{\mathbf{0}\}$, and J^{-1} is bijective we get $h \in \text{Ker } R_r \Rightarrow h = \mathbf{0}$ so Ker $R_r = \{\mathbf{0}\}$. The second claim of the lemma and the fact that P_y^r is self-adjoint for all $y \in \mathbb{R}$ show that R_r is self-adjoint as well. For M > 0, since $m \in C_0(\mathbb{R})$ (by (Condition 1)) we have $\int_{[-M,M]} m(y) ||P_y^r|| dy \leq 2M ||m||_{\infty} < \infty$. We can thus define an operator $R_{r,M} \in \mathcal{L}(L^2((-1,1)))$ by $R_{r,M} := \int_{[-M,M]} m(y)P_y^r dy$, where the integral is in the sense of Bochner. Using (2.36) with $R_r - R_{r,M}$ in place of R_r and (2.35) with $m\mathbf{1}_{[-M,M]^c}$ in place of m we get $||R_r - R_{r,M}|| \leq \sqrt{C} ||m\mathbf{1}_{[-M,M]^c}||_{L^2(\mathbb{R})}$ so $R_{r,M}$ converges to R_r in $\mathcal{L}(L^2((-1,1)))$ as M goes to infinity. By properties of Bochner integrals, $R_{r,M}$ (and therefore R_r) can be approximated by finite linear combinations of operators P_y^r . Since each operator P_y^r has rank one, R_r is the limit of a sequence of finite rank operators and is thus compact. The existence of a Hilbert basis of $L^2((-1, 1))$ consisting of eigenfunctions of R_r then follows from the spectral theorem. This concludes the proof of the third claim of the lemma.

That the eigenvalues of R_r are real and form a sequence converging to 0 follow from R_r being self-adjoint and from the spectral theorem. From (2.37) we get that each eigenvalue of R_r is non-negative. Since Ker $R_r = \{0\}$, the eigenvalues of R_r are even positive. This concludes the proof of the fourth claim of the lemma.

Remark 21. The compactness of R_r alternatively follows from R_r being a Hilbert-Schmidt operator, which follows from the first part of Lemma 23 below.

2.2. Traces of compositions of R_r . The following lemma prepares the ground to work with the traces of compositions of R_r .

Lemma 22. Assume that r > 0 and (Condition 2) holds. Let $(u_k)_{k\geq 1}$ be a Hilbert basis of $L^2((-1,1))$. For any $y \in \mathbb{R}$ and any bounded operator H we have

$$\sum_{j\geq 1} |\langle H.P_y^r.u_j, u_j \rangle_{L^2((-1,1))}| \leq |||H||| \text{ and } \sum_{j\geq 1} \langle H.P_y^r.u_j, u_j \rangle_{L^2((-1,1))} = \langle H.\phi_y^r, \phi_y^r \rangle_{L^2((-1,1))}$$

Proof. By definition of P_y^r we can see that we have $\langle H.P_y^r.u_j, u_j \rangle_{L^2((-1,1))} = \langle H.\phi_y^r, u_j \rangle_{L^2((-1,1))} \times \overline{\langle \phi_y^r, u_j \rangle_{L^2((-1,1))}}$ so the first claim follows from Cauchy-Schwartz inequality for sums and Parseval's identity while the second claim follows from Parseval's identity for the inner product.

Recall that R_r^n denotes the composition of R_r by itself *n* times. The following lemma shows that the traces of compositions of R_r are well-defined and relate their asymptotic behaviors to λ_1^R .

Lemma 23. Assume that r > 0 and (Condition 1) and (Condition 2) hold. For any $n \ge 2$, and any Hilbert basis $(u_k)_{k\ge 1}$ of $L^2((-1,1))$ we have $\sum_{j\ge 1} |\langle R_r^n.u_j, u_j \rangle_{L^2((-1,1))}| < \infty$ and the quantity $\sum_{j\ge 1} \langle R_r^n.u_j, u_j \rangle_{L^2((-1,1))}$ does not depend on the choice of the Hilbert basis $(u_k)_{k\ge 1}$ so we denote it by $Tr(R_r^n)$. Moreover we have

$$Tr(R_r^n) \underset{n \to \infty}{\sim} N \times (\lambda_1^R)^n,$$
 (2.38)

where N denotes the multiplicity of the eigenvalue λ_1^R of R_r .

Remark 24. The proof of Lemma 31 below will show that the eigenvalue λ_1^R of R_r has multiplicity 1. A consequence will be that we have actually N = 1 in (2.38).

Proof of Lemma 23. Fix $n \ge 2$. Using the second point of Lemma 20 and Fubini's theorem we get

$$\sum_{j\geq 1} |\langle R_r^n . u_j, u_j \rangle_{L^2((-1,1))}| \leq \int_{\mathbb{R}^n} m(y_1) \dots m(y_n) \left(\sum_{j\geq 1} |\langle P_{y_n}^r \dots P_{y_1}^r . u_j, u_j \rangle_{L^2((-1,1))}| \right) dy_1 \dots dy_n =: I_n.$$
(2.39)

Using the definition of P_y^r we get that $\langle P_{y_n}^r \dots P_{y_1}^r u_j, u_j \rangle_{L^2((-1,1))}$ equals

$$\langle u_j, \phi_{y_1}^r \rangle_{L^2((-1,1))} \times \langle \phi_{y_1}^r, \phi_{y_2}^r \rangle_{L^2((-1,1))} \times \dots \times \langle \phi_{y_{n-1}}^r, \phi_{y_n}^r \rangle_{L^2((-1,1))} \times \langle \phi_{y_n}^r, u_j \rangle_{L^2((-1,1))}.$$
(2.40)

We now compute the terms $\langle \phi_{y_1}^r, \phi_{y_2}^r \rangle_{L^2((-1,1))}$. Using the definition of ϕ_y^r , the substitution $x = \varphi_r^{-1}(u)$, and Lemma 50, we get

$$\langle \phi_{y_1}^r, \phi_{y_2}^r \rangle_{L^2((-1,1))} = \frac{1}{2} \int_{-1}^{1} e^{2i\pi(y_2 - y_1)\varphi_r^{-1}(u)} du = \frac{1}{v_{\xi}^r(0)} \int_{\mathbb{R}} \frac{e^{2i\pi(y_2 - y_1)x}}{-\psi_{\xi}(2\pi x) + r} dx$$

$$= \frac{1}{v_{\xi}^r(0)} \mathcal{F}^{-1} \left(\frac{1}{-\psi_{\xi}(2\pi \cdot) + r}\right) (y_2 - y_1) = \frac{v_{\xi}^r(y_2 - y_1)}{v_{\xi}^r(0)}.$$

$$(2.41)$$

The combination of (2.40) and (2.41) yields that $\sum_{j\geq 1} |\langle P_{y_n}^r \dots P_{y_1}^r ... P_{y_1}^r ... u_j \rangle_{L^2((-1,1))}|$ is smaller than

$$\begin{split} & \frac{v_{\xi}^{r}(y_{2}-y_{1})\cdots v_{\xi}^{r}(y_{n}-y_{n-1})}{v_{\xi}^{r}(0)^{n-1}}\sum_{j\geq 1}|\langle u_{j},\phi_{y_{1}}^{r}\rangle_{L^{2}((-1,1))}|\times|\langle\phi_{y_{n}}^{r},u_{j}\rangle_{L^{2}((-1,1))}|\\ & \leq \frac{v_{\xi}^{r}(y_{2}-y_{1})\cdots v_{\xi}^{r}(y_{n}-y_{n-1})}{v_{\xi}^{r}(0)^{n-1}}\sqrt{\sum_{j\geq 1}|\langle u_{j},\phi_{y_{1}}^{r}\rangle_{L^{2}((-1,1))}|^{2}}\sqrt{\sum_{j\geq 1}|\langle\phi_{y_{n}}^{r},u_{j}\rangle_{L^{2}((-1,1))}|^{2}}\\ & = \frac{v_{\xi}^{r}(y_{2}-y_{1})\cdots v_{\xi}^{r}(y_{n}-y_{n-1})}{v_{\xi}^{r}(0)^{n-1}}\times\|\phi_{y_{1}}^{r}\|_{L^{2}((-1,1))}\times\|\phi_{y_{n}}^{r}\|_{L^{2}((-1,1))} = \frac{v_{\xi}^{r}(y_{2}-y_{1})\cdots v_{\xi}^{r}(y_{n}-y_{n-1})}{v_{\xi}^{r}(0)^{n-1}}, \end{split}$$

where we have used Cauchy-Schwartz inequality for sums, Parseval's identity, and $\|\phi_y^r\|_{L^2((-1,1))} = 1$. Plugging this in the definition of I_n in (2.39) we get

$$\begin{aligned}
I_n &\leq \frac{1}{v_{\xi}^r(0)^{n-1}} \int_{\mathbb{R}^n} m(y_1) \dots m(y_n) v_{\xi}^r(y_2 - y_1) \cdots v_{\xi}^r(y_n - y_{n-1}) dy_1 \dots dy_n \\
&= \frac{1}{v_{\xi}^r(0)^{n-1}} \int_{\mathbb{R}} m(y_n) \left(\dots \left(\int_{\mathbb{R}} m(y_2) v_{\xi}^r(y_3 - y_2) \left(\int_{\mathbb{R}} m(y_1) v_{\xi}^r(y_2 - y_1) dy_1 \right) dy_2 \right) \dots \right) dy_n \\
&= \frac{1}{v_{\xi}^r(0)^{n-1}} \int_{\mathbb{R}} m(y) f_{n-1}(y) dy,
\end{aligned}$$
(2.42)

where we have set $f_0(y) := 1$ and $f_{k+1}(y) := \int_{\mathbb{R}} m(z) v_{\xi}^r(y-z) f_k(z) dz$ for $k \ge 0$.

In order to prove that $I_n < \infty$, let us first show that for any function f,

$$nf \in L^{2}(\mathbb{R}) \Rightarrow \int_{\mathbb{R}} m(z) v_{\xi}^{r}(\cdot - z) f(z) dz \in L^{2}(\mathbb{R}).$$
(2.43)

Lemma 50 shows that $v_{\mathcal{E}}^r(\cdot) \in L^1(\mathbb{R}) \cap \mathcal{C}_0(\mathbb{R}) \subset L^2(\mathbb{R})$. Using Plancherel's identity we get

$$\int_{\mathbb{R}} m(z) v_{\xi}^{r}(y-z) f(z) dz = \int_{\mathbb{R}} \mathcal{F}^{-1}(mf)(x) \mathcal{F}^{-1}(v_{\xi}^{r}(y-\cdot))(x) dx = \int_{\mathbb{R}} e^{2i\pi yx} \mathcal{F}^{-1}(mf)(x) \mathcal{F}(v_{\xi}^{r}(\cdot))(x) dx.$$

By Plancherel's isometry we have $\mathcal{F}^{-1}(mf) \in L^2(\mathbb{R})$ and $\mathcal{F}(v_{\xi}^r(\cdot)) \in L^2(\mathbb{R})$ so the product of the two function is in $L^1(\mathbb{R})$ by Cauchy-Schwartz inequality. We can thus recognize the last term as the inverse Fourier transform of a function in $L^1(\mathbb{R})$ and get $\int_{\mathbb{R}} m(z)v_{\xi}^r(y-z)f(z)dz = \mathcal{F}^{-1}(\mathcal{F}^{-1}(mf) \times \mathcal{F}(v_{\xi}^r(\cdot)))(y)$. Since, by Lemma 50, $\mathcal{F}(v_{\xi}^r(\cdot)) = (-\psi_{\xi}(2\pi \cdot) + r)^{-1}$ which is bounded by 1/r we get that the product $\mathcal{F}^{-1}(mf) \times \mathcal{F}(v_{\xi}^r(\cdot))$ is actually in $L^1(\mathbb{R}) \cap L^2(\mathbb{R})$. By Plancherel's isometry, its Fourier transform is in $L^2(\mathbb{R})$ and this concludes the proof of (2.43).

Since $m \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$ by (Condition 1), (2.43) shows that $f_1 \in L^2(\mathbb{R})$ and then shows by induction that $f_k \in L^2(\mathbb{R})$ for all $k \ge 1$. Therefore, $I_n < \infty$ follows from applying Cauchy-Schwartz inequality in the right-hand side of (2.42). Combining this with (2.39) yields the claim $\sum_{j\ge 1} |\langle R_r^n . u_j, u_j \rangle_{L^2((-1,1))}| < \infty$.

Then, using the second point of Lemma 20, Fubini's theorem, and Lemma 22 we get

$$\sum_{j\geq 1} \langle R_r^n . u_j, u_j \rangle_{L^2((-1,1))} = \sum_{j\geq 1} \int_{\mathbb{R}^n} m(y_1) \dots m(y_n) \langle P_{y_n}^r \dots P_{y_1}^r . u_j, u_j \rangle_{L^2((-1,1))} dy_1 \dots dy_n$$

$$= \int_{\mathbb{R}^n} m(y_1) \dots m(y_n) \left(\sum_{j\geq 1} \langle P_{y_n}^r \dots P_{y_1}^r . u_j, u_j \rangle_{L^2((-1,1))} \right) dy_1 \dots dy_n$$

$$= \int_{\mathbb{R}^n} m(y_1) \dots m(y_n) \langle P_{y_n}^r \dots P_{y_2}^r . \phi_{y_1}^r, \phi_{y_1}^r \rangle_{L^2((-1,1))} dy_1 \dots dy_n,$$
(2.44)

where the use of Fubini's theorem is allowed since $I_n < \infty$. This proves the independence with respect to the choice of the Hilbert basis $(u_k)_{k\geq 1}$. $Tr(R_r^n)$ is thus well-defined.

Choosing, for $(u_k)_{k\geq 1}$, the Hilbert basis of eigenfunctions of R_r we get $Tr(R_r^n) = \sum_{j\geq 1} (\lambda_j^R)^n$ for all $n\geq 2$ so in particular,

$$\sum_{j\geq 1} (\lambda_j^R)^2 < \infty. \tag{2.45}$$

For $n \geq 2$ we have

$$N \times (\lambda_1^R)^n \le Tr(R_r^n) = \sum_{j\ge 1} (\lambda_j^R)^n = (\lambda_1^R)^n \left(N + \sum_{j>N} \left(\frac{\lambda_j^R}{\lambda_1^R}\right)^n\right) \le (\lambda_1^R)^n \left(N + \frac{1}{(\lambda_1^R)^2} \left(\frac{\lambda_{N+1}^R}{\lambda_1^R}\right)^{n-2} \sum_{j>N} (\lambda_j^R)^2\right).$$

$$(2.46)$$

Since $\lambda_{N+1}^R < \lambda_1^R$, the combination of (2.45) and (2.46) yields (2.38).

The following lemma slightly improves Lemma 23 when the stronger assumption (Condition 1') holds.

Lemma 25. Assume that r > 0 and (Condition 1') and (Condition 2) hold. For any $n \ge 1$, and any Hilbert basis $(u_k)_{k\ge 1}$ of $L^2((-1,1))$ we have $\sum_{j\ge 1} |\langle R_r^n.u_j, u_j \rangle_{L^2((-1,1))}| < \infty$ and the quantity $\sum_{j\ge 1} \langle R_r^n.u_j, u_j \rangle_{L^2((-1,1))}$ does not depend on the choice of the Hilbert basis $(u_k)_{k\ge 1}$ so we denote it by $Tr(R_r^n)$. Moreover we have $Tr(R_r) = \|m\|_{L^1(\mathbb{R})}$ and for any $n \ge 1, Tr(R_r^n) = \sum_{j\ge 1} \langle \lambda_j^R \rangle^n < \infty$.

Note that for $n \ge 2$ the result is a consequence of Lemma 23.

Proof. Fix $n \ge 1$. Note that for any $y \in \mathbb{R}$, $|||P_y^r||| \le 1$ because P_y^r is an orthogonal projection. Proceeding as in (2.39) and using Lemma 22 and $m \in L^1(\mathbb{R})$ (from (Condition 1')) we get

$$\begin{split} \sum_{j\geq 1} |\langle R_r^n . u_j, u_j \rangle_{L^2((-1,1))}| &\leq \int_{\mathbb{R}^n} m(y_1) \dots m(y_n) \left(\sum_{j\geq 1} |\langle P_{y_n}^r \dots P_{y_1}^r . u_j, u_j \rangle_{L^2((-1,1))}| \right) dy_1 \dots dy_n \\ &\leq \int_{\mathbb{R}^n} m(y_1) \dots m(y_n) dy_1 \dots dy_n = \|m\|_{L^1(\mathbb{R})}^n < \infty. \end{split}$$

We thus get $\sum_{j\geq 1} |\langle R_r^n.u_j, u_j \rangle_{L^2((-1,1))}| < \infty$. Moreover the above finiteness allows to use Fubini's theorem so (2.44) holds true in our case (for all $n \geq 1$). This proves the independence with respect to the choice of the Hilbert basis $(u_k)_{k\geq 1}$. $Tr(R_r^n)$ is thus well-defined. Choosing, for $(u_k)_{k\geq 1}$, the Hilbert basis of eigenfunctions of R_r we get $Tr(R_r^n) = \sum_{j\geq 1} (\lambda_j^R)^n < \infty$. Applying (2.44) with n = 1 we get $Tr(R_r) = \int_{\mathbb{R}} m(y) \|\phi_y^r\|_{L^2((-1,1))}^2 dy = \|m\|_{L^1(\mathbb{R})}$.

2.3. A duality with $X^{m,\xi,r}$.

Proposition 26. Assume that r > 0 and (Condition 1) and (Condition 2) hold. For any $h \in L^2((-1,1))$,

(

$$y \mapsto \langle R_r.\phi_y^r, h \rangle_{L^2((-1,1))}) \in \mathcal{D}(\mathcal{A}_{X^m,\xi,r}),$$
(2.47)

$$\forall y \in \mathbb{R}, \ \mathcal{A}_{X^{m,\xi,r}} \langle R_r.\phi_{\cdot}^r,h \rangle_{L^2((-1,1))}(y) = \frac{-1}{v_{\xi}^r(0)} \langle \phi_y^r,h \rangle_{L^2((-1,1))}.$$
(2.48)

Proof. Using the second point of Lemma 20, the definition of P_z^r , and (2.41) we get

$$\langle R_r.\phi_y^r,h\rangle_{L^2((-1,1))} = \int_{\mathbb{R}} m(z)\langle P_z^r.\phi_y^r,h\rangle_{L^2((-1,1))}dz = \int_{\mathbb{R}} m(z)\langle \phi_y^r,\phi_z^r\rangle_{L^2((-1,1))}\langle \phi_z^r,h\rangle_{L^2((-1,1))}dz$$

$$= \frac{1}{v_{\xi}^r(0)}\int_{\mathbb{R}} v_{\xi}^r(z-y)m(z)\langle \phi_z^r,h\rangle_{L^2((-1,1))}dz = \frac{1}{v_{\xi}^r(0)}\int_{\mathbb{R}} v_{\xi}^r(u)G_h(y+u)du,$$

$$(2.49)$$

where we have set $G_h(z) := m(z)\langle \phi_z^r, h \rangle_{L^2((-1,1))}$. Let us prove that $(z \mapsto \langle \phi_z^r, h \rangle_{L^2((-1,1))}) \in \mathcal{C}_0(\mathbb{R})$. We set $M_h(z) := h(\varphi_r(z))/(-\psi_{\xi}(2\pi z)+r)$. Since $h \in L^2((-1,1)) \subset L^1((-1,1))$ we have $\int_{\mathbb{R}} |M_h(u)| du = v_{\xi}^r(0) \int_{(-1,1)} |h(x)| dx/2 < \infty$ so $M_h \in L^1(\mathbb{R})$. Therefore $\mathcal{F}^{-1}M_h \in \mathcal{C}_0(\mathbb{R})$. Then we have

$$(\mathcal{F}^{-1}M_h)(z) = \int_{\mathbb{R}} e^{2i\pi u z} \frac{h(\varphi_r(u))}{-\psi_{\xi}(2\pi u) + r} du = \frac{v_{\xi}^r(0)}{2} \int_{(-1,1)} h(x) e^{2i\pi z \varphi_r^{-1}(x)} dx = \frac{v_{\xi}^r(0)}{\sqrt{2}} \overline{\langle \phi_z^r, h \rangle_{L^2((-1,1))}}.$$

Therefore $(z \mapsto \langle \phi_z^r, h \rangle_{L^2((-1,1))}) \in \mathcal{C}_0(\mathbb{R})$. Since $m \in \mathcal{C}_0(\mathbb{R})$ (by (Condition 1)) we get $G_h \in \mathcal{C}_0(\mathbb{R})$. $V_{\xi}^r G_h$ is thus well-defined (where V_{ξ}^r is the resolvent operator at r, as in Section 1.6) and, using Lemma 50, we get

$$V_{\xi}^{r}G_{h}(y) = \int_{\mathbb{R}} G_{h}(y+u)V_{\xi}^{r}(du) = \int_{\mathbb{R}} v_{\xi}^{r}(u)G_{h}(y+u)du = v_{\xi}^{r}(0)\langle R_{r}.\phi_{y}^{r},h\rangle_{L^{2}((-1,1))},$$

where we have used (2.49) for the last equality. Therefore $(y \mapsto \langle R_r.\phi_y^r,h\rangle_{L^2((-1,1))}) \in V_{\xi}^r(\mathcal{C}_0(\mathbb{R}))$. According to Lemma 1.27 in [6] we have $V_{\xi}^r(\mathcal{C}_0(\mathbb{R})) \subset \mathcal{D}(\mathcal{A}_{\xi})$ and for any $f \in \mathcal{C}_0(\mathbb{R}), \mathcal{A}_{\xi}V_{\xi}^rf - rV_{\xi}^rf = -f$. We thus get $(y \mapsto \langle R_r.\phi_y^r,h\rangle_{L^2((-1,1))}) \in \mathcal{D}(\mathcal{A}_{\xi}) = \mathcal{D}(\mathcal{A}_{\xi^r})$ and for all $y \in \mathbb{R}$,

$$\mathcal{A}_{\xi^r} \langle R_r.\phi_{\cdot}^r,h \rangle_{L^2((-1,1))}(y) = \frac{1}{v_{\xi}^r(0)} \left(\mathcal{A}_{\xi} V_{\xi}^r G_h(y) - r V_{\xi}^r G_h(y) \right) = -\frac{G_h(y)}{v_{\xi}^r(0)} = -\frac{m(y) \langle \phi_y^r,h \rangle_{L^2((-1,1))}}{v_{\xi}^r(0)}.$$
 (2.50)

Since $(z \mapsto \langle \phi_z^r, h \rangle_{L^2((-1,1))}) \in \mathcal{C}_0(\mathbb{R})$ we have $\frac{1}{m} \mathcal{A}_{\xi^r} \langle R_r.\phi_\cdot^r, h \rangle_{L^2((-1,1))} \in \mathcal{C}_0(\mathbb{R})$ so, by Lemma 45, we get (2.47) and $\mathcal{A}_{X^m,\xi,r} \langle R_r.\phi_\cdot^r, h \rangle_{L^2((-1,1))}(y) = \frac{1}{m(y)} \mathcal{A}_{\xi^r} \langle R_r.\phi_\cdot^r, h \rangle_{L^2((-1,1))}(y)$. Combining this with (2.50) we get (2.48).

15

3. Decomposition of the survival probability of $X^{m,\xi,r}$

3.1. A basis of eigenfunctions of the generator. According to Lemma 20 we can, under the assumptions of that lemma, choose an orthonormal Hilbert basis $(a_n)_{n\geq 1}$ of $L^2((-1,1))$ consisting of eigenfunctions of R_r such that λ_n^R is the eigenvalue associated with a_n . For any $n \geq 1$ and $y \in \mathbb{R}$ let us define

$$q_n(y) := \frac{1}{\sqrt{\lambda_n^R}} \langle \phi_y^r, a_n \rangle_{L^2((-1,1))}.$$
(3.51)

The following result shows that the family $(q_n)_{n\geq 1}$ diagonalizes the generator $\mathcal{A}_{X^{m,\xi,r}}$. It is a consequence of the duality established in Proposition 26.

Proposition 27. Assume that r > 0 and (Condition 1) and (Condition 2) hold. For any $n \ge 1$, $q_n \in C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$ and $||q_n||_{L^2(\mathbb{R})} \le \sqrt{C/\lambda_n^R}$, where C is as in Lemma 19. $(q_n)_{n\ge 1}$ is an orthonormal Hilbert basis of $L^2(m)$. Moreover for any $n \ge 1$, $q_n \in \mathcal{D}(\mathcal{A}_{X^m,\xi,r})$ and

$$\mathcal{A}_{X^{m,\xi,r}}q_n = \frac{-1}{\lambda_n^R v_\xi^r(0)} q_n. \tag{3.52}$$

Proof. $q_n \in L^2(\mathbb{R})$ and the bound $||q_n||_{L^2(\mathbb{R})} \leq \sqrt{C/\lambda_n^R}$ are a consequence of (3.51) and Lemma 19. Using (3.51), that a_n is an eigenfunction of R_r , and that R_r is self-adjoint we get

$$\forall y \in \mathbb{R}, \ q_n(y) = \frac{1}{\sqrt{\lambda_n^R}} \langle \phi_y^r, a_n \rangle_{L^2((-1,1))} = \frac{1}{(\lambda_n^R)^{3/2}} \langle \phi_y^r, R_r.a_n \rangle_{L^2((-1,1))} = \frac{1}{(\lambda_n^R)^{3/2}} \langle R_r.\phi_y^r, a_n \rangle_{L^2((-1,1))}.$$
(3.53)

By (2.47) we get $q_n \in \mathcal{D}(\mathcal{A}_{X^{m,\xi,r}}) \subset \mathcal{C}_0(\mathbb{R})$. Then the combination of (3.53), (2.48) and (3.51) yields

$$\mathcal{A}_{X^{m,\xi,r}}q_n = \frac{1}{(\lambda_n^R)^{3/2}} \mathcal{A}_{X^{m,\xi,r}} \langle R_r.\phi_{\cdot}^r, a_n \rangle_{L^2((-1,1))}(y) = \frac{-1}{(\lambda_n^R)^{3/2} v_{\xi}^r(0)} \langle \phi_y^r, a_n \rangle_{L^2((-1,1))} = \frac{-1}{\lambda_n^R v_{\xi}^r(0)} q_n(y),$$

which is (3.52). Then for any $m, n \ge 1$, using (3.51), the definition of P_y^r , the second point of Lemma 20, and that a_m is an eigenfunction of R_r we get

$$\begin{split} \sqrt{\lambda_n^R \lambda_m^R} \langle q_n, q_m \rangle_{L^2(m)} &= \int_{\mathbb{R}} m(y) \langle \phi_y^r, a_n \rangle_{L^2((-1,1))} \langle a_m, \phi_y^r \rangle_{L^2((-1,1))} dy = \int_{\mathbb{R}} m(y) \langle P_y^r. a_m, a_n \rangle_{L^2((-1,1))} dy \\ &= \langle R_r. a_m, a_n \rangle_{L^2((-1,1))} = \lambda_m^R \langle a_m, a_n \rangle_{L^2((-1,1))}. \end{split}$$

Therefore $(q_n)_{n\geq 1}$ is an orthonormal family of $L^2(m)$. Now let $f \in L^2(m)$ be such that $\langle q_n, f \rangle_{L^2(m)} = 0$ for all $n \geq 1$ and let us prove that f = 0. For any $g \in L^2((-1,1))$ we set $q_g(y) := \langle \phi_y^r, g \rangle_{L^2((-1,1))}$. Note from Lemma 19 that $q_g \in L^2(\mathbb{R}) \subset L^2(m)$. Moreover, if $g \in Span < a_1, a_2, \dots >$ then q_g is a linear combination of finitely many q_n 's so $\langle q_g, f \rangle_{L^2(m)} = 0$. For an arbitrary $g \in L^2((-1,1))$, let $(g_n)_{n\geq 1}$ a sequence in $Span < a_1, a_2, \dots >$ that converges to g in $L^2((-1,1))$. Then, using Lemma 19, we get

$$\|q_g - q_{g_n}\|_{L^2(m)}^2 \le \|m\|_{\infty} \|q_g - q_{g_n}\|_{L^2(\mathbb{R})}^2 = \|m\|_{\infty} \int_{\mathbb{R}} |\langle \phi_y^r, g - g_n \rangle_{L^2((-1,1))}|^2 dy \le C \|m\|_{\infty} \|g - g_n\|_{L^2((-1,1))}^2,$$

with C as in Lemma 19. Therefore q_{g_n} converges to q_g in $L^2(m)$ so $\langle q_g, f \rangle_{L^2(m)} = 0$. For any $g \in L^2((-1,1))$ let us define two linear forms from $L^2(m)$ to \mathbb{R} :

$$\forall h \in L^{2}(m), \ A_{g}(h) := \langle q_{g}, h \rangle_{L^{2}(m)}, \ B_{g}(h) := \frac{1}{\sqrt{2}} \langle \mathcal{F}(\overline{h}m)(\varphi_{r}^{-1}(\cdot)), g \rangle_{L^{2}((-1,1))}.$$

 A_g is clearly continuous on $L^2(m)$. Then for any $h \in L^2(m)$ we have $\|\overline{h}m\|_{L^2(\mathbb{R})}^2 \leq \|m\|_{\infty} \|h\|_{L^2(m)}^2$ so $(h \mapsto \overline{h}m)$ is continuous from $L^2(m)$ to $L^2(\mathbb{R})$. \mathcal{F} is continuous from $L^2(\mathbb{R})$. For any $k \in L^2(\mathbb{R})$,

$$\|k(\varphi_r^{-1}(\cdot))\|_{L^2((-1,1))}^2 \leq \frac{1}{r} \int_{(-1,1)} |k(\varphi_r^{-1}(x))|^2 (-\psi_{\xi}(2\pi\varphi_r^{-1}(x)) + r) dx = \frac{2}{rv_{\xi}^r(0)} \int_{\mathbb{R}} |k(y)|^2 dy = \frac{2}{rv_{\xi}^r(0)} \|k\|_{L^2(\mathbb{R})}^2,$$

so $(k \mapsto k(\varphi_r^{-1}(\cdot)))$ is continuous from $L^2(\mathbb{R})$ to $L^2((-1,1))$. Therefore B_g is continuous on $L^2(m)$. Let $h \in L^2(\mathbb{R})$. Using $m \in L^2(\mathbb{R})$ (from (Condition 1)), two times Cauchy-Schwartz inequality, and $\|\phi_y^r\|_{L^2((-1,1))} = 1$ we get

$$\int_{\mathbb{R}} \int_{(-1,1)} m(y) \times |h(y)| \times |\phi_y^r(x)| \times |g(x)| dx dy \le \|m\|_{L^2(\mathbb{R})} \times \|h\|_{L^2(\mathbb{R})} \times \|g\|_{L^2((-1,1))} < \infty$$

We can thus use Fubini's theorem and get

$$\int_{\mathbb{R}} m(y)\overline{h(y)} \left(\int_{(-1,1)} \phi_y^r(x)\overline{g(x)}dx \right) dy = \int_{(-1,1)} \left(\int_{\mathbb{R}} m(y)\overline{h(y)}\phi_y^r(x)dy \right) \overline{g(x)}dx.$$
(3.54)

On the one hand, the left-hand side of (3.54) equals $\int_{\mathbb{R}} m(y)\overline{h(y)}\langle \phi_y^r, g \rangle_{L^2((-1,1))} dy = \int_{\mathbb{R}} q_g(y)\overline{h(y)}m(y)dy = A_g(h)$. On the other hand, the right-hand side of (3.54) equals

$$\frac{1}{\sqrt{2}}\int_{(-1,1)} \left(\int_{\mathbb{R}} m(y)\overline{h(y)}e^{-2i\pi y\varphi_r^{-1}(x)}dy\right)\overline{g(x)}dx = \frac{1}{\sqrt{2}}\int_{(-1,1)} \mathcal{F}(\overline{h}m)(\varphi_r^{-1}(x))\overline{g(x)}dx = B_g(h).$$

Therefore, for any $g \in L^2((-1,1))$, the linear forms A_g and B_g coincide on the subspace $L^2(\mathbb{R})$ that is dense in $L^2(m)$. Since A_g and B_g are continuous on $L^2(m)$ we get $A_g = B_g$. We have already shown that $A_g(f) = 0$ for all $g \in L^2((-1,1))$ so $B_g(f) = 0$ for all $g \in L^2((-1,1))$. We deduce that $\mathcal{F}(\overline{f}m)(\varphi_r^{-1}(\cdot)) = 0$ a.e. on (-1,1) so $\mathcal{F}(\overline{f}m) = 0$ a.e. on \mathbb{R} so $\overline{f}m = 0$ a.e. on \mathbb{R} and since m is positive on \mathbb{R} (by (Condition 1)), f = 0 a.e. on \mathbb{R} . Therefore the orthonormal family $(q_n)_{n\geq 1}$ is total in $L^2(m)$ so it is an Hilbert basis. This concludes the proof. \Box

Under the assumptions of Proposition 27, let $(\lambda_n^X)_{n\geq 1}$ be the sequence defined by

$$\forall n \ge 1, \ \lambda_n^X := \frac{1}{\lambda_n^R v_{\varepsilon}^r(0)}.$$
(3.55)

Proposition 27 shows that $(\lambda_n^X)_{n\geq 1}$ is the sequence of eigenvalues of $-\mathcal{A}_{X^m,\xi,r}$ associated with the eigenfunctions $(q_n)_{n\geq 1}$.

Lemma 28. Assume that r > 0 and (Condition 1) and (Condition 2) hold. For any $\alpha \ge 5/2$ we have

$$\sum_{n\geq 1} (\lambda_n^X)^{-\alpha} (1 \vee ||q_n||_\infty) < \infty.$$
(3.56)

In particular, for any $\beta \geq 0$ and t > 0 we have

$$\sum_{n\geq 1} (\lambda_n^X)^{\beta} e^{-t\lambda_n^X} (1 \vee ||q_n||_{\infty}) < \infty.$$
(3.57)

Proof. (3.57) easily follows from (3.56) so we only prove the later. From (3.51) and Cauchy-Schwartz inequality we see that for any $n \ge 1$ and $y \in \mathbb{R}$ we have

$$|q_n(y)| \le \frac{1}{\sqrt{\lambda_n^R}} \times \|\phi_y^r\|_{L^2((-1,1))} \times \|a_n\|_{L^2((-1,1))} = \frac{1}{\sqrt{\lambda_n^R}},$$

where, for the last inequality, we have used that $(a_n)_{n\geq 1}$ is an orthonormal family and that $\|\phi_y^r\|_{L^2((-1,1))} = 1$, which is easily seen from the definition of ϕ_y^r in Section 2.1. We thus get $\|q_n\|_{\infty} \leq 1/\sqrt{\lambda_n^R}$. We see from this and (3.55) that we only need to show $\sum_{n\geq 1} (\lambda_n^X)^{-2} < \infty$, but this is a consequence of (3.55) and (2.45). This concludes the proof.

3.2. Decomposition of the semigroup in the basis, and the first eigenfunction. The following proposition provides a decomposition of P_t f in the basis $(q_n)_{n\geq 1}$ for $f \in C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$.

Proposition 29. Assume that r > 0 and (Condition 1) and (Condition 2) hold. For any $f \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$ and t > 0 we have

$$\forall x \in \mathbb{R}, \ P_t.f(x) = \sum_{n \ge 1} e^{-t\lambda_n^X} \langle f, q_n \rangle_{L^2(m)} q_n(x),$$
(3.58)

where the series of functions in the right-hand side converges absolutely in $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$.

Proof. Let $f \in L^2(m)$ and t > 0. According to Proposition 27, $(q_n)_{n\geq 1}$ is an orthonormal family so, by Cauchy-Schwartz inequality, we have $|\langle f, q_n \rangle_{L^2(m)}| \leq ||f||_{L^2(m)}$ for any $n \geq 1$. Moreover Proposition 27 shows that $q_n \in C_0(\mathbb{R})$ for any $n \geq 1$. Combining with Lemma 28 we get that the series of functions in the right-hand side of (3.58) converges absolutely in $(C_0(\mathbb{R}), \|\cdot\|_{\infty})$. In particular the right-hand side of (3.58) is continuous in x, let us denote it by $S_t.f(x)$. Note that there is a constant $C_t > 0$ such that $||S_t.f||_{\infty} \leq C_t ||f||_{L^2(m)}$.

By Proposition 27 we have $||q_n||_{L^2(\mathbb{R})} \leq \sqrt{C/\lambda_n^n}$ for some C > 0. Combining this with $|\langle f, q_n \rangle_{L^2(m)}| \leq ||f||_{L^2(m)}$ and Lemma 28 (together with (3.55)) we get that $S_t \cdot f \in L^2(\mathbb{R}) \subset L^2(m)$. We now compute $\langle S_t \cdot f, q_n \rangle_{L^2(m)}$ for $n \geq 1$. By Proposition 27, $q_n \in L^2(\mathbb{R})$ so, by Cauchy-Schwartz inequality, $\overline{q_n}m \in L^1(\mathbb{R})$. Combining with the absolute convergence in $(\mathcal{C}_0(\mathbb{R}), || \cdot ||_{\infty})$ of the series of functions defining $S_t \cdot f$ and using that $(q_m)_{m\geq 1}$ is an orthonormal Hilbert basis of $L^2(m)$ by Proposition 27 we get for any $n \geq 1$,

$$\langle S_t.f, q_n \rangle_{L^2(m)} = \int_{\mathbb{R}} S_t.f(x)\overline{q_n(x)}m(x)dx = \sum_{m \ge 1} e^{-t\lambda_m^X} \langle f, q_m \rangle_{L^2(m)} \langle q_m, q_n \rangle_{L^2(m)} = e^{-t\lambda_n^X} \langle f, q_n \rangle_{L^2(m)}.$$
 (3.59)

17

We now make the restriction $f \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$ (and recall that $C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$). According to Proposition 1 the semigroup is Feller so we have $P_t f \in C_0(\mathbb{R})$. By Lemma 46 we have $P_t f \in L^2(m)$. We now compute $\langle P_t f, q_n \rangle_{L^2(m)}$ for $n \geq 1$. By Proposition 27, $q_n \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$. Therefore, using Lemma 49 we get for any $n \geq 1$,

$$\langle P_t.f, q_n \rangle_{L^2(m)} = \langle f, P_t.q_n \rangle_{L^2(m)} = e^{-t\lambda_n^{\Lambda}} \langle f, q_n \rangle_{L^2(m)},$$

where the last equality comes from $P_t q_n = e^{-t\lambda_n^X} q_n$, which follows from Proposition 27, (3.55) and the differentiation rule for the semigroup (see Section 1.6).

We thus get $\langle P_t.f, q_n \rangle_{L^2(m)} = \langle S_t.f, q_n \rangle_{L^2(m)}$ for all $n \ge 1$. Since $(q_n)_{n\ge 1}$ is an orthonormal Hilbert basis of $L^2(m)$ we get $P_t.f = S_t.f$ in $L^2(m)$ so, since m is positive on \mathbb{R} by (Condition 1), $P_t.f(x) = S_t.f(x)$ for almost every $x \in \mathbb{R}$. As both functions are continuous in x, (3.58) follows.

Remark 30. Let t > 0. Combining (3.59) with Parseval's identity we get $||S_t.f||_{L^2(m)} \leq ||f||_{L^2(m)}$ so the operator S_t from the above proof is a contraction of $L^2(m)$. Also, it satisfies $S_t(L^2(m)) \subset C_0(\mathbb{R})$ and it is continuous from $L^2(m)$ to $(C_0(\mathbb{R}), ||\cdot||_{\infty})$. Moreover, S_t coincides with P_t on the subspace $C_0(\mathbb{R}) \cap L^2(\mathbb{R})$ (in the sense that, for $f \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$, we have $P_t.f(x) = S_t.f(x)$ for all $x \in \mathbb{R}$).

The first eigenfunction is commonly called ground state and enjoys special properties. They are gathered in the following lemma that builds on Lemma 44 from Section 6.3 and they will allow to show the positivity of the constant $K_{m,\xi,r}(x)$ in Theorem 5, and to relate that constant with the infinite-volume Gibbs states of the spin system from Section 1.2.

Lemma 31. Assume that r > 0 and (Condition 1) and (Condition 2) hold. There is $\theta \in [0, 2\pi)$ such that $q_1 = e^{i\theta}|q_1|$ and we have $|q_1(x)| > 0$ for all $x \in \mathbb{R}$. Moreover, the eigenvalue λ_1^X of $-\mathcal{A}_{X^m,\xi,r}$ has multiplicity 1 (in other words, $\lambda_2^X > \lambda_1^X$).

Proof. Since R_r is a compact operator by Lemma 20, its largest eigenvalue λ_1^R has a finite multiplicity that we denote by N. From (3.55) we get $\lambda_1^X = \cdots = \lambda_N^X$ and $\lambda_k^X > \lambda_1^X$ for k > N.

Let us first show that $Span\{q_1, \ldots, q_N\}$ admits an orthonormal basis (for the inner product in $L^2(m)$) made of real eigenfunctions of operators P_t . For any $k \in \{1, \ldots, N\}$ we have $q_k \in C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$ by Proposition 27 so $Re(q_k), Im(q_k) \in C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$ and, by (1.3), $P_t.Re(q_k)$ and $P_t.Im(q_k)$ are real functions for any t > 0. We have $P_t.q_k = e^{-t\lambda_1^X}q_k$ from Proposition 27, (3.55) and the differentiation rule for the semigroup (see Section 1.6). We thus get that for any $t > 0, P_t.Re(q_k) + iP_t.Im(q_k) = P_t.q_k = e^{-t\lambda_1^X}q_k = e^{-t\lambda_1^X}Re(q_k) + ie^{-t\lambda_1^X}Im(q_k)$ so $P_t.Re(q_k) = e^{-t\lambda_1^X}Re(q_k)$ and $P_t.Im(q_k) = e^{-t\lambda_1^X}Im(q_k)$. Therefore $Re(q_k), Im(q_k) \in Span\{q_1, \ldots, q_N\}$ and $\{Re(q_1), Im(q_1), \ldots, Re(q_N), Im(q_N)\}$ is a spanning family of $Span\{q_1, \ldots, q_N\}$ that only contains real eigenfunctions of operators P_t . Extracting a linearly independent family and applying Gram,ÄiSchmidt process we obtain an orthonormal basis $\{\tilde{q}_1, \ldots, \tilde{q}_N\}$ of $Span\{q_1, \ldots, q_N\}$, where each \tilde{q}_k lies in $C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$ and is a real eigenfunction of all operators P_t . Each \tilde{q}_k is not identically 0 so, replacing \tilde{q}_k by $-\tilde{q}_k$ if necessary, we can assume that

$$\forall k \in \{1, \dots, N\}, \ \exists x_k \in \mathbb{R} \text{ s.t. } \tilde{q}_k(x_k) > 0.$$
(3.60)

Let us fix $k \in \{1, \ldots, N\}$. We have clearly $|\tilde{q}_k| \in C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$ so Proposition 29 yields that for all t > 0 and $x \in \mathbb{R}$, $P_t.|\tilde{q}_k|(x) = \sum_{n \ge 1} e^{-t\lambda_n^X} \langle |\tilde{q}_k|, q_n \rangle_{L^2(m)} q_n(x)$ where the series of functions converges absolutely in $(C_0(\mathbb{R}), \|\cdot\|_{\infty})$. Proceeding as in (3.59) we get $\langle P_t.|\tilde{q}_k|, q_n \rangle_{L^2(m)} = e^{-t\lambda_n^X} \langle |\tilde{q}_k|, q_n \rangle_{L^2(m)}$ for any $n \ge 1$. Therefore, by Parseval's identity we get

$$\|P_{t}.|\tilde{q}_{k}|\|_{L^{2}(m)}^{2} = \sum_{n\geq 1} e^{-2t\lambda_{n}^{X}} |\langle |\tilde{q}_{k}|, q_{n}\rangle_{L^{2}(m)}|^{2}.$$
(3.61)

Now note that for any $x \in \mathbb{R}$, $|P_t.\tilde{q}_k(x)| \le P_t.|\tilde{q}_k|(x)$ so $||P_t.\tilde{q}_k||_{L^2(m)} \le ||P_t.|\tilde{q}_k||_{L^2(m)}$. Using $||\tilde{q}_k||_{L^2(m)} = 1$, $P_t.\tilde{q}_k = e^{-t\lambda_1^X}\tilde{q}_k$, $||P_t.\tilde{q}_k||_{L^2(m)} \le ||P_t.|\tilde{q}_k||_{L^2(m)}$ and (3.61) we get

$$e^{-2t\lambda_1^X} = e^{-2t\lambda_1^X} \|\tilde{q}_k\|_{L^2(m)}^2 = \|P_t.\tilde{q}_k\|_{L^2(m)}^2 \le \|P_t.|\tilde{q}_k\|_{L^2(m)}^2 = \sum_{n\ge 1} e^{-2t\lambda_n^X} |\langle |\tilde{q}_k|, q_n\rangle_{L^2(m)}|^2.$$

Since $\||\tilde{q}_k\|\|_{L^2(m)} = \|\tilde{q}_k\|_{L^2(m)} = 1$ we have $\sum_{n\geq 1} |\langle |\tilde{q}_k|, q_n \rangle_{L^2(m)}|^2 = 1$. The above thus implies $\langle |\tilde{q}_k|, q_n \rangle_{L^2(m)} = 0$ for n > N. Therefore, $|\tilde{q}_k| \in Span\{q_1, \ldots, q_N\}$ so $|\tilde{q}_k|$ is an eigenfunction of P_t with eigenvalue $e^{-t\lambda_1^X}$, and so is $\tilde{q}_k + |\tilde{q}_k|$. By (3.60) and by continuity there is an open set $\mathcal{U} \subset \mathbb{R}$ such that $\tilde{q}_k(x) > 0$ for all $x \in \mathcal{U}$. Lemma 44 shows that $\mathbb{P}(X_x^{m,\xi,r}(1) \in \mathcal{U}) > 0$ for all $x \in \mathbb{R}$ so, since $\tilde{q}_k + |\tilde{q}_k|$ is positive on \mathcal{U} and non-negative on $\mathbb{R} \setminus \mathcal{U}$.

 $P_1.(\tilde{q}_k+|\tilde{q}_k|)(x) > 0.$ For any $x \in \mathbb{R}$ we have $\tilde{q}_k(x)+|\tilde{q}_k(x)| = e^{\lambda_1^X} P_1.(\tilde{q}_k+|\tilde{q}_k|)(x)$. We thus get $\tilde{q}_k(x)+|\tilde{q}_k(x)| > 0$ for all $x \in \mathbb{R}$ so there is no $x \in \mathbb{R}$ such that $\tilde{q}_k(x) < 0$. Therefore, \tilde{q}_k is positive on \mathcal{U} and non-negative on $\mathbb{R} \setminus \mathcal{U}$ so the same argument again shows that $\tilde{q}_k(x) > 0$ for all $x \in \mathbb{R}$. This and the orthonormality of $(\tilde{q}_k)_{k \in \{1,...,N\}}$ implies N = 1 so $\lambda_2^X > \lambda_1^X$ and $q_1 = e^{i\theta}\tilde{q}_1$ for some $\theta \in [0, 2\pi)$. This concludes the proof.

3.3. Decomposition of the survival probability of $X^{m,\xi,r}$. The following result is based on the decomposition from Proposition 29 and provides a representation for $\mathbb{P}(\zeta_x > t)$.

Proposition 32. Assume that r > 0 and (Condition 1) and (Condition 2) hold. We have for any t > 0,

$$\forall x \in \mathbb{R}, \ \mathbb{P}(\zeta_x > t) = \sum_{n \ge 1} e^{-t\lambda_n^X} \langle m, q_n \rangle_{L^2(\mathbb{R})} q_n(x), \tag{3.62}$$

where the series of functions in the right-hand side converges absolutely in $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$.

Proof. Let us fix t > 0. For M > 0 let

$$f_M(x) := \begin{cases} 1 \text{ if } x \in [-M, M], \\ 1 - (|x| - M) \text{ if } x \in (M, M + 1) \cup (-M - 1, -M), \\ 0 \text{ if } |x| \ge M + 1. \end{cases}$$
(3.63)

We have clearly $f_M \in \mathcal{C}_0(\mathbb{R}) \cap L^2(\mathbb{R})$ so, by Proposition 29, we get

$$\forall x \in \mathbb{R}, \ P_t.f_M(x) = \sum_{n \ge 1} e^{-t\lambda_n^X} \langle f_M, q_n \rangle_{L^2(m)} q_n(x).$$
(3.64)

For any $n \geq 1$, $\langle f_M, q_n \rangle_{L^2(m)} = \int_{\mathbb{R}} f_M(x) \overline{q_n(x)} m(x) dx$. For every $x \in \mathbb{R}$, $f_M(x)$ converges to 1 as M goes to infinity. Moreover $|f_M(x)| \leq 1$ and $\overline{q_n}m \in L^1(\mathbb{R})$ by Cauchy-Schwartz inequality since $q_n \in L^2(\mathbb{R})$ by Proposition 27 and $m \in L^2(\mathbb{R})$ by (Condition 1). We thus get by dominated convergence that $\langle f_M, q_n \rangle_{L^2(m)}$ converges to $\langle m, q_n \rangle_{L^2(\mathbb{R})}$ as M goes to infinity. Then, using Cauchy-Schwartz inequality and Proposition 27, $|\langle f_M, q_n \rangle_{L^2(m)}| \leq \int_{\mathbb{R}} |q_n(x)|m(x)dx \leq ||q_n||_{L^2(\mathbb{R})} \times ||m||_{L^2(\mathbb{R})} \leq ||m||_{L^2(\mathbb{R})} \sqrt{C/\lambda_n^R}$ for some C > 0. This and Lemma 28 (together with (3.55)) allow to use dominated convergence to show that the series in the right-hand side of (3.64) converges to the right-hand side of (3.62) as M goes to infinity. Then, for any $x \in \mathbb{R}$ we see from (1.3) and monotone convergence that the left-hand side of (3.64) converges to $\mathbb{P}(\zeta_x > t)$ as M goes to infinity. (3.62) follows. Since $q_n \in C_0(\mathbb{R})$ (by Proposition 27) the claim about absolute convergence in $(C_0(\mathbb{R}), \|\cdot\|_{\infty})$ follows from $|\langle q_n, m \rangle_{L^2(\mathbb{R})}| \leq ||m||_{L^2(\mathbb{R})} \sqrt{C/\lambda_n^R}$ and Lemma 28 (together with (3.55)).

The following proposition provides an equivalent for $\mathbb{P}(\zeta_x > t)$ as t goes to infinity.

Proposition 33. Assume that r > 0 and (Condition 1) and (Condition 2) hold. For every $x \in \mathbb{R}$, we have $\langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x) > 0$ and

$$\mathbb{P}(\zeta_x > t) \underset{t \to \infty}{\sim} \langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x) e^{-t\lambda_1^X}.$$
(3.65)

The proof builds on the representation of $\mathbb{P}(\zeta_x > t)$ from Proposition 32 and on Lemma 31.

Proof. From Proposition 32 we get

$$\forall x \in \mathbb{R}, \ e^{t\lambda_1^X} \mathbb{P}(\zeta_x > t) - \langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x) = \sum_{n>1} e^{-t(\lambda_n^X - \lambda_1^X)} \langle m, q_n \rangle_{L^2(\mathbb{R})} q_n(x).$$
(3.66)

By Lemma 31 we have $\lambda_n^X - \lambda_1^X > 0$ for all n > 1 so each term in the above series converges to 0 as t goes to infinity. Since λ_n^R converges to 0 as n goes to infinity (see Lemma 20) we see from (3.55) that $\lambda_n^X \sim \lambda_n^X - \lambda_1^X$ as n goes to infinity. Therefore, reasoning as in the proof of Lemma 28 we get $\sum_{n \ge 1} (\lambda_n^X)^{1/2} e^{-(\lambda_n^X - \lambda_1^X)} (||q_n||_{\infty} \vee 1) < \infty$. Since moreover $|\langle m, q_n \rangle_{L^2(\mathbb{R})}| \le ||m||_{L^2(\mathbb{R})} (Cv_{\xi}^r(0)\lambda_n^X)^{1/2}$ (by Cauchy-Schwartz inequality, Proposition 27 and (3.55)), we get by dominated convergence that the right-hand side of (3.66) converges to 0 as t goes to infinity. Therefore $e^{t\lambda_1^X} \mathbb{P}(\zeta_x > t)$ converges to $\langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x)$ as t goes to infinity. We are only left to show that $\langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x) > 0$. For this recall that, by Lemma 31 and the differentiation rule for the semigroup (see Section 1.6), $|q_1|$ is an eigenfunction of P_t associated with the eigenvalue $e^{-t\lambda_1^X}$ and is positive. We thus get

$$0 < |q_1(x)| = e^{t\lambda_1^X} P_t . |q_1|(x) \le ||q_1||_{\infty} e^{t\lambda_1^X} P_t . 1(x) = ||q_1||_{\infty} e^{t\lambda_1^X} \mathbb{P}(\zeta_x > t) \xrightarrow[t \to \infty]{} ||q_1||_{\infty} \langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x)$$

This proves that $\langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x) > 0$, concluding the proof.

3.4. Relation with traces of compositions of R_r . The following proposition connects the traces of compositions of R_r with the partition function Z_n .

Proposition 34. Assume that r > 0 and (Condition 1) and (Condition 2) hold. Let Z_n be the partition function defined by (1.7) with the choice $V_{int} := -\log(v_{\xi}^r(\cdot))$ and $U_{mass} := -\log(m(\cdot))$ and let \mathcal{E} be the associated normalized free energy (defined in (1.8)). For any $n \ge 2$, Z_n is well-defined and we have $Tr(R_r^n) = Z_n/v_{\xi}^r(0)^n$. Moreover, \mathcal{E} is well-defined and we have $\mathcal{E} = \log(\lambda_1^X)$.

Proof. Fix $n \ge 2$. The finiteness of the integral defining Z_n in (1.7) is a consequence of the finiteness of the integrals in (2.42), in the proof of Lemma 23. Z_n is thus well-defined. Recall that the well-definedness of $Tr(R_r^n)$ is proved by Lemma 23. Using (2.44) from the proof of Lemma 23 and the definition of P_y^r , we get

$$Tr(R_{r}^{n}) = \int_{\mathbb{R}^{n}} m(y_{1}) \dots m(y_{n}) \langle \phi_{y_{1}}^{r}, \phi_{y_{2}}^{r} \rangle_{L^{2}((-1,1))} \times \dots \times \langle \phi_{y_{n-1}}^{r}, \phi_{y_{n}}^{r} \rangle_{L^{2}((-1,1))}$$

$$\times \langle \phi_{y_{n}}^{r}, \phi_{y_{1}}^{r} \rangle_{L^{2}((-1,1))} dy_{1} \dots dy_{n}.$$
(3.67)

Plugging (2.41) into (3.67) and comparing with (1.7) yields that $Tr(R_r^n) = Z_n/v_{\xi}^r(0)^n$ for each $n \ge 2$. We have clearly $Z_n > 0$ for any $n \ge 2$. Then, combining $Tr(R_r^n) = Z_n/v_{\xi}^r(0)^n$ with (2.38) we get that \mathcal{E} exists and equals $-\log(v_{\xi}^r(0)\lambda_1^R)$. Combining with (3.55) we obtain $\mathcal{E} = \log(\lambda_1^R)$. This concludes the proof.

The following proposition will be the main ingredient in the proof of (1.18).

Proposition 35. Assume that r > 0 and (Condition 1') and (Condition 2) hold. We have $1/||m||_{L^1(\mathbb{R})} v_{\xi}^r(0) \le \lambda_1^X \le ||m||_{L^1(\mathbb{R})} v_{\xi}^r(0)/Z_2$, with Z_n as in Proposition 34.

Proof. Using Lemma 25 and that all eigenvalues of R_r are positive (from Lemma 20) we get

$$\begin{split} \|m\|_{L^{1}(\mathbb{R})} &= Tr(R_{r}) = \sum_{k \ge 1} \lambda_{k}^{R} \ge \lambda_{1}^{R} = \lambda_{1}^{R} \times \frac{Tr(R_{r})}{\|m\|_{L^{1}(\mathbb{R})}} = \frac{1}{\|m\|_{L^{1}(\mathbb{R})}} \sum_{k \ge 1} \lambda_{1}^{R} \lambda_{k}^{R} \\ &\ge \frac{1}{\|m\|_{L^{1}(\mathbb{R})}} \sum_{k \ge 1} (\lambda_{k}^{R})^{2} = \frac{Tr(R_{r}^{2})}{\|m\|_{L^{1}(\mathbb{R})}} = \frac{Z_{2}}{\|m\|_{L^{1}(\mathbb{R})} \times v_{\xi}^{r}(0)^{2}}, \end{split}$$

where the last equality comes from Proposition 34. Combining the bound $||m||_{L^1(\mathbb{R})} \ge \lambda_1^R \ge Z_2/||m||_{L^1(\mathbb{R})} v_{\xi}^r(0)^2$ with (3.55) we get $1/||m||_{L^1(\mathbb{R})} v_{\xi}^r(0) \le \lambda_1^X \le ||m||_{L^1(\mathbb{R})} v_{\xi}^r(0)/Z_2$.

3.5. Infinite-volume Gibbs states. Recall the sequence of probability measures $(L_n^k)_{n\geq 2\vee k}$ defined in Section 1.2. The following proposition relates the first eigenfunction of the semigroup $(P_t)_{t\geq 0}$ with the infinite-volume Gibbs states of the spin system from Section 1.2.

Proposition 36. Assume that r > 0 and (Condition 1) and (Condition 2) hold. Under the choice $V_{int} := -\log(v_{\xi}^r(\cdot))$ and $U_{mass} := -\log(m(\cdot))$, the infinite-volume Gibbs states \mathcal{L}_k defined in (1.11) is well-defined for any $k \ge 1$ and we have

$$\mathcal{L}_{k}(dy_{1}\dots dy_{k}) = (\lambda_{1}^{X})^{k-1} \times m(y_{1})\dots m(y_{k}) \times |q_{1}(y_{1})| v_{\xi}^{r}(y_{2}-y_{1})\dots v_{\xi}^{r}(y_{k}-y_{k-1})|q_{1}(y_{k})| dy_{1}\dots dy_{k}.$$
 (3.68)
In particular we have

$$\mathcal{L}_1(dy) = m(y) |q_1(y)|^2 dy.$$
(3.69)

Moreover, for any $y \in \mathbb{R}$ we have

$$l_1^n(y) \underset{n \to \infty}{\longrightarrow} m(y) |q_1(y)|^2. \tag{3.70}$$

(3.69) says that, if we consider a fixed particle in the infinite volume limit of the spin system from Section 1.2, its interactions with the infinitely many other particles in the system are coded in the factor $|q_1(y)|^2$.

Proof. Let $k \ge 1$ and $f_1, \ldots, f_k \in C_b(\mathbb{R})$ be real non-negative functions. Proposition 34 shows that, for $n \ge 2 \lor k$, Z_n is finite so, in particular, the probability measures $(L_n^k)_{n\ge 2\lor k}$ are well-defined. We study the convergence of the quantity $L_n^k(f_1, \ldots, f_k)$ defined in (1.9). For this, when $j \in \{1, \ldots, k\}$, let us define the operator $R_{r,j}$: $L^2((-1,1)) \to L^2((-1,1))$ by $R_{r,j}.h := \int_{\mathbb{R}} f_j(y)m(y)P_y^r.hdy$, where the integral is in the sense of Bochner. By (Condition 1) we have $mf_j \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$ so the operator $R_{r,j}$ also satisfies the properties established for R_r in the first two points of Lemma 20 (in particular it is well-defined). The reasoning from the proof of Lemma 23 can be repeated and yields that for any $n \ge 2 \lor k$, and any Hilbert basis $(u_j)_{j\ge 1}$ of $L^2((-1,1))$ we have $\sum_{j\ge 1} |\langle R_{r,1} \ldots R_{r,k}.R_r^{n-k}.u_j, u_j \rangle_{L^2((-1,1))}| < \infty$ and the quantity $\sum_{j\ge 1} \langle R_{r,1} \ldots R_{r,k}.R_r^{n-k}.u_j, u_j \rangle_{L^2((-1,1))}|$

not depend on the choice of the Hilbert basis $(u_j)_{j\geq 1}$ so we denote it by $Tr(R_{r,1}\ldots R_{r,k}.R_r^{n-k})$. Recall that $(a_j)_{j\geq 1}$ is the orthonormal Hilbert basis of $L^2((-1,1))$ consisting of eigenfunctions of R_r , chosen a little before (3.51). Since $|\langle R_{r,1}\ldots R_{r,k}.a_j, a_j \rangle_{L^2((-1,1))}| \leq ||R_{r,1}\ldots R_{r,k}||$ for all $j \geq 1$, the reasoning from the proof of Lemma 23, together with Remark 24, yields that

$$\frac{Tr(R_{r,1}\dots R_{r,k}.R_r^{n-k})}{(\lambda_1^R)^{n-k}} \xrightarrow[n \to \infty]{} \langle R_{r,1}\dots R_{r,k}.a_1, a_1 \rangle_{L^2((-1,1))}.$$
(3.71)

Proceeding as in (2.44) from the proof of Lemma 23 we get for any $n \ge 2 \lor k$,

$$Tr(R_{r,1}\dots R_{r,k}.R_r^{n-k}) = \int_{\mathbb{R}^n} f_1(y_1)\dots f_k(y_k)m(y_1)\dots m(y_n) \langle P_{y_n}^r \dots P_{y_2}^r.\phi_{y_1}^r, \phi_{y_1}^r \rangle_{L^2((-1,1))} dy_1\dots dy_n.$$

Then using the definition of P_y^r and (2.41) we obtain

$$Tr(R_{r,1}...R_{r,k}.R_r^{n-k}) = v_{\xi}^r(0)^{-n} \int_{\mathbb{R}^n} f_1(y_1)...f_k(y_k)m(y_1)...m(y_n)v_{\xi}^r(y_2 - y_1) \times \cdots \times v_{\xi}^r(y_n - y_{n-1}) \times v_{\xi}^r(y_1 - y_n)dy_1...dy_n.$$

Comparing with (1.9) we get that for any $n \ge 2 \lor k$,

$$Tr(R_{r,1}...R_{r,k}.R_r^{n-k}) = Z_n \times L_n^k(f_1,...,f_k) / v_{\xi}^r(0)^n.$$
(3.72)

Then the combination of (3.72), (3.71), Proposition 34, Lemma 23 and Remark 24 yields

$$L_n^k(f_1,\ldots,f_k) \xrightarrow[n\to\infty]{} \frac{1}{(\lambda_1^R)^k} \langle R_{r,1}\ldots R_{r,k}.a_1,a_1 \rangle_{L^2((-1,1))}.$$
(3.73)

Using the second point of Lemma 20 (but for the $R_{r,j}$'s in place of R_r), the identities (2.40), (2.41) and (3.51), Lemma 31, and (3.55) we get that the right-hand side of (3.73) equals

$$\frac{1}{(\lambda_1^R)^k} \int_{\mathbb{R}^k} (f_1 m)(y_1) \dots (f_k m)(y_k) \langle P_{y_1}^r \dots P_{y_k}^r.a_1, a_1 \rangle_{L^2((-1,1))} dy_1 \dots dy_k \tag{3.74}$$

$$= \frac{1}{(\lambda_1^R)^k} \int_{\mathbb{R}^k} (f_1 m)(y_1) \dots (f_k m)(y_k) (\lambda_1^R)^{1/2} \overline{q_1(y_1)} \frac{v_{\xi}^r(y_2 - y_1)}{v_{\xi}^r(0)} \dots \frac{v_{\xi}^r(y_k - y_{k-1})}{v_{\xi}^r(0)} (\lambda_1^R)^{1/2} q_1(y_k) dy_1 \dots dy_k \tag{3.74}$$

$$= (\lambda_1^X)^{k-1} \int_{\mathbb{R}^k} (f_1 m)(y_1) \dots (f_k m)(y_k) |q_1(y_1)| v_{\xi}^r(y_2 - y_1) \dots v_{\xi}^r(y_k - y_{k-1}) |q_1(y_k)| dy_1 \dots dy_k.$$

Combining with (3.73) we get

$$L_{n}^{k}(f_{1},\ldots,f_{k}) \xrightarrow[n\to\infty]{} (\lambda_{1}^{X})^{k-1} \int_{\mathbb{R}^{k}} (f_{1}m)(y_{1})\ldots(f_{k}m)(y_{k})|q_{1}(y_{1})|v_{\xi}^{r}(y_{2}-y_{1})\ldots v_{\xi}^{r}(y_{k}-y_{k-1})|q_{1}(y_{k})|dy_{1}\ldots dy_{k}.$$
(3.75)

According to Proposition 27, $(q_j)_{j\geq 1}$ is an orthonormal family of $L^2(m)$, so $\int_{\mathbb{R}} m(y)|q_1(y)|^2 dy = 1$. The measure appearing in the right-hand side of (3.75) is thus a probability measure when k = 1. For k > 1, the calculation in the proof of Proposition 13 below will show that the measure in the right-hand side of (3.75) is the law of a random vector, so in particular it is a probability measure. This and (3.75) thus imply that for each k > 1 (resp. k = 1), we have the convergence in law of L_n^k toward a limit \mathcal{L}_k with expression specified in (3.68) (resp. (3.69)). This concludes the proof of the first part of the proposition. To prove (3.70), fix $y \in \mathbb{R}$ and repeat the above argument leading to (3.75) (with k = 1), but with $L_n^1(f_1)$ replaced by $l_1^n(y)$ and $R_{r,1}$ replaced by $\tilde{R}_r := m(y)P_y^r$ (formally, this amounts to replacing the function f_1 by the distribution δ_y). This concludes the proof.

4. Proof of main results on $X^{m,\xi,r}$ from Section 1.3

4.1. Asymptotic of survival probability: Proof of Theorem 5. Proposition 33 shows that (1.15) holds with $\gamma_{m,\xi,r} = \lambda_1^X$ and $K_{m,\xi,r}(x) = \langle m, q_1 \rangle_{L^2(\mathbb{R})} q_1(x)$, and that $K_{m,\xi,r}(x) > 0$. We recall that the positivity of λ_1^X comes from the combination of Lemma 20 with (3.55). Proposition 34 (together with $\gamma_{m,\xi,r} = \lambda_1^X$) yields the well-definedness of the partition function Z_n and of the normalized free energy \mathcal{E} , and (1.16). Using Lemma 31 we see that we have $K_{m,\xi,r}(x) = \langle m, |q_1| \rangle_{L^2(\mathbb{R})} |q_1(x)|$ so (1.17) and ℓ_1 being the point-wise limit of $(l_1^n)_{n\geq 2}$ both follow from Proposition 36. In particular, $\ell_1/m = |q_1|^2$ so the claims that $\ell_1/m \in \mathcal{C}_0(\mathbb{R}) \cap L^1(\mathbb{R})$ and that ℓ_1 is positive on \mathbb{R} follow respectively from Proposition 27 and Lemma 31. Under (Condition 1'), Proposition 35 (together with $\gamma_{m,\xi,r} = \lambda_1^X$) shows that (1.18) holds. This completes the proof. Remark 37. Once (1.15) has been established, the moment approach discussed in Section 1.5 yields an alternative proof of the lower bound in (1.18). Indeed, using $\zeta_x = A_x(e_r)$ and Khas'minskii's condition from [13] we get $\mathbb{E}[e^{\theta\zeta_x}] < \infty$ for all $\theta < 1/\|G_m\|_{\infty}$ where $G_m(y) := \int_{\mathbb{R}} v_{\xi}^r(y-z)m(z)dz$. By Lemma 50 and (Condition 1') we get $\|G_m\|_{\infty} \leq \|m\|_{L^1(\mathbb{R})}v_{\xi}^r(0)$. Therefore, by Chernoff's inequality we get $\mathbb{P}(\zeta_x > t) << e^{-\theta t}$ for all $\theta < 1/\|m\|_{L^1(\mathbb{R})}v_{\xi}^r(0)$. Combining with (1.15) we obtain $1/\|m\|_{L^1(\mathbb{R})}v_{\xi}^r(0) \leq \gamma_{m,\xi,r}$, which is the lower bound in (1.18).

4.2. Asymptotic behavior of $\gamma_{m,\xi,r}$: Proof of Corollary 7. We first justify the following lemma that determines the asymptotic behavior of $v_{\xi}^{r}(0)$ when r goes to 0.

Lemma 38. Assume that (Condition 2) holds, so that the potential density $v_{\xi}^{r}(\cdot)$ exists (see Lemma 50). We assume that there are $\alpha \geq 1$ and c > 0 such that $-\psi_{\xi}(y) \sim c|y|^{\alpha}$ as y goes to 0. Then we have

$$v_{\xi}^{r}(0) \underset{r \to 0}{\sim} r^{-\frac{\alpha-1}{\alpha}} \frac{1}{\pi c^{\frac{1}{\alpha}}} \int_{0}^{\infty} \frac{1}{1+u^{\alpha}} du \quad \text{if } \alpha > 1,$$
(4.76)

$$v_{\xi}^{r}(0) \underset{r \to 0}{\sim} \frac{1}{c\pi} \log\left(\frac{1}{r}\right) \quad if \ \alpha = 1.$$

$$(4.77)$$

Remark 39. If ξ is a symmetric α -stable Lévy process on \mathbb{R} with $\alpha > 1$ then $-\psi_{\xi}(y) = c|y|^{\alpha}$ for some c > 0 (see for example Theorem 14.15 in [46]) and $v_{\xi}^{r}(0) = r^{-(\alpha-1)/\alpha}v_{\xi}^{1}(0)$ (see for example the discussion after Theorem II.19 in [4]) so (4.76) is even an equality and Lemma 38 allows to identify the constant $v_{\xi}^{1}(0)$ with $(\pi c^{1/\alpha})^{-1} \int_{0}^{\infty} 1/(1+u^{\alpha}) du$.

Proof of Lemma 38. For $\delta > 0$ let

$$q_{\delta}(r) := \frac{1}{2\pi} \int_{-\delta}^{\delta} \frac{1}{r - \psi_{\xi}(z)} dz, \quad \text{and} \quad C_{\delta} := \frac{1}{2\pi} \int_{[-\delta, \delta]^c} \frac{1}{-\psi_{\xi}(z)} dz.$$
(4.78)

Note that (Condition 2) ensures that $C_{\delta} < \infty$. From Lemma 50 we see that, for any $\delta > 0$ and r > 0, $q_{\delta}(r) \le v_{\xi}^{r}(0) \le q_{\delta}(r) + C_{\delta}$. From the assumption $-\psi_{\xi}(y) \sim c|y|^{\alpha}$, $q_{\delta}(r)$ and $v_{\xi}^{r}(0)$ converge to infinity as r goes to zero. We thus have that for any $\delta > 0$,

$$v_{\xi}^r(0) \underset{r \to 0}{\sim} q_{\delta}(r).$$
 (4.79)

Since $-\psi_{\xi}(y) \sim c|y|^{\alpha}$, we can write $-\psi_{\xi}(y) = c|y|^{\alpha}B(y)$ where B(y) converges to 1 as y goes to 0. Let us fix $\epsilon \in (0, 1)$. Then, there is $\delta > 0$ such that $B(y) \in (1 - \epsilon, 1 + \epsilon)$ for all $y \in (0, \delta]$. From now onward we fix such a $\delta > 0$. Using (4.79), the symmetry of $\psi_{\xi}(\cdot)$, and $-\psi_{\xi}(y) = c|y|^{\alpha}B(y)$, we get

$$v_{\xi}^{r}(0) \sim_{r \to 0} \frac{1}{\pi} \int_{0}^{\delta} \frac{1}{r + cy^{\alpha} B(y)} dy.$$
 (4.80)

Then,

$$r^{\frac{\alpha-1}{\alpha}} \int_0^{\delta} \frac{1}{r+cy^{\alpha}B(y)} dy = r^{-\frac{1}{\alpha}} \int_0^{\delta} \frac{1}{1+\frac{c}{r}y^{\alpha}B(y)} dy = c^{-\frac{1}{\alpha}} \int_0^{\delta(c/r)^{\frac{1}{\alpha}}} \frac{1}{1+u^{\alpha}B((r/c)^{\frac{1}{\alpha}}u)} du.$$

We thus get that

$$c^{-\frac{1}{\alpha}} \int_0^{\delta(c/r)^{\frac{1}{\alpha}}} \frac{1}{1 + (1+\epsilon)u^{\alpha}} du \le r^{\frac{\alpha-1}{\alpha}} \int_0^{\delta} \frac{1}{r + cy^{\alpha}B(y)} dy \le c^{-\frac{1}{\alpha}} \int_0^{\delta(c/r)^{\frac{1}{\alpha}}} \frac{1}{1 + (1-\epsilon)u^{\alpha}} du$$

Let us now assume that $\alpha > 1$. Combining the above with (4.80) we get

$$\frac{c^{-\frac{1}{\alpha}}}{\pi} \int_0^\infty \frac{1}{1 + (1+\epsilon)u^\alpha} du \le \liminf_{r \to 0} r^{\frac{\alpha-1}{\alpha}} v_\xi^r(0) \le \limsup_{r \to 0} r^{\frac{\alpha-1}{\alpha}} v_\xi^r(0) \le \frac{c^{-\frac{1}{\alpha}}}{\pi} \int_0^\infty \frac{1}{1 + (1-\epsilon)u^\alpha} du.$$

Letting ϵ go to 0 and using monotone convergence we get (4.76). If $\alpha = 1$ then a similar reasoning yields (4.77).

We now assume that the assumptions from Corollary 7 hold and prove the corollary. We first show that, under the choice $V_{int} := -\log(v_{\xi}^r(\cdot))$ and $U_{mass} := -\log(m(\cdot))$, we have

$$Z_2 \underset{r \to 0}{\sim} \|m\|_{L^1(\mathbb{R})}^2 \times v_{\xi}^r(0)^2.$$
(4.81)

Recall from Remark 4 that \hat{m} denotes the characteristic function of m(x)dx. Let us fix $\epsilon \in (0, 1)$. Since $\hat{m}(0) = \|m\|_{L^1(\mathbb{R})}$ and \hat{m} is continuous at 0, there is $\delta > 0$ such that $|\hat{m}(x)|^2 / \|m\|_{L^1(\mathbb{R})}^2 \in (1 - \epsilon, 1]$ for all $x \in [-2\delta, 2\delta]$. According to (1.14) we have

$$\frac{Z_2}{\|m\|_{L^1(\mathbb{R})}^2} \ge \frac{1}{(2\pi)^2} \int_{-\delta}^{\delta} \int_{-\delta}^{\delta} \frac{1-\epsilon}{(r-\psi_{\xi}(z_1)) \times (r-\psi_{\xi}(z_2))} dz_1 dz_2 = (1-\epsilon)q_{\delta}(r)^2, \tag{4.82}$$

where $q_{\delta}(r)$ is defined as in (4.78). Since $|\hat{m}(x)|^2 \leq ||m||^2_{L^1(\mathbb{R})}$ for all x,

$$\frac{Z_2}{\|m\|_{L^1(\mathbb{R})}^2} \le \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} \frac{1}{(r - \psi_{\xi}(z_1)) \times (r - \psi_{\xi}(z_2))} dz_1 dz_2 = v_{\xi}^r(0)^2.$$
(4.83)

Since $q_{\delta}(r)$ and $v_{\varepsilon}^{r}(0)$ converges to infinity as r goes to infinity we get from (4.82), (4.83) and (4.79) that

$$1 - \epsilon \le \liminf_{r \to 0} \frac{Z_2}{\|m\|_{L^1(\mathbb{R})}^2 \times v_{\xi}^r(0)^2} \le \limsup_{r \to 0} \frac{Z_2}{\|m\|_{L^1(\mathbb{R})}^2 \times v_{\xi}^r(0)^2} \le 1.$$

Since ϵ can be chosen arbitrary small we get (4.81).

The combination of (4.81) with (1.18) yields

$$\gamma_{m,\xi,r} \sim_{r \to 0} \frac{1}{\|m\|_{L^1(\mathbb{R})}} \times \frac{1}{v_{\xi}^r(0)}.$$
(4.84)

The combination of (4.84) with Lemma 38 yields (1.20) and (1.21), completing the proof.

Remark 40. The assumption "there are $\alpha \geq 1$ and c > 0 such that $-\psi_{\xi}(y) \sim c|y|^{\alpha}$ as y goes to 0" implies that the un-killed Lévy process ξ is recurrent (see for example Theorem 37.5 in [46]). If we only assume that (Condition 1') and (Condition 2) hold and that ξ is recurrent (which occurs, in particular, if $\mathbb{E}[|\xi(1)|] < \infty$, see Theorem 36.7 in [46]), then (4.84) is still satisfied, which can be seen as a weak version of Corollary 7. Indeed, Theorem 37.5 in [46] shows that $q_{\delta}(r)$ converges to infinity as r goes to zero so (4.79) still holds true. Then the above proof of (4.81) still holds and we get that (4.84) holds true.

4.3. Further spectral properties of $X^{m,\xi,r}$: Proof of Theorem 11. For t > 0, let S_t be the operator from Remark 30. According to that remark, S_t is a contraction of $L^2(m)$ that satisfies $S_t(L^2(m)) \subset C_0(\mathbb{R})$ and that is continuous from $L^2(m)$ to $(C_0(\mathbb{R}), \|\cdot\|_{\infty})$. For any $x \in \mathbb{R}$ we define the linear forms $p_{t,x} : B_b(\mathbb{R}) \to \mathbb{C}$ and $s_{t,x} : L^2(m) \to \mathbb{C}$ by $p_{t,x}.f = P_t.f(x)$ and $s_{t,x}.f = S_t.f(x)$. By (1.3), $p_{t,x}$ corresponds to the restriction to \mathbb{R} of the finite measure $\mathbb{P}(X_x^{m,\xi,r}(t) \in dz)$. Since $s_{t,x}$ is continuous, by Riesz representation theorem, there is $h_{t,x} \in L^2(m)$ such that, for $f \in L^2(m)$, $s_{t,x}.f = \langle f, h_{t,x} \rangle_{L^2(m)}$. According to Remark 30, $s_{t,x}$ coincides with $p_{t,x}$ on $C_0(\mathbb{R}) \cap L^2(\mathbb{R})$ so it is a positive linear form on that subspace. Since any non-negative function of $L^2(m)$ can be approximated by a sequence of non-negative functions from $C_0(\mathbb{R}) \cap L^2(\mathbb{R})$, $s_{t,x}$ is a positive linear form on $L^2(m)$ so the function $mh_{t,x}$ is real and almost everywhere non-negative. Moreover, by (Condition 1), for any compact set $K \subset \mathbb{R}$ we have $\mathbf{1}_K \in L^2(m)$ so $\int_K h_{t,x}(z)m(z)dz = \langle \mathbf{1}_K, h_{t,x}\rangle_{L^2(m)} < \infty$. Therefore, $s_{t,x}$ corresponds to a sigma-finite measure on \mathbb{R} . Since $p_{t,x}$ and $s_{t,x}$ coincide on the space $C_0(\mathbb{R}) \cap L^2(\mathbb{R})$, the corresponding measures on \mathbb{R} are equal. Therefore, for any $f \in \mathcal{B}_b(\mathbb{R}) \cap L^2(m)$ and $x \in \mathbb{R}$ we have $P_t.f(x) = S_t.f(x)$. This shows that S_t is indeed an extension of P_t on $L^2(m)$. Since $\mathcal{B}_b(\mathbb{R}) \cap L^2(m)$ is dense in $L^2(m)$, such a contractile extension is unique. Using (3.59) and Parseval's identity we get

$$||S_t \cdot f - f||^2_{L^2(m)} = \sum_{n \ge 1} (1 - e^{-t\lambda_n^X})^2 |\langle f, q_n \rangle_{L^2(m)}|^2.$$

Since $\sum_{n\geq 1} |\langle f, q_n \rangle_{L^2(m)}|^2 = ||f||^2_{L^2(m)} < \infty$ we get by dominated convergence that the right-hand side converges to 0 as t goes to 0. This shows that S_t is strongly continuous in $L^2(m)$, concluding the proof of the first claim.

Let $h_n := q_n$ for $n \ge 2$ and $h_1 := |q_1|$ where $(q_n)_{n\ge 1}$ is the family of functions provided by Proposition 27. Note from Lemma 31 that $h_1 = e^{-i\theta}q_1$ for some $\theta \in [0, 2\pi)$. By Proposition 27, $(h_n)_{n\ge 1}$ is an orthonormal Hilbert basis of $L^2(m)$ such that for any $n \ge 1$, $h_n \in C_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$. Proposition 27 and (3.55), together with the differentiation rule for the semigroup (see Section 1.6) yield that $P_t.h_n = e^{-t\lambda_n^X}h_n$ for any $n \ge 1$. The combination of Lemma 20 and (3.55) yields $\lambda_1^X > 0$ and that $(\lambda_n^X)_{n\ge 1}$ is non-decreasing. (1.22) follows from the definition of $S_t.f(x)$ in the proof of Proposition 29 and from the fact, shown above, that the unique contractile extension of P_t to $L^2(m)$ equals S_t . This proves the second claim. Lemma 31 yields $\lambda_2^X > \lambda_1^X$. Finally, the two equalities from (1.23) have already been proved in Section 4.1. This proves the third claim. Under (Condition 1') we have $m \in L^1(\mathbb{R})$ so $\mathcal{B}_b(\mathbb{R}) \subset L^2(m)$ and the strong Feller property follows from the first claim.

4.4. **Proof of Example 16.** Let us set $f(x) := \sqrt{2/3}(1+|x|)e^{-|x|}$ then we have clearly $f \in \mathcal{C}_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$ and $||f||_{L^2(m)} = 1$. One can also see that $f \in \mathcal{C}_0^2(\mathbb{R})$ (one can check that f and f' have right and left derivatives at 0 and that those right and left derivatives at 0 coincide). By Theorem 31.5 of [46] we get $f \in \mathcal{D}(\mathcal{A}_{\xi})$ and $\mathcal{A}_{\xi}f = f''/2$. Since $\mathcal{D}(\mathcal{A}_{\xi}) = \mathcal{D}(\mathcal{A}_{\xi r})$ and $\mathcal{A}_{\xi r}f = \mathcal{A}_{\xi}f - rf$ we get $f \in \mathcal{D}(\mathcal{A}_{\xi r})$ and, after a straightforward calculation, $\frac{1}{m}\mathcal{A}_{\xi r}f = -f$. In particular we have $\frac{1}{m}\mathcal{A}_{\xi r}f \in \mathcal{C}_0(\mathbb{R})$ so by Lemma 45 we get $f \in \mathcal{D}(\mathcal{A}_{Xm,\xi,r})$ and $\mathcal{A}_{Xm,\xi,r}f = -f$. By the differentiation rule for the semigroup (see Section 1.6) we get $P_t.f = e^{-t}f$ for all $t \geq 0$. We have $P_t.h_1 = e^{-t\lambda_1^X}h_1$ from Theorem 11. If we had $\lambda_1^X \neq 1$ then f and h_1 would be eigenfunctions of P_t associated with different eigenvalues so, applying Lemma 49 with f and h_1 , we would get $\langle f, h_1 \rangle_{L^2(m)} = 0$. Since f and h_1 are both positive on \mathbb{R} (for h_1 this comes from (1.23) and Theorem 5) we have $\langle f, h_1 \rangle_{L^2(m)} > 0$. Therefore $\lambda_1^X = 1$. By Theorem 11, the eigenspace of λ_1^X has dimension 1. Since both f and h_1 lie in this eigenspace, $\|f\|_{L^2(m)} = 1 = \|h_1\|_{L^2(m)}$ and f and h_1 are both positive, we get $f = h_1$.

5. Ground state transform of $X^{m,\xi,r}$ and relation with the spin system

5.1. Existence and uniqueness of $\tilde{X}^{m,\xi,r}$: Proof of Proposition 12. Let us first justify that $(\tilde{p}_t(x,dy))_{t\geq 0,x\in\mathbb{R}}$ defines a transition function on \mathbb{R} . From Theorem 11 we have $h_1 \in C_0(\mathbb{R})$ and $P_t.h_1 = e^{-t\lambda_1^X}h_1$. We thus deduce from (1.24) that, for any $t \geq 0$ and $x \in \mathbb{R}$, $\tilde{p}_t(x,dy)$ is a probability measure on \mathbb{R} . Since h_1 is continuous on \mathbb{R} and $X^{m,\xi,r}$ is a Feller process, we easily get that for any Borel set $B \subset \mathbb{R}$, the function $(x \mapsto \tilde{p}_t(x,B))$ is measurable for all $t \geq 0$. Finally, the semigroup property for $(P_t)_{t\geq 0}$ (proved in Lemma 43) easily leads to the Chapman-Kolmogorov relation $\int_{\mathbb{R}} \tilde{p}_t(x,dz)\tilde{p}_s(z,dy) = \tilde{p}_{t+s}(x,dy)$, for any $t,s \geq 0$ and $x \in \mathbb{R}$. Therefore, $(\tilde{p}_t(x,dy))_{t\geq 0,x\in\mathbb{R}}$ defines a transition function on \mathbb{R} . By Theorem 1.5 in Chapter 3 of [42] we deduce the existence and uniqueness of $\tilde{X}^{m,\xi,r}$.

5.2. Representation of \mathcal{L}_k in terms of $\tilde{X}^{m,\xi,r}$: Proof of Proposition 13. Let $f_0, \ldots, f_k \in \mathcal{C}_b(\mathbb{R})$ be real non-negative functions. Using (1.24), the combination of (6.99) and Lemma 50, and Fubini's theorem

$$\mathbb{E}\left[f_{0}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{0}))\dots f_{k}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k}))\right] \tag{5.85}$$

$$=\mathbb{E}\left[f_{0}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{0}))\dots f_{k-1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))\frac{\lambda_{1}^{X}}{h_{1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))}\int_{0}^{\infty}P_{t}.(h_{1}f_{k})(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))dt\right] \\
=\mathbb{E}\left[f_{0}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{0}))\dots f_{k-1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))\frac{\lambda_{1}^{X}}{h_{1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))}U_{0}(h_{1}f_{k})(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))\right] \\
=\mathbb{E}\left[f_{0}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{0}))\dots f_{k-1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))\frac{\lambda_{1}^{X}}{h_{1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))}\int_{\mathbb{R}}(f_{k}h_{1}m)(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1})+y)v_{\xi}^{r}(y)dy\right] \\
=\lambda_{1}^{X}\int_{\mathbb{R}}\mathbb{E}\left[f_{0}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{0}))\dots f_{k-1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))\frac{v_{\xi}^{r}(y_{k}-\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))}{h_{1}(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k-1}))}\right]f_{k}(y_{k})h_{1}(y_{k})m(y_{k})dy_{k}.$$

Iterating this calculation we get that the left-hand side of (5.85) equals

where we have used $h_1 = |q_1|$ (by definition of h_1), $\tilde{X}_{\mathcal{L}_1}^{m,\xi,r}(T_0) \sim \mathcal{L}_1$, and (3.69) for the penultimate equality, and (3.68) for the last equality.

5.3. Spectral gap of $\tilde{X}^{m,\xi,r}$ and correlation decay in the spin system: Proof of Theorem 15. Let $f \in C_b(\mathbb{R})$. We have $q_1 f \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$ thanks to Proposition 27. Recall also from the proof of Theorem 11 that h_1 is a multiple of q_1 . Therefore, combining (1.24) with Proposition 29 we get that for any t > 0 and $x \in \mathbb{R}$,

$$F_t(x) := \mathbb{E}\left[f\left(\tilde{X}_x^{m,\xi,r}(t)\right)\right] = \frac{e^{t\lambda_1^A}}{q_1(x)} P_t.(q_1f)(x) = \langle q_1f, q_1 \rangle_{L^2(m)} + \frac{1}{q_1(x)} \sum_{n \ge 2} e^{-t(\lambda_n^X - \lambda_1^X)} \langle q_1f, q_n \rangle_{L^2(m)} q_n(x),$$
(5.86)

where the series of functions in the right-hand side converges absolutely in $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$. For any $x \in \mathbb{R}$, we get by dominated convergence that the series of functions in the right-hand side of (5.86) converges to 0 as t goes to infinity. We thus get that for any $x \in \mathbb{R}$,

$$\mathbb{E}\left[f\left(\tilde{X}_x^{m,\xi,r}(t)\right)\right] \xrightarrow[t\to\infty]{} \langle q_1f,q_1\rangle_{L^2(m)} = \int_{\mathbb{R}} f(y)\mathcal{L}_1(dy),$$

where the last equality comes from (3.69). This proves that $\tilde{X}^{m,\xi,r}$ is ergodic with stationary distribution $\mathcal{L}_1(dy)$.

It is easily seen from (3.69) and Proposition 27 that $(q_n/q_1)_{n\geq 1}$ is an orthonormal family of $L^2(\mathcal{L}_1)$. Now let $g \in L^2(\mathcal{L}_1)$ such that g is orthogonal to $(q_n/q_1)_{n\geq 1}$ in $L^2(\mathcal{L}_1)$. Then, by (3.69), $gq_1 \in L^2(m)$ and gq_1 is orthogonal to $(q_n)_{n\geq 1}$ in $L^2(m)$. Proposition 27 yields $gq_1 = 0$ and, by Lemma 31, we deduce that g = 0. Therefore, $(q_n/q_1)_{n\geq 1}$ is an orthonormal Hilbert basis of $L^2(\mathcal{L}_1)$. For our fixed $f \in \mathcal{C}_b(\mathbb{R})$ and $n \geq 2$ we have $||q_n/q_1||_{L^2(\mathcal{L}_1)} = 1$ and $|\langle q_1 f, q_n \rangle_{L^2(m)}| \leq ||f||_{\infty} \langle |q_1|, |q_n| \rangle_{L^2(m)} \leq ||f||_{\infty}$. Combining with Lemma 28 we get that the right-hand side of (5.86) is an absolutely convergent series in $L^2(\mathcal{L}_1)$. In particular $F_t \in L^2(\mathcal{L}_1)$, and proceeding as in (3.59) we get for any $n \geq 1$,

$$\langle F_t, q_n/q_1 \rangle_{L^2(\mathcal{L})} = \langle q_1 F_t, q_n \rangle_{L^2(m)} = e^{-t(\lambda_n^X - \lambda_1^X)} \langle q_1 f, q_n \rangle_{L^2(m)} = e^{-t(\lambda_n^X - \lambda_1^X)} \langle f, q_n/q_1 \rangle_{L^2(\mathcal{L})}.$$

By Parseval's identity we thus get

$$\|F_t - \langle f, 1 \rangle_{L^2(\mathcal{L})}\|_{L^2(\mathcal{L}_1)}^2 = \sum_{n \ge 2} e^{-2t(\lambda_n^X - \lambda_1^X)} |\langle f, q_n/q_1 \rangle_{L^2(\mathcal{L})}|^2 \le e^{-2t(\lambda_2^X - \lambda_1^X)} \|f - \langle f, 1 \rangle_{L^2(\mathcal{L})}\|_{L^2(\mathcal{L}_1)}^2.$$
(5.87)

We get (1.26) with $c := \lambda_2^X - \lambda_1^X$. To show the optimality of this choice of c, take f_M as in (3.63) and note that $f_M q_2/q_1 \in \mathcal{C}_b(\mathbb{R})$. For M large enough we have $|\langle f_M q_2/q_1, q_2/q_1 \rangle_{L^2(\mathcal{L})}|^2 > 0$ (since the latter converges to 1 as M goes to infinity). Applying (5.87) with the choice $f := f_M q_2/q_1$ we thus get $||F_t - \langle f, 1 \rangle_{L^2(\mathcal{L})}||^2_{L^2(\mathcal{L}_1)} \ge e^{-2t(\lambda_2^X - \lambda_1^X)} |\langle f, q_2/q_1 \rangle_{L^2(\mathcal{L})}|^2$ with $|\langle f, q_2/q_1 \rangle_{L^2(\mathcal{L})}|^2 > 0$, so (1.26) can hold for all $f \in \mathcal{C}_b(\mathbb{R})$ only if $c \le \lambda_2^X - \lambda_1^X$. Finally, it will be proved in Section 5.4 below that the coefficient \mathcal{C} from Corollary 14 exists and equals $\log(\lambda_2^X/\lambda_1^X)$. The combination of this with Proposition 34 will yield $c = e^{\mathcal{E}}(e^{\mathcal{C}} - 1)$, completing the proof.

5.4. Correlation decay in the spin system: Proof of Corollary 14. Let us fix $k \ge 3$. Using Proposition 13, the decomposition (5.86) and (3.69) we get

$$C_{k}(f,g) = \mathbb{E}\left[f(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(0))\left(g(\tilde{X}_{\mathcal{L}_{1}}^{m,\xi,r}(T_{k})) - \int_{\mathbb{R}}g(y)\mathcal{L}_{1}(dy)\right)\right]$$
$$= \int_{\mathbb{R}}f(x)\int_{0}^{\infty}\left(\mathbb{E}\left[g(\tilde{X}_{x}^{m,\xi,r}(t))\right] - \int_{\mathbb{R}}g(y)\mathcal{L}_{1}(dy)\right)\mathbb{P}(T_{k}\in dt)\mathcal{L}_{1}(dx)$$
$$= \int_{\mathbb{R}}m(x)f(x)\overline{q_{1}(x)}\int_{0}^{\infty}\left(\sum_{n\geq 2}e^{-t(\lambda_{n}^{X}-\lambda_{1}^{X})}\langle q_{1}g,q_{n}\rangle_{L^{2}(m)}q_{n}(x)\right)\mathbb{P}(T_{k}\in dt)dx.$$
(5.88)

Then, using that $|\langle q_1 g, q_n \rangle_{L^2(m)}| \leq ||g||_{\infty} \langle |q_1|, |q_n| \rangle_{L^2(m)} \leq ||g||_{\infty}$ we get

$$\int_{0}^{\infty} \left(\sum_{n \ge 2} e^{-t(\lambda_{n}^{X} - \lambda_{1}^{X})} \left| \langle q_{1}g, q_{n} \rangle_{L^{2}(m)} q_{n}(x) \right| \right) \mathbb{P}(T_{k} \in dt) = \sum_{n \ge 2} \left(\frac{\lambda_{1}^{X}}{\lambda_{n}^{X}} \right)^{k} \left| \langle q_{1}g, q_{n} \rangle_{L^{2}(m)} q_{n}(x) \right|$$

$$\leq \|g\|_{\infty} \sum_{n \ge 2} \left(\frac{\lambda_{1}^{X}}{\lambda_{n}^{X}} \right)^{k} \|q_{n}\|_{\infty}.$$
(5.89)

Since $k \ge 3$, Lemma 28 shows that the right-hand side is finite. We can thus use Fubini's theorem in (5.88) and get

$$C_k(f,g) = \int_{\mathbb{R}} m(x)f(x)\overline{q_1(x)} \left(\sum_{n\geq 2} \left(\frac{\lambda_1^X}{\lambda_n^X} \right)^k \langle q_1g, q_n \rangle_{L^2(m)} q_n(x) \right) dx.$$
(5.90)

We have $m \in L^2(\mathbb{R})$ by (Condition 1), $q_1 \in L^2(\mathbb{R})$ by Proposition 27 and $f \in C_b(\mathbb{R})$ so $mf\overline{q_1} \in L^1(\mathbb{R})$ by Cauchy-Schwartz inequality. Also, (5.89) shows that the convergence is the series in the above expression is uniform in x. We can thus intervene the sum and the series in (5.90) and get

$$\begin{split} C_k(f,g) &= \sum_{n\geq 2} \left(\frac{\lambda_1^X}{\lambda_n^X}\right)^k \overline{\langle q_1 \overline{f}, q_n \rangle_{L^2(m)}} \langle q_1 g, q_n \rangle_{L^2(m)} \\ &= \left(\frac{\lambda_1^X}{\lambda_2^X}\right)^k \left(\sum_{j=2}^{M_2+1} \overline{\langle q_1 \overline{f}, q_j \rangle_{L^2(m)}} \langle q_1 g, q_j \rangle_{L^2(m)} + \sum_{n>M_2+1} \left(\frac{\lambda_2^X}{\lambda_n^X}\right)^k \overline{\langle q_1 \overline{f}, q_n \rangle_{L^2(m)}} \langle q_1 g, q_n \rangle_{L^2(m)} \right), \end{split}$$

where M_2 denotes the multiplicity of the eigenvalue λ_2^X (ie the number of indices j such that $\lambda_j^X = \lambda_2^X$). Note that $|\langle q_1 f, q_n \rangle_{L^2(m)} \langle q_1 g, q_n \rangle_{L^2(m)}| \leq ||f||_{\infty} ||g||_{\infty}$ so, by dominated convergence, we get that the term $\sum_{n>M_2+1} \cdots$ converges to 0 as k goes to infinity. This yields that (1.25) holds with $\mathcal{C} = \log(\lambda_2^X/\lambda_1^X)$ and $B(f,g) = \sum_{j=2}^{M_2+1} \overline{\langle q_1 \overline{f}, q_j \rangle_{L^2(m)}} \langle q_1 g, q_j \rangle_{L^2(m)}$. We have $|B(f,g)| \leq M_2 ||f||_{\infty} ||g||_{\infty}$ so $B(\cdot, \cdot)$ is continuous. To see that $B(\cdot, \cdot)$ is non-zero, take f_M as in (3.63) and note that $f_M q_2/q_1 \in \mathcal{C}_b(\mathbb{R})$. Then it is easy to see that $B(f_M \overline{q_2}/\overline{q_1}, f_M q_2/q_1)$ converges to 1 as M goes to infinity, so $B(\cdot, \cdot)$ is non-zero.

6. Some facts about $X^{m,\xi,r}$, its semigroup, and its generator

6.1. Properties of the random time-change. In this subsection we prove some properties of the random time-change $A_x^{-1}(\cdot)$ defined in (1.1).

Lemma 41. Assume that (Condition 1) holds. If ξ is recurrent (which occurs, in particular, if $\mathbb{E}[|\xi(1)|] < \infty$, see Theorem 36.7 in [46]) then $A_x(\infty) = \infty$ almost surely.

Proof. Since *m* is continuous and positive (by (Condition 1)) we have $\inf_{y \in [-1,1]} m(y) > 0$. Note that $A_x(\infty) \ge (\inf_{y \in [-1,1]} m(y)) \int_0^\infty \mathbf{1}_{|\xi(s)| \le 1} ds$. If ξ is recurrent then by Theorem 35.4 in [46] we have almost surely $\int_0^\infty \mathbf{1}_{|\xi(s)| \le 1} ds = \infty$, from which we conclude that $A_x(\infty) = \infty$.

Lemma 42. Assume that (Condition 1) holds.

- (1) For any $x \in \mathbb{R}$, $A_x(\cdot)$ and $A_x^{-1}(\cdot)$ are almost surely continuous and increasing on the intervals $[0, \infty)$ and $[0, A_x(\infty))$ respectively.
- (2) Let $t \ge 0$, $x \in \mathbb{R}$, and $(x_n)_{n\ge 1}$ be a sequence converging to x. Then $A_{x_n}^{-1}(t)$ converges to $A_x^{-1}(t)$ almost surely (regardless of $A_x^{-1}(t) < \infty$ or $A_x^{-1}(t) = \infty$).
- (3) Let $t \ge 0$ and $x \in \mathbb{R}$. Then $\xi(\cdot)$ is continuous at $A_x^{-1}(t)$ almost surely on $\{A_x^{-1}(t) < \infty\}$.

Proof. The first point was justified a little after (1.1). Let us prove the second point. Let $t \ge 0, x \in \mathbb{R}$, and $(x_n)_{n\ge 1}$ be a sequence converging to x. We fix a realization of ξ such that $A_x(\cdot)$ is continuous and increasing on $[0, \infty)$. Assume first that $A_x^{-1}(t) < \infty$ and that there is $\epsilon > 0$ such that $A_{xn}^{-1}(t) - A_x^{-1}(t) > \epsilon$ for infinitely many indices n (this includes indices n such that $A_{xn}^{-1}(t) = \infty$). Let $(n(k))_{k\ge 1}$ be the enumeration of those indices. For any $k \ge 1$ we have $t \ge A_{xn(k)}(A_x^{-1}(t) + \epsilon)$. Let $\rho(\cdot)$ denote the modulus of continuity of m, ie $\rho(\delta) := \sup\{|m(y) - m(z)|, y, z \in \mathbb{R} \text{ s.t. } |y-z| \le \delta\}$. Since $m \in C_0(\mathbb{R})$ (by (Condition 1)), m is uniformly continuous on \mathbb{R} so $\rho(\delta)$ converges to 0 as δ goes to 0. Note from (1.1) that for any $T \ge 0$, $|A_x(T) - A_{xn(k)}(T)| \le T\rho(|x-x_{n(k)}|)$, so $A_{xn(k)}(A_x^{-1}(t) + \epsilon)$ converges to $A_x(A_x^{-1}(t) + \epsilon)$ as k goes to infinity. We thus get $t \ge A_x(A_x^{-1}(t) + \epsilon)$. Since $A_x(\cdot)$ is increasing on $[0, \infty)$ and $A_x^{-1}(t) < \infty$ we get $A_x(A_x^{-1}(t) + \epsilon) > A_x(A_x^{-1}(t)) = t$, which is a contradiction. Still in the case $A_x^{-1}(t) < \infty$, if there is $\epsilon > 0$ such that $A_{xn}^{-1}(t) - A_x^{-1}(t) < -\epsilon$ for infinitely many indices n we obtain a similar contradiction. In conclusion, if $A_x^{-1}(t) - A_x^{-1}(t) < -\epsilon$ for infinitely many indices n we as that $A_x^{-1}(t) < \infty$ and $|A_{xn}^{-1}(t)| \le \epsilon$. This proves the claimed result for any fixed realization of ξ such that $A_x(M) < t$ so $A_{xn}(M) < t$ so $A_{xn}(M) < t$. Using $|A_x(M) - A_{xn}(M)| \le M\rho(|x - x_n|)$ we get that for all large n, $A_{xn}(M) < t$ so $A_{xn}^{-1}(t) \ge M$. Therefore the claimed result is now also proved for any fixed realization of ξ such that $A_x(K)$ is continuous and increasing on $[0, \infty)$, and $A_x^{-1}(t) = \infty$. Combining with the first point of the lemma, we get the second point.

We now prove the third point. Recall that $A_x^{-1}(t)$ is a \mathcal{F}^{ξ} -stopping time, where \mathcal{F}^{ξ} is the filtration defined in Section 1.6. By the first point, on $\{A_x^{-1}(t) < \infty\}$, we have almost surely that the sequence of \mathcal{F}^{ξ} -stopping times $(A_x^{-1}(t-1/n))_{n\geq 1}$ increases to $A_x^{-1}(t)$ and that $A_x^{-1}(t-1/n) < A_x^{-1}(t)$ for any $n \geq 1$. By Proposition I.7 of [4] we get that ξ is continuous at $A_x^{-1}(t)$ almost surely on $\{A_x^{-1}(t) < \infty\}$. This proves the third point.

6.2. Markov property, Feller property.

Lemma 43. Assume that r > 0 and (Condition 1) holds. The process $X^{m,\xi,r}$ defined by (1.2) is an homogeneous Markovian process. In particular the family $(P_t)_{t\geq 0}$ (defined by (1.3)) is a semigroup (i.e. it satisfies $P_{t+s} f = P_s P_t f$ for any bounded measurable f and $t, s \geq 0$).

Proof. We fix $x \in \mathbb{R}$. Recall the filtrations \mathcal{F}^{ξ} and \mathcal{F}^{X} defined in Section 1.6. Note that for any $t \ge 0$, $A_{x}^{-1}(t)$ is a (possibly infinite) \mathcal{F}^{ξ} -stopping time and that $\mathcal{F}_{t}^{X} = \mathcal{F}_{A_{x}^{-1}(t)}^{\xi}$. Let us fix $t \ge 0$ and prove the Markov property for $X_{x}^{m,\xi,r}$ at time t. We have clearly $X^{m,\xi,r}(t+s) = \dagger$ for any $s \ge 0$ if we are on $\{X_{x}^{m,\xi,r}(t) = \dagger\} = \{A_{x}^{-1}(t) \ge e_{r}\}$ so let assume we are on $\{X_{x}^{m,\xi,r}(t) \neq \dagger\} = \{A_{x}^{-1}(t) < e_{r}\}$. By the strong Markov property for ξ , the process $\hat{\xi} := \xi(A_{x}^{-1}(t) + \cdot) - \xi(A_{x}^{-1}(t))$ has same law as ξ and is independent of $\mathcal{F}_{A_{x}^{-1}(t)}^{\xi} = \mathcal{F}_{t}^{X}$. Let also $\hat{e}_{r} := e_{r} - A_{x}^{-1}(t)$. For $y \in \mathbb{R}$, let \hat{A}_{y} and $\hat{X}_{y}^{m,\xi,r}$ be defined by (1.1) and (1.2), but with ξ and e_{r} replaced by $\hat{\xi}$ and \hat{e}_{r} respectively.

We see that, conditionally on \mathcal{F}_t^X and on $\{X_x^{m,\xi,r}(t) \neq \dagger\}$, $\hat{X}_y^{m,\xi,r}$ has the same law as $X_y^{m,\xi,r}$. Let us choose $y = X_x^{m,\xi,r}(t) = x + \xi(A_x^{-1}(t))$. For any $s \in [0, \hat{A}_y(\infty))$,

$$s = \int_{0}^{\hat{A}_{y}^{-1}(s)} m(y + \hat{\xi}(u)) du = \int_{0}^{\hat{A}_{y}^{-1}(s)} m(x + \xi(A_{x}^{-1}(t) + u)) du$$
$$= \int_{A_{x}^{-1}(t)}^{A_{x}^{-1}(t) + \hat{A}_{y}^{-1}(s)} m(x + \xi(u)) du = A_{x}(A_{x}^{-1}(t) + \hat{A}_{y}^{-1}(s)) - t.$$

We thus get $A_x^{-1}(t+s) < \infty$ and $A_x^{-1}(t+s) = A_x^{-1}(t) + \hat{A}_y^{-1}(s)$. For $s \ge \hat{A}_y(\infty)$, we have $s > \int_0^M m(y + \hat{\xi}(u)) du$ for any M > 0, thanks to the positivity of m. Proceeding as above we get $s > A_x(A_x^{-1}(t) + M) - t$ for any M > 0 so $A_x^{-1}(t+s) = \infty$. In conclusion $A_x^{-1}(t+s) < \infty \Leftrightarrow s < \hat{A}_y(\infty) \Leftrightarrow \hat{A}_y^{-1}(s) < \infty$ and in that case we have $A_x^{-1}(t+s) < e_r \Leftrightarrow \hat{A}_y^{-1}(s) < \hat{e}_r$. Therefore,

$$X_x^{m,\xi,r}(t+s) = x + \xi(A_x^{-1}(t+s)) = x + \xi(A_x^{-1}(t) + \hat{A}_y^{-1}(s)) = y + \hat{\xi}(\hat{A}_y^{-1}(s)) = \hat{X}_y^{m,\xi,r}(s).$$

We have thus obtained that $X_x^{m,\xi,r}(t+\cdot)$ equals $\hat{X}_y^{m,\xi,r}$ with $y = X_x^{m,\xi,r}(t)$ and we know that, conditionally on \mathcal{F}_t^X and on $\{X_x^{m,\xi,r}(t) \neq \dagger\}$, the later is distributed as a version of $X^{m,\xi,r}$ starting at $X_x^{m,\xi,r}(t)$. This shows that $X^{m,\xi,r}$ is Markovian and homogeneous. The semigroup property for $(P_t)_{t\geq 0}$ trivially follows.

We now prove the Feller property for $X^{m,\xi,r}$.

Proof of Proposition 1. From (1.3) and Lemma 43 we see that $(P_t)_{t\geq 0}$ is a sub-Markov semigroup in the sense of Definition 1.1 of [6]. In order to check that $(P_t)_{t\geq 0}$ satisfies Definition 1.2 of [6] we need to show that $P_t f \in C_0(\mathbb{R})$ for any $f \in C_0(\mathbb{R})$ and t > 0 (Feller property) and that $||P_t f - f||_{\infty}$ converges to 0 as t goes to 0 for any $f \in C_0(\mathbb{R})$ (strong continuity).

We first prove the strong continuity. Let $f \in C_0(\mathbb{R})$, $\epsilon > 0$ and let us prove the existence of $t_0 > 0$ such that $t < t_0 \Rightarrow ||P_t \cdot f - f||_{\infty} \le \epsilon$. If f is the null function the claim trivially holds so we assume that f is not identically 0. Since $f \in C_0(\mathbb{R})$ is uniformly continuous, there is $\delta > 0$ such that $|x - y| \le \delta \Rightarrow |f(x) - f(y)| \le \epsilon/4$. Let us fix $\delta > 0$ small enough and M > 0 large enough such that

$$\mathbb{P}\left(e_r \le \tilde{\delta}\right) \le \frac{\epsilon}{8\|f\|_{\infty}}, \ \mathbb{P}\left(\sup_{s \in [0,\tilde{\delta}]} |\xi(s)| > \delta\right) \le \frac{\epsilon}{8\|f\|_{\infty}}, \ \mathbb{P}\left(\sup_{s \in [0,e_r]} |\xi(s)| > M\right) \le \frac{\epsilon}{8\|f\|_{\infty}}, \ \sup_{|x| \ge M} |f(x)| \le \frac{\epsilon}{3}.$$
(6.91)

We set $t_0 := \tilde{\delta} \times \inf_{z \in [-3M, 3M]} m(z)$. Since *m* is positive and continuous by (Condition 1) we have $t_0 > 0$. We now prove that $\|P_t.f - f\|_{\infty} \leq \epsilon$ for all $t \in (0, t_0)$. First let $t \in (0, t_0)$ and $|x| \leq 2M$. Note from (1.3) that

$$|P_{t}.f(x) - f(x)| \le \mathbb{E}\left[\left| f(x + \xi(A_x^{-1}(t))) - f(x) \right| \mathbf{1}_{A_x^{-1}(t) < e_r} + |f(x)| \mathbf{1}_{A_x^{-1}(t) \ge e_r} \right].$$
(6.92)

If $\sup_{s \in [0, e_r]} |\xi(s)| \leq M$ and $e_r > \tilde{\delta}$ then using (1.1) and the definition of t_0 ,

$$A_x(\tilde{\delta}) = \int_0^{\delta} m(x+\xi(u)) du \ge \tilde{\delta} \times \inf_{u \in [0,\tilde{\delta}]} m(x+\xi(u)) \ge \tilde{\delta} \times \inf_{u \in [0,e_r]} m(x+\xi(u)) \ge \tilde{\delta} \times \inf_{z \in [-3M,3M]} m(z) = t_0 > t.$$

Therefore $A_x^{-1}(t) \leq \tilde{\delta} < e_r$. If moreover $\sup_{s \in [0, \tilde{\delta}]} |\xi(s)| \leq \delta$ then $|\xi(A_x^{-1}(t))| \leq \delta$ so $|f(x+\xi(A_x^{-1}(t)))-f(x)| \leq \epsilon/4$. Plugging into (6.92) and then using (6.91) we get that for $t \in (0, t_0)$ and $|x| \leq 2M$,

$$|P_t \cdot f(x) - f(x)| \le \epsilon/4 + 2||f||_{\infty} \left(\mathbb{P}\left(\sup_{s \in [0, e_r]} |\xi(s)| > M \right) + \mathbb{P}\left(e_r \le \tilde{\delta} \right) + \mathbb{P}\left(\sup_{s \in [0, \tilde{\delta}]} |\xi(s)| > \delta \right) \right) \le \epsilon.$$
(6.93)

Now let t > 0 and $|x| \ge 2M$. We have from (1.3) that

$$|P_t \cdot f(x)| \le \mathbb{E}\left[\left| f(x + \xi(A_x^{-1}(t))) \right| \mathbf{1}_{A_x^{-1}(t) < e_r} \right].$$

If $\sup_{s \in [0, e_r]} |\xi(s)| \le M$, we have $|x + \xi(A_x^{-1}(t))| \ge M$ so, by (6.91), $|f(x + \xi(A_x^{-1}(t)))| \le \epsilon/3$. Therefore, for any t > 0 and $|x| \ge 2M$,

$$|P_t \cdot f(x)| \le \epsilon/3 + ||f||_{\infty} \times \mathbb{P}\left(\sup_{s \in [0, e_r]} |\xi(s)| > M\right) \le 2\epsilon/3,$$
(6.94)

where the last equality comes from (6.91). We also have $|f(x)| \leq \epsilon/3$ by (6.91). Combining with (6.94) we get $|P_t f(x) - f(x)| \leq \epsilon$ for all t > 0 and $|x| \geq 2M$. The combination of this with (6.93) yields $||P_t f(x) - f||_{\infty} \leq \epsilon$ for all $t \in (0, t_0)$, proving the strong continuity.

Let us fix $f \in C_0(\mathbb{R})$, t > 0 and now prove that $P_t.f \in C_0(\mathbb{R})$. Let $x \in \mathbb{R}$ and $(x_n)_{n\geq 1}$ be a sequence converging to x. Recall from (1.3) that $P_t.f(x_n) = \mathbb{E}[f(x_n + \xi(A_{x_n}^{-1}(t)))\mathbf{1}_{A_{x_n}^{-1}(t) < e_r}]$. Using the second and third points of Lemma 42 and $f \in C_0(\mathbb{R})$ we see that what is inside the expectation converges almost surely to $f(x + \xi(A_x^{-1}(t)))\mathbf{1}_{A_x^{-1}(t) < e_r}$ (the convergence may not hold on $\{A_x^{-1}(t) = e_r\}$ but that event has probability 0). Moreover, what is inside the expectation is bounded by $\|f\|_{\infty}$ so, by dominated convergence, we get that $P_t.f(x_n)$ converges to $P_t.f(x)$. It follows that $P_t.f$ is continuous. Then fix $\epsilon > 0$ and note from (6.94) that there is M > 0 such that $|x| \ge 2M \Rightarrow |P_t.f(x)| \le \epsilon$. Therefore $P_t.f(x) \longrightarrow_{|x|\to\infty} 0$, so $P_t.f \in C_0(\mathbb{R})$. This concludes the proof.

6.3. Support of $X^{m,\xi,r}$. The following lemma is crucial to prove Lemma 31 which plays an important role in showing the positivity of the constant $K_{m,\xi,r}(x)$ in Theorem 5, and in relating it with the infinite-volume Gibbs states of the spin system from Section 1.2.

Lemma 44. Assume that r > 0 and (Condition 1) and (Condition 2) hold. For any t > 0 and $x \in \mathbb{R}$ we have $Supp(X_x^{m,\xi,r}(t)) = \mathbb{R} \cup \{\dagger\}.$

Proof. Let us fix t > 0 and $x \in \mathbb{R}$. To show that $\mathbb{R} \subset Supp(X_x^{m,\xi,r}(t))$, we prove that for any $z \in \mathbb{R}$ and $\epsilon > 0$ we have $\mathbb{P}(X_x^{m,\xi,r}(t) \in [z - \epsilon, z + \epsilon]) > 0$. Let us fix $\delta \in (0, t/||m||_{\infty})$ and $T > \delta + t/\min_{y \in [z - x - \epsilon, z - x + \epsilon]} m(y)$. We define the event

$$\mathcal{E} := \{e_r > T\} \cap \{\xi(\delta) \in (z - x - \epsilon/2, z - x + \epsilon/2)\} \cap \left\{ \sup_{s \in [\delta, T]} |\xi(s) - \xi(\delta)| < \epsilon/2 \right\}.$$

From (1.1) we see that on this event we have

$$A_x(\delta) \le \delta \|m\|_{\infty} < t < (T-\delta) \times \min_{y \in [z-x-\epsilon, z-x+\epsilon]} m(y) \le A_x(T) < A_x(e_r),$$

and $x + \xi^r(s) \in [z - \epsilon, z + \epsilon]$ for all $s \in [\delta, T]$. In conclusion, we have

$$\mathcal{E} \subset \left\{ \delta < A_x^{-1}(t) < T < e_r \right\} \cap \left\{ \forall s \in [\delta, T], \ x + \xi^r(s) \in [z - \epsilon, z + \epsilon] \right\} \subset \left\{ X_x^{m, \xi, r}(t) \in [z - \epsilon, z + \epsilon] \right\},$$

where the last inclusion comes from the definition of $X_x^{m,\xi,r}(t)$ in (1.2).

We are now left to prove that $\mathbb{P}(\mathcal{E}) > 0$. Using that e_r is independent of ξ and that $\xi(\delta)$ and $(\xi(s) - \xi(\delta))_{s \ge \delta}$ are independent we get

$$\mathbb{P}(\mathcal{E}) = e^{-rT} \times \mathbb{P}\left(\xi(\delta) \in (z - x - \epsilon/2, z - x + \epsilon/2)\right) \times \mathbb{P}\left(\sup_{s \in [\delta, T]} |\xi(s) - \xi(\delta)| < \epsilon/2\right).$$
(6.95)

By Lemma 51, $Supp(\xi(\delta)) = \mathbb{R}$, so the second factor in (6.95) is positive. Lemma 52 shows that $\mathbb{P}(\sup_{s \in [0, T-\delta]} |\xi(s)| < \epsilon/2) > 0$. By the Markov property at time δ , we get that the third factor in (6.95) is positive. This concludes the proof of $\mathbb{R} \subset Supp(X_x^{m,\xi,r}(t))$.

To show that $\dagger \in Supp(X_x^{m,\xi,r}(t))$, note that the event $\{e_r < t/||m||_{\infty}\}$ has positive probability and is included into $\{A_x^{-1}(t) \ge e_r\} = \{X_x^{m,\xi,r}(t) = \dagger\}$. Therefore $\dagger \in Supp(X_x^{m,\xi,r}(t))$, so the proof is complete. \Box

6.4. Properties of the generator.

Lemma 45. Assume that r > 0 and (Condition 1) holds true. If $f \in \mathcal{D}(\mathcal{A}_{\xi^r})$ is such that $\frac{1}{m}\mathcal{A}_{\xi^r}f \in \mathcal{C}_0(\mathbb{R})$ then $f \in \mathcal{D}(\mathcal{A}_{X^m,\xi,r})$ and we have $\mathcal{A}_{X^m,\xi,r}f(x) = \frac{1}{m(x)}\mathcal{A}_{\xi^r}f(x)$.

Proof. Let $f \in \mathcal{D}(\mathcal{A}_{\xi^r})$ and $g(x) := \frac{1}{m(x)}\mathcal{A}_{\xi^r}f(x)$. It is assumed that $g \in \mathcal{C}_0(\mathbb{R})$ so, according to Theorem 1.33 in [6], we only need to prove

$$\forall x \in \mathbb{R}, \ t^{-1}(P_t f(x) - f(x)) \xrightarrow{}_{t \to 0} g(x).$$
(6.96)

For this, let us proceed similarly as for the proof of Corollary 4.2 in [6]. This requires some carefulness since we are not in their setting, as 1/m is not a bounded function. Since $f \in \mathcal{D}(\mathcal{A}_{\xi^r})$ we have that $(M_t^f)_{t\geq 0}$ defined by $M_t^f := f(x+\xi(t))\mathbf{1}_{t< e_r} - f(x) - \int_0^{t\wedge e_r} \mathcal{A}_{\xi^r} f(x+\xi(s)) ds$ is a \mathcal{F}^{ξ} -martingale (see for example Theorem 1.36 in [6]),

where \mathcal{F}^{ξ} is the filtration defined in Section 1.6. Therefore, using that $dA_x(u) = m(x + \xi(u))du$, the definition of g, and making the substitution $s = A_x(u)$ we get that for any $t \ge 0$ and $n \ge 1$,

$$f(x+\xi(A_x^{-1}(t)\wedge n))\mathbf{1}_{A_x^{-1}(t)\wedge n < e_r} - f(x) - M_{A_x^{-1}(t)\wedge n}^f = \int_0^{A_x^{-1}(t)\wedge n \wedge e_r} \mathcal{A}_{\xi^r} f(x+\xi(u)) du$$
$$= \int_0^{A_x^{-1}(t)\wedge n \wedge e_r} g(x+\xi(u)) dA_x(u) = \int_0^{t\wedge A_x(n)\wedge A_x(e_r)} g(x+\xi(A_x^{-1}(s))) ds.$$

Since $A_x^{-1}(t) \wedge n$ is a bounded \mathcal{F}^{ξ} -stopping time, we get from the optional stopping theorem

$$\mathbb{E}[f(x+\xi(A_x^{-1}(t)\wedge n))\mathbf{1}_{A_x^{-1}(t)\wedge n< e_r}] - f(x) = \mathbb{E}\left[\int_0^{t\wedge A_x(n)\wedge A_x(e_r)} g(x+\xi(A_x^{-1}(s)))ds\right].$$
(6.97)

From the third point of Lemma 42 and the continuity of f, what is inside the expectation in the left-hand side of (6.97) converges to $f(x + \xi(A_x^{-1}(t)))\mathbf{1}_{A_x^{-1}(t) < e_r}$ as n increases to infinity. Since f and g are bounded functions, we get by dominated convergence that

$$\mathbb{E}[f(x+\xi(A_x^{-1}(t)))\mathbf{1}_{A_x^{-1}(t)< e_r}] - f(x) = \mathbb{E}\left[\int_0^{t\wedge A_x(e_r)} g(x+\xi(A_x^{-1}(s)))ds\right]$$

By (1.3), this translates into

$$t^{-1}(P_t.f(x) - f(x)) = \mathbb{E}\left[t^{-1} \int_0^{t \wedge A_x(e_r)} g(x + \xi(A_x^{-1}(s)))ds\right].$$
(6.98)

From the first point of Lemma 42, $A_x^{-1}(\cdot)$ is almost surely continuous on $[0, A_x(e_r))$ and $A_x(e_r)$ is almost surely positive. Moreover ξ is almost surely right-continuous by definition of a Lévy process and g is continuous. We deduce that what is inside the expectation in the right-hand side of (6.98) converges almost surely to g(x) as t goes to 0. Moreover it is bounded by $||g||_{\infty}$. By dominated convergence we obtain (6.96), which concludes the proof.

6.5. Properties of the semigroup and resolvent.

Lemma 46. Assume that r > 0 and (Condition 1) holds. We have $P_t(\mathcal{C}_0(\mathbb{R})) \subset L^2(\mathbb{R}) \subset L^2(m)$ for all t > 0.

Proof. Let t > 0 and $f \in C_0(\mathbb{R})$. By (1.3), Markov inequality, two times Jensen inequality, (1.1), Fubini's theorem and $m \in L^2(\mathbb{R})$ (from (Condition 1)) we get

$$\begin{split} \int_{\mathbb{R}} |P_t f(x)|^2 dx &\leq \|f\|_{\infty}^2 \int_{\mathbb{R}} \mathbb{P}(A_x^{-1}(t) < e_r)^2 dx = \|f\|_{\infty}^2 \int_{\mathbb{R}} \mathbb{P}(A_x(e_r) > t)^2 dx \leq \frac{\|f\|_{\infty}^2}{t^2} \int_{\mathbb{R}} \mathbb{E}[A_x(e_r)]^2 dx \\ &\leq \frac{\|f\|_{\infty}^2}{t^2} \int_{\mathbb{R}} \mathbb{E}\left[A_x(e_r)^2\right] dx \leq \frac{\|f\|_{\infty}^2}{t^2} \int_{\mathbb{R}} \mathbb{E}\left[e_r \int_0^{e_r} m(x + \xi(u))^2 du\right] dx \\ &= \frac{\|f\|_{\infty}^2}{t^2} \mathbb{E}\left[e_r \int_0^{e_r} \int_{\mathbb{R}} m(x + \xi(u))^2 dx du\right] = \frac{\|f\|_{\infty}^2 \|m\|_{L^2(\mathbb{R})}^2}{t^2} \mathbb{E}[e_r^2] = \frac{2\|f\|_{\infty}^2 \|m\|_{L^2(\mathbb{R})}^2}{r^2 t^2} < \infty. \end{split}$$

The above proof even shows that the operator P_t is continuous from $(\mathcal{C}_0(\mathbb{R}), \|\cdot\|_{\infty})$ to $L^2(\mathbb{R})$ and $L^2(m)$.

Remark 47. Applying the above proof to the constant function equal to 1, instead of $f \in \mathcal{C}_0(\mathbb{R})$, one gets that $P_t.1 = (x \mapsto \mathbb{P}(\zeta_x > t)) \in L^2(\mathbb{R}) \subset L^2(m)$.

Before proving the next lemma, let us recall that for any $\alpha \geq 0$, the resolvent operator at α associated with the semigroup $(P_t)_{t\geq 0}$ is defined by $U_{\alpha}f := \int_0^{\infty} e^{-\alpha t} P_t f dt$ for $f \in C_0(\mathbb{R})$ (see Definition 1.21 in [6]). Assume that r > 0 and (Condition 1) holds. Using (1.3), Fubini's theorem, the change of variable $s = A_x^{-1}(t)$, and again Fubini's theorem, we get that for any $x \in \mathbb{R}$,

$$U_{\alpha}f(x) = \mathbb{E}\left[\int_{0}^{A_{x}(e_{r})} e^{-\alpha t} f(x + \xi(A_{x}^{-1}(t)))dt\right] = \mathbb{E}\left[\int_{0}^{e_{r}} e^{-\alpha A_{x}(s)}(fm)(x + \xi(s))ds\right]$$
(6.99)
$$= \mathbb{E}\left[\int_{0}^{\infty} \mathbf{1}_{s \leq e_{r}} e^{-\alpha A_{x}(s)}(fm)(x + \xi(s))ds\right] = \int_{0}^{\infty} e^{-rs} \mathbb{E}\left[e^{-\alpha A_{x}(s)}(fm)(x + \xi(s))\right]ds.$$

Lemma 48. Assume that r > 0 and (Condition 1) holds. For any $\alpha > 0$ we have $U_{\alpha}(\mathcal{C}_0(\mathbb{R})) \subset \mathcal{C}_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$. Moreover, we have $\langle U_{\alpha}f, g \rangle_{L^2(m)} = \langle f, U_{\alpha}g \rangle_{L^2(m)}$ for all $\alpha > 0$ and $f, g \in \mathcal{C}_0(\mathbb{R}) \cap L^2(\mathbb{R})$.

Proof. From Lemma 1.27 in [6] we get $U_{\alpha}(\mathcal{C}_0(\mathbb{R})) \subset \mathcal{D}(\mathcal{A}_{X^{m,\xi,r}}) \subset \mathcal{C}_0(\mathbb{R})$. Let $f \in \mathcal{C}_0(\mathbb{R})$. Using (6.99), two times Jensen's inequality, Fubini's theorem, and $m \in L^2(\mathbb{R})$ (from (Condition 1)) we get

$$\int_{\mathbb{R}} |U_{\alpha}f(x)|^2 dx \leq \frac{1}{r} \int_{\mathbb{R}} \int_0^{\infty} e^{-rs} \mathbb{E} \left[\left| e^{-\alpha A_x(s)}(fm)(x+\xi(s)) \right|^2 \right] ds dx \\ \leq \frac{\|f\|_{\infty}^2}{r} \int_0^{\infty} e^{-rs} \mathbb{E} \left[\int_{\mathbb{R}} |m(x+\xi(s))|^2 dx \right] ds = \frac{\|f\|_{\infty}^2 \|m\|_{L^2(\mathbb{R})}^2}{r^2} < \infty$$

Therefore $U_{\alpha}(\mathcal{C}_0(\mathbb{R})) \subset L^2(\mathbb{R})$.

Now let $f, g \in \mathcal{C}_0(\mathbb{R}) \cap L^2(\mathbb{R})$. Therefore $U_\alpha f, U_\alpha g, f$ and g are all in $L^2(\mathbb{R}) \subset L^2(m)$. Using (6.99) we have

$$\langle U_{\alpha}f,g\rangle_{L^{2}(m)} = \int_{\mathbb{R}} U_{\alpha}f(x)\overline{g(x)}m(x)dx = \int_{\mathbb{R}} \int_{0}^{\infty} e^{-rs}\mathbb{E}\left[e^{-\alpha A_{x}(s)}(fm)(x+\xi(s))(\overline{g}m)(x)\right]dsdx.$$
(6.100)

By $m \in L^2(\mathbb{R})$ and Cauchy-Schwartz inequality, $\overline{g}m \in L^1(\mathbb{R})$, so $\int_{\mathbb{R}} \int_0^\infty e^{-rs} \mathbb{E}[|e^{-\alpha A_x(s)}(fm)(x+\xi(s))(\overline{g}m)(x)|] ds dx \leq ||f||_\infty ||m||_\infty ||\overline{g}m||_{L^1(\mathbb{R})}/r < \infty$. We can thus use Fubini's theorem in (6.100) and get

$$\langle U_{\alpha}f,g\rangle_{L^{2}(m)} = \int_{0}^{\infty} e^{-\tau s} \mathbb{E}\left[\int_{\mathbb{R}} e^{-\alpha A_{x}(s)}(fm)(x+\xi(s))(\overline{g}m)(x)dx\right] ds.$$
(6.101)

For any s > 0 let us define the process ξ^s on [0, s] by $\xi^s(s) := -\xi(s)$ and $\xi^s(u) := \xi((s - u) - (\xi - u)) - \xi(s)$ when $u \in [0, s)$. By Lemma II.2 of [4], $(\xi^s(u))_{u \in [0, s]}$ is a Lévy process equal in law to $(-\xi(u))_{u \in [0, s]}$ which, by the symmetry of ξ , is equal in law to $(\xi(u))_{u \in [0, s]}$. Using (1.1), the change of variable v = s - u, that ξ is continuous at almost every time, and the definition of ξ^s , we see that we have a.s. for all $x \in \mathbb{R}$,

$$A_x(s) = \int_0^s m(x + \xi(s) + \xi((s - u) -) - \xi(s)) du = \int_0^s m(x - \xi^s(s) + \xi^s(u)) du = \tilde{A}_{x - \xi^s(s)}(s),$$

where we have set $\tilde{A}_y(t) := \int_0^t m(y + \xi^s(u)) du$ for any $y \in \mathbb{R}$ and $t \in [0, s]$. Plugging this in (6.101) we get

$$\langle U_{\alpha}f,g\rangle_{L^{2}(m)} = \int_{0}^{\infty} e^{-rs} \mathbb{E}\left[\int_{\mathbb{R}} e^{-\alpha \tilde{A}_{x-\xi^{s}(s)}(s)}(fm)(x-\xi^{s}(s))(\overline{g}m)(x)dx\right] ds$$

$$= \int_{0}^{\infty} e^{-rs} \mathbb{E}\left[\int_{\mathbb{R}} e^{-\alpha \tilde{A}_{y}(s)}(fm)(y)(\overline{g}m)(y+\xi^{s}(s))dy\right] ds.$$

$$(6.102)$$

Since $(\xi^s(u))_{u \in [0,s]}$ is equal in law to $(\xi(u))_{u \in [0,s]}$ we have $\mathbb{E}[F(\tilde{A}(s), \xi^s(s))] = \mathbb{E}[F(A(s), \xi(s))]$ for any integrable function F. (6.102) thus becomes

$$\langle U_{\alpha}f,g\rangle_{L^{2}(m)} = \int_{0}^{\infty} e^{-rs} \mathbb{E}\left[\int_{\mathbb{R}} e^{-\alpha A_{y}(s)}(fm)(y)(\overline{g}m)(y+\xi(s))dx\right] ds = \langle f,U_{\alpha}g\rangle_{L^{2}(m)},$$

where, for the last equality, we have used (6.101) but where the roles of f and g are switched.

The following lemma shows that P_t is self-adjoint for $\langle \cdot, \cdot \rangle_{L^2(m)}$ on $\mathcal{C}_0(\mathbb{R}) \cap L^2(\mathbb{R}) \subset L^2(m)$.

Lemma 49. Assume that r > 0 and (Condition 1) holds. We have $\langle P_t.f,g \rangle_{L^2(m)} = \langle f,P_t.g \rangle_{L^2(m)}$ for all $t \ge 0$ and $f,g \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$.

Proof. The claim is obvious for t = 0 so let us assume t > 0 and fix $f, g \in C_0(\mathbb{R}) \cap L^2(\mathbb{R})$. By Lemma 48 we get that for any $n \ge 1$,

$$\int_{\mathbb{R}} \left(\left(\frac{n}{t} U_{n/t}\right)^n f \right)(x) \overline{g(x)} m(x) dx = \int_{\mathbb{R}} f(x) \left(\left(\frac{n}{t} U_{n/t}\right)^n \overline{g} \right)(x) m(x) dx.$$
(6.103)

Since f and \overline{g} are in $C_0(\mathbb{R})$, by (1.31) in [6] we have that $(\frac{n}{t}U_{n/t})^n f$ and $(\frac{n}{t}U_{n/t})^n \overline{g}$ converge to respectively $P_t.f$ and $P_t.\overline{g}$ in $(C_0(\mathbb{R}), \|\cdot\|_{\infty})$. Since, by $m \in L^2(\mathbb{R})$ (from (Condition 1)) and Cauchy-Schwartz inequality, $\overline{g}m$ and fm are in $L^1(\mathbb{R})$ we can let n go to infinity on both sides of (6.103) and get $\int_{\mathbb{R}} (P_t.f)(x)\overline{g(x)}m(x)dx = \int_{\mathbb{R}} f(x)(P_t.\overline{g})(x)m(x)dx$. By Lemma 46, $P_t.f$ and $P_t.g$ are in $L^2(m)$ and, by assumption f and g are in $L^2(\mathbb{R}) \subset L^2(m)$. Therefore the last equality can be re-written as $\langle P_t.f, g \rangle_{L^2(m)} = \langle f, P_t.g. \rangle_{L^2(m)}$.

Appendix A. Some facts about the Lévy process ξ

Recall that ξ is a real symmetric Lévy process and that $\psi_{\xi}(\cdot)$ denotes its characteristic exponent. We prove below some properties of ξ , sometimes assuming that (Condition 2) is satisfied.

Lemma 50. Under (Condition 2) we have that, for any r > 0, the potential measure $V_{\xi}^{r}(dx)$ (defined in Section 1.6) has a continuous density $v_{\xi}^{r}(\cdot) \in L^{1}(\mathbb{R}) \cap C_{0}(\mathbb{R})$, and for any $x \in \mathbb{R}$,

$$v_{\xi}^{r}(x) = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{ixy}}{r - \psi_{\xi}(y)} dy = \mathcal{F}^{-1}\left(\frac{1}{r - \psi_{\xi}(2\pi \cdot)}\right)(x).$$
(A.104)

In particular, for any $x \in \mathbb{R}$ we have $0 \le v_{\xi}^{r}(x) \le \|(-\psi_{\xi}(2\pi \cdot) + r)^{-1}\|_{L^{1}(\mathbb{R})} = v_{\xi}^{r}(0).$

As mentioned in the Introduction, the existence, under (Condition 2), of a continuous density for the potential measure is a consequence of the combination of Remark 43.6 and Theorem 43.5 from [46]. However, let us provide a short and direct proof of Lemma 50 for the sake of completeness.

Proof of Lemma 50. According to Proposition 37.4 of [46] we have for any r > 0 and $x \in \mathbb{R}$,

$$\int_{\mathbb{R}} e^{izx} V_{\xi}^{r}(dz) = \frac{1}{r - \psi_{\xi}(x)}.$$
(A.105)

Note that the convention used in [46] for the Fourier transform is different from the one we use here, which is why we do not state (A.105) in term of \mathcal{F} . From (Condition 2) we see that for any r > 0 the right hand side of (A.105) is in $L^1(\mathbb{R})$ so V_{ξ}^r has a continuous density $v_{\xi}^r(\cdot)$ that satisfies

$$v_{\xi}^{r}(x) = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{ixy}}{r - \psi_{\xi}(y)} dy = \int_{\mathbb{R}} \frac{e^{2i\pi xz}}{r - \psi_{\xi}(2\pi z)} dz = \mathcal{F}^{-1}\left(\frac{1}{r - \psi_{\xi}(2\pi \cdot)}\right)(x),$$

where we have used the symmetry of ξ . This is precisely (A.104). Finally, since $v_{\xi}^{r}(\cdot)$ is the density of a finite measure we have $v_{\xi}^{r}(\cdot) \in L^{1}(\mathbb{R})$ and, by (A.104), $v_{\xi}^{r}(\cdot) \in \mathcal{F}^{-1}(L^{1}(\mathbb{R}))$ so $v_{\xi}^{r}(\cdot) \in \mathcal{C}_{0}(\mathbb{R})$.

Lemma 51. Under (Condition 2) we have $Supp(\xi(s)) = \mathbb{R}$ for any s > 0.

Proof. As explained a little below the statement of (Condition 2), that condition implies that ξ is of type C in the sense of Definition 11.9 of [46]. By Theorem 24.10 of [46] we get $Supp(\xi(s)) = \mathbb{R}$ for all s > 0.

Lemma 52. For any t > 0 and $\epsilon > 0$ we have $\mathbb{P}(\sup_{s \in [0,t]} |\xi(s)| < \epsilon) > 0$.

Proof. Let us fix $\eta > 0$. By the Lévy-Ito decomposition we have

$$\xi(s) = \sqrt{A_{\xi}}W(s) + \int_{0}^{s} \int_{[-\eta,\eta]} z\tilde{M}_{1}(ds, dz) + \int_{0}^{s} \int_{\mathbb{R}\setminus[-\eta,\eta]} zM_{2}(ds, dz),$$
(A.106)

where the three terms are independent. W is a standard Brownian motion. $M_1(ds, dz)$ (resp. $M_2(ds, dz)$) is a Poisson random measure on $[0, \infty) \times [-\eta, \eta]$ (resp. $[0, \infty) \times (\mathbb{R} \setminus [-\eta, \eta])$) with intensity measure $\mathbf{1}_{z \in [-\eta, \eta]} ds \times \Pi_{\xi}(dz)$ (resp. $\mathbf{1}_{z \notin [-\eta, \eta]} ds \times \Pi_{\xi}(dz)$), and $\tilde{M}_1(ds, dz) := M_1(ds, dz) - \mathbf{1}_{z \in [-\eta, \eta]} ds \times \Pi_{\xi}(dz)$.

It is well-known that $\mathbb{P}(\sup_{s \in [0,t]} |W(s)| < \epsilon/2\sqrt{A_{\xi}}) > 0$. For the second term in the right-hand side of (A.106), Doob's martingale inequality yields

$$\mathbb{E}\left[\left(\sup_{s\in[0,t]}\int_0^s\int_{[-\eta,\eta]}z\tilde{M}_1(ds,dz)\right)^2\right] \le 4\mathbb{E}\left[\left(\int_0^t\int_{[-\eta,\eta]}z\tilde{M}_1(ds,dz)\right)^2\right] = 4t\int_{[-\eta,\eta]}u^2\Pi_{\xi}(du).$$

Since $\int_{\mathbb{R}} (1 \wedge u^2) \Pi_{\xi}(du) < \infty$, the right-hand side goes to 0 as η goes to 0. For η chosen small enough we thus have $\mathbb{P}(\sup_{s \in [0,t]} | \int_0^s \int_{[-\eta,\eta]} z \tilde{M}_1(ds, dz) | < \epsilon/2) > 0$. Finally, with probability $e^{-t \Pi_{\xi}(\mathbb{R} \setminus [-\eta,\eta])}$, the third term in the right-hand side of (A.106) is null for all $s \in [0,t]$. We conclude that $\mathbb{P}(\sup_{s \in [0,t]} |\xi(s)| < \epsilon) > 0$.

Appendix B. Fourier duality for the partition function: Proof of Remark 4

We work under the assumptions of Remark 4 and the choice $V_{int} := -\log(v_{\xi}^r(\cdot))$ and $U_{mass} := -\log(m(\cdot))$. $v_{\xi}^r(\cdot) \in \mathcal{C}_0(\mathbb{R})$ by Lemma 50 and $m \in L^1(\mathbb{R})$ so, for all $n \geq 2$, Z_n is well-defined and we have

$$Z_n = \int_{\mathbb{R}^n} v_{\xi}^r (y_2 - y_1) \dots v_{\xi}^r (y_n - y_{n-1}) v_{\xi}^r (y_1 - y_n) m(y_1) \dots m(y_n) dy_1 \dots dy_n.$$
(B.107)

Using (B.107), Lemma 50, Fubini's theorem, and the definitions of $\hat{m}(\cdot)$ and \hat{Z}_n we get

$$Z_{n} = \frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} \left(\int_{\mathbb{R}^{n}} \frac{e^{i(y_{2}-y_{1})z_{1}} \times \dots \times e^{i(y_{n}-y_{n-1})z_{n-1}} \times e^{i(y_{1}-y_{n})z_{n}}}{(r-\psi_{\xi}(z_{1})) \times \dots \times (r-\psi_{\xi}(z_{n}))} m(y_{1}) \dots m(y_{n}) dz_{1} \dots dz_{n} \right) dy_{1} \dots dy_{n}$$

$$= \frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} \left(\int_{\mathbb{R}^{n}} \frac{e^{-i(z_{1}-z_{n})y_{1}} \times e^{-i(z_{2}-z_{1})y_{2}} \times \dots \times e^{-i(z_{n}-z_{n-1})y_{n}}}{(r-\psi_{\xi}(z_{1})) \times \dots \times (r-\psi_{\xi}(z_{n}))} m(y_{1}) \dots m(y_{n}) dy_{1} \dots dy_{n} \right) dz_{1} \dots dz_{n}$$

$$= \frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} \frac{\hat{m}(z_{1}-z_{n}) \times \hat{m}(z_{2}-z_{1}) \times \dots \times \hat{m}(z_{n}-z_{n-1})}{(r-\psi_{\xi}(z_{1})) \times \dots \times (r-\psi_{\xi}(z_{n}))} dz_{1} \dots dz_{n} = \frac{\hat{Z}_{n}}{(2\pi)^{n}}.$$

Appendix C. Removal of periodic boundary condition: Proof of Remark 6

We assume that the assumptions of Theorem 5 are satisfied and set $V_{int} := -\log(v_{\xi}^r(\cdot))$ and $U_{mass} := -\log(m(\cdot))$ in the definitions (1.5) and (1.19). From Lemma 50 we have $0 \le v_{\xi}^r(y) \le v_{\xi}^r(0)$ for any $y \in \mathbb{R}$ so $Z_n \le v_{\xi}^r(0)Z_n^f$. Recall from the combination of Lemma 23, Remark 24, and Proposition 34 that $Z_n \sim (v_{\xi}^r(0)\lambda_1^R)^n$ for large n. We thus get that $\liminf_{n\to\infty} Z_n^f/(v_{\xi}^r(0)\lambda_1^R)^n > 0$.

Let us define $g_0 := \int_{\mathbb{R}} m(y) \phi_y^r dy$, where ϕ_y^r is defined in Section 2.1. We also define the operator $H_r \in \mathcal{L}(L^2((-1,1)))$ by $H_r.f := \langle f, g_0 \rangle_{L^2((-1,1))} g_0$. Let $n \ge 4$ and repeat the proof of Lemma 23 to compute $Tr(H_r.R_r^{n-2})$. We get $Tr(H_r.R_r^{n-2}) = \sum_{j\ge 1} (\lambda_j^R)^{n-2} |\langle g_0, a_j \rangle_{L^2((-1,1))}|^2$, with $(a_j)_{j\ge 1}$ as in Section 3.1. Repeating the proof of Proposition 34 we get $Tr(H_r.R_r^{n-2}) = Z_n^f/v_\xi^r(0)^{n-1}$. Combining both expressions we get $Z_n^f/v_\xi^r(0)^{n-1} = \sum_{j\ge 1} (\lambda_j^R)^{n-2} |\langle g_0, a_j \rangle_{L^2((-1,1))}|^2$. Recall from Remark 24 that the eigenvalue λ_1^R of R_r has multiplicity 1 so, if $|\langle g_0, a_1 \rangle_{L^2((-1,1))}|^2 = 0$ then, proceeding as in the proof of Lemma 23 we get $Z_n^f/v_\xi^r(0)^{n-1} = o((\lambda_1^R)^n)$, which contradicts $\liminf_{n\to\infty} Z_n^f/(v_\xi^r(0)\lambda_1^R)^n > 0$. Therefore $|\langle g_0, a_1 \rangle_{L^2((-1,1))}|^2 > 0$ so, proceeding as in the proof of Lemma 23 we get $Z_n^f/v_\xi^r(0)^{n-1} = v_\xi^r(0)(\lambda_1^R)^n/2 + |\langle g_0, a_1 \rangle_{L^2((-1,1))}|^2 = (0,\infty)$, and the result follows.

Acknowledgments: This paper is supported by NSFC grant No. 11688101. The author is grateful to Professor Fuzhou Gong and to Eric Endo for interesting discussions and references.

References

- D. Applebaum. Lévy Processes and Stochastic Calculus. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2 edition, 2009.
- [2] S. Baguley, L. Doering, and A. Kyprianou. General path integrals and stable sdes. Preprint arXiv:2012.07406, 2020.
- [3] C. J. K. Batty. Asymptotic stability of schrödinger semigroups: path integral methods. *Mathematische Annalen*, 292:457 492, 1992.
- [4] J. Bertoin. Lévy Processes. Tracts in Mathematics, Cambridge, 1996.
- [5] J. Bertoin and M. Yor. Exponential functionals of Lévy processes. Probab. Surveys, 2:191–212, 2005.
- [6] B. Bottcher, R. Schilling, and J. Wang. Lévy Matters III. Lecture Notes in Mathematics. Springer, Cham, 2013.
- [7] T. Brox. A one-dimensional diffusion process in a Wiener medium. Ann. Probab., 14(4): 1206–1218, 1986.
- [8] M.-F. Chen. Eigenvalues, Inequalities, and Ergodic Theory. Probability and Its Applications. Springer, London, 2005.
- B. David, J. Fröhlich, and T. Spencer. The random walk representation of classical spin systems and correlation inequalities. *Communications in Mathematical Physics*, 83(1):123 – 150, 1982.
- [10] D.A. Dawson and Z. Li. Skew convolution semigroups and affine markov processes. The Annals of Probability, 34(3):1103–1142, 2006.
- [11] J.-D. Deuschel, G. Giacomin, and D. Ioffe. Large deviations and concentration properties for ∇φ interface models. Probability Theory and Related Fields, 117:49–111, 2000.

- [12] R. Fernández, J Frohlich, and A.D. Sokal. Random Walks, Critical Phenomena, and Triviality in Quantum Field Theory. Springer Berlin, Heidelberg, 1992.
- [13] P.J. Fitzsimmons and J. Pitman. Kac's moment formula and the Feynman-Kac formula for additive functionals of a Markov process. Stochastic Processes and their Applications, 79(1):117 – 134, 1999.
- [14] S. Friedli and Y. Velenik. Statistical Mechanics of Lattice Systems: A Concrete Mathematical Introduction. Cambridge University Press, 2017.
- [15] F.-Z. Gong, Y. Liu, Y. Liu, and D.-J. Luo. Spectral gaps of Schrodinger operators and diffusion operators on abstract Wiener spaces. *Journal of Functional Analysis*, 266(9):5639–5675, 2014.
- [16] F.-Z. Gong and L. Wu. Spectral gap of positive operators and applications. Journal de Mathématiques Pures et Appliquées, 85(2):151–191, 2006.
- [17] D.R. Grey. Asymptotic behaviour of continuous time, continuous state-space branching processes. Journal of Applied Probability, 11(4):669–677, 1974.
- [18] H. He, Z. Li, and W. Xu. Continuous-state branching processes in Lévy random environments. Journal of Theoretical Probability, 31: 1952–1974, 2018.
- [19] A. Lindner J. Bertoin and R. Maller. On continuity properties of the law of integrals of Lévy processes. Séminaire de Probabilités XLI, 1934:137–159, 2008.
- [20] K. Kawazu and H. Tanaka. A diffusion process in a Brownian environment with drift. J. Math. Soc. Japan, 49: 189–211, 1997.
- [21] M. Kolb and M. Savov. A characterization of the finiteness of perpetual integrals of Lévy processes. Bernoulli, 26(2):1453 – 1472, 2020.
- [22] F. Kühn. Perpetual integrals via random time changes. Bernoulli, 25(3):1755 1769, 2019.
- [23] A. Kuznetsov and J. C. Pardo. Fluctuations of stable processes and exponential functionals of hypergeometric lévy processes. Acta Applicandae Mathematicae, 123(1):113–139, 2013.
- [24] A. Kyprianou. Stable lévy processes, self-similarity and the unit ball. ALEA, 15(1):617–690, 2018.
- [25] A. Lambert. Population dynamics and random genealogies. Stochastic Models, 24:163 45, 2008.
- [26] J. Lamperti. Continuous state branching processes. Bulletin of the American Mathematical Society, 73(3):382 – 386, 1967.
- [27] J. Lamperti. Semi-stable Markov processes. I. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, 22: 205–225, 1972.
- [28] J.-F. Le Gall. Some properties of planar Brownian motion. In Paul Louis Hennequin, editor, Ecole d'Eté de Probabilités de Saint-Flour XX - 1990, pages 111–229, Berlin, Heidelberg, 1992. Springer Berlin Heidelberg.
- [29] P.-S. Li. A continuous-state polynomial branching process. *Stochastic Process. Appl.*, 2018.
- [30] P.-S. Li, X. Yang, and X. Zhou. A general continuous-state nonlinear branching process. The Annals of Applied Probability, 29(4):2523 – 2555, 2019.
- [31] K. Maulik and B. Zwart. Tail asymptotics for exponential functionals of Lévy processes. Stochastic Processes and their Applications, 116(2):156 – 177, 2006.
- [32] F. Petit P. Carmona and M. Yor. On the distribution and asymptotic results for exponential functionals of Lévy processes. In *Exponential functionals and principal values related to Brownian motion*, Bibl. Rev. Mat. Iberoamericana, pages 73–130. Rev. Mat. Iberoamericana, Madrid, 1997.
- [33] S. Palau and J. C. Pardo. Branching processes in a Lévy random environment. Acta Applicandae Mathematicae, 153: 55–79, 2018.
- [34] J. C. Pardo, P. Patie, and M. Savov. A Wiener-Hopf type factorization for the exponential functional of Lévy processes. Journal of the London Mathematical Society, 86(3):930, 2012.
- [35] J.C. Pardo and V. Rivero. Self-similar Markov processes. Boletin de la Sociedad Matematica Mexicana.
- [36] J.C. Pardo, V. Rivero, and K. van Schaik. On the density of exponential functionals of Lévy processes. Bernoulli, 19(5A):1938–1964, 11 2013.
- [37] P. Patie. Infinite divisibility of solutions to some self-similar integro-differential equations and exponential functionals of Lévy processes. Ann. Inst. H. Poincaré Probab. Statist., (3):667–684, 08 2009.
- [38] P. Patie. Law of the absorption time of some positive self-similar Markov processes. Ann. Probab., 40(2):765– 787, 2012.
- [39] P. Patie and M. Savov. Extended factorizations of exponential functionals of Lévy processes. *Electron. J. Probab.*, 17:22, 2012.
- [40] P. Patie and M. Savov. Exponential functional of Lévy processes: Generalized Weierstrass products and wiener-hopf factorization. *Comptes Rendus Mathematique*, 351(9-10):393–396, 2013.
- [41] P. Patie and M. Savov. Bernstein-gamma functions and exponential functionals of Lévy processes. *Electronic Journal of Probability*, 23(none):1 101, 2018.
- [42] Daniel Revuz and Marc Yor. Continuous martingales and Brownian motion. Number 293. Springer, 3. ed edition, 1999.

- [43] V. Rivero. Recurrent extensions of self-similar Markov processes and Cramér's condition. Bernoulli, 11(3):471–509, 2005.
- [44] V. Rivero. Recurrent extensions of self-similar Markov processes and Cramér's condition II. Bernoulli, 13(4):1053–1070, 2007.
- [45] V. Rivero. Tail asymptotics for exponential functionals of Lévy processes: The convolution equivalent case. Ann. Inst. H. Poincaré Probab. Statist., 48(4): 1081–1102, 2012.
- [46] K. Sato. Lévy Processes and Infinitely Divisible Distributions. Cambridge studies in advanced mathematics, 1999.
- [47] A. Singh. Rates of convergence of a transient diffusion in a spectrally negative Lévy potential. Ann. Probab., 36: 279–318, 2008.
- [48] G. Véchambre. Exponential functionals of spectrally one-sided Lévy processes conditioned to stay positive. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 55(2):620–660, 2019.
- [49] G. Véchambre. General self-similarity properties for markov processes and exponential functionals of Lévy processes. Journal of Theoretical Probability, 35(4):2083–2144, 2022.
- [50] G. Véchambre. Almost sure behavior for the local time of a diffusion in a spectrally negative Lévy environment. Journal of Theoretical Probability, 36(2):876–925, 2023.
- [51] H. von Dreifus, A. Klein, and J.F. Perez. Taming Griffiths' singularities: Infinite differentiability of quenched correlation functions. *Commun. Math. Phys.*, 170:21–39, 1995.
- [52] F.-Y. Wang. Functional Inequalities, Markov Semigroups and Spectral Theory. The Science Series of the Contemporary Elite Youth. Elsevier, San Diego, 2005.
- [53] J. Wang. A simple approach to functional inequalities for non-local Dirichlet forms. ESAIM: PS, 18:503–513, 2014.

¹Academy of Mathematics and Systems Science, Chinese Academy of Sciences, No. 55, Zhongguancun East Road, Haidian District, Beijing, China

Email address: vechambre@amss.ac.cn