Statistics of Random Binning Based on Tsallis Divergence

1

Masoud Kavian, Mohammad M. Mojahedian, Mohammad H. Yassaee, Mahtab Mirmohseni, Mohammad Reza Aref

Abstract

Random binning is a widely utilized tool in information theory, finding applications in various domains. In this paper, we focus on the output statistics of random binning (OSRB) using the Tsallis divergence T_{α} . Our investigation encompasses all values of α within the range of $(0,\infty)$. The proofs provided in this paper cover both the achievability and converse aspects. To accommodate the unbounded nature of T_{∞} , we analyze the OSRB framework using the Rényi's divergence criterion with the order of infinity, denoted as D_{∞} . During our exploration of OSRB, we encounter a specific form of Rényi's conditional entropy and delve into its properties. Additionally, we demonstrate the effectiveness of this framework in establishing achievability results for wiretap channel, where Tsallis divergence serves as a security measure.

I. Introduction

The concepts of distance between probability measures and correlation metrics are closely intertwined and widely utilized in information theory and machine learning. These metrics play a crucial role in various applications, such as quantifying security levels [1]–[9] and bounding generalization errors [10]–[13]. Several well-known distance measures are commonly employed, including the total variation distance, KL divergence, Rényi divergence, and Tsallis divergence. The total variation distance is a popular metric defined as

$$||p(x) - q(x)||_{\text{TV}} = \frac{1}{2} \sum_{x} |p(x) - q(x)|.$$
 (1)

KL divergence, another widely used measure, is defined as

$$D(p(x) \parallel q(x)) = \sum_{x} p(x) \log\left(\frac{p(x)}{q(x)}\right). \tag{2}$$

Rényi divergence generalizes KL divergence and is defined as

$$D_{\alpha}(p(x) \parallel q(x)) = \frac{1}{\alpha - 1} \log \left(\sum_{x} p(x)^{\alpha} q(x)^{1 - \alpha} \right), \tag{3}$$

where α is a parameter that determines the order of the divergence.

Tsallis divergence is another generalization of relative entropy, given by

$$T_{\alpha}(p(x) \parallel q(x)) = \frac{1}{\alpha - 1} \left(\sum_{x} p(x)^{\alpha} q(x)^{1 - \alpha} - 1 \right).$$
 (4)

Correlation measures also play a significant role in various scientific domains. Two well-known measures are Shannon's mutual information and Rényi mutual information. Shannon's mutual information is defined as

$$I(X;Y) = \sum_{x,y} p(x,y) \log \frac{p(x,y)}{p(x)p(y)}.$$

Rényi mutual information of order α , proposed by Sibson, is defined as

$$I_{\alpha}(X;Y) = \frac{\alpha}{\alpha - 1} \log \left(\sum_{y} \left[\sum_{x} p(x)p(y|x)^{\alpha} \right]^{1/\alpha} \right),$$

where α belongs to the range $(0,1) \cup (1,\infty)$.

These different metrics exhibit various relationships with each other, with some metrics being stronger or weaker than others. Bounding one metric often leads to bounding another. For instance, Pinsker's inequality states that mutual information is a stronger measure than total variation distance, as shown by the inequality:

$$||p(x,y) - p(x)p(y)||_{\text{TV}} \le \sqrt{\frac{1}{2}I(X;Y)}.$$
 (5)

Moreover, Rényi mutual information is a non-decreasing function of its order, implying that any upper bound on Rényi mutual information with an order greater than 1 yields a bound on mutual information.

Random binning has long been recognized as a valuable tool in information theory, particularly for establishing achievability results. One essential technique for analyzing random binning is the method introduced in [14]. This work investigates the output statistics of random binning (OSRB) and demonstrates that, when the binning rate is below a certain threshold, the bin index variable becomes asymptotically independent from other variables as the block length increases, based on the total variation criterion. In [15], the asymptotic analysis of OSRB is extended to a stronger measure of mutual information.

This paper adopts Tsallis divergence as a metric, which is a stronger measure compared to Rényi divergence, Rényi mutual information, and total variation distance. The asymptotic analysis of OSRB is conducted based on this measure. In this context, the binning rate is constrained by a conditional Rényi entropy. Various definitions of conditional Rényi entropy have been proposed [16]-[21]. The conditional Rényi entropy introduced in this paper is based on the framework of OSRB analysis using Tsallis divergence, giving it operational significance. Furthermore, the properties of the introduced conditional Rényi entropy are thoroughly investigated in this work. We explore the OSRB framework by considering T_{α} across the entire range of $\alpha \in (0, \infty)$, providing both achievability and converse proofs. For the case of $\alpha = \infty$, where T_{α} is unbounded, we utilize D_{∞} as a substitute for the OSRB analysis. We present our OSRB theorems in three distinct scenarios. Firstly, we consider a scenario where the binned sequence is generated independently and identically distributed (i.i.d.). Secondly is randomly chosen from an ϵ -typical set. Lastly, we investigate a scenario where the binned sequence is mapped form an ϵ -typical set and then passed from non-memoryless virtual channel. We analyze the second and third cases as determinstic and stochastic encoders, respectively. By utilizing the OSRB theorem for determinstic and stochastic cases, we are able to achieve higher achievable rates. In particular, we focus on the wiretap channel and derive a secure rate using the Tsallis divergence and D_{∞} as the security criterion. Our results can be seen as an extension of work presented in [2], where the wiretap secure rate is computed using Rényi divergence with a parameter $\alpha \in (0,2] \cup \{\infty\}$ the measure of secrecy. Moreover, we present an applications of our OSRB framework. The application is in establishing the achievability rate region for the wiretap channel using Tsallis divergence and D_{∞} as a secrecy criterion.

The remaining sections of this paper are structured as follows. Section II provides the necessary preliminaries. In Section III, we introduce a specific type of conditional Rényi entropy and investigate its properties. Theorems for asymptotic OSRB based on the Tsallis measure criterion are presented in Section IV. Section V explores the applications of Tsallis-based OSRB in analyzing the achievability rate regions of the wiretap channel problem. Finally, Section VI presents the concluding remarks of the paper.

II. PRELIMINARIES

This section begins by introducing the relevant notations and providing a definition of random binning. We then identify a counting problem and discuss the necessary properties of Tsallis divergence.

A. Notations and Definitions

Random variables are denoted by capital letters and their values by lowercase letters. Alphabet sets of random variables are shown in calligraphic font. The n-ary Cartesian power of a set \mathcal{X} is denoted as \mathcal{X}^n . The notation $p^U(x)$ represents the uniform distribution over the set \mathcal{X} . We use $[\ell]$ to denote the set $\{1, 2, \ldots, \ell\}$. We use $\mathbb{I}\{\cdot\}$ to denote the indicator function; it is equal to one if the condition inside $\{\cdot\}$ holds; otherwise, it is zero. For a real number x, its integer part and fractional part are denoted by $\lfloor x \rfloor$ and $\{x\}$, respectively. All the logarithms in this paper are in base two.

Definition 1 (Random binning). In the process of random binning, each realization of a random variable is randomly assigned to a bin index. Therefore, random binning can be viewed as a random function denoted as $\mathcal{B}: \mathcal{X} \to [M]$, where each symbol $x \in \mathcal{X}$ is uniformly and independently mapped to a symbol $b \in [M]$. Consider two dependent random variables X and Z with a joint probability mass function p(x, z). By applying random binning, we map the set \mathcal{X} to [M]. As a result, the induced random probability mass function on the set $\mathcal{X} \times \mathcal{Z} \times [M]$ can be expressed as

$$P(x, z, b) = p_{X,Z}(x, z) \, \mathbb{1}\{\mathcal{B}(x) = b\}. \tag{6}$$

Consequently, we obtain the conditional probability mass function as

$$P(b|z) = \sum_{x} p(x|z) \, \mathbb{1}\{\mathcal{B}(x) = b\}.$$
 (7)

Here, the use of capital letter P indicates that the induced probability mass function on x, z, b is random.

Definition 2. A channel p(x|z) is referred to as a singleton channel if, for any x_1, x_2, z_1, z_2 such that $p(z_1)p(z_2) > 0$, the following condition holds.

$$p(x_1|z_1) = p(x_2|z_2). (8)$$

On the other hand, if the above condition is not satisfied, the channel is classified as a non-singleton channel.

B. Counting

The count of positive integer solutions for the linear equation

$$x_1 + x_2 + \ldots + x_\ell = \alpha, \tag{9}$$

is given by

$$\binom{\alpha - 1}{\ell - 1} = \frac{(\alpha - 1)!}{(\ell - 1)!(\alpha - \ell)!}.$$
 (10)

It is evident that the following inequality holds.

$$\binom{\alpha - 1}{\ell - 1} \le \alpha^{\alpha}.$$
 (11)

C. Some useful properties of Tsallis divergence

By using the inequality $\log x \le x - 1$, we can establish the following inequalities for Tsallis and Rényi divergences.

$$T_{\alpha}(p(x) \parallel q(x)) \ge D_{\alpha}(p(x) \parallel q(x)) \qquad \alpha \in (1, \infty)$$
(12)

$$D_{\alpha}(p(x) \parallel q(x)) \ge T_{\alpha}(p(x) \parallel q(x)) \qquad \alpha \in (0, 1). \tag{13}$$

Moreover, the limit values of the Tsallis divergence are given by

$$\lim_{\alpha \to 1} T_{\alpha}(p(x) \parallel q(x)) = D(p(x) \parallel q(x)), \tag{14}$$

and

$$\lim_{\alpha \to \infty} T_{\alpha}(p(x) \parallel q(x)) = \infty, \tag{15}$$

for two different distributions p(x) and q(x).

Due to the unbounded nature of the Tsallis divergence as α approaches infinity, it is less appealing for investigation. Therefore, the paper focuses on analyzing D_{∞} instead, defined as

$$D_{\infty}(p(x) \parallel q(x)) = \log\left(\max_{x} \frac{p(x)}{q(x)}\right). \tag{16}$$

In comparing the definition of T_{α} and D_{α} in equations (3-4) for two different distributions on X^n , it can be observed that they exhibit similar behavior for $\alpha \in (0, \infty)$ as n approaches to ∞ . This observation suggests that if one of them converges to 0 or ∞ , the other also converges accordingly.

III. CONDITIONAL RÉNYI ENTROPY

In this section, we present a candidate for the conditional Rényi entropy and thoroughly examine its properties. This particular entropy emerges naturally when bounding the output of random binning using the Tsallis divergence. Since the properties of this entropy will be utilized in the subsequent theorems, we introduce it earlier. Although the conditional Rényi entropy we introduce in this paper has been previously discussed in [16] and some of its properties have been investigated in [21], we provide a comprehensive overview of all its properties here for the sake of completeness.

Lemma 1. Consider the function

$$\tilde{H}_{\alpha}(X|Z) = \frac{1}{1-\alpha} \log \left(\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z) \right). \tag{17}$$

Then it has the following properties

1) Let p(x,z) = p(x)p(z), then

$$\tilde{H}_{\alpha}(X|Z) = H_{\alpha}(X), \tag{18}$$

where $H_{\alpha}(X)$ is the Rényi entropy of order α .

- 2) $\tilde{H}_{\alpha}(X|Z)$ is decreasing in $\alpha \in \mathbb{R}$. To be more precise, it is strictly decreasing for non-singleton channel p(x|z).
- 3) $\lim_{\alpha \to 1} \tilde{H}_{\alpha}(X|Z) = H(X|Z).$
- 4) Data processing inequality for $\alpha \in (1, \infty)$,

$$\tilde{H}_{\alpha}(X|Y) \le \tilde{H}_{\alpha}(X|Z),$$
(19)

such that X - Y - Z forms a Markov chain.

5) Suppose $\alpha \in (1, \infty)$ and $\lambda \in [0, 1]$, then $\tilde{H}_{\alpha}(X|Z)$ for fixed p(x|z) is a convex function of p(z). More precisely, we have

$$\frac{1}{1-\alpha} \log \left(\sum_{z} \left(\lambda p_{\lambda}(z) + \bar{\lambda} p_{\bar{\lambda}}(z) \right) \cdot \sum_{x} p^{\alpha}(x|z) \right) \\
\leq \frac{\lambda}{1-\alpha} \log \left(\sum_{z} p_{\lambda}(z) \sum_{x} p^{\alpha}(x|z) \right) + \frac{\bar{\lambda}}{1-\alpha} \log \left(\sum_{z} p_{\bar{\lambda}}(z) \sum_{x} p^{\alpha}(x|z) \right), \quad (20)$$

where $\bar{\lambda} = 1 - \lambda$ and $p_{\lambda}(z)$ and $p_{\bar{\lambda}}(z)$ are two different probability mass functions over the alphabet \mathcal{Z} .

6) In the limit as $\alpha \to \pm \infty$,

$$\lim_{\alpha \to \infty} \tilde{H}_{\alpha}(X|Z) = \log\left(\frac{1}{\max_{x,z} p(x|z)}\right)$$
 (21)

$$\lim_{\alpha \to -\infty} \tilde{H}_{\alpha}(X|Z) = \log \left(\frac{1}{\min_{x,z} p(x|z)} \right). \tag{22}$$

7) Let (X^n, Z^n) be i.i.d. random variables distributed according to p(x, z), i.e.,

$$p(x^n, z^n) = \prod_{i=1}^n p(x_i, z_i).$$
(23)

Then,

$$\tilde{H}_{\alpha}(X^n|Z^n) = \sum_{i=1}^n \tilde{H}_{\alpha}(X_i|Z_i) = n\tilde{H}_{\alpha}(X|Z). \tag{24}$$

- 8) $\tilde{H}_{\alpha}(X|Z)$ is non-negative.
- 9) For singleton channel p(x|z) and any $\alpha_1 \neq \alpha_2 \in \mathbb{R}$, we have

$$\tilde{H}_{\alpha_1}(X|Z) = \tilde{H}_{\alpha_2}(X|Z). \tag{25}$$

The proofs of the properties are given in Appendix A.

Lemma 2 (Harris' inequality). Let $f, g : \mathbb{R} \to \mathbb{R}$ be non-decreasing functions, and let X be a real-valued random variable taking values in \mathbb{R} . Then

$$\mathbb{E}[f(X)g(X)] \ge \mathbb{E}[f(X)]\,\mathbb{E}[g(X)],\tag{26}$$

which by induction and paying attention to the fact that the multiplication of non-decreasing and non-negative functions is non-decreasing, we have

$$\mathbb{E}\left[\prod_{i=1}^{\ell} f_i(X)\right] \ge \prod_{i=1}^{\ell} \mathbb{E}[f_i(X)]$$
(27)

when $f_i(X)$ is non-decreasing and non-negative.

IV. STATISTICS OF RANDOM BINNING

In the subsequent analysis, we examine the output statistics of random binning based on the Tsallis divergence criterion. We consider two separate cases, namely $\alpha \in (0,1)$ and $\alpha \in (1,\infty)$, and present two theorems that encompass both achievability and converse proofs. Additionally, we provide a theorem specifically for the case of $\alpha = \infty$, where we utilize D_{∞} as a measure.

A. Statistics of random binning over i.i.d random variable

Theorem 1. Consider the scenario where $\alpha \in (1, \infty)$ and $\mathcal{B} : \mathcal{X}^n \to [2^{nR}]$ represents the set of all random mappings with a rate of R, satisfying

$$R < \tilde{H}_{\alpha}(X|Z). \tag{28}$$

As n approaches infinity, we observe that

$$\mathbb{E}_{\mathcal{B}}\left[T_{\alpha}\left(P(b,z^{n}) \parallel p^{U}(b) p(z^{n})\right)\right] \to 0. \tag{29}$$

Furthermore, if $R > \tilde{H}_{\alpha}(X|Z)$, then we have

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\Big(P(b,z^n) \parallel p^{U}(b) p(z^n)\Big)\Big] \to \infty. \tag{30}$$

Proof. We begin by proving the theorem for $\alpha \in \mathbb{N}$, and then we proceed to generalize it. It is important to note that the proof initially focuses on the single-shot scenario and later extends to the asymptotic regime. By considering the definition of Tsallis divergence in equation (4), we obtain

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\Big(P(b,z) \parallel p^{U}(b) p(z)\Big)\Big] = \frac{1}{\alpha - 1} \left(\mathbb{E}_{\mathcal{B}}\left[M^{\alpha - 1} \sum_{b,z} P^{\alpha}(b|z) p(z)\right] - 1\right)$$
(31)

$$= \frac{1}{\alpha - 1} \left(\sum_{z} p(z) \left(M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[P^{\alpha}(b = 1|z) \right] \right) - 1 \right), \tag{32}$$

where (32) is due to the symmetry and linearity of expectation.

Using the multinomial expansion, we now compute the expectation by substituting (7) in (32).

$$M^{\alpha} \mathbb{E}_{\mathcal{B}} \Big[p(b=1|z)^{\alpha} \Big] \tag{33}$$

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\left(\sum_{x} p(x|z) \mathbb{1} \{ \mathcal{B}(x) = 1 \} \right)^{\alpha} \right]$$
(34)

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\sum_{x_1, \dots, x_{\alpha}} \prod_{i=1}^{\alpha} p(x_i|z) \mathbb{1} \{ \mathcal{B}(x_i) = 1 \} \right]$$
(35)

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\sum_{x_1} p^{\alpha}(x_1|z) \mathbb{1} \{ \mathcal{B}(x_1) = 1 \} \right]$$

$$+ \left(\sum_{x_1 \neq x_2} p^{\alpha - 1}(x_1|z) p(x_2|z) \cdot \prod_{i=1}^{2} \mathbb{1} \{ \mathcal{B}(x_i) = 1 \} \right]$$

$$+ \sum_{x_1 \neq x_2} p^{\alpha - 2}(x_1|z) p^2(x_2|z) \cdot \prod_{i=1}^{2} \mathbb{1} \{ \mathcal{B}(x_i) = 1 \} + \dots \right]$$

$$+ \left(\sum_{x_1 \neq x_2 \neq x_3} p^{\alpha - 2}(x_1 | z) p(x_2 | z) p(x_3 | z) \cdot \prod_{i=1}^{3} \mathbb{1} \{ \mathcal{B}(x_i) = 1 \} \right)$$

$$+ \sum_{x_1 \neq \dots \neq x_3} \prod_{i=1}^{\alpha} p(x_i | z) \mathbb{1} \{ \mathcal{B}(x_i) = 1 \}$$

$$+ \sum_{x_1 \neq \dots \neq x_3} \prod_{i=1}^{\alpha} p(x_i | z) \mathbb{1} \{ \mathcal{B}(x_i) = 1 \}$$

$$= M^{\alpha - 1} \sum_{x_1} p^{\alpha}(x_1 | z)$$

$$+ M^{\alpha - 2} \Big[\sum_{x_1 \neq x_2} p^{\alpha - 1}(x | z) p(x_2 | z) + \sum_{x_1 \neq x_2} p^{\alpha - 2}(x_1 | z) p^2(x_2 | z^n) + \cdots \Big]$$

$$+ M^{\alpha - 3} \Big[\sum_{x_1 \neq x_2 \neq x_3} p^{\alpha - 2}(x | z) p(x_2 | z) p(x_3 | z)$$

$$+ \sum_{x_1 \neq x_2 \neq x_3} p^{\alpha - 3}(x | z) p^2(x_2 | z) p(x_3 | z) + \cdots \Big]$$

$$+ \cdots + \sum_{x_1 \neq x_2 \neq \dots \neq x_n} \prod_{i=1}^{\alpha} p(x_i | z)$$

$$\leq M^{\alpha - 1} \sum_{x} p^{\alpha}(x | z)$$

$$+ \left[M^{\alpha - 2} \Big(\sum_{x} p^{\alpha - 1}(x | z) \Big) + M^{\alpha - 2} \Big(\sum_{x} p^{\alpha - 2}(x | z) \Big) \Big(\sum_{x} p^2(x | z) \Big) + \cdots \Big]$$

$$+ \left[M^{\alpha - 2} \Big(\sum_{x} p^{\alpha - 2}(x | z) \Big) + M^{\alpha - 3} \Big(\sum_{x} p^{\alpha - 3}(x | z) \Big) \Big(\sum_{x} p^2(x | z) \Big) + \cdots \Big]$$

$$+ \cdots + 1,$$

$$(39)$$

where (36) comes from the fact that the set \mathcal{X}^{α} can be partitioned as $\bigcup_{\ell=1}^{\alpha} \mathcal{A}_{\ell}$ where \mathcal{A}_{ℓ} is the set of α -tuples in the following form

$$\mathcal{A}_{\ell} = \left\{ (\underbrace{x_1, \dots, x_1}_{\alpha_1}, \underbrace{x_2, \dots, x_2}_{\alpha_2}, \dots, \underbrace{x_{\ell}, \dots, x_{\ell}}_{\alpha_{\ell}}) : x_j \in \mathcal{X}, j \in [\ell], x_i \neq x_j, i \neq j \right\}, \tag{40}$$

with $\alpha_1 + \ldots + \alpha_\ell = \alpha$. In other words, \mathcal{A}_ℓ means that we divide the set \mathcal{X}^α into ℓ parts and the data inside each part are equal, while they are not equal to the data of other parts.

Counting how the power α is divided into ℓ parts yields an upper bound of (11) for the cardinality of the set A_{ℓ} . The equation (37) is a consequence of the fact that random binning

maps independently and uniformly, or equivalently

$$\mathbb{E}_{\mathcal{B}} \left[\prod_{i=1}^{k} \mathbb{1} \{ \mathcal{B}(x_i) = 1 \} \right] = \frac{1}{M^k}.$$
 (41)

(38) arises from

$$\sum_{x_1 \neq x_2 \neq \dots \neq x_\ell} \prod_{i=1}^{\ell} p^{\alpha_i}(x_i|z) \le \prod_{i=1}^{\ell} \sum_{x} p^{\alpha_i}(x|z).$$
 (42)

In addition, we benefited from

$$\sum_{x_1 \neq \dots \neq x_{\alpha}} \prod_{i=1}^{\alpha} p(x_i|z) \le \sum_{x_1, \dots, x_{\alpha}} \prod_{i=1}^{\alpha} p(x_i|z) = 1.$$

$$(43)$$

The terms in (39), by Harris' inequality, can be further bounded as

$$M^{\alpha-\ell} \cdot \prod_{i=1}^{\ell} \left(\sum_{x} p^{\alpha_i}(x|z) \right) = M^{\alpha-\ell} \cdot \prod_{i=1}^{\ell} \mathbb{E}_{p_{X|Z=z}} \left[p^{\alpha_i - 1}(X|z) \right]$$
(44)

$$\leq M^{\alpha-\ell} \cdot \mathbb{E}_{p_{X|Z=z}} \left[\prod_{i=1}^{\ell} p^{\alpha_i - 1}(X|z) \right] \tag{45}$$

$$= M^{\alpha - \ell} \sum_{x} p^{\alpha - \ell + 1}(x|z), \tag{46}$$

where $\sum_{i=1}^{\ell} \alpha_i = \alpha$. By substituting these upper bounds into (32), we obtain the following upper-bound terms for Tsallis divergence.

$$M^{\alpha-\ell} \sum_{z} p(z) \sum_{x} p^{\alpha-\ell+1}(x|z), \qquad \ell = 1, \dots, \alpha - 1.$$
 (47)

Let (X^n, Z^n) be i.i.d. random variables distributed according to p(x, z). Suppose that we randomly (and uniformly) bin the set \mathcal{X}^n into $M = 2^{nR}$ bins. Then, with the increase of n, each term forming the upper bound in (47) tends to zero if the rate applies in the following inequality.

$$R < \tilde{H}_{\alpha-\ell+1}(X|Z), \qquad \ell = 1, \dots, \alpha - 1, \tag{48}$$

or equivalently

$$R < \min_{\ell} \tilde{H}_{\alpha - \ell + 1}(X|Z) \stackrel{(a)}{=} \tilde{H}_{\alpha}(X|Z), \tag{49}$$

which (a) results from the non-increasing property of $\tilde{H}_{\alpha}(X|Z)$.

It should be noted that for the singleton case, according to property 9, the value of $\tilde{H}_{\alpha}(X|Z)$ is constant.

Therefore, with the increase of n, the upper bound (39) tends to 1 and subsequently (32) tends to zero. By letting n go to infinity, the proof of the theorem for $\alpha > 1 \in \mathbb{N}$ is complete.

The generalization of the achievability proof to real values of α and the converse part of the proof can be found in Appendix B.

Remark 1. Inequality (12) shows that considering D_{α} instead of T_{α} in the case of $\alpha \in (1, \infty)$, the result of Theorem 1 will be valid.

Remark 2. In case of $\alpha = 1$, Theorem 1 shows that if R < H(X|Z) holds, then

$$\mathbb{E}_{\mathcal{B}}[I(Z;B)] \to 0. \tag{50}$$

This case can also be deduced from the results of [14], [15]. More specifically, it has been proven in [14] that R < H(X|Z) gives results

$$\mathbb{E}_{\mathcal{B}} \| p(z, b) - p(z) p^{U}(b) \|_{\text{TV}} \to 0.$$
 (51)

Furthermore, in [15], it is shown that (51) approaches zero exponentially fast, which subsequently results in (50).

The following theorem holds for $\alpha \in (0, 1)$.

Theorem 2. Let $\alpha \in (0,1)$. Then

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\Big(P(b,z^n) \parallel p^{U}(b) p(z^n)\Big)\Big] \to 0$$
 (52)

holds if and only if the binning rate satisfies R < H(X|Z).

The proof is provided in Appendix C.

B. Statistics of random binning for special type

Theorem 3. Let consider

$$\mathcal{T}_{\epsilon}^{n}(p_X) = \left\{ x^n : \left| \frac{1}{n} \sum_{i=1}^{n} \mathbb{1}\{x_i = x\} - p(x) \right| < \epsilon \right\}$$
 (53)

and X^n , which X^n distributed over $\mathcal{T}_{\epsilon}^n(p_X)$ with distribution of

$$\tilde{p}(x^n) = \frac{\prod_{i=1}^n p(x_i)}{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \prod_{i=1}^n p(x_i)}$$
(54)

$$1 - \delta_n \le \sum_{x^n \in \mathcal{T}_n^n(p_X)} \prod_{i=1}^n p(x_i)$$

$$\tag{55}$$

and

$$p_{Z^n|X^n} = \prod_{i=1}^n p_{Z_i|X_i} \tag{56}$$

consider the scenario where $\alpha \in (1, \infty)$ and $\mathcal{B} : \mathcal{T}_{\epsilon}^n(p_X) \to [2^{nR}]$ represents the set of all random mappings on a special type with a rate of R, satisfying

$$R < H(X) - \sum_{x} p(x) D_{\alpha}(p(z|x) \parallel p(z)).$$
 (57)

As n approaches infinity and $\epsilon \to 0$, we observe that

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\Big(P(b,z^n) \parallel p^{U}(b) q(z^n)\Big)\Big] \to 0 \tag{58}$$

which $p(z) = \sum_{x} p(x)p(z|x)$ and $q(z^n) = \prod_{i=1}^{n} p(z_i)$.

Proof. Similar the steps in proof of Theorem 1 we start the proof with $\alpha \in \mathbb{N}$, then we generalize that. Unlike the proof of Theorem 1 we represent this proof in asymptotic regime. Most of the steps in this part are similar to 1 based on expansion. Let consider the definition of Tsallis divergence in (4), so we have

$$\mathbb{E}_{\mathcal{B}}\left[T_{\alpha}\left(P(b,z^{n}) \parallel p^{U}(b)q(z^{n})\right)\right] = \frac{1}{\alpha-1}\left(\mathbb{E}_{\mathcal{B}}\left[M^{\alpha-1}\sum_{b,z^{n}}q(z^{n})\left(\frac{p(b,z^{n})}{q(z^{n})}\right)^{\alpha}\right] - 1\right)$$
(59)

$$= \frac{1}{\alpha - 1} \left(\sum_{z^n} q(z^n) M^{\alpha} \mathbb{E}_{\mathcal{B}} \left(\frac{p(1, z^n)}{q(z^n)} \right)^{\alpha} - 1 \right), \tag{60}$$

where (60) comes from symmetry and linearity of expectation.

Multinomial expansion combining by substituting (7) in (60) implises that

$$M^{\alpha} \mathbb{E}_{\mathcal{B}} \left(\frac{p(1, z^n)}{q(z^n)} \right)^{\alpha} \tag{61}$$

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\left(\frac{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} p(x^n, z^n) \mathbb{1} \{ \mathcal{B}(x) = 1 \}}{q(z^n)} \right)^{\alpha} \right]$$
 (62)

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\left(\frac{\sum_{x_1^n, \dots, x_{\alpha}^n} \prod_{i=1}^{\alpha} p(x_i^n, z^n) \mathbb{1} \{ \mathcal{B}(x_i^n) = 1 \}}{q^{\alpha}(z^n)} \right) \right]$$

$$(63)$$

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\sum_{x_{1}^{n} \in \mathcal{T}_{\epsilon}^{n}(p_{X})} \frac{p^{\alpha}(x_{1}^{n}, z^{n})}{q^{\alpha}(z^{n})} \mathbb{1} \{ \mathcal{B}(x_{1}^{n}) = 1 \} + \left(\sum_{x_{1}^{n} \neq x_{1}^{n}} \frac{p^{\alpha-1}(x_{1}^{n}, z^{n}) p(x_{2}^{n}, z^{n}) \cdot \prod_{i=1}^{2} \mathbb{1} \{ \mathcal{B}(x_{i}^{n}) = 1 \}}{q^{\alpha}(z^{n})} \right]$$

(68)

$$+ \sum_{x_{1}^{n} \neq x_{2}^{n}} \frac{p^{\alpha-2}(x_{1}^{n}, z^{n})p^{2}(x_{2}^{n}, z^{n}) \cdot \prod_{i=1}^{2} \mathbb{1}\{\mathcal{B}(x_{i}^{n}) = 1\}}{q^{\alpha}(z^{n})} + \dots$$

$$+ \left(\sum_{x_{1}^{n} \neq x_{2}^{n} \neq x_{3}^{n}} \frac{p^{\alpha-2}(x_{1}^{n}, z^{n})p(x_{2}^{n}, z^{n})p(x_{3}^{n}, z^{n}) \cdot \prod_{i=1}^{3} \mathbb{1}\{\mathcal{B}(x_{i}^{n}) = 1\}}{q^{\alpha}(z^{n})} \right)$$

$$+ \frac{\sum_{x_{1}^{n} \neq x_{2}^{n} \neq x_{3}^{n}} p^{\alpha-3}(x_{1}^{n}, z^{n})p^{2}(x_{2}^{n}, z^{n})p(x_{3}^{n}, z^{n}) \cdot \prod_{i=1}^{3} \mathbb{1}\{\mathcal{B}(x_{i}^{n}) = 1\}}{q^{\alpha}(z^{n})} \right)$$

$$+ \frac{\sum_{x_{1}^{n} \neq x_{2}^{n} \neq x_{3}^{n}} \prod_{i=1}^{\alpha} p(x_{i}^{n}, z^{n})\mathbf{1}\{\mathcal{B}(x_{i}^{n}) = 1\}}{q^{\alpha}(z^{n})} \right]$$

$$= M^{\alpha-1} \sum_{x_{1}^{n} \neq x_{2}^{n}} \frac{p^{\alpha}(x_{1}^{n}, z^{n})}{q^{\alpha}(z^{n})}$$

$$+ M^{\alpha-2} \left[\sum_{x_{1}^{n} \neq x_{2}^{n} \neq x_{3}^{n}} \frac{p^{\alpha-1}(x_{1}^{n}, z^{n})p(x_{2}^{n}, z^{n})}{q^{\alpha}(z^{n})} + \sum_{x_{1}^{n} \neq x_{2}^{n}} \frac{p^{\alpha-2}(x_{1}^{n}, z^{n})p^{2}(x_{2}^{n}, z^{n})}{q^{\alpha}(z^{n})} + \cdots \right]$$

$$+ M^{\alpha-3} \left[\sum_{x_{1}^{n} \neq x_{2}^{n} \neq x_{3}^{n}} \frac{p^{\alpha-2}(x^{n}, z^{n})p(x_{2}^{n}, z^{n})p(x_{3}^{n}, z^{n})}{q^{\alpha}(z^{n})} + \cdots \right]$$

$$+ \sum_{x_{1}^{n} \neq x_{2}^{n} \neq x_{3}^{n}} \frac{p^{\alpha-3}(x^{n}, z^{n})p^{2}(x_{2}^{n}, z^{n})p(x_{3}^{n}, z^{n})}{q^{\alpha}(z^{n})} + \cdots \right]$$

$$+ \cdots + \sum_{x_{1}^{n} \neq x_{2}^{n} \neq x_{3}^{n}} \frac{\prod_{i=1}^{\alpha} p(x_{i}^{n}, z^{n})}{q^{\alpha}(z^{n})}$$

$$\leq M^{\alpha-1} \sum_{x_{1}^{n} \in T_{2}^{n}(p_{X})} \frac{p^{\alpha}(x^{n}, z^{n})}{q^{\alpha}(z^{n})}$$

$$\leq M^{\alpha-1} \sum_{x_{1}^{n} \in T_{2}^{n}(p_{X})} \frac{p^{\alpha}(x^{n}, z^{n})}{q^{\alpha}(z^{n})}$$

$$\leq M^{\alpha-2} \left(\sum_{x_{1}^{n} \in T_{2}^{n}(p_{X})} \frac{p^{\alpha-1}(x^{n}, z^{n})}{q^{\alpha-1}(z^{n})} \right) \left(\sum_{x_{1}^{n} \in T_{2}^{n}(p_{X})} \frac{p^{2}(x^{n}, z^{n})}{q^{2}(z^{n})} \right) + \cdots \right]$$

$$M^{\alpha-2} \left(\sum_{x_{1}^{n} \in T_{2}^{n}(p_{X})} \frac{p^{\alpha-2}(x^{n}, z^{n})}{q^{\alpha-2}(z^{n})} \right) \left(\sum_{x_{1}^{n} \in T_{2}^{n}(p_{X})} \frac{p^{2}(x^{n}, z^{n})}{q^{2}(z^{n})} \right) + \cdots \right]$$

$$+ \left[(1+2\delta_n)^2 M^{\alpha-3} \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p^{\alpha-2}(x^n, z^n)}{q^{\alpha-2}(z^n)} \right) \right]$$

$$\tag{69}$$

$$+ (1 + 2\delta_n) M^{\alpha - 3} \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p^{\alpha - 3}(x^n, z^n)}{q^{\alpha - 3}(z^n)} \right) \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p^2(x^n, z^n)}{q^2(z^n)} \right) + \cdots \right]$$

$$+\cdots + (1+2\delta_n)^{\alpha},\tag{70}$$

equation (64) is due to the fact that the set \mathcal{X}^{α} can be partitioned as (40) in proof of Theorem 1 for $\alpha \in \mathbb{N}$. The equation (65) is a consequence of the fact that random binning maps independently and uniformly and uses (41-43) for x_i^n instead of x_i .

(68-70) is due to

$$\sum_{\substack{x_1^n \neq \dots \neq x_l^n}} \frac{\prod_{i=1}^l p^{\alpha_i}(x_i^n, z^n)}{q^{\alpha}(z^n)} \le \prod_{i=1}^l \left(\sum_{\substack{x^n \in \mathcal{T}_{\epsilon}^n(p_X)}} \frac{p^{\alpha_i}(x_i^n, z^n)}{q^{\alpha_i}(z^n)} \right)$$
(71)

combining with special case of $(\alpha_i = 1)$

$$\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p(x^n, z^n)}{q(z^n)} = \frac{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \tilde{p}(x^n) p(z^n | x^n)}{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} q(x^n) p(z^n | x^n)}$$
(72)

$$\leq \frac{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \tilde{p}(x^n) p(z^n | x^n)}{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} q(x^n) p(z^n | x^n)} \leq \frac{1}{1 - \delta_n} \leq 1 + 2\delta_n \tag{73}$$

with $q(x^n) = \prod_{i=1}^n p(x_i)$, and $\delta_n \to 0$ for $n \to \infty$.

This is clear that the upper bound obtained in (70), terms are created as follows

$$M^{\alpha-k-1} \sum_{z^n} q(z^n) \sum_{x^n \in \mathcal{T}^n(p_X)} \frac{p^{\alpha-k}(x^n, z^n)}{q^{\alpha-k}(z^n)}, \qquad k = 0, 1, \dots, \alpha - 2.$$
 (74)

Let consider (X^n, Z^n) such as mentioned in Theorem 3. Suppose that we randomly (and uniformly) bin the set \mathcal{X}^n into $M = 2^{nR}$ bins. Then, Lemma (3) implies that with the increase of n, each term forming the upper bound in (74) tends to zero if the rate applies in the following inequality.

$$R < H(X) - \sum_{x} p(x) D_{\alpha - k} (p(z|x) \parallel p(z)), \qquad k = 0, 1, \dots, \alpha - 2,$$
 (75)

or equivalently

$$R < \min_{k} H(X) - D_{\alpha - k} \left(p(z|x) \parallel p(z) \right) \tag{76}$$

$$\stackrel{(a)}{=} H(X) - D_{\alpha} (p(z|x) \parallel p(z)), \tag{77}$$

which (a) results from the increasing property of $D_{\alpha}(p(z|x) \parallel p(z))$ in α .

for correctess of the

$$\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \left[\frac{p(x^n, z^n)}{q(z^n)} \prod_{i=1}^{\ell} \left(\frac{p(x^n, z^n)}{q(z^n)} \right)^{\alpha_i - 1} \right] \ge \left(\frac{1}{1 + 2\delta_n} \right)^{\ell - 1} \prod_{i=1}^{\ell} \left[\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \left(\frac{p(x^n, z^n)}{q(z^n)} \right)^{\alpha_i} \right]$$

$$(78)$$

with $\sum_{i=1}^{\ell} \alpha_i = \alpha$ for every realization of $Z^n = z^n$ let assume $\nu_n(x)$ measure be defined as

$$\nu_n(x^n) = \frac{\frac{p(x^n, z^n)}{q(z^n)}}{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p(x^n, z^n)}{q(z^n)}} = \frac{\frac{p(x^n, z^n)}{q(z^n)}}{\frac{p(z^n)}{q(z^n)}}$$
(79)

HARRIS inequality implies that for $f_i(x) = x^{\alpha_i - 1}$, we have

$$\left(\frac{q(z^n)}{p(z^n)}\right) \sum_{x^n \in \mathcal{T}_n^n(p_X)} \left[\frac{p(z^n, z^n)}{q(z^n)} \prod_{i=1}^{\ell} \left(\frac{p(x^n, z^n)}{q(z^n)} \right)^{\alpha_i - 1} \right] = \mathbb{E}_{\nu_n(x^n)} \left[\prod_{i=1}^{\ell} f_i \left(\frac{p(X^n, z^n)}{q(z^n)} \right) \right]$$
(80)

$$\geq \prod_{i=1}^{\ell} \mathbb{E}_{\nu_n(x^n)} \left[f_i \left(\frac{p(X^n, z^n)}{q(z^n)} \right) \right] = \left(\frac{q(z^n)}{p(z^n)} \right)^{\ell} \prod_{i=1}^{\ell} \left[\sum_{x^n \mathcal{T}_e^n(p_X)} \left(\frac{p(x^n, z^n)}{q(z^n)} \right)^{\alpha_i} \right]$$
(81)

since $\frac{q(z^n)}{p(z^n)} \ge \frac{1}{1+2\delta_n}$ concluded in (73) with $\delta_n \to 0$ for $n \to \infty$.

Therefore, with the increase of n, the upper bound (70) tends to 1 and subsequently (60) tends to zero. By letting n go to infinity and $\epsilon \to 0$, the proof of the theorem for $\alpha > 1 \in \mathbb{N}$ is complete.

The proof for real α 's is in a way similar to the generalization proof of Theorem 1 and hence is omitted.

Remark 3. Let consider $q(z^n) = \prod_{i=1}^n p(z_i)$, converse part of proof for lemma 3 could be concluded exactly by repeating the steps (283-285), replacing $(H(p_X) + \delta_n(\epsilon))$ in (334) instead of $(H(p_X) - \delta_n(\epsilon))$, and

$$\sum_{z^n, x^n \in \mathcal{T}_{\epsilon}^n(p_X)} q(z^n) \tilde{p}(x^n) \frac{p^{\alpha - k}(z^n | x^n)}{q^{\alpha - k}(z^n)} \ge \min_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \sum_{z^n} q(z^n) \frac{p^{\alpha - k}(z^n | x^n)}{q^{\alpha - k}(z^n)}$$
(82)

$$= \sum_{z^n} q(z^n) \frac{p^{\alpha-k}(z^n|\bar{x}^n)}{q^{\alpha-k}(z^n)}$$
(83)

$$\geq \log \prod_{x} \left(\sum_{z} p(z) \frac{p^{\alpha - k}(z|x)}{p^{\alpha - k}(z)} \right)^{n(p(x) - \epsilon)}$$
(84)

$$= \sum_{x} n(p(x) - \epsilon) \log \left(\sum_{z} p(z) \frac{p^{\alpha - k}(z|x)}{p^{\alpha - k}(z)} \right)$$
(85)

instead of (340-341) for simplification respectively.

Remark 4. Achievability and converse part for special type on $\alpha \in (0,1)$ has no gain because is coincided with i.i.d random variable on similar interval hence is omitted.

Theorem 4. As $\alpha = \infty$, consider X^n distributed over $\mathcal{T}_{\epsilon}^n(p_X)$ with distribution of

$$\tilde{p}(x^n) = \frac{\prod_{i=1}^n p(x_i)}{\sum_{x^n \in \mathcal{T}^n(p_X)} \prod_{i=1}^n p(x_i)}$$
(86)

$$1 - \delta_n \le \sum_{x^n \in \mathcal{T}^n(p_X)} \prod_{i=1}^n p(x_i) \tag{87}$$

and

$$p_{Z^n|X^n} = \prod_{i=1}^n p_{Z_i|X_i} \tag{88}$$

if the binning rate satisfies the following condition

$$R < H(X) - \sum_{x} p(x) D_{\infty}(p(z|x) \parallel p(z)),$$
 (89)

then as n approaches infinity and $\epsilon \to 0$, we observe that

$$\mathbb{E}_{\mathcal{B}}\left[D_{\infty}\left(P(b, z^{n}) \parallel p^{U}(b) q(z^{n})\right)\right] \to 0 \tag{90}$$

which $q(z^n) = \prod_{i=1}^n p(z_i)$ and $p(z) = \sum_x p(x)p(z|x)$.

Proof. Similar to Theorem 3, the proof is given for asymptotic regime case by considering the special type of $\mathcal{T}_{\epsilon}^{n}(p_{X})$ versions of the variables.

For any

$$0 < \beta \le \frac{1}{M(1 + 2\delta_n) \exp(-n(H(X) - \delta_n(\epsilon)) + n \sum_{x} (p(x) + \epsilon) D_{\infty}(p(z|x) \parallel p(z)))}, \quad (91)$$

we have

$$\mathbb{E}_{\mathcal{B}} \Big[D_{\infty} \Big(p(b, z^n) \parallel p^{U}(b) \, q(z^n) \Big) \Big] \tag{92}$$

$$= \mathbb{E}_{\mathcal{B}} \left[\log \left(\max_{b, z^n} M \frac{p(b, z^n)}{q(z^n)} \right) \right] \tag{93}$$

$$\leq \log \left(\mathbb{E}_{\mathcal{B}} \left[\max_{b, z^n} M \frac{p(b, z^n)}{q(z^n)} \right] \right)$$
(94)

$$\leq \log \left(\frac{1}{\beta} \log \left(\sum_{b,z^n} \mathbb{E}_{\mathcal{B}} \left[\exp \left(\beta M \frac{p(b,z^n)}{q(z^n)} \right) \right] \right) \right)$$
 (95)

$$= \log \left(\frac{1}{\beta} \log \left(M \sum_{z^n} \mathbb{E}_{\mathcal{B}} \left[\exp \left(\beta M \frac{p(1, z^n)}{q(z^n)} \right) \right] \right) \right)$$
 (96)

$$= \log \left(\frac{1}{\beta} \log \left(M \sum_{z^n} \mathbb{E}_{\mathcal{B}} \left[\exp \left(\beta M \frac{\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} p(x^n, z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \}}{q(z^n)} \right) \right] \right) \right)$$
(97)

$$= \log \left(\frac{1}{\beta} \log \left(M \sum_{z^n} \mathbb{E}_{\mathcal{B}} \left[\exp \left(\beta M \sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p(x^n, z^n)}{q(z^n)} \mathbb{1} \left\{ \mathcal{B}(x^n) = 1 \right\} \right) \right] \right) \right)$$
(98)

$$\leq \log \left(\frac{1}{\beta} \log \left(M | \mathcal{Z}^n | \max_{z^n} \left[\prod_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \left(\frac{\exp\left(M \beta \frac{p(x^n, z^n)}{q(z^n)} \right)}{M} + 1 - \frac{1}{M} \right) \right] \right) \right)$$
(99)

$$\leq \log \left(\frac{1}{\beta} \log \left(M | \mathcal{Z}^n | \exp \left(\max_{z^n} \sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{1}{M} \left(\exp \left(M \beta \frac{p(x^n, z^n)}{q(z^n)} \right) - 1 \right) \right) \right)$$
 (100)

$$\leq \log \left(\frac{1}{\beta} \log(M|\mathcal{Z}^n|) + \frac{1}{\beta} \max_{z^n} \left[\beta \sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p(x^n, z^n)}{q(z^n)} + \beta^2 M \sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p^2(x^n, z^n)}{q^2(z^n)} \right] \right)$$

$$\tag{101}$$

$$\leq \log\left(\frac{1}{\beta}\log(M|\mathcal{Z}^n|) + \frac{1}{\beta}\left[\beta \max_{z^n} \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \frac{p(x^n, z^n)}{q(z^n)}\right)\right)$$
(102)

$$+\beta^{2} M \max_{x^{n} \in \mathcal{T}_{\epsilon}^{n}(p_{X}), z^{n}} \left(\frac{p(x^{n}, z^{n})}{q(z^{n})} \right) \sum_{x^{n} \in \mathcal{T}_{\epsilon}^{n}(p_{X})} \frac{p(x^{n}, z^{n})}{q(z^{n})} \right]$$
 (103)

$$= \log \left(\frac{1}{\beta} \log(M|\mathcal{Z}^n|) + \frac{1}{\beta} \left[\beta \max_{z^n} \left(\frac{p(z^n)}{q(z^n)} \right) \right]$$
 (104)

$$+ \beta^{2} M \max_{x^{n} \in \mathcal{T}_{\epsilon}^{n}(p_{X}), z^{n}} \left(\tilde{p}(x^{n}) \frac{p(z^{n}|x^{n})}{q(z^{n})} \right) \left(\sum_{x^{n} \in \mathcal{T}_{\epsilon}^{n}(p_{X})} \frac{p(x^{n}, z^{n})}{q(z^{n})} \right) \right]$$

$$(105)$$

$$\leq \log\left(1 + 2\delta_n + \frac{1}{\beta}\log(M|\mathcal{Z}^n|)\right) \tag{106}$$

$$+ \left[\beta M \exp\left(-n(H(X) - \delta_n(\epsilon)) + D_{\infty}(p(z^n|x^n) \parallel q(z^n))\right) (1 + 2\delta_n) \right] \right)$$
(107)

 $\leq 2\delta_n + \frac{n}{\beta}\log|\mathcal{Z}| + \frac{1}{\beta}\log M \tag{108}$

$$+ M\beta(1 + 2\delta_n) \exp\left(-n(H(X) - \delta_n(\epsilon)) + n\sum_{x} (p(x) + \epsilon)D_{\infty}(p(z|x) \parallel p(z))\right)$$
 (109)

where

- (94) comes from Jensen's inequality for $f(x) = \log x$.
- (95) follows from the following standard inequality for bounding the expectation of maxi-

mum,

$$\mathbb{E}\left[\max_{k=1,\dots,K} Z_k\right] \le \frac{1}{\beta} \log \left(\sum_{k=1}^K \mathbb{E}\left[\exp(\beta Z_k)\right]\right), \quad \forall \beta > 0.$$
 (110)

- (96) is the consequence of symmetry.
- Putting (7) in (96) and because of that we have an independent and uniform mapping, (99) is the result.
- (100) is due to $1 + x \le \exp(x)$, and (101) utilizes

$$e^x \le 1 + x + x^2, \quad 0 \le x \le 1.$$
 (111)

• (107) uses $\sum_{x^n \in \mathcal{T}_\epsilon^n(p_X)} \frac{p(x^n, z^n)}{q(z^n)} = \frac{p(z^n)}{q(z^n)} \le 1 + 2\delta_n$ concluded in (72-73) combining with

$$-H(X) - \delta_n(\epsilon) \le \frac{1}{n} \log(\tilde{p}(x^n)) \le -H(X) + \delta_n(\epsilon), \qquad x^n \in \mathcal{T}_{\epsilon}^n(p_X)$$
 (112)

and $\delta_n, \delta_n(\epsilon), \epsilon \to 0$ as $n \to \infty$.

• (109) is because of $\log(1+x) \le x$ and

$$D_{\infty}\left(p(z^n|x^n) \parallel q(z^n)\right) = \max_{z^n} \log\left(\frac{p(z^n|x^n)}{q(z^n)}\right)$$
(113)

$$\leq \max_{x^n \in \mathcal{T}_{\epsilon}^n(p_X), z^n} \left(\frac{p(z^n | x^n)}{q(z^n)} \right) \tag{114}$$

$$= \log \prod_{x} \left(\max_{z} \frac{p(z|x)}{p(z)} \right)^{n(p(x)+\epsilon)}$$
 (115)

$$= n \sum_{x} (p(x) + \epsilon) D_{\infty} (p(z|x) \parallel p(z)), \qquad (116)$$

for $x^n \in \mathcal{T}^n_{\epsilon}(p_X)$.

Taking the minimum of (109) over

$$\beta \in [0, \frac{1}{M(1+2\delta_n)} \exp(n(H(X) - \delta_n(\epsilon)) - n \sum_{x} (p(x) + \epsilon) D_{\infty}(p(z|x) \parallel p(z)))]$$
 (117)

we get the following upper bound,

$$\mathbb{E}_{\mathcal{B}}\Big[D_{\infty}\Big(p(b,z^n) \parallel p^{U}(b) \, q(z^n)\Big)\Big] \tag{118}$$

$$\leq \delta_n + \sqrt{2(1+2\delta_n)M\log(M|\mathcal{Z}|^n)\exp\left(-n(H(X)-\delta_n(\epsilon)) + n\sum_{x}(p(x)+\epsilon)D_{\infty}(p(z|x) \parallel p(z))\right)}$$
(119)

provided that

$$\sqrt{\frac{(1+2\delta_n)\log(M|\mathcal{Z}|^n)\exp\left(n(H(X)-\delta_n(\epsilon))-n\sum_x(p(x)+\epsilon)D_\infty(p(z|x)\parallel p(z))\right)}{M}}$$
(120)

$$<\frac{1}{M(1+2\delta_n)}\exp\Big(n(H(X)-\delta_n(\epsilon))-n\sum_{x}(p(x)+\epsilon)D_{\infty}\big(p(z|x)\parallel p(z)\big)\Big),\tag{121}$$

which is true if

$$M(1+2\delta_n)\log(M|\mathcal{Z}|^n)\exp\left(-n(H(X)+\delta_n(\epsilon))+n\sum_{x}(p(x)+\epsilon)D_{\infty}(p(z|x)\parallel p(z))\right)<1.$$
(122)

Turning to the asymptotic regime, we consider (X^n, Z^n) and we set $M = 2^{nR}$. Then if

$$R < \left(H(X) - \delta_n(\epsilon) - \sum_{x} (p(x) + \epsilon) D_{\infty} (p(z|x) \parallel p(z))\right), \tag{123}$$

the condition (122) is satisfied and we have,

$$\mathbb{E}_{\mathcal{B}}\Big[D_{\infty}\Big(p(b,z^n) \parallel p^{U}(b)\,p(z^n)\Big)\Big] \tag{124}$$

$$\leq 2\delta_n + \sqrt{2n(1+2\delta_n)(R+\log(|\mathcal{Z}|))} \exp\left(nR - n\left(H(X) - \delta_n(\epsilon) - \sum_{x} (p(x) + \epsilon)D_{\infty}(p(z|x) \parallel p(z))\right)\right)$$
(125)

which for $n \to \infty$ and $\epsilon \to$, we have $\delta_n, \delta_n(\epsilon) \to 0$, therefore (125) tending exponentially fast to zero.

C. Random binning as stochastic encoder

Theorem 5. Let consider random variables (U, X) with distribution p(u, x), in addition assume joint distribution

$$\tilde{p}(u^n, x^n, z^n) = \tilde{p}(u^n)\tilde{p}(x^n|u^n) \prod_{i=1}^n p(z_i|x_i)$$
(126)

over all realizations (u^n, x^n) with

$$\mathcal{T}_{\epsilon}^{n}(p_{UX}) = \{(u^{n}, x^{n}) : u^{n} \in \mathcal{T}_{\epsilon}^{n}(p_{U}), x^{n} \in \mathcal{T}_{\epsilon}^{n}(X|u^{n})\}$$
(127)

which

$$\mathcal{T}_{\epsilon}^{n}(X|u^{n}) = \left\{ x^{n} : \left| \frac{1}{n} \sum_{i=1}^{n} \mathbb{1}\{(x_{i}, u_{i}) = (x, u)\} - p(x, u) \right| < 2\epsilon \right\}$$
 (128)

$$\mathcal{T}_{\epsilon}^{n}(p_{U}) = \left\{ u^{n} : \left| \frac{1}{n} \sum_{i=1}^{n} \mathbb{1}\{(u_{i}) = (u)\} - p(u) \right| < \epsilon \right\}$$
 (129)

with

$$\tilde{p}(u^n, x^n) = \frac{\prod_{i=1}^n p(u_i) p(x_i | u_i)}{\left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \prod_{i=1}^n p(u_i)\right) \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^n(X | u^n)} \prod_{i=1}^n p(x_i | u_i)\right)}$$
(130)

and

$$R'_{\alpha}(p_{X}, p_{Z|X}, p_{U}) = \max_{\substack{T_{Z|XU} \\ \sum_{z} t(x|u,z) t(z|u) p(u) = p(u,x)}} \left[-\frac{\alpha}{\alpha - 1} D(t(z|u, x) \parallel p(z|x) | p(x, u)) + D(t(z|u) \parallel p(z) | p(u)) \right],$$
(131)

for $(u^n, x^n) \in \mathcal{T}_{\epsilon}^n(p_{UX})$, and $\alpha \in (1, \infty)$ if

$$R < H(U) - R'_{\alpha}(p_X, p_{Z|X}, p_U)$$
 (132)

then as $n \to \infty$ and $\epsilon \to 0$ we have

$$\mathbb{E}_{\mathcal{B}}\left[T_{\alpha}\left(P(b, z^{n}) \parallel p^{U}(b)q(z^{n})\right)\right] \to 0,\tag{133}$$

for $u^n \in \mathcal{T}^n_{\epsilon}(p_U)$ random binning $\{\mathcal{B}(u^n): \mathcal{U}^n \to [2^{nR}]\}$. Which $p(z) = \sum_x p(x)p(z|x)$ and $q(z^n) = \prod_{i=1}^n p(z_i)$ with $|\mathcal{U}| \leq |\mathcal{X}| + 1$.

Proof. (126) implies that joint distribution between variables in this setup is as following

$$P(b, u^n, x^n, z^n) = \tilde{p}(u^n, x^n) \mathbb{1} \{ \mathcal{B}(u^n) = b \} \prod_{i=1}^n p(z_i | x_i)$$
 (134)

$$P(b, z^n) = \sum_{(u^n, x^n) \in \mathcal{T}^n(p_{UX})} \tilde{p}(u^n, x^n) \mathbb{1} \{ \mathcal{B}(u^n) = b \} \prod_{i=1}^n p(z_i | x_i).$$
 (135)

Therefore for $\alpha \in \mathbb{N}$, we can write

$$\mathbb{E}_{\mathcal{B}}[(\alpha - 1)T_{\alpha}(P(b, z^n) \parallel p^U(b)q(z^n))]$$
(136)

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\sum_{b,z^n} \frac{q(z^n)}{M} \left(\sum_{(u^n,x^n) \in \mathcal{T}_{\epsilon}^n(p_{UX})} \tilde{p}(u^n,x^n) \mathbb{1} \{ \mathcal{B}(u)^n = b \} \prod_{i=1}^n p(z_i|x_i) \right)^{\alpha} q^{-\alpha}(z^n) \right] - 1$$

$$(137)$$

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\sum_{z^n} q(z^n) \left(\sum_{(u^n, x^n) \in \mathcal{T}_{\epsilon}^n(p_{UX})} \tilde{p}(u^n, x^n) \mathbb{1} \{ \mathcal{B}(u^n) = 1 \} \prod_{i=1}^n p(z_i | x_i) \right)^{\alpha} q^{-\alpha}(z^n) \right] - 1$$

$$(138)$$

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\sum_{z^n} q(z^n) \left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \tilde{p}(u^n) \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^n(X|u^n)} \tilde{p}(x^n|u^n) \prod_{i=1}^n p(z_i|x_i) \right) \mathbb{1} \{ \mathcal{B}(u^n) = 1 \} \right)^{\alpha} q^{-\alpha}(z^n) \right] - 1.$$

$$(139)$$

Let define

$$S(z^n, u_i^n) = \sum_{x^n \in \mathcal{T}_{\epsilon}^n(X|u_i^n)} \tilde{p}(x^n|u_i^n) \prod_{i=1}^n p(z_i|x_i),$$
(140)

rewrite (139) with the expansion as following, we have

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\sum_{z^n} q(z^n) \left(\sum_{u_1^n \in \mathcal{T}_{\epsilon}^n(p_U)} \tilde{p}^{\alpha}(u_1^n) S^{\alpha}(z^n, u_1^n) q^{-\alpha}(z^n) \mathbb{1} \{ \mathcal{B}(u_1^n) = 1 \} \right]$$

$$(141)$$

$$+\sum_{(u_1^n \neq u_2^n) \in \mathcal{T}_{\epsilon}^n(p_U)} \tilde{p}^{\alpha-1}(u_1^n) \tilde{p}(u_2^n) S^{\alpha-1}(z^n, u_1^n) S(z^n, u_2^n) q^{-\alpha}(z^n) \prod_{i=1}^2 \mathbb{1} \{\mathcal{B}(u_i^n) = 1\} + \cdots$$
 (142)

$$+ \sum_{\substack{(u_1^n \neq u_2^n \neq u_3^n) \in \mathcal{T}_{\epsilon}^n(p_U)}} \tilde{p}^{\alpha-2}(u_1^n) \tilde{p}(u_2^n) \tilde{p}(u_3^n) S^{\alpha-2}(z^n, u_1^n) S(z^n, u_2^n) S(z^n, u_3^n) q^{-\alpha}(z^n) \prod_{i=1}^3 \mathbb{1} \{\mathcal{B}(u_i^n) = 1\} + \cdots$$

(143)

$$\cdots$$
 (144)

$$+ \sum_{\substack{(u_1^n \neq \dots \neq u_{\alpha}^n) \in \mathcal{T}_{\epsilon}^n(p_U)}} \left(\prod_{i=1}^{\alpha} \tilde{p}(u_i^n) S(z^n, u_i^n) \mathbb{1} \{ \mathcal{B}(u_i^n) = 1 \} \right) q^{-\alpha}(z^n) \right) - 1$$

$$(145)$$

$$= M^{\alpha - 1} \sum_{z^n} q(z^n) \left(\sum_{\substack{u_1^n \in \mathcal{T}^n(p_U) \\ q^{\alpha}(z^n)}} \frac{\tilde{p}^{\alpha}(u_1^n) S^{\alpha}(z^n, u_1^n)}{q^{\alpha}(z^n)} \right)$$

$$(146)$$

$$+ M^{\alpha-2} \sum_{z^{n}} q(z^{n}) \left(\sum_{(u_{1}^{n} \neq u_{2}^{n}) \in \mathcal{T}_{\epsilon}^{n}(p_{U})} \left(\frac{\tilde{p}^{\alpha-1}(u_{1}^{n}) S^{\alpha-1}(z^{n}, u_{1}^{n})}{q^{\alpha-1}(z^{n})} \right) \left(\frac{\tilde{p}(u_{2}^{n}) S(z^{n}, u_{2}^{n})}{q(z^{n})} \right) + \cdots \right)$$
(147)

$$+ M^{\alpha - 3} \sum_{z^{n}} q(z^{n}) \left(\sum_{(u_{1}^{n} \neq u_{2}^{n} \neq u_{3}^{n}) \in \mathcal{T}_{\epsilon}^{n}(p_{U})} \left(\frac{\tilde{p}^{\alpha - 2}(u_{1}^{n}) S^{\alpha - 2}(z^{n}, u_{1}^{n})}{q^{\alpha - 2}(z^{n})} \right) \left(\frac{\tilde{p}(u_{2}^{n}) S(z^{n}, u_{2}^{n})}{q(z^{n})} \right) \left(\frac{\tilde{p}(u_{3}^{n}) S(z^{n}, u_{3}^{n})}{q(z^{n})} \right) + \cdots \right)$$

$$(148)$$

$$+\sum_{z^n} q(z^n) \sum_{\substack{(u_1^n \neq \dots \neq u_{\alpha}^n) \in \mathcal{T}_{\epsilon}^n(p_U)}} \prod_{i=1}^{\alpha} \left(\frac{\tilde{p}(u_i^n) S(z^n, u_i^n)}{q(z^n)} \right) - 1$$

$$(150)$$

$$\leq M^{\alpha-1} \sum_{z^n} q(z^n) \left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}^{\alpha}(u^n) S^{\alpha}(z^n, u^n)}{q^{\alpha}(z^n)} \right) \tag{151}$$

$$+ M^{\alpha-2} \sum_{z^n} q(z^n) \left[\left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}^{\alpha-1}(u^n) S^{\alpha-1}(z^n, u^n)}{q^{\alpha-1}(z^n)} \right) \left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}(u^n) S(z^n, u^n)}{q(z^n)} \right) + \cdots \right]$$

$$(152)$$

$$\cdots$$
 (153)

$$+\sum_{z^n} q(z^n) \prod_{i=1}^{\alpha} \left[\left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}(u^n) S(z^n, u^n)}{q(z^n)} \right) \right] - 1$$

$$(154)$$

$$\leq M^{\alpha-1} \sum_{z^n} q(z^n) \left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}^{\alpha}(u^n) S^{\alpha}(z^n, u^n)}{q^{\alpha}(z^n)} \right) \tag{155}$$

$$+ M^{\alpha-2} \sum_{z^n} q(z^n) \left[\left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}^{\alpha-1}(u^n) S^{\alpha-1}(z^n, u^n)}{q^{\alpha-1}(z^n)} \right) (1 + 2\delta_n) + \cdots \right]$$

$$(156)$$

$$\cdots$$
 (157)

$$+\left(1+2\delta_{n}\right)^{\alpha}-1.\tag{158}$$

(151-154) follows from

$$\sum_{(u_1 \neq \dots \neq u_\ell) \in \mathcal{T}_{\epsilon}^n(p_U)} \prod_{i=1}^{\ell} \frac{\tilde{p}^{\alpha_i}(u_i^n) S^{\alpha_i}(z^n, u_i^n)}{q^{\alpha_i}(z^n)} \le \prod_{i=1}^{\ell} \left(\sum_{u \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}^{\alpha_i}(u^n) S^{\alpha_i}(z^n, u^n)}{q^{\alpha_i}(z^n)} \right). \tag{159}$$

(155-158) is due to

$$\sum_{u^n \in \mathcal{T}^n(p_U)} \frac{\tilde{p}(u^n) S(z^n, u^n)}{q(z^n)} \le \frac{\sum_{(u^n, x^n) \in \mathcal{T}^n_{\epsilon}(p_{UX})} \tilde{p}(u^n, x^n) \prod_{i=1}^n p(z_i | x_i)}{q(z^n)}$$
(160)

$$= \frac{\sum_{(u^n, x^n) \in \mathcal{T}_{\epsilon}^n(p_{UX})} \tilde{p}(u^n, x^n) \prod_{i=1}^n p(z_i | x_i)}{\sum_{x^n, u^n} \prod_{i=1}^n p(u_i, x_i) \prod_{i=1}^n p(z_i | x_i)}$$
(161)

$$\leq \frac{\sum_{(u^n, x^n) \in \mathcal{T}_{\epsilon}^n(p_{UX})} \tilde{p}(u^n, x^n) \prod_{i=1}^n p(z_i | x_i)}{\sum_{(x^n, u^n) \in \mathcal{T}_{\epsilon}^n(p_{UX})} \prod_{i=1}^n p(u_i, x_i) \prod_{i=1}^n p(z_i | x_i)}$$
(162)

$$\leq \frac{1}{1 - \delta_n} \leq 1 + 2\delta_n,\tag{163}$$

and (163) is due to (130) combining with

$$\left(\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \prod_{i=1}^n p(u_i)\right) \left(\sum_{x^n \in \mathcal{T}_{\epsilon}^n(X|u^n)} \prod_{i=1}^n p(x_i|u_i)\right) \ge 1 - \delta_n \tag{164}$$

for $u^n \in \mathcal{T}^n_{\epsilon}(p_U)$, when n is large enough and δ_n is small.

Let assume $Z^n = z^n$ be fixed, in addition consider measure

$$\nu(u^n) = \frac{\frac{\tilde{p}(u^n)S(u^n, z^n)}{q(z^n)}}{\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \frac{\tilde{p}(u^n)S(u^n, z^n)}{q(z^n)}} = \frac{\tilde{p}(u^n)S(u^n, z^n)}{\sum_{u^n \in \mathcal{T}_{\epsilon}^n} \tilde{p}(u^n)S(u^n, z^n)}$$
(165)

Harris' inequality for non-decreasing function $f_i(x) = x^{\alpha_i - 1}$ with $\sum_{i=1}^{\ell} \alpha_i = \alpha$ implies that

$$\left(\frac{q(z^n)}{\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \tilde{p}(u^n) S(z^n, u^n)}\right) \left[\sum_{u^n \in \mathcal{T}_{\epsilon}^n} \frac{\tilde{p}(u^n) S(u^n, z^n)}{q(z^n)} \prod_{i=1}^{\ell} \left(\frac{\tilde{p}(u^n) S(u^n, z^n)}{q(z^n)}\right)^{\alpha_i - 1}\right]$$
(166)

$$= \mathbb{E}_{\nu(u^n)} \left[\prod_{i=1}^{\ell} f_i \left(\frac{\tilde{p}(U^n) S(U^n, z^n)}{q(z^n)} \right) \right]$$
(167)

$$\geq \prod_{i=1}^{\ell} \mathbb{E}_{\nu(u^n)} \left[f_i \left(\frac{\tilde{p}(U^n) S(U^n, z^n)}{q(z^n)} \right) \right] \tag{168}$$

$$= \left(\frac{q(z^n)}{\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \tilde{p}(u^n) S(z^n, u^n)}\right)^{\ell} \prod_{i=1}^{\ell} \left[\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \left(\frac{\tilde{p}(u^n) S(z^n, u^n)}{q(z^n)}\right)^{\alpha_i} \right]$$
(169)

which we can conclude for every fixed $Z^n = z^n$

$$\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \left[\frac{\tilde{p}(u^n) S(u^n, z^n)}{q(z^n)} \prod_{i=1}^{\ell} \left(\frac{\tilde{p}(u^n) S(u^n, z^n)}{q(z^n)} \right)^{\alpha_i - 1} \right] \ge \left(\frac{1}{1 + 2\delta_n} \right)^{\ell - 1} \prod_{i=1}^{\ell} \left[\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \left(\frac{\tilde{p}(u^n) S(u^n, z^n)}{q(z^n)} \right)^{\alpha_i} \right]. \tag{170}$$

This implied that for obtaining the binning rate that is enough to put for $k=0,1,\cdots,\alpha-2$,

$$R < \min_{k} \frac{1}{n(\alpha - k - 1)} \log \left(\frac{1}{\sum_{z^{n}} q(z^{n}) \sum_{u^{n} \in \mathcal{T}_{\epsilon}^{n}(p_{U})} \left(\frac{\tilde{p}^{\alpha - k}(u^{n})S^{\alpha - k}(u^{n}, z^{n})}{q^{\alpha - k}(z^{n})} \right)} \right), \tag{171}$$

which decreasing property of R in α implies that k=0, therefore via simplification in lemma 4 completes the proof.

The proof for real α 's is in a way similar to the generalization proof of Theorem 1 and hence is omitted.

Remark 5. Converse part is similar to Theorem (3), and uses

$$2^{-n(H(U)+\delta'_n(\epsilon))} \le \tilde{p}(u^n) \le (1+2\delta_n(\epsilon))2^{-n(H(U)-\delta'_n(\epsilon))}$$
(172)

since

$$2^{-n(H(U)+\delta'_n(\epsilon))} \le \prod_{i=1}^n p(u_i) \le \tilde{p}(u^n) = \frac{\prod_{i=1}^n p(u_i)}{\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \prod_{i=1}^n p(u_i)}$$
(173)

$$\leq (1 + 2\delta_n(\epsilon)) \prod_{i=1}^n p(u_i) \tag{174}$$

$$\leq (1 + 2\delta_n(\epsilon))2^{-n(H(U) - \delta_n'(\epsilon))} \tag{175}$$

combining with

$$\sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \prod_{i=1}^n p(u_i) \ge 1 - \delta_n(\epsilon)$$
(176)

when $u^n \in \mathcal{T}^n_{\epsilon}(p_U)$ with $\epsilon \to 0$.

Remark 6. Achievability and converse part for stochastic encoder in the case of $\alpha \in (0,1)$ are similar to Theorem 2 and we have R < H(U|Z) condition on random binning rate instead of H(X|Z).

Remark 7. Obtaining the random binning rate in case of D_{∞} has steps coincided to Theorem 4 and we have

$$R < H(U) - R'_{\infty}(p_X, p_{Z|X}, p_U),$$
 (177)

which

$$R'_{\infty}(p_{X}, p_{Z|X}, p_{U}) = \max_{\substack{T_{Z|XU} \\ \sum_{z} p(x)p(z|x) = p(z) \\ \sum_{z} t(x|u, z)t(z|u)p(u) = p(u, x)}} \left[-D\left(t(z|u, x) \parallel p(z|x)|p(x, u)\right) + D\left(t(z|u) \parallel p(z)|p(u)\right) \right].$$
(178)

V. APPLICATIONS

In this section, we explore the application of Tsallis-based OSRB in analyzing the achievability rate regions for the wiretap channel problem.

The wiretap channel is modeled as a conditional distribution $p_{YZ|X}$, so that X is the channel's input, Y is the information received by the legitimate receiver, and Z is the leakage output received by the eavesdropper. Assuming the message set is $\mathcal{M} = [2^{nR}]$, the goal is to send the maximum amount of information securely to the legal receiver. Considering mutual information as a measure of security, it is well known that the rate R < I(X;Y) - I(X;Z) can be achieved for any input distribution p(x). Here, we assume Tsallis divergence as a security measure and provide an asymptotic analysis of the achievable rate.

A. Wiretap channel with deterministic encoder

Theorem 6. Let $p_{YZ|X}$ be an arbitrary wiretap channel, in addition assume $q(z^n) = \prod_{i=1}^n p(z_i)$ with $p(z) = \sum_x p(x)p(z|x)$, $\mathcal{M} = [2^{nR}]$ be the message set, and $\alpha \in (1, \infty)$. Then, for any arbitrary distribution \tilde{p}_X over $\mathcal{T}^n_{\epsilon}(p_X)$ with

$$\tilde{p}(x^n) = \frac{\prod_{i=1}^n p(x_i)}{\sum_{x^n \in \mathcal{T}_i^n(p_X)} \prod_{i=1}^n p(x_i)}$$
(179)

and $\epsilon \to 0$ there exists a code that can send a message M with rate R, reliably with vanishing error and strongly securely

$$\lim_{n \to \infty} T_{\alpha} \Big(p(m, z^n) \parallel p^{U}(m) q(z^n) \Big) = 0.$$
 (180)

Provided that the rate satisfies

$$R < I(X;Y) - \sum_{x} p(x) D_{\alpha} (p(z|x) \parallel p(z)).$$
 (181)

Proof. Let $\beta = (m, f): X^n \to [2^{nR_1}] \times [2^{nR_2}]$ be a random binning that uniformly and independently maps each sequence x^n to two bin indices, $m \in [2^{nR_1}]$ and $f \in [2^{nR_2}]$. More precisely, if the condition $R_1 + R_2 < H(X)$ holds [14, Theorem 1], then we have

$$\lim_{T \to \infty} \mathbb{E}_{\mathcal{B}} [\|P(m, f) - p^{U}(m) p^{U}(f)\|_{\text{TV}}] = 0$$
 (182)

In addition, if the condition $R_2 > H(X|Y)$ is met, the decoder, having (F, Y^n) , can reliably decode the sequence X^n . More specifically, we have

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{B}} \left[\mathbb{P} \left(\hat{X}^n \neq X^n \right) \right] = 0, \tag{183}$$

by taking the expectation over all random binnings \mathcal{B} . Finally, using Theorem 1, the constraint $R_1 + R_2 < H(X) - \sum_x p(x) D_{\alpha} \big(p(z|x) \parallel p(z) \big)$ results that

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{B}} \Big[T_{\alpha} \Big(P(m, f, z^n) \parallel p^{U}(m) p^{U}(f) q(z^n) \Big) \Big] = 0.$$
 (184)

On the other hand, using Lemma 5 in Appendix F, we obtain that for at least half of the (\mathcal{B}, F) pairs

$$\lim_{n \to \infty} T_{\alpha} \Big(p(m, z^n | f) \parallel p^{U}(m) q(z^n) \Big) = 0.$$

$$(185)$$

Therefore, by choosing the positive real numbers R_1, R_2 so that

$$R_1 + R_2 < H(X) (186)$$

$$R_2 > H(X|Y) \tag{187}$$

$$R_1 + R_2 < H(X) - \sum_{x} p(x) D_{\alpha} (p(z|x) \parallel p(z)),$$
 (188)

the reliability and security constraints are established, and M and F become independent. Based on property 2 of Lemma 1, it can be concluded that $H(X) - \sum_x p(x) D_{\alpha} \big(p(z|x) \parallel p(z) \big) \le H(X|Z) \le H(X)$, which shows that satisfying condition (188) leads to satisfying condition (186). Therefore, condition (186) can be ignored.

Therefore, using (183), (185) and Markov's inequality together with a union bound, for any $\epsilon' > 0$ and sufficiently large n, there exists $f \in [2^{nR_2}]$ and a random labeling β_0 such that

$$\mathbb{P}\Big[\hat{X}^n \neq X^n \big| \beta_0, F = f\Big] \le \epsilon', \tag{189}$$

and

$$T_{\alpha}\Big(p(m, z^n | F = f) \parallel p^{U}(m) q(z^n)\Big) \to 0.$$

$$\tag{190}$$

Now we construct the code as follows. We consider M as the message that is uniformly distributed according to (182), and then we transmit the codeword $X^n = \beta_0^{-1}(M, f)$. Based on (189), the legitimate receiver decodes the message with an asymptotically vanishing error and the eavesdropper does not obtain any information from the message according to (190).

Take note that our coding strategy establishes the distribution as

$$p(m|f) p(x^n|m, f) p(z^n|x^n).$$
 (191)

However, it is imperative that condition (190) is satisfied with the ensuing distribution.

$$p^{U}(m) p(x^{n}|m, f) p(z^{n}|x^{n}).$$
 (192)

This assurance is provided by Lemma 6, which guarantees that a vanishing decrease in message rate results in an asymptotically diminishing Tsallis divergence for the distribution in (192). This concludes the proof.

Corollary 1. For $\alpha = 1$, the rate R = I(X;Y) - I(X;Z) can be achieved with an asymptotically vanishing error and with a strong security measure of $I(M;Z^n) \leq \epsilon$.

Proof. The proof according to property 3 of Lemma 1 is straightforward.

Remark 8. Inequality (12) suggests that all the achievability proofs for T_{α} as a security measure also holds for D_{α} when $\alpha \in (1, \infty)$.

Remark 9. Suppose we are in the $\alpha \in (0,1)$ case. Using Theorem 2 and similar to the proof of Theorem 6, the achievable secrecy rate with secrecy criterion T_{α} is equal to R < H(X|Z) - H(X|Y). In [2], taking into account the stronger criterion D_{α} , the same achievable region has been obtained.

As already mentioned, due to the infinity of T_{∞} , we consider D_{∞} instead. Therefore, using Theorem 4, we have the following secure rate region.

Theorem 7. For an arbitrary wiretap channel assume $q(z^n) = \prod_{i=1}^n p(z_i)$ with $p(z) = \sum_x p(x)p(z|x)$, there exists a code that can send a message M with rate R, reliably with vanishing error and strongly securely with

$$\lim_{n \to \infty} D_{\infty} \Big(p(m, z^n) \parallel p^{U}(m) \, q(z^n) \Big) = 0.$$
 (193)

Provided that the rate satisfies

$$R < I(X;Y) - \sum_{x} p(x) D_{\infty} (p(z|x) \parallel p(z)).$$
 (194)

Proof. The proof steps are similar to Theorem 6. We choose positive reals R_1 , R_2 such that

$$R = R_1 \tag{195}$$

$$R_2 > H(X|Y) \tag{196}$$

$$R_1 + R_2 < H(X) - \sum_{x} p(x) D_{\infty} (p(z|x) \parallel p(z)).$$
 (197)

 R_2 guarantees negligible-error decodability and based on Theorem 4, condition (197) satisfies strong security. Eliminating R_2 from (196) and (197) and setting the message rate equal to R_1 proves the theorem.

Corollary 2. If the binned sequences are generated independently and identically distributed (i.i.d.), then by repeating all the steps in Theorems (6-7) and combining them with Theorem 1, the achievable secure rates are

$$R < \tilde{H}_{\alpha}(X|Z) - H(X|Y) \tag{198}$$

for T_{α} and

$$R < \tilde{H}_{\infty}(X|Z) - H(X|Y), \tag{199}$$

for D_{∞} respectively, where $\alpha \in (1, \infty)$. In the case of random binning over ϵ -typical sets, the achievable rate is higher because

$$\tilde{H}_{\alpha}(X|Z) \le H(X) - \sum_{x} p(x) D_{\alpha}(p(z|x) \parallel p(z)) \tag{200}$$

and

$$\tilde{H}_{\infty}(X|Z) \le H(X) - \sum_{x} p(x) D_{\infty}(p(z|x) \parallel p(z)). \tag{201}$$

Therefore, the results for i.i.d. sequences are omitted. The proof of equation (200) for T_{α} with $\alpha \in (1, \infty)$ are presented below.

For T_{α} with $\alpha \in (1, \infty)$ presented such as below.

Proof.

$$H(X) - \sum_{x} p(x) D_{\alpha}(p(z|x) \parallel p(z))$$
 (202)

$$\sum_{x} p(x) \log \frac{1}{p(x)} - \frac{1}{\alpha - 1} \sum_{x} p(x) \log \left(\sum_{z} p(z) \left(\frac{p(z|x)}{p(z)} \right)^{\alpha} \right)$$
 (203)

$$= \frac{1}{\alpha - 1} \sum_{x} p(x) \log \left(\frac{1}{p(x)} \right)^{\alpha - 1} - \frac{1}{\alpha - 1} \sum_{x} p(x) \log \left(\sum_{z} p(z) \left(\frac{p(z|x)}{p(z)} \right)^{\alpha} \right) \tag{204}$$

$$= \frac{1}{\alpha - 1} \sum_{x} p(x) \log \left(\frac{1}{p^{\alpha - 1}(x) \sum_{z} p(z) \left(\frac{p(z|x)}{p(z)} \right)^{\alpha}} \right)$$
 (205)

$$= \frac{1}{1-\alpha} \left(\sum_{x} p(x) \log \left(\sum_{z} p^{\alpha-1}(x) p^{1-\alpha}(z) p^{\alpha}(z|x) \right) \right)$$
 (206)

$$\geq \frac{1}{1-\alpha} \left(\log \left(\sum_{x,z} p^{\alpha}(x) p^{1-\alpha}(z) p^{\alpha}(z|x) \right) \right) \tag{207}$$

$$= \frac{1}{1-\alpha} \left(\log \sum_{x,z} p(z) \left(\frac{p(x,z)}{p(z)} \right)^{\alpha} \right) \tag{208}$$

$$= \frac{1}{1-\alpha} \log \left(\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z) \right)$$
 (209)

$$=\tilde{H}_{\alpha}(X|Z). \tag{210}$$

Equation (207) follows from Jensen's inequality for $f(x) = \log x$.

Proof of equation (201) for case D_{∞} is such as following

$$H(X) - \sum_{z} p(z) D_{\infty}(p(z|x) \parallel p(z))$$
 (211)

$$= \sum_{x} p(x) \log \frac{1}{p(x)} - \sum_{x} p(x) \log \left(\max_{z} \frac{p(z|x)}{p(z)} \right)$$
 (212)

$$= \sum_{x} p(x) \log \left(\frac{1}{p(x) \left(\max_{z} \frac{p(z|x)}{p(z)} \right)} \right)$$
 (213)

$$= \sum_{x} p(x) \log \left(\frac{1}{\max_{z} \frac{p(x,z)}{p(z)}} \right) \tag{214}$$

$$\geq \log\left(\frac{1}{\max_{x,z} \frac{p(x,z)}{p(z)}}\right) \tag{215}$$

$$=\tilde{H}_{\infty}(X|Z). \tag{216}$$

B. Wiretap channel with stochastic encoder

In this section we derive an optimal rate for wiretap channel when the encoder is stochastic instead of deterministic encoder. In the achievability part we uses OSRB technique for variable (U, X) derived in Theorem 5.

Theorem 8. For a wiretap channel with conditional distribution $p_{YZ|X}$ and joint distribution p(x,u) such that $U \sim p(u)$ and $|\mathcal{U}| \leq |\mathcal{X}| + 1$, there exist a code with rate

$$R < I(U,Y) - R'_{\alpha}(p_X, p_{Z|X}, p_U)$$
 (217)

which message could be sent with vanishing error and strong secrecy under Tsallis divergence for $\alpha \in (1, \infty)$. i.e.

$$\lim_{n \to \infty} T_{\alpha} \left(p(m, z^n) \parallel p^U(m) q(z^n) \right) = 0 \tag{218}$$

with
$$q(z^n) = \prod_{i=1}^n p(z_i)$$
 and $p(z) = \sum_x p(x)p(z|x)$ which $R'_{\alpha}(p_X, p_{Z|X}, p_U)$ defined in (131).

Proof. Similar wiretap channel with deterministic encoder the proof structure included two parts Secrecy and reliability condition.

Let consider an arbitrary random variable U^n over $\mathcal{T}^n_\epsilon(p_U)$ with distribution

$$\tilde{p}(u^n) = \frac{\prod_{i=1}^n p(u_i)}{\sum_{u^n \in \mathcal{T}^n(p_U)} \prod_{i=1}^n p(u_i)}.$$
(219)

 $\beta = (m, f) : U^n \to [2^{nR_1}] \times [2^{nR_2}]$ is a random binning that maps uniformly and independently each sequences u^n to bin indices.

• Secrecy condition: Based on Theorem 5 secrecy condition could be satisfies if

$$R = R_1 \tag{220}$$

$$R_1 + R_2 < H(U) - R'_{\alpha}(p_X, p_{Z|X}, p_U)$$
(221)

more precisely under condition (221) we have

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{B}} \left[T_{\alpha} \left(p(m, f, z^n) \parallel p^U(m) p^U(f) p(z^n) \right) \right] = 0. \tag{222}$$

• Reliability condition: For satisfying the reliability condition let consider $R_2 > H(U|Y)$ therefore having (F, Y^n) the sequence U^n could be decoded with vanish error, so we have

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{B}} \left[\mathbb{P} \left(\hat{U}^n \neq U^n \right) \right] = 0 \tag{223}$$

which $\mathbb{E}_{\mathcal{B}}$ is expectation over all of the random mappings.

Lemma F implies that there exsist a fixed realization of $(\mathcal{B}, \mathcal{F})$ such that

$$\lim_{n \to \infty} T_{\alpha} \left(p(m, z^n | f) \parallel p^U(m) q(z^n) \right) = 0$$
(224)

$$\lim_{n \to \infty} \mathbb{P}\Big[\hat{U}^n \neq U^n \big| \beta_0, F = f\Big] = 0. \tag{225}$$

Similar to deterministic encoder our coding strategy establishes the distribution as

$$p(m|f)\,\tilde{p}(u^n|m,\,f)\,\tilde{p}(x^n|u^n)\,p(z^n|x^n).$$
 (226)

However, it is imperative that condition (224) is satisfied with the ensuing distribution

$$p^{U}(m)\,\tilde{p}(u^{n}|m,f)\,\tilde{p}(x^{n}|u^{n})\,p(z^{n}|x^{n}).$$
 (227)

This assurance is provided by Lemma 6 and Remark 16, which guarantees that a vanishing decrease in message rate results in an asymptotically diminishing Tsallis divergence for the distribution in (227). Let consider $\epsilon \to 0$ this concludes the proof.

Remark 10. The proof of secrecy rate in case of D_{∞} for stochastic encoder follows similar steps with rates

$$R = R_1 \tag{228}$$

$$R_1 + R_2 < H(U) - R'_{\infty}(p_X, p_{Z|X}, p_U)$$
(229)

$$R_2 > H(U|Y) \tag{230}$$

which $R'_{\infty}(p_X, p_{Z|X}, p_U)$ defined in (178).

Remark 11. The achievable secure rate with OSRB method for $\alpha \in (0,1)$ under Tsallis divergence is I(U;Y) - I(U;Z), similar to mutual information criterion so is omitted.

Remark 12. In [2], the secure rate is computed for $\alpha \in (0,2] \cup \{\infty\}$ for both of deterministic and stochastic encoder. The secure rate computed in Theorems (6-8) for $\alpha \in (0,\infty) \cup \{\infty\}$ coincides with the results in [2, Theorem 5-6] for $\alpha \in [0,2]$ and extends it for $\alpha \in (2,\infty)$.

Remark 13. T_{α} and D_{α} have similar behavior for $\alpha \in (0, \infty)$ when $n \to \infty$, demonstrating that the secure capacity with the D_{α} is equal to that with T_{α} for $\alpha \in (0, \infty)$. The achievable rate provided with T_{α} in (6) and (8), which is equal to one as presented in [2] with D_{α} (also proven to be capacity), implies that the computed Tsallis-based secure rate provided in this paper is also the capacity.

VI. CONCLUSION

In this paper, we conducted an asymptotic analysis of OSRB utilizing the Tsallis divergence criterion. Specifically, we examined the conditions on the binning rate, denoted as R, that would satisfy the following expectation over all random binnings.

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{B}} \Big[T_{\alpha} \Big(p(m, z^n) \parallel p^U(m) q(z^n) \Big) \Big] = 0, \tag{231}$$

where X and Z are random variables with a joint distribution p(x,z), and $\beta: \mathcal{X}^n \to [M]$ is a random binning function. Our analysis covers the entire range of $\alpha \in (0,\infty) \cup \{\infty\}$ and provides both achievability and converse proofs. Additionally, we investigated the binning rate by employing D_{∞} as an alternative to T_{∞} . By leveraging the established theorems for asymptotic OSRB analysis, we were able to examine the achievable rate region for wiretap channel.

Generated by IEEEtran.bst, version: 1.12 (2007/01/11)

REFERENCES

- [1] M. M. Mojahedian, S. Beigi, A. Gohari, M. H. Yassaee, and M. R. Aref, "A correlation measure based on vector-valued *l_p*-norms," *IEEE Transactions on Information Theory*, vol. 65, no. 12, pp. 7985–8004, 2019.
- [2] L. Yu and V. Y. Tan, "Rényi resolvability and its applications to the wiretap channel," *IEEE Transactions on Information Theory*, vol. 65, no. 3, pp. 1862–1897, 2018.
- [3] I. Issa and A. B. Wagner, "Measuring secrecy by the probability of a successful guess," *IEEE Transactions on Information Theory*, vol. 63, no. 6, pp. 3783–3803, June 2017.
- [4] I. Issa, S. Kamath, and A. B. Wagner, "An operational measure of information leakage," in 2016 Annual Conference on Information Science and Systems (CISS), March 2016, pp. 234–239.
- [5] C. T. Li and A. E. Gamal, "Maximal correlation secrecy," in 2015 IEEE International Symposium on Information Theory (ISIT), June 2015, pp. 2939–2943.
- [6] C. Schieler and P. Cuff, "Rate-distortion theory for secrecy systems," *IEEE Transactions on Information Theory*, vol. 60, no. 12, pp. 7584–7605, Dec 2014.
- [7] N. Weinberger and N. Merhav, "A large deviations approach to secure lossy compression," *IEEE Transactions on Information Theory*, vol. 63, no. 4, pp. 2533–2559, April 2017.
- [8] M. Bellare, S. Tessaro, and A. Vardy, "Semantic security for the wiretap channel," in Advances in Cryptology-CRYPTO 2012. Springer, 2012, pp. 294–311.
- [9] Y. Dodis and A. Smith, "Entropic security and the encryption of high entropy messages," in *Theory of Cryptography Conference*. Springer, 2005, pp. 556–577.
- [10] A. Xu and M. Raginsky, "Information-theoretic analysis of generalization capability of learning algorithms," *Advances in Neural Information Processing Systems*, vol. 30, 2017.
- [11] T. Steinke and L. Zakynthinou, "Reasoning about generalization via conditional mutual information," in *Conference on Learning Theory*. PMLR, 2020, pp. 3437–3452.
- [12] A. R. Esposito, M. Gastpar, and I. Issa, "Generalization error bounds via rényi-, f-divergences and maximal leakage," *IEEE Transactions on Information Theory*, vol. 67, no. 8, pp. 4986–5004, 2021.
- [13] B. Rodríguez Gálvez, G. Bassi, R. Thobaben, and M. Skoglund, "Tighter expected generalization error bounds via wasserstein distance," Advances in Neural Information Processing Systems, vol. 34, pp. 19109–19121, 2021.
- [14] M. H. Yassaee, M. R. Aref, and A. Gohari, "Achievability proof via output statistics of random binning," *IEEE Transactions on Information Theory*, vol. 60, no. 11, pp. 6760–6786, 2014.
- [15] M. M. Mojahedian, A. Gohari, and M. R. Aref, "On the equivalency of reliability and security metrics for wireline networks," *arXiv:1609.04586*, 2016.
- [16] M. Hayashi, "Exponential decreasing rate of leaked information in universal random privacy amplification," *IEEE Transactions on Information Theory*, vol. 57, no. 6, pp. 3989–4001, 2011.
- [17] B. Škorić, C. Obi, E. Verbitskiy, and B. Schoenmakers, "Sharp lower bounds on the extractable randomness from non-uniform sources," *Information and Computation*, vol. 209, no. 8, pp. 1184–1196, 2011.
- [18] S. Fehr and S. Berens, "On the conditional rényi entropy," *IEEE Transactions on Information Theory*, vol. 60, no. 11, pp. 6801–6810, 2014.
- [19] A. Rényi et al., "On measures of entropy and information," in *Proceedings of the fourth Berkeley symposium on mathematical statistics and probability*, vol. 1, no. 547-561. Berkeley, California, USA, 1961.
- [20] Y. Sakai and V. Y. Tan, "Asymptotic expansions of smooth Rényi entropies and their applications," *arXiv preprint* arXiv:2003.05545, 2020.

[21] M. Iwamoto and J. Shikata, "Revisiting conditional rényi entropies and generalizing shannon's bounds in information theoretically secure encryption," *Cryptology ePrint Archive*, 2013.

APPENDIX

A. Proof of Lemma 1

1) Let X and Z be two independent random variables or equivalently p(x,z)=p(x)p(z), hence

$$\tilde{H}_{\alpha}(X|Z) = \frac{1}{1-\alpha} \log \left(\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z) \right)$$
 (232)

$$= \frac{1}{1-\alpha} \log \left(\sum_{z} p(z) \sum_{x} p^{\alpha}(x) \right)$$
 (233)

$$= \frac{1}{1-\alpha} \log \left(\sum_{x} p^{\alpha}(x) \right) \tag{234}$$

$$=H_{\alpha}(X). \tag{235}$$

2) To prove it, it suffices to show that $\frac{d\tilde{H}_{\alpha}(X|Z)}{d\alpha} < 0$ holds for $\alpha \in (1, \infty)$.

$$\frac{d\tilde{H}_{\alpha}(X|Z)}{d\alpha} = \frac{1}{(\alpha - 1)^2} \log \left(\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z) \right) + \frac{1}{1 - \alpha} \frac{\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z) \log p(x|z)}{\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z)} \right)$$
(236)

$$= \frac{1}{(\alpha - 1)^2} \log \left(\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z) \right)$$

$$-\frac{1}{(\alpha-1)^2} \frac{\sum_z p(z) \sum_x p^{\alpha}(x|z) \log p_{x|z}^{\alpha-1}}{\sum_z p(z) \sum_x p^{\alpha}(x|z)}$$
(237)

$$= \frac{\mathbb{E}\left[p_{x|z}^{\alpha-1}\right] \log \mathbb{E}\left[p_{x|z}^{\alpha-1}\right] - \mathbb{E}\left[p_{x|z}^{\alpha-1} \log p_{x|z}^{\alpha-1}\right]}{(\alpha-1)^2 \mathbb{E}\left[p_{x|z}^{\alpha-1}\right]}$$
(238)

$$\leq 0. \tag{239}$$

(238) is obtained by taking the common denominator and the following

$$\mathbb{E}\Big[p_{x|z}^{\alpha-1}\Big] = \sum_{x,z} p(x,z) \, p(x|z)^{\alpha-1} \tag{240}$$

$$= \sum_{z} p(z) \sum_{x} p^{\alpha}(x|z), \tag{241}$$

where the expectation is over p(x, z). Moreover, (239) follows from Jensen's inequality for the convex function $g(c) = c \log c$. Further, "=" occurs when the channel is singleton.

3) Using L'Hôpital's rule, we get

$$\lim_{\alpha \to 1} \tilde{H}_{\alpha}(X|Z) = -\frac{d}{d\alpha} \left(\log \sum_{z} p(z) \sum_{x} p_{x|z}^{\alpha} \right) \Big|_{\alpha=1}$$
(242)

$$= -\frac{\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z) \log p(x|z)}{\sum_{z} p(z) \sum_{x} p^{\alpha}(x|z)} \bigg|_{\alpha=1}$$

$$(243)$$

$$= -\mathbb{E}_{X,Z} \Big[\log p(x|z) \Big]$$
 (244)

$$=H(X|Z). (245)$$

4) Data processing inequality

$$\tilde{H}_{\alpha}(X|Y) = \frac{1}{1-\alpha} \log \left(\sum_{y} p(y) \sum_{x} p^{\alpha}(x|y) \right)$$
(246)

$$= \frac{1}{1-\alpha} \log \left(\mathbb{E}_Y \left[\sum_x p^{\alpha}(x|Y) \right] \right) \tag{247}$$

$$= \frac{1}{1 - \alpha} \log \left(\mathbb{E}_{Y,Z} \left[\sum_{x} p^{\alpha}(x|Y) \right] \right)$$
 (248)

$$= \frac{1}{1 - \alpha} \log \left(\mathbb{E}_Z \mathbb{E}_{Y|Z} \left[\sum_x p^{\alpha}(x|Y) \right] \right)$$
 (249)

$$\leq \frac{1}{1-\alpha} \log \left(\mathbb{E}_Z \left[\sum_{x} p^{\alpha}(x|Z) \right] \right) \tag{250}$$

$$=\tilde{H}_{\alpha}(X|Z). \tag{251}$$

(250) follows from Jensen's inequality for $g(c) = c^{\alpha}, \alpha > 1$ and $p_{x|Z} = \mathbb{E}_{Y|Z}[p(x|Y)]$.

- 5) The proof is straightforward using Jensen's inequality for the concave function $g(c) = \log c$.
- 6) By tending α to infinity and keeping the maximum term, the result is

$$\lim_{\alpha \to \infty} \tilde{H}_{\alpha}(X|Z) = \lim_{\alpha \to \infty} \frac{1}{1 - \alpha} \log \left(\sum_{z,x} p(z) p^{\alpha}(x|z) \right)$$
 (252)

$$= \lim_{\alpha \to \infty} \frac{1}{1 - \alpha} \log \left(p(z) \cdot \left(\max_{x, z} p(x|z) \right)^{\alpha} \right)$$
 (253)

$$= \lim_{\alpha \to \infty} \frac{1}{1 - \alpha} \log p(z)$$

$$+\lim_{\alpha \to \infty} \frac{\alpha}{1-\alpha} \log \left(\max_{x,z} p(x|z) \right) \tag{254}$$

$$= \log \left(\frac{1}{\max_{x,z} p(x|z)} \right). \tag{255}$$

7) Let $p(x^n, z^n) = \prod_{i=1}^n p(x_i, z_i)$ then

$$\tilde{H}_{\alpha}(X^n|Z^n) = \frac{1}{1-\alpha} \log \left(\sum_{z^n} p(z^n) \sum_{x^n} p^{\alpha}(x^n|z^n) \right)$$
(256)

$$= \frac{1}{1-\alpha} \log \left(\sum_{z^n} \prod_{i=1}^n p(z_i) \sum_{x^n} \prod_{i=1}^n p^{\alpha}(x_i|z_i) \right)$$
 (257)

$$= \frac{1}{1-\alpha} \log \prod_{i=1}^{n} \left(\sum_{z_i} p(z_i) \sum_{x_i} p^{\alpha}(x_i|z_i) \right)$$
 (258)

$$=\sum_{i=1}^{n} \tilde{H}_{\alpha}(X_i|Z_i) \tag{259}$$

$$= n\tilde{H}_{\alpha}(X|Z). \tag{260}$$

8) According to the property of decreasing $\tilde{H}_{\alpha}(X|Z)$ with respect to α (property 2) and non-negativity of $\tilde{H}_{\infty}(X|Z)$, the proof is complete.

$$\tilde{H}_{\alpha}(X|Z) \ge \tilde{H}_{\infty}(X|Z) = \log\left(\frac{1}{\max_{x,z} p(x|z)}\right) \ge 0.$$
 (261)

9) According to the definition of the singleton channel, we have

$$\max_{x,z} p(x,z) = \min_{x,z} p(x,z),$$
(262)

which subsequently, based on property 6, leads to

$$\tilde{H}_{\infty}(X|Z) = \tilde{H}_{-\infty}(X|Z). \tag{263}$$

The decreasing property of $\tilde{H}_{\alpha}(X|Z)$ completes the proof.

B. Proof of Theorem 1 for real α 's

Achievability part of the proof

Starting from (34) for real $\alpha \in (1, \infty)$, we have

Let consider $\alpha \in (1, \infty)$,

$$M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\left(\sum_{x} p(x|z) \mathbb{1} \{ \mathcal{B}(x) = 1 \} \right)^{\alpha} \right]$$

$$= M^{\alpha} \mathbb{E}_{\mathcal{B}} \left[\left(\sum_{x} p(x|z) \mathbb{1} \{ \mathcal{B}(x) = 1 \} \right)^{\lfloor \alpha \rfloor} \left(\sum_{\bar{x}} p(\bar{x}|z) \mathbb{1} \{ \mathcal{B}(\bar{x}) = 1 \} \right)^{\{\alpha\}} \right]$$
(264)

$$\leq M^{\alpha} \sum_{\alpha_1 + \dots + \alpha_{\ell} = |\alpha|} \left(\sum_{x_1 \neq \dots \neq x_{\ell}} \mathbb{E}_{\mathcal{B}(x_1), \dots, \mathcal{B}(x_{\ell})} \left[\prod_{i=1}^{\ell} p^{\alpha_i}(x_i|z) \mathbb{1} \{ \mathcal{B}(x_i) = 1 \} \right] \right)$$
(265)

$$\cdot \left(\mathbb{E}_{\mathcal{B}(\bar{x})|\mathcal{B}(x_1),\dots,\mathcal{B}(x_\ell)} \left[\sum_{\bar{x}} p(\bar{x}|z) \mathbb{1} \{ \mathcal{B}(\bar{x}) = 1 \} \right] \right)^{\{\alpha\}}$$
 (266)

$$= M^{\alpha - \ell} \sum_{\alpha_1 + \dots + \alpha_\ell = \lfloor \alpha \rfloor} \sum_{x_1 \neq \dots \neq x_\ell} \left(\prod_{i=1}^\ell p^{\alpha_i}(x_i|z) \right) \cdot \left(\frac{1}{M} + \sum_{i=1}^\ell p(x_i|z) \right)^{\{\alpha\}}$$
(267)

$$\leq M^{\alpha-\ell} \sum_{\alpha_1 + \dots + \alpha_\ell = |\alpha|} \sum_{x_1 \neq \dots \neq x_\ell} \left(\prod_{i=1}^\ell p^{\alpha_i}(x_i|z) \right) \cdot \left(M^{\lfloor \alpha \rfloor - \alpha} + \sum_{i=1}^\ell p^{\{\alpha\}}(x_i|z) \right) \tag{268}$$

$$= \sum_{\alpha_1 + \dots + \alpha_{\ell} = \lfloor \alpha \rfloor} \left(M^{\lfloor \alpha \rfloor - \ell} \sum_{x_1 \neq \dots \neq x_{\ell}} \prod_{i=1}^{\ell} p^{\alpha_i}(x_i | z) + \sum_{j=1}^{\ell} M^{\alpha - \ell} \sum_{x_1 \neq \dots \neq x_{\ell}} \left(p^{\alpha_j + \{\alpha\}}(x_j | z) \prod_{\substack{i=1 \ j \neq j}}^{\ell} p^{\alpha_i}(x_i | z) \right) \right)$$

$$(269)$$

$$\leq \sum_{\alpha_{1}+\dots+\alpha_{\ell}=\lfloor\alpha\rfloor} \left(M^{\lfloor\alpha\rfloor-\ell} \left(\prod_{i=1}^{\ell} \sum_{x_{i}} p^{\alpha_{i}}(x_{i}|z) \right) + \sum_{j=1}^{\ell} M^{\alpha-\ell} \left(\prod_{\substack{i=1\\i\neq j}}^{\ell} \sum_{x_{i}} p^{\alpha_{i}}(x_{i}|z) \right) \cdot \left(\sum_{x_{j}} p^{\alpha_{j}+\{\alpha\}}(x_{j}|z) \right) \right) \tag{270}$$

$$= \sum_{\alpha_1 + \dots + \alpha_\ell = \lfloor \alpha \rfloor} M^{\lfloor \alpha \rfloor - \ell} \left(\prod_{i=1}^{\ell} \left(\sum_{x_i} p^{\alpha_i}(x_i|z) \right) + \sum_{j=1}^{\ell} M^{\alpha - \ell} \left(\sum_{x_j} p^{\alpha_j + \{\alpha\}}(x_j|z) \right) \prod_{\substack{i=1\\i \neq j}}^{\ell} \left(\sum_{x_i} p^{\alpha_i}(x_i|z) \right) \right), \tag{271}$$

where (265) is the result of Jensen's inequality for the concave function $f(x) = x^{\{\alpha\}}$. Inequality (270) comes from

$$\sum_{x_1 \neq x_2 \neq \dots \neq x_\ell} \prod_{i=1}^{\ell} p^{\alpha_i}(x_i|z) \le \prod_{i=1}^{\ell} \sum_{x_i} p^{\alpha_i}(x_i|z).$$
 (272)

Moreover, (268) arises out of

$$\left(\sum_{i} x_{i}\right)^{\{\alpha\}} \leq \sum_{i} x_{i}^{\{\alpha\}}.$$
(273)

Substituting (271) into (32), considering (X^n, Z^n) and $M = [2^{nR}]$, the rate R must satisfy the following inequality.

$$R < \min_{\substack{i \in [\ell] \\ 1 \le \alpha_i \le \lfloor \alpha \rfloor}} \{ \tilde{H}_{\alpha_i}(X|Z), \tilde{H}_{\alpha_i + \{\alpha\}}(X|Z) \} = \tilde{H}_{\alpha}(X|Z), \tag{274}$$

where equality results from the decreasing property of $\tilde{H}_{\alpha}(X|Z)$ with α . Let fix Z=z and nondecreasing function of $f_i(x)=x^{\alpha_i-1}$, for correctness of

$$\sum_{x} p^{\lfloor \alpha \rfloor - \ell + 1}(x|z) = \mathbb{E}_{p_{X|Z=z}} \left[p^{\lfloor \alpha \rfloor - \ell}(X|z) \right]$$
 (275)

$$= \mathbb{E}_{p_{X|Z=z}} \left[\prod_{i=1}^{\ell} f_i(p(X|z)) \right] \ge \prod_{i=1}^{\ell} \mathbb{E}_{p_{X|Z=z}} \left[f_i(p(X|z)) \right]$$
 (276)

$$= \prod_{i=1}^{\ell} \left[\sum_{x} p^{\alpha_i}(x|z) \right] \tag{277}$$

we used HARRIS inequality for $f_i(x)$ with $\sum_{i=1}^{\ell} a_i = \lfloor \alpha \rfloor$.

For

$$\sum_{x} p^{\alpha - \ell + 1}(x|z) \ge \left(\sum_{x} p^{a_j + \{\alpha\} + 1}(x|z)\right) \prod_{\substack{i=1\\i \ne j}}^{\ell} \left(\sum_{x} p^{a_i + 1}(x|z)\right) \tag{278}$$

let consider $f_i(x) = x^{\alpha_i - 1}, g_j(x) = x^{\alpha_j + \{\alpha\} - 1}$

$$\mathbb{E}_{p_{X|Z=z}}\left[p^{\alpha-\ell}(X|z)\right] = \mathbb{E}_{p_{X|Z=z}}\left[g_j(p(X|z))\prod_{\substack{i=1\\i\neq j}}^{\ell}f_i(p(X|z))\right]$$
(279)

$$\geq \mathbb{E}_{p_{X|Z=z}} \left[g_j(p(X|z)) \right] \prod_{\substack{i=1\\i\neq j}}^{\ell} \mathbb{E}_{p_{X|Z=z}} \left[f_i(p(X|z)) \right]$$
(280)

$$= \left(\sum_{x} p^{a_j + \{\alpha\}}(x|z)\right) \prod_{\substack{i=1\\i \neq j}}^{\ell} \left[\sum_{x} p^{\alpha_i}(x|z)\right]$$
(281)

with $\sum_{i=1}^{\ell} \alpha_i = \lfloor \alpha \rfloor$. With this, we conclude the achievability proof.

Converse part of the proof

Our objective is to demonstrate that for $\alpha \in (1, \infty)$, if $R > \tilde{H}_{\alpha}(X|Z)$, then it follows that

$$\mathbb{E}_{\mathcal{B}}\left[T_{\alpha}\left(P(b, z^{n}) \parallel p^{U}(b)p(z^{n})\right)\right] \to \infty. \tag{282}$$

By putting (34) in (32), we have

$$\mathbb{E}_{\mathcal{B}}\left[T_{\alpha}\left(P(b,z) \parallel p^{U}(b) p(z)\right)\right]$$

$$= \frac{1}{\alpha - 1} \left(M^{\alpha} \mathbb{E}_{Z} \mathbb{E}_{\mathcal{B}}\left[\left(\sum_{x} p(x|Z) \mathbb{1}\{\mathcal{B}(x) = 1\}\right)^{\alpha}\right] - 1\right)$$
(283)

$$\geq \frac{M^{\alpha}}{\alpha - 1} \mathbb{E}_{Z} \mathbb{E}_{\mathcal{B}} \left[\sum_{x} p^{\alpha}(x|Z) \mathbb{1} \{ \mathcal{B}(x) = 1 \} \right] - \frac{1}{\alpha - 1}$$
 (284)

$$=\frac{M^{\alpha-1}}{\alpha-1}\sum_{z}p(z)\sum_{x}p^{\alpha}(x|z)-\frac{1}{\alpha-1}.$$
(285)

(284) is the result of the following inequality for $\alpha > 1$.

$$\left(\sum_{i} x_{i}\right)^{\alpha} \ge \sum_{i} x_{i}^{\alpha}. \tag{286}$$

Let (X^n, Z^n) and $M = [2^{nR}]$, then we have

$$\mathbb{E}_{\mathcal{B}}\left[T_{\alpha}\left(P(b,z^n) \parallel p^{U}(b) p(z^n)\right)\right] \ge 2^{(\alpha-1)n(R-\tilde{H}_{\alpha}(X|Z))} - \frac{1}{\alpha-1},\tag{287}$$

which with $R > \tilde{H}_{\alpha}(X|Z)$, the right side tends to infinity as n increases, and the converse part of the proof is complete.

C. Proof of Theorem 2

Proof.

Achievability part of the proof

$$(\alpha - 1) \cdot \mathbb{E}_{\mathcal{B}} \left[T_{\alpha} \left(P(b, z) \parallel p^{U}(b) p(z) \right) \right]$$

$$= \mathbb{E}_{\mathcal{B}} \left[\sum_{b, z} \frac{p(z)}{M} \cdot \left(M^{\alpha} P^{\alpha}(b|z) - 1 \right) \right]$$
(288)

$$= \mathbb{E}_{\mathcal{B}} \left[\sum_{z} p(z) \left(M^{\alpha} P^{\alpha} (b=1|z) - 1 \right) \right]$$
 (289)

$$= \mathbb{E}_Z \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x} p(x|z) \mathbb{1} \{ \mathcal{B}(x) = 1 \} \right)^{\alpha} - 1 \right]$$
 (290)

$$= \mathbb{E}_{Z} \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \frac{\sum_{x} p(x|z) \mathbb{1} \{ \mathcal{B}(x) = 1 \}}{\left(\sum_{\bar{x}} p(\bar{x}|z) \mathbb{1} \{ \mathcal{B}(\bar{x}) = 1 \} \right)^{1-\alpha}} - 1 \right]$$

$$(291)$$

$$= \mathbb{E}_{Z} \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x} p(x|z) \frac{\mathbb{1} \{ \mathcal{B}(x) = 1 \}}{(\sum_{\bar{x}} p(\bar{x}|z) \mathbb{1} \{ \mathcal{B}(\bar{x}) = 1 \})^{1-\alpha}} \right) - 1 \right]$$
 (292)

$$= \mathbb{E}_{Z} \mathbb{E}_{\mathcal{B}(x)} \mathbb{E}_{\mathcal{B}(\bar{x})|\mathcal{B}(x)} \left[M^{\alpha} \left(\sum_{x} p(x|z) \frac{\mathbb{1} \{ \mathcal{B}(x) = 1 \}}{(\sum_{\bar{x}} p(\bar{x}|z) \mathbb{1} \{ \mathcal{B}(\bar{x}) = 1 \})^{1-\alpha}} \right) - 1 \right]$$
(293)

$$\geq \mathbb{E}_{Z} \mathbb{E}_{\mathcal{B}(x)} \left[M^{\alpha} \left(\sum_{x} p(x|z) \frac{\mathbb{1}\{\mathcal{B}(x) = 1\}}{\mathbb{E}_{\mathcal{B}(\bar{x})|\mathcal{B}(x)} \left[\left(\sum_{\bar{x}} p(\bar{x}|z) \mathbb{1}\{\mathcal{B}(\bar{x}) = 1\} \right)^{1-\alpha} \right]} \right) - 1 \right]$$
(294)

$$\geq \mathbb{E}_{Z} \mathbb{E}_{\mathcal{B}(x)} \left[M^{\alpha} \left(\sum_{x} p(x|z) \frac{\mathbb{1} \{ \mathcal{B}(x) = 1 \}}{\left(\mathbb{E}_{\mathcal{B}(\bar{x})|\mathcal{B}(x)} \left[\sum_{\bar{x}} p(\bar{x}|z) \mathbb{1} \{ \mathcal{B}(\bar{x}) = 1 \} \right] \right)^{1-\alpha}} \right) - 1 \right]$$
(295)

$$\geq \mathbb{E}_{Z} \left[M^{\alpha} \left(\sum_{x} p(x|z) \frac{\frac{1}{M}}{\left(\frac{1}{M} + p(x|z)\right)^{1-\alpha}} \right) - 1 \right]$$
 (296)

$$= \mathbb{E}_{X,Z} \left[\left(\frac{1}{1 + Mp(x|z)} \right)^{1-\alpha} - 1 \right], \tag{297}$$

where (294) and (295) are obtained by applying Jensen's inequality on functions $f(x) = \frac{1}{x}$ and $g(x) = x^{1-\alpha}$, respectively.

Let $M = [2^{nR}]$ and (X^n, Z^n) , it suffices to prove that by increasing n

$$\mathbb{E}_{X,Z}\left[\left(\frac{1}{1+Mp(x|z)}\right)^{1-\alpha}\right] \to 1. \tag{298}$$

Using (297), We define the following typical set.

$$\mathcal{A}_{\epsilon}^{n} = \{ (x^{n}, z^{n}) : \frac{1}{n} h(x^{n}|z^{n}) \ge H(X|Z) - \epsilon \}, \tag{299}$$

which $h(x|z) = \log \frac{1}{p(x|z)}$, and ϵ is an arbitrary positive number. By the weak law of large numbers, we have $\lim_{n\to\infty} p(\mathcal{A}^n_{\epsilon}) = 1$, so we can write

$$\mathbb{E}_{X,Z} \left[\left(\frac{1}{1 + 2^{nR - h(X^n | Z^n)}} \right)^{1 - \alpha} \right] \ge \mathbb{E}_{X,Z} \left[\left(\frac{1}{1 + 2^{nR - h(X^n | Z^n)}} \right)^{1 - \alpha} \mathbb{1} \{ (x^n, z^n) \in \mathcal{A}_{\epsilon}^n \} \right]$$
(300)

$$\geq \mathbb{E}_{X,Z} \left[\left(\frac{1}{1 + 2^{n(R - H(X|Z) + \epsilon)}} \right)^{1 - \alpha} \mathbb{1} \left\{ (x^n, z^n) \in \mathcal{A}_{\epsilon}^n \right\} \right]$$
(301)

$$= p(\mathcal{A}_{\epsilon}^{n}) \left[\left(\frac{1}{1 + 2^{n(R - H(X|Z) + \epsilon)}} \right)^{1 - \alpha} \right]$$
 (302)

$$\rightarrow \left[\left(\frac{1}{1 + 2^{n(R - H(X|Z) + \epsilon)}} \right)^{1 - \alpha} \right]. \tag{303}$$

Therefore, with the increase of n, the condition $R < H(X|Z) - \epsilon$ makes (303) tend to 1 and the achievability proof is complete.

Converse part of the proof

Starting from the definition of T_{α} in (4), we have

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\big(P(b,z^n) \parallel p^U(b)p(z^n)\big)\Big]$$
(304)

$$= \frac{1}{\alpha - 1} \mathbb{E}_{\mathcal{B}} \left[\sum_{b, z^n} M^{\alpha - 1} p(z^n) P^{\alpha}(b|z^n) - 1 \right]$$
(305)

$$= \frac{1}{\alpha - 1} \mathbb{E}_{\mathcal{B}} \left[\sum_{z^n} M^{\alpha} p(z^n) P^{\alpha}(b = 1|z^n) - 1 \right]$$
(306)

$$= \frac{1}{\alpha - 1} \mathbb{E}_{\mathcal{B}} \left[\sum_{z^n} p(z^n) \left(M^{\alpha} P^{\alpha} (b = 1|z^n) - 1 \right) \right]$$
(307)

$$= \frac{1}{\alpha - 1} \sum_{z^n} p(z^n) \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^n} p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \} \right)^{\alpha} - 1 \right], \tag{308}$$

(306) is on account of symmetry, and (308) is the result of substituting (7). Suppose that \mathcal{A}^n_{ϵ} is the ϵ -typical set of z^n sequences defined as

$$\mathcal{A}_{\epsilon}^{n} = \left\{ z^{n} : \left| \frac{1}{n} \sum_{i=1}^{n} \mathbb{1} \left\{ z_{i} = z \right\} - p(z) \right| < \epsilon \right\}. \tag{309}$$

Decomposing the sum into typical and non-typical z^n sequences in (308) gives

$$(308) = \frac{1}{\alpha - 1} \sum_{z^n \in \mathcal{A}_{\epsilon}^n} p(z^n) \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^n} p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \} \right)^{\alpha} - 1 \right]$$

$$+ \frac{1}{\alpha - 1} \sum_{z^n \notin \mathcal{A}_{\epsilon}^n} p(z^n) \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^n} p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \} \right)^{\alpha} - 1 \right]$$

$$\geq \frac{1}{\alpha - 1} \sum_{z^n \in \mathcal{A}_{\epsilon}^n} p(z^n) \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^n} p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \} \right)^{\alpha} - 1 \right]$$

$$+ \frac{1}{\alpha - 1} \sum_{z^n \notin \mathcal{A}_{\epsilon}^n} p(z^n) \left[\left(M \mathbb{E}_{\mathcal{B}} \sum_{x^n} p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \} \right)^{\alpha} - 1 \right]$$

$$= \frac{1}{\alpha - 1} \sum_{z^n \in \mathcal{A}_{\epsilon}^n} p(z^n) \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^n} p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \} \right)^{\alpha} - 1 \right] ,$$

$$(312)$$

where (311) is a direct consequence of applying Jensen's inequality to the concave function $f(x) = x^{\alpha}$, $\alpha \in (0,1)$. Given z^{n} , we define the conditional ϵ' -typical set as

$$\nu_n^* = \left\{ x^n : \left| \frac{1}{n} \sum_{i=1}^n \mathbb{1}\{(x_i, z_i) = (x, z)\} - p(x, z) \right| < \epsilon' \right\}.$$
 (313)

Once again, the approach involves dividing the sum over x^n into two parts: one for typical sequences and the other for non-typical sequences. Building upon (312), we can obtain an upper bound for the expectation term as follows.

$$\mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^{n}} p(x^{n}|z^{n}) \mathbb{I} \left\{ \mathcal{B}(x^{n}) = 1 \right\} \right)^{\alpha} - 1 \right] \tag{314}$$

$$\leq \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^{n} \in \nu_{n}^{*}} p(x^{n}|z^{n}) \mathbb{I} \left\{ \mathcal{B}(x^{n}) = 1 \right\} \right)^{\alpha} - 1 \right]$$

$$+ \mathbb{E}_{\mathcal{B}} \left[\left(M \sum_{x^{n} \notin \nu_{n}^{*}} p(x^{n}|z^{n}) \mathbb{I} \left\{ \mathcal{B}(x^{n}) = 1 \right\} \right)^{\alpha} \right]$$

$$\leq \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^{n} \in \nu_{n}^{*}} p(x^{n}|z^{n}) \mathbb{I} \left\{ \mathcal{B}(x^{n}) = 1 \right\} \right)^{\alpha} - 1 \right]$$

$$+ \left(M \mathbb{E}_{\mathcal{B}} \left[\sum_{x^{n} \notin \nu_{n}^{*}} p(x^{n}|z^{n}) \mathbb{I} \left\{ \mathcal{B}(x^{n}) = 1 \right\} \right)^{\alpha} - 1 \right]$$

$$= \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^{n} \in \nu_{n}^{*}} p(x^{n}|z^{n}) \mathbb{I} \left\{ \mathcal{B}(x^{n}) = 1 \right\} \right)^{\alpha} - 1 \right] + \left(\sum_{x^{n} \notin \nu_{n}^{*}} p(x^{n}|z^{n}) \right)^{\alpha}$$

$$\leq \left(\sum_{x^{n} \in \nu_{n}^{*}} \frac{p(x^{n}|z^{n})}{\left(Mp(x^{n}|z^{n}) \right)^{1-\alpha}} - 1 \right) + \left(\sum_{x^{n} \notin \nu_{n}^{*}} p(x^{n}|z^{n}) \right)^{\alpha},$$
(318)

where (315) is derived from the inequality

$$(x+y)^{\alpha} \le x^{\alpha} + y^{\alpha},\tag{319}$$

which holds for $\alpha \in (0,1)$. On the other hand, (316) follows from Jensen's inequality applied to the concave function $f(x) = x^{\alpha}$ with $\alpha \in (0,1)$.

By taking into account the sequences $x^n \in \nu_n^*$ and conditioning on the knowledge of their binning index, while employing the law of total expectation, it is possible to establish an upper bound for the expectation term in (317). Consequently, this leads to the derivation of (318) in the following manner.

$$\mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^n \in \nu_n^*} p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \} \right)^{\alpha} \right]$$
 (320)

$$= \mathbb{E}_{\mathcal{B}} \left[M^{\alpha} \left(\sum_{x^{n} \in \nu_{n}^{*}} \frac{p(x^{n}|z^{n}) \mathbb{1} \{ \mathcal{B}(x^{n}) = 1 \}}{\left(\sum_{\bar{x}^{n} \in \nu_{n}^{*}} p(\bar{x}^{n}|z^{n}) \mathbb{1} \{ \mathcal{B}(\bar{x}^{n}) = 1 \} \right)^{1-\alpha}} \right) \right]$$
(321)

$$= \mathbb{E}_{\mathcal{B}(\nu_n^*)} \mathbb{E}_{\mathcal{B}|\mathcal{B}(\nu_n^*)} \left[M^{\alpha} \left(\sum_{x^n \in \nu_n^*} \frac{p(x^n | z^n) \mathbb{1} \{ \mathcal{B}(x^n) = 1 \}}{\left(\sum_{\bar{x}^n \in \nu_n^*} p(\bar{x}^n | z^n) \mathbb{1} \{ \mathcal{B}(\bar{x}^n) = 1 \} \right)^{1-\alpha}} \right) \right]$$
(322)

$$= \mathbb{E}_{\mathcal{B}(\nu_n^*)} \left[\frac{1}{M} \sum_{x^n \in \nu_n^*} \frac{M^{\alpha} p(x^n | z^n)}{\left(p(x^n | z^n) + \sum_{x^n \neq \bar{x}^n \in \nu_n^*} p(\bar{x}^n | z^n) \mathbb{1} \{ \mathcal{B}(\bar{x}^n) = 1 \} \right)^{1-\alpha}} + \frac{M-1}{M} \times 0 \right]$$
(323)

$$= \mathbb{E}_{\mathcal{B}(\nu_n^*)} \left| \sum_{x^n \in \nu_n^*} \frac{p(x^n | z^n)}{M^{1-\alpha} \Big(p(x^n | z^n) + \sum_{x^n \neq \bar{x}^n \in \nu_n^*} p(\bar{x}^n | z^n) \mathbb{1} \{ \mathcal{B}(\bar{x}^n) = 1 \} \Big)^{1-\alpha}} \right|$$
(324)

$$\leq \sum_{x^n \in \nu_n^*} \frac{p(x^n | z^n)}{\left(Mp(x^n | z^n)\right)^{1-\alpha}}.$$
(325)

(321) is simply a restatement of (320). In (322), we utilize the law of total expectation by taking into account that expectations are applied to each $x^n \in \nu_n^*$ individually. Then, by applying $\mathbb{E}_{\mathcal{B}|\mathcal{B}(x^n)}$ step-by-step to each sequence $x^n \in \nu_n^*$ and considering that \mathcal{B} is a randomly uniformly mapping, we arrive at (323). Further simplification and the elimination of the summation term in the denominator yield equation (325).

By combining equations (312) and (318), and considering that α is within the range of (0,1), we obtain the following lower bound for Tsallis divergence.

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\big(P(b,z^n) \parallel p^U(b)p(z^n)\big)\Big] \ge \frac{1}{\alpha-1} \sum_{z^n \in \mathcal{A}_{\epsilon}^n} p(z^n) \left(\sum_{x^n \in \nu_n^*} \frac{p(x^n|z^n)}{\left(Mp(x^n|z^n)\right)^{1-\alpha}} - 1\right)$$
(326)

$$+\frac{1}{\alpha-1}\sum_{z^n\in\mathcal{A}_{\epsilon}^n}p(z^n)\left(\sum_{x^n\not\in\nu_n^*}p(x^n|z^n)\right)^{\alpha}.$$
 (327)

For every sequence $x^n \in \nu_n^*$, we have the inequality $p(x^n|z^n) \geq 2^{-n(H(X|Z)+\epsilon'')}$. Hence, when we set $M=2^{nR}$ and choose $R>H(X|Z)+\epsilon''$, the right side of (326) converges to $1/(1-\alpha)$, which is strictly positive. This establishes the completion of the proof.

D. An asymptotic analysis on OSRB rate with special type

Lemma 3. Let p(x,z) be an arbitrary joint probability distribution. Let X^n distributed over $\mathcal{T}_{\epsilon}^n(p_X)$ with distribution of

$$\tilde{p}(x^n) = \frac{\prod_{i=1}^n p(x_i)}{\sum_{x^n \in \mathcal{T}_e^n(p_X)} \prod_{i=1}^n p(x_i)}$$
(328)

$$1 - \delta_n \le \sum_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \prod_{i=1}^n p(x_i)$$
 (329)

and

$$p_{Z^n|X^n} = \prod_{i=1}^n p_{Z_i|X_i} \tag{330}$$

Then for any $\alpha \in (1, \infty)$ and $\epsilon \to 0$ we have

$$\lim_{n \to \infty} \frac{1}{n(\alpha - k - 1)} \log \left(\sum_{z^n, x^n \in \mathcal{T}_{\epsilon}^n(p_X)} q(z^n) \frac{p^{\alpha - k}(x^n, z^n)}{q^{\alpha - k}(z^n)} \right)$$
(331)

$$\leq -H(X) + \sum_{x} (p(x)) D_{\alpha-k}(p(z|x) \parallel p(z))$$
 (332)

which $q(z^n) = \prod_{i=1}^n p(z_i)$.

Proof.

$$\log \left(\sum_{z^n, x^n \in \mathcal{T}_{\epsilon}^n(p_X)} q(z^n) \frac{p^{\alpha - k}(x^n, z^n)}{q^{\alpha - k}(z^n)} \right)$$
(333)

$$\leq -n(\alpha - k - 1)(H(X) - \delta_n(\epsilon)) + \log \left(\sum_{z^n, x^n \in \mathcal{T}_{\epsilon}^n(p_X)} q(z^n) \tilde{p}(x^n) \frac{p^{\alpha - k}(z^n | x^n)}{q^{\alpha - k}(z^n)} \right)$$
(334)

$$\leq -n(\alpha - k - 1)(H(X) - \delta_n(\epsilon)) + \log\left(\sum_{z^n} q(z^n) \frac{p^{\alpha - k}(z^n | \bar{x}^n)}{q^{\alpha - k}(z^n)}\right)$$
(335)

$$= -n(\alpha - k - 1)(H(X) - \delta_n(\epsilon)) + \log\left(\sum_{z^n} \prod_{i=1}^n p(z_i) \frac{p^{\alpha - k}(z_i|\bar{x}_i)}{p^{\alpha - k}(z_i)}\right)$$
(336)

$$\leq -n(\alpha - k - 1)(H(X) - \delta_n(\epsilon)) + \log \prod_{x} \left(\sum_{z} p(z) \frac{p^{\alpha - k}(z|x)}{p^{\alpha - k}(z)} \right)^{n(p(x) + \epsilon)}$$
(337)

$$= -n(\alpha - k - 1)(H(X) - \delta_n(\epsilon)) + n\sum_{x} (p(x) + \epsilon) \log \left(\sum_{z} p(z) \frac{p^{\alpha - k}(z|x)}{p^{\alpha - k}(z)} \right)$$
(338)

$$= -n(\alpha - k - 1)(H(X) - \delta_n(\epsilon)) + n(\alpha - k - 1) \sum_{x} (p(x) + \epsilon) D_{\alpha - k}(p(z|x) \parallel p(z)).$$
 (339)

(334) uses $\frac{1}{n}\log(\tilde{p}(x^n)) \leq -H(X) + \delta_n(\epsilon)$ for $x^n \in \mathcal{T}^n_{\epsilon}(p_X)$. In (335) (\bar{x}^n) fixed since

$$\sum_{z^n, x^n \in \mathcal{T}_{\epsilon}^n(p_X)} q(z^n) \tilde{p}(x^n) \frac{p^{\alpha - k}(z^n | x^n)}{q^{\alpha - k}(z^n)} \le \max_{x^n \in \mathcal{T}_{\epsilon}^n(p_X)} \sum_{z^n} q(z^n) \frac{p^{\alpha - k}(z^n | x^n)}{q^{\alpha - k}(z^n)}$$
(340)

$$= \sum_{z^n} q(z^n) \frac{p^{\alpha-k}(z^n|\bar{x}^n)}{q^{\alpha-k}(z^n)}$$
(341)

finally let $n \to \infty$ and $\epsilon \to 0$, this implies that $\delta_n(\epsilon), \delta_n \to 0$ which completes the proof.

E. An asymptotic analysis on OSRB rate with special type in stochastic encoder

Lemma 4. For $\alpha \in (1, \infty)$ in (171) we have

$$\lim_{n \to \infty} \frac{1}{n(\alpha - 1)} \log \left(\sum_{z^n} q(z^n) \sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \left(\frac{\tilde{p}^{\alpha}(u^n) S^{\alpha}(u^n, z^n)}{q^{\alpha}(z^n)} \right) \right)$$
(342)

$$\leq -H(U) + R'_{\alpha}(p_X, p_{Z|X}, p_U) \tag{343}$$

which $R'_{\alpha}(p_X, p_{Z|X}, p_U)$ defined in (131) and $\epsilon \to 0$.

Proof.

$$\lim_{n \to \infty} \frac{1}{n(\alpha - 1)} \log \left(\sum_{z^n} q(z^n) \sum_{u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \left(\frac{\tilde{p}^{\alpha}(u^n) S^{\alpha}(u^n, z^n)}{q^{\alpha}(z^n)} \right) \right)$$
(344)

$$= \lim_{n \to \infty} \frac{1}{n(\alpha - 1)} \log \left(\sum_{z^n, u^n \in \mathcal{T}_{\epsilon}^n(p_U)} q(z^n) \left(\frac{\tilde{p}^{\alpha}(u^n) S^{\alpha}(u^n, z^n)}{q^{\alpha}(z^n)} \right) \right)$$
(345)

$$\leq \lim_{n \to \infty} \frac{1}{n(\alpha - 1)} \left(\log \left[\max_{u^n \in \mathcal{T}_{\epsilon}(p_U)} \tilde{p}^{\alpha - 1}(u^n) \right] + \log \left[\sum_{z^n, u^n \in \mathcal{T}_{\epsilon}^n(p_U)} q(z^n) \left(\frac{\tilde{p}(u^n) S^{\alpha}(u^n, z^n)}{q^{\alpha}(z^n)} \right) \right] \right)$$
(346)

$$\leq \lim_{n \to \infty} \left[-(H(U) - \delta'_n(\epsilon)) + \frac{1}{n(\alpha - 1)} \log \left((1 + 2\delta_n(\epsilon))^{\alpha - 1} \sum_{z^n, u^n \in \mathcal{T}_{\epsilon}^n(p_U)} q(z^n) \left(\frac{\tilde{p}(u^n) S^{\alpha}(u^n, z^n)}{q^{\alpha}(z^n)} \right) \right) \right]$$
(347)

$$= -H(U) + \lim_{n \to \infty} \frac{1}{n(\alpha - 1)} \log \left(\sum_{z^n, u^n \in \mathcal{T}_{\epsilon}^n(p_U)} \tilde{p}(u^n) S^{\alpha}(u^n, z^n) q^{1-\alpha}(z^n) \right)$$
(348)

Equation (347) follows from (172) with $\delta'_n(\epsilon)$, $\delta_n(\epsilon) \to 0$, when $n \to \infty$ and $\epsilon \to 0$. For correctness of last step in (349), follow (450-455) in [2].

F. T_{α} and D_{∞} on conditional probability distributions

In the following Lemma, we show how T_{α} and D_{∞} on the conditional distribution is related to T_{α} on the joint distribution.

Lemma 5. Let consider arbitrary distribution of $q(z^n)$, let $\alpha \in (1, \infty)$ and assume that

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\Big(P(m,f,z^n) \parallel p^{U}(m) p^{U}(f) q(z^n)\Big)\Big] \le \epsilon, \tag{350}$$

then with probability at least $\frac{1}{2}$ over all pair (\mathcal{B}, F) ,

$$T_{\alpha}\Big(P(m, z^{n}|f) \parallel p^{U}(m) q(z^{n})\Big) \le \eta(\epsilon), \tag{351}$$

where $\eta(\epsilon)$ is a constant depending only on ϵ and it vanishes as ϵ vanishes.

Proof. According to the contractivity property of T_{α} for $\alpha \geq 1$, the following result is obtained from (350).

$$\mathbb{E}_{\mathcal{B}}\Big[T_{\alpha}\Big(P(f) \parallel p^{U}(f)\Big)\Big] \le \epsilon. \tag{352}$$

Then the non-decreasing property of T_{α} with α results

$$\mathbb{E}_{\mathcal{B}}\Big[D\Big(P(f) \parallel p^{U}(f)\Big)\Big] \le \epsilon. \tag{353}$$

In addition, by applying Pinsker's inequality and Jensen's inequality, we have

$$\mathbb{E}_{\mathcal{B}}\Big[\|P(f) - p^{U}(f)\|_{\text{TV}}\Big] \le c\sqrt{\epsilon},\tag{354}$$

where c is a constant and it is equal to $\sqrt{\frac{\ln 2}{2}}$ if we take two as the base of logarithm. From here, with a probability of at least $\frac{7}{8}$, we have

$$||P(f) - p^{U}(f)||_{\text{TV}} \le 8c\sqrt{\epsilon},\tag{355}$$

Further, the assumption (350) implies that with probability at least $\frac{7}{8}$, we have

$$T_{\alpha}\Big(p(m,f,z^n) \parallel p^{U}(m) p^{U}(f) q(z^n)\Big) \le 8\epsilon. \tag{356}$$

Thus with with probability at least $\frac{3}{4}$ both the inequalities (355) and (356) are simultaneously satisfied. Let β is one of those codebooks for which both the inequalities (355) and (356) are simultaneously satisfied.

Representing the total variation distance as a f-divergence, we can rewrite (355) as below,

$$\mathbb{E}_F\left[\left|\frac{p^U(F)}{P(F)} - 1\right|\right] \le 8c\sqrt{\epsilon},\tag{357}$$

which with a probability of at least $1 - \sqrt{8c\sqrt{\epsilon}}$ (over random variable F conditioned on fixed binning β) gives

$$\left| \frac{p^{U}(f)}{p(f)} - 1 \right| \le \sqrt{8c\sqrt{\epsilon}}.\tag{358}$$

In other words, by defining $\delta(\epsilon):=\sqrt{8c\sqrt{\epsilon}}$, we have with probability $\geq 1-\delta(\epsilon)$,

$$1 - \delta(\epsilon) \le \frac{p^{U}(f)}{p(f)} \le 1 + \delta(\epsilon). \tag{359}$$

Let \mathcal{G} be the set of all f for which (359) holds. Further (356) implies

$$1 + 8(\alpha - 1)\epsilon \ge \sum_{m, z^n, f} \frac{p^{\alpha}(m, z^n, f)}{\left(\frac{1}{M}p^{U}(f)q(z^n)\right)^{\alpha - 1}}.$$
 (360)

Using the definition of \mathcal{G} , we have

$$1 + 8(\alpha - 1)\epsilon \ge \sum_{m, z^n, f} \frac{p^{\alpha}(m, z^n, f)}{\left(\frac{1}{M}p^{U}(f)q(z^n)\right)^{\alpha - 1}}$$

$$(361)$$

$$\geq \sum_{(m,z^n,f):f\in\mathcal{G}} \frac{p^{\alpha}(m,z^n,f)}{\left(\frac{1}{M}p^{U}(f)q(z^n)\right)^{\alpha-1}}$$
(362)

$$= \sum_{(m,z^n,f):f\in\mathcal{G}} p(f) \frac{p^{\alpha}(m,z^n|f)}{\left(\frac{1}{M}q(z^n)\right)^{\alpha-1}} \cdot \left(\frac{p(f)}{p^{U}(f)}\right)^{\alpha-1}$$
(363)

$$\geq \frac{1}{(1+\delta(\epsilon))^{\alpha-1}} \sum_{(m,z^n,f):f \in \mathcal{G}} p(f) \frac{p^{\alpha}(m,z^n|f)}{\left(\frac{1}{M}q(z^n)\right)^{\alpha-1}}$$
(364)

$$= \frac{1}{(1+\delta(\epsilon))^{\alpha-1}} \mathbb{E}_F \left[\sum_{(m,z^n)} \frac{p^{\alpha}(m,z^n|F)}{\left(\frac{1}{M}q(z^n)\right)^{\alpha-1}} \mathbb{1}\left\{ F \in \mathcal{G} \right\} \right]$$
(365)

$$= \frac{1}{(1+\delta(\epsilon))^{\alpha-1}} \mathbb{E}_F \Big[\Big(1 + (\alpha - 1) T_\alpha \Big(p(m, z^n | F) \parallel p^U(m) q(z^n) \Big) \Big) \mathbb{1} \{ F \in \mathcal{G} \} \Big]$$
(366)

Define $\tilde{\delta}(\epsilon) := (1 + \delta(\epsilon))^{\alpha - 1}(1 + 8(\alpha - 1)\epsilon) - 1$. Observe that $\tilde{\delta}(\epsilon)$ vanishes as ϵ vanishes. Rearranging (366) implies,

$$(\alpha - 1)\mathbb{E}_{F}\left[\left(T_{\alpha}\left(p(m, z^{n}|F) \parallel p^{U}(m) q(z^{n})\right)\right)\mathbb{1}\left\{F \in \mathcal{G}\right\}\right] \leq 1 + \tilde{\delta}(\epsilon) - \mathbb{P}[F \in \mathcal{G}] \leq \delta(\epsilon) + \tilde{\delta}(\epsilon)$$
(367)

Thus with probability at least $\frac{3}{4} - \mathbb{P}[F \notin \mathcal{G}] \geq \frac{2}{3}$ (for small enough ϵ), we have

$$T_{\alpha}\Big(p(m, z^{n}|F) \parallel p^{U}(m) q(z^{n})\Big) \le \eta(\epsilon)$$
(368)

where $\eta(\epsilon) := \frac{4}{(\alpha - 1)}(\delta(\epsilon) + \tilde{\delta}(\epsilon))$. Finally the probability of occurring (368) over the pair (\mathcal{B}, F) is at least the probability that (355), (356) and (368) occurring simultaneously. By Bayes rule, this probability is at least $\frac{3}{4} \cdot \frac{2}{3} = \frac{1}{2}$.

This statement concludes the proof.

Remark 14. Similar to Lemma 5, another result could be concluded by repeating the steps for D_{∞} instead of T_{α} with

$$\eta'(\epsilon) = 8\epsilon + \log(1 + \delta(\epsilon)),$$
 (369)

where $\eta'(\epsilon)$ replaced of $\eta(\epsilon)$, and $\delta(\epsilon) = \sqrt{8c\sqrt{\epsilon}}$ defined in Lemma 5.

Lemma 6. Let consider arbitrary distribution of $q(z^n)$, suppose there are two distributions over the variables (m, f, x^n, z^n) given by

①
$$p(m|f) p(x^n|m, f) p(z^n|x^n)$$

Assume that the following condition holds for distribution 1.

$$T_{\alpha}\Big(p_1(m,z^n|f) \parallel p^U(m) q(z^n)\Big) \leq \xi.$$

Then, there exists $M' \in \mathcal{M}'$ with $|\mathcal{M}'| \ge |\mathcal{M}|/2$, such that for distribution in 2,

$$T_{\alpha}\Big(p_2(m',z^n|f) \parallel p^U(m') q(z^n)\Big) \leq \delta(\xi).$$

Moreover, the function $\delta(\xi)$ converges to zero as ξ approaches zero.

Proof. Given that

$$T_{\alpha}\Big(p_{1}(m, z^{n}|f) \parallel p^{U}(m) q(z^{n})\Big) = \frac{1}{\alpha - 1} \mathbb{E}_{M, Z^{n}} \left[\left(\frac{p_{1}(M, Z^{n}|F = f)}{p^{U}(M) q(Z^{n})} \right)^{\alpha} - 1 \right] \leq \xi, \quad (370)$$

we can deduce, by employing Markov's inequality, that a minimum of 3/4 of the instances $m \in \mathcal{M}$, satisfy

$$\mathbb{E}_{Z^n} \left[\left(\frac{p_1(m, Z^n | F = f)}{p^U(m) q(Z^n)} \right)^{\alpha} \right] \le 1 + 4(\alpha - 1)\xi.$$

$$(371)$$

Furthermore, it can be deduced from (370), by combining the contractivity property of the Tsallis divergence with Pinsker's inequality, that

$$\mathbb{E}_{M} \left[\left| \frac{p(M|F = f)}{p^{U}(M)} - 1 \right| \right] = \left\| p(m|F = f) - p^{U}(m) \right\|_{\text{TV}} \le c\sqrt{\xi}, \tag{372}$$

where $c = \sqrt{\ln 2/2}$. Thus, employing Markov's inequality once again, it can be inferred that for over 3/4 of realizations of $m \in \mathcal{M}$, the following inequality holds.

$$1 - 4c\sqrt{\xi} \le \frac{p(m|F = f)}{p^U(m)} \le 1 + 4c\sqrt{\xi}$$
(373)

Hence, for a minimum of half of the \mathcal{M} members, both constraints (371) and (373) are met. By selecting these qualified members and assigning them to the set \mathcal{M}' with $|\mathcal{M}'| \geq |\mathcal{M}|/2$, we obtain

$$T_{\alpha}\left(p_{2}(m', z^{n}|f) \parallel p^{U}(m') q(z^{n})\right)$$

$$= \frac{1}{\alpha - 1} \mathbb{E}_{M', Z^{n}} \left[\left(\frac{p_{2}(M', Z^{n}|F = f)}{p_{1}(M', Z^{n}|F = f)}\right)^{\alpha} \left(\frac{p_{1}(M', Z^{n}|F = f)}{p^{U}(M') q(Z^{n})}\right)^{\alpha} - 1 \right]$$
(374)

$$= \frac{1}{\alpha - 1} \left[\mathbb{E}_{M', Z^n} \left[\left(\frac{p^U(M')}{p(M'|F = f)} \right)^{\alpha} \left(\frac{p_1(M', Z^n|F = f)}{p^U(M') \, q(Z^n)} \right)^{\alpha} \right] - 1 \right]$$
(375)

$$\leq \frac{1}{\alpha - 1} \left[(1 + 4(\alpha - 1)\xi) \left(\frac{1}{1 - 4c\sqrt{\xi}} \right)^{\alpha} - 1 \right] = \delta(\xi), \tag{376}$$

where (376) are derived from (373) and (371), respectively.

Remark 15. Similar to Lemma 6, holds for D_{∞} instead of T_{α} with

$$\delta'(\xi) = \xi + \log\left(\frac{1}{1 - 2c\sqrt{\xi}}\right),\tag{377}$$

replaced of $\delta(\xi)$, and $c = \sqrt{\frac{\ln 2}{2}}$.

Remark 16. For generalization of Lemma 6 from deterministic encoder to stochastic encoder consider new distributions

②
$$p^{U}(m)\,\tilde{p}(u^{n}|m,f)\,\tilde{p}(x^{n}|u^{n})\,p(z^{n}|x^{n}),$$
 (379)

over variables (m, f, u^n, x^n, z^n) , and repeat the steps similar to lemma 6.