

ON THE RATE OF GENERIC GORENSTEIN K -ALGEBRAS

MATS BOIJ, EMANUELA DE NEGRI*, ALESSANDRO DE STEFANI*, AND MARIA EVELINA ROSSI*

ABSTRACT. The rate of a standard graded K -algebra A is a measure of the growth of the shifts in a minimal free resolution of K as an A -module. In particular A has rate one if and only if it is Koszul. It is known that a generic Artinian Gorenstein algebra of embedding dimension $n \geq 3$ and socle degree $s = 3$ is Koszul. We prove that a generic Artinian Gorenstein algebra with $n \geq 4$ and $s \geq 3$ has rate $\lfloor \frac{s}{2} \rfloor$. In the process we show that such an algebra is generated in degree $\lfloor \frac{s}{2} \rfloor + 1$. This gives a partial positive answer to a longstanding conjecture stated by the first author on the minimal free resolution of a generic Artinian Gorenstein ring of odd socle degree.

1. INTRODUCTION AND PRELIMINARY RESULTS

Let K be a field and $A = \bigoplus_{i \in \mathbb{N}} A_i$ be a finitely generated \mathbb{N} -graded K -algebra. We recall that A is said to be standard graded if $A_0 = K$ and A is generated (as a K -algebra) by elements of degree 1, that is, $A = K[A_1]$. We denote by \mathfrak{m} the homogeneous maximal ideal $\bigoplus_{i \geq 1} A_i$ of A . We let $\text{edim}(A) = \dim_K(A_1)$ be the embedding dimension of A , $\text{soc}(A) = 0 :_A \mathfrak{m}$ be its socle, which is a homogeneous ideal of A , and $\text{socdeg}(A) = \sup\{i \in \mathbb{N} \mid \text{soc}(A)_i \neq 0\}$. Observe that, if A is Artinian, then $\text{socdeg}(A) = \max\{i \in \mathbb{N} \mid A_i \neq 0\}$. Consider a minimal graded free resolution

$$\mathbb{F} : \cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_0 \rightarrow K \rightarrow 0$$

of $K = A/\mathfrak{m}$ as an A -module. It is well-known that \mathbb{F} plays an important role in the study of the homological properties of A . For instance, by the Auslander-Buchsbaum-Serre's Theorem, $F_i = 0$ for $i \gg 0$ if and only if A is regular, i.e., A is isomorphic to a polynomial ring over K .

There are several invariants attached to \mathbb{F} . An important one is the Backelin rate, which measures the growth of the shifts in \mathbb{F} ; we now recall its definition. For any integer $i \geq 0$ we let $t_i^A(K) = \sup\{j \mid \text{Tor}_i^A(K, K)_j \neq 0\}$ where, by convention, $t_i^A(K) = -\infty$ whenever $\text{Tor}_i^A(K, K) = 0$. We observe that $\text{Tor}_i^A(K, K)$ is a finitely generated graded K -vector space, therefore the integer $t_i^A(K)$ is indeed well-defined. It readily follows from the given definitions and the Auslander-Buchsbaum-Serre's Theorem that, if A is not regular, then $t_i^A(K) \geq i$ for all $i \geq 0$. In the following we assume A not regular, hence \mathbb{F} is an infinite resolution. The Backelin rate of A (see [5]) is defined as

$$\text{rate}(A) = \sup_{i \geq 2} \left\{ \frac{t_i^A(K) - 1}{i - 1} \right\}.$$

It turns out that the rate of any standard graded K -algebra is finite, see [1, 2]. Moreover, it follows from our previous considerations that $\text{rate}(A) \geq 1$ always holds, and one has that $\text{rate}(A) = 1$ if and only if A is Koszul. In this sense, the rate of A can be regarded as a measure of how far is A from being Koszul. Consider a minimal presentation of A as a quotient of a polynomial ring, i.e. $A \cong R/I$ where $R = K[x_1, \dots, x_n]$ is a polynomial ring and I is an ideal generated by homogeneous

* The three authors were partially supported by the PRIN 2020 project 2020355B8Y "Squarefree Gröbner degenerations, special varieties and related topics".

elements of degree > 1 (or, equivalently, $n = \text{edim}(A)$). Let $m(I)$ be the maximum degree of a minimal homogeneous generator of I . It follows from (the graded version of) [7, Thm. 2.3.2] that $t_2(K) = m(I)$, and therefore one has

$$\text{rate}(R/I) \geq m(I) - 1.$$

In [5] Backelin proved that $\text{rate}(R/I) = m(I) - 1$ for any monomial ideal I (see also [15]). Another interesting class of standard graded K -algebras with minimal rate is given by generic toric rings, as proved by Gasharov, Peeva and Welker in [17]. Extending a result of Kempf [21], Conca, De Negri and Rossi in [11] proved that, under suitable conditions, the coordinate ring of generic points in \mathbb{P}^n has minimal rate. It is an interesting problem to provide further classes of algebras of minimal rate.

Artinian Gorenstein K -algebras of fixed embedding dimension n and socle degree s provide a good setting for studying this question, as they can be parametrized by points in a suitable projective space via the correspondence given by Macaulay's inverse system. In fact they are in bijective correspondence with polynomials in n variables which are homogeneous of degree s . Artinian Gorenstein K -algebras which have maximal length in this family are called compressed; it is known that, when A is a generic element of such a family, A is compressed [19]. Conca, Rossi and Valla proved that a generic Artinian Gorenstein algebra A of socle degree three and $n \geq 3$ is Koszul, see [13]. The result stimulated several interesting questions and recent results on being $A = R/I$ Koszul, provided I is generated by quadrics (see for instance [25, 26]). Observe that if the socle degree of A is two, then quadratic always implies Koszul. The main goal of this paper is to prove the analogue of Conca, Rossi and Valla's result for higher socle degrees in terms of the rate. We prove the following result.

Theorem A. *Let A be a generic Artinian Gorenstein graded K -algebra of socle degree $s \geq 3$, and assume that $\text{edim}(A) \geq 4$. Then $\text{rate}(A) = \lfloor \frac{s}{2} \rfloor$.*

A first important step for proving Theorem A is to provide an upper bound on $t_i^A(K)$ for all $i \geq 0$ in terms of $m(I)$, see Proposition 3.5. Several interesting bounds on $t_i^A(K)$ have been proved in the literature, see for instance [3, 4, 14]. However, the one which is needed for our purposes is obtained from an explicit computation of the bigraded Poincaré series of K , see Proposition 3.4. The proof is based on the graded version of an argument due to Rossi and Şega, used in [28] in the local case. As a consequence of the above results, we prove that if $A = R/I$ is Gorenstein compressed with socle degree $s \geq 4$ and if I is generated by homogeneous polynomials of the same degree, then the rate is minimal, that is $\lfloor \frac{s}{2} \rfloor$. Then the key step in the proof of our main result is to show that a K -algebra satisfying the assumptions of Theorem A has generators in the same degree.

To this end, recall that a generic Artinian Gorenstein K -algebra A of socle degree $s = 3$ and embedding dimension $n \geq 3$ is generated by quadrics. An Artinian compressed Gorenstein algebra of even socle degree $s = 2(t - 1) \geq 2$ has almost linear resolution; as a consequence, it is minimally generated only in degree t , see [19, Example 4.7]. On the other hand, an Artinian compressed Gorenstein K -algebra $A = R/I$ of odd socle degree $s = 2t - 1 \geq 3$ always has generators in degree t , and it has no generators in degrees different than t or $t + 1$ [6, Proposition 3.2]. If $A = R/I$ is a generic Artinian Gorenstein K -algebras of odd socle degree $s = 2t - 1 \geq 3$ and $n \geq 4$, the graded Betti numbers in a minimal free resolution of A as R -module were conjectured by Boij [6, Conjecture 3.13] in terms of n and t . For every homological position there exists a Zariski open set (depending on the coefficients of the form F of degree s) where the Artinian compressed

Gorenstein algebras have the same graded Betti numbers. The conjecture in particular predicts that I should be minimally generated in degree t .

In this article we give a positive answer to this part of the conjecture. In order to do this, we need to show that the Zariski open set parametrizing the Artinian compressed Gorenstein algebras generated in the same degree t is not empty.

Theorem B. *For all integers $t \geq 2$ and $n \geq 4$ there exists an Artinian compressed Gorenstein K -algebra $A = R/I$ with socle degree $s = 2t - 1$ and embedding dimension n such that I is generated in degree t .*

The first step in our proof is to produce an Artinian level algebra with socle degree s defined by monomials of degree s , see Lemma 4.2. The required Artinian compressed Gorenstein K -algebra is then obtained from it using a tricky construction, see Theorem 4.3.

We point out that Boij's conjecture fails in general for generic Artinian Gorenstein algebras but in the middle of the resolution, not at the level of generators, consistently with Theorem B. In fact, counterexamples were exhibited by Kunte [23], see Corollary 5.6 and Remark 5.7.

2. LG_t -ALGEBRAS AND RATE

In this section we present some basic results on the rate of standard graded K -algebras. Consider a minimal presentation of A as a quotient of a polynomial ring, i.e. $A \cong R/I$ where $R = K[x_1, \dots, x_n]$ is a polynomial ring and $I \neq 0$ is a homogeneous ideal generated in degree at least two.

Let τ be a monomial order on R . Denote by $\text{in}_\tau(I)$ the initial ideal of I with respect to τ . In [8, Cor. 2.4, Thm.2.2] it is proved that:

$$\text{rate}(A) \leq \text{rate}(R/\text{in}_\tau(I)).$$

It is known that if $0 \neq J \subseteq R$ is any monomial ideal, then $\text{rate}(R/J) = m(J) - 1$ (see [5, 15]). Putting all these facts together one has that

$$m(I) - 1 \leq \text{rate}(A) \leq m(\text{in}_\tau(I)) - 1.$$

It follows that if I is generated by a Gröbner basis of forms of degree at most $m(I)$ with respect to some coordinate system and some monomial order, then the rate is minimal, that is, $\text{rate}(A) = m(I) - 1$. Another interesting class of standard graded K -algebras with minimal rate is given by generic toric rings, see [17].

Motivated by the previous observations, we make the following definitions.

Definition 2.1. Let A be a standard graded K -algebra. For an integer $t > 1$ we say that A is a G_t -algebra if it has a defining ideal I with respect to some minimal presentation $A \cong R/I$ possessing a Gröbner basis of forms of degree at most t . We say that A is an LG_t -algebra if there exists a graded G_t -algebra S and a regular sequence y_1, \dots, y_r of linear forms of S such that $A \cong S/(y_1, \dots, y_r)$.

Observe that being a G_2 -algebra (resp. an LG_2 -algebra) is equivalent to being G -quadratic (resp. LG -quadratic). LG -quadratic algebras are Koszul, see [10, 12]. The following proposition extends this result to the case $t \geq 2$.

Proposition 2.2. *Let A be an LG_t -algebra. Then $\text{rate}(A) \leq t - 1$.*

Proof. First, we claim that if $A = S/(y)$ where $y \in S$ is a regular element of degree one, then $\text{rate}(A) = \text{rate}(S)$. By the graded version of [29, Satz 1] we have that $P_K^S(u, v) = (1 + uv)P_K^A(u, v)$. In particular, for all $i \geq 2$ we have that $\beta_{ij}^S(K) = \beta_{i-1,j}^A(K) + \beta_{ij}^A(K)$. Thus, for all $i \geq 2$ one has $t_i^S(K) = \max\{t_i^A(K), t_{i-1}^A(K) + 1\}$. Note that for every $i \geq 3$ one has $t_{i-1}^A(K)/(i-1) \leq (t_{i-1}^A(K) - 1)/(i-2)$. Indeed, this is equivalent to the fact that $t_{i-1}^A(K) \geq i-1$, which holds true since A is assumed to be not regular. This together with the fact that $t_1^A(K) = 1$ gives

$$\text{rate}(S) = \sup_{i \geq 2} \left\{ \frac{t_1^A(K) - 1}{i-1}, \frac{t_{i-1}^A(K)}{i-1} \right\} = \sup_{i \geq 2} \left\{ \frac{t_i^A(K) - 1}{i-1} \right\} = \text{rate}(A).$$

Now assume that A is an LG_t -algebra, i.e., there exists a G_t -algebra S such that $A \cong S/(y_1, \dots, y_r)$, with y_1, \dots, y_r a regular sequence of linear forms. A repeated application of the above claim together with the fact that G_t -algebras have rate at most $t-1$ gives that $\text{rate}(A) \leq t-1$. \square

A standard application of Proposition 2.2 gives a proof of the following generalization of a result of Tate [30], which states that every quadratic complete intersection is Koszul.

Corollary 2.3. *Let A be a standard graded K -algebra which is a complete intersection. Assume that the defining ideal of A is generated by a regular sequence of degrees at most t . Then $\text{rate}(A) = t-1$.*

Proof. Consider a minimal presentation $A \cong R/I$ with $R = K[x_1, \dots, x_n]$ a standard graded polynomial ring, and $I = (f_1, \dots, f_c)$ an ideal generated by a regular sequence with $m(I) = t$. Let $d_i = \deg(f_i)$. Let $S = R[y_1, \dots, y_c]$, $g_i = y_i^{d_i} - f_i$ for $i = 1, \dots, c$, and set $J = (g_1, \dots, g_c)$. Fix a monomial order τ on S such that $y_1 > y_2 > \dots > y_c > x_1 > \dots > x_n$. Then the leading terms $\text{in}_\tau(g_1) = y_1^{d_1}, \dots, \text{in}_\tau(g_c) = y_c^{d_c}$ form a regular sequence, and thus g_1, \dots, g_c are a Gröbner basis of forms of degree at most t . As y_1, \dots, y_c are a regular sequence of linear forms in S , it follows that $A \cong S/(y_1, \dots, y_c)$ is an LG_t -algebra, and the result follows from Proposition 2.2 and the general fact that $\text{rate}(A) \geq m(I) - 1 = t-1$. \square

3. THE GRADED POINCARÉ SERIES OF A COMPRESSED GORENSTEIN K -ALGEBRA

Let A be a standard graded K -algebra and M be a finitely generated \mathbb{Z} -graded A -module. The graded Poincaré series of M is defined as

$$P_M^A(u, v) = \sum_{i \geq 0} \left(\sum_{j \in \mathbb{Z}} \beta_{ij}^A(M) v^j \right) u^i \in \mathbb{Z}[v, v^{-1}][[u]],$$

where $\beta_{ij}^A(M) = \dim_K(\text{Tor}_i^A(M, K)_j)$. Note that, fixed an integer $i \geq 0$, one has that $\text{Tor}_i^A(M, K)_j = 0$ for all but finitely many $j \in \mathbb{Z}$. Therefore each sum $\sum_{j \in \mathbb{Z}} \beta_{ij}^A(M) v^j$ is finite.

Given a minimal presentation $A = R/I$ with $R = K[x_1, \dots, x_n]$, there is a point-wise inequality

$$(1) \quad P_K^A(u, v) \leq \frac{P_K^R(u, v)}{1 - u(P_A^R(u, v) - 1)}.$$

Definition 3.1. A standard graded K -algebra A is said to be Golod in the graded sense if equality holds in (1).

Remark 3.2. Since $\beta_i^A(K) = \beta_i^{A_{\mathfrak{m}}}(A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}})$ for all i , and because one always has a point-wise inequality between graded Betti numbers in (1), it turns out that A is Golod in the graded sense if and only if $A_{\mathfrak{m}}$ is a Golod local ring.

With the aim of bounding $t_A^i(K)$ for every $i > 0$, the goal of this section is to compute the graded Poincaré series of $K = A/\mathfrak{m}$ for any Artinian compressed Gorenstein algebra A .

Assume that A is an Artinian Gorenstein standard graded K -algebra. Then we can write $A = \bigoplus_{i=0}^s A_i$ with $A_s \cong K$. The integer s is called the socle degree of A . The Hilbert function of A is by definition the function $\text{HF}_A : \mathbb{N} \rightarrow \mathbb{N}$, defined as $\text{HF}_A(i) = \dim_K(A_i)$.

Artinian Gorenstein algebras of fixed embedding dimension n and socle degree s can be parametrized by points in projective space via the Macaulay inverse system and from this one can deduce that for every integer i

$$\text{HF}_A(i) \leq \min \left\{ \binom{n+i-1}{i}, \binom{n+s-i-1}{s-i} \right\}.$$

Definition 3.3. An Artinian Gorenstein K -algebra A of socle degree s is said to be *compressed* if for all $i \in \{0, \dots, s\}$ one has

$$\text{HF}_A(i) = \min \left\{ \binom{n+i-1}{i}, \binom{n+s-i-1}{s-i} \right\}.$$

There exists a non-empty Zariski open set of Artinian Gorenstein algebras with maximal Hilbert function and several authors investigated Artinian Gorenstein compressed algebras because they provide a good setting for studying interesting questions (see [6, 18–20, 28]).

The following result is a graded version of [28, Proposition 6.2].

Proposition 3.4. *Let A be an Artinian compressed Gorenstein K -algebra of embedding dimension $n \geq 2$ and socle degree s , with $2 \leq s \neq 3$. Then*

$$P_K^A(u, v) = \frac{(1+uv)^n}{1 - u(P_A^R(u, v) - 1) + u^{n+1}v^{n+s}(1+u)}.$$

Proof. By [24], the natural map $A \rightarrow T = A/\text{soc}(A)$ is Golod, therefore

$$P_K^T(u, v) = \frac{P_K^A(u, v)}{1 - u(P_T^A(u, v) - 1)}.$$

From the graded short exact sequence $0 \rightarrow \text{soc}(A) \cong K(-s) \rightarrow A \rightarrow T \rightarrow 0$, we see that $P_T^A(u, v) = 1 + uv^s P_K^A(u, v)$, and therefore

$$P_K^T(u, v) = \frac{P_K^A(u, v)}{1 - u^2 v^s P_K^A(u, v)},$$

from which we conclude

$$P_K^A(u, v) = \frac{P_K^T(u, v)}{1 + u^2 v^s P_K^T(u, v)}.$$

Since $A_{\mathfrak{m}}$ satisfies the assumptions of [28, Theorem 5.1], we have that $T_{\mathfrak{m}}$ is a Golod ring (see the paragraph before [28, Proposition 6.3]). Thus, we have that T is Golod in the graded sense by

Remark 3.2, and therefore

$$P_K^T(u, v) = \frac{P_K^R(u, v)}{1 - u(P_T^R(u, v) - 1)} = \frac{(1 + uv)^n}{1 - u(P_T^R(u, v) - 1)}.$$

Consider again the graded short exact sequence $0 \rightarrow \text{soc}(A) \cong K(-s) \rightarrow A \rightarrow T \rightarrow 0$. As the map $\text{Tor}_i^R(\text{soc}(A), K) \rightarrow \text{Tor}_i^R(A, K)$ is zero for $i \neq n$, we obtain the following exact sequences

$$0 \longrightarrow \text{Tor}_i^R(A, K) \longrightarrow \text{Tor}_i^R(T, K) \longrightarrow \text{Tor}_{i-1}^R(K, K)(-s) \longrightarrow 0$$

for all $0 < i < n$, and

$$0 \longrightarrow \text{Tor}_n^R(K, K)(-s) \longrightarrow \text{Tor}_n^R(A, K) \longrightarrow \text{Tor}_n^R(T, K) \longrightarrow \text{Tor}_{n-1}^R(K, K)(-s) \longrightarrow 0.$$

Therefore for $0 < i < n$ we have

$$\beta_{i,j}^R(T) = \beta_{i,j}^R(A) + \beta_{i-1, j-s}^R(K),$$

while for $i = n$ we have

$$\beta_{n,j}^R(T) = \beta_{n,j}^R(A) + \beta_{n-1, j-s}^R(K) - \beta_{n, j-s}^R(K).$$

Putting all together we have

$$\begin{aligned} P_T^R(u, v) &= \sum_{i,j} \beta_{i,j}^R(T) u^i v^j \\ &= \sum_{\substack{0 \leq i \leq n \\ j \in \mathbb{Z}}} \beta_{i,j}^R(A) u^i v^j + uv^s \sum_{\substack{1 \leq i \leq n \\ j \in \mathbb{Z}}} \beta_{i-1, j-s}^R(K) u^{i-1} v^{j-s} - \sum_{j \in \mathbb{Z}} \beta_{n, j-s}^R(K) \\ &= P_A^R(u, v) + uv^s P_K^R(u, v) - u^n \sum_{j \in \mathbb{Z}} (\beta_{n, j-s}^R(K) uv^j + \beta_{n, j-s}^R(K) v^j) \\ &= P_A^R(u, v) + uv^s P_K^R(u, v) - u^n (u+1) v^{n+s} = P_A^R(u, v) + uv^s (1+uv)^n - u^n (u+1) v^{n+s} \end{aligned}$$

where we used that $\beta_{n, j-s}^R(K) = \begin{cases} 1 & \text{if } j-s = n \\ 0 & \text{otherwise.} \end{cases}$

Substituting the identities obtained so far in the formula for $P_K^A(u, v)$, we finally get that

$$\begin{aligned} P_K^A(u, v) &= \frac{P_K^T(u, v)}{1 + u^2 v^s P_K^T(u, v)} \\ &= \frac{(1 + uv)^n}{1 - u(P_T^R(u, v) - 1) + u^2 v^s (1 + uv)^n} \\ &= \frac{(1 + uv)^n}{1 - u(P_A^R(u, v) - 1) + u^{n+1} v^{n+s} (1 + u)}. \end{aligned} \quad \square$$

Proposition 3.5. *Let A be an Artinian compressed Gorenstein graded K -algebra of embedding dimension $n \geq 2$ and socle degree $s \geq 4$, and let $A = R/I$ be a minimal presentation. Then $t_i^A(K) \leq (m(I) - 1)(i - 1) + 1$.*

Proof. Let $t = \lfloor \frac{s}{2} \rfloor + 1$. Since A is a compressed Gorenstein K -algebra, by [16] (see also [6]) a minimal free resolution of A over R has the following shape:

$$0 \longrightarrow R(-(n+s)) \longrightarrow \bigoplus_{j=1}^{n-1} R(-(n+t-2))^{a_{n-1}} \longrightarrow \dots \longrightarrow \bigoplus_{j=1}^{n-1} R(-(n+t-1))^{b_{n-1}} \longrightarrow R \longrightarrow A \longrightarrow 0.$$

By Proposition 3.4 we therefore obtain

$$\begin{aligned} P_K^A(u, v) &= \frac{(1+uv)^n}{1 - u(\sum_{j=1}^{n-1} (a_j + b_j v) u^j v^{t+j-1}) + u^n v^{n+s} + u^{n+1} v^{n+s} (1+u)} \\ &= \frac{(1+uv)^n}{1 - \sum_{j=1}^{n-1} (a_j + b_j v) u^{j+1} v^{t+j-1} + u^{n+2} v^{n+s}}. \end{aligned}$$

Write $P_K^A(u, v) = \sum_{i \geq 0} \beta_i(v) u^i$, where $\beta_i(v) \in \mathbb{Z}[v]$. For simplicity, for all $i \in \mathbb{N}$ we let $\delta_i = \deg(\beta_i(v)) = t_i^A(K)$, where $\deg(-)$ denotes the degree of a polynomial in the variable v . By clearing denominators and equating the coefficients of u^i in both sides of the previous identity we deduce that

$$(2) \quad \beta_i = \begin{cases} 1 & \text{if } i = 0 \\ nv & \text{if } i = 1 \\ \binom{n}{i} v^i + \sum_{j=1}^{i-1} \beta_{i-j-1} (a_j + b_j v) v^{t+j-1} & \text{if } 2 \leq i \leq n \\ \sum_{j=1}^{n-1} \beta_{n-j+1} (a_j + b_j v) v^{t+j-1} + v^{n+s} & \text{if } i = n+2 \\ \sum_{j=1}^{n-1} \beta_{i-j-1} (a_j + b_j v) v^{t+j-1} & \text{otherwise} \end{cases}$$

We distinguish two cases. If $b_1 = 0$, then by induction on $i \geq 1$ we prove that $\delta_i \leq (t-1)(i-1) + 1$. Note that for $i = 1, 2$ the inequality is satisfied since $\delta_1 = 1$ and $\delta_2 = t$; thus we may assume $i \geq 3$. For $2 \leq j = i-1 \leq n-1$ we have that $\deg(\beta_0(a_{i-1} + b_{i-1}v)v^{t+i-2}) \leq t+i-1 \leq (t-1)(i-1) + 1$. For $1 \leq j \leq n-1$ with $j \neq i-1$, if we assume that $\delta_{i-j-1} \leq (t-1)(i-j-2) + 1$ then

$$(3) \quad \deg(\beta_{i-j-1}(a_j + b_j v)v^{t+j-1}) \leq (t-1)(i-j-2) + 1 + t + j \leq (t-1)(i-1) + 1,$$

where the latter holds since $t \geq 2$ in our assumptions. One can check directly that $i \leq (t-1)(i-1) + 1$ holds. Finally, since $s \leq 2t-1$, $n \geq 2$ and $t \geq 3$, we also have that $n+s \leq (t-1)(n+1) - 1$. All these inequalities, together with (2) and the induction hypothesis which allows to use (3) for appropriate values of j , give that $\delta_i \leq (t-1)(i-1) + 1$. Note that the assumption $b_1 = 0$ guarantees that $m(I) = t$, and since $\delta_i = t_i^A(K)$ we proved that $t_i^A(K) \leq (m(I) - 1)(i-1) + 1$, as desired.

Now assume that $b_1 \neq 0$, so that $m(I) = t+1$. In this case, we prove by induction on $i \geq 1$ that $\delta_i \leq t(i-1) + 1$. The strategy is similar to the one used in the previous case. Since $\delta_1 = 1$ the claimed inequality is true for the base case, and we may assume $i \geq 2$. For $1 \leq j = i-1 \leq n-1$ we have that $\deg(\beta_0(a_{i-1} + b_{i-1}v)v^{t+i-2}) \leq t+i-1 \leq t(i-1) + 1$. Observe that, for $1 \leq j \leq n-1$ with $j \neq i-1$, if we assume that $\delta_{i-j-1} \leq t(i-j-2) + 1$ then

$$(4) \quad \deg(\beta_{i-j-1}(a_j + b_j v)v^{t+j-1}) \leq \delta_{i-j-1} + t + j \leq t(i-j-2) + 1 + t + j \leq t(i-1) + 1.$$

Finally, we have that $i \leq t(i-1) + 1$ for all $i \geq 2$ and $n+s \leq t(n+1) - 1$ since $s \leq 2t-1$. These inequalities, together with (2) and the induction hypothesis which allows to use (4) for all appropriate values of j , give that $t_i^A(K) = \delta_i \leq t(i-1) + 1 = (m(I) - 1)(i-1) + 1$. \square

In [13, Question 6.10] the authors ask if any quadratic Gorenstein algebra of socle degree three is Koszul. However, for any embedding dimension $n \geq 8$ there are examples of Gorenstein quadratic algebras of regularity three which are not Koszul, see [25, 27].

A surprising consequence of Proposition 3.5 is that compressed Gorenstein algebras with $s \geq 4$ and generated in one degree have minimal rate.

Corollary 3.6. *Let A be an Artinian compressed Gorenstein graded K -algebra of socle degree $s \geq 4$, and let $A = R/I$ be a minimal presentation. Assume I is generated in degree $\lfloor \frac{s}{2} \rfloor + 1$. Then $\text{rate}(A) = \lfloor \frac{s}{2} \rfloor$.*

4. THE RATE OF A GENERIC GORENSTEIN K -ALGEBRA

Every Artinian Gorenstein graded algebra $A = R/I$ of socle degree s , corresponds, up to scalars, to a form F of degree s in another set of variables as follows. Let $R = K[x_1, \dots, x_n]$ and $B = K[y_1, \dots, y_n]$. Regard B as a R -module via the action $x_i \circ F = \partial F / \partial y_i$; this action can be extended to every element $G \in R$ acting as a differential operator on the elements of B . Given a form F of degree s , we denote by I_F the ideal of the elements of R which annihilate F :

$$I_F = \{G \in R \mid G \circ F = 0\}.$$

Set $A_F = R/I_F$. Then A_F is a standard Artinian Gorenstein algebra of socle degree s . Moreover, every ideal $I \subset R$ which defines a standard graded Artinian Gorenstein algebra of socle degree s arises in this way. This correspondence is known as Macaulay's Inverse System. For more information we refer to the book of Iarrobino and Kanev [20].

Definition 4.1. Assume that K is infinite. We say that an Artinian Gorenstein K -algebra A of socle degree s is generic if $A = A_F$ with $F \in B$ a generic form of degree s .

In particular generic Artinian Gorenstein K -algebras are compressed, see [20, Proposition 3.12].

4.1. Generators of generic Artinian Gorenstein algebras. It is known that Artinian compressed Gorenstein K -algebras of even socle degree $s = 2(t - 1) > 1$ have almost linear resolution (see [19, 4.7]). That is, if $n = \text{edim}(A)$ and $A = R/I$ is a minimal presentation, then a minimal free resolution of A over R is almost linear:

$$0 \rightarrow R(-(n+s)) \rightarrow R^{a_{n-1}}(-(n+t-2)) \rightarrow \dots \rightarrow R(-(t+1)) \rightarrow R^{a_1}(-t) \rightarrow R \rightarrow A \rightarrow 0.$$

In particular

$$m(I) = t = s/2 + 1.$$

If $s = 2t - 1$ and $n = 3$ the Betti numbers of generic Gorenstein algebras are known, see for instance [6, Example 3.16], and the following holds:

$$m(I) = \begin{cases} t & \text{if } t \text{ is even} \\ t+1 & \text{if } t \text{ is odd} \end{cases}.$$

The values of the graded Betti numbers of generic Artinian Gorenstein algebras with $n > 3$ and odd socle degree $s = 2t - 1$ were conjectured by Boij [6, Conjecture 3.13]. In fact, Boij proved that there exists a Zariski open set of Artinian compressed Gorenstein algebras all sharing the same numerical resolution, and he conjectured that such a set is non-empty. Computer generated evidence was given in [6, Section 4], but the conjecture was later disproved for special values of

n and s by Kunte [23, Corollary 5.6, Remark 5.7]. However, in all known counterexamples the failure is in the middle of the resolution. The aim of this section is to prove that Boij's conjecture holds true at the beginning of the resolution. Namely, we prove that generic Artinian Gorenstein algebras of socle degree $s = 2t - 1$ are minimally generated in degree t .

We recall that an Artinian K -algebra A of socle degree s is called level if $\text{soc}(A) \subseteq A_s$. The construction of the following lemma will be crucial in what follows.

Lemma 4.2. *For all integers $t \geq 2$ and $m \geq 3$ there exists a monomial ideal $J \subseteq S = K[x_1, \dots, x_m]$ such that $m(J) = t$ and $A = S/J$ is an Artinian level algebra with socle degree t .*

Proof. We start by considering the even case $t = 2u$ with $u \geq 1$. Given $m \geq 3$, we let j be an integer such that $1 \leq j < m$. We set $X = (x_1, \dots, x_j)$, $Y = (x_{j+1}, \dots, x_m)$ and $\mathfrak{m} = X + Y$. We claim that the ideal $J = (X^2 + Y^2)^u$ satisfies the desired conditions. It is clear that J is a monomial ideal generated in degree $2u = s$. In order to show that $A = S/J$ is level of socle degree s , we first set up some notation: we let $\mathbf{x} = x_1 \cdots x_j$, and given $\mathbf{a} = (a_1, \dots, a_j) \in \mathbb{N}^j$ and we set $\mathbf{x}^{\mathbf{a}} = x_1^{a_1} \cdots x_j^{a_j}$. Similarly we let $\mathbf{y} = x_{j+1} \cdots x_m$, and given $\mathbf{b} = (b_{j+1}, \dots, b_m) \in \mathbb{N}^{m-j}$ we set $\mathbf{y}^{\mathbf{b}} = x_{j+1}^{b_{j+1}} \cdots x_m^{b_m}$. Note that S is bigraded by setting $\deg(x_i) = (1, 0)$ for all $x_i \in X$, and $\deg(x_i) = (0, 1)$ for all $x_i \in Y$. In this way, if we let $a = |\mathbf{a}| = \sum_{i=1}^j a_i$ and $b = |\mathbf{b}| = \sum_{i=j+1}^m b_i$, then the monomial $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}$ is bigraded of degree (a, b) . Observe the generators of J are precisely the monomials $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}$ of degree $a + b = s$ and such that both a and b are even.

To show that A is level with $\text{socdeg}(A) = t$ we first prove that $\mathfrak{m}^{t+1} \subseteq J$. Let $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}$ be a monomial of degree $a + b \geq t + 1 = 2u + 1$. It is easy to see that there exist vectors $\mathbf{a}' \leq \mathbf{a}$ and $\mathbf{b}' \leq \mathbf{b}$ (where the inequality is intended point-wise) such that $|\mathbf{a}'| = 2h$ and $|\mathbf{b}'| = 2(u - h)$ for some $h \in \mathbb{N}$. It follows that $\mathbf{x}^{\mathbf{a}} \in (\mathbf{x}^{\mathbf{a}'}) \subseteq X^{2h} = (X^2)^h$ and $\mathbf{y}^{\mathbf{b}} \in (\mathbf{y}^{\mathbf{b}'}) \subseteq Y^{2(u-h)} = (Y^2)^{u-h}$. Therefore, we conclude that $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}} \in (X^2)^h(Y^2)^{u-h} \subseteq J$. This shows that $\text{socdeg}(A) \leq t$. Now we show that $\text{socdeg}(A) \geq t$. Since J is a monomial ideal, so is $J : \mathfrak{m}$. Therefore, by degree considerations we only have to show that no monomial of degree $t - 1 = 2u - 1$ belongs to $J : \mathfrak{m}$. But if $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}$ has degree $a + b = t - 1$, then either a is odd and b is even or viceversa. Assume without loss of generality that we are in the first case; if by contradiction we assume that $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}} \in J : \mathfrak{m}$, then $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}} \cdot x_{j+1} = \mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}'}$ where $\mathbf{b}' = \mathbf{b} + (1, 0, \dots, 0)$. By degree considerations, $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}'}$ must then be one of the minimal generators of J ; however, this contradicts our previous characterization of such generators, since a is odd. This concludes the proof in the case s is even.

We now focus on the case in which $t = 2u + 1$ is odd, with $u \geq 1$. Given $m \geq 3$, we let j, k be integers such that $1 \leq j < k \leq m$, and we let $X = (x_1, \dots, x_j)$, $Y = (x_{j+1}, \dots, x_k)$, and $Z = (x_{k+1}, \dots, x_m)$. We also let $\mathfrak{m} = X + Y + Z$. We claim that the ideal $J = X(X^2 + Y^2)^u + Y(Y^2 + Z^2)^u + Z(Z^2 + X^2)^u + XYZ((X + Y)^2 + Z^2)^{u-1}$ satisfies the desired conditions. It is again clear that J is a monomial ideal generated in degree t .

To show that A is level of socle degree t , we consider a trigrading on S by setting $\deg(x_i) = (1, 0, 0)$ for $x_i \in X$, $\deg(x_i) = (0, 1, 0)$ for $x_i \in Y$, and $\deg(x_i) = (0, 0, 1)$ for $x_i \in Z$. We let $\mathbf{x} = x_1 \cdots x_j$, $\mathbf{y} = x_{j+1} \cdots x_k$ and $\mathbf{z} = x_{k+1} \cdots x_m$. Given vectors \mathbf{a} , \mathbf{b} and \mathbf{c} of appropriate length and with non-negative entries, we set $\mathbf{x}^{\mathbf{a}}$, $\mathbf{y}^{\mathbf{b}}$ and $\mathbf{z}^{\mathbf{c}}$ in analogy with the previous case.

Observe that the minimal monomial generators of J are all the monomials $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}\mathbf{z}^{\mathbf{c}}$ of degree $a + b + c = 2u + 1$ which satisfy one of the following four conditions:

- (1) $c = 0$, a is odd and b is even.

- (2) $a = 0$, b is odd and c is even.
- (3) $b = 0$, c is odd and a is even.
- (4) $abc \neq 0$, $a + b$ is even and c is odd.

We start by showing that $\mathfrak{m}^{t+1} \subseteq J$. Assume that $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}\mathbf{z}^{\mathbf{c}}$ has degree $a + b + c \geq t + 1 = 2(u + 1)$; we distinguish some cases: if $abc \neq 0$, then we can find vectors $\mathbf{a}' \leq \mathbf{a}$, $\mathbf{b}' \leq \mathbf{b}$ and $\mathbf{c}' \leq \mathbf{c}$ with $|\mathbf{a}'| + |\mathbf{b}'| + |\mathbf{c}'| = 2u - 1$ and such that $\mathbf{x}^{\mathbf{a}'}\mathbf{y}^{\mathbf{b}'}\mathbf{z}^{\mathbf{c}'}$ is a minimal generator of J . We have already shown in the even case that $\mathfrak{m}^{2u-1} \subseteq ((X+Y)^2 + Z^2)^{u-1}$, and in particular we have that $\mathbf{x}^{\mathbf{a}'}\mathbf{y}^{\mathbf{b}'}\mathbf{z}^{\mathbf{c}'}$ is a minimal generator of J . It follows that $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}\mathbf{z}^{\mathbf{c}} \in J$ in this case. Now assume that $abc = 0$. If $a = 0$, then since $b + c \geq 2(u + 1)$ we can find $\mathbf{b}' \leq \mathbf{b}$ and $\mathbf{c}' \leq \mathbf{c}$ such that $|\mathbf{b}'|$ is odd, $|\mathbf{c}'|$ is even, and $|\mathbf{b}'| + |\mathbf{c}'| = 2u + 1$. It follows that $\mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}'}\mathbf{z}^{\mathbf{c}'} \in (Y^2 + Z^2)^u \subseteq J$. The remaining cases in which either $b = 0$ or $c = 0$ are handled similarly. This concludes the proof that $\text{socdeg}(A) \leq t$.

To see that equality holds, as above it suffices to show that no monomial $w = \mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}\mathbf{z}^{\mathbf{c}}$ of degree $a + b + c = t - 1 = 2u$ belongs to $J : \mathfrak{m}$. Assume by way of contradiction that such a monomial w belongs to $J : \mathfrak{m}$. If two of the exponents a, b, c are equal to zero, for instance if $a = 2u$ and $b = c = 0$, then observe that $w \cdot x_{j+1} \in J$ must be a minimal monomial generator of J , by degree considerations. But $w \cdot x_{j+1} = \mathbf{x}^{\mathbf{a}} \cdot x_{j+1} = \mathbf{x}^{\mathbf{a}'}\mathbf{y}^{\mathbf{b}'}\mathbf{z}^{\mathbf{c}'}$ with $\mathbf{b}' = (1, 0, \dots, 0)$ does not belong to J , thanks to the above characterization of the minimal generators of J . This contradicts our choice of $w \in J : \mathfrak{m}$. The cases in which $a = b = 0$ or $a = c = 0$ are handled similarly. Now assume that only one among a, b, c is equal to zero. For instance, say $a = 0$ and $b + c = 2u$. If b is even, then we consider $w \cdot x_{k+1} = \mathbf{x}^{\mathbf{0}}\mathbf{y}^{\mathbf{b}'}\mathbf{z}^{\mathbf{c}'}$ with $\mathbf{c}' = \mathbf{c} + (1, 0, \dots, 0)$, which must be a minimal generator of J , necessarily of type (2). However b is even, again a contradiction. If b is odd, then we consider $w \cdot x_{j+1} = \mathbf{x}^{\mathbf{0}}\mathbf{y}^{\mathbf{b}'}\mathbf{z}^{\mathbf{c}'}$ with $|\mathbf{b}'| = b + 1$ odd, and we reach again a similar contradiction. The other cases in which only one of the three exponents is equal to zero are tackled in a similar fashion.

Finally, assume that $abc \neq 0$. If c is odd, then $w \cdot x_{k+1} = \mathbf{x}^{\mathbf{a}}\mathbf{y}^{\mathbf{b}}\mathbf{z}^{\mathbf{c}'}$ with $|\mathbf{c}'| = c + 1$ must be a minimal generator of J , necessarily of type (4); but $c + 1$ is even, a contradiction. If c is even, then so is $a + b = 2u - c$. Then $w \cdot x_1 = \mathbf{x}^{\mathbf{a}'}\mathbf{y}^{\mathbf{b}}\mathbf{z}^{\mathbf{c}}$ with $|\mathbf{a}'| = a + 1$ must be a minimal generator of J of type (4); however $|\mathbf{a}'| + |\mathbf{b}| = a + 1 + b$ is odd, a contradiction. \square

Theorem 4.3. *For all odd integers $s = 2t - 1 \geq 3$ and all $n \geq 4$ there exists an Artinian compressed Gorenstein algebra $A = K[x_1, \dots, x_n]/I$ with socle degree s and such that $m(I) = t$.*

Proof. From Lemma 4.2 we can construct a monomial ideal $J \subseteq S = K[x_1, \dots, x_{n-1}]$ with $m(J) = t$ and $\text{soc}(S/J)$ concentrated only in degree t . From the construction by Hartshorne, there is a reduced set of points X in \mathbb{P}^{n-1} such that the Artinian reduction $K[x_1, \dots, x_n]/(I_X + (x_n))$ is isomorphic to S/J . Now, the canonical module ω_X can be embedded as an ideal in $R_X = K[x_1, \dots, x_n]/I_X$. This can be done in all degrees i where $\dim_k((R_X)_i) = \deg(X)$ holds [22]. Since t is the socle degree of the Artinian reduction S/J of R_X we have the equality $\dim_k((R_X)_t) = \deg(X)$, and therefore such an embedding can be obtained in degree t . Let \mathfrak{a} be such a canonical ideal. If we let $I \subseteq R = K[x_1, \dots, x_n]$ be the ideal such that $I/I_X = \mathfrak{a}$, then since both $\mathfrak{a} \subseteq R_X$ and $I_X \subseteq R$ are generated in degree t we conclude that I is generated in degree t as well. Thus $A = R_X/\mathfrak{a} \cong R/I$ is an Artinian Gorenstein K -algebra with defining ideal I generated only in degree t . Note that $\text{socdeg}(A) = 2t - 1$. In fact, ω_X has regularity t , and since \mathfrak{a} is a canonical ideal generated in degree t we have that $\dim_k(\mathfrak{a}_i) = \deg(X)$ if and only if $i \geq 2t$. It follows that $A_i = 0$ for all $i \geq 2t$, while $A_{2t-1} \neq 0$. Finally, since I has no generators in degree less than t , we conclude that A is compressed. \square

Remark 4.4. The construction of Theorem 4.3 also works for $n = 3$ and s an even integer. However, we point out that for $s = 2t - 1$ with $t \geq 3$ odd there is no Artinian Gorenstein compressed algebra $K[x_1, x_2, x_3]/I$ generated only in degree t . In fact, if such an algebra existed, the ideal I would have an even number of generators, contradicting the structure theorem of Buchsbaum and Eisenbud for ideals of codimension three defining Gorenstein rings [9], Corollary 2.2. For this reason we assume $n \geq 4$.

As a consequence of the above results, we may conclude that Boij's conjecture holds true for the first Betti number. Since generic Artinian Gorenstein graded K -algebras are compressed, it is enough to observe that, by Theorem 4.3, the Zariski open set parameterizing the Artinian compressed Gorenstein algebras generated in one degree is non-empty.

Corollary 4.5. *A generic Artinian Gorenstein K -algebra A with $\text{edim}(A) \geq 4$ and socle degree $s \geq 3$ is generated in degree $\lfloor \frac{s}{2} \rfloor + 1$.*

4.2. Rate of of generic Artinian Gorenstein algebras. Thanks to Corollary 4.5 we have the following result.

Theorem 4.6. *Let A be a generic Artinian Gorenstein graded K -algebra of socle degree $s \geq 3$, and assume that $\text{edim}(A) \geq 4$. Then $\text{rate}(A) = \lfloor \frac{s}{2} \rfloor$.*

Proof. Since generic Artinian Gorenstein graded K -algebras are compressed, by Proposition 3.5 we have that the generic Artinian Gorenstein K -algebra A satisfies $\text{rate}(A) \leq m(I) - 1$, with $A = R/I$ a minimal presentation. On the other hand, the reverse inequality always holds, and therefore $\text{rate}(A) = m(I) - 1$. Finally, by Theorem 4.3, $m(I) = \lfloor \frac{s}{2} \rfloor + 1$, and the result follows. \square

Remark 4.7. Artinian compressed Gorenstein K -algebras of even socle degree $s = 2(t - 1)$ have almost linear resolution and, in particular, they are generated in degree t . Hence the conclusion of Theorem 4.6 holds for any Artinian Gorenstein compressed K -algebra of even socle degree, not necessarily generic.

Remark 4.8. When $A = R/I$ is a generic Artinian Gorenstein K -algebra of odd socle degree $s = 2t - 1$ with $\text{edim}(A) = 3$ we have that

$$\text{rate}(A) = \begin{cases} t - 1 & \text{if } t \text{ is even} \\ t & \text{if } t \text{ is odd} \end{cases}.$$

In fact, if $s = 3$ then A is Koszul by [13]. If $s \geq 5$ then in the first case I is generated in degree t , while in the second there is one generator in degree $t + 1$. In both cases we conclude by Proposition 3.5 since A is compressed.

REFERENCES

- [1] D. J. Anick, *On the homology of associative algebras*, Trans. Amer. Math. Soc. **296** (1986), no. 2, 641–659. MR846601
- [2] A. Aramova, Ş. Bărcănescu, and J. Herzog, *On the rate of relative Veronese submodules*, Rev. Roumaine Math. Pures Appl. **40** (1995), no. 3-4, 243–251. MR1409310
- [3] L. L. Avramov, A. Conca, and S. B. Iyengar, *Free resolutions over commutative Koszul algebras*, Math. Res. Lett. **17** (2010), no. 2, 197–210. MR2644369
- [4] ———, *Subadditivity of syzygies of Koszul algebras*, Math. Ann. **361** (2015), no. 1-2, 511–534. MR3302628

- [5] J. Backelin, *On the rates of growth of the homologies of Veronese subrings*, Algebra, algebraic topology and their interactions (Stockholm, 1983), 1986, pp. 79–100. MR846440
- [6] M. Boij, *Betti numbers of compressed level algebras*, J. Pure Appl. Algebra **134** (1999), no. 2, 111–131. MR1663785
- [7] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1993. MR1251956
- [8] W. Bruns, J. Herzog, and U. Vetter, *Syzygies and walks*, Commutative algebra (Trieste, 1992), 1994, pp. 36–57. MR1421076
- [9] D. A. Buchsbaum and D. Eisenbud, *Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension 3*, Amer. J. Math. **99** (1977), no. 3, 447–485. MR453723
- [10] A. Conca, *Koszul algebras*, Boll. Unione Mat. Ital. (9) **1** (2008), no. 2, 429–437. MR2424302
- [11] A. Conca, E. De Negri, and M. E. Rossi, *On the rate of points in projective spaces*, Israel J. Math. **124** (2001), 253–265. MR1856518
- [12] ———, *Koszul algebras and regularity*, Commutative algebra, 2013, pp. 285–315. MR3051376
- [13] A. Conca, M. E. Rossi, and G. Valla, *Gröbner flags and Gorenstein algebras*, Compositio Math. **129** (2001), no. 1, 95–121. MR1856025
- [14] W. A. da Silva, H. Hassanzadeh, and A. Simis, *Bounds for the degree and Betti sequences along a graded resolution*, Preprint available on the arXiv:2201.09994 (2022).
- [15] D. Eisenbud, A. Reeves, and B. Totaro, *Initial ideals, Veronese subrings, and rates of algebras*, Adv. Math. **109** (1994), no. 2, 168–187. MR1304751
- [16] R. Fröberg and D. Laksov, *Compressed algebras*, Complete intersections (Acireale, 1983), 1984, pp. 121–151. MR775880
- [17] V. Gasharov, I. Peeva, and V. Welker, *Rationality for generic toric rings*, Math. Z. **233** (2000), no. 1, 93–102. MR1738353
- [18] A. Iarrobino and J. Emsalem, *Some zero-dimensional generic singularities; finite algebras having small tangent space*, Compositio Math. **36** (1978), no. 2, 145–188. MR515043
- [19] A. Iarrobino, *Compressed algebras: Artin algebras having given socle degrees and maximal length*, Trans. Amer. Math. Soc. **285** (1984), no. 1, 337–378. MR748843
- [20] A. Iarrobino and V. Kanev, *Power sums, Gorenstein algebras, and determinantal loci*, Lecture Notes in Mathematics, vol. 1721, Springer-Verlag, Berlin, 1999. Appendix C by Iarrobino and Steven L. Kleiman. MR1735271
- [21] G. R. Kempf, *Syzygies for points in projective space*, J. Algebra **145** (1992), no. 1, 219–223. MR1144669
- [22] M. Kreuzer, *On the canonical ideal of a set of points*, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) **3** (2000), no. 1, 221–261. MR1755711
- [23] M. Kunte, *Gorenstein modules of finite length*, Math. Nachr. **284** (2011), no. 7, 899–919. MR2815960
- [24] G. L. Levin and L. L. Avramov, *Factoring out the socle of a Gorenstein ring*, J. Algebra **55** (1978), no. 1, 74–83. MR515760
- [25] M. Mastroeni, H. Schenck, and M. Stillman, *Quadratic Gorenstein rings and the Koszul property I*, Trans. Amer. Math. Soc. **374** (2021), no. 2, 1077–1093. MR4196387
- [26] ———, *Quadratic Gorenstein Rings and the Koszul Property II*, Int. Math. Res. Not. IMRN **2** (2023), 1461–1482. MR4537330
- [27] J. McCullough and A. Seceleanu, *Quadratic Gorenstein algebras with many surprising properties*, Arch. Math. (Basel) **115** (2020), no. 5, 509–521. MR4154566
- [28] M. E. Rossi and L. M. Şega, *Poincaré series of modules over compressed Gorenstein local rings*, Adv. Math. **259** (2014), 421–447. MR3197663
- [29] G. Scheja, *Über die Bettizahlen lokaler Ringe*, Math. Ann. **155** (1964), 155–172. MR162819
- [30] J. Tate, *Homology of Noetherian rings and local rings*, Illinois J. Math. **1** (1957), 14–27. MR86072

DEPARTMENT OF MATHEMATICS, KTH - ROYAL INSTITUTE OF TECHNOLOGY, SE-100 44 STOCKHOLM, SWEDEN

Email address: boij@kth.se

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, VIA DODECANESO 35, 16146 GENOVA, ITALY

Email address: denegri@dim.unige.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, VIA DODECANESO 35, 16146 GENOVA, ITALY

Email address: destefani@dim.unige.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, VIA DODECANESO 35, 16146 GENOVA, ITALY

Email address: rossim@dim.unige.it