ON THE THIRD AND FOURTH BETTI NUMBERS OF A HOMOGENEOUS SPACE OF A LIE GROUP

INDRANIL BISWAS, PRALAY CHATTERJEE, AND CHANDAN MAITY

ABSTRACT. In [\[BCM\]](#page-41-0), computable explicit descriptions of the second and first cohomology groups of a general homogeneous space of a Lie group are given, extending an earlier result in [\[BC\]](#page-41-1). They turned out to be crucial in our computations related to the cohomology of nilpotent orbits as done in [\[BC\]](#page-41-1) and [\[BCM\]](#page-41-0). In this paper, we describe the third and fourth cohomologies of a general homogeneous space along the line mentioned above. We also draw various consequences of our main results in numerous special settings. In particular, we obtain an interesting invariant by showing that for a large class of homogeneous spaces, the difference between the third and fourth Betti numbers coincides with the difference in the numbers of simple factors of the ambient group and the associated closed subgroup.

CONTENTS

1. Introduction

1.1. Context and motivation. Let L be a connected Lie group and $M \subset L$ a closed subgroup with finitely many connected components. Let G and H be maximal compact subgroups of L and M respectively, with $H \subset G$. Then using a minor variant of a result due to Mostow (see Theorem [4.12\)](#page-27-0) one concludes that the compact homogeneous space G/H embeds in the homogeneous space L/M as a deformation retraction. Thus finding the topological invariants, such as cohomology of a homogeneous space, which is the coset space of a closed subgroup with finitely many connected components in a general connected Lie group, boils down to the set-up where the Lie groups involved are all compact. If G is a connected compact Lie group with $H \subset G$ a closed subgroup, and

²⁰²⁰ Mathematics Subject Classification. 57T15.

Key words and phrases. Homogeneous space, de Rham cohomology, Koszul complex, homotopy invariant.

if g and h are the respective Lie algebras, then it follows that the cohomology of G/H is the relative Lie algebra cohomology of the chain complex of forms on $\mathfrak g$ which vanish on h and which are invariant under the natural action of H . When H is connected, the Cartan's Theorem (see $\lbrack Ca, Theorem 5\rbrack$, $\lbrack Bo, Theorem 25.2\rbrack$ gives an alternative description of the cohomology of the above cochain complex in terms of the cohomology of a certain associated Koszul complex (defined in Section [2.2.4\)](#page-9-0). See [\[Bo\]](#page-41-4), [\[Ba\]](#page-41-5), [\[Wo\]](#page-42-0), and, more recently, [\[Fr\]](#page-42-1) for other viewpoints on Cartan's theorem, many general results, and applications in various special settings. We also refer to [\[GS\]](#page-42-2) and [\[Tu\]](#page-42-3) for more expositions on the Cartan's theorem and its ramifications.

While the above results provide a foundational basis, they do not seem to provide computational means which can be used in finding the cohomologies of homogeneous spaces in a broad context. In [\[BCM,](#page-41-0) Theorem 3.3, Theorem 3.6] a simple and computable description of the second and first cohomologies of general compact connected homogeneous spaces is obtained, extending an earlier result in [\[BC,](#page-41-1) Theorem 3.3]. Such descriptions turned out to be extremely useful in [\[BC,](#page-41-1) [BCM\]](#page-41-0), where second cohomology groups of nilpotent orbits are computed for classical Lie algebras. See also [\[CM\]](#page-41-6) for some more applications on the second cohomology of nilpotent orbits in exceptional Lie algebras along the same line.

Surprisingly enough, apart from the results in [\[BC,](#page-41-1) [BCM\]](#page-41-0) on the first and second cohomologies, there seems to be hardly any literature describing cohomologies in explicit ways, despite the availability of the above remarkable Cartan's theorem. Thus from computational point of view, it would be desirable to obtain simple descriptions of the cohomology groups, as in $[BC, Theorem 3.3]$ and $[BCM, Theorem 3.3]$ which should be based entirely on Lie theoretic data such as the Lie algebras of the associated groups, how the closed subgroup or the corresponding subalgebra is positioned in the ambient Lie group or the Lie algebra, and the component group of the closed subgroup. Towards this, we obtain explicit formulas for the lower dimensional cohomologies, namely, the third and the fourth ones; see Theorem [1.1](#page-1-0) and Theorem [1.2](#page-2-0) appearing below. As the application of our results, we also obtain several interesting consequences in special cases; see § [1.3,](#page-2-1) § [4.1](#page-22-1) and § [4.2.](#page-27-1) This includes a new way of computing a certain topological invariant when the ambient compact group and the closed subgroup are semisimple; see Corollary [1.6.](#page-3-0)

1.2. Description of the main results. We now describe our main results and some of their corollaries.

In this paper by a semisimple (respectively, simple) Lie group we mean a connected non-trivial Lie group with semisimple (respectively, simple) Lie algebra. Our first result describes the third cohomology of a connected compact homogeneous space. We refer to Definition [3.11](#page-15-0) for the notion $\mathcal{N}_{q,b}$, and § [2.1](#page-4-1) for other relevant notation.

Theorem 1.1. Let G be a compact connected Lie group, and H a closed subgroup of G. Then there is an isomorphism:

$$
H^3(G/H,\,\mathbb{R})\,\simeq\,\big((\mathfrak{z}(\mathfrak{h})\cap [\mathfrak{g},\,\mathfrak{g}])^*\big)^{H/H^0}\otimes \Big(\frac{\mathfrak{g}}{[\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h}}\Big)^*\,\bigoplus\,\mathcal{N}_{\mathfrak{g},\mathfrak{h}}\,\bigoplus\,\wedge^3\Big(\frac{\mathfrak{g}}{[\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h}}\Big)^*\,.
$$

Our second result is on the fourth cohomology of a connected compact homogeneous space. See Definition [3.11](#page-15-0) for $C_{\mathfrak{g},H}$.

Theorem 1.2. Let G be a compact connected Lie group, and H a closed subgroup of G. Let $r_0 := \dim_{\mathbb{R}} \left(\frac{\mathfrak{g}}{[\mathfrak{a}, \mathfrak{a}]} \right)$ $\frac{\mathfrak{g}}{[\mathfrak{g},\mathfrak{g}]+\mathfrak{h}}$). Then there is an isomorphism:

$$
H^{4}(G/H, \mathbb{R}) \simeq ((\mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}])^{*})^{H/H^{0}} \otimes \wedge^{2} \Big(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\Big)^{*} \bigoplus (\mathcal{N}_{\mathfrak{g}, \mathfrak{h}})^{r_{0}}
$$

$$
\bigoplus \mathcal{C}_{\mathfrak{g}, H} \bigoplus \wedge^{4} \Big(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\Big)^{*}.
$$

It should be mentioned that the notions $\mathcal{N}_{\mathfrak{g},\mathfrak{h}}$ and $\mathcal{C}_{\mathfrak{g},H}$ (see Definition [3.11\)](#page-15-0) depend directly on how H acts on the space $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}]$ and on how certain bilinear forms on \mathfrak{g} , constructed using the Killing forms on the simple factors of the semisimple part $[g, g]$, behave when restricted to the subalgebra $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}]$. Thus all the quantities involved in the above two results are explicit, computable, and in fact, depending on the following: the Lie algebra $\mathfrak g$, how H (respectively $\mathfrak h$) is positioned in G (respectively $\mathfrak g$), and the action of the component group H/H^0 .

1.3. Applications of our results and an interesting new interpretation of a topological invariant. We next mention several applications of our main theorems to compact homogeneous spaces; see also $\S 4.1$. In $\S 4.2$ $\S 4.2$ we derive consequences in the set-up of homogeneous spaces which are not-necessarily compact. In specific contexts, we obtain more precise information on the Betti numbers which are interesting on their own right. The two important special cases, namely when G is semisimple or when H^0 is semisimple, give stronger results in Corollary [1.3](#page-2-2) and Corollary [1.4.](#page-3-1) We also obtain Corollary [1.6,](#page-3-0) which follows immediately from Corollary [1.3](#page-2-2) and gives rise to an elegant way of computing a topological invariant associated to a large class of compact homogeneous spaces.

For a semisimple Lie algebra \mathfrak{k} , let $\#\mathfrak{k}$ denote the number of simple factors in \mathfrak{k} . See Definition [4.1](#page-22-2) for the notation $\#([\mathfrak{h}, \mathfrak{h}], H)$, and [\(2.1\)](#page-4-2), [\(2.3\)](#page-5-0) for $S^2(\mathfrak{z}(\mathfrak{h})^*)^{H/H^0}$. As above, we refer to Definition [3.11](#page-15-0) for the notions $\mathcal{N}_{\mathfrak{g},\mathfrak{h}}$ and $\mathcal{C}_{\mathfrak{g},H}$.

Corollary 1.3. Let G be a compact semisimple Lie group and H a closed subgroup of G such that dim $H > 0$. Then the following hold:

- (1) $H^3(G/H, \mathbb{R}) \simeq \mathcal{N}_{\mathfrak{g},\mathfrak{h}}$. Thus, $\dim_{\mathbb{R}} H^3(G/H, \mathbb{R}) \leq \# \mathfrak{g} - 1$.
	- (a) Moreover assume $\mathfrak{g}_1,\ldots,\mathfrak{g}_s$ are distinct simple ideals of \mathfrak{g} such that $\mathfrak{h}\subset\mathfrak{g}_1\oplus\mathfrak{g}_2$ $\cdots \oplus \mathfrak{g}_s$ and that $\mathfrak{h} \cap \mathfrak{g}_i \neq 0$ for all $i = 1, \ldots, s$. Then $\dim_{\mathbb{R}} H^3(G/H, \mathbb{R}) =$ $\#\mathfrak{a}-s.$
	- (b) In particular if G is simple, then $H^3(G/H, \mathbb{R}) = 0$.
- (2) $H^4(G/H, \mathbb{R}) \simeq C_{\mathfrak{g},H}$. Thus, $\dim_{\mathbb{R}} H^4(G/H, \mathbb{R}) \leq \dim_{\mathbb{R}} S^2(\mathfrak{z}(\mathfrak{h})^*)^{H/H^0} + \#([\mathfrak{h}, \mathfrak{h}], H)$ −1. In particular, for specific G and H we have the following bound.
	- (a) If H^0 is semisimple, then $\dim_{\mathbb{R}} H^4(G/H, \mathbb{R}) \leq #(\mathfrak{h}, H) 1$.
	- (b) If G is simple and H⁰ is semisimple, then $\dim_{\mathbb{R}} H^4(G/H, \mathbb{R}) = \#(\mathfrak{h}, H)$ -1.
	- (c) If G is semisimple and H^0 is simple, then $\dim_{\mathbb{R}} H^4(G/H, \mathbb{R}) = 0$.
- (3) The equality,

$$
\dim_{\mathbb{R}} H^{4}(G/H, \mathbb{R}) - \dim_{\mathbb{R}} H^{3}(G/H, \mathbb{R}) = \dim_{\mathbb{R}} S^{2}(\mathfrak{z}(\mathfrak{h})^{*})^{H/H^{0}} + \#([\mathfrak{h}, \mathfrak{h}], H) - (\#\mathfrak{g})
$$

holds.

(4) If moreover H^0 is semisimple, the equality

$$
\dim_{\mathbb{R}} H^4(G/H, \mathbb{R}) - \dim_{\mathbb{R}} H^3(G/H, \mathbb{R}) = \#(\mathfrak{h}, H) - \# \mathfrak{g}
$$

holds. Furthermore, if either H is connected or all the simple factors of H are pairwise non-isomorphic then we have

$$
\dim_{\mathbb{R}} H^4(G/H, \mathbb{R}) - \dim_{\mathbb{R}} H^3(G/H, \mathbb{R}) = \# \mathfrak{h} - \# \mathfrak{g}.
$$

(5) If moreover H is a toral subgroup, then

$$
\dim_{\mathbb{R}} H^4(G/H,\,\mathbb{R})\, - \dim_{\mathbb{R}} H^3(G/H,\,\mathbb{R})\,=\, \frac{\dim_{\mathbb{R}} \mathfrak{h}(\dim_{\mathbb{R}} \mathfrak{h} + 1)}{2} \,-\,(\#\mathfrak{g})\,.
$$

The Corollary $1.3(1b)$ $1.3(1b)$ was proved earlier in $[Az]$.

In the next result, we specialize to the situation where H^0 is assumed to be semisimple.

Corollary 1.4. Let G be a compact, connected Lie group $H \subset G$ be a closed subgroup such that H^0 is a semisimple Lie group. Let $l := \dim_{\mathbb{R}} \mathfrak{z}(\mathfrak{g})$, and $r := \#[\mathfrak{g}, \mathfrak{g}]$. Let $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_r$, where \mathfrak{g}_i 's are simple ideals in $[\mathfrak{g}, \mathfrak{g}]$. Moreover assume that $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$ for some $s \leq \#[\mathfrak{g}, \mathfrak{g}]$ and that $\mathfrak{h} \cap \mathfrak{g}_i \neq 0$ for all $i = 1, \ldots, s$. Then

(1) dim<sub>$$
\mathbb{R}
$$</sub>(H³(G/H, \mathbb{R})) = r - s + $\binom{l}{3}$,
(2) dim _{\mathbb{R}} (H⁴(G/H, \mathbb{R})) = l(r - s) + #(\mathfrak{h} , H) - s + $\binom{l}{4}$.

Remark 1.5. In this remark we will include a vanishing result which follows from [\[BC,](#page-41-1) [BCM\]](#page-41-0), and Corollary [1.3\(](#page-2-2)[1a\)](#page-2-4) and [\(2b\)](#page-2-5) : Let G be a simple Lie group, $H \subset G$ be a closed subgroup such that H^0 is also simple. Then $H^i(G/H,\mathbb{R}) = 0$ for $1 \leq i \leq 4$.

As a consequence of Corollary [1.3](#page-2-2)[\(4\)](#page-3-2), for the class of homogeneous spaces obtained by quotienting a semisimple Lie group by a semisimple subgroup, the difference between the number of simple factors of the ambient group and that of the subgroup gives rise to an interesting topological invariant.

Corollary 1.6. Let G and G_1 be compact semisimple Lie groups. Let H and H_1 be closed semisimple subgroups of G and G_1 , respectively. Let $\mathfrak{g}, \mathfrak{g}_1, \mathfrak{h}$ and \mathfrak{h}_1 be the Lie algebras of G, G_1, H and H_1 , respectively. Assume that

$$
\# \mathfrak{h}_1 - \#\mathfrak{g}_1 \ \neq \ \#\mathfrak{h} - \#\mathfrak{g} \, .
$$

Then G_1/H_1 is not homotopically equivalent to G/H .

The next result is another interesting consequence of Corollary [1.3](#page-2-2)[\(4\)](#page-3-2).

Corollary 1.7. Let G be a connected compact Lie group, and let H be a closed subgroup which is not necessarily connected. Assume that G and H^0 are both semisimple with $\dim_{\mathbb{R}} G/H = 7$. Further if either H is connected or all the simple factors of H are pairwise non-isomorphic, then $\#\mathfrak{g} = \#\mathfrak{h}$.

It should be noted that Corollary [1.7](#page-3-3) also follows from the classification of seven dimensional compact homogeneous spaces as in [\[Go,](#page-42-4) Theorem 2].

1.4. Our strategy in proving the main results. In the proofs of our main results, we use a version of Cartan's theorem on cohomologies of homogeneous spaces; see Theorem [6.15](#page-41-8) and Appendix [6](#page-33-0) for details. Although bringing out simple computable expressions of the cohomologies in higher dimensions from the Cartan's theorem seem prohibitively intricate we are able to do it in the third and fourth dimensions using techniques that involve breaking up suitably certain spaces related to the lower dimensional parts of the Koszul complex, as well as an intricate analysis of the action of the component group H/H^0 on certain relevant subalgebras associated to the pair $(\mathfrak{g}, \mathfrak{h})$. An explicit choice of transgression plays a vital role in determining certain differential maps in the Koszul complex; see § [3.2.](#page-13-0)

1.5. Organization of the paper. This paper is organized as follows. In Section [2](#page-4-0) we recall some background needed in the proofs of our main results. Section [3](#page-11-0) is devoted to deriving our main results, namely Theorems [1.1](#page-1-0) and [1.2.](#page-2-0) In Section [3.4,](#page-22-3) we indicate simpler proofs of earlier obtained results on the first and second cohomologies. We obtain numerous applications in Section [4.1](#page-22-1) and Section [4.2.](#page-27-1) As an appendix, in Section [5,](#page-29-0) we clarify certain issues that may occur with regard to the relation between maximal compact subgroups of a semisimple Lie group and those of the simple factors. In Section [5](#page-29-0) we also list the simple Lie groups according to their types. Although it is expected that Theorem [6.14](#page-41-9) and Theorem [6.15](#page-41-8) are known to experts but proofs could not be found in the literature, and hence, for the sake of completeness, proofs are given in the Appendix [6.](#page-33-0)

2. NOTATION AND BACKGROUND

The purpose of this section is to recall certain technical but unavoidable background required in proving our main results.

2.1. Notation. Here we fix some notation that will be used throughout this paper. Subsequently, a few specialized notations are introduced as and when they occur.

Let $\mathcal{P}(r)$ denote the group of permutations of the set $\{1, \ldots, r\}$. Let V be a vector space over $\mathbb R$ and k a positive integer. We denote $V^r := V \oplus \cdots \oplus V$ r -many . Let $S^r(V^*)$

(respectively, $\wedge^r V^*$) denote the space of symmetric r-linear maps (respectively, space of alternating r-linear maps)

$$
f: \underbrace{V \times \cdots \times V}_{r-\text{many}} \longrightarrow \mathbb{R}.
$$
 (2.1)

For $f \in S^k(V^*)$ and $g \in S^l(V^*)$, we define a multiplication as follows (see [\[GHV-II,](#page-42-5) p. 5]):

$$
(f \vee g)(v_1, \ldots, v_{k+l}) := \frac{1}{k! \, l!} \sum_{\sigma \in \mathcal{P}(k+l)} f(v_{\sigma(1)}, \ldots, v_{\sigma(k)}) \, g(v_{\sigma(k+1)}, \ldots, v_{\sigma(k+l)}).
$$

Similarly, for $\omega \in \wedge^k V^*$ and $\delta \in \wedge^l V^*$, we define a multiplication as follows (see [\[GHV-II,](#page-42-5) p. 5]) :

$$
(\omega \wedge \delta)(v_1, \ldots, v_{k+l}) := \frac{1}{k! \, l!} \sum_{\sigma \in \mathcal{P}(k+l)} \text{Sgn } \sigma \, \omega(v_{\sigma(1)}, \ldots, v_{\sigma(k)}) \, \delta(v_{\sigma(k+1)}, \ldots, v_{\sigma(k+l)}).
$$

In the above definition Sgn σ denotes the sign of the permutation $\sigma \in \mathcal{P}(k+l)$.

With respect to the above multiplication " \vee " the direct sum $S(V^*) := \sum_{i \geq 0} S^i(V^*)$ is an algebra isomorphic to the $\mathbb R$ -algebra of polynomials on V. Analogously, with respect to the above multiplication " \wedge " the direct sum $\wedge V^* := \sum_{j\geq 0} \wedge^j V^*$ is an algebra.

Let $E = \sum_i E^i$ be a graded algebra. We denote

$$
E^+ \, := \, \sum_{i>0} E^i \, .
$$

The graded algebra E is called an *anticommutative algebra* if

$$
e^i \cdot e^j = (-1)^{ij} e^j \cdot e^i
$$

for all $e^i \in E^i$, $e^j \in E^j$. For graded algebras E and F, the tensor product $E \otimes F$ is a bigraded algebra with the following anticommutative multiplication rule :

$$
(e \otimes f^{i})(e^{j} \otimes f) := (-1)^{ij}(e \cdot e^{j} \otimes f^{i} \cdot f).
$$
\n(2.2)

We will always use this algebra structure on the tensor product of graded algebras unless mentioned otherwise.

Let Γ be a group acting linearly on a vector space V. The subspace of V fixed pointwise by the action of Γ is denoted by V^{Γ} . If $\Gamma_1 \subset \Gamma$ is a normal subgroup acting trivially on V then Γ/Γ_1 has a natural action on V, and moreover, $V^{\Gamma} = V^{\Gamma/\Gamma_1}$.

The Lie groups will be denoted by the capital letters, while the Lie algebra of a Lie group will be denoted by the corresponding lower case German letter, unless a different notation is explicitly mentioned. Sometimes, for notational convenience, the Lie algebra of a Lie group G is also denoted by $Lie(G)$. The connected component of G containing the identity element is denoted by G^0 . The *center* of a group G is denoted by $Z(G)$, while the *center* of a Lie algebra $\mathfrak g$ is denoted by $\mathfrak z(\mathfrak g)$.

If $\mathfrak k$ is a Lie algebra, W is a R-vector space and $\psi : \mathfrak k \longrightarrow \text{End}_{\mathbb R}W$ is a representation of $\mathfrak k$, then define

$$
W^{\mathfrak{k}} := \{ w \in W \mid \psi(x)w = 0 \text{ for all } x \in \mathfrak{k} \}. \tag{2.3}
$$

For a semisimple Lie algebra $\mathfrak k$ over $\mathbb R$ (respectively, $\mathbb C$), let $\#\mathfrak k$ denotes the *number of* simple factors of $\mathfrak k$ over $\mathbb R$ (respectively, $\mathbb C$).

It should be noted that although various notions from the book [\[GHV-III\]](#page-42-6) are used in Sections [3,](#page-11-0) [6,](#page-33-0) the corresponding notation in [\[GHV-III\]](#page-42-6) differ from that in this paper; for instance, $W^{\mathfrak{k}}$ in [\(2.3\)](#page-5-0) corresponds to $W_{\psi=0}$ in [\[GHV-III\]](#page-42-6).

2.2. Background. Here we recall some background required for the next sections.

2.2.1. Multilinear algebra. Let V, W be vector spaces, and let $T: V \longrightarrow W$ be a linear map. Define another linear map T_i as follows :

$$
T_i: \wedge^i V \longrightarrow T(V) \otimes (\wedge^{i-1} V), \qquad i \geq 2
$$

$$
T_i(v_1 \wedge \cdots \wedge v_i) := \sum_{j=1}^i (-1)^{j-1} T(v_j) \otimes (v_1 \wedge \cdots \wedge \widehat{v}_j \wedge \cdots \wedge v_i).
$$

Let $V_0 := \ker T$. Let $V' \subset V$ be a complementary subspace of V_0 , such that

$$
V = V' \oplus V_0.
$$

Let $n_0 = \dim V_0$ and $n' = \dim V'$. For $1 \le r \le i$, set

$$
A_r := \operatorname{Span} \left\{ \sum_{j=1}^r (-1)^{j-1} T(v_j) \otimes (v_1 \wedge \cdots \wedge \hat{v_j} \wedge \cdots \wedge v_r \wedge \cdots \wedge v_i) \middle| v_l \in \begin{cases} V' & l \leq r \\ V_0 & l > r \end{cases} \right\}.
$$

The following statements are straight-forward to check :

(1) dim
$$
A_r = {n_0 \choose i-r} {n' \choose r}
$$
.
\n(2) Note that $A_r \cap A_l = \{0\}$ for $r \neq l$, and $A_r \subseteq \text{Im } T_i$ for $1 \leq r \leq i$. Hence
\n
$$
\dim \text{Im } T_i \geq \sum_{1 \leq r \leq i} {n_0 \choose i-r} {n' \choose r}.
$$
\n(3) $\wedge^i V_0 \subseteq \text{ker } T_i$. Thus dim ker $T_i \geq {n_0 \choose i}$.
\n(4)
$$
\sum_{0 \leq r \leq i} {n_0 \choose i-r} {n' \choose r} = {n_0 + n' \choose i}.
$$

Lemma 2.1. The following two hold :

$$
(1) \wedge^i V_0 = \ker T_i.
$$

$$
(2) \sum_{1 \le r \le i} A_r = \operatorname{Im} T_i.
$$

Let V, W, Z be three vector spaces such that V, $W \subseteq Z$ and $V \cap W = \{0\}$. Define an operator T on the vector space $V \otimes ((V \oplus W) \wedge (V \oplus W))$ as follows :

$$
T: V \otimes ((V \oplus W) \wedge (V \oplus W)) \longrightarrow V \otimes ((V \oplus W) \wedge (V \oplus W))
$$

\n
$$
v \otimes ((v_1 + w_1) \wedge (v_2 + w_2)) \longmapsto v \otimes ((v_1 + w_1) \wedge (v_2 + w_2)) - v_1 \otimes (v \wedge (v_2 + w_2))
$$

\n
$$
+ v_2 \otimes (v \wedge (v_1 + w_1)).
$$
\n(2.4)

Note that $V \otimes (W \wedge V)$ is T-invariant. Let $T_1 := T|_{V \otimes (W \wedge V)}$. Then $T_1^2 - 2T_1 = 0$. Let

$$
E_2 := \mathrm{Span}\{v \otimes (w \wedge u) - u \otimes (w \wedge v) \mid v, u \in V, w \in W\},\
$$

and $E_0 := (T_1 - 2 \cdot \text{Id})(V \otimes (W \wedge V))$. Then it follows that

$$
E_0 = \mathrm{Span}\{v \otimes (w \wedge u) + u \otimes (w \wedge v) \mid v, u \in V, w \in W\}.
$$

The vector spaces E_2 and E_0 are the eigenspaces of T_1 for the eigenvalues 2 and 0, respectively. Thus, we have the following direct sum decomposition :

$$
V \otimes (W \wedge V) = E_2 \oplus E_0. \tag{2.5}
$$

Next consider another linear operator T_2 on $V \otimes (V \wedge V)$ as follows:

$$
T_2: V \otimes (V \wedge V) \longrightarrow V \otimes (V \wedge V) \tag{2.6}
$$

$$
v_1 \otimes (v_2 \wedge v_3) \longmapsto v_1 \otimes (v_2 \wedge v_3) - v_2 \otimes (v_1 \wedge v_3) + v_3 \otimes (v_1 \wedge v_2).
$$

Note that the vector space $V \otimes (V \wedge V)$ is invariant under the linear map T in [\(2.4\)](#page-6-0). We may consider T_2 as the restriction of T in an obvious way, i.e., $T_2 := T|_{V \otimes (V \wedge V)}$. Note that $T_2^2 = 3T_2$. Let $E'_3 := T_2(V \otimes (V \wedge V))$, and $E'_0 := (T_2 - 3 \cdot \text{Id})(V \otimes (V \wedge V))$. Then E'_{3} and E'_{0} are eigenspaces of T_{2} for the eigenvalues 3 and 0, respectively. Thus we have

$$
V \otimes (V \wedge V) = E'_3 \oplus E'_0. \tag{2.7}
$$

We will use a minor variant of [\(2.5\)](#page-6-1) and [\(2.7\)](#page-7-0). Let $\lambda: V \longrightarrow V'$ be a linear isomor-phism. Following the map T as in [\(2.4\)](#page-6-0), define \widetilde{T} as follows :

$$
\tilde{T}: V' \otimes ((V \oplus W) \wedge (V \oplus W)) \longrightarrow V' \otimes ((V \oplus W) \wedge (V \oplus W))
$$
\n
$$
v' \otimes ((v_1 + w_1) \wedge (v_2 + w_2)) \longmapsto v' \otimes ((v_1 + w_1) \wedge (v_2 + w_2))
$$
\n
$$
-\lambda(v_1) \otimes (\lambda^{-1}(v') \wedge (v_2 + w_2)) + \lambda(v_2) \otimes (\lambda^{-1}(v') \wedge (v_1 + w_1)).
$$
\n(2.8)

Define the following two subspaces E_2 and E_0 of $V' \otimes (W \wedge V)$

$$
\widetilde{E}_2 := \operatorname{Span} \{ v' \otimes (w \wedge v) - \lambda(v) \otimes (w \wedge \lambda^{-1}(v')) \mid v' \in V', v \in V, w \in W \},\
$$

\n
$$
\widetilde{E}_0 := \operatorname{Span} \{ v' \otimes (w \wedge v) + \lambda(v) \otimes (w \wedge \lambda^{-1}(v')) \mid v' \in V', v \in V, w \in W \}.
$$

Similarly, the two subspaces E'_3 and E'_0 of $V' \otimes (V \wedge V)$ are as defined to be

$$
\widetilde{E}'_3 := \operatorname{Span} \left\{ \begin{array}{l} v' \otimes (v_1 \wedge v_2) - \lambda(v_1) \otimes (\lambda^{-1}(v') \wedge v_2) \\ + \lambda(v_2) \otimes (\lambda^{-1}(v') \wedge v_1) \end{array} \middle| v_1, v_2 \in V, v' \in V' \right\},
$$

$$
\widetilde{E}'_0 := \operatorname{Span} \left\{ \begin{array}{l} -2v' \otimes (v_1 \wedge v_2) - \lambda(v_1) \otimes (\lambda^{-1}(v') \wedge v_2) \\ + \lambda(v_2) \otimes (\lambda^{-1}(v') \wedge v_1) \end{array} \middle| v_1, v_2 \in V, v' \in V' \right\}.
$$

Then it follows from (2.5) , and (2.7) that

$$
\lambda(V) \otimes (W \wedge V) = \overline{E}_2 \oplus \overline{E}_0, \qquad (2.9)
$$

$$
\lambda(V) \otimes (V \wedge V) = \widetilde{E}'_3 \oplus \widetilde{E}'_0. \tag{2.10}
$$

2.2.2. Primitive elements. Let $\mathfrak g$ be a reductive Lie algebra. Consider the homomorphism μ_{\wedge} : \wedge g $\otimes \wedge$ g $\longrightarrow \wedge$ g defined by $X \otimes Y \longmapsto X \wedge Y$. The ad-representation of g naturally provides another representation $\theta : \mathfrak{g} \longrightarrow \text{End } \wedge \mathfrak{g}$, and with respect to this representation we may form the subspace $(\wedge \mathfrak{g})^{\mathfrak{g}}$ of $\wedge \mathfrak{g}$. It is clear that $\mu_{\wedge}((\wedge \mathfrak{g})^{\mathfrak{g}} \otimes (\wedge \mathfrak{g})^{\mathfrak{g}}) \subset (\wedge \mathfrak{g})^{\mathfrak{g}}$. Let

$$
(\mu_\wedge)^{\mathfrak{g}}\,:\;(\wedge\mathfrak{g})^{\mathfrak{g}}\otimes(\wedge\mathfrak{g})^{\mathfrak{g}}\,\longrightarrow\,(\wedge\mathfrak{g})^{\mathfrak{g}}
$$

denote the restriction of the map μ_{\wedge} to $({\wedge} \mathfrak{g})^{\mathfrak{g}} \otimes ({\wedge} \mathfrak{g})^{\mathfrak{g}} \subset {\wedge} \mathfrak{g} \otimes {\wedge} \mathfrak{g}$. Let $\gamma_{\mathfrak{g}}:$ $({\wedge} \mathfrak{g}^*)^{\mathfrak{g}} \longrightarrow$ $(\wedge \mathfrak{g}^*)^{\mathfrak{g}} \otimes (\wedge \mathfrak{g}^*)^{\mathfrak{g}}$ be the dual of the homomorphism $(\mu_\wedge)^{\mathfrak{g}}$. An element $\phi \in (\wedge^+\mathfrak{g}^*)^{\mathfrak{g}}$ is called primitive if

$$
\gamma_{\mathfrak{g}}(\phi) = \phi \otimes 1 + 1 \otimes \phi.
$$

The *subspace of primitive elements* will be denoted by $P_{\mathfrak{g}}$. Then $P_{\mathfrak{g}}$ forms a graded subspace $P_{\mathfrak{g}} = \sum_{j} P_{\mathfrak{g}}^{j}$ of $(\wedge^{+} \mathfrak{g}^{*})^{\mathfrak{g}}$.

2.2.3. Transgression and the Cartan map. In what follows we use the notion of the (invariant) Weil algebra $W(\mathfrak{g})^{\mathfrak{g}}$ for which we refer to Definition [6.2.](#page-35-0)

Definition 2.2. A *transgression* in the differential algebra $W(\mathfrak{g})^{\mathfrak{g}}$ is a linear map

$$
\tau:\,P_{\mathfrak{g}}\,\longrightarrow\,S(\mathfrak{g}^*)^{\mathfrak{g}}
$$

satisfying the following two conditions (see [\[GHV-III,](#page-42-6) $\S 6.13$, p. 241]):

- (1) τ is a graded map of degree 1.
- (2) For every $\phi \in P_{\mathfrak{g}}$, there is an element $\omega \in (W(\mathfrak{g})^+)^{\mathfrak{g}}$ such that

$$
\delta_W(\omega) = \tau \phi \otimes 1 \quad \text{and} \quad 1 \otimes \phi - \omega \in \sum_{j \ge 1} (S^j(\mathfrak{g}^*) \otimes \wedge \mathfrak{g}^*)^{\mathfrak{g}}.
$$

We denote by Trans(\mathfrak{g}) the *space of all transgressions* for the Lie algebra \mathfrak{g} .

Definition 2.3. Let $\mathcal{P}(r)$ denote the group of permutations of the set $\{1, \ldots, r\}$. For a Lie algebra g define a linear graded map $\rho_{\mathfrak{g}}: (S(\mathfrak{g}^*)^+)^{\mathfrak{g}} \longrightarrow (\wedge \mathfrak{g}^*)^{\mathfrak{g}}$ of degree -1 (this means that $\rho_{\mathfrak{g}}(S^p(\mathfrak{g}^*)^{\mathfrak{g}}) \subset (\wedge^{2p-1}\mathfrak{g}^*)^{\mathfrak{g}}$ for all $p > 0$) by

$$
(\rho_{\mathfrak{g}}\psi)(X_1,\ldots,X_{2p-1}) = \frac{(-1)^{p-1}(p-1)!}{2^{p-1}(2p-1)!} \sum_{\sigma \in \mathcal{P}(2p-1)} \text{sgn}(\sigma)\psi(X_{\sigma(1)},[X_{\sigma(2)},X_{\sigma(3)}],\ldots,[X_{\sigma(2p-2)},X_{\sigma(2p-1)}]),
$$

for all $\psi \in S^p(\mathfrak{g}^*)^{\mathfrak{g}}, X_i \in \mathfrak{g}$. This $\rho_{\mathfrak{g}}$ is called the *Cartan map* for \mathfrak{g} .

Remark 2.4. The Definition [2.3](#page-8-0) of Cartan map is an equivalent formulation of the one given in $[GHV-III, § 6.7, p. 231]$, and the reason behind our choice is that it demands much less background compared to the latter one in $[GHV-III, § 6.7, p. 231]$.

Lemma 2.5 ([\[GHV-III,](#page-42-6) p. 242, Lemma VII]). Let $\tau: P_{\mathfrak{g}} \longrightarrow S(\mathfrak{g}^*)^{\mathfrak{g}}$ be a linear map. Assume further that τ is graded and of degree 1. Then τ is a transgression if and only if $\rho_\mathfrak{g} \circ \tau \, = \, \mathrm{Id}_{P_\mathfrak{g}}.$

Remark 2.6.

(1) Any transgression τ can be written as $\tau = \bigoplus_i \tau^{2i-1}$, where $\tau^{2i-1} : P_{\mathfrak{g}}^{2i-1} \longrightarrow$ $S^{i}(\mathfrak{g}^*)^{\mathfrak{g}}$ is a linear map satisfying the condition $\rho_{\mathfrak{g}} \circ \tau^{2i-1} = \text{Id}.$

Conversely, suppose that, for each i, there is a linear map τ^{2i-1} : $P_{\mathfrak{g}}^{2i-1}$ \longrightarrow $S^i(\mathfrak{g}^*)^{\mathfrak{g}}$ satisfying $\rho_{\mathfrak{g}} \circ \tau^{2i-1} = \text{Id}$. Let us define $\tau := \bigoplus_i \tau^{2i-1}$. Then τ is a linear graded map satisfying $\rho_{\mathfrak{g}} \circ \tau = \text{Id}$. Hence τ is a transgression.

- (2) As transgressions exist, it now follows from the above remark that, for a given $j \geq 0$, any linear map $\zeta : P_{\mathfrak{g}}^{2j-1} \longrightarrow S^{j}(\mathfrak{g}^*)^{\mathfrak{g}}$ satisfying $\rho_{\mathfrak{g}} \circ \zeta = \text{Id}$ can be extended to a transgression.
- (3) Let $\tau \in \text{Hom}(P_{\mathfrak{g}}, S(\mathfrak{g}^*)^{\mathfrak{g}})$ be any transgression. Then from Lemma [2.5](#page-8-1) it follows that Trans(\mathfrak{g}) is an affine space in Hom($P_{\mathfrak{g}}$, $S(\mathfrak{g}^*)^{\mathfrak{g}}$) obtained by translating the subspace $\text{Hom}(P_{\mathfrak{g}}, \text{ker } \rho_{\mathfrak{g}})$ using τ . More precisely, we have

$$
\text{Trans}(\mathfrak{g}) = \tau + \text{Hom}(P_{\mathfrak{g}}, \ker \rho_{\mathfrak{g}}).
$$

Furthermore, the kernel of $\rho_{\mathfrak{g}}$ is given by (see [\[GHV-III,](#page-42-6) p. 244, Theorem II]) $\ker \rho_{\mathfrak{g}} = \sum_{r,s>0} S^r(\mathfrak{g}^*)^{\mathfrak{g}} \vee S^s(\mathfrak{g}^*)^{\mathfrak{g}}.$

2.2.4. Koszul Complex. Let (g, h) be reductive Lie algebra pair with the inclusion map $j : \mathfrak{h} \longrightarrow \mathfrak{g}$. Let

$$
j^* : \mathfrak{g}^* \longrightarrow \mathfrak{h}^* \quad \text{and} \quad j^* : S(\mathfrak{g}^*) \longrightarrow S(\mathfrak{h}^*)
$$
 (2.11)

denote the dual map from \mathfrak{g}^* to \mathfrak{h}^* , and the corresponding natural extension from the symmetric algebra $S(\mathfrak{g}^*)$ to $S(\mathfrak{h}^*)$, respectively (we continue to denote both the maps by the same notation as long as there are no confusions). Then $j^*(S(\mathfrak{g}^*)^{\mathfrak{g}}) \subset S(\mathfrak{h}^*)^{\mathfrak{h}}$. Let $\tau: P_{\mathfrak{g}} \longrightarrow S(\mathfrak{g}^*)^{\mathfrak{g}}$ be a transgression. Define

$$
\sigma := j^* \circ \tau : P_{\mathfrak{g}} \longrightarrow S(\mathfrak{h}^*)^{\mathfrak{h}}. \tag{2.12}
$$

Consider the Koszul complex $(S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes (\wedge P_{\mathfrak{g}}), \nabla_{\sigma})$ associated to the transgression τ , where the differential ∇_{σ} is given by (see [\[GHV-III,](#page-42-6) § 10.8, p. 420])

$$
\nabla_{\sigma}(\psi \otimes (\phi_0 \wedge \cdots \wedge \phi_p)) := \sum_{j=0}^p (-1)^j (\psi \vee \sigma(\phi_j)) \otimes (\phi_0 \wedge \cdots \wedge \widehat{\phi_j} \wedge \cdots \wedge \phi_p), \quad (2.13)
$$

for all $\psi \in S(\mathfrak{h}^*)^{\mathfrak{h}}, \phi_i \in P_{\mathfrak{g}}$. It is important to keep in mind that the Koszul complex is a graded algebra with respect to the total degree, deg $\psi = 2k$ for all non-zero $\psi \in S^k(\mathfrak{h}^*)^{\mathfrak{h}},$ and the elements of $P_{\rm g}$ have the usual degrees.

It can be shown that $\nabla^2_\sigma = 0$. Thus the Koszul complex $(S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes (\wedge P_{\mathfrak{g}}), \nabla_\sigma)$ is indeed a complex. The differential ∇_{σ} depends on the choice of the transgression τ . However, the cohomology of the complex is independent of the choice of τ .

2.2.5. Initial terms of the Koszul Complex $S(\mathfrak{h}^*)^H \otimes \wedge P_{\mathfrak{g}}$. To compute the de Rham cohomology of the homogeneous space G/H it suffices to consider the co-chain complex $(\mathscr{C}^*, \nabla) := (S(\mathfrak{h}^*)^H \otimes \wedge P_{\mathfrak{g}}, \nabla_{\sigma}).$ We have the following :

•
$$
\mathscr{C}^1 = 1 \otimes P^1_{\mathfrak{g}}
$$
.

•
$$
\mathscr{C}^2 = (\mathfrak{h}^*)^H \otimes 1 \oplus 1 \otimes \wedge^2 P_{\mathfrak{g}}^1
$$
.

- $\mathscr{C}^3 = (\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \oplus 1 \otimes P_{\mathfrak{g}}^3 \oplus 1 \otimes \wedge^3 P_{\mathfrak{g}}^1.$
- $\bullet \ \mathscr{C}^4 \ = \ S^2(\mathfrak{h}^*)^H \otimes 1 \ \bigoplus \ (\mathfrak{h}^*)^H \otimes \wedge^2 P_\mathfrak{g}^1 \ \bigoplus \ 1 \otimes P_\mathfrak{g}^3 \wedge P_\mathfrak{g}^1 \ \bigoplus \ 1 \otimes \wedge^4 P_\mathfrak{g}^1.$
- $\mathscr{C}^5 = S^2(\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus (\mathfrak{h}^*)^H \otimes (\wedge^3 P_{\mathfrak{g}}^1 \oplus P_{\mathfrak{g}}^3) \bigoplus 1 \otimes (\wedge^5 P_{\mathfrak{g}}^1 \oplus (P_{\mathfrak{g}}^3 \wedge P_{\mathfrak{g}}^1 \wedge P_{\mathfrak{g}}^1) \oplus P_{\mathfrak{g}}^5).$

Next consider the differential map ∇^i : $\mathscr{C}^i \longrightarrow \mathscr{C}^{i+1}$ for $i = 1, 2, 3, 4$:

$$
(1) \nabla^{1}: \mathscr{C}^{1} = 1 \otimes P_{\mathfrak{g}}^{1} \longrightarrow (\mathfrak{h}^{*})^{H} \otimes 1 \text{ is given by}
$$
\n
$$
\nabla^{1}(1 \otimes f) := f|_{\mathfrak{h}} \otimes 1, \qquad f \in P_{\mathfrak{g}}^{1}.
$$
\n
$$
(2) \nabla^{2}: (\mathfrak{h}^{*})^{H} \otimes 1 \bigoplus 1 \otimes \wedge^{2} P_{\mathfrak{g}}^{1} \longrightarrow (\mathfrak{h}^{*})^{H} \otimes P_{\mathfrak{g}}^{1} \text{ is given by}
$$
\n
$$
\nabla^{2}(\psi \otimes 1) = 0, \qquad \psi \in (\mathfrak{h}^{*})^{H},
$$
\n
$$
\nabla^{2}(1 \otimes f_{1} \wedge f_{2}) = f_{1}|_{\mathfrak{h}} \otimes f_{2} - f_{2}|_{\mathfrak{h}} \otimes f_{1}, \qquad f_{1}, f_{2} \in P_{\mathfrak{g}}^{1}.
$$
\n
$$
(3) \nabla^{3}: (\mathfrak{h}^{*})^{H} \otimes P_{\mathfrak{g}}^{1} \bigoplus 1 \otimes P_{\mathfrak{g}}^{3} \bigoplus 1 \otimes \wedge^{3} P_{\mathfrak{g}}^{1} \longrightarrow S^{2}(\mathfrak{h}^{*})^{H} \otimes 1 \bigoplus (\mathfrak{h}^{*})^{H} \otimes \wedge^{2} P_{\mathfrak{g}}^{1} \text{ is given by}
$$
\n
$$
\nabla^{3}(\psi \otimes f) = \psi \vee f|_{\mathfrak{h}} \otimes 1, \qquad f \in P_{\mathfrak{g}}^{1}, \qquad \qquad
$$
\n(2.14)

$$
\nabla^3 (1 \otimes \rho(\widetilde{B}_i)) = \widetilde{B}_i|_{\mathfrak{h} \times \mathfrak{h}} \otimes 1, \quad J \in P_{\mathfrak{g}},
$$

$$
\nabla^3 (1 \otimes \rho(\widetilde{B}_i)) = \widetilde{B}_i|_{\mathfrak{h} \times \mathfrak{h}} \otimes 1, \quad \text{see (3.12) and Lemma 3.9.}
$$

$$
\nabla^3 (1 \otimes f_1 \wedge f_2 \wedge f_3) = f_1|_{\mathfrak{h}} \otimes f_2 \wedge f_3 - f_2|_{\mathfrak{h}} \otimes f_1 \wedge f_3 + f_3|_{\mathfrak{h}} \otimes f_1 \wedge f_2, f_i \in P_{\mathfrak{g}}^1.
$$
 (2.15)

(4) The map ∇^4 is given as below:

$$
\nabla^{4}: (S^{2}(\mathfrak{h}^{*})^{H} \otimes 1 \bigoplus (\mathfrak{h}^{*})^{H} \otimes \wedge^{2} P_{\mathfrak{g}}^{1} \bigoplus 1 \otimes (P_{\mathfrak{g}}^{3} \wedge P_{\mathfrak{g}}^{1}) \bigoplus 1 \otimes \wedge^{4} P_{\mathfrak{g}}^{1})
$$

\n
$$
\longrightarrow S^{2}(\mathfrak{h}^{*})^{H} \otimes P_{\mathfrak{g}}^{1} \bigoplus (\mathfrak{h}^{*})^{H} \otimes P_{\mathfrak{g}}^{3} \bigoplus (\mathfrak{h}^{*})^{H} \otimes \wedge^{3} P_{\mathfrak{g}}^{1}
$$

\n
$$
\nabla^{4}(S^{2}(\mathfrak{h}^{*})^{H} \otimes 1) = 0,
$$

\n
$$
\nabla^{4}(\psi \otimes f_{1} \wedge f_{2}) = \psi \vee f_{1}|_{\mathfrak{h}} \otimes f_{2} - \psi \vee f_{2}|_{\mathfrak{h}} \otimes f_{1}, \qquad f_{1}, f_{2} \in P_{\mathfrak{g}}^{1},
$$

\n
$$
\nabla^{4}(1 \otimes \rho(\widetilde{B}_{i}) \wedge f) = \widetilde{B}_{i}|_{\mathfrak{h} \times \mathfrak{h}} \otimes f - f|_{\mathfrak{h}} \otimes \rho(\widetilde{B}_{i}),
$$

\n
$$
\nabla^{4}(1 \otimes f_{1} \wedge f_{2} \wedge f_{3} \wedge f_{4}) = f_{1}|_{\mathfrak{h}} \otimes f_{2} \wedge f_{3} \wedge f_{4} - f_{2}|_{\mathfrak{h}} \otimes f_{1} \wedge f_{3} \wedge f_{4}
$$

\n
$$
+ f_{3}|_{\mathfrak{h}} \otimes f_{1} \wedge f_{2} \wedge f_{4} - f_{4}|_{\mathfrak{h}} \otimes f_{1} \wedge f_{2} \wedge f_{3}, \quad f_{i} \in P_{\mathfrak{g}}^{1}.
$$

\n(2.16)

The following diagram is useful in our context to keep track of the maps:

$$
\mathscr{C}^1 := 1 \otimes P_{\mathfrak{g}}^1
$$

\n
$$
\downarrow \nabla^1
$$

\n
$$
\mathscr{C}^2 := (\mathfrak{h}^*)^H \otimes 1 \bigoplus 1 \otimes \wedge^2 P_{\mathfrak{g}}^1
$$

\n
$$
\downarrow \nabla^2 \swarrow
$$

\n
$$
\mathscr{C}^3 := (\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus 1 \otimes P_{\mathfrak{g}}^3 \bigoplus 1 \otimes \wedge^3 P_{\mathfrak{g}}^1
$$

\n
$$
\downarrow \nabla^3 \swarrow \swarrow \nabla^3
$$

\n
$$
\mathscr{C}^4 := S^2(\mathfrak{h}^*)^H \otimes 1 \bigoplus (\mathfrak{h}^*)^H \otimes \wedge^2 P_{\mathfrak{g}}^1 \bigoplus 1 \otimes (P_{\mathfrak{g}}^3 \wedge P_{\mathfrak{g}}^1) \bigoplus 1 \otimes \wedge^4 P_{\mathfrak{g}}^1
$$

\n
$$
\nabla^4 \downarrow \swarrow \nabla^4 \downarrow \vee \nabla^4 \downarrow \nabla^4 \downarrow
$$

\n
$$
\mathscr{C}^5 \supseteq \qquad 0 \bigoplus S^2(\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus (\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^3 \bigoplus (\mathfrak{h}^*)^H \otimes \wedge^3 P_{\mathfrak{g}}^1
$$

A full description of the vector space \mathscr{C}^5 is not required; we have mentioned those subspaces which are needed for the above maps.

2.2.6. Invariant decomposition of associated a Lie algebra and a subalgebra. Here we list certain decompositions associated to a Lie algebra and a subalgebra which will be used later.

Let G be a compact connected Lie group and H a closed subgroup. As before, let $\mathfrak{g} := \mathrm{Lie}(G)$ and $\mathfrak{h} := \mathrm{Lie}(H)$. Since \mathfrak{g} and \mathfrak{h} are reductive, we have

 $\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus [\mathfrak{g}, \mathfrak{g}], \quad \mathfrak{h} = \mathfrak{z}(\mathfrak{h}) \oplus [\mathfrak{h}, \mathfrak{h}] \quad \text{and} \quad [\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{h} = [\mathfrak{h}, \mathfrak{h}] \oplus (\mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}])$.

Consider a G-invariant inner product $\langle \cdot, \cdot \rangle$ on the R-vector space g. We decompose [g, g], $\mathfrak h$ and $[\mathfrak g, \mathfrak g] + \mathfrak h$ as follows :

$$
\mathfrak{h} = (([\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{h})^{\perp} \cap \mathfrak{h}) \bigoplus ([\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{h})
$$

= (([\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{h})^{\perp} \cap \mathfrak{h}) \bigoplus [\mathfrak{h}, \mathfrak{h}] \bigoplus (\mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}]), \qquad (2.17)

$$
[\mathfrak{g},\mathfrak{g}]+\mathfrak{h}=[\mathfrak{g},\mathfrak{g}]\bigoplus (([\mathfrak{g},\mathfrak{g}]\cap\mathfrak{h})^{\perp}\cap\mathfrak{h})\ .
$$
 (2.18)

For notational convenience, we set

$$
\begin{aligned}\n\mathfrak{a} &:= [\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{z}(\mathfrak{h}), \\
\mathfrak{b} &:= ([\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{h})^{\perp} \cap \mathfrak{h}.\n\end{aligned}
$$
\n(2.19)

It may be observed that

$$
\mathfrak{b} = (([\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{z}(\mathfrak{h})) + [\mathfrak{h}, \mathfrak{h}])^{\perp} \cap \mathfrak{h} = ([\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{z}(\mathfrak{h}))^{\perp} \cap \mathfrak{z}(\mathfrak{h}).
$$

Then we have

$$
\mathfrak{z}(\mathfrak{h}) = \mathfrak{a} \oplus \mathfrak{b}.\tag{2.20}
$$

$$
\mathfrak{h} = \mathfrak{a} \oplus [\mathfrak{h}, \mathfrak{h}] \oplus \mathfrak{b}. \tag{2.21}
$$

It should be noted that all the subspaces of $\mathfrak g$ appearing above are H-invariant.

3. Proofs of the main results

In this section, we will prove Theorem [1.1](#page-1-0) and Theorem [1.2.](#page-2-0) We will use an equivariant version of Cartan's theorem in the proofs, namely Theorem [6.15.](#page-41-8) Towards the end of this section, we take the opportunity to use Theorem [6.15](#page-41-8) again and record alternative proofs of results in [\[BCM\]](#page-41-0) describing the first and second cohomologies of homogeneous spaces.

3.1. Some preparatory results. We begin with a couple of useful general observations.

Lemma 3.1. Let $\mathfrak g$ be a Lie algebra over R. Assume that $\mathfrak g = \mathfrak g_1 \oplus \mathfrak g_2$ where $\mathfrak g_1, \mathfrak g_2$ are ideals in $\mathfrak g$ and $\mathfrak g_1 = [\mathfrak g_1, \mathfrak g_1]$. Let $\theta : \mathfrak g \times \mathfrak g \longrightarrow \mathbb R$ be a $\mathfrak g$ -invariant bilinear form on $\mathfrak g$. Then \mathfrak{g}_1 and \mathfrak{g}_2 are mutually orthogonal with respect to θ .

Proof. Let $X, Y \in \mathfrak{g}_1$, and $Z \in \mathfrak{g}_2$. Since θ is a $\mathfrak{g}\text{-invariant bilinear form, it}$ follows that $\theta([X, Y], Z) = \theta(X, [Y, Z])$. As \mathfrak{g}_1 and \mathfrak{g}_2 are ideals in \mathfrak{g} , we know that $[Y, Z] \in \mathfrak{g}_1 \cap \mathfrak{g}_2$. Thus $[Y, Z] = 0$. This completes the proof.

As the Lie algebra of a compact Lie group is reductive, we have an immediate corollary.

Corollary 3.2. Let G be a compact Lie group with Lie algebra \mathfrak{g} . Consider the adjoint action of G on $\mathfrak g$, and let θ be a G-invariant bilinear form on $\mathfrak g$. Then $\mathfrak z(\mathfrak g)$ and $[\mathfrak g, \mathfrak g]$ are mutually orthogonal with respect to θ . Furthermore, if $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_r$ with \mathfrak{g}_i a simple ideals in \mathfrak{g} for every $i = 1, \ldots, r$, then $\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_r$ is in fact a decomposition into subspaces which are mutually orthogonal with respect to θ . In particular, if θ is an inner product, then $[\mathfrak{g}, \mathfrak{g}]^{\perp} = \mathfrak{z}(\mathfrak{g}).$

It is important to keep in mind that the following convention and notation will be used in rest of this section.

C.1 In what follows whenever we have a vector space V and subspaces V_1, \ldots, V_k of it such that $V = V_1 \oplus \cdots \oplus V_k$, for every $i = 1, \ldots, k$ the dual V_i^* is identified with the subspace of V^* consisting all elements that vanish on V_j for every $j \neq i$. With this identification, we have the direct sum decomposition $V^* = V_1^* \oplus \cdots \oplus V_k^*$. Further, for all $1 \leq i, j \leq k$, let $V_i^* \vee V_j^* \subset S^2(V^*)$ be the subspace defined by $V_i^* \vee V_j^* := \text{Span}\{f \vee g \mid f \in V_i^*, g \in V_j^*\}.$ Then we have $S^2(V^*) =$ $\bigoplus_{i\leq j}V_i^*\vee V_j^*$. We will use this as well as the analogous decompositions of $S^r(V^*)$ and $\Lambda^r V^*$.

- C.2 In this section we will consider a general connected compact Lie group G and a closed subgroup H , which is not necessarily connected. We fix a G -invariant inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g} . Recall that $P_{\mathfrak{g}}^1 = \{ f \in \mathfrak{g}^* \mid f|_{[\mathfrak{g},\mathfrak{g}]} = 0 \}$. It follows that the coadjoint action of G on \mathfrak{g}^* keeps $P_{\mathfrak{g}}^1$ invariant. We now choose a G-invariant inner product $\langle \cdot, \cdot \rangle^*$ on $P_{\mathfrak{g}}^1$ (the latter inner product may be obtained as the restriction of a G-invariant inner product on \mathfrak{g}^*).
- **C**.3 Let $j^* : \mathfrak{g}^* \longrightarrow \mathfrak{h}^*$ be the map as in [\(2.11\)](#page-9-1). Observe that $j^*(P_{\mathfrak{g}}^1) \subset (\mathfrak{h}^*)^H$. Let $R_{\mathfrak{g},\mathfrak{h}}: P_{\mathfrak{g}}^1 \longrightarrow (\mathfrak{h}^*)^H$ denote the restriction of j^* to $P_{\mathfrak{g}}^1$. Thus $R_{\mathfrak{g},\mathfrak{h}}$ maps $f \longmapsto f|_{\mathfrak{h}}$ for all $f \in P^1_{\mathfrak{g}}$.

As the coadjoint action of H on \mathfrak{g}^* keeps $P_{\mathfrak{g}}^1$ invariant, it is easy to see that ker $R_{\mathfrak{g},\mathfrak{h}}$ is a H-invariant subspace of $P_{\mathfrak{g}}^1$. Moreover, we have the following H-invariant orthogonal decomposition:

$$
P_{\mathfrak{g}}^1 = \ker \mathcal{R}_{\mathfrak{g}, \mathfrak{h}} \oplus (\ker \mathcal{R}_{\mathfrak{g}, \mathfrak{h}})^\perp.
$$
 (3.1)

Consider the map ∇^1 :

$$
\nabla^1: 1 \otimes P_{\mathfrak{g}}^1 \longrightarrow (\mathfrak{h}^*)^H \otimes 1 \quad \text{ given by} \quad \nabla^1(1 \otimes f) := f|_{\mathfrak{h}} \otimes 1. \tag{3.2}
$$

Observe that $\ker \nabla^1 \simeq \ker R_{\mathfrak{g},\mathfrak{h}} = \{ f \in \mathfrak{g}^* \mid f|_{[\mathfrak{g},\mathfrak{g}]+\mathfrak{h}} = 0 \} \simeq \left(\frac{\mathfrak{g}}{[\mathfrak{g},\mathfrak{g}]+\mathfrak{h}} \right)$ ∗ (3.3)

Lemma 3.3. Let ∇^1 : $1 \otimes P^1_{\mathfrak{g}} \longrightarrow (\mathfrak{h}^*)^H \otimes 1$ be as in [\(3.2\)](#page-12-0). Then

 $\dim_\mathbb{R} \mathrm{Im} \nabla^1 \,=\, \dim_\mathbb{R} P_\mathfrak{g}^1 - \dim_\mathbb{R} \ker \nabla^1 \,=\, \dim_\mathbb{R} ([\mathfrak{g},\,\mathfrak{g}] + \mathfrak{h}) - \dim_\mathbb{R} [\mathfrak{g},\,\mathfrak{g}] \,=\, \dim_\mathbb{R} \mathfrak{b}.$ In particular, $\text{Im }\nabla^1 = (\mathfrak{b}^*)^H \otimes 1$ and $(\mathfrak{b}^*)^H = \mathfrak{b}^*$.

Proof. Note that $\dim_{\mathbb{R}} \text{Im}\nabla^1 = \dim_{\mathbb{R}} P^1_{\mathfrak{g}} - \dim_{\mathbb{R}} \ker \nabla^1 = \dim_{\mathbb{R}} \mathfrak{z}(\mathfrak{g}) - \dim_{\mathbb{R}} \mathfrak{g} +$ $\dim_\mathbb{R}([\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h})\,=\,\dim_\mathbb{R}([\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h})-\dim_\mathbb{R}[\mathfrak{g},\,\mathfrak{g}]\,=\,\dim_\mathbb{R}\mathfrak{b},\,(\text{see}\;(2.18)).$

Note that $\text{Im }\nabla^1 \subseteq (\mathfrak{b}^*)^H \otimes 1$. Thus $\dim_{\mathbb{R}} \text{Im }\nabla^1 \leq \dim_{\mathbb{R}} (\mathfrak{b}^*)^H$. As $\dim_{\mathbb{R}} \text{Im }\nabla^1 =$ dim_R **b**, it follows that $\text{Im }\nabla^1 = (\mathfrak{b}^*)^H \otimes 1$. This completes the proof.

From the above result, it follows that, although H is not necessarily connected, it fixes all the elements of \mathfrak{b}^* . This fact also follows from the following surprising lemma, which may be of independent interest.

Lemma 3.4. Let G be a connected compact Lie group, and $H \subset G$ be a closed subgroup which is not necessarily connected. Let $\mathfrak{p} \subset \mathfrak{g}$ be a subspace such that $\text{Ad}(H)(\mathfrak{p}) \subset \mathfrak{p}$. Assume that $\langle \cdot, \cdot \rangle$ is a G-invariant inner product on g. Then the Lie group H acts *trivially on the space* $(\mathfrak{p} \cap [\mathfrak{g}, \mathfrak{g}])^{\perp} \cap \mathfrak{p}$.

Proof. Recall that $\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus [\mathfrak{g}, \mathfrak{g}]$. For any $Z \in (\mathfrak{p} \cap [\mathfrak{g}, \mathfrak{g}])^{\perp} \cap \mathfrak{p}$, we can write $Z =$ $X + Y$ where $X \in \mathfrak{z}(\mathfrak{g})$ and $Y \in [\mathfrak{g}, \mathfrak{g}]$. Let $h \in H$. Then $\text{Ad}(h)Z - Z \in (\mathfrak{p} \cap [\mathfrak{g}, \mathfrak{g}])^{\perp} \cap \mathfrak{p}$. On the other hand $\text{Ad}(h)Y - Y \in [\mathfrak{g}, \mathfrak{g}]$. Since $\text{Ad}(h)Z - Z = \text{Ad}(h)Y - Y$, we also have $\text{Ad}(h)Z - Z \in \mathfrak{p} \cap [\mathfrak{g}, \mathfrak{g}]$. Thus $\text{Ad}(h)Z - Z = 0$. This completes the proof. \Box

We will now return to the direct sum decompositions as in Section [2.2.6](#page-10-1) and observe some more facts associated with actions of H on some of the summands. These facts will be used later in this section.

The following simple yet useful fact on the representations of compact Lie groups, which we state below, skipping the proof, will be needed later.

Proposition 3.5. Let K be a compact Lie group which is not necessarily connected and V be a finite-dimensional vector space over R. Assume that $\phi : K \longrightarrow GL(V)$ is a smooth representation of K. Let V^K be the space of fixed points in V under the action of K, that is, $V^K = \{v \mid \phi(g)(v) = v \text{ for all } g \in K\}$ and let $(K-1)V$ be the R-span of the elements of the set $\{\phi(q)v - v \mid q \in K, v \in V\}.$

- (1) Then $(K-1)V$ is the unique K-invariant complement of V^K . In particular, $V = V^K \oplus (K-1)V.$
- (2) Let B be any K-invariant bilinear form on V. Then $V = V^K \oplus (K 1)V$ is an orthogonal direct sum with respect to B.

Recall that all the direct summands in Section [2.2.6](#page-10-1) are H-invariant under the adjoint action. It is now easy to observe that $\mathfrak{h}^H \subset \mathfrak{z}(\mathfrak{h})$. Thus $\mathfrak{h}^H = \mathfrak{z}(\mathfrak{h})^H$ and $[\mathfrak{h}, \mathfrak{h}]^H =$ $0, (H-1)[\mathfrak{h}, \mathfrak{h}] = [\mathfrak{h}, \mathfrak{h}].$ As $\mathfrak{b} = \mathfrak{b}^H \oplus (H-1)\mathfrak{b}$ and $\mathfrak{b} = \mathfrak{b}^H$, it follows that $(H-1)\mathfrak{b} = 0$. Thus using (2.21) we obtain

$$
\mathfrak{h} = \mathfrak{a}^H \oplus (H - 1)\mathfrak{a} \oplus \mathfrak{b} \oplus [\mathfrak{h}, \mathfrak{h}]. \tag{3.4}
$$

Moreover, we have

$$
(H-1)\mathfrak{h} = (H-1)\mathfrak{a} \oplus [\mathfrak{h}, \mathfrak{h}], \qquad (3.5)
$$

$$
(H-1)\mathfrak{z}(\mathfrak{h}) = (H-1)\mathfrak{a} \text{ and } \mathfrak{h}^H = \mathfrak{z}(\mathfrak{h})^H = \mathfrak{a}^H \oplus \mathfrak{b}.
$$
 (3.6)

It may be further observed that the decompositions as in (3.4) , (3.5) and (3.6) are Hinvariant and orthogonal with respect to the (G-invariant) inner product $\langle \cdot, \cdot \rangle$ introduced in Section [2.2.6.](#page-10-1)

3.2. Description of third cohomology. Here we will prove Theorem [1.1.](#page-1-0) To do this we first compute ker ∇^2 and Im ∇^2 . Let

$$
\mathbf{A}_1 := \text{Span}\{f_1|_{\mathfrak{h}} \otimes f_2 - f_2|_{\mathfrak{h}} \otimes f_1 \mid f_1, f_2 \in (\text{ker } R_{\mathfrak{g},\mathfrak{h}})^{\perp}\},\tag{3.7}
$$

$$
\mathbf{S}_1 := \text{Span}\{f_1|_{\mathfrak{h}} \otimes f_2 + f_2|_{\mathfrak{h}} \otimes f_1 \mid f_1, f_2 \in (\text{ker } R_{\mathfrak{g},\mathfrak{h}})^{\perp}\}.
$$
 (3.8)

Lemma 3.6. Let A_1, S_1 be defined as above. Then the following hold :

(1) ker $\nabla^i|_{1\otimes(\wedge^i P^1_{\mathfrak{g}})}\simeq 1\otimes(\wedge^i \ker R_{\mathfrak{g},\mathfrak{h}})\simeq \wedge^i\left(\frac{\mathfrak{g}}{[\mathfrak{g},\mathfrak{g}]}\right)$ $[\mathfrak{g},\mathfrak{g}]{+\mathfrak{h}}$ $\big)$ ^{*}, for $i = 2, 3, 4$. (2) dim_R Im ∇^2 = dim_R($\wedge^2 P_g^1$) – dim_R(\wedge^2 ker R_{g,b}) $= (\dim_{\mathbb{R}} P_{\mathfrak{g}}^1 - \dim_{\mathbb{R}} \ker R_{\mathfrak{g},\mathfrak{h}}) (\dim_{\mathbb{R}} P_{\mathfrak{g}}^1 + \dim_{\mathbb{R}} \ker R_{\mathfrak{g},\mathfrak{h}} - 1)/2.$ (3) Im $\nabla^2 = (\mathfrak{b}^* \otimes \ker R_{\mathfrak{g},\mathfrak{h}}) \bigoplus A_1.$ (4) $\mathfrak{b}^* \otimes (\text{ker } R_{\mathfrak{g},\mathfrak{h}})^{\perp} = \mathbf{A}_1 \oplus \mathbf{S}_1$. (5) $\operatorname{Im} \nabla^2 \bigoplus (\mathfrak{a}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus \mathbf{S}_1 = (\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1$.

Proof. The proof of (1) , (2) , (3) follows from (3.3) , and Lemma [2.1](#page-6-2) by replacing the linear map T by $R_{\mathfrak{g},\mathfrak{h}}$.

To prove (4), we will use the linear isomorphism $R_{\mathfrak{g},\mathfrak{h}}$: (ker $R_{\mathfrak{g},\mathfrak{h}}$)^{\perp} \longrightarrow \mathfrak{b}^* . Finally, in view of the convention mentioned at $(C.1)$ $(C.1)$ it follows from the decomposition (3.6) that $(\mathfrak{h}^*)^H = (\mathfrak{a}^*)^H \oplus \mathfrak{b}^*$. Now (5) follows from [\(3.1\)](#page-12-2), (3) and (4).

Recall that for a semisimple Lie algebra \mathfrak{k} , the number of simple factors of \mathfrak{k} is denoted $by \#E.$

Proposition 3.7. Let G be a compact connected Lie group. Then

$$
\dim_{\mathbb{R}} H^3(G,\,\mathbb{R})\,=\,\,\#[\mathfrak{g},\,\mathfrak{g}]+\dim_{\mathbb{R}}\wedge^3\mathfrak{z}(\mathfrak{g})\,.
$$

Proof. See [\[GHV-III,](#page-42-6) Corollary I, p. 205].

Corollary 3.8. Let G be a compact connected Lie group, and $P_{\mathfrak{g}}^3$ be the space of primitive three-forms. Then

$$
\dim_{\mathbb{R}} P_{\mathfrak{g}}^3 = \#[\mathfrak{g}, \mathfrak{g}].
$$

Proof. The proof follows from Proposition [3.7](#page-14-2) and Hopf's Theorem which says that for any compact connected Lie group G, the cohomology algebra $H^*(G,\mathbb{R}) \simeq \wedge P_{\mathfrak{g}}$. \Box

Let G be a compact connected Lie group, and $r := \#[\mathfrak{g}, \mathfrak{g}]$. Let

$$
\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_r, \qquad (3.9)
$$

where \mathfrak{g}_i 's are simple factors in $[\mathfrak{g}, \mathfrak{g}]$. Let B_i denotes the Killing form on \mathfrak{g}_i . Define B_i on g by

$$
B_i(X, Y) := B_i(X_i, Y_i), \qquad (3.10)
$$

where $X = X_0 + \sum_i X_i$, $Y = Y_0 + \sum_i Y_i$, $X_0, Y_0 \in \mathfrak{z}(\mathfrak{g})$ and $X_i, Y_i \in \mathfrak{g}_i$. Then $\mathbf{B}_i \in S^2(\mathfrak{g}^*)^{\mathfrak{g}}$. Define a map ρ as follows.

$$
\rho: S^2(\mathfrak{g}^*)^{\mathfrak{g}} \longrightarrow (\wedge^3 \mathfrak{g}^*)^{\mathfrak{g}}; \quad \rho(\eta)(x, y, z) := \eta([x, y], z). \tag{3.11}
$$

Then we observe from the Definition [2.3](#page-8-0) that $\rho = -2\rho_{\rm g}$ where $\rho_{\rm g}$ is the Cartan map. It is also clear that $\widetilde{B}_i \not\in \text{ker } \rho$ for all $1 \leq i \leq r$.

Lemma 3.9. Let $r := #[\mathfrak{g}, \mathfrak{g}]$. Let $P_{\mathfrak{g}}^3$ be the space of primitive three-forms. Then the set $\{\rho(\mathbf{B}_i) \mid 1 \leq i \leq r\}$ forms a basis of $P^3_{\mathfrak{g}}$.

Proof. As the Killing form B_i is non-degenerate on \mathfrak{g}_i , $\rho(B_i) \neq 0$ for $1 \leq i \leq r$. In view of [\[GHV-III,](#page-42-6) Theorem II, p. 244], $\rho(\dot{B}_i) \in P^3_{\mathfrak{g}}$. It also follows from the definitions of ρ and B_i that $\{\rho(B_1), \ldots, \rho(B_r)\}\$ are linearly independent. Let $\lambda_i, i = 1, \ldots, r$ be scalars in R such that $\sum_i \lambda_i \rho(\widetilde{B}_i) = 0$. Let $X_i, Y_i, Z_i \in \mathfrak{g}_i, i = 1, \ldots, r$ satisfying $B_i([X_i, Y_i], Z_i) \neq 0$. If we substitute X_i, Y_i, Z_i in the above relation, we will get $\lambda_i = 0$ for all *i*. Now the proof follows as dim_R $P_{\mathfrak{g}}^3 = r$; see Corollary [3.8.](#page-14-3)

Now we define an explicit transgression which will be useful in the computation of the third and fourth cohomology groups. Using the basis in the Lemma [3.9](#page-14-1) above, we first define a linear map

$$
\widetilde{\tau}^3 \colon P^3_{\mathfrak{g}} \longrightarrow S^2(\mathfrak{g}^*)^{\mathfrak{g}}; \quad \widetilde{\tau}^3(\rho(\widetilde{\mathbf{B}}_i)) := \widetilde{\mathbf{B}}_i. \tag{3.12}
$$

In view of Remark [2.6\(](#page-8-2)2), we may choose a transgression $\tau = \bigoplus_i \tau^i$ such that $\tau^3 := \tilde{\tau}^3$. As $\{\rho(\mathbf{B}_i) \mid 1 \leq i \leq \#[\mathfrak{g}, \mathfrak{g}]\}$ is a basis of $P_{\mathfrak{g}}^3$, we may further define

$$
\nabla^3(1\otimes\rho(\widetilde{\mathrm{B}}_i))\,:=\,\widetilde{\mathrm{B}}_i|_{\mathfrak{h}\times\mathfrak{h}}\otimes 1\,.
$$

Lemma 3.10. Let G be a connected compact Lie group, and $H \subseteq G$ be a closed subgroup which is not necessarily connected. Let $\mathfrak g$ be the Lie algebra of G and $\mathfrak h \subset \mathfrak g$ be the Lie subalgebra corresponding to H . Let B be a G -invariant symmetric bilinear form on \mathfrak{g} . Let $\mathfrak{a} := \mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}]$ as in [\(2.19\)](#page-10-2). Then the following are equivalent.

 (1) B $|_{({\mathfrak{h}} \cap [{\mathfrak{g}}, {\mathfrak{g}}]) \times ({\mathfrak{h}} \cap [{\mathfrak{g}}, {\mathfrak{g}}])} = 0.$ $\left(2\right) B\right|_{\left(\left(3\left(\mathfrak{h}\right) \cap \left[\mathfrak{g},\mathfrak{g}\right] \right) + \left[\mathfrak{h},\mathfrak{h}\right] \right)} \times \left(\left(3\left(\mathfrak{h}\right) \cap \left[\mathfrak{g},\mathfrak{g}\right] \right) + \left[\mathfrak{h},\mathfrak{h}\right] \right) = 0.$ (3) $B|_{\mathfrak{a} \times \mathfrak{a}} = 0$ and $B|_{[\mathfrak{h}, \mathfrak{h}] \times [\mathfrak{h}, \mathfrak{h}]} = 0.$ (4) B $|_{\mathfrak{a}^H \times \mathfrak{a}^H} = 0$, B $|_{(H-1)\mathfrak{a} \times (H-1)\mathfrak{a}} = 0$ and B $|_{[\mathfrak{h}, \mathfrak{h}] \times [\mathfrak{h}, \mathfrak{h}]} = 0$. (5) B $|_{\mathfrak{a}^H \times \mathfrak{a}^H} = 0$ and B $|_{(H-1)\mathfrak{h} \times (H-1)\mathfrak{h}} = 0$.

Proof. Observe that $(H - 1)\mathfrak{h} = (H - 1)\mathfrak{z}(\mathfrak{h}) \oplus [\mathfrak{h}, \mathfrak{h}]$ is a Lie subalgebra of \mathfrak{h} . As $B|_{\mathfrak{h}\times\mathfrak{h}}$ is H-invariant and $\mathfrak{a}^H\subset\mathfrak{z}(\mathfrak{h})^H$ it follows using Lemma [3.1](#page-11-3) and Proposition [3.5](#page-13-4) that \mathfrak{a}^H , $(H-1)\mathfrak{a}$ and $[\mathfrak{h},\mathfrak{h}]$ are mutually orthogonal with respect to B. This completes the \Box

We next define a couple of spaces that are naturally associated to a pair of a connected compact Lie group G and a closed subgroup H , which will be required in the description of the third and fourth cohomologies of the homogeneous space G/H .

Definition 3.11. Let G be a compact connected Lie group, and $H \subseteq G$ be a closed subgroup which is not necessarily connected. Let $\Psi_{\mathfrak{g},\mathfrak{h}} : S^2([\mathfrak{g}, \mathfrak{g}]^*)^G \longrightarrow S^2((\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}])^*)^H$ be the restriction map defined by

$$
\Psi_{\mathfrak{g},\mathfrak{h}}(\phi) := \phi|_{(\mathfrak{h} \cap [\mathfrak{g},\mathfrak{g}]) \times (\mathfrak{h} \cap [\mathfrak{g},\mathfrak{g}])}.
$$
\n(3.13)

We then set

$$
\mathcal{N}_{\mathfrak{g},\mathfrak{h}} := \ker \Psi_{\mathfrak{g},\mathfrak{h}}, \text{ and } \mathcal{C}_{\mathfrak{g},H} := \frac{S^2((\mathfrak{h} \cap [\mathfrak{g},\mathfrak{g}])^*)^H}{\text{Im } \Psi_{\mathfrak{g},\mathfrak{h}}} = \text{coker } \Psi_{\mathfrak{g},\mathfrak{h}}.
$$

Remark 3.12. One may use Lemma [3.10](#page-15-1) to have alternative descriptions of $\mathcal{N}_{\mathfrak{a},\mathfrak{h}}$ involving $\mathfrak{z}(\mathfrak{h})\cap[\mathfrak{g},\mathfrak{g}]$ and $[\mathfrak{h},\mathfrak{h}]$. Note that the definition of $\mathcal{N}_{\mathfrak{g},\mathfrak{h}}$ does not involve H/H^0 and involves only data related to the Lie algebras g and h. On the other hand, the definition of $C_{\mathfrak{q},H}$ does involve the Lie group H (which is not necessarily connected) along with the Lie algebras $\mathfrak{g}, \mathfrak{h}$. However, if H^0 is a semisimple Lie group and all the simple factors of H are pairwise non-isomorphic, then the component group $H/H⁰$ does not play any role in the definition of the vector space $\mathcal{C}_{\mathfrak{g},H}$; see Lemma [4.4.](#page-24-0)

The next result will be used in the proof of Theorem [1.1.](#page-1-0)

Lemma 3.13. There is an isomorphism between the following vector spaces:

 $\mathcal{N}_{\mathfrak{g}, \mathfrak{h}} \, \simeq \, \ker \nabla^3|_{(\mathfrak{a}^*)^H \otimes (\ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}})^\perp} \oplus \mathbf{S}_1 \oplus \mathbf{1} \otimes P_{\mathfrak{g}}^3 \, .$

Proof. We begin by fixing some more notations which will be used in this proof as well as in the proof of Lemma 3.15 .

We first use the convention mentioned at $(C.1)$ $(C.1)$ and (3.4) to identify the dual spaces $(\mathfrak{a}^H)^*$, $((H-1)\mathfrak{a})^*$, \mathfrak{b}^* , $[\mathfrak{h}, \mathfrak{h}]^*$ as subspaces of \mathfrak{h}^* . Under this identification, we have

$$
\mathfrak{h}^* = (\mathfrak{a}^H)^* \oplus ((H-1)\mathfrak{a})^* \oplus \mathfrak{b}^* \oplus [\mathfrak{h}, \mathfrak{h}]^*.
$$
 (3.14)

We next fix some more notations, which will be used here and in the proof of Lemma [3.15.](#page-19-0)

- **D**.1 Let $r_{\mathfrak{a}^H} := \dim_{\mathbb{R}} \mathfrak{a}^H$, $r_{\mathfrak{b}} := \dim_{\mathbb{R}} \mathfrak{b}$. We fix bases $\{A_1, \ldots, A_{r_{\mathfrak{a}^H}}\}$ and $\{D_1, \ldots, D_{r_{\mathfrak{b}}}\}$ of \mathfrak{a}^H and \mathfrak{b} , respectively. Let $\{A_1^*, \ldots, A_{r_{\mathfrak{a}H}}^*\}$ be a basis of $(\mathfrak{a}^H)^*$ dual to the basis $\{A_1, \ldots, A_{r_{\mathfrak{a}H}}\}$ and $\{D_1^*, \ldots, D_{r_{\mathfrak{b}}}^*\}$ be a basis of \mathfrak{b}^* dual to the basis $\{D_1, \ldots, D_{r_{\mathfrak{b}}}\}.$ Thus $A_p^*, D_i^* \in \mathfrak{h}^*$ for all $1 \leq p \leq r_{\mathfrak{a}^H}$, $1 \leq i \leq r_{\mathfrak{b}}$. Moreover, $A_p^*(A_q) =$ $\delta_{pq}, A^*_{p}([H-1)a + b + [b, b]) = 0$, for all $1 \leq p, q \leq r_{a^H}$ and $D_i^*(D_j) =$ $\delta_{ij}, D_i^*(\mathfrak{a} + [\mathfrak{h}, \mathfrak{h}]) = 0$, for all $1 \leq i, j \leq r_{\mathfrak{b}}$. As $\mathfrak{b}^H = \mathfrak{b}$ it should be noted that elements of $(\mathfrak{a}^H)^*$ and \mathfrak{b}^* remain fixed under the action of H. Thus H fixes $A_p^*, D_i^* \in \mathfrak{h}^*$ for all $1 \leq p \leq r_{\mathfrak{a}^H}, 1 \leq i \leq r_{\mathfrak{b}}.$
- **D.**2 Recall that the restriction map $f \mapsto f|_{\mathfrak{b}}$ from $(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}$ to \mathfrak{b}^* is a linear isomorphism of vector spaces. Using this map, we choose a basis $\{\lambda_1, \dots, \lambda_{r_b}\}\$ of $(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}$ such that $\lambda_i|_{\mathfrak{b}} = D_i^*|_{\mathfrak{b}}$ for all $1 \leq i \leq r_{\mathfrak{b}}$. Moreover as $\lambda_i \in P_{\mathfrak{g}}^1$, it follows that $\lambda_i(\mathfrak{a} + [\mathfrak{h}, \mathfrak{h}]) = 0$ for all $1 \leq i \leq r_{\mathfrak{b}}$. Thus $\lambda_i|_{\mathfrak{h}} = D_i^*$ for all $1 \leq i \leq r_{\mathfrak{b}}$.

Now write any arbitrary element of $(\mathfrak{a}^*)^H \otimes (\text{ker } R_{\mathfrak{g},\mathfrak{h}})^\perp \bigoplus \mathbf{S}_1 \bigoplus 1 \otimes P_\mathfrak{g}^3$ with respect to the bases described above and in Lemma [3.9,](#page-14-1) say,

$$
\sum_{1 \leq i \leq r_{\mathfrak{a}} H \atop 1 \leq j \leq r_{\mathfrak{b}}} \alpha_{ij} A_i^* \otimes \lambda_j + \sum_{1 \leq l \leq k \leq r_{\mathfrak{b}}} \beta_{lk} (D_l^* \otimes \lambda_k + D_k^* \otimes \lambda_l) + \sum_{1 \leq i \leq \#[\mathfrak{g},\mathfrak{g}]} 1 \otimes \eta_i \rho(\widetilde{B}_i).
$$

Then apply ∇^3 on it:

$$
\nabla^3 \left(\sum_{i,j} \alpha_{ij} A_i^* \otimes \lambda_j + \sum_{l \le k} \beta_{lk} (D_l^* \otimes \lambda_k + D_k^* \otimes \lambda_l) + \sum_i 1 \otimes \eta_i \rho(\widetilde{B}_i) \right)
$$

=
$$
\left(\sum_{i,j} \alpha_{ij} A_i^* \vee D_j^* + \sum_{l \le k} 2\beta_{lk} D_l^* \vee D_k^* + \sum_i \eta_i \widetilde{B}_i |_{\mathfrak{h} \times \mathfrak{h}} \right) \otimes 1
$$
 (3.15)

Thus using [\(3.15\)](#page-16-0) it is immediate that the kernel of ∇^3 restricted to the subspace $(\mathfrak{a}^*)^H \otimes$ $(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \bigoplus \mathbf{S}_1 \bigoplus 1 \otimes P_{\mathfrak{g}}^3$ consists of elements $\sum_{i,j} \alpha_{ij} A_i^* \otimes \lambda_j + \sum_{l,k} \beta_{lk} (D_l^* \otimes \lambda_k + D_k^* \otimes \lambda_k)$ λ_l) + $\sum_i \eta_i \rho(\widetilde{B}_i)$ such that the symmetric bilinear form

$$
\sum_{i,j} \alpha_{ij} A_i^* \vee D_j^* + \sum_{l \le k} 2\beta_{lk} D_l^* \vee D_k^* + \sum_i \eta_i \widetilde{B}_i |_{\mathfrak{h} \times \mathfrak{h}}
$$
(3.16)

is identically zero on $\mathfrak{h} \times \mathfrak{h}$.

Observe that the form appearing in (3.16) is H-invariant. Thus by Proposition [3.5,](#page-13-4) the subspaces \mathfrak{h}^H and $(H-1)\mathfrak{h}$ are orthogonal with respect to this form. Furthermore, by Lemma [3.1,](#page-11-3) as $(H-1)\mathfrak{h} = (H-1)\mathfrak{z}(\mathfrak{h}) + [\mathfrak{h}, \mathfrak{h}],$ the spaces $(H-1)\mathfrak{z}(\mathfrak{h}) = (H-1)\mathfrak{a}$ and $[\mathfrak{h}, \mathfrak{h}]$ are orthogonal with respect to the form in [\(3.16\)](#page-16-1). Recall that $\mathfrak{h}^H = \mathfrak{a}^H + \mathfrak{b}$. Thus it follows that the form as in (3.16) is zero on $\mathfrak{h} \times \mathfrak{h}$ if and only if it is zero on each of the spaces $\mathfrak{a}^H \times \mathfrak{a}^H$, $\mathfrak{a}^H \times \mathfrak{b}$, $\mathfrak{b} \times \mathfrak{b}$, $(H-1)\mathfrak{h} \times (H-1)\mathfrak{h}$.

We next evaluate the form as in [\(3.16\)](#page-16-1) at (D_j, D_j) and (D_i, D_j) for $i \neq j$, respectively:

$$
4\beta_{jj} + \sum_{l} \eta_l \widetilde{B}_l(D_j, D_j) = 0, \qquad (3.17)
$$

$$
2\beta_{ij} + \sum_{l} \eta_l \widetilde{B}_l(D_i, D_j) = 0, \quad \text{for } i \neq j.
$$
 (3.18)

Similarly, by evaluating the form as in (3.16) at (A_i, D_j) we have :

$$
\alpha_{ij} + \sum_{l} \eta_l \widetilde{\mathbf{B}}_l(A_i, D_j) = 0. \qquad (3.19)
$$

It is immediate that the form as in [\(3.16\)](#page-16-1) is zero on each of the spaces $a^H \times \mathfrak{b}$, $\mathfrak{b} \times \mathfrak{b}$ if and only if (3.17) , (3.18) and (3.19) hold. Moreover, the form as in (3.16) is zero on each of the spaces $\mathfrak{a}^H \times \mathfrak{a}^H$, $(H-1)\mathfrak{h} \times (H-1)\mathfrak{h}$ if and only if

$$
\sum_{i} \eta_i \widetilde{B}_i \big|_{\left(\mathfrak{a}^H + (H - 1)\mathfrak{h}\right) \times \left(\mathfrak{a}^H + (H - 1)\mathfrak{h}\right)} = 0. \tag{3.20}
$$

In view of (3.18) , (3.17) , (3.19) , (3.20) and Lemma [3.10](#page-15-1) we conclude that $\mathcal{N}_{\mathfrak{g},\mathfrak{h}}$ and ker ∇^3 restricted to the subspace $(\mathfrak{a}^*)^H \otimes (\text{ker} R_{\mathfrak{g},\mathfrak{h}})^{\perp} \bigoplus \mathbf{S}_1 \bigoplus \mathbf{1} \otimes P_{\mathfrak{g}}^3$ are isomorphic.

Proof of Theorem [1.1.](#page-1-0) Recall the Koszul complex $(\mathscr{C}^* := S(\mathfrak{h}^*)^H \otimes \wedge P_{\mathfrak{g}}, \nabla)$ as defined in Section [2.2.4.](#page-9-0) As G is compact connected, the cohomology of G/H is given by the cohomology of the Koszul complex $(S(\mathfrak{h}^*)^H \otimes \wedge P_{\mathfrak{g}}, \nabla)$; see Theorem [6.15.](#page-41-8) Therefore,

$$
H^3(G/H) \simeq \frac{\ker \nabla^3}{\operatorname{Im} \nabla^2},\tag{3.21}
$$

where the differentials ∇^3 and ∇^2 are defined in [\(2.15\)](#page-9-2) and [\(2.14\)](#page-9-3), respectively. The numerator and the denominator in [\(3.21\)](#page-17-2) will be identified. Recall that

$$
\mathscr{C}^3 = (\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus 1 \otimes P_{\mathfrak{g}}^3 \bigoplus 1 \otimes \wedge^3 P_{\mathfrak{g}}^1,
$$

and Im $\nabla^2 \subset (\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1$. In fact, Im $\nabla^2 \bigoplus (\mathfrak{a}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus \mathbf{S}_1 = (\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1$; see Lemma [3.6\(](#page-13-5)5). Also recall that $\nabla^2((\mathfrak{a}^*)^H \otimes \ker R_{\mathfrak{g},\mathfrak{h}}) = 0$. Now

$$
\begin{aligned} \frac{\ker \nabla^3}{\mathrm{Im}\,\nabla^2}&\simeq\,\frac{\ker \nabla^3\big|_{\big((\mathfrak{h}^*)^H\otimes P_\mathfrak{g}^1\big)}\oplus 1\otimes P_\mathfrak{g}^3}{(\mathrm{Im}\,\nabla^2)}\bigoplus\ker \nabla^3\big|_{1\otimes\wedge^3 P_\mathfrak{g}^1}\\ &\simeq\,\ker \nabla^3\big|_{(\mathfrak{a}^*)^H\otimes\ker R_{\mathfrak{g},\mathfrak{h}}\oplus(\mathfrak{a}^*)^H\otimes(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}\oplus S_1\oplus 1\otimes P_\mathfrak{g}^3}\bigoplus 1\otimes\wedge^3\ker R_{\mathfrak{g},\mathfrak{h}}\\ &\simeq \big(\mathfrak{a}^*\big)^H\otimes\ker R_{\mathfrak{g},\mathfrak{h}}\bigoplus\ker \nabla^3\big|_{(\mathfrak{a}^*)^H\otimes(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}\oplus S_1\oplus 1\otimes P_\mathfrak{g}^3}\bigoplus 1\otimes\wedge^3\ker R_{\mathfrak{g},\mathfrak{h}}\\ &\simeq \big((\mathfrak{z}(\mathfrak{h})\cap [\mathfrak{h},\mathfrak{h}])^*\big)^H\otimes\Big(\frac{\mathfrak{g}}{[\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h}}\Big)^*\bigoplus \mathcal{N}_{\mathfrak{g},\mathfrak{h}}\bigoplus \wedge^3\Big(\frac{\mathfrak{g}}{[\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h}}\Big)^*. \end{aligned}
$$

The last isomorphism follows from (3.3) , Lemma [3.13,](#page-15-2) and Lemma [3.6\(](#page-13-5)1).

3.3. Description of fourth cohomology. Here we will prove Theorem [1.2.](#page-2-0) Now we will decompose $(\mathfrak{h}^*)^H \otimes (\wedge^2 P_{\mathfrak{g}}^1)$ into various subspaces using [\(3.6\)](#page-13-3) and [\(3.1\)](#page-12-2) as follows :

$$
(\mathfrak{h}^*)^H \otimes \wedge^2 P_{\mathfrak{g}}^1 = (\mathfrak{a}^*)^H \otimes (\wedge^2 P_{\mathfrak{g}}^1) \bigoplus \mathfrak{b}^* \otimes (\wedge^2 P_{\mathfrak{g}}^1)
$$

= $(\mathfrak{a}^*)^H \otimes \wedge^2 \ker R_{\mathfrak{g},\mathfrak{h}} \bigoplus (\mathfrak{a}^*)^H \otimes ((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge \ker R_{\mathfrak{g},\mathfrak{h}}) \bigoplus (\mathfrak{a}^*)^H \otimes \wedge^2 (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}\bigoplus \mathfrak{b}^* \otimes \wedge^2 \ker R_{\mathfrak{g},\mathfrak{h}} \bigoplus \mathfrak{b}^* \otimes ((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge \ker R_{\mathfrak{g},\mathfrak{h}}) \bigoplus \mathfrak{b}^* \otimes \wedge^2 (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}.$ \n(3.22)

We will next define two subspaces $\mathbf{A_2}$ and $\mathbf{S_2}$ of $\mathfrak{b}^* \otimes (\ker R_{\mathfrak{g},\mathfrak{h}} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp})$ as follows :

$$
\mathbf{A_2} := \operatorname{Span} \{ f_1|_{\mathfrak{b}} \otimes (g \wedge f_2) - f_2|_{\mathfrak{b}} \otimes (g \wedge f_1) \mid f_1, f_2 \in (\ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}})^{\perp}, g \in \ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}} \}, (3.23)
$$

\n
$$
\mathbf{S_2} := \operatorname{Span} \{ f_1|_{\mathfrak{b}} \otimes (g \wedge f_2) + f_2|_{\mathfrak{b}} \otimes (g \wedge f_1) \mid f_1, f_2 \in (\ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}})^{\perp}, g \in \ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}} \}. (3.24)
$$

Similarly, we define another two subspaces $\mathbf{A_3}$ and $\mathbf{S_3}$ of $\mathfrak{b}^* \otimes ((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp})$ as follows:

$$
\mathbf{A_3} := \text{Span}\left\{\n\begin{array}{l}\nf_1|_{\mathfrak{b}} \otimes (f_2 \wedge f_3) - f_2|_{\mathfrak{b}} \otimes (f_1 \wedge f_3) \\
+ f_3|_{\mathfrak{b}} \otimes (f_1 \wedge f_2)\n\end{array}\n\middle|\n\begin{array}{l} f_1, f_2, f_3 \in (\ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}})^\perp\n\end{array}\n\right\},\n\tag{3.25}
$$

$$
\mathbf{S_3} := \mathrm{Span} \left\{ \begin{array}{c} -2f_1|_{\mathfrak{b}} \otimes (f_2 \wedge f_3) - f_2|_{\mathfrak{b}} \otimes (f_1 \wedge f_3) \\ + f_3|_{\mathfrak{b}} \otimes (f_1 \wedge f_2) \end{array} \middle| f_1, f_2, f_3 \in (\ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}})^\perp \right\}.
$$
 (3.26)

Lemma 3.14. Let A_2 , S_2 , A_3 and S_3 be defined as above. Then the following hold :

- $(1) \nabla^3 (1 \otimes (\ker R_{\mathfrak{g},\mathfrak{h}} \wedge \ker R_{\mathfrak{g},\mathfrak{h}} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^\perp)) = \mathfrak{b}^* \otimes (\wedge^2 \ker R_{\mathfrak{g},\mathfrak{h}}).$
- (2) $\nabla^3\big(1\otimes(\ker R_{\mathfrak{g},\mathfrak{h}}\wedge(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}\wedge(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp})\big) = \mathbf{A}_2$ and $\nabla^3\big(1\otimes((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}\wedge(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}\wedge(\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp})\big)\,=\,\mathbf{A}_3.$
- $(\textit{3}) \ \nabla^3 (1 \otimes \wedge^3 P_{\mathfrak{g}}^1) \ = \ \mathfrak{b}^* \otimes \wedge^2 \ker \mathrm{R}_{\mathfrak{g}, \mathfrak{h}} \ \bigoplus \ \mathbf{A_2} \ \bigoplus \ \mathbf{A_3}.$
- (4) $\mathbf{A}_2 \oplus \mathbf{S}_2 = \mathfrak{b}^* \otimes (\ker R_{\mathfrak{g},\mathfrak{h}} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}).$
- (5) $\mathbf{A}_3 \oplus \mathbf{S}_3 = \mathfrak{b}^* \otimes ((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}).$
- $(6) \nabla^3 (1 \otimes \wedge^3 P_{\mathfrak{g}}^1) \bigoplus \mathbf{S_2} \bigoplus \mathbf{S_3} \bigoplus (\mathfrak{a}^*)^H \otimes (\wedge^2 P_{\mathfrak{g}}^1) = (\mathfrak{h}^*)^H \otimes (\wedge^2 P_{\mathfrak{g}}^1).$

Proof. Proof of (1) follows from the fact that $R_{\mathfrak{g},\mathfrak{h}}$: (ker $R_{\mathfrak{g},\mathfrak{h}}$)^{\perp} \longrightarrow \mathfrak{b}^* is a linear isomorphism. The statement in (2) is immediate from the definition of the map ∇^3 . Now (3) follows from (1) and (2). The proof of (4) follows form [\(2.9\)](#page-7-1). Similarly, the proof of (5) follows from (2.10) . Finally, (6) follows from (3) , (4) , (5) and (3.22) .

Recall that in view of Lemma $3.14(6)$ $3.14(6)$, the following isomorphism in (3.27) holds. Since $\nabla^4((\mathfrak{a}^*)^H \otimes (\ker R_{\mathfrak{g},\mathfrak{h}} \wedge \ker R_{\mathfrak{g},\mathfrak{h}})) = 0$, the isomorphism in [\(3.28\)](#page-18-2) follows from the direct sum decomposition in [\(3.22\)](#page-17-3).

$$
\frac{\ker \nabla^4|_{(\mathfrak{h}^*)^H \otimes (\wedge^2 P_\mathfrak{g}^1) \bigoplus 1 \otimes (P_\mathfrak{g}^3 \wedge P_\mathfrak{g}^1)}{\nabla^3 (1 \otimes \wedge^3 P_\mathfrak{g}^1)} \simeq \ker \nabla^4|_{\mathbf{S}_2 \bigoplus \mathbf{S}_3 \bigoplus (\mathfrak{a}^*)^H \otimes (\wedge^2 P_\mathfrak{g}^1) \bigoplus 1 \otimes (P_\mathfrak{g}^3 \wedge P_\mathfrak{g}^1)}\n\tag{3.27}
$$

$$
\simeq (\mathfrak{a}^*)^H \otimes (\ker R_{\mathfrak{g}, \mathfrak{h}} \wedge \ker R_{\mathfrak{g}, \mathfrak{h}}) \bigoplus \qquad (3.28)
$$

$$
\ker \nabla^4|_{(\mathfrak{a}^*)^H \otimes ((\ker \mathrm{R}_{\mathfrak{g},\mathfrak{h}})^\perp \wedge \ker \mathrm{R}_{\mathfrak{g},\mathfrak{h}}) \bigoplus (\mathfrak{a}^*)^H \otimes \wedge^2 (\ker \mathrm{R}_{\mathfrak{g},\mathfrak{h}})^\perp \bigoplus \mathrm{S}_2 \bigoplus \mathrm{S}_3 \bigoplus 1 \otimes (P_\mathfrak{g}^3 \wedge P_\mathfrak{g}^1)
$$

Recall that for a vector space V and a positive integer $k, V^k := V \oplus \cdots \oplus V$ \overline{k} -many .

.

Lemma 3.15. The following isomorphisms hold :

$$
\ker \nabla^{4} \Big|_{(\mathfrak{a}^{*})^{H} \otimes \left((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge \ker R_{\mathfrak{g},\mathfrak{h}} \right) \bigoplus (\mathfrak{a}^{*})^{H} \otimes \wedge^{2} (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \bigoplus S_{2} \bigoplus S_{3} \bigoplus 1 \otimes (P_{\mathfrak{g}}^{3} \wedge P_{\mathfrak{g}}^{1})
$$
\n
$$
\simeq \ker \nabla^{4} \Big|_{(\mathfrak{a}^{*})^{H} \otimes \left((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge \ker R_{\mathfrak{g},\mathfrak{h}} \right) \bigoplus S_{2} \bigoplus 1 \otimes (P_{\mathfrak{g}}^{3} \wedge \ker R_{\mathfrak{g},\mathfrak{h}})
$$
\n
$$
\bigoplus \ker \nabla^{4} \Big|_{(\mathfrak{a}^{*})^{H} \otimes \wedge^{2} (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \bigoplus S_{3} \bigoplus 1 \otimes (P_{\mathfrak{g}}^{3} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp})
$$
\n
$$
\simeq (\mathcal{N}_{\mathfrak{g},\mathfrak{h}})^{\dim_{\mathbb{R}} \ker R_{\mathfrak{g},\mathfrak{h}}.
$$
\n(3.30)

Proof. Observe that

$$
(\mathfrak{a}^*)^H \otimes ((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge \ker R_{\mathfrak{g},\mathfrak{h}}) \bigoplus (\mathfrak{a}^*)^H \otimes \wedge^2 (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \bigoplus S_2 \bigoplus S_3 \bigoplus 1 \otimes (P_{\mathfrak{g}}^3 \wedge P_{\mathfrak{g}}^1)
$$

=
$$
((\mathfrak{a}^*)^H \otimes ((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge \ker R_{\mathfrak{g},\mathfrak{h}}) \bigoplus S_2 \bigoplus 1 \otimes (P_{\mathfrak{g}}^3 \wedge \ker R_{\mathfrak{g},\mathfrak{h}})\big)
$$

$$
\bigoplus (\mathfrak{a}^*)^H \otimes \wedge^2 (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \bigoplus S_3 \bigoplus 1 \otimes (P_{\mathfrak{g}}^3 \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp})\big).
$$

In view of (2.16) , it follows that

$$
\nabla^{4} ((\mathfrak{a}^{*})^{H} \otimes (\ker R_{\mathfrak{g},\mathfrak{h}} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}) \bigoplus S_{2} \bigoplus 1 \otimes (P_{\mathfrak{g}}^{3} \wedge \ker R_{\mathfrak{g},\mathfrak{h}})\n\n\subseteq S^{2}(\mathfrak{h}^{*})^{H} \otimes \ker R_{\mathfrak{g},\mathfrak{h}}, \text{ and}
$$
\n
$$
\nabla^{4} ((\mathfrak{a}^{*})^{H} \otimes ((\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}) \bigoplus S_{3} \bigoplus 1 \otimes (P_{\mathfrak{g}}^{3} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp})\n\n\subseteq S^{2}(\mathfrak{h}^{*})^{H} \otimes (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp} \bigoplus \mathfrak{b}^{*} \otimes P_{\mathfrak{g}}^{3}.
$$
\n(3.32)

Since the right hand sides of [\(3.31\)](#page-19-1) and [\(3.32\)](#page-19-2) are distinct direct summands of $S^2(\mathfrak{h}^*)^H \otimes$ $P_{\mathfrak{g}}^1 \bigoplus \mathfrak{b}^* \otimes P_{\mathfrak{g}}^3$, [\(3.29\)](#page-19-3) holds.

Let $r_0 := \dim_{\mathbb{R}} \ker R_{\mathfrak{g},\mathfrak{h}} = \dim_{\mathbb{R}} \frac{\mathfrak{g}}{\lceil \mathfrak{g}, \mathfrak{g} \rceil}$ $\frac{\mathfrak{g}}{[\mathfrak{g},\mathfrak{g}]+\mathfrak{h}}$. Let $\{\nu_1,\cdots,\nu_{r_0}\}$ be a basis of ker $R_{\mathfrak{g},\mathfrak{h}}$. We now adhere to the notations used in $(D.1)$ $(D.1)$ and $(D.2)$ $(D.2)$ $(D.2)$. Let \mathbf{A}_2 , \mathbf{S}_2 , \mathbf{A}_3 and \mathbf{S}_3 be defined as in [\(3.23\)](#page-18-3), [\(3.24\)](#page-18-4),[\(3.25\)](#page-18-5) and [\(3.26\)](#page-18-6), respectively. Then observe that $\{D_i^* \otimes (\nu_k \wedge \lambda_l)$ – $D_l^* \otimes (\nu_k \wedge \lambda_i)$ | $1 \leq i < l \leq r_b$, $1 \leq k \leq r_0$ } and $\{D_i^* \otimes (\nu_k \wedge \lambda_l) + D_l^* \otimes (\nu_k \wedge \lambda_i)$ | $1 \leq i \leq l \leq r_{\mathfrak{b}}, 1 \leq k \leq r_0$ form a basis of $\mathbf{A_2}$ and $\mathbf{S_2}$, respectively. Similarly, $\{D_i^* \otimes (\lambda_k \wedge \lambda_l) - D_k^* \otimes (\lambda_i \wedge \lambda_l) + D_l^* \otimes (\nu_k \wedge \lambda_i) \mid 1 \leq i \leq k \leq l \leq r_{\mathfrak{b}}\},\ \{-2D_i^* \otimes$ $(\lambda_k \wedge \lambda_l) - D_k^* \otimes (\lambda_i \wedge \lambda_l) + D_l^* \otimes (\lambda_i \wedge \lambda_k) \mid 1 \leq k \leq i \leq r_{\mathfrak{b}}, k < l \leq r_{\mathfrak{b}}\}$ form a basis of A_3 , S_3 , respectively.

We divide the rest of the proof into two steps.

Step 1. In this step we will prove

$$
\ker \nabla^4|_{(\mathfrak{a}^*)^H \otimes (\wedge^2(\ker \mathrm{R}_{\mathfrak{g},\mathfrak{h}})^\perp) \bigoplus \mathbf{S}_3 \bigoplus 1 \otimes (P_\mathfrak{g}^3 \wedge (\ker \mathrm{R}_{\mathfrak{g},\mathfrak{h}})^\perp) = 0. \tag{3.33}
$$

We begin by applying ∇^4 on any arbitrary element of $(\mathfrak{a}^*)^H \otimes (\wedge^2 (\ker R_{\mathfrak{g},\mathfrak{h}})^\perp) \bigoplus \mathbf{S}_3 \bigoplus 1 \otimes$ $(P_{\mathfrak{g}}^3 \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}), \text{ say},$

$$
\sum_{1 \leq i \leq r_{\mathfrak{a}H} \atop 1 \leq j < l \leq r_{\mathfrak{b}}} \mu_{ijl} A_i^* \otimes (\lambda_j \wedge \lambda_l) \bigoplus \sum_{1 \leq j \leq i \leq r_{\mathfrak{b}} \atop j < l \leq r_{\mathfrak{b}}} \alpha_{ijl} \Big(-2D_i^* \otimes (\lambda_j \wedge \lambda_l) - D_j^* \otimes (\lambda_i \wedge \lambda_l) + D_l^* \otimes (\lambda_i \wedge \lambda_j) \Big) \bigoplus \sum_{1 \leq i \leq \#[\mathfrak{g}, \mathfrak{g}]} 1 \otimes \zeta_{ij} (\rho(\widetilde{\mathbf{B}}_i) \wedge \lambda_j)
$$
(3.34)

and we obtain

$$
\nabla^{4}\Big(\sum_{\substack{i\\j
+ $D_{l}^{*}\otimes(\lambda_{i}\wedge\lambda_{j})\Big)\bigoplus\sum_{i,j}\mathbb{1}\otimes\zeta_{ij}(\rho(\widetilde{B}_{i})\wedge\lambda_{j})\Big) =$

$$
\sum_{\substack{i\\j

$$
-(D_{i}^{*}\vee D_{j}^{*})\otimes\lambda_{l}\big)+\sum_{i,j}\zeta_{ij}(\widetilde{B}_{i}|_{\mathfrak{h}\times\mathfrak{h}}\otimes\lambda_{j}-D_{j}^{*}\otimes\rho(\widetilde{B}_{i})).
$$
 (3.35)
$$
$$

We will now equate the right hand side of (3.35) to zero, and show that this will force all the coefficients $\mu_{ijl}, \alpha_{ijl}, \zeta_{ij}$ in [\(3.34\)](#page-19-4) to be zero.

Since $D_j^* \otimes \rho(\dot{B}_i)$ is a part of basis of $(\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^3$, it follows that $\zeta_{ij} = 0$ for all i, j . Next consider the term of the form $(- \vee -) \otimes \lambda_{r_{\mathfrak{b}}}$ in [\(3.35\)](#page-20-0). Then

$$
\Big(\sum_{\substack{i\\j
$$

From the above equation it follows by substituting at dual basis element that $\mu_{ijr_b} = 0$ and $\alpha_{ijr_b} = 0$. Next we will consider the term of the form $(- \vee -) \otimes \lambda_{r_b-1}$ in (3.35) . We will get the following relation:

$$
\Big(\sum_{\substack{i\\j
$$

Similarly, we conclude that $\mu_{ijr_b-1} = 0$ and $\alpha_{ijr_b-1} = 0$. Proceeding in this way we can conclude that all the coefficients in [\(3.34\)](#page-19-4) are 0. This proves [\(3.33\)](#page-19-5).

Step 2. In this step we will prove the following isomorphism

$$
\ker \nabla^4|_{(\mathfrak{a}^*)^H \otimes (\ker R_{\mathfrak{g},\mathfrak{h}} \wedge (\ker R_{\mathfrak{g},\mathfrak{h}})^{\perp}) \oplus \mathbf{S}_2 \oplus \mathbf{1} \otimes (P_{\mathfrak{g}}^3 \wedge \ker R_{\mathfrak{g},\mathfrak{h}})} \simeq (\mathcal{N}_{\mathfrak{g},\mathfrak{h}})^{\dim_{\mathbb{R}} \ker R_{\mathfrak{g},\mathfrak{h}}}.
$$
 (3.36)

As in Step 1 we will apply the map ∇^4 on any arbitrary element of $(\mathfrak{a}^*)^H \otimes (\ker R_{\mathfrak{g},\mathfrak{h}} \wedge$ $(\ker R_{\mathfrak{g},\mathfrak{h}})^\perp\big)\bigoplus \mathbf{S}_2\, \bigoplus 1\otimes (P_\mathfrak{g}^3\wedge \ker \mathrm{R}_{\mathfrak{g},\mathfrak{h}}).$

$$
\nabla^{4}\Big(\sum_{i,j,l}\mu_{ijl}A_{i}^{*}\otimes(\nu_{l}\wedge\lambda_{j})+\sum_{i\leq j}\beta_{ijl}\big(D_{i}^{*}\otimes(\nu_{l}\wedge\lambda_{j})+D_{j}^{*}\otimes(\nu_{l}\wedge\lambda_{i})\big)+\sum_{i,j}1\otimes\xi_{ij}(\rho(\widetilde{B}_{i})\wedge\nu_{j})\Big)=\sum_{i,j,l}-\mu_{ijl}(A_{i}^{*}\vee D_{j}^{*})\otimes\nu_{l}+\sum_{i\leq j}-2\beta_{ijl}(D_{i}^{*}\vee D_{j}^{*})\otimes\nu_{l}+\sum_{i,j}\xi_{ij}\widetilde{B}_{i}|_{\mathfrak{h}\times\mathfrak{h}}\otimes\nu_{j}=\sum_{k}\Big(\sum_{i,l}-\mu_{ilk}(A_{i}^{*}\vee D_{l}^{*})+\sum_{i\leq j}-2\beta_{ijk}(D_{i}^{*}\vee D_{j}^{*})+\sum_{i}\xi_{ik}\widetilde{B}_{i}|_{\mathfrak{h}\times\mathfrak{h}}\Big)\otimes\nu_{k}
$$
\n(3.37)

Since $\{\nu_k \mid 1 \leq k \leq r_0\}$ is a basis of ker $R_{\mathfrak{g},\mathfrak{h}}$, it follows that the left hand side of (3.37) is zero if and only if $\sum_{i,l} -\mu_{ilk}(A_i^* \vee D_l^*) + \sum_{i \leq j} 2\beta_{ijk}(D_i^* \vee D_j^*) + \sum_i \xi_{ik}\widetilde{B}_i|_{\mathfrak{h} \times \mathfrak{h}} = 0$, for all $k = 1, \ldots, r_0$. Now the isomorphism (3.36) follows from (3.15) and Lemma [3.13.](#page-15-2)

The isomorphism in (3.30) follows from (3.33) and (3.36) . This completes the proof. \Box

Lemma 3.16. Let $\mathcal{C}_{\mathfrak{g},H}$ be as in Definition [3.11.](#page-15-0) Then there is an isomorphism between the following two vector spaces.

$$
\frac{S^2(\mathfrak{h}^*)^H \otimes 1}{\nabla^3((\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \oplus 1 \otimes P_{\mathfrak{g}}^3)} \simeq \mathcal{C}_{\mathfrak{g},H}.
$$

Proof. Recall that $\mathfrak{a} = \mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}], \mathfrak{h}^H = \mathfrak{a}^H \oplus \mathfrak{b}, \mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}] = \mathfrak{a} + [\mathfrak{h}, \mathfrak{h}],$ and $(H - 1)\mathfrak{h} = (H - 1)\mathfrak{a} + [\mathfrak{h}, \mathfrak{h}]$; see [\(3.5\)](#page-13-2), [\(3.6\)](#page-13-3). Now using Proposition [3.5](#page-13-4) and Lemma [3.1,](#page-11-3) we have the following decomposition.

$$
S^{2}(\mathfrak{h}^{*})^{H} = S^{2}((\mathfrak{h}^{H})^{*})^{H} \oplus S^{2}(((H-1)\mathfrak{h})^{*})^{H}
$$

= $S^{2}((\mathfrak{h}^{H})^{*}) \oplus S^{2}(((H-1)\mathfrak{h})^{*})^{H}$
= $S^{2}((\mathfrak{a}^{H})^{*}) \oplus S^{2}(\mathfrak{b}^{*}) \oplus (\mathfrak{a}^{H})^{*} \vee \mathfrak{b}^{*} \oplus S^{2}(((H-1)\mathfrak{a})^{*})^{H} \oplus S^{2}([\mathfrak{h}, \mathfrak{h}]^{*})^{H}$
= $S^{2}((\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}])^{*})^{H} \oplus S^{2}(\mathfrak{b}^{*}) \oplus (\mathfrak{a}^{H})^{*} \vee \mathfrak{b}^{*}.$

Let $r := \#[\mathfrak{g}, \mathfrak{g}],$ B_i be as in [\(3.10\)](#page-14-4), and ∇^3 be as in [\(2.15\)](#page-9-2). Since \mathfrak{a}^H , $(H-1)\mathfrak{a}, [\mathfrak{h}, \mathfrak{h}]$ are mutually orthogonal with respect to \hat{B}_i for all $i = 1, ..., r$, and $R_{\mathfrak{g},\mathfrak{h}}(P_{\mathfrak{g}}^1) = \mathfrak{b}^*$, we also have

$$
\nabla^3 ((\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus 1 \otimes P_{\mathfrak{g}}^3) =
$$

$$
((\mathfrak{a}^H)^* \vee \mathfrak{b}^* \oplus S^2(\mathfrak{b}^*) \oplus \left\{ \sum_i \eta_i \widetilde{B}_i |_{(\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}]) \times (\mathfrak{h} \cap [\mathfrak{g} \mathfrak{g}])} \mid (\eta_1, \ldots, \eta_r) \in \mathbb{R}^r \right\} \otimes 1.
$$

Now the proof follows from the fact that $\left\{ \sum_i \eta_i \widetilde{B}_i |_{(\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}]) \times (\mathfrak{h} \cap [\mathfrak{g} \mathfrak{g}])} \mid (\eta_1, \ldots, \eta_r) \in \mathbb{R}^r \right\} \subset$ $S^2((\mathfrak{h} \cap [\mathfrak{g},\,\mathfrak{g}])^*)^H$. The contract of the contract of the contract of \Box

Proof of Theorem [1.2.](#page-2-0) Recall the Koszul complex $(\mathscr{C}^* := S(\mathfrak{h}^*)^H \otimes \wedge P_{\mathfrak{g}}, \nabla)$ as defined in Section [2.2.4.](#page-9-0) Using Theorem [6.15,](#page-41-8) we have the following isomorphism:

$$
H^4(G/H, \mathbb{R}) \simeq \frac{\ker \nabla^4}{\operatorname{Im} \nabla^3},\tag{3.38}
$$

where the differentials ∇^4 and ∇^3 are as in [\(2.16\)](#page-10-3) and [\(2.15\)](#page-9-2), respectively. The numerator and the denominator in [\(3.38\)](#page-21-0) will be identified. Recall that

$$
\mathscr{C}^4 = S^2(\mathfrak{h}^*)^H \otimes 1 \bigoplus (\mathfrak{h}^*)^H \otimes \wedge^2 P_{\mathfrak{g}}^1 \bigoplus 1 \otimes P_{\mathfrak{g}}^3 \wedge P_{\mathfrak{g}}^1 \bigoplus 1 \otimes \wedge^4 P_{\mathfrak{g}}^1,
$$

$$
{}^*M \otimes P_{\mathfrak{g}}^1 \bigoplus 1 \otimes P_{\mathfrak{g}}^3 \otimes P_{\mathfrak{g}}^2 \otimes P_{\mathfrak{g}}^2 \otimes P_{\mathfrak{g}}^1 \otimes P_{\mathfrak{g}}^1 \otimes P_{\mathfrak{g}}^1 \otimes P_{\mathfrak{g}}^2 \otimes P_{\mathfrak{g}}^1,
$$

 $\nabla^3((\mathfrak{h}^*)^H \otimes P_{\mathfrak{g}}^1 \bigoplus 1 \otimes P_{\mathfrak{g}}^3) \subset S^2(\mathfrak{h}^*)^H \otimes 1$, and $\nabla^3(1 \otimes (\wedge^3 P_{\mathfrak{g}}^1)) \subset (\mathfrak{h}^*)^H \otimes (\wedge^2 P_{\mathfrak{g}}^1)$.

Thus, the right hand side of (3.38) is isomorphic to the following vector space:

$$
\frac{S^2(\mathfrak{h}^*)^H\otimes 1}{\nabla^3\big((\mathfrak{h}^*)^H\otimes P_\mathfrak{g}^1\oplus 1\otimes P_\mathfrak{g}^3\big)}\bigoplus\frac{\ker \nabla^4\big|_{(\mathfrak{h}^*)^H\otimes \big(\wedge^2 P_\mathfrak{g}^1\big)\oplus 1\otimes (P_\mathfrak{g}^3\wedge P_\mathfrak{g}^1)}{\nabla^3(1\otimes \wedge^3 P_\mathfrak{g}^1)}\bigoplus\ker \nabla^4\big|_{\big(1\otimes (\wedge^4 P_\mathfrak{g}^1)\big)}\,.
$$

Now the proof of the theorem follows from Lemma [3.16;](#page-21-1) [\(3.27\)](#page-18-1), [\(3.28\)](#page-18-2), Lemma [3.15;](#page-19-0) and Lemma $3.6(1)$ $3.6(1)$.

3.4. Second and first cohomologies revisited. We end this section by recording short proofs of our earlier results on the first and second cohomology of compact homogeneous spaces; see [\[BCM,](#page-41-0) Theorem 3.6, Theorem 3.3].

First we will reprove [\[BCM,](#page-41-0) Theorem 3.6]. In view of Theorem [6.15,](#page-41-8) we have

$$
H^1(G/H) \simeq \ker \nabla^1 \simeq \left(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\right)^*, \qquad \text{(using (3.3))}.
$$

Next we will reprove [\[BCM,](#page-41-0) Theorem 3.3].

Theorem 3.17. Let G be a compact connected Lie group, and H be a closed subgroup of G. Then

$$
H^2(G/H, \mathbb{R}) \simeq ((\mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}])^*)^{H/H^0} \bigoplus \wedge^2 \Big(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\Big)^*.
$$

Proof. Recall that $(\mathfrak{h}^*)^H = (\mathfrak{a}^*)^H \oplus \mathfrak{b}^*$ and $\text{Im } \nabla^1 = \mathfrak{b}^* \otimes 1 \subset (\mathfrak{h}^*)^H \otimes 1$; see Lemma [3.3.](#page-12-3) Then using Theorem [6.15,](#page-41-8) the following is deduced.

$$
H^2(G/H, \mathbb{R}) \simeq \frac{\ker \nabla^2}{\operatorname{Im} \nabla^1} = \frac{(\mathfrak{h}^*)^H \otimes 1}{\operatorname{Im} \nabla^1} \bigoplus \ker (\nabla^2|_{1 \otimes \wedge^2 P^1_{\mathfrak{g}}}) \simeq (\mathfrak{a}^*)^H \bigoplus \ker (\nabla^2|_{1 \otimes \wedge^2 P^1_{\mathfrak{g}}}).
$$

Since H^0 acts trivially on $\mathfrak{z}(\mathfrak{h})$, the theorem follows from [\(2.19\)](#page-10-2) and Lemma [3.6\(](#page-13-5)1). \Box

4. Applications of the main results

In this section, we apply our main results, namely, Theorem [1.1,](#page-1-0) and Theorem [1.2](#page-2-0) in various special cases of homogeneous spaces which occur naturally and record some interesting consequences.

4.1. Compact homogeneous spaces. First we will consider homogeneous spaces of connected compact Lie groups. The proofs of most of these results are fairly straightforward from our main theorems, Lemma [4.4](#page-24-0) and Proposition [4.5](#page-25-0) below. Hence they are either briefly sketched or omitted.

The notations $\mathcal{N}_{\mathfrak{g},\mathfrak{h}}$ and $\mathcal{C}_{\mathfrak{g},H}$ as in Definition [3.11](#page-15-0) will be used in this section. We next introduce some additional notation which will also be used.

Definition 4.1. Let G be a compact Lie group, which is not necessarily connected. Let $\mathfrak{a} \subset [\mathfrak{g}, \mathfrak{g}]$ be an ideal of \mathfrak{g} . We denote the set of simple factors of \mathfrak{a} by $\mathcal{I}(\mathfrak{a})$, and the number of simple factors in α by $\#\alpha$. Moreover, if α is invariant under the adjoint action of G then we let $\#(\mathfrak{a}, G)$ denote the number of non-zero minimal G-invariant ideals of α .

Corollary 4.2. Let G be a compact connected Lie group, and H a closed subgroup of G .

(1) Assume that H is a subgroup of
$$
(G, G)
$$
, the commutator subgroup of G. Then

$$
H^3(G/H, \mathbb{R}) \simeq (\mathfrak{z}(\mathfrak{h})^*)^{H/H^0} \otimes \mathfrak{z}(\mathfrak{g})^* \bigoplus \mathcal{N}_{\mathfrak{g},\mathfrak{h}} \bigoplus \wedge^3 \mathfrak{z}(\mathfrak{g})^*,
$$

$$
H^4(G/H, \mathbb{R}) \simeq (\mathfrak{z}(\mathfrak{h})^*)^{H/H^0} \otimes \wedge^2 \mathfrak{z}(\mathfrak{g})^* \bigoplus (\mathcal{N}_{\mathfrak{g},\mathfrak{h}})^{\dim_{\mathbb{R}} \mathfrak{z}(\mathfrak{g})} \bigoplus \mathcal{C}_{\mathfrak{g},H} \bigoplus \wedge^4 \mathfrak{z}(\mathfrak{g})^*.
$$

(2) Recall that $Z(H)$ denotes the center of the Lie group H. Let $r_0 := \dim_{\mathbb{R}} \left(\frac{\mathfrak{g}}{\lceil \mathfrak{a}, \mathfrak{a} \rceil} \right)$ $\frac{\mathfrak{g}}{[\mathfrak{g},\mathfrak{g}]+\mathfrak{h}}\Big).$ Assume that $Z(H) \cap (G, G)$ is a finite group. Then we have the following isomorphisms.

$$
H^{3}(G/H, \mathbb{R}) \simeq \mathcal{N}_{\mathfrak{g},\mathfrak{h}} \bigoplus \wedge^{3} \left(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\right)^{*},
$$

$$
H^{4}(G/H, \mathbb{R}) \simeq (\mathcal{N}_{\mathfrak{g},\mathfrak{h}})^{r_{0}} \bigoplus \mathcal{C}_{\mathfrak{g},H} \bigoplus \wedge^{4} \left(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\right)^{*}.
$$

(3) Assume that $\dim_{\mathbb{R}} \mathfrak{z}(\mathfrak{g}) \leq 1$. Then we have the following isomorphisms.

$$
H^3(G/H, \mathbb{R}) \simeq ((\mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}])^*)^{H/H^0} \otimes \left(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\right)^* \bigoplus \mathcal{N}_{\mathfrak{g}, \mathfrak{h}},
$$

$$
H^4(G/H, \mathbb{R}) \simeq \begin{cases} \mathcal{C}_{\mathfrak{g}, H} & \text{if } [\mathfrak{g}, \mathfrak{g}] + \mathfrak{h} = \mathfrak{g} \\ \mathcal{N}_{\mathfrak{g}, \mathfrak{h}} \bigoplus \mathcal{C}_{\mathfrak{g}, H} & \text{if } [\mathfrak{g}, \mathfrak{g}] + \mathfrak{h} \subsetneqq \mathfrak{g} .\end{cases}
$$

Remark 4.3. In this remark, we will indicate a natural situation where Corollary [4.2\(](#page-22-4)[3\)](#page-23-0) can be applied. Let L be a connected simple Lie group which is not necessarily compact, and M a closed subgroup with finitely many connected components. Let G be a maximal compact subgroup of L such that $H := G \cap M$ is a maximal compact subgroup of M. Then $H^{i}(L/M,\mathbb{R}) \simeq H^{i}(G/H,\mathbb{R})$ for all *i*; see [\(4.2\)](#page-27-2). It is well-known that maximal compact subgroups of a simple real Lie group need not be semisimple, but the center of a maximal compact subgroup is of dimension at the most one; see [\[He,](#page-42-7) Proposition 6.2, p. 382]. Thus $\dim_{\mathbb{R}} \mathfrak{z}(\mathfrak{g}) \leq 1$, and hence to calculate $H^{i}(L/M, \mathbb{R})$, $i = 3, 4$ we may invoke Corollary [4.2\(](#page-22-4)[3\)](#page-23-0).

Let G be a compact Lie group which is not necessarily connected. Then recall that the adjoint action of G on $\mathfrak g$ keeps the center $\mathfrak z(\mathfrak h)$ and the commutator $[\mathfrak g, \mathfrak g]$ invariant. As in [\(3.9\)](#page-14-5) we assume that $r := \#[\mathfrak{g}, \mathfrak{g}]$ and that $\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_r$ where $\mathfrak{g}_i, i = 1, \ldots, r$ are all the simple factors of $[\mathfrak{g}, \mathfrak{g}]$. Let $\mathfrak{p}_1, \ldots, \mathfrak{p}_k$ denote all the *isotypical* (semisimple) factors of $[\mathfrak{g}, \mathfrak{g}]$ (for some $k \leq r$) then \mathfrak{p}_i also remain invariant under the adjoint action of G, for all $j = 1, \ldots, k$. Without loss of generality we may further assume that there exists integers $0 < l_1 < \cdots < l_{k-1} < l_k = r$ such that

$$
\mathfrak{p}_1 = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_{l_1}, \, \cdots, \, \mathfrak{p}_k = \mathfrak{g}_{l_{k-1}+1} \oplus \cdots \oplus \mathfrak{g}_{l_k}.
$$

Then the adjoint action of G on \mathfrak{p}_i permutes all the simple factors thereof, and thus, induces a natural action on the finite set $\mathcal{I}(\mathfrak{p}_i)$. Further observe that the restricted action of G^0 on $\mathcal{I}(\mathfrak{p}_i)$ is trivial, and hence the finite group G/G^0 has a natural action on $\mathcal{I}(\mathfrak{p}_i)$ for all $j = 1, ..., k$. One may see that, for all $j = 1, ..., k$, the number $\#(\mathfrak{p}_j, G)$ coincides with the number of distinct G/G^0 -orbits in $\mathcal{I}(\mathfrak{p}_j)$, for all $j = 1, \ldots, r$. One also has

$$
\#([\mathfrak{g},\mathfrak{g}],G) = \#(\mathfrak{p}_1,G) + \cdots + \#(\mathfrak{p}_k,G). \tag{4.1}
$$

Note that when G is either connected or all the simple factors of $[\mathfrak{g}, \mathfrak{g}]$ are pairwise nonisomorphic then $\#([\mathfrak{g}, \mathfrak{g}], G) = \#[\mathfrak{g}, \mathfrak{g}]$. It may be further recalled that the adjoint action of G on g also induces a natural action on each of the spaces $S^2(\mathfrak{z}(\mathfrak{g})^*), S^2(\mathfrak{p}_1^*), \ldots, S^2(\mathfrak{p}_k^*).$ Let B_i , $i = 1, \ldots, r$ be defined as in [\(3.10\)](#page-14-4). In the next lemma, we observe certain facts on the space of invariant symmetric bilinear forms on the Lie algebra of a compact Lie group.

- (1) $S^2(\mathfrak{g}^*)^{G^0} = S^2(\mathfrak{z}(\mathfrak{g})^*) \bigoplus S^2(\mathfrak{g}_1^*)^{G^0} \oplus \cdots \oplus S^2(\mathfrak{g}_r^*)^{G^0}.$ (a) Moreover, for all $i = 1, \ldots, r$, the space $S^2(\mathfrak{g}_i^*)^{G^0}$ is the R-span of the Killing form on \mathfrak{g}_i and hence $\dim_{\mathbb{R}} S^2(\mathfrak{g}_i^*)^{G^0} = 1$.
	- (b) Let B_i be as in [\(3.10\)](#page-14-4). Then $\{B_i \mid 1 \leq i \leq \#[\mathfrak{g},\mathfrak{g}]\}$ form a basis of $S^2([{\mathfrak g},\,{\mathfrak g}]^*)^{G^0}.$
- (2) $S^2(\mathfrak{g}^*)^G = S^2(\mathfrak{z}(\mathfrak{g})^*)^{G/G^0} \bigoplus S^2(\mathfrak{p}_1^*)^G \oplus \cdots \oplus S^2(\mathfrak{p}_k^*)^G$, and $\dim_{\mathbb{R}} S^2(\mathfrak{p}_j^*)^G = \#(\mathfrak{p}_j, G)$, for all $i = 1, \ldots, k$.
- (3) dim $S^2([\mathfrak{g}, \mathfrak{g}]^*)^G = \#([\mathfrak{g}, \mathfrak{g}], G)$, and dim $S^2(\mathfrak{g}^*)^G = \dim S^2(\mathfrak{z}(\mathfrak{g})^*)^{G/G^0} + \#([\mathfrak{g}, \mathfrak{g}], G)$.
- (4) Let G be either connected or all the simple factors of $[\mathfrak{g}, \mathfrak{g}]$ are pairwise nonisomorphic. Then $\dim S^2(\mathfrak{g}^*)^G = \dim S^2(\mathfrak{z}(\mathfrak{g})^*)^{G/G^0} + \#[\mathfrak{g}, \mathfrak{g}]$. In particular, for simple G^0 , dim $S^2(\mathfrak{g}^*)^G = 1$.

Proof. We first prove (1). Observe that if $B \in S^2(\mathfrak{g}^*)^{G^0}$ then B is also \mathfrak{g} -invariant. As \mathfrak{g}_i is a simple ideal for all i, by Lemma [3.1,](#page-11-3) it is immediate that $B(\mathfrak{g}_i, \mathfrak{g}_j) = 0 = B(\mathfrak{z}(\mathfrak{g}), \mathfrak{g}_k)$, for all i, j with $i \neq j$ and for all k. Thus we have $S^2(\mathfrak{g}^*)^{G^0} = S^2(\mathfrak{z}(\mathfrak{g})^*) \bigoplus S^2(\mathfrak{g}_1^*)^{G^0} \oplus \cdots \oplus S^2(\mathfrak{g}_n^*)^{G^0}$ $S^2(\mathfrak{g}_r^*)^{G^0}.$

To see a proof of [\(1a\)](#page-24-1), first observe that compactness of G forces the commutator $[\mathfrak{g}, \mathfrak{g}]$ to be a compact Lie algebra. Thus the simple factor \mathfrak{g}_i is also compact, and this in turn implies \mathfrak{g}_i to be absolutely simple, that is, $\mathbb{C} \otimes_{\mathbb{R}} \mathfrak{g}_i$ is a simple Lie algebra over \mathbb{C} . Thus by Schur's lemma any G^0 -invariant symmetric bilinear form on \mathfrak{g}_i is a (real) scalar multiple of the Killing form on \mathfrak{g}_i . The proof now follows from the fact that the Killing form B_i on \mathfrak{g}_i is G^0 -invariant.

The proof of $(1b)$ follows from (1) and $(1a)$.

We next prove (2). To prove the first part of (2) we use Lemma [3.1](#page-11-3) and the proof follows analogously as the proof of the first part of (1). We next prove the second part of (2), that is, $\dim_{\mathbb{R}} S^2(\mathfrak{p}_j^*)^G = \#(\mathfrak{p}_j, G)$, for all $j = 1, \ldots, k$. It is enough to prove the above assertion when $j = 1$, that is, $\dim_{\mathbb{R}} S^2(\mathfrak{p}_1^*)^G = \#(\mathfrak{p}_1, G)$. We have $\mathcal{I}(\mathfrak{p}_1) := {\mathfrak{g}_1, \ldots, \mathfrak{g}_{l_1}}$. As mentioned before, the adjoint action of G on $\mathfrak g$ induces an action of G on the set $\mathcal{I}(\mathfrak{p}_1)$. Thus one has an action of G on the set $\{1,\ldots,l_1\}$ such that $\text{Ad}(g)\mathfrak{g}_i = \mathfrak{g}_{g \cdot i}$, for all $i = 1, \ldots, l_1$ and for all $g \in G$. It is clear that $n := \#(\mathfrak{p}_1, G)$ is the number of orbits of the action of G on $\{1, \ldots, l_1\}$. Let S_1, \ldots, S_n denote all the G-orbits in $\{1, \ldots, l_1\}$. The assertion now follows from the claim:

$$
S^{2}(\mathfrak{p}_{1}^{*})^{G} = \left\{ \lambda_{1} \sum_{i_{1} \in S_{1}} \widetilde{B}_{i_{1}} + \cdots + \lambda_{n} \sum_{i_{n} \in S_{n}} \widetilde{B}_{i_{n}} \mid \lambda_{1}, \ldots, \lambda_{n} \in \mathbb{R} \right\}.
$$

To prove the claim we first fix a compact Lie algebra a, which is simple, such that there is an isomorphism $\delta_i : \mathfrak{g}_i \to \mathfrak{a}$ for all $i = 1, \ldots, l_1$. Let $\pi_i : \mathfrak{g} \to \mathfrak{g}_i$ be the *i*-th projection with respect to the decomposition $\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \bigoplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_r$. Let B be the Killing form on **a**. Let $\pi'_i := \delta_i \circ \pi_i$, for all $i = 1, \ldots, l_1$. Then it follows that $\overline{B}_i = B \circ (\pi'_i \times \pi'_i)$, for all $i = 1, \ldots, l_1$. Using this it is now straightforward to complete the proof of the claim.

The proof of (3) is immediate from (2).

The proof of (4) is immediate from (3) once we recall that G acts on all the sets $\mathcal{I}(\mathfrak{p}_1), \ldots, \mathcal{I}(\mathfrak{p}_k)$ trivially either if G is connected or if all the sets $\mathcal{I}(\mathfrak{p}_1), \ldots, \mathcal{I}(\mathfrak{p}_k)$ are \Box singletons.

Proposition 4.5. Let G be a compact, connected Lie group, $H \subset G$ a closed subgroup. Let $r := #[\mathfrak{g}, \mathfrak{g}],$ and let $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_r$, where \mathfrak{g}_i 's are simple ideals in $[\mathfrak{g}, \mathfrak{g}].$

- (1) If $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}] \neq 0$, then $\dim_{\mathbb{R}} \mathcal{N}_{\mathfrak{g},\mathfrak{h}} \leq \#[\mathfrak{g}, \mathfrak{g}]-1$. Furthermore, if H^0 is simple then $\dim_{\mathbb{R}}\mathcal{N}_{\mathfrak{g},\mathfrak{h}} = \#[\mathfrak{g},\,\mathfrak{g}]-1,\ and\ \dim_{\mathbb{R}}\mathcal{C}_{\mathfrak{g},H} = 0.$
- (2) If $k := #\{i \mid \mathfrak{h} \cap \mathfrak{g}_i \neq 0\}$, then $\dim_{\mathbb{R}} \mathcal{N}_{\mathfrak{g},\mathfrak{h}} \leq #[\mathfrak{g}, \mathfrak{g}] k$.
- (3) If $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$ for some $s \leq \#[\mathfrak{g}, \mathfrak{g}]$, then $\dim_{\mathbb{R}} \mathcal{N}_{\mathfrak{g}, \mathfrak{h}} \geq \#[\mathfrak{g}, \mathfrak{g}] - s$.
- (4) If $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$, and $\mathfrak{h} \cap \mathfrak{g}_i \neq 0$ for all $i = 1, \ldots, s$, then $(a) \dim_{\mathbb{R}} \mathcal{N}_{\mathfrak{g},\mathfrak{h}} = \#[\mathfrak{g}, \mathfrak{g}] - s.$ (b) $\dim_{\mathbb{R}} \text{Im } \Psi_{\mathfrak{g},\mathfrak{h}} = s$. In particular, $s \leq \dim S^2((\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}])^*)^H$.

Proof. Proof of (1). The proof follows from the fact that the map $\Psi_{\mathfrak{a},\mathfrak{h}}$ as in [\(3.13\)](#page-15-3) is a non-zero map and the Killing form on $[g, g]$ is negative-definite.

To prove the second part of (1) first observe that, as h is simple and $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}] \neq 0$, $[\mathfrak{g}, \mathfrak{g}] \neq 0$. Further, as H^0 is simple, by Lemma [4.4](#page-24-0) (4), dim $S^2(\mathfrak{h}^*)^H = 1$. Let B be the Killing from of $[\mathfrak{g}, \mathfrak{g}]$. Let $0 \neq X \in \mathfrak{h}$. As B is negative definite, $B(X, X) \neq 0$. This forces $\Psi_{\mathfrak{g},\mathfrak{h}}$ to be non-zero and hence a surjective map. Thus $\dim_{\mathbb{R}}\mathcal{N}_{\mathfrak{g},\mathfrak{h}} = \#[\mathfrak{g},\mathfrak{g}] - 1$, and dim_R $C_{\mathfrak{g},H} = 0$.

Proof of (2). Recall that ${B_i \mid 1 \leq i \leq \#[g, g]}$ form a basis of $S^2([\mathfrak{g}, \mathfrak{g}]^*)^G$ where B_i is as in [\(3.10\)](#page-14-4); see Lemma [4.4\(](#page-24-0)[1b\)](#page-24-2). Let $\mathfrak{g}_{i_l}, l = 1, \ldots, k$ be distinct ideals of $[\mathfrak{g}, \mathfrak{g}]$ such that $\mathfrak{h} \cap \mathfrak{g}_{i_l} \neq 0$. We choose $0 \neq X_{i_l} \in \mathfrak{h} \cap \mathfrak{g}_{i_l}$ for $1 \leq l \leq k$. Since B_{i_l} is negative-definite on \mathfrak{g}_{i_l} we have that $B_{i_l}(X_{i_l}, X_{i_l}) = B_{i_l}(X_{i_l}, X_{i_l}) \neq 0$. Thus $\dim_{\mathbb{R}} \ker \Psi_{\mathfrak{g},\mathfrak{h}} \leq r - k$.

Proof of (3). Since $\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$ for $s \leq r$, we have $B_j|_{(\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}]) \times (\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}])} = 0$ for $j = s + 1, \ldots, r$. Now the first conclusion follows from Lemma [4.4\(](#page-24-0)[1b\)](#page-24-2).

Proof of (4). The statements in (4) follows from (2), (3) and Definition [3.11.](#page-15-0)

Example 4.6. The *n*-dimensional sphere \mathbb{S}^n is diffeomorphic to the homogeneous space $SO(n+1)/SO(n)$. Recall that $SO(3)$, $SO(n)$ for $n > 5$ are simple, where as $SO(4)$ is not simple $(\mathfrak{so}(4) \simeq \mathfrak{su}(2) \oplus \mathfrak{su}(2))$. Thus using Corollary [4.2\(](#page-22-4)[1\)](#page-22-5) and Proposition [4.5,](#page-25-0) it follows that for $i = 3, 4$, $\dim H^i(\mathbb{S}^n) = 1$ if $i = n$, and $\dim H^i(\mathbb{S}^n) = 0$ if $i \neq n$.

Proof of Corollary [1.3.](#page-2-2) The proofs of [\(1\)](#page-2-6) follows from Theorem [1.1,](#page-1-0) and Proposition [4.5\(](#page-25-0)1). The proof of $(1a)$ follows from Proposition [4.5](#page-25-0)[\(4\)](#page-25-1).

The first part of [\(2\)](#page-2-7) follows from Theorem [1.2.](#page-2-0) Second part follows using Lemma [4.4\(](#page-24-0)4) and the fact that $\Psi_{\mathfrak{g},\mathfrak{h}}$ as in [\(3.13\)](#page-15-3) is a non-zero map. [\(2a\)](#page-2-8) follows from above as \mathfrak{h} is semisimple. For [\(2b\)](#page-2-5), use the fact that dim Im $\Psi_{\mathfrak{g},\mathfrak{h}} = 1$, and dim $S^2(\mathfrak{h}^*)^H = \#(\mathfrak{h}, H)$. For $(2c)$, use Proposition [4.5\(](#page-25-0)1).

The proofs of (3) and (5) follow from (1) , (2) and Lemma [4.4.](#page-24-0) The proof of (4) follows from (3) .

If G is a general compact connected Lie group and H is a closed subgroup, then the finite group H/H^0 has a natural right action on G/H^0 turning it into a H/H^0 principal bundle over G/H . Thus one has the following inequality of the Betti numbers in all dimensions: dim_R $H^i(G/H,\mathbb{R}) = \dim_{\mathbb{R}} H^i(G/H^0,\mathbb{R})^{H/H^0} \leq \dim_{\mathbb{R}} H^i(G/H^0,\mathbb{R})$ for all i. One may observe, in view of Theorem [1.1](#page-1-0) and Theorem [1.2,](#page-2-0) that the equality does not hold even in dimensions three and four; see Example [4.7](#page-26-0) below.

Example 4.7. Let $G := U(2) \times \mathbb{S}^1$, and $H := O(2) \times 1 \subset G$. Then $\mathfrak{z}(\mathfrak{h}) = \mathfrak{so}(2) \subset [\mathfrak{g}, \mathfrak{g}],$ and $\overline{O(2)}/SO(2) = \mathbb{Z}/2\mathbb{Z}$ acts non-trivially on $\mathfrak{so}(2)$. In this case dim $\mathcal{N}_{\mathfrak{g},\mathfrak{h}} = 0$ $\dim \mathcal{C}_{\mathfrak{g},H}$. Using Theorem [1.1](#page-1-0) and Theorem [1.2,](#page-2-0) we conclude that $\dim H^3(G/H, \mathbb{R}) =$ $0 \neq \dim H^3(G/H^0, \, \mathbb{R}) \, = \, 2, \text{ and } \dim H^4(G/H, \, \mathbb{R}) \, = \, 0 \, \neq \dim H^4(G/H^0, \, \mathbb{R}) \, = \, 1. \quad \Box$

However, if H is further assumed to be finite, it is not difficult to see that the equality of Betti numbers does hold: If H is finite then $H^0 = e$, and by the connectedness of G it follows that $H^i(G/H^0, \mathbb{R}) \simeq H^i(G, \mathbb{R}) \simeq H^i(G, \mathbb{R})^H \simeq H^i(G/H, \mathbb{R})$. Interestingly, when G is semisimple, in the next corollary we can draw analogous conclusions regarding the three and four dimensional Betti numbers of G/H and $G/H⁰$.

Corollary 4.8. Let G be a compact semisimple Lie group, and H be a closed subgroup. Then the following hold:

- (1) $H^3(G/H, \mathbb{R}) \simeq H^3(G/H^0, \mathbb{R}).$
- (2) If H^0 is a semisimple Lie group and all the simple factors of H^0 are pairwise non-isomorphic, then $H^4(G/H, \mathbb{R}) \simeq H^4(G/H^0, \mathbb{R})$.

Proof. The proof of (1) follows immediately from Corollary [1.3\(](#page-2-2)1). The proof of (2) follows from Corollary [1.3\(](#page-2-2)2) and Lemma [4.4\(](#page-24-0)4).

Proof of Corollary [1.4.](#page-3-1) Using Lemma [4.4,](#page-24-0) it follows that $\dim_{\mathbb{R}} S^2([\mathfrak{g}, \mathfrak{g}]^*)^G = r$ and $\dim_{\mathbb{R}} S^2(\mathfrak{h}^*)^H = \#(\mathfrak{h}, H)$. In view of Proposition [4.5](#page-25-0)[\(4\)](#page-25-1), we have $\dim_{\mathbb{R}} \mathcal{N}_{\mathfrak{g}, \mathfrak{h}} = r - s$. Thus $\dim_{\mathbb{R}} C_{\mathfrak{g},H} = \#(\mathfrak{h},H) - s$. Now proof of the results follows from Theorems [1.1,](#page-1-0) [1.2.](#page-2-0)

Remark 4.9. In this remark, we will illustrate a natural situation where Corollary [1.4](#page-3-1) can be applied. Let G be as in Corollary [1.4.](#page-3-1) Let $H \subset G$ be a closed subgroup. Suppose there is a s with $s \leq r$ such that \mathfrak{h} can be decomposed as $\mathfrak{h} = \mathfrak{h}_1 \oplus \cdots \oplus \mathfrak{h}_s$ where $\mathfrak{h}_i, i = 1, \ldots, s$ are semisimple (not necessarily simple) ideals of \mathfrak{h} with $0 \neq \mathfrak{h}_i \subset \mathfrak{g}_i$ for all $i = 1, \ldots, s$. Then one may apply Corollary [1.4](#page-3-1) to obtain the third and fourth Betti numbers of G/H .

The next result follows immediately from Corollary [1.4.](#page-3-1)

Corollary 4.10. Let G be a compact, connected Lie group, $H \subset G$ a closed subgroup so that H^0 is a simple Lie group. Let $l := \dim_{\mathbb{R}} \mathfrak{z}(\mathfrak{g})$, and $r := \#[\mathfrak{g}, \mathfrak{g}]$. Then

(1)
$$
\dim_{\mathbb{R}}(H^3(G/H, \mathbb{R})) = r - 1 + {l \choose 3},
$$

(2) $\dim_{\mathbb{R}}(H^4(G/H, \mathbb{R})) = l(r - 1) + {l \choose 4}.$

Proof of Corollary [1.7.](#page-3-3) Observe that G/H^0 is compact orientable and $\dim_{\mathbb{R}} G/H^0 =$ 7. Using Poincaré duality, we have $\dim_{\mathbb{R}} H^3(G/H^0) = \dim_{\mathbb{R}} H^4(G/H^0)$. Now we apply Corollary [1.3\(](#page-2-2)4) to conclude that $\#\mathfrak{g} = \#\mathfrak{h}$. **Remark 4.11.** Let G be a compact connected Lie group and H be a closed subgroup of G so that G/H is orientable. Assume further that $\dim_{\mathbb{R}}(G/H) \leq 9$. Then using [\[BCM,](#page-41-0) Theorem 3.6, Theorem 3.3, Theorem [1.1,](#page-1-0) Theorem [1.2,](#page-2-0) and Poincaré duality one may compute dimensions of all the cohomologies of G/H .

4.2. Homogeneous spaces which are not necessarily compact. In this section, we draw some consequences in the setting of homogeneous spaces of Lie groups which are not necessarily compact. We need a result due to Mostow using which computation of the cohomologies of a general connected homogeneous space boils down to that of certain compact homogeneous (sub)space.

Theorem 4.12 ([\[Mo\]](#page-42-8)). Let L be a connected Lie group, and let $M \subset L$ be a closed subgroup with finitely many connected components. Let G be a maximal compact subgroup of L such that $G \cap M$ is a maximal compact subgroup of M. Then the image of the natural embedding $G/(G \cap M) \hookrightarrow L/M$ is a deformation retraction of L/M .

Theorem [4.12](#page-27-0) is proved in $[M_0, p. 260,$ Theorem 3.1 under the assumption that M is connected. However, as mentioned in $[BC]$, using $[Ho, p. 180, Theorem 3.1]$, the proof as in $[M_o]$ goes through when H has finitely many connected components.

Let L, M, G be as in Theorem [4.12,](#page-27-0) and let $H := G \cap M$. As $G/H \hookrightarrow L/M$ is a deformation retraction by Theorem [4.12,](#page-27-0) we have

$$
H^{i}(L/M, \mathbb{R}) \simeq H^{i}(G/H, \mathbb{R}) \quad \text{for all } i. \tag{4.2}
$$

In view of Theorem [4.12](#page-27-0) we next recast Theorem [1.1](#page-1-0) and Theorem [1.2](#page-2-0) in the setting of a general connected homogeneous space which is not necessarily compact.

Theorem 4.13. Let L be a connected Lie group and $M \subset L$ a closed subgroup with finitely many connected components. Let G be a maximal compact subgroup of L, and H a maximal compact subgroup of M with $H \subset G$. Then there is an isomorphism:

$$
H^3(L/M,\,\mathbb{R})\,\simeq\,\big((\mathfrak{z}(\mathfrak{h})\cap [\mathfrak{g},\,\mathfrak{g}])^*\big)^{H/H^0}\otimes\Big(\frac{\mathfrak{g}}{[\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h}}\Big)^*\,\bigoplus\,\mathcal{N}_{\mathfrak{g},\mathfrak{h}}\,\bigoplus\,\wedge^3\Big(\frac{\mathfrak{g}}{[\mathfrak{g},\,\mathfrak{g}]+\mathfrak{h}}\Big)^*\,.
$$

Theorem 4.14. Let L, M, G and H be as in Theorem [4.13.](#page-27-3) Let $r_0 := \dim_{\mathbb{R}} \left(\frac{\mathfrak{g}}{\lceil \mathfrak{a}, \mathfrak{a} \rceil} \right)$ $\frac{\mathfrak{g}}{[\mathfrak{g},\mathfrak{g}]+\mathfrak{h}}\Big).$ Then there is an isomorphism:

$$
H^{4}(L/M, \mathbb{R}) \simeq ((\mathfrak{z}(\mathfrak{h}) \cap [\mathfrak{g}, \mathfrak{g}])^{*})^{H/H^{0}} \otimes \wedge^{2} \Big(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\Big)^{*} \bigoplus (\mathcal{N}_{\mathfrak{g}, \mathfrak{h}})^{r_{0}}
$$

$$
\bigoplus \mathcal{C}_{\mathfrak{g}, H} \bigoplus \wedge^{4} \Big(\frac{\mathfrak{g}}{[\mathfrak{g}, \mathfrak{g}] + \mathfrak{h}}\Big)^{*}.
$$

We will now restrict our attention to certain special types of (non-compact) semisimple Lie groups to obtain better results. We first define a notion associated to a real simple Lie group which will facilitate formulating the next set of corollaries. In addition to the foregoing fact regarding the dimension of the center of any maximal compact subgroup, we also need to recall that in a simple Lie group the semisimple part of a maximal compact subgroup is not necessarily simple. However, the number of simple factors of the semisimple part of a maximal compact subgroup in a simple Lie group is at the most two; see [\[Kn,](#page-42-10) Appendix C].

Definition 4.15. Let L be a (connected) simple Lie group. Then L is said to be of Type-I if a maximal compact subgroup of L is a simple Lie group, and L is said to be of Type-II if a maximal compact subgroup of L is a product of two simple Lie groups. See $\S5$ $\S5$ for a complete list of all the Type-I and Type-II simple Lie groups. \Box

Corollary 4.16. Let L be a semisimple group, and M be a semisimple subgroup such that all the simple factors of both L and M are either of Type-I or of Type-II. Assume that L admits exactly r_1 -many simple factors of Type-I and s_1 -many simple factors of Type-II.

- (1) Then dim_R $H^3(L/M, \mathbb{R}) \le (r_1 + 2s_1)$.
- (2) Moreover, if M admits exactly r_2 -many simple factors of Type-I and s_2 -many simple factors of Type-II, then $\dim_{\mathbb{R}} H^4(L/M, \mathbb{R}) \le (r_1 + r_2) + 2(s_1 + s_2) - 1$.

Proof. Let H be a maximal compact subgroup of M, and G be a maximal compact subgroup of L containing H. It is clear from Proposition [5.3](#page-31-0) that $\dim \mathfrak{z}(\mathfrak{g}) \leq 1$, $\dim \mathfrak{z}(\mathfrak{h}) \leq$ 1. Moreover, we have that $\#[\mathfrak{g}, \mathfrak{g}] = r_1 + 2s_1$, and $\#[\mathfrak{h}, \mathfrak{h}] = r_2 + 2s_2$. We now use Proposition [4.5](#page-25-0) (1) to conclude that $\dim \mathcal{N}_{\mathfrak{g},\mathfrak{h}} \leq r_1 + 2s_1 - 1$, and Lemma [4.4](#page-24-0) (4) to conclude that $\dim S^2((\mathfrak{h} \cap [\mathfrak{g}, \mathfrak{g}])^*)^H \leq 1 + \#[\mathfrak{h}, \mathfrak{h}]$. Thus $\dim \mathcal{C}_{\mathfrak{g},H} \leq r_2 + 2s_2$; see Definition [3.11.](#page-15-0) We now use Theorem [4.12](#page-27-0) and Corollary [4.2](#page-22-4) [\(3\)](#page-23-0) to see that

$$
\dim H^3(L/M, \mathbb{R}) \le 1 + \dim \mathcal{N}_{\mathfrak{g},\mathfrak{h}} \le r + 2s, \text{ and}
$$

$$
\dim_{\mathbb{R}} H^4(L/M, \mathbb{R}) \le \dim \mathcal{N}_{\mathfrak{g},\mathfrak{h}} + \dim \mathcal{C}_{\mathfrak{g},H} \le r_1 + r_2 + 2(s_1 + s_2) - 1.
$$

We will now derive vanishing results of the third and fourth cohomologies of homogeneous space in certain non-compact setting. In view of Theorem [4.12,](#page-27-0) the next result follows either from $[Az]$ or from Corollary [1.3\(](#page-2-2)[1b\)](#page-2-3).

Corollary 4.17. Let L be a simple Lie group of Type-I. Let M be a closed subgroup of L with finitely many components. Assume that M contains a compact subgroup of positive dimension. Then $H^3(L/M, \mathbb{R}) = 0$.

In the next corollary we generalize the foregoing one.

Corollary 4.18. Let L be a semisimple Lie group, and $M \subset L$ be a closed semisimple subgroup. Assume that both L and M admit semisimple maximal compact subgroups. Let L_1, \ldots, L_r be the simple factors of L, and each L_i is of Type-I. Suppose that $L_i \cap M$ contains a compact subgroup of positive dimension for each $i = 1, \ldots, r$. Then

$$
H^3(L/M,\mathbb{R})\,=\,0\,.
$$

Proof. Let $C_i \subset L_i \cap M$ be a compact subgroup of positive dimension. Then $C :=$ $C_1 \cdots C_r$ is a compact subgroup in M. Let H be a maximal compact subgroup of M containing C , and G be a maximal compact subgroup of L containing H . Then by Corollary [5.4,](#page-32-0) $G = G_1 \cdots G_r$ where G_i is a maximal compact subgroup of L_i for each $i = 1, \ldots, r$. Since L_i is of Type-I, G_i is simple for all i. Next we claim that $\mathfrak{c}_i \subset \mathfrak{g}_i \ \forall i$. To see this note that $\mathfrak{c}_i \subset \mathfrak{g} = \mathfrak{g}_1 + \cdots + \mathfrak{g}_r$. Let $X \in \mathfrak{c}_i$ and $\mathfrak{c}_i \subset \mathfrak{l}_i$. Then $X = X_1 + \cdots + X_r$ for some $X_i \in \mathfrak{g}_i \subset \mathfrak{l}_i$. Thus

$$
X - X_i = X_1 + \cdots X_{i-1} + X_{i+1} + \cdots + X_r.
$$

But $X - X_i \in \mathfrak{l}_i$ and $X_1 + \cdots + X_{i-1} + X_{i+1} + \cdots + X_r \in \mathfrak{l}_1 \oplus \cdots \oplus \mathfrak{l}_{i-1} \oplus \mathfrak{l}_{i+1} \oplus \mathfrak{l}_r$. Hence $X - X_i = 0$ which proves the claim. Thus $c_i \subset g_i \cap \mathfrak{h}$, and hence dim $g_i \cap \mathfrak{h} > 0$. Now in view of Theorem [4.12,](#page-27-0) the proof follows from Corollary [1.4.](#page-3-1)

Corollary 4.19. Let L be a semisimple Lie group, and $M \subset L$ be a closed semisimple subgroup. Assume that both L and M admit semisimple maximal compact subgroups. Let L_1, \ldots, L_r be the simple factors of L, and M_1, \ldots, M_s be the simple factors of M such that $r \geq s$. Assume that $M_i \subset L_i$ and both L_i , M_i are of Type-I for all $i = 1, \ldots, s$. Then

$$
H^4(L/M,\mathbb{R})=0.
$$

Proof. Let $H_i \subset M_i$ be a maximal compact subgroup. Since M_i is of Type-I, H_i is simple for all i. Let $G_i \subset L_i$ be a maximal compact subgroup containing H_i . Set $H := H_1 \cdots H_s \subset M$ and $G := G_1 \cdots G_r \subset L$. Then using Corollary [5.4,](#page-32-0) it follows that H is a maximal compact subgroup of M and G is a maximal compact subgroup of L containing H. Here $\mathfrak{h} \subset \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$ and $\mathfrak{h} \cap \mathfrak{g}_i \neq 0$ for all $i = 1, \ldots s$. Since H is connected and semisimple, $\#(\mathfrak{h}, H) = \# \mathfrak{h} = s$. Now in view of Theorem [4.12,](#page-27-0) the proof follows from Corollary [1.4.](#page-3-1)

Corollary 4.20. Let L be a connected Lie group which admits a semisimple maximal compact subgroup. Let $M \subset L$ be a closed subgroup with finitely many connected components, and M^0 be a simple Lie group of Type-I. Then $H^4(L/M, \mathbb{R}) = 0$.

Proof. In view of Theorem [4.12,](#page-27-0) the proof follows from Corollary [1.3](#page-2-2)[\(2c\)](#page-2-9). \Box

5. Appendix : Maximal compact subgroups and types of simple Lie groups

In this appendix we deal with two points which are required in $\S4.2$ $\S4.2$: in the first part we describe certain subtle issues regarding relations between a maximal compact subgroup of a semisimple groups and those of its simple factors, and in the second part we list all the Type-I, Type-II simple Lie groups.

Lemma 5.1. Let G be a connected Lie group.

- (1) Let A be a connected compact subgroup. Then the coset space G/A is contractible if and only if A is a maximal compact subgroup.
- (2) Let B be a connected closed subgroup such that the coset space G/B is contractible. Then a maximal compact subgroup in B is a maximal compact subgroup in G .

Proof. We first prove (1). If A is a maximal compact subgroup, then by $[H_0,$ Theorem 3.1., pp. 180–181, G/A is contractible. Now suppose that G/A is contractible. We will show that A is a maximal compact subgroup in G . As A is connected compact, there is a maximal compact subgroup H of G such that $A \subset H$. Then by [H₀, Theorem 3.1., pp. 180–181, G/H is contractible. Observe that G/A is a fiber bundle over G/H with fiber H/A . As the base G/H is contractible it follows that $G/A \simeq G/H \times H/A$ as fiber bundles. Thus G/A has the same homotopy type as H/A , and hence H/A is contractible. Recall that a compact manifold without boundary can not be contractible if it is of positive dimension (this is because if M is a compact manifold without boundary and if $n := \dim M$ the *n*-th singular homology with coefficient in \mathbb{Z}_2 satisfies $H_n(M, \mathbb{Z}_2) = \mathbb{Z}_2$. Thus we have that dim $H/A = 0$ which forces $A = H$ as H is connected. Hence A is a maximal compact subgroup in G.

We now prove (2). Let $K \subset B$ be a maximal compact subgroup of B. Note that G/K is a fiber bundle over G/B with fiber B/K . As the base G/B is contractible it follows, as above, that $G/K \simeq G/B \times B/K$ as fiber bundles. As K is a maximal compact subgroup by $[H_0,$ Theorem 3.1., pp. 180–181 B/K is contractible. As G/B is contractible, so is G/K . As K is compact, by (1), it follows that K is a maximal compact subgroup of $G.$

The next lemma is formulated in a slightly more general context, but we need only a special case of it in the proof of Proposition [5.3.](#page-31-0)

Lemma 5.2. Let C be a connected compact Lie group, and let V be a finite dimensional vector space over $\mathbb R$. Let $n := \dim_{\mathbb R} V$. Let $H := V \times C$ be the direct product of the vector group V and C. Let $\Gamma \subset H$ be a discrete central subgroup of H, and let $\theta : H \to H/\Gamma$ be the quotient homomorphism.

- (1) Then Γ is a finitely generated abelian group, and moreover, rank $\Gamma \leq n$.
- (2) Suppose rank $\Gamma = m$. Then there exists closed subgroups $W, T \subset H/\Gamma$ such that W is isomorphic to \mathbb{R}^{n-m} , T is isomorphic to the m-dimensional compact torus $(\mathbb{S}^1)^m$ and $H/\Gamma = W T \theta(C)$. Moreover, any pair of subgroups among the subgroups $W, T, \theta(C)$ commute and intersect in a discrete subgroup. In particular, H/Γ is isomorphic to $W \times (T\theta(C))$, and, consequently, $T\theta(C)$ is a maximal compact subgroup of H/Γ .

Proof. We prove (1) first. As Γ is discrete and central it follows that Γ is finitely generated. Let π : $H = V \times C \rightarrow V$ denote the projection to the first factor. As C is compact, π is a proper map, and hence $\pi(\Gamma)$ is a discrete subgroup of V. Thus rank $\pi(\Gamma) \leq n$. Now it is easy to see that the group of torsion elements of Γ is ker $\pi \cap \Gamma$, and hence rank $\Gamma = \text{rank } \Gamma / (\text{ker } \pi) \cap \Gamma = \text{rank } \pi(\Gamma) \leq n$.

We now prove (2). As rank $\Gamma = \text{rank } \pi(\Gamma) = m$ we may choose elements in Γ , say, $(v_1, c_1), \ldots, (v_m, c_m)$ such that $\{v_1, \ldots, v_m\}$ is a Z-basis of $\pi(\Gamma)$. Since $\pi(\Gamma)$ is a discrete subgroup of V it follows that $\{v_1, \ldots, v_m\}$ is R-linearly independent. Let $W_1 := \mathbb{R}v_1 +$ $\cdots + \mathbb{R}v_m$, and let W be a subspace of V complementary to W_1 . Let T be a maximal torus of the compact Lie group C. As $c_1, \ldots, c_m \in Z(C)$ it follows that $c_1, \ldots, c_m \in T$. Choose $X_1, \ldots, X_m \in \text{Lie}(T)$ such that $\exp(X_i) = c_i$ for all i. We now set a map $\phi : \mathbb{R}^m \longrightarrow V \times C$ by

$$
\phi((t_1,\ldots,t_m)) := (t_1v_1 + \cdots + t_mv_m, \exp(t_1X_1 + \cdots + t_mX_m)), (t_1,\ldots,t_m) \in \mathbb{R}^m.
$$

As the Lie algebra $Lie(T)$ is abelian, the map ϕ is a (smooth) group homomorphism. We now show that $\phi((t_1,\ldots,t_m)) \in \Gamma$ if and only if $(t_1,\ldots,t_m) \in \mathbb{Z}^m$. It is easy to see that $\phi(\mathbb{Z}^m) \subset \Gamma$. Now if $(t_1v_1 + \cdots + t_mv_m, \exp(t_1X_1 + \cdots + t_mX_m)) \in \Gamma$ then $t_1v_1 + \cdots + t_mv_m \in \pi(\Gamma)$. As $\{v_1, \ldots, v_m\}$ is R-linearly independent and is a Z-basis for $\pi(\Gamma)$ it follows that $t_1, \ldots, t_m \in \mathbb{Z}$. As $\mathbb{R}^m/\mathbb{Z}^m$ is compact one has the embedding of groups, say, $\phi' : \mathbb{R}^m/\mathbb{Z}^m \to W \times C/\Gamma$ which is induced from ϕ . Let $T := \phi'(\mathbb{R}^m/\mathbb{Z}^m)$. Then $T \simeq (\mathbb{S}^1)^m$. We next show that $T \cap \theta(C)$ is finite. For this one notices that if $(t_1v_1 + \cdots + t_mv_m, \exp(t_1X_1 + \cdots + t_mX_m)) = (0, c) \mod \Gamma.$ Then $t_1v_1 + \cdots + t_mv_m \in \pi(\Gamma).$ Thus $t_i \in \mathbb{Z}$ for all i. Thus $(t_1v_1 + \cdots + t_mv_m, \exp(t_1X_1 + \cdots + t_mX_m)) \in \Gamma$, and hence $(0, c) \in 0 \times C \cap \Gamma$. Thus $T \cap \theta(C) \subset \theta(C \cap \Gamma)$, and hence $T \cap \theta(C)$ is finite.

In this section we also require facts associated to the Cartan decomposition in a semisimple Lie group for which we refer to \overline{Kn} , p. 362, Theorem 6.31 for details. If L is a semisimple Lie group and if $\sigma : L \longrightarrow L$ is a covering homomorphism then we may choose Cartan involutions $\Theta: L \longrightarrow L$, $\widetilde{\Theta}: \widetilde{L} \longrightarrow \widetilde{L}$, aligning with each other, that is, $\Theta \circ \sigma = \sigma \circ \Theta$. If G is a semisimple group and K denotes the group of fixed points of a Cartan involution then $Z(G) \subset K$. Moreover, it follows from [\[Kn,](#page-42-10) p. 362, Theorem 6.31] that $K/Z(G)$ is a maximal compact subgroup of $G/Z(G)$. Thus, one may observe from the facts above that K need not be a maximal compact subgroup in G but any maximal compact subgroup of K is a maximal compact subgroup of G. However, the Lie algebra $\mathfrak k$ is a compact Lie algebra (hence it is reductive). We conclude that any maximal compact subgroup of a simply connected semisimple group is either semisimple or the trivial group. We also need Weyl's theorem which says that the universal cover of a compact semisimple group is compact; see $[Kn, p. 268, Theorem 4.69].$

If L is a connected Lie group and if L is an almost direct product of the distinct closed connected subgroups L_1, \ldots, L_n then L_i and L_j commute if $i \neq j$. Moreover, $Z(L) = Z(L_1) \cdots Z(L_n)$. The kernel of the homomorphism $\phi : L_1 \times \cdots \times L_n \to L$ defined by $\phi(x_1,\ldots,x_n) := x_1 \cdots x_n$ is isomorphic to $L_1 \cap L_2 \cdots L_n \simeq L_i \cap L_1 \cdots L_i \cdots L_n$ (here A means A is omitted). Moreover, for all $i, L_i \cap L_1 \cdots L_i \cdots L_n = Z(L_1) \cap$ $Z(L_1)\cdots\overline{Z}(L_i)\cdots Z(L_n).$

Proposition 5.3. Let L be a semisimple Lie group. Suppose L_1, \ldots, L_r are all the simple factors of L. Let $\Gamma := L_1 \cap L_2 \cdots L_r$. Then the following hold:

- (1) If $|\Gamma| < \infty$, then for any maximal compact subgroup K of L there exist maximal compact subgroup K_i in L_i for all i such that $K = K_1 \cdots K_r$.
- (2) If $|\Gamma| = \infty$, then Γ is a finitely generated (abelian) group of rank 1. One also has that, for all i, if K_i is a the subgroup of fixed points corresponding to a (global) Cartan involution of L_i then $K_i \simeq \mathbb{R} \times [K_i, K_i]$ and $[K_i, K_i]$ is a maximal compact subgroup of L_i . Moreover, for all i, either K_i is abelian (this happens only when $L_i \simeq \text{SL}_2(\mathbb{R})$, the simply connected cover of $\text{SL}_2(\mathbb{R})$) or $[K_i, K_i]$ is compact semisimple. Further, for any maximal compact subgroup K of L there exist Cartan involutions of L_i , for all i such that if K_i is the corresponding subgroup of fixed points then $K = T[K_1, K_1] \cdots [K_r, K_r]$ where T is a central (compact) torus of K with dim $T = 1$. In particular, any maximal compact subgroup of L is not semisimple and the center is one dimensional.

Proof. We begin by claiming that rank $Z(L_1) \cap Z(L_2) \cdots Z(L_n) \leq 1$. Recall that $Z(L_1) \cap Z(L_2) \cdots Z(L_n)$ is a discrete finitely generated abelian subgroup of $Z(L)$. Let K_1 be the fixed point subgroup of a Cartan involution of L_1 . Then $Z(L_1) \subset K_1$; see [\[Kn,](#page-42-10) Theorem 6.31, p. 362]. Thus $Z(L_1) \subset Z(K_1)$, and it is well-known that dim $Z(K_1) \leq 1$. Thus either dim $Z(K_1) = 0$ or $Z(K_1) \simeq \mathbb{S}^1$ or $Z(K_1) \simeq \mathbb{R}$. Since $Z(L_1) \cap Z(L_2) \cdots Z(L_n)$ is a discrete subgroup the claim follows.

For the proof we set the homomorphism $\phi: L_1 \times \cdots \times L_n \to L$ defined by $\phi(x_1, \ldots, x_n) :=$ $x_1 \cdots x_n$. Let $\Gamma := \ker \phi$. Then $\Gamma \simeq L_1 \cap L_2 \cdots L_n$.

If $|L_1 \cap L_2 \cdots L_n| < \infty$ then the homomorphism ϕ is a finite covering map (hence proper). Let $K \subset L$ be a maximal compact then $\phi^{-1}(K)$ is a compact subgroup of

 $L_1 \times \cdots \times L_n$. Let M be a maximal compact of $L_1 \times \cdots \times L_n$ such that $M \subset \phi^{-1}(K)$. It is easy to see that there exists maximal compact subgroups K_i of L_i for all i such that $M = K_1 \times \cdots \times K_n$. Now as $K \subset K_1 \cdots K_n$ by maximality it follows that $K = K_1 \cdots K_n$. Let $|L_1 \cap L_2 \cdots L_n| = \infty$. Let K_i be the fixed point subgroup of a Cartan involution of L_i , for all i. Recall that $Z(L_i) \subset K_i$ for all i. It follows that $|L_i \cap L_1 \cdots L_i \cdots L_n| =$ ∞ for all *i*. As $L_i \cap L_1 \cdots \widehat{L_i} \cdots L_n = Z(L_i) \cap Z(L_1) \cdots \widehat{Z(L_i)} \cdots Z(L_n)$ it follows that $Z(K_i) \simeq \mathbb{R}$ for all i. Observe that $\Gamma \subset Z(L_1) \times \cdots \times Z(L_n) \subset Z(K_1) \times \cdots \times Z(K_n)$. Since $L_1 \times \cdots \times L_n/K_1 \times \cdots \times K_n$ is contractible, by Lemma [5.1](#page-29-1) (2), a maximal compact of $K_1 \times \cdots \times K_n/\Gamma$ is a maximal compact of $L_1 \times \cdots \times L_n/\Gamma$ (note that $K_1 \times \cdots \times K_n/\Gamma$ naturally embeds in $L_1 \times \cdots \times L_n/\Gamma$). Now recall by Weyl's theorem that $[K_i, K_i]$ is either a trivial group or a compact semisimple group. As rank $\Gamma = 1$ and $K_1 \times \cdots \times K_n \simeq \mathbb{R}^n \times C$ where C is a compact group, by Lemma [5.2,](#page-30-0) the results follows. \square

The next result follows immediately from Proposition [5.3.](#page-31-0)

Corollary 5.4. Let L be a semisimple Lie group such that a maximal compact subgroup of L is semisimple. Let L_1, \ldots, L_r be the simple factors of L. Then any maximal compact subgroup of L is of the form $G_1 \cdots G_r$ where G_i is a maximal compact subgroup of L_i for each $i = 1, \ldots, r$.

Remark 5.5. We remark here that semisimplicity of maximal compact subgroups in each simple factor of a semisimple group L does not force semisimplicity of a maximal compact subgroup in L. One can construct an example as in the following way. Let $SU(p, q)$ be the universal cover of $SU(p, q)$. Then one can see that the fixed point subgroup of any global Cartan involution in $SU(p, q)$ is isomorphic to $\mathbb{R} \times SU(p) \times SU(q)$. Thus any maximal compact subgroup in $SU(p, q)$ is isomorphic to $SU(p) \times SU(q)$, and hence it is semisimple. It is clear that there is a discrete central subgroup $\Lambda \subset SU(p, q)$ such that $\Lambda \simeq \mathbb{Z}$. Let $\Gamma := \{(\gamma, \gamma) \mid \gamma \in \Lambda\}$. Now consider $L := \mathrm{SU}(p, q) \times \mathrm{SU}(p, q)/\Gamma$, and let θ : $\widetilde{\mathrm{SU}(p, q)} \times \widetilde{\mathrm{SU}(p, q)} \to L$ be the quotient homomorphism. Let $I \in \mathrm{SU}(p, q)$ be the identity element, $L_1 := \theta(\mathrm{SU}(p, q) \times \{I\})$, and $L_2 := \theta(\{I\} \times \mathrm{SU}(p, q))$. Then $L_1 \simeq \text{SU}(p, q) \simeq L_2$, and L_1, L_2 are the simple factors of L. Thus both the simple factors of L have semisimple maximal compact subgroups, but by Proposition [5.3](#page-31-0) (2), maximal compact subgroups of L are not semisimple. \Box

Our next goal is to list all the simple real groups according to their types; see Definition [4.15.](#page-28-0) We first make the following observations. Let G be a simple Lie group and K be the group of fixed points of a Cartan involution. When $\dim_{\mathbb{R}} Z(K)^0 = 1$ there are only two possibilities, namely, $Z(K)^0 \simeq \mathbb{S}^1$ or $Z(K)^0 \simeq \mathbb{R}$. We list below various equivalent conditions for these possibilities to hold, which are easily verifiable.

The following conditions are equivalent to $Z(K)^0 \simeq \mathbb{R}$:

- $K \simeq \mathbb{R} \times [K, K]$, $|\pi_1(G, e)| < \infty$, $|Z(G)| = \infty$,
- K is non-compact, [K, K] is a maximal compact subgroup in G.

Similarly, the following is a bunch of equivalent conditions for the situation $Z(K)^0 \simeq \mathbb{S}^1$:

- $\mathbb{S}^1 \times [K, K]$ is a (finite) cover of K, $|\pi_1(G, e)| = \infty$, $|Z(G)| < \infty$,
- K is compact, K is a maximal compact subgroup in G , $\mathbb{Z} \subset \pi_1(G, e)$.

In view of the foregoing observations we list below the simple real groups with respect to their types. We follow the notation of real simple Lie algebras as in $[Kn]$. Chapter VI, $\S10$] and the data associated with them as in [\[Kn,](#page-42-10) Appendix C]. Let us now assume that G is a simple Lie group. Let $\mathfrak k$ be the Lie algebra of K. If $\mathfrak k$ is semisimple then G is either of Type-I or of Type-II (and in this case K is compact).

When $\mathfrak k$ is simple we list below the Type-I and the Type-II simple Lie groups.

- (1) The simple Lie group G is of Type-I if and only if G is a compact simple Lie group or G is a complex simple Lie group or $\mathfrak g$ is one of the Lie algebras $\mathfrak{sl}(n,\mathbb R)$ for $n > 2$, $\mathfrak{sl}(n, \mathbb{H})$, EI, EIV, EV, EVIII, FII.
- (2) The simple Lie group G is of Type-II if and only if the Lie algebra $\mathfrak g$ is isomorphic to one of the Lie algebras $\mathfrak{sp}(p,q)$, $\mathfrak{so}(p,q)$ for $p \geq q > 2$, EII, EVI, EIX, FI, G.

We now assume dim_R $\mathfrak{z}(\mathfrak{k}) = 1$. Then we have the following:

- (1) If $\mathfrak{g} \simeq \mathfrak{sl}_2(\mathbb{R})$ then G is neither of Type-I nor of Type-II. In this case a maximal compact subgroup is either isomorphic to \mathbb{S}^1 or the trivial group according as $|Z(G)| < \infty$ or $|Z(G)| = \infty$.
- (2) Let **g** be isomorphic to one of the Lie algebras $\mathfrak{sp}(n,\mathbb{R})$, $\mathfrak{so}^*(2n)$, $\mathfrak{so}(n,2)$ for $n > 2$, EIII, EVII. In this case $[\mathfrak{k}, \mathfrak{k}]$ is simple. Thus G is of Type-I if and only if $|Z(G)| = \infty$, and the subgroup K is a maximal compact with a one dimensional center if and only if $|Z(G)| < \infty$.
- (3) Let $\mathfrak{g} \simeq \mathfrak{su}(p,q)$. In this case $[\mathfrak{k}, \mathfrak{k}]$ has two simple factors. Thus G is of Type-II if and only if $|Z(G)| = \infty$, and the subgroup K is a maximal compact with a one dimensional center if and only if $|Z(G)| < \infty$.

6. Appendix : The equivariant Cartan's Theorem

The celebrated Cartan's theorem (see Theorem [6.10,](#page-38-0) [\[GHV-III,](#page-42-6) p. 462, Theorem II], [\[Ca\]](#page-41-3)) gives an alternate description of the cohomologies of a homogeneous space in terms of cohomologies of the associated Koszul complex. In the proofs of our main theorems, we have used a version of the above Cartan's theorem, namely, Theorem [6.15.](#page-41-8) This moreover follows from a slightly more general version, namely Theorem [6.14,](#page-41-9) which we call the equivariant Cartan's Theorem. Although Theorem [6.14](#page-41-9) is expected to be known but a proof could not be found in the current literature. Hence in this appendix we include a proof of Theorem [6.14,](#page-41-9) which is essentially a technical modification of the proof of the original Cartan's theorem as in [\[GHV-III,](#page-42-6) p. 462, Theorem II].

6.1. Preliminaries for the equivariant Cartan's Theorem. For the convenience of the reader we begin by recalling briefly some of the basic notions and required results.

6.1.1. The differential algebra $\wedge \mathfrak{g}^*$. Let G be a connected Lie group, and let $H \subset G$ be a closed, but not necessarily connected, subgroup. Let $\mathfrak{g} := \mathrm{Lie}(G)$ and $\mathfrak{h} := \mathrm{Lie}(H)$.

For any $X \in \mathfrak{g}$ there are uniquely determined operators $i_{\mathfrak{g}}(X), \theta_{\Lambda \mathfrak{g}^*}(X)$ acting on $\Lambda \mathfrak{g}^*$ which are given by

$$
(i_{\mathfrak{g}}(X)\omega)(X_1, \ldots, X_{p-1}) = \omega(X, X_1, \ldots, X_{p-1})
$$
\n(6.1)

$$
(\theta_{\wedge \mathfrak{g}^*}(X)\omega)(X_1, \ldots, X_p) = -\sum_{i=1}^p \omega(X_1, \ldots, [X, X_i], \ldots, X_p).
$$
 (6.2)

Moreover, one may define the *exterior derivative* $\delta_{\mathfrak{g}}$ on $\wedge \mathfrak{g}^*$ by

$$
(\delta_{\mathfrak{g}}\omega)(X_0, X_1, \ldots, X_p) = \sum_{i < j} (-1)^{i+j} \omega([X_i, X_j], X_0, \ldots, \widehat{X_i}, \ldots, \widehat{X_j}, \ldots, X_p). \tag{6.3}
$$

See [\[GHV-II,](#page-42-5) § 4.6, p. 156, Proposition III] for more details. Let

$$
(\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0} := \{ \omega \in \wedge \mathfrak{g}^* \mid i_{\mathfrak{g}}(Y)\omega = 0, \text{ for all } Y \in \mathfrak{h} \}.
$$

We next set $(\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}^{\mathfrak{h}} := (\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0} \cap (\wedge \mathfrak{g}^*)^{\mathfrak{h}}.$

We now recall a standard result which will be used later.

Theorem 6.1 ($\boxed{\text{Sp}, \text{p. 310}}$, Theorem 30, $\boxed{\text{Sp}, \text{p. 313}}$). Let G be a compact connected Lie group, and let $H \subset$ be a closed subgroup with identity component H^0 . Then the following isomorphisms hold :

$$
(1) H^*(G/H^0, \mathbb{R}) \simeq H^*((\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}, \delta_{\mathfrak{g}}).
$$

$$
(2) H^*(G/H, \mathbb{R}) \simeq H^*((\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}, \delta_{\mathfrak{g}})^{H/H^0}.
$$

6.1.2. The Weil algebra of a pair. Let $\mathfrak g$ be a reductive Lie algebra and $\mathfrak h \subset \mathfrak g$ a reductive subalgebra. The *Weil algebra of the pair* (a, b) is defined by

$$
W(\mathfrak{g}, \mathfrak{h}) := S(\mathfrak{h}^*) \otimes \wedge \mathfrak{g}^* \,. \tag{6.4}
$$

As both $S(\mathfrak{h}^*)$ and $\wedge \mathfrak{g}^*$ are graded algebras so is $W(\mathfrak{g}, \mathfrak{h})$ with respect to the total degree, where, very importantly, $\deg \psi = 2k$ for all non-zero $\psi \in S^k(\mathfrak{h}^*)$, and the elements of ∧g [∗] have the usual degree; see [\[GHV-III,](#page-42-6) § 10.9, p. 422]. We will use the anticommutative product operation on $W(\mathfrak{g}, \mathfrak{h})$ as in (2.2) .

Consider the coadjoint representation $-\text{ad}^* : \mathfrak{h} \longrightarrow \text{End } \mathfrak{h}^*$ defined by $X \longmapsto -(\text{ad } X)^*$. The linear transformation $-(ad X)^*$ extends to a derivation $\theta_{S(\mathfrak{h}^*)}(X)$ on the algebra $S(\mathfrak{h}^*)$. Let $\theta_{W(\mathfrak{g},\mathfrak{h})}$: $\mathfrak{h} \longrightarrow \text{End } W(\mathfrak{g},\mathfrak{h})$ be the representation of \mathfrak{h} defined by

$$
\theta_{W(\mathfrak{g},\mathfrak{h})}(Y) \, := \, \theta_{S(\mathfrak{h}^*)}(Y) \otimes 1 + 1 \otimes \theta_{\wedge \mathfrak{g}^*}(Y), \qquad Y \, \in \, \mathfrak{h}\,,
$$

where $\theta_{\Lambda g^*}(Y)$ is defined in [\(6.2\)](#page-34-0). Let $W(\mathfrak{g}, \mathfrak{h})^{\mathfrak{h}} \subset W(\mathfrak{g}, \mathfrak{h})$ be the *invariant subalgebra* with respect to the above representation $\theta_{W(\mathfrak{g},\mathfrak{h})}$; see [\(2.3\)](#page-5-0). We next define a map $h_{W(\mathfrak{g},\mathfrak{h})}$ on $W(\mathfrak{g}, \mathfrak{h})$ by

$$
h_{W(\mathfrak{g},\mathfrak{h})}(\psi \otimes (x_0^* \wedge \cdots \wedge x_p^*)) := \sum_{i=0}^p (-1)^i ((x_i^*|_{\mathfrak{h}}) \vee \psi) \otimes (x_0^* \wedge \cdots \wedge \hat{x_i^*} \wedge \cdots \wedge x_p^*),
$$

\n
$$
h_{W(\mathfrak{g},\mathfrak{h})}(\psi \otimes 1) := 0, \quad \text{for all } \psi \in S(\mathfrak{h}^*), \ x_0^*, \dots, x_p^* \in \mathfrak{g}^*.
$$
 (6.5)

The map $h_{W(\mathfrak{g},\mathfrak{h})}$ in (6.5) is an anti-derivation, and it is a graded map of degree 1. Moreover, from the identity

$$
h_{W(\mathfrak{g},\mathfrak{h})}\circ \theta_{W(\mathfrak{g},\mathfrak{h})}(Y) \,=\, \theta_{W(\mathfrak{g},\mathfrak{h})}(Y)\circ h_{W(\mathfrak{g},\mathfrak{h})}\,,\qquad Y\,\in\,\mathfrak{h}\,,
$$

it follows that the space $W(\mathfrak{g}, \mathfrak{h})^{\mathfrak{h}}$ remains invariant under the operator $h_{W(\mathfrak{g}, \mathfrak{h})}$. Hence the latter restricts to an operator on $W(\mathfrak{g},\mathfrak{h})^{\mathfrak{h}}$. We next define another operator D on $W(\mathfrak{g}, \mathfrak{h})$ by

$$
D := 1 \otimes \delta_{\mathfrak{g}} - h_{W(\mathfrak{g}, \mathfrak{h})} \tag{6.6}
$$

(see [\(6.3\)](#page-34-2) for the definition of $\delta_{\mathfrak{g}}$). It follows that $(W(\mathfrak{g}, \mathfrak{h})^{\mathfrak{h}}, D)$ is a graded differential algebra (see [\[GHV-III,](#page-42-6) p. 422]).

Definition 6.2. The Weil algebra $W(\mathfrak{g})$ of a reductive Lie algebra \mathfrak{g} is defined by (see [\(6.4\)](#page-34-3))

$$
W(\mathfrak{g}) := W(\mathfrak{g}, \mathfrak{g}) = S(\mathfrak{g}^*) \otimes \wedge \mathfrak{g}^*.
$$

We set $W(\mathfrak{g})^{\mathfrak{g}} := W(\mathfrak{g}, \mathfrak{g})^{\mathfrak{g}}$ where $W(\mathfrak{g}, \mathfrak{g})^{\mathfrak{g}}$ is the invariant subalgebra of $W(\mathfrak{g}, \mathfrak{g})$ in sense above. Then $\delta_W := 1 \otimes \delta_{\mathfrak{g}} - h_{W(\mathfrak{g}, \mathfrak{g})}$ is a differential on $W(\mathfrak{g})^{\mathfrak{g}}$; see [\(6.6\)](#page-35-1).

6.1.3. Curvature and the Chern-Weil homomorphism. Let G be a connected Lie group and $H \subset G$ a compact subgroup which is not necessarily connected. Let $\mathfrak{g} := \mathrm{Lie}(G)$, and $\mathfrak{h} := \text{Lie}(H)$. Let \mathfrak{m} be a H-invariant subspace in g such that

$$
\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}. \tag{6.7}
$$

Let $\pi_{\mathfrak{h}} : \mathfrak{g} \longrightarrow \mathfrak{h}$ and $\pi_{\mathfrak{m}} : \mathfrak{g} \longrightarrow \mathfrak{m}$ be the natural projections corresponding to the decomposition in [\(6.7\)](#page-35-2).

Definition 6.3. Let $\Omega^2(G, \mathfrak{h})$ be the space of smooth \mathfrak{h} -valued two-forms on G.

Theorem 6.4 ($|KN, p. 103, Theorem 11.1$). Let G and H be as above. Then the following hold :

- (1) The projection π_h produces a connection on the principal H-bundle $G \longrightarrow G/H$; this connection is invariant under the left translation action of G.
- (2) Conversely, any G-invariant connection on the principal H-bundle $G \longrightarrow G/H$ is given by a H-invariant decomposition as in [\(6.7\)](#page-35-2).
- (3) The curvature form $\kappa \in \Omega^2(G, \mathfrak{h})$ corresponding to the invariant connection in statement (1) is a G-invariant two from and it is given by

$$
\kappa(e)(X, Y) = -\pi_{\mathfrak{h}}([\pi_{\mathfrak{m}}(X), \pi_{\mathfrak{m}}(Y)]), \qquad X, Y \in \mathfrak{g},
$$

where $e \in G$ is the identity element.

Remark 6.5. In [\[KN,](#page-42-12) p. 103, Theorem 11.1] the expression for the curvature κ evaluated at the identity $e \in G$ differs from the above by the scaling factor $1/2$. This deviation happens due to the presence of an additional scaling factor in the definition of the exterior derivative in [\[KN,](#page-42-12) p. 36, Proposition 3.11].

Definition 6.6. Recall that $\pi_{\mathfrak{m}} : \mathfrak{g} \longrightarrow \mathfrak{m}$ is the projection map with respect to the decomposition in [\(6.7\)](#page-35-2). The map π^{hor} : $\wedge \mathfrak{g}^* \longrightarrow (\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}|\mathfrak{h}}=0}$ defined by

$$
\pi^{hor}(\omega)(X_1, \ldots, X_k) := \omega(\pi_{\mathfrak{m}}(X_1), \ldots, \pi_{\mathfrak{m}}(X_k)),
$$

where $X_i \in \mathfrak{g}, 1 \leq i \leq k$, and $\omega \in \Delta \mathfrak{g}^*$, is called the *horizontal projection* associated to (6.7) .

Let $\kappa \in \Omega^2(G, \mathfrak{h})$ be the *curvature* of the invariant connection $\pi_{\mathfrak{h}}$, as in Theorem [6.4](#page-35-3) (3) on the principal H-bundle $G \longrightarrow G/H$. Let $e \in G$ be the identity element. Then, for $X, Y \in \mathfrak{g}$ and $f \in \mathfrak{h}^*$, we have

$$
\pi^{hor} \circ \delta_{\mathfrak{g}} \circ \pi_{\mathfrak{h}}^*(f)(X, Y) = -f(\pi_{\mathfrak{h}}([\pi_{\mathfrak{m}}(X), \pi_{\mathfrak{m}}(Y)])) = f(\kappa(e))(X, Y), \tag{6.8}
$$

where $\delta_{\mathfrak{g}}$ and π^{hor} are constructed in [\(6.3\)](#page-34-2) and in Definition [6.6,](#page-35-4) respectively. The map $\mathfrak{h}^* \longrightarrow \wedge^2(\mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}$ defined by $f \longmapsto \pi^{hor} \circ \delta_{\mathfrak{g}} \circ \pi_{\mathfrak{h}}^*(f)$ extends to an algebra homomorphism

$$
\kappa_{\vee}: S(\mathfrak{h}^*) \longrightarrow (\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}|_{\mathfrak{h}}=0}}.\tag{6.9}
$$

Let $\mathrm{Ad}^*_{\wedge}: G \longrightarrow \mathrm{GL}(\wedge \mathfrak{g}^*)$ be the representation naturally obtained from the coadjoint representation Ad^* : $G \longrightarrow \mathrm{GL}(\mathfrak{g}^*)$. Observe that the subspace $(\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0} \subset \wedge \mathfrak{g}^*$ remains invariant under the action of the operator $\mathrm{Ad}^*_{\wedge}(h)$ for all $h \in H$. Analogously, let $\mathrm{Ad}^*_{\vee}: H \longrightarrow \mathrm{GL}(S(\mathfrak{h}^*))$ be the representation naturally obtained from the coadjoint representation $\text{Ad}^* : H \longrightarrow \text{GL}(\mathfrak{h}^*)$. Note that this representation is infinite dimensional but H acts on $S(\mathfrak{h}^*)$ locally finitely.

In view of the above notation it may be observed that the homomorphism in [\(6.9\)](#page-36-0) is also H-equivariant, that is,

$$
\mathrm{Ad}^*_{\wedge}(h) \circ \kappa_{\vee} = \kappa_{\vee} \circ \mathrm{Ad}^*_{\vee}(h) \quad \text{for all } h \in H.
$$

This induces the following algebra homomorphisms :

$$
S(\mathfrak{h}^*)^H \longrightarrow (\wedge \mathfrak{g}^*)^H_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0} \longrightarrow H((\wedge \mathfrak{g}^*)^H_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}, \delta_{\mathfrak{g}}).
$$

The above composition of homomorphisms

$$
S(\mathfrak{h}^*)^H \longrightarrow H((\wedge \mathfrak{g}^*)^H_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}, \delta_{\mathfrak{g}})
$$

is known as the *Chern-Weil homomorphism*; see [\[GHV-III,](#page-42-6) § 8.26, p. 355].

6.1.4. Cartan's theorem for connected H . We shall now discuss Cartan's theorem on cohomology of compact connected homogeneous space G/H , where the group H is connected. The complex $((\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}\mid\mathfrak{h}}=0}^{\mathfrak{h}}, \delta_{\mathfrak{g}})$ is a subcomplex of $((S(\mathfrak{h}^*)\otimes \wedge \mathfrak{g}^*)^{\mathfrak{h}}, D)$, where the differential D is defined as in (6.6) . The inclusion map

$$
\varepsilon: (\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}|\mathfrak{h}}=0} \longrightarrow (S(\mathfrak{h}^*) \otimes \wedge \mathfrak{g}^*)^{\mathfrak{h}}; \ \ \phi \longmapsto 1 \otimes \phi \tag{6.10}
$$

is a homomorphism of graded differential algebras. Next consider the map

$$
\alpha_{\kappa}:\;S(\mathfrak{h}^*)\otimes\wedge \mathfrak{g}^*\;\longrightarrow\;(\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}\mid \mathfrak{h}}=0}
$$

defined by

$$
\sum_i \psi_i \otimes \phi_i \; \longmapsto \; \sum_i (\kappa \vee \psi_i) \wedge (\pi^{hor} \phi_i) \, , \quad \text{ for all } \psi_i \, \in \, S(\mathfrak{h}^*) \text{ and } \phi_i \, \in \, \wedge \mathfrak{g}^* \, ,
$$

where the map κ_{\vee} is constructed in [\(6.9\)](#page-36-0).

Lemma 6.7 ([\[GHV-III,](#page-42-6) Lemma III, p. 347]). The following hold :

 $(1) \alpha_{\kappa} \circ \theta_{W(\mathfrak{g},\mathfrak{h})} = \theta_{\wedge \mathfrak{g}^*|\mathfrak{h}} \circ \alpha_{\kappa}.$ (2) $\alpha_{\kappa} \circ D = \pi^{\text{hor}} \circ \delta_{\mathfrak{g}} \circ \alpha_{\kappa}.$

Consequently, the map

$$
\alpha_{\kappa}: (S(\mathfrak{h}^*) \otimes \wedge \mathfrak{g}^*)^{\mathfrak{h}} \longrightarrow (\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}|\mathfrak{h}}=0} \tag{6.11}
$$

is a graded algebra homomorphism which commutes with the differentials.

The next result constitutes one of the important parts in the proof of the Cartan's Theorem (Theorem [6.10\)](#page-38-0). In fact it ensures that the homomorphism in one side of the triangle in Figure [1](#page-38-1) induces an isomorphism of the cohomologies.

Proposition 6.8 ([\[GHV-III,](#page-42-6) p. 344, Theorem IV], [\[GHV-III,](#page-42-6) p. 348, Proposition IX]). The homomorphisms of cohomologies ε^* and α^*_κ induced by the homomorphisms ε and α_κ , respectively, (see (6.10) and (6.11))

$$
H(W(\mathfrak{g},\mathfrak{h})^{\mathfrak{h}},\,D)\overset{\alpha_{\kappa}^*}{\underset{\varepsilon^*}{\rightleftarrows}}H((\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}^{\mathfrak{h}},\,\delta_{\mathfrak{g}})
$$

are isomorphism and inverse of each other.

6.1.5. The Chevalley homomorphism. Recall from Section [2.2.2](#page-7-3) that $P_{\mathfrak{g}} = P_{\mathfrak{g}} \cap (\wedge^{2i-1} \mathfrak{g}^*)^{\mathfrak{g}}$ is a graded subspace. We will consider $W(\mathfrak{g})$ as a graded algebra with respect to the total degree. Let $\tau: P_{\mathfrak{g}} \longrightarrow S(\mathfrak{g}^*)^{\mathfrak{g}}$ be a transgression in $W(\mathfrak{g})^{\mathfrak{g}}$. There is a graded linear map

$$
\alpha \, : \, P_{\mathfrak{g}} \, \longrightarrow \, (W(\mathfrak{g})^+)^{\mathfrak{g}} \tag{6.12}
$$

of degree 0 such that

$$
\delta_W \alpha(\phi) = \tau \phi \otimes 1 \quad \text{and} \quad 1 \otimes \phi - \alpha(\phi) \in (S(\mathfrak{g}^*)^+ \otimes \wedge \mathfrak{g}^*)^{\mathfrak{g}}.
$$

As $S(\mathfrak{g}^*)$ and $\wedge \mathfrak{g}^*$ are associative algebras, so is the Weil algebra $W(\mathfrak{g}^*)$ with respect to the anticommutative product in [\(2.2\)](#page-5-1). We extend α to a homomorphism $\alpha_{\wedge} : \wedge P_{\mathfrak{g}} \longrightarrow$ $W(\mathfrak{g})^{\mathfrak{g}}$. It can be easily seen that $j^* \otimes 1$ restricts to a homomorphism

 $j^* \otimes 1 : W(\mathfrak{g})^{\mathfrak{g}} \longrightarrow W(\mathfrak{g}, \, \mathfrak{h})^{\mathfrak{h}}$,

where $j^*: S(\mathfrak{g}^*) \longrightarrow S(\mathfrak{h}^*)$ is induced by the dual of the inclusion map $j: \mathfrak{h} \longrightarrow \mathfrak{g}$ (see (2.11)). Next define a homomorphism of graded algebras,

$$
\vartheta_{\tau}: S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes \wedge P_{\mathfrak{g}} \longrightarrow W(\mathfrak{g}, \mathfrak{h})^{\mathfrak{h}} \tag{6.13}
$$

by

$$
\vartheta_\tau(\psi \otimes \phi) = (\psi \otimes 1) \cdot (j^* \otimes 1)(\alpha \wedge \phi), \qquad \psi \in S(\mathfrak{h}^*)^{\mathfrak{h}}, \ \phi \in \wedge P_{\mathfrak{g}}.
$$

The map ϑ_{τ} satisfies the relation

$$
\vartheta_{\tau} \circ (-\nabla_{\sigma}) = D \circ \vartheta_{\tau};\tag{6.14}
$$

see [\[GHV-III,](#page-42-6) p.424,(10.4)]. The map ϑ_{τ} in [\(6.13\)](#page-37-1) is called the *Chevalley homomorphism*.

As in Proposition [6.8](#page-37-2) the next result is another step in the proof of the Cartan's theorem. It gives an isomorphism of cohomologies induced by the map in another side of the triangle in Figure [1.](#page-38-1)

Proposition 6.9 ([\[GHV-III,](#page-42-6) p. 424, Proposition IV]). Let τ be a transgression as in Definition [2.2.](#page-8-3) Let σ be the map as in [\(2.12\)](#page-9-4) associated to τ . Consider the Koszul complex $(S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes \wedge P_{\mathfrak{g}}, \nabla_{\sigma})$ (see § [2.2.4\)](#page-9-0). Then the Chevalley homomorphism ϑ_{τ} corresponding to τ induces an isomorphism of cohomologies

$$
\vartheta_{\tau}^* : H(S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes \wedge P_{\mathfrak{g}}, -\nabla_{\sigma}) \longrightarrow H(W(\mathfrak{g}, \mathfrak{h})^{\mathfrak{h}}, D).
$$

This isomorphism does not depend on the choice of the transgression.

FIGURE 1.

We need the well-known Cartan's theorem which is stated next.

Theorem 6.10 ([\[GHV-III,](#page-42-6) p. 462, Theorem II], [\[Ca\]](#page-41-3)). Let G be a compact connected Lie group and $H \subset G$ a closed subgroup. Let $H^0 \subset H$ be the connected component containing the identity element. Consider the map $\phi_{G/H}$ defined by

$$
\phi_{G/H^0} := \alpha_{\kappa} \circ \vartheta_{\tau} : (S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes \wedge P_{\mathfrak{g}}, -\nabla_{\sigma}) \longrightarrow ((\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}, \delta_{\mathfrak{g}}),
$$

where α_{κ} and ϑ_{τ} be the algebra homomorphisms defined in [\(6.11\)](#page-37-0) and [\(6.13\)](#page-37-1), respectively. Then the induced map of cohomologies

$$
\phi_{G/H^0}^* \,=\, \alpha_\kappa^* \circ \vartheta_\tau^* :\; H^*\big(S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes \wedge P_{\mathfrak{g}},\, -\nabla_\sigma\big) \,\stackrel{\cong}{\longrightarrow}\, H^*\big((\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0},\, \delta_{\mathfrak{g}}\big)
$$

is an isomorphism. Hence, $H^*(G/H^0, \mathbb{R}) \simeq H^*(S(\mathfrak{h}^*)^{\mathfrak{h}} \otimes \wedge P_{\mathfrak{g}}, -\nabla_{\sigma}).$

Proof. This follows from Proposition [6.9,](#page-37-3) Proposition [6.8](#page-37-2) and Theorem [6.1\(](#page-34-4)1). \Box

The commutative diagram in Figure [1](#page-38-1) represents the maps of the complexes involved in Theorem [6.10,](#page-38-0) and it will be useful in the proof of Theorem [6.14.](#page-41-9)

6.2. The equivariant Cartan's Theorem. In this section, we will prove Theorem [6.14](#page-41-9) and Theorem [6.15.](#page-41-8) We shall use notation and the background sketched in Section [2.](#page-4-0)

Let G be a compact connected Lie group and let $\mathfrak{g} := \mathrm{Lie}(G)$. Let Γ be a group acting on **g** by Lie algebra automorphisms. Then Γ induces an action on $S(\mathfrak{g}^*)^{\mathfrak{g}}$, as well as an action on $(\Lambda \mathfrak{g}^*)^{\mathfrak{g}}$ that leaves $P_{\mathfrak{g}}$ invariant (see Section [2.2.2](#page-7-3) for the definition of $P_{\mathfrak{g}}$). Let Trans(\mathfrak{g}) be the space of all transgressions for the Lie algebra \mathfrak{g} as introduced in Definition [2.2.](#page-8-3) The group Γ has a natural action on Trans(g) by the rule $\gamma \cdot \tau := \gamma \circ \tau \circ \gamma^{-1}$ for $\gamma \in \Gamma$.

Lemma 6.11. There exists an element in Trans(\mathfrak{g}) which is fixed by Γ.

Proof. Considering the distinguished transgression τ_{g} defined in [\[GHV-III,](#page-42-6) p. 236], the lemma follows form [\[GHV-III,](#page-42-6) p. 240, Proposition VIII].

A straightforward alternative proof is as follows. Recall that $Trans(g)$ is an affine subspace of $\text{Hom}(P_{\mathfrak{g}}, S(\mathfrak{g}^*)^{\mathfrak{g}})$; see Remark [2.6\(](#page-8-2)2). Also, recall that Aut \mathfrak{g} keeps Trans(\mathfrak{g}) invariant while $(\text{Aut } \mathfrak{g})^0$ acts trivially on Trans (\mathfrak{g}) . Note that the triviality of the action of $(\text{Aut } \mathfrak{g})^0$ on Trans (\mathfrak{g}) follows immediately from the fact that action of $(\text{Aut } \mathfrak{g})^0$ is trivial on both $P_{\mathfrak{g}}$ and $S(\mathfrak{g}^*)^{\mathfrak{g}}$. It is easy to see that any finite group acting on an affine space has a fixed point. Thus, $\text{Aut } \mathfrak{g} / (\text{Aut } \mathfrak{g})^0$ being a finite group, there is a point of Trans (\mathfrak{g}) , say λ, which is fixed by the action of Aut g. This λ will remain fixed by Γ for any Γ acting on $\mathfrak g$ via automorphisms.

6.2.1. Γ-invariant connection and curvature. Let G and Γ be as at the beginning of this section. Let $H \subset G$ be a closed connected subgroup with Lie algebra \mathfrak{h} . Assume that the action of Γ on g preserves the subalgebra h. We now fix a Γ -invariant and H -invariant subspace $\mathfrak m$ in $\mathfrak g$ (such a subspace $\mathfrak m$ exists) such that

$$
\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m} \,.
$$

In view of Theorem [6.4\(](#page-35-3)1), the projection $\pi_{\mathfrak{h},\Gamma}$: $\mathfrak{g} \longrightarrow \mathfrak{h}$ for the decomposition in [\(6.15\)](#page-39-0) produces a G-invariant connection on the principal H-bundle $G \longrightarrow G/H$. Let $\pi_{\mathfrak{m},\Gamma} : \mathfrak{g} \longrightarrow \mathfrak{m}$ be the projection onto \mathfrak{m} with respect the decomposition in [\(6.15\)](#page-39-0). As in Definition [6.6,](#page-35-4) the horizontal projection map π_{Γ}^{hor} : $\wedge \mathfrak{g}^* \longrightarrow (\wedge \mathfrak{g}^*)_{i_{\mathfrak{gl}(p)}=0}$ is defined by

$$
\pi_{\Gamma}^{hor}(\omega)(X_1,\ldots,X_k) := \omega(\pi_{\mathfrak{m},\Gamma}(X_1),\ldots,\pi_{\mathfrak{m},\Gamma}(X_k)). \qquad (6.16)
$$

Let

$$
\kappa_{\Gamma} \in \Omega^2(G, \mathfrak{h})
$$

be the curvature of the connection on the principal H-bundle $G \longrightarrow G/H$ given by $\pi_{h,\Gamma}$. Let $e \in G$ be the identity element. Then using Theorem [6.4](#page-35-3) (3) it follows that

$$
\kappa_{\Gamma}(e)(X, Y) = -\pi_{\mathfrak{h},\Gamma}([\pi_{\mathfrak{m},\Gamma}(X), \pi_{\mathfrak{m},\Gamma}(Y)]), \qquad X, Y \in \mathfrak{g}.
$$
 (6.17)

For X, $Y \in \mathfrak{g}$, and $f \in \mathfrak{h}^*$, we have

$$
\pi_{\Gamma}^{hor} \circ \delta_{\mathfrak{g}} \circ \pi_{\mathfrak{h},\Gamma}^*(f)(X,Y) = -f(\pi_{\mathfrak{h},\Gamma}([\pi_{\mathfrak{m},\Gamma}(X), \pi_{\mathfrak{m},\Gamma}(Y)])) = (f(\kappa_{\Gamma}))(e)(X,Y).
$$

The map $\mathfrak{h}^* \longrightarrow \wedge^2(\mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}$ defined by $f \longmapsto \pi_{\Gamma}^{hor} \circ \delta_{\mathfrak{g}} \circ \pi_{\mathfrak{h},\Gamma}^*(f)$ extends to an algebra homomorphism

$$
(\kappa_{\Gamma})_{\vee}: S(\mathfrak{h}^*) \longrightarrow (\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}|_{\mathfrak{h}}=0}}.
$$
\n(6.18)

Lemma 6.12. Let $(\kappa_{\Gamma})_{\vee}$ be the algebra homomorphism as in [\(6.18\)](#page-39-1). Let Γ be as above, and let us consider the natural actions of Γ on $S(\mathfrak{h}^*)$ and on $\wedge \mathfrak{g}^*$. Observe that $(\wedge \mathfrak{g}^*)_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}$ remains invariant under Γ. For an element $\gamma \in \Gamma$ let the associated automorphisms on each of the above spaces be denoted again by γ . Then the algebra homomorphism $(\kappa_{\Gamma})_{\vee}$ is Γ-equivariant, that is,

$$
\gamma \circ (\kappa_{\Gamma})_{\vee} = (\kappa_{\Gamma})_{\vee} \circ \gamma, \qquad \forall \gamma \in \Gamma.
$$

Proof. Recall that the curvature form κ_{Γ} at $e \in G$ is given by [\(6.17\)](#page-39-2). First note that $\kappa_{\Gamma}(e)(\gamma \cdot X, \gamma \cdot Y) = \gamma \cdot \kappa_{\Gamma}(e)(X, Y)$ for all $X, Y \in \mathfrak{g}, \gamma \in \Gamma$. Let $f \in S^p(\mathfrak{h}^*)$, and $X_1, \cdots, X_{2p} \in \mathfrak{g}.$ Now

$$
\gamma \circ (\kappa_{\Gamma})_{\vee}(f)(X_1, \cdots, X_{2p}) = \gamma \circ f(\underbrace{\kappa_{\Gamma}(e), \cdots, \kappa_{\Gamma}(e)}_{p-\text{many}})(X_1, \cdots, X_{2p})
$$

$$
= f(\kappa_{\Gamma}, \cdots, \kappa_{\Gamma})(e)(\gamma^{-1} \cdot X_1, \cdots, \gamma^{-1} \cdot X_{2p})
$$

$$
= \frac{1}{2^p} \sum_{\sigma \in \mathcal{P}(2p)} (\operatorname{sgn} \sigma) f(\kappa_{\Gamma}(e)(\gamma^{-1} \cdot X_{\sigma(1)}, \gamma^{-1} \cdot X_{\sigma(2)}), \cdots, \kappa_{\Gamma}(e)(\gamma^{-1} \cdot X_{\sigma(2p-1)}, \gamma^{-1} \cdot X_{\sigma(2p)}))
$$

\n
$$
= \frac{1}{2^p} \sum_{\sigma \in \mathcal{P}(2p)} (\operatorname{sgn} \sigma) f(\gamma^{-1} \cdot \kappa_{\Gamma}(e)(X_{\sigma(1)}, X_{\sigma(2)}), \cdots, \gamma^{-1} \cdot \kappa_{\Gamma}(e)(X_{\sigma(2p-1)}, X_{\sigma(2p)}))
$$

\n
$$
= \frac{1}{2^p} \sum_{\sigma \in \mathcal{P}(2p)} (\operatorname{sgn} \sigma)(\gamma \cdot f)(\kappa_{\Gamma}(e)(X_{\sigma(1)}, X_{\sigma(2)}), \cdots, \kappa_{\Gamma}(e)(X_{\sigma(2p-1)}, X_{\sigma(2p)}))
$$

\n
$$
= (\gamma \cdot f)(\kappa_{\Gamma}, \cdots, \kappa_{\Gamma})(e)(X_1, \cdots, X_{2p})
$$

\n
$$
= (\kappa_{\Gamma})_{\vee}(\gamma \cdot f)(X_1, \cdots, X_{2p}).
$$

This completes the proof. \Box

The algebra homomorphism in [\(6.11\)](#page-37-0) for the curvature form κ_{Γ}

$$
\alpha_{\kappa_{\Gamma}}: (S(\mathfrak{h}^*) \otimes \wedge \mathfrak{g}^*)^{\mathfrak{h}} \longrightarrow (\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}|\mathfrak{h}}=0} \tag{6.19}
$$

is given by

$$
\sum_i \psi_i \otimes \phi_i \longmapsto \sum_i ((\kappa_{\Gamma})_{\vee} \psi_i) \wedge (\pi_{\Gamma}^{hor} \phi_i), \qquad \psi_i \in S(\mathfrak{h}^*) \text{ and } \phi_i \in \wedge \mathfrak{g}^*,
$$

where $(\kappa_{\Gamma})_{\vee}$ is constructed in [\(6.18\)](#page-39-1). For the Chevalley homomorphism $\vartheta_{\tau_{\mathfrak{g}}}$ we will consider the transgression $\tau_{\mathfrak{g}}$ as in Lemma [6.11.](#page-38-2) Recall the definition of $\delta_{\mathfrak{g}}$ in [\(6.3\)](#page-34-2), D in (6.6) , and ∇_{σ} in (2.13) .

Lemma 6.13. Consider the curvature form κ_{Γ} and a transgression $\tau_{\mathfrak{g}}$ which is fixed under the Γ -action (see Lemma [6.11\)](#page-38-2). Then the following two hold:

- (1) The differentials $\delta_{\mathfrak{g}}, D, \nabla_{\sigma}$ are Γ -equivariant.
- (2) The algebra homomorphisms $\alpha_{\kappa_{\Gamma}}$, and $\vartheta_{\tau_{\mathfrak{g}}}$ are Γ -equivariant.

Proof. Proof of (1): Since $\Gamma \subset \text{Aut } \mathfrak{g}$, the differentials $\delta_{\mathfrak{g}}$ and D are Γ-equivariant. To prove that ∇_{σ} is Γ-equivariant, it is enough to check that the map σ in [\(2.12\)](#page-9-4) is Γequivariant. Recall that the transgression $\tau_{\mathfrak{g}}$ is fixed under the Γ-action. Thus the map σ is Γ-equivariant.

Proof of (2): It can be shown that the horizontal projection π_{Γ}^{hor} in [\(6.16\)](#page-39-3) satisfies the condition

$$
\pi_{\Gamma}^{hor}(\gamma \cdot \omega) = \gamma \cdot \pi_{\Gamma}^{hor}(\omega), \quad \text{ for all } \gamma \in \Gamma, \, \omega \in \Lambda \mathfrak{g}^*.
$$

To prove this, enough to consider $\omega \in \wedge^k(\mathfrak{g}^*)$. Now

$$
\pi_{\Gamma}^{hor}(\gamma \cdot \omega)(X_1, \cdots, X_k) = (\gamma \cdot \omega)(\pi_{\mathfrak{m},\Gamma}(X_1), \cdots, \pi_{\mathfrak{m},\Gamma}(X_k))
$$

\n
$$
= \omega(\gamma^{-1} \cdot \pi_{\mathfrak{m},\Gamma}(X_1), \cdots, \gamma^{-1} \cdot \pi_{\mathfrak{m},\Gamma}(X_k))
$$

\n
$$
= \omega(\pi_{\mathfrak{m},\Gamma}(\gamma^{-1} \cdot X_1), \cdots, \pi_{\mathfrak{m},\Gamma}(\gamma^{-1} \cdot X_k))
$$

\n
$$
= \pi_{\Gamma}^{hor}(\omega)((\gamma^{-1} \cdot X_1), \cdots, (\gamma^{-1} \cdot X_k))
$$

\n
$$
= \gamma \cdot \pi_{\Gamma}^{hor}(\omega)(X_1, \cdots, X_k).
$$

Next we will use the above relation and Lemma [6.12](#page-39-4) to show that the map $\alpha_{\kappa_{\Gamma}}$ is Γ equivariant:

$$
\gamma(\alpha_{\kappa_{\Gamma}}(\sum_{i} \psi_{i} \otimes \phi_{i})) = \gamma\Big(\sum_{i} ((\kappa_{\Gamma})_{\vee} \psi_{i}) \wedge (\pi_{\Gamma}^{hor} \phi_{i})\Big)
$$

$$
= \sum_{i} (\gamma \cdot ((\kappa_{\Gamma})_{\vee} \psi_{i})) \wedge (\gamma \cdot (\pi_{\Gamma}^{hor} \phi_{i}))
$$

$$
= \sum_{i} ((\kappa_{\Gamma})_{\vee}(\gamma \cdot \psi_{i})) \wedge (\pi_{\Gamma}^{hor} (\gamma \cdot \phi_{i}))
$$

$$
= \alpha_{\kappa_{\Gamma}} (\gamma \cdot (\sum_{i} \psi_{i} \otimes \phi_{i})).
$$

Finally, to prove the algebra homomorphism $\vartheta_{\tau_{\mathfrak{g}}}$ is Γ -equivariant, it is enough to show that the map $\alpha_{\tau_{\mathfrak{g}}}$ as in [\(6.12\)](#page-37-4) is Γ-equivariant, and this follows from Lemma [6.11.](#page-38-2) \Box

Theorem 6.14. Let G be a compact connected Lie group, and let $H \subset G$ be a closed connected subgroup. Suppose $\Gamma := \{ \gamma \in \text{Aut } \mathfrak{g} \mid \gamma(\mathfrak{h}) = \mathfrak{h} \}.$ Assume that there is an H-invariant subspace $\mathfrak m$ of $\mathfrak g$ such that $\mathfrak g = \mathfrak h \oplus \mathfrak m$, and $\gamma(\mathfrak m) = \mathfrak m$ for all $\gamma \in \Gamma$. Then the maps $\vartheta_{\tau_{\mathfrak{g}}}$ (as in [\(6.13\)](#page-37-1)) and α_{κ} (as in [\(6.11\)](#page-37-0)) are Γ -equivariant. Moreover, the Γ -fixed points induce an isomorphism

$$
H^*\big((S(\mathfrak{h}^*)^{\mathfrak{h}}\otimes \wedge P_{\mathfrak{g}})^{\Gamma}, -\nabla_{\sigma}\big)\,\simeq\, H^*\big(((\wedge \mathfrak{g}^*)^{\mathfrak{h}}_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0})^{\Gamma}, \,\delta_{\mathfrak{g}}\big)\,.
$$

Proof. This follows from Theorem [6.10](#page-38-0) and Lemma [6.13.](#page-40-0)

6.2.2. Cartan's theorem for disconnected H . Let G be a compact connected Lie group and $H \subset G$ a closed, but not necessarily connected subgroup. The next result follows from Theorem [6.14](#page-41-9) (set $\Gamma = H/H^0$) combined with Theorem [6.1\(](#page-34-4)2).

Theorem 6.15. Let G, H be as above. The following vector spaces are isomorphic:

$$
H^*\bigl(G/H, \mathbb{R}\bigr) \simeq H^*\bigl((\wedge \mathfrak{g}^*)^H_{i_{\mathfrak{g}}|_{\mathfrak{h}}=0}, \delta_{\mathfrak{g}}\bigr) \simeq H^*\bigl((S(\mathfrak{h}^*)^H \otimes \wedge P_{\mathfrak{g}}), -\nabla_{\sigma}\bigr).
$$

Acknowledgements. I. Biswas is supported by a J. C. Bose Fellowship. P. Chatterjee acknowledges support from the SERB-DST MATRICS project: MTR/2020/000007. C. Maity is supported by an NBHM PDF during the course of this work.

REFERENCES

[Az] H. Azad, On the third Betti number of some compact homogeneous manifolds, Studia Sci. Math. Hungar., 25 (1990), no. 1–2, 1–2. [4,](#page-3-5) [29](#page-28-1) [Ba] P. Baum, On the cohomology of homogeneous spaces, Topology, 7 (1968), 15–38. [2](#page-1-1) [Bo] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, Ann. of Math. 57 (1953), 115–207. [2](#page-1-1) [BC] I. Biswas and P. Chatterjee, On the exactness of Kostant-Kirillov form and the second cohomology of nilpotent orbits, Internat. J. Math. 23 (2012), no. 8, 1250086. [1,](#page-0-1) [2,](#page-1-1) [4,](#page-3-5) [28](#page-27-4) [BCM] I. Biswas, P. Chatterjee, C. Maity, The Second cohomology of nilpotent orbits in classical Lie algebras, Kyoto J. Math. 60 (2020), 717–799. [1,](#page-0-1) [2,](#page-1-1) [4,](#page-3-5) [12,](#page-11-4) [23,](#page-22-6) [28](#page-27-4) [Ca] H. Cartan, La transgression dans un groupe de Lie et dans un espace fibr´e principal. (French) Colloque de topologie (espaces fibrés), Bruxelles, 1950, pp. 57–71. [2,](#page-1-1) [34,](#page-33-1) [39](#page-38-3) [CM] P. Chatterjee and C. Maity, On the second cohomology of nilpotent orbits in exceptional Lie algebras, Bull. Sci. Math. 141 (2017), no. 1, 10–24. [2](#page-1-1)

- [Fr] M. Franz, The cohomology rings of homogeneous spaces. J. Topol., 14 (2021), no. 4, 1396– 1447. [2](#page-1-1)
- [Go] V. V. Gorbatsevich, Compact homogeneous manifolds of dimension at most 7 up to a finite-sheeted covering. (Russian) Izv. Ross. Akad. Nauk Ser. Mat. 76 (2012), no. 4, 27–40; translation in Izv. Math. 76 (2012), no. 4, 669–680. [4](#page-3-5)
- [GHV-II] W. Greub, S. Halperin and R. Vanstone, Connections, curvature, and cohomology. Vol. II: Lie groups, principal bundles, and characteristic classes, Pure and Applied Mathematics, Vol. 47-II. Academic Press, New York-London, 1973. [5,](#page-4-3) [6,](#page-5-2) [35](#page-34-5)
- [GHV-III] W. Greub, S. Halperin and R. Vanstone, Connections, curvature, and cohomology, Volume III: Cohomology of principal bundles and homogeneous spaces, Pure and Applied Mathematics, Vol. 47-III. Academic Press, New York-London, 1976. [6,](#page-5-2) [9,](#page-8-4) [10,](#page-9-6) [15,](#page-14-6) [34,](#page-33-1) [35,](#page-34-5) [36,](#page-35-5) [37,](#page-36-2) [38,](#page-37-5) [39](#page-38-3)
- [GS] V. Guillemin and S. Sternberg, Supersymmetry and equivariant de Rham theory, Springer-Verlag, Berlin, 1999. [2](#page-1-1)
- [He] S. Helgason, Differential Geometry, Lie Groups, and Symmetric Spaces, Academic Press, New York, 1978. [24](#page-23-1)
- [Ho] G. Hochschild, The structure of Lie groups, Holden-Day, Inc., San Francisco, 1965. [28,](#page-27-4) [30,](#page-29-2) [31](#page-30-1)
- [Kn] A. W. Knapp, Lie groups beyond an introduction, Second edition, Progress in Mathematics, Vol. 140, Birkhäusar, Basel, 2002. [28,](#page-27-4) [32,](#page-31-1) [34](#page-33-1)
- [KN] S. Kobayashi and K. Nomizu, Foundations of differential geometry, Vol. I. Reprint of the 1963 original. Wiley Classics Library. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1996. [36](#page-35-5)
- [Mo] G. D. Mostow, On covariant fiberings of Klein spaces, Amer. Jour. Math. 77 (1955), 247– 278. [28](#page-27-4)
- [Sp] M. Spivak, A comprehensive introduction to differential geometry. Vol. IV, Publish or Perish, Inc., Boston, Mass., 1975. [35](#page-34-5)
- [Tu] Loring W. Tu, Introductory lectures on equivariant cohomology, with appendices by Tu and Alberto Arabia. Annals of Mathematics Studies, 204. Princeton University Press, Princeton, NJ, 2020. [2](#page-1-1)
- [Wo] J. Wolf, The cohomology of homogeneous spaces, Amer. J. Math. 99 (1977), no. 2, 312–340. [2](#page-1-1)

School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India

Email address: indranil@math.tifr.res.in

The Institute of Mathematical Sciences, HBNI, CIT Campus, Tharamani, Chennai 600113, India

Email address: pralay@imsc.res.in

Indian Institute of Science Education and Research (IISER) Mohali, Knowledge City, Sector 81, S.A.S. Nagar 140306, Punjab, India

 $\emph{Email address:}$ maity.chandan1@gmail.com , cmaity@iisermohali.ac.in