




Unpacking merger jets: a Bayesian analysis of GW170817, GW190425 and electromagnetic observations of short gamma-ray bursts

FERGUS HAYES ¹, IK SIONG HENG ¹, GAVIN LAMB ^{2,3}, EN-TZU LIN ⁴, JOHN VEITCH ¹ AND MICHAEL J. WILLIAMS ¹

¹*SUPA, School of Physics and Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK*

²*Astrophysics Research Institute, Liverpool John Moores University, IC2 Liverpool Science Park, 146 Brownlow Hill, Liverpool, L3 5RF, UK*

³*School of Physics and Astronomy, University of Leicester, University Road, Leicester, LE1 7RH, UK*

⁴*Institute of Astronomy, National Tsing Hua University, Hsinchu 30013, Taiwan*

ABSTRACT

We present a novel fully Bayesian analysis to constrain short gamma-ray burst jet structures associated with cocoon, wide-angle and simple top-hat jet models, as well as the binary neutron star merger rate. These constraints are made given the distance and inclination information from GW170817, observed flux of GRB 170817A, observed rate of short gamma-ray bursts detected by Swift, and the neutron star merger rate inferred from LIGO’s first and second observing runs. A separate analysis is conducted where a fitted short gamma-ray burst luminosity function is included to provide further constraints. The jet structure models are further constrained using the observation of GW190425 and we find that the assumption that it produced a GRB 170817-like short gamma-ray burst that went undetected due to the jet geometry is consistent with previous observations. We find and quantify evidence for low luminosity and wide-angled jet structuring in the short gamma-ray burst population, independently from afterglow observations, with log Bayes factors of 0.45–0.55 for such models when compared to a classical top-hat jet. Slight evidence is found for a Gaussian jet structure model over all others when the fitted luminosity function is provided, producing log Bayes factors of $0.25–0.9 \pm 0.05$ when compared to the other models. However without considering GW190425 or the fitted luminosity function, the evidence favours a cocoon-like model with log Bayes factors of 0.14 ± 0.05 over the Gaussian jet structure. We provide new constraints to the binary neutron star merger rates of $1–1300 \text{ Gpc}^{-3} \text{ yr}^{-1}$ or $2–680 \text{ Gpc}^{-3} \text{ yr}^{-1}$ when a fitted luminosity function is assumed.

Keywords: Gamma-ray bursts (629), Gravitational waves (678)

1. INTRODUCTION

The joint detection of both gravitational wave (GW) GW170817 (Abbott et al. 2017a), and counterpart short gamma-ray burst (sGRB) GRB 170817A (Goldstein et al. 2017; Savchenko et al. 2017), followed by the detection of kilonova AT 2017gfo (McCully et al. 2017; Evans et al. 2017) not only solidified the belief that sGRBs are produced from the merger of binary neutron star (BNS) systems, but also began the era of GW multimessenger astronomy (Abbott et al. 2017a). The combination of both the and GW data gave insight into various problems that a detection through a single data channel could not provide; ranging from cosmology (Abbott et al. 2017b), the origin of the abundance

of heavy elements in the universe (Tanvir et al. 2017), tests of general relativity and the speed of gravity (Abbott et al. 2017c) among others.

However the detection of the event not only provided answers but also provoked questions, as it was inferred through GW parameter inference that the event was exceptionally nearby at only 40 Mpc (Abbott et al. 2019), giving an observed isotropic luminosity of the event of $10^{47} \text{ erg s}^{-1}$, three orders of magnitude lower than that observed for any other sGRB (Abbott et al. 2017c). It was also inferred through GW parameter inference that the event was viewed at a wide angle of $14–41^\circ$ from the central axis (Abbott et al. 2019). This led to the hypothesis that the jet of GRB 170817A exhibited some wide-angle structure to produce the observed flux, and that it may still have had a typically luminous central jet component. The long duration observations of the event’s afterglow across the electromagnetic (EM) spec-

trum provided further evidence for this claim (Troja et al. 2017; Margutti et al. 2018; Lyman et al. 2018; D’Avanzo et al. 2018; Mooley et al. 2018; Ruan et al. 2018; Troja et al. 2018; Alexander et al. 2018; Troja et al. 2020).

While evidence for this wide-angled structure is provided by the multitude of afterglow observations, the functional form of the luminosity profile over viewing angle remains in question (e.g. Granot et al. 2017; Duffell et al. 2018; Gottlieb et al. 2018; Lamb & Kobayashi 2018; Mooley et al. 2018; Troja et al. 2018; Ioka & Nakamura 2019; Beniamini et al. 2019; Fraija et al. 2019; Lamb et al. 2019; Salafia et al. 2019; Biscoveanu et al. 2020; Lamb et al. 2020; Takahashi & Ioka 2021). Accurate modelling of this jet structure is important in both understanding the astrophysics of the event, and in preventing systematic biases in any multimessenger analysis that must make assumptions about the jet geometry (Nakar & Piran 2021; Lamb et al. 2021), which include constraints on the Hubble constant (Abbott et al. 2017b) and the BNS merger rate (Wanderman & Piran 2015).

With the promise of future GW BNS merger detections (Abbott et al. 2020b), a jet structure model that best represents the data of joint detection events should be discerned (Hayes et al. 2020). However, current analyses are limited to the data provided by GW170817/GRB 170817A (including AT 2017gfo), GW BNS merger event GW190425 (Abbott et al. 2020a), as well as the population of sGRBs detected independently of GW detection (Poolakkil et al. 2021; Lien et al. 2016). Previous similar work has considered constraining jet structure models with joint GW and EM detections using a Bayesian analysis (Biscoveanu et al. 2020; Hayes et al. 2020; Farah et al. 2020). The historical rate of detected sGRB and GWs has also been used in Bayesian analyses to constrain the jet structure (Williams et al. 2018; Sarin et al. 2022). Work by Mogushi et al. (2019) and Tan & Yu (2020) combine the detection rates of sGRB and GWs with the prompt emission data of GRB 170817A and the parameter inference results of GW170817, along with assuming a luminosity function fitted by short gamma-ray burst events with known redshift. The jet structure has been constrained given the weak gamma-ray emission detected by the *INTEGRAL* detector coincident with the gravitational wave event GW190425 in the work by Saleem et al. (2020). In this work, we put forward a comprehensive Bayesian framework that combines the prompt emission data from GRB 170817A, GW parameter inference posteriors of GW170817 and GW190425, GW informed BNS merger rate, as well as the detection rate of sGRBs by the Neil Gehrels Swift Observatory (Swift)

detector. This analysis is then further combined with the information provided by a luminosity function fitted by sGRB events with known redshifts to provide tighter constraints, with the caveat of also introducing bias into the analysis. We provide parameter constraints and model comparison results between a classical top-hat jet structure and three different jet structures with wide-angled structuring: a Gaussian, power-law and double Gaussian jet. These results are presented alongside constraints on the intrinsic luminosity (when it is not fitted by the luminosity function), and the merger rate for both cases.

The physical model assumed is detailed in Section 2, before the analysis method and data are laid out in Section 3. In Section 4 we report the results of the analysis, both for when the fitted luminosity function is incorporated and when it is not. In Section 5 the implications of the results are discussed and a conclusion provided in Section 6.

2. BACKGROUND

The sGRB data consists of the observed T90 integrated flux F as well as the number of observed sGRBs N_{EM} . The average T90 integrated flux \hat{F} is related to the isotropic equivalent luminosity L_{iso} at a given viewing angle θ_v , as well as a redshift z dependent luminosity distance d_L and k -correction k :

$$\hat{F} = \frac{L_{\text{iso}}(\theta_v)}{4\pi d_L(z)^2 k(z)}. \quad (1)$$

The mean number of sGRBs \hat{N}_{EM} observed by a detector within a duration T that covers an area of sky equal to $\Delta\Omega$ depends on the redshift, viewing angle and intrinsic luminosity $\Lambda = \{z, \theta_v, L_0\}$ through the number density:

$$\hat{N}_{\text{EM}} = \int \frac{d\hat{N}_{\text{EM}}}{d\Lambda} \left[\int p(D_{\text{EM}}|\Lambda) dD_{\text{EM}} \right] d\Lambda, \quad (2)$$

where D_{EM} denotes the selection effects of the detector, such that $D_{\text{EM}} = 1$ if a detection is made and $D_{\text{EM}} = 0$ otherwise.

We consider sGRBs detected by the Swift instrument, which has a detector response determined empirically in (Lien et al. 2014) to fit:

$$\int p(D_{\text{EM}}|\Lambda) dD_{\text{EM}} = \begin{cases} 0 & \text{if } \hat{F} < F_{\text{thr}}, \\ a \frac{b+c\hat{F}/F_0}{1+\hat{F}/dF_0} & \text{else,} \end{cases} \quad (3)$$

where $a = 0.47$, $b = -0.05$, $c = 1.46$, $d = 1.45$, $F_0 = 6 \times 10^{-6} \text{ erg s}^{-1} \text{ cm}^{-2}$ and $F_{\text{thr}} = 5.5 \times 10^{-9} \text{ erg s}^{-1} \text{ cm}^{-2}$.

The relation between the number density and the physical parameters of Λ is:

$$\frac{d\hat{N}_{\text{EM}}}{d\Lambda} = T \frac{\Delta\Omega}{8\pi} \frac{\mathcal{R}_{\text{GRB}}(z)}{1+z} \frac{dV(z)}{dz} p(L_0|\Sigma) \sin\theta_v, \quad (4)$$

where $V(z)$ is the co-moving volume, \mathcal{R}_{GRB} is the rate of sGRBs and $p(L_0|\Sigma)$ is the intrinsic luminosity function given the hyperparameter Σ .

2.1. Short gamma-ray burst rate

The rate of sGRBs is assumed to be in the form:

$$\mathcal{R}_{\text{GRB}}(z) = R_{\text{BNS}} R_{\text{GRB}}(z), \quad (5)$$

where R_{BNS} is the local rate of BNS mergers and $R_{\text{GRB}}(z)$ is defined so that $R_{\text{GRB}}(0) = 1$. This assumes that every BNS merger results in a sGRB, and that the number of sGRBs produced by neutron star-black hole mergers is negligible.

The form of $R_{\text{GRB}}(z)$ can be assumed to follow the star formation rate $R_*(z)$ convolved with the probability distribution of the delay time between the system formation and the eventual merger that leads to the sGRB $P(t)$ (Wanderman & Piran 2015):

$$R_{\text{GRB}}(z) \propto \int_{t_{\text{min}}}^{T(\infty)-T(z)} \frac{R_*(z_*)}{1+z_*} P(t) dt, \quad (6)$$

where $z_* = z(T(z) + t)$ is the redshift when the system was formed, $T(z)$ is the *look-back time* and t_{min} is the minimum delay time. This minimum delay time is set to 20 Myr and $P(t) \propto 1/t$ according to (Guetta & Piran 2006). The star formation rate is assumed to be of the form (Cole et al. 2001):

$$R_*(z) \propto \frac{a + bz}{1 + (z/c)^d} H(z), \quad (7)$$

where the parameter values are taken from (Hopkins & Beacom 2006) to be $a = 0.017$, $b = 0.13$, $c = 3.3$ and $d = 5.3$.

2.2. Cosmology

A flat, vacuum dominated universe is assumed. The co-moving volume distribution over redshift is defined:

$$\frac{dV(z)}{dz} = 4\pi \frac{c}{H(z)} \left(\frac{d_L(z)}{1+z} \right)^2. \quad (8)$$

For a flat cosmology, the luminosity distance is related to the redshift by:

$$d_L(z) = (1+z) \frac{c}{H_0} \int_0^z \frac{H_0}{H(z')} dz', \quad (9)$$

where H_0 is the Hubble constant and $H(z)$ is equal to:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}. \quad (10)$$

Here $\Omega_m = 0.308$ and $\Omega_\Lambda = 0.692$ are the matter density and dark energy density respectively (Hogg 1999), with values taken from (Adam et al. 2016) along with $H_0 = 67.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

The look-back time, defined as the time between when a source emits light at redshift z and the time it is detected, is then:

$$T(z) = \frac{1}{H_0} \int_0^z \frac{H_0}{(1+z')H(z')} dz'. \quad (11)$$

For a flat, vacuum dominated universe, the inverse function has an analytical expression (Petrillo et al. 2013):

$$z(T) = \left(\frac{\Omega_\Lambda}{\Omega_m} \right)^{1/3} \left[\left(\frac{1+W(T)}{1-W(T)} \right)^2 - 1 \right]^{1/3} - 1, \quad (12)$$

where:

$$W(T) = \exp \left[\ln \left(\frac{1+\sqrt{\Omega_\Lambda}}{1-\sqrt{\Omega_\Lambda}} \right) - 3H_0\sqrt{\Omega_\Lambda}T \right]. \quad (13)$$

The k -correction accounts for the cosmological redshifting in the intrinsic sGRB spectrum with respect to the detector's spectrum (Bloom et al. 2001):

$$k(z) = \frac{\int_{\nu_{s,1}/(1+z)}^{\nu_{s,2}/(1+z)} \nu f(\nu) d\nu}{\int_{\nu_1}^{\nu_2} \nu f(\nu) d\nu}. \quad (14)$$

Here we assume the form of $f(\nu)$ follows the Band function described in (Band et al. 1993).

2.3. Intrinsic and isotropic equivalent luminosity

The luminosity structure of a gamma-ray burst is defined to be:

$$L(\theta) = L_0 y_L(\theta), \quad (15)$$

where L_0 is the intrinsic luminosity at $\theta = 0$. It is assumed that the distribution of intrinsic luminosity L_0 follows a Schechter function:

$$p(L_0|\Sigma = \{L_0^*, \gamma\}) \propto \left(\frac{L_0}{L_0^*} \right)^{-\gamma} e^{-L_0/L_0^*}, \quad (16)$$

for $L_{\text{min}} \leq L_0 \leq L_{\text{max}}$ where $L_{\text{min}} = 10^{-3}L_0^*$ and $L_{\text{max}} = 10^2L_0^*$.

Similarly the Lorentz factor's dependence over angle follows:

$$\Gamma(\theta) = (\Gamma_0 - 1)y_\Gamma(\theta) + 1, \quad (17)$$

with Γ_0 being the Lorentz factor of the jet at $\theta = 0$. Given these definitions, both y_L and y_Γ are defined to equal 1 at $\theta = 0$.

The Lorentz factor determines the degree of relativistic beaming, which for the luminosity is governed by:

$$\mathcal{B}_L(\theta_v, \theta) = \frac{1}{4\Gamma(\theta)^6(A^2 - B^2)^2} \left(5 \left(\frac{A}{\sqrt{A^2 - B^2}} \right)^3 + 3 \left(\frac{A}{\sqrt{A^2 - B^2}} \right) \right), \quad (18)$$

with $A = 1 - \beta \cos \theta \cos \theta_v$ and $B = -\beta \sin \theta \sin \theta_v$ where $\beta = \sqrt{1 - \Gamma(\theta)^{-2}}$.

The apparent isotropic equivalent luminosity, for an observer at θ_v from the jet axis, can then be related to the intrinsic luminosity via the beaming function by combining Eqn. 15 and Eqn. 18:

$$L_{\text{iso}}(\theta_v) = L_0 \int_0^{\theta_j} \frac{1}{2} \mathcal{B}(\theta_v, \theta) y_L(\theta) \sin \theta d\theta, \quad (19)$$

where θ_j is the maximum angle for which the beaming of gamma-rays occurs. A conservative maximum outer jet angle for the emission of gamma-rays is approximated by considering scattering by electrons accompanying baryons within the jet. The condition is given by (Matsumoto et al. 2019; Lamb et al. 2022),

$$\Gamma(\theta_j) \simeq 19.1 \left(\frac{L(\theta_j)}{10^{51} \text{ erg/s}} \right)^{1/6}, \quad (20)$$

Beyond this limit, $\theta > \theta_j$, the jet becomes opaque to gamma-rays.

2.4. Jet structures

The implications of structuring within compact stellar merger jets for the EM counterparts from GW detected systems has been highlighted in the literature (Lamb & Kobayashi 2017; Lazzati et al. 2017; Kathirgamaraju et al. 2018; Beniamini et al. 2020); here we choose a sample of fiducial jet structure models that are representative of the literature diversity.

The top-hat jet (TH) is the simplest structure, where the beam is uniform until the jet opening angle θ_j where the jet sharply cuts off:

$$y_L(\theta) = \begin{cases} 1 & \text{if } 0 \leq \theta \leq \theta_j, \\ 0 & \text{else.} \end{cases} \quad (21)$$

$$y_\Gamma(\theta) = \begin{cases} 1 & \text{if } 0 \leq \theta \leq \theta_j, \\ 0 & \text{else.} \end{cases}$$

We note that the condition expressed in Eqn. 20 is not enforced for this case, as Γ_0 is above the Eqn. 20 limit at all points within the jet.

Wide-angle structure can be introduced with a Gaussian jet (GJ) structure, described by a single width parameter θ_σ (e.g. Rossi et al. 2002, 2004; Zhang & Meszaros 2002; Kumar & Granot 2003):

$$y_L(\theta) = e^{-\frac{1}{2} \left(\frac{\theta}{\theta_\sigma} \right)^2}, \quad y_\Gamma(\theta) = e^{-\frac{1}{2} \left(\frac{\theta}{\theta_\sigma} \right)^2}. \quad (22)$$

An alternative to the Gaussian profile has the wide-angle emission expressed as a three parameter power-law jet (PL) structure (e.g. Kumar & Granot 2003; Zhang et al. 2004; Rossi et al. 2004), where the jet can be described by some uniform core out to width θ_c , and then the intrinsic luminosity structure falls off at wide angles according to power s and the Lorentz factor with a :

$$y_L(\theta) = \begin{cases} 1 & \text{if } 0 \leq \theta \leq \theta_c, \\ \left(\frac{\theta}{\theta_c} \right)^{-s} & \text{else.} \end{cases} \quad (23)$$

$$y_\Gamma(\theta) = \begin{cases} 1 & \text{if } 0 \leq \theta \leq \theta_c, \\ \left(\frac{\theta}{\theta_c} \right)^{-a} & \text{else.} \end{cases}$$

Finally, let us consider a two component, or double Gaussian jet (DG), with emission from both an inner core described by a Gaussian structure of width θ_{in} and an outer cocoon described by width θ_{out} (Salafia et al. 2020):

$$y_L(\theta) = (1 - \mathcal{C}) e^{-\frac{1}{2} \left(\frac{\theta}{\theta_{\text{in}}} \right)^2} + \mathcal{C} e^{-\frac{1}{2} \left(\frac{\theta}{\theta_{\text{out}}} \right)^2},$$

$$y_\Gamma(\theta) = \frac{(1 - \mathcal{C}) e^{-\frac{1}{2} \left(\frac{\theta}{\theta_{\text{in}}} \right)^2} + \mathcal{C} e^{-\frac{1}{2} \left(\frac{\theta}{\theta_{\text{out}}} \right)^2}}{(1 - \frac{\mathcal{C}}{\mathcal{A}}) e^{-\frac{1}{2} \left(\frac{\theta}{\theta_{\text{in}}} \right)^2} + \frac{\mathcal{C}}{\mathcal{A}} e^{-\frac{1}{2} \left(\frac{\theta}{\theta_{\text{out}}} \right)^2}}. \quad (24)$$

The luminosity of the outer cocoon is equal to $\mathcal{C}L_0$ and the Lorentz factor $\mathcal{A}(\Gamma_0 - 1) + 1$.

We do not consider hollow-cone jet structure models in our study (see e.g., Nathanail et al. 2021; Takahashi & Ioka 2021); we expect that the combination of our intrinsic luminosity distribution (Eqn. 16) and the beaming (Eqn. 19) will wash-out the effect of any hollow-cone structuring within the core. For this study, the structure outside of the jet's core is the critical component.

3. BAYESIAN FRAMEWORK

Constraints are placed on model parameters λ of a model M when given data \mathcal{D} in Bayesian data analysis by determining the posterior distribution using *Bayes theorem*:

$$p(\lambda|\mathcal{D}, M) = \frac{\mathcal{L}(\mathcal{D}|\lambda)\pi(\lambda)}{p(\mathcal{D}|M)}, \quad (25)$$

where \mathcal{L} is the likelihood, π is the prior and the normalisation term $p(\mathcal{D}|M)$ is the evidence. Consider comparing two models M_1 and M_2 when given data \mathcal{D} . In the

context of Bayesian data analysis, the statistic used to compare two models is the *posterior odds* defined:

$$\mathcal{O}_{12} = \frac{p(M_1|\mathcal{D})}{p(M_2|\mathcal{D})} = \frac{p(M_1)p(\mathcal{D}|M_1)}{p(M_2)p(\mathcal{D}|M_2)}. \quad (26)$$

Normally we are interested in cases where the *a priori* probability of either model being correct is comparable, and therefore the posterior odds is dominated by the *Bayes factor*:

$$\mathcal{B}_{12} = \frac{p(\mathcal{D}|M_1)}{p(\mathcal{D}|M_2)}, \quad (27)$$

which quantifies the contribution to the posterior odds given by the data \mathcal{D} . A value of $\ln \mathcal{B}_{12} > 0$ favours M_1 , while $\ln \mathcal{B}_{12} < 0$ favours M_2 .

The analysis is performed by applying the model described in Section 2 with an assumed jet structure from Section 2.4 given both GW and sGRB prompt emission data.

Table 1 lists the notation used in the following section. The data can be split into that produced by a GW-triggered event, denoted with the subscript ‘GW’, and that produced from an EM trigger, denoted with the subscript ‘EM’.

The data from the N_{GW} GW-triggered events consists of the GW strain x_{GW} and the flux of the counterpart F_{GW} . The GW-triggered events may not necessarily require a counterpart to be considered for the analysis. If the sky localisation of the source coincides with the sky coverage of gamma-ray burst detectors then we can assume that it was not detected due to its distance and orientation to us. The current events that meet this criteria are both GW170817 with GRB 170817A as well as GW190425 and the non-detection of its counterpart, under the assumption that a sGRB was produced, given the Fermi detector covered 50% of the sky localisation and Konus-*Wind* covered the entire sky (Hosseinzadeh et al. 2019).

The EM-triggered events are simply the number of sGRB detections that Swift made within a 10 year operational period N_{EM} .

The likelihood can be decoupled into two terms, one of which considers GW-triggered events and the other EM-triggered:

$$\mathcal{L} = \mathcal{L}_{\text{EM}}\mathcal{L}_{\text{GW}}. \quad (28)$$

The likelihood of the EM-triggered events is a Poisson distribution with a mean given in Eqn. 2:

$$\mathcal{L}_{\text{EM}} = \frac{\hat{N}_{\text{EM}}(\Theta, R_{\text{BNS}}, \Sigma)^{N_{\text{EM}}} e^{-\hat{N}_{\text{EM}}(\Theta, R_{\text{BNS}}, \Sigma)}}{N_{\text{EM}}!}. \quad (29)$$

The mean is evaluated over a regular grid of shape $(z, (2\theta/\pi)^{1/3}, \log_{10} L_0) = (\times 50, \times 100, \times 1000)$. The angular grid points were chosen to be distributed over a

Variable	Description
x_{GW}	GW detector data
F_{GW}	sGRB detector data
N_{GW}	Number of detected GWs
N_{EM}	Number of detected sGRBs
Σ	Luminosity function hyperparameters
R_{BNS}	BNS merger rate
Θ	Jet structure parameters
Φ	$\{\theta_v, d_L\}$
L_0	Intrinsic on-axis luminosity

Table 1. Shorthand notation of the GW and EM data as well as sets of parameters of interest.

Data set	Data
\mathcal{D}_{170817}	$\{x_{170817}, F_{170817}\}$
\mathcal{D}_{190425}	$\{x_{190425}, F_{190425}\}$
\mathcal{D}_R	$\{N_{\text{EM}}, N_{\text{GW}}\}$
$\mathcal{D}_{170817+R}$	$\{\mathcal{D}_{170817}, \mathcal{D}_R\}$
\mathcal{D}_{all}	$\{\mathcal{D}_{170817}, \mathcal{D}_{190425}, \mathcal{D}_R\}$

Table 2. Summary of the data used in the analysis.




Figure 1. High-level Bayesian network of the model described in Section 2. The variable names are defined in Table 1.

power-law so as to populate low θ areas of the parameter space with grid points, while also maintaining a relatively high density of points at wider angles where emission from some jet structures is still significant.

The GW-triggered events likelihood is the product of each of the N_{GW} events:

$$\mathcal{L}_{\text{GW}} \propto \prod_{i=1}^{N_{\text{GW}}} \sum_{j=1}^S p(F_{i,\text{GW}}|\Theta, \Phi_{i,j}, L_{j,0}), \quad (30)$$

where S samples are taken of $\Phi_{i,j}$ and $L_{j,0}$ from $p(\Phi_i, L_0|x_{i,\text{GW}}, \Sigma)$. The parameters can be sampled from separate distributions $p(\Phi_i|x_{i,\text{GW}})$ and $p(L_0|\Sigma)$ respectively, where $p(\Phi_i|x_{i,\text{GW}})$ are samples from the posteriors produced from GW parameter estimation for each event. The likelihood of the prompt emission of the GW-triggered events is assumed to be a Gaussian distribution of width σ_F about a mean described in Eqn. 1:

$$p(F_{\text{GW}}|\Theta, \Phi, L_0) = \frac{1}{\sqrt{2\pi\sigma_F^2}} \exp\left(-\frac{(F_{\text{GW}} - F)^2}{2\sigma_F^2}\right). \quad (31)$$

The priors for the model are specified in Table 3 for each of the gamma-ray burst rate, luminosity function and jet structure parameters. A normal distribution is denoted $\mathcal{N}(\mu, \sigma)$ with a mean of μ and standard deviation of σ , a uniform distribution as $\mathcal{U}(A, B)$ with lower bound of A and upper bound of B , a Gamma distribution and inverse Gamma distribution as $\Gamma(\alpha, \beta)$ and $\Gamma^{-1}(\alpha, \beta)$ with a shape of α and a scale of β . We assume that every BNS merger results in a gamma-ray burst so that $\epsilon_{\text{BNS}} = 1$.

Model	Parameter	Prior
	$\log_{10} R'_{\text{BNS}}$	$\mathcal{N}(-6.6, 0.77)$
	$\log_{10} L_0^*$	$\mathcal{N}(51.6, 1)$
	γ	$\mathcal{U}(0, 1)$
	$\log_{10} \Gamma_0$	$\text{Gamma}^{-1}(2, 2.5 \times 10^{-3})$
TH	θ_j	$\mathcal{U}(0, \pi/2)$
GJ	θ_σ	$\mathcal{U}(0, \pi/2)$
	θ_c	$\mathcal{U}(0, \pi/2)$
PL	s	$\text{Gamma}(2, 4)$
	a	$\text{Gamma}(2, 1)$
	θ_{in}	$\mathcal{U}(0, \pi/2)$
DG	θ_{out}	$\mathcal{U}(0, \theta_{\text{out}})$
	$\log_{10} \mathcal{C}$	$\mathcal{U}(-6, 0)$
	$\log_{10} \mathcal{A}$	$\mathcal{U}(-6, 0)$

Table 3. Assumed prior distributions for each parameter. The analysis assumes one jet structure out of the top-hat (TH), Gaussian (GJ), power-law (PL) and double Gaussian (DG) models. Some parameters are made unit-less so that $R'_{\text{BNS}} = R_{\text{BNS}}/\text{Mpc}^{-3} \text{yr}^{-1}$ and $L_0^* = L_0^*/\text{erg s}^{-1}$.

The T90 integrated flux of GRB170817A in the Fermi detector's 50 – 300 keV band is set at $F_{170817} = 1.4 \times 10^{-7} \text{ erg s}^{-1} \text{ cm}^{-2}$ with an uncertainty of $\sigma_{170817} = 3.64 \times 10^{-8} \text{ erg s}^{-1} \text{ cm}^{-2}$ (Goldstein et al. 2017). For the unobserved counterpart of GW190425, it is assumed that the T90 integrated flux takes a value of zero with an uncertainty of $\sigma_{190425} = 10^{-8} \text{ erg s}^{-1} \text{ cm}^{-2}$ as a conservative upper bound to the Fermi detector's detection

threshold (Tan & Yu 2020). The distance and viewing angle posteriors of GW170817 and GW190425 are each represented by 500 samples taken from the their respective parameter estimation data releases. In this work we consider an observing period of approximately 9.8 years by the Swift detector in which it observed $N_{\text{EM}} = 107$ sGRBs as recorded by Lien et al. (2016), given a sky coverage of $\Delta\Omega = 0.1$. The log prior on the rate of BNS mergers of $\mathcal{N}(-6.6, 0.77)$ is roughly chosen to reflect the constraints imposed to the rates by GWTC-2 (Abbott et al. 2020c). The log prior on L_0^* is centred around the fitted value taken from Mogushi et al. (2019) with a standard deviation set to span one order of magnitude. This is chosen to reflect some prior information in the allowed luminosity from prior observations, but with a width to allow for flexibility into higher or lower luminosity regimes.

Posteriors and Bayes factors are calculated from Eqn. 25 and Eqn. 27 by assigning \mathcal{D} and λ as the variables in Table 1. We collect the data into three sets: one only given the number of Swift detections N_{EM} and GW detections N_{GW} called \mathcal{D}_R , another with the combined GW170817 GW x_{170817} and EM data F_{170817} called \mathcal{D}_{170817} , and the other with GW190425 GW data x_{190425} and the flux from the non-detection F_{190425} called \mathcal{D}_{190425} . The analysis is performed on five combinations of these data sets: $\mathcal{D}_{170817} = \{x_{170817}, F_{170817}\}$, $\mathcal{D}_{190425} = \{x_{190425}, F_{190425}\}$, $\mathcal{D}_R = \{N_{\text{EM}}, N_{\text{GW}}\}$, $\mathcal{D}_{170817+R} = \{\mathcal{D}_{170817}, \mathcal{D}_R\}$, and $\mathcal{D}_{\text{all}} = \{\mathcal{D}_{170817}, \mathcal{D}_{190425}, \mathcal{D}_R\}$. We allow that $\lambda = \{\Sigma, \mathcal{R}_0, \Theta_M\}$, where Θ_M are the jet structure model parameters dependent on jet structure model M . The three analyses are repeated for each of the jet structure models: $M = \text{TH, GJ, PL and DG}$.

The posterior samples and evidence for each case are calculated via the nested sampling algorithm NESSAI, that utilises machine learning techniques to drastically reduce the number of evaluations of the expensive likelihood function (Williams et al. 2021).

4. RESULTS

The analysis that is described in the previous section is applied to all three sets of data. The full corner plots for each jet structure model are shown in the appendix, where Figure A1 shows the results for the top-hat jet, Figure A2 the Gaussian jet, Figure A3 the power-law jet and Figure A4 the double Gaussian jet structure model. The posteriors for each of the data sets is overlaid upon one another where \mathcal{D}_{170817} is shown in red, \mathcal{D}_{190425} in orange, \mathcal{D}_R in violet, $\mathcal{D}_{170817+R}$ in blue and \mathcal{D}_{all} in black. The log evidence $\ln p(\mathcal{D}|M)$ that corresponds to each posterior is shown in Table A1 for each of the five data




Figure 2. Posterior distributions of (a) the rate of BNS mergers and (b) the mean intrinsic luminosity \hat{L}_0 assuming the top-hat jet (TH, pink), Gaussian jet (GJ, green), power-law jet (PL, orange), and double Gaussian jet (DG, blue) models when given \mathcal{D}_{all} . The solid vertical lines represent the minimum and maximum of each distribution while the thickness of the fill in between represents the probability density. The dashed vertical lines represent 90% credible intervals. The median is shown by the middle solid line. The top distribution represents the prior distribution taken from Table 3 for $\log_{10} R_{\text{BNS}}$, L_0 and γ respectively. Similar posterior distributions are recovered for models with wide-angled structuring in comparison to the top-hat model case.

$\ln \mathcal{B}_{\text{row,col}}$	Top-hat	Gaussian	Power-law	Double Gaussian
Top-hat	0	-0.54 / -0.91	-0.46 / -0.45	-0.55 / -0.64
Gaussian	0.54 / 0.91	0	0.08 / 0.46	-0.01 / 0.27
Power-law	0.46 / 0.45	-0.08 / -0.46	0	-0.09 / -0.19
Double Gaussian	0.55 / 0.64	0.01 / -0.27	0.09 / 0.19	0

Table 4. The log Bayes factor $\ln \mathcal{B}$ between each model when given \mathcal{D}_{all} w/o/with the fitted luminosity function. A positive value is evidence towards the model of the row while a negative is evidence that favours the column model. Slight evidence is provided for models with wide-angled structuring over the top-hat model while little evidence distinguishes between the power-law, Gaussian and double Gaussian models without the fitted luminosity function. When the fitted luminosity function is included, slight evidence is further provided in favour of the Gaussian jet over the double Gaussian and power-law jet structures while the top-hat remains least favoured. All values can be assumed to have uncertainties of ± 0.05 .

sets over the four jet structure models. The log Bayes factors between the different models given the same data

set can simply be calculated by taking the difference between entries of the same row.

Much of the discussion in this section concerns the posterior constraints when all of the data is considered $\mathcal{D} = \mathcal{D}_{\text{all}}$, however the outcomes given the other subsets are considered to explain these results and provide further insight.

The log Bayes factors between the jet structure models are given as the left-hand entries of Table 4 when given the \mathcal{D}_{all} data. A positive value indicates that the data supports the model of the row while a negative value supports the column model. Evidently the top-hat model is less favourable than the models with wide-angled jet structuring, with Bayes factors of -0.54 , -0.46 and -0.55 between it and the Gaussian, power-law and Double Gaussian jet structures respectively. The log Bayes factors between the power-law, Gaussian and the double Gaussian is slight, with only insignificant evidence in favour of the Gaussian and double Gaussian model of log Bayes factors of less than 0.1, and negligibly small log Bayes factors between the two.

The constraints on the rate of BNS mergers when given \mathcal{D}_{all} are shown in Figure 4 for the four jet structure models. These constraints take the form of posterior distributions that are represented in the violin plots, where the outermost solid vertical lines indicate the minimum and maximum sample value while the fill in between represents the probability density. The 90% narrowest credible intervals are shown by the vertical dashed lines which enclose the median indicated by the middle solid line. The posterior distributions are compared to samples from the prior distribution at the top of the figure. For all cases, the posterior places tighter constraints on the merger rate than the prior distribution. The cases with wide-angled structuring (Gaussian, power-law and double Gaussian models) produce similar posterior distributions to one another, centred around a value of $\sim 10^{-7} \text{ Mpc}^{-3} \text{ yr}^{-1}$ consistent with the mean of the prior. The Gaussian jet structure produces the narrowest constraints with a 90% credible interval of $\log_{10} R_{\text{BNS}}/\text{Mpc}^{-3} \text{ yr}^{-1} = -6.7^{+0.7}_{-0.8}$, compared to the power-law and double Gaussian models of $-6.8^{+0.7}_{-0.9}$ and $-6.6^{+0.9}_{-0.9}$ respectively. The top-hat model favours lower rates of BNS mergers, and even pushes the lower bound on the 90% credible interval to lower values of that of the prior, constraining it between $\log_{10} R_{\text{BNS}}/\text{Mpc}^{-3} \text{ yr}^{-1} = -7^{+0.9}_{-1.0}$.

The median intrinsic luminosity posteriors determined for each model when given \mathcal{D}_{all} is shown in Figure 4. The mean intrinsic luminosity \hat{L}_0 is determined by drawing L_0 and γ from the respective posterior distribution and then drawing 1000 samples from the corresponding Schechter function of Eqn. 16 before finding the ensemble median. This process is then repeated for 2000

median intrinsic luminosity samples. These posteriors take a form similar to the rates posteriors in Figure 4 as violin plots where the shaded probability density is contained within the outermost maximum and minimum values indicated by the solid vertical lines, while the median is marked by the middle solid line. The median is enclosed by the narrowest 90% credible intervals displayed as dashed vertical lines. A distribution of prior samples of \hat{L}_0 is also plotted at the top of the figure, which is determined by sampling from the individual $\log_{10} L_0^*$ and γ priors defined in Table 3. The top-hat jet structure resembles the prior in width, but shifts to favour lower luminosity and exhibits some bimodality as the probability density pinches at the median. This is due to the bimodality of the L_0 posterior distribution in Figure A1 given $\mathcal{D}_{170817+R}$, which shall be discussed later in Section 5. The models with wide-angled structure tend towards lower mean intrinsic luminosity values, with the Gaussian model constrained to $\log_{10} \hat{L}_0/\text{ergs}^{-1} = 50.4^{+1.6}_{-1.3}$, power-law model $50.1^{+1.3}_{-1.6}$ and double Gaussian model of $50.2^{+1.5}_{-1.5}$. The top-hat model is constrained to $\log_{10} \hat{L}_0/\text{ergs}^{-1} = 50.2^{+1.7}_{-1.7}$ which we can compare to the prior of $51^{+1.8}_{-1.8}$.

Model	Parameter	Constraints	
		w/o fitted LF	Fitted LF
TH	θ_j	$14.9^{+46.0}_{-14.3}$	$9.2^{+17.1}_{-7.9}$
GJ	θ_σ	$5.9^{+28.2}_{-5.4}$	$4.2^{+5.2}_{-3.2}$
PL	θ_c	$10.6^{+32.9}_{-10.0}$	$6.0^{+9.2}_{-4.9}$
	s	$6.7^{+8.1}_{-5.2}$	$6.4^{+7.9}_{-4.3}$
	a	$1.5^{+2.2}_{-1.4}$	$1.5^{+2.1}_{-1.4}$
DG	θ_{in}	$6.3^{+24.1}_{-6.2}$	$3.6^{+6.2}_{-2.8}$
	θ_{out}	$49.3^{+40.7}_{-38.0}$	$46.2^{+43.1}_{-36.3}$
	$\log_{10} \mathcal{C}$	$-3.8^{+2.9}_{-2.2}$	$-4.1^{+1.9}_{-1.9}$
	$\log_{10} \mathcal{A}$	$-3.0^{+2.8}_{-2.9}$	$-3.1^{+2.9}_{-2.7}$

Table 5. Constraints placed on each of the variables for each model given data set \mathcal{D}_{all} with and without the fitted luminosity function (LF). The median of each posterior distribution is quoted along with upper and lower bounds placed by 90% credible intervals.

The constraints from the posteriors on the jet structure parameters given each jet structure model are shown in Table 5. The median is quoted along with the upper and lower bounds placed by the 90% narrowest credible intervals.

4.1. Fitted luminosity function

The analysis is repeated but instead of assuming a prior distribution on the luminosity scale and shape, a luminosity function fitted from the observed isotropic

equivalent luminosity of sGRBs with associated redshifts. The values of L_0 and γ are taken from the mean fitted Schechter function in Mogushi et al. (2019) of $\log_{10} L_0/\text{ergs}^{-1} = 51.6$ and $\gamma = 0.55$, fitted to the isotropic equivalent luminosity of 35 sGRBs.

The log evidence between each of the jet structure models and the different data sets are shown in Table A2, while the respective Bayes factors when given \mathcal{D}_{all} between each of the jet structure models are shown on the right-hand entries of Table 4.




Figure 3. Posterior distributions of the rate of BNS mergers assuming the top-hat jet (TH, pink), Gaussian jet (GJ, green), power-law jet (PL, orange) and double Gaussian jet (DG, blue) models when given \mathcal{D}_{all} and the fitted luminosity function. The solid vertical lines represent the minimum and maximum of each distribution while the thickness of the fill in between represents the probability density. The dashed vertical lines represent 90% credible intervals. The median is shown by the middle solid line. The posterior distributions are compared to samples from the prior of $\mathcal{N}(-6.6, 0.77)$ at the top. All models recover similar constraints that narrow about the mean value assigned to the prior.

The posteriors on the local rate of BNS merger when given the fitted luminosity function and \mathcal{D}_{all} are shown in Figure 3 in the same format as Figure 4, where the

widths of the violin plots indicate the probability density, the maximum and minimum sample is indicated by the extreme solid vertical lines, and median with the middle solid vertical line. The narrowest 90% credible intervals are indicated by the dashed vertical lines and are $\log_{10} R_{\text{BNS}}/\text{Mpc}^{-3} \text{yr}^{-1} = -7.0^{+0.7}_{-0.6}$, $-7.2^{+0.7}_{-0.8}$, $-6.8^{+0.8}_{-0.9}$ for the Gaussian, power-law and double Gaussian models respectively. For the top-hat jet model, the rate is constrained to $\log_{10} R_{\text{BNS}}/\text{Mpc}^{-3} \text{yr}^{-1} = -7.3^{+1.0}_{-0.9}$.

The constraints on the jet structure models when given the fitted luminosity function and \mathcal{D}_{all} are presented in Table 5 for each jet structure models, with the upper and lower bounds representing the narrowest 90% credible intervals.

5. DISCUSSION

5.1. Wide-angle jet structuring

The log Bayes factors are greater than 0.45 for all models with wide-angle jet structure when compared to the top-hat model. This is due to the top-hat jets failing to resolve the number of observed sGRBs with the flux of GRB 170817A when assuming that the event had a typical event opening angle. Given the assumed star formation rate and the constraints on the BNS merger rate from gravitational-wave detections, to obtain a Swift sGRB detection rate of 11yr^{-1} , the jets are either predicted to have narrow opening angles and high luminosities or wide opening angles and low luminosities. This constraint can be seen in the bottom left corner plot panel of Figure A1 in the violet posterior, where much of the probability density is concentrated in the low luminosity and wide opening angle area of the parameter space. In contrast, the constraints made by GW170817 and GRB 170817A favour a wide opening angle of $\theta_j \lesssim \theta_v$ and a high luminosity event, as seen in the bottom left-hand panel of Figure A1 in the red posterior. This is as emission from an event from a top-hat jet structure when viewed at wide angles can only come from Doppler beaming, which falls off very sharply with increased viewing angles. As $\theta_v > \theta_j$ is more probable than $\theta_v \gtrsim \theta_j$, then a high luminosity event is deemed more probable. The two constraints produce posteriors that share very little overlap in the parameter space, leading to the top-hat model providing a smaller evidence than the other models. This contradiction also manifests in the bimodality of the mean luminosity posterior for the top-hat jet model, as seen in Figure 4, as the lower luminosity high density region corresponds to the constraint produced from the observed number of gamma-ray bursts, while the higher luminosity high

density region corresponds to the constraints made by GW170817 and GRB 170817A.

Jet structure models with wide-angle structuring are only favoured when the data from GW170817/GRB 170817A of \mathcal{D}_{170817} is combined with the event rate information from \mathcal{D}_R . When these data sets are considered individually the evidence for a top-hat jet structure is comparable or higher than the other models in most cases, as seen in the Table A1. The inclusion of the GW190425 event provides evidence against jets with wide-angle structuring. This can be seen by comparing the difference in log evidence between the top-hat jet and the other models given $\mathcal{D}_{170817+R}$, and the difference when given \mathcal{D}_{all} , where there is relatively less evidence between the models when \mathcal{D}_{190425} is included. As it is assumed that GW190425 produced a counterpart that went undetected due to its distance and viewing angle, the event places an upper-bound on the luminosity and jet width. This upper bound on the jet structure limits the possible wide-angle emission, which makes wide-angle jet structuring unnecessary to explain the event.

5.2. Cocoon emission

The double Gaussian jet structure provides a stand-in for a jet structure with cocoon-like emission, where the outer Gaussian provides a secondary component for the emission contribution from an energetic cocoon. Interestingly, comparing the log evidence given the double Gaussian jet model to the other models shows weak evidence for the double Gaussian jet structure when considering \mathcal{D}_{170817} , \mathcal{D}_R , $\mathcal{D}_{170817+R}$ for all cases (with the exception of $\ln p(\mathcal{D}_{170817+R}|\text{GJ})$ given the fitted Schechter luminosity function), suggesting it is the favourable model when considering both GW170817/GRB 170817A and the observed rate data. As discussed in the previous section, GW190425 places an upper-bound on the wide-angled emission and provides support for the top-hat and power-law jet structure with sharper cut-offs. Given that the suitability of GW190425 in the analysis is not as clear-cut as an event like GW170817 due to the uncertainty of the EM coverage of the event, this result should not be disregarded. While this may not provide convincing evidence for the observation of cocoon emission, it suggests that with the inclusion of future events, the necessity for the cocoon-like component can be better assessed.

5.3. Rate of binary neutron star mergers

The narrowest 90% credible intervals of the rate of BNS mergers are constrained within $1-1300 \text{ Gpc}^{-3} \text{ yr}^{-1}$ independent of the jet structure model considered, im-

proving upon the constraints imposed by GWTC-3 (Abbott et al. 2023). The rate is further constrained to the interval of $2-680 \text{ Gpc}^{-3} \text{ yr}^{-1}$ when the fitted Schechter luminosity function is assumed. Future BNS detections will provide tighter constraints on their merger rate. These constraints will allow for a tighter prior to be placed on the rate of mergers, allowing the possible jet structures to be distinguished.

5.4. Luminosity function

Two different cases are explored in the analysis: one where the luminosity function is fitted in advance of the analysis, and the other where priors are placed on the luminosity function parameters L_0^* and γ . When priors are placed on the luminosity function, the luminosity function generally favours low luminosities for all models assumed. This is apparent in Figure 4 where the \hat{L}_0 posterior for all jet structure models shifts to low luminosity when compared to the prior distribution, and mean values of $\log_{10}(L_0^*/\text{erg s}^{-1})$ shift to 51–51.25 in comparison to the value of 51.6 taken from Mogushi et al. (2019) and used as the mean of the prior.

The inclusion of the fitted luminosity function informs the analysis of the prompt emission of all sGRBs that are used in the fit — information that is excluded from the case where the luminosity priors are placed. This allows for narrower constraints on the jet structure model parameters, as seen by comparing the left to right hand-side of the last column of Table 5. Similarly, this also leads to tighter constraints in the BNS merger rate posteriors as seen by comparing Figure 4 to Figure 3. Interestingly, fitting the luminosity function provides slight evidence for the Gaussian jet structure model over all other jet structures given all the data, as seen by the right-hand log Bayes factors shown in Table 4. However, fitting the luminosity function requires assumptions about the jet structure to be made, which will lead to biases in this analysis. In Mogushi et al. (2019) which the fitted luminosity function is taken from, the fit is produced by assuming that all 35 sGRB prompt emission observations with associated redshifts are seen on-axis. However, if some of the events used in fitting the luminosity function were observed at an angle, then the observed variability in their observed isotropic luminosity would be wrongly attributed to variability in the intrinsic luminosity. Assuming a wider distribution to the intrinsic luminosity would favour wider jet structures. To avoid this bias, a future analysis should adjust the likelihood to accommodate the flux data of all observed sGRBs along with their associated redshifts while placing priors on the luminosity function parameters. This would allow for the luminosity function to

be fitted internally within the analysis without having to make the additional jet structure assumptions in a pre-processing step.

5.5. *GW190425*

The inclusion of GW190425 in the analysis provides an upper bound to the wide-angled jet structure emission, due to the absence of an EM detection. The viewing angle posterior of the event exhibits a similar distribution as that of GW170817, while the distance to the event is notably larger at a distance of approximately 160 Mpc compared to GW170817's distance of 40 Mpc. The event is close enough in proximity that, if observed on-axis and is of typical luminosity, would produce a flux tens or hundreds of times greater than GRB 170817A. However, there are assumptions about the event that are made by including it in this way. Firstly, it implies that the event produced a sGRB. This assumption is made explicitly in the analysis when incorporating the observed rate of merger, where every BNS merger is assumed to produce a sGRB in Eqn. 5. However as the prior on the local rate of BNS mergers is relatively wide and covers multiple orders of magnitude, this assumption should not affect the analysis when considering the whole population as long as BNS mergers do typically produce sGRBs. This assumption has a much greater impact when analysing individual events where wrongly asserting a particular event produced a sGRB leads to false conclusions. Secondly, it is assumed that the event would be observed given a wider jet structure or higher luminosity. While the event was within the field of view of the *Konus-Wind* satellite, the incomplete sky coverage of the event by the more sensitive detectors such as the *Fermi*-GBM detector and *Swift*-BAT bring the detectability of the event into question. Despite the validity of these assumptions, and that the event produces evidence against wide-angle jet structuring, it is found that GW190425 is still compatible with the jet structure models given the rest of the data. This can be assessed by the comparison of $\ln p(\mathcal{D}_{\text{all}}|M)$ to $\ln p(\mathcal{D}_{190425}|M) + \ln p(\mathcal{D}_{170817+R}|M)$ for each of the jet structure models M . For all models, the value of $\ln p(\mathcal{D}_{\text{all}}|M) > \ln p(\mathcal{D}_{190425}|M) + \ln p(\mathcal{D}_{170817+R}|M)$, suggesting that the observation of GW190425 is informative to the analysis in all cases, and does not conflict with the constraints imposed to the model given by the detection of GW170817/GRB 170817A and the rate of observed sGRBs. This result suggests that it is feasible for GW190425 to have had a typical sGRB counterpart with the same jet structure as GRB 170817A that would have remained undetectable to our instrumentation even given full sky coverage. This observation is consistent

with the result obtained in Saleem et al. (2020) where it was concluded that such a structured jet is consistent with the observed flux of the *INTEGRAL* detector given the detector's flux upper limit.

6. CONCLUSION

We provide an extensive Bayesian analysis that constrains the jet structure, intrinsic luminosity function and rate of BNS mergers as well as providing a comparison between competing jet structure models. This is achieved by combining four data avenues: 1. the parameter inference posteriors from a GW trigger, 2. the sGRB flux when a counterpart is detected or the detector flux upper limit otherwise, 3. the observation rate of detected sGRBs, 4. the merger rate informed from GW observations. We perform this analysis using the GW triggers GW170817 and GW190425, GRB 170817A, the non-detection of a GW190425 counterpart, the rate of sGRB detections by the Swift detector within a 9.8 year observation period and a merger rate consistent with the constraints imposed by GWTC-2 (Abbott et al. 2020c). This provides us with the following results:

- The rate of BNS mergers is constrained within $1 - 1300 \text{ Gpc}^{-3} \text{ yr}^{-1}$, improving upon the results of GWTC-3.
- Wide-angled jet structures prove more compatible with the given model than top-hat jet in explaining the observed number of sGRBs in the wake of the low observed isotropic luminosity of GRB 170817A.
- Slight evidence is provided for a cocoon-like wide-angled jet structure when considering the observed rate of sGRBs and GRB 170817A. However, the evidence becomes awash across all wide-angled jet structures when GW190425 is included in the analysis.
- While providing evidence against wide-angled structuring, the hypothesis that GW190425 had a typical sGRB counterpart with a GRB 170817A-like jet structure and would remain undetectable to the Fermi detector given full-sky coverage is feasible given the model.

The analysis was extended to consider a fitted intrinsic luminosity function to further incorporate the detected flux and estimated redshifts of past sGRB detections. This provides the results:

- The rate of BNS mergers is further constrained to $2 - 680 \text{ Gpc}^{-3} \text{ yr}^{-1}$.

- Slight evidence for the Gaussian jet structure is provided, unless GW190425 is excluded in which the cocoon-like double Gaussian jet structure is equally favoured.

However, we note that the fitting of the luminosity function requires strong assumptions about the jet structure and therefore introduces a bias towards jet structures with wide central components. Interestingly, this bias does not appear to manifest in the resulting Bayes factors where the top-hat jet loses favour over the wide-angled jet structures. A future analysis will work to incorporate the flux measurements and redshift estimations of detected sGRBs directly, and therefore avoid introducing this bias. Future work would also include

incorporating afterglow data into the analysis for events that coincide with a GW detection (Lin et al. 2021).

7. ACKNOWLEDGEMENTS

We are grateful for computational resources provided by Cardiff University, and funded by an STFC (grant no. ST/I006285/1) supporting UK Involvement in the Operation of Advanced LIGO. The authors thank Shiho Kobayashi for fruitful discussions. F.H. was supported by Science and Technology Research Council (STFC) (grant no. ST/N504075/1). J.V., I.S.H. and M.J.W. are supported by STFC (grant no. ST/V005634/1). M.J.W. was also supported by STFC (grant no. 2285031). G.P.L. is supported by a Royal Society Dorothy Hodgkin Fellowship (grant no. DHF-R1-221175 and DHFERE-221005). E.T.L is supported by National Science and Technology Council (NSTC) of Taiwan (grant no. 111-2112-M-007-020).

REFERENCES

- Abbott, B., Abbott, R., Abbott, T., et al. 2019, *Physical Review X*, 9, 011001
- . 2020a, *The Astrophysical Journal*, 892, L3
- Abbott, B. P., Bloemen, S., Canizares, P., et al. 2017a, *The Astrophysical Journal Letters*, 848, L12, doi: [10.3847/2041-8213/aa91c9](https://doi.org/10.3847/2041-8213/aa91c9)
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017b, *Nature*, 551, 85, doi: [10.1038/nature24471](https://doi.org/10.1038/nature24471)
- Abbott, B. P., Abbott, R., Abbott, T., et al. 2017c, *The Astrophysical Journal Letters*, 848, L13
- . 2017a, *Physical Review Letters*, 119, 161101
- . 2020b, *Living reviews in relativity*, 23, 1
- Abbott, R., Abbott, T., Abraham, S., et al. 2020c, *The Astrophysical Journal Letters*, 913, L7
- Abbott, R., Abbott, T., Acernese, F., et al. 2023, *Physical Review X*, 13, 011048
- Adam, R., Ade, P. A., Aghanim, N., et al. 2016, *Astronomy & Astrophysics*, 594, A1
- Alexander, K., Margutti, R., Blanchard, P., et al. 2018, *The Astrophysical Journal Letters*, 863, L18
- Band, D., Matteson, J., Ford, L., et al. 1993, *The Astrophysical Journal*, 413, 281
- Beniamini, P., Granot, J., & Gill, R. 2020, *Monthly Notices of the Royal Astronomical Society*, 493, 3521
- Beniamini, P., Petropoulou, M., Barniol Duran, R., & Giannios, D. 2019, *MNRAS*, 483, 840, doi: [10.1093/mnras/sty3093](https://doi.org/10.1093/mnras/sty3093)
- Biscoveanu, S., Thrane, E., & Vitale, S. 2020, *The Astrophysical Journal*, 893, 38
- Bloom, J. S., Frail, D. A., & Sari, R. 2001, *The Astronomical Journal*, 121, 2879
- Cole, S., Norberg, P., Baugh, C. M., et al. 2001, *Monthly Notices of the Royal Astronomical Society*, 326, 255
- Duffell, P. C., Quataert, E., Kasen, D., & Klion, H. 2018, *ApJ*, 866, 3, doi: [10.3847/1538-4357/aae084](https://doi.org/10.3847/1538-4357/aae084)
- D’Avanzo, P., Campana, S., Salafia, O. S., et al. 2018, *Astronomy & Astrophysics*, 613, L1
- Evans, P., Cenko, S., Kennea, J., et al. 2017, *Science*, 358, 1565
- Farah, A., Essick, R., Doctor, Z., Fishbach, M., & Holz, D. E. 2020, *The Astrophysical Journal*, 895, 108
- Fraija, N., De Colle, F., Veres, P., et al. 2019, *ApJ*, 871, 123, doi: [10.3847/1538-4357/aaf564](https://doi.org/10.3847/1538-4357/aaf564)
- Goldstein, A., Veres, P., Burns, E., et al. 2017, *The Astrophysical Journal Letters*, 848, L14
- Gottlieb, O., Nakar, E., Piran, T., & Hotokezaka, K. 2018, *MNRAS*, 479, 588, doi: [10.1093/mnras/sty1462](https://doi.org/10.1093/mnras/sty1462)
- Granot, J., Guetta, D., & Gill, R. 2017, *ApJL*, 850, L24, doi: [10.3847/2041-8213/aa991d](https://doi.org/10.3847/2041-8213/aa991d)
- Guetta, D., & Piran, T. 2006, *Astronomy & Astrophysics*, 453, 823
- Hayes, F., Heng, I. S., Veitch, J., & Williams, D. 2020, *The Astrophysical Journal*, 891, 124
- Hogg, D. W. 1999, arXiv preprint astro-ph/9905116
- Hopkins, A. M., & Beacom, J. F. 2006, *The Astrophysical Journal*, 651, 142
- Hosseinzadeh, G., Cowperthwaite, P., Gomez, S., et al. 2019, *The Astrophysical Journal Letters*, 880, L4

- Ioka, K., & Nakamura, T. 2019, *Monthly Notices of the Royal Astronomical Society*, 487, 4884
- Kathirgamaraju, A., Barniol Duran, R., & Giannios, D. 2018, *MNRAS*, 473, L121, doi: [10.1093/mnras/slx175](https://doi.org/10.1093/mnras/slx175)
- Kumar, P., & Granot, J. 2003, *The Astrophysical Journal*, 591, 1075
- Lamb, G. P., & Kobayashi, S. 2017, *Monthly Notices of the Royal Astronomical Society*, 472, 4953
- . 2018, *Monthly Notices of the Royal Astronomical Society*, 478, 733
- Lamb, G. P., Levan, A. J., & Tanvir, N. R. 2020, *ApJ*, 899, 105, doi: [10.3847/1538-4357/aba75a](https://doi.org/10.3847/1538-4357/aba75a)
- Lamb, G. P., Nativi, L., Rosswog, S., et al. 2022, *Universe*, 8, 612, doi: [10.3390/universe8120612](https://doi.org/10.3390/universe8120612)
- Lamb, G. P., Lyman, J. D., Levan, A. J., et al. 2019, *ApJL*, 870, L15, doi: [10.3847/2041-8213/aaf96b](https://doi.org/10.3847/2041-8213/aaf96b)
- Lamb, G. P., Fernández, J. J., Hayes, F., et al. 2021, *Universe*, 7, 329, doi: [10.3390/universe7090329](https://doi.org/10.3390/universe7090329)
- Lazzati, D., López-Cámara, D., Cantiello, M., et al. 2017, *The Astrophysical Journal Letters*, 848, L6
- Lien, A., Sakamoto, T., Gehrels, N., et al. 2014, *The Astrophysical Journal*, 783, 24
- Lien, A., Sakamoto, T., Barthelmy, S. D., et al. 2016, *The Astrophysical Journal*, 829, 7
- Lin, E.-T., Hayes, F., Lamb, G. P., et al. 2021, *Universe*, 7, 349
- Lyman, J., Lamb, G., Levan, A., et al. 2018, *Nature Astronomy*, 2, 751
- Margutti, R., Alexander, K., Xie, X., et al. 2018, *The Astrophysical Journal Letters*, 856, L18
- Matsumoto, T., Nakar, E., & Piran, T. 2019, *MNRAS*, 486, 1563, doi: [10.1093/mnras/stz923](https://doi.org/10.1093/mnras/stz923)
- McCully, C., Hiramatsu, D., Howell, D. A., et al. 2017, *The Astrophysical Journal Letters*, 848, L32
- Mogushi, K., Cavaglia, M., & Siellez, K. 2019, *The Astrophysical Journal*, 880, 55
- Mooley, K., Nakar, E., Hotokezaka, K., et al. 2018, *Nature*, 554, 207
- Nakar, E., & Piran, T. 2021, *The Astrophysical Journal*, 909, 114
- Nathanail, A., Gill, R., Porth, O., Fromm, C. M., & Rezzolla, L. 2021, *MNRAS*, 502, 1843, doi: [10.1093/mnras/stab115](https://doi.org/10.1093/mnras/stab115)
- Petrillo, C. E., Dietz, A., & Cavaglia, M. 2013, *The Astrophysical Journal*, 767, 140
- Poolakkil, S., Preece, R., Fletcher, C., et al. 2021, *The Astrophysical Journal*, 913, 60
- Rossi, E., Lazzati, D., & Rees, M. J. 2002, *Monthly Notices of the Royal Astronomical Society*, 332, 945
- Rossi, E. M., Lazzati, D., Salmonson, J. D., & Ghisellini, G. 2004, *Monthly Notices of the Royal Astronomical Society*, 354, 86
- Ruan, J. J., Nynka, M., Haggard, D., Kalogera, V., & Evans, P. 2018, *The Astrophysical Journal Letters*, 853, L4
- Salafia, O. S., Barbieri, C., Ascenzi, S., & Toffano, M. 2020, *A&A*, 636, A105, doi: [10.1051/0004-6361/201936335](https://doi.org/10.1051/0004-6361/201936335)
- Salafia, O. S., Ghirlanda, G., Ascenzi, S., & Ghisellini, G. 2019, *A&A*, 628, A18, doi: [10.1051/0004-6361/201935831](https://doi.org/10.1051/0004-6361/201935831)
- Saleem, M., Resmi, L., Arun, K., & Mohan, S. 2020, *The Astrophysical Journal*, 891, 130
- Sarin, N., Lasky, P. D., Vivanco, F. H., et al. 2022, *Physical Review D*, 105, 083004
- Savchenko, V., Ferrigno, C., Kuulkers, E., et al. 2017, *The Astrophysical Journal Letters*, 848, L15
- Takahashi, K., & Ioka, K. 2021, *MNRAS*, 501, 5746, doi: [10.1093/mnras/stab032](https://doi.org/10.1093/mnras/stab032)
- Tan, W.-W., & Yu, Y.-W. 2020, *The Astrophysical Journal*, 902, 83
- Tanvir, N. R., Levan, A., González-Fernández, C., et al. 2017, *The Astrophysical Journal Letters*, 848, L27
- Troja, E., Piro, L., Van Eerten, H., et al. 2017, *Nature*, 551, 71
- Troja, E., Piro, L., Ryan, G., et al. 2018, *Monthly Notices of the Royal Astronomical Society: Letters*, 478, L18
- Troja, E., van Eerten, H., Zhang, B., et al. 2020, *Monthly Notices of the Royal Astronomical Society*, 498, 5643
- Wanderman, D., & Piran, T. 2015, *Monthly Notices of the Royal Astronomical Society*, 448, 3026
- Williams, D., Clark, J. A., Williamson, A. R., & Heng, I. S. 2018, *The Astrophysical Journal*, 858, 79
- Williams, M. J., Veitch, J., & Messenger, C. 2021, *Physical Review D*, 103, 103006
- Zhang, B., Dai, X., Lloyd-Ronning, N. M., & Mészáros, P. 2004, *The Astrophysical Journal Letters*, 601, L119
- Zhang, B., & Meszaros, P. 2002, *The Astrophysical Journal*, 571, 876

APPENDIX

A. EVIDENCES

Data \mathcal{D}	Model M			
	Top hat	Gaussian	Power-law	Double Gaussian
\mathcal{D}_{170817}	11.87 ± 0.02	11.8 ± 0.03	11.79 ± 0.03	12.08 ± 0.03
\mathcal{D}_{190425}	17.29 ± 0.01	16.3 ± 0.03	16.41 ± 0.02	16.42 ± 0.03
\mathcal{D}_R	-7.67 ± 0.04	-7.8 ± 0.04	-7.73 ± 0.04	-7.53 ± 0.04
$\mathcal{D}_{170817+R}$	4.45 ± 0.05	5.23 ± 0.05	5.15 ± 0.05	5.37 ± 0.05
\mathcal{D}_{all}	22.12 ± 0.05	22.66 ± 0.05	22.58 ± 0.05	22.67 ± 0.05

Table A1. The log evidence $\ln p(\mathcal{D}|M)$ given the five different data sets when assuming each of the four models.

Data \mathcal{D}	Model M			
	Top hat	Gaussian	Power-law	Double Gaussian
\mathcal{D}_{170817}	11.48 ± 0.02	11.6 ± 0.03	11.45 ± 0.03	11.62 ± 0.03
\mathcal{D}_{190425}	17.28 ± 0.01	16.23 ± 0.03	16.29 ± 0.03	16.21 ± 0.03
\mathcal{D}_R	-8.08 ± 0.05	-8.53 ± 0.05	-8.23 ± 0.05	-7.84 ± 0.05
$\mathcal{D}_{170817+R}$	3.86 ± 0.05	4.97 ± 0.05	4.74 ± 0.05	4.91 ± 0.05
\mathcal{D}_{all}	21.87 ± 0.05	22.78 ± 0.05	22.32 ± 0.05	22.51 ± 0.05

Table A2. The log evidence $\ln p(\mathcal{D}|M)$ given the five different data sets when assuming each of the four models given the fitted Schechter luminosity function.

B. POSTERIOR




Figure A1. Parameter posterior for all data subsets when given the top-hat jet structure model.




Figure A2. Parameter posterior for all data subsets when given the Gaussian jet structure model.




Figure A3. Parameter posterior for all data subsets when given the power-law jet structure model.




Figure A4. Parameter posterior for all data subsets when given the double Gaussian jet structure model.