

An output-polynomial time algorithm to determine all supported efficient solutions for multi-objective integer network flow problems

David Könen¹ and Michael Stiglmayr¹

¹University of Wuppertal, School of Mathematics and Natural Sciences, Optimization Group, Gaußstraße 20, 42103 Wuppertal, Germany,

¹E-Mail: koenen@uni-wuppertal.de, stiglmayr@uni-wuppertal.de

This paper addresses the problem of enumerating all *supported efficient solutions* for a linear multi-objective integer minimum cost flow problem (MOIMCF). First, we highlight an inconsistency in various definitions of supported nondominated vectors for multi-objective integer linear programs (MOILP). Several characterizations for supported nondominated vectors/efficient solutions are used in the literature, which are equivalent in the non-integer case. However, they may lead to different sets of supported nondominated vectors/efficient solutions for MOILPs. This motivates us to summarize equivalent definitions and characterizations for supported efficient solutions and to distinguish between *supported* and *weakly supported* efficient solutions. In this paper we derive an output-polynomial time algorithm to determine all supported efficient solutions for MOIMCF problems. This is the first approach that solves this general problem in output-polynomial time. Moreover, we prove that the existence of an output-polynomial time algorithm to determine all weakly supported nondominated vectors (or all weakly supported efficient solutions) for a MOIMCF problem with a fixed number of $d \geq 3$ objectives can be excluded, unless $\mathbf{P} = \mathbf{NP}$.

Keywords: minimum cost flow, multi-objective integer linear programming, multi-objective network flow, complexity theory, weakly supported, output-polynomial algorithm

1 Introduction

The minimum cost integer network flow problem is a fundamental, well-studied problem in combinatorial optimization [1, 2]. The problem has many real-world applications in industry, and decision-making—among others the important *transportation problem* and *assignment problem* reduce to the minimum cost network flow problem. As an example, Chapter 19 of [1] specifies applications from inventory planning, data scaling, lot sizing, location problems, DNA sequence alignment, and project management.

Given a directed graph with n nodes $V = \{v_1, \dots, v_n\}$ and m arcs $A = \{e_1, \dots, e_m\}$, non-negative arc costs and non-negative integer capacities, and an integer-valued supply/demand b_v for each node $v \in V$, the task is to determine a b -flow, or referred to as flow, with minimum total cost, i.e., a network flow that respects the capacities on all arcs $a \in A$ and has flow balance exactly b_v in each node $v \in V$.

For the single-objective version of this problem, various polynomial algorithms exist. We refer to [1] for a comprehensive overview. However, real-world problems often involve multiple conflicting objectives, such that no solution simultaneously optimizes all objectives. In such cases, one is interested in finding solutions with the property that none of the objectives can be improved without deterioration of at least one other objective. Such a solution is called *efficient solution* and its image is called *nondominated vector*. Finding all or a suitable subset of the efficient solutions/nondominated vectors is the goal of multi-objective optimization (see [27] for a summary of solution concepts in multi-objective optimization). Such a subset could be the set of all *supported efficient solutions*, which are those efficient solutions that can be obtained as optimal solutions of a weighted sum scalarization with weights strictly greater zero. *Extreme supported nondominated solutions* can be characterized as extreme points of the *upper image* of feasible outcome vectors, which is defined as the polyhedron given by the convex hull of feasible outcome vectors plus the (closed) positive orthant.

Several characterizations exist for supported nondominated vectors/efficient solutions, which are all equivalent in the non-integer case [7]. However, they may lead to different sets of supported nondominated vectors/efficient solutions and have different properties in the integer case. While convex optimization problems only contain supported efficient solutions, integer problems can contain *unsupported* efficient solutions, namely efficient solutions that are not supported. In the literature, supported nondominated vectors are often characterized as nondominated vectors on the boundary of the upper image and that they only lie on the *nondominated frontier*, which is defined as the union of all *maximally nondominated faces*, whereas unsupported solutions are characterized as nondominated vectors that lie in the interior of the upper image [10, 20]. Hereby, a face of the upper image is called a *nondominated face* if all its points are nondominated. A face F is called *maximally nondominated* if there is no other nondominated face G such that $F \subset G$, this would imply that the dimension of G is greater than the dimension of F . However, there may exist unsupported nondominated vectors that lie on the boundary of the upper image or supported nondominated vectors that lie on non-maximally nondominated faces, i.e., *weakly nondominated faces* and thus do not lie on the nondominated frontier. This motivates us to distinguish between *supported*

efficient solutions and *weakly supported* efficient solutions. A efficient solution is denoted as weakly supported if it is an optimal solution of a weighted sum scalarization with non-negative weights. In contrast to the definition of supportedness, weakly supportedness allows single weights to have a value of zero. In order to avoid confusion, note that weakly supported efficient solutions are efficient. However, their images may lie on *weakly nondominated faces*, which could be dominated in the non-integer case. An unsupported efficient solutions is then defined as an efficient solution which is neither supported nor weakly supported. Then the following characterizations hold: While the weakly supported nondominated vectors lie on the boundary of the upper image, the supported nondominated vectors lie only on the nondominated frontier, i. e., only on maximally nondominated faces. The unsupported nondominated vectors lie in the interior of the of the upper image. We will lay out an example where the set of supported efficient solutions is a proper subset of all weakly efficient solutions in Section 3.

Multi-objective integer minimum cost flow (MOIMCF) problems are much harder to solve as compared to the single-objective case. MOIMCF had been reviewed in [15], where the authors comment on the lack of efficient algorithms. However, since then there still is a lack of efficient algorithms for determining supported efficient solutions for MOIMCF problems and we give an overview of new results below.

While its linear relaxation, the *multi-objective minimum cost flow problem* (MOMCF), has only *supported efficient solutions*, the *multi-objective integer minimum cost flow* (MOIMCF) problem, on the other hand, can have supported, weakly supported and unsupported efficient solutions.

There are several algorithms to determine efficient solutions for bi-objective integer minimum cost flow (BOIMCF) problems, e.g., [9, 11, 21, 24, 25, 26]. Raith and Sedeño-Noda introduced an enhanced parametric approach to determine all extreme efficient solutions for BOIMCF problems [22]. However, there are only quite a few specific methods designed to determine all (or subsets) of the nondominated vectors in the objective space (nor the corresponding efficient solutions in the decision space) for MOIMCF problems [10, 13, 29].

Eusébio and Figueria [10] introduced an algorithm that enumerates all supported nondominated vectors/efficient solutions for MOIMCF problems (assuming extreme supported solutions and corresponding weight vectors are given), based on zero-cost cycles in the incremental graph associated with the corresponding parametric network flow problems. They conclude, that their proposed algorithm is the first step in developing further zero-cost cycle algorithms for solving MOIMCF problems. However, they do not provide a specific method for determining those zero-cost cycles. As mentioned before, different definitions exist for supported efficient solutions, which lead to different sets of supported efficient solutions, namely supported- and weaklysupported efficient solutions. Using the definition in [10] would also include the weakly supported efficient solutions. However, it can be shown that their proposed algorithm can only determine all supported efficient solutions. We refer to the example in Section 2.

This paper proves that there is no *output-polynomial* time algorithm to determine all weaklysupported nondominated vectors (or weaklysupported efficient solutions) for a MOIMCF problem with a fixed number of objectives. In contrast, we derive *output-*

polynomial enumeration algorithms to determine all supported efficient solutions, first for the bi-objective case and afterwards for every fixed number of objectives. Here, output-polynomial time refers to a computation time that can be bounded by a polynomial in the input and output size. We refer to the paper of Johnson, Papadimitriou and Yannakakis [16] for a detailed survey of output-sensitive complexity.

The approach consists of two phases: First, the algorithm determines all extreme points of the upper image and the associated weight vector for each maximally non-dominated face. Hence, the approach successively determines all efficient solutions for each maximally nondominated face by determining all optimal solutions for the linear weighted-sum scalarization (single-objective parametric network flow) problem with corresponding weight vector using the previously presented algorithm to determine all optimum integer flows in a network [18]. The method successively searches for *proper zero-cost cycles* in linear time by using a modified depth-first search technique.

Given a BOIMCF problem and using the enhanced parametric network approach [22] to determine all N extreme nondominated points in $\mathcal{O}(N \cdot n(m + n \log n))$ time in a first step, results in an $\mathcal{O}(N \cdot n(m + n \log n) + \mathcal{S}(m + n))$ time algorithm to determine all \mathcal{S} supported efficient solutions to a BOIMCF problem.

Given a MOIMCF problem with a fixed number of d objectives, the dual Benson's algorithm [8] can be used for the first phase. In addition, while the lexicographic MCFP can be solved in polynomial time [15], we can use the lexicographic dual Benson's algorithm presented in [4], which determines all extreme nondominated vectors, all facets of the *upper image* and the *weight space decomposition* in $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor}(\text{poly}(n, m) + N \log N))$. Note that in phase two, the corresponding image of some solutions may lie in more than one maximally nondominated face and therefore are included in more than one face. The approach yields an $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor}(\text{poly}(n, m) + N \log N + N^{\lfloor \frac{d}{2} \rfloor} \mathcal{S}(m + n + N^{\lfloor \frac{d}{2} \rfloor} d)))$ time algorithm to determine all \mathcal{S} supported efficient solutions for a d -objective MOIMCF problem. To our best knowledge, this is the first output-polynomial time algorithm to determine the complete set of supported efficient solutions.

The following table summarizes the contribution of our article and the existing results from literature on the existence of output-polynomial time algorithms for the MOIMCF problem w.r.t. different solution concepts. In the table, a check-mark indicates existence, a cross indicates that the existence of such an algorithm can be ruled out, and the question mark indicates that this problem remains an open question.

	extreme supported	supported	weakly supported	all
nondominated vectors	✓ [4, 8]	?	✗	✗ [3]
efficient solutions	✓	✓	✗	✗

Note that an output-polynomial time algorithm for the determination of all supported efficient solutions is not sufficient for the computation of all nondominated supported vectors in output-polynomial time since there may be exponentially many solutions

mapping to the same vectors. It is easy to check, that all extreme supported efficient solutions can be determined in output-polynomial time, due to the existence of the output-polynomial time algorithm for the extreme supported nondominated vectors. Note, that the algorithm of Eusébio and Figueria [10] would also determine all supported efficient solutions in output-polynomial time, when all maximally nondominated faces and corresponding weight vectors are already given.

The remainder of the paper is structured as follows. Section 2 presents the notations and preliminaries. Section 3 introduces and compares the definitions of supported and weaklysupported nondominated vectors, and Section 4 presents the algorithms to determine all supported efficient solutions for BOIMCF problems as well as all supported efficient solutions for MOIMCF problems. Section 5 summarizes the article and gives an outlook on future research.

2 Multi-objective integer network flows

This section formally introduces the multi-objective integer minimum cost flow problem along with some of the most important results and properties of network flows. For a comprehensive introduction to the graph theoretic foundations and the basics of network flow theory we refer to [1, 5].

Let $D = (V, A)$ be a directed graph with $|V| = n$ nodes and $|A| = m$ arcs together with integer-valued, non-negative, finite lower and upper capacity bounds l_{ij} and u_{ij} , respectively, for each arc $(i, j) \in A$. Moreover, let b_i be the integer-valued *flow balance* of the node $i \in V$ and let $c: \mathbb{R}^m \rightarrow \mathbb{R}_{\geq}^d$ be the (vector valued) objective function and $C \in \mathbb{R}^{d \times m}$ the corresponding cost matrix, i. e., $c(f) = C \cdot f$. Furthermore, let $\mathbb{R}_{\geq}^d := \{x \in \mathbb{R}^d: x \geq 0\}$ denote the non-negative orthant of \mathbb{R}^d . Its interior $\mathbb{R}_{>}^d$ is defined accordingly. In our notation, we also use the Minkowski sum and the Minkowski product of two sets $A, B \subseteq \mathbb{R}^d$, which are defined as $A + B := \{a + b: a \in A, b \in B\}$ and $A \cdot B := \{a \cdot b: a \in A, b \in B\}$, respectively. Then the *multi-objective integer minimum cost flow* (MOIMCF) problem can be formulated as:

$$\begin{aligned} \min \quad & c(f) \\ \text{s.t.} \quad & \sum_{j:(i,j) \in A} f_{ij} - \sum_{j:(j,i) \in A} f_{ji} = b_i \quad i \in V, \quad (\text{flow balance constraint}) \\ & l_{ij} \leq f_{ij} \leq u_{ij} \quad (i, j) \in A, \quad (\text{capacity constraint}) \\ & f_{ij} \in \mathbb{Z}_{\geq} \quad (i, j) \in A. \end{aligned}$$

Where we denote a feasible solution f as *flow* and define the *cost* of a flow f as

$$c(f) := (c^1(f), \dots, c^d(f))^{\top} := \left(\sum_{(i,j) \in A} c_{ij}^1 f_{ij}, \dots, \sum_{(i,j) \in A} c_{ij}^d f_{ij} \right)^{\top} = C \cdot f.$$

The (*continuous*) *multi-objective minimum cost flow* (MOMCF) problem can be formulated as the LP-relaxation of MOIMCF. From now on, we assume that D is connected (see [1]) and that there is at least one feasible flow.




Figure 1: Illustration of the upper image $\mathcal{P} = \text{conv}(\mathcal{Y}) + \mathbb{R}_{\geq}^2$ and the different solution types.

Then, we denote the set of feasible outcome vectors in objective space by $\mathcal{Y} := \{Cf : f \in \mathcal{X}\}$, where \mathcal{X} is the set of all feasible flows. We assume that the objective functions are conflicting, which implies that we exclude the existence of an *ideal* solution that minimizes all objectives simultaneously. Throughout this article, we will use the *Pareto concept of optimality*, which is based on the componentwise order:

$$c(f) \leq c(f') \iff c^k(f) \leq c^k(f') \quad \forall k \in \{1, \dots, d\} \quad \text{and} \quad c(f) \neq c(f')$$

A vector $c(f) \in \mathbb{R}^d$ is called *dominated* by $c(f') \in \mathbb{R}^d$ if $c(f') \leq c(f)$. Accordingly, a feasible flow f is called an *efficient flow* if there exists no other feasible flow f' such that $c(f') \leq c(f)$. The image $c(f)$ of an efficient flow f is called a *nondominated vector*. The set of efficient flows is denoted by $\mathcal{X}_E \subset \mathcal{X}$ and the set of nondominated vectors by $\mathcal{Y}_N \subset \mathcal{Y}$. Moreover, a feasible flow f is called weakly efficient if there is no other flow f' such that $c(f') < c(f)$ where “<” denotes the strict componentwise order:

$$c(f) < c(f') \iff c^k(f) < c^k(f') \quad \forall k \in \{1, \dots, d\}$$

The polyhedron $\mathcal{P} = \text{conv}(\mathcal{Y}) + \mathbb{R}_{\geq}^d$ is called the *upper image* of \mathcal{Y} . Then, we denote an efficient flow *supported efficient*, if it is on the *nondominated frontier* defined as the set $\{y \in \text{conv}(\mathcal{Y}_N) : \text{conv}(\mathcal{Y}_N) \cap (y - \mathbb{R}_{\geq}^d) = \{y\}\}$. All efficient flows that lie in the interior of the upper image are called *unsupported*. A feasible flow is called *extreme (supported efficient) flow* if its image is an extreme point of the upper image \mathcal{P} . Such an image point in the outcome space is called an *extreme (supported nondominated) vector*. Note that the MOIMCF problem has in general also unsupported efficient solutions. Figure 1 illustrates supported extreme, supported and unsupported nondominated points as well as the upper image in the bi-objective case.

Due to the total unimodularity of MOMCF, there exists for each extreme supported nondominated point of MOMCF an integer flow mapping to it, if u and b are integral

(see [1]). In other words, the sets of extreme supported nondominated points of MOMCF and MOIMCF, and thus the respective upper images, coincide. In the remainder of this paper, only integer flows are considered, and from now on, flow always refers to an integer flow.

Let $\|x\|_1$ denote the 1-norm of $x \in \mathbb{R}^d$, i. e., $\|x\|_1 := \sum_{i=1}^d |x_i|$. For the weighted sum method, we define the set of *normalized weighting vectors* as the set $\Lambda_d = \{\lambda \in \mathbb{R}_{>}^d : \|\lambda\|_1 = 1\}$ or $\Lambda_d^0 = \{\lambda \in \mathbb{R}_{\geq}^d : \|\lambda\|_1 = 1\}$ if weights equal to zero are included. Then, the *weighted-sum linear program* w. r. t. $\lambda \in \Lambda_d$ or $\lambda \in \Lambda_d^0$ is defined as the parametric program

$$\begin{aligned} \min \quad & \lambda^\top C f \\ \text{s.t.} \quad & f \in \mathcal{X} \end{aligned} \tag{P_\lambda}$$

Note, that the problem (P_λ) is as easy to solve as the associated single objective problem, as long as the encoding lengths of the components of λ are not too large. (Consider that λ is not part of the original input.) However, on a positive note, it is shown that these lengths can be bounded by $\mathcal{O}(\text{poly}(n, m))$ [4]. If $\lambda \in \Lambda_d^0$ every optimal solution to (P_λ) is a weakly efficient solution. Moreover, every optimal solution of (P_λ) is efficient, if $\lambda \in \Lambda_d$ [7].

Since the capacities on each arc are bounded, \mathcal{X} is a polytope, and therefore, \mathcal{Y} is also a polytope. We define the concepts of face, facet, maximal, and maximally nondominated faces similarly as in ([10], [28]). We call $F \subset \mathcal{Y}$ a *face* of \mathcal{Y} if there exists a supporting hyperplane H such that $H \cap \mathcal{Y} = F$. A face F of \mathcal{Y} is denoted as *r-face* if F is of dimension r . Extreme points are 0-faces and 1-faces are edges. A *r-face* $F \neq \mathcal{Y}$ is denoted as *maximal face* or *facet* of \mathcal{Y} if there does not exist another *s-face* G , such that $F \subset G$ and $r < s$. Note, that the dimension of a facet F is $\dim(F) = \dim(\mathcal{Y}) - 1$. A face is called *nondominated* if there exists a corresponding weight vector to the face with weights strictly greater than zero. A *r-face* F , is a *maximally nondominated face* if there is no nondominated *s-face* G , such that $F \subset G$ and $r < s$. Let $F_{\mathcal{X}}$ be the set of the preimage of a maximally nondominated face $F_{\mathcal{Y}}$ of polytope \mathcal{Y} , i. e., all solutions whose image lies in $F_{\mathcal{Y}}$. We use the expression *maximally efficient face* for $F_{\mathcal{X}}$. Even if we use the expression *face* for $F_{\mathcal{X}}$, it must not really be a face of the decision space, since more than one feasible solution could have a vertex of \mathcal{Y} as image and they may lie in different faces of \mathcal{X} .

3 Supported and weaklysupported nondominated vectors in MOILP

There exist several definitions and characterizations of supported nondominated vectors in the literature: (1) a nondominated vector is called supported if it lies on the boundary of the convex hull of the feasible region $\text{conv}(\mathcal{Y})$ in the outcome space [10, 23], (2) a nondominated vector is called supported if it lies on the boundary of $\mathcal{Y}^{\geq} := \text{conv}(\mathcal{Y}_N + \mathbb{R}_{\geq}^d)$ [9, 10], i. e., the boundary of the upper image, (3) a nondominated vector is called supported if it is on the *nondominated frontier* defined as the set $\{y \in \text{conv}(\mathcal{Y}_N) : \text{conv}(\mathcal{Y}_N) \cap (y - \mathbb{R}_{\geq}^d) = \{y\}\}$ [15], and (4) supported nondominated

vector is the image of a supported efficient solution which are those efficient solutions that can be obtained as optimal solutions of a weighted sum scalarization with weights strictly greater zero [7, 20, 21, 22].

While all definitions are equivalent in the non-integer case [7], this does not hold for the integer case in general. In the literature, supported nondominated vectors are often characterized as nondominated vectors on the boundary of the upper image and that they only lie on the maximally nondominated faces, i.e, the nondominated frontier, whereas the unsupported are characterized as nondominated vectors that lie in the interior of the upper image, e.g., [10, 20]. However, if a supported vector is defined in the way of (3) and (4), there may exist unsupported vectors on the boundary of the upper image. However, if a supported nondominated vector is defined as in (1) and (2), there may exist supported nondominated vectors on the boundary of the upper image which are not lying on the nondominated frontier, i.e., which lie on weakly nondominated faces. This motivates us to develop new consistent definitions of supported and weakly supported nondominated vectors.

Definition 3.1 *An efficient solution is called a weakly supported efficient solution if it is an optimal solution of a weighted-sum scalarization P_λ for some weight $\lambda \in \Lambda_d^0$. Moreover, if the weight is strictly positive $\lambda \in \Lambda_d$ it is called supported efficient solution. The corresponding image is called weakly supported or supported (nondominated) vector, respectively.*

The weaklysupported nondominated vectors lie on the boundary of the upper image $\partial\mathcal{P}$ while the supported lies only on the nondominated frontier, i.e., on maximally nondominated faces. Definition (1) and (2) will include the weakly supported nondominated vectors while definition (3) and (4) only contains the supported nondominated vectors. Let \mathcal{Y}_S and \mathcal{Y}_{wS} be the set of all supported nondominated vectors and the set of all weakly supported nondominated vectors, respectively. According, \mathcal{X}_S (\mathcal{X}_{wS}) is the set of all (weakly) supported efficient solutions. We denote the cardinality of \mathcal{X}_S by $|\mathcal{X}_S| = \mathcal{S}$.

Theorem 3.2 *Let \mathcal{Y}_S and \mathcal{Y}_{wS} be the sets of a multi-objective integer linear program. Then it holds that $\mathcal{Y}_S \subseteq \mathcal{Y}_{wS}$ and there exist instances where $\mathcal{Y}_S \subset \mathcal{Y}_{wS}$, i.e., the set of supported nondominated vectors is a proper subset of the set of the supported nondominated vectors.*

Proof. The inclusion $\mathcal{Y}_S \subseteq \mathcal{Y}_{wS}$ holds per definition. The MOIMCF problem shown in Figure 2 has three extreme nondominated vectors $y^1 = (8, 16, 6)^\top$, $y^2 = (12, 12, 6)^\top$, $y^3 = (16, 8, 10)^\top$ and in addition the following nondominated vectors $s^1 = (9, 15, 7)^\top$, $s^2 = (10, 14, 8)^\top$, $s^3 = (11, 13, 9)^\top$, $s^4 = (13, 11, 7)^\top$, $s^5 = (14, 10, 8)^\top$, $s^6 = (15, 9, 9)^\top$ and as well one dominated vector $d^1 = (12, 12, 10)^\top$. The different vectors in objective space and their convex hull are illustrated in Figure 3. All vectors s^i with $i \in \{1, \dots, 6\}$ are weakly supported nondominated vectors, since their preimages are optimal solutions of the corresponding weighted sum problem P_λ with $\lambda = (0.5, 0.5, 0)^\top$. The vectors s^4, s^5, s^6 are also supported nondominated vectors since they are optimal solutions of




Figure 2: The Graph corresponds to a tri-objective MOIMCF problem. Thereby, $l_{ij} = 0$ and $u_{ij} = 4$ for all arcs $(i, j) \in A$. The arcs are labeled with their cost coefficients $(c_{ij}^1, c_{ij}^2, c_{ij}^3)$ and the nodes are labeled with their supply/demand b_i .

(P_{λ^2}) with $\lambda^2 = (0.25, 0.5, 0.25)^\top$. However s^1, s^2, s^3 are not optimal for any weighted sum problem P_λ with $\lambda \in \Lambda_d$. Hence, the set of weakly supported nondominated vectors is $\mathcal{Y}_w S = \{y^1, y^2, y^3, s^1, s^2, s^3, s^4, s^5, s^6\}$. While the set of supported nondominated vectors is $\mathcal{Y}_S = \{y^1, y^2, y^3, s^4, s^5, s^6\}$. Thus, in this example the set of supported nondominated vectors is a proper subset of the set of weakly supported nondominated vectors $\mathcal{Y}_S \subset \mathcal{Y}_w S$. \square

It is easy to see that there do not exist weights $\lambda \in \Lambda_d$ such that s_1, s_2 , or s_3 are optimal solutions for P_λ using the *weight space decomposition* which will be formally introduced in Section 4.2. Figure 6 shows the weight space decomposition of the *upper image* of the MOIMCF given in Figure 2.

Note that in the biobjective case $\mathcal{Y}_S = \mathcal{Y}_w S$ as the following lemma shows.

Lemma 3.3 *Every weakly supported nondominated point of a biobjective integer optimization problem is supported nondominated.*

Proof. Let $y = f(x) \in \mathbb{R}^2$ be a weakly supported but not supported nondominated point of a biobjective integer optimization problem and x the corresponding preimage. Then there is a weighting vector $\lambda \in \Lambda_0$ such that x is an optimal solution of the weighted sum problem P_λ . Since y is not supported nondominated, one of the components of λ must be zero. W.l.o.g. let $\lambda = (1, 0)^\top$.

Since y is nondominated there does not exist a feasible outcome vector \bar{y} , with $\bar{y}_1 = y_1$ and $\bar{y}_2 < y_2$. Thus, x is also an optimal solution of the weighted sum problem $P_{\lambda'}$ with $\lambda' = (1 - \varepsilon, \varepsilon)^\top$ for $\varepsilon > 0$ sufficiently small, which makes y supported nondominated. \square

The clear distinction between the sets of supported nondominated and weakly supported nondominated solutions (Definition 3.1) is also necessary as the corresponding problems differ in their output time complexity. Eusébio and Figueria [10] rely on the definition (2) which is equivalent to what we define as weakly supported nondominated vectors. However, their proposed algorithm, computes only for supported efficient solutions which images lie in the maximally nondominated faces, i.e., supported efficient solutions. In this paper, an output-polynomial algorithm is developed to determine all supported efficient solutions to the MOIMCF problem with a fixed number of objectives.




Figure 3: The figure shows the convex hull $\text{conv}(\mathcal{Y})$ of the given problem in Figure 2 in blue, all its integer vectors and the hyperplane $h = \{y \in \mathcal{Y} : 0.5y_1 + 0.5y_2 = 12\}$ in gray. The red dots on the edge between (y^2, y^3) are the vectors s^4, s^5 , and s^6 . The light red dots on the edge between the edge (y^1, d^1) are the vectors s^1, s^2 , and s^3 , which would be dominated in the non-integer case by the points on the edge (y^1, y^2) . The maximally nondominated faces are the edges (y^1, y^2) and (y^2, y^3) . Note that s^1, s^2 , and s^3 would lie on $\text{conv}(\mathcal{Y})$ but not in any of the maximally nondominated faces. The vectors s^1, s^2 , and s^3 would also lie on the boundary of the upper image.

However, unless $\mathbf{P} = \mathbf{NP}$, this work proves that an output-polynomial algorithm to determine all weakly supported nondominated vectors (or all weakly supported efficient solutions) for a MOIMCF problem with a fixed number of more than two objectives can be excluded. This is proven by showing that there cannot exist an output-polynomial algorithm for the multi-objective s - t -path problem (MOSP) with more than two objectives.

Lemma 3.4 ([3]) *There is no output-polynomial algorithm for determining all nondominated vectors for the MO s - t -path problem, unless $\mathbf{P} = \mathbf{NP}$.*

The proof of Lemma 3.4 is given by showing that the finished decision variant $\text{MOSP}_y^{\text{Fin}}$ of the MOSP is **co-NP**-hard, by a reduction of the complement of the Knapsack problem:

$$\{(c^1, c^2, k_1, k_2) : c^{1\top} x \leq k_1, c^{2\top} x \geq k_2, x \in \{0, 1\}^n\}. \quad (\text{KP})$$

This problem is NP-complete [17]. A finished decision problem E^{Fin} for an enumeration problem $E = (I, C)$ is hereby defined as the problem: Given an instance $x \in I$ of the enumeration problem and a subset $M \subseteq C(x)$ of the configuration set, the goal is to decide if $M = C(x)$, i.e., we want to decide if we already have found all configurations. If the enumeration problem E can be solved in output-polynomial time then $E^{\text{Fin}} \in \mathbf{P}$ [19].

The determination of all nondominated vectors of a MOSP as well as the determination of all efficient solutions of a MOSP can be formulated as an enumeration problem [3]. We denote the finished decision problem for the determination of all nondominated vectors of a MOSP as $\text{MOSP}_y^{\text{Fin}}$ and the determination of all efficient solutions of a MOSP as $\text{MOSP}_x^{\text{Fin}}$, respectively.

The same reduction that is used in the proof of Lemma 3.4 given in [3] can be extended to show that the $\text{MOSP}_x^{\text{Fin}}$ is also **co-NP**-hard and thus there cannot exist an output-polynomial time algorithm to determine all efficient solutions for the MOSP. However, we have to adjust the costs of some weights. In Figure 4 an example of the reduction is given, which is similar to the one used in the proof in [3] in order to show the following Lemma. For the sake of simplicity, we do not give a formal proof here. For more details we refer to [3].

Lemma 3.5 *There is no output-polynomial time algorithm to determine all efficient solutions for the MO s - t -path problem, unless $\mathbf{P} = \mathbf{NP}$.*

Theorem 3.6 *Unless $\mathbf{P} = \mathbf{NP}$, there is no output-polynomial algorithm to determine all weakly supported nondominated vectors for the MO shortest s - t -path problem with a fixed number of $d \geq 3$ objectives.*

Proof. Assume an output-polynomial time algorithm exists to determine all weakly supported nondominated vectors for the MOSP. The existence of an output-polynomial time algorithm for determining all nondominated vectors in a MOSP with d objectives can be excluded, even in the case of two objectives [3]. Let M_d be a MOSP with d objectives.




Figure 4: Showing the reduction in the proof of Lemma 3.4 presented by Bökler et al. [3] with modified costs in order to proof Lemma 3.5. Here $\epsilon = 1/(n + 1)$ and arcs with no label have cost $\mathbf{0}$. Determining another efficient s - t path to the two with cost equal to $(k_1 + 1, 0)$ and $(0, c^{2^\top} \mathbf{1} - k_2 + 1)$ would be an instance of the (KP) problem.

However, if we add an artificial objective $c_{ij}^{d+1} = 0$ for all $(i, j) \in A$ to our MOSP and denote it by M_{d+1} , we obtain a weakly efficient facet for M_{d+1} , where all nondominated vectors for M_d are weakly supported nondominated vectors for M_{d+1} . Therefore, even the unsupported nondominated vectors for M_d are weakly supported nondominated vectors for M_{d+1} , since they are part of the boundary of the upper image for M_{d+1} . So, if an output-polynomial time algorithm exist to determine all weakly supported nondominated vectors for a MOSP there also exists an output-polynomial time algorithm to determine all nondominated vectors, which raises a contradiction. \square

Replacing nondominated vectors by efficient solutions and using Lemma 3.5 instead of Lemma 3.4 proves the following.

Theorem 3.7 *Unless $\mathbf{P} = \mathbf{NP}$, there is no output-polynomial algorithm to determine all weakly supported efficient solutions for the MO s - t -path problem with a fixed number of $d \geq 3$ objectives.*

Note, that we can easily transfer the proofs of Lemma 3.4 and Lemma 3.5 to MOIMCF problems, by transferring the shortest path problem into an flow problem. Note that the multiobjective shortest path problem is only a special case of MOIMCF when all edge costs/lengths are strictly positive or (as in Figure 4) the considered graph is acyclic, as optimal solutions of MOIMCF may in general contain dicycle flows. Showing that there exists another nondominated vector or efficient flow would solve the complement of the knapsack problem.

Theorem 3.8 *Unless $\mathbf{P} = \mathbf{NP}$, there is no output-polynomial algorithm to determine all weakly supported nondominated vectors (or supported efficient solutions) for the*

MOIMCF problem with a fixed number of $d \geq 3$ objectives.

This interrelation between the computation of all non-dominated vectors/all efficient solutions for an problem with d objectives and the determination of all supported non-dominated vectors/efficient solutions of an associated problem with $d+1$ objectives holds also for general MOILPs and can be shown by an analogous construction as in the proof of Theorem 3.6.

Theorem 3.9 *The determination of all weaklysupported nondominated vectors (all supported efficient solutions) of an MOILP with $d \geq 3$ objectives is as hard as the determination of all nondominated vectors (efficient solutions) with $d - 1$ objectives.*

4 Finding all supported efficient flows

In this section, output-polynomial time algorithms are derived that determine all supported efficient flows for MOIMCF. It is based on the determination of all optimal flows for a sequence of single-objective parametric network flow problems, each corresponding to a maximally nondominated face. The approach consists of two phases and relies on the following widely-known fact for any integer supported flow, see e.g., [14].

Theorem 4.1 *A flow f is contained in $F_{\mathcal{X}}$, i. e., its image of $c(f)$ lies on a maximally nondominated face $F_{\mathcal{Y}}$ w. r. t. an associated weight vector $\lambda \in \Lambda_d$, if f is an optimal solution to the parametric network flow program (P_{λ}) .*

Any image of a supported flow must lie in at least one maximally nondominated face, and any integer point in a maximally nondominated face corresponds to a supported integer flow. Assuming that for each maximally nondominated face $F_i \in \{F_1, \dots, F_t\}$ one optimal solution f^i and the corresponding weighting vectors λ^i are given the problem of determining all supported flows reduces to determining all optimal flows for each parametric single-objective problem (P_{λ^i}) . These optimal solutions can be determined by using the algorithm for determining all optimum flows for single-objective minimum cost flow problems presented in [18], which we refer to as the all optimum flow (AOF) algorithm. The AOF algorithm successively searches for so-called *proper zero-cost cycles* efficiently by using a modified depth-first search technique.

Theorem 4.2 ([18]) *Given an initial optimal integer flow f , we can determine all optimal integer flows in $\mathcal{O}(\mathcal{F}(m+n) + mn)$ time for a single-objective minimum cost flow problem, where \mathcal{F} is the number of all optimal integer flows.*

We, therefore, divide the approach into two phases: In phase one, we determine all extreme nondominated vectors and one weighting vector for each maximally nondominated face. In phase two, we apply the AOF algorithm to the corresponding weighted-sum program for each maximally nondominated face.

The extreme nondominated vectors and the weighting vectors for each maximally nondominated face can be determined much easier in the case of two objectives. Thus,

we start by deriving an algorithm for BOIMCF problems and consider afterwards the general case of MOIMCF problems.

4.1 Bi-objective minimum cost flow problem

Note, that in the biobjective case, the set of supported flows is equal to the set of weakly supported flows, since every weakly nondominated face contains exactly one nondominated vector (namely an extreme point), which dominates the complete face.

First, we determine all N extreme points and precisely one corresponding extreme flow by using the enhanced parametric programming approach in $\mathcal{O}(M + Nn(m + n \log n))$ time [22], where M denotes the time required to solve a given single-objective minimum cost flow problem. Also, the algorithm stores one extreme flow for each extreme nondominated point.

Note, that in the biobjective case every maximally nondominated face $F_{\mathcal{Y}}$ of $\text{conv}(\mathcal{Y})$ is a line segment connecting two adjacent extreme supported points if there is more than one nondominated point ($|\mathcal{N}| > 1$). A maximally nondominated face can only have dimension zero if there is only one extreme nondominated point, which implies that there is only one nondominated point (or in other words the ideal point is feasible).

In the following, we will derive a procedure to determine the complete set of all supported efficient flows. For that, we will determine all supported flows which images lie on the maximally nondominated edges. Let y^1, \dots, y^N be the extreme supported points obtained by the enhanced parametric programming approach [22] and let f^1, \dots, f^N be a set of corresponding extreme supported flows each mapping to one extreme supported point. Moreover, we sort the set of extreme supported points and flows $\{y^i = (c^1(f^i), c^2(f^i)), f^i : i \in \{1, \dots, N\}\}$ by non-decreasing values of c^1 . For each pair of consecutive extreme points y^i and y^{i+1} , we determine the weighting vector $\lambda^i \in \Lambda$ that corresponds to the normal of maximally nondominated facet F_i connecting the extreme points y^i and y^{i+1} :

$$\lambda^i := \frac{1}{\sqrt{\lambda_1^2 + \lambda_2^2}} (\lambda_1, \lambda_2)^\top \quad \text{where} \quad \lambda_1^i := c^2(f^i) - c^2(f^{i+1})$$

$$\lambda_2^i := c^1(f^{i+1}) - c^1(f^i).$$

Then f^i and f^{i+1} are both optimal flows for the single-objective weighted-sum (MCF) program (P_{λ^i}) [9]. Hence, determining all optimal solutions for (P_{λ^i}) gives all supported efficient flows whose image lie in between F_i . Figure 5 illustrates the objective function of the weighted-sum problem (P_{λ^i}) and the maximally nondominated face between two consecutive extreme points in the outcome space.

Theorem 4.3 *Given the directed network $(D, l, u, b, (c^1, c^2)^\top)$, Algorithm 1 determines the set of all supported flows \mathcal{X}_S in $\mathcal{O}(Nn(m + n \log n) + \mathcal{S}(m + n))$ time.*

Proof. Any supported point must lie at least on one maximally nondominated edge and only extreme supported points y^i for $i = 2, \dots, N - 1$ lie in two maximally nondominated edges, namely F_i and F_{i+1} . According to Theorem 4.1, all supported flows can be




Figure 5: Illustration of two neighboring extreme points y^i and y^{i+1} , the maximally nondominated edge F_i in blue and the associated weight vector λ^i .

Algorithm 1: FindAllSupportedEfficientFlowsBiObjective

Data: (D, l, u, b, c^1, c^2)

Result: The set of all supported flows \mathcal{X}_S

$\mathcal{X}_S \leftarrow \emptyset;$

// Determine all extreme supported points y^i and for each one
corresponding extreme flow f^i , sorted non-decreasingly in $c^1(f)$

$\{(y^i, f^i) : i \in \{1, \dots, N\}\} \leftarrow \text{EnhancedParametricNetworkAlgorithm}(D);$

for $i = 1, \dots, N - 1$ **do**

$\lambda_1^i \leftarrow c^2(f^i) - c^2(f^{i+1}); \quad \lambda_2^i \leftarrow c^1(f^{i+1}) - c^1(f^i);$

$\lambda^i \leftarrow \frac{1}{\sqrt{\lambda_1^2 + \lambda_2^2}} (\lambda_1, \lambda_2)^\top;$

$\mathcal{X}_S \leftarrow \mathcal{X}_S \cup \text{FindAllOptimalFlows}(P_{\lambda^i}, f^i);$

// Return only flows with $c^1(f) \neq c^1(f^i)$ to avoid repetitions.

determined as optimal flows for the different weighted-sum problems (P_{λ^i}) . Moreover, unsupported flows correspond to suboptimal solutions for all weighed-sum problems. Since we only store flows for $i = 2, \dots, N - 1$ where $c^1(f) \neq c^1(f^i)$ no supported flow is stored twice. Thus, Algorithm 1 determines the complete set of all supported flows.

The enhanced parametric network approach [22] requires $\mathcal{O}(Nn(m+n \log n) + M)$ time, where M is the time required to solve a single-objective minimum cost flow problem. Since the algorithm determines the extreme points in a decreasing order of $c^1(f)$ we do not need additional time to sort the extreme points. Defining the weight vectors λ^i for $i = 1, \dots, N - 1$ and building the network with the corresponding cost function takes $\mathcal{O}(N(n + m))$ time. Determining all \mathcal{F}_i optimal flows for one weighted-sum problem (P_{λ^i}) using the algorithm FindAllOptimalFlows($D_{i, i+1}$) from [18], requires $\mathcal{O}(\mathcal{F}_i(m + n) + mn)$ time. Since the image of every supported efficient solution lies at most on

two maximally nondominated faces, it holds that $\sum_{i=1}^{N-1} \mathcal{F}_i < 2\mathcal{S}$. We have to consider $N-1$ of these single-objective minimum cost flow problems corresponding the maximally nondominated faces. Hence, Algorithm 1 requires overall $\mathcal{O}(Nn(m+n \log n) + \mathcal{S}(m+n))$ time. \square

Note, that the determination of all supported efficient flows could easily be integrated during the enhanced parametric network approach [22]. Whenever a new extreme non-dominated vector is found, determine all optimal flows to (P_{λ_i}) with the AOF algorithm.

4.2 Multi-objective minimum cost flow problems

In the following, we derive an algorithm to determine the complete set of all supported efficient flows and hence all supported nondominated vectors to the multiobjective integer minimum cost flow problem.

First, we need to determine the set of extreme supported nondominated vectors and the associated weight space decomposition. We can determine all extreme nondominated vectors using the dual Benson's algorithm [8]. However, since the lexicographic version of a MOIMCF problem can be solved in polynomial time [15], we may also use the lexicographic dual Benson's algorithm recently presented by Bökler and Mutzel in [4]. Both versions work with the upper image $\mathcal{P} := \text{conv}(\mathcal{Y} + \mathbb{R}_{\geq}^d)$ and its dual polyhedron, or *lower image* \mathcal{D} .

While we work with normalized weight vectors $\lambda \in \Lambda_d$, it suffices to consider the so-called *projected weight space* $\{(\lambda_1, \dots, \lambda_{d-1}) \in \Lambda_{d-1} : \sum_{i=1}^{d-1} \lambda_i < 1\}$ and calculate the normalized weighting vector $\ell(v) := (v_1, \dots, v_{d-1}, \sum_{i=1}^{d-1} v_i)$ of a projected weight v when needed. The dual problem (D_{λ}) of the weighted sum-problem (P_{λ}) is given by

$$\begin{aligned} \max \quad & b^{\top} u \\ \text{s.t.} \quad & A^{\top} u = C^{\top} \lambda \\ & u \in \mathbb{R}_{\geq}^m. \end{aligned} \tag{D_{\lambda}}$$

The dual polyhedron \mathcal{D} then consists of vectors $(\lambda_1, \dots, \lambda_{d-1}, b^{\top} u)$ with $\lambda \in \Lambda_d^0$ and solutions u of (D_{λ}) . Following the duality theory of polyhedra there exists a bijective mapping Ψ between the set of all faces of \mathcal{P} and the set of all faces of \mathcal{D} such that Ψ is *order reversing*, i. e., if two faces F_1 and F_2 of \mathcal{P} satisfy $F_1 \subseteq F_2$ then $\Psi(F_1) \supseteq \Psi(F_2)$ and $\Psi(F_1)$ and $\Psi(F_2)$ are faces of \mathcal{D} , see e.g., [23]. Thus, an extreme point of \mathcal{D} corresponds to a facet of \mathcal{P} and an extreme point of \mathcal{P} corresponds to a facet of \mathcal{D} . The dual Benson's algorithm solves an MOLP by computing the extreme points of \mathcal{D} . For more details on the dual Benson's algorithm or its lexicographic version we refer to [4, 8].

Thus, we obtain all extreme nondominated vectors and one corresponding extreme efficient solution for each of the extreme nondominated vectors, as well as all facets of \mathcal{P} . On this basis, we yield the *weight space decomposition* using the dual (lexicographic) Benson's algorithm. The set of weighting vectors associated with a point $y \in \mathcal{Y}$ is given by

$$\mathcal{W}(y) := \left\{ w \in \Lambda_d^0 : w^{\top} y \leq w^{\top} y' \text{ for all } y' \in \mathcal{P} \right\}.$$

Note, that the facets of \mathcal{P} may only be *weakly* nondominated, i. e., they might contain dominated (integer feasible) vectors. Recall that all supported nondominated vectors can be determined by a parametric MCF problem (P_λ) for some weight vector $\lambda \in \Lambda_d$. However, the weight vectors corresponding to the facets of \mathcal{P} might have components equal to zero $\lambda_i = 0$ for $i \in \{1, \dots, d\}$, i. e., $\lambda \in \Lambda_d^0$. In the following we describe a recursive algorithm to obtain the weight vectors for all maximally nondominated faces.

Let U be the set of all extreme points in the lower image \mathcal{D} and $\{\lambda^u : u \in U\}$ the set of corresponding weight vectors. Let, furthermore, be F_u the facet of \mathcal{P} corresponding to $u \in U$. Then, we call two extreme points u and u' of \mathcal{D} *adjacent*, iff $\dim(F_u \cap F_{u'}) = d-2$. In the following, we will denote the set of adjacent extreme points for $u \in U$ by $Q_u \subseteq U$. Recall that the intersection of k adjacent facets yields a $d-k$ dimensional face. For each $\lambda^u \in \Lambda_d$ (i. e., $\lambda^u > 0$) we know that all vectors on the facet F_u are supported nondominated vectors. Thus, we only have to solve the all optimum flow problem on (P_{λ^u}). Since some solutions may lie in the same sub-faces of adjacent facets, we have to ensure that no solution is stored twice. In order to do so, we keep track of the neighbouring extreme points during Benson's algorithm and store all already processed adjacent extreme points of $u \in U$ in a list δ_u .

There may exist maximally nondominated faces (with dimension less than $d-1$), which are intersections of a number of facets for which the corresponding weight vector equals zero in at least one component. We call these facets *weakly nondominated facets* of \mathcal{P} . In order to determine all supported efficient solutions we investigate nondominated faces which are intersections of weakly supported facets. With $U_{>} := \{u \in U : \lambda^u > 0\}$ we denote the set of extreme points of \mathcal{D} corresponding to nondominated facets and with $U_0 := \{u \in U : \lambda^u \geq 0\}$ the set of extreme points corresponding to all weakly nondominated facets ($U_{>} \subseteq U_0$). Note that weakly nondominated faces can contain supported nondominated points only at its (relative) boundary, while unsupported nondominated points can be located also in its (relative) interior.

Figure 6 presents the weight space decomposition for the example given in Section 2 and illustrated in Figure 3. Any point in the weight space decomposition corresponds to an λ^u for each $u \in U$. Here $U_{>} = \emptyset$. However, there do exist weights in the lines in the interior connecting two adjacent extreme points of \mathcal{D} which correspond to the maximally nondominated faces $[y^1, y^2]$ and $[y^2, y^3]$.

Theorem 4.4 *Given the directed network (D, l, u, b, c) , Algorithm 2 determines the complete set of all supported flows in $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor}(\text{poly}(n, m) + N \log N + N^{\lfloor \frac{d}{2} \rfloor} \mathcal{S}(m+n+N^{\lfloor \frac{d}{2} \rfloor} d)))$ time.*

Proof. *Correctness:* Any supported efficient flow must lie in at least one maximally efficient face. However a supported efficient flow can lie in more than one face. Due to Theorem 4.1, all supported flows are found by determining all optimal flows for each weight vectors λ^i corresponding to a maximally nondominated face. Moreover, no weakly supported flow can be optimal for a parametric network flow problem with one of these cost functions. While the algorithm iterates through all maximally nondominated faces only flows are stored which have not been considered yet. Thus, Algorithm 2 determines




Figure 6: Weight space decomposition to the upper image of Figure 3. Here it holds $\lambda_3 = 1 - \lambda_2 - \lambda_1$.

Algorithm 2: FindAllSupportedEfficientFlows

Data: (D, l, u, b, c)

Result: The complete set of all supported efficient flows

```

{U, Q_u, F_u, lambda_u, f^u : u in U} ← BensonLex(D, l, u, b, c) ;
// Determine all extreme points of D the corresponding facets of P
// and weight vectors.
delta_u = empty set for all u in U;
for u in U_> do
    FindAllOptimalFlows(P_{lambda^u}, f^u);
    // In the FindAllOptimalFlows algorithm only store flows f for
    // which <lambda^u, C f> != min{<lambda^{u'}, C f'> : f' in X} for any u' in delta_u
    for u' in Q_u do
        | delta_{u'} = delta_{u'} union {u}
w_u = {lambda^{u'} : u' in delta_u} for all u in U;
B = empty set;
for u in U_0 \setminus U_> do
    | Tilde_U = {u};
    ConsiderSubFaces(Tilde_U, U_0 \setminus U_>, B, Q_u, delta_u, lambda^u, f^u, w_u for all u in U);
    B = B union {u}

```

the complete set of all supported efficient solutions.

Run-time: Benson's Algorithm requires $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor} (\text{poly}(n, m) + N \log N))$ time [4]. Thereafter, we consider each face at most one. The number of all faces can be bounded by $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor})$ [4]. For each weakly nondominated face we check if a strict convex combination with adjacent weight vectors yields a weight vector $\lambda > 0$ componentwise strictly greater than zero. In this case, we call Algorithm 3. Note, that λ is not part

Algorithm 3: ConsiderSubFaces

Data: $(\tilde{U}, U_0 \setminus U_{>}, B, Q_u, \delta_u, \lambda^u, f^u, w_u) \quad \forall u \in U$

Result: The complete set of all unseen supported efficient flows in the maximally nondominated sub-faces of the corresponding facet F_u of \mathcal{P} .

Let $\bar{U} := \{u' \in (\bigcap_{u \in \tilde{U}} Q_u \cap U_0 \setminus U_{>}) \setminus (B \cup \tilde{U})\}$;

for $u \in \bar{U}$ **do**

$\tilde{U} = \tilde{U} \cup \{u\}$;

if $|\tilde{U}| \leq d - 1$ **then**

$\lambda^k := \sum_{i=1}^{|\tilde{U}|} \ell_i \lambda^{\tilde{u}_i}$ for an

$\ell \in \{\sum_{i=1}^{|\tilde{U}|} \ell_i = 1, \ell_i > 0 \quad \forall i \in \{1, \dots, |\tilde{U}|\}\}, \tilde{u}_i \in \tilde{U}$;

if $\lambda^k > 0$ **then**

 FindAllOptimalFlows($P_{\lambda^k}, f^{\tilde{u}_1}$);

 // Only store flows f for which

$\langle \lambda^k, Cf \rangle \neq \{\min\{\langle \lambda^{u'}, Cf' \rangle : f' \in \mathcal{X}\} : \lambda^{u'} \in \bigcup_{u \in \tilde{U}} w_u\}$

for $u' \in \bigcup_{u \in \tilde{U}} Q_u$ **do**

$w_{u'} = w_{u'} \cup \{\lambda^k\}$;

else

 ConsiderSubFaces($\tilde{U}, U_0 \setminus U_{>}, B, Q_u, \delta_u, \lambda^u, f^u, w_u \quad \forall u \in U$);

$B = B \cup \{u\}$

of the input of Algorithm 2. However, it can be shown that these encoding lengths can be bounded by $\mathcal{O}(\text{poly}(n, m))$ [4]. The convex combination can be computed in $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor} d)$. For each maximally nondominated face, which is not a facet, we first must also create the weight vector through a strictly convex combination. Afterwards we solve the AOF problem for (P_λ) for all of these maximally nondominated faces in time $\mathcal{O}(F_i(m+n+N^{\lfloor \frac{d}{2} \rfloor} d) + mn)$, where F_i is the number of optimal solutions for the current weighted sum problem (P_λ) . Additionally, it takes $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor} d)$ time to check if the flow is also optimal for an adjacent already considered maximally nondominated face. Since each flow may be contained in all faces we obtain the bound $\sum F_i \leq \mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor} \mathcal{S})$, where \mathcal{S} is the number of all supported efficient flows. Hence, Algorithm 2 requires overall $\mathcal{O}(N^{\lfloor \frac{d}{2} \rfloor} (\text{poly}(n, m) + N \log N + N^{\lfloor \frac{d}{2} \rfloor} \mathcal{S}(m+n+N^{\lfloor \frac{d}{2} \rfloor} d)))$ time. \square

5 Conclusion

Some previous papers use inconsistent characterizations of supported nondominated vectors for multi-objective integer linear programs (MOILP). In this paper we propose the definition of weakly supported nondominated vectors and distinguish clearly between weakly supported and supported nondominated vectors, which is particularly important for multiobjective integer programming with more than three objectives.

Then, we can conclude that there is no output-polynomial algorithm for a MOIMCF

problem with a fixed number of d objectives that determines all weakly supported nondominated vectors, unless $\mathbf{P} = \mathbf{NP}$. In contrast, this paper presents output-polynomial time algorithms for determining all supported efficient solutions for BOIMCF problems and general MOIMCF problems with a fixed number of objectives. First, the approach determines all extreme supported nondominated vectors and the weighting vectors for each maximally nondominated face. Then, it successively determines all supported efficient solutions in the preimage of each maximally nondominated face by determining all optimal solutions for the corresponding single-objective parametric network flow problem using the all optimum flow algorithm recently presented in [18].

However, it might be that many supported efficient flows may be mapped to the same vector in the objective space. Consider a directed graph with $\{1, \dots, n\}$ transshipment nodes (flow balance equal to zero), a node s and t with flow balance n and $-n$, respectively. The graph contains the arcs (s, i) and (i, t) for all $i \in \{1, \dots, n\}$ with upper capacity n . The cost of all arcs is equal to $(1, 1)^\top \in \mathbb{R}^2$. Then we have $\binom{2n-1}{n}$ supported efficient solutions but all map to the same extreme nondominated point. Thus, often a minimal complete set (all nondominated vectors and one (efficient) preimage for each of them) is considered as solution of a multiobjective optimization problem [27]. An open question remains if there exist an output-polynomial time algorithm to determine all supported nondominated vectors for a MOIMCF problem with a fixed number of objectives.

Even though, an output-polynomial time algorithm to determine all nondominated vectors for MOIMCF problems does not exist, even for the bi-objective case [3], future research could focus on new approaches to compute also unsupported nondominated vectors/efficient solutions in bi- or even multi-objective MCF problems. It is well known that unsupported solutions may be good compromise solutions and should thus not be neglected completely. Note, that the difficulty to compute unsupported solutions is not a specific property multiobjective integer network flow problems but arises in many integer and combinatorial optimization problems, and is one reason for their computational complexity, in general [6, 12]. One way to overcome this computational burden—at least to a certain degree—could be to determine unsupported solutions only in regions of the Pareto front which are not well represented by the set of supported nondominated points.

Acknowledgements. We would like to thank Fritz Bökler for his very kind and in depth discussion on the properties of the dual Benson’s algorithm and on the complexity of multi-objective combinatorial optimization problems. Moreover, we would like to thank Kathrin Klamroth for many fruitful discussions on our work.

References

- [1] R. K. Ahuja, T. L. Magnanti, and J. B. Orlin. *Network Flows: Theory, Algorithms, and Applications*. Prentice Hall, 1993.
- [2] D. P. Bertsekas. *Network optimization*. Athena Scientific, 1998.

- [3] F. Böklér, M. Ehrgott, C. Morris, and P. Mutzel. Output-sensitive complexity of multiobjective combinatorial optimization. *Journal of Multi-Criteria Decision Analysis*, 24(1-2):25–36, 2017.
- [4] F. Böklér and P. Mutzel. Output-sensitive algorithms for enumerating the extreme nondominated points of multiobjective combinatorial optimization problems. In N. Bansal and I. Finocchi, editors, *Algorithms - ESA 2015*, pages 288–299, Berlin, Heidelberg, 2015. Springer Berlin Heidelberg.
- [5] R. Diestel. *Graphentheorie*. Springer, Berlin, 2006.
- [6] M. Ehrgott. Hard to say it’s easy – four reasons why combinatorial multiobjective programmes are hard. In Y. Y. Haimes and R. E. Steuer, editors, *Research and Practice in Multiple Criteria Decision Making*, volume 487 of *Lecture Notes in Economics and Mathematical Systems*, pages 69–80. Springer Berlin Heidelberg, 2000.
- [7] M. Ehrgott. *Multicriteria optimization*, volume 491. Springer Science & Business Media, 2005.
- [8] M. Ehrgott, A. Löhne, and L. Shao. A dual variant of Benson’s “outer approximation algorithm” for multiple objective linear programming. *Journal of Global Optimization*, 52(4):757–778, 2012.
- [9] A. Eusébio and J. R. Figueira. Finding non-dominated solutions in bi-objective integer network flow problems. *Computers & Operations Research*, 36:2554–2564, 2009.
- [10] A. Eusébio and J. R. Figueira. On the computation of all supported efficient solutions in multi-objective integer network flow problems. *European Journal of Operational Research*, 199(1):68–76, 2009.
- [11] A. Eusébio, J. R. Figueira, and M. Ehrgott. On finding representative non-dominated points for bi-objective integer network flow problems. *Computers & Operations Research*, 48:1–10, 2014.
- [12] J. R. Figueira, C. M. Fonseca, P. Halffmann, K. Klamroth, L. Paquete, S. Ruzika, B. Schulze, M. Stiglmayr, and D. Willems. Easy to say they’re hard, but hard to see they’re easy—towards a categorization of tractable multiobjective combinatorial optimization problems. *Journal of Multi-Criteria Decision Analysis*, 24(1-2):82–98, 2017.
- [13] M. Fonseca, J. Figueira, and M. Resende. Solving scalarized multi-objective network flow problems using an interior point method. *International Transactions in Operational Research*, 17:607–636, 2010.
- [14] T. Gal. A general method for determining the set of all efficient solutions to a linear vectormaximum problem. *European Journal of Operational Research*, 1:307–322, 1977.

- [15] H. W. Hamacher, C. R. Pedersen, and S. Ruzika. Multiple objective minimum cost flow problems: A review. *European Journal of Operational Research*, 176(3):1404–1422, 2007.
- [16] D. S. Johnson, M. Yannakakis, and C. H. Papadimitriou. On generating all maximal independent sets. *Information Processing Letters*, 27(3):119–123, 1988.
- [17] H. Kellerer, U. Pferschy, and D. Pisinger. *Knapsack Problems*. Springer, Berlin, Germany, 2004.
- [18] D. Könen, D. Schmidt, and C. Spisla. Finding all minimum cost flows and a faster algorithm for the k best flow problem. *Discrete Applied Mathematics*, 321:333–349, 2022.
- [19] E. L. Lawler, J. K. Lenstra, and A. H. G. Rinnooy Kan. Generating all maximal independent sets: Np-hardness and polynomial-time algorithms. *SIAM Journal on Computing*, 9(3):558–565, 1980.
- [20] A. Przybylski, X. Gandibleux, and M. Ehrgott. A recursive algorithm for finding all nondominated extreme points in the outcome set of a multiobjective integer programme. *INFORMS Journal on Computing*, 22:371–386, 2010.
- [21] A. Raith and M. Ehrgott. A two-phase algorithm for the biobjective integer minimum cost flow problem. <http://www.esc.auckland.ac.nz/research/tech/esc-tr-661.pdf>, 36, June 2009.
- [22] A. Raith and A. Sedeño-Noda. Finding extreme supported solutions of biobjective network flow problems: An enhanced parametric programming approach. *Computers & Operations Research*, 82, 2017.
- [23] B. Schulze, K. Klamroth, and M. Stiglmayr. Multi-objective unconstrained combinatorial optimization: a polynomial bound on the number of extreme supported solutions. *Journal of Global Optimization*, 74(3):495–522, 2019.
- [24] A. Sedeño-Noda. An algorithm for the biobjective integer minimum cost flow problem. *Computers & Operations Research*, 28:139–156, 2001.
- [25] A. Sedeño-Noda and C. Gonzalez-Martin. Biobjective minimum cost flow problem. *European Journal of Operational Research*, 124:591–600, 2000.
- [26] A. Sedeño-Noda and C. González-Martin. An alternative method to solve the biobjective minimum cost flow problem. *Asia-Pacific Journal of Operational Research*, 2003.
- [27] P. Serafini. Some considerations about computational complexity for multi objective combinatorial problems. In J. Jahn and W. Krabs, editors, *Recent Advances and Historical Development of Vector Optimization*, volume 294 of *Lecture Notes in Economics and Mathematical Systems*, pages 222–232. Springer, 1987.

- [28] R. E. Steuer. Multiple criteria optimization: Theory, computation, and application. John Wiley, New York, 2008.
- [29] M. Sun. Finding integer efficient solutions for multiple objective network programming problems. *Networks*, 57:362–375, July 2011.