

Deformed Newton's (s, t) -Binomial Series and Generating Functions of Generalized Central Binomial Coefficients and Generalized Catalan Numbers

Ronald Orozco López

August 2, 2024

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 2 |
| 2 | Preliminaries | 3 |
| 2.1 | Generalized Fibonacci polynomials | 3 |
| 2.2 | Generalized (s, t) -binomial coefficients | 5 |
| 2.3 | Derivative on generalized Fibonacci polynomials | 6 |
| 3 | Deformed Newton (s, t)-binomial series | 7 |
| 3.1 | Definition | 7 |
| 3.2 | Properties | 9 |
| 3.3 | Proportional functional difference equation | 12 |
| 4 | The central (s, t)-binomial coefficients | 13 |
| 5 | Generating functions of central (s, t)-binomial coefficients | 15 |
| 5.1 | Deformed (s, t) -binomial series with rational powers | 15 |
| 5.2 | Generating function of generalized Catalan numbers | 17 |

Abstract

We define the deformed (s, t) -binomial formula and the deformed Newton (s, t) -binomial series, and we will use it to establish the generating functions of the generalized central binomial coefficients and the generalized Catalan numbers.

2020 *Mathematics Subject Classification*: Primary 05A15. Secondary 11B39; 11B65; 05A10.

Keywords: Generalized Fibonacci polynomial, generalized simplicial polytopical numbers, generalized central binomial coefficients, generalized Catalan numbers, Newton series.

1 Introduction

The Catalan numbers are defined by

$$C_n = \frac{1}{n+1} \binom{2n}{n}$$

for $n \geq 0$. It is a known fact that the generating function of Catalan's numbers is

$$C(x) = \frac{1 - \sqrt{1 - 4x}}{2x}. \quad (1)$$

The generalized Catalan numbers [7, 14, 2] are defined by

$$C_{\{n\}} = \frac{1}{\{n+1\}_{s,t}} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}_{s,t}. \quad (2)$$

The generalized Catalan numbers are polynomials in s, t with nonnegative integral coefficients [7], and its combinatorial interpretation is given in [2]. However, a generating function, analogous to Eq.(1), for the polynomials $C_{\{n\}}$ is not known. This is the main goal of this paper. Therefore all the results are original and useful.

In this paper, we introduce the deformed (s, t) -analog of the binomial theorem

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$$

as the equation given by

$$(x \oplus_{u,v} y)^{(n)} = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{s,t} u^{\binom{n-k}{2}} v^{\binom{k}{2}} x^{n-k} y^k. \quad (3)$$

When make $s = u = q$, $t = v = 1$, $a = t$ and $x = 1$, $y = t$ in Eq.(3), we obtain the q -theorem binomial

$$\prod_{k=0}^{n-1} (1 + q^k t) = \sum_{k=0}^n q^{k(k-1)/2} \binom{n}{k}_q t^k$$

where $\binom{n}{k}_q$ are the Gaussian binomial coefficients [4, 5].

On the other hand, if in Eq.(3) we extend the defining range of n to complex numbers, then we obtain the (u, v) -deformed Newton's (s, t) -binomial series

$$(x \oplus_{u,v} y)^{(\alpha)} = \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} u^{\binom{\alpha-k}{2}} v^{\binom{k}{2}} x^{\alpha-k} y^k. \quad (4)$$

This article is divided as follows. Section 2 addresses basic and necessary results for developing this article. Section 3 introduces the deformed Newton (s, t) -binomial series and establishes its algebraic properties. In Section 4, we express the fibonomial coefficients $\left\{ \begin{matrix} 1/2 \\ n \end{matrix} \right\}$ and $\left\{ \begin{matrix} -1/2 \\ n \end{matrix} \right\}$ in term of central (s, t) -binomial coefficients. Finally, we conclude our paper by focusing on finding the generating functions of the generalized binomial coefficients introduced above.

2 Preliminaries

2.1 Generalized Fibonacci polynomials

The generalized Fibonacci numbers on the parameters s, t are defined by

$$\{n+2\}_{s,t} = s\{n+1\}_{s,t} + t\{n\}_{s,t} \quad (5)$$

with initial values $\{0\}_{s,t} = 0$ and $\{1\}_{s,t} = 1$, where $s \neq 0$ and $t \neq 0$. In [14] this sequence is called the generalized Lucas sequence. In this paper, the generalized Lucas numbers $\langle n \rangle_{s,t}$ on the parameters s, t are defined recursively by

$$\langle n+2 \rangle_{s,t} = s\langle n+1 \rangle_{s,t} + t\langle n \rangle_{s,t}$$

for $n \geq 2$, with the initial conditions $\langle 0 \rangle_{s,t} = 2$ and $\langle 1 \rangle_{s,t} = s$. The Lucas sequence $L_n = (2, 1, 3, 4, 7, 11, 18, 29, \dots)$ is obtained when $s = 1, t = 1$ and is not a special case of the sequence in Eq.(5). Therefore, we will insist on calling to $\{n\}_{s,t}$ the generalized Fibonacci numbers and we will reserve the name of generalized Lucas numbers for the sequence $\langle n \rangle_{s,t}$.

Below are some important specializations of generalized Fibonacci and Lucas numbers.

1. When $s = 2, t = -1$, then $\{n\}_{2,-1} = n$, the positive integer and $\langle n \rangle_{2,-1} = 2$, for all $n \in \mathbb{N}$.
2. When $s = 1, t = 1$, then $\{n\}_{1,1} = F_n$, the Fibonacci numbers and $\langle 2 \rangle_{1,1} = L_n$, the Lucas numbers.
3. When $s = 2, t = 1$, then $\{n\}_{2,1} = P_n$, where $P_n = (0, 1, 2, 5, 12, 29, 70, 169, 408, \dots)$ are the Pell numbers and $\langle n \rangle_{2,1} = Q_n$, where $Q_n = (2, 2, 6, 14, 34, 82, 198, \dots)$ are the Pell-Lucas numbers.
4. When $s = 1, t = 2$, then $\{n\}_{1,2} = J_n$, where $J_n = (0, 1, 1, 2, 3, 5, 11, 21, 43, 85, \dots)$ are the Jacobsthal numbers and $\langle n \rangle_{1,2} = j_n$, where $j_n = (2, 1, 5, 7, 17, 31, 65, 127, \dots)$ are the Jacobsthal-Lucas numbers.
5. When $s = p + q, t = -pq$, then $\{n\}_{p+q,-pq} = [n]_{p,q} = \frac{p^n - q^n}{p - q}$ are the (p, q) -numbers and $\langle n \rangle_{p+q,-pq} = [2n]_{p,q} / [n]_{p,q}$.
6. When $s = 2t, t = -1$, then $\{n\}_{2t,-1} = U_{n-1}(t)$, where $U_n(t)$ are the Chebyshev polynomials of the second kind, with $U_{-1}(t) = 0$ and $\langle n \rangle_{2t,-1} = 2T_n(t)$, where $T_n(t)$ are the Chebyshev polynomials of the first kind.
7. When $s = 3, t = -2$, then $\{n\}_{3,-2} = M_n$, where $M_n = 2^n - 1$ are the Mersenne numbers and $\langle n \rangle_{3,-2} = 2^n + 1$.
8. When $s = P, t = -Q$, then $\{n\}_{P,-Q} = U_n(P, -Q)$, where $U_n(P, -Q)$ is the $(P, -Q)$ -Fibonacci sequence, with P, Q integer numbers, and $\langle n \rangle_{P,-Q} = V(n)$ is the $(P, -Q)$ -Lucas sequence. If $Q = -1$, then the sequence $U_n(P, -1)$ reduces to the P -Fibonacci sequence. If $s = x$ and $t = 1$, we obtain the Fibonacci polynomials

The (s, t) -Fibonacci constant is the ratio toward which adjacent (s, t) -Fibonacci polynomials tend. This is the only positive root of $x^2 - sx - t = 0$. We will let $\varphi_{s,t}$ denote this constant, where

$$\varphi_{s,t} = \frac{s + \sqrt{s^2 + 4t}}{2}$$

and

$$\varphi'_{s,t} = s - \varphi_{s,t} = -\frac{t}{\varphi_{s,t}} = \frac{s - \sqrt{s^2 + 4t}}{2}.$$

Some Generalized Fibonacci and Lucas polynomials are given in the table following.

| n | $\{n\}_{s,t}$ | $\langle n \rangle_{s,t}$ |
|-----|-----------------------|---------------------------------|
| 0 | 0 | 2 |
| 1 | 1 | s |
| 2 | s | $s^2 + 2t$ |
| 3 | $s^2 + t$ | $s^3 + 3st$ |
| 4 | $s(s^2 + 2t)$ | $s^4 + 4s^2t + 2t^2$ |
| 5 | $s^4 + 3s^2t + t^2$ | $s^5 + 5s^3t + 5st^2$ |
| 6 | $s^5 + 4s^3t + 3st^2$ | $s^6 + 6s^4t + 11s^2t^2 + 2t^3$ |

It is a known fact [14] that

$$\{2n\}_{s,t} = \langle n \rangle_{s,t} \{n\}_{s,t}. \quad (6)$$

Definition 1. For all $\alpha \in \mathbb{C}$ we define the generalized Fibonacci functions $\{\alpha\}_{s,t}$ as

$$\{\alpha\}_{s,t} = \frac{\varphi_{s,t}^\alpha - \varphi'_{s,t}{}^\alpha}{\varphi_{s,t} - \varphi'_{s,t}} \quad (7)$$

and the generalized Lucas functions $\langle \alpha \rangle_{s,t}$ as

$$\langle \alpha \rangle_{s,t} = \varphi_{s,t}^\alpha + \varphi'_{s,t}{}^\alpha. \quad (8)$$

Example 1. For all $s, t \in \mathbb{R}$,

$$\left\{\frac{1}{2}\right\}_{s,t} = \frac{1}{\sqrt{\frac{s + \sqrt{s^2 + 4t}}{2}} + \sqrt{\frac{s - \sqrt{s^2 + 4t}}{2}}}$$

and

$$\left\langle\frac{1}{2}\right\rangle_{s,t} = \sqrt{\frac{s + \sqrt{s^2 + 4t}}{2}} + \sqrt{\frac{s - \sqrt{s^2 + 4t}}{2}}.$$

The negative (s, t) -Fibonacci functions are given by

$$\{-\alpha\}_{s,t} = -(-t)^{-\alpha} \{\alpha\}_{s,t} \quad (9)$$

for all $\alpha \in \mathbb{R}$.

2.2 Generalized (s, t) -fibonomial coefficients

The (s, t) -fibonomial coefficients are define by

$$\begin{Bmatrix} n \\ k \end{Bmatrix}_{s,t} = \frac{\{n\}_{s,t}!}{\{k\}_{s,t}! \{n-k\}_{s,t}!},$$

where $\{n\}_{s,t}! = \{1\}_{s,t} \{2\}_{s,t} \cdots \{n\}_{s,t}$ is the (s, t) -factorial or generalized fibotorial.

The (s, t) -fibonomial coefficients satisfy the following Pascal recurrence relationships. For $1 \leq k \leq n-1$ it holds that

$$\begin{aligned} \begin{Bmatrix} n+1 \\ k \end{Bmatrix}_{s,t} &= \varphi_{s,t}^k \begin{Bmatrix} n \\ k \end{Bmatrix}_{s,t} + \varphi_{s,t}^{(n+1-k)} \begin{Bmatrix} n \\ k-1 \end{Bmatrix}_{s,t}, \\ &= \varphi_{s,t}^{(k)} \begin{Bmatrix} n \\ k \end{Bmatrix}_{s,t} + \varphi_{s,t}^{n+1-k} \begin{Bmatrix} n \\ k-1 \end{Bmatrix}_{s,t}. \end{aligned}$$

Definition 2. For all $\alpha \in \mathbb{C}$ the generalized (s, t) -fibonomial coefficient is

$$\begin{Bmatrix} \alpha \\ k \end{Bmatrix}_{s,t} = \frac{\{\alpha\}_{s,t} \{\alpha-1\}_{s,t} \cdots \{\alpha-k+1\}_{s,t}}{\{k\}_{s,t}!}. \quad (10)$$

Equally, the generalized (s, t) -fibonomial coefficient satisfy the following relationships:

$$\begin{Bmatrix} \alpha+1 \\ k \end{Bmatrix}_{s,t} = \varphi_{s,t}^k \begin{Bmatrix} \alpha \\ k \end{Bmatrix}_{s,t} + \varphi_{s,t}^{(\alpha+1-k)} \begin{Bmatrix} \alpha \\ k-1 \end{Bmatrix}_{s,t}, \quad (11)$$

$$= \varphi_{s,t}^{(k)} \begin{Bmatrix} \alpha \\ k \end{Bmatrix}_{s,t} + \varphi_{s,t}^{\alpha+1-k} \begin{Bmatrix} \alpha \\ k-1 \end{Bmatrix}_{s,t}. \quad (12)$$

Proposition 1. For all nonzero $a \in \mathbb{C}$,

$$\begin{Bmatrix} \alpha \\ n \end{Bmatrix}_{as,a^2t} = a^{n(\alpha-n)} \begin{Bmatrix} \alpha \\ n \end{Bmatrix}_{s,t}. \quad (13)$$

Proof. As $\{n\}_{as,a^2t} = a^{n-1} \{n\}_{s,t}$, then $\{n\}_{as,a^2t}! = a^{\binom{n}{2}} \{n\}_{s,t}!$. Therefore,

$$\begin{aligned} \begin{Bmatrix} \alpha \\ n \end{Bmatrix}_{as,a^2t} &= \frac{\{\alpha\}_{as,a^2t} \{\alpha-1\}_{as,a^2t} \cdots \{\alpha-n+1\}_{as,a^2t}}{\{n\}_{as,a^2t}!} \\ &= \frac{a^{\alpha-1+\alpha-2+\cdots+\alpha-n} \{\alpha\}_{s,t} \{\alpha-1\}_{s,t} \cdots \{\alpha-n+1\}_{s,t}}{a^{\binom{n}{2}} \{n\}_{s,t}!} \\ &= \frac{a^{n\alpha - \binom{n+1}{2}} \{\alpha\}_{s,t} \{\alpha-1\}_{s,t} \cdots \{\alpha-n+1\}_{s,t}}{a^{\binom{n}{2}} \{n\}_{s,t}!} \\ &= a^{n(\alpha-n)} \frac{\{\alpha\}_{s,t} \{\alpha-1\}_{s,t} \cdots \{\alpha-n+1\}_{s,t}}{\{n\}_{s,t}!}. \end{aligned}$$

The proof is completed. □

Proposition 2. The negative (s, t) -fibonomial polynomial is

$$\begin{Bmatrix} -\alpha \\ k \end{Bmatrix}_{s,t} = (-1)^k (-t)^{-\alpha k - \binom{k}{2}} \begin{Bmatrix} \alpha+k-1 \\ k \end{Bmatrix}_{s,t}.$$

Proof. By using Eq.(9)

$$\begin{aligned}
\binom{-\alpha}{k}_{s,t} &= \frac{\{-\alpha\}_{s,t} \{-\alpha-1\}_{s,t} \cdots \{-\alpha-k+1\}_{s,t}}{\{k\}_{s,t}!} \\
&= \frac{[-(-t)^{-\alpha} \{\alpha\}_{s,t}] [-(-t)^{-(\alpha+1)} \{\alpha+1\}_{s,t}] \cdots [-(-t)^{-(\alpha+k-1)} \{\alpha+k-1\}_{s,t}]}{\{k\}_{s,t}!} \\
&= \frac{(-1)^k (-t)^{-k\alpha - \binom{k}{2}} \{\alpha+k-1\}_{s,t} \{\alpha+k-2\}_{s,t} \cdots \{\alpha+1\}_{s,t} \{\alpha\}_{s,t}}{\{k\}_{s,t}!} \\
&= (-1)^k (-t)^{-k\alpha - \binom{k}{2}} \binom{\alpha+k-1}{k}_{s,t}.
\end{aligned}$$

The proof is completed. \square

2.3 Derivative on generalized Fibonacci polynomials

Definition 3. Set $s, t \in \mathbb{R}$, $s \neq 0$, $t \neq 0$. If $s^2 + 4t \neq 0$, define the (s, t) -derivative $\mathbf{D}_{s,t}$ of the function $f(x)$ as

$$(\mathbf{D}_{s,t}f)(x) = \begin{cases} \frac{f(\varphi_{s,t}x) - f(\varphi'_{s,t}x)}{(\varphi_{s,t} - \varphi'_{s,t})x}, & \text{if } x \neq 0; \\ f'(0), & \text{if } x = 0 \end{cases} \quad (14)$$

provided $f(x)$ differentiable at $x = 0$. If $s^2 + 4t = 0$, $t < 0$, define the $(\pm i\sqrt{t}, t)$ -derivative of the function $f(x)$ as

$$(\mathbf{D}_{\pm i\sqrt{t}, t}f)(x) = f'(\pm i\sqrt{t}x). \quad (15)$$

If $(\mathbf{D}_{s,t}f)(x)$ exist at $x = a$, then $f(x)$ is (s, t) -differentiable at a .

Example 2. For all $s, t \in \mathbb{R}$, $s \neq 0$, $t \neq 0$

$$\mathbf{D}_{s,t}x^n = \frac{(\varphi_{s,t}x)^n - (\varphi'_{s,t}x)^n}{(\varphi_{s,t} - \varphi'_{s,t})x} = \{n\}_{s,t}x^{n-1}, \quad n \in \mathbb{N}$$

and x^n is (s, t) -differentiable in \mathbb{R} .

Example 3. For all $s, t \in \mathbb{R}$, $s \neq 0$, $t \neq 0$ and $x \neq 0$ the (s, t) -derivative of $f(x) = e^x$ is

$$\mathbf{D}_{s,t}e^x = \frac{e^{\varphi_{s,t}x} - e^{\varphi'_{s,t}x}}{(\varphi_{s,t} - \varphi'_{s,t})x} = \begin{cases} 2 \frac{e^{sx/2}}{\sqrt{s^2+4t}} \sinh\left(\frac{\sqrt{s^2+4t}}{2}x\right), & \text{if } s^2 + 4t > 0; \\ e^{\pm i\sqrt{t}x}, & \text{if } s^2 + 4t = 0, t < 0; \\ 2 \frac{e^{sx/2}}{\sqrt{-s^2-4t}} \sin\left(\frac{\sqrt{-s^2-4t}}{2}x\right), & \text{if } s^2 + 4t < 0. \end{cases}$$

As $f'(0) = 1$, then $\mathbf{D}_{s,t}e^x$ is continuous in \mathbb{R} and (s, t) -differentiable in \mathbb{R} .

Proposition 3. For all $\alpha, \beta \in \mathbb{R}$,

1. $\mathbf{D}_{s,t}(\alpha f + \beta g) = \alpha \mathbf{D}_{s,t}f + \beta \mathbf{D}_{s,t}g$.
2. $\mathbf{D}_{s,t}(f(x)g(x)) = f(\varphi_{s,t}x)\mathbf{D}_{s,t}g(x) + g(\varphi'_{s,t}x)\mathbf{D}_{s,t}f(x)$.
3. $\mathbf{D}_{s,t}(f(x)g(x)) = f(\varphi'_{s,t}x)\mathbf{D}_{s,t}g(x) + g(\varphi_{s,t}x)\mathbf{D}_{s,t}f(x)$.

4.

$$\mathbf{D}_{s,t} \left(\frac{f(x)}{g(x)} \right) = \frac{g(\varphi_{s,t}x) \mathbf{D}_{s,t} f(x) - f(\varphi_{s,t}x) \mathbf{D}_{s,t} g(x)}{g(\varphi_{s,t}x) g(\varphi'_{s,t}x)}.$$

5.

$$\mathbf{D}_{s,t} \left(\frac{f(x)}{g(x)} \right) = \frac{g(\varphi'_{s,t}x) \mathbf{D}_{s,t} f(x) - f(\varphi'_{s,t}x) \mathbf{D}_{s,t} g(x)}{g(\varphi_{s,t}x) g(\varphi'_{s,t}x)}.$$

3 Deformed Newton (s, t) -binomial series

3.1 Definition

Definition 4. For all $\alpha \in \mathbb{C}$ and $s \neq 0, t \neq 0$ define the (u, v) -deformed (s, t) -binomial series, for x, y commuting, as

$$(x \oplus_{u,v} y)_{s,t}^{(\alpha)} = \sum_{n=0}^{\infty} \left\{ \alpha \right\}_{s,t}^{(n)} u^{\binom{\alpha-n}{2}} v^{\binom{n}{2}} x^{\alpha-n} y^n. \quad (16)$$

We define the (u, v) -deformed (s, t) -analogue of $(x - y)^\alpha$ by

$$(x \ominus_{u,v} y)_{s,t}^{(\alpha)} \equiv (x \oplus_{u,v} (-y))_{s,t}^{(\alpha)}.$$

Definition 5. For all nonzero complex numbers $u, v, u, v \in \mathbb{C}$, define the (u, v) -deformed (s, t) -power, for x, y commuting, as

$$(x \oplus_{u,v} y)_{s,t}^{(n)} = \sum_{k=0}^n \left\{ n \right\}_{s,t}^{(k)} u^{\binom{n-k}{2}} v^{\binom{k}{2}} x^{n-k} y^k \quad n \geq 0.$$

Example 4.

$$\begin{aligned} (x \oplus_{u,v} y)_{s,t}^{(0)} &= 1, \\ (x \oplus_{u,v} y)_{s,t}^{(1)} &= x + y, \\ (x \oplus_{u,v} y)_{s,t}^{(2)} &= ux^2 + \{2\}_{s,t} xy + vy^2, \\ (x \oplus_{u,v} y)_{s,t}^{(3)} &= u^3 x^3 + \{3\}_{s,t} ux^2 y + \{3\}_{s,t} vxy^2 + v^3 y^3, \\ (x \oplus_{u,v} y)_{s,t}^{(4)} &= u^6 x^4 + \{4\}_{s,t} u^3 x^3 y + \{3\}_{s,t} \langle 2 \rangle_{s,t} uvx^2 y^2 + \{4\}_{s,t} v^3 xy^3 + v^6 y^4. \end{aligned}$$

In post quantum calculus, or (p, q) -calculus, the (p, q) -powers are defined by Sadjang in [12] by

$$(x \ominus a)_{p,q}^n = \begin{cases} 1, & \text{if } n = 0; \\ \prod_{k=0}^{n-1} (p^k x - q^k a), & \text{if } n \geq 1. \end{cases} \quad (17)$$

$$(x \oplus a)_{p,q}^n = \begin{cases} 1, & \text{if } n = 0; \\ \prod_{k=0}^{n-1} (p^k x + q^k a), & \text{if } n \geq 1. \end{cases} \quad (18)$$

In q -calculus the q -shifted factorial is

$$(a; q)_n = \begin{cases} 1, & \text{if } n = 0; \\ \prod_{k=0}^{n-1} (1 - aq^k), & \text{if } n \geq 1. \end{cases}$$

When $u = \varphi$ and $v = \varphi'$, then

$$(x \ominus_{\varphi, \varphi'} a)_{s,t}^{(n)} = (x \ominus a)_{\varphi, \varphi'}^n = \varphi_{s,t}^{(2)} x^n (a/x; q)_n$$

where $x \neq 0$ and $q = \varphi'_{s,t}/\varphi_{s,t}$. Therefore, the (u, v) -deformed (s, t) -power generalize the (p, q) -power. Also, the (u, v) -deformed (s, t) -power generalize the (p, q) -Gauss Binomial formula

$$(x \ominus a)_{p,q}^n = \sum_{k=0}^n \binom{n}{k}_{p,q} p^{\binom{n-k}{2}} q^{\binom{k}{2}} x^{n-k} (-a)^k, \quad (19)$$

the Jackson-Hanh-Cigler q -addition [10, 3, 8]

$$(x \boxplus_q y)^n = \sum_{k=0}^n \binom{n}{k}_q q^{\binom{k}{2}} x^k y^{n-k},$$

the Ward-Al-Salam q -addition [1, 15]

$$(x \oplus_q y)^n = \sum_{k=0}^n \binom{n}{k}_q x^k y^{n-k},$$

and the Golden Binomial formula [13]

$$(x + y)_F^n = \sum_{k=0}^n \binom{n}{k}_F (-1)^{\binom{k}{2}} x^k y^{n-k}.$$

For $s = 2$ and $t = -1$, the (u, v) -deformed n -th power of $x + y$ is

$$(x \oplus_{u,v} y)^{(n)} = \sum_{k=0}^n \binom{n}{k} u^{\binom{n-k}{2}} v^{\binom{k}{2}} x^{n-k} y^k. \quad (20)$$

Eqs.(17),(18) are extended to

$$\begin{aligned} (x \ominus_{\varphi, \varphi'} y)_{s,t}^{(\infty)} &= \prod_{k=0}^{\infty} (\varphi_{s,t}^k x - \varphi'_{s,t}{}^k y), \\ (x \oplus_{\varphi, \varphi'} y)_{s,t}^{(\infty)} &= \prod_{k=0}^{\infty} (\varphi_{s,t}^k x + \varphi'_{s,t}{}^k y). \end{aligned}$$

It is easy to verify that

$$(x \oplus_{\varphi, \varphi'} y)_{s,t}^{(n)} = \frac{(x \oplus_{\varphi, \varphi'} y)_{s,t}^{(\infty)}}{(x \varphi_{s,t}^n \oplus_{\varphi, \varphi'} y \varphi_{s,t}^n)_{s,t}^{(\infty)}} \quad (21)$$

for $n \in \mathbb{N}$. Then Eq.(21) can be extended for all complex number as

$$(x \oplus_{\varphi, \varphi'} y)_{s,t}^{(\alpha)} = \frac{(x \oplus_{\varphi, \varphi'} y)_{s,t}^{(\infty)}}{(x \varphi_{s,t}^\alpha \oplus_{\varphi, \varphi'} y \varphi_{s,t}^\alpha)_{s,t}^{(\infty)}}, \quad (22)$$

$\alpha \in \mathbb{C}$.

3.2 Properties

The following are properties satisfied by the formula (u, v) -deformed (s, t) -binomial.

Theorem 1. *For all $\alpha, x, y, z, u, v \in \mathbb{C}$ the (u, v) -deformed (s, t) -binomial series has the following properties*

1. $(x \oplus_{u,v} y)_{s,t}^{(\alpha+1)} = x(ux \oplus_{u,v} \varphi_{s,t} y)_{s,t}^{(\alpha)} + y(\varphi'_{s,t} x \oplus_{u,v} vy)_{s,t}^{(\alpha)},$
2. $(x \oplus_{u,v} y)_{s,t}^{(\alpha+1)} = x(ux \oplus_{u,v} \varphi'_{s,t} y)_{s,t}^{(\alpha)} + y(\varphi_{s,t} x \oplus_{u,v} vy)_{s,t}^{(\alpha)},$
3. $z^\alpha(x \oplus_{u,v} y)_{s,t}^{(\alpha)} = (zx \oplus_{u,v} zy)_{s,t}^{(\alpha)},$
4. $(x \oplus_{au,av} y)_{as,a^2t}^{(\alpha)} = a^{\binom{\alpha}{2}}(x \oplus_{u,v} y)_{s,t}^{(\alpha)},$
5. $(x \oplus_{u,v} y)_{s,t}^{(\alpha)} = (y \oplus_{v,u} x)_{s,t}^{(\alpha)},$
6. $(x \oplus_{u,v} 0)^{(\alpha)} = u^{\binom{\alpha}{2}} x^\alpha, \text{ for all } x \in \mathbb{C}.$
7. $(0 \oplus_{u,v} y)^{(\alpha)} = v^{\binom{\alpha}{2}} y^\alpha, \text{ for all } y \in \mathbb{C}.$

Proof. By Eq.(16)

$$(x \oplus_{u,v} y)_{s,t}^{(\alpha+1)} = \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha+1 \\ k \end{matrix} \right\}_{s,t} u^{\binom{\alpha+1-k}{2}} v^{\binom{k}{2}} x^{\alpha+1-k} y^k.$$

By extracting the first summand and by using Eq.(11), we obtain

$$\begin{aligned} & (x \oplus_{u,v} y)_{s,t}^{(\alpha+1)} \\ &= u^{\binom{\alpha+1}{2}} x^{\alpha+1} + \sum_{k=1}^{\infty} \left(\varphi_{s,t}^k \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} + \varphi'_{s,t}{}^{(\alpha-k+1)} \left\{ \begin{matrix} \alpha \\ k-1 \end{matrix} \right\}_{s,t} \right) v^{\binom{k}{2}} u^{\binom{\alpha+1-k}{2}} y^k x^{\alpha+1-k} \\ &= u^{\binom{\alpha+1}{2}} x^{\alpha+1} + \sum_{k=1}^{\infty} \varphi_{s,t}^k \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v^{\binom{k}{2}} u^{\binom{\alpha+1-k}{2}} y^k x^{\alpha+1-k} \\ &\quad + \sum_{k=1}^{\infty} \varphi'_{s,t}{}^{(\alpha-k+1)} \left\{ \begin{matrix} \alpha \\ k-1 \end{matrix} \right\}_{s,t} v^{\binom{k}{2}} u^{\binom{\alpha+1-k}{2}} y^k x^{\alpha+1-k} \\ &= u^{\binom{\alpha+1}{2}} x^{\alpha+1} + x \sum_{k=1}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v^{\binom{k}{2}} u^{\binom{\alpha-k}{2}} (\varphi_{s,t} y)^k (ux)^{\alpha-k} \\ &\quad + \sum_{k=1}^{\infty} \varphi'_{s,t}{}^{(\alpha-k+1)} \left\{ \begin{matrix} \alpha \\ k-1 \end{matrix} \right\}_{s,t} v^{\binom{k}{2}} u^{\binom{\alpha+1-k}{2}} y^k x^{\alpha+1-k}. \end{aligned}$$

In addition, we rearrange the second summation for k

$$\begin{aligned}
(x \oplus_{u,v} y)_{s,t}^{(\alpha+1)} &= u \binom{\alpha+1}{2} x^{\alpha+1} + x \sum_{k=1}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{k}{2} u \binom{\alpha-k}{2} (\varphi_{s,t} y)^k (ux)^{\alpha-k} \\
&\quad + \sum_{k=0}^{\infty} \varphi'_{s,t} \binom{\alpha}{k} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{k+1}{2} u \binom{\alpha-k}{2} y^{k+1} x^{\alpha-k} \\
&= u \binom{\alpha+1}{2} x^{\alpha+1} + x \sum_{k=1}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{k}{2} u \binom{\alpha-k}{2} (\varphi_{s,t} y)^k (ux)^{\alpha-k} \\
&\quad + y \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{k}{2} u \binom{\alpha-k}{2} (vy)^k (\varphi'_{s,t} x)^{\alpha-k}.
\end{aligned}$$

Then

$$\begin{aligned}
&(x \oplus_{u,v} y)_{s,t}^{(\alpha+1)} \\
&= x \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{k}{2} u \binom{\alpha-k}{2} (\varphi_{s,t} y)^k (ux)^{\alpha-k} + y \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{k}{2} u \binom{\alpha-k}{2} (vy)^k (\varphi'_{s,t} x)^{\alpha-k} \\
&= x(ux \oplus_{u,v} \varphi_{s,t} y)_{s,t}^{(\alpha)} + y(\varphi'_{s,t} x \oplus_{u,v} vy)_{s,t}^{(\alpha)}
\end{aligned}$$

and thus we obtain the first statement. By using (s, t) -Pascal formula, Eq.(12), we obtain

2. The proof of 3 is trivial. From Proposition 1

$$\begin{aligned}
(x \oplus_{au,av} y)_{as,a^2t}^{(\alpha)} &= \sum_{n=0}^{\infty} \left\{ \begin{matrix} \alpha \\ n \end{matrix} \right\}_{as,a^2t} (au)^{\binom{\alpha-n}{2}} (av)^{\binom{n}{2}} x^{\alpha-n} y^n \\
&= \sum_{n=0}^{\infty} a^{n(\alpha-1) + \binom{\alpha-n}{2} + \binom{n}{2}} \left\{ \begin{matrix} \alpha \\ n \end{matrix} \right\}_{s,t} u^{\binom{\alpha-n}{2}} v^{\binom{n}{2}} x^{\alpha-n} y^n \\
&= a^{\binom{\alpha}{2}} \sum_{n=0}^{\infty} \left\{ \begin{matrix} \alpha \\ n \end{matrix} \right\} u^{\binom{\alpha-n}{2}} v^{\binom{n}{2}} x^{\alpha-n} y^n = a^{\binom{\alpha}{2}} (x \oplus_{u,v} y)_{s,t}^{(\alpha)}.
\end{aligned}$$

Which leads us to the truth of 4. By changing u to v and x to y in the Eq.(16) we find that

$$\begin{aligned}
(x \oplus_{u,v} y)_{s,t}^{(\alpha)} &= \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{k}{2} u \binom{\alpha-k}{2} y^k x^{\alpha-k} \\
&= \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} v \binom{\alpha-k}{2} u \binom{k}{2} y^{\alpha-k} y^k \\
&= (y \oplus_{v,u} x)_{s,t}^{(\alpha)}.
\end{aligned}$$

Then it proves 5. The statement 6 and 7 are trivial. \square

We will now give a result on the convergence of Eq.(16). Set $x \neq 0$. As $(x \oplus_{u,v} y)_{s,t}^{(\alpha)} = x^\alpha (1 \oplus_{u,v} (y/x))_{s,t}^{(\alpha)}$, then it is sufficient to prove our result for the series representation of $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$.

Theorem 2. *Set $s \neq 0$, $t \neq 0$ and pick $u \neq 0$, $|q| = |\varphi'_{s,t}/\varphi_{s,t}| \neq 1$. For all $\alpha \in \mathbb{C}$ we have that*

1. If $|uv| < |t|$ and $|q| \neq 1$, then $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$ is a function entire.
2. If $|uv| = |t|$, then $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$ is convergent in $|x| < |(\varphi_{s,t}/\varphi'_{s,t})(u/\varphi'_{s,t})^{\alpha-1}|$ if $0 < |q| < 1$ or convergent in $|x| < |(\varphi'_{s,t}/u)(u/\varphi_{s,t})^\alpha|$ if $|q| > 1$.
3. If $|uv| > |t|$ and $|q| \neq 1$, then $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$ converge in $x = 0$.

Proof. By applying the criteria of D'Alembert to the series representation of $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$ we have

$$\begin{aligned}
\lim_{k \rightarrow \infty} \left| \frac{\{\alpha\}_{k+1} v^{\binom{k+1}{2}} u^{\binom{\alpha-k-1}{2}} x^{k+1}}{\{\alpha\}_k v^{\binom{k}{2}} u^{\binom{\alpha-k}{2}} x^k} \right| &= \lim_{k \rightarrow \infty} \left| \frac{x \{\alpha - k\}_{s,t} v^k}{\{k+1\}_{s,t} u^{\alpha-k-1}} \right| \\
&= \left| \frac{x}{u^{\alpha-1}} \right| \lim_{k \rightarrow \infty} |uv|^k \left| \frac{\{\alpha - k\}_{s,t}}{\{k+1\}_{s,t}} \right| \\
&= \left| \frac{x}{u^{\alpha-1}} \right| \lim_{k \rightarrow \infty} |uv|^k \left| \frac{\varphi_{s,t}^{\alpha-k} - \varphi_{s,t}^{(\alpha-k)}}{\varphi_{s,t}^{k+1} - \varphi_{s,t}^{(k+1)}} \right| \\
&= \left| \frac{x}{u^{\alpha-1}} \right| \lim_{k \rightarrow \infty} |uv|^k |\varphi_{s,t}^{\alpha-2k-1}| \left| \frac{1 - q^{\alpha-k}}{1 - q^{k+1}} \right| \\
&= \left| \frac{x \varphi_{s,t}^{\alpha-1}}{u^{\alpha-1}} \right| \lim_{k \rightarrow \infty} \left| \frac{uv}{\varphi_{s,t}^2} \right|^k \left| \frac{1 - q^{\alpha-k}}{1 - q^{k+1}} \right|.
\end{aligned}$$

Set $A = uv/\varphi_{s,t}^2$ and define

$$H_k = |A|^k \left| \frac{1 - q^{\alpha-k}}{1 - q^{k+1}} \right| = \left| \frac{A}{q} \right|^k \left| \frac{1}{q^{-k} - q} - \frac{q^\alpha}{1 - q^{k+1}} \right|.$$

If $|A| < |q|$ and if $0 < |q| < 1$ or if $|q| > 1$, then $H_k \rightarrow 0$ as $k \rightarrow \infty$ and therefore $x \in \mathbb{C}$ for each $\alpha \in \mathbb{C}$. If $|A| = |q|$, then $H_k \rightarrow |q|^\alpha$ if $|q| < 1$ and $H_k \rightarrow |q|^{-1}$ if $|q| > 1$ as $k \rightarrow \infty$. Finally, if $|A| > |q|$, then $H_k \rightarrow \infty$ as $k \rightarrow \infty$. The three statements follow from the above. \square

Theorem 3. Set $s \neq, t \neq 0$ and pick $u \neq 0$. For all $\alpha \in \mathbb{C}$ we have that

1. If $|uv| < 1$, then $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$ is a function entire.
2. If $|uv| = 1$, then $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$ is convergent in $|x| < |u^{\alpha-1}|$.
3. If $|uv| > 1$, then $(1 \oplus_{u,v} x)_{s,t}^{(\alpha)}$ converge in $x = 0$.

The analytical importance of research into series representation with deformations lies in their convergence. For example, $f(z) = 1/(1-z)$ has a convergent series representation when $|z| < 1$, $z \in \mathbb{C}$. While $(1 \ominus_{1,v} z)^{(-1)}$, the $(1, v)$ -deformed analog of $f(z)$, has a convergent series representation in all \mathbb{C} whenever $|v| < 1$.

Theorem 4. For all $\alpha \in \mathbb{C}$

1. $\mathbf{D}_{s,t}(x \oplus_{u,v} a)_{s,t}^{(\alpha)} = \{\alpha\}_{s,t}(ux \oplus_{u,v} a)_{s,t}^{(\alpha-1)}$.

$$2. \mathbf{D}_{s,t}(a \oplus_{u,v} x)_{s,t}^{(\alpha)} = \{\alpha\}_{s,t}(a \oplus_{u,v} vx)_{s,t}^{(\alpha-1)}.$$

$$3. \mathbf{D}_{s,t}(a \ominus_{u,v} x)_{s,t}^{(\alpha)} = -\{\alpha\}_{s,t}(a \oplus_{u,v} vx)_{s,t}^{(\alpha-1)}.$$

Proof. We prove assertion 1. The other assertions are proved similarly.

$$\begin{aligned} \mathbf{D}_{s,t}(x \oplus_{u,v} a)_{s,t}^{(\alpha)} &= \mathbf{D}_{s,t} \left(\sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} u^{\binom{k}{2}} v^{\binom{\alpha-k}{2}} x^k a^{\alpha-k} \right) \\ &= \sum_{k=1}^{\infty} \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} u^{\binom{k}{2}} v^{\binom{\alpha-k}{2}} \{k\}_{s,t} x^{k-1} a^{\alpha-k} \\ &= \sum_{k=0}^{\infty} \left\{ \begin{matrix} \alpha \\ k+1 \end{matrix} \right\}_{s,t} u^{\binom{k}{2}} u^k v^{\binom{\alpha-k-1}{2}} \{k+1\}_{s,t} x^k a^{\alpha-k-1} \\ &= \sum_{k=0}^{\infty} \frac{\{\alpha\}_{s,t} \{\alpha-1\}_{s,t} \cdots \{\alpha-k\}_{s,t}}{\{k\}_{s,t}!} u^{\binom{k}{2}} v^{\binom{\alpha-1-k}{2}} (ux)^k a^{\alpha-1-k} \\ &= \{\alpha\}_{s,t} \sum_{k=0}^{\infty} \frac{\{\alpha-1\}_{s,t} \cdots \{\alpha-1-k+1\}_{s,t}}{\{k\}_{s,t}!} u^{\binom{k}{2}} v^{\binom{\alpha-1-k}{2}} (ux)^k a^{\alpha-1-k} \\ &= \{\alpha\}_{s,t} (ux \oplus_{u,v} a)_{s,t}^{(\alpha-1)}. \end{aligned}$$

□

When set $u = \varphi_{s,t}$ and $v = \varphi'_{s,t}$ in Theorem 4, then

$$\mathbf{D}_{s,t}(x \oplus_{\varphi,\varphi'} \alpha)_{s,t}^{(\alpha)} = \{\alpha\}_{s,t}(\varphi_{s,t}x \oplus_{\varphi,\varphi'} \alpha)_{s,t}^{(\alpha-1)}.$$

Theorem 5. For all $\alpha, u, v, a \in \mathbb{C}$

$$\begin{aligned} \mathbf{D}_{s,t}^k(x \oplus_{u,v} a)_{s,t}^{(\alpha)} &= u^{\binom{k}{2}} \{k\}_{s,t}! \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} (u^k x \oplus_{u,v} a)_{s,t}^{(\alpha-k)}, \\ \mathbf{D}_{s,t}^k(a \oplus_{u,v} x)_{s,t}^{(\alpha)} &= u^{\binom{k}{2}} \{k\}_{s,t}! \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} (a \oplus_{u,v} v^k x)_{s,t}^{(\alpha-k)}, \\ \mathbf{D}_{s,t}^k(a \ominus_{u,v} x)_{s,t}^{(\alpha)} &= (-1)^k v^{\binom{k}{2}} \{k\}_{s,t}! \left\{ \begin{matrix} \alpha \\ k \end{matrix} \right\}_{s,t} (a \ominus_{u,v} v^k x)_{s,t}^{(\alpha-k)}. \end{aligned}$$

Proof. They are obtained by induction on k .

□

3.3 Proportional functional difference equation

Remark 1. As

$$(1 \oplus_{u,v} x)_{s,t}^{(\alpha)} = u^{\binom{\alpha}{2}} (1 \oplus_{1,uv} u^{-\alpha+1} x)_{s,t}^{(\alpha)}, \quad (23)$$

then it is sufficient to research the $(1, v)$ -deformed (s, t) -binomial series $(1 \oplus_{1,v} x)_{s,t}^{(\alpha)}$.

Theorem 6. The function $f(x) = (1 \oplus_{1,v} x)_{s,t}^{(\alpha)}$ is solution of the proportional functional difference equation

$$(\mathbf{D}_{s,t} f)(\varphi_{s,t}x/v) + \varphi'^{(\alpha-1)}_{s,t} x (\mathbf{D}_{s,t} f)(x/\varphi'_{s,t}) = \{\alpha\}_{s,t} f(x), \quad (24)$$

with initial value $f(0) = 1$. If $v = \varphi_{s,t}\varphi'_{s,t}$, then

$$(\mathbf{D}_{s,t}f)(x/\varphi'_{s,t}) = \frac{\{\alpha\}_{s,t}f(x)}{1 + \varphi'^{(\alpha-1)}_{s,t}x}. \quad (25)$$

If $s = 2$ and $t = -1$, then $f(x) = (1 \oplus_{1,v} x)^{(\alpha)}$ is solution of the functional differential equation with proportional delay

$$f'(x/v) + xf'(x) = \alpha f(x). \quad (26)$$

Proof. Use the Theorems 1 and 4. □

The solution of Eq.(25) is

$$f(x) = \varphi_{s,t}^{-\binom{n}{2}} \frac{(1 \oplus_{\varphi,\varphi'} \varphi_{s,t}^{\alpha-1} x)^{(\infty)}}{(\varphi_{s,t}^{\alpha} \oplus_{\varphi,\varphi'} (-t)^{\alpha} \varphi_{s,t}^{-1} x)^{(\infty)}}.$$

The above theorem shows another justification for using deformed series representation. There is growing research on differential equations with proportional delay of the form $x'(t) = f(t, x, x(qt))$, or pantograph equations, [6], [9], [11]. Eq.(26) is part of the more general differential equation with proportional delay

$$f'(x) = F(x, f, f(qx), f'(qx)).$$

As Eq.(24) is the (s, t) -analog of Eq.(26), then it opens the way to research proportional difference equations defined on generalized Fibonacci polynomials.

4 The central (s, t) -binomial coefficients

The first thing to do is to establish some properties of the generalized central binomial coefficient $\left\{ \begin{smallmatrix} 2n \\ n \end{smallmatrix} \right\}_{s,t}$ and in particular to find an expression for the coefficients $\left\{ \begin{smallmatrix} 1/2 \\ n \end{smallmatrix} \right\}_{s,t}$ and $\left\{ \begin{smallmatrix} -1/2 \\ n \end{smallmatrix} \right\}_{s,t}$.

Definition 6. For all $n \in \mathbb{N}$ we define the functions $L_n(s, t)$ as

$$L_n(s, t) = \frac{(-t)^{-\frac{n^2}{2}} 4^n}{\langle n \rangle_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi,\varphi'} \sqrt{\varphi'_{s,t}})^{\binom{n}{2}}_{s,t}} \quad (27)$$

The function $L_n(s, t)$ reduces to unity when $s = 2$ and $t = -1$, that is,

$$L_n(2, -1) = \frac{4^n}{\langle n \rangle_{2,-1}! 2^n} = 1$$

for all $n \geq 0$. For the case $s = 1 + q$, $t = -q$, we get

$$L_n(q) \equiv L_n(1 + q, -q) = \frac{q^{-\frac{n^2}{2}} 4^n}{(-q; q)_n (-\sqrt{q}; q)_n}.$$

Proposition 4. Set $q = \varphi'_{s,t}/\varphi_{s,t}$. For all $n \in \mathbb{N}$

$$L_{n+1}(s, t) = \frac{4(-t)^{-(n+1/2)}}{\langle n+1 \rangle_{s,t} (\varphi_{s,t}^{n+1/2} + \varphi'^{(n+1/2)}_{s,t})} L_n(s, t) \quad (28)$$

Proof. From Definition 6,

$$\begin{aligned}
L_{n+1}(s, t) &= \frac{(-t)^{-\frac{(n+1)^2}{2}} 4^{n+1}}{\langle n+1 \rangle_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n+1)}} \\
&= \frac{4(-t)^{-(n+1/2)}}{\langle n+1 \rangle_{s,t} (\varphi_{s,t}^{n+1/2} + \varphi'_{s,t}{}^{(n+1/2)})} \frac{(-t)^{-\frac{n^2}{2}} 4^n}{\langle n \rangle_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n)}} \\
&= \frac{4(-t)^{-(n+1/2)}}{\langle n+1 \rangle_{s,t} (\varphi_{s,t}^{n+1/2} + \varphi'_{s,t}{}^{(n+1/2)})} L_n(s, t)
\end{aligned}$$

□

Proposition 5. For all $n \geq 1$

$$\left\{ \begin{matrix} 1/2 \\ n \end{matrix} \right\}_{s,t} = \left\{ \begin{matrix} 1 \\ 2 \end{matrix} \right\}_{s,t} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}_{s,t} \frac{(-1)^{n+1}}{4^{n-1} \langle n \rangle_{s,t} \{2n-1\}_{s,t}} L_{n-1}(s, t). \quad (29)$$

Proof. From Definition 2 and Eq.(9),

$$\begin{aligned}
\left\{ \begin{matrix} 1/2 \\ n \end{matrix} \right\}_{s,t} &= \frac{\{1/2\}_{s,t} \{-1/2\}_{s,t} \cdots \{-(2n-3)/2\}_{s,t}}{\{n\}_{s,t}!} \\
&= \frac{\{1/2\}_{s,t} (-(-t)^{-1/2} \{1/2\}_{s,t}) \cdots (-(-t)^{-(2n-3)/2} \{(2n-3)/2\}_{s,t})}{\{n\}_{s,t}!} \\
&= \frac{(-1)^{n-1} (-t)^{-\left(\frac{1}{2} + \frac{3}{2} + \cdots + \frac{2n-3}{2}\right)} \{1/2\}_{s,t}^2 \{3/2\}_{s,t} \cdots \{(2n-3)/2\}_{s,t}}{\{n\}_{s,t}!}.
\end{aligned}$$

As $\left\{ \frac{2n+1}{2} \right\}_{s,t} = \frac{\{2n+1\}_{s,t}}{\langle (2n+1/2) \rangle_{s,t}}$, then

$$\begin{aligned}
\left\{ \begin{matrix} 1 \\ 2 \end{matrix} \right\}_{s,t} \left\{ \begin{matrix} 3 \\ 2 \end{matrix} \right\}_{s,t} \cdots \left\{ \begin{matrix} 2n-3 \\ 2 \end{matrix} \right\}_{s,t} &= \frac{\{1\}_{s,t} \{3\}_{s,t} \cdots \{2n-3\}_{s,t}}{\langle 1/2 \rangle_{s,t} \langle 3/2 \rangle_{s,t} \cdots \langle (2n-3)/2 \rangle_{s,t}} \\
&= \frac{\{1\}_{s,t} \{3\}_{s,t} \cdots \{2n-3\}_{s,t}}{(\varphi_{s,t}^{1/2} + \varphi'_{s,t}{}^{1/2})(\varphi_{s,t}^{3/2} + \varphi'_{s,t}{}^{3/2}) \cdots (\varphi_{s,t}^{(2n-3)/2} + \varphi'_{s,t}{}^{(2n-3)/2})} \\
&= \frac{\{1\}_{s,t} \{3\}_{s,t} \cdots \{2n-3\}_{s,t}}{(\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n-1)}}
\end{aligned}$$

and

$$\left\{ \begin{matrix} 1/2 \\ n \end{matrix} \right\}_{s,t} = \left\{ \begin{matrix} 1 \\ 2 \end{matrix} \right\}_{s,t} \frac{(-1)^{n-1} (-t)^{-\frac{(n-1)^2}{2}} \{1\}_{s,t} \{3\}_{s,t} \cdots \{2n-3\}_{s,t}}{\{n\}_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n-1)}}.$$

On the other hand,

$$\{2\}_{s,t} \{4\}_{s,t} \cdots \{2n\}_{s,t} = \langle 1 \rangle_{s,t} \{1\}_{s,t} \langle 2 \rangle_{s,t} \{2\}_{s,t} \cdots \langle n \rangle_{s,t} \{n\}_{s,t} = \langle n \rangle_{s,t}! \{n\}_{s,t}!.$$

Therefore,

$$\begin{aligned}
\begin{Bmatrix} 1/2 \\ n \end{Bmatrix}_{s,t} &= \begin{Bmatrix} 1 \\ 2 \end{Bmatrix}_{s,t} \frac{(-1)^{n-1} (-t)^{-\frac{(n-1)^2}{2}} \{2n\}_{s,t}!}{\langle n \rangle_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n-1)} \{n\}_{s,t}! \{n\}_{s,t}! \{2n-1\}_{s,t}} \\
&= \begin{Bmatrix} 1 \\ 2 \end{Bmatrix}_{s,t} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} \frac{(-1)^{n-1} (-t)^{-\frac{(n-1)^2}{2}}}{\langle n \rangle_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n-1)} \{2n-1\}_{s,t}} \\
&= \begin{Bmatrix} 1 \\ 2 \end{Bmatrix}_{s,t} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} \frac{(-1)^{n-1}}{4^{n-1} \langle n \rangle_{s,t} \{2n-1\}_{s,t}} L_{n-1}(s, t).
\end{aligned}$$

The proof is completed. \square

Proposition 6. For all $n \in \mathbb{N}$,

$$\begin{Bmatrix} -1/2 \\ n \end{Bmatrix}_{s,t} = \frac{(-1)^n}{4^n} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} L_n(s, t). \quad (30)$$

Proof. From Definition 2 and Eq.(9),

$$\begin{aligned}
\begin{Bmatrix} -1/2 \\ n \end{Bmatrix}_{s,t} &= \frac{\{-1/2\}_{s,t} \{-3/2\}_{s,t} \cdots \{-(2n-1)/2\}_{s,t}}{\{n\}_{s,t}!} \\
&= \frac{(-(-t)^{-1/2} \{1/2\}_{s,t}) (-(-t)^{-3/2} \{3/2\}_{s,t}) \cdots (-(-t)^{-(2n-1)/2} \{(2n-1)/2\}_{s,t})}{\{n\}_{s,t}!} \\
&= \frac{(-1)^n (-t)^{-\left(\frac{1}{2} + \frac{3}{2} + \cdots + \frac{2n-1}{2}\right)} \{1/2\}_{s,t} \{3/2\}_{s,t} \cdots \{(2n-1)/2\}_{s,t}}{\{n\}_{s,t}!} \\
&= \frac{(-1)^n (-t)^{-\frac{n^2}{2}} \{1\}_{s,t} \{3\}_{s,t} \cdots \{2n-1\}_{s,t}}{\{n\}_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n)}} \\
&= \frac{(-1)^n (-t)^{-\frac{n^2}{2}} \{2n\}_{s,t}!}{(\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n)} \langle n \rangle_{s,t}! \{n\}_{s,t}! \{n\}_{s,t}!} \\
&= \frac{(-1)^n}{4^n} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} L_n(s, t).
\end{aligned}$$

The expected result is reached. \square

5 Generating functions of central (s, t) -binomial coefficients

5.1 Deformed (s, t) -binomial series with rational powers

Definition 7. For all non-zero real number v , $v \in \mathbb{R}$, such that $|v| \leq |t|$ and for all $\frac{n}{m} \in \mathbb{Q}$, $\frac{n}{m} > 0$, we define the $(1, v)$ -deformed (s, t) -binomial series with rational powers as

$$\sqrt[m]{(1 \oplus_{1,v} x)_{s,t}^{(n)}} \equiv (1 \oplus_{1,v} x)_{s,t}^{(n/m)} = \sum_{k=0}^{\infty} \begin{Bmatrix} n/m \\ k \end{Bmatrix}_{s,t} v^{(k)} x^k \quad (31)$$

and we define the "reciprocal" $(1, v)$ -deformed (s, t) -binomial series as

$${}^{(m)}\sqrt{(1 \oplus_{1,v} x)_{s,t}^{(-n)}} \equiv (1 \oplus_{1,v} x)_{s,t}^{(-n/m)} = \sum_{k=0}^{\infty} \left\{ \begin{matrix} -n/m \\ k \end{matrix} \right\}_{s,t} v^{\binom{k}{2}} x^k. \quad (32)$$

Proposition 7. For all $\frac{n}{m} \in \mathbb{Q}$,

1. ${}^{(n)}\sqrt{(1 \oplus_{1,v} x)_{s,t}^{(n)}} = 1 + x.$
2. ${}^{(m)}\sqrt{(1 \oplus_{1,v} x)_{s,t}^{(nm)}} = (1 \oplus_{1,v} x)_{s,t}^{(n)}.$
3. ${}^{(m)}\sqrt{(1 \oplus_{1,v} x)_{s,t}^{(n+m)}} = {}^{(m)}\sqrt{(1 \oplus_{1,v} \varphi_{s,t} x)_{s,t}^{(n)}} + x {}^{(m)}\sqrt{(\varphi'_{s,t} \oplus_{1,v} v x)_{s,t}^{(n)}}.$

From Theorem 5 we have the following results.

Theorem 7. For all $k \in \mathbb{N}$,

$$\mathbf{D}_{s,t}^k {}^{(m)}\sqrt{(1 \oplus_{1,v} x)_{s,t}^{(n)}} = v^{\binom{k}{2}} \{k\}_{s,t}! \left\{ \begin{matrix} n/m \\ k \end{matrix} \right\}_{s,t} {}^{(m)}\sqrt{(1 \oplus_{1,v} v^k x)_{s,t}^{(n-mk)}}.$$

Theorem 8. For $|v| \leq |t|$, the v -deformed series representation of ${}^{(2)}\sqrt{(1 \oplus_{1,v} 4x)_{s,t}}$ is

$${}^{(2)}\sqrt{(1 \oplus_{1,v} 4x)_{s,t}} = 1 + 4 \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=1}^{\infty} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}_{s,t} \frac{(-1)^{n-1} L_{n-1}(s, t)}{\langle n \rangle_{s,t} \{2n-1\}_{s,t}} v^{\binom{n}{2}} x^n. \quad (33)$$

Proof. By Definition 7 and Proposition 5

$$\begin{aligned} {}^{(2)}\sqrt{(1 \oplus_{1,v} x)_{s,t}} &= 1 + \sum_{n=1}^{\infty} \left\{ \frac{1}{2} \right\}_{s,t} v^{\binom{n}{2}} x^n \\ &= 1 + \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=1}^{\infty} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}_{s,t} \frac{(-1)^{n-1} (-t)^{-\frac{(n-1)^2}{2}} v^{\binom{n}{2}}}{\langle n \rangle_{s,t}! (\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{(n-1)} \{2n-1\}_{s,t}} x^n. \end{aligned}$$

Finish by changing x with $4x$. □

If we put $x = 1/4$ in Eq.(33), we get the series for the $(1, v)$ -deformed (s, t) -analog of $\sqrt{2}$

$${}^{(2)}\sqrt{(1 \oplus_{1,v} 1)_{s,t}} = 1 + 4 \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=1}^{\infty} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}_{s,t} \frac{(-1)^{n-1}}{\langle n \rangle_{s,t} \{2n-1\}_{s,t}} L_{n-1}(s, t) v^{\binom{n}{2}}.$$

The $(1, q)$ -deformed q -analog is

$${}^{(2)}\sqrt{(1 \oplus_{1,q} 1)_q} = 1 + 4 \left[\frac{1}{2} \right]_q \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(q^2; q^2)_n (q; q^2)_n}{(q; q)_n^2 (-q; q)_n (-\sqrt{q}; q)_n} \frac{q^{\frac{n-1}{2}} (1-q)}{1 - q^{2n-1}}$$

and the (s, t) -analog is

$${}^{(2)}\sqrt{(1 \oplus_{1,-t} 1)_{s,t}} = \varphi_{s,t}^{1/8} \frac{(1 \oplus_{\varphi, \varphi'} \varphi_{s,t}^{-1/2})_{s,t}^{(\infty)}}{(\sqrt{\varphi_{s,t}} \oplus_{\varphi, \varphi'} \sqrt{\varphi'_{s,t}})_{s,t}^{-1/2} (\infty)}.$$

Theorem 9. For $|v| \leq |t|$, the v -deformed series representation of $\sqrt[2]{(1 \ominus_{1,v} 4x)_{s,t}^{(-1)}}$ is

$$\sqrt[2]{(1 \ominus_{1,v} 4x)_{s,t}^{(-1)}} = \sum_{n=0}^{\infty} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} L_n(s, t) v^{\binom{n}{2}} x^n. \quad (34)$$

Proof. By Definition 7 and Proposition 6

$$\begin{aligned} \sqrt[2]{(1 \ominus_{1,v} 4x)_{s,t}^{(-1)}} &= \sum_{n=0}^{\infty} (-1)^n \begin{Bmatrix} -1/2 \\ n \end{Bmatrix}_{s,t} v^{\binom{n}{2}} x^n \\ &= \sum_{n=0}^{\infty} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} L_n(s, t) v^{\binom{n}{2}} x^n. \end{aligned}$$

The proof is reached. \square

If we put $x = 1/8$ in Eq.(34), we get another series for the v -deformed (s, t) -analog of $\sqrt{2}$

$$\sqrt[2]{(1 \ominus_{1,v} (1/2))_{s,t}^{(-1)}} = \sum_{n=0}^{\infty} \frac{1}{8^n} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} L_n(s, t) v^{\binom{n}{2}}.$$

If we put $x = -1/8$, we get another series for the v -deformed (s, t) -analog of $\sqrt{2/3}$

$$\sqrt[2]{(1 \oplus_{1,v} (1/2))_{s,t}^{(-1)}} = \sum_{n=0}^{\infty} \frac{(-1)^n}{8^n} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} L_n(s, t) v^{\binom{n}{2}}.$$

Averaging the two above series we get the v -deformed (s, t) -analog of $\frac{3\sqrt{2}+\sqrt{6}}{6}$

$$\sqrt[2]{(1 \oplus_{1,v} (1/2))_{s,t}^{(-1)}} + \sqrt[2]{(1 \ominus_{1,v} (1/2))_{s,t}^{(-1)}} = \sum_{n=0}^{\infty} \frac{1}{64^n} \begin{Bmatrix} 4n \\ 2n \end{Bmatrix}_{s,t} L_{2n}(s, t) v^{\binom{2n}{2}}.$$

5.2 Generating function of generalized Catalan numbers

Theorem 10. Set $|v| \leq |t|$. The v -deformed generating function of the Catalan polynomials $C_{\{n\}}$ is

$$\frac{1 - \sqrt[2]{(1 \ominus_{1,v} 4x)_{s,t}}}{2x} = 2 \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=0}^{\infty} C_{\{n\}} L_n(s, t) v^{\binom{n+1}{2}} x^n. \quad (35)$$

Proof. From Eq.(33),

$$\begin{aligned} \frac{1 - \sqrt[2]{(1 \ominus_{1,v} 4x)_{s,t}}}{2x} &= \frac{2}{x} \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=1}^{\infty} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} \frac{(-1)^n L_{n-1}(s, t)}{\langle n \rangle_{s,t} \{2n-1\}_{s,t}} v^{\binom{n}{2}} (-1)^n x^n \\ &= 2 \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=0}^{\infty} \begin{Bmatrix} 2n+2 \\ n+1 \end{Bmatrix}_{s,t} \frac{L_n(s, t)}{\langle n+1 \rangle_{s,t} \{2n+1\}_{s,t}} v^{\binom{n+1}{2}} x^n \\ &= 2 \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=0}^{\infty} \begin{Bmatrix} 2n \\ n \end{Bmatrix}_{s,t} \frac{L_n(s, t)}{\{n+1\}_{s,t}} v^{\binom{n+1}{2}} x^n. \end{aligned}$$

The proof is completed. \square

Then

$$\sum_{n=0}^{\infty} \frac{C_{\{n\}}}{4^n} L_n(s, t) v^{\binom{n+1}{2}} = \frac{1 - \sqrt[2]{(1 \ominus_{1,v} 1)_{s,t}}}{\{1/2\}_{s,t}}$$

is the v -deformed (s, t) -analog of $\sum_{n=0}^{\infty} \frac{C_n}{4^n} = 2$.

Theorem 11. Set $|v| \leq |t|$. The v -deformed generating function of the polynomials $\{n\}_{s,t} C_{\{n\}}$ is

$$\begin{aligned} & \frac{4\{1/2\}_{s,t} \varphi_{s,t} x - (1 - \sqrt[2]{(1 \ominus_{1,v} 4\varphi_{s,t} x)_{s,t}}) \sqrt[2]{(1 \ominus_{1,v} 4vx)_{s,t}}}{2\varphi_{s,t} \varphi'_{s,t} \sqrt[2]{(1 \ominus_{s,t} 4vx)_{s,t}}} \\ & = 2 \left\{ \frac{1}{2} \right\}_{s,t} \sum_{n=0}^{\infty} C_{\{n\}} \{n\}_{s,t} L_n(s, t) v^{\binom{n+1}{2}} x^n. \end{aligned}$$

Proof. Apply the operator $x\mathbf{D}_{s,t}$ to both sides of Eq.(35). \square

Theorem 12. Set $|v| \leq |t|$. The v -deformed generating function of the polynomials $C_{\{n\}_{s,t}} \{2n+1\}_{s,t} \langle n+1 \rangle_{s,t}$ is

$$\frac{(1 \ominus_{1,v} 4x)_{s,t}^{(-1/2)} - 1}{2x} = \frac{1}{2} \sum_{n=0}^{\infty} C_{\{n\}_{s,t}} \{2n+1\}_{s,t} \langle n+1 \rangle_{s,t} L_{n+1}(s, t) v^{\binom{n+1}{2}} x^n. \quad (36)$$

Proof. From Eq.(34)

$$\begin{aligned} \frac{(1 \ominus_{1,v} 4x)_{s,t}^{(-1/2)} - 1}{2x} &= \frac{1}{2x} \sum_{n=1}^{\infty} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}_{s,t} L_n(s, t) v^{\binom{n}{2}} x^{n-1} \\ &= \frac{1}{2} \sum_{n=0}^{\infty} \left\{ \begin{matrix} 2n+2 \\ n+1 \end{matrix} \right\}_{s,t} L_{n+1}(s, t) v^{\binom{n+1}{2}} x^n \\ &= \frac{1}{2} \sum_{n=0}^{\infty} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}_{s,t} \frac{\{2n+1\}_{s,t} \langle n+1 \rangle_{s,t}}{\{n+1\}_{s,t}} L_{n+1}(s, t) v^{\binom{n+1}{2}} x^n. \end{aligned}$$

The proof is completed. \square

References

- [1] W.A. Al-Salam, q -Bernoulli Numbers and Polynomials, *Math. Nachr* **17**(1959), 239–260.
- [2] C. Bennet, J. Carrillo, J. Machacek, and B. Sagan, Combinatorial Interpretations of Lucas Analogues of Binomial Coefficients and Catalan Numbers, *Annal of Combinatorics*, **24** (2020), 503–530.
- [3] J. Cigler, Operatormethoden für q -Identitäten. *Monatshefte Math.* **88**, 87–105 (1979)
- [4] C.F. Gauss, *Werke* 2, (1876), 9–45.
- [5] C.F. Gauss, *Werke* 3, (1876).

- [6] A. Ebaid, H.K Al-Jeaid, On the exact solution of the functional differential equation $y'(t) = ay(t) + by(-t)$, *Adv. Differ. Equ. Control Process*, **26**, (2022), 39–49.
- [7] S. Ekhad, The Sagan-Savage Lucas-Catalan polynomials have positive coefficients. Preprint, January 21, 2011. Available at <http://arxiv.org/abs/1101.4060>.
- [8] W. Hahn, Beitrage zur Theorie der Heineschen Reihen. *Math. Nachr.* **2**, 340–379 (1949)
- [9] A. Iserles, On the generalized pantograph functional-differential equation, *European J. Appl. Math.*, **4**, (1993), 1–38.
- [10] F.H. Jackson, On basic double hypergeometric functions, *Q. J. Math.*, **13**, (1942), 69–82.
- [11] T. Kato, J.B. McLeod, The Functional-Differential Equation $y'(x) = ay(\lambda x) + by(x)$, *Bulletin of the American Mathematical Society*, **77** 6, (1971), 891–937.
- [12] P. Njionou Sadjang, On the fundamental Theorem of (p, q) -Calculus and Some (p, q) -Taylor Formulas. *Results Math* **73**, 39 (2018) <https://doi.org/10.1007/s00025-018-0783-z>
- [13] O.K Pashaev, S. Nalci, Golden Quantum Oscillator and Binet-Fibonacci Calculus, *J. Phys. A: Math.Theor.*, **45**, 015303 (2012)
- [14] B. Sagan and C. Savage, Combinatorial interpretations of binomial coefficient analogs related to Lucas sequences, *Integers*, **10** (2010), A52, 697–703.
- [15] M. Ward, A calculus of sequences, *Am. J. Math.* **58**(1936) 255–266.