Strong gravitational lensing as a probe of dark matter

S.Vegetti¹, S. Birrer², G. Despali^{3,4}, C. D. Fassnacht⁵, D. Gilman⁶, Y. Hezaveh^{7,8,9,10,11}, L. Perreault Levasseur^{7,8,9,10,11}, J. P. McKean^{12,13}, D. M. Powell¹, C. M. O'Riordan¹, G. Vernardos¹⁴

Received: date / Accepted: date

Abstract Dark matter structures within strong gravitational lens galaxies and along their line of sight leave a gravitational imprint on the multiple images of lensed sources. Strong gravitational lensing provides, therefore, a key test of different dark matter models in a way that is independent of the baryonic content of matter structures on subgalactic scales. In this chapter, we describe how galaxy-scale strong gravitational lensing observations are sensitive to the physical nature of dark matter. We provide a historical perspec-

1. Max Planck Institut für Astrophysik, Karl-Schwarzschild-Strasse 1, D-85748 Garching bei München, Germany

- 2. Department of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794, USA
- 3. Dipartimento di Fisica e Astronomia "A. Righi", Alma Mater Studiorum Università di Bologna, Via Piero Gobetti 93/2, I-40129 Bologna, Italy
- 4. Institut für Theoretische Astrophysik, Zentrum für Astronomie, Heidelberg Universität, Albert-Ueberle-Str. 2, 69120, Heidelberg, Germany
- 5. Dept. of Physics and Astronomy, University of California, Davis, 1 Shields Ave. Davis, CA 95616, USA
- 6. Department of Astronomy and Astrophysics, University of Toronto, Toronto, ON, M5S 3H4, Canada
- 7. Department of Physics, Université de Montréal, Montréal, Canada
- 8. Mila Quebec Artificial Intelligence Institute, Montréal, Canada
- 9. Ciela Montreal Institute for Astrophysical Data Analysis and Machine Learning, Montréal, Canada
- 10. Center for Computational Astrophysics, Flatiron Institute, 162 5th Avenue, 10010, New York, NY, USA
- 11. Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada, N2L 2Y5
- 12. Kapteyn Astronomical Institute, University of Groningen, PO Box 800, NL-9700 AV Groningen, The Netherlands
- ASTRON, Netherlands Institute for Radio Astronomy, PO Box 2, NL-7990 AA Dwingeloo, The Netherlands
- 14. Institute of Physics, Laboratory of Astrophysique, École Polytechnique Fédérale de Lausanne (EPFL), Observatoire de Sauverny, 1290 Versoix, Switzerland

E-mail: svegetti@mpa-garching.mpg.de

tive of the field, and review its current status. We discuss the challenges and advances in terms of data, treatment of systematic errors and theoretical predictions, that will enable one to deliver a stringent and robust test of different dark matter models in the near future. With the advent of the next generation of sky surveys, the number of known strong gravitational lens systems is expected to increase by several orders of magnitude. Coupled with high-resolution followup observations, these data will provide a key opportunity to constrain the properties of dark matter with strong gravitational lensing.

Keywords gravitational lensing: strong; haloes: number, structure; dark matter

Contents

1	Introduction
2	Dark matter models
3	Strong lensing as a probe of dark matter
4	Lens modelling
5	Degeneracies and systematic errors
6	Historical perspective
7	Dark matter constraints
8	Future prospects
9	Summary and conclusions

1 Introduction

In the standard cosmological model, 85 per cent of the total amount of matter in the Universe is made of non-baryonic dark matter (Planck Collaboration et al., 2020b). Evidence of the existence of dark matter spans a wide range of independent astronomical observations: the cosmic micro-wave background (Planck Collaboration et al., 2020a), the rotation curves of disk galaxies (e.g. van Albada et al., 1985; Rubin, 1991), the motion of galaxies within clusters (e.g. Zwicky, 1933), gravitational lensing by galaxies and galaxy clusters (e.g Treu and Koopmans, 2004; Mandelbaum et al., 2006; Clowe et al., 2006; Auger et al., 2009; Barnabè et al., 2011; Sonnenfeld et al., 2022; Shajib et al., 2022), hot gas in galaxy clusters (e.g. Ettori et al., 2013), weak lensing and galaxy clustering (e.g. Alam et al., 2017; Heymans et al., 2021; Abbott et al., 2022). To this day, the physical nature of dark matter remains an unsolved problem. There is no dark matter particle within the standard model of particle physics. So far, several candidates, spanning 90 orders of magnitude in mass, have been proposed (Bertone and Tait, 2018). In the 1980s, weakly interactive massive particles (WIMPs) emerged as the favourite candidate for cold dark matter. The success of WIMPs stems from the fact that their existence is predicted by Supersymmetry theories (Fox, 2018, and references therein), according to which they are thermally produced with a self-interaction cross-section that leads to the correct amount of dark matter at the present day. However, decades of particle physics experiments have failed to produce a WIMP detection (Arcadi et al., 2018).

Other viable dark matter candidates include warm (WDM), self-interacting (SIDM) and fuzzy (FDM) dark matter. We refer the reader to Section 2 for a more detailed description. An interesting aspect of these alternative models is that they predict a different distribution of dark matter on subgalactic scales.

Strong gravitational lensing is sensitive to the distribution of matter and dark matter between the observer and the source. It is a purely gravitational probe and does not rely on the presence and distribution of baryons. It therefore provides a channel to observationally constrain the physical nature of dark matter. Other probes include the Lyman- α forest (e.g. Villasenor et al., 2022), the satellite galaxies of the Milky-Way (e.g. Nadler et al., 2021) and stellar streams in the Local Group (e.g. Erkal et al., 2017).

This chapter focuses on galaxy-scale strong gravitational lensing as a probe of dark matter and is organized as follows. In Section 2, we give a description of the dark matter models that have so far been tested with strong gravitational lensing observations. We expect this list to increase in the future as theoretical predictions for structure formation in more dark matter models become available. In Section 3, we discuss how lensing observables (i.e. image positions, fluxes, and time delays) are affected by the distribution of dark matter on small scales. In Section 4, we describe the process of lens modelling and how one can constrain the properties of dark matter from strong gravitational lensing observations as an inference problem. Another important aspect is the role played by degeneracies, systematic errors and unknowns, all of which are discussed in Section 5. The chapter then continues with an historical perspective of the field (Section 6), its current status (Section 7) and how it is likely to evolve in the near future (Section 8). We acknowledge that the last

two sections may soon be obsolete. Therefore, we encourage the interested reader to complement this text with the latest relevant publications. Finally, we present our concluding remarks in Section 9.

2 Dark matter models

2.1 Cold Dark Matter

Dark matter models where the particle is non-relativistic are described as cold. Possible candidates include WIMPs and axions (e.g. Feng, 2010). At the same time, dark matter does not have to be a particle and primordial black holes have also been proposed as a CDM candidate (e.g. Green and Kavanagh, 2021). One of the fundamental predictions of CDM is the fact that dark matter structures form hierarchically and bottom-up: low-mass haloes form first and subsequently merge into structures of increasing mass and size. The structure and evolution of CDM haloes has been extensively studied and precisely characterised using numerical simulations (see Zavala and Frenk, 2019, for a review). The collisionless nature of CDM and hierarchical structure formation result in a population of haloes with well defined properties.

Halo mass function: at the low-mass end, the number density n of haloes is well-described, to first approximation, by a power-law distribution:

$$\frac{dn}{dM} \propto M^{-\alpha} \,. \tag{1}$$

For halo masses smaller than a mass of $M \sim 10^{11} M_{\odot}$ the slope is $\alpha = 1.9$, and the number of structures increases with decreasing halo mass. The shape of the halo mass function is well understood from statistical arguments. These are based on the properties of the initial density field of fluctuations and the gravitational collapse process that leads to the formation of virialised haloes. This is at the basis of the formalism first introduced by Press and Schechter (1974), and subsequently extended and improved by Sheth and Tormen (1999), Sheth et al. (2001), Tinker et al. (2008) and Despali et al. (2016).

Subhalo mass function: the number density of haloes that have been accreted onto larger ones is also a power-law with a normalisation that depends on the host halo mass and redshift (Giocoli et al., 2008; Despali and Vegetti, 2017). Due to the interaction between the subhaloes and the host, the former are tidally stripped and sometimes destroyed. As a consequence of this process, the total number of subhaloes in hydrodynamical simulations is reduced by a factor between 20 and 50 per cent, depending on the galaxy formation model, relative to dark-matter-only simulations (Sawala et al., 2017; Despali and Vegetti, 2017). The number density of subhaloes also changes as a function of distance from the halo centre according to an Einasto profile. However, at



Fig. 1 Power spectra (left) and mass functions (right) in warm and self-interacting dark matter models. In the left panel, the CDM P(k) is shown in black, while different colours represent four WDM models and one SIDM model (ETHOS, see Vogelsberger et al., 2016) in which the power-spectrum is modified. In the WDM case, the suppression is shown both for sterile neutrino models (solid) and their thermal-relic closest counterpart (dashed). The right panel shows instead the suppression in the halo mass function with respect to CDM, calculated with the extended Press-Schechter (EPS) formalism, or measured in simulations (Lovell et al., 2014; Despali et al., 2018). The figures are adapted from Figure 1 and 2 in Lovell (2020), respectively.

fixed subhalo mass, the projected number density distribution is constant with radius (Xu et al., 2015; Despali and Vegetti, 2017).

Halo mean mass density profile: the mass density $\rho(r)$ of a dark matter halo as a function of radius r is described by the Navarro, Frenk and White profile (NFW, Navarro et al., 1996):

$$\rho(r) = \frac{\rho_s}{\frac{r}{r_s} \left(1 + \frac{r}{r_s}\right)^2} \,. \tag{2}$$

Here, r_s is the scale radius and ρ_s is the density normalization. The NFW profile can also be defined in terms of the halo virial mass $M_{\rm vir}$ (i.e. the mass within the radius, $r_{\rm vir}$, that encloses a virial overdensity $\Delta_{\rm vir}$, defined following Bryan and Norman, 1998), and the virial concentration $c_{\rm vir} = r_{\rm vir}/r_s$.

Subhalo mean mass density profile: due to the tidal interactions between the subhaloes and the host halo, the mass density profiles of the former tend to deviate from a standard NFW profile and are significantly more concentrated than isolated haloes of the same mass (e.g. Moliné et al., 2017). For this reason, their properties are better described in terms of the peak circular velocity V_{max} and the corresponding R_{max} radius.

In Section 5.5, we discuss current uncertainties relative to the subhalo mass function and mass density profile, and how these affect the robustness of dark matter constraints from strong gravitational lensing observations.

Alternative dark matter models can affect both the number and the structural properties of haloes and subhaloes and, as a consequence, the lensing signal that they produce. In the rest of this section, we describe the most studied ones and how they differs from CDM.

2.2 Warm Dark Matter

Dark matter particles with (close to) relativistic free-streaming velocities in the early Universe, for example, light neutrinos, are commonly described as Hot Dark Matter candidates (Doroshkevich et al., 1981). A universe predominantly made of HDM has already been ruled out by observations of the clustering of galaxies on large scales (White et al., 1983). Between CDM and HDM lies a class of dark matter models known as warm dark matter (WDM), whose candidates include the gravitino and sterile neutrinos (e.g. Boyarsky et al., 2009; Boyarsky et al., 2019). They are non-relativistic, but have a non-negligibile free-streaming velocity at early times. This property leads to the suppression of the powerspectrum of the mass-density fluctuations on scales smaller than the half-mode scale $\lambda_{\rm hm}$ (see Fig. 1). The corresponding suppression in the number density of low-mass haloes relative to CDM can be expressed as follows (Schneider et al., 2012; Lovell, 2020):

$$\frac{n_{\rm WDM}}{n_{\rm CDM}} = \left(1 + \gamma \frac{M_{\rm hm}}{M}\right)^{\beta} . \tag{3}$$

 $M_{\rm hm}$ is the *half-mode mass* and is defined as the mass-scale at which the square root of the WDM linear matter power-spectrum is 50 per cent smaller than in CDM. In practice, the half-mode mass is inversely proportional to the dark matter particle mass m_{DM} : the lighter the particle candidate, the stronger the suppression in the mass function at small scales.

As a result of the increased particle velocity in WDM, structure formation is delayed to a time when the Universe is less dense. As a consequence, haloes of a given mass are not only less numerous, but also less concentrated (e.g. Ludlow et al., 2016). By suppressing both halo abundance and halo concentration, the latter quantity determining the lensing efficiency of haloes, WDM models predict less smallscale perturbations to the strong gravitationally lensed images than one would expect from CDM.

2.3 Self-interacting Dark Matter

Self-interacting dark matter (SIDM) postulates that dark matter particles are not collisionless, but have non-gravitational interactions in which they exchange energy and momentum. The term SIDM refers to a variety of models that include elastic or inelastic scattering and a constant or velocity dependent interaction cross-section (Vogelsberger et al., 2012; Rocha et al., 2013; Kaplinghat et al., 2014; Vogelsberger et al., 2016; Sameie et al., 2018; Robertson et al., 2018; Lovell et al., 2019; Kaplinghat et al., 2019; Vogelsberger et al., 2019; Despali et al., 2019; Sameie et al., 2020; Kaplinghat et al., 2020; Robertson et al., 2021; Correa, 2021).

The distinctive signature of SIDM is a modification of the central mass density profile of haloes and subhaloes. The interactions between dark matter particles in high-density regions lead to a transfer of heat and the formation of a density core with a depth and size that is related to the strength of the self-interaction cross-section σ . In some cases, however, the halo and subhalo subsequently undergo a runaway contraction, also known as gravo-thermal catastrophe or core collapse. This phenomenon results in the formation of a cuspy density profile. It is accelerated by the presence of baryons (e.g. Feng et al., 2021) and, in the case of subhaloes, by tidal stripping (Nishikawa et al., 2020). From a strong gravitational lensing perspective, these diverse changes to the central mass density distribution imply that certain (sub-)haloes will be more efficient lenses than others (Despali et al., 2019; Robertson et al., 2019; Gilman et al., 2021, 2022).

In addition to altering the internal structure of haloes, SIDM can also suppress the abundance of subhaloes relative to CDM through ram-pressure stripping of the subhaloes while they are accreted by the host galaxy (see 1) and Nadler et al., 2020; Zeng et al., 2022). As ram-pressure stripping occurs due to self-interactions between dark matter particles bound to the subhaloes and those bound to the host, the efficiency of this mechanism depends on the amplitude of the self-interaction cross-section. For example, in velocity-independent SIDM models with elastic scattering, a very high cross-section ($\sigma \sim 10$) is required. These models are currently ruled out by observations of galaxy clusters (Sagunski et al., 2021; Andrade et al., 2022). Strong gravitational lensing, being sensitive to the amount and concentration of low-mass haloes (i.e. with a central velocity dispersion from less than $\sim 10 \text{ km s}^{-1}$ to $\sim 50 \text{ km s}^{-1}$), provides, therefore, an independent avenue to constrain SIDM models. It allows one to constrain the self-interaction crosssection at low velocities and complements the constraints derived from galaxies and galaxy clusters (e.g. Loudas et al., 2022).

2.4 Fuzzy Dark Matter

Fuzzy dark matter is a particular form of dark matter made of *ultra-light* bosons, i.e., $m_{DM}c^2 \sim 10^{-22}$ eV. This particle mass is orders of magnitude smaller than that of WIMPs and WDM models. As a result, the de Broglie wavelength is larger than the inter-particle separation and waves better describe the behaviour of the FDM field. This effect leads to a series of distinctive phenomenologies with respect to the other dark matter models so far considered (see Hui, 2021, for a review). For example, numerical simulations, which model the full non-linear evolution (Schive et al., 2014a; May and Springel, 2021), show that the wave-like behaviour (e.g. interference effects) of FDM leads to the formation of a soliton core at the centre of haloes and density granules on scales smaller than a kpc.

Similarly to WDM, there is a cut-off in the FDM transfer function at small scales, though via a different mechanism that is dependent on the de Broglie wavelength rather than a free-streaming length¹. As a consequence, from a strong gravitational lensing perspective, FDM models are expected to lead to fewer perturbations of the lensed images compared to CDM. However, the granular structure in the halo density profile of lens galaxies has been shown to lead to a distinct new source of small-scale perturbations of the lensed images, which is unique to FDM models (see Fig. 2 and Chan et al., 2020; Laroche et al., 2022; Powell et al., 2023; Amruth et al., 2023).

3 Strong lensing as a probe of dark matter

Changes to the matter distribution on subgalactic scales within lens galaxies and along their line of sight leave a gravitational imprint on the strong lensing observable. Here, we briefly describe the nature and strength of this effect. In particular, we discuss the changes that low-mass haloes induce on the lensing potential, its first and second derivative, and how they affect the observed time-delay, image positions

¹ Note, however, that the wave-like nature of ultra-light dark matter particles results in the FDM power-spectrum to briefly exceeds that in CDM on O(kpc) scales that correspond to the mean de Broglie wavelength.



Fig. 2 Lensed image multiplicity in FDM. The left panel shows the caustic structure of a galaxy-scale lens in FDM. The colour indicates the number of lensed images for a point source located at different positions on the source plane. The right panel shows the corresponding lensed image position on the observed plane. Rare hexad and octad images become more likely in FDM models. The figure is taken from Chan et al. (2020).

and magnifications, respectively. For an historical perspective of the field, we refer the reader to Section 6.

Image magnification: The largest effect produced by local fluctuations in the lensing mass density distribution (either in the form of subhaloes, field haloes or FDM granules) is a change in the relative magnification of unresolved lensed images (see Fig. 3 for an example), and the creation of so called flux-ratio anomalies. These anomalies could also be related to micro-lensing by stars (see Chapter 5) or propagation effects such as free-free absorption at long wavelength (Mittal et al., 2007) and dust extinction (Elíasdóttir et al., 2006) in the lens galaxy. The observed ratios are also affected by intrinsic and extrinsic variability of the background source (Koopmans et al., 2003; Biggs, 2023). As such, they can only be used as a probe of dark matter with observations at wavelengths where the angular size of the lensed object is larger than the scale of the micro-lenses, and at which dust and free electron absorption is low. Since flux-ratio anomalies are related to a local change of the second derivative of the lensing potential, they are an effective tool to detect some of the lowest mass perturbations. Lenses with small opening angles in the fold and cusp configurations (i.e. where the source lies on the cusp or fold of the caustic curves) are the most sensitive to the perturbations of the lensing potential and least sensitive to intrinsic variability. In these cases, the following R_{fold} and R_{cusp} relations can be used to quantify the strengths of the anomaly:

$$R_{\text{fold}} = \frac{\mu_A + \mu_B}{|\mu_A| + |\mu_B|}, \qquad (4)$$

and

$$R_{\rm cusp} = \frac{\mu_B + \mu_A + \mu_c}{|\mu_A| + |\mu_B| + |\mu_C|} \,. \tag{5}$$

Here, $\mu_{A,B}$ in R_{fold} and $\mu_{A,B,C}$ in R_{cusp} are the magnifications of the merging double and triplet images, respectively. As the opening angles between the images A and C ($\Delta \phi$) in the cusp configuration and A and B (ϕ_1) in the fold configuration approach zero, so do R_{fold} and R_{cusp} , when the lens mass distribution is smooth and the background source is a point-source. However, it is worth noting that astrophysical sources of emission, whether they be the accretion disk, the narrow emission line region or the warm (> 50 K) dust torus around a black hole or the relativistic jet that they produce are not point-like, but have some angular scale.

Image positions: The second largest effect is a local change of the relative positions of the multiple lensed images and the creation of so called astrometric perturbations. These anomalies are related to a local change of the first derivative of the lensing potential. Therefore, they can only be produced by a gravitational perturbation and cannot be caused by micro-lensing nor dust. In the case of extended sources, astrometric anomalies appear as perturbations to the surface brightness distribution of highly magnified arcs and Einstein rings, and are sometimes referred to as surface-brightness anomalies (see Fig. 3 for an example). The level of perturbations that can be detected using a surface-brightness anomaly is set by the quality of the data and structure of the source surface brightness distribution (see Section 7 and Despali et al., 2022). For example, a subhalo with a mass of $10^6 M_{\odot}$ will produce distortions on angular scales of a couple of milli-arcsecond.

Image time delay: The weakest effect is a change of the lensing potential itself. In the case of multiply imaged quasars with a flux-varying source, this phenomenon is observed as a perturbation to the time delay (see Chapter 7) between the multiple images (Keeton and Moustakas, 2009; Cyr-Racine et al., 2016; Gilman et al., 2020). Unlike flux-ratio anomalies, time-delay anomalies are not affected by dust extinction in the lens galaxy. However, they suffer from stellar microlensing which induces time-delay changes of the order of days (Tie and Kochanek, 2018). Time-delay anomalies due to subhaloes are typically of the order of a fraction of a day. At present, this is smaller than the typical time-delay uncertainty (equal or larger than a day, e.g., Fassnacht et al., 2002). To be an effective probe of the nature of dark matter, time-delay anomalies require, therefore, sensitive measurements with a high observing cadence.

Image polarisation angle: The presence of a field of Axion-Like Particles (ALPs) and its interaction with photons leads to the so-called phenomenon of birefringence whereby the polarisation angle of a linearly polarised source gets rotated. If the field oscillates over time, so does the change in the polarisation angle in a way that is related to the mass of the particles. Due to gravitational time delay, the multiple images of strongly lensed sources experience a different level of rotation, leading to differential birefringence, which can be used to probe the particle mass and it coupling with photons (Basu et al., 2021).

4 Lens modelling

The process of constraining the properties of dark matter with strong gravitational lensing is best understood as a (hierarchical) Bayesian inference problem with the following unknowns: the intrinsic properties of the source \vec{s} , the parameters of the main lens(es) mass distribution $\vec{\eta}$, and the amount and properties of low-mass haloes or fuzzy dark matter granules $\vec{\eta}_{pert}$. These have to be simultaneously inferred from the observed data \vec{d} to which they are related in a statistical sense via the following posterior distribution:

$$P(\vec{s}, \vec{\eta}, \vec{\eta}_{\text{pert}} | \vec{d}) = \frac{P(\vec{d} | \vec{s}, \vec{\eta}, \vec{\eta}_{\text{pert}}) P(\vec{s}, \vec{\eta}, \vec{\eta}_{\text{pert}})}{P(\vec{d} | M)} \,. \tag{6}$$

In the numerator, the first term is the Likelihood function and the second is the prior on the parameters of the model M. The term in the denominator is the marginalised Likelihood (also known as Bayesian evidence). A detailed description of each ingredient is provided in the following sections. The equation above is an exact expression of the inference problem at hand, and it encodes how the different components are related to each other. However, depending on the modelling approach (see Section 4.4), one may not make direct use of this posterior probability and instead adopt simplifying assumptions to make the inference problem more tractable.

4.1 The Likelihood

The Likelihood function returns the probability of observing the data, given a choice of model. In the context of strongly lensed unresolved sources, the data \vec{d} is the set of image fluxes and positions. For extended sources, the data is the surface brightness distribution in each image pixel in optical observations and a set of visibilities for interferometric ones. In the case of radio polarised emission, the data is given by the visibilities of the coherency vector. In studies that combine multiple data-sets of the same lens system at different wavelengths, the data can be thought of as a concatenation of the different observations, which originate from the same lensing potential, but with a different surface brightness distribution of the source. In many cases, the Likelihood function is well approximated by a Gaussian distribution with uncorrelated noise².

4.2 The source

Sources in gravitational lens systems can be of a large variety. In the following, we simply classify them into two categories: compact and extended sources. AGN and QSOs are examples of compact sources. Extended sources include galaxies and radio jets.

Compact sources: Traditionally, for gravitationally lensed sources that appear unresolved, either a point or a Gaussian source is assumed. Free parameters of the model are therefore the source position and flux, and in the latter case also the source size. Both assumptions can introduce systematic errors in the dark matter inference that are discussed in more detail below.

Extended sources: Typically, one of the following methods is used to describe the surface brightness distribution on the source plane: an analytical (e.g. a Sérsic) profile, a pixellated model, a basis function regression (e.g. starlets and shapelets) or a deep generative model. Here, we briefly discuss the main strengths and weaknesses of each approach.

The advantage of analytical models is that the number of free parameters is small, and the modelling procedure is fast. The main drawback is that such simplistic sources are an unlikely representation of lensed star-forming galaxies, which are often clumpy and irregular (e.g. Ritondale et al., 2019b).

Pixellated sources have enough freedom to fit complicated light distributions well. However, as the lensing problem is poorely constrained, they require a regularising prior (e.g. Suyu et al., 2006; Birrer et al., 2015; Vernardos and Koopmans, 2022; Galan et al., 2022). The choice of prior

² However, optical images may have correlated noise as a result of the pixel re-sampling using, for example, the drizzle image processing.



Fig. 3 Two examples of galaxy-scale lenses with a compact (left) and extended (right) background source. On the left panel, the gravitational lens system B2045+265 shows a strong flux-ratio anomaly which is inconsistent with a smooth lens mass distribution (Fassnacht et al., 1999). On the right, the gravitational lens system SDSSJ120602.09+514229.5, also known as the Clone, displays a distorted arc due to the presence of a small luminous satellite galaxy (Vegetti et al., 2010).

is non-trivial and may not necessarily be physically motivated. A possible solution is the introduction of hyper-priors (Rizzo et al., 2018; Vernardos and Koopmans, 2022), which have the advantage of retaining the freedom of a pixellated source while imposing physically-meaningful constraints. Another limitation of free-form models is that they are challenging from a computational perspective, especially for highresolution interferometric data (Hezaveh et al., 2016; Powell et al., 2021). One more aspect to consider is that these models require constructing a regular or adaptive grid on the source plane. We refer the reader to Tagore and Keeton (2014) and Nightingale and Dye (2015) for a detailed discussion of biases related to different choices of source discretisation and regularisation schemes.

Recently, machine learning techniques have been introduced as a model for source galaxies (e.g. Chianese et al., 2020; Adam et al., 2022; Karchev et al., 2022). These approaches can overcome some of the above limitations. However, their performance is sensitive to the choice of training data. It is currently unclear how to generate large training samples of realistic lensed galaxies at high angular resolution. It is, for example, unlikely that nearby galaxies are a good description of the high-redshift population. Similarly, lensed and unlensed galaxies at the same redshift are not observed on the same angular scales. Holzschuh et al. (2022), have shown how generative models can be used to create arbitrarily large samples of source galaxies from hydrodynamical simulations. However, it is unclear how well these simulated galaxies represent the population of lensed sources. As for pixellated sources, deep learning techniques also suffer from biases related to the data discretisation.

Independently of the chosen model, a certain level of degeneracy exists between structures in the source and structures in the lensing potential. Therefore, any assumption on the source light has important consequences for the inference of dark matter with strong gravitational lensing. We provide a more detailed discussion of this issue in Section 5.

4.3 The lensing potential

In this chapter, we focus our attention on galaxy-scale lenses. While galaxy clusters could also provide constraints on smallscale fluctuations in the dark matter distribution, most studies so far have focused on smaller scale deflectors. This choice is related to the challenge of modelling the complicated mass distribution of galaxy clusters with enough precision.

4.3.1 Main lens

For galaxy scale lenses, the main deflector is typically a single galaxy, most commonly but not limited to a massive early-type. For many years, the most common parametrization for its mass density distribution was a single power law (SPL) or a singular isothermal (SIE) elliptical profile, plus external shear. This choice was motivated by the analysis of relatively large samples of lens galaxies (e.g. Koopmans et al., 2009; Barnabè et al., 2011). However, it is worth noting that no existing sample of strong lenses is representative of the actual distribution of observable lenses in the Universe.

An important question is the level of complexity in the mass distribution of lens galaxies and how it affects the con-

straints on dark matter. For example, deep Near-Infrared (NIR) observations have revealed the presence of a disk component in two lenses that were previously assumed to be purely elliptical. It was also shown that these disks have a nonnegligible lensing effect and account for most of the observed flux ratio anomaly (Hsueh et al., 2016, 2017). Similarly, Spingola et al. (2018) and Powell et al. (2022) have shown with VLBI observations that an SPL is a good description of the lens mass distribution of MG J0751+2716 only down to scales of a few milli-arcseconds. Below these scales more complex angular and radial structures become important. In light of these and other results (see Section 5), more recent analyses have included the effect of multipole mass moments, such as boxiness and diskyness, as well as that of nearby satellite galaxies. The degeneracy between different forms of complexity in the lensing potential (e.g. subhaloes versus disks or multipoles) is a source of systematic error in the inference of dark matter. We discuss this problem and possible solutions in Section 5.

4.3.2 Low-mass haloes

In the following, we refer to subhaloes and field haloes collectively as low-mass haloes. The parameters describing this population are: the number of objects, their masses, positions, redshifts and mass density profiles. Different assumptions regarding these quantities can be found in the literature. Here, we provide a description of those that are most frequently used.

Analytical: most commonly, low-mass haloes are modelled as spherical systems with a Pseudo-Jaffe (PJ, e.g. Dalal and Kochanek, 2002) or a (truncated) NFW (e.g. Gilman et al., 2019; Hsueh et al., 2020) profile. The mass-concentration relation of the NFW depends on the dark matter properties and is either taken from numerical simulations (e.g. Gilman et al., 2019) or is a free parameter of the model (e.g. Gilman et al., 2020a). In fully forward models, the number of objects as a function of their redshift and mass are drawn from a Poisson distribution with an expectation value derived from the halo and subhalo mass functions. The latter are set by the dark matter model and are taken from numerical simulations (see Section 2). On each redshift plane, the projected positions of low-mass haloes are generally assumed to be uniformly distributed, as motivated by numerical simulations (Xu et al., 2015).

Pixellated: low-mass haloes are described as linear local corrections to the lensing potential (Koopmans, 2005; Vegetti and Koopmans, 2009). Individually detected objects are identified as positive convergence corrections of the otherwise smooth lens mass distribution. As for pixellated sources, a regularising prior for the potential corrections must be defined (e.g. Vernardos and Koopmans, 2022). Due to its freeform nature, this approach does not make a priori assumptions on the number and mass density profile of the lowmass haloes. Typically, the connection to specific dark matter models is done a posteriori as described in Section 4.4. Gaussian Random Field: low-mass haloes are represented by a Gaussian Random Field (GRF, e.g. Hezaveh et al., 2016; Diaz Rivero et al., 2018; Díaz Rivero et al., 2018; Chatterjee and Koopmans, 2018; Cyr-Racine et al., 2019; Bayer et al., 2023a,b). The corresponding power-spectrum, which is by construction well represented by a power-law, then carries information on the low-mass haloes abundance, mass function and density profile. Hence, these quantities do not have to be assumed a priori. As the assumption of Gaussianity only holds for the very low mass haloes that appear in great number, the more rare and massive objects have to be individually detected and separately treated. The connection to the properties of dark matter is done a posteriori and in terms of the power-spectrum (see Section 4.4).

4.3.3 Fuzzy Dark matter granules and subhaloes

In FDM cosmologies, the small-scale structure of galaxyscale haloes and their subhalo populations are markedly different from their CDM or WDM analogues. The main difference is that the main dark matter halo exhibits wave intereference on ~kpc scales due to the ultra-low mass of the dark matter particle; this gives rise to O(1) fluctuations in the halo density, which are commonly termed "granules". FDM haloes that are self-consistent with regard to the governing Schrödinger-Poisson equations can be obtained via numerical simulation (Schive et al., 2014b; May and Springel, 2022) or direct construction of wave-function eigen-modes (Yavetz et al., 2022). However, for the practical purpose of gravitational lens modeling, a faster analytic prescription is often preferred. To this end, Chan et al. (2020) and Kawai et al. (2022) derive statistical properties of FDM granules that can be used to quickly generate perturbations to a smooth lensing potential, which are consistent with an FDM halo. Laroche et al. (2022) implement this approach by randomly placing a large population of Gaussian density profiles in the lens, while Powell et al. (2023) use a Fourier-space approach to achieve a similar result.

4.4 Modelling approaches

4.4.1 Semi-linear

Warren and Dye (2003) presented a lens modelling approach for the analysis of data with an extended source in which the latter is pixellated (see Section 4.2). The most probable a posteriori (MAP) source and main lens parameters are inferred from the posterior probability in Eq. (6). In the case of quadratic prior distributions, the MAP source is obtained by solving a linear system (hence the term semi-linear - see Birrer et al., 2017; Galan et al., 2022, for examples of a semilinear techinique with a basis function regression model for the source). The Bayesian evidence (denominator in Eq. 6) is used to compare different models for the lensing potential and different choices of regularizations.

Koopmans (2005) introduced the so-called gravitational imaging technique for the detection of low-mass haloes with galaxy-galaxy lensing. In this approach, the source is pixellated and the low-mass haloes are described as linear, pixellated corrections to the analytical main lensing potential (see Section 4.3.2). The methodology was fully embedded in the framework of Bayesian statistics with an adaptive source by Vegetti and Koopmans (2009) and extended to the 3D (one frequency and two spatial dimensions) and interferometric domain respectively by Rizzo et al. (2018) and Powell et al. (2021). Recently, Vernardos and Koopmans (2022) have further extended the original method by Koopmans (2005) to include physically motivated priors. Due to the pixellated nature of the potential corrections, the gravitational imaging methodology does not require any assumption on the number, mass and position of low-mass haloes. Indeed, Vegetti and Koopmans (2009) have explicitly shown that more than one subhalo can be identified (provided that they have an effect on the lensed images), and Dhanasingham et al. (2022) have introduced a formalism to differentiate between subhaloes and field haloes based on the two-point function of the effective deflection angle field. Moreover, the potential corrections are not limited to capturing the effect of lowmass haloes. As shown by Barnabè et al. (2009), Ritondale et al. (2019a) and Galan et al. (2022) they can be used to identify components in the lensing potential that are not captured by the main parametric lens model. Indeed, the freedom allowed to the lensing potential is one of the main advantages of this approach as one can directly identify and differentiate different forms of complexity. One disadvantage is that the method is not fully forward, and two more steps are required to derive constraints on the properties of the dark matter: assessing the statistical relevance of detections and non-detections, and the interpretation of these in the context of theoretical predictions.

Detections: Vegetti et al. (2010), Vegetti et al. (2012) and Ritondale et al. (2019a) have introduced the following criteria to define the detection of a low-mass halo as statistically robust. (i) The mass and position of the pixellated convergence corrections have to be consistent with those inferred from an analytical description of the low-mass halo. The latter is inferred from the posterior distribution with a non-informative prior on the object mass and position, and a given choice of mass density profile. (ii) The model that includes low-mass haloes is preferred over the smooth one with a Bayes factor of at least 50. Under the assumption of statistical Gaussian errors, this difference in Bayesian evi9

dence corresponds roughly to a $\geq 10-\sigma$ detection. While this may sound overly conservative, this choice is made due to the presence of unaccounted-for systematic errors that can result in false positives at lower significance levels. For example, Ritondale et al. (2019a) have found that the rate of false positives changes from 40 per cent with a 4- σ level threshold to 20 per cent at the 5- σ level. At the same time, the rate of false negatives decreases from 60 per cent with a 4- σ level threshold to 30 per cent at 6σ . Recently, this issue was more systematically quantified by Nightingale et al. (2022). We discuss their work in more detail in Section 5. It should be noted that false detections are not limited to this lens modelling approach. They are intrinsic to the general problem of detecting low-mass haloes with lensing and their degeneracy with other aspects of the lens inference problem. If anything, requiring that the free-form and the analytical model are consistent with each other can significantly mitigate some of these issues and reduce the incidence of false positives.

Non detections: Vegetti et al. (2014) and Ritondale et al. (2019a) quantify the statistical relevance of the non-detections using the so called sensitivity function. For each pixel on the image plane the sensitivity function returns the lowest subhalo mass that could have been detected with a Bayes factor of 50. One of the main draw-backs of the sensitivity function is that, depending on the size of the data, it can be computationally expensive to evaluate. For each pixel on the image plane one has to calculate the Bayesian evidence of the model with a subhalo of a given mass. Hence marginalizing over redshift, number, mass and mass density profile could be practically unfeasible. O'Riordan et al. (2023) have recently shown that in principle one can successfully overcome these computational limitations by calculating the sensitivity function with machine-learning approaches. In Section 7 we will discuss in more detail what sets the sensitivity of a given data set to the presence of low-mass haloes of a given mass.

Dark matter constraints: From the sensitivity function, one can then interpret detections and non-detections within a given dark matter model. This is obtained by calculating the posterior probability of the dark matter particle mass assuming that the number of low-mass haloes has a Poisson distribution with expectation value given by the halo and subhalo mass function from that specific dark matter model (Vegetti et al., 2014, 2018; Ritondale et al., 2019a; Enzi et al., 2021).

Hezaveh et al. (2016) have introduced a new formalism in which the presence of subhaloes in the lens galaxy is modelled as a GRF (see Section 4.3.2). From the lensing observations, one constrains the power spectrum of projected density fluctuations, which amplitude and shape can then be a-posteriori compared to predictions from different dark matter models. The advantage of this approach is that no a priori assumption is made on the properties of the lowmass haloes as these can be inferred from the analysis itself. Indeed Diaz Rivero et al. (2018) have shown that the amplitude and shape of the power-spectrum are sensitive to the abundance, mass density profile, and concentration of subhaloes. The main disadvantage is that the assumption of a GRF only holds for the lowest-mass objects $M < 5 \times 10^7 M_{\odot}$ (Hezaveh et al., 2016). Hence, the larger ones have to be first individually identified and then explicitly included in the mass model. Moreover, in its current implementation, this approach does not allow one to identify other forms of complexity in the lens mass distribution (these are unlikely to be well described by a GRF), which then introduce a systematic bias on the dark matter inference (e.g. Bayer et al., 2023b).

4.4.2 ABC

Approximate Bayesian Computation (ABC, Rubin, 1984; Diggle and Gratton, 2018; Tavaré et al., 1997; Turner and Van Zandt, 2012; Liepe et al., 2014) is an algorithm rooted in Bayesian statistics to circumvent the direct calculation of intractable Likelihood functions. As such, it enables inference analysis based on simulated data sets computed in a forward fashion. ABC methods follow a common general process: (i) emulate the data many times with different underlying target parameters as well as noise realizations; (ii) compress the difference between the simulation and the data in a set of summary statistics to provide a metric distance between the simulation and the data; (iii) accept the proposed simulations and their underlying parameters if the distance metric is within a certain threshold ϵ and (iv) the accepted samples can be interpreted as posterior distributions with the prior being the draws of the simulator. If the acceptance criteria converges to the identical matching of the simulation and data, $\epsilon \to 0$, the accepted sample is identical to the posterior from the exact Likelihood expression applied on the summary statistic.

An accurate ABC inference is made of three main components. The first one is accurate simulations including all relevant aspects affecting the data (or more specifically the summary statistic). The second ingredient is a summary statistic that captures significant aspects of the signal of interest. Finally, one needs a sufficient number of simulations such that a narrow acceptance criteria (small ϵ) can be chosen that leads to convergence and an accurate posterior prediction. One of the main advantages of an ABC approach is that, in contrast to some machine-learning-based methods (see Section 4.4.4 for more details), the summary statistic is explicit and is a means of understanding the impact of possible sources of systematic errors. However, the choice of summary statistic is arbitrary, potentially leading to a significant loss of information and constraining power. Specifically, the choice of summary statistic and level of data compression determine the number of simulations needed for the ABC process to converge, potentially resulting in a loss of computational efficiency relative to machine learning approaches.

In the context of inferring the properties of dark matter with strong gravitational lensing, Birrer et al. (2017) were the first to propose an ABC method. In particular, they made use of an elaborate summary statistics consisting of a substructure scanning approach and the map of relative Likelihood (Section 4.1) values in the reconstruction with and without a substructure. This scanning mechanism was meant to filter signal and not absorb other spurious effects between the simulations and the data. The methodology was designed for strong gravitational lens systems with an extended source, which was modelled with a basis function regression method in the form of shapelets. We refer the reader to Enzi et al. (2020), He et al. (2023), Bayer et al. (2023a) and Bayer et al. (2023b) for other examples of summary statistics commonly used when modelling strongly lensed galaxies. Gilman et al. (2018), Gilman et al. (2019) and Gilman et al. (2020b) have made use of an ABC approach to analyse the fluxes and positions of strongly lensed quasars. In this case, as the size of the data \vec{d} was small, no compression was required and the summary statistics was set to the difference in the flux ratio of the multiple images between the simulation and the data.

As the ABC is a forward method, subhaloes and field haloes are typically described by analytical mass profiles and have properties statistically drawn as described in Section 4.3.2.

4.4.3 Trans-dimensional

One of the challenges in constraining the properties of dark matter with strong gravitational lensing is that the number of low-mass haloes in any given lens system is unknown. As a consequence, the model describing the low-mass halo population and its properties needs to have a variable and a-priori unknown number of free parameters. An additional challenge is that most of the low-mass haloes will be at or below the detection limit. Brewer et al. (2016) and Daylan et al. (2018) have proposed the use of trans-dimensional Bayesian inference approaches (Green, 1995, 2003) to overcome these problems.

These methods apply probabilistic cataloging to images of strongly lensed systems (Hogg and Lang, 2010; Brewer et al., 2013, 2016; Daylan et al., 2017; Portillo et al., 2017) and output an ensemble of probability-weighted (sub-)halo catalogs providing a good fit to data. The prior distribution for the properties of the low-mass haloes is specified hierarchically, so that their mass function is a natural output of the method. Sampling of the posterior can be done with reversible jump Markov Chain Monte Carlo with differences in the explicit sampling methods, such as Diffusive Nested Sampling (DSN, Brewer et al., 2016) or PCAT (Daylan et al., 2018). One can also evaluate the marginal Likelihood of the model, including over the unknown number of low-mass haloes, and the source and lens properties. In most applications, both the source and the lens are described by analytical models. However, there is in principle no limitation to couple a trans-dimensional treatment of low-mass haloes with, for example, pixellated sources.

4.4.4 Machine Learning

In recent years, it has been shown that neural networks can efficiently estimate the parameters of strong lens models directly from observations (Hezaveh et al., 2017; Perreault Levasseur et al., 2017; Schuldt et al., 2021). Motivated by these results, a number of works have explored machine learning methods for identifying subhaloes in mock observations (Ostdiek et al., 2020; Yao-Yu Lin et al., 2020; Diaz Rivero and Dvorkin, 2020). Others have shown that subhalo summary statistics can be directly estimated from mock observations (Brehmer et al., 2019; Alexander et al., 2021; Varma et al., 2020; Alexander et al., 2021; Vattis et al., 2022; Ostdiek et al., 2022). However, these works do not provide any rigorous uncertainty estimates.

To address this issue, Brehmer et al. (2019) proposed an approach relying on Neural Ratio Estimators (NREs). NREs transform the problem of inferring the value of a continuous variable into a classification problem between two sets. Using this, it is possible to calculate the posterior of a parameter as the ratio of the Likelihood, multiplied by the prior, and the evidence. This allows for better uncertainty estimates. However, the approach of Brehmer et al. (2019) is extremely data-hungry, making its scaling to realistic data non-trivial. A number of variants of NREs have been proposed to solve this problem (Cranmer et al., 2020; Coogan et al., 2020; Montel et al., 2022).

Uncertainties on the subhalo population can also be obtained similarly to those of the macro-model as proposed by Perreault Levasseur et al. (2017). Wagner-Carena et al. (2023) showed that it is possible to infer the parameters of the subhalo mass function over a population of gravitational lenses, in a Bayesian hierarchical formalism. They trained a neural network that directly predicts a parameterized distribution approximating the posterior of the low-dimensional macro-model parameters and subhalo mass function normalization (see also Vernardos et al., 2020). The issue of how to consistently incorporate realistic sources of biases in this inference such as complex selection functions, for example, selection from a lens-finding neural networks, has been discussed by Legin et al. (2022).

Together, these works have shown that neural networks can extract low dimensional information about the density fields, that is, the parameters of the mass function or m_{WDM} .

The main limitation of these methods, however, is that they are only tractable for the inference of low-dimensional representations, implicitly marginalizing over all nuisance parameters that are not explicitly estimated. At first glance, implicit marginalization appears appealing because the marginal posterior of the dark matter model parameters is the desired outcome. However, it hinders reproducibility and confirmation of the results by alternative, more traditional approaches. Machine learning models do not predict, at least for now, the actual distribution of density in the foreground, or the surface brightness of the background source. One cannot therefore compare their predictions with the data by raytracing through the model favoured by the neural network.

If, for example, excess power is detected in the mass function, it would be difficult to identify the specific feature in the data that accounts for it. Investigating whether a neural network's preference for a given dark matter model comes from a massive subhalo or a population of low mass subhaloes would be intractable in most machine learning frameworks developed to date. The lack of explicit predictions of physical features like density or brightness distributions, makes it difficult to verify measurements of lowdimensional substructure statistics by neural networks using more traditional methods.

One of the central difficulties in modelling the physical features of strong gravitational lenses, either with traditional analysis methods or with deep learning, has been the difficulty of defining priors over such high dimensional spaces. Recent advances in denoising diffusion-based models are now making this previously-intractable problem possible (Song et al., 2020). Further, Adam et al. (2022) have shown that a neural network can learn the score of the prior over background source images, learned from unlensed highresolution images of galaxies. By adding the Likelihood score to this learnt prior score and using a reverse-time stochastic differential equation solver, the authors can obtain samples from the posterior over background source pixels. Generalizing the framework of denoising score-based models to solve highly non-linear problems such as the reconstruction of density maps in strong gravitational lenses is an active area of research.

5 Degeneracies and systematic errors

When assessing the robustness of strong gravitational lensing constraints on dark matter it is important to consider the effect of degeneracies and systematic errors. Some of these are related to the measurement at hand or unsolved theoretical questions and affect both compact and extended sources. Others are lens modelling dependent. All of them are discussed in this section.

5.1 Degeneracy with complex macro-models

For any analytical model of the main deflector, a degeneracy exists between the macro-model parameters (e.g. Einstein radius and external shear) and the presence and properties of low-mass haloes. Its strength depends on the data signal-to-noise ratio (SNR) and angular resolution, and the properties of the low-mass haloes themselves. For low-mass haloes that can be individually detected, the more concentrated their density profile and the closer to the lensed emission their position, the smaller the degeneracy with the lens properties (Despali et al., 2022).

At the same time, real galaxies are unlikely to be simple elliptical objects and are expected to have complex radial and angular structures (e.g. Spingola et al., 2018; Powell et al., 2022; Van de Vyvere et al., 2022). Unmodelled components in the lensing potential are likely to lead to an overestimation of the amount of low-mass haloes or FDM granules, resulting in a biased inference on the properties of dark matter. For example, Xu et al. (2015) have shown that the level of flux-ratio anomalies in eight strongly lensed quasars from the CLASS survey (Myers et al., 2003; Browne et al., 2003) is too large to be solely explained by low-mass haloes from CDM. They concluded that other forms of departure from a power-law macro-model may be partly contributing to the observed signal. Thanks to deep NIR observations of the CLASS gravitational lens systems B1555+375 and B0712+472, Hsueh et al. (2016) and Hsueh et al. (2017) were able to show that previously undetected stellar disks are responsible for most of the observed flux-ratio anomaly in these systems. Indeed, using mock simulated data, Gilman et al. (2017) and Hsueh et al. (2018) have reported that baryonic structures in the main lens can lead to an increase of the probability of large flux-ratio anomalies of between 10 and 20 per cent, depending on the lens galaxy morphology (see Fig. 4). Similarly, from the analysis of the the Cosmic Horseshoe lens system, Brewer et al. (2016) concluded that the inferred substructure population might have partially mimicked larger scale components not included in the primary lens model. He et al. (2023) have found that deviations from an elliptical shape that are not explicitly included in the macro-model lead to a biased characterisation of correctly identified subhaloes as well as false positive detections.

Nightingale et al. (2022) have performed a systematic study of the degeneracies between the macro-model and isolated subhaloes. From the analysis of a sample of fifty-four gravitationally lensed galaxies from the SLACS and BELLS GALLERY surveys (Bolton et al., 2008; Shu et al., 2016), they initially reported thirty-four subhalo detections at low significance. Upon further analysis, it is found that sixteen of these thirty-four are due to a degeneracy with the lens light (see Section 5.3) and five are due to insufficient source plane

resolution. The remaining thirteen false positives are macromodel dependent with eight due to an overly simple lens mass model: a subhalo is favoured because it can replicate the effect of the missing complexity. These false positives can be accounted for by including in the lens mass models more radial (e.g. the broken power-law model) and angular (e.g. multipoles) structure in two and six cases, respectively. In general, this analysis by Nightingale et al. (2022) demonstrates that no one lens is like another and more complicated mass distributions can both remove and add a subhalo candidate depending on the lens system and the type of model considered (e.g. multipoles versus composite mass distribution). The situation is further confounded by the fact that the nature of the macro-model is unknown a priori. While stellar disks may be visually identified with appropriate observations (Hsueh et al., 2016, 2017; Nierenberg et al., 2020), this is not necessarily the case for other forms of radial and angular structure. Moreover, depending on the spatial scales affected, the latter may be identified and characterized only with very-high angular-resolution data (Powell et al., 2022). These results highlight the importance of allowing for models other than a smooth SPL and modelling each lens to great detail.

While this degeneracy is intrinsic to the measurement itself and affects any type of lens modelling technique and data, it is particularly thorny for compact sources and models that solely rely on analytical macro-models³. In the first case, the data (at most, eight positions and four flux measurements) provide only limited information on the lens model. In the second case, for each lens system, one needs to marginalise over a wide range of macro-model assumptions. Extended sources, where the potential is modelled with a free-form approach (Section 4.4) have in this respect an advantage: the data has more constraining power and the freedom allowed to the potential makes it possible to detect and potentially differentiate various mass components from each other (e.g. Galan et al., 2022).

In reality, the degeneracy between the lens macro-model and dark matter fluctuations, does not act in isolation. Its net effect is the result of its interplay with another degeneracy, that between the lensing potential and the source light distribution. We discuss the latter in the following section.

5.2 Degeneracy with the source

5.2.1 Compact sources

The angular size of the background source relative to that of a low-mass halo of a given mass sets the level of perturbation

³ These also include the case where low-mass haloes are described by a GRF, as the latter is an unlikely characterisation of (baryonic) structures in the lens mass distribution



Fig. 4 Strength of the flux-ratio anomaly for a cusp (left) and fold (right) configuration arising from different mass distributions: an elliptical smooth power-law (top), an edge-on stellar disk (middle) and an early-type galaxy (bottom) selected from the Illustris simulation (Vogelsberger et al., 2014b). The curves represent 1, 5, 10, 20, and 50 per cent probabilities to find values of $|R_{fold}|$ and $|R_{cusp}|$ larger than a given value for fixed opening angles ϕ_1 (\circ) and $\Delta \phi$ (\circ). The panels are taken from Figures 5, 6 and 7 by Hsueh et al. (2017).

to the lensed images magnification. This effect introduces a degeneracy between the amount of low-mass haloes, micro-lensing by stars in the main deflector and the unknown size of the source. Micro-lensing by stars (see Chapter 5) becomes problematic when the image flux is gathered from a region surrounding the background quasar that is less than ~ 0.1 pc in diameter. It is responsible for an overestimation of the amount of low-mass haloes and hence a bias in favour of colder dark matter models. One can mitigate or even remove its effect with observations at wavelengths that are known to originate from an extended region around the source or by monitoring the system for a sufficiently large amount of time.

Provided that the source is large enough to avoid microlensing, uncertainties in the intrinsic size of the emission region translates into uncertainties in the amount of low-mass haloes. An overestimation (underestimation) of the former leads to an underestimation of the latter and a bias in favour of warmer (colder) dark matter models. Typically, one tries to account for this degeneracy by adopting observationally motivated priors (e.g. Koopmans et al., 2003; Chiba et al., 2005; Müller-Sánchez et al., 2011) and marginalising over the size of the source.

Another source of systematic error is related to flux variations in the emission region over time. Flux changes appear with a time delay between different positions on the image plane. As a result, single-epoch measurements sample the intrinsic light curve of the source at different times for the different lensed images. This effect leads to flux measurement errors that can be as large as 20 per cent (Dalal and Kochanek, 2002; Koopmans et al., 2003). It can be mitigated by monitoring and averaging the observed flux over a long period of time, or correcting for the time-delay.

At radio wavelengths, propagation effects, such as freefree absorption and scatter broadening can alter the measured properties of the different lensed images. As these effects have a strong wavelength dependence, they can be identified and corrected for with multi-wavelength observations (e.g. Winn et al. 2004; Biggs et al. 2003; Mittal et al. 2007).

5.2.2 Extended sources

The detection of low-mass haloes with extended sources is based on the idea that the many pixels on the image plane provide redundant information about the source surface brightness distribution, allowing one to separate structures in the background object (which appear two or four times in the data) and structures in the lensing potential (which produce a relatively localised effect on the lensed images). In practice, however, due to the presence of noise in the data and the smoothing effect of the telescope point spread function, a degeneracy exists.

The extent of this degeneracy is strongly dependent on how the source and lensing potential are modelled. Vernardos and Koopmans (2022) have studied in detail the case of a pixellated source and potential corrections. They concluded that the two can partly absorb each other's complex structures in a way that depends on the form of the regularisation, the pixellatisation resolution and the actual complexity in the data (see also Bayer et al., 2023a). For example, when modelling a lens system, which is well described by a smooth analytical lens mass distribution and a complex source, the potential corrections can partly absorb fluctuations in the latter that are not well captured by the grid resolution and regularisation. Similarly, the source can adapt to compensate for the lack of structure in the model for the lensing potential. Interestingly, Vernardos and Koopmans (2022) find that the degeneracy between complexity in the source and in the potential mainly results in a biased lens macro-model and source, while the statistical properties of the potential corrections are well recovered. This suggests that, at least in the free-form approach, this degeneracy should not result in biased constraints on the properties of dark matter. However, a detailed investigation of this issue as a function of data type (e.g. optical or interferometric) and quality (e.g. angular resolution, signal-to-noise ratio and uv-coverage) is still lacking.

At the other end of the spectrum, models with analytical sources and potentials are likely to result in dark-matter constraints that are biased towards models that are colder or have a lower FDM particle mass. As the only complexity allowed in the analysis is in the form of low-mass haloes or FDM granules, these are then likely to absorb structures in the source as well as in the macro-model. Identifying suitable priors for the lensing potential and source light is therefore a key ingredient to infer the properties of dark matter with strong gravitational lensing.

5.3 Degeneracy with the lens light

Optical data includes light emission from the lens galaxy. This contribution is either pre-subtracted from the data (e.g. Vegetti et al., 2010) or inferred during the lens modelling analysis (e.g. Ritondale et al., 2019b,a; Nightingale et al., 2022). Typically, it is described via a two dimensional B-spline function (e.g. Vegetti et al., 2014) or an analytical profile such as a (combination of) Sérsic model (e.g. Ritondale et al., 2019a; Nightingale et al., 2022). As for the lens mass distribution, simplistic models for the lens light may fail to reproduce complex radial and angular structures, as for example, boxy or disky isophotes and dust lanes. As a result, false positive detections of low-mass haloes may be obtained (Nightingale et al., 2022). It is therefore important to test candidate detections against different assumptions for the lens light (e.g. Vegetti et al., 2012; Nightingale

et al., 2022) or, if possible, with observations at different frequencies⁴. For example, one can use observations in two or more wavelengths and the information carried by the multiple lensed images to correct for the effect of dust in the lens galaxy (e.g., Suyu et al., 2009).

5.4 Some considerations on interferometric data

At present, radio interferometetry (e.g. Smirnov, 2011) provides the highest angular resolution available for strong gravitational lensing observations. In principle, this makes it the most sensitive probe of the halo and subhalo mass functions, with very long baseline interferometry (VLBI) arrays capable of individual halo detections down to $10^6 M_{\odot}$ (McKean et al., 2015). However, modeling radio interferometric observations of strong gravitational lenses is challenging given the very large data sizes, and is an area of active research.

It is preferable to model radio interferometric observations directly in the native visibility space (a visibility is a Fourier component of the sky brightness as measured by two antennas at a given time and frequencey interval). In principle, it is possible to model a gravitational lens observation by first imaging the data using some established deconvolution technique (e.g. CLEAN, Högbom 1974; or maximumentropy, Cornwell and Evans 1985), then applying standard lens modelling techniques to the deconvolved image. However, on the space of the CLEANed images, the noise is correlated and not well characterised, and the CLEANing process may introduce artefacts in the surface brightness distribution that mimic the effect of low-mass haloes. Sourceplane deconvolution algorithms for radio observations of gravitational lenses (such as LensClean; Kochanek and Narayan 1992; Ellithorpe et al. 1996; Wucknitz 2004) were the first to be explored as a solution to this issue. Direct χ^2 fitting in the visibility plane was then applied by several authors (Bussmann et al., 2012, 2013; Hezaveh et al., 2013) to observations of lensed dusty star-forming galaxies (DSFGs) taken with the Sub-Millimetre Array (SMA) and the Atacama Large Millimetre Array (ALMA). A fully Bayesian treatment for fitting interferometric observations, which included prior information on the source surface brightness, was finally realized by Rybak et al. (2015) and Hezaveh et al. (2016).

Amplitude and phase calibration errors (e.g. Pearson and Readhead, 1984) in interferometric data can potentially masquerade as false-positive detections of low-mass dark matter haloes. This was investigated using ALMA observations of the DSFG SDP.81 by Hezaveh et al. (2016), with a test of the data sensitivity to the subhalo detection both with and without a treatment for phase errors. Hezaveh et al. (2013) and Hezaveh et al. (2016) addressed this issue by including 15

a single antenna-based phase correction as a free parameter in the lens modeling, self-consistently incorporating uncertainties due to this systematic effect into the model posterior. It is expected that the problem of calibration errors is less relevant for cm-wavelength VLBI observations as the atmosphere is more stable over time and the antenna receivers are more sensitive. Strongly lensed sources observed with VLBI are typically radio-bright jets containing both extended and compact features, which help to provide robust phase calibration solutions prior to the lens modeling step. This is in contrast to mm-wavelength observations of DSFGs, which feature rather diffuse, low-surface-brightness emission; in this case modeling phase errors as part of the lens modeling pipeline likely yields more robust calibration solutions than a priori self-calibration. While it is clear that residual phase calibration errors have an effect on the rate of falsepositive detections and sensitivity to low-mass dark matter haloes, there is yet to be a systematic study of this effect.

The size of the data is important when it comes to modeling antenna phases directly; the ALMA observation to which this phase-correction model was applied contains a small enough number of visibilities (5×10^5 after some additional binning) that a standard linear-algebra solver framework could be applied to the source inversion step (Warren and Dye, 2003; Suyu et al., 2006). For cm-wavelength VLBI data, for which averaging can introduce additional systematics that degrades sensitivity to low-mass dark haloes, an FFT-based iterative solver is required in order to be computationally tractable (Powell et al., 2021, 2022).

5.5 Theoretical Unknowns

5.5.1 Resolution, halo and subhalo mass functions

Theoretical predictions for the abundance and structure of CDM haloes and subhaloes initially came from analytical models, such as perturbation theory (Zel'dovich, 1970), the Press-Schechter model (Press et al., 1992), the statistics of peaks in Gaussian random fields (Bardeen et al., 1986) and the excursion set approach (Bond et al., 1991; Sheth and Lemson, 1999). Predictions from these theoretical models have then been compared to the results from N-body (i.e. dark-matter-only) numerical simulations. This comparison has led to precise fitting functions for the halo mass function that take into account the non-linear evolution of structure formation (Sheth et al., 2001; Giocoli et al., 2008; Tinker et al., 2008; Despali et al., 2016). Similarly, the functional form of the NFW profile (Navarro et al., 1996) reflects the expectations for the density profile of a dark-matter-dominated structure that forms via hierarchical accretion. CDM N-body simulations agree with each other to the per cent level, and the impact of numerical effects and (sub) halo identifica-

⁴ Ideally at frequencies at which the lens does not emit any light.

tion methods has been widely studied and is well understood (Knebe et al., 2011, 2013; Okabe et al., 2013).

However, some uncertainties remain. These are mainly related to the resolution limit of the numerical simulations and the consequent fact that analytical predictions have only been tested down to a finite scale. Numerical effects are especially problematic for the subhalo population. For example, the artificial disruption of subhaloes related to the limited spatial resolution of the simulations (Green and van den Bosch, 2019; Green et al., 2021) can lead to an underestimation of the number of such objects on scales of the resolution limit of a factor of 10 to 20 per cent. Moreover, while the density profile of isolated haloes is well understood, subhaloes are affected by tidal disruption and stripping inside the main halo and thus show a larger variety of profiles (Sawala et al., 2017; Moliné et al., 2017), which are not all well described by the same functional form.

WDM N-body simulations, in which the power spectrum cutoff is resolved, are known to undergo artificial fragmentation in filaments producing spurious clumps that, close to the resolution limit, can outnumber real structures. One challenge is thus to correctly identify and remove them from the (sub)halo catalogues: this can be achieved by studying the shapes of the Lagrangian initial regions that correspond to the final structures, and eliminating those that are very elongated (Lovell et al., 2012). Alternatively, Stücker et al. (2020) have developed a method to smooth the density field using phase-sheet methods and a high-resolution force calculation, in order to completely circumvent the issue and avoid the formation of spurious subhaloes.

FDM theories predict differences in the subhalo population of a lens, with a suppression of the halo mass function at low masses due to the large de Broglie wavelength of the dark matter particle. Characterizing the low-mass halo population in FDM cosmologies is an area of active research. Analytic approximations to the FDM halo mass function in FDM have been attempted (Marsh and Silk, 2014; Kulkarni and Ostriker, 2022), but at this time only mass functions obtained from numerical simulations (Schive et al., 2016; May and Springel, 2022) have been applied in a lens modeling context (Laroche et al., 2022). The mass-concentration relation for FDM haloes is also highly uncertain; Laroche et al. (2022) modelled it using an extended Press-Schechter formalism (Schneider, 2015) assuming some correspondence between the gravitational collapse time-scales of FDM and CDM haloes. Additionally, FDM haloes contain a characteristic soliton core that alters their density profiles (Schive et al., 2014b; Chan et al., 2022). An accurate model for subhaloes around a lens galaxy is important to consider for inferences based on flux-ratio anomalies of unresolved compact sources, as subhaloes and granules can produce similar observational signatures. However, for FDM particle masses lower than ~ 5×10^{-21} eV, subhaloes are too few and too diffuse to impart a small-scale signature on the observed source morphology, and can be absorbed into a sufficiently complex macro-model. This leaves the presence of granules as the main source of constraint on the particle mass.

5.5.2 The effect of baryons

In the past few years, numerical simulations, and especially those with a CDM cosmology, have made significant progress. Large-scale structure simulations are now able to reproduce realistic galaxy morphologies and the observed scaling relations (Vogelsberger et al., 2014b; Schave et al., 2015; Pillepich et al., 2018; Dubois et al., 2021), reducing some of the tensions between CDM and observations (Brooks et al., 2013). Despite these successes, uncertainties, which may affect the interpretation of strong gravitational lensing studies within a given model, persist. Feedback processes, that cause a loss of baryonic mass, alter the total halo mass in a non-trivial way that depends on the halo mass and the galaxy formation model (Sawala et al., 2015; Despali and Vegetti, 2017; Pillepich et al., 2018; Garrison-Kimmel et al., 2019). As a result, the number density of low-mass haloes $(10^7 M_{\odot} <$ $M < 10^{10} M_{\odot}$) is suppressed in hydrodynamical simulations by a factor between 10 and 40 per cent. This suppression is smaller than the one seen in most WDM models in the same mass range, as the inclusion of baryons further suppresses the halo and subhalo mass function (Lovell et al., 2019; Despali and Vegetti, 2017). However, due to the lack of hydrodynamical simulations with high-enough mass resolution, it is at present unclear how these mass functions are affected by the baryon and dark matter physics below the probed mass limit.

Baryonic physics also affects the inner mass density profile of galaxies in a way that depends on the feedback model (e.g. Mukherjee et al., 2021) and its interplay with the dark matter model. For example, Robertson et al. (2019), Despali et al. (2019) and Shen et al. (2022) have found that, in hydrosimulations with elastic SIDM, haloes have a larger variety of density profiles (with respect to the non-elastic SIDM case) and that the shape of haloes is much closer to the CDM hydrodynamical case, than initially inferred from dark-matteronly simulations. Mocz et al. (2019) studied the interplay between baryons and fuzzy dark matter, finding that the first stars can form in filamentary structures along the cosmic web, instead of only in the collapsed haloes. These results, could have interesting implication for the analysis of strong gravitational lens galaxies. However, the limited spatial resolution available in simulations at the scale of massive galaxies prevents us from a robust comparison between observations and simulations. For example, simulated galaxies display a central density core, which is large enough to produce strongly lensed image configurations that are not observed. While these uncertainties affect simulations in all

dark models, they are currently more problematic for alternative non-CDM models. The vast majority of simulations where dark matter is different from CDM does not include baryonic physics (Lovell et al., 2012; Vogelsberger et al., 2014a) and is not yet at the level of the CDM case, in terms of resolution, number of objects and volume (Adhikari et al., 2022). Moreover, even when baryons are included, the subgrid physics processes that describe their behaviour often remain calibrated on CDM simulations. Hence, it is unclear whether non-CDM hydro-dynamical simulations provide a correct description of the interplay between dark-matter and baryons. Obtaining large samples of numerical simulations with different galaxy formation and dark matter physics, as well as the volume and resolution required for a meaningful comparison with strong gravitational lensing observations is a fundamental step for future works.

6 Historical perspective

6.1 Compact sources

The use of strong gravitational lensing as a probe of dark matter was first proposed by Mao and Schneider (1998). They showed that anomalies in the flux ratios (i.e. ratios that deviated from those predicted by smooth mass models) of lensed quasars could be accounted for by subhaloes in the lens galaxy. In particular, they focused on the two and three brightest images in a fold and cusp configuration, respectively. For a smooth lens mass distribution, the fluxes of these lensed quasar images should satisfy the asymptotic relations in Eqs. (4) and (5). Discrepancies between the observed flux ratios and the generic predictions are therefore an indication of some type of small-scale structure in the mass distribution. Mao and Schneider (1998) also pointed out that flux-ratio anomalies observed in optical data could easily be due to micro-lensing by stars in the lensing galaxy. Focusing on radio observations was, therefore, a better way to investigate the presence of subhaloes. At these frequencies, the lensed emission had a large enough angular extent as to be insensitive to micro-lensing by stars (although see Koopmans and de Bruyn, 2000). The paper investigated both compact mass distributions (globular clusters with masses of $\sim 10^{6} M_{\odot}$) and smoother fluctuations such as spiral arms in the lensing galaxy.

Mao and Schneider (1998) thus truly set the stage for subsequent investigations of perturbations by dark matter haloes by: (1) considering perturbations by larger-scale structures than stars and, (2) pointing out that the observations had to be conducted using sources that had angular scales that were large compared to the Einstein ring radii of the stars in the lensing galaxy. This second requirement ruled out using emission from the accretion disks and broad-line regions associated with the lensed AGN, which are sensitive

to micro-lensing. However, observations at radio or mid-IR wavelengths could be used to investigate dark matter since the regions in the background objects that produce emission at these wavelengths are large enough to be mostly unaffected by micro-lensing by stars in the lensing galaxy, but small enough to be sensitive to perturbation by relatively low-mass dark matter haloes. Unfortunately, it was difficult to obtain large samples at these wavelengths, since only ~ 10 per cent of AGN are radio loud, and ground-based observations at mid-IR wavelengths are extraordinarily difficult due to thermal emission from the Earth's atmosphere. Thus, for many years the sample of lenses that were useful for dark matter investigations was on the order of 10 systems, primarily discovered in the MG (Bennett et al., 1986; Langston et al., 1990; Griffith et al., 1990, 1991), JVAS (Patnaik et al., 1992; Browne et al., 1998; Wilkinson et al., 1998), and CLASS (Myers et al., 2003; Browne et al., 2003) radio surveys. Initial investigations focused on individual lenses, particularly radio-loud four-image systems that strongly violated the standard relationships for merging images (Fassnacht et al., 1999; Marlow et al., 1999; Trotter et al., 2000; Biggs et al., 2004), and included some work that incorporated numerical simulations to understand subhaloes (Bradač et al., 2002).

The seminal paper by Dalal and Kochanek (2002) was the first analysis of flux-ratio measurements in a statistically significant sample, with a goal of testing the Λ CDM model. They examined the flux ratios in a sample of seven lens systems and found broad consistency with CDM. Although this analysis had a high impact for many years, it did have several shortcomings in the context of current approaches to using flux-ratio statistics to draw inferences on the nature of dark matter. These include using CDM-only simulations, using fairly simple lens models, only considering subhaloes within the halo of the primary lensing galaxy while not including line-of-sight haloes, using PJ mass profiles for the subhaloes, and being restricted to using the somewhat uncertain flux-ratio measurements that were available at the time.

For nearly two decades following this analysis, no new lens systems that had high-sensitivity radio or mid-IR flux ratio measurements were discovered, so subsequent investigations had to focus on extending or improving the analysis rather than working with larger samples. One improvement came from a monitoring program to look for extrinsic variability in radio-loud lenses, which had the additional benefit of providing high-precision flux-ratio measurements after correcting for any time delays in the systems (Koopmans et al., 2003). Other work investigated astrometric shifts and parity dependence as methods for determining the abundance of subhaloes (Chen et al., 2007; Chen, 2009).

Another major effort addressed the problem by conducting ray-tracing analyses through high-resolution numerical simulations, which allowed the inclusion of lower subhalo masses and, thus, more thoroughly to explore the lensing effects of dark-matter subhaloes and other possible perturbers (Mao et al., 2004; Xu et al., 2009, 2010; Richardson et al., 2022). These studies incorporated not only the number of anomalies in the observed radio lenses, but also how much the observed flux ratios deviated from the predictions of smooth mass models.

Interestingly, several of these studies found that CDM subhaloes were not sufficient to explain the observations fully (Xu et al., 2009, 2015), except maybe when including lineof-sight haloes (Xu et al., 2012), suggesting that more complex macro models were needed. Earlier work had reached a related conclusion from simulated data, finding that an edgeon stellar disk could cause violations of the standard cusp relation for the lensed image magnifications (Bradač et al., 2004). Subsequent work discovered exactly this type of situation in observed lens systems. In some systems with the most extreme flux-ratio anomalies, high-resolution imaging from ground-based adaptive optics and HST data revealed edge-on disk components of the lensing galaxies. These additional baryonic components could explain the observed flux-ratio anomalies without needing to resort to either subhaloes or line-of-sight haloes (Hsueh et al., 2016, 2017). Further investigations used simulated galaxies to confirm the importance of baryonic structures in the lensing galaxies (Gilman et al., 2017; Hsueh et al., 2018).

In parallel with the work on extending and improving the analyses was an effort to increase the sample size of observed four-image lenses by pushing past the traditional radio and mid-IR observations that had provided the primary samples for the analyses above. In particular, Moustakas and Metcalf (2003) proposed a spectroscopic technique based on the differential magnification of several different emitting regions in the lensed quasar, i.e., the continuum, broad-line region, and narrow-line region (NLR). This approach allowed them to separate contributions from the smooth lens model, micro-lensing, and lensing by subhaloes. This method was applied to a single very low-redshift lens that was observed with an integral field unit (IFU) spectrograph, and seemed promising (Metcalf et al., 2004). It is difficult to apply this technique to higher redshift systems using the seeing-limited ground-based observations used in Metcalf et al. (2004), due to angular resolution issues. However, a similar approach has been successful when using either ground-based adaptive optics IFUs (Nierenberg et al., 2014) or HST grism observations (Nierenberg et al., 2017). This technique has led to the first major increase in the sample size of four-image systems with measurements that are useful for flux-ratio investigations (Nierenberg et al., 2020).

Vegetti

6.2 Extended sources

Koopmans (2005) was the first to propose the use of galaxygalaxy lensing observations to detect subhaloes within the lens using the semi-linear pixellated approach described in Section 4.4. A few years later, Vegetti and Koopmans (2009) extended the original idea to be fully embedded within the context of Bayesian statistics.

Then, Vegetti and Koopmans (2009) proposed a statistical framework for the interpretation of both detections and non detections within the context of specific theoretical predictions. They also showed how the achievable level of constraints on the subhalo mass function parameters is related to the number of lenses in the sample and the sensitivity of the data. Recently, Despali et al. (2022) have provided a detailed quantification of how the latter depends on the data properties. We discuss their findings and relative implications in Section 7.

Vegetti et al. (2010) and Vegetti et al. (2012) were the first one to apply the pixellated gravitational imaging technique to real data taken with the HST and Keck-AO. They reported a 16- σ and 12- σ detection of two individual subhaloes in the gravitational lens systems SDSSJ0946+1006 (also known as the double ring or the Jackpot) and B1938+666, respectively. These detections, obtained in a pixellated fashion, were then modelled with a PJ profile. From the latter, a total mass of ~ $10^9 M_{\odot}$ and ~ $10^8 M_{\odot}$ was inferred for each system, respectively. Both detections have been independently confirmed (Minor et al., 2021; Sengül et al., 2022). However, Despali et al. (2018) and Sengül et al. (2022) have found that the detection in the system B1938+666 is more likely a field halo. Whether the concentration of these two objects is consistent with CDM predictions is currently under investigation (e.g Minor et al., 2021; Şengül and Dvorkin, 2022). From the analysis of the ALMA long baseline campaign data for the lens system SDP.81, Hezaveh et al. (2016) reported the detection at the 5- σ level a subhalo with a total PJ mass of ~ $10^9 M_{\odot}$. Inoue et al. (2016) also found a detection in this lens system. However, the inferred subhalo position and lens macro-model are inconsistent between the two analyses. Further investigations are required to understand the origin of this discrepancy.

Focusing, for the first time, on a larger number of strong gravitational lens systems, Vegetti et al. (2014) found no additional subhaloes in a sample of ten SLACS lenses, in the mass regime probed by the data. Recently, Nightingale et al. (2022) have searched for the presence of subhaloes in fifty-four lens systems (the largest number considered so far) from the SLACS and BELLS gallery samples. They reported two candidate detections, one of which matches the one by Vegetti et al. (2010) in the Jackpot lens.

While early studies considered the subhalo population only, Li et al. (2017), Despali et al. (2018) and Amorisco

et al. (2022) have shown the contribution from field haloes to be important and in some cases dominant. This is a significant result. Unlike for subhaloes the properties and number of field haloes are better understood from a theoretical perspective (see Section 5.5), and the resulting increase in the number of detectable objects per lens allows for stronger constraints on dark matter with fewer systems. Allowing for the contributions of both populations, Ritondale et al. (2019a) concluded that their lack of detection in twenty-one lens systems from the HST BELLS gallery sample is consistent with the CDM model.

Dark-matter constraints from strongly lensed extended sources have been mainly limited by the amount of available data with enough angular resolution. In recent years, efforts have been made to increase the number of known strong gravitational lens systems (e.g. Lanusse et al., 2018; Huang et al., 2021; Petrillo et al., 2019; Cañameras et al., 2021; Rezaei et al., 2022), as well as the number of observations with improved angular resolution with, for example, Keck-AO (Lagattuta et al., 2012) in the NIR, ALMA in the sub-mm (Spilker et al., 2016) and VLBI at cm-wavelenght (Spingola et al., 2019). At same time, several new lens modelling approaches have been developed (see Section 4.4 and references therein).

Existing data-sets as well as simulated ones, have been used to improve our knowledge of systematic errors and degeneracies (see Section 5 and references therein), and our understanding of the signal under study. For example, Amorisco et al. (2022) showed that an improvement on the level of dark matter constraints can be obtained by taking into account the low-mass haloes mass-concentration relation (and how it changes with the dark-matter model) and its scatter. Similarly, ray-tracing through numerical simulations has been instrumental to quantify the lensing effect arising from smaller- and larger-scale structures (e.g. Enzi et al., 2020; He et al., 2023) in different dark matter models (Despali et al., 2019, 2020).

7 Dark matter constraints

7.1 Compact sources

Since 2019, four new analyses (Hsueh et al., 2020; Gilman et al., 2020b,a; Laroche et al., 2022) have characterized the properties of low-mass haloes using a sample of quadruplyimaged quasars. These works interpret strong lensing data in the context of a variety of scenarios, including classes of dark matter such as warm dark matter, ultra-light dark matter, and self-interacting dark matter, as well as early Universe physics that alters the power spectrum of primordial density fluctuations.

Hsueh et al. (2020) and Gilman et al. (2020b) analyzed a sample of strong lens systems in the context of warm dark

matter, presenting evidence for an unseen population of dark matter haloes with masses around $10^8 M_{\odot}$. This inference places an upper limit on the free-streaming length of dark matter equivalent to that of a 5 to 6 keV thermal relic dark matter particle, and constrains a variety of sterile neutrino models with varying production mechanisms (Zelko et al., 2022). Gilman et al. (2020a) used the sensitivity of the relative magnifications among the lensed images to infer the concentrations of CDM haloes, an analysis that was later generalized by Gilman et al. (2022) to make a direct connection between the abundance and internal structure of dark matter haloes and the primordial matter power spectrum on small scales $(k > 10 \text{ Mpc}^{-1})$. The resulting inference on the concentration-mass relation and power spectrum agreed with the CDM prediction. Laroche et al. (2022) interpreted the same sample of eleven four-image lenses as analysed by Gilman et al. (2022) in the context of ultra-light dark matter, showing that the granular structure of the host halo density profile that arises from quantum wave interference effects can impact image flux ratios in a similar manner to dark matter haloes. Finally, building on work by Gilman et al. (2021), Gilman et al. (2022) interpreted the same sample of eleven lenses in the context of self-interacting dark matter and showed that existing data disfavors SIDM models with large amplitudes at low speeds, such as those that can arise from resonances in the self-interaction cross section, assuming the large amplitude of self-interaction cross section drives low-mass haloes to core-collapse. By combining observations of the Milky-Way satellites with the sample of strongly lensed compact sourses from Gilman et al. (2022), Nadler et al. (2021) have derived a limit on the half-mode mass of $M_{\rm hm} < 10^7 M_{\odot}$ (i.e., $m_{\rm WDM} > 9.7$ keV) at the 95 per cent confidence level (see Fig. 5).

These analyses have incorporated the latest theoretical understanding of structure formation in CDM and alternative dark matter models, such as the halo mass function and concentation-mass relation in warm dark matter, the process of core-collapse that is expected to occur in self-interacting dark matter, and the phenomenon of wave-interference unique to ultra-light dark matter. At the same time, Basu et al. (2021) have introduced a novel approach to constrain Axion-Like Particles dark matter models from the differential birefringence effect imparted on the strongly lensed images. Using broad-band polarisation observations of the lens system B1152+199 from the Karl G. Jansky Very Large Array (VLA), they derived an upper bound on the ALP-photon coupling between $g_{ay} \leq 9.2 \times 10^{-11}$ and 7.7×10^{-8} eV at the 95 per cent confidence limit for an ALP mass between $m_a = 3.6 \times 10^{-21}$ eV and 4.6×10^{-18} eV.

Two recent innovations have expanded the scope of strong lensing of compact sources as a probe of fundamental dark matter physics. First, the sample size of lenses suitable for a subhalo inference doubled with measurements of relative

 10^{-1} 10^{1} Expected detections per lens with $f_{sub} = 10^{-2}$ 2σ Expected detections per lens $(1/f_{sub})$ CDM 3σ 10^{0} 10 4σ N21 E21 H20 10 10^{12} 10^{4} 10^{5} 10^{6} 10^{7} 10^{8} 10^{9} 10^{10} 10^{11} $M_{\rm HM}~(M_{\odot})$

Fig. 5 Expected number of subhaloes per lens as a function of the half-mode mass from Euclid-like observations. Each curve (with 64 per cent, 95 per cent, and 99 per cent confidence areas) is for a different level of significance of subhalo detection. The horizontal dotted lines display the expected number of detectable subhaloes in CDM (see O'Riordan et al., 2023, for more details). The vertical dotted lines show the current 95 per cent upper limits on the half-mode mass from Nadler et al. (2021) and Hsueh et al. (2020) together with the 1/20th of the maximum Likelihood from Enzi et al. (2021).

image fluxes from narrow-line emission around the background quasar. Nuclear narrow-line emission, which emanates from an area in the source plane with a typical size between 10 to 100 pc, subtends angular scales on the sky much larger than a micro-arcsecond. This renders the relative brightness of lensed images immune to micro-lensing, while retaining sensitivity to milli-lensing by dark matter haloes. Second, open-source software to model gravitational lens systems and to generate populations of dark matter haloes for lensing computations have enabled ray-tracing simulations to be performed in parallel on computing clusters with realistic background sources and full populations of subhaloes. At the same time, recent analyses (from 2021 onwards) have made some efforts to mitigate potential systematic uncertainties associated with the lens macro-model by including the contribution of multipole mass moments.

7.2 Extended sources

Most studies based on extended lensed sources are so far consistent with the CDM model. For example, Vegetti et al. (2014) and Hezaveh et al. (2016) have inferred a dark matter fraction in subhaloes that is consistent with the subhalo population in CDM simulations (Springel et al., 2008; Xu et al., 2015; Despali and Vegetti, 2017). From an analysis

(which only include the contribution of subhaloes) of the lens system RXJ1131–1231, Birrer et al. (2017) have derived a 2- σ lower limit on the particle of a thermal relic dark matter model of $m_{\text{th}} > 2$ keV. On the other hand, Bayer et al. (2023b) have found an upper limit on the power-spectrum of mass density fluctuations in the lens system SDSS J0252+0039 that exceeds the value expected from CDM. Their result is likely driven by their choice of macro-model (SPL) and the lack of other form complexity beyond subhaloes.

Allowing for the contribution of field haloes, Vegetti et al. (2018) have derived constraints on sterile neutrinos that show a preference for colder dark matter models: $\log M_{\rm hm}[M_{\odot}] <$ 12.0 at the 2- σ level. This result excludes sterile neutrino models with neutrino masses $m_s < 0.8$ keV at any value of the lepton asymmetry L_6 . Ritondale et al. (2019a) reported zero detection of low-mass haloes in a sample of twenty-one strong gravitational lens systems from the BELLS survery, in agreement with CDM given the quality of the data. Recently, Enzi et al. (2021) have derived constraints on thermal relic and sterile neutrino dark matter models by combining the lensing results by Vegetti et al. (2018) and Ritondale et al. (2019a) with observations of the Lyman- α forest and the Milky-Way satellite galaxies. They derived a joint limit on the thermal relic mass of $m_{\rm th} > 6.048$ keV (i.e. $M_{\rm hm} < 3 \times 10^7 M_{\odot} h^{-1}$) at the 95 per cent confidence level



Fig. 6 Bounds on the FDM particle mass from different observational probes. The solid black curve and arrow show the fully marginalized posterior odds ratio and relative lower bound from Powell et al. (2023), respectively. These were derived from the analysis of mas-resolution observations of the lens system MG J0751+2716 taken with the Global Very Long Baseline Interferometry array. The results by Laroche et al. (2022, blue arrow) were obtained from a sample of eleven strongly lensed QSOs. Also shown are the constraints from the Lyman- α forest (Iršič et al., 2017, purple arrow), the Milky-Way satellites (Nadler et al., 2021, green arrow), stellar streams in the Milky-Way (Banik et al., 2021, dark red arrow). The limit by Marsh and Niemeyer (2019, red dashed arrow) lie beyond the limit of the plot, and were derived from observations of star clusters within the Milky-Way Ultra-faint Dwarf satellite galaxy Eridanus II.

(see Fig. 5). This result is mainly set by the Milky-Way and the Lyman- α forest. The lensing measurements alone lead to a 95 per cent confidence level lower limit of $m_{\rm th} > 0.6$ keV and $m_{\rm th} > 0.1$ keV, for the SLACS and BELLS samples, respectively.

While most published analyses of extended sources show a preference for CDM, they do not rule out alternative and still viable dark matter models yet. This lack of constraints is partly due to the low number of lens systems available and the relatively limited quality of the data: signal-to-noise ratio and in particular angular resolution. Using realistic mock observations of varying data quality, Despali et al. (2022) have recently quantified how the sensitivity to subhaloes depends on the properties of the lens system (e.g. source structure and position relative to the caustics) and the data (i.e. signal-to-noise ratio and angular resolution). They concluded that the lowest detectable subhalo mass decreases linearly with signal-to-noise ratio, and more strongly with the angular resolution. An increase in the latter by a factor of two leads on average to an increase in mass sensitivity of a similar factor (for a fixed signal-to-noise ratio, lensing configuration and source properties). For example, existing HST observations (< SNR >~ 3.5 and PSF_{FWHM} = 0.09 mas) have a maximum sensitivity of $M_{\rm vir}^{\rm NFW} = 10^9 M_{\odot}$, leading to an expected number of detectable subhaloes per lens in CDM

consistent with the current number of objects detected so far (Vegetti et al., 2014; Ritondale et al., 2019a; Nightingale et al., 2022). These effects also explain why the constraints are, at present, less stringent (though possibly more robust) than those obtained with compact sources: predictions from different dark matter models mostly differ at halo masses smaller than currently probed.

Observations at very high-angular resolution are therefore key to probe the nature of dark matter with extended sources. For example, from observations of the gravitational lens system MG J0751+2716 taken with the Global Very Long Baseline Interferometry array, Powell et al. (2023) have ruled out scalar FDM models with a particle mass $m_{\gamma} \leq$ 4.4×10^{-21} eV. Their constraints, which were obtained from a single lens system with milli-arcsecond angular resolution, are more stringent than previously published results from larger samples of lensed quasars (see Fig. 6). Their work, together with the results by Despali et al. (2022) and O'Riordan et al. (2023), clearly demonstrates how a smaller number of high-angular resolution observations can be more effective at constraining the properties of dark matter than many lens systems with lower quality data. We will discuss this further in the context of future surveys in the following section.

8 Future prospects

At present, the level of constraints that can be obtained on the properties of dark matter from both extended and compact sources has mainly been limited by the amount of available data. Strong gravitational lensing is a rare phenomenon and the number of known systems with suitable data (e.g. high enough signal-to-noise ratio and angular resolution for extended sources and the right frequency coverage for compact sources) is small (of the order of a few tens).

Thanks to ongoing wide-sky surveys the number of known strong gravitational lens system has significantly increased in recent years (e.g. Storfer et al., 2022). With the advent of the next generation of sky surveys with, e.g., Euclid, the Low-Frequency Array (LOFAR), the Square Kilometre Array (SKA) and the Vera Rubin Observatory, this number is expected to increase even further (e.g. Oguri and Marshall, 2010; McKean et al., 2015). The size of these samples is unprecedented. Coupled with follow-up observations they will provide robust and meaningful constraints on dark matter.

8.1 Compact sources

Increasing the sample size for four-image systems will reduce the statistical uncertainties of measurements made with these types of system. Provided that all sources of systematic error can be accounted for, this increased constraining power can lead to stronger constraints over various dark matter models, such as warm and self-interacting dark matter (Gilman et al., 2019, 2021; Hsueh et al., 2020). The existing analysis techniques applied to interpret data from multiply-imaged quasars can scale to match the increased sample size. Thus, computational costs will not limit the scientific output achievable with forthcoming data.

Follow-up observations will be required to precisely measure the positions and relative magnifications among the unresolved images of the lensed source. These observations can be performed with spaced-based observatories, such as HST and JWST (while they remain operational), but most likely with ground-based facilities with adaptive optics for optical and infrared-bright quasars (Nierenberg et al., 2014), and with the Very Large Array (VLA), the enhanced Multi Element Remotely Linked Interferometer Network (e-MERLIN) source of systematic error. Moreover, in those cases withand VLBI for compact lensed radio sources. Upcoming observational facilities, such as the Keck All sky Precision Adaptive optics program (Wizinowich et al., 2022) and thirtymetre class telescopes, like the European Extremely Large Telescope (ELT), will provide exquisite astrometry and sensitivity with which to measure the relative image fluxes and positions. Likewise, the SKA will provide high resolution imaging at a resolution of 30 to 70 mas, which, when combined with VLBI, can provide astrometric measurements at sub-mas-arcsecond precision (e.g. Spingola and Barnacka 2020).

The JWST will soon deliver precise measurements of relative image fluxes in the mid-infrared at >20 μ m (Nierenberg et al., 2021), emission that emanates from a more compact region (1 to 10 pc) around the background source than the nuclear narrow-line emission (10 to 100 pc) that has been measured with the HST and Keck-AO (Nierenberg et al., 2020). The spatial extent of the mid-IR emission renders these data immune to stellar micro-lensing and variability of the background quasar, systematic effects that one could conflate with the perturbation by low-mass haloes ⁵. The more compact mid-IR emission that is measurable with the JWST increases sensitivity to perturbations by low-mass haloes relative to existing data because the minimum deflection angle that can affect the data scales with the angular size of the lensed source (Dobler and Keeton, 2006). Mid-IR flux ratios measured with the JWST have anticipated sensitivity to populations of haloes less massive than $10^7 M_{\odot}$ (Nierenberg et al., 2021). Moreover, the compact background sources measurable with JWST will increase sensitivity to the internal structure of low-mass haloes. This improved sensitivity, in combination with the expanded sample size of strong lens systems, will enable searches for populations of core-

collapsed haloes, such as structures with logarithmic central density profile slopes steeper than -2 (Balberg et al., 2002; Turner et al., 2021). If detected, the existence of these objects would suggest dark matter has a velocity-dependent cross section with strong self-interactions at low speeds that triggers core collapse (Gilman et al., 2021; Turner et al., 2021; Yang et al., 2022; Gilman et al., 2022). Increased sensitivity to the internal structure of low-mass haloes will also enable stronger constraints on the concentration-mass relation of low-mass haloes (Gilman et al., 2020a). This type of measurement can be interpreted in the context of the primordial matter power spectrum on scales $k > 10 \text{ Mpc}^{-1}$ (Zentner and Bullock, 2003; Gilman et al., 2022). However, as the background sources will remain unresolved, their unknown surface brightness distribution will still stand as a potential out any extended emission from the quasar-host galaxy, the limited number of observational constraints provided by the data may lead to a potential systematic error in the macromodel.

At radio wavelengths (between around 4 and 18 GHz), large fractional bandwidths with, for example, the ngVLA and the SKA, will detect the emission from pc-scale radio sources, which are expected to be immune from microlensing (but see, e.g., Koopmans and de Bruyn 2000; Biggs 2023). The orders of magnitude improvement in sensitivity will allow monitoring of the lensed radio sources for cosmology (in the case of any intrinsic variability) or for darkmatter studies, when the radio source is found not to vary. While the large fractional bandwidths will allow radio propagation effects, such as free-free absorption or scattering, to be identified, which can either corrected for or used to create samples of lensed radio sources that do not show such issues. However, the main contribution that radio-selected samples can provide is our ability to characterise the source structure on mas-scales through observations with VLBI (e.g McKean et al., 2007; Spingola and Barnacka, 2020). In general, it is worth noting that combining the data from multiple emission regions (mid-IR-bright torus, NRL, radio-jets) will allow various systematics associated with the source model to be quantified and possibly corrected for. This may include better constraints to the macro-model via large scale source properties, such as the optical/infrared emission from the stellar component of the quasar host galaxy or through high resolution imaging of the kpc-scale dust emission with ALMA (e.g. Stacey et al. 2021).

Finally, high-sensitivity observations with the LOFAR2.0 and SKA-MID of large samples of polarised and strongly lensed sources are expected to improve upon current bounds on the ALP-photon coupling by one to two orders of magnitude, depending on the ALP mass range (see Basu et al., 2021, and Fig. 7). Providing, thereby, competitive constraints on dark-matter models made of ALPs.

⁵ Scattering of light into our field of view by the spatially-extended nuclear narrow-line and mid-IR emission regions acts as a low-pass filter that washes out variability in the quasar light curves on timescales less than the light crossing time.



Fig. 7 Limits on the ALP-photon coupling from strong gravitational lensing. The dashed area shows current bounds from the gravitational lens system B1152+199. The dash-dotted blue line shows the expected constraint that can be obtained from monitoring B1152+199 over the course of ~ 5 year. The solid and dashed green lines show the predicted bounds for a sample of 100 and 1000 gravitational lens systems for different values of the maximum time delay (Δt_{max}) on the observed plane. The dotted black line shows the parameter space that can be probed with the SKA. The figure is taken from Basu et al. (2021).

8.2 Extended sources

The larger number of strong gravitational lens systems with an extended source that will be discovered by wide-sky surveys will represent a unique opportunity. However, followup observations at high angular resolution will be necessary to obtain meaningful constraints on the properties of dark matter. Recently, O'Riordan et al. (2023) calculated the sensitivity function for 16000 Euclid-VIS-like mock observations (see Fig. 5). They found most of the lenses in the sample to be completely insensitive to the presence of subhaloes with a mass lower than $M_{\text{max}}^{\text{NFW}} = 10^{11} M_{\odot}$. The most sensitive pixels yield a lowest detectable mass of $M_{\text{max}}^{\text{NFW}}$ = $10^{8.8\pm0.2}M_{\odot}$. Assuming CDM and a dark matter fraction in subhaloes of 0.01, this sensitivity leads to one detectable object in every seventy lenses. These results are in line with Despali et al. (2022) and the current number of detected objects (see previous section). Hence, Euclid will allow one to detect a number of subhaloes that is significantly larger than what has been possible so far, and potentially deliver tighter constraints on the amount of dark matter in subhaloes (i.e. the normalisation of the subhalo mass function) than currently possible. However, most of these objects will be relatively massive, and will probe (thermal relic) dark matter models with a half-mode mass of $M_{\rm hm} > 10^8 M_{\odot}$. These

models have already been ruled out by other observations, including strongly lensed quasars. Similarly, if not slightly worse results are expected from the Vera Rubin Observatory, given its slightly worse angular resolution.

While the sensitivity to the presence of low-mass haloes increases linearly with decreasing signal-to-noise ratio, the angular resolution of the observation sets a hard limit on the lowest detectable halo and the spatial scales on which the dark matter distribution can be probed (Despali et al., 2022; O'Riordan et al., 2023). For this reason, follow-up observations with Keck-AO, HST and ALMA are likely to lead to an increase in sensitivity (relatively to Euclid) between two to three orders in magnitude in halo mass. With an angular resolution of a few milli-arcseconds, Global VLBI observations at 1.6 to 15 GHz, will allow one to detect masses as low as $10^6 M_{\odot}$ (McKean et al., 2015), and probe the general spatial distribution of dark matter on sub-kpc scales (e.g. Powell et al., 2023). Moreover, searches for milli-arcsecond scale separation images with VLBI will potentially reveal the presence of super-critical low-mass haloes, which existence is predicted by certain SIDM models (Casadio et al., 2021; Loudas et al., 2022). In the future, 30-metre class optical telescopes such as the ELT are likely to provide a similar level of sensitivity, while interferometric observations in the 10 to 100 GHz range with SKA-VLBI or the ngVLA will lead to an even stronger increase. This significant improvement in data quality will also result in better constraints on the properties of the lens macro-model. This effect will significantly reduce the role of one of the main sources of systematic uncertainty and will potentially have interesting implications for other scientific applications of strong gravitational lensing. A relatively small number of high-resolution observations is, therefore, expected to potentially deliver stronger constraints on the properties of dark matter than the several orders of magnitude larger sample of Euclid and Vera Rubin lenses. Note, however, that these facilities and also LOFAR and the SKA will be vital for finding the lenses needed for high-resolution follow-up observations.

9 Summary and conclusions

Strong gravitational lensing provides a unique channel to constrain the dark matter distribution on subgalactic scales, and hence provide a key test of several dark matter models. In this chapter, we described the process of inferring the properties of dark matter from strong lensing observations in terms of a Bayesian inference problem, and showed how it can be approached from different perspectives (Section 4). For each possible prior choice and lens modelling technique, we then discussed their advantages and limitations (Section 4.4). In Section 5 we provided an extensive discussion of the degeneracies and theoretical unknowns. In particular, we highlighted how better numerical simulations are needed to answer important and still open theoretical questions (Section 5.5).

While we only focused on galaxy-scale lenses, all of the approaches discussed above can potentially be extended to cluster-scale lenses. These type of studies would allows one to probe larger cosmological volumes and hence potentially require fewer observations to reach an equal level of constraints on dark matter. However, the lensing potential in galaxy clusters is significantly more complex than those of galaxies and it remains yet to be demonstrated that one can constrain their mass distribution with enough precision to avoid contamination by complex macro-models (see Section 5.1 for a discussion).

Currently, constraints on dark matter from lensing are based on relative small samples of systems (about ten for compact sources and about 50 for extended ones). Ongoing and upcoming wide-sky surveys, with for example Euclid, the Vera Rubin Telescope, LOFAR and the SKA are expected to deliver orders of magnitudes more lensed galaxies and quasars. On the other hand, high-resolution followup with, for example, ALMA, the Global VLBI Network and the ELT will likely provide the data quality necessary to probe the dark matter distribution on small scales with extended sources as well as obtain better constraints on the macro-model for compact sources. We can expect, therefore, that in the next five to ten years, the field of strong gravitational lensing will provide robust and statistically meaningful constraints on the nature of dark matter.

Acknowledgements S. V. thanks the Max Planck Society for support through a Max Planck Lise Meitner Group and funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (LEDA: grant agreement No 758853). G. V. has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodovska-Curie grant agreement No 897124. G. D. was supported by a Gliese Fellowship. C. D. F acknowledges support for this work from the National Science Foundation under Grant No. AST-1715611. D. G. was partially supported by a HQP grant from the McDonald Institute (reference number HQP 2019-4-2). Y. H. and L. P. are supported by a generous donation by Eric and Wendy Schmidt with the recommendation of the Schmidt Futures Foundation, as well as the National Sciences and Engineering Council of Canada, the Fonds de recherche du Québec, and the Canada Research Chairs Program. J. P. M. acknowledges support from the Netherlands Organization for Scientific Research (NWO, project number 629.001.023) and the Chinese Academy of Sciences (CAS, project number 114A11KYSB20170054).

References

T.M.C. Abbott, M. Aguena, A. Alarcon, S. Allam, O. Alves, A. Amon, F. Andrade-Oliveira, J. Annis, S. Avila, D. Bacon, E. Baxter, K. Bechtol, M.R. Becker, G.M. Bernstein, S. Bhargava, S. Birrer, J. Blazek, A. Brandao-Souza, S.L. Bridle, D. Brooks, E. Buckley-Geer, D.L. Burke, H. Camacho, A. Campos, A. Carnero Rosell, M. Carrasco Kind, J. Carretero, F.J. Castander, R. Cawthon, C. Chang, A. Chen, R. Chen, A. Choi, C. Conselice, J. Cordero, M. Costanzi, M. Crocce, L.N. da Costa, M.E. da Silva Pereira, C. Davis, T.M. Davis, J. De Vicente, J. DeRose, S. Desai, E. Di Valentino, H.T. Diehl, J.P. Dietrich, S. Dodelson, P. Doel, C. Doux, A. Drlica-Wagner, K. Eckert, T.F. Eifler, F. Elsner, J. Elvin-Poole, S. Everett, A.E. Evrard, X. Fang, A. Farahi, E. Fernandez, I. Ferrero, A. Ferté, P. Fosalba, O. Friedrich, J. Frieman, J. García-Bellido, M. Gatti, E. Gaztanaga, D.W. Gerdes, T. Giannantonio, G. Giannini, D. Gruen, R.A. Gruendl, J. Gschwend, G. Gutierrez, I. Harrison, W.G. Hartley, K. Herner, S.R. Hinton, D.L. Hollowood, K. Honscheid, B. Hoyle, E.M. Huff, D. Huterer, B. Jain, D.J. James, M. Jarvis, N. Jeffrey, T. Jeltema, A. Kovacs, E. Krause, R. Kron, K. Kuehn, N. Kuropatkin, O. Lahav, P.-F. Leget, P. Lemos, A.R. Liddle, C. Lidman, M. Lima, H. Lin, N. MacCrann, M.A.G. Maia, J.L. Marshall, P. Martini, J. McCullough, P. Melchior, J. Mena-Fernández, F. Menanteau, R. Miquel, J.J. Mohr, R. Morgan, J. Muir, J. Myles, S. Nadathur, A. Navarro-Alsina, R.C. Nichol, R.L.C. Ogando, Y. Omori, A. Palmese, S. Pandey, Y. Park, F. Paz-Chinchón, D. Petravick, A. Pieres, A.A. Plazas Malagón, A. Porredon, J. Prat, M. Raveri, M. Rodriguez-Monroy, R.P. Rollins, A.K. Romer, A. Roodman, R. Rosenfeld, A.J. Ross, E.S. Rykoff, S. Samuroff, C. Sánchez, E. Sanchez, J. Sanchez, D. Sanchez Cid, V. Scarpine, M. Schubnell, D. Scolnic, L.F. Secco, S. Serrano, I. Sevilla-Noarbe, E. Sheldon, T. Shin, M. Smith, M. Soares-Santos, E. Suchyta, M.E.C. Swanson, M. Tabbutt, G. Tarle, D. Thomas, C. To, A. Troja, M.A. Troxel, D.L. Tucker, I. Tutusaus, T.N. Varga, A.R. Walker, N. Weaverdyck, R. Wechsler, J. Weller, B. Yanny, B. Yin, Y. Zhang, J. Zuntz, DES Collaboration, Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and weak lensing. PRD 105(2), 023520 (2022). doi:10.1103/PhysRevD.105.023520

- A. Adam, A. Coogan, N. Malkin, R. Legin, L. Perreault-Levasseur, Y. Hezaveh, Y. Bengio, Posterior samples of source galaxies in strong gravitational lenses with score-based priors. arXiv e-prints, 2211–03812 (2022). doi:10.48550/arXiv.2211.03812
- S. Adhikari, A. Banerjee, K.K. Boddy, F.-Y. Cyr-Racine, H. Desmond, C. Dvorkin, B. Jain, F. Kahlhoefer, M. Kaplinghat, A. Nierenberg, A.H.G. Peter, A. Robertson, J. Sakstein, J. Zavala, Astrophysical Tests of Dark Matter Self-Interactions. arXiv e-prints, 2207–10638 (2022). doi:10.48550/arXiv.2207.10638
- S. Alam, M. Ata, S. Bailey, F. Beutler, D. Bizyaev, J.A. Blazek, A.S. Bolton, J.R. Brownstein, A. Burden, C.-H. Chuang, et al., The clustering of galaxies in the completed sdss-iii baryon oscillation spectroscopic survey: cosmo-

logical analysis of the dr12 galaxy sample. Monthly Notices of the Royal Astronomical Society **470**(3), 2617– 2652 (2017)

- S. Alexander, S. Gleyzer, H. Parul, P. Reddy, M.W. Toomey, E. Usai, R. Von Klar, Decoding Dark Matter Substructure without Supervision. arXiv e-prints, 2008–12731 (2020a). doi:10.48550/arXiv.2008.12731
- S. Alexander, S. Gleyzer, E. McDonough, M.W. Toomey, E. Usai, Deep Learning the Morphology of Dark Matter Substructure. Astrophys. J. 893(1), 15 (2020b). doi:10.3847/1538-4357/ab7925
- S. Alexander, S. Gleyzer, P. Reddy, M. Tidball, M.W. Toomey, Domain Adaptation for Simulation-Based Dark Matter Searches Using Strong Gravitational Lensing. arXiv e-prints, 2112–12121 (2021)
- N.C. Amorisco, J. Nightingale, Q. He, A. Amvrosiadis, X. Cao, S. Cole, A. Etherington, C.S. Frenk, R. Li, R. Massey, A. Robertson, Halo concentration strengthens dark matter constraints in galaxy-galaxy strong lensing analyses. Mon. Notices of the Royal Astron. Soc. 510(2), 2464–2479 (2022). doi:10.1093/mnras/stab3527
- A. Amruth, T. Broadhurst, J. Lim, M. Oguri, G.F. Smoot, J.M. Diego, E. Leung, R. Emami, J. Li, T. Chiueh, H.-Y. Schive, M.C.H. Yeung, S. Kei Li, Anomalies in Gravitational-Lensed Images Revealing Einstein Rings Modulated by Wavelike Dark Matter. arXiv e-prints, 2304–09895 (2023). doi:10.48550/arXiv.2304.09895
- K.E. Andrade, J. Fuson, S. Gad-Nasr, D. Kong, Q. Minor, M.G. Roberts, M. Kaplinghat, A stringent upper limit on dark matter self-interaction cross-section from cluster strong lensing. Mon. Notices of the Royal Astron. Soc. 510(1), 54–81 (2022). doi:10.1093/mnras/stab3241
- G. Arcadi, M. Dutra, P. Ghosh, M. Lindner, Y. Mambrini, M. Pierre, S. Profumo, F.S. Queiroz, The waning of the WIMP? A review of models, searches, and constraints. European Physical Journal C 78(3), 203 (2018). doi:10.1140/epjc/s10052-018-5662-y
- M.W. Auger, T. Treu, A.S. Bolton, R. Gavazzi, L.V.E. Koopmans, P.J. Marshall, K. Bundy, L.A. Moustakas, The Sloan Lens ACS Survey. IX. Colors, Lensing, and Stellar Masses of Early-Type Galaxies. Astrophys. J. **705**(2), 1099–1115 (2009). doi:10.1088/0004-637X/705/2/1099
- S. Balberg, S.L. Shapiro, S. Inagaki, Self-Interacting Dark Matter Halos and the Gravothermal Catastrophe. Astrophys. J. 568(2), 475–487 (2002). doi:10.1086/339038
- N. Banik, J. Bovy, G. Bertone, D. Erkal, T.J.L. de Boer, Novel constraints on the particle nature of dark matter from stellar streams. J. of Cosm. and Astroparticle Phys. **2021**(10), 043 (2021). doi:10.1088/1475-7516/2021/10/043
- U. Banik, F.C. van den Bosch, M. Tremmel, A. More, G. Despali, S. More, S. Vegetti, J.P. McKean, Constraining the mass density of free-floating black holes using razor-

thin lensing arcs. Mon. Notices of the Royal Astron. Soc. **483**(2), 1558–1573 (2019). doi:10.1093/mnras/sty3267

- J.M. Bardeen, J.R. Bond, N. Kaiser, A.S. Szalay, The statistics of peaks of Gaussian random fields. Astrophys. J. 304, 15–61 (1986). doi:10.1086/164143
- M. Barnabè, C. Nipoti, L.V.E. Koopmans, S. Vegetti, L. Ciotti, Crash-testing the CAULDRON code for joint lensing and dynamics analysis of early-type galaxies. Mon. Notices of the Royal Astron. Soc. **393**(4), 1114–1126 (2009). doi:10.1111/j.1365-2966.2008.14208.x
- M. Barnabè, O. Czoske, L.V.E. Koopmans, T. Treu, A.S. Bolton, Two-dimensional kinematics of SLACS lenses III. Mass structure and dynamics of early-type lens galaxies beyond z ~ 0.1. Mon. Notices of the Royal Astron. Soc. 415(3), 2215–2232 (2011). doi:10.1111/j.1365-2966.2011.18842.x
- A. Basu, J. Goswami, D.J. Schwarz, Y. Urakawa, Searching for Axionlike Particles under Strong Gravitational Lenses. PRL **126**(19), 191102 (2021). doi:10.1103/PhysRevLett.126.191102
- D. Bayer, L.V.E. Koopmans, J.P. McKean, S. Vegetti, T. Treu, C.D. Fassnacht, K. Glazebrook, Probing subgalactic mass structure with the power spectrum of surface-brightness anomalies in high-resolution observations of galaxy-galaxy strong gravitational lenses - I. Power-spectrum measurement and feasibility study. Mon. Notices of the Royal Astron. Soc. 523(1), 1326–1345 (2023a). doi:10.1093/mnras/stad1403
- D. Bayer, S. Chatterjee, L.V.E. Koopmans, S. Vegetti, J.P. McKean, T. Treu, C.D. Fassnacht, K. Glazebrook, Probing sub-galactic mass structure with the power spectrum of surface-brightness anomalies in high-resolution observations of galaxy-galaxy strong gravitational lenses. II. Observational constraints on the subgalactic matter power spectrum. Mon. Notices of the Royal Astron. Soc. 523(1), 1310–1325 (2023b). doi:10.1093/mnras/stad1402
- C.L. Bennett, C.R. Lawrence, B.F. Burke, J.N. Hewitt, J. Mahoney, The MIT-Green Bank (MG) 5 GHz Survey. Astrophys. J. Supplement **61**, 1 (1986). doi:10.1086/191108
- G. Bertone, T.M.P. Tait, A new era in the search for dark matter. Nature **562**(7725), 51–56 (2018). doi:10.1038/s41586-018-0542-z
- A.D. Biggs, A VLA monitoring study of JVAS B1422+231: investigation of time delays and detection of extrinsic variability. Mon. Notices of the Royal Astron. Soc. 522(1), 426–437 (2023). doi:10.1093/mnras/stad870
- A.D. Biggs, I.W.A. Browne, N.J. Jackson, T. York, M.A. Norbury, J.P. McKean, P.M. Phillips, Radio, optical and infrared observations of CLASS B0128+437. Mon. Notices of the Royal Astron. Soc. **350**(3), 949–961 (2004). doi:10.1111/j.1365-2966.2004.07701.x
- S. Birrer, A. Amara, A. Refregier, Gravitational Lens Mod-

eling with Basis Sets. Astrophys. J. **813**(2), 102 (2015). doi:10.1088/0004-637X/813/2/102

- S. Birrer, A. Amara, A. Refregier, Lensing substructure quantification in RXJ1131-1231: a 2 keV lower bound on dark matter thermal relic mass. J. of Cosm. and Astroparticle Phys. **2017**(5), 037 (2017). doi:10.1088/1475-7516/2017/05/037
- A.S. Bolton, S. Burles, L.V.E. Koopmans, T. Treu, R. Gavazzi, L.A. Moustakas, R. Wayth, D.J. Schlegel, The Sloan Lens ACS Survey. V. The Full ACS Strong-Lens Sample. Astrophys. J. 682(2), 964–984 (2008). doi:10.1086/589327
- J.R. Bond, S. Cole, G. Efstathiou, N. Kaiser, Excursion set mass functions for hierarchical Gaussian fluctuations. Astrophys. J. 379, 440–460 (1991). doi:10.1086/170520
- A. Boyarsky, M. Drewes, T. Lasserre, S. Mertens, O. Ruchayskiy, Sterile neutrino Dark Matter. Progress in Particle and Nuclear Physics 104, 1–45 (2019). doi:10.1016/j.ppnp.2018.07.004
- A. Boyarsky, O. Ruchayskiy, M. Shaposhnikov, The role of sterile neutrinos in cosmology and astrophysics. Annual Review of Nuclear and Particle Science 59(1), 191– 214 (2009). doi:10.1146/annurev.nucl.010909.083654. https://doi.org/10.1146/annurev.nucl.010909.083654
- M. Bradač, P. Schneider, M. Steinmetz, M. Lombardi, L.J. King, R. Porcas, B1422+231: The influence of mass substructure on strong lensing. Astron. Astrophys. 388, 373– 382 (2002). doi:10.1051/0004-6361:20020559
- M. Bradač, P. Schneider, M. Lombardi, M. Steinmetz, L.V.E. Koopmans, J.F. Navarro, The signature of substructure on gravitational lensing in the *A*CDM cosmological model. Astron. Astrophys. **423**, 797–809 (2004). doi:10.1051/0004-6361:20040168
- J. Brehmer, S. Mishra-Sharma, J. Hermans, G. Louppe, K. Cranmer, Mining for dark matter substructure: Inferring subhalo population properties from strong lenses with machine learning. The Astrophysical Journal 886(1), 49 (2019). doi:10.3847/1538-4357/ab4c41. https://dx.doi.org/10.3847/1538-4357/ab4c41
- B.J. Brewer, D. Foreman-Mackey, D.W. Hogg, Probabilistic Catalogs for Crowded Stellar Fields. Astron. J. 146(1), 7 (2013). doi:10.1088/0004-6256/146/1/7
- B.J. Brewer, D. Huijser, G.F. Lewis, Trans-dimensional Bayesian inference for gravitational lens substructures. Mon. Notices of the Royal Astron. Soc. 455(2), 1819– 1829 (2016). doi:10.1093/mnras/stv2370
- A.M. Brooks, M. Kuhlen, A. Zolotov, D. Hooper, A Baryonic Solution to the Missing Satellites Problem. Astrophys. J. 765(1), 22 (2013). doi:10.1088/0004-637X/765/1/22
- I.W.A. Browne, et al., The Cosmic Lens All-Sky Survey
 II. Gravitational lens candidate selection and followup. Mon. Notices of the Royal Astron. Soc. 341, 13–32

(2003). doi:10.1046/j.1365-8711.2003.06257.x

- I.W.A. Browne, P.N. Wilkinson, A.R. Patnaik, J.M. Wrobel, Interferometer phase calibration sources - II. The region 0 deg<=delta_B1950<=+20 deg. Mon. Notices of the Royal Astron. Soc. 293(3), 257–287 (1998). doi:10.1046/j.1365-8711.1998.01072.x
- G.L. Bryan, M.L. Norman, Statistical Properties of X-Ray Clusters: Analytic and Numerical Comparisons. Astrophys. J. 495, 80 (1998). doi:10.1086/305262
- R.S. Bussmann, M.A. Gurwell, H. Fu, D.J.B. Smith, S. Dye, R. Auld, M. Baes, A.J. Baker, D. Bonfield, A. Cava, D.L. Clements, A. Cooray, K. Coppin, H. Dannerbauer, A. Dariush, G. De Zotti, L. Dunne, S. Eales, J. Fritz, R. Hopwood, E. Ibar, R.J. Ivison, M.J. Jarvis, S. Kim, L.L. Leeuw, S. Maddox, M.J. Michałowski, M. Negrello, E. Pascale, M. Pohlen, D.A. Riechers, E. Rigby, D. Scott, P. Temi, P.P. Van der Werf, J. Wardlow, D. Wilner, A. Verma, A Detailed Gravitational Lens Model Based on Submillimeter Array and Keck Adaptive Optics Imaging of a Herschel-ATLAS Submillimeter Galaxy at z = 4.243. Astrophys. J. 756(2), 134 (2012). doi:10.1088/0004-637X/756/2/134
- R.S. Bussmann, I. Pérez-Fournon, S. Amber, J. Calanog, M.A. Gurwell, H. Dannerbauer, F. De Bernardis, H. Fu, A.I. Harris, M. Krips, A. Lapi, R. Maiolino, A. Omont, D. Riechers, J. Wardlow, A.J. Baker, M. Birkinshaw, J. Bock, N. Bourne, D.L. Clements, A. Cooray, G. De Zotti, L. Dunne, S. Dye, S. Eales, D. Farrah, R. Gavazzi, J. González Nuevo, R. Hopwood, E. Ibar, R.J. Ivison, N. Laporte, S. Maddox, P. Martínez-Navajas, M. Michalowski, M. Negrello, S.J. Oliver, I.G. Roseboom, D. Scott, S. Serjeant, A.J. Smith, M. Smith, A. Streblyanska, E. Valiante, P. van der Werf, A. Verma, J.D. Vieira, L. Wang, D. Wilner, Gravitational Lens Models Based on Submillimeter Array Imaging of Herschel-selected Strongly Lensed Sub-millimeter Galaxies at z > 1.5. Astrophys. J. 779(1), 25 (2013). doi:10.1088/0004-637X/779/1/25
- R. Cañameras, S. Schuldt, Y. Shu, S.H. Suyu, S. Taubenberger, T. Meinhardt, L. Leal-Taixé, D.C.-Y. Chao, K.T. Inoue, A.T. Jaelani, A. More, HOLISMOKES. VI. New galaxy-scale strong lens candidates from the HSC-SSP imaging survey. Astron. Astrophys. 653, 6 (2021). doi:10.1051/0004-6361/202141758
- C. Casadio, D. Blinov, A.C.S. Readhead, I.W.A. Browne, P.N. Wilkinson, T. Hovatta, N. Mandarakas, V. Pavlidou, K. Tassis, H.K. Vedantham, J.A. Zensus, V. Diamantopoulos, K.E. Dolapsaki, K. Gkimisi, G. Kalaitzidakis, M. Mastorakis, K. Nikolaou, E. Ntormousi, V. Pelgrims, K. Psarras, SMILE: Search for MIlli-LEnses. Mon. Notices of the Royal Astron. Soc. 507(1), 6–10 (2021). doi:10.1093/mnrasl/slab082
- H.Y.J. Chan, E.G.M. Ferreira, S. May, K. Hayashi, M. Chiba, The diversity of core-halo structure in the fuzzy

dark matter model. Mon. Notices of the Royal Astron. Soc. **511**(1), 943–952 (2022). doi:10.1093/mnras/stac063

- J.H.H. Chan, H.-Y. Schive, S.-K. Wong, T. Chiueh, T. Broadhurst, Multiple Images and Flux Ratio Anomaly of Fuzzy Gravitational Lenses. PRL **125**(11), 111102 (2020). doi:10.1103/PhysRevLett.125.111102
- S. Chatterjee, L.V.E. Koopmans, The inner mass power spectrum of galaxies using strong gravitational lensing: beyond linear approximation. Mon. Notices of the Royal Astron. Soc. 474(2), 1762–1772 (2018). doi:10.1093/mnras/stx2674
- J. Chen, Parity dependence in strong lens systems as a probe of dark matter substructure. Astron. Astrophys. 498(1), 49–60 (2009). doi:10.1051/0004-6361/200811134
- J. Chen, E. Rozo, N. Dalal, J.E. Taylor, Astrometric Perturbations in Substructure Lensing. Astrophys. J. 659(1), 52–68 (2007). doi:10.1086/512002
- M. Chianese, A. Coogan, P. Hofma, S. Otten, C. Weniger, Differentiable strong lensing: uniting gravity and neural nets through differentiable probabilistic programming. Mon. Notices of the Royal Astron. Soc. 496(1), 381–393 (2020). doi:10.1093/mnras/staa1477
- M. Chiba, T. Minezaki, N. Kashikawa, H. Kataza, K.T. Inoue, Subaru Mid-Infrared Imaging of the Quadruple Lenses PG 1115+080 and B1422+231: Limits on Substructure Lensing. Astrophys. J. 627(1), 53–61 (2005). doi:10.1086/430403
- D. Clowe, M. Bradač, A.H. Gonzalez, M. Markevitch, S.W. Randall, C. Jones, D. Zaritsky, A Direct Empirical Proof of the Existence of Dark Matter. Astrophys. J. Letters 648(2), 109–113 (2006). doi:10.1086/508162
- A. Coogan, K. Karchev, C. Weniger, Targeted Likelihood-Free Inference of Dark Matter Substructure in Strongly-Lensed Galaxies. arXiv e-prints, 2010–07032 (2020)
- T.J. Cornwell, K.F. Evans, A simple maximum entropy deconvolution algorithm. Astron. Astrophys. 143(1), 77–83 (1985)
- C.A. Correa, Constraining velocity-dependent selfinteracting dark matter with the Milky Way's dwarf spheroidal galaxies. Mon. Notices of the Royal Astron. Soc. **503**(1), 920–937 (2021). doi:10.1093/mnras/stab506
- K. Cranmer, J. Brehmer, G. Louppe, The frontier of simulation-based inference. Proceedings of the National Academy of Science **117**(48), 30055–30062 (2020). doi:10.1073/pnas.1912789117
- A.Ç. Şengül, C. Dvorkin, Probing dark matter with strong gravitational lensing through an effective density slope.
 Mon. Notices of the Royal Astron. Soc. 516(1), 336–357 (2022). doi:10.1093/mnras/stac2256
- F.-Y. Cyr-Racine, C.R. Keeton, L.A. Moustakas, Beyond subhalos: Probing the collective effect of the Universe's small-scale structure with grav-

itational lensing. PRD **100**(2), 023013 (2019). doi:10.1103/PhysRevD.100.023013

- F.-Y. Cyr-Racine, L.A. Moustakas, C.R. Keeton, K. Sigurdson, D.A. Gilman, Dark census: Statistically detecting the satellite populations of distant galaxies. PRD 94(4), 043505 (2016). doi:10.1103/PhysRevD.94.043505
- N. Dalal, C.S. Kochanek, Direct Detection of Cold Dark Matter Substructure. Astrophys. J. 572(1), 25–33 (2002). doi:10.1086/340303
- T. Daylan, S.K.N. Portillo, D.P. Finkbeiner, Inference of Unresolved Point Sources at High Galactic Latitudes Using Probabilistic Catalogs. Astrophys. J. 839(1), 4 (2017). doi:10.3847/1538-4357/aa679e
- T. Daylan, F.-Y. Cyr-Racine, A. Diaz Rivero, C. Dvorkin, D.P. Finkbeiner, Probing the Small-scale Structure in Strongly Lensed Systems via Transdimensional Inference. Astrophys. J. 854(2), 141 (2018). doi:10.3847/1538-4357/aaaa1e
- G. Despali, S. Vegetti, The impact of baryonic physics on the subhalo mass function and implications for gravitational lensing. Mon. Notices of the Royal Astron. Soc. 469, 1997–2010 (2017). doi:10.1093/mnras/stx966
- G. Despali, C. Giocoli, R.E. Angulo, G. Tormen, R.K. Sheth, G. Baso, L. Moscardini, The universality of the virial halo mass function and models for nonuniversality of other halo definitions. Mon. Notices of the Royal Astron. Soc. 456, 2486–2504 (2016). doi:10.1093/mnras/stv2842
- G. Despali, M. Sparre, S. Vegetti, M. Vogelsberger, J. Zavala, F. Marinacci, The interplay of self-interacting dark matter and baryons in shaping the halo evolution. Mon. Notices of the Royal Astron. Soc. 484, 4563–4573 (2019). doi:10.1093/mnras/stz273
- G. Despali, S. Vegetti, S.D.M. White, C. Giocoli, F.C. van den Bosch, Modelling the line-of-sight contribution in substructure lensing. Mon. Notices of the Royal Astron. Soc. 475(4), 5424–5442 (2018). doi:10.1093/mnras/sty159
- G. Despali, M. Lovell, S. Vegetti, R.A. Crain, B.D. Oppenheimer, The lensing properties of subhaloes in massive elliptical galaxies in sterile neutrino cosmologies. Mon. Notices of the Royal Astron. Soc. 491(1), 1295–1310 (2020). doi:10.1093/mnras/stz3068
- G. Despali, S. Vegetti, S.D.M. White, D.M. Powell, H.R. Stacey, C.D. Fassnacht, F. Rizzo, W. Enzi, Detecting low-mass haloes with strong gravitational lensing I: the effect of data quality and lensing configuration. Mon. Notices of the Royal Astron. Soc. 510(2), 2480–2494 (2022). doi:10.1093/mnras/stab3537
- B. Dhanasingham, F.-Y. Cyr-Racine, A.H.G. Peter, A. Benson, D. Gilman, Interlopers speak out: studying the dark universe using small-scale lensing anisotropies. Mon. Notices of the Royal Astron. Soc. (2022).

doi:10.1093/mnras/stac2993

- A. Diaz Rivero, C. Dvorkin, Direct detection of dark matter substructure in strong lens images with convolutional neural networks. PRD 101(2), 023515 (2020). doi:10.1103/PhysRevD.101.023515
- A. Diaz Rivero, F.-Y. Cyr-Racine, C. Dvorkin, Power spectrum of dark matter substructure in strong gravitational lenses. PRD 97(2), 023001 (2018). doi:10.1103/PhysRevD.97.023001
- A. Díaz Rivero, C. Dvorkin, F.-Y. Cyr-Racine, J. Zavala, M. Vogelsberger, Gravitational lensing and the power spectrum of dark matter substructure: Insights from the ETHOS N -body simulations. PRD **98**(10), 103517 (2018). doi:10.1103/PhysRevD.98.103517
- P.J. Diggle, R.J. Gratton, Monte Carlo Methods of Inference for Implicit Statistical Models. Journal of the Royal Statistical Society: Series B (Methodological) 46(2), 193– 212 (2018). doi:10.1111/j.2517-6161.1984.tb01290.x
- G. Dobler, C.R. Keeton, Finite source effects in strong lensing: implications for the substructure mass scale. Mon. Notices of the Royal Astron. Soc. 365(4), 1243–1262 (2006). doi:10.1111/j.1365-2966.2005.09809.x
- A.G. Doroshkevich, M.I. Khlopov, R.A. Sunyaev, A.S. Szalay, I.B. Zeldovich, Cosmological impact of the neutrino rest mass. Annals of the New York Academy of Sciences **375**, 32–42 (1981). doi:10.1111/j.1749-6632.1981.tb33688.x
- Y. Dubois, R. Beckmann, F. Bournaud, H. Choi, J. Devriendt, R. Jackson, S. Kaviraj, T. Kimm, K. Kraljic, C. Laigle, G. Martin, M.-J. Park, S. Peirani, C. Pichon, M. Volonteri, S.K. Yi, Introducing the NEWHORIZON simulation: Galaxy properties with resolved internal dynamics across cosmic time. Astron. Astrophys. 651, 109 (2021). doi:10.1051/0004-6361/202039429
- Á. Elíasdóttir, J. Hjorth, S. Toft, I. Burud, D. Paraficz, Extinction Curves of Lensing Galaxies out to z = 1. Astrophys. J. Supplement 166(2), 443–469 (2006). doi:10.1086/507131
- J.D. Ellithorpe, C.S. Kochanek, J.N. Hewitt, Visibility LensClean and the Reliability of Deconvolved Radio Images. Astrophys. J. 464, 556 (1996). doi:10.1086/177346
- W. Enzi, S. Vegetti, G. Despali, J.-W. Hsueh, R.B. Metcalf, Systematic errors in strong gravitational lensing reconstructions, a numerical simulation perspective. Mon. Notices of the Royal Astron. Soc. 496(2), 1718–1729 (2020). doi:10.1093/mnras/staa1224
- W. Enzi, R. Murgia, O. Newton, S. Vegetti, C. Frenk, M. Viel, M. Cautun, C.D. Fassnacht, M. Auger, G. Despali, J. McKean, L.V.E. Koopmans, M. Lovell, Joint constraints on thermal relic dark matter from strong gravitational lensing, the Ly α forest, and Milky Way satellites. Mon. Notices of the Royal Astron. Soc. **506**(4), 5848–5862 (2021). doi:10.1093/mnras/stab1960

- D. Erkal, S.E. Koposov, V. Belokurov, A sharper view of Pal 5's tails: discovery of stream perturbations with a novel non-parametric technique. Mon. Notices of the Royal Astron. Soc. **470**(1), 60–84 (2017). doi:10.1093/mnras/stx1208
- S. Ettori, A. Donnarumma, E. Pointecouteau, T.H. Reiprich, S. Giodini, L. Lovisari, R.W. Schmidt, Mass Profiles of Galaxy Clusters from X-ray Analysis. Space Science Reviews 177(1-4), 119–154 (2013). doi:10.1007/s11214-013-9976-7
- C.D. Fassnacht, R.D. Blandford, J.G. Cohen, K. Matthews, T.J. Pearson, A.C.S. Readhead, D.S. Womble, S.T. Myers, I.W.A. Browne, N.J. Jackson, D.R. Marlow, P.N. Wilkinson, L.V.E. Koopmans, A.G. de Bruyn, R.T. Schilizzi, M. Bremer, G. Miley, B2045+265: A New Four-Image Gravitational Lens from CLASS. Astron. J. **117**(2), 658–670 (1999). doi:10.1086/300724
- C.D. Fassnacht, E. Xanthopoulos, L.V.E. Koopmans, D. Rusin, A Determination of H₀ with the CLASS Gravitational Lens B1608+656. III. A Significant Improvement in the Precision of the Time Delay Measurements. Astrophys. J. **581**(2), 823–835 (2002). doi:10.1086/344368
- J.L. Feng, Dark Matter Candidates from Particle Physics and Methods of Detection. Annual Review of Astronomy and Astrophysics 48, 495–545 (2010). doi:10.1146/annurevastro-082708-101659
- W.-X. Feng, H.-B. Yu, Y.-M. Zhong, Seeding Supermassive Black Holes with Self-interacting Dark Matter: A Unified Scenario with Baryons. Astrophys. J. Letters **914**(2), 26 (2021). doi:10.3847/2041-8213/ac04b0
- P.J. Fox, TASI Lectures on WIMPs and Supersymmetry, in *TASI 2018 - Theory in an Era of Data*, 2018, p. 5. doi:10.22323/1.333.0005
- A. Galan, G. Vernardos, A. Peel, F. Courbin, J.-L. Starck, Using wavelets to capture deviations from smoothness in galaxy-scale strong lenses. Astron. Astrophys. 668, 155 (2022). doi:10.1051/0004-6361/202244464
- S. Garrison-Kimmel, P.F. Hopkins, A. Wetzel, J.S. Bullock, M. Boylan-Kolchin, D. Kereš, C.-A. Faucher-Giguère, K. El-Badry, A. Lamberts, E. Quataert, R. Sanderson, The Local Group on FIRE: dwarf galaxy populations across a suite of hydrodynamic simulations. Mon. Notices of the Royal Astron. Soc. 487(1), 1380–1399 (2019). doi:10.1093/mnras/stz1317
- D. Gilman, S. Birrer, T. Treu, TDCOSMO. III. Dark matter substructure meets dark energy. The effects of (sub)halos on strong-lensing measurements of H_0 . Astron. Astrophys. **642**, 194 (2020). doi:10.1051/0004-6361/202038829
- D. Gilman, Y.-M. Zhong, J. Bovy, Constraining resonant dark matter self-interactions with strong gravitational lenses. arXiv e-prints, 2207–13111 (2022). doi:10.48550/arXiv.2207.13111

- D. Gilman, A. Agnello, T. Treu, C.R. Keeton, A.M. Nierenberg, Strong lensing signatures of luminous structure and substructure in early-type galaxies. Mon. Notices of the Royal Astron. Soc. 467(4), 3970–3992 (2017). doi:10.1093/mnras/stx158
- D. Gilman, S. Birrer, T. Treu, C.R. Keeton, A. Nierenberg, Probing the nature of dark matter by forward modelling flux ratios in strong gravitational lenses. Mon. Notices of the Royal Astron. Soc. 481(1), 819–834 (2018). doi:10.1093/mnras/sty2261
- D. Gilman, S. Birrer, T. Treu, A. Nierenberg, A. Benson, Probing dark matter structure down to 10⁷ solar masses: flux ratio statistics in gravitational lenses with line-ofsight haloes. Mon. Notices of the Royal Astron. Soc. 487(4), 5721–5738 (2019). doi:10.1093/mnras/stz1593
- D. Gilman, X. Du, A. Benson, S. Birrer, A. Nierenberg, T. Treu, Constraints on the mass-concentration relation of cold dark matter halos with 11 strong gravitational lenses. Mon. Notices of the Royal Astron. Soc. **492**(1), 12–16 (2020a). doi:10.1093/mnrasl/slz173
- D. Gilman, S. Birrer, A. Nierenberg, T. Treu, X. Du, A. Benson, Warm dark matter chills out: constraints on the halo mass function and the free-streaming length of dark matter with eight quadruple-image strong gravitational lenses. Mon. Notices of the Royal Astron. Soc. **491**(4), 6077–6101 (2020b). doi:10.1093/mnras/stz3480
- D. Gilman, J. Bovy, T. Treu, A. Nierenberg, S. Birrer, A. Benson, O. Sameie, Strong lensing signatures of self-interacting dark matter in low-mass haloes. Mon. Notices of the Royal Astron. Soc. 507(2), 2432–2447 (2021). doi:10.1093/mnras/stab2335
- D. Gilman, A. Benson, J. Bovy, S. Birrer, T. Treu, A. Nierenberg, The primordial matter power spectrum on subgalactic scales. Mon. Notices of the Royal Astron. Soc. 512(3), 3163–3188 (2022). doi:10.1093/mnras/stac670
- C. Giocoli, L. Pieri, G. Tormen, Analytical approach to subhalo population in dark matter haloes. Mon. Notices of the Royal Astron. Soc. 387, 689–697 (2008). doi:10.1111/j.1365-2966.2008.13283.x
- A.M. Green, B.J. Kavanagh, Primordial black holes as a dark matter candidate. Journal of Physics G Nuclear Physics 48(4), 043001 (2021). doi:10.1088/1361-6471/abc534
- P.J. Green, Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. Biometrika 82(4), 711–732 (1995). doi:10.1093/biomet/82.4.711
- P. Green, Trans-dimensional Markov chain Monte Carlo (Chapter 6), ed. by P. Green, N. Hjort, S. Richardson (Oxford University Press, United Kingdom, 2003), pp. 179– 198
- S.B. Green, F.C. van den Bosch, The tidal evolution of dark matter substructure - I. subhalo density profiles. Mon. No-

tices of the Royal Astron. Soc. **490**(2), 2091–2101 (2019). doi:10.1093/mnras/stz2767

- S.B. Green, F.C. van den Bosch, F. Jiang, The tidal evolution of dark matter substructure - II. The impact of artificial disruption on subhalo mass functions and radial profiles. Mon. Notices of the Royal Astron. Soc. 503(3), 4075– 4091 (2021). doi:10.1093/mnras/stab696
- M. Griffith, G. Langston, M. Heflin, S. Conner, J. Lehar, B. Burke, The Third MIT–Green Bank 5 GHz Survey. Astrophys. J. Supplement 74, 129 (1990). doi:10.1086/191495
- M. Griffith, G. Langston, M. Heflin, S. Conner, B. Burke, The Fourth MIT–Green Bank 5 GHz Survey. Astrophys. J. Supplement 75, 801 (1991). doi:10.1086/191549
- Q. He, J. Nightingale, A. Robertson, A. Amvrosiadis, S. Cole, C.S. Frenk, R. Massey, R. Li, N.C. Amorisco, R.B. Metcalf, X. Cao, A. Etherington, Testing strong lensing subhalo detection with a cosmological simulation. Mon. Notices of the Royal Astron. Soc. **518**(1), 220–239 (2023). doi:10.1093/mnras/stac2779
- C. Heymans, T. Tröster, M. Asgari, C. Blake, H. Hildebrandt, B. Joachimi, K. Kuijken, C.-A. Lin, A.G. Sánchez, J.L. van den Busch, A.H. Wright, A. Amon, M. Bilicki, J. de Jong, M. Crocce, A. Dvornik, T. Erben, M.C. Fortuna, F. Getman, B. Giblin, K. Glazebrook, H. Hoekstra, S. Joudaki, A. Kannawadi, F. Köhlinger, C. Lidman, L. Miller, N.R. Napolitano, D. Parkinson, P. Schneider, H. Shan, E.A. Valentijn, G. Verdoes Kleijn, C. Wolf, KiDS-1000 Cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering constraints. Astron. Astrophys. **646**, 140 (2021). doi:10.1051/0004-6361/202039063
- Y.D. Hezaveh, D.P. Marrone, C.D. Fassnacht, J.S. Spilker, J.D. Vieira, J.E. Aguirre, K.A. Aird, M. Aravena, M.L.N. Ashby, M. Bayliss, B.A. Benson, L.E. Bleem, M. Bothwell, M. Brodwin, J.E. Carlstrom, C.L. Chang, S.C. Chapman, T.M. Crawford, A.T. Crites, C. De Breuck, T. de Haan, M.A. Dobbs, E.B. Fomalont, E.M. George, M.D. Gladders, A.H. Gonzalez, T.R. Greve, N.W. Halverson, F.W. High, G.P. Holder, W.L. Holzapfel, S. Hoover, J.D. Hrubes, K. Husband, T.R. Hunter, R. Keisler, A.T. Lee, E.M. Leitch, M. Lueker, D. Luong-Van, M. Malkan, V. McIntyre, J.J. McMahon, J. Mehl, K.M. Menten, S.S. Meyer, L.M. Mocanu, E.J. Murphy, T. Natoli, S. Padin, T. Plagge, C.L. Reichardt, A. Rest, J. Ruel, J.E. Ruhl, K. Sharon, K.K. Schaffer, L. Shaw, E. Shirokoff, B. Stalder, Z. Staniszewski, A.A. Stark, K. Story, K. Vanderlinde, A. Weiß, N. Welikala, R. Williamson, ALMA Observations of SPT-discovered, Strongly Lensed, Dusty, Star-forming Galaxies. Astrophys. J. 767(2), 132 (2013). doi:10.1088/0004-637X/767/2/132
- Y.D. Hezaveh, L. Perreault Levasseur, P.J. Marshall, Fast automated analysis of strong gravitational lenses with convolutional neural networks. Nature 548(7669), 555–557

(2017). doi:10.1038/nature23463

- Y.D. Hezaveh, N. Dalal, D.P. Marrone, Y.-Y. Mao, W. Morningstar, D. Wen, R.D. Blandford, J.E. Carlstrom, C.D. Fassnacht, G.P. Holder, A. Kemball, P.J. Marshall, N. Murray, L. Perreault Levasseur, J.D. Vieira, R.H. Wechsler, Detection of Lensing Substructure Using ALMA Observations of the Dusty Galaxy SDP.81. Astrophys. J. 823, 37 (2016). doi:10.3847/0004-637X/823/1/37
- Y. Hezaveh, N. Dalal, G. Holder, M. Kuhlen, D. Marrone, N. Murray, J. Vieira, Dark Matter Substructure Detection Using Spatially Resolved Spectroscopy of Lensed Dusty Galaxies. Astrophys. J. **767**(1), 9 (2013). doi:10.1088/0004-637X/767/1/9
- Y. Hezaveh, N. Dalal, G. Holder, T. Kisner, M. Kuhlen, L. Perreault Levasseur, Measuring the power spectrum of dark matter substructure using strong gravitational lensing. J. of Cosm. and Astroparticle Phys. **2016**(11), 048 (2016). doi:10.1088/1475-7516/2016/11/048
- J.A. Högbom, Aperture Synthesis with a Non-Regular Distribution of Interferometer Baselines. Astron. Astrophy. Supplement 15, 417 (1974)
- D.W. Hogg, D. Lang, Telescopes don't make catalogues!, in *EAS Publications Series*. EAS Publications Series, vol. 45, 2010, pp. 351–358. doi:10.1051/eas/1045059
- B.J. Holzschuh, C.M. O'Riordan, S. Vegetti, V. Rodriguez-Gomez, N. Thuerey, Realistic galaxy images and improved robustness in machine learning tasks from generative modelling. Mon. Notices of the Royal Astron. Soc. 515(1), 652–677 (2022). doi:10.1093/mnras/stac1188
- J.-W. Hsueh, C.D. Fassnacht, S. Vegetti, J.P. McKean, C. Spingola, M.W. Auger, L.V.E. Koopmans, D.J. Lagattuta, SHARP - II. Mass structure in strong lenses is not necessarily dark matter substructure: a flux ratio anomaly from an edge-on disc in B1555+375. Mon. Notices of the Royal Astron. Soc. 463(1), 51–55 (2016). doi:10.1093/mnrasl/slw146
- J.-W. Hsueh, L. Oldham, C. Spingola, S. Vegetti, C.D. Fassnacht, M.W. Auger, L.V.E. Koopmans, J.P. McKean, D.J. Lagattuta, SHARP - IV. An apparent flux-ratio anomaly resolved by the edge-on disc in B0712+472. Mon. Notices of the Royal Astron. Soc. 469(3), 3713–3721 (2017). doi:10.1093/mnras/stx1082
- J.-W. Hsueh, W. Enzi, S. Vegetti, M.W. Auger, C.D. Fassnacht, G. Despali, L.V.E. Koopmans, J.P. McKean, SHARP - VII. New constraints on the dark matter free-streaming properties and substructure abundance from gravitationally lensed quasars. Mon. Notices of the Royal Astron. Soc. **492**(2), 3047–3059 (2020). doi:10.1093/mnras/stz3177
- J.-W. Hsueh, G. Despali, S. Vegetti, D. Xu, C.D. Fassnacht, R.B. Metcalf, Flux-ratio anomalies from discs and other baryonic structures in the Illustris simulation. Mon. Notices of the Royal Astron. Soc. 475(2), 2438–2451 (2018).

doi:10.1093/mnras/stx3320

- X. Huang, C. Storfer, A. Gu, V. Ravi, A. Pilon, W. Sheu, R. Venguswamy, S. Banka, A. Dey, M. Landriau, D. Lang, A. Meisner, J. Moustakas, A.D. Myers, R. Sajith, E.F. Schlafly, D.J. Schlegel, Discovering New Strong Gravitational Lenses in the DESI Legacy Imaging Surveys. Astrophys. J. 909(1), 27 (2021). doi:10.3847/1538-4357/abd62b
- L. Hui, Wave Dark Matter. Annual Review of Astronomy and Astrophysics **59**, 247–289 (2021). doi:10.1146/annurev-astro-120920-010024
- K.T. Inoue, T. Minezaki, S. Matsushita, M. Chiba, ALMA imprint of intergalactic dark structures in the gravitational lens SDP.81. Mon. Notices of the Royal Astron. Soc. 457(3), 2936–2950 (2016). doi:10.1093/mnras/stw168
- V. Iršič, M. Viel, M.G. Haehnelt, J.S. Bolton, G.D. Becker, First Constraints on Fuzzy Dark Matter from Lyman- α Forest Data and Hydrodynamical Simulations. PRL **119**(3), 031302 (2017). doi:10.1103/PhysRevLett.119.031302
- M. Kaplinghat, R.E. Keeley, T. Linden, H.-B. Yu, Tying Dark Matter to Baryons with Self-Interactions. Physical Review Letters **113**(2), 021302 (2014). doi:10.1103/PhysRevLett.113.021302
- M. Kaplinghat, T. Ren, H.-B. Yu, Dark matter cores and cusps in spiral galaxies and their explanations. J. of Cosm. and Astroparticle Phys. 2020(6), 027 (2020). doi:10.1088/1475-7516/2020/06/027
- M. Kaplinghat, M. Valli, H.-B. Yu, Too big to fail in light of Gaia. Mon. Notices of the Royal Astron. Soc. 490(1), 231–242 (2019). doi:10.1093/mnras/stz2511
- K. Karchev, A. Coogan, C. Weniger, Strong-lensing source reconstruction with variationally optimized Gaussian processes. Mon. Notices of the Royal Astron. Soc. 512(1), 661–685 (2022). doi:10.1093/mnras/stac311
- H. Kawai, M. Oguri, A. Amruth, T. Broadhurst, J. Lim, An Analytic Model for the Subgalactic Matter Power Spectrum in Fuzzy Dark Matter Halos. Astrophys. J. 925(1), 61 (2022). doi:10.3847/1538-4357/ac39a2
- C.R. Keeton, L.A. Moustakas, A New Channel for Detecting Dark Matter Substructure in Galaxies: Gravitational Lens Time Delays. Astrophys. J. 699(2), 1720–1731 (2009). doi:10.1088/0004-637X/699/2/1720
- A. Knebe, S.R. Knollmann, S.I. Muldrew, F.R. Pearce, M.A. Aragon-Calvo, Y. Ascasibar, P.S. Behroozi, D. Ceverino, et al., Haloes gone MAD: The Halo-Finder Comparison Project. Mon. Notices of the Royal Astron. Soc. 415, 2293–2318 (2011). doi:10.1111/j.1365-2966.2011.18858.x
- A. Knebe, F.R. Pearce, H. Lux, Y. Ascasibar, P. Behroozi, J. Casado, C.C. Moran, J. Diemand, et al., Structure finding in cosmological simulations: the state of affairs. Mon. Notices of the Royal Astron. Soc. 435, 1618–1658 (2013).

doi:10.1093/mnras/stt1403

- C.S. Kochanek, R. Narayan, LensClean: an Algorithm for Inverting Extended, Gravitationally Lensed Images with Application to the Radio Ring Lens PKS 1830-211. Astrophys. J. 401, 461 (1992). doi:10.1086/172078
- L.V.E. Koopmans, Gravitational imaging of cold dark matter substructures. Mon. Notices of the Royal Astron. Soc. **363**(4), 1136–1144 (2005). doi:10.1111/j.1365-2966.2005.09523.x
- L.V.E. Koopmans, A.G. de Bruyn, Microlensing of multiply-imaged compact radio sources. Evidence for compact halo objects in the disk galaxy of B1600+434. Astron. Astrophys. **358**, 793–811 (2000). doi:10.48550/arXiv.astro-ph/0004112
- L.V.E. Koopmans, A. Biggs, R.D. Blandford, I.W.A. Browne, N.J. Jackson, S. Mao, P.N. Wilkinson, A.G. de Bruyn, J. Wambsganss, Extrinsic Radio Variability of JVAS/CLASS Gravitational Lenses. Astrophys. J. 595(2), 712–718 (2003). doi:10.1086/377434
- L.V.E. Koopmans, A. Bolton, T. Treu, O. Czoske, M.W. Auger, M. Barnabè, S. Vegetti, R. Gavazzi, L.A. Moustakas, S. Burles, The Structure and Dynamics of Massive Early-Type Galaxies: On Homology, Isothermality, and Isotropy Inside One Effective Radius. Astrophys. J. Letters **703**(1), 51–54 (2009). doi:10.1088/0004-637X/703/1/L51
- M. Kulkarni, J.P. Ostriker, What is the halo mass function in a fuzzy dark matter cosmology? Mon. Notices of the Royal Astron. Soc. 510(1), 1425–1430 (2022). doi:10.1093/mnras/stab3520
- D.J. Lagattuta, S. Vegetti, C.D. Fassnacht, M.W. Auger, L.V.E. Koopmans, J.P. McKean, SHARP - I. A highresolution multiband view of the infrared Einstein ring of JVAS B1938+666. Mon. Notices of the Royal Astron. Soc. 424(4), 2800–2810 (2012). doi:10.1111/j.1365-2966.2012.21406.x
- G.I. Langston, M.B. Heflin, S.R. Conner, J. Lehar, C.L. Carilli, B.F. Burke, The Second MIT–Green Bank 5 GHz Survey. Astrophys. J. Supplement 72, 621 (1990). doi:10.1086/191427
- F. Lanusse, Q. Ma, N. Li, T.E. Collett, C.-L. Li, S. Ravanbakhsh, R. Mandelbaum, B. Póczos, CMU DeepLens: deep learning for automatic image-based galaxy-galaxy strong lens finding. Mon. Notices of the Royal Astron. Soc. 473(3), 3895–3906 (2018). doi:10.1093/mnras/stx1665
- A. Laroche, D. Gilman, X. Li, J. Bovy, X. Du, Quantum fluctuations masquerade as haloes: bounds on ultra-light dark matter from quadruply imaged quasars. Mon. Notices of the Royal Astron. Soc. 517(2), 1867–1883 (2022). doi:10.1093/mnras/stac2677
- R. Legin, C. Stone, Y. Hezaveh, L. Perreault-Levasseur, Population-Level Inference of Strong Gravita-

tional Lenses with Neural Network-Based Selection Correction. arXiv e-prints, 2207–04123 (2022). doi:10.48550/arXiv.2207.04123

- R. Li, C.S. Frenk, S. Cole, Q. Wang, L. Gao, Projection effects in the strong lensing study of subhaloes. Mon. Notices of the Royal Astron. Soc. 468(2), 1426–1432 (2017). doi:10.1093/mnras/stx554
- J. Liepe, P. Kirk, S. Filippi, T. Toni, C.P. Barnes, M.P.H. Stumpf, A framework for parameter estimation and model selection from experimental data in systems biology using approximate bayesian computation. Nature protocols 9, 439–456 (2014). doi:10.1038/nprot.2014.025
- N. Loudas, V. Pavlidou, C. Casadio, K. Tassis, Discriminating power of milli-lensing observations for dark matter models. Astron. Astrophys. 668, 166 (2022). doi:10.1051/0004-6361/202244978
- M.R. Lovell, The halo mass function in alternative dark matter models. Mon. Notices of the Royal Astron. Soc. 493(1), 11–15 (2020). doi:10.1093/mnrasl/slaa005
- M.R. Lovell, V. Eke, C.S. Frenk, L. Gao, A. Jenkins, T. Theuns, J. Wang, S.D.M. White, A. Boyarsky, O. Ruchayskiy, The haloes of bright satellite galaxies in a warm dark matter universe. Mon. Notices of the Royal Astron. Soc. **420**, 2318–2324 (2012). doi:10.1111/j.1365-2966.2011.20200.x
- M.R. Lovell, Toward a General Parameterization of the Warm Dark Matter Halo Mass Function. Astrophys. J. 897(2), 147 (2020). doi:10.3847/1538-4357/ab982a
- M.R. Lovell, J. Zavala, M. Vogelsberger, ETHOS an effective theory of structure formation: formation of the first haloes and their stars. Mon. Notices of the Royal Astron. Soc. **485**(4), 5474–5489 (2019). doi:10.1093/mnras/stz766
- M.R. Lovell, C.S. Frenk, V.R. Eke, A. Jenkins, L. Gao, T. Theuns, The properties of warm dark matter haloes. Mon. Notices of the Royal Astron. Soc. 439(1), 300–317 (2014). doi:10.1093/mnras/stt2431
- A.D. Ludlow, S. Bose, R.E. Angulo, L. Wang, W.A. Hellwing, J.F. Navarro, S. Cole, C.S. Frenk, The massconcentration-redshift relation of cold and warm dark matter haloes. Mon. Notices of the Royal Astron. Soc. 460, 1214–1232 (2016). doi:10.1093/mnras/stw1046
- R. Mandelbaum, U. Seljak, G. Kauffmann, C.M. Hirata, J. Brinkmann, Galaxy halo masses and satellite fractions from galaxy-galaxy lensing in the Sloan Digital Sky Survey: stellar mass, luminosity, morphology and environment dependencies. Mon. Notices of the Royal Astron. Soc. 368(2), 715–731 (2006). doi:10.1111/j.1365-2966.2006.10156.x
- S. Mao, P. Schneider, Evidence for substructure in lens galaxies? Mon. Notices of the Royal Astron. Soc. 295(3), 587–594 (1998). doi:10.1046/j.1365-8711.1998.01319.x
- S. Mao, Y. Jing, J.P. Ostriker, J. Weller, Anomalous Flux

Ratios in Gravitational Lenses: For or against Cold Dark Matter? Astrophys. J. Letters **604**(1), 5–8 (2004). doi:10.1086/383413

- D.R. Marlow, S.T. Myers, D. Rusin, N. Jackson, I.W.A. Browne, P.N. Wilkinson, T. Muxlow, C.D. Fassnacht, L. Lubin, T. Kundić, R.D. Blandford, T.J. Pearson, A.C.S. Readhead, L. Koopmans, A.G. de Bruyn, CLASS B1555+375: A New Four-Image Gravitational Lens System. Astron. J. **118**(2), 654–658 (1999). doi:10.1086/300987
- D.J.E. Marsh, J.C. Niemeyer, Strong Constraints on Fuzzy Dark Matter from Ultrafaint Dwarf Galaxy Eridanus II. PRL **123**(5), 051103 (2019). doi:10.1103/PhysRevLett.123.051103
- D.J.E. Marsh, J. Silk, A model for halo formation with axion mixed dark matter. Mon. Notices of the Royal Astron. Soc. **437**(3), 2652–2663 (2014). doi:10.1093/mnras/stt2079
- S. May, V. Springel, Structure formation in large-volume cosmological simulations of fuzzy dark matter: impact of the non-linear dynamics. Mon. Notices of the Royal Astron. Soc. 506(2), 2603–2618 (2021). doi:10.1093/mnras/stab1764
- S. May, V. Springel, The halo mass function and filaments in full cosmological simulations with fuzzy dark matter. arXiv e-prints, 2209–14886 (2022). doi:10.48550/arXiv.2209.14886
- J.P. McKean, L.V.E. Koopmans, C.E. Flack, C.D. Fassnacht, D. Thompson, K. Matthews, R.D. Blandford, A.C.S. Readhead, B.T. Soifer, High-resolution imaging of the anomalous flux ratio gravitational lens system CLASS B2045+265: dark or luminous satellites? Mon. Notices of the Royal Astron. Soc. **378**(1), 109–118 (2007). doi:10.1111/j.1365-2966.2007.11744.x
- J. McKean, N. Jackson, S. Vegetti, M. Rybak, S. Serjeant, L.V.E. Koopmans, R.B. Metcalf, C. Fassnacht, P.J. Marshall, M. Pandey-Pommier, Strong Gravitational Lensing with the SKA, in *Advancing Astrophysics with the Square Kilometre Array (AASKA14)*, 2015, p. 84
- R.B. Metcalf, L.A. Moustakas, A.J. Bunker, I.R. Parry, Spectroscopic Gravitational Lensing and Limits on the Dark Matter Substructure in Q2237+0305. Astrophys. J. 607(1), 43–59 (2004). doi:10.1086/383243
- Q. Minor, S. Gad-Nasr, M. Kaplinghat, S. Vegetti, An unexpected high concentration for the dark substructure in the gravitational lens SDSSJ0946+1006. Mon. Notices of the Royal Astron. Soc. 507(2), 1662–1683 (2021). doi:10.1093/mnras/stab2247
- R. Mittal, R. Porcas, O. Wucknitz, Free-free absorption in the gravitational lens JVAS B0218+357. Astron. Astrophys. 465(2), 405–415 (2007). doi:10.1051/0004-6361:20066127
- P. Mocz, A. Fialkov, M. Vogelsberger, F. Becerra, M.A.

Amin, S. Bose, M. Boylan-Kolchin, P.-H. Chavanis, L. Hernquist, L. Lancaster, F. Marinacci, V.H. Robles, J. Zavala, First Star-Forming Structures in Fuzzy Cosmic Filaments. PRL **123**(14), 141301 (2019). doi:10.1103/PhysRevLett.123.141301

- Á. Moliné, M.A. Sánchez-Conde, S. Palomares-Ruiz, F. Prada, Characterization of subhalo structural properties and implications for dark matter annihilation signals. Mon. Notices of the Royal Astron. Soc. 466, 4974–4990 (2017). doi:10.1093/mnras/stx026
- N.A. Montel, A. Coogan, C. Correa, K. Karchev, C. Weniger, Estimating the warm dark matter mass from strong lensing images with truncated marginal neural ratio estimation. Mon. Notices of the Royal Astron. Soc. (2022). doi:10.1093/mnras/stac3215
- L.A. Moustakas, R.B. Metcalf, Detecting dark matter substructure spectroscopically in strong gravitational lenses. Mon. Notices of the Royal Astron. Soc. **339**(3), 607–615 (2003). doi:10.1046/j.1365-8711.2003.06055.x
- S. Mukherjee, L.V.E. Koopmans, R.B. Metcalf, C. Tortora, M. Schaller, J. Schaye, G. Vernardos, F. Bellagamba, SEAGLE - II. Constraints on feedback models in galaxy formation from massive early-type strong-lens galaxies. Mon. Notices of the Royal Astron. Soc. **504**(3), 3455– 3477 (2021). doi:10.1093/mnras/stab693
- F. Müller-Sánchez, M.A. Prieto, E.K.S. Hicks, H. Vives-Arias, R.I. Davies, M. Malkan, L.J. Tacconi, R. Genzel, Outflows from Active Galactic Nuclei: Kinematics of the Narrow-line and Coronal-line Regions in Seyfert Galaxies. Astrophys. J. **739**(2), 69 (2011). doi:10.1088/0004-637X/739/2/69
- S.T. Myers, N.J. Jackson, I.W.A. Browne, A.G. de Bruyn, T.J. Pearson, A.C.S. Readhead, P.N. Wilkinson, A.D. Biggs, R.D. Blandford, C.D. Fassnacht, L.V.E. Koopmans, D.R. Marlow, J.P. McKean, M.A. Norbury, P.M. Phillips, D. Rusin, M.C. Shepherd, C.M. Sykes, The Cosmic Lens All-Sky Survey - I. Source selection and observations. Mon. Notices of the Royal Astron. Soc. 341, 1–12 (2003). doi:10.1046/j.1365-8711.2003.06256.x
- E.O. Nadler, A. Drlica-Wagner, K. Bechtol, S. Mau, R.H. Wechsler, V. Gluscevic, K. Boddy, A.B. Pace, T.S. Li, M. McNanna, A.H. Riley, J. García-Bellido, Y.-Y. Mao, G. Green, D.L. Burke, A. Peter, B. Jain, T.M.C. Abbott, M. Aguena, S. Allam, J. Annis, S. Avila, D. Brooks, M. Carrasco Kind, J. Carretero, M. Costanzi, L.N. da Costa, J. De Vicente, S. Desai, H.T. Diehl, P. Doel, S. Everett, A.E. Evrard, B. Flaugher, J. Frieman, D.W. Gerdes, D. Gruen, R.A. Gruendl, J. Gschwend, G. Gutierrez, S.R. Hinton, K. Honscheid, D. Huterer, D.J. James, E. Krause, K. Kuehn, N. Kuropatkin, O. Lahav, M.A.G. Maia, J.L. Marshall, F. Menanteau, R. Miquel, A. Palmese, F. Paz-Chinchón, A.A. Plazas, A.K. Romer, E. Sanchez, V. Scarpine, S. Serrano, I. Sevilla-Noarbe, M. Smith, M.

Soares-Santos, E. Suchyta, M.E.C. Swanson, G. Tarle, D.L. Tucker, A.R. Walker, W. Wester, Constraints on dark matter properties from observations of milky way satellite galaxies. Phys. Rev. Lett. **126**, 091101 (2021). doi:10.1103/PhysRevLett.126.091101

- E.O. Nadler, A. Banerjee, S. Adhikari, Y.-Y. Mao, R.H. Wechsler, Signatures of Velocity-dependent Dark Matter Self-interactions in Milky Way-mass Halos. Astrophys. J. 896(2), 112 (2020). doi:10.3847/1538-4357/ab94b0
- E.O. Nadler, S. Birrer, D. Gilman, R.H. Wechsler, X. Du, A. Benson, A.M. Nierenberg, T. Treu, Dark Matter Constraints from a Unified Analysis of Strong Gravitational Lenses and Milky Way Satellite Galaxies. Astrophys. J. 917(1), 7 (2021). doi:10.3847/1538-4357/abf9a3
- J.F. Navarro, C.S. Frenk, S.D.M. White, The Structure of Cold Dark Matter Halos. Astrophys. J. 462, 563 (1996). doi:10.1086/177173
- A.M. Nierenberg, T. Treu, S.A. Wright, C.D. Fassnacht, M.W. Auger, Detection of substructure with adaptive optics integral field spectroscopy of the gravitational lens B1422+231. Mon. Notices of the Royal Astron. Soc. 442(3), 2434–2445 (2014). doi:10.1093/mnras/stu862
- A.M. Nierenberg, T. Treu, G. Brammer, A.H.G. Peter, C.D. Fassnacht, C.R. Keeton, C.S. Kochanek, K.B. Schmidt, D. Sluse, S.A. Wright, Probing dark matter substructure in the gravitational lens HE 0435-1223 with the WFC3 grism. Mon. Notices of the Royal Astron. Soc. 471(2), 2224–2236 (2017). doi:10.1093/mnras/stx1400
- A.M. Nierenberg, D. Gilman, T. Treu, G. Brammer, S. Birrer, L. Moustakas, A. Agnello, T. Anguita, C.D. Fassnacht, V. Motta, A.H.G. Peter, D. Sluse, Double dark matter vision: twice the number of compact-source lenses with narrow-line lensing and the WFC3 grism. Mon. Notices of the Royal Astron. Soc. 492(4), 5314–5335 (2020). doi:10.1093/mnras/stz358810.48550/arXiv.1908.06344
- A. Nierenberg, V.N. Bennert, A. Benson, S. Birrer, S.G. Djorgovski, X. Du, C. Fassnacht, D. Gilman, S.F. Hoenig, A. Kusenko, M.A. Malkan, V. Motta, L.A. Moustakas, D. Sluse, D.K. Stern, T.L. Treu, A definitive test of the dark matter paradigm on small scales, JWST Proposal. Cycle 1, ID. #2046, 2021
- J.W. Nightingale, S. Dye, Adaptive semi-linear inversion of strong gravitational lens imaging. Mon. Notices of the Royal Astron. Soc. 452(3), 2940–2959 (2015). doi:10.1093/mnras/stv1455
- J.W. Nightingale, Q. He, X. Cao, A. Amvrosiadis, A. Etherington, C.S. Frenk, R.G. Hayes, A. Robertson, S. Cole, S. Lange, R. Li, R. Massey, Scanning For Dark Matter Subhalos in Hubble Space Telescope Imaging of 54 Strong Lenses. arXiv e-prints, 2209–10566 (2022)
- H. Nishikawa, K.K. Boddy, M. Kaplinghat, Accelerated core collapse in tidally stripped self-interacting dark matter halos. Physical Review D **101**(6), 063009 (2020)

- M. Oguri, P.J. Marshall, Gravitationally lensed quasars and supernovae in future wide-field optical imaging surveys.
 Mon. Notices of the Royal Astron. Soc. 405(4), 2579– 2593 (2010). doi:10.1111/j.1365-2966.2010.16639.x
- N. Okabe, G.P. Smith, K. Umetsu, M. Takada, T. Futamase, LoCuSS: The Mass Density Profile of Massive Galaxy Clusters at z = 0.2. Astrophys. J. Letters **769**, 35 (2013). doi:10.1088/2041-8205/769/2/L35
- C.M. O'Riordan, G. Despali, S. Vegetti, M.R. Lovell, Á. Moliné, Sensitivity of strong lensing observations to dark matter substructure: a case study with Euclid. Mon. Notices of the Royal Astron. Soc. 521(2), 2342–2356 (2023). doi:10.1093/mnras/stad650
- B. Ostdiek, A. Diaz Rivero, C. Dvorkin, Extracting the Subhalo Mass Function from Strong Lens Images with Image Segmentation. arXiv e-prints, 2009–06639 (2020)
- B. Ostdiek, A. Diaz Rivero, C. Dvorkin, Image segmentation for analyzing galaxy-galaxy strong lensing systems. Astron. Astrophys. 657, 14 (2022). doi:10.1051/0004-6361/202142030
- A.R. Patnaik, I.W.A. Browne, P.N. Wilkinson, J.M. Wrobel, Interferometer phase calibration sources - I. The region 35deg <= Dec. <= 75deg. Mon. Notices of the Royal Astron. Soc. 254, 655 (1992). doi:10.1093/mnras/254.4.655
- T.J. Pearson, A.C.S. Readhead, Image Formation by Self-Calibration in Radio Astronomy. Annual Review of Astronomy and Astrophysics **22**, 97–130 (1984). doi:10.1146/annurev.aa.22.090184.000525
- L. Perreault Levasseur, Y.D. Hezaveh, R.H. Wechsler, Uncertainties in Parameters Estimated with Neural Networks: Application to Strong Gravitational Lensing. Astrophys. J. Letters 850(1), 7 (2017). doi:10.3847/2041-8213/aa9704
- C.E. Petrillo, C. Tortora, G. Vernardos, L.V.E. Koopmans, G. Verdoes Kleijn, M. Bilicki, N.R. Napolitano, S. Chatterjee, G. Covone, A. Dvornik, T. Erben, F. Getman, B. Giblin, C. Heymans, J.T.A. de Jong, K. Kuijken, P. Schneider, H. Shan, C. Spiniello, A.H. Wright, LinKS: discovering galaxy-scale strong lenses in the Kilo-Degree Survey using convolutional neural networks. Mon. Notices of the Royal Astron. Soc. **484**(3), 3879–3896 (2019). doi:10.1093/mnras/stz189
- A. Pillepich, V. Springel, D. Nelson, S. Genel, J. Naiman, R. Pakmor, L. Hernquist, P. Torrey, M. Vogelsberger, R. Weinberger, F. Marinacci, Simulating galaxy formation with the IllustrisTNG model. Mon. Notices of the Royal Astron. Soc. **473**(3), 4077–4106 (2018). doi:10.1093/mnras/stx2656
- Planck Collaboration, N. Aghanim, Y. Akrami, F. Arroja, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A.J. Banday, R.B. Barreiro, N. Bartolo, S. Basak, R. Battye, K. Benabed, J.-P. Bernard, M. Bersanelli, P. Bielewicz, J.J. Bock, J.R. Bond, J. Borrill, F.R. Bouchet, F. Boulanger,

M. Bucher, C. Burigana, R.C. Butler, E. Calabrese, J.-F. Cardoso, J. Carron, B. Casaponsa, A. Challinor, H.C. Chiang, L.P.L. Colombo, C. Combet, D. Contreras, B.P. Crill, F. Cuttaia, P. de Bernardis, G. de Zotti, J. Delabrouille, J.-M. Delouis, F.-X. Désert, E. Di Valentino, C. Dickinson, J.M. Diego, S. Donzelli, O. Doré, M. Douspis, A. Ducout, X. Dupac, G. Efstathiou, F. Elsner, T.A. Enßlin, H.K. Eriksen, E. Falgarone, Y. Fantaye, J. Fergusson, R. Fernandez-Cobos, F. Finelli, F. Forastieri, M. Frailis, E. Franceschi, A. Frolov, S. Galeotta, S. Galli, K. Ganga, R.T. Génova-Santos, M. Gerbino, T. Ghosh, J. González-Nuevo, K.M. Górski, S. Gratton, A. Gruppuso, J.E. Gudmundsson, J. Hamann, W. Handley, F.K. Hansen, G. Helou, D. Herranz, S.R. Hildebrandt, E. Hivon, Z. Huang, A.H. Jaffe, W.C. Jones, A. Karakci, E. Keihänen, R. Keskitalo, K. Kiiveri, J. Kim, T.S. Kisner, L. Knox, N. Krachmalnicoff, M. Kunz, H. Kurki-Suonio, G. Lagache, J.-M. Lamarre, M. Langer, A. Lasenby, M. Lattanzi, C.R. Lawrence, M. Le Jeune, J.P. Leahy, J. Lesgourgues, F. Levrier, A. Lewis, M. Liguori, P.B. Lilje, M. Lilley, V. Lindholm, M. López-Caniego, P.M. Lubin, Y.-Z. Ma, J.F. Macías-Pérez, G. Maggio, D. Maino, N. Mandolesi, A. Mangilli, A. Marcos-Caballero, M. Maris, P.G. Martin, M. Martinelli, E. Martínez-González, S. Matarrese, N. Mauri, J.D. McEwen, P.D. Meerburg, P.R. Meinhold, A. Melchiorri, A. Mennella, M. Migliaccio, M. Millea, S. Mitra, M.-A. Miville-Deschênes, D. Molinari, A. Moneti, L. Montier, G. Morgante, A. Moss, S. Mottet, M. Münchmeyer, P. Natoli, H.U. Nørgaard-Nielsen, C.A. Oxborrow, L. Pagano, D. Paoletti, B. Partridge, G. Patanchon, T.J. Pearson, M. Peel, H.V. Peiris, F. Perrotta, V. Pettorino, F. Piacentini, L. Polastri, G. Polenta, J.-L. Puget, J.P. Rachen, M. Reinecke, M. Remazeilles, C. Renault, A. Renzi, G. Rocha, C. Rosset, G. Roudier, J.A. Rubiño-Martín, B. Ruiz-Granados, L. Salvati, M. Sandri, M. Savelainen, D. Scott, E.P.S. Shellard, M. Shiraishi, C. Sirignano, G. Sirri, L.D. Spencer, R. Sunyaev, A.-S. Suur-Uski, J.A. Tauber, D. Tavagnacco, M. Tenti, L. Terenzi, L. Toffolatti, M. Tomasi, T. Trombetti, J. Valiviita, B. Van Tent, L. Vibert, P. Vielva, F. Villa, N. Vittorio, B.D. Wandelt, I.K. Wehus, M. White, S.D.M. White, A. Zacchei, A. Zonca, Planck 2018 results. I. Overview and the cosmological legacy of Planck. Astron. Astrophys. 641, 1 (2020a). doi:10.1051/0004-6361/201833880

Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A.J. Banday, R.B. Barreiro, N. Bartolo, S. Basak, R. Battye, K. Benabed, J.-P. Bernard, M. Bersanelli, P. Bielewicz, J.J. Bock, J.R. Bond, J. Borrill, F.R. Bouchet, F. Boulanger, M. Bucher, C. Burigana, R.C. Butler, E. Calabrese, J.-F. Cardoso, J. Carron, A. Challinor, H.C. Chiang, J. Chluba, L.P.L. Colombo, C. Combet, D. Contreras, B.P. Crill, F. Cuttaia, P. de Bernardis, G. de Zotti, J. Delabrouille,

J.-M. Delouis, E. Di Valentino, J.M. Diego, O. Doré, M. Douspis, A. Ducout, X. Dupac, S. Dusini, G. Efstathiou, F. Elsner, T.A. Enßlin, H.K. Eriksen, Y. Fantaye, M. Farhang, J. Fergusson, R. Fernandez-Cobos, F. Finelli, F. Forastieri, M. Frailis, A.A. Fraisse, E. Franceschi, A. Frolov, S. Galeotta, S. Galli, K. Ganga, R.T. Génova-Santos, M. Gerbino, T. Ghosh, J. González-Nuevo, K.M. Górski, S. Gratton, A. Gruppuso, J.E. Gudmundsson, J. Hamann, W. Handley, F.K. Hansen, D. Herranz, S.R. Hildebrandt, E. Hivon, Z. Huang, A.H. Jaffe, W.C. Jones, A. Karakci, E. Keihänen, R. Keskitalo, K. Kiiveri, J. Kim, T.S. Kisner, L. Knox, N. Krachmalnicoff, M. Kunz, H. Kurki-Suonio, G. Lagache, J.-M. Lamarre, A. Lasenby, M. Lattanzi, C.R. Lawrence, M. Le Jeune, P. Lemos, J. Lesgourgues, F. Levrier, A. Lewis, M. Liguori, P.B. Lilje, M. Lilley, V. Lindholm, M. López-Caniego, P.M. Lubin, Y.-Z. Ma, J.F. Macías-Pérez, G. Maggio, D. Maino, N. Mandolesi, A. Mangilli, A. Marcos-Caballero, M. Maris, P.G. Martin, M. Martinelli, E. Martínez-González, S. Matarrese, N. Mauri, J.D. McEwen, P.R. Meinhold, A. Melchiorri, A. Mennella, M. Migliaccio, M. Millea, S. Mitra, M.-A. Miville-Deschênes, D. Molinari, L. Montier, G. Morgante, A. Moss, P. Natoli, H.U. Nørgaard-Nielsen, L. Pagano, D. Paoletti, B. Partridge, G. Patanchon, H.V. Peiris, F. Perrotta, V. Pettorino, F. Piacentini, L. Polastri, G. Polenta, J.-L. Puget, J.P. Rachen, M. Reinecke, M. Remazeilles, A. Renzi, G. Rocha, C. Rosset, G. Roudier, J.A. Rubiño-Martín, B. Ruiz-Granados, L. Salvati, M. Sandri, M. Savelainen, D. Scott, E.P.S. Shellard, C. Sirignano, G. Sirri, L.D. Spencer, R. Sunyaev, A.-S. Suur-Uski, J.A. Tauber, D. Tavagnacco, M. Tenti, L. Toffolatti, M. Tomasi, T. Trombetti, L. Valenziano, J. Valiviita, B. Van Tent, L. Vibert, P. Vielva, F. Villa, N. Vittorio, B.D. Wandelt, I.K. Wehus, M. White, S.D.M. White, A. Zacchei, A. Zonca, Planck 2018 results. VI. Cosmological parameters. Astron. Astrophys. 641, 6 (2020b). doi:10.1051/0004-6361/201833910

- S.K.N. Portillo, B.C.G. Lee, T. Daylan, D.P. Finkbeiner, Improved Point-source Detection in Crowded Fields Using Probabilistic Cataloging. Astron. J. **154**(4), 132 (2017). doi:10.3847/1538-3881/aa8565
- D.M. Powell, S. Vegetti, J.P. McKean, C. Spingola, H.R. Stacey, C.D. Fassnacht, A lensed radio jet at milliarcsecond resolution I: Bayesian comparison of parametric lens models. Mon. Notices of the Royal Astron. Soc. 516(2), 1808–1828 (2022). doi:10.1093/mnras/stac2350
- D.M. Powell, S. Vegetti, J.P. McKean, S.D.M. White, E.G.M. Ferreira, S. May, C. Spingola, A lensed radio jet at milli-arcsecond resolution II: Constraints on fuzzy dark matter from an extended gravitational arc. arXiv e-prints, 2302–10941 (2023). doi:10.48550/arXiv.2302.10941
- D. Powell, S. Vegetti, J.P. McKean, C. Spingola, F. Rizzo, H.R. Stacey, A novel approach to visibility-

space modelling of interferometric gravitational lens observations at high angular resolution. Mon. Notices of the Royal Astron. Soc. **501**(1), 515–530 (2021). doi:10.1093/mnras/staa2740

- W.H. Press, P. Schechter, Formation of Galaxies and Clusters of Galaxies by Self-Similar Gravitational Condensation 187, 425–438 (1974). doi:10.1086/152650
- W.H. Press, S.A. Teukolsky, W.T. Vetterling, B.P. Flannery, *Numerical recipes in C. The art of scientific computing* 1992
- S. Rezaei, J.P. McKean, M. Biehl, W. de Roo, A. Lafontaine, A machine learning based approach to gravitational lens identification with the International LOFAR Telescope. Mon. Notices of the Royal Astron. Soc. 517(1), 1156– 1170 (2022). doi:10.1093/mnras/stac2078
- T.R.G. Richardson, J. Stücker, R.E. Angulo, O. Hahn, Nonhalo structures and their effects on gravitational lensing. Mon. Notices of the Royal Astron. Soc. 511(4), 6019– 6032 (2022). doi:10.1093/mnras/stac493
- E. Ritondale, S. Vegetti, G. Despali, M.W. Auger, L.V.E. Koopmans, J.P. McKean, Low-mass halo perturbations in strong gravitational lenses at redshift z ~ 0.5 are consistent with CDM. Mon. Notices of the Royal Astron. Soc. 485(2), 2179–2193 (2019a). doi:10.1093/mnras/stz464
- E. Ritondale, M.W. Auger, S. Vegetti, J.P. McKean, Resolving on 100 pc scales the UV-continuum in Lyman-α emitters between redshift 2 and 3 with gravitational lensing. Mon. Notices of the Royal Astron. Soc. **482**(4), 4744–4762 (2019b). doi:10.1093/mnras/sty2833
- F. Rizzo, S. Vegetti, F. Fraternali, E. Di Teodoro, A novel 3D technique to study the kinematics of lensed galaxies. Mon. Notices of the Royal Astron. Soc. 481(4), 5606– 5629 (2018). doi:10.1093/mnras/sty2594
- A. Robertson, R. Massey, V. Eke, S. Tulin, H.-B. Yu, Y. Bahé, D.J. Barnes, R.G. Bower, R.A. Crain, C. Dalla Vecchia, S.T. Kay, M. Schaller, J. Schaye, The diverse density profiles of galaxy clusters with self-interacting dark matter plus baryons. Mon. Notices of the Royal Astron. Soc. 476, 20–24 (2018). doi:10.1093/mnrasl/sly024
- A. Robertson, D. Harvey, R. Massey, V. Eke, I.G. Mc-Carthy, M. Jauzac, B. Li, J. Schaye, Observable tests of self-interacting dark matter in galaxy clusters: cosmological simulations with SIDM and baryons. Mon. Notices of the Royal Astron. Soc. 488(3), 3646–3662 (2019). doi:10.1093/mnras/stz1815
- A. Robertson, R. Massey, V. Eke, J. Schaye, T. Theuns, The surprising accuracy of isothermal Jeans modelling of self-interacting dark matter density profiles. Mon. Notices of the Royal Astron. Soc. 501(3), 4610–4634 (2021). doi:10.1093/mnras/staa3954
- M. Rocha, A.H.G. Peter, J.S. Bullock, M. Kaplinghat, S. Garrison-Kimmel, J. Oñorbe, L.A. Moustakas, Cosmological simulations with self-interacting dark matter -

I. Constant-density cores and substructure. Mon. Notices of the Royal Astron. Soc. **430**, 81–104 (2013). doi:10.1093/mnras/sts514

- D.B. Rubin, Bayesianly justifiable and relevant frequency calculations for the applied statistician. The Annals of Statistics **12** (1984). doi:10.1214/aos/1176346785
- V.C. Rubin, Evidence for dark matter from rotation curves: Ten years later, in *After the first three minutes*, ed. by S.S. Holt, C.L. Bennett, V. Trimble American Institute of Physics Conference Series, vol. 222, 1991, pp. 371–380. doi:10.1063/1.40398
- M. Rybak, J.P. McKean, S. Vegetti, P. Andreani, S.D.M. White, ALMA imaging of SDP.81 - I. A pixelated reconstruction of the far-infrared continuum emission. Mon. Notices of the Royal Astron. Soc. 451, 40–44 (2015). doi:10.1093/mnrasl/slv058
- L. Sagunski, S. Gad-Nasr, B. Colquhoun, A. Robertson, S. Tulin, Velocity-dependent self-interacting dark matter from groups and clusters of galaxies. J. of Cosm. and Astroparticle Phys. **2021**(1), 024 (2021). doi:10.1088/1475-7516/2021/01/024
- O. Sameie, P. Creasey, H.-B. Yu, L.V. Sales, M. Vogelsberger, J. Zavala, The impact of baryonic discs on the shapes and profiles of self-interacting dark matter haloes. Mon. Notices of the Royal Astron. Soc. 479, 359–367 (2018). doi:10.1093/mnras/sty1516
- O. Sameie, H.-B. Yu, L.V. Sales, M. Vogelsberger, J. Zavala, Self-Interacting Dark Matter Subhalos in the Milky Way's Tides. PRL **124**(14), 141102 (2020). doi:10.1103/PhysRevLett.124.141102
- T. Sawala, C.S. Frenk, A. Fattahi, J.F. Navarro, R.G. Bower, R.A. Crain, C. Dalla Vecchia, M. Furlong, A. Jenkins, I.G. McCarthy, Y. Qu, M. Schaller, J. Schaye, T. Theuns, Bent by baryons: the low-mass galaxy-halo relation. Mon. Notices of the Royal Astron. Soc. 448, 2941–2947 (2015). doi:10.1093/mnras/stu2753
- T. Sawala, P. Pihajoki, P.H. Johansson, C.S. Frenk, J.F. Navarro, K.A. Oman, S.D.M. White, Shaken and stirred: the Milky Way's dark substructures. Mon. Notices of the Royal Astron. Soc. 467, 4383–4400 (2017). doi:10.1093/mnras/stx360
- J. Schaye, R.A. Crain, R.G. Bower, M. Furlong, M. Schaller, T. Theuns, C. Dalla Vecchia, C.S.e.a. Frenk, The EAGLE project: simulating the evolution and assembly of galaxies and their environments. Mon. Notices of the Royal Astron. Soc. 446, 521–554 (2015). doi:10.1093/mnras/stu2058
- H.-Y. Schive, T. Chiueh, T. Broadhurst, Cosmic structure as the quantum interference of a coherent dark wave. Nature Physics 10(7), 496–499 (2014a). doi:10.1038/nphys2996
- H.-Y. Schive, M.-H. Liao, T.-P. Woo, S.-K. Wong, T. Chiueh, T. Broadhurst, W.-Y.P. Hwang, Understanding the Core-Halo Relation of Quantum Wave Dark Mat-

ter from 3D Simulations. PRL **113**(26), 261302 (2014b). doi:10.1103/PhysRevLett.113.261302

- H.-Y. Schive, T. Chiueh, T. Broadhurst, K.-W. Huang, Contrasting Galaxy Formation from Quantum Wave Dark Matter, ψDM, with ΛCDM, using Planck and Hubble Data. Astrophys. J. 818(1), 89 (2016). doi:10.3847/0004-637X/818/1/89
- A. Schneider, R.E. Smith, A.V. Macciò, B. Moore, Nonlinear evolution of cosmological structures in warm dark matter models. Mon. Notices of the Royal Astron. Soc. 424, 684–698 (2012). doi:10.1111/j.1365-2966.2012.21252.x
- A. Schneider, Structure formation with suppressed small-scale perturbations. Mon. Notices of the Royal Astron. Soc. 451(3), 3117–3130 (2015). doi:10.1093/mnras/stv1169
- S. Schuldt, S.H. Suyu, T. Meinhardt, L. Leal-Taixé, R. Cañameras, S. Taubenberger, A. Halkola, HOLISMOKES. IV. Efficient mass modeling of strong lenses through deep learning. Astron. Astrophys. 646, 126 (2021). doi:10.1051/0004-6361/202039574
- A.Ç. Sengül, C. Dvorkin, B. Ostdiek, A. Tsang, Substructure detection reanalysed: dark perturber shown to be a line-of-sight halo. Mon. Notices of the Royal Astron. Soc. 515(3), 4391–4401 (2022). doi:10.1093/mnras/stac1967
- A.J. Shajib, G. Vernardos, T.E. Collett, V. Motta, D. Sluse, L.L.R. Williams, P. Saha, S. Birrer, C. Spiniello, T. Treu, Strong Lensing by Galaxies. arXiv e-prints, 2210–10790 (2022). doi:10.48550/arXiv.2210.10790
- X. Shen, T. Brinckmann, D. Rapetti, M. Vogelsberger, A. Mantz, J. Zavala, S.W. Allen, X-ray morphology of cluster-mass haloes in self-interacting dark matter. Mon. Notices of the Royal Astron. Soc. 516(1), 1302–1319 (2022). doi:10.1093/mnras/stac2376
- R.K. Sheth, H.J. Mo, G. Tormen, Ellipsoidal collapse and an improved model for the number and spatial distribution of dark matter haloes. Mon. Notices of the Royal Astron. Soc. **323**, 1–12 (2001). doi:10.1046/j.1365-8711.2001.04006.x
- R.K. Sheth, G. Lemson, The forest of merger history trees associated with the formation of dark matter haloes. Mon. Notices of the Royal Astron. Soc. **305**(4), 946–956 (1999). doi:10.1046/j.1365-8711.1999.02477.x
- R.K. Sheth, G. Tormen, Large-scale bias and the peak background split. Mon. Notices of the Royal Astron. Soc. **308**(1), 119–126 (1999). doi:10.1046/j.1365-8711.1999.02692.x
- Y. Shu, A.S. Bolton, S. Mao, C.S. Kochanek, I. Pérez-Fournon, M. Oguri, A.D. Montero-Dorta, M.A. Cornachione, R. Marques-Chaves, Z. Zheng, J.R. Brownstein, B. Ménard, The BOSS Emission-line Lens Survey. IV. Smooth Lens Models for the BELLS GALLERY Sample. Astrophys. J. 833(2), 264 (2016). doi:10.3847/1538-

4357/833/2/264

- O.M. Smirnov, Revisiting the radio interferometer measurement equation. I. A full-sky Jones formalism. Astron. Astrophys. 527, 106 (2011). doi:10.1051/0004-6361/201016082
- Y. Song, J. Sohl-Dickstein, D.P. Kingma, A. Kumar, S. Ermon, B. Poole, Score-Based Generative Modeling through Stochastic Differential Equations. arXiv e-prints, 2011–13456 (2020). doi:10.48550/arXiv.2011.13456
- A. Sonnenfeld, C. Tortora, H. Hoekstra, M. Asgari, M. Bilicki, C. Heymans, H. Hildebrandt, K. Kuijken, N.R. Napolitano, N. Roy, E. Valentijn, A.H. Wright, The dark matter halo masses of elliptical galaxies as a function of observationally robust quantities. Astron. Astrophys. 662, 55 (2022). doi:10.1051/0004-6361/202142511
- J.S. Spilker, D.P. Marrone, M. Aravena, M. Béthermin, M.S. Bothwell, J.E. Carlstrom, S.C. Chapman, T.M. Crawford, C. de Breuck, C.D. Fassnacht, A.H. Gonzalez, T.R. Greve, Y. Hezaveh, K. Litke, J. Ma, M. Malkan, K.M. Rotermund, M. Strandet, J.D. Vieira, A. Weiss, N. Welikala, ALMA Imaging and Gravitational Lens Models of South Pole Telescope—Selected Dusty, Star-Forming Galaxies at High Redshifts. Astrophys. J. **826**(2), 112 (2016). doi:10.3847/0004-637X/826/2/112
- C. Spingola, J.P. McKean, M.W. Auger, C.D. Fassnacht, L.V.E. Koopmans, D.J. Lagattuta, S. Vegetti, SHARP
 V. Modelling gravitationally lensed radio arcs imaged with global VLBI observations. Mon. Notices of the Royal Astron. Soc. 478(4), 4816–4829 (2018). doi:10.1093/mnras/sty1326
- C. Spingola, J.P. McKean, M. Lee, A. Deller, J. Moldon, A novel search for gravitationally lensed radio sources in wide-field VLBI imaging from the mJIVE-20 survey. Mon. Notices of the Royal Astron. Soc. 483(2), 2125– 2153 (2019). doi:10.1093/mnras/sty3189
- C. Spingola, A. Barnacka, Constraining VLBI-optical offsets in high redshift galaxies using strong gravitational lensing. Mon. Notices of the Royal Astron. Soc. 494(2), 2312–2326 (2020). doi:10.1093/mnras/staa870
- V. Springel, J. Wang, M. Vogelsberger, A. Ludlow, A. Jenkins, A. Helmi, J.F. Navarro, C.S. Frenk, S.D.M. White, The Aquarius Project: the subhaloes of galactic haloes. Mon. Notices of the Royal Astron. Soc. **391**(4), 1685– 1711 (2008). doi:10.1111/j.1365-2966.2008.14066.x
- H.R. Stacey, J.P. McKean, D.M. Powell, S. Vegetti, F. Rizzo, C. Spingola, M.W. Auger, R.J. Ivison, P.P. van der Werf, The rocky road to quiescence: compaction and quenching of quasar host galaxies at $z \sim 2$. Mon. Notices of the Royal Astron. Soc. **500**(3), 3667–3688 (2021). doi:10.1093/mnras/staa3433
- C. Storfer, X. Huang, A. Gu, W. Sheu, S. Banka, A. Dey, A. Jain, J. Kwon, D. Lang, V. Lee, A. Meisner, J. Moustakas, A.D. Myers, S. Tabares-Tarquinio,

E.F. Schlafly, D.J. Schlegel, New Strong Gravitational Lenses from the DESI Legacy Imaging Surveys Data Release 9. arXiv e-prints, 2206–02764 (2022). doi:10.48550/arXiv.2206.02764

- J. Stücker, O. Hahn, R.E. Angulo, S.D.M. White, Simulating the complexity of the dark matter sheet I: numerical algorithms. Mon. Notices of the Royal Astron. Soc. 495(4), 4943–4964 (2020). doi:10.1093/mnras/staa1468
- S.H. Suyu, P.J. Marshall, M.P. Hobson, R.D. Blandford, A Bayesian analysis of regularized source inversions in gravitational lensing. Mon. Notices of the Royal Astron. Soc. **371**(2), 983–998 (2006). doi:10.1111/j.1365-2966.2006.10733.x
- S.H. Suyu, P.J. Marshall, R.D. Blandford, C.D. Fassnacht, L.V.E. Koopmans, J.P. McKean, T. Treu, Dissecting the Gravitational Lens B1608+656. I. Lens Potential Reconstruction. Astrophys. J. 691(1), 277–298 (2009). doi:10.1088/0004-637X/691/1/277
- A.S. Tagore, C.R. Keeton, Statistical and systematic uncertainties in pixel-based source reconstruction algorithms for gravitational lensing. Mon. Notices of the Royal Astron. Soc. 445(1), 694–710 (2014). doi:10.1093/mnras/stu1671
- S. Tavaré, D.J. Balding, R.C. Griffiths, P. Donnelly, Inferring Coalescence Times From DNA Sequence Data. Genetics 145(2), 505–518 (1997). doi:10.1093/genetics/145.2.505
- S.S. Tie, C.S. Kochanek, Microlensing makes lensed quasar time delays significantly time variable. Mon. Notices of the Royal Astron. Soc. **473**(1), 80–90 (2018). doi:10.1093/mnras/stx2348
- J. Tinker, A.V. Kravtsov, A. Klypin, K. Abazajian, M. Warren, G. Yepes, S. Gottlöber, D.E. Holz, Toward a Halo Mass Function for Precision Cosmology: The Limits of Universality. Astrophys. J. 688, 709–728 (2008). doi:10.1086/591439
- T. Treu, L.V.E. Koopmans, Massive Dark Matter Halos and Evolution of Early-Type Galaxies to z ~1. Astrophys. J. 611(2), 739–760 (2004). doi:10.1086/422245
- C.S. Trotter, J.N. Winn, J.N. Hewitt, A Multipole-Taylor Expansion for the Potential of the Gravitational Lens MG J0414+0534. Astrophys. J. **535**(2), 671–691 (2000). doi:10.1086/308861
- B. Turner, T. Van Zandt, A tutorial on approximate bayesian computation. Journal of Mathematical Psychology 56, 69–85 (2012). doi:10.1016/j.jmp.2012.02.005
- H.C. Turner, M.R. Lovell, J. Zavala, M. Vogelsberger, The onset of gravothermal core collapse in velocitydependent self-interacting dark matter subhaloes. Mon. Notices of the Royal Astron. Soc. 505(4), 5327–5339 (2021). doi:10.1093/mnras/stab1725
- T.S. van Albada, J.N. Bahcall, K. Begeman, R. Sancisi, Distribution of dark matter in the spiral galaxy NGC 3198. Astrophys. J. 295, 305–313 (1985). doi:10.1086/163375

- L. Van de Vyvere, M.R. Gomer, D. Sluse, D. Xu, S. Birrer, A. Galan, G. Vernardos, TDCOSMO. VII. Boxyness/discyness in lensing galaxies: Detectability and impact on H_0 . Astron. Astrophys. **659**, 127 (2022). doi:10.1051/0004-6361/202141551
- S. Varma, M. Fairbairn, J. Figueroa, Dark Matter Subhalos, Strong Lensing and Machine Learning. arXiv e-prints, 2005–05353 (2020). doi:10.48550/arXiv.2005.05353
- K. Vattis, M.W. Toomey, S.M. Koushiappas, Deep learning the astrometric signature of dark matter substructure. PRD **104**(12), 123541 (2021). doi:10.1103/PhysRevD.104.123541
- S. Vegetti, L.V.E. Koopmans, Bayesian strong gravitationallens modelling on adaptive grids: objective detection of mass substructure in Galaxies. Mon. Notices of the Royal Astron. Soc. **392**(3), 945–963 (2009). doi:10.1111/j.1365-2966.2008.14005.x
- S. Vegetti, L.V.E. Koopmans, A. Bolton, T. Treu, R. Gavazzi, Detection of a dark substructure through gravitational imaging. Mon. Notices of the Royal Astron. Soc. 408(4), 1969–1981 (2010). doi:10.1111/j.1365-2966.2010.16865.x
- S. Vegetti, D.J. Lagattuta, J.P. McKean, M.W. Auger, C.D. Fassnacht, L.V.E. Koopmans, Gravitational detection of a low-mass dark satellite galaxy at cosmological distance. Nature 481(7381), 341–343 (2012). doi:10.1038/nature10669
- S. Vegetti, L.V.E. Koopmans, M.W. Auger, T. Treu, A.S. Bolton, Inference of the cold dark matter substructure mass function at z = 0.2 using strong gravitational lenses. Mon. Notices of the Royal Astron. Soc. **442**(3), 2017–2035 (2014). doi:10.1093/mnras/stu943
- S. Vegetti, G. Despali, M.R. Lovell, W. Enzi, Constraining sterile neutrino cosmologies with strong gravitational lensing observations at redshift $z \sim 0.2$. Mon. Notices of the Royal Astron. Soc. **481**(3), 3661–3669 (2018). doi:10.1093/mnras/sty2393
- S. Vegetti, L.V.E. Koopmans, Statistics of mass substructure from strong gravitational lensing: quantifying the mass fraction and mass function. Mon. Notices of the Royal Astron. Soc. 400(3), 1583–1592 (2009). doi:10.1111/j.1365-2966.2009.15559.x
- S. Vegetti, O. Czoske, L.V.E. Koopmans, Quantifying dwarf satellites through gravitational imaging: the case of SDSSJ120602.09+514229.5. Mon. Notices of the Royal Astron. Soc. 407(1), 225–231 (2010). doi:10.1111/j.1365-2966.2010.16952.x
- G. Vernardos, L.V.E. Koopmans, The very knotty lenser: Exploring the role of regularization in source and potential reconstructions using Gaussian process regression. Mon. Notices of the Royal Astron. Soc. 516(1), 1347– 1372 (2022). doi:10.1093/mnras/stac1924
- G. Vernardos, G. Tsagkatakis, Y. Pantazis, Quantifying

the structure of strong gravitational lens potentials with uncertainty-aware deep neural networks. Mon. Notices of the Royal Astron. Soc. **499**(4), 5641–5652 (2020). doi:10.1093/mnras/staa3201

- B. Villasenor, B. Robertson, P. Madau, E. Schneider, New Constraints on Warm Dark Matter from the Lyman-α Forest Power Spectrum. arXiv e-prints, 2209–14220 (2022)
- M. Vogelsberger, J. Zavala, A. Loeb, Subhaloes in selfinteracting galactic dark matter haloes. Mon. Notices of the Royal Astron. Soc. 423, 3740–3752 (2012). doi:10.1111/j.1365-2966.2012.21182.x
- M. Vogelsberger, J. Zavala, C. Simpson, A. Jenkins, Dwarf galaxies in CDM and SIDM with baryons: observational probes of the nature of dark matter. Mon. Notices of the Royal Astron. Soc. 444, 3684–3698 (2014a). doi:10.1093/mnras/stu1713
- M. Vogelsberger, S. Genel, V. Springel, P. Torrey, D. Sijacki, D. Xu, G. Snyder, D. Nelson, L. Hernquist, Introducing the Illustris Project: simulating the coevolution of dark and visible matter in the Universe. Mon. Notices of the Royal Astron. Soc. 444, 1518–1547 (2014b). doi:10.1093/mnras/stu1536
- M. Vogelsberger, J. Zavala, F.-Y. Cyr-Racine, C. Pfrommer, T. Bringmann, K. Sigurdson, ETHOS - an effective theory of structure formation: dark matter physics as a possible explanation of the small-scale CDM problems. Mon. Notices of the Royal Astron. Soc. 460, 1399–1416 (2016). doi:10.1093/mnras/stw1076
- M. Vogelsberger, J. Zavala, K. Schutz, T.R. Slatyer, Evaporating the Milky Way halo and its satellites with inelastic self-interacting dark matter. Mon. Notices of the Royal Astron. Soc. 484(4), 5437–5452 (2019). doi:10.1093/mnras/stz340
- S. Wagner-Carena, J. Aalbers, S. Birrer, E.O. Nadler, E. Darragh-Ford, P.J. Marshall, R.H. Wechsler, From Images to Dark Matter: End-to-end Inference of Substructure from Hundreds of Strong Gravitational Lenses. Astrophys. J. 942(2), 75 (2023). doi:10.3847/1538-4357/aca525
- S.J. Warren, S. Dye, Semilinear Gravitational Lens Inversion. Astrophys. J. **590**(2), 673–682 (2003). doi:10.1086/375132
- S.D.M. White, C.S. Frenk, M. Davis, Clustering in a neutrino-dominated universe. Astrophys. J. Letters 274, 1–5 (1983). doi:10.1086/184139
- P.N. Wilkinson, I.W.A. Browne, A.R. Patnaik, J.M. Wrobel, B. Sorathia, Interferometer phase calibration sources III. The regions +20<=delta_B1950<=+35 deg and +75<=delta_B1950<=+90 deg. Mon. Notices of the Royal Astron. Soc. **300**(3), 790–816 (1998). doi:10.1046/j.1365-8711.1998.01941.x
- P. Wizinowich, J.R. Lu, S. Cetre, J. Chin, C. Correia, J.-R. Delorme, L. Gers, S. Lilley, J. Lyke, E. Marin, S. Ragland,

P. Richards, A. Surendran, E. Wetherell, C.-F. Chen, D. Chu, T. Do, C. Fassnacht, M. Freeman, A. Gautam, A. Ghez, L. Hunter, T. Jones, M.C. Liu, D. Mawet, C. Max, M. Morris, M. Phillips, J.-B. Ruffio, N.-E. Rundquist, S. Sabhlok, S. Terry, T. Treu, S. Wright, Keck All sky Precision Adaptive optics program overview, in *Adaptive Optics Systems VIII*, ed. by L. Schreiber, D. Schmidt, E. Vernet Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, vol. 12185, 2022, p. 121850. doi:10.1117/12.2628275

- O. Wucknitz, LENSCLEAN revisited. Mon. Notices of the Royal Astron. Soc. **349**(1), 1–13 (2004). doi:10.1111/j.1365-2966.2004.07513.x
- D.D. Xu, S. Mao, J. Wang, V. Springel, L. Gao, S.D.M. White, C.S. Frenk, A. Jenkins, G. Li, J.F. Navarro, Effects of dark matter substructures on gravitational lensing: results from the Aquarius simulations. Mon. Notices of the Royal Astron. Soc. **398**(3), 1235–1253 (2009). doi:10.1111/j.1365-2966.2009.15230.x
- D.D. Xu, S. Mao, A.P. Cooper, J. Wang, L. Gao, C.S. Frenk, V. Springel, Substructure lensing: effects of galaxies, globular clusters and satellite streams. Mon. Notices of the Royal Astron. Soc. **408**(3), 1721–1729 (2010). doi:10.1111/j.1365-2966.2010.17235.x
- D.D. Xu, S. Mao, A.P. Cooper, L. Gao, C.S. Frenk, R.E. Angulo, J. Helly, On the effects of line-of-sight structures on lensing flux-ratio anomalies in a ACDM universe. Mon. Notices of the Royal Astron. Soc. **421**(3), 2553– 2567 (2012). doi:10.1111/j.1365-2966.2012.20484.x
- D. Xu, D. Sluse, L. Gao, J. Wang, C. Frenk, S. Mao, P. Schneider, V. Springel, How well can cold dark matter substructures account for the observed radio fluxratio anomalies. Mon. Notices of the Royal Astron. Soc. 447(4), 3189–3206 (2015). doi:10.1093/mnras/stu2673
- D. Yang, E.O. Nadler, H.-b. Yu, Strong Dark Matter Selfinteractions Diversify Halo Populations Within and Surrounding the Milky Way. arXiv e-prints, 2211–13768 (2022). doi:10.48550/arXiv.2211.13768
- J. Yao-Yu Lin, H. Yu, W. Morningstar, J. Peng, G. Holder, Hunting for Dark Matter Subhalos in Strong Gravitational Lensing with Neural Networks. arXiv e-prints, 2010– 12960 (2020). doi:10.48550/arXiv.2010.12960
- T.D. Yavetz, X. Li, L. Hui, Construction of wave dark matter halos: Numerical algorithm and analytical constraints. PRD **105**(2), 023512 (2022). doi:10.1103/PhysRevD.105.023512
- J. Zavala, C.S. Frenk, Dark Matter Haloes and Subhaloes. Galaxies 7(4), 81 (2019). doi:10.3390/galaxies7040081
- Y.B. Zel'dovich, Gravitational instability: An approximate theory for large density perturbations. Astron. Astrophys. 5, 84–89 (1970)
- I.A. Zelko, T. Treu, K.N. Abazajian, D. Gilman, A.J. Benson, S. Birrer, A.M. Nierenberg, A. Kusenko,

Constraints on Sterile Neutrino Models from Strong Gravitational Lensing, Milky Way Satellites, and the Lyman- α Forest. PRL **129**(19), 191301 (2022). doi:10.1103/PhysRevLett.129.191301

- Z.C. Zeng, A.H.G. Peter, X. Du, A. Benson, S. Kim, F. Jiang, F.-Y. Cyr-Racine, M. Vogelsberger, Corecollapse, evaporation, and tidal effects: the life story of a self-interacting dark matter subhalo. Mon. Notices of the Royal Astron. Soc. **513**(4), 4845–4868 (2022). doi:10.1093/mnras/stac1094
- A.R. Zentner, J.S. Bullock, Halo Substructure and the Power Spectrum. Astrophys. J. **598**(1), 49–72 (2003). doi:10.1086/378797
- F. Zwicky, Die Rotverschiebung von extragalaktischen Nebeln. Helvetica Physica Acta **6**, 110–127 (1933)