# Deconfined quantum critical points: a review

T. Senthil\*

Department of Physics, Massachusetts Institute of Technology, Cambridge, MA, 02139 (Dated: July 7, 2023)

Continuous phase transitions in equilibrium statistical mechanics were successfully described 50 years ago with the development of the renormalization group framework. This framework was initially developed in the context of phase transitions whose universal properties are captured by the long wavelength (and long time) fluctuations of a Landau order parameter field. Subsequent developments include a straightforward generalization to a class of T = 0 phase transitions driven by quantum fluctuations. In the last 2 decades it has become clear that there is a vast landscape of quantum phase transitions where the physics is not always usefully (or sometimes cannot be) formulated in terms of fluctuations of a Landau order parameter field. A wide class of such phase transitions - dubbed deconfined quantum critical points - involve the emergence of fractionalized degrees of freedom coupled to emergent gauge fields. Here I review some salient aspects of these deconfined critical points.

To be published in "50 years of the renormalization group", dedicated to the memory of Michael E. Fisher, edited by Amnon Aharony, Ora Entin-Wohlman, David Huse, and Leo Radzihovsky, World Scientific.

<sup>\*</sup>Electronic address: senthil@mit.edu

#### I. INTRODUCTION: THE LANDAU-GINZBURG-WILSON-FISHER PARADIGM

The development of the theory of phase transitions in critical phenomena in the 1960s and 1970s is one of the towering achievements in modern physics. Michael Fisher played a leading role in this development. The resulting understanding, associated primarily with the names of Landau, Ginzburg, Wilson, and Fisher, has come to be known as the LGWF theory (or sometimes the LGWF paradigm). This theory - which we briefly review below - provided a conceptual framework to discuss equilibrium phase transition phenomena. Combined with renormalization group methods and the dimensionality expansion introduced by Wilson and Fisher, the theory enables computations of critical exponents and other universal properties in a systematic approximation. The LGWF theory was generalized in many different directions, eg, to deal with dynamical critical phenomena, and more pertinently to this chapter, to quantum phase transitions at T = 0.

Phase transition theory begins by first asking how to distinguish distinct phases of matter. A crucial idea (due to Landau) is to focus on the symmetries of the microscopic Hamiltonian describing the many body system. The equilibrium state in a range of parameters (or the ground state if, at T = 0 may not have the symmetries of the microscopic Hamiltonian, *i.e* the symmetry is spontaneously broken. Then this state is sharply distinct from a symmetry preserving state that may exist at other parameter values. The evolution from the broken symmetry phase to one where the symmetry is restored must then involve (at least one) phase transition. Landau also introduced the notion of an order parameter to quantify the extent of broken symmetry. The Landau order parameter appears as a new thermodynamic variable needed to fully characterize the macroscopic state of the symmetry broken phase. The universal properties of the phase transition are then associated with the long wavelength (and long time, if dealing with quantum or dynamical critical phenomena) fluctuations of the order parameter field. The energy (or action, for quantum critical points) is written as a spatial integral of sums of an energy density that is expanded in powers of the order parameter field and its gradients. This is a coarse grained continuum description (an 'effective field theory') of the original microscopic model. The statistical mechanics of the resulting action (the Ginzburg-Landau action) then provides a description of the universal physics of the critical point. In practice, this can be analysed within the framework of the renormalization group with the Wilson-Fisher  $\epsilon$  expansion (or other expansions) as a tool to access the fixed points.

These beautiful set of ideas are sufficient to describe phase transitions in a large number of systems. The classic examples are the classical Ising or O(N) magnets. However, in the last few

decades, it has become clear that they turn out to not be the full story in a number of situations, particularly at quantum critical points. In this chapter we will survey a few different examples of how the LGWF paradigm breaks down. Nevertheless the framework of effective field theory and the renormalization group will remain central. What is different at these 'beyond LGWF' transitions is that the effective field theory will not be the naive one written in terms of just the order parameter field.

As we will see, there are a number of distinct routes for 'beyond LGWF' phase transitions. First, there are phases of matter which cannot be simply characterized in terms of the usual broken symmetry paradigm and hence do not admit a description in terms of a Landau order parameter. Obviously, their phase transitions will not be described by the LGWF paradigm. More surprisingly, we can have Landau-forbidden quantum critical points between phases that themselves are described as broken symmetry phases with a Landau order parameter. Suppose the microscopic symmetry of some system is described by a group G, and consider two phases of matter where the symmetry G is broken to a subgroup H in one phase, and to a different subgroup H' in another phase. In standard Landau theory. a second order phase transition between the two phases is possible only if one of the two groups H, H' is a subgroup of the other. Remarkably, we have learnt that this rule can be broken at a variety of quantum crtical points [1, 2]. The resulting critical theory is most usefully described in terms of a continuum field theory with emergent gauge fields coupled to matter fields carrying fractional quantum numbers of the microscopic global symmetry. These emergent fractionalized fields and associated gauge fields are not simply associated with quasiparticle excitations of either phase of matter; rather they rear their head at the critical point as useful variables to access the critical fixed point. Thus they have been dubbed 'deconfined quantum critical points'.

There are a number of examples of deconfined quantum critical points and related phenomena that have been studied in recent years. For instance, examples have been found where the same phase transition allows multiple universality classes[3]. These include situations[4] in which a Landau-allowed phase transition has both the conventional universality class as well as an unconventional one that is not described by the LGWF theory. Thus even when the phase transition occurs between a trivial gapped phase and a broken symmetry one described by a Landau order parameter, the effective field theory of the critical point may not be captured by the LGWF action. Even more striking are "unnecessary" critical points[3] which do not separate distinct phases of matter but rather live suspended within a single phase of matter. It is possible to find a path in parameter space which avoids the critical point completely. Together these results underscore the idea that we have very little intuition for whether a phase transition is allowed to be second order, and if so, what its field theoretic description will be. As such they embolden us to contemplate the possibility that many mysterious quantum critical phenomena observed in metallic systems might find explanations in a form that is beyond a conventional description (known as the Hertz-Millis paradigm) in terms of electronic quasiparticles coupled to a critically fluctuating bosonic order parameter field.

#### A. Phases of matter beyond Landau

For orientation, let us consider the different kinds of non-Landau phases of matter that are known to exist. The focus of this article will be on T = 0 quantum phases of matter, *i.e.*, ground states of quantum many body systems. A familiar example of phases whose distinction is not captured by the concept of broken symmetry are electronic band insulators as compared to band metals. The discovery of the integer and fractional quantum hall phases in the 1980s - which clearly do not break any microscopic symmetries - emphasized the possibility of new kinds of quantum ground states which fall under the rubric of 'topological quantum order'<sup>1</sup>. Since then, a large variety of 'beyond-Landau' phases have been shown to be possible theoretically, and some have been discovered in experiment. These include other topologically ordered states (such as gapped quantum spin liquids in insulating magnets), symmetry protected topological states (such as topological band insulators and the Haldane spin-1 chain), and various gapless phases of matter where the gaplessness is not due to Goldstone fluctuations associated with a broken symmetry. The most famous example is the Landau fermi liquid itself. Other variants include Dirac materials such as graphene. Most novel are phases of matter with no quasiparticle excitations at all: examples include some gapless quantum

<sup>&</sup>lt;sup>1</sup> In recent years, it has become popular to say that some such phases can actually be captured by the Landau paradigm if one generalizes the notion of symmetry. For a clear presentation of this point of view, see Ref. [5]. For instance topologically ordered states of matter are considered to spontaneously break 'higher-form' symmetries rather than ordinary (0-form) symmetries. This point of view is interesting and useful. In applying it to the systems typically of interest in condensed matter physics, we must recognize that these higher form (or other generalized) symmetries are not present in the microscopic system. They are emergent symmetries in particular phases of matter that may then be spontaneously broken. This is different from the conventional Landau paradigm which dictates that we distinguish phases by asking whether *microscopic* symmetries are present/absent in the equilibrium (ground) state. Thus, in my view, incorporating topological ordered or other "non-Landau" phases into the Landau paradigm requires not only generalizing the notion of symmetry but also generalizing what is meant by the Landau paradigm itself. Thus, in this article, I will refer to such phases as non-Landau as has been common for years.



FIG. 1: Varieties of quantum critical points. (a) The only case where the traditional LGWF paradigm is usually applicable. (b) Unless the unbroken symmetry group of one phase is a subgroup of the unbroken symmetry of the other phase, the transition is Landau-forbidden (c) Non-Landau order, if present in at least one of the two phases guarantees a transition beyond the LGWF paradigm.

spin liquids, and metallic non-fermi liquid phases.

Obviously, if the phase itself is not described simply by specifying a Landau order parameter, the phase transition out of it will not be described by an LGWF action in terms of any fluctuating order parameter field. In Fig 1 we sketch the variety of quantum phase transitions that accomodate this variety of phases of matter. The LGWF paradigm in its original form is only applicable to the situation where the symmetry preserving phase is gapped and trivial, and the symmetry broken phase is trivial apart from the broken symmetry. In all other cases, we should expect to find other descriptions of the phase transition.

#### B. Landau ordering transitions: a brief review

We begin with a quick review of Landau ordering quantum phase transitions. The canonical example is the transverse field Ising model on a *d*-dimensional lattice with global  $Z_2$  symmetry. This has a disordered phase that preserves the global  $Z_2$  symmetry, and an ordered phase that spontaneously breaks the symmetry. The quantum phase transition between these two phases is second order, and is in the universality class of the d + 1-dimensional classical Ising model. The universal critical singularities are captured by an LGWF action of the form

$$S = \int d^d x d\tau \left(\frac{\partial \phi}{\partial \tau}\right)^2 + (\nabla \phi)^2 + r\phi^2 + u\phi^4 \tag{1}$$

where  $\phi$  is the coarse-grained Ising order parameter field;  $\tau$  is the imaginary-time coordinate, and **x** is the spatial coordinate.

A simple generalization is to models with a global O(N) symmetry which (for d > 1) have a second order phase transition between a symmetry preserving trivial gapped phase, and a symmetry broken phase where the global O(N) symmetry is broken to the subgroup O(N-1). The transition is described by an action of the same form as in eqn. 1 but with  $\phi$  replaced by an N-component order parameter field **N** that transforms as a vector under the global O(N) symmetry.

When might we expect the LGWF continuum field theory to correctly describe the phase transition out of the broken symmetry phase? Note that the LGWF theory should really be understood as an expansion about the disordered symmetry preserving phase which is implicitly assumed to be trivial and gapped. There are quasiparticle excitations charged under the global O(N) symmetry. The most elementary such excitations transform under the vector representation of O(N). As the transition is approached, the gap for these excitations closes.

Thus we should expect LGWF actions of the form of Eqn. 1 to fail to describe the transition if the phase where the symmetry is restored is non-trivial in some way. For instance it may not be a trivial gapped phase and may have topological or other quantum order. Alternately it may break some other symmetry spontaneously. Clearly such cases are beyond the purview of the standard LGWF theory.

An important and interesting class of examples arises in situations where a trivial gapped symmetry preserving phase is forbidden on general grounds. A classic example is in quantum many body systems of interacting particles with conserved particle number (corresponding to a global U(1) symmetry) on a translation invariant lattice. If the lattice filling  $\nu$  is not an integer, then a trivial symmetric gapped ground state is not possible through a (generalization[6, 7] of) theorem by Lieb, Schultz, and Mattis (LSM). Such LSM restrictions apply very generally also to a number of translation-invariant models on a lattice where the degrees of freedom at each unit cell transform projectively under the global internal symmetry. For instance, in a spin system with global SO(3)symmetry, if we have a spin-1/2 (or other half-integer spin) moment at each unit cell, then a trivial gapped ground state is forbidden by the LSM restriction.

It follows that phase transitions in systems obeying LSM restrictions are never to be expected to be described by the LGWF theory.

A closely related situation arises for phase transitions that occur at the boundary of a higher dimensional Symmetry Protected Topological[8] (SPT) phase<sup>2</sup>. An SPT phase has a ground state that cannot be adiabatically deformed through any path that preserves the global symmetries into a completely trivial ground state. Examples of SPT phases include topological band insulators in diverse dimensions and the spin-1 Haldane spin chain. In the last decade a great deal has been understood about the classification and physical properties of such SPT phases. A defining feature is that it is not possible for the boundary of such an SPT phase to be trivially gapped. Thus phase transitions at SPT boundaries are also not expected to be described by the LGWF theory. Indeed the connection[10] of the theory of deconfined quantum criticality to SPT boundaries has been very effective in understanding the latter.

# **II. LGWF\* PHASE TRANSITIONS**

We begin by describing a simple example of a non-Landau transition, one that involves a critical point between a symmetry broken phase and one where the symmetry is restored but there is topological quantum order. Consider models of interacting bosons on a lattice with spatial dimension d = 2. The total boson number is conserved and there is a corresponding global U(1) symmetry. At every lattice filling, a superfluid phase where the global U(1) symmetry is spontaneously broken is clearly possible. As is well known, if the average number of bosons per unit cell is an integer, a Mott insulating phase is also possible where both the global U(1) and translation symmetry are preserved. A physical picture is that in the Mott insulator, there are an integer number of bosons localized to each unit cell of the lattice. This Mott insulator is a symmetric trivial gapped phase.

<sup>&</sup>lt;sup>2</sup> As emphasized in Ref. [9] we can regard a system satisfying the LSM restriction itself as the boundary of a specific case of an STP phase in one higher dimension, one involving lattice translation as well as internal symmetries.

The corresponding superfluid-Mott phase transition is described by LGWF theory, and is in the universality class of the classical XY model in 3 dimensions.

Next consider the situation when the number of bosons per unit cell is 1/2. Then a trivially gapped Mott insulator is forbidden. Nevertheless a symmetry preserving gapped ground state can exist so long as it has topological order and associated fractionally charged excitations. This state may be dubbed a fractionalized Mott insulator. The simplest example is when the topological order is that of a deconfined  $Z_2$  gauge theory. These can also be thought of as a quantum spin liquid ground state of a spin-1/2 magnet; for reviews, see Ref. [11, 12]. This theory has gapped quasiparticles in distinct topological superselection sectors denoted 1, e, m and  $\epsilon$ . The corresponding e and m quasiparticles have bosonic self-statistics, and a mutual braiding statistics of  $\pi$  (*i.e.*, when one of e, m is taken on a loop around the other, there is a phase of  $\pi$ ). They may be identified with the  $Z_2$  gauge charge and the  $Z_2$  gauge flux of the  $Z_2$  gauge theory<sup>3</sup>. Though the e and m are both good quasiparticles in this state of matter, their non-local statistical interaction implies that neither can be created locally, *i.e* by acting with local operators in the physical Hilbert space. This allows them to carry fractional quantum numbers of the global symmetry. Though a single e or m particle is not a local excitation,  $e^2$  or  $m^2$  (where a pair of e or m particles are created) are local. Thus we can create, say, a pair of e particles locally. The non-trivial statement is that the individual e-particles can be moved apart with a finite energy cost. Thus  $e^2$ ,  $m^2$  cannot carry fractional quantum numbers of the microscopic symmetries. It follows that e can have a quantum number 1/2 under the global U(1) symmetry. (In the presence of time reversal, it can be shown [10, 13, 14] that the *m* particle cannot also have fractional U(1) quantum numbers but will fractionalize lattice symmetries [15, 16].) Indeed it is possible to construct concrete models where precisely such 1/2-charged e particles, and neutral m particles exist [17–19]. Now if the microscopic boson model is at a lattice filling of 1/2a boson per unit cell, we can contemplate a fractionalized Mott insulator where there is a single eparticle that is localized in each unit cell.

The phase transition between the superfluid and this fractionalized Mott insulator is conveniently described as a condensation of the e particles while the gap to m stays non-zero. At energy scales well below the m gap, we can describe this transition through the action

$$S = \int d^d x d\tau \left| \frac{\partial \phi}{\partial \tau} \right|^2 + |\nabla \phi|^2 + r|\phi|^2 + u|\phi|^4 \tag{2}$$

<sup>&</sup>lt;sup>3</sup> Their bound state  $\epsilon$  is a fermion, and has  $\pi$  braiding statistics with both e and with m; the superselection sector 1 is simply associated with excitations that can be created by the action of local operators

where  $\phi$  is a coarse-grained complex field whose quanta (in the disordered phase) describe the *e* particles. This looks just the same as the standard LGWF action for the superfluid-Mott transition at integer filling but there is an important subtlety. As the  $\phi$  field creates the *e*-particle, it should not be directly identified with the superfluid order parameter, or indeed with any local observable in the original boson model. Relatedly this action does not know about the gapped *m*-sector which is part of the universal physics of both the fractionalized Mott insulator and the critical point. These subtleties can be handled by explicitly coupling the  $\phi$  field to a  $Z_2$  gauge field whose flux represents the *m*-particles. In that description, the  $\phi$  field is not gauge invariant under  $Z_2$  gauge transformations (though  $\phi^2$  is).

The action in Eqn. 2 is nevertheless useful to read off many universal critical properties. We simply note that it describes the gapless sector as the 3DXY universality class except that operators which are not  $Z_2$  gauge invariant are thrown out. The correlation length exponent  $\nu$  (which is determined by the scaling dimension of  $|\phi|^2$  which is gauge invariant) is the same as in the classical 3D XY model. However the scaling dimension of the physical superfluid order parameter  $\psi$  is different. This is because we can identify  $\psi = \phi^2$ . It thus has the anomalous exponent (introduced by Michael Fisher)  $\eta_{\Psi} \approx 1.49$ . This large anomalous exponent must be contrasted with that of the usual superfluid-Mott transition at integer filling where  $\eta \approx 0.03$ . For numerical studies of this transition, see Ref. [20].

Thus the superfluid-fractionalized Mott insulator transition is beyond the standard LGWF paradigm, albeit with a simple relation to the standard LGWF theory. For these reasons, this kind of transition has come to be known as the LGWF\* theory. For a sample of discussions of other LGWF\* transitions, see Ref. [21–23].

Theories with a different structure also arise at certain phase transitions between topologically ordered and symmetry broken phases, for instance between fractional quantum Hall and superfluid phases of bosons on a periodic lattice. Unlike in the example above, there is usually no anyon in the fractional quantum Hall state that can be directly condensed to both break the global U(1)symmetry and simultaneously destroy the topological order. Nevertheless, as argued in Ref. [24], a continuous transition is possible and can be formulated in terms of emergent fermionic degrees of freedom coupled to a U(1) gauge field. These fermions can be roughly thought of as composite fermions used to great effect in describing quantum Hall phenomena.



FIG. 2: The Neel and VBS states of a square lattice spin-1/2 magnet.

# III. LANDAU-FORBIDDEN QUANTUM CRITICAL POINTS BETWEEN LANDAU ALLOWED PHASES

We now turn to phase transitions between two phases which can both be characterized in terms of Landau order parameters corresponding to distinct broken symmetries. We will show that, contrary to naive expectations, such a phase transition can be second order. The primary example - which we review below - involves phase transitions in two dimensional insulating square lattice quantum antiferromagnets. These are described by models of systems of interacting quantum spins on the lattice. A number of other examples in two space dimension have also been presented. (We postpone the discussion of other dimensions to later sections). An interesting example involves a phase transition in a system of interacting fermions, as we also review below.

#### A. Neel-VBS transition on the square lattice

Consider a system of SU(2) spins on the square lattice with short range interactions. More precisely we will take the spin symmetry to be  $SO(3) = \frac{SU(2)}{Z_2}$ . and place spins that transform in the spin-1/2 (projective) representation of SO(3) at each site. We consider situations where the microscopic Hamiltonian has global SU(2), time reversal, and square lattice space group symmetries.

The standard ground state of such a magnet is the Néel state that breaks spin SO(3) and some lattice symmetries. Specifically it preserves lattice rotation symmetry, and a combination of unit lattice translation and time reversal. A different symmetry breaking state is a Valence Bond Solid (VBS) that preserves spin SO(3), and time reversal but breaks unit lattice translations in one direction, and lattice rotational symmetry. Clearly these phases break distinct symmetries, and the remnant symmetry in neither phase is a subgroup of the other. Naively we might expect the transition between these two phases to either be first order, or go through a region where both broken symmetries coexist, or go through an intermediate phase with no broken symmetry. Note that LSM restrictions applied to these spin models tell us that the last possibility is only allowed if such an intermediate symmetric phase has topological or some other non-trivial quantum order. In particular a trivially gapped symmetric phase is not possible. Thus, as discussed above, a theory for the destruction of either the Néel or the VBS order will necessarily be 'beyond LGWF'.

In Ref. [1, 2] it was argued that a direct second order transition is potentially possible and is described by the field theory

$$\mathcal{L}_0 = \sum_{\alpha=1,2} |D_b z_\alpha|^2 + \left(|z_1|^2 + |z_2|^2\right)^2.$$
(3)

Here  $z_{\alpha}$  ( $\alpha = 1, 2$ ) are scalar fields (known as 'bosonic spinons') that are coupled to a dynamical U(1)gauge field b, and  $D_{b,\mu} = \partial_{\mu} - ib_{\mu}$  is the covariant derivative. (This Euclidean action and subsequent similar actions are short-hands for the appropriate strongly coupled Wilson-Fisher critical theory where a background gauge field has been promoted to a dynamical field.) The model has a global SO(3) symmetry under which  $z_{\alpha}$  transforms as a spinor.<sup>4</sup> In the microsopic lattice spin model, this corresponds to the SO(3) spin rotation. It also has a global U(1) symmetry associated with the conservation of the flux of b. In the microsopic lattice spin model, this is not an exact symmetry. Consequently monopole operators (which pick up a phase under a U(1) rotation) must be added to the Lagrangian. However, it is known that lattice symmetries ensure that the minimal allowed monopole operator (with continuum angular momentum  $\ell = 0$ ) has strength 4.

Analytic arguments [1, 2] and numerical calculations [25–27] strongly support the possibility that these monopoles are irrelevant at the putative critical fixed point of Eq. (3). The Néel phase is obtained when  $z_{\alpha}$  is condensed, and the VBS phase when  $z_{\alpha}$  is gapped. The Néel phase breaks SO(3) to a U(1) subgroup while the VBS phase breaks the U(1) flux conservation symmetry. The Néel order parameter is simply  $\mathbf{N} = z^{\dagger}\sigma z$  ( $\sigma$  are Pauli matrices), and the VBS order parameter is the strength 1 monopole operator  $\mathcal{M}_b$  which creates  $2\pi$  flux of b.

The field theory in Eqn. 3 in the absence of monopoles is known as the Non-Compact  $CP^1$ ( $NCCP^1$ ) model. It describes[28] the phase transition in O(3) models where hedgehog topological defects have been suppressed by hand. These hedgehog defects correspond precisely to monopoles in the  $CP^1$  formulation of the O(3) non-linear sigma model.

<sup>&</sup>lt;sup>4</sup> Though  $z_{\alpha}$  transforms as a spinor under SU(2), rotations in the center of SU(2) can be compensated by a U(1) gauge transformation so that the spin rotation symmetry of the model is  $\frac{SU(2)}{Z_2} = SO(3)$ .

12

A heroic body of numerical work on specific quantum magnets and related systems [25–27, 29–42] shows a striking apparently continuous transition<sup>5</sup> between the N'eel and VBS phases with properties consistent with expectations based on the field theory in Eqn. 3.

What makes such a second order transition possible? The key physics is the properties of topological defects of the order parameters of either symmetry broken phase. To see this most simply, consider starting in the VBS phase which breaks the discrete symmetries of lattice translation and lattice rotation. There are four degenerate VBS ground states which are related to each other by  $Z_4$  lattice rotations about a site. Naively we might then expect a phase transition associated with the disordering of the VBS order to be in the universality class of the phase transition of a  $Z_4$  clock model in 2 + 1 - D. As is well known, this is the same as the 2 + 1 - D XY universality class: four-fold anisotropy of the XY order parameter is *dangerously irrelevant*<sup>6</sup> at the 3D XY fixed point.

This naive expectation is in fact not correct due to non-trivial structure of the topological defects of the VBS order parameter. Before exploring this further, it is useful to first recall the physics of topological defects at the usual 2 + 1 - D XY phase transition. For an XY order parameter, the topological defects are of course vortices. The transition out of the XY ordered phase requires the proliferation of these vortices. The most famous example is in the BKT transition of the *classical* 2d XY model but it remains true for the quantum 2 + 1 - D XY transition as well. Note that the 2 + 1 - D XY transition is described by the Wilson-Fisher fixed point with an action of the form in Eqn. 2. Equivalently we can describe it with a 'dual' field theory in terms of a complex scalar field  $\phi_v$  (identified physically with the vortex field) coupled to a dynamical U(1) gauge field:

$$\mathcal{L}_{v} = |D_{b}\phi_{v}|^{2} + \left(|\phi_{v}|^{2}\right)^{2}.$$
(4)

This the celebrated particle-vortex duality [47–49]. The disordered phase of the XY model is identified with the phase in which  $\phi_v$  is condensed. This gaps out the gauge field b through the Anderson-Higgs mechanism and we get a trivial gapped phase. The ordered phase of the XY model is identified with the phase in which  $\phi_v$  is gapped. At low energies, we integrate out  $\phi_v$  and get the free Maxwell action for the gauge field b. The corresponding 'photon' is then identified with the Goldstone mode of the broken XY symmetry. The XY order parameter itself is identified with the monopole operator  $\mathcal{M}_b$ . If the microscopic system has four-fold anisotropy on the XY order parameter, then we

<sup>&</sup>lt;sup>5</sup> It is however not yet clear whether the transition is truly second order or whether it displays only 'quasiuniversal' behavior up to a very large but finite length scale [27, 31–34, 37, 43–46]. We will discuss this below.

<sup>&</sup>lt;sup>6</sup> Another concept first introduced by Michael Fisher.

are allowed to add the operator  $\mathcal{M}_b^4 + c.c$  to the dual action.

Armed with this understanding, let us now return to the task of disordering the VBS order in the square lattice quantum magnet. For more detail, please see Ref. [50]. Naturally the topological defects are domain walls. Four such elementary domain walls can end at a vortex which is a point defect. It is easy to see that these VBS vortices necessarily involve a site of the lattice that is not paired with any other sites through a valence bond. Thus this site carries an unpaired spin-1/2 moment. This is a key difference with an ordinary  $Z_4$  clock model. Now consider quantum disordering the VBS order by proliferating (*i.e condensing*) these vortices. The spin-1/2 moment carried by these vortices will then lead to the breaking of spin rotation symmetry. It can be argued that the resulting state is precisely the Néel state. Indeed, following the principles of particle-vortex duality, the action in Eqn. 3 can be very simply understood as an effective field theory for the VBS vortices which are identified with the  $z_{\alpha}$  field<sup>7</sup>.

An alternate perspective is obtained by starting from the Néel state. The Néel order parameter **N** transforms in the vector representation of SO(3). It thus admits skyrmion defects corresponding to  $\Pi_2(S^2) = Z$ . In the original lattice system, the conservation of skyrmions can potentially be violated by 'hedgehog' events in space-time. However for spin-1/2 magnets on the square lattice, it can be shown that single skyrmions transform non-trivially under lattice symmetries[1, 2, 55, 56]. Thus a single skyrmion cannot be created or destroyed while preserving microscopic symmetries. In fact, the symmetry transformation of the skyrmion creation operator is identical to that of the VBS order parameter. If we now disorder the Néel order by proliferating skyrmions, the resulting paramagnetic phase will have VBS order.

The structure of the topological defects also reveals a problem with a naive Ginzburg-Landau action to describe the competition between the Néel and VBS orders. Indeed, a description of this competition will focus on the Néel order parameter  $\mathbf{N}$  and the VBS order parameter  $\psi$ , and write an action for their fluctuations. To that end it is convenient to introduce a 5-component unit vector

<sup>&</sup>lt;sup>7</sup> From a modern point of view, the structure of the topological defects implies that the  $NCCP^1$  theory has a mixed ' t Hooft anomaly between the global U(1) and SO(3) symmetries[51]. Specifically, if we introduce a background gauge field A for the global U(1) symmetry, the monopole operators in A will transform in the spinor representation of SO(3) which is a reflection of the anomaly. Similarly, the hypothesized emergent SO(5) symmetry discussed below of the low energy theory has its own 't Hooft anomaly which reduces to the mixed U(1) - SO(3) anomaly upon restricting to that subgroup[51]. The anomaly perspective provides a powerful tool to obtain some nonperturbative constraints on the RG flows of the theory. For a discussion on the relationship between the anomalies of the low energy theory and LSM constraints of the lattice spin model, see Ref. [52–54].

 $n^a$  (a = 1, ..., 5) such that  $n^{3,4,5}$  correspond to the three components of the Néel vector **N**, and  $n^{1,2}$  to the two real components of the VBS order parameter  $\psi$ . A naive Ginzburg-Landau action will then take the form of a non-linear sigma model

$$S_{naive} = \frac{1}{2g} \int d^3x \, (\partial n^a)^2 + \lambda \left( \mathbf{N}^2 - |\psi|^2 \right) + \dots$$
(5)

The first term is fully O(5) symmetric under rotations of  $n^a$ ; the second term introduces anisotropy between the 5 components of  $n^a$  so that the symmetry is reduced to  $O(3) \times O(2)$ . This term can also be used to tune between Néel and VBS ordered states. The ellipses include terms that further break these symmetries to those of the lattice spin model. This kind of Ginzburg-Landau action would usually suggest that the Néel-VBS transition is either first order, or goes through an intermediate phase with neither order. Further in this naive sigma model, this intermediate phase will be a trivially gapped phase<sup>8</sup> However since such a phase is forbidden in the lattice spin-1/2 magnet, clearly the naive sigma model needs to be modified in some way.

The crucial point is that the naive Ginzburg-Landau model does not capture the structure of the topological defects described above. Consider for instance the VBS ordered state where  $n^a$ points in the  $n^4$ ,  $n^5$  plane, and construct a vortex configuration of  $\psi = n^4 + in^5$ . At the core of the vortex,  $|\psi| = 0$  and so the unit vector  $n^a$  will point along the Néel direction. The dynamics of the Néel vector in the vortex core will be described by an SO(3) rotor model and will have quantum states transforming in integer representations of SO(3). However we saw above that the VBS vortex in our physical setting must transform as spin-1/2 under SO(3). This leads to a 'quantum intertwinement' of the two orders that must be incorporated into the Ginzburg-Landau theory. The correct action[57, 58] turns out to require inclusion of a Wess-Zumino-Witten (WZW) term at level-1 for the dynamics of the SO(5) vector  $n^a$ :

$$S = \frac{1}{2g} \int d^3x \, (\partial n^a)^2 + 2\pi\Gamma\left[n^a\right]. \tag{6}$$

The WZW term  $\Gamma$  is defined in the standard way: the field  $n^a$  defines a map from spacetime  $S^3$  to the target space  $S^4$ , and  $\Gamma$  is the ratio of the volume in  $S^4$  traced out by  $n_a$  to the total volume of  $S^4$ . If  $n^a(x, u)$  is any smooth extension of  $n^a(x)$  such that  $n^a(x, 0) = (0, 0, 0, 0, 1)$  and  $n^a(x, 1) = n^a(x)$ , then

$$\Gamma = \frac{\epsilon_{abcde}}{\text{Area}(S^4)} \int_0^1 d\, u \int d^3x \, n^a \partial_x n^b \partial_y n^c \partial_t n^d \partial_u n^e.$$
(7)

<sup>&</sup>lt;sup>8</sup> A direct second-order Neel-VBS transition can then only be expected at a multicritical point.

The WZW term correctly captures the non-trivial quantum numbers of the topological defects and is responsible for the non-Landau physics. Repeating the calculation of the structure of the VBS vortex, it is easy to see that the WZW term reduces - at the vortex core - to the standard spin Berry phase as appropriate for a spin-1/2 degree of freedom deescribed by N.

The sigma model is well-defined as a continuum field theory only in the weak coupling limit, where it is ordered. Here there is a clear semiclassical picture for the effect of the WZW term on the topological defects in the ordered state. For the transition itself — driven by the  $\lambda$  term — this ordered state corresponds to a first order phase transition. In particular the sigma model formulation does not by itself make it at all natural that the Néel-VBS transition can be second order. To directly say anything about the putative second-order Neel-VBS phase transition in this formulation requires extending the sigma model to strong coupling, and looking for a symmetry preserving critical phase (see below for discussion of a recent novel numerical study[59]). At strong coupling the sigma model theory is non-renormalizable and requires an alternative formulation as a continuum field theory. Physically, disordered phases of the sigma model (defined with an explicit UV cutoff) correspond to phases where topological defects of the order parameter have proliferated. Thus a modification of the topological defects leads to a modification of the corresponding disordered phases. The sigma model formulation thus exposes the seed, in the ordered phase, of the impending non-Landau physics of the disordered critical regime.

The field theory in Eqn.3 thus raises the possibility that the Neel-VBS transition may be second order. Whether it is actually so or not depends on the IR fate of the  $NCCP^1$  model. This is however hard to study. To get a handle, it is extremely useful to consider the general case of interacting SU(N) quantum spins. The proposed field theory for the non-Landau transition can be studied analytically in the limit of large-N through a 1/N expansion. A second order transition described by a conformal field theory is obtained. Thus the lattice spin model for sufficiently large-N provides a solvable and, hence important, example of a non-Landau deconfined quantum critical point. Extrapolating to small values of N, the critical fixed point has large values of the anomalous exponent  $\eta$  for both the N'eel and VBS order parameters. The four-fold monopoles are irrelevant at the critical fixed point, but of course they are important in the VBS ordered phase where they pin the orientation of the VBS order parameter to one of four degenerate values. This dangerous irrelevance of four-fold monopoles leads to the presence of two divergent length scales: one is the spin correlation length  $\xi$ , and the other - which may be identified as a scale associated with the pinning of the VBS order -  $\xi_{VBS}$  diverges as a power of  $\xi$ . The field theory for the N = 2 model can only be studied numerically, and its precise fate remains murky. In particular, many studies [27, 31–34, 37, 43–45, 60, 61] notice vexing problems (such as drifts with increasing system size in some critical properties).

Further studies [35] show the emergence of an SO(5) symmetry that rotates the Néel and VBS order parameters into one another. Such a symmetry is of course suggested by the sigma model formulation of the intertwinement between Neel and VBS orders, which however is not directly suited to describe the QCP as discussed above. This symmetry is however not at all apparent in the  $NCCP^1$  formulation. Ref. [51] provided an understanding of the emergence of the SO(5) symmetry from many points of view. The field theory admits multiple ('dual') descriptions. In any one of these descriptions only a subgroup of the full symmetry is apparent but it is a different subgroup in different descriptions. Thus by combining the information from these multiple descriptions we can get an understanding of the full emergent SO(5) symmetry.

A manifestly SO(5) invariant effective field theory[51] of the intertwinement of the Néel and VBS orders is obtained by starting with a fermionic spinon description of the square lattice spin-1/2 magnet. This naturally leads to a low energy theory of two flavors of massless Dirac fermions coupled to a dynamical SU(2) gauge field — a theory denoted  $N_f = 2 \ QCD_3$ . It is expected<sup>9</sup> that this theory shares the same universal infrared physics as the  $NCCP^1$  theory.

The possibility that the putative fixed point theory may have full SO(5) symmetry enables another check on its existence as a conformal field theory using conformal bootstrap methods. Unfortunately an SO(5) symmetric CFT with exponents in the range expected from numerical calculations of lattice spin models is not found[45]. Combined with the drifts in critical exponents mentioned above, it appears possible that for N = 2, the Neel-VBS transition is weakly first order but with a very long correlation length leading to the appearance of a continuous transition

A possible explanation of these facts (an unnaturally weak first order transition together with the emergence of the SO(5) symmetry) is provided by the idea that the transition in the N = 2 model is 'pseudocritical'[51, 62]. Pseudocriticality refers to a situation where the transition is second order 'close' to the physical model (in a generalized space including spatial dimensionality d, number of components N, etc) but the fixed points move slightly away from the real axis for the physical model. Then along the renormalization group flow the coupling constants linger ('walk') for a long time when they are closest to the fixed point but never really reach it, and ultimately flow away.

<sup>&</sup>lt;sup>9</sup> It is a bosonic theory with global SO(5) symmetry, and the same anomaly.

Pseudocriticality commonly occurs when a stable fixed point merges with an unstable one, and they annihilate each other. A famous example[63], is the 5-state classical Potts model in two dimensions. For other examples and a general discussion, see Ref. [64]. For the Néel-VBS transition, the pseudocritical scenario finds support[65, 66] in an epsilon expansion near space dimension d = 1 of the WZW non-linear sigma model.

An alternate suggestion [67–71] is that at N = 2, there is (in a parameter range) a quantum spin liquid intervening between the N'eel and VBS phases. As the parameters are varied, the quantum spin liquid region shrinks and is eventually replaced by a first order N'eel-VBS transition. There is an associated multicritical point when the spin liquid terminates and the line of first order transitions begins, and this is identified [70] with the deconfined quantum critical point.

Very recently, a beautiful study[59] provides strong support for the emergence of approximate conformal invariance and pseudocriticality at this phase transition. This work studied the SO(5)WZW model regulated as a fermion model on a "fuzzy sphere" (a sphere with a magnetic monopole at the center, which leads to the formation of Landau levels for the fermions). As shown in a previous work[72] on the 2 + 1 - D Ising transition, such a fuzzy sphere regularization enables examining the emergence of conformal symmetry by studying the spectrum of eigenstates through exact diagonalization. For conformal field theories, these energies are related to operator scaling dimensions, and the presence of conformal symmetry leads to many restrictions on the spectrum. For the regulated SO(5) WZW model, the fuzzy sphere calculation finds a spectrum nicely consistent with the emergence of conformal symmetry. However with increasing system size, an SO(5) singlet operator becomes weakly relevant. As we emphasized above, for the Néel-VBS transition, it is necessary that the SO(5) WZW model at strong coupling have a critical phase (the phase transition is then tuned by an anisotropy term that explicitly breaks SO(5) to  $SO(3) \times O(2)$ ). The presence of a weakly relevant SO(5) singlet is thus nicely consistent with the conjectured pseudocriticality at the Néel-VBS transition.

Clearly further clarifying the precise fate of the N = 2 model is an important goal for the near future. Nevertheless we emphasize that the existence of a second order Néel-VBS transition for large enough N establishes the matter-of-principle question on whether such transitions can exist at all.

#### **B.** Other realizations

A different realization of a deconfined critical point is a phase transition in a system of interacting electrons. The continuous global internal symmetry is U(2) corresponding to conservation of the total electronic charge and spin. In additon there is time reversal symmetry under which the electrons transform as Kramers doublets. Consider a phase where SU(2) spin rotation symmetry is broken spontaneously to a U(1) subgroup but the charge U(1) and time reversal are preserved. The corresponding order parameter is a three component field that transforms in the spin-1 representation of the SU(2) spin rotation symmetry. Now let us further specialize to a situation where - at the mean field level - there is a quantum spin Hall effect[73, 74]. In the bulk of such a system, the electronic spectrum is gapped but there will be gapless edge states at a boundary to the vacuum.

It was shown in Ref. [75] that fat skyrmion defects of the order parameter carry electrical charge-2. Restoring the broken SO(3) symmetry by condensing skyrmion defects then leads to superconducting order<sup>10</sup>. *i.e.*, to a spontaneous breakdown of the global U(1) associated with charge conservation. The phase transition between these two states is described at low energies by the  $NCCP^{1}$  field theory, and hence is another realization of the deconfined quantum critical point. In the  $NCCP^{1}$  formulation, the flux of the U(1) gauge field corresponds to the physical charge density, and hence is conserved exactly. Thus monopole operators of the U(1) gauge field correspond to Cooper pair operators that change the charge by 2. Obviously these monopole operators are not allowed in the action (unlike in the quantum magnetism realization)<sup>11</sup>.

The absence of monopoles of any strength implies that (in contrast to the Neel-VBS example), here there is only a single diverging length scale at the transition. Fermionic lattice models that realize this transition have been written down and are amenable to large scale Monte Carlo studies[78, 82] which find evidence for a second order transition.

<sup>&</sup>lt;sup>10</sup> This 'skyrmion condensation' mechanism of superconductivity has, in recent years, been proposed to be realized in twisted bilayer graphene[76, 77]. It has been demonstrated numerically both on lattice models[78], and on continuum models similar to those appropriate for twisted bilayer graphene[79].

<sup>&</sup>lt;sup>11</sup> The astute reader will recall that the low energy  $NCCP^1$  field theory has a mixed t' Hooft anomaly between the global U(1) and the global SO(3) symmetries. While we are certainly used to emergent symmetries having 't Hooft anomalies, it may be surprising that there is such an anomaly in a system where these symmetries are well-defined as 'on-site' symmetries of a microscopic lattice model. The resolution is that the anomaly is only present if the theory is regarded as a bosonic theory. In the presence of the (gapped) electronic degrees of freedom, the anomaly is trivialized, and this allows an on-site realization of the  $SO(3) \times U(1)$  symmetry. This has the consequence[80, 81] that in a system with boundaries, the fermion will be gapless at the edge even in the presence of the gapless critical bulk.

Deconfined critical field theories also arise in the context of phase transitions between trivial and SPT phases [10, 83, 84]. These are often related by duality transformations to the theories of the Landau-forbidden transitions[51].

#### IV. DQCP IN 1D

In the last few years, there have been a number of interesting studies [85–90] on one dimensional lattice quantum models that possess deconfined quantum critical points. In contrast to their 2-dimensional counterparts, these one dimensional models can be studied reliably with field theoretic and numerical methods. Models have even been constructed which are exactly solvable and show a deconfined quantum critical point.

Consider a 1*d* lattice model[85] of spin-1/2 moments with the Hamiltonian

$$\mathcal{H} = -\sum_{j} \left( J_x \sigma_j^x \sigma_{j+1}^x + J_z \sigma_j^z \sigma_{j+1}^z \right) + \sum_{j} \left( K_x \sigma_j^x \sigma_{j+2}^x + K_z \sigma_j^z \sigma_{j+2}^z \right) \tag{8}$$

The model has two global  $Z_2$  symmetries corresponding to  $\pi$  rotations of all the spins about either the x or the z-axis. These are denoted  $Z_2^z$  and  $Z_2^x$  respectively. It also has lattice translation symmetry, and the (anti-unitary) time reversal symmetry. By tuning the various parameters  $J_{x,z}, K_{x,z}$ , a number of phases can be accessed. For instance, a z - FM phase spontaneously breaks the  $Z_2^x$ and time reversal symmetries but preserves  $Z_2^z$  and translation. A distinct VBS phase preserves both  $Z_2$  symmetries and time reversal but breaks unit lattice translation symmetry. The phase transition between these two phases can be shown to be second order. The structure of the resulting theory has many similarities with that of the Neel-VBS transition in two space dimensions[85]. The Landau-forbidden phase transition is enabled by the structure of the topological defects. In the VBS phase, a domain wall leaves behind an unpaired site with a single spin-1/2 moment. This unpaired spin transforms projectively under the  $Z_2^x \times Z_2^y$  symmetry. Destroying the Neel order by proliferating these domain walls then necessarily breaks  $Z_2^x \times Z_2^y$  symmetry.

Ref. [88] constructed an exactly solvable 1*d* model with global  $Z_2 \times Z_2$  symmetry which features a continuous phase transition between two phases which break the two different  $Z_2$  symmetries. The symmetry action is however anomalous (one of the  $Z_2$  symmetries is not an on-site symmetry). Thus this system is usefully viewed as living at the boundary of a 2*d* Symmetry Protected Topological phase of spins with the same  $Z_2 \times Z_2$  symmetry. The model could be exactly mapped to a  $Z_4$ clock model coupled to a  $Z_2$  gauge field. In 1*d* the presence of the  $Z_2$  gauge field simply implies global selection rules on physical states when the system is placed on a ring. The Landau-forbidden transition can then be related to the ordering transition of the  $Z_4$  model which is continuous.

#### V. DQCP IN 3D: SOLVABLE EXAMPLES

A fruitful approach to generating a wealth of examples of deconfined critical points whose universal IR physics is solvable is described in Ref. [3]. The idea is to study deconfined critical theories that are free in the IR but become strongly coupled and confine upon moving away from criticality. To illustrate this, consider a quantum field theory in 3 + 1 spacetime dimensions described by  $N_f$ massless Dirac fermions coupled to, say, an SU(2) gauge field. The one-loop beta function for the gauge coupling g is well known, and takes the form

$$\beta(g^2) = \frac{dg^2}{dl} = \frac{g^3}{24\pi^2} (N_f - 11) \tag{9}$$

The theory is thus IR-free for  $N_f > 11$ . The gauge coupling g is irrelevant at the IR fixed point. The low energy physics is then given by a theory of free massless Dirac fermions ('quarks'), and a separate set of free bosons ('gluons'). Now consider perturbing the theory by adding a mass mfor the Dirac fermions. If we demand time reversal symmetry, then there is a unique (real) mass term that is allowed. However once the fermions are massive, they can be integrated out and the low energy theory is that of pure SU(2) gauge theory; in this phase, g flows to strong coupling, and the theory is confining. The RG flows are shown in Fig. 3. The massless point m = 0 can be interpreted as a quantum critical point in the phase diagram. The nature of the two phases depends on the value of  $N_f$ . For even  $N_f$ , the m = 0 point describes a phase transition between a trivial gapped phase and an SPT phase<sup>12</sup> (we discuss odd  $N_f$  below). Though this model may seem esoteric to condensed matter physics, it has the advantage that it realizes a particularly simple and solvable (in the IR) example of deconfined quantum criticality.

#### VI. OTHER NOVEL QCPS

In this section, we briefly discuss work in the last few years demonstrating a number of interesting critical phenomena. These were first found in the context of IR-free deconfined critical theories of

<sup>&</sup>lt;sup>12</sup> This field theory should be regarded as describing a microscopic system of interacting bosons: all local, *i.e* gauge invariant, operators in the theory are bosons. See Ref. [3] for identification of the protecting symmetries and further details.



FIG. 3: Renormalization group flows of the IR-free non-abelian gauge theory.

the kind described above (by considering different gauge groups and different matter content). More natural models (from a condensed matter perspective) that realize some of these phenomena have since been found.

## A. Multiple universality classes

In traditional phase transition theory, it is common to assume that, given two phases that are separated by a second order transition, there is a unique universality class for the phase transition. This is not generally true. For instance in disordered systems (eg, the  $\pm J$  Ising spin glass), it has been known for a long time that there may be multiple universality classes for the same phase transition. It is perhaps to be expected that this phenomenon of multiple universality classes also occurs in clean systems but the present author had not seen examples till recently. The IR-free non-abelian gauge theories naturally lead to many examples of multiple (deconfined critical) universality

classes for the same trivial-SPT phase transition. For a simpler example in a d = 1 spin systems, see Ref. [91].

#### B. Landau beyond Landau

The examples of multi-universality mentioned above were for transitions between gapped symmetry preserving phases. These of course are not to be described within the LGWF paradigm. It is thus interesting that it also possible to find examples where a Landau ordering phase transition that admits a universality class within the LGWF paradigm also allows a distinct universality class that is beyond LGWF. An example of this phenomenon is found in the model of the IR-free non-abelian gauge theory. Specifically, the SU(2) gauge theory with  $N_f$  massless fermions, with  $N_f$  odd and sufficiently large, is a (deconfined) critical point between a trivial gapped phase and a phase that spontaneously breaks time reversal symmetry but is otherwise gapped. The transition is driven by changing the sign of the fermion mass. This transition can clearly also be in the usual Ising universality class. The deconfined critical universality class has (many) more gapless degrees of freedom which further transform non-trivially under some global symmetries<sup>13</sup> - other than time reversal - present in this system. There is thus no formulation of this new universality class in terms of a theory written just in terms of the Ising order parameter field; it thus represents a dramatic breakdown of the LGWF paradigm.

At the time this article was written, there are no other examples of this 'Landau beyond Landau' phenomenon in simpler models in condensed matter physics, and finding one such is a fascinating future endeavor.

## C. "Unnecessary" critical points

It is often stated - in discussions on quantum materials - that the most fundamental question about a quantum critical point is to ask about what distinguishes the phases on either side. However, Ref. [3] described an interesting phenomenon where a second order critical line lives "inside" a single phase of matter. This was dubbed an "unnecessary" quantum critical point. In this case there is no

<sup>&</sup>lt;sup>13</sup> These global symmetries are unbroken on either side of the phase transition; in the usual Ising universality class, excitations charged under these other symmetries will be gapped. The deconfined universality class however has gapless excitations that are charged under these other global symmetries.

sharp distinction between the two sides of the transition and it is possible to go around it smoothly. It is analogous to the liquid-gas transition except that it is second order. Furthermore in the initial example described in Ref. [3], this unnecessary critical point had emergent deconfined gapless degrees of freedom though the phase it is submerged in is a trivial gapped phase. This example was once again constructed by studying the phase diagram of perturbed IR-free non-abelian gauge theories.

Subsequent work has found a number of examples in simpler systems. These include spin chains [80, 91] in 1d, and free fermion systems [3, 92] in 2d or 3d. It appears that unnecessary criticality is not an uncommon phenomenon.

#### VII. DISCUSSION

As reviewed here, there are a number of phase transitions where the strategy of focusing on long wavelength, long time fluctuations of a Landau order parameter field is not adequate. Nevertheless the framework of effective field theory and the renormalization group remain crucial. This is arguably one of the most important insights of the theory of critical phenomena developed in the 1960s and 1970s. At the deconfined critical points described in this review, useful effective field theories can be constructed in terms of emergent gapless fields that are charged under an emergent gauge field.

There are many further developments and applications that we have not discussed. For instance, recent work[93] studies the dynamics of the Neel-VBS transition in a driven 2d quantum magnet through numerical simulations. The deconfinement at the putative critical point impacts the annealing process, and leads to dynamics distinct from that described by the usual Kibble-Zurek scaling of conventional phase transitions.

An outgrowth of the theory of deconfined criticality is the understanding that there can be stable gapless 'deconfined' phases in two dimensional quantum magnets whose low energy theories involve fermions coupled to gauge fields[94]. An interesting example is a phase known as an algebraic (or Dirac) spin liquid where there are emergent gapless fermions coupled to a dynamical U(1) gauge field. Interesting generalizations of these states have been proposed[95] where it is not clear that there is *any* Lagrangian-based field theoretic description at all. If such states do exist, they may pose a challenge even to the paradigm of writing down an action for an effective field theory to describe the low energy physics. At the time of writing, there are relatively few experimental realizations of deconfined quantum critical points. For the Neel-VBS transition, very recent experiments on the material  $SrCu_2(BO_3)_2$  have been interpreted in terms of a deconfined critical point[96]. A provocative recent paper suggests[97] that a deconfined critical point may underlie the onset of superconductivity in the monolayer material  $WTe_2$ . A realization of a 1*d* deconfined quantum critical point in Rydberg atom simulators has been proposed[98].

One of the prime motivations for the original development of the theory of deconfined criticality in Ref. [1, 2] was the observation of non-fermi liquid physics at the transition between two fermi liquid phases in the heavy fermion materials[99, 100]. These critical points are usually associated with the onset of magnetic order out of the heavy fermi liquid but there may be a concomitant destruction[101, 102] of the Kondo screening responsible for the formation of the heavy fermi liquid in the first place. There is of course a qualitative resemblance to the simpler deconfined critical points discussed above. However a theory of these kinds of heavy fermion critical points must deal with hard questions on the fate of the fermi surfaces in either phase. Despite years of effort, these critical points remain mysterious. Nevertheless it is encouraging that the study of simpler models (without fermi surfaces) already demonstrates that Landau order parameter fluctuations may not be the full story at phase transitions, and, even worse, may distract from the essential physics of some phase transitions.

This work was supported by NSF grant DMR-2206305, and partially through a Simons Investigator Award from the Simons Foundation. Part of this article was written at the KITP and supported in part by the National Science Foundation under Grant No. NSF PHY-1748958. I am honored to be part of an academic lineage which places me as Michael Fisher's academic 'great grandson' (through David Nelson and Subir Sachdev). A different connection is through my postdoctoral advisor (Matthew Fisher) - one of Michael's physicist sons. Reading an article on critical phenomena and the renormalization group by Michael was one of the reasons I got attracted to condensed matter physics as an undergraduate. It is thus a pleasure to contribute to this memorial volume.

T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, Deconfined quantum critical points, *Science*. **303**, 1490, (2004).

- T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher, Quantum criticality beyond the landau-ginzburg-wilson paradigm, *Phys. Rev. B.* 70, 144407 (Oct, 2004). doi: 10.1103/PhysRevB. 70.144407. URL http://link.aps.org/doi/10.1103/PhysRevB.70.144407.
- [3] Z. Bi and T. Senthil, Adventure in topological phase transitions in 3+ 1-d: non-abelian deconfined quantum criticalities and a possible duality, *Physical Review X.* 9(2), 021034, (2019).
- [4] Z. Bi, E. Lake, and T. Senthil, Landau ordering phase transitions beyond the landau paradigm, *Physical Review Research.* 2(2), 023031, (2020).
- [5] J. McGreevy, Generalized symmetries in condensed matter, Annual Review of Condensed Matter Physics. 14, 57–82, (2023).
- [6] M. Oshikawa, Commensurability, excitation gap, and topology in quantum many-particle systems on a periodic lattice, *Physical review letters*. 84(7), 1535, (2000).
- [7] M. B. Hastings, Lieb-schultz-mattis in higher dimensions, *Phys. Rev. B.* 69, 104431 (Mar, 2004). doi: 10.1103/PhysRevB.69.104431. URL https://link.aps.org/doi/10.1103/PhysRevB.69.104431.
- [8] T. Senthil, Symmetry-protected topological phases of quantum matter, Annu. Rev. Condens. Matter Phys. 6(1), 299–324, (2015).
- [9] M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, Translational symmetry and microscopic constraints on symmetry-enriched topological phases: A view from the surface, *Physical Review X.* 6(4), 041068, (2016).
- [10] A. Vishwanath and T. Senthil, Physics of Three-Dimensional Bosonic Topological Insulators: Surface-Deconfined Criticality and Quantized Magnetoelectric Effect, *Physical Review X.* 3(1):011016 (Jan., 2013). doi: 10.1103/PhysRevX.3.011016.
- [11] L. Savary and L. Balents, Quantum spin liquids: a review, Reports on Progress in Physics. 80(1), 016502, (2016).
- [12] C. Broholm, R. Cava, S. Kivelson, D. Nocera, M. Norman, and T. Senthil, Quantum spin liquids, *Science.* 367(6475), eaay0668, (2020).
- [13] C. Wang and T. Senthil, Boson topological insulators: A window into highly entangled quantum phases, *Phys. Rev. B.* 87, 235122 (Jun, 2013). doi: 10.1103/PhysRevB.87.235122. URL http://link.aps.org/doi/10.1103/PhysRevB.87.235122.
- M. A. Metlitski, C. L. Kane, and M. P. A. Fisher, Bosonic topological insulator in three dimensions and the statistical witten effect, *Phys. Rev. B.* 88, 035131 (Jul, 2013). doi: 10.1103/PhysRevB.88.035131. URL http://link.aps.org/doi/10.1103/PhysRevB.88.035131.

- [15] R. A. Jalabert and S. Sachdev, Spontaneous alignment of frustrated bonds in an anisotropic, threedimensional ising model, *Physical Review B.* 44(2), 686, (1991).
- [16] T. Senthil and M. P. Fisher, Z 2 gauge theory of electron fractionalization in strongly correlated systems, *Physical Review B.* 62(12), 7850, (2000).
- [17] L. Balents, M. P. Fisher, and S. M. Girvin, Fractionalization in an easy-axis kagome antiferromagnet, *Physical Review B.* 65(22), 224412, (2002).
- [18] T. Senthil and O. Motrunich, Microscopic models for fractionalized phases in strongly correlated systems, *Physical Review B.* 66(20), 205104, (2002).
- [19] O. Motrunich and T. Senthil, Exotic order in simple models of bosonic systems, *Physical review letters*. 89(27), 277004, (2002).
- [20] S. V. Isakov, R. G. Melko, and M. B. Hastings, Universal signatures of fractionalized quantum critical points, *Science.* 335(6065), 193–195, (2012).
- [21] A. V. Chubukov, T. Senthil, and S. Sachdev, Universal magnetic properties of frustrated quantum antiferromagnets in two dimensions, *Physical review letters*. 72(13), 2089, (1994).
- [22] Y.-C. Wang, M. Cheng, W. Witczak-Krempa, and Z. Y. Meng, Fractionalized conductivity and emergent self-duality near topological phase transitions, *Nature Communications*. 12(1), 5347, (2021).
- [23] Y.-H. Zhang, Z. Zhu, and A. Vishwanath, Xy\* transition and extraordinary boundary criticality from fractional exciton condensation in quantum hall bilayer, arXiv preprint arXiv:2302.03703. (2023).
- [24] M. Barkeshli and J. McGreevy, Continuous transition between fractional quantum hall and superfluid states, *Physical Review B.* 89(23), 235116, (2014).
- [25] A. W. Sandvik, Evidence for deconfined quantum criticality in a two-dimensional heisenberg model with four-spin interactions, *Phys. Rev. Lett.* 98, 227202 (Jun, 2007). doi: 10.1103/PhysRevLett.98. 227202. URL http://link.aps.org/doi/10.1103/PhysRevLett.98.227202.
- [26] J. Lou, A. W. Sandvik, and N. Kawashima, Antiferromagnetic to valence-bond-solid transitions in two-dimensional SU(n) heisenberg models with multispin interactions, *Phys. Rev. B.* 80, 180414 (Nov, 2009). doi: 10.1103/PhysRevB.80.180414. URL http://link.aps.org/doi/10.1103/ PhysRevB.80.180414.
- [27] A. Nahum, J. T. Chalker, P. Serna, M. Ortuño, and A. M. Somoza, Deconfined quantum criticality, scaling violations, and classical loop models, *Phys. Rev. X.* 5, 041048 (Dec, 2015). doi: 10.1103/PhysRevX.5.041048. URL http://link.aps.org/doi/10.1103/PhysRevX.5.041048.
- [28] O. I. Motrunich and A. Vishwanath, Emergent photons and transitions in the o (3) sigma model

with hedgehog suppression, *Physical Review B.* **70**(7), 075104, (2004).

- [29] R. G. Melko and R. K. Kaul, Scaling in the fan of an unconventional quantum critical point, *Phys. Rev. Lett.* 100, 017203 (Jan, 2008). doi: 10.1103/PhysRevLett.100.017203. URL http://link.aps.org/doi/10.1103/PhysRevLett.100.017203.
- [30] A. Banerjee, K. Damle, and F. Alet, Impurity spin texture at a deconfined quantum critical point, *Phys. Rev. B.* 82, 155139 (Oct, 2010). doi: 10.1103/PhysRevB.82.155139. URL http://link.aps. org/doi/10.1103/PhysRevB.82.155139.
- [31] A. W. Sandvik, Continuous quantum phase transition between an antiferromagnet and a valence-bond solid in two dimensions: Evidence for logarithmic corrections to scaling, *Phys. Rev. Lett.* 104, 177201 (Apr, 2010). doi: 10.1103/PhysRevLett.104.177201. URL http://link.aps.org/doi/10.1103/PhysRevLett.104.177201.
- [32] K. Harada, T. Suzuki, T. Okubo, H. Matsuo, J. Lou, H. Watanabe, S. Todo, and N. Kawashima, Possibility of deconfined criticality in su(n) heisenberg models at small n, Phys. Rev. B. 88, 220408 (Dec, 2013). doi: 10.1103/PhysRevB.88.220408. URL http://link.aps.org/doi/10.1103/PhysRevB. 88.220408.
- [33] F.-J. Jiang, M. Nyfeler, S. Chandrasekharan, and U.-J. Wiese, From an antiferromagnet to a valence bond solid: evidence for a first-order phase transition, *Journal of Statistical Mechanics: Theory and Experiment.* 2008(02), P02009, (2008). URL http://stacks.iop.org/1742-5468/2008/i=02/a= P02009.
- [34] K. Chen, Y. Huang, Y. Deng, A. B. Kuklov, N. V. Prokof'ev, and B. V. Svistunov, Deconfined criticality flow in the heisenberg model with ring-exchange interactions, *Phys. Rev. Lett.* 110, 185701 (May, 2013). doi: 10.1103/PhysRevLett.110.185701. URL http://link.aps.org/doi/10.1103/PhysRevLett.110.185701.
- [35] A. Nahum, P. Serna, J. T. Chalker, M. Ortuño, and A. M. Somoza, Emergent so(5) symmetry at the néel to valence-bond-solid transition, *Phys. Rev. Lett.* 115, 267203 (Dec, 2015). doi: 10.1103/ PhysRevLett.115.267203. URL http://link.aps.org/doi/10.1103/PhysRevLett.115.267203.
- [36] O. I. Motrunich and A. Vishwanath, Comparative study of Higgs transition in one-component and two-component lattice superconductor models, ArXiv e-prints (May. 2008).
- [37] A. B. Kuklov, M. Matsumoto, N. V. Prokof'ev, B. V. Svistunov, and M. Troyer, Deconfined criticality: Generic first-order transition in the su(2) symmetry case, *Phys. Rev. Lett.* 101, 050405 (Aug, 2008). doi: 10.1103/PhysRevLett.101.050405. URL http://link.aps.org/doi/10.1103/

PhysRevLett.101.050405.

- [38] L. Bartosch, Corrections to scaling in the critical theory of deconfined criticality, *Phys. Rev. B.* 88, 195140 (Nov, 2013). doi: 10.1103/PhysRevB.88.195140. URL http://link.aps.org/doi/10.1103/PhysRevB.88.195140.
- [39] D. Charrier, F. Alet, and P. Pujol, Gauge theory picture of an ordering transition in a dimer model, *Phys. Rev. Lett.* 101, 167205 (Oct, 2008). doi: 10.1103/PhysRevLett.101.167205. URL http: //link.aps.org/doi/10.1103/PhysRevLett.101.167205.
- [40] G. Chen, J. Gukelberger, S. Trebst, F. Alet, and L. Balents, Coulomb gas transitions in threedimensional classical dimer models, *Phys. Rev. B.* 80, 045112 (Jul, 2009). doi: 10.1103/PhysRevB. 80.045112. URL http://link.aps.org/doi/10.1103/PhysRevB.80.045112.
- [41] D. Charrier and F. Alet, Phase diagram of an extended classical dimer model, *Phys. Rev. B.* 82, 014429 (Jul, 2010). doi: 10.1103/PhysRevB.82.014429. URL http://link.aps.org/doi/10.1103/PhysRevB.82.014429.
- [42] G. J. Sreejith and S. Powell, Scaling dimensions of higher-charge monopoles at deconfined critical points, *Phys. Rev. B.* 92, 184413 (Nov, 2015). doi: 10.1103/PhysRevB.92.184413. URL http://link.aps.org/doi/10.1103/PhysRevB.92.184413.
- [43] H. Shao, W. Guo, and A. W. Sandvik, Quantum criticality with two length scales, Science. 352 (6282), 213-216, (2016). ISSN 0036-8075. doi: 10.1126/science.aad5007. URL http://science.sciencemag.org/content/352/6282/213.
- [44] D. Simmons-Duffin. unpublished.
- [45] Y. Nakayama and T. Ohtsuki, Necessary condition for emergent symmetry from the conformal bootstrap, *Phys. Rev. Lett.* **117**, 131601 (Sep, 2016). doi: 10.1103/PhysRevLett.117.131601. URL http://link.aps.org/doi/10.1103/PhysRevLett.117.131601.
- [46] Z. Wang, M. P. Zaletel, R. S. Mong, and F. F. Assaad, Phases of the (2+1) dimensional so (5) nonlinear sigma model with topological term, *Physical Review Letters*. **126**(4), 045701, (2021).
- [47] M. E. Peskin, Mandelstam-'t hooft duality in abelian lattice models, Annals of Physics. 113(1), 122–152, (1978).
- [48] C. Dasgupta and B. I. Halperin, Phase transition in a lattice model of superconductivity, Phys. Rev. Lett. 47, 1556-1560 (Nov, 1981). doi: 10.1103/PhysRevLett.47.1556. URL http://link.aps.org/ doi/10.1103/PhysRevLett.47.1556.
- [49] M. P. A. Fisher and D. H. Lee, Correspondence between two-dimensional bosons and a bulk supercon-

ductor in a magnetic field, *Phys. Rev. B.* **39**, 2756–2759 (Feb, 1989). doi: 10.1103/PhysRevB.39.2756. URL http://link.aps.org/doi/10.1103/PhysRevB.39.2756.

- [50] M. Levin and T. Senthil, Deconfined quantum criticality and Néel order via dimer disorder, Phys. Rev. B. 70(22):220403 (Dec., 2004). doi: 10.1103/PhysRevB.70.220403.
- [51] C. Wang, A. Nahum, M. A. Metlitski, C. Xu, and T. Senthil, Deconfined quantum critical points: symmetries and dualities, *Physical Review X*. 7(3), 031051, (2017).
- [52] M. A. Metlitski and R. Thorngren, Intrinsic and emergent anomalies at deconfined critical points, *Physical Review B.* 98(8), 085140, (2018).
- [53] Z. Komargodski, T. Sulejmanpasic, and M. Ünsal, Walls, anomalies, and deconfinement in quantum antiferromagnets, *Physical Review B.* 97(5), 054418, (2018).
- [54] W. Ye, M. Guo, Y.-C. He, C. Wang, and L. Zou, Topological characterization of lieb-schultz-mattis constraints and applications to symmetry-enriched quantum criticality, *SciPost Physics*. 13(3), 066, (2022).
- [55] F. D. M. Haldane, O(3) nonlinear σ model and the topological distinction between integer- and halfinteger-spin antiferromagnets in two dimensions, *Phys. Rev. Lett.* **61**, 1029–1032 (Aug, 1988). doi: 10.1103/PhysRevLett.61.1029. URL http://link.aps.org/doi/10.1103/PhysRevLett.61.1029.
- [56] N. Read and S. Sachdev, Spin-peierls, valence-bond solid, and néel ground states of low-dimensional quantum antiferromagnets, *Phys. Rev. B.* 42, 4568–4589 (Sep, 1990). doi: 10.1103/PhysRevB.42. 4568. URL http://link.aps.org/doi/10.1103/PhysRevB.42.4568.
- [57] A. Tanaka and X. Hu, Many-body spin berry phases emerging from the π-flux state: Competition between antiferromagnetism and the valence-bond-solid state, *Phys. Rev. Lett.* 95, 036402 (Jul, 2005).
   doi: 10.1103/PhysRevLett.95.036402. URL http://link.aps.org/doi/10.1103/PhysRevLett.95.036402.
- [58] T. Senthil and M. P. A. Fisher, Competing orders, nonlinear sigma models, and topological terms in quantum magnets, *Phys. Rev. B.* 74, 064405 (Aug, 2006). doi: 10.1103/PhysRevB.74.064405. URL http://link.aps.org/doi/10.1103/PhysRevB.74.064405.
- [59] Z. Zhou, L. Hu, W. Zhu, and Y.-C. He, The so(5) deconfined phase transition under the fuzzy sphere microscope: Approximate conformal symmetry, pseudo-criticality, and operator spectrum, arXiv preprint arXiv:2306.16435. (2023).
- [60] J. Zhao, Y.-C. Wang, Z. Yan, M. Cheng, and Z. Y. Meng, Scaling of entanglement entropy at deconfined quantum criticality, *Physical Review Letters.* 128(1), 010601, (2022).

- [61] Y. Da Liao, G. Pan, W. Jiang, Y. Qi, and Z. Y. Meng, The teaching from entanglement: 2d deconfined quantum critical points are not conformal, arXiv preprint arXiv:2302.11742. (2023).
- [62] A. Nahum, J. Chalker, P. Serna, M. Ortuño, and A. Somoza, Deconfined quantum criticality, scaling violations, and classical loop models, *Physical Review X*. 5(4), 041048, (2015).
- [63] B. Nienhuis, A. N. Berker, E. K. Riedel, and M. Schick, First- and second-order phase transitions in potts models: Renormalization-group solution, *Phys. Rev. Lett.* 43, 737–740 (Sep, 1979). doi: 10.1103/PhysRevLett.43.737. URL http://link.aps.org/doi/10.1103/PhysRevLett.43.737.
- [64] V. Gorbenko, S. Rychkov, and B. Zan, Walking, weak first-order transitions, and complex cfts, Journal of High Energy Physics. 2018(10), 1–49, (2018).
- [65] R. Ma and C. Wang, Theory of deconfined pseudocriticality, *Physical Review B.* 102(2), 020407, (2020).
- [66] A. Nahum, Note on wess-zumino-witten models and quasiuniversality in 2+ 1 dimensions, *Physical Review B.* 102(20), 201116, (2020).
- [67] B. Zhao, J. Takahashi, and A. W. Sandvik, Multicritical deconfined quantum criticality and lifshitz point of a helical valence-bond phase, *Physical Review Letters*. **125**(25), 257204, (2020).
- [68] W.-Y. Liu, J. Hasik, S.-S. Gong, D. Poilblanc, W.-Q. Chen, and Z.-C. Gu, Emergence of gapless quantum spin liquid from deconfined quantum critical point, *Physical Review X.* 12(3), 031039, (2022).
- [69] D.-C. Lu, C. Xu, and Y.-Z. You, Self-duality protected multicriticality in deconfined quantum phase transitions, *Physical Review B.* 104(20), 205142, (2021).
- [70] J. Yang, A. W. Sandvik, and L. Wang, Quantum criticality and spin liquid phase in the shastrysutherland model, *Physical Review B.* 105(6), L060409, (2022).
- [71] H. Shackleton, A. Thomson, and S. Sachdev, Deconfined criticality and a gapless z 2 spin liquid in the square-lattice antiferromagnet, *Physical Review B.* 104(4), 045110, (2021).
- [72] W. Zhu, C. Han, E. Huffman, J. S. Hofmann, and Y.-C. He, Uncovering conformal symmetry in the 3d ising transition: State-operator correspondence from a quantum fuzzy sphere regularization, *Phys. Rev. X.* 13, 021009 (Apr, 2023). doi: 10.1103/PhysRevX.13.021009. URL https://link. aps.org/doi/10.1103/PhysRevX.13.021009.
- [73] C. L. Kane and E. J. Mele, Quantum spin hall effect in graphene, *Physical Review Letter*. 95, 226801, (2005).
- [74] C. L. Kane and E. J. Mele, Z<sub>2</sub> topological order and the quantum spin hall effect, *Physical Review*

Letter. 95, 146802, (2005).

- [75] T. Grover and T. Senthil, Topological spin hall states, charged skyrmions, and superconductivity in two dimensions, *Physical review letters*. 100(15), 156804, (2008).
- [76] E. Khalaf, S. Chatterjee, N. Bultinck, M. P. Zaletel, and A. Vishwanath, Charged skyrmions and topological origin of superconductivity in magic-angle graphene, *Science advances.* 7(19), eabf5299, (2021).
- [77] M. Christos, S. Sachdev, and M. S. Scheurer, Superconductivity, correlated insulators, and wess– zumino–witten terms in twisted bilayer graphene, *Proceedings of the National Academy of Sciences*. 117(47), 29543–29554, (2020).
- [78] Y. Liu, Z. Wang, T. Sato, M. Hohenadler, C. Wang, W. Guo, and F. F. Assaad, Superconductivity from the condensation of topological defects in a quantum spin-hall insulator, *Nature Communications.* 10(1), 2658, (2019).
- [79] S. Chatterjee, M. Ippoliti, and M. P. Zaletel, Skyrmion superconductivity: Dmrg evidence for a topological route to superconductivity, *Physical Review B*. 106(3), 035421, (2022).
- [80] R. Thorngren, A. Vishwanath, and R. Verresen, Intrinsically gapless topological phases, *Physical Review B*. 104(7), 075132, (2021).
- [81] R. Ma, L. Zou, and C. Wang, Edge physics at the deconfined transition between a quantum spin hall insulator and a superconductor, *SciPost Physics*. 12(6), 196, (2022).
- [82] M. Hohenadler, Y. Liu, T. Sato, Z. Wang, W. Guo, and F. F. Assaad, Thermodynamic and dynamical signatures of a quantum spin hall insulator to superconductor transition, *Physical Review B*. 106(2), 024509, (2022).
- [83] K. Slagle, Y.-Z. You, and C. Xu, Exotic quantum phase transitions of strongly interacting topological insulators, *Phys. Rev. B* . 91(11):115121 (Mar., 2015). doi: 10.1103/PhysRevB.91.115121.
- [84] Y.-Y. He, H.-Q. Wu, Y.-Z. You, C. Xu, Z. Y. Meng, and Z.-Y. Lu, Bona fide interaction-driven topological phase transition in correlated symmetry-protected topological states, *Phys. Rev. B*. 93 (11):115150 (Mar., 2016). doi: 10.1103/PhysRevB.93.115150.
- [85] S. Jiang and O. Motrunich, Ising ferromagnet to valence bond solid transition in a one-dimensional spin chain: Analogies to deconfined quantum critical points, *Physical Review B.* 99(7), 075103, (2019).
- [86] B. Roberts, S. Jiang, and O. I. Motrunich, Deconfined quantum critical point in one dimension, *Physical Review B.* 99(16), 165143, (2019).

- [87] R.-Z. Huang, D.-C. Lu, Y.-Z. You, Z. Y. Meng, and T. Xiang, Emergent symmetry and conserved current at a one-dimensional incarnation of deconfined quantum critical point, *Physical Review B*. 100(12), 125137, (2019).
- [88] C. Zhang and M. Levin, Exactly solvable model for a deconfined quantum critical point in 1d, *Physical Review Letters.* 130(2), 026801, (2023).
- [89] C. Mudry, A. Furusaki, T. Morimoto, and T. Hikihara, Quantum phase transitions beyond landauginzburg theory in one-dimensional space revisited, *Physical Review B*. 99(20), 205153, (2019).
- [90] L. Su, Boundary criticality via gauging finite subgroups: a case study on the clock model, arXiv preprint arXiv:2306.02976. (2023).
- [91] A. Prakash, M. Fava, and S. Parameswaran, Multiversality and unnecessary criticality in one dimension, arXiv preprint arXiv:2209.00037. (2022).
- [92] C.-M. Jian and C. Xu, Generic unnecessary quantum critical points with minimal degrees of freedom, *Physical Review B.* 101(3), 035118, (2020).
- [93] Y.-R. Shu, S.-K. Jian, A. W. Sandvik, and S. Yin, Equilibration of topological defects at the deconfined quantum critical point, arXiv preprint arXiv:2305.04771. (2023).
- [94] M. Hermele, T. Senthil, M. P. Fisher, P. A. Lee, N. Nagaosa, and X.-G. Wen, Stability of u (1) spin liquids in two dimensions, *Physical Review B*. 70(21), 214437, (2004).
- [95] L. Zou, Y.-C. He, and C. Wang, Stiefel liquids: Possible non-lagrangian quantum criticality from intertwined orders, *Physical Review X.* 11(3), 031043, (2021).
- [96] Y. Cui, L. Liu, H. Lin, K.-H. Wu, W. Hong, X. Liu, C. Li, Z. Hu, N. Xi, S. Li, et al., Proximate deconfined quantum critical point in srcu2 (bo3) 2, *Science*. 380(6647), eadc9487, (2023).
- [97] T. Song, Y. Jia, G. Yu, Y. Tang, P. Wang, R. Singha, X. Gui, A. J. Uzan, M. Onyszczak, K. Watanabe, T. Taniguchi, R. J. Cava, L. M. Schoop, N. P. Ong, and S. Wu. Unconventional superconducting quantum criticality in monolayer wte2, (2023).
- [98] J. Y. Lee, J. Ramette, M. A. Metlitski, V. Vuletic, W. W. Ho, and S. Choi, Landau-forbidden quantum criticality in rydberg quantum simulators, arXiv preprint arXiv:2207.08829. (2022).
- [99] P. Coleman, C. Pépin, Q. Si, and R. Ramazashvili, Topical review: How do fermi liquids get heavy and die?, *Journal of Physics Condensed Matter.* 13(35), R723–R738, (2001).
- [100] H. v. Löhneysen, A. Rosch, M. Vojta, and P. Wölfle, Fermi-liquid instabilities at magnetic quantum phase transitions, *Reviews of Modern Physics*. **79**(3), 1015, (2007).
- [101] Q. Si, S. Rabello, K. Ingersent, and J. L. Smith, Locally critical quantum phase transitions in strongly

correlated metals, *Nature*. **413**(6858), 804–808, (2001).

[102] T. Senthil, M. Vojta, and S. Sachdev, Weak magnetism and non-fermi liquids near heavy-fermion critical points, *Physical Review B.* 69(3), 035111, (2004).