

A PROOF OF GUO-WANG'S CONJECTURE ON THE UNIQUENESS OF POSITIVE HARMONIC FUNCTIONS IN THE UNIT BALL

PINGXIN GU AND HAIZHONG LI

ABSTRACT. Guo-Wang [*Calc. Var. Partial Differential Equations*, **59** (2020)] conjectured that for $1 < q < \frac{n}{n-2}$ and $0 < \lambda \leq \frac{1}{q-1}$, the positive solution $u \in C^\infty(\bar{B})$ to the equation

$$\begin{cases} \Delta u = 0 & \text{in } B^n, \\ u_\nu + \lambda u = u^q & \text{on } S^{n-1}, \end{cases}$$

must be constant. In this paper, we give a proof of this conjecture.

1. INTRODUCTION

In the past decades, a great deal of mathematical effort in best constant of Sobolev inequality has been devoted. For $n \geq 3$, a well-known subject is to figure out the minimum constant of Sobolev trace inequalities

$$\|u\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)} \leq C \|\nabla u\|_{L^2(\mathbb{R}_+^n)}, \quad \forall u \in C_0^\infty(\bar{\mathbb{R}}_+^n). \quad (1.1)$$

A key issue for this study is to investigate the extreme value of Sobolev quotient. Escobar [3] showed by conformal transformations that the best constant equals to

$$Q(B^n) := \inf_{u \in C^\infty(\bar{B}^n)} \frac{\int_{B^n} |\nabla u|^2 + \frac{n-2}{2} \int_{S^{n-1}} u^2}{\left(\int_{S^{n-1}} |u|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{n-1}}}. \quad (1.2)$$

Lions [9] proved that (1.2) can be achieved by a positive u satisfying the Euler-Lagrange equation

$$\begin{cases} \Delta u = 0 & \text{in } B^n, \\ u_\nu + \frac{n-2}{2} u = u^{\frac{n}{n-2}} & \text{on } S^{n-1}, \end{cases} \quad (1.3)$$

where ν is the unit outer normal vector on S^{n-1} . With this conclusion, Escobar [3] classified all positive solutions of (1.3) by an integral method and hence [4] proved that

$$|S^{n-1}|^{\frac{1}{n-1}} \left(\int_{S^{n-1}} u^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{n-1}} \leq \frac{2}{n-2} \int_{B^n} |\nabla u|^2 + \int_{S^{n-1}} u^2, \quad \forall u \in C^\infty(\bar{B}^n). \quad (1.4)$$

Different from Escobar, using harmonic analysis, Beckner [1] derived a family of inequalities

$$|S^{n-1}|^{\frac{q-1}{q+1}} \left(\int_{S^{n-1}} u^{q+1} \right)^{\frac{2}{q+1}} \leq (q-1) \int_{B^n} |\nabla u|^2 + \int_{S^{n-1}} u^2, \quad \forall u \in C^\infty(\bar{B}^n), \quad (1.5)$$

provided $1 < q < \infty$, if $n = 2$, and $1 < q \leq \frac{n}{n-2}$, if $n \geq 3$. The corresponding Euler-Lagrange equation to (1.5) is

$$\begin{cases} \Delta u = 0 & \text{in } B^n, \\ u_\nu + \frac{1}{q-1} u = u^q & \text{on } S^{n-1}. \end{cases} \quad (1.6)$$

2020 *Mathematics Subject Classification*. 58J90, 35B33.

Key words and phrases. Uniqueness, Positive harmonic function, Sobolev inequality, Obata type identity, Pohozaev identity.

It is apparent that the case $n \geq 3$ and $q = \frac{n}{n-2}$ of (1.5) and (1.6) are just (1.4) and (1.3) respectively. Also, in the same paper, Beckner [1] confirmed

$$|S^{n-1}|^{\frac{q-1}{q+1}} \left(\int_{S^{n-1}} u^{q+1} \right)^{\frac{2}{q+1}} \leq \frac{q-1}{n-1} \int_{S^{n-1}} |\nabla u|^2 + \int_{S^{n-1}} u^2, \quad \forall u \in C^\infty(S^{n-1}), \quad (1.7)$$

provided $1 < q < \infty$, if $n = 2$ or 3 , and $1 < q \leq \frac{n+1}{n-3}$, if $n \geq 4$. By considering the Euler-Lagrange equation and using integral method, Bidaut-Véron and Véron [2] gave a new proof of (1.7).

A natural question is: Now that (1.7) can be proved by the method of integration, can one prove (1.5) with the same strategy? Inspired by the arguments, Guo-Wang [7] proposed the following conjecture.

Conjecture ([7]). *If $u \in C^\infty(\bar{B}^n)$ is positive solution of the following equation*

$$\begin{cases} \Delta u = 0 & \text{in } B^n, \\ u_\nu + \lambda u = u^q & \text{on } S^{n-1}. \end{cases}$$

Then u is constant provided $1 < q < \frac{n}{n-2}$ and $0 < \lambda \leq \frac{1}{q-1}$.

In recent years, there are some partial results about the conjecture, see [5, 7, 8]. A remarkable one is that Guo-Hang-Wang [5] confirmed the conjecture in $n = 2$. In this paper, we investigate satisfactory Obata type identities. Combine with auxiliary functions, we give a proof of the conjecture when $n \geq 3$.

Theorem 1.1. *For $n \geq 3$, suppose $u \in C^\infty(\bar{B}^n)$ is a positive solution of the following equation*

$$\begin{cases} \Delta u = 0 & \text{in } B^n, \\ u_\nu + \lambda u = u^q & \text{on } S^{n-1}. \end{cases} \quad (1.8)$$

If $1 < q \leq \frac{n}{n-2}$ and $0 < \lambda \leq \frac{1}{q-1}$, then u must be constant $\lambda^{\frac{1}{q-1}}$ unless $q = \frac{n}{n-2}$ and $\lambda = \frac{1}{q-1}$, u is given by the following formula

$$u_\xi(x) = \left(\frac{n-2}{2} \frac{1 - |\xi|^2}{1 + |\xi|^2|x|^2 - 2\langle \xi, x \rangle} \right)^{\frac{n-2}{2}}, \quad (1.9)$$

for some $\xi \in B^n$.

From Theorem 1.1, we immediately get

Corollary 1.2. *The conjecture holds for $n \geq 3$.*

The paper is organized as follows. In Sect.2, we establish a Pohozaev identity by calculating divergence of a given tensor field with two parameters, and designate one of the parameters to make Pohozaev identity works. In Sect.3, we give an Obata type identity and determine the other parameter to ensure the effectiveness of Obata's skill. In Sect.4, we introduce auxiliary functions involving the length of position vector with one parameter to obtain improved identities. In Sect.5, we give the proof of Theorem 1.1 and a new proof of Beckner's inequality (1.5) by integral method.

2. A POHOZAEV IDENTITY

In this section, we use divergence theorem to prove an identity, and use Pohozaev identity to simplify it. During the simplification, we will determine the parameter as what we have hoped.

2.1. Preliminary. To begin with, let $u = v^{-\frac{1}{q-1}}$ in (1.8), then v satisfies

$$\begin{cases} \Delta v = \frac{q}{q-1} \frac{|\nabla v|^2}{v} & \text{in } B^n, \\ v_\nu = (q-1)(\lambda v - 1) & \text{on } S^{n-1}. \end{cases} \quad (2.1)$$

In this case, the boundary condition is concise, which only involves normal derivatives and linear expression of the solution v . In the text that follows, we define

$$M := \operatorname{div}(v^a \nabla_{\nabla v} \nabla v), \quad N := \operatorname{div}(v^a \Delta v \nabla v), \quad (2.2)$$

and focus on the quantity $M - bN$, where parameters $a, b \in \mathbb{R}$ are to be determined later.

In order to integrate by parts and handle the boundary term, we choose an orthonormal frame $\{e_i\}_{i=1}^n$ such that $e_n = \nu$ is the unit outward normal vector on S^{n-1} and the second fundamental form of S^{n-1} equals to identity. Thus on S^{n-1} , by use of Reilly's formula, see [10], we have

$$v_{\alpha n} = (\lambda(q-1) - 1)v_\alpha, \quad \forall 1 \leq \alpha \leq n-1, \quad (2.3)$$

$$\sum_{\alpha=1}^{n-1} v_{\alpha\alpha} = \Delta_{S^{n-1}} v + (n-1)v_n, \quad (2.4)$$

and

$$v_{nn} = \Delta v - \sum_{\alpha=1}^{n-1} v_{\alpha\alpha} = \frac{q}{q-1} \frac{|\nabla v|^2}{v} - \Delta_{S^{n-1}} v - (n-1)v_n. \quad (2.5)$$

For simplicity of presentation, we always set subscripts $1 \leq \alpha \leq n-1$ and $1 \leq i, j \leq n$. Einstein summation convention for these subscripts is always used in what follows.

2.2. Choice of parameter a. With these preliminaries, we integrate M and N in B^n respectively. Combining with (2.3), (2.5) and (2.1), we obtain by use of divergence theorem

$$\begin{aligned} \int_{B^n} M &= \int_{B^n} (v^a v_{ij} v_i)_j = \int_{S^{n-1}} v^a v_{nn} v_n + \int_{S^{n-1}} v^a v_{\alpha n} v_\alpha \\ &= \frac{q}{q-1} \int_{S^{n-1}} v^{a-1} |\nabla v|^2 v_n - \int_{S^{n-1}} v^a \Delta_{S^{n-1}} v v_n - (n-1) \int_{S^{n-1}} v^a v_n^2 \\ &\quad + (\lambda(q-1) - 1) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2, \end{aligned} \quad (2.6)$$

$$\int_{B^n} N = \int_{B^n} (v^a \Delta v v_i)_i = \int_{S^{n-1}} v^a \Delta v v_n = \frac{q}{q-1} \int_{S^{n-1}} v^{a-1} |\nabla v|^2 v_n. \quad (2.7)$$

Now we deal the term $\int_{S^{n-1}} v^a \Delta_{S^{n-1}} v v_n$ in (2.6) by applying divergence theorem on S^{n-1} :

$$\begin{aligned} \int_{S^{n-1}} v^a \Delta_{S^{n-1}} v v_n &= (q-1)\lambda \int_{S^{n-1}} v^{a+1} \Delta_{S^{n-1}} v - (q-1) \int_{S^{n-1}} v^a \Delta_{S^{n-1}} v \\ &= -(q-1)(a+1)\lambda \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 + (q-1)a \int_{S^{n-1}} v^{a-1} |\nabla_{S^{n-1}} v|^2. \end{aligned} \quad (2.8)$$

Combining formulas (2.6), (2.7) through (2.8) and splitting $|\nabla v|^2$ into $|\nabla_{S^{n-1}} v|^2 + v_n^2$ on S^{n-1} , we derive

$$\begin{aligned} \int_{B^n} (M - bN) &= (1-b) \frac{q}{q-1} \int_{S^{n-1}} v^{a-1} |\nabla_{S^{n-1}} v|^2 v_n + (1-b) \frac{q}{q-1} \int_{S^{n-1}} v^{a-1} v_n^3 \\ &\quad + \lambda(q-1)(a+1) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 - (q-1)a \int_{S^{n-1}} v^{a-1} |\nabla_{S^{n-1}} v|^2 \\ &\quad - (n-1) \int_{S^{n-1}} v^a v_n^2 + (\lambda(q-1) - 1) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2. \end{aligned} \quad (2.9)$$

Use (2.1) to eliminate v_n in the term $(1-b)\frac{q}{q-1}\int_{S^{n-1}}v^{a-1}|\nabla_{S^{n-1}}v|^2v_n$ and one of the three v_n 's in the term $(1-b)\frac{q}{q-1}\int_{S^{n-1}}v^{a-1}v_n^3$, (2.9) becomes

$$\begin{aligned} \int_{B^n}(M-bN) &= \lambda q(1-b)\int_{S^{n-1}}v^a|\nabla_{S^{n-1}}v|^2 - q(1-b)\int_{S^{n-1}}v^{a-1}|\nabla_{S^{n-1}}v|^2 \\ &\quad + \lambda q(1-b)\int_{S^{n-1}}v^av_n^2 - q(1-b)\int_{S^{n-1}}v^{a-1}v_n^2 \\ &\quad + \lambda(q-1)(a+1)\int_{S^{n-1}}v^a|\nabla_{S^{n-1}}v|^2 - (q-1)a\int_{S^{n-1}}v^{a-1}|\nabla_{S^{n-1}}v|^2 \\ &\quad - (n-1)\int_{S^{n-1}}v^av_n^2 + (\lambda(q-1)-1)\int_{S^{n-1}}v^a|\nabla_{S^{n-1}}v|^2. \end{aligned} \quad (2.10)$$

We obtain from (2.10) that

$$\begin{aligned} \int_{B^n}(M-bN) &= (\lambda q(1-b) + \lambda(q-1)(a+2) - 1)\int_{S^{n-1}}v^a|\nabla_{S^{n-1}}v|^2 \\ &\quad - (q(1-b) + a(q-1))\int_{S^{n-1}}v^{a-1}|\nabla_{S^{n-1}}v|^2 \\ &\quad + (\lambda q(1-b) - (n-1))\int_{S^{n-1}}v^av_n^2 \\ &\quad - q(1-b)\int_{S^{n-1}}v^{a-1}v_n^2. \end{aligned} \quad (2.11)$$

The term $\int_{S^{n-1}}v^{a-1}|\nabla_{S^{n-1}}v|^2$ in (2.11) appears for the reason that the boundary condition of (2.1) is not homogeneous. It is desirable to eliminate it with some equalities. The key idea is to confirm Pohozaev identities from conditions (2.1). Only if we choose $a = -\frac{q+1}{q-1}$ makes it work.

Proposition 2.1. *Let v be a positive solution of (2.1). For $a = -\frac{q+1}{q-1}$, we derive the following Pohozaev identity*

$$\int_{S^{n-1}}v^{a-1}|\nabla_{S^{n-1}}v|^2 = \int_{S^{n-1}}v^{a-1}v_n^2 + (n-2)\int_{S^{n-1}}v^av_n^2 - (n-2)(q-2)\lambda\int_{B^n}v^a|\nabla v|^2. \quad (2.12)$$

Proof. We note $x_{ij} = \delta_{ij}$ in B^n . Thus, we have the following calculation

$$\begin{aligned} \int_{B^n}\operatorname{div}(v^{a-1}|\nabla v|^2x) &= \\ &= (a-1)\int_{B^n}v^{a-2}|\nabla v|^2x_iv_i + 2\int_{B^n}v^{a-1}v_{ij}x_iv_j + n\int_{B^n}v^{a-1}|\nabla v|^2. \end{aligned} \quad (2.13)$$

On the other hand, take (2.1) into consideration, we obtain

$$\begin{aligned} \int_{B^n}\operatorname{div}(v^{a-1}\langle\nabla v, x\rangle\nabla v) &= \\ &= \left(a-1+\frac{q}{q-1}\right)\int_{B^n}v^{a-2}|\nabla v|^2x_iv_i + \int_{B^n}v^{a-1}v_{ij}x_iv_j + \int_{B^n}v^{a-1}|\nabla v|^2. \end{aligned} \quad (2.14)$$

Note the condition $a = -\frac{q+1}{q-1}$ is the only choice to satisfy

$$\frac{a-1}{a-1+\frac{q}{q-1}} = 2.$$

Then (2.13) $-2\times$ (2.14) implies

$$(n-2)\int_{B^n}v^{a-1}|\nabla v|^2 = \int_{B^n}\operatorname{div}(v^{a-1}|\nabla v|^2x) - 2\int_{B^n}\operatorname{div}(v^{a-1}\langle\nabla v, x\rangle\nabla v). \quad (2.15)$$

Notice on S^{n-1} , we have $x = \nu$, thus $\langle x, \nu \rangle = 1$ and $\langle \nabla v, x \rangle = \langle \nabla v, \nu \rangle = v_n$. By divergence theorem, we have from (2.15) that

$$\begin{aligned} (n-2) \int_{B^n} v^{a-1} |\nabla v|^2 &= \int_{S^{n-1}} v^{a-1} |\nabla v|^2 \langle x, \nu \rangle - 2 \int_{S^{n-1}} v^{a-1} \langle \nabla v, x \rangle \langle \nabla v, \nu \rangle \\ &= \int_{S^{n-1}} v^{a-1} |\nabla v|^2 - 2 \int_{S^{n-1}} v^{a-1} v_n^2 \\ &= \int_{S^{n-1}} v^{a-1} |\nabla_{S^{n-1}} v|^2 - \int_{S^{n-1}} v^{a-1} v_n^2. \end{aligned} \quad (2.16)$$

Now it remains to deal with the term $\int_{B^n} v^{a-1} |\nabla v|^2$ on the left-hand side of (2.16). A key observation follows from divergence theorem and $a = -\frac{q+1}{q-1}$ that

$$\int_{S^{n-1}} v^a v_n = \int_{B^n} \operatorname{div}(v^a \nabla v) = \left(a + \frac{q}{q-1}\right) \int_{B^n} v^{a-1} |\nabla v|^2 = -\frac{1}{q-1} \int_{B^n} v^{a-1} |\nabla v|^2, \quad (2.17)$$

and

$$\int_{S^{n-1}} v^{a+1} v_n = \int_{B^n} \operatorname{div}(v^{a+1} \nabla v) = \left(a + 1 + \frac{q}{q-1}\right) \int_{B^n} v^a |\nabla v|^2 = \frac{q-2}{q-1} \int_{B^n} v^a |\nabla v|^2. \quad (2.18)$$

Using the fact that the boundary condition of (2.1) yields

$$\int_{S^{n-1}} v^a v_n^2 = (q-1) \int_{S^{n-1}} v^a v_n (\lambda v - 1) = \lambda(q-1) \int_{S^{n-1}} v^{a+1} v_n - (q-1) \int_{S^{n-1}} v^a v_n. \quad (2.19)$$

Putting (2.17) and (2.18) into (2.19), we conclude

$$\int_{S^{n-1}} v^a v_n^2 = \lambda(q-2) \int_{B^n} v^a |\nabla v|^2 + \int_{B^n} v^{a-1} |\nabla v|^2. \quad (2.20)$$

Then (2.12) follows from substituting (2.20) into (2.16) to eliminate the term $\int_{B^n} v^{a-1} |\nabla v|^2$. \square

By means of Proposition 2.1, we eliminate $\int_{S^{n-1}} v^{a-1} |\nabla_{S^{n-1}} v|^2$ in (2.11) and derive

Corollary 2.2. *Let v be a positive solution of (2.1). The quantities M, N are defined as (2.2). Then for $a = -\frac{q+1}{q-1}$,*

$$\begin{aligned} \int_{B^n} (M - bN) &= \left(\lambda q(1-b) + \lambda(q-3) - 1\right) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 \\ &\quad + \left(q(1-b) - (q+1)\right) (n-2)(q-2)\lambda \int_{B^n} v^a |\nabla v|^2 \\ &\quad + \left(\lambda q(1-b) + (n-2)qb - 1\right) \int_{S^{n-1}} v^a v_n^2 \\ &\quad - \left(2q(1-b) - (q+1)\right) \int_{S^{n-1}} v^{a-1} v_n^2. \end{aligned} \quad (2.21)$$

In the next section, we will determine the value of b , which is a rather subtle issue. One can guess that we should control the coefficient of $\int_{S^{n-1}} v^{a-1} v_n^2$ as

$$2q(1-b) - (q+1) \geq 0, \quad (2.22)$$

since we have no other ways to decompose it. In fact, we will make the equality sign in (2.22) hold. More logical reasons for such choice will be explained later.

3. ON OBATA'S SKILL

3.1. Obata type identities. It is time to expand $M - bN$, which gives

$$\begin{aligned} M - bN &= (v^a v_{ij} v_i)_j - b(v^a \Delta v v_j)_j \\ &= \left(v^a v_{ij} v_{ij} + av^{a-1} v_{ij} v_i v_j + v^a (\Delta v)_i v_i \right) - b \left(av^{a-1} \Delta v |\nabla v|^2 + v^a (\Delta v)_j v_j + v^a (\Delta v)^2 \right) \\ &= v^a v_{ij} v_{ij} + av^{a-1} v_{ij} v_i v_j + (1-b)v^a (\Delta v)_i v_i - abv^{a-1} \Delta v |\nabla v|^2 - bv^a (\Delta v)^2. \end{aligned} \quad (3.1)$$

Recall that Δv satisfies (2.1), so

$$v^a (\Delta v)_i v_i = \frac{q}{q-1} v^a \left(\frac{|\nabla v|^2}{v} \right)_i v_i = \frac{2q}{q-1} v^{a-1} v_{ij} v_i v_j - \frac{q}{q-1} v^{a-2} |\nabla v|^4. \quad (3.2)$$

If we put (3.2) and (2.1) into (3.1), we will obtain

$$\begin{aligned} M - bN &= v^a v_{ij} v_{ij} + \left(a + (1-b) \frac{2q}{q-1} \right) v^{a-1} v_{ij} v_i v_j \\ &\quad - \left((1-b) \frac{q}{q-1} + ab \frac{q}{q-1} + b \left(\frac{q}{q-1} \right)^2 \right) v^{a-2} |\nabla v|^4. \end{aligned} \quad (3.3)$$

For simplicity, we define d as

$$d := \frac{a}{2} + (1-b) \frac{q}{q-1} = \frac{2q(1-b) - (q+1)}{2(q-1)}.$$

After putting $a = -\frac{q+1}{q-1}$, (3.3) becomes

$$\begin{aligned} M - bN &= v^a v_{ij} v_{ij} + 2dv^{a-1} v_{ij} v_i v_j - \left(\frac{q}{q-1} d + \frac{q}{2(q-1)} \right) v^{a-2} |\nabla v|^4 \\ &= v^a \left| v_{ij} + d \frac{v_i v_j}{v} \right|^2 - \left(d^2 + \frac{q}{q-1} d + \frac{q}{2(q-1)} \right) v^{a-2} |\nabla v|^4. \end{aligned} \quad (3.4)$$

By means of the technique of Obata, we define a trace-free 2-tensor

$$E_{ij} := v_{ij} + d \frac{v_i v_j}{v} - \frac{1}{n} \left(\frac{q}{q-1} + d \right) \frac{|\nabla v|^2}{v} \delta_{ij}. \quad (3.5)$$

The tensor satisfies

$$\left| v_{ij} + d \frac{v_i v_j}{v} \right|^2 = \left| E_{ij} + \frac{1}{n} \left(\frac{q}{q-1} + d \right) \frac{|\nabla v|^2}{v} \delta_{ij} \right|^2 = |E|^2 + \frac{1}{n} \left(\frac{q}{q-1} + d \right)^2 \frac{|\nabla v|^4}{v^2}. \quad (3.6)$$

As what we have anticipated, (3.4) can be expressed as Obata type identities

Proposition 3.1. *Let v be a positive solution of (2.1). The quantities M, N are defined as (2.2). Then for $a = -\frac{q+1}{q-1}$ and b, d satisfies $d = \frac{2q(1-b) - (q+1)}{2(q-1)}$, we have*

$$M - bN = v^a |E|^2 + \left(\frac{1}{n} \left(\frac{q}{q-1} + d \right)^2 - \left(d^2 + \frac{q}{q-1} d + \frac{q}{2(q-1)} \right) \right) v^{a-2} |\nabla v|^4. \quad (3.7)$$

3.2. Choice of parameter b . As what we have mentioned, (2.22) hopes $d \geq 0$. However, to make Obata's skill work, we apply

Proposition 3.2. *For $n \geq 3$, $1 < q < \frac{n}{n-2}$, we have*

$$\frac{1}{n} \left(\frac{q}{q-1} + d \right)^2 - \left(d^2 + \frac{q}{q-1} d + \frac{q}{2(q-1)} \right) \geq 0, \quad (3.8)$$

provided

$$-\frac{2q}{(q-1)(q+1)} \leq d \leq 0. \quad (3.9)$$

Proof. Based on the observation that the left-hand side of (3.8) is decreasing on n . It suffices to show the case when $n \rightarrow \frac{2q}{q-1}$, i.e.

$$\frac{q-1}{2q} \left(\frac{q}{q-1} + d \right)^2 - \left(d^2 + \frac{q}{q-1}d + \frac{q}{2(q-1)} \right) \geq 0,$$

equivalently,

$$-\frac{q+1}{2q}d^2 - \frac{1}{q-1}d \geq 0.$$

we complete the proof. \square

Combining with (2.22) and Proposition 3.2, it is sure that we need $d = 0$, i.e. $b = \frac{q-1}{2q}$. In this case, we may both eliminate the term $\int_{S^{n-1}} v^{a-1} v_n^2$ in (2.21) and make sure the coefficient of $v^{a-2} |\nabla v|^4$ in (3.7) is positive. To conclude what we have stated, from (3.7) and (2.21),

Corollary 3.3. *Let v be a positive solution of (2.1). The quantities M, N are defined as (2.2). Then for $a = -\frac{q+1}{q-1}$ and $b = \frac{q-1}{2q}$, we have*

$$M - bN = v^a |E|^2 + \frac{q}{2n(q-1)^2} (n - (n-2)q) v^{a-2} |\nabla v|^4 \geq 0, \quad (3.10)$$

and

$$\begin{aligned} \int_{B^n} (M - bN) &= \frac{\lambda(3q-5) - 2}{2} \int_{S^{n-1}} v^a |\nabla_{S^{n-1}}|^2 \\ &\quad - (n-2)(q-2) \frac{q+1}{2} \lambda \int_{B^n} v^a |\nabla v|^2 \\ &\quad + \left(\lambda \frac{q+1}{2} + \frac{(n-2)q-n}{2} \right) \int_{S^{n-1}} v^a v_n^2. \end{aligned} \quad (3.11)$$

Remark 3.4. The proof of Proposition 3.2 also suggests that for $a = -\frac{q+1}{q-1}$ and any $b' \in \mathbb{R}$ satisfies

$$\frac{q-1}{2q} \leq b' \leq \frac{q^2 + 4q - 1}{2q(q+1)}, \quad (3.12)$$

we always have

$$M - b'N \geq 0, \quad (3.13)$$

where v is a positive solution of (2.1) and M, N are defined as (2.2). In fact, if we define

$$d' := \frac{2q(1-b') - (q+1)}{2(q-1)}, \quad (3.14)$$

then correspondingly, the range of d' is precisely

$$-\frac{2q}{(q-1)(q+1)} \leq d' \leq 0. \quad (3.15)$$

It is easy to check that (3.15) is equivalent to (3.12).

4. AUXILIARY FUNCTION AND IMPROVED IDENTITIES

To deal with more complicate situations, an introduction of auxiliary function ϕ here is essential. Set

$$\phi(x) = \frac{|x|^2 + c}{2} \quad (4.1)$$

with $c > 0$ to be determined. It is evident to see

$$\phi_i = x_i, \quad \phi_{ij} = \delta_{ij} \quad \text{and} \quad \phi(x) \in \left[\frac{c}{2}, \frac{1+c}{2} \right) \quad \text{in } B^n, \quad (4.2)$$

and

$$\phi(x) \equiv \frac{1+c}{2}, \quad \text{and} \quad \nabla \phi = \nu, \quad \text{on } S^{n-1}. \quad (4.3)$$

4.1. Improved inequalities involving auxiliary function. We define

$$P := \operatorname{div}(v^a |\nabla v|^2 \nabla \phi), \quad Q := \operatorname{div}(v^a \langle \nabla v, \nabla \phi \rangle \nabla v), \quad (4.4)$$

where $a = -\frac{q+1}{q-1}$. Direct computations lead

$$P = 2v^a v_{ij} v_i x_j - \frac{q+1}{q-1} v^{a-1} |\nabla v|^2 v_i x_i + n v^a |\nabla v|^2, \quad (4.5)$$

$$Q = v^a v_{ij} v_i x_j - \frac{1}{q-1} v^{a-1} |\nabla v|^2 v_i x_i + v^a |\nabla v|^2. \quad (4.6)$$

If we focus on the quantity $(q-3)P + 4Q$, then by (4.5)-(4.6),

$$(q-3)P + 4Q = 2(q-1)v^a v_{ij} v_i x_j - (q-1)v^{a-1} |\nabla v|^2 v_i x_i + (4 - (3-q)n)v^a |\nabla v|^2. \quad (4.7)$$

Recall the definition of M, N in (2.2), and note clearly $\phi \equiv \frac{1+c}{2}$ on S^{n-1} . If we multiply ϕ on M, N and integrate them on B^n respectively, with the help of divergence theorem and (4.2)-(4.3), we will obtain

$$\begin{aligned} \int_{B^n} M\phi &= \int_{B^n} (v^a v_{ij} v_i)_j \phi = \int_{B^n} (v^a v_{ij} v_i \phi)_j - \int_{B^n} v^a v_{ij} v_i x_j \\ &= \int_{S^{n-1}} v^a v_{in} v_i \phi - \int_{B^n} v^a v_{ij} v_i x_j \\ &= \frac{1+c}{2} \int_{S^{n-1}} v^a v_{in} v_i - \int_{B^n} v^a v_{ij} v_i x_j \\ &= \frac{1+c}{2} \int_{B^n} M - \int_{B^n} v^a v_{ij} v_i x_j, \end{aligned} \quad (4.8)$$

and

$$\begin{aligned} \int_{B^n} N\phi &= \int_{B^n} (v^a \Delta v v_i)_i \phi = \int_{B^n} (v^a \Delta v v_i \phi)_i - \int_{B^n} v^a \Delta v v_i x_i \\ &= \int_{S^{n-1}} v^a \Delta v v_n \phi - \frac{q}{q-1} \int_{B^n} v^{a-1} |\nabla v|^2 v_i x_i \\ &= \frac{1+c}{2} \int_{S^{n-1}} v^a \Delta v v_n - \frac{q}{q-1} \int_{B^n} v^{a-1} |\nabla v|^2 v_i x_i \\ &= \frac{1+c}{2} \int_{B^n} N - \frac{q}{q-1} \int_{B^n} v^{a-1} |\nabla v|^2 v_i x_i. \end{aligned} \quad (4.9)$$

Putting (4.8)-(4.9) into the integration of (4.7) and noting $b = \frac{q-1}{2q}$, we arrive at a neat result that

$$\int_{B^n} \left((q-3)P + 4Q \right) = 2(q-1) \int_{B^n} (M - bN) \left(\frac{1+c}{2} - \phi \right) - \left((3-q)n - 4 \right) \int_{B^n} v^a |\nabla v|^2. \quad (4.10)$$

On the other hand, if we right use divergence theorem on the integration of P, Q in B^n , we will derive by use of (4.3) that

$$\int_{B^n} P = \int_{S^{n-1}} v^a |\nabla v|^2 \langle \nabla \phi, \nu \rangle = \int_{S^{n-1}} v^a |\nabla v|^2 = \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 + \int_{S^{n-1}} v^a v_n^2, \quad (4.11)$$

$$\int_{B^n} Q = \int_{S^{n-1}} v^a \langle \nabla v, \nabla \phi \rangle \langle \nabla v, \nu \rangle = \int_{S^{n-1}} v^a v_n^2. \quad (4.12)$$

Combine with (4.11) and (4.12), we conclude

$$\int_{B^n} \left((q-3)P + 4Q \right) = (q-3) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 + (q+1) \int_{S^{n-1}} v^a v_n^2. \quad (4.13)$$

Thus (4.10) and (4.13) imply

$$\begin{aligned} \int_{B^n} (M - bN) \phi &= \frac{1+c}{2} \int_{B^n} (M - bN) + \frac{3-q}{2(q-1)} \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 \\ &\quad - \frac{(3-q)n-4}{2(q-1)} \int_{B^n} v^a |\nabla v|^2 - \frac{q+1}{2(q-1)} \int_{S^{n-1}} v^a v_n^2. \end{aligned} \quad (4.14)$$

Putting (3.11) into (4.14), we obtain an improved identity

Corollary 4.1. *Let v be a positive solution of (2.1). The quantities M, N are defined as (2.2) where $a = -\frac{q+1}{q-1}$, $b = \frac{q-1}{2q}$. Set $\phi(x) = \frac{|x|^2+c}{2}$ with $c > 0$ to be determined. Then we have*

$$\begin{aligned} \int_{B^n} (M - bN) \phi &= \left(\frac{1+c}{2} \left(\frac{\lambda(3q-5)-2}{2} \right) + \frac{3-q}{2(q-1)} \right) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 \\ &\quad - \left(\frac{1+c}{2} (n-2)(q-2) \frac{q+1}{2} \lambda + \frac{(3-q)n-4}{2(q-1)} \right) \int_{B^n} v^a |\nabla v|^2 \\ &\quad + \left(\frac{1+c}{2} \left(\lambda \frac{q+1}{2} + \frac{(n-2)q-n}{2} \right) - \frac{q+1}{2(q-1)} \right) \int_{S^{n-1}} v^a v_n^2. \end{aligned} \quad (4.15)$$

5. PROOFS OF THEOREM 1.1

After having addressed all the preceding conclusions, we are now in a position to establish the proof of Theorem 1.1. Integrating (4.6) in B^n and combining with (4.12), (4.8) and (4.9), it is obvious that

$$\int_{S^{n-1}} v^a v_n^2 = \int_{B^n} Q = \int_{B^n} \left(M - \frac{1}{q} N \right) \left(\frac{1+c}{2} - \phi \right) + \int_{B^n} v^a |\nabla v|^2. \quad (5.1)$$

Choose $c = \frac{2}{\lambda(q-1)} - 1 \geq 1$ in Corollary 4.1. In this case, (4.15) becomes

$$\begin{aligned} \int_{B^n} (M - bN) \phi &= \left(1 - \frac{1}{\lambda(q-1)} \right) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 \\ &\quad - \frac{1}{2} \left((n-2)q - n \right) \int_{B^n} v^a |\nabla v|^2 \\ &\quad + \frac{(n-2)q-n}{2\lambda(q-1)} \int_{S^{n-1}} v^a v_n^2. \end{aligned} \quad (5.2)$$

Putting (5.1) into (5.2) to eliminate $\int_{B^n} v^a |\nabla v|^2$, we obtain

$$\begin{aligned} \int_{B^n} (M - bN)\phi &= \frac{1}{2}((n-2)q - n) \int_{B^n} (M - \frac{1}{q}N)(\frac{1+c}{2} - \phi) \\ &\quad + \left(1 - \frac{1}{\lambda(q-1)}\right) \int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 \\ &\quad - \frac{(n - (n-2)q)(1 - \lambda(q-1))}{2\lambda(q-1)} \int_{S^{n-1}} v^a v_n^2. \end{aligned} \quad (5.3)$$

Recall when $n \geq 3$, we have $1 < q \leq \frac{n}{n-2} \leq 3$, Thus

$$\frac{q-1}{2q} \leq \frac{1}{q} \leq \frac{q^2 + 4q - 1}{2q(q+1)}.$$

By Remark 3.4, we conclude that

$$M - \frac{1}{q}N \geq 0.$$

So the conditions $1 < q \leq \frac{n}{n-2}$ and $0 < \lambda \leq \frac{1}{q-1}$ imply the right-hand side of (5.3) is no greater than 0. Combining with (3.10) that the left-hand side of (5.3) is no less than 0, we conclude both sides of (5.3) equal to 0. Then the continuity of non-negative function $(M - bN)\phi$ implies

$$(M - bN)\phi \equiv 0, \quad \forall x \in B^n. \quad (5.4)$$

Taking into account that the range of ϕ strictly exceeds 0, we attain

$$M - bN \equiv 0, \quad \forall x \in B^n. \quad (5.5)$$

Again by (3.10), if $1 < q < \frac{n}{n-2}$, we derive $|\nabla v| \equiv 0$ in B^n . Thus v is constant.

On the other hand, if $0 < \lambda < \frac{1}{q-1}$, then the right-hand side of (5.3) equals to 0 forces

$$\int_{S^{n-1}} v^a |\nabla_{S^{n-1}} v|^2 \equiv 0.$$

Thus v is constant on S^{n-1} . So u is a harmonic function with constant boundary value, which must be constant by maximum principle.

It remains to discuss the case $q = \frac{n}{n-2}$ and $\lambda = \frac{1}{q-1}$. In this case, (5.5) and (3.10) only implies $|E| = 0$. Putting $a = -\frac{q+1}{q-1}$ and $b = \frac{q-1}{2q}$ into the definition (3.5) of E , we have for any $1 \leq i, j \leq n$,

$$v_{ij} = \frac{\Delta v}{n} \delta_{ij}. \quad (5.6)$$

By taking derivative on (5.6) and taking a sum, we derive for any $1 \leq i \leq n$,

$$(\Delta v)_i = \sum_{j=1}^n v_{jji} = \sum_{j=1}^n v_{ijj} = \sum_{j=1}^n \left(\frac{\Delta v}{n} \delta_{ij}\right)_j = \frac{1}{n} \sum_{j=1}^n (\Delta v)_j \delta_{ij} = \frac{1}{n} (\Delta v)_i.$$

Since $n \geq 3$, we have $(\Delta v)_i = 0$ for any $1 \leq i \leq n$. Thus Δv is constant. Suppose $\Delta v \equiv 2nr$ for some $r \in \mathbb{R}$, then from (5.6), we obtain

$$v(x) = r|x|^2 + \langle \zeta, x \rangle + s \quad \forall x \in B^n, \quad (5.7)$$

for some $s \in \mathbb{R}$ and $\zeta \in \mathbb{R}^n$. Direct computations imply

$$\nabla v = 2rx + \zeta \quad \forall x \in B^n. \quad (5.8)$$

Putting (5.7) and (5.8) into (2.1), we get the relations between r, s, ζ that

$$4rs = |\zeta|^2, \quad (5.9)$$

$$r = s - \frac{2}{n-2}. \quad (5.10)$$

Since v is a positive function, necessarily, we have $v(0) > 0$, i.e. $s > 0$. Combining with (5.9) and (5.10), we solve

$$r = \sqrt{\left(\frac{1}{n-2}\right)^2 + \frac{1}{4}|\zeta|^2} - \frac{1}{n-2}, \quad s = \sqrt{\left(\frac{1}{n-2}\right)^2 + \frac{1}{4}|\zeta|^2} + \frac{1}{n-2}. \quad (5.11)$$

Using the fact that for any $\zeta \in \mathbb{R}^n$, there is a unique $\xi \in B^n$, s.t.

$$\zeta = -\frac{4}{n-2} \frac{\xi}{1-|\xi|^2}. \quad (5.12)$$

Putting (5.12) into (5.11), we derive

$$r = \frac{2}{n-2} \frac{|\xi|^2}{1-|\xi|^2}, \quad s = \frac{2}{n-2} \frac{1}{1-|\xi|^2}. \quad (5.13)$$

Then putting (5.12) and (5.13) into (5.7), we obtain

$$v(x) = \frac{2}{n-2} \frac{1 + |\xi|^2|x|^2 - 2\langle \xi, x \rangle}{1 - |\xi|^2}, \quad (5.14)$$

i.e.

$$u(x) = \left(\frac{n-2}{2} \frac{1 - |\xi|^2}{1 + |\xi|^2|x|^2 - 2\langle \xi, x \rangle} \right)^{\frac{n-2}{2}}, \quad (5.15)$$

for some $\xi \in B^n$. We complete Theorem 1.1. □

Combing Guo-Hang-Wang [5] in $n = 2$ and Theorem 1.1 in $n \geq 3$, the conjecture is true. And thus we obtain a new proof of Beckner's inequality (1.5) by integral method.

Corollary 5.1. *The following inequalities hold*

$$|S^{n-1}|^{\frac{q-1}{q+1}} \left(\int_{S^{n-1}} u^{q+1} \right)^{\frac{2}{q+1}} \leq (q-1) \int_{B^n} |\nabla u|^2 + \int_{S^{n-1}} u^2, \quad \forall u \in C^\infty(\bar{B}^n),$$

provided $1 < q < \infty$, if $n = 2$, and $1 < q \leq \frac{n}{n-2}$, if $n \geq 3$.

Proof. We define the Sobolev quotient of a function $u \in H^1(B^n) - \{0\}$ as

$$Q_{\lambda,q}(u) := \frac{\int_{B^n} |\nabla u|^2 + \lambda \int_{S^{n-1}} u^2}{\left(\int_{S^{n-1}} |u|^{q+1} \right)^{\frac{2}{q+1}}}. \quad (5.16)$$

For the case $1 < q < \frac{n}{n-2}$, the trace operator $H^1(B^n) \rightarrow L^{q+1}(S^{n-1})$ is compact. Thus the minimization problem

$$S_{\lambda,q} := \inf_{u \in H^1(B^n)} Q_{\lambda,q}(u) \quad (5.17)$$

is achieved by smooth positive functions which satisfies (1.8). By use of the conjecture, such minimizer u must be

$$u(x) \equiv \lambda^{\frac{1}{q-1}}, \quad \forall x \in B^n. \quad (5.18)$$

Putting (5.18) into (5.16), since the constant achieves $S_{\lambda,q}$, we have for any $u \in C^\infty(\bar{B}^n)$,

$$\frac{\int_{B^n} |\nabla u|^2 + \lambda \int_{S^{n-1}} u^2}{\left(\int_{S^{n-1}} |u|^{q+1} \right)^{\frac{2}{q+1}}} \geq \lambda |S^{n-1}|^{\frac{q-1}{q+1}}, \quad \forall 0 < \lambda \leq \frac{1}{q-1}. \quad (5.19)$$

Let $\lambda = \frac{1}{q-1}$, we obtain (1.5) while $1 < q < \frac{n}{n-2}$.

For the critical case $q = \frac{n}{n-2}$, we follow the method of continuity of Aubin and Trudinger, see for example [6]. It suffices to show the function $q \mapsto S_{\lambda,q}$ is continuous on the left at $q = \frac{n}{n-2}$. To prove, for any $\epsilon > 0$, $\exists u_1 \in H^1(B^n)$, s.t.

$$S_{\lambda, \frac{n}{n-2}} \geq Q_{\lambda, \frac{n}{n-2}}(u_1) - \frac{\epsilon}{2}. \quad (5.20)$$

Since the function $q \mapsto Q_{\lambda,q}(u_1)$ is continuous in $(1, \frac{n}{n-2}]$. $\exists \delta > 0$, s.t. $\forall q' \in (\frac{n}{n-2} - \delta, \frac{n}{n-2}]$, we have

$$Q_{\lambda, \frac{n}{n-2}}(u_1) \geq Q_{\lambda, q'}(u_1) - \frac{\epsilon}{2}. \quad (5.21)$$

Combining with (5.20) and (5.21), we obtain

$$S_{\lambda, \frac{n}{n-2}} \geq Q_{\lambda, \frac{n}{n-2}}(u_1) - \frac{\epsilon}{2} \geq Q_{\lambda, q'}(u_1) - \epsilon \geq S_{\lambda, q'} - \epsilon. \quad (5.22)$$

On the other hand, assume $u_2 \in H^1(B^n)$ satisfies

$$S_{\lambda, q'} \geq Q_{\lambda, q'}(u_2) - \frac{\epsilon}{2}. \quad (5.23)$$

By Hölder inequality, we have

$$Q_{\lambda, q'}(u_2) = \frac{\int_{B^n} |\nabla u_2|^2 + \lambda \int_{S^{n-1}} u_2^2}{\left(\int_{S^{n-1}} |u_2|^{q'+1} \right)^{\frac{2}{q'+1}}} \geq \frac{\int_{B^n} |\nabla u_2|^2 + \lambda \int_{S^{n-1}} u_2^2}{\left(\int_{S^{n-1}} |u_2|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{n-1}}} |S^{n-1}|^{\left(\frac{n-2}{n-1} - \frac{2}{q'+1} \right)}. \quad (5.24)$$

Since the function $q' \mapsto |S^{n-1}|^{\left(\frac{n-2}{n-1} - \frac{2}{q'+1} \right)}$ is continuous in q' , we may decrease the above $\delta > 0$, s.t. $\forall q' \in (\frac{n}{n-2} - \delta, \frac{n}{n-2}]$, the following inequalities hold

$$\frac{\int_{B^n} |\nabla u_2|^2 + \lambda \int_{S^{n-1}} u_2^2}{\left(\int_{S^{n-1}} |u_2|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{n-1}}} |S^{n-1}|^{\left(\frac{n-2}{n-1} - \frac{2}{q'+1} \right)} \geq S_{\lambda, \frac{n}{n-2}} |S^{n-1}|^{\left(\frac{n-2}{n-1} - \frac{2}{q'+1} \right)} \geq S_{\lambda, \frac{n}{n-2}} - \frac{\epsilon}{2}. \quad (5.25)$$

Combine with (5.23), (5.24) and (5.25), we obtain

$$S_{\lambda, q'} \geq S_{\lambda, \frac{n}{n-2}} - \epsilon. \quad (5.26)$$

By (5.22), (5.26) and the arbitrariness of ϵ , the function $q \mapsto S_{\lambda,q}$ is continuous on the left at $q = \frac{n}{n-2}$. Thus

$$S_{\lambda, \frac{n}{n-2}} = \lim_{q \rightarrow (\frac{n}{n-2})^-} S_{\lambda, q} = \lim_{q \rightarrow (\frac{n}{n-2})^-} \lambda |S^{n-1}|^{\frac{q-1}{q+1}} = \lambda |S^{n-1}|^{\frac{1}{n-1}}. \quad (5.27)$$

We finish the new proof of Beckner's inequality. □

Acknowledgment. The authors were partially supported by NSFC Grant No.11831005. We would like to express our thanks to Yao Wan for his helpful discussions and suggestions.

REFERENCES

- [1] William Beckner, *Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality*, Ann. of Math. (2) **138** (1993), no. 1, 213–242.
- [2] Marie-Françoise Bidaut-Véron and Laurent Véron, *Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations*, Invent. Math. **106** (1991), no. 3, 489–539.
- [3] José F. Escobar, *Sharp constant in a Sobolev trace inequality*, Indiana Univ. Math. J. **37** (1988), no. 3, 687–698.
- [4] ———, *Uniqueness theorems on conformal deformation of metrics, Sobolev inequalities, and an eigenvalue estimate*, Comm. Pure Appl. Math. **43** (1990), no. 7, 857–883.

- [5] Qianqiao Guo, Fengbo Hang, and Xiaodong Wang, *Liouville type theorems on manifolds with nonnegative curvature and strictly convex boundary*, Math. Res. Lett. **28** (2021), no. 5, 1419–1439.
- [6] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, 2nd ed., Vol. 224, Springer-Verlag, Berlin, 1983.
- [7] Qianqiao Guo and Xiaodong Wang, *Uniqueness results for positive harmonic functions on $\overline{\mathbb{B}^n}$ satisfying a nonlinear boundary condition*, Calc. Var. Partial Differential Equations **59** (2020), no. 5, Paper No. 146, 8.
- [8] Daowen Lin and Qianzhong Ou, *Liouville type theorems for positive harmonic functions on the unit ball with a nonlinear boundary condition*, Calc. Var. Partial Differential Equations **62** (2023), no. 1, Paper No. 34, 13.
- [9] P.-L. Lions, *The concentration-compactness principle in the calculus of variations. The limit case. II*, Rev. Mat. Iberoamericana **1** (1985), no. 2, 45–121.
- [10] Robert C. Reilly, *Applications of the Hessian operator in a Riemannian manifold*, Indiana Univ. Math. J. **26** (1977), no. 3, 459–472.
- [11] Xiaodong Wang, *On compact Riemannian manifolds with convex boundary and Ricci curvature bounded from below*, J. Geom. Anal. **31** (2021), no. 4, 3988–4003.

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, P.R. CHINA

Email address: gpx21@mails.tsinghua.edu.cn

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, P.R. CHINA

Email address: lih2@tsinghua.edu.cn