L^2 TO L^p BOUNDS FOR SPECTRAL PROJECTORS ON THIN INTERVALS IN RIEMANNIAN MANIFOLDS

PIERRE GERMAIN

ABSTRACT. Given a Riemannian manifold with Laplace-Beltrami operator Δ , consider the projector on a thin interval $[\lambda - \delta, \lambda + \delta]$ associated to the self-adjoint operator $\sqrt{-\Delta}$. Viewed as an operator from $L^2(M)$ to $L^p(M)$, what is its operator norm? For $\delta = 1$, this question is fully answered by a celebrated theorem of Sogge, which applies to any manifold. For $\delta < 1$, the global geometry of the manifold comes into play, and connections to a number of mathematical fields (such as Differential Geometry, Combinatorics, Number Theory) appear, but the problem remains mostly open. The aim of this article is to review known results, focusing on cases exhibiting symmetry (where the regime where δ is polynomially small in λ becomes approachable) and emphasizing harmonic analytic rather than microlocal methods.

Contents

1.	Introduction	l
2.	Sogge's universal estimates for $\delta = 1$	2
3.	The Euclidean space and its quotients	6
4.	The hyperbolic space and its quotients	10
5.	Classical estimates implying spectral projector bounds	13
6.	$L^2 \to L^p$ projector bounds beyond complete smooth Riemannian manifolds	15
7.	Some interesting questions	15
References		17

1. INTRODUCTION

Consider a (smooth, complete) Riemannian manifold M, of dimension d, with Laplace-Beltrami operator Δ . Define the spectral projector $P_{\lambda,\delta}$ through functional calculus by

$$P_{\lambda,\delta} = \mathbf{1}_{[\lambda-\delta,\lambda+\delta]} \left(\sqrt{-\Delta}\right).$$

It is the orthogonal projector on the (generalized) eigenmodes of $\sqrt{-\Delta}$ with eigenvalue in $[\lambda - \delta, \lambda + \delta]$. We ask the following question:

What is the operator norm of $P_{\lambda,\delta}$ from $L^2(M)$ to $L^p(M)$?

A more classical question is to ask for L^p bounds of L^2 -normalized eigenfunctions on compact manifolds, which would correspond to $\delta = 0$. Similarly, the question above could be formulated in terms of L^p bounds for L^2 -normalized *almost*-eigenfunctions.

Why is it meaningful to consider the case $\delta > 0$? First, this has the advantage of applying to any manifold, be it compact or not, with discrete or continuous spectrum, and this more general viewpoint is helpful. Second, in the case of compact manifolds, the case $\delta = 0$ seems in general completely out of reach, while the case $\delta = 1$ is fully understood, as we will see; thus, interpolating

 $^{2000 \} Mathematics \ Subject \ Classification. \ \dots$

Key words and phrases. Riemannian manifolds, L^p estimates, spectral projectors, eigenfunctions, quasimodes.

between these two situations seems natural. Finally, considering a window of positive width encodes information on the distribution of the spectrum, together with the size of the eigenfunctions.

Before reviewing the theory, let us observe that the question of estimating $||P_{\lambda,\delta}||_{L^2 \to L^p}$ can be asked in different ways. First, an immediate TT^* argument shows that

(1.1)
$$\|P_{\lambda,\delta}\|_{L^2 \to L^p}^2 = \|P_{\lambda,\delta}\|_{L^{p'} \to L^p}.$$

Second, it is essentially equivalent to estimate the operator norm of $P_{\lambda,\delta}$, or that of the smoothed out version

$$P_{\lambda,\delta}^{\flat} = \chi\left(\frac{\sqrt{-\Delta} - \lambda}{\delta}\right)$$

where χ is a cutoff function; we will use indiscriminately all these formulations¹.

A beautiful textbook presentation of the theory for $\delta = 1$ can be found in [92], see also [91, 105, 106]. In this review note, we will try to give an overview of progress on the case $\delta < 1$, in particular in the case where δ is polynomially small in λ .

A very closely connected topic is that of Quantum Unique Ergodicity for negatively curved manifolds, or for general manifolds the description of possible weak- L^2 limits of eigenfunctions. This is in many ways an L^2 version of the L^p -problems which will occupy us in the present review, with much of the same underlying phenomenology. We refer to [2, 81] for surveys on Quantum Unique Ergodicity.

2. Sogge's universal estimates for $\delta = 1$

2.1. Sogge's theorem. In this subsection, we consider general compact manifolds M, without any further restriction. Denoting by (λ_j) and (φ_j) the eigenvalues and eigenvectors of the Laplacian on M, Hörmander's bound [55], see also [5, 74] states that

(2.1)
$$\|\varphi_j\|_{L^{\infty}} \lesssim \lambda_j^{\frac{d-1}{2}} \|\varphi_j\|_{L^2}$$

(this estimate arose in connection with the Weyl law, to which we will come back in Section 5.1).

This classical bound was subsumed in the following much stronger result of Sogge: it covers any p > 2, allows for spectral projectors instead of eigenfunctions, and it is not only providing an upper bound, but also a lower bound. It will be convenient to denote

(2.2)
$$\gamma(p) = \max\left[\frac{d-1}{2} - \frac{d}{p}, \frac{d-1}{2}\left(\frac{1}{2} - \frac{1}{p}\right)\right],$$

since this exponent will appear throughout the present review.

Theorem 2.1 (Sogge). If M is a complete Riemannian manifold with bounded geometry, then

(2.3)
$$\|P_{\lambda,1}\|_{L^2 \to L^p} \lesssim \lambda^{\gamma(p)} \quad \text{if } 2 \le p \le \infty.$$

Furthermore, there exists a constant R_0 such that: for any λ_0 , there exists λ with $|\lambda - \lambda_0| < R_0$ and

$$||P_{\lambda,1}||_{L^2 \to L^p} \gtrsim \lambda^{\gamma(p)} \qquad \text{if } 2 \le p \le \infty.$$

The exponent $\gamma(p)$ is defined as a maximum in (2.2), and which of the two terms in this maximum is dominant depends on how p compares compares to the Stein-Tomas exponent defined by

$$p_{ST} = \frac{2(d+1)}{d-1}.$$

More precisely

¹Up to logarithmic factors, it is also equivalent to consider boundedness of the resolvent operator $(\Delta - z)^{-1}$, $\Im m z \neq 0$.



FIGURE 1. Consider a coordinate patch on a Riemannian manifold. One can then contruct spectrally localized functions $(P_{\lambda,\delta}f = f)$ which concentrate their L^p mass in a neighborhood of a given point (red), or a given geodesic (blue). These functions saturate the value $\lambda^{\gamma(p)}$ for the operator norm in (2.3) up to a bounded factor in the regimes $p > p_{ST}$ and $p < p_{ST}$, respectively.

- For $p \ge p_{ST}$, $P_{\lambda,1}$ is in the *point-focusing regime*: the term $\lambda^{\frac{d-1}{2}-\frac{d}{p}}$ is dominant, and examples of functions which focus at a point can be shown to achieve this value (up to a bounded factor) for $\frac{\|P_{\lambda,1}f\|_{L^p}}{\|f\|_{L^2}}$.
- For $2 \le p \le p_{ST}$, $P_{\lambda,1}$ is in the geodesic-focusing regime: the term $\lambda^{\frac{d-1}{2}\left(\frac{1}{2}-\frac{1}{p}\right)}$ is dominant. Functions can be constructed which focus on a geodesic and achieve this value (up to a bounded factor) for $\frac{\|P_{\lambda,1}f\|_{L^p}}{\|f\|_{L^2}}$.

Proof of Theorem 2.3. A detailed proof can be found in [88, 92] in the compact case, see also [?] for the simple extension to the complete case. A sketch of the upper bound is as follows.

First, one can reduce matters to an operator $\tilde{P}_{\lambda,1}$ which is a slightly modified version of $P_{\lambda,1}$, simpler to estimate since it is local. Through pseudodifferential calculus, one can then obtain a very precise description of the kernel of $\tilde{P}_{\lambda,1}$:

(2.4)
$$\widetilde{P}_{\lambda,1}f(x) = \int_M K_{\lambda,1}(x,y)f(y)\,dx, \qquad K_{\lambda,1}(x,y) = \lambda^{\frac{d-1}{2}}a(x,y)e^{i\lambda\psi(x,y)} + O(\lambda^{-\infty}),$$

where $a \in \mathcal{C}_0^{\infty}$. The desired result follows by interpolation between the following Lebesgue exponents

- For p = 2, the bound O(1) is an immediate consequence of the definition of $\widetilde{P}_{\lambda,1}$.
- For $p = \infty$, the bound $O(\lambda^{\frac{d-1}{2}})$ follows immediately from the formula (2.4) and the Cauchy-Schwarz inequality.
- Finally, $p = p_{ST}$ is the critical exponent. The properties of the phase $\psi(x, y)$ now come into play: indeed, it satisfies the Carleson-Sjölin condition, which, roughly speaking, requires that its Hessian has rank n - 1. The $L^2 \to L^p$ bounds on Carleson-Sjölin operators (due in dimension 2 to Carleson-Sjölin [32] and Hörmander [56], and in higher dimension to Stein [98]) then give the desired result. These bounds can be thought of as a variable coefficient generalization of the Stein-Tomas theorem, which will presented in Section 3.

As observed in [3], a consequence of the optimality of the Sogge estimates is the universal lower bound

(2.5)
$$\|P_{\lambda,\delta}\|_{L^2 \to L^p} \gtrsim \lambda^{\gamma(p)} \delta^{\frac{1}{2}} \quad \text{if } 2 \le p \le \infty$$

(more precisely: for any λ_0, δ , there exists λ such that $|\lambda - \lambda_0| < R_0$, and such that the above holds).

2.2. The case of the sphere. As we saw, the universal Sogge estimate (2.3) corresponds to the case $\delta = 1$, and is sharp for any manifold. Is it possible to improve this bound by taking δ smaller? The example of the sphere shows that the answer is negative in general. Indeed, the spectrum of the Laplace-Beltrami operator on \mathbb{S}^d is given by $\{k(k+d-1), k \in \mathbb{Z}\}$, so that

$$P_{\lambda,\delta} = P_{\lambda,\frac{1}{2}}$$
 if $\delta < \frac{1}{2}$, $\lambda = \sqrt{k(k+d-1)}$, $k \in \mathbb{Z}$

(and λ sufficiently big). This implies that operator bounds for $P_{\lambda,\delta}$ and $P_{\lambda,\frac{1}{2}}$ are equal and that the universal Sogge estimate (2.3) will be saturated by eigenfunctions; the two following examples appear in [87] and are fundamental.

- In the point-focusing regime, zonal spherical harmonics saturate (2.3). By definition, zonal functions can be written as functions of the geodesic distance to a point on the sphere. Each eigenspace contains a zonal function, unique up to the symmetry group SO(d + 1). Normalizing this function in L², it reaches a size ~ λ^{d-1}/₂ on a set of diameter ~ λ⁻¹ around the pole, leading to the estimate ||f||_{L^p} ~ λ^{d-1}/₂ ^d/_p ||f||_{L²} for p > p_{ST}.
 In the geodesic-focusing regime, highest weight spherical harmonics saturate (2.3). Viewing
- In the geodesic-focusing regime, highest weight spherical harmonics saturate (2.3). Viewing the sphere \mathbb{S}^{d-1} as embedded in \mathbb{R}^{d+1} , they can be taken to be the trace on the sphere of $(x_1+ix_2)^k$. Normalizing this function in L^2 , it reaches a size $\sim \lambda^{\frac{d-1}{4}}$ on a neighbourhood of size $\sim \lambda^{-\frac{1}{2}}$ around the great circle $x_1^2 + x_2^2 = 1$, leading to the estimate $||f||_{L^p} \sim \lambda^{\frac{d-1}{2}(\frac{1}{2}-\frac{1}{p})}$ for $p < p_{ST}$.

2.3. When are Sogge's estimates optimal for eigenfunctions? We saw that, in the case of the sphere, the Sogge bound (2.3) agrees with the optimal bound for eigenfunctions, or in other words that the spectral projectors for $\delta = 0$ and $\delta = 1$ have comparable operator norms. Is it possible to characterize all manifolds for which this phenomenon occurs?

2.3.1. Point-focusing regime. A line of research initiated by Sogge and Zelditch [94, 95, 96, 90] and pursued by Canzani and Galkowski [29, 31] seeks conditions on a compact manifold M under which there exists, or not, a sequence of eigenfunctions φ_j associated to eigenvalues λ_j exhibiting maximum eigenfunction growth, namely

(2.6)
$$\|\varphi_j\|_{L^{\infty}} \sim \lambda_j^{\frac{d-1}{2}} \|\varphi_j\|_{L^2}.$$

Of course, this implies that $\|P_{\lambda,\delta}\|_{L^2 \to L^\infty} \gtrsim \lambda^{\frac{d-1}{2}}$ for any $\delta > 0$, along a sequence of λ .

While very powerful and precise results have been proved along these lines, we will give here a simple example of such a statement, which can be found in [93]. For $x \in M$, define $\mathcal{L}_x \subset S_x^*M$ (unit cotangent bundle) to be the set of directions $\xi \in T_x^*M$ such that the geodesic γ with initial data $\gamma(0) = x$, $\dot{\gamma}(0) = \xi$ returns to x. Then there cannot exists a sequence of eigenfunctions with maximal eigenfunction growth unless \mathcal{L}_x has positive measure for some $x \in M$.

This is a strong constraint, of course reminiscent of the case of the sphere; but it is not stable under perturbation. Indeed, it was showed [94] that for generic metrics, $\|\varphi_j\|_{L^{\infty}} = o(\lambda_j^{\frac{d-1}{2}})$.



FIGURE 2. On the sphere, zonal spherical harmonics concentrate most of their L^p mass close to the North pole (in red), and highest weight spherical harmonics concentrate most of their L^p mass close to the equator (in blue). These eigenfunctions of the Laplacian saturate the Sogge bound (2.3) for $p > p_{ST}$ and $p < p_{ST}$ respectively. Of course, the North pole and the equator do not play a distinguished role, since the Laplacian is invariant under O(d+1), and this figure can be rotated.

Finally, we mention the original point of view in [100], which relates poinwise L^{∞} growth to correlations between eigenfunctions at different points of the manifold.

2.3.2. Geodesic-focusing regime. We call a geodesic stable if its associated Poincaré map only has eigenvalues of modulus 1. If there exists a stable closed geodesic, one can construct a sequence (λ_j) going to infinity and associated quasimodes (f_j) such that

$$\left\| \left[\Delta + \lambda_j^2 \right] f_j \right\|_{L^2} \lesssim_N \lambda_j^{-N} \quad \text{for any } j, N \in \mathbb{N};$$

furthermore, these quasimodes concentrate within $\frac{1}{\sqrt{\lambda_j}}$ of the geodesic, and are bounded pointwise

by $\lambda_j^{\frac{d-1}{4}}$. To the best of our knowledge, this exact statement does not appear in the literature, but should follow from the construction of quasimodes supported on stable geodesics [51, 77, 40, 1].

As a consequence of the existence of these quasimodes, the spectral projectors can be bounded from below by

$$\|P_{\lambda_j,\delta}\|_{L^2 \to L^p} \gtrsim_N \lambda_j^{\frac{d-1}{2}\left(\frac{1}{2} - \frac{1}{p}\right)} \quad \text{if } \delta > \lambda_j^{-N}.$$

Note that the right-hand side matches the universal bound of Sogge in the regime $p < p_{ST}$.

The preceding discussion suggests that, for "most" manifolds, the bounds on $||P_{\lambda,\delta}||_{L^2 \to L^p}$ should improve as δ decreases from $\delta = 1$. In the remainder of this review article, we will try and characterize this improvement in some specific examples.

3. The Euclidean space and its quotients

3.1. The Euclidean space. Consider the extension operator mapping functions on the sphere \mathbb{S}^{d-1} to functions on the whole space \mathbb{R}^d through the formula

$$f \mapsto \mathcal{F}\left[f \, d\sigma_{|\mathbb{S}^{d-1}}\right],$$

where \mathcal{F} is the Fourier transform and $d\sigma_{|\mathbb{S}^{d-1}}$ is the surface measure on \mathbb{S}^{d-1} . The Stein-Tomas theorem establishes the boundedness of this operator from $L^2(\mathbb{S}^{d-1})$ to $L^p(\mathbb{R}^d)$ if $p \ge p_{ST}$. Recently, several works have investigated the existence and structure of functions which saturate the operator norm [85, 42]. The Stein-Tomas theorem can be reformulated in terms of our spectral projectors:

Theorem 3.1 (Stein-Tomas [99, 102]). For any $\lambda > 1$, $\delta < 1$,

(3.1)
$$||P_{\lambda,\delta}||_{L^2 \to L^p} \sim \lambda^{\frac{d-1}{2} - \frac{d}{p}} \delta^{\frac{1}{2}} + \lambda^{\frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{p}\right)} \delta^{\frac{d+1}{2} \left(\frac{1}{2} - \frac{1}{p}\right)} \quad \text{if } 2 \le p \le \infty.$$

Before sketching the proof of this theorem, we remark that the Stein-Tomas estimate above can be extended to asymptotically conic, non-trapping manifolds [50]; extensions to non-elliptic, constant coefficients operators can also be considered [68].

Proof. By scaling, this theorem is a consequence of the universal Sogge estimate (2.3); but it came first, and can be proved by simpler means. We sketch the proof in dimension 3, where formulas are cleaner. The smoothed out operator $P_{\lambda,\delta}^{\chi}$ defined as $\chi\left(\frac{\sqrt{-\Delta}-\lambda}{\delta}\right)$ (where $\chi \in \mathcal{C}_0^{\infty}$) can be expressed through a radial convolution kernel

(3.2)
$$P_{\lambda,\delta}^{\chi}f(x) = \int_{\mathbb{R}^3} K_{\lambda,\delta}(|x-y|)f(y)\,dy, \quad \text{with} \quad K_{\lambda,\delta}(r) = \delta\widehat{\chi}(\delta r)\lambda\frac{\sin(\lambda r)}{r}.$$

The proof consists in splitting $K_{\lambda,\delta}$ into

$$K_{\lambda,\delta}(r) = \sum_{j} K^{j}_{\lambda,\delta}(r),$$
 and accordingly $P_{\lambda,\delta} = \sum_{j} P^{j}_{\lambda,\delta}$

where each $K_{\lambda,\delta}^j$ is smoothly localized where $r \sim 2^j$. Then the operator norm of $P_{\lambda,\delta}^j$ can be estimated in two ways

• $\|P_{\lambda,\delta}^{j}\|_{L^{1}\to L^{\infty}} \lesssim \|K_{\lambda,\delta}^{j}\|_{\infty}$ • $\|P_{\lambda,\delta}^{j}\|_{L^{2}\to L^{2}} \lesssim \|m_{\lambda,\delta}^{j}\|_{\infty}$, where $m_{\lambda,\delta}^{j}$ is the Fourier symbol of $P_{\lambda,\delta}^{j}$.

By interpolation, this gives a bound for $||P_{\lambda,\delta}^j||_{L^{p'}\to L^p}$, and, after filling out the estimates and summing over j, the desired bound for $||P_{\lambda,\delta}||_{L^{p'}\to L^p}$ if $p \neq p_{ST}$. The critical point p_{ST} requires complex interpolation.

Just like for Sogge's estimate (2.3), one can distinguish two regimes in (3.1):

- In the point-focusing regime $p \ge p_{ST}$, the dominant term is $\lambda^{\frac{d-1}{2} \frac{d}{p}} \delta^{\frac{1}{2}}$. It is achieved up to a bounded factor for the *spherical example* f (the terminology coming from the radial symmetry of f) given in Fourier by $\hat{f}(\xi) = \mathbf{1}_{[\lambda \delta, \lambda + \delta]}(|x|)$, which has a sharp peak at the origin.
- In the geodesic-focusing regime $p \leq p_{ST}$, the dominant term is $\lambda^{\frac{d-1}{2}\left(\frac{1}{2}-\frac{1}{p}\right)}\delta^{\frac{d+1}{2}\left(\frac{1}{2}-\frac{1}{p}\right)}$; it can be achieved up to a bounded factor by the so-called *Knapp example* which we now describe. It is the function defined, in Fourier space, by $\hat{f} = \mathbf{1}_R$, where R is a rectangle of dimensions $\sim (\sqrt{\lambda\delta})^{d-1} \times \delta$ contained in the annulus $\{\xi, \lambda \delta < |\xi| < \lambda + \delta\}$. In physical space, it is mostly supported on the dual rectangle \hat{R} , of dimensions $((\lambda\delta)^{-\frac{1}{2}})^{d-1} \times \delta$.

These two examples are depicted in Figure 3, both in physical space and in Fourier space.



FIGURE 3. The spherical example (red) and the Knapp example (blue) for the Euclidean space represented in frequency and physical space: the Fourier transform exchanges the left and right columns. In frequency space, the supports are contained within the annulus with inner and outer radii $\lambda - \delta$ and $\lambda + \delta$, either the full annulus or a rectangle contained in it. In physical space, the L^p mass concentrates at a point or on the dual rectangle, respectively.

3.2. Tori. We now turn to quotients of the Euclidean space, namely tori. As in the whole space, we can rely on Fourier analysis (Fourier series rather than the Fourier transform) to carry out the analysis, but as we shall see, it becomes much more challenging.

Choosing a family of independent vectors (e_i) of \mathbb{R}^d generating the lattice $\Lambda = \mathbb{Z}e_1 + \cdots + \mathbb{Z}e_d$, we will denote $\mathbb{T}^d = \mathbb{T}^d_{\Lambda}$ for the quotient $\mathbb{R}^d/(\mathbb{Z}e_1 + \cdots + \mathbb{Z}e_d)$.

3.2.1. The case $p = \infty$. A basis of eigenfunctions of the Laplacian on \mathbb{T}^d is provided by complex exponentials of the type $e^{ik \cdot x}$, where k belongs to the dual lattice Λ^* . It is then easy to see that

$$\|P_{\lambda,\delta}\|_{L^1 \to L^\infty} = \#\{k \in \Lambda^*, \, \lambda - \delta < |k| < \lambda + \delta\},\$$

which brings us to a lattice point counting problem. This is a variant of the more classical problem of estimating the number of lattive points in a sphere of radius λ :

$$N(\lambda) = \#\{k \in \Lambda^*, |k| < \lambda\}.$$

To leading order, $N(\lambda)$ equals $C_{\Lambda}\lambda^d$, and the problem is to bound the error $E(\lambda) = |P(\lambda) - C_{\Lambda}\lambda^d|$. A full understanding of optimal bounds on $E(\lambda)$ is still missing in dimensions 2 and 3; we refer the reader to [49, 61, 73, 76] for entry points into the literature.

Though bounds on $E(\lambda)$ imply bounds on $\|P_{\lambda,\delta}\|_{L^1\to L^\infty}$ by the above characterization, this cannot lead to an optimal answer, and the questions of estimating $E(\lambda)$ and $\|P_{\lambda,\delta}\|_{L^1\to L^\infty}$ are not equivalent.

3.2.2. The case $\delta = 0$ (eigenfunctions). The earliest L^p bound on eigenfunctions of the Laplacian on the torus is the famous result of Cooke-Zygmund [35, 107] that, on \mathbb{T}^2 , an eigenfunction φ of the Laplacian satisfies

$$\|\varphi\|_{L^4} \lesssim \|\varphi\|_{L^2}.$$

Whether this estimate can be extended to any p > 4 has remained an open problem. If $\Lambda = \mathbb{Z}^d$, Bourgain [18] conjectured the general bound for an eigenfunction φ associated to the eigenvalue λ

(3.3)
$$\|\varphi\|_{L^p} \lesssim \lambda^{\frac{d}{2}-1-\frac{d}{p}} \|\varphi\|_{L^2} \quad \text{for } p > p^* = \frac{2d}{d-2}$$

This conjecture was the subject of a number of subsequent papers [19, 20, 21] culminating in the ℓ^2 decoupling estimate of Bourgain and Demeter [22]. Though the ℓ^2 decoupling theorem was a landmark result in harmonic analysis, it only proved the conjecture in the range $p \geq \frac{2(d-1)}{d-2}$.

3.2.3. General values of p and δ . The conjecture (3.3) gives L^p bounds on eigenfunctions of the Laplacian on the torus. This is meaningful if $\Lambda = \mathbb{Z}^d$, but this question becomes trivial on generic tori, for which eigenspaces of the Laplacian have uniformly bounded dimension. A natural generalization consists in asking the question which is at the heart of the present review article, namely that of estimating $\|P_{\lambda,\delta}\|_{L^2\to L^p}$. The author and Myerson [45] proposed the following conjecture²

(3.4)
$$\|P_{\lambda,\delta}\|_{L^2 \to L^p} \lesssim \lambda^{\frac{d-1}{2} - \frac{d}{p}} \delta^{\frac{1}{2}} + (\lambda \delta)^{\frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{p}\right)} \quad \text{if } 2 \le p \le \infty \text{ and } \delta > \lambda^{-1}.$$

As in the case $\delta = 1$, and in the case of the Euclidean space, the two summands on the right-hand side are achieved up to a bounded factor for eigenfunctions focusing at a point, or on a geodesic. However, the dominant term does not depend on how p compares to p_{ST} ; it is rather a condition depending on p, λ , and δ .

- In the point-focusing regime, the dominant term is $\lambda^{\frac{d-1}{2}-\frac{d}{p}}\delta^{\frac{1}{2}}$. This operator norm for $P_{\lambda,\delta}$ can be achieved up to a bounded factor for functions which are evenly spread amongst all Fourier modes $k \in \Lambda^*$ such that $\lambda - \delta < |k| < \lambda + \delta$: this is the torus analog of the spherical example.
- In the geodesic-focusing regime, the dominant term is $(\lambda \delta)^{\frac{d-1}{2} \left(\frac{1}{2} \frac{1}{p}\right)}$. This can be achieved up to a bounded factor for functions which, in physical space, are focusing on a closed geodesic of \mathbb{T}^d : this is the torus analog of the Knapp example.

The spherical and Knapp examples for the torus are depicted in Figure 4, both in physical and in frequency space.

In the following theorem, we do not attempt to gather all the cases for which the conjecture is known to be true, but rather record two relevant instances where it holds. For the current state of the art, the interested reader can consult [37, 46, 54], besides the papers which have already been mentioned.

Theorem 3.2. The conjecture (3.4) is verified

(i) for $\Lambda = \mathbb{Z}^d$, if $p \ge \frac{2(d-1)}{d-3}$ and $d \ge 4$ (ii) for general Λ , if $p \le p_{ST}$.

²Notice that it implies the conjecture (3.3) by choosing $\Lambda = \mathbb{Z}^d$ and $\delta \sim \lambda^{-1}$.



FIGURE 4. The spherical example (red) and the Knapp example (blue) for the torus represented in frequency and physical space: the Fourier series exchange the left and right columns. In frequency space, the supports are contained within the annulus with inner and outer radii $\lambda - \delta$ and $\lambda + \delta$, either the full annulus or a rectangle contained in it. In physical space, the L^p mass concentrates close at a point or on the dual rectangle, respectively.

Proof. The first assertion is due to Bourgain and Demeter [22]; its proof combines the ℓ^2 decoupling theorem with arithmetic properties of the lattice \mathbb{Z}^d .

The second assertion turns out to be a rather direct consequence of ℓ^2 decoupling. To simplify notations, we focus on the case d = 2, for which $p_{ST} = 6$, and choose the lattice to be $\Lambda = \mathbb{Z}^2$. By interpolation with the trivial case p = 2, it suffices to bound the L^6 norm of a function $f \in L^2(\mathbb{T}^2)$ which is Fourier supported on an annulus of inner radius $\lambda - \delta$ and outer radius $\lambda + \delta$. We write f as the sum of its Fourier series $f = \sum a_k e^{2\pi i k \cdot x}$ and rescale variables to $X = \lambda x$ and $K = k/\lambda$ in order to apply the ℓ^2 decoupling theorem:

$$\|f\|_{L^6(\mathbb{T}^2)} = \left\|\sum_{k\in\mathbb{Z}^2} a_k e^{2\pi i k\cdot x}\right\|_{L^6(\mathbb{T}^2)} \lesssim \left(\frac{\delta}{\lambda}\right)^{\frac{1}{3}} \left\|\phi\left(\frac{\delta X}{\lambda}\right)\sum_{K\in\mathbb{Z}^2/\lambda} a_{\lambda K} e^{2\pi i K\cdot X}\right\|_{L^6(\mathbb{R}^2)}$$

...

where the cutoff function ϕ can be chosen to have compactly supported Fourier transform. As a result, the Fourier transform of the function on the right-hand side is supported on a δ/λ neighborhood of \mathbb{S}^{d-1} . It can be written as a sum of functions which are supported in Fourier

on caps θ with dimensions $\sim \frac{\delta}{\lambda} \times \frac{\delta^{\frac{1}{2}}}{\lambda^{\frac{1}{2}}}$. These caps are almost disjoint rectangles which cover the annulus, and to which we associate a smooth partition of unity (χ_{θ}) . By ℓ^2 decoupling, the L^6 norm above is bounded by

$$\lesssim_{\epsilon} \left(\frac{\delta}{\lambda}\right)^{\frac{1}{3}-\epsilon} \left(\sum_{\theta} \left\| \phi\left(\frac{\delta X}{\lambda}\right) \sum_{K \in \mathbb{Z}^2/\lambda} \chi_{\theta}(K) a_{\lambda K} e^{2\pi i K \cdot X} \right\|_{L^6(\mathbb{R}^2)}^2 \right)^{\frac{1}{2}}$$

At this point, we use the inequality $\|g\|_{L^p(\mathbb{R}^2)} \lesssim \|g\|_{L^2(\mathbb{R}^2)} |\operatorname{Supp} \widehat{g}|^{\frac{1}{2}-\frac{1}{p}}$, which is valid for $p \geq 2$ and follows by applying successively the Hausdorff-Young, Hölder, and the Plancherel inequalities. We use this inequality for $g_{\theta}(X) = \phi\left(\frac{\delta X}{\lambda}\right) \sum_{K} \chi_{\theta}(K) a_{\lambda K} e^{2\pi i K \cdot X}$. Since each cap contains at most $\lambda^{\frac{1}{2}} \delta^{\frac{1}{2}}$ rescaled lattice points, its Fourier transform is supported on the union of at most $\lambda^{\frac{1}{2}} \delta^{\frac{1}{2}}$ balls of radius $O(\delta/\lambda)$, giving $|\operatorname{Supp} \widehat{f}| \leq \lambda^{-\frac{1}{2}} \delta^{\frac{3}{2}}$. Thus the L^6 norm we are trying to bound is less than

$$\lesssim \left(\frac{\delta}{\lambda}\right)^{\frac{1}{3}-\epsilon} (\lambda^{-\frac{1}{2}}\delta^{\frac{3}{2}})^{\frac{1}{3}} \left(\sum_{\theta} \left\|\phi\left(\frac{\delta X}{\lambda}\right)\sum_{K\in\mathbb{Z}^{2}/\lambda}\chi_{\theta}(\lambda K)a_{\lambda K}e^{2\pi i K\cdot X}\right\|_{L^{2}(\mathbb{R}^{2})}^{2}\right)^{\frac{1}{2}}$$

By almost orthogonality and periodicity of the Fourier series, this is bounded by $\lambda^{\epsilon}(\lambda\delta)^{\frac{1}{6}} ||f||_{L^2(\mathbb{T}^2)}$, which is the desired estimate.

Before completing this section on Euclidean tori, we mention two related directions. First, the two-dimensional Euclidean cylinder is considered in [44]; in this case, there are no difficulties related to number theoretical questions, and the conjecture (3.4) could be proved. Second, Euclidean tori are the simplest examples of manifolds with integrable geodesic flow; this broader class was investigated in [17, 103].

4. The hyperbolic space and its quotients

4.1. Eigenfunctions of compact manifolds of nonpositive curvature. Manifolds of nonpositive curvature form a much broader class than the hyperbolic space and its quotients (it even includes Euclidean tori), and a natural context in which improvements to the Hörmander bound (2.1) are expected. A long line of research [8, 53, 52, 11, 12, 13] focused on this question through microlocal analysis, ultimately resulting in the following theorem.

Theorem 4.1. If the manifold M is compact and has nonnegative curvature, then, denoting its eigenfunctions and eigenvalues by (φ_i, λ_i) , there holds

$$\|\varphi_j\|_{L^p} \lesssim \lambda_j^{\gamma(p)} (\log \lambda_j)^{-\kappa},$$

where $\kappa = \kappa(p) = \frac{1}{2}$ for $p > p_{ST}$, and $\kappa > 0$ for $p \in (2, p_{ST}]$.

The above theorem is fundamental in the analysis of the spectral resolution of the Laplacian: it deals with the hardest case, that of compact manifolds, and it does so without any symmetry assumptions, under the single geometric condition of nonpositive curvature. Similar improvements can actually be obtained in even more general contexts, see [31, 28, 30].

However, stronger bounds are expected: under chaoticity assumptions for the geodesic flow, the random wave model of Berry [9] leads to predicting a bound $\sqrt{\log \lambda_j}$ (see for instance [100] for the heuristic argument, and [4] for numerics which appear to support this conjecture).

Such a bound seems utterly out of reach of current analytic methods. However, in the hope of achieving some progress, we will now shift the focus and consider a more restricted class of examples, namely the hyperbolic space and its quotients, allowing also for infinite volume. 4.2. The hyperbolic space. The question of estimating $||P_{\lambda,\delta}||_{L^2 \to L^p}$ can be fully answered for the hyperbolic space. After partial results were obtained in [57], this was done in [33], and the optimal dependence of the bounds on p was proved in [43].

Theorem 4.2. If M is the hyperbolic space \mathbb{H}^d ,

$$||P_{\lambda,\delta}||_{L^2 \to L^p} \lesssim [(p-2)^{-1}+1]\lambda^{\gamma(p)}\delta^{\frac{1}{2}}.$$

The reader should by now not be surprised that the upper bound in this theorem comprises two regimes, as reflected in the definition of $\gamma(p)$ in 2.2: these are of course the point-focusing and the geodesic-focusing regime. Just like in the Euclidean case, a spherical example and a Knapp example can be constructed, which nearly saturate the bounds in the point-focusing and geodesic-focusing regimes respectively. These examples are depicted in Figure 5, both in physical space (hyperbolic space) and in frequency space (the appropriate analog of the Fourier transform being provided by the Helgason-Fourier transform).

Comparing this bound to the Euclidean case or the sphere, it appears that the strongest bound is reached in the hyperbolic case, the weakest in the spherical case, with the Euclidean case being a middle ground. This can be explained, at least partially, in relation to the stability of geodesics: the more stable geodesics are, the easier it is for almost-eigenfunctions to focus on them, and therefore the weaker the bounds for $||P_{\lambda,\delta}||_{L^2 \to L^p}$.

Proof of Theorem 4.2. We sketch the proof in dimension 3, where formulas are cleaner. The operator $P_{\lambda,\delta}$ can then be expressed through a radial convolution kernel

$$P_{\lambda,\delta}f(x) = \int_{\mathbb{H}^3} K_{\lambda,\delta}(\operatorname{dist}(x,y))f(y)\,dy, \quad \text{with} \quad K_{\lambda,\delta}(r) = \delta\widehat{\chi}(\delta r)\lambda \frac{\sin(\lambda r)}{\sinh r}$$

This is quite similar to the Euclidean case (3.2) over small distances, but over large distances, the kernel decays exponentially. Nevertheless, it is possible to follow closely the scheme of the Euclidean proof: split the kernel dyadically, estimate all summands in the $L^2 \to L^p$ topology, and then conclude through real or complex interpolation.

4.3. Infinite area quotients. Only the case of dimension 2 has been explored so far, with the following theorem, which essentially asserts that the bounds are identical to the case of the hyperbolic space, up to possible suppolynomial factors.

Theorem 4.3 ([3]). Consider a geometrically finite hyperbolic quotient $M = \Gamma \setminus \mathbb{H}^2$ of the hyperbolic space by a Fuchsian subgroup. For any $\epsilon, N > 0$,

$$\|P_{\lambda,\delta}\|_{L^2 \to L^p} \lesssim_{p,\epsilon} \lambda^{\gamma(p)+\epsilon} \delta^{\frac{1}{2}} \quad if \ \delta > \lambda^{-N}.$$

Furthermore, ϵ and N can be taken equal to 0 and ∞ respectively if the exponent of convergence of N is $<\frac{1}{2}$.

Proof. Decomposition into compact and unbounded parts. By a structure theorem for hyperbolic surfaces [14], M can be written as the union of a compact part, hyperbolic ends (identical to hyperbolic cylinders) and cusps. It is easy to see that cusps prevent boundedness of the projectors, and thus we only consider the case when they are absent of the decomposition. We now consider a partition of unity

$$\chi_c(x) + \sum_{j=1}^N \chi_{hf}^j(x) = 1 \quad \text{for all } x \in M,$$

where χ_c is compactly supported, the supports of the χ_{hf}^j are disjoint, and M reduces to a hyperbolic funnel (halves of hyperbolic cylinders) on the support of each χ_{hf}^j .



FIGURE 5. The spherical example (red) and the Knapp example (blue) for the hyperbolic space represented in frequency and physical space. On the physical side, we use the Poincaré disk model to represent the hyperbolic space; and on the frequency side, it is the Fourier-Helgason transform which provides the dual picture. In frequency space, the supports are contained within the annulus with inner and outer radii $\lambda - \delta$ and $\lambda + \delta$, either the full annulus or a rectangle contained in it. In physical space, the L^p mass concentrates at a point, or along a geodesic; notice how the instability of geodesics on the hyperbolic space leads to spreading at the ends of the Knapp example.

At this point, it is convenient to replace $P_{\lambda,\delta}$ by a smoothed-out version of this projector (which is indifferent as far as $L^2 \to L^p$ bounds are concerned), and to express it through the Schrödinger group

(4.1)
$$P_{\lambda,\delta}^{\flat} = \chi_0 \left(\frac{\Delta + \lambda^2}{\lambda \delta}\right) = \frac{\lambda \delta}{\sqrt{2\pi}} \int \widehat{\chi_0}(\lambda \delta t) e^{i\lambda^2 t} e^{it\Delta} dt,$$

where χ_0 is a smooth cutoff function. Combining this formula with the partition of unity defined earlier, we will rely on the decomposition (simplifying slightly the formulas)

$$P_{\lambda,\delta}^{p} = P_{\lambda,1}\chi_{c}(x)P_{\lambda,\delta}^{p} + P_{\lambda,1}\chi_{hf}(x)P_{\lambda,\delta}^{p}.$$

Bounding the compact part. The crucial ingredient is the local L^2 smoothing estimate, which follows from Bourgain-Dyatlov [23]. As explained in Subsection 5.2 below, this results in the local L^2 TO L^p BOUNDS FOR SPECTRAL PROJECTORS ON THIN INTERVALS IN RIEMANNIAN MANIFOLDS $\ 13$

smoothing bound

$$\|\chi_c(x)e^{it\Delta}f\|_{L^2_tL^2_x} \lesssim \lambda^{-\frac{1}{2}+\epsilon}\|f\|_{L^2_x}$$

(for any $\epsilon > 0$). Combining this bound with formula (4.1), and the Cauchy-Schwarz inequality, we see that

$$\|\chi_c(x)P_{\lambda,\delta}^{\flat}\|_{L^2} \lesssim \lambda \delta \left\|\chi_c(x) \int \widehat{\chi_0}(\lambda \delta t) e^{i\lambda^2 t} P_{\lambda} e^{it\Delta} dt\right\|_{L^2} \lesssim (\lambda \delta)^{\frac{1}{2}} \|\chi_c(x) e^{it\Delta} f\|_{L^2_t L^2_x} \lesssim \lambda^{\epsilon} \delta^{\frac{1}{2}} \|f\|_{L^2_x}.$$

Using in addition the Sogge theorem, we can bound

$$\left\| P_{\lambda,1}\chi_c(x)P_{\lambda,\delta}^{\flat}f \right\|_{L^p} \lesssim \lambda^{\gamma(p)} \left\| \chi_c(x)P_{\lambda,\delta}^{\flat}f \right\|_{L^2} \lesssim \lambda^{\gamma(p)+\epsilon}\delta^{\frac{1}{2}}.$$

Bounding the hyperbolic part. To treat the contribution of the hyperbolic ends, the idea is to write

$$\chi_{hf}(x)\chi_0\left(\frac{\Delta+\lambda^2}{\lambda\delta}\right) = \frac{\lambda\delta}{\sqrt{2\pi}}\int\widehat{\chi_0}(\lambda\delta t)e^{i\lambda^2 t}P_\lambda\chi_{hf}(x)e^{it\Delta}\,dt.$$

Then, the function $u = \chi_{hf}^{j}(x)e^{it\Delta}$ is supported on the *j*-th funnel where it satisfies the Schrödinger equation

$$i\partial_t u - \Delta u = [\Delta, \chi^j_{hf}]e^{it\Delta}f.$$

The source term (on the right) is controlled by the local smoothing estimates; in order to control u, there remains to derive sufficiently sharp Strichartz, smoothing, and dispersive estimates on the funnel.

4.4. Finite area quotients. In the case where $\Gamma \setminus \mathbb{H}^2$ has finite area, Iwaniec and Sarnak [63], see also the review [81], conjectured the following: any eigenfunction ϕ_{λ_j} associated to the eigenvalue λ_j satisfies, for K compact and any $\epsilon > 0$

$$\|\varphi_{\lambda_j}\|_{L^{\infty}(K)} \lesssim \lambda_j^{\epsilon} \|\phi_{\lambda}\|_{L^2}$$

(which reduces to $\|\varphi_{\lambda}\|_{L^{\infty}(X)} \lesssim \lambda^{\epsilon}$ on a compact surface, for a normalized eigenfunction).

This conjecture should be regarded as very hard, since, for $p = \infty$ and in the particular case of the modular surface, it implies the Lindelöf hypothesis [80]. We refer to [27, 59, 60] for progress on this problem in the case of arithmetic manifolds.

Note that a markedly different behavior can occur in dimension d = 3, as evidenced by the result in [79].

5. Classical estimates implying spectral projector bounds

In this section, we examine how estimates for the spectrum of the Schrödinger operator (Weyl law), or the decay of the Schrödinger group (smoothing estimates, dispersive estimates, Strichartz estimates) imply bounds for the spectral projectors $P_{\lambda,\delta}$ in various regimes.

5.1. Weyl law. If M is compact, let (λ_j^2) denote the eigenvalues and (φ_j) the normalized eigenfunctions of the Laplace-Beltrami operator. The counting function enjoys the following asymptotics [62, 91]

$$N(\lambda) = \#\{\lambda_i^2 \text{ eigenvalue of } -\Delta \text{ with } \lambda_j < \lambda\} = C\lambda^d + O(\lambda^{d-1})$$

(for a constant C corresponding to the volume of the unit cotangent bundle). This estimate is sharp for spheres and Zoll manifolds, but the error term is expected to be smaller than $O(\lambda^{d-1})$ for "most" manifolds. Known improvements are as follows: $o(\lambda^{d-1})$ if the set of closed geodesics has measure zero [41], $O(\lambda^{d-1}(\log \lambda)^{-1})$ if M has negative curvature [8], and $O(\lambda^{d-1-\delta})$ (for some $\delta > 0$ depending on M) if M is integrable [36].

Local Weyl laws refer to (pointwise) estimates on the function

$$N_x(\lambda) = \sum_{\lambda_j < \lambda} |\varphi_j(x)|^2$$
, with $x \in M$ and $\lambda > 0$.

For any compact manifold [55, 94],

(5.1)
$$N_x(\lambda) = C\lambda^d + O(\lambda^{d-1}) \qquad \text{(uniformly in } x\text{)}.$$

To see the connection to operator norms of $P_{\lambda,\delta}$, observe that the kernel of $P_{\lambda,\delta}$ is given by $K_{\lambda,\delta}(x,y) = \sum_{\lambda-\delta < \lambda_j < \lambda+\delta} \varphi_j(x)\varphi_j(y)$. By the Cauchy-Schwarz inequality,

$$|K_{\lambda,\delta}(x,y)| \le \sqrt{N_x(\lambda+\delta) - N_x(\lambda-\delta)} \sqrt{N_y(\lambda+\delta) - N_y(\lambda-\delta)}$$

Using in addition the bound (5.1) gives the theorem of Sogge (2.3) for $p = \infty$, namely

$$\|P_{\lambda,1}\|_{L^1 \to L^\infty} \lesssim \lambda^{d-1}$$

Any improvement of the error term in (5.1) leads to an improved estimate for $||P_{\lambda,\delta}||_{L^1\to L^\infty}$.

5.2. Local smoothing estimates. Assume that the resolvent $R(\lambda) = (\Delta - \lambda^2 - i0)^{-1}$ enjoys the local smoothing bound

$$\|\chi R(\lambda)\chi\|_{L^2\to L^2} \lesssim \lambda^{-1+\epsilon}$$

here, $\epsilon \ge 0$ and χ is a localized function which can be compactly supported or decay at a given rate depending on the context. It follows from Kato's theorem that the Schrödinger group also exhibits local smoothing:

(5.2)
$$\left\|\chi P_{\lambda,1}e^{-it\Delta}f\right\|_{L^2_t L^2(M)} \lesssim \lambda^{-\frac{1}{2}+\epsilon} \|f\|_{L^2}.$$

Bounds of this type were proved for various infinite volume manifolds, in particular asymptotically Euclidean or hyperbolic, basic references include [34, 38, 39, 66].

It is then possible to follow ideas similar to those presented in Subsection 4.3.

5.3. Dispersive estimates. Assume that dispersive estimates hold on M with the same decay rate as on Euclidean space

(5.3)
$$\|e^{it\Delta}\|_{L^{p'}\to L^p} \lesssim |t|^{\frac{d}{p}-\frac{d}{2}}, \qquad p \ge 2.$$

(see the review [82] and also [83, 84]). Expressing again the smoothed-out operator $P_{\lambda,\delta}^{\flat}$ in terms of the Schrödinger group, it can be decomposed into

$$P_{\lambda,\delta}^{\flat} = \frac{\lambda\delta}{\sqrt{2\pi}} \int \widehat{\chi}(\lambda\delta t) e^{i\lambda^2 t} e^{it\Delta} dt \sim \frac{\lambda\delta}{\sqrt{2\pi}} \left[\int_{|t| \lesssim \frac{1}{\lambda}} + \int_{\frac{1}{\lambda} \lesssim |t| \lesssim \frac{1}{\lambda\delta}} \right] e^{i\lambda^2 t} e^{it\Delta} dt = P_{small} + P_{large}$$

(here, integration bounds such as $|t| \leq \frac{1}{\lambda}$ should be understood in terms of smooth cutoffs).

Adjusting the cutoff function, the operator P_{small} can be written as $\delta P_{\lambda,1}^{\flat}$, to which Sogge's theorem applies: $\|P_{small}\|_{L^{p'}\to L^p} \lesssim \lambda^{d-1-\frac{2d}{p}}\delta$ if $p \geq p_{ST}$. Turning to P_{large} , it can be bounded through the dispersive estimates (5.3) to obtain, if $p \geq p^* = \frac{2d}{d-2}$,

$$\left\|\int_{\frac{1}{\lambda}\lesssim |t|\lesssim \frac{1}{\lambda\delta}}\lambda\delta\widehat{\chi}(\lambda\delta t)e^{i\lambda^{2}t}e^{it\Delta}\,dt\right\|_{L^{p'}\to L^{p}}\lesssim \lambda\delta\int_{\frac{1}{\lambda}\lesssim |t|\lesssim \frac{1}{\lambda\delta}}|t|^{\frac{d}{p}-\frac{d}{2}}\,dt\lesssim \lambda^{\frac{d}{2}-\frac{d}{p}}\delta\lesssim \lambda^{d-1-\frac{2d}{p}}\delta.$$

Overall, we found that, if the dispersive estimates (5.3) hold, the $L^2 \to L^p$ norm of $P_{\lambda,\delta}$ enjoys the optimal bound $\lambda^{\frac{d-1}{2}-\frac{d}{p}}\delta^{\frac{1}{2}}$ as soon as $p \ge p^* = \frac{2d}{d-2}$.

5.4. Strichartz estimates. Strichartz estimates on \mathbb{R}^d are given by

$$||e^{it\Delta}f||_{L^p_t L^q_x} \lesssim ||f||_{L^2}, \text{ if } p, q \ge 2, \quad \frac{2}{p} + \frac{d}{q} = \frac{d}{2}.$$

This estimate was first proved [47, 48, 104] for p > 2, and the endpoint result for $p = p^* = \frac{2d}{d-2}$

$$\|e^{it\Delta}f\|_{L^2_t L^{p^*}_r} \lesssim \|f\|_{L^2}$$

(which will be of particular relevance for us) was then obtained by Keel and Tao [67].

Strichartz estimates were subsequently extended to perturbations of the Euclidean space: either short range perturbations yielding a global in time estimate [97, 78], or long range perturbations yielding local in time estimates [25, 15]

Assuming that this endpoint estimate holds on M, we can apply it together with the Cauchy-Schwarz inequality to the mollified spectral projector expressed through the Schrödinger group (4.1) and obtain

$$\|P_{\lambda,\delta}^{\flat}f\|_{L^{p^*}} \lesssim \lambda\delta(\lambda\delta)^{-\frac{1}{2}} \|e^{it\Delta}f\|_{L^2([-\frac{1}{\lambda\delta},\frac{1}{\lambda\delta}],L^{p^*}(M))} \lesssim \lambda^{\frac{1}{2}}\delta^{\frac{1}{2}} \|f\|_{L^2};$$

(neglecting tails of $\widehat{\chi_0}$). This shows that the endpoint Strichartz estimate up to time $\sim \frac{1}{\lambda\delta}$ implies the optimal $L^2 \to L^{p^*}$ estimate for $P_{\lambda,\delta}$.

6. $L^2 \rightarrow L^p$ projector bounds beyond complete smooth Riemannian manifolds

While the present review focuses on bounds for spectral projectors on smooth complete Riemannian manifolds, many interesting variants can be considered. In the list below, we try to summarize all generalizations which have been investigated, with the relevant references.

- Non-smooth metric [69, 70]
- Manifold with boundary [89, 86]
- Sub-Riemannian manifolds [75, 6]
- Pseudo-differential operators [72]
- Schrödinger operators [65, 71, 64, 10, 58]. In the asymptotically Euclidean case, a related question is the boundedness of wave operators [82], which can be combined with the Stein-Tomas theorem to prove a natural analog.
- Restriction of eigenfunctions to lower-dimensional submanifolds [26, 24, 32, 101]
- Finite graphs (size N), for which one should substitute the infinite size limit $N \to \infty$ to the high frequency limit $\lambda \to \infty$ which is of interest for manifolds. Nearly optimal pointwise bounds for eigenvectors could be proved for random graphs [7].

7. Some interesting questions

We conclude with some interesting questions

7.1. Geometry of maximizers. In all known examples, the functions at which the operator norm of $P_{\lambda,\delta}$ is nearly achieved concentrate asymptotically (as $\lambda \to \infty$) either at a point, or along a geodesic. Could other possibilities arise?

7.2. Infinite volume manifolds. In all known examples, $||P_{\lambda,\delta}||_{L^2 \to L^p} \lesssim \lambda^{\frac{d-1}{2} - \frac{d}{p}} \delta^{\frac{1}{2}} + \lambda^{\frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{p}\right)}$, where the first term comes from point focusing, and the second from geodesic focusing on a stable closed geodesic. Is this bound indeed universal? At least for all manifolds with a simple behavior at infinity (e.g. flat, conic, hyperbolic)?

For $p < p_{ST}$, very different behaviors can occur for $||P_{\lambda,\delta}||_{L^2 \to L^p}$. One finds that this is $\sim \lambda^{\frac{d-1}{2}\left(\frac{1}{2}-\frac{1}{p}\right)}\delta^{\alpha}$, with $\alpha = 0$ if there exists a stable closed geodesic, $\alpha = \frac{d+1}{2}\left(\frac{1}{2}-\frac{1}{p}\right)$ on the Euclidean space, $\alpha = \frac{d-1}{2}\left(\frac{1}{2}-\frac{1}{p}\right)$ on Euclidean cylinders, and $\alpha = \frac{1}{2}$ on the hyperbolic space (these values of α can be interpreted in relation to the stability of geodesics). Are other values of α possible?

7.3. Equidistribution of eigenfunctions and of the spectrum. In the compact case, is it possible to establish a relation between equidistribution of eigenfunctions on the manifold, and equidistribution of the spectrum over the reals? For instance, denoting η for a quantity related to the scale over which eigenvalues are equidistributed, could there hold an estimate such as $\|\varphi\|_{L^{\infty}} \lesssim \lambda^{\frac{d-1}{2} - \frac{d}{p}} \eta^{\frac{1}{2}}$ for any eigenfunction φ with eigenvalue λ , and for p sufficiently large? On the one hand, such an estimate seems to fail for surfaces of revolution, whose spectrum can be equidistributed, but which admit exceptional eigenfunctions concentrating at the poles. On the other hand, such an estimate is verified on spheres and rational or irrational tori. An analogy can also be made with random matrices, for which a connection between quantum unique ergodicy and GUE statistics for the spectrum can be established in some cases [16].

7.4. Ignoring stable closed geodesics. Stable closed geodesics are the obstacle to an improvement over the sphere case for p close to 2. Would an improvement become possible by localizing away from stable geodesics?

7.5. Link with the Fourier restriction problem. The question of estimating $||P_{\lambda,\delta}||_{L^2 \to L^p}$ provided a generalization of the Fourier restriction problem to arbitrary Riemannian manifolds, when the density in Fourier space is L^2 . Is it possible (and interesting) to obtain such a generalization for the analog of L^p density in Fourier space?

The formulation of this question depends on a choice of a distinguished basis of (generalized) eigenfunctions. If the manifold is of infinite volume and for instance asymptotically Euclidean, such a distinguished basis could be provided by scattering theory. If the manifold is compact, then the eigenspaces of the Laplacian are generically non-degenerate, and this provides naturally an orthonormal basis of L^2 , which will be denoted (φ_k) . It is tempting to ask about the $\ell^q(\mathbb{N}) \to L^p(M)$ boundedness properties of $(c_k)_{k \in \mathbb{N}} \mapsto \sum_{|\lambda_k - \lambda| < \delta} c_k \varphi_k$ (denoting φ_k for the k-th eigenfunction). In the case of the torus, it was noticed by Bourgain and Demeter [20] that the case of general q is an immediate consequence of q = 2. Does this remain true for other manifolds?

7.6. Nonsymmetric and nonintegrable compact manifolds. We conclude with this most ambitious question: if the manifold under consideration is neither symmetric, nor integrable, there does not exist, to the best of our knowledge, any result giving more than logarithmic improvements over the universal Sogge estimate. Could that be obtained in a single example?

Acknowledgements. The author is grateful to Tristan Léger for his suggestions to improve a preliminary version, and to Paul Bourgade, Isabelle Gallagher, Chris Sogge, Stefan Steinerberger, Daniel Tataru, Laurent Thomann and Maciej Zworski for enlightening conversations during the writing of this review article.

When working on this article, the author was supported by a start-up grant from Imperial College and a Wolfson fellowship L^2 TO L^p BOUNDS FOR SPECTRAL PROJECTORS ON THIN INTERVALS IN RIEMANNIAN MANIFOLDS $\ 17$

References

- P. Albin, H. Christianson, J. L. Marzuola, and L. Thomann. Construction of nonlinear quasimodes near elliptic periodic orbits. *Physica D: Nonlinear Phenomena*, 241(4):409–425, 2012.
- [2] N. Anantharaman. Delocalization of Schrödinger eigenfunctions. In Proceedings of the International Congress of Mathematicians: Rio de Janeiro 2018, pages 341–375. World Scientific, 2018.
- [3] J.-P. Anker, P. Germain, and T. Léger. Spectral projectors on hyperbolic surfaces. arXiv preprint arXiv:2306.12827, 2023.
- [4] R. Aurich, A. Bäcker, R. Schubert, and M. Taglieber. Maximum norms of chaotic quantum eigenstates and random waves. *Physica D: Nonlinear Phenomena*, 129(1-2):1–14, 1999.
- [5] V. G. Avakumović. über die eigenfunktionen auf geschlossenen riemannschen mannigfaltigkeiten. Mathematische ZeitSchrift, 65(1):327–344, 1956.
- [6] H. Bahouri, D. Barilari, and I. Gallagher. Strichartz estimates and fourier restriction theorems on the Heisenberg group. Journal of Fourier Analysis and Applications, 27(2):1–41, 2021.
- [7] R. Bauerschmidt, J. Huang, and H.-T. Yau. Local kesten-mckay law for random regular graphs. Communications in Mathematical Physics, 369:523-636, 2019.
- [8] P. H. Bérard. On the wave equation on a compact Riemannian manifold without conjugate points. Mathematische ZeitSchrift, 155(3):249–276, 1977.
- [9] M. V. Berry. Regular and irregular semiclassical wavefunctions. Journal of Physics A: Mathematical and General, 10(12):2083, 1977.
- [10] M. D. Blair, X. Huang, Y. Sire, and C. D. Sogge. Uniform Sobolev estimates on compact manifolds involving singular potentials. *Revista Matemática Iberoamericana*, 2021.
- [11] M. D. Blair and C. D. Sogge. Refined and microlocal kakeya-nikodym bounds of eigenfunctions in higher dimensions. *Communications in Mathematical Physics*, 356(2):501–533, 2017.
- [12] M. D. Blair and C. D. Sogge. Concerning toponogov's theorem and logarithmic improvement of estimates of eigenfunctions. Journal of Differential Geometry, 109(2):189–221, 2018.
- [13] M. D. Blair and C. D. Sogge. Logarithmic improvements in l^p bounds for eigenfunctions at the critical exponent in the presence of nonpositive curvature. *Inventiones mathematicae*, 217(2):703–748, 2019.
- [14] D. Borthwick. Spectral theory of infinite-area hyperbolic surfaces. Springer, 2007.
- [15] J.-M. Bouclet and N. Tzvetkov. Strichartz estimates for long range perturbations. American journal of mathematics, 129(6):1565–1609, 2007.
- [16] P. Bourgade. Random band matrices. In Proceedings of the International Congress of Mathematicians: Rio de Janeiro 2018, pages 2759–2783. World Scientific, 2018.
- [17] J. Bourgain. Eigenfunction bounds for compact manifolds with integrable geodesic flow. Institut des Hautes Etudes Scientifique, 1993.
- [18] J. Bourgain. Eigenfunction bounds for the laplacian on the n-torus. International Mathematics Research Notices, 1993(3):61–66, 1993.
- [19] J. Bourgain. Moment inequalities for trigonometric polynomials with spectrum in curved hypersurfaces. Israel Journal of Mathematics, 193(1):441–458, 2013.
- [20] J. Bourgain and C. Demeter. Improved estimates for the discrete Fourier restriction to the higher dimensional sphere. *Illinois Journal of Mathematics*, 57(1):213–227, 2013.
- [21] J. Bourgain and C. Demeter. New bounds for the discrete Fourier restriction to the sphere in 4D and 5D. Int. Math. Res. Not. IMRN, (11):3150–3184, 2015.
- [22] J. Bourgain and C. Demeter. The proof of the l 2 decoupling conjecture. Annals of mathematics, pages 351–389, 2015.
- [23] J. Bourgain and S. Dyatlov. Spectral gaps without the pressure condition. Annals of Mathematics, 187(3):825– 867, 2018.
- [24] J. Bourgain and Z. Rudnick. Restriction of toral eigenfunctions to hypersurfaces and nodal sets. Geometric and Functional Analysis, 22(4):878–937, 2012.
- [25] N. Burq. Estimations de strichartz pour des perturbations à longue portée de l'operateur de Schrodinger. Séminaire Équations aux dérivées partielles (Polytechnique) dit aussi" Séminaire Goulaouic-Schwartz", pages 1-8, 2001.
- [26] N. Burq, P. Gérard, and N. Tzvetkov. Restrictions of the laplace-beltrami eigenfunctions to submanifolds. Duke Mathematical Journal, 138(3):445–486, 2007.
- [27] J. Buttcane and R. Khan. On the fourth moment of Hecke–Maass forms and the random wave conjecture. Compositio Mathematica, 153(7):1479–1511, 2017.
- [28] Y. Canzani and J. Galkowski. Improvements for eigenfunction averages: an application of geodesic beams. arXiv preprint arXiv:1809.06296, 2018.

- [29] Y. Canzani and J. Galkowski. On the growth of eigenfunction averages: microlocalization and geometry. Duke Mathematical Journal, 168(16):2991–3055, 2019.
- [30] Y. Canzani and J. Galkowski. Growth of high l^p norms for eigenfunctions: an application of geodesic beams. arXiv preprint arXiv:2003.04597, 2020.
- [31] Y. Canzani and J. Galkowski. Eigenfunction concentration via geodesic beams. Journal f
 ür die reine und angewandte Mathematik (Crelles Journal), 2021(775):197–257, 2021.
- [32] L. Carleson and P. Sjölin. Oscillatory integrals and multiplier problem for the disc. Studia Mathematica, 3(44):287–299, 1972.
- [33] X. Chen and A. Hassell. Resolvent and spectral measure on non-trapping asymptotically hyperbolic manifolds ii: spectral measure, restriction theorem, spectral multipliers. In Annales de l'Institut Fourier, volume 68, pages 1011–1075, 2018.
- [34] P. Constantin and J.-C. Saut. Local smoothing properties of schrödinger equations. Indiana University mathematics journal, pages 791–810, 1989.
- [35] R. Cooke. A cantor-lebesgue theorem in two dimensions. Proceedings of the American Mathematical Society, 30(3):547–550, 1971.
- [36] Y. C. de Verdiere. Quasi-modes sur les variétés riemanniennes. Inventiones mathematicae, 43(1):15–52, 1977.
- [37] C. Demeter and P. Germain. L2 to lp bounds for spectral projectors on the euclidean two-dimensional torus. arXiv preprint arXiv:2306.14286, 2023.
- [38] S.-i. Doi. Remarks on the cauchy problem for schrödinger-type equations. Communications in Partial Differential Equations, 21(1-2):163–178, 1996.
- [39] S.-i. Doi. Smoothing effects of schrödinger evolution groups on riemannian manifolds. Duke Mathematical Journal, 82(3):679–706, 1996.
- [40] H. Donnelly. Bounds for eigenfunctions of the laplacian on compact Riemannian manifolds. Journal of Functional Analysis, 187(1):247–261, 2001.
- [41] J. J. Duistermaat. On the spectrum of positive elliptic operators and periodic bicharacteristics. In Fourier Integral Operators and Partial Differential Equations, pages 15–22. Springer, 1975.
- [42] D. Foschi and D. Oliveira e Silva. Some recent progress on sharp Fourier restriction theory. Analysis Mathematica, 43(2):241–265, 2017.
- [43] P. Germain and T. Léger. Spectral projectors, resolvent, and Fourier restriction on the hyperbolic space. arXiv preprint arXiv:2104.04126, 2021.
- [44] P. Germain and S. Myerson. Bounds for spectral projectors on the Euclidean cylinder. Comptes Rendus. Mathématique, 360(G11):1257–1262, 2022.
- [45] P. Germain and S. L. R. Myerson. Bounds for spectral projectors on tori. In *Forum of Mathematics, Sigma*, volume 10. Cambridge University Press, 2022.
- [46] P. Germain and S. L. Rydin Myerson. Bounds for spectral projectors on generic tori. Mathematische Annalen, pages 1–37, 2022.
- [47] J. Ginibre and G. Velo. The global cauchy problem for the non linear Schrödinger equation revisited. In Annales de l'Institut Henri Poincaré C, Analyse non linéaire, volume 2, pages 309–327. Elsevier, 1985.
- [48] J. Ginibre and G. Velo. Smoothing properties and retarded estimates for some dispersive evolution equations. Communications in mathematical physics, 144(1):163–188, 1992.
- [49] F. Götze. Lattice point problems and values of quadratic forms. Invent. Math., 157(1):195–226, 2004.
- [50] C. Guillarmou, A. Hassell, and A. Sikora. Restriction and spectral multiplier theorems on asymptotically conic manifolds. Analysis & PDE, 6(4):893–950, 2013.
- [51] V. Guillemin and A. Weinstein. Eigenvalues associated with a closed geodesic. Bulletin of the American Mathematical Society, 82(1):92–94, 1976.
- [52] A. Hassell and M. Tacy. Improvement of eigenfunction estimates on manifolds of nonpositive curvature. In Forum Mathematicum, volume 27, pages 1435–1451. De Gruyter, 2015.
- [53] H. Hezari and G. Rivière. Lp norms, nodal sets, and quantum ergodicity. *Advances in Mathematics*, 290:938–966, 2016.
- [54] J. Hickman. Uniform l^p resolvent estimates on the torus. Mathematics Research Reports, 1:31–45, 2020.
- [55] L. Hörmander. The spectral function of an elliptic operator. Acta mathematica, 121:193–218, 1968.
- [56] L. Hörmander. Oscillatory integrals and multipliers onfl p. Arkiv för Matematik, 11(1):1–11, 1973.
- [57] S. Huang and C. D. Sogge. Concerning lp resolvent estimates for simply connected manifolds of constant curvature. *Journal of Functional Analysis*, 267(12):4635–4666, 2014.
- [58] X. Huang and C. D. Sogge. Uniform Sobolev estimates in \mathbb{R}^n involving singular potentials. The Journal of Geometric Analysis, 32(2):1–36, 2022.
- [59] P. Humphries. Equidistribution in shrinking sets and l^4 -norm bounds for automorphic forms. *Mathematische* Annalen, 371(3):1497–1543, 2018.

 L^2 TO L^p BOUNDS FOR SPECTRAL PROJECTORS ON THIN INTERVALS IN RIEMANNIAN MANIFOLDS 19

- [60] P. Humphries and R. Khan. l^p-norm bounds for automorphic forms via spectral reciprocity. preprint arXiv:2208.05613, 2022.
- [61] M. N. Huxley. Exponential sums and lattice points. III. Proc. London Math. Soc. (3), 87(3):591-609, 2003.
- [62] V. Ivrii. 100 years of weyl's law. Bulletin of Mathematical Sciences, 6(3):379–452, 2016.
- [63] H. Iwaniec and P. Sarnak. L^{∞} norms of eigenfunctions of arithmetic surfaces. Annals of Mathematics, 141(2):301–320, 1995.
- [64] E. Jeong, S. Lee, and J. Ryu. Endpoint eigenfunction bounds for the Hermite operator. arXiv preprint arXiv:2205.03036, 2022.
- [65] G. Karadzhov. Riesz summability of multiple Hermite series. Comptes Rendus-Académie des Sciences Paris Série 1, 317:1023–1023, 1993.
- [66] T. Kato and K. Yajima. Some examples of smooth operators and the associated smoothing effect. Reviews in Mathematical Physics, 1(04):481–496, 1989.
- [67] M. Keel and T. Tao. Endpoint strichartz estimates. American Journal of Mathematics, 120(5):955–980, 1998.
- [68] C. E. Kenig, A. Ruiz, and C. D. Sogge. Uniform Sobolev inequalities and unique continuation for second order constant coefficient differential operators. *Duke Mathematical Journal*, 55(2):329–347, 1987.
- [69] H. Koch, H. F. Smith, and D. Tataru. Sharp l^q bounds on spectral clusters for holder metrics. Mathematical Research Letters, 14(1):77–85, 2007.
- [70] H. Koch, H. F. Smith, and D. Tataru. Sharp l p bounds on spectral clusters for lipschitz metrics. American Journal of Mathematics, 136(6):1629–1663, 2014.
- [71] H. Koch and D. Tataru. L p eigenfunction bounds for the Hermite operator. Duke Mathematical Journal, 128(2):369–392, 2005.
- [72] H. Koch, D. Tataru, and M. Zworski. Semiclassical l p estimates. In Annales Henri Poincaré, volume 8, pages 885–916. Springer, 2007.
- [73] E. Krätzel. Analytische Funktionen in der Zahlentheorie, volume 139 of Teubner-Texte zur Mathematik [Teubner Texts in Mathematics]. B. G. Teubner, Stuttgart, 2000.
- [74] B. M. Levitan. On the asymptotic behavior of the spectral function of a self-adjoint differential equation of the second order. *Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya*, 16(4):325–352, 1952.
- [75] D. Muller. A restriction theorem for the Heisenberg group. Annals of mathematics, 131(3):567–587, 1990.
- [76] W. G. Nowak. Integer Points in Large Bodies, pages 583–599. Springer International Publishing, Cham, 2014.
- [77] J. Ralston. Approximate eigenfunctions of the laplacian. Journal of Differential Geometry, 12(1):87–100, 1977.
- [78] L. Robbiano and C. Zuily. Strichartz estimates for Schrödinger equations with variable coefficients. Mémoires de la SMF, 101:102, 2005.
- [79] Z. Rudnick and P. Sarnak. The behaviour of eigenstates of arithmetic hyperbolic manifolds. Communications in Mathematical Physics, 161:195–213, 1994.
- [80] P. Sarnak. Arithmetic quantum chaos. Blyth Lectures. Toronto, 1993.
- [81] P. Sarnak. Recent progress on the quantum unique ergodicity conjecture. Bull. Amer. Math. Soc, 48:211–228, 2012.
- [82] W. Schlag. Intertwining wave operators, Fourier restriction, and Wiener theorems. arXiv preprint arXiv:1802.01982, 2018.
- [83] W. Schlag, A. Soffer, and W. Staubach. Decay for the wave and schrödinger evolutions on manifolds with conical ends, part i. *Transactions of the American Mathematical Society*, 362(1):19–52, 2010.
- [84] W. Schlag, A. Soffer, and W. Staubach. Decay for the wave and schrödinger evolutions on manifolds with conical ends, part ii. *Transactions of the American Mathematical Society*, 362(1):289–318, 2010.
- [85] S. Shao. On existence of extremizers for the tomas-stein inequality for s1. Journal of Functional Analysis, 270(10):3996-4038, 2016.
- [86] H. F. Smith and C. D. Sogge. On the lp norm of spectral clusters for compact manifolds with boundary. Acta mathematica, 198(1):107–153, 2007.
- [87] C. D. Sogge. Oscillatory integrals and spherical harmonics. Duke Mathematical Journal, 53(1):43–65, 1986.
- [88] C. D. Sogge. Concerning the lp norm of spectral clusters for second-order elliptic operators on compact manifolds. *Journal of functional analysis*, 77(1):123–138, 1988.
- [89] C. D. Sogge. Eigenfunction and Bochner-Riesz estimates on manifolds with boundary. Mathematical Research Letters, 9, 2002.
- [90] C. D. Sogge. Riemannian manifolds with maximal eigenfunction growth. Séminaire Equations aux dérivées partielles (Polytechnique) dit aussi" Séminaire Goulaouic-Schwartz", pages 1–16, 2002.
- [91] C. D. Sogge. Hangzhou lectures on eigenfunctions of the laplacian (am-188). In Hangzhou Lectures on Eigenfunctions of the Laplacian (AM-188). Princeton University Press, 2014.
- [92] C. D. Sogge. Fourier integrals in classical analysis, volume 210. Cambridge University Press, 2017.
- [93] C. D. Sogge, J. A. Toth, and S. Zelditch. About the blowup of quasimodes on Riemannian manifolds. Journal of Geometric Analysis, 21(1):150–173, 2011.

- [94] C. D. Sogge and S. Zelditch. Riemannian manifolds with maximal eigenfunction growth. Duke Mathematical Journal, 114(3):387–432, 2002.
- [95] C. D. Sogge and S. Zelditch. Focal points and sup-norms of eigenfunctions. Revista Matemática Iberoamericana, 32(3):971–994, 2016.
- [96] C. D. Sogge and S. Zelditch. Focal points and sup-norms of eigenfunctions ii: the two-dimensional case. Revista matemática iberoamericana, 32(3):995–999, 2016.
- [97] G. Staffilani and D. Tataru. Strichartz estimates for a Schrödinger operator with nonsmooth coefficients. Communications in Partial Differential Equations, 27(7-8):1337–1372, 2002.
- [98] E. M. Stein. Oscillatory integrals in Fourier analysis. Beijing lectures in harmonic analysis, 112:307–355, 1986.
- [99] E. M. Stein and T. S. Murphy. Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals, volume 3. Princeton University Press, 1993.
- [100] S. Steinerberger. Quantum entanglement and the growth of laplacian eigenfunctions. arXiv preprint arXiv:2111.12625, 2021.
- [101] M. Tacy. Semiclassical l p estimates of quasimodes on submanifolds. Communications in Partial Differential Equations, 35(8):1538–1562, 2010.
- [102] P. Tomas et al. A restriction theorem for the Fourier transform. Bull. Amer. Math. Soc, 81(s1), 1975.
- [103] J. A. Toth and S. Zelditch. L p norms of eigenfunctions in the completely integrable case. In Annales Henri Poincaré, volume 4, pages 343–368. Springer, 2003.
- [104] K. Yajima. Existence of solutions for Schrödinger evolution equations. Communications in Mathematical Physics, 110(3):415–426, 1987.
- [105] S. Zelditch. Local and global analysis of eigenfunctions: Advanced lectures in mathematics, vol. 7. In Handbook of Geometric Analysis, No. 1: Advanced Lectures in Mathematics, Vol. 7, pages 545–658. International Press, 2008.
- [106] S. Zelditch. Eigenfunctions of the Laplacian on a Riemannian manifold, volume 125. American Mathematical Soc., 2017.
- [107] A. Zygmund. On Fourier coefficients and transforms of functions of two variables. Studia Mathematica, 50(2):189–201, 1974.

PIERRE GERMAIN, DEPARTMENT OF MATHEMATICS, HUXLEY BUILDING, SOUTH KENSINGTON CAMPUS, IMPERIAL COLLEGE LONDON, LONDON SW7 2AZ, UNITED KINGDOM

Email address: pgermain@ic.ac.uk