Edge-coloring a graph G so that every copy of a graph H has an odd color class

Patrick Bennett * H

Emily Heath [†]

Shira Zerbib[‡]

August 22, 2023

Abstract

Recently, Alon [1] introduced the notion of an *H*-code for a graph *H*: a collection of graphs on vertex set [*n*] is an *H*-code if it contains no two members whose symmetric difference is isomorphic to *H*. Let $D_H(n)$ denote the maximum possible cardinality of an *H*-code, and let $d_H(n) = D_H(n)/2^{\binom{n}{2}}$. Alon observed that a lower bound on $d_H(n)$ can be obtained by attaining an upper bound on the number of colors needed to edge-color K_n so that every copy of *H* has an odd color class.

Motivated by this observation, we define g(G, H) to be the minimum number of colors needed to edge-color a graph G so that every copy of H has an odd color class. We prove $g(K_n, K_5) \leq n^{o(1)}$ and $g(K_{n,n}, C_4) = n/2 + o(n)$. The first result shows $d_{K_5}(n) \geq \frac{1}{n^{o(1)}}$ and was obtained independently in [12].

1 Introduction

Given a graph G and a subgraph H, let g(G, H) be the minimum number of colors needed to edge-color G so that every copy of H sees some color an odd number of times. The problem of determining g(G, H) is a natural question motivated by recent work of Alon [1] introducing the related notion of graph-codes.

Let V = [n] and let \mathcal{H} be a family of graphs on the set of vertices [n] which is closed under isomorphism. A collection of graphs \mathcal{F} on [n] is called an \mathcal{H} -code if it contains no two members whose symmetric difference is a graph in \mathcal{H} . For the special case that \mathcal{H} contains all copies of a single graph H on [n] this is called an \mathcal{H} -code. Let $D_{\mathcal{H}}(n)$ denote the maximum possible cardinality of an \mathcal{H} -code, and let

$$d_{\mathcal{H}}(n) = \frac{D_{\mathcal{H}}(n)}{2^{\binom{n}{2}}}$$

be the maximum possible fraction of the total number of graphs on [n] in an \mathcal{H} -code. If \mathcal{H} consists of all graphs isomorphic to one graph H, denote $d_{\mathcal{H}}(n)$ by $d_H(n)$.

In the case where \mathcal{H} consists of all graphs with independence number at most 2, a result of Ellis, Filmus and Friedgut [10] shows that $d_{\mathcal{H}}(n) = 1/8$ for $n \geq 3$. Berger and Zhao [5] proved the analogous result for the family \mathcal{H} of graphs with independence number at most 3, showing that $d_{\mathcal{H}}(n) = 1/64$ for all $n \geq 4$. In [2], Alon, Gujgiczer, Körner, Milojević, and Simonyi studied $D_{\mathcal{H}}(n)$ and $d_{\mathcal{H}}(n)$ for several families \mathcal{H} , such as disconnected graphs, graphs that are not 2-connected, non-Hamiltonian graphs, graphs that contain or do not contain a spanning star, graphs that contain an induced or non-induced copy of a fixed graph T, and graphs that do not contain such a subgraph. Alon [1] also studied the cases for cliques, stars, and matchings.

In [1], Alon mentions that the case $\mathcal{H} = \mathcal{K}$, where \mathcal{K} is the family of all cliques, is of particular interest. This case is motivated by a conjecture of Gowers (see [14, 1]). Further, he comments that $d_{K_4}(n) \geq \frac{1}{n^{o(1)}}$.

^{*}Department of Mathematics, Western Michigan University, Kalamazoo, MI. patrick.bennett@wmich.edu.

[†]Department of Mathematics, Iowa State University, Ames, IA. eheath@iastate.edu. Research partially supported by NSF RTG Grant DMS-1839918.

[‡]Department of of Mathematics, Iowa State University, Ames, IA. zerbib@iastate.edu. Research partially supported by NSF Grant DMS-1953929.

This result follows from the existence of an edge-coloring of K_n by $n^{o(1)}$ colors with no copy of K_4 in which every color appears an even number of times; that is, $g(K_n, K_4) \leq n^{o(1)}$. A coloring with this property was given in [7], modifying constructions in [8, 16]. In our first result, we show that the same coloring has the property that there exists no copy of K_5 in which every color appears an even number of times. We prove:

Theorem 1.1. We have $g(K_n, K_5) \le n^{o(1)}$.

This, together with a similar argument to that in [1], implies the following.

Theorem 1.2. We have $d_{K_5}(n) \ge \frac{1}{n^{o(1)}}$.

Very recently, Ge, Xu, and Zhang [12] also independently obtained this result using a similar method. They further note that $g(K_n, K_5) \ge \Omega(\log n)$.

As is observed in [1], if every member of \mathcal{H} has an odd number of edges then $d_{\mathcal{H}}(n) \geq \frac{1}{2}$, as the family of all graphs on [n] with an even number of edges forms an \mathcal{H} -code. Thus, when considering cliques, the next interesting case is the case of $H = K_8$. We do not know if the same coloring, or some generalization of it, has the property that there exists no copy of K_8 in which every color appears an even number of times. As our method relies on case analysis, it will be hard to generalize it to the case of K_8 .

An upper bound on $g(K_n, K_p)$ for larger p follows from recent work of Bennett, Delcourt, Li, and Postle [3] on the generalized Ramsey number f(n, p, q), that is, the minimum number of colors needed to color $E(K_n)$ so that every copy of K_p sees at least q colors. Until recently, the best upper bound on f(n, p, q) for general p, q was the original bound of Erdős and Gyárfás [11] obtained using the Lovász Local Lemma. Bennett, Dudek, and English [4] used a random greedy process to improve this bound by a logarithmic factor for values of q and p with $q \leq (p^2 - 26p + 55)/4$. In [3], their result is extended to all values of p and q except at the values $q = {p \choose 2} - p + 2$ and $q = {p \choose 2} - \lfloor \frac{p}{2} \rfloor + 2$, where the local lemma bound is known to be tight.

Theorem 1.3 (Bennett, Delcourt, Li, Postle [3]). For fixed positive integers p, q with p-2 not divisible by $\binom{p}{2} - q + 1$, we have

$$f(n, p, q) = O\left(\left(\frac{n^{p-2}}{\log n}\right)^{\frac{1}{\binom{p}{2}-q+1}}\right)$$

In fact, Theorem 1.3 is generalized in [3] to give an analogous upper bound in a list-coloring setting for any subgraph (not only cliques) and in hypergraphs with higher uniformity. Note that Theorem 1.3 with $q = \binom{p}{2}/2 + 1$ implies that

$$g(K_n, K_p) = O\left(\left(\frac{n^{p-2}}{\log n}\right)^{2/\binom{p}{2}}\right).$$

Indeed, any $(p, \frac{1}{2} {p \choose 2} + 1)$ -coloring of K_n has the stronger condition that some color must appear exactly once in each *p*-clique.

Our second result focuses on the question of determining $g(K_{n,n}, C_4)$. In order to prove an upper bound, we will construct a coloring using the "forbidden submatching method" recently introduced by Delcourt and Postle [9] and independently by Glock, Joos, Kim, Kühn, and Lichev [13] as "conflict-free hypergraph matchings." In [15], Joos and Mubayi use the variant of the method from [13] along with a probabilistic argument to show that $f(K_{n,n}, C_4, 3) = \frac{2}{3}n + o(n)$. Note that this implies $g(K_{n,n}, C_4) \leq \frac{2}{3}n + o(n)$. We use a similar approach to show the stronger upper bound of $g(K_{n,n}, C_4) \leq \frac{1}{2}n + o(n)$. As such, combined with a short proof that $g(K_{n,n}, C_4) > n/2$, we obtain the following result.

Theorem 1.4. We have $g(K_{n,n}, C_4) = \frac{1}{2}n + o(n)$.

The rest of the paper is organized as follows. In Section 2, we prove Theorem 1.2. In Section 3, we give the necessary preliminaries to state the main tool for the proof of Theorem 1.4, namely the theorem on forbidden submatchings given in [13]. Finally, in Section 4, we prove Theorem 1.4.

2 Proof of Theorem 1.1

A (p,q)-coloring of K_n is an edge-coloring in which each *p*-clique receives at least *q* colors. In 2015, Conlon, Fox, Lee, and Sudakov [8] constructed a (p, p-1)-coloring with $n^{o(1)}$ colors. In 2017, Cameron and Heath [6] used a modified version of their (5, 4)-coloring as part of their construction of a (5, 5)-coloring with $n^{1/3+o(1)}$ colors. We will prove that this "Modified CFLS" (5, 4)-coloring using $n^{o(1)}$ colors guarantees that every K_5 contains some color an odd number of times.

Let $n = 2^{m^2}$ for some positive integers m. We will view the vertices of K_n as binary strings of length m^2 ; that is, $V = \{0, 1\}^{m^2}$. For each $v \in V$, let $v^{(i)}$ denote the *i*th block of bits of length m in v, so

$$v = \left(v^{(1)}, v^{(2)}, \dots, v^{(m)}\right),$$

where each $v^{(i)} \in \{0, 1\}^m$.

Note that we can assign a linear order to the vertices by considering each to be an integer represented in binary and taking the standard ordering of these integers. That is, x < y if and only if the first bit of difference between x and y is zero in x and one in y. Furthermore, each m-block of our vertices can be viewed as a binary representation of an integer from 0 to $2^m - 1$, so the m-blocks can be ordered in the same way.

Let $x, y \in V$ such that x < y. Let *i* be the first index for which $x^{(i)} \neq y^{(i)}$, and for each $k \in [m]$, let i_k be the first index at which a bit of $x^{(k)}$ differs from the corresponding bit of $y^{(k)}$, or $i_k = 0$ if $x^{(k)} = y^{(k)}$. In addition, for each $k \in [m]$, let

$$\delta_k = \begin{cases} +1 & x^{(k)} \le y^{(k)}, \\ -1 & x^{(k)} > y^{(k)}. \end{cases}$$

The Modified CFLS coloring assigns to the edge xy the color

$$\varphi(xy) = \left(\left(i, \{x, y\} \right), i_1, i_2, \dots, i_m, \delta_1, \delta_2, \dots, \delta_m \right).$$

It was shown in [6] that the number of colors in the Modified CFLS coloring is $2^{O(\sqrt{\log n} \log \log n)}$. We will now show that if K_n is edge-colored by the Modified CFLS coloring then every copy of K_5 has an odd color class.

Let $E(K_n)$ be colored by the Modified CFLS coloring φ given by Cameron and Heath [6]. Then every copy of K_5 contains at least 4 distinct colors. If a copy of K_5 contains at least six colors, then clearly some color appears in this K_5 exactly once, giving us the odd color we desire. Therefore, we only need to consider copies of K_5 which contain four or five colors under φ . The case where K_5 contains exactly four colors is easy to understand; indeed, Cameron and Heath [7] characterized the colorings of K_5 with exactly four colors which can appear under φ .

Theorem 2.1. Let $p \ge 3$ and let the edges of K_n be colored with the Modified CFLS (p, p-1)-coloring. The only p-cliques that contain exactly p-1 distinct edge-colors are isomorphic (as edge-colored graphs) to one of the edge-colored p-cliques given in the definition below.

Definition 2.2. Given an edge-coloring $f : E(K_n) \to C$, we say that a subset $S \subseteq V(K_n)$ has a leftover structure under f if either |S| = 1 or there exists a bipartition of S into nonempty sets A and B for which

- A and B each have a leftover structure under f;
- $f(A) \cap f(B) = \emptyset$; and
- there is a fixed color $\alpha \in C$ such that $f(a,b) = \alpha$ for all $a \in A$ and all $b \in B$, and $\alpha \notin f(A)$ and $\alpha \notin f(B)$.

In particular, there are only three such leftover configurations in the p = 5 setting, shown in Figure 1 below. Note that in each case, there is a color which appears exactly once. Thus, it suffices to consider colorings of K_5 with exactly five colors.



Figure 1: The three leftover colorings of K_5 with four colors under φ



Figure 2: Forbidden configurations under φ with four vertices

Assume towards a contradiction that φ permits some coloring of K_5 with exactly 5 colors, each appearing an even number of times. Call these colors $\alpha, \beta, \gamma, \pi$, and θ . Then each color class is either a matching of two disjoint edges or a path with two adjacent edges. We say that such colorings of K_5 are 2-2-2-2 colorings.

Lemma 2.3. No vertex in a 2-2-2-2 coloring of K_5 can be incident to edges of only two colors.

Proof. If there was such a vertex, then the other four vertices would form a copy of K_4 with three colors appearing twice each. But, each such coloring of K_4 is one of the configurations shown in Figure 2, which were shown to be forbidden under φ in [6].

Now we will prove a series of lemmas showing that several other common configurations (in Figure 3) are forbidden under φ . Throughout these proofs, when referring to some color α , we will let α_0 denote the 0-coordinate of the color (of the form $(i, \{x, y\})$) and let α_k denote the k-coordinate of the color, that is, the index of the first bit of difference between the kth blocks. Furthermore, if $x, y \in \{0, 1\}^{m^2}$ are two vectors and $x^{(k)} = y^{(k)}$ (where $x^{(k)}$ is the kth m-block of x), then we say that x and y agree at k; otherwise, we say x and y disagree at k.

Lemma 2.4. The configuration with five distinct vertices $a, b, c, d, e \in V$ for which $\varphi(be) = \varphi(cd)$, $\varphi(ab) = \varphi(de)$, and $\varphi(bc) = \varphi(ae)$, as in Figure 3a, is forbidden.

Proof. Assume towards a contradiction that for distinct vertices a, b, c, d, e, we have $\varphi(be) = \varphi(cd) = \alpha$, $\varphi(ab) = \varphi(de) = \beta$, and $\varphi(bc) = \varphi(ae) = \gamma$. Let $\alpha_0 = (i, \{x, y\})$. First suppose $b^{(i)} = c^{(i)} = x$ and $e^{(i)} = d^{(i)} = y$. Then $\beta_i = 0$ since d and e agree at i, and $\gamma_i = 0$ since b and c agree at i. But then $a^{(i)} = x$ (because $\varphi(ab) = \beta$, $\beta_i = 0$ and $b^{(i)} = x$), and similarly $a^{(i)} = y$ (because $\varphi(ae) = \gamma$, $\gamma_i = 0$ and $e^{(i)} = y$), a contradiction since $x \neq y$. So, instead suppose that $b^{(i)} = d^{(i)} = x$ and $c^{(i)} = e^{(i)} = y$. Then $\beta_i = t$, the first index of difference between $d^{(i)} = x$ and $e^{(i)} = y$. Similarly, $\gamma_i = t$. Let $a^{(i)} = w$. Then since $\varphi(ab) = \beta$ and $\beta_i = t$, we know $x_j = w_j$ for all $1 \leq j \leq t - 1$ and $x_t \neq w_t$. Similarly, since $\varphi(ae) = \gamma$ and $\gamma_i = t$, we have $y_j = w_j$ for all $1 \leq j \leq t - 1$ and $y_t \neq w_t$. But since $\{x_t, y_t\} = \{0, 1\}$, this leaves no choice for w_t , a contradiction.

Lemma 2.5. The configuration with five distinct vertices $a, b, c, d, e \in V$ for which $\varphi(ac) = \varphi(bc), \varphi(ab) = \varphi(bd), \varphi(ad) = \varphi(de), and \varphi(ae) = \varphi(ec), as in Figure 3b, is forbidden.$



Figure 3: Forbidden configurations under φ with five vertices

Proof. Assume for contradiction that for distinct vertices a, b, c, d, e, we have $\varphi(ac) = \varphi(bc) = \alpha$, $\varphi(ab) = \varphi(bd) = \beta$, $\varphi(ad) = \varphi(de) = \gamma$, and $\varphi(ae) = \varphi(ec) = \pi$. Let $\alpha_0 = (i, \{x, y\})$, and let $c^{(i)} = x$ and $a^{(i)} = b^{(i)} = y$. Then $\beta_i = 0$ since a and b agree at i, which implies $d^{(i)} = y$ as well. Now we conclude $\gamma_i = 0$ since a and d agree at i, which implies $e^{(i)} = y$ too. But now $\pi_i = 0$ since a and e agree at i, while also $\pi_i \neq 0$ since c and e disagree at i, a contradiction.

Lemma 2.6. Any 2-2-2-2 coloring of K_5 with a rainbow 5-cycle and at least one matching, as in Figure 3c, is forbidden under φ .

Proof. Assume towards a contradiction that for distinct vertices $a, b, c, d, e \in V$, we have $\varphi(be) = \varphi(cd) = \alpha$, $\varphi(ab) = \beta$, $\varphi(bc) = \gamma$, $\varphi(de) = \pi$, and $\varphi(ea) = \theta$. Note that since this is a 2-2-2-2-2 coloring of K_5 , we have $\{\varphi(bd), \varphi(ce)\} = \{\beta, \theta\}$, since otherwise we would find a copy of the forbidden configuration shown in Figure 2a using the colors $\{\alpha, \gamma\}$ or $\{\alpha, \pi\}$. This implies $\{\varphi(ac), \varphi(ad)\} = \{\pi, \gamma\}$.

Suppose first that $\varphi(bd) = \beta$ and $\varphi(ce) = \theta$. If $\varphi(ac) = \pi$ and $\varphi(ad) = \gamma$, we find the forbidden configuration shown in Figure 3a in colors $\{\alpha, \gamma, \pi\}$. If instead $\varphi(ac) = \gamma$ and $\varphi(ad) = \pi$, then we find the forbidden configuration shown in Figure 3b in colors $\{\beta, \gamma, \pi, \theta\}$. So, we must have $\varphi(bd) = \theta$ and $\varphi(ce) = \beta$, and we get the forbidden configuration shown in Figure 3a in colors $\{\alpha, \beta, \theta\}$, a contradiction.

Lemma 2.7. The configuration with five distinct vertices $a, b, c, d, e \in V$ for which $\varphi(bc) = \varphi(de)$, $\varphi(ad) = \varphi(cd)$, and $\varphi(ab) = \varphi(ac)$ as in Figure 3d, is forbidden.

Proof. Assume towards a contradiction that for distinct vertices a, b, c, d, e, we have $\varphi(bc) = \varphi(de) = \alpha$, $\varphi(ad) = \varphi(cd) = \beta$, and $\varphi(ab) = \varphi(ac) = \gamma$. Let $\alpha_0 = (i, \{x, y\})$. First suppose $b^{(i)} = e^{(i)} = x$ and $c^{(i)} = d^{(i)} = y$. Then $\beta_i = 0$ since b and c agree at i, and hence $a^{(i)} = y$ as well. But now $\gamma_i = 0$ since a and c agree at i while also $\gamma_i \neq 0$ since a and b disagree at i, a contradiction. So, it must be the case that $b^{(i)} = d^{(i)} = x$ and $c^{(i)} = e^{(i)} = y$.

Now $\beta_i = t$, the first index of difference between $c^{(i)} = y$ and $d^{(i)} = x$. Say $a^{(i)} = w$. Since $\varphi(ad) = \beta$, we know that $x_j = w_j$ for all $1 \le j \le t - 1$ and $x_t \ne w_t$. Then since $x_t \ne y_t$ as well, we have $w_t = y_t$. Furthermore, $\beta_i = t$ implies that $w_j = x_j = y_j$ for all $1 \le j \le t - 1$, so the first index of difference between w and y must be greater than t. Now consider γ_i . Since $\varphi(ac) = \gamma$, we know $\gamma_i > t$. But $\varphi(ab) = \gamma$ implies that $\gamma_i = t$, a contradiction.

Lemma 2.8. The configuration with five distinct vertices $a, b, c, d, e \in V$ for which $\varphi(ab) = \varphi(be), \varphi(bc) = \varphi(ac), \varphi(cd) = \varphi(bd), \varphi(de) = \varphi(ce), and \varphi(ae) = \varphi(ad), as in Figure 3e, is forbidden.$

Proof. Assume towards a contradiction that for distinct vertices a, b, c, d, e, we have $\varphi(ab) = \varphi(be) = \alpha$, $\varphi(bc) = \varphi(ac) = \beta$, $\varphi(cd) = \varphi(bd) = \gamma$, $\varphi(de) = \varphi(ce) = \pi$, and $\varphi(ae) = \varphi(ad) = \theta$. Let $\alpha_0 = (i, \{x, y\})$, and let $a^{(i)} = e^{(i)} = x$ and $b^{(i)} = y$. Then $\theta_i = 0$ since a and e agree at i, which implies $d^{(i)} = x$ as well. Similarly, $\pi_i = 0$ since d and e agree at i, which forces $c^{(i)} = x$. But now $\gamma_i = 0$ because c and d agree at i, while $\gamma_i \neq 0$ since b and d disagree at i, a contradiction.

Finally, we are ready to show that every 2-2-2-2 coloring of K_5 is forbidden by φ . We will do this by splitting our proof into cases based on the number of matchings and paths in a 2-2-2-2 configuration.

Lemma 2.9. There is no 2-2-2-2-2 configuration in φ in which all five color classes are paths of length 2.

Proof. Assume for contradiction that such a coloring of K_5 exists under φ . First, we will show that this K_5 must contain a rainbow cycle. To this end, note that each monochromatic path of length 2 is contained in two 5-cycles in our K_5 . So if all the 5 colors are length 2 paths, there are at most 10 cycles in K_5 that are not rainbow. But since there are 12 distinct 5-cycles in K_5 , there must be at least two 5-cycles containing no monochromatic path; that is, there must be some rainbow cycle.

Consider this rainbow cycle and suppose $\varphi(ab) = \alpha$, $\varphi(bc) = \beta$, $\varphi(cd) = \gamma$, $\varphi(de) = \pi$, and $\varphi(ea) = \theta$. Since the color class of α forms a length 2 path, there are two cases up to symmetry: either $\varphi(be) = \alpha$, or $\varphi(bd) = \alpha$.

First suppose $\varphi(be) = \alpha$. Note that $\varphi(bd) \in \{\gamma, \pi\}$, because by Lemma 2.3, b must be adjacent to edges of at least three colors, and we know each color class is a path. This forces $\varphi(ad) = \theta$ because d must be adjacent to edges of at least three colors by Lemma 2.3, and since β forms a path. If $\varphi(bd) = \pi$, then $\{\varphi(ac), \varphi(ce)\} = \{\beta, \gamma\}$, contradicting Lemma 2.3 at c. So, it must be the case that $\varphi(bd) = \gamma$ and $\{\varphi(ac), \varphi(ce)\} = \{\beta, \pi\}$. Since π forms a path, this implies $\varphi(ac) = \beta$ and $\varphi(ce) = \pi$. However, this is the forbidden configuration in Figure 3e, so we reach a contradiction.

Now instead suppose $\varphi(bd) = \alpha$. By Lemma 2.3, and because γ must form a path, we have $\varphi(be) \in \{\pi, \theta\}$. Then $\varphi(ce) \in \{\beta, \gamma\}$ since Lemma 2.3 requires at least three colors on the edges incident to e. This forces $\varphi(ac) = \theta$, which in turns implies $\varphi(be) = \pi$. Finally, since β must be a path, we have $\varphi(ad) = \gamma$ and $\varphi(ce) = \beta$. But now we again have the forbidden configuration in Figure 3e, a contradiction.

Lemma 2.10. There is no 2-2-2-2-2 configuration in φ in which one color classes is a matching and the rest are paths.

Proof. Assume towards a contradiction that distinct vertices $a, b, c, d, e \in V$ form a 2-2-2-2 coloring of K_5 where one color class is a matching and four are paths. Let $\varphi(ab) = \varphi(cd) = \alpha$. Consider the path in color β . If e is incident to two edges of color β , then by Lemma 2.3, the other two edges incident to e must receive distinct colors, say γ and π . Now the fifth color θ must appear on a path in the subgraph induced by $\{a, b, c, d\}$. But this forms a forbidden configuration as in Figure 2a in colors θ and α , a contradiction. Since the same argument applies to γ , π , and θ , we know e is incident to edges of four distinct colors. However, there is only one such configuration in which the color classes of β, γ, π , and θ are all paths, and it is the forbidden configuration shown in Figure 3b, so we again reach a contradiction.

Lemma 2.11. There is no 2-2-2-2-2 configuration in φ in which two color classes are matchings and three are paths.

Proof. Assume for contradiction that distinct vertices $a, b, c, d, e \in V$ form a 2-2-2-2 coloring of K_5 with matchings in colors α and β and paths in colors γ, π, θ . If α and β form an alternating path, say $\varphi(ab) = \varphi(cd) = \alpha$ and $\varphi(bc) = \varphi(de) = \beta$, then consider the edge bd. Without loss of generality, $\varphi(bd) = \gamma$. If $\varphi(be) = \gamma$ or $\varphi(ad) = \gamma$, then there is a copy of the forbidden configuration in Figure 2a. So, the other edge

of color γ must be one of *ac*, *ce*, or *ae*. But in these cases, γ would form a matching rather than a path, a contradiction.

Now suppose α and β form an alternating cycle, say $\varphi(ab) = \varphi(cd) = \alpha$ and $\varphi(bc) = \varphi(ad) = \beta$. Since all remaining edges receives colors in $\{\gamma, \pi, \theta\}$, we know that the two edges incident to e must receive the same color. Without loss of generality, say $\varphi(ae) = \varphi(be) = \gamma$. Lemma 2.3 implies that π and θ each appear on one of the edges ce and de. Furthermore, since the color classes of π and θ are paths, we may assume that $\varphi(ac) = \varphi(ce) = \pi$ and $\varphi(bd) = \varphi(de) = \theta$. But now the colors α, γ , and π form a forbidden configuration as in Figure 3d, a contradiction.

Lemma 2.12. There is no 2-2-2-2-2 configuration in φ in which three color classes are matchings and two are paths.

Proof. Assume for contradiction that distinct vertices $a, b, c, d, e \in V$ form a 2-2-2-2 coloring of K_5 with three color classes that are matchings and two color classes that are paths. Let $\varphi(ab) = \varphi(bc) = \alpha$ and $\varphi(ac) = \beta$. Then either β is a path, or a matching.

First suppose β forms a path. Without loss of generality, $\varphi(cd) = \beta$. Then *ad* receives some third color, say $\varphi(ad) = \gamma$, and γ forms a matching. We cannot have $\varphi(ec) = \gamma$ without forming a forbidden configuration as in Figure 2a with colors $\{\beta, \gamma\}$, so $\varphi(be) = \gamma$ is forced. Now the remaining three edges incident to *e* should be colored with the two colors π and θ , contradicting the fact that π and θ both form a matching each.

Therefore, β must form a matching. Since we must avoid the forbidden configuration in Figure 2a, the other edge colored β cannot be incident to b, which implies $\varphi(de) = \beta$. Consider where the second monochromatic path, say in color γ , can occur. If b is not incident to an edge of color γ , then we find a forbidden configuration as in Figure 2a in colors $\{\beta, \gamma\}$. So, b must be incident to an edge of color γ , say $\varphi(bd) = \gamma$. By Lemma 2.3, $\varphi(be) \neq \gamma$, so either cd or ad also receive color γ .

In both cases, there are three remaining edges incident to e which must be colored with $\{\pi, \theta\}$, but both π and θ form matchings in the K_5 , a contradiction.

Lemma 2.13. There is no 2-2-2-2-2 configuration in φ in which four color classes are matchings and one is a path.

Proof. Assume towards a contradiction that distinct vertices $a, b, c, d, e \in V$ form a 2-2-2-2 coloring of K_5 with a path in color α and four matchings. Let $\varphi(ab) = \varphi(bc) = \alpha$, and say $\varphi(bc) = \beta$. Then β cannot appear on the edges bd or be, otherwise there is a copy of the forbidden configuration in Figure 2a. So, since β forms a matching, we must have $\varphi(de) = \beta$. Now the remaining six edges in this K_5 form a copy of $K_{2,3}$ on vertex set $\{a, b, c\} \cup \{d, e\}$ which must be colored with three colors, but this cannot occur when all three color classes are matchings of size two.

Lemma 2.14. There is no 2-2-2-2-2 configuration in φ in which all five color classes are matchings.

Proof. Assume towards a contradiction that distinct vertices $a, b, c, d, e \in V$ form a 2-2-2-2 coloring of K_5 where each color class is a matching. Without loss of generality, assume $\varphi(ab) = \varphi(cd) = \alpha$ and $\varphi(bc) = \beta$. Then up to symmetry, there are two cases: either $\varphi(ad) = \beta$ or $\varphi(de) = \beta$.

Consider the case $\varphi(ad) = \beta$. Let $\varphi(ae) = \gamma$. Then since γ is a matching, we must have $\varphi(bd) = \gamma$ as well. But then the remaining two colors cannot both form matchings, since two of the edges *be*, *ce*, and *de* must receive the same color by pigeon-hole principle.

Thus, we may assume $\varphi(de) = \beta$ instead. Let $\varphi(ae) = \gamma$. Then since γ must form a matching, we have $\varphi(bd) = \gamma$ as well. However, this forms the forbidden configuration shown in Figure 3a, a contradiction.

Combining the results of Lemmas 2.9-2.14 shows that φ forbids all 2-2-2-2 colorings of K_5 . Therefore, any K_5 which appears under φ must contain some color an odd number of times.

3 Forbidden submatching method

We will use the simplified version presented in [15] of the conflict-free hypergraph matching theorem from [13] to prove Theorem 1.4. In order to state this theorem, we will need to introduce some terminology and notation.

Given a hypergraph \mathcal{H} and a vertex $v \in V(\mathcal{H})$, its degree $\deg_{\mathcal{H}}(v)$ is the number of edges in \mathcal{H} containing v. The maximum degree and minimum degree of \mathcal{H} are denoted by $\Delta(\mathcal{H})$ and $\delta(\mathcal{H})$, respectively. For $j \geq 2$, $\Delta_j(\mathcal{H})$ denotes the maximum number of edges in \mathcal{H} which contain a particular set of j vertices, over all such sets.

In addition, for a (not necessarily uniform) hypergraph C and an integer k, let $C^{(k)}$ be the set of edges in C of size k. For a vertex $u \in V(C)$, let C_u denote the hypergraph $\{C \setminus \{u\} \mid C \in E(C), u \in C\}$.

Given a hypergraph \mathcal{H} , a hypergraph \mathcal{C} is a *conflict system* for \mathcal{H} if $V(\mathcal{C}) = E(\mathcal{H})$. A set of edges $E \subset \mathcal{H}$ is \mathcal{C} -free if E contains no subset $C \in \mathcal{C}$. Given integers $d \geq 1$, $\ell \geq 3$, and $\varepsilon \in (0, 1)$, we say \mathcal{C} is (d, ℓ, ε) -bounded if \mathcal{C} satisfies the following conditions:

(C1) $3 \leq |C| \leq \ell$ for all $C \in \mathcal{C}$;

(C2) $\Delta(\mathcal{C}^{(j)}) \leq \ell d^{j-1}$ for all $3 \leq j \leq \ell$;

(C3) $\Delta_{j'}(\mathcal{C}^{(j)}) \leq d^{j-j'-\varepsilon}$ for all $2 \leq j' < j \leq \ell$.

Finally, given a (d, ℓ, ε) -bounded conflict system \mathcal{C} for a hypergraph \mathcal{H} , we will define a type of weight function which can be used to guarantee that the almost-perfect matching given by Theorem 3.1 below satisfies certain quasirandom properties. We say a function $w : \binom{\mathcal{H}}{j} \to [0, \ell]$ for $j \in \mathbb{N}$ is a *test function* for \mathcal{H} if w(E) = 0 whenever $E \in \binom{\mathcal{H}}{j}$ is not a matching, and we say w is *j*-uniform. For a function $w : A \to \mathbb{R}$ and a finite set $X \subset A$, let $w(X) := \sum_{x \in X} w(x)$. If w is a *j*-uniform test function, then for each $E \subset \mathcal{H}$, let $w(E) = w(\binom{E}{j})$. Given $j, d \in \mathbb{N}, \varepsilon > 0$, and a conflict system \mathcal{C} for hypergraph \mathcal{H} , we say a *j*-uniform test function w for \mathcal{H} is $(d, \varepsilon, \mathcal{C})$ -trackable if w satisfies the following conditions:

(W1)
$$w(\mathcal{H}) \ge d^{j+\varepsilon};$$

(W2)
$$w(\{E \in \binom{\mathcal{H}}{i} : E \supseteq E'\}) \le w(\mathcal{H})/d^{j'+\varepsilon}$$
 for all $j' \in [j-1]$ and $E' \in \binom{\mathcal{H}}{i'}$;

(W3) $|(\mathcal{C}_e)^{(j')} \cap (\mathcal{C}_f)^{(j')}| \leq d^{j'-\varepsilon}$ for all $e, f \in \mathcal{H}$ with $w(\{E \in \binom{\mathcal{H}}{i} : e, f \in E\}) > 0$ and all $j' \in [\ell-1];$

(W4) w(E) = 0 for all $E \in \binom{\mathcal{H}}{i}$ that are not \mathcal{C} -free.

Theorem 3.1 ([13], Theorem 3.3). For all $k, \ell \geq 2$, there exists $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, there exists d_0 such that the following holds for all $d \geq d_0$. Suppose \mathcal{H} is a k-regular hypergraph on $n \leq \exp(d^{\varepsilon^3})$ vertices with $(1 - d^{-\varepsilon})d \leq \delta(\mathcal{H}) \leq \Delta(\mathcal{H}) \leq d$ and $\Delta_2(\mathcal{H}) \leq d^{1-\varepsilon}$. Suppose \mathcal{C} is a (d, ℓ, ε) -bounded conflict system for \mathcal{H} , and suppose \mathcal{W} is a set of $(d, \varepsilon, \mathcal{C})$ -trackable test functions for \mathcal{H} of uniformity at most ℓ with $|\mathcal{W}| \leq \exp(d^{\varepsilon^3})$. Then, there exists a \mathcal{C} -free matching $\mathcal{M} \subset \mathcal{H}$ of size at least $(1 - d^{-\varepsilon^3})n/k$ with $w(\mathcal{M}) = (1 \pm d^{-\varepsilon^3})d^{-j}w(\mathcal{H}))$ for all j-uniform $w \in \mathcal{W}$.

We will say that a hypergraph \mathcal{H} with $(1 - d^{-\varepsilon})d \leq \delta(\mathcal{H}) \leq \Delta(\mathcal{H}) \leq d$ is almost d-regular.

4 Proof of Theorem 1.4

First, observe that $g(K_{n,n}, C_4) > n/2$. Indeed, let $V(K_{n,n}) = X \cup Y$ and suppose the edges of $K_{n,n}$ are colored with at most n/2 colors. Then for each vertex $x \in X$, there are at least n/2 pairs $u, v \in Y$ such that xu and xv receive the same color. Since there are $\binom{n}{2}$ pairs of vertices in Y, the pigeonhole principle implies that there must be two vertices $x, z \in X$ such that xu and xv share a color and zu and zv share a color, giving a copy of C_4 with no color appearing an odd number of times.

From now on, we remove the ceiling function notation in order to make the proof easier to read. However, whenever n is odd, when we write $\frac{n}{2}$ we mean $\lceil \frac{n}{2} \rceil$.

In order to prove the upper bound in Theorem 1.4, we will construct a coloring of $K_{n,n}$ in two stages. The coloring in the first stage will use k = n/2 colors to color a majority of the edges of $K_{n,n}$. To determine this coloring, we define appropriate hypergraphs \mathcal{H} and \mathcal{C} for which a \mathcal{C} -free matching in \mathcal{H} corresponds to a partial coloring of $K_{n,n}$ with monochromatic copies of C_6 . Each copy of C_4 in the resulting coloring sees some color an odd number of times.

Theorem 4.1 below guarantees that our choices of \mathcal{H} and \mathcal{C} satisfy the requirements for applying the forbidding submatching method to find the desired matching. In the second stage of the coloring, we then randomly color the remaining uncolored edges using a new set of o(n) colors.

Theorem 4.1. There exists $\delta > 0$ such that for all sufficiently large n in terms of δ , there is an edge-coloring of a subgraph $F \subset K_{n,n}$ with at most n/2 colors and the following properties:

- 1. Every color class is a union of vertex-disjoint copies of C_6 .
- 2. Every copy of C_4 in F intersects some color class in an odd number of edges.
- 3. The graph $L = K_{n,n} \setminus E(F)$ has maximum degree at most $n^{1-\delta}$.
- 4. Let $V(K_{n,n}) = X \cup Y$. For each $(x, y) \in X \times Y$, the number of edges $x'y' \in E(L)$ with $x' \in X \setminus \{x\}$, $y' \in Y \setminus \{y\}$ such that xy' and yx' receive the same color is at most $4n^{1-\delta}$.

Proof. Let $U = \binom{X \cup Y}{2}$ and $V = \bigcup_{i \in [\frac{n}{2}]} V_i$, where each V_i is a copy of $X \cup Y$. We will denote the copy of $v \in X \cup Y$ in V_i by v^i . Let \mathcal{H} be the 18-uniform hypergraph with vertex set $U \cup V$ and edges e_i for each copy e of C_6 in $K_{n,n}$ and $i \in [\frac{n}{2}]$ defined as follows. Given a 6-cycle $e = x_1y_1x_2y_2x_3y_3$ with $x_j \in X$ and $y_j \in Y$, let \mathcal{H} contain the edge

$$e_i = \{x_1y_1, y_1x_2, x_2y_2, y_2x_3, x_3y_3, y_3x_1\} \cup \{x_1x_2, x_2x_3, x_1x_3, y_1y_2, y_2y_3, y_1y_3\} \cup \{x_1^i, x_2^i, x_3^i, y_1^i, y_2^i, y_3^i\}$$

Note that a matching in \mathcal{H} corresponds to a set of edge-disjoint monochromatic copies of C_6 in $K_{n,n}$. Furthermore, no vertex in the matching appears in two cycles of the same color. Thus, a matching in \mathcal{H} yields an edge-coloring of a subgraph of $K_{n,n}$ with at most $\frac{n}{2}$ colors which satisfies property (1) above.

The hypergraph \mathcal{H} is essentially *d*-regular for $d = \frac{1}{2}n^5$. Indeed, we can check that for each vertex u in \mathcal{H} , $d - O(n^4) \leq \deg_{\mathcal{H}}(u) \leq d$. First, suppose $u \in V_i$ for some i. There are $n^3\binom{n}{2} - O(n^4)$ ways to pick an edge in \mathcal{H} containing u, since there are $\binom{n}{2}$ choices for the neighbors of u and n - O(1) choices for each of the remaining vertices along the cycle. If instead u = xy for some $x \in X$ and $y \in Y$, then there are $\frac{1}{2}n$ ways to pick a color of an edge containing xy and $n^4 - O(n^3)$ ways to pick the remaining vertices along the cycle. Similarly, if u = xx' for $x, x' \in X$ (or yy' for $y, y' \in Y$), there are $\frac{1}{2}n^5 - O(n^4)$ edges in \mathcal{H} containing the edge. Thus, in each case, we have $\deg_{\mathcal{H}}(u) = \frac{1}{2}n^5 - O(n^4)$.

Furthermore, we have $\Delta_2(\mathcal{H}) \leq d^{1-\varepsilon}$ for all $\varepsilon \in (0, \frac{1}{6})$, since each pair of vertices $u, v \in V(\mathcal{H})$ is contained in at most $O(n^4)$ edges in \mathcal{H} .

Next we define a conflict system C for \mathcal{H} with edges of size 3 and 4. Let $V(\mathcal{C}) = E(\mathcal{H})$, and let the edges of C correspond to copies of C_4 in $K_{n,n}$ formed by two monochromatic matchings of size 2. We will call these edges of C conflicts, and they correspond to sets of three or four monochromatic cycles in $K_{n,n}$. That is, given four vertices $x, x' \in X$ and $y, y' \in Y$, colors $i, j \in [\frac{n}{2}]$ and edges $e_i, e'_i, f_j, f'_j \in E(\mathcal{H})$ (not necessarily distinct) with $xy \in e_i, x'y' \in e'_i, xy' \in f_j$, and $x'y \in f'_j$, we have $\{e_i, e'_i, f_j, f'_j\} \in E(\mathcal{C})$. Note that we do not need to consider conflicts of size 2: it is possible to have two monochromatic copies e, e' of C_6 that form a copy of C_4 with alternating colors, however such e, e' would have to share two vertices in X (and two vertices in Y), and thus cannot appear together in a matching of \mathcal{H} .

We will show that $\Delta(\mathcal{C}^{(3)}) = O(d^2)$ and $\Delta(\mathcal{C}^{(4)}) = O(d^3)$. To this end, fix $e_i \in V(\mathcal{C})$. Note that any conflict in $\mathcal{C}^{(3)}$ containing e_i must either have the form $\{e_i, e'_i, f_j\}$ or $\{e_i, f_j, f'_j\}$ for some copies e', f, f' of C_6 in $K_{n,n}$ and some color $j \in [\frac{n}{2}] \setminus \{i\}$. In either case, there are O(n) ways to pick j. If the edge has the form $\{e_i, e'_i, f_j\}$, then there are $O(n^6)$ ways to pick e' and then $O(n^2)$ ways to pick f. If instead the edge has the form $\{e_i, f_j, f'_j\}$, then there are $O(n^4)$ ways to pick f and $O(n^4)$ ways to pick f'. Thus, in either case, $d_{\mathcal{C}}(e_i) = O(n^9)$, and we have $\Delta(\mathcal{C}^{(3)}) = O(d^2)$. Similarly, any conflict in $\mathcal{C}^{(4)}$ containing e_i must have the form $\{e_i, e'_i, f_j, f'_j\}$ for some copies e, e', f, f' of C_6 and some color $j \in [\frac{n}{2}]$, so there are O(n) choices for j, $O(n^6)$ choices for e', and $O(n^4)$ choices for each of f and f'. Thus, $\Delta(\mathcal{C}^{(4)}) = O(n^{15}) = O(d^3)$.

In addition, we have $\Delta_2(\mathcal{C}^{(3)}) < d^{1-\varepsilon}$ for all $\varepsilon \in (0, \frac{1}{6})$. Indeed, the number of conflicts in $\mathcal{C}^{(3)}$ containing two fixed vertices $e_i, e'_i \in V(\mathcal{C})$ is $O(n^3)$, since there are O(n) ways to pick another color and $O(n^2)$ ways to pick two more vertices for the third edge in the conflict. Similarly, the number of conflicts in $\mathcal{C}^{(3)}$ containing two fixed vertices $e_i, f_j \in V(\mathcal{C})$ is $O(n^4)$, which is the number of ways to pick four more vertices for the third edge in the conflict. Similarly, we have $\Delta_2(\mathcal{C}^{(4)}) < d^{2-\varepsilon}$ for all $\varepsilon \in (0, \frac{1}{6})$. The number of conflicts in $\mathcal{C}^{(4)}$ containing two fixed vertices $e_i, e'_i \in V(\mathcal{C})$ is $O(n^9)$ since there are O(n) choices for a second color and $O(n^4)$ choices for each of f and f' to complete the conflict, and the number of conflicts in $\mathcal{C}^{(4)}$ containing two fixed vertices $e_i, f_j \in V(\mathcal{C})$ is $O(n^9)$ since there are $O(n^5)$ choices for e' and $O(n^4)$ choices for f' to complete the conflict. Finally, $\Delta_3(\mathcal{C}^{(4)}) < d^{1-\varepsilon}$, because given three vertices $e_i, e'_i, f_j \in V(\mathcal{C})$, there are $O(n^4)$ choices for f' to complete a conflict. Therefore, \mathcal{C} is a $(d, O(1), \varepsilon)$ -bounded conflict system for \mathcal{H} for all $\varepsilon \in (0, \frac{1}{6})$.

Note that we could apply Theorem 3.1 at this point to obtain a conflict-free matching M in \mathcal{H} which would correspond to a coloring of a subgraph F of $K_{n,n}$ satisfying properties (1) and (2) of Theorem 4.1. In order to obtain a coloring which also satisfies properties (3) and (4) of Theorem 4.1, we now introduce appropriate test functions.

First, we define a set of 1-uniform test functions as in [15]. For each $v \in X \cup Y$, let $S_v \subset U$ be the set of edges incident to v in $K_{n,n}$. Let $w_v : E(\mathcal{H}) \to \{0,2\}$ be the weight function which assigns to each edge $e_i \in E(\mathcal{H})$ the size of its intersection with S_v . Then we have

$$w_v(\mathcal{H}) = \sum_{e_i \in \mathcal{H}} w_v(e_i) = \sum_{f \in S_v} \deg_{\mathcal{H}}(f) = nd - O(n^5) \ge d^{1+\varepsilon},$$

satisfying condition (W1). In addition, conditions (W2)-(W4) are trivially satisfied since w_v is 1-uniform, so w_v is a $(d, \varepsilon, \mathcal{C})$ -trackable test function for all $\varepsilon \in (0, \frac{1}{6})$ and $v \in X \cup Y$.

Let $\delta < \varepsilon^3 \log_n(\frac{1}{2}n^5)$. Note that applying Theorem 3.1 with these test functions would yield a C-free matching $M \subset \mathcal{H}$ such that for each $v \in X \cup Y$,

$$w_v(M) > (1 - d^{-\varepsilon^3})d^{-1}w_v(\mathcal{H}) > (1 - n^{-\delta})n.$$

Thus, for each $v \in X \cup Y$, there are at most $n^{1-\delta}$ edges in $K_{n,n}$ incident to v which do not belong to any edge in M. Hence, property (3) is satisfied, as the graph L containing the uncolored edges has maximum degree $\Delta(L) \leq n^{1-\delta}$.

In order to guarantee that property (4) is also satisfied, we will define four more types of test functions. The first two will help us to show that for each $(x, y) \in X \times Y$, there are at most $n^{1-\delta}$ edges $x'y' \in E(L)$ with $x' \in X \setminus \{x\}$ and $y' \in Y \setminus \{y\}$ such that xy' and yx' are in distinct edges of the same color in the matching $M \subset \mathcal{H}$. The next two will help us to show that for each $(x, y) \in X \times Y$, there are at most $n^{1-\delta}$ edges in the matching that contain both x and y, and hence at most $3n^{1-\delta}$ graph edges $x'y' \in E(L)$ with $x' \in X \setminus \{x\}$ and $y' \in Y \setminus \{y\}$ such that xy are contained in the same edge of the matching.

For each pair of vertices $(x, y) \in X \times Y$, let

$$\mathcal{P}_{x,y} = \left\{ \{e_i, e_i'\} : i \in \left[\frac{n}{2}\right], V(e) \cap V(e') = \emptyset, x \in V(e), y \in V(e') \right\}.$$

Note that

$$|\mathcal{P}_{x,y}| = \frac{n}{2} \cdot \frac{1}{2} n^5 \cdot \frac{1}{2} n^5 \pm O(n^{10}) = \frac{1}{8} n^{11} \pm O(n^{10}) > d^{2+\varepsilon}.$$

For each $(x, y) \in X \times Y$, define $w_{x,y}$ to be the indicator weight function for the pairs in $\mathcal{P}_{x,y}$. Since each pair in $\mathcal{P}_{x,y}$ is a matching of size 2 in \mathcal{H} , $w_{x,y}$ is a 2-uniform test function. We will show that $w_{x,y}$ is $(d, \varepsilon, \mathcal{C})$ -trackable for all $\varepsilon \in (0, \frac{1}{6})$, but first we will define a set of 3-uniform test functions.

For each pair of vertices $(x, y) \in X \times Y$, let

$$\mathcal{T}_{x,y} = \left\{ \{e_i, e'_i, f_j\} : \{e_i, e'_i\} \in \mathcal{P}_{x,y}, j \in \left[\frac{n}{2}\right] \setminus \{i\}, |V(f) \cap V(e) \cap Y| = 1, |V(f) \cap V(e') \cap X| = 1 \right\}.$$

Thus, the triples in $\mathcal{T}_{x,y}$ are matchings of size 3 in \mathcal{H} . Note that

$$|\mathcal{T}_{x,y}| = 9(d \pm O(n^4))|\mathcal{P}_{x,y}|$$

Similarly to above, define for each $(x, y) \in X \times Y$ the function $w'_{x,y}$ to be the indicator weight function for triples in $\mathcal{T}_{x,y}$. This is a 3-uniform test function for \mathcal{H} .

Assuming that these test functions $w_{x,y}$ and $w'_{x,y}$ are $(d, \varepsilon, \mathcal{C})$ -trackable, we can apply Theorem 3.1 to obtain a matching M in \mathcal{H} such that

$$w_{x,y}(M) = \left| \binom{M}{2} \cap \mathcal{P}_{x,y} \right| \le (1 + d^{-\varepsilon^3}) d^{-2} |\mathcal{P}_{x,y}| \le (1 + n^{-2\delta}) \frac{n}{2}$$

and

$$w'_{x,y}(M) = \left| \binom{M}{3} \cap \mathcal{T}_{x,y} \right| \ge (1 - d^{-\varepsilon^3}) d^{-3} |\mathcal{T}_{x,y}| \ge (1 - n^{-2\delta}) \frac{9n}{2}.$$

Note that each edge $x'y' \in E(L)$ with xy' and x'y appearing in distinct edges of \mathcal{H} of the same color corresponds to an edge of \mathcal{H} in a triple of $\mathcal{T}_{x,y}$ extending some pair from $\mathcal{P}_{x,y}$. Thus, we can bound the number of such x'y' as follows:

$$9\left|\binom{M}{2}\cap\mathcal{P}_{x,y}\right|-\left|\binom{M}{3}\cap\mathcal{T}_{x,y}\right|\leq n^{1-\delta}.$$

Therefore, in order to prove the first case of property (4), we must verify that $w_{x,y}$ and $w'_{x,y}$ are (d, ε, C) -trackable for $\varepsilon \in (0, \frac{1}{6})$. For this we must check conditions (W1)-(W4) for both functions.

Condition (W1) for $w_{x,y}$ is satisfied since $w_{x,y}(\mathcal{H}) = |\mathcal{P}_{x,y}| > d^{2+\varepsilon}$. To check condition (W2), fix an edge $e_i \in \mathcal{H}$. The number of edges e'_i which form a pair in $\mathcal{P}_{x,y}$ with e_i is at most $O(n^5) < n^{11}/d^{1+\varepsilon}$. For condition (W3), note that the only pairs of edges $\{e, f\} \in \binom{\mathcal{H}}{2}$ for which $w_{x,y}(\{e, f\}) > 0$ are pairs $\{e_i, e'_i\}$ with $x \in e_i$ and $y \in e'_i$. Given such a pair, the number of triples of edges in \mathcal{H} which can form a conflict with e_i or e'_i is at most $O(n \cdot n^6 \cdot n^3 \cdot n^3) = O(n^{13}) < d^{3-\varepsilon}$. Finally, condition (W4) is vacuously true, so $w_{x,y}$ is $(d, \varepsilon, \mathcal{C})$ -trackable.

We now verify that $w'_{x,y}$ is $(d, \varepsilon, \mathcal{C})$ -trackable for $\varepsilon \in (0, \frac{1}{6})$ as well. Condition (W1) holds because $w'_{x,y}(\mathcal{H}) = |\mathcal{T}_{x,y}| = \Theta(n^{16}) > d^{3+\varepsilon}$. To see that condition (W2) holds, fix $e_i \in \mathcal{H}$. There are $O(n^{11}) < w'_{x,y}(\mathcal{H})/d^{1+\varepsilon}$ triples in $\mathcal{T}_{x,y}$ containing e_i since we must either pick a second color j, six more vertices for e_i , and four more vertices for f_j to form a triple $\{e_i, e_i', f_j\}$ or a second color j, five more vertices for f_j , and five more vertices for f_j' to form a triple $\{e_i, f_j, f_j'\}$. And, for each pair of edges $\{e_i, e_i'\}$ or $\{e_i, f_j\}$ in \mathcal{H} , there are $O(n^5) < w'_{x,y}(\mathcal{H})/d^{2+\varepsilon}$ triples containing this pair. For condition (W3), fix a pair of edges $\{e, f\} \in \binom{\mathcal{H}}{2}$ which are in at least one triple together in $\mathcal{T}_{x,y}$. Either we have $\{e, f\} = \{e_i, e_i'\}$ or $\{e, f\} = \{e_i, f_j\}$. In the first case, as above, there are $O(n^{13})$ triples of edges in \mathcal{H} which form a conflict with either e_i or e_i' . In the second case, there are at most $O(n^{14}) < d^{3-\varepsilon}$ conflicts containing e_i which use only the colors i and j, and this bounds the number of triples which form a conflict with either e_i or f_j . Thus, condition (W3) holds. Finally, condition (W4) is vacuously true.

We now define for each $(x, y) \in X \times Y$ another pair of test functions to guarantee that there are at most $n^{1-\delta}$ edges in the matching $M \in \mathcal{H}$ which contain both x and y. Set

$$S_x = \left\{ e_i : i \in \left[\frac{n}{2}\right], x \in V(e) \right\}$$

and

$$\mathcal{D}_{x,y} = \left\{ e_i : i \in \left[\frac{n}{2}\right], x \in V(e), y \notin V(e) \right\}.$$

Let u_x and $u'_{x,y}$ be the indicator functions for S_x and $\mathcal{D}_{x,y}$, respectively. Note that $u_x(\mathcal{H}) = |S_x| = \frac{1}{2}n \cdot \frac{1}{2}n^5 \pm O(n^5)$, and similarly $u'_{x,y} = |\mathcal{D}_{x,y}| = \frac{1}{4}n^6 \pm O(n^5)$. Thus, property (W1) holds for both u_x and $u'_{x,y}$. Furthermore, (W2)-(W4) are vacuously true since these are 1-uniform test functions, hence both are $(d, \varepsilon, \mathcal{C})$ -trackable for $\varepsilon \in (0, \frac{1}{6})$.

Applying Theorem 3.1 with all of these test functions gives a C-free matching $M \subset \mathcal{H}$ such that for each $x \in X$,

$$u_x(M) \le (1 + d^{-\varepsilon^3})d^{-1}u_x(\mathcal{H}) \le (1 + n^{-2\delta})\frac{n}{2},$$

and for each $y \in Y$,

$$u'_{x,y}(M) \ge (1 - d^{-\varepsilon^3})d^{-1}|\mathcal{D}_{x,y}| \ge (1 - n^{-2\delta})\frac{n}{2}.$$

Therefore, the number of edges in M which contain both x and y is at most $u_x(M) - u'_{x,y}(M) \le n^{1-\delta}$, as desired. Thus, the coloring obtained from M satisfies property (4).

We now proceed with the proof of Theorem 1.4. Applying Theorem 4.1 with 2δ in place of δ , we obtain a partial coloring of $K_{n,n}$ using at most n/2 colors for which the uncolored subgraph L of $K_{n,n}$ has maximum degree at most $n^{1-2\delta}$. To color these remaining edges, we randomly color with a new set P of $k = n^{1-\delta}$ colors, with each edge in L receiving a color in P with equal probability 1/k, independently of the other edges. We will show using the symmetric local lemma that such a coloring exists which creates no 2-colored C_4 , and hence, $g(K_{n,n}, C_4) \leq n/2 + n^{1-\delta}$.

To this end, we define three types of "bad events," without which every C_4 in our coloring will contain some color exactly once. First, for each pair of adjacent edges $e, f \in E(L)$ and each color $i \in P$, let $A_{e,f,i}$ be the event that e and f receive the same color i. Note that $\mathbb{P}[A_{e,f,i}] = k^{-2}$. Next, for each 4-cycle D in L, let B_D be the event that D is properly-colored with two colors from P. Then $\mathbb{P}[B_D] \leq 2k^{-2}$. Finally, for each color $i \in P$ and each 4-cycle D = xyx'y' in $K_{n,n}$ with $xy, x'y' \in E(L)$ and $xy', x'y \in E(F)$ where xy' and x'y received the same color in the first coloring, let $C_{D,i}$ be the event that xy and x'y' both receive color i. Then $\mathbb{P}[C_{D,i}] = k^{-2}$. Let \mathcal{E} be the collection of all bad events of the types $A_{e,f,i}, B_D$, and $C_{D,i}$.

Fix an event $E \in \mathcal{E}$. Note that E is mutually independent from the set of all events E' which do not involve any of the same edges as E in L. We call such events edge-disjoint from E. There are at most $8\Delta(L)k \leq 8k^2n^{-\delta}$ events of the form $A_{e,f,i}$ which are not edge-disjoint from E. In addition, there are at most $4(\Delta(L))^2 \leq k^2n^{-\delta}$ events B_D and, by property (4) of Theorem 4.1, at most $4(4n^{1-2\delta})k \leq 16k^2n^{-\delta}$ events $C_{D,i}$ which are not edge-disjoint from E. Thus, in total, E is mutually independent of all but at most $25k^2n^{-\delta}$ events in \mathcal{E} . By the local lemma, since $25k^2n^{-\delta} \cdot 2k^{-2} \leq 1/4$, there is a coloring of L which contains none of the events in \mathcal{E} , as desired.

References

- [1] Noga Alon. Graph-codes. arXiv:2301.13305, 2023.
- [2] Noga Alon, Anna Gujgiczer, János Körner, Aleksa Milojević, and Gábor Simonyi. Structured codes of graphs. SIAM Journal on Discrete Mathematics, 2022.
- [3] Patrick Bennett, Michelle Delcourt, Lina Li, and Luke Postle. On generalized Ramsey numbers in the sublinear regime. arXiv:2212.10542, 2022.
- [4] Patrick Bennett, Andrzej Dudek, and Sean English. A random coloring process gives improved bounds for the Erdős–Gyárfás problem on generalized Ramsey numbers. arXiv:2212.06957, 2022.
- [5] Aaron Berger and Yufei Zhao. k₄-intersecting families of graphs. arxiv:2103.12671, 2021.
- [6] Alex Cameron and Emily Heath. A (5,5)-colouring of K_n with few colours. Combinatorics, Probability and Computing, 27(6):892–912, 2018.
- [7] Alex Cameron and Emily Heath. New upper bounds for the Erdős–Gyárfás problem on generalized Ramsey numbers. Combinatorics, Probability and Computing, 32(2):349–362, 2023.
- [8] David Conlon, Jacob Fox, Choongbum Lee, and Benny Sudakov. The Erdős–Gyárfás problem on generalized Ramsey numbers. *Proceedings of the London Mathematical Society*, 110(1):1–18, 2015.
- [9] Michelle Delcourt and Luke Postle. Finding an almost perfect matching in a hypergraph avoiding forbidden submatchings. arXiv:2204.08981, 2022.

- [10] David Ellis, Yuval Filmus, and Ehud Friedgut. Triangle-intersecting families of graphs. Journal of the European Mathematical Society, 14:841–885, 2010.
- [11] Paul Erdős and András Gyárfás. A variant of the classical Ramsey problem. Combinatorica, 17(4):459–467, 1997.
- [12] Gennian Ge, Zixiang Xu, and Yixuan Zhang. A new variant of the Erdős–Gyárfás problem on k_5 . arXiv:2306.14682, 2023.
- [13] Stefan Glock, Felix Joos, Jaehoon Kim, Marcus Kühn, and Lyuben Lichev. Conflict-free hypergraph matchings. In Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), pages 2991–3005. SIAM, 2023.
- [14] W. T. Gowers. https://gowers.wordpress.com/2009/11/14/the-first-unknown-case-of-polynomial-dhj/.
- [15] Felix Joos and Dhruv Mubayi. Ramsey theory constructions from hypergraph matchings. arXiv:2208.12563, 2022.
- [16] Dhruv Mubayi. Edge-coloring cliques with three colors on all 4-cliques. Combinatorica, 18(2):293–296, 1998.