

BETTER BOUNDS ON THE MINIMAL LEE DISTANCE

JESSICA BARIFFI^{1,2} AND VIOLETTA WEGER³

ABSTRACT. This paper provides new and improved Singleton-like bounds for Lee metric codes over integer residue rings. We derive the bounds using various novel definitions of generalized Lee weights based on different notions of a support of a linear code. In this regard, we introduce three main different support types for codes in the Lee metric and analyze their utility to derive bounds on the minimum Lee distance. Eventually, we propose a new point of view to generalized weights and give an improved bound on the minimum distance of codes in the Lee metric for which we discuss the density of maximum Lee distance codes with respect to this novel Singleton-like bound.

1. INTRODUCTION

The Lee metric was introduced in 1958 by Lee [24] to cope with phase modulation in communication. It provides an interesting alternative to the Hamming and rank metric which are considered for orthogonal modulation and network coding, respectively. The Lee metric is most known for the celebrated result in [18] where the authors showed that some optimal non-linear binary codes can be represented as linear codes over $\mathbb{Z}/4\mathbb{Z}$ endowed with the Lee metric. Recently, there is a renewed interest in the Lee metric due to its similarities with the Euclidean norm used in lattice-based cryptography. In fact, the Lee metric was introduced to cryptography in [20] and its hard problems were further studied in [33]. Recently, a first Lee-metric primitive [28] has been submitted to the reopened NIST standardization process for digital signature schemes.

Although the Lee metric is one of the oldest metrics and has interesting properties and applications, it did not receive as much attention as other metrics which is visible in the lacking of a well understood algebraic foundation of Lee-metric codes. Indeed, only recently it was discovered that Lee-metric codes attain the Gilbert-Varshamov bound with high probability [4], for the length of the code tending to infinity. This aligns with famous and well studied results in the Hamming [3] and the rank metric [25]. In addition, the characterization of constant Lee weight codes, initiated by Wood [35], has only recently been completed in [6].

The study of optimal codes is a classical direction in algebraic coding theory and the most famous bound within this direction is the Singleton bound. The bound gives an upper bound on the minimum distance of a code, given other parameters of the code. Thus, codes attaining this bound have the maximal possible minimum distance and in turn the largest error correction capacity.

In the Hamming metric, the Singleton bound was introduced by Singleton [32] and was already studied by Komamiya [21]. Codes in the Hamming metric that

2020 *Mathematics Subject Classification.* 94B05,94B65.

Key words and phrases. Ring-linear code, Lee distance, Generalized Weights.

are attaining the Singleton bound are called Maximum Distance Separable (MDS) codes and are constituting some of the most used and studied codes in coding theory. It is well known that codes attain the Hamming-metric Singleton bound with high probability when letting the size of the finite field q tend to infinity. Instead, if one lets the length n tend to infinity, the probability for a code to attain the Singleton bound tends to 0.

When changing the underlying metric, this behaviour can drastically change. This is for example the case in the rank metric. While the rank metric is younger than the Lee metric, first being introduced by Delsarte [8] in 1978 and reintroduced by Gabidulin [12] and Roth [29], its Singleton bound was already given in [12] in 1985 and its optimal codes, called Maximum Rank Distance (MRD) codes, have been well studied since. In fact, we know that for \mathbb{F}_{q^m} -linear codes, MRD codes are dense when letting q or m tend to infinity [26]. For \mathbb{F}_q linear codes, however MRD codes are sparse when letting q tend to infinity [17], except for some special cases, where m or n are 2 [2, 5, 13].

The situation in the Lee metric is completely different. Indeed, the first Singleton bound in the Lee metric has only been stated in 2000 by Shiromoto [31]. This bound is tight, as there exists a code attaining it. However, the recent paper [6] revealed that this example is in fact the only non-trivial linear code that is optimal with respect to this Lee-metric Singleton bound. Thus, such optimal codes are "extremely" sparse and show the need of a more thorough study of bounds in the Lee metric and their optimal codes. The used puncturing argument to derive the classical Singleton bound and thus also the Lee-metric alternative in [31] proves to be not suitable for the Lee metric. One thus requires other techniques to derive a Singleton-like bound which appear to be more tailored to the Lee metric setting. In fact, one possible technique is through generalized weights, first introduced in [19].

In this paper we introduce several possible definitions of Lee supports of a code, which allows us to define generalized Lee weights. We compare the resulting Lee-metric Singleton bounds and compute the density of their optimal codes. The main contribution of this paper is a new Lee-metric Singleton bound for which the optimal codes are not sparse.

This paper is structured as follows. In Section 2 we provide the preliminary basics on ring-linear codes and the Lee metric, which will be helpful for the remainder of the paper. We also provide the already known Singleton-like bounds for the Lee metric and the densities of their optimal codes. Section 3 serves as a recap of generalized Hamming weights for finite fields and for finite integer rings. We restate there the main properties and definitions and discuss how to derive bounds on the minimum Lee distance using generalized weights. In Sections 4 and 5 we introduce two new support definitions in the Lee metric deriving from the ideas used in the Hamming metric. For both of them we are able to derive new bounds on the minimum Lee distance defining generalized Lee weights with respect to the new supports. Additionally, we show that the newly proposed generalized Lee weights are invariant under isometries in the Lee metric, and we discuss the density of optimal codes. Even though these new bounds are sharper than the existing bounds on the minimum Lee distance, they are still not sufficiently tight. Therefore, we introduce new generalized weights based on filtrations of a code in Section 6. Together with some additional parameters for a generator matrix of a filtration of a code we are able

to derive an improved bound on the minimum Lee distance. Again, we discuss the density of optimal codes and the invariance under isometries for the new generalized weights. In Section 7 we compare all the bounds for several parameters. Finally, we conclude the paper in Section 8.

2. PRELIMINARIES

We start this section by introducing the main definitions, notions and results used in the course of this paper. Throughout this paper we denote by p a prime number, by s a positive integer, and by $\mathbb{Z}/p^s\mathbb{Z}$ the integer residue ring. Furthermore, for any integer $i \in \{0, \dots, s-1\}$, we write $\langle p^i \rangle$ to denote either the ideal $p^i(\mathbb{Z}/p^s\mathbb{Z})$ or the submodule $p^i(\mathbb{Z}/p^s\mathbb{Z})^n$.

2.1. Ring-Linear Codes.

Classical coding theory considers finite fields \mathbb{F}_q with q elements and a linear code \mathcal{C} is a subspace of the vector space \mathbb{F}_q^n . Thus, \mathcal{C} has a dimension k , which determines its size $|\mathcal{C}| = q^k$ and the minimum number of generators. Such a code \mathcal{C} can then be represented through a generator matrix $G \in \mathbb{F}_q^{k \times n}$ or a parity-check matrix $H \in \mathbb{F}_q^{(n-k) \times n}$, which have the code as image, respectively as kernel.

In this paper, we focus on codes over $\mathbb{Z}/p^s\mathbb{Z}$.

Definition 1. A linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ is a $\mathbb{Z}/p^s\mathbb{Z}$ -submodule of $(\mathbb{Z}/p^s\mathbb{Z})^n$.

Due to the fundamental theorem of finite abelian groups, a linear code is isomorphic to the following direct sum of $\mathbb{Z}/p^s\mathbb{Z}$ -modules

$$\mathcal{C} \cong \bigoplus_{i=1}^K (\mathbb{Z}/p^s\mathbb{Z}) / \langle p \rangle^{\lambda_i}.$$

The type of a module \mathcal{C} is then defined as the partition

$$\lambda = \underbrace{(s, \dots, s)}_{k_0} \underbrace{(s-1, \dots, s-1)}_{k_1} \dots \underbrace{(1, \dots, 1)}_{k_{s-1}}$$

or equivalently $\lambda = (s^{k_0}(s-1)^{k_1} \dots 1^{k_{s-1}})$. Instead of using this well-known notation from the theory of modules, we prefer the notation (k_0, \dots, k_s) , called *subtype* of the code \mathcal{C} , due to its simplicity.

We call n the *length* of the code \mathcal{C} and its elements *codewords*. We can define the $\mathbb{Z}/p^s\mathbb{Z}$ -dimension of a linear code \mathcal{C} as

$$k := \log_{p^s}(|\mathcal{C}|).$$

The *rank* K of a code is given by the number of generators, i.e., $K := \sum_{i=0}^{s-1} k_i$. Additionally, k_0 is called the *free rank* of the code \mathcal{C} . We denote by the $\mathbb{Z}/p^s\mathbb{Z}$ -dimension of the code \mathcal{C} the following

$$k = \log_{p^s}(|\mathcal{C}|).$$

Note that the $\mathbb{Z}/p^s\mathbb{Z}$ -dimension of a code is not necessarily an integer. In fact, k is determined by the subtype as

$$k = \sum_{i=0}^{s-1} \frac{s-i}{s} k_i.$$

In general it holds that $0 \leq k_0 \leq k \leq K \leq n$. If the rank and the $\mathbb{Z}/p^s\mathbb{Z}$ -dimension of a code coincide, we call the code *free*. It has been shown in [4] that free codes are dense as $p \rightarrow \infty$ but they are neither dense nor sparse as the length n or s tend to infinity.

As in the finite field case, ring-linear codes are represented through a generator matrix or a parity-check matrix.

Definition 2. Consider a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K . A matrix $G \in (\mathbb{Z}/p^s\mathbb{Z})^{K \times n}$ is called a generator matrix of \mathcal{C} if the rows of G span the code. A *parity-check matrix* H is an $(n - K) \times n$ matrix over $\mathbb{Z}/p^s\mathbb{Z}$ whose its null-space coincides with \mathcal{C} .

Usually it is helpful to consider these matrices in their systematic form.

Proposition 3. Let \mathcal{C} be a linear code in $(\mathbb{Z}/p^s\mathbb{Z})^n$ of subtype (k_0, \dots, k_{s-1}) and rank K . Then \mathcal{C} is permutation equivalent to a code having a generator matrix $G_{\text{sys}} \in (\mathbb{Z}/p^s\mathbb{Z})^{K \times n}$ of the form

$$(2.1) \quad G_{\text{sys}} = \begin{pmatrix} \mathbb{I}_{k_0} & A_{1,2} & A_{1,3} & \cdots & A_{1,s} & A_{1,s+1} \\ 0 & p\mathbb{I}_{k_1} & pA_{2,3} & \cdots & pA_{2,s} & pA_{2,s+1} \\ 0 & 0 & p^2\mathbb{I}_{k_2} & \cdots & p^2A_{3,s} & p^2A_{3,s+1} \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & p^{s-1}\mathbb{I}_{k_{s-1}} & p^{s-1}A_{s,s+1} \end{pmatrix},$$

where $A_{i,s+1} \in (\mathbb{Z}/p^{s+1-i}\mathbb{Z})^{k_{i-1} \times (n-K)}$, $A_{i,j} \in (\mathbb{Z}/p^{s+1-i}\mathbb{Z})^{k_{i-1} \times k_j}$ for $j \leq s$. In addition, the code \mathcal{C} is permutation equivalent to a code having a parity-check matrix $H \in (\mathbb{Z}/p^s\mathbb{Z})^{(n-k_0) \times n}$ of the form

$$(2.2) \quad H_{\text{sys}} = \begin{pmatrix} B_{1,1} & B_{1,2} & \cdots & B_{1,s-1} & B_{1,s} & \mathbb{I}_{n-K} \\ pB_{2,1} & pB_{2,2} & \cdots & pB_{2,s-1} & p\mathbb{I}_{k_{s-1}} & 0 \\ p^2B_{3,1} & p^2B_{3,2} & \cdots & p^2\mathbb{I}_{k_{s-2}} & 0 & 0 \\ \vdots & \vdots & & \vdots & \vdots & \vdots \\ p^{s-1}B_{s,1} & p^{s-1}\mathbb{I}_{k_1} & \cdots & 0 & 0 & 0 \end{pmatrix},$$

where $B_{1,j} \in (\mathbb{Z}/p^s\mathbb{Z})^{(n-K) \times k_{j+1}}$, $B_{i,j} \in (\mathbb{Z}/p^{s+1-i}\mathbb{Z})^{k_{s-i+1} \times k_{j+1}}$ for $i > 1$.

We call the forms in (2.1) and (2.2) the *systematic form* of a generator matrix and a parity-check matrix, respectively.

Additionally to the subtype of a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$, we can define a similar parameter going over the columns of a generator matrix of \mathcal{C} .

Definition 4. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K . For each $j \in \{1, \dots, n\}$ consider the j -th coordinate map

$$\pi_j : \begin{array}{ccc} (\mathbb{Z}/p^s\mathbb{Z})^n & \longrightarrow & \mathbb{Z}/p^s\mathbb{Z} \\ (c_1, \dots, c_n) & \longmapsto & c_j \end{array}.$$

The *support subtype* of \mathcal{C} is defined to be an $(s+1)$ -tuple $(n_0(\mathcal{C}), \dots, n_s(\mathcal{C}))$, where $n_i(\mathcal{C})$ counts the number coordinates $j \in \{1, \dots, n\}$ belonging to ideal $\langle p^i \rangle$, i.e.,

$$n_i(\mathcal{C}) := |\{j \in \{1, \dots, n\} \mid \langle \pi_j(\mathcal{C}) \rangle = \langle p^i \rangle\}|.$$

A code with $n_s(\mathcal{C}) = 0$ is called *non-degenerate*.

We will simply write n_i instead of $n_i(\mathcal{C})$ if the code \mathcal{C} is clear from the context.

2.2. Lee Metric.

In this paper we will focus on the Lee metric introduced in [24]. This metric was introduced to cope with phase modulation in communications. However, we will often refer and compare the Lee metric to the Hamming metric.

Thus, recall that for two n -tuples $x, y \in (\mathbb{Z}/p^s\mathbb{Z})^n$ their *Hamming distance* is defined to be the number of positions where they differ, i.e.,

$$d_H(x, y) = |\{i \in \{1, \dots, n\} \mid x_i \neq y_i\}|.$$

For a single element $x \in (\mathbb{Z}/p^s\mathbb{Z})^n$ its *Hamming weight* is the Hamming difference to zero, i.e., the number of nonzero entries of x , and we denote it by $\text{wt}_H(x) := d_H(x, 0)$. The *minimum Hamming distance* of a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ is then defined as

$$d_H(\mathcal{C}) := \min \{\text{wt}_H(x) \mid x \in \mathcal{C} \setminus \{0\}\}.$$

Let us now introduce the Lee metric.

Definition 5. Given an integer residue ring $\mathbb{Z}/p^s\mathbb{Z}$ and consider an element $a \in \mathbb{Z}/p^s\mathbb{Z}$ interpreted as an integer in $\{0, \dots, p^s - 1\}$. The *Lee weight* $a \in \mathbb{Z}/p^s\mathbb{Z}$ is given by

$$\text{wt}_L(a) := \min \{a, |p^s - a|\}.$$

For $x \in (\mathbb{Z}/p^s\mathbb{Z})^n$ its Lee weight is defined in an additive fashion. That is,

$$\text{wt}_L(x) = \sum_{i=1}^n \text{wt}_L(x_i).$$

Similarly to the Hamming metric, the Lee weight induces a distance which we refer to as *Lee distance*. In fact, for $x, y \in (\mathbb{Z}/p^s\mathbb{Z})^n$ their Lee distance is defined to be

$$d_L(x, y) = \text{wt}_L(x - y).$$

Note that the Hamming weight of an element $x \in (\mathbb{Z}/p^s\mathbb{Z})^n$ is a natural lower bound on its Lee weight. Additionally, the Lee weight of each entry x_i can never exceed $M := \lfloor p^s/2 \rfloor$. Hence, we have

$$0 \leq \text{wt}_H(x) \leq \text{wt}_L(x) \leq \text{wt}_H(x)M \leq nM.$$

Considering the Lee metric for linear codes \mathcal{C} over $\mathbb{Z}/p^s\mathbb{Z}$, we are able to introduce the minimum Lee distance of \mathcal{C} .

Definition 6. Given a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$, we define its *minimum Lee distance* as

$$d_L(\mathcal{C}) := \min \{\text{wt}_L(c) \mid c \in \mathcal{C} \setminus \{0\}\}.$$

Again, we easily observe that

$$d_H(\mathcal{C}) \leq d_L(\mathcal{C}) \leq M d_H(\mathcal{C}).$$

The linear isometries of the Lee metric are $\{-1, 1\} \times S_n$, that is we can permute the entries and multiply each entry with either 1 or -1 .

2.3. Singleton-like Bounds over $\mathbb{Z}/p^s\mathbb{Z}$.

One of the major research directions in coding theory is to bound the minimum distance of a code. This will then, in turn, bound the error-correction capability of a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$. In fact, the higher the minimum distance of \mathcal{C} , the more errors can be corrected.

The task of bounding the minimum distance, clearly depends on the metric used to endow the ambient space. For the Hamming metric over finite fields, the best-known bound is the Singleton bound describing the trade-off between the minimum Hamming distance of a code and its dimension.

Theorem 7 (Singleton Bound). *Given a linear code $\mathcal{C} \subseteq \mathbb{F}_q^n$ of dimension k over \mathbb{F}_q , its minimum Hamming distance is upper bounded by*

$$d_H(\mathcal{C}) \leq n - k + 1.$$

Codes achieving this bound are called *maximum distance separable* (MDS). It is well known that for q tending to infinity a random linear code will attain the Singleton bound. Hence, MDS codes are dense for q tending to infinity with high probability. On the other hand, if we let the length n grow, MDS codes are sparse. Note that this also follows immediately from the famous MDS conjecture [30], which states that if q is odd, then an MDS code must have $n \leq q + 1$. In the case where $q = 2^s$, and $k = 3$ or $k = q - 1$, in which case $n \leq q + 2$. A very analogous bound can be established over finite integer residue rings $\mathbb{Z}/p^s\mathbb{Z}$ for the Hamming metric.

Proposition 8. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of $\mathbb{Z}/p^s\mathbb{Z}$ -dimension k , then*

$$d_H(\mathcal{C}) \leq n - k + 1.$$

This Singleton-like bound does hold for non-linear codes as well, stating that $|\mathcal{C}| \leq q^{n-d_H(\mathcal{C})+1}$. For linear codes only, this bound has been further tightened:

Proposition 9 ([9, 11]). *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K , then*

$$d_H(\mathcal{C}) \leq n - K + 1.$$

To differentiate codes achieving the bound in Proposition 9 from MDS codes, we will refer to the first as *maximum distance codes with respect to the rank* (or MDR codes for short). However, MDS codes and MDR codes are closely related. In fact, any linear MDS code is always an MDR code. Vice versa we can say that a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ is MDR if and only if the socle $\mathcal{C} \cap \langle p^{s-1} \rangle$ can be identified with an MDS over \mathbb{F}_p . We can therefore directly apply the results from finite fields which means that by the MDS conjecture, MDR codes are sparse as n grows large and, as usual, they are dense as p grows large.

Note that the relation

$$d_H(\mathcal{C}) \leq d_L(\mathcal{C}) \leq M d_H(\mathcal{C})$$

immediately gives rise to a bound for codes in the Lee metric, i.e.,

$$d_L(\mathcal{C}) \leq M(n - \lfloor k \rfloor + 1).$$

Apart from this obvious bound, the first Lee-metric Singleton-like bound was introduced by Shiromoto in 2000 [31].

Theorem 10 ([31]). *Consider a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of $\mathbb{Z}/p^s\mathbb{Z}$ -dimension k . Then the following bound holds*

$$\left\lfloor \frac{d_L(\mathcal{C}) - 1}{M} \right\rfloor \leq n - k.$$

Note that this bound implies $d_L(\mathcal{C}) \leq M(n - k) + \alpha$ for some $\alpha \in \{1, \dots, M\}$, and can be generalized easily to $d_L(\mathcal{C}) \leq M(n - \lfloor k \rfloor) + \alpha$, again for some $\alpha \in \{1, \dots, M\}$.

The bound from Theorem 10 can be derived from the simple fact that $d_L(\mathcal{C}) \leq M d_H(\mathcal{C})$ for a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ or by a puncturing argument, as in the classical case. That is, given a code of size $|\mathcal{C}|$ and minimum Lee distance $d_L(\mathcal{C})$, we can puncture the code in only $\left\lfloor \frac{d_L(\mathcal{C}) - 1}{M} \right\rfloor$ coordinates, to make sure that the punctured code \mathcal{C}' still has size $|\mathcal{C}|$. In fact, every pair of codewords in \mathcal{C} has Lee distance at least $d_L(\mathcal{C})$, by puncturing in one position, one has to assume that their distance decreased by the maximal possible value, i.e., by M . The claim then follows easily as

$$\mathcal{C}' \subseteq (\mathbb{Z}/p^s\mathbb{Z})^{n - \left\lfloor \frac{d_L(\mathcal{C}) - 1}{M} \right\rfloor}.$$

Shiromoto also provided a code which attains the bound, showing its tightness.

Example 11. Let $\mathcal{C} = \langle 1, 2 \rangle \subseteq (\mathbb{Z}/5\mathbb{Z})^2$. Then, since $d_L(\mathcal{C}) = 3$, this code attains the bound of Theorem 10, as

$$\left\lfloor \frac{d_L(\mathcal{C}) - 1}{M} \right\rfloor = \frac{3 - 1}{2} = 1 = n - k = 2 - 1.$$

However, in [6], it was observed that this is actually the only non-trivial linear code that attains the bound in Theorem 10. The Lee-metric Singleton bound can be further improved, as shown in [6], e.g. by employing the rank K .

Corollary 12. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K , then*

$$d_L(\mathcal{C}) \leq M(n - K + 1).$$

Alderson and Huntemann in [1] provided a similar bound to Corollary 12 by restricting k to a positive integer bounded by n .

Theorem 13 ([1]). *For any code \mathcal{C} in $(\mathbb{Z}/p^s\mathbb{Z})^n$ of $\mathbb{Z}/p^s\mathbb{Z}$ -dimension k a positive integer, $1 < k < n$ we have that*

$$d_L(\mathcal{C}) \leq M(n - k).$$

However, in [6], the authors characterized all codes attaining the above Lee-metric Singleton bounds, with the result that their optimal codes are sparse in both cases, i.e., when n, p or $s \rightarrow \infty$.

Using the newly introduced parameter for the code, i.e., the support subtype, one can easily derive an improved Lee-metric Singleton bound from the puncturing argument.

Theorem 14. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K and support subtype $(n_0, \dots, n_{s-1}, 0)$. Define for all $i \in \{0, \dots, s\}$*

$$M_i = \left\lfloor \frac{p^{s-i}}{2} \right\rfloor p^i, \quad B_j = \sum_{i=j}^{s-1} n_i, \quad A_j = \sum_{i=j}^{s-1} n_i M_i.$$

Let $j \in \{1, \dots, s-1\}$ be the smallest positive integer such that $A_j < d_L(\mathcal{C})$, then

$$K \leq n - B_j - \left\lfloor \frac{d_L(\mathcal{C}) - A_j - 1}{M_{j-1}} \right\rfloor.$$

Proof. We start by puncturing the code in the positions of smallest possible Lee weight. To identify these positions, we use the support subtype. Clearly, in the ideal $\langle p^i \rangle$, we have as largest possible Lee weight $M_i = \lfloor \frac{p^{s-1}}{2} \rfloor p^i$, and thus we would start puncturing in the positions, where all codewords live in $\langle p^{s-1} \rangle$, i.e., in the positions belonging to the support subtype n_{s-1} . We hence assume that the minimum distance between two distinct tuples decreased by $A_{s-1} = n_{s-1}M_{s-1}$. If this is still smaller than the minimum Lee distance, we can continue puncturing in the next ideal, namely $\langle p^{s-2} \rangle$. We continue in this fashion, every time puncturing in $n_i M_i$ positions, until $A_j = \sum_{i=j}^{s-1} n_i M_i$ has reached the minimum Lee distance. We are left with codewords that are at least $d_L(\mathcal{C}) - A_j$ apart, thus we can continue puncturing in $\left\lfloor \frac{d_L(\mathcal{C}) - A_j - 1}{M_{j-1}} \right\rfloor$ positions living in $\langle p^{j-1} \rangle$, i.e., belonging to the support subtype n_{j-1} , and still be sure that the punctured code has the same size as the original code. In this case, we have the new length of the punctured code, being $n - B_j - \left\lfloor \frac{d_L(\mathcal{C}) - A_j - 1}{M_{j-1}} \right\rfloor$, for $B_j = \sum_{i=j}^{s-1} n_i$. \square

Example 15. Let us consider $\mathcal{C} \subseteq (\mathbb{Z}/9\mathbb{Z})^4$ generated by

$$G = \begin{pmatrix} 1 & 0 & 2 & 3 \\ 0 & 3 & 6 & 0 \\ 0 & 0 & 3 & 6 \end{pmatrix}.$$

The $\mathbb{Z}/9\mathbb{Z}$ -dimension of this code is $k = 2$. The minimum Lee distance of this code is $d_L(\mathcal{C}) = 6$. The code also has support subtype $(2, 2, 0)$. Thus, we would identify $j = s = 2$, as we cannot puncture in both positions belonging to $n_1 = 2$, namely the second and the last column, as we would get $n_1 M_1 = 6 \not< d_L(\mathcal{C})$. However, we can puncture in one of these two columns. In fact, $\left\lfloor \frac{d_L(\mathcal{C}) - 0 - 1}{3} \right\rfloor = 1$. That is, the bound in Theorem 14 is attained as

$$K = 3 = 4 - 0 - \left\lfloor \frac{6 - 0 - 1}{3} \right\rfloor = n - B_j - \left\lfloor \frac{d_L(\mathcal{C}) - A_j - 1}{M_{j-1}} \right\rfloor.$$

The bound from Theorem 10 would instead give

$$\left\lfloor \frac{d_L(\mathcal{C}) - 1}{M} \right\rfloor = \left\lfloor \frac{6 - 1}{4} \right\rfloor = 1 < 2 = n - k.$$

Since we are also in the case where k is an integer strictly larger than 1, we can also apply the bound from Theorem 13, and get

$$d_L(\mathcal{C}) = 6 < 8 = (4 - 2) \cdot 4 = (n - k)M.$$

We can rewrite the bound from Theorem 14 as upper bound on the minimum Lee distance as, for j the smallest positive integer with $A_j < d_L(\mathcal{C})$ we have

$$d_L(\mathcal{C}) \leq M_{j-1} \left(\sum_{i=0}^{j-1} n_i - K \right) + \sum_{i=j}^{s-1} n_i M_i + \alpha$$

for some $\alpha \in \{1, \dots, M_{j-1}\}$. However, the condition to find the smallest j such that $A_j < d_L(\mathcal{C})$, renders the bound impractical, as usually one does not know the minimum Lee distance of the code and thus wants to bound it.

3. GENERALIZED HAMMING WEIGHTS

In this section we propose a new definition of generalized Lee weights. We base the definition on the Hamming weight counterparts. Generalized Hamming weights have originally been introduced in [19] and were then rediscovered by Wei in [34]. Generalized weights in the Hamming metric have been studied in various areas [7, 10, 15, 16, 27]. In [14] the authors defined and the generalized Hamming weights of ring-linear codes by considering the join-Hamming support of a code.

Let us recap the definition of a Hamming support of a vector and a code. For this, consider a finite field \mathbb{F}_q of q elements and a positive integer n . The *Hamming support* of $x \in \mathbb{F}_q^n$ is defined to be the set of indices where x is nonzero, i.e.,

$$\text{supp}_H(x) := \{i \in \{1, \dots, n\} \mid x_i \neq 0\}.$$

Note here, that the cardinality of the support of x corresponds to the Hamming weight of x , that is

$$(3.1) \quad |\text{supp}_H(x)| = \text{wt}_H(x).$$

Let us consider now a linear code $\mathcal{C} \subset \mathbb{F}_q^n$ of length n and dimension K . For the definition of the Hamming support of \mathcal{C} we take into account every codeword $c \in \mathcal{C}$ and for each index, we figure out whether a codeword exists which is nonzero in this position, i.e.,

$$(3.2) \quad \text{supp}_H(\mathcal{C}) = \{i \in \{1, \dots, n\} \mid \exists c \in \mathcal{C}, c_i \neq 0\}.$$

Analogously to the definition of the weight of a vector x in (3.1), we can define the weight of a code \mathcal{C} to be the size of its support, i.e.,

$$(3.3) \quad \text{wt}(\mathcal{C}) = |\text{supp}(\mathcal{C})|.$$

The goal is to generalize these notions to other metrics and ambient spaces. In particular, we are interested in the Lee metric defined over rings. In order to do so, we will follow the approach of [14]. Let \mathcal{R} denote a finite unitary ring. Let us consider a weight function

$$\text{wt} : \mathcal{R} \rightarrow \mathbb{N},$$

which is such that

1. $\text{wt}(0) = 0$ and $\text{wt}(x) > 0$ for all $x \neq 0$,
2. $\text{wt}(x) = \text{wt}(-x)$,
3. $\text{wt}(x + y) \leq \text{wt}(x) + \text{wt}(y)$.

Note the difference to the definition used in [14, Definition 2.4] which corresponds to a norm, as they also require the absolute homogeneity property. That is, for any $\lambda \in \mathcal{R} \setminus \{0\}$, we have that

4. $\text{wt}(\lambda x) = \text{wt}(x)$.

This property holds for the Hamming weight and many other weights, however it is not a requirement for a weight function. The properties 1.-3. are enough to

induce a distance. In fact, for a weight function wt with properties 1.-3. we can define a distance as

$$\begin{aligned} d : \mathcal{R} \times \mathcal{R} &\rightarrow \mathbb{N}, \\ (x, y) &\mapsto \text{wt}(x - y). \end{aligned}$$

By abuse of notation, we will also denote their coordinate-wise extension by wt and d , that is for $x, y \in \mathcal{R}^n$ we have

$$\text{wt}(x) = \sum_{i=1}^n \text{wt}(x_i) \text{ and } d(x, y) = \sum_{i=1}^n d(x_i, y_i).$$

We call such weight functions *additive weights*. Given a weight function, one can then define the *support* of $x \in \mathcal{R}^n$ as an n -tuple

$$\text{supp}(x) := (\text{wt}(x_1), \dots, \text{wt}(x_n)).$$

As we are now dealing with an n -tuple instead of a subset of $\{1, \dots, n\}$, we require some additional definitions. Let $s, t \in \mathbb{N}^n$ be n -tuples. The size of s is given by the sum of its entries, that is $|s| = \sum_{i=1}^n s_i$. The *join* of s and t , denoted by $s \vee t$ is given by taking the maximum in each position, that is

$$s \vee t = (\max\{s_1, t_1\}, \dots, \max\{s_n, t_n\}).$$

The *meet* of s and t , denoted by $s \wedge t$ is given by taking the minimum in each position, that is

$$s \wedge t = (\min\{s_1, t_1\}, \dots, \min\{s_n, t_n\}).$$

Since the weight is additive, we have that

$$|\text{supp}(x)| = \sum_{i=1}^n \text{wt}(x_i) = \text{wt}(x).$$

The Hamming support is thus such an example, where instead of considering the support as subset of $\{1, \dots, n\}$, the support is considered as the n -tuple

$$\text{supp}_{\text{H}}(x) = (\text{wt}_{\text{H}}(x_1), \dots, \text{wt}_{\text{H}}(x_n)).$$

In order to extend this to the support of codes, we have several options. One of those, is the join-support, as considered in [14]: for $\mathcal{C} \subseteq \mathcal{R}^n$ a linear code, we define its *join-support* as

$$\text{supp}_{\text{join}}(\mathcal{C}) := \left(\max_{c \in \mathcal{C}} \text{wt}(c_1), \dots, \max_{c \in \mathcal{C}} \text{wt}(c_n) \right) = \bigvee_{c \in \mathcal{C}} \text{supp}(c).$$

Note that another possibility would be to define the *meet-support*, as follows

$$\begin{aligned} \text{supp}_{\text{meet}}(\mathcal{C}) &:= \left(\min_{c \in \mathcal{C}} \{\max\{\text{wt}(c_1), 0\}\}, \dots, \min_{c \in \mathcal{C}} \{\max\{\text{wt}(c_n), 0\}\} \right) \\ &= \bigwedge_{c \in \mathcal{C}} (\text{supp}(c) \vee 0). \end{aligned}$$

As the Hamming weight of nonzero elements equals one, we observe that the join-support coincides with the meet-support of a code \mathcal{C} in the Hamming metric, i.e.,

$$\text{supp}_{\text{H,join}}(\mathcal{C}) = \text{supp}_{\text{H,meet}}(\mathcal{C}).$$

Example 16. Let us consider the code over \mathbb{F}_3 generated by

$$G = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}.$$

With the usual definition of the Hamming support in (3.2), we have that

$$\text{supp}_H(\mathcal{C}) = \{1, 2, 3, 4\}.$$

With the join-support, we are considering the maximal value of the weight of the entries of a codeword in each position, that is

$$\text{supp}_{H,\text{join}}(\mathcal{C}) = (1, 1, 1, 1, 0).$$

For the meet-support, we take the minimum nonzero value of the weight of the entries of a codeword in each position, which also gives $(1, 1, 1, 1, 0)$.

By applying definition of the weight of a code (3.3) we observe that all the three support definitions of \mathcal{C} yield the same weight

$$\begin{aligned} \text{wt}_H(\mathcal{C}) &= |\text{supp}_H(\mathcal{C})| = 4, \\ \text{wt}_{H,\text{join}}(\mathcal{C}) &= \text{wt}_{H,\text{meet}}(\mathcal{C}) = 4. \end{aligned}$$

In the classical case, one defines the r -th generalized weights as follows.

Definition 17. Let $\mathcal{C} \subseteq \mathbb{F}_q^n$ be a linear code of dimension k . Then for any $r \in \{1, \dots, k\}$ the r -th generalized weight is given by

$$d^r(\mathcal{C}) = \min\{\text{wt}(\mathcal{D}) \mid \mathcal{D} \leq \mathcal{C}, \dim(\mathcal{D}) = r\}.$$

For the generalized weights, we want the following properties to hold. Let \mathcal{C} be a linear code of dimension k . Then we have

1. $d(\mathcal{C}) = d^1(\mathcal{C})$,
2. $d^r(\mathcal{C}) < d^{r+1}(\mathcal{C})$ for every $1 \leq r < k$,
3. $d^k(\mathcal{C}) = \text{wt}(\mathcal{C})$.

For the Hamming support and the rank support these properties have been showed in [34, 22, 23]. In [14] they get similar properties, with the exception of $d^r(\mathcal{C}) \leq d^{r+1}(\mathcal{C})$, instead of the strict inequality.

The strict inequality is important to us, however, as it then leads to neat Singleton bounds, that is as

$$d(\mathcal{C}) = d^1(\mathcal{C}) < d^2(\mathcal{C}) < \dots < d^{k-1}(\mathcal{C}) < d^k(\mathcal{C}) = \text{wt}(\mathcal{C}),$$

we get

$$d(\mathcal{C}) \leq \text{wt}(\mathcal{C}) - k + 1.$$

Note that for non-degenerate codes we have that $\text{wt}_H(\mathcal{C}) = n$, and thus we retrieve the classical Singleton bound

$$d_H(\mathcal{C}) \leq n - k + 1.$$

Since we move from the classical case of finite fields to rings, we have to exchange the fixed dimension of the subcodes with a ring-analogue parameter. A natural choice would be the $\mathbb{Z}/p^s\mathbb{Z}$ -dimension, but as this value is not necessarily an integer and there might not exist subcodes of \mathcal{C} of certain fixed smaller rational number as the $\mathbb{Z}/p^s\mathbb{Z}$ -dimension, we choose to discard this option.

In [10], the authors chose to exchange the dimension with the subtype. In fact, in the same paper the authors defined generalized Lee weights for $\mathbb{Z}/4\mathbb{Z}$. This particular case is, however, not of interest for us, as the Lee-metric Singleton bound over $\mathbb{Z}/4\mathbb{Z}$ directly follows from the Gray isometry.

Following the idea of [10], a first attempt on defining generalized weights over $\mathbb{Z}/p^s\mathbb{Z}$ would be the following.

Definition 18. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of subtype (k_0, \dots, k_{s-1}) . Then for any (r_0, \dots, r_{s-1}) with $r_i \leq k_i$ for all $i \in \{1, \dots, s-1\}$ the (r_0, \dots, r_{s-1}) -th generalized weight is given by

$$d^{(r_0, \dots, r_{s-1})}(\mathcal{C}) = \min\{\text{wt}(\mathcal{D}) \mid \mathcal{D} \leq \mathcal{C}, \mathcal{D} \text{ has subtype } (r_0, \dots, r_{s-1})\}.$$

Note that this definition is not considering all possible subcodes or all possible subtypes of subcodes.

To allow for a comparison between two different subtypes (r_0, \dots, r_{s-1}) and (r'_0, \dots, r'_{s-1}) which might have $r_i < r'_i$ for some i but $r_j > r'_j$ for some j , a natural choice is to impose a lexicographical order, i.e., we consider the order

$$(k_0, \dots, k_{s-1}) > (k_0 - 1, \dots, k_{s-1}) > \dots > (0, k_1, \dots, k_{s-1}) > \dots > (0, \dots, 0, 1).$$

However, then the property $d(\mathcal{C}) = d^{(0, \dots, 0, 1)}(\mathcal{C})$ is not guaranteed.

In fact, a minimum Lee weight codeword will live in a subcode having subtype one of the standard vectors e_i . Thus, we have $d(\mathcal{C}) = d^{e_i}(\mathcal{C})$ for some i . Observing that this just means to fix the rank of the subcode as 1, we choose to directly fix the rank instead.

Definition 19. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K . Then for any $r \in \{1, \dots, K\}$ the r -th generalized weight is given by

$$d^{(r)}(\mathcal{C}) = \min\{\text{wt}(\mathcal{D}) \mid \mathcal{D} \leq \mathcal{C}, \text{rk}(\mathcal{D}) = r\}.$$

4. JOIN-SUPPORT IN THE LEE METRIC

Now we turn our focus on exchanging the Hamming weight with the Lee weight. We want to define the Lee support and hence the generalized Lee weights in a similar fashion. For the Lee support of a vector $x \in (\mathbb{Z}/p^s\mathbb{Z})^n$ we view the Lee support as an n -tuple and define it analogous to the Hamming support, i.e.,

$$\text{supp}_L(x) := (\text{wt}_L(x_1), \dots, \text{wt}_L(x_n)).$$

We now want to define the Lee support and the generalized Lee weights of a code $\mathcal{C} \subseteq \mathbb{Z}/p^s\mathbb{Z}^n$, according to the two possibilities: the join-support and the meet-support. However, in the Lee metric, the meet-support is not practical to derive bounds on the minimum distance for the code. Let us quickly argue why.

Definition 20. For a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ we define the *Lee meet-support* as the minimal (if possible) nonzero Lee weight in each position among all codewords, meaning that

$$\text{supp}_{L, \text{meet}}(\mathcal{C}) := \left(\min_{c \in \mathcal{C}} \{\max\{\text{wt}_L(c_1), 0\}\}, \dots, \min_{c \in \mathcal{C}} \{\max\{\text{wt}_L(c_n), 0\}\} \right).$$

Proposition 21. For $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of support subtype (n_0, \dots, n_s) , we have that

$$|\text{supp}_{\text{L,meet}}(\mathcal{C})| = \text{wt}_{\text{L}}(\mathcal{C}) = \sum_{i=0}^s n_i p^i.$$

Proof. The Lee meet-support asks to take the smallest nonzero Lee weight in position j and then to sum over all entries $j \in \{1, \dots, n\}$. Since any position belonging to the support subtype n_i is living in the ideal $\langle p^i \rangle$, this position has as smallest nonzero Lee weight p^i . \square

We can then define the r -th generalized meet-Lee weights.

Definition 22. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K . For $i \in \{1, \dots, K\}$ define the i th generalized meet-Lee weight as

$$d_{\text{L,meet}}^i(\mathcal{C}) = \min\{|\text{supp}_{\text{L,meet}}(\mathcal{D})| \mid \mathcal{D} \leq \mathcal{C}, \text{rk}(\mathcal{D}) = i\}.$$

Unfortunately, the meet-support in the Lee metric does not generally fulfill the property

$$d_{\text{L}}(\mathcal{C}) \leq d_{\text{L,meet}}^1(\mathcal{C}).$$

As an easy example for $d_{\text{L}}(\mathcal{C}) > d_{\text{L,meet}}^1(\mathcal{C})$, consider $\mathcal{C} = \langle (1, 2) \rangle \subseteq \mathbb{Z}/9\mathbb{Z}^n$. The minimum Lee distance of this code is 3, however, the first generalized meet-Lee weight is 2, as $\text{supp}_{\text{L,meet}}\langle (1, 2) \rangle = (1, 1)$ is the minimal meet-Lee support. This will then not lead to a Singleton bound and is thus discarded.

Instead we will now focus on the join-Lee support, as also promoted in [14].

Definition 23. For a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ its *join-Lee support* is defined as the maximal possible Lee weight in each position among all codewords, i.e.,

$$\text{supp}_{\text{L,join}}(\mathcal{C}) := \left(\max_{c \in \mathcal{C}} \text{wt}_{\text{L}}(c_1), \dots, \max_{c \in \mathcal{C}} \text{wt}_{\text{L}}(c_n) \right).$$

Proposition 24. For $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of support subtype (n_0, \dots, n_s) , we have that

$$|\text{supp}_{\text{L,join}}(\mathcal{C})| = \text{wt}_{\text{L,join}}(\mathcal{C}) = \sum_{i=0}^s n_i M_i.$$

Proof. In each index $j \in \{1, \dots, n\}$, we can check in which ideal this coordinate of the code lives. Let us assume that this is $\langle p^i \rangle$, for some $i \in \{0, \dots, s\}$. Since the support of the code takes the maximum over all codewords in the code, we will reach in this entry the maximal Lee weight of the ideal $\langle p^i \rangle$, which is given by $M_i = \lfloor \frac{p^{s-i}}{2} \rfloor p^i$. Since we know the support subtype of the code, we have n_i many of this entries. \square

The r -th generalized join-Lee weight is then defined as follows.

Definition 25. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K . For $i \in \{1, \dots, K\}$ we define the i th generalized join-Lee weight as

$$d_{\text{L,join}}^r(\mathcal{C}) = \min\{\text{wt}_{\text{L,join}}(\mathcal{D}) \mid \mathcal{D} \leq \mathcal{C}, \text{rk}(\mathcal{D}) = r\}.$$

Let us consider an example, which also perfectly shows the differences between the meet-Lee support and the join-Lee support.

Example 26. Let us consider the code $\mathcal{C} \subseteq \mathbb{Z}/9\mathbb{Z}^4$ generated by

$$G = \begin{pmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 3 & 3 \end{pmatrix},$$

which has support subtype $(4, 0, 0)$ and minimum Lee distance 2, for example $(1, 0, 0, 8)$ is a minimal Lee weight codeword. For the generalized meet-Lee weights we have that

$$d_{\mathcal{L}}(\mathcal{C}) \geq d_{\mathcal{L},\text{meet}}^1(\mathcal{C}) \leq d_{\mathcal{L},\text{meet}}^2(\mathcal{C}) = d_{\mathcal{L},\text{meet}}^3(\mathcal{C}) = \text{wt}_{\mathcal{L},\text{meet}}(\mathcal{C}).$$

Since

$$\begin{aligned} d_{\mathcal{L},\text{meet}}^1(\mathcal{C}) &= \text{wt}_{\mathcal{L},\text{meet}}(\langle(0, 1, 2, 0)\rangle) = 2 \\ d_{\mathcal{L},\text{meet}}^2(\mathcal{C}) &= \text{wt}_{\mathcal{L},\text{meet}}(\langle(1, 0, 3, 2), (0, 1, 2, 0)\rangle) = 4 \\ d_{\mathcal{L},\text{meet}}^3(\mathcal{C}) &= \text{wt}_{\mathcal{L},\text{meet}}(\langle G \rangle) = 4 = \text{wt}_{\mathcal{L},\text{meet}}(\mathcal{C}). \end{aligned}$$

For the generalized join-Lee weights we have that

$$d_{\mathcal{L}}(\mathcal{C}) \leq d_{\mathcal{L},\text{join}}^1(\mathcal{C}) < d_{\mathcal{L},\text{join}}^2(\mathcal{C}) < d_{\mathcal{L},\text{join}}^3(\mathcal{C}) \leq \text{wt}_{\mathcal{L},\text{join}}(\mathcal{C}).$$

Since

$$\begin{aligned} d_{\mathcal{L},\text{join}}^1(\mathcal{C}) &= \text{wt}_{\mathcal{L},\text{join}}(\langle(0, 0, 3, 3)\rangle) = 6 \\ d_{\mathcal{L},\text{join}}^2(\mathcal{C}) &= \text{wt}_{\mathcal{L},\text{join}}(\langle(0, 0, 3, 3), (3, 0, 0, 6)\rangle) = 9 \\ d_{\mathcal{L},\text{join}}^3(\mathcal{C}) &= \text{wt}_{\mathcal{L},\text{join}}(\mathcal{C} \cap \langle 3 \rangle) = 12 \\ \text{wt}_{\mathcal{L},\text{join}}(\mathcal{C}) &= 16. \end{aligned}$$

Proposition 27. *The subcodes which attain the r -th generalized join-Lee weights all live in the socle.*

Proof. By contradiction, assume that $\mathcal{D} \leq \mathcal{C}$ of rank r achieves the r -th generalized Lee weight $d_{\mathcal{L},\text{join}}^r(\mathcal{C})$ and \mathcal{D} does not live in the socle. That is, if \mathcal{D} has support subtype (n_0, \dots, n_s) , then for some $i < s - 1$ we have $n_i \neq 0$. Thus,

$$d_{\mathcal{L},\text{join}}^r(\mathcal{C}) = |\text{supp}_{\mathcal{L},\text{join}}(\mathcal{D})| \leq \sum_{i=1}^{s-1} n_i M_i.$$

By considering the subcode $\mathcal{D}_0 = \mathcal{D} \cap \langle p^{s-1} \rangle$, we observe that its support subtype is $(0, \dots, 0, n_0 + \dots + n_{s-1}, n_s)$. Furthermore,

$$\text{wt}_{\mathcal{L},\text{join}}(\mathcal{D}_0) = M_{s-1}(n_0 + \dots + n_{s-1}) < \sum_{i=1}^{s-1} n_i M_i,$$

since $M_{s-1} < M_i$ for all $i < s - 1$. This gives a contradiction to the minimality of the subcode \mathcal{D} . \square

Thus, it is enough to only consider the generalized join-Lee weights of the socle $\mathcal{C} \cap \langle p^{s-1} \rangle$.

Corollary 28. *Let $\mathcal{C} \subseteq \mathbb{Z}/p^s\mathbb{Z}^n$ be a linear code of rank K . Then for all $r \in \{1, \dots, K\}$ we have*

$$d_{\mathcal{L},\text{join}}^r(\mathcal{C}) = d_{\mathcal{L},\text{join}}^r(\mathcal{C} \cap \langle p^{s-1} \rangle).$$

This property gives us an immediate relation to the generalized Hamming weights. In fact, the socle can be considered as a code over \mathbb{F}_p and the subcodes which attain the minimal join-Lee support are then those which attain the minimal Hamming support.

Corollary 29. *Let $\mathcal{C} \subseteq \mathbb{Z}/p^s\mathbb{Z}^n$ be a linear code of rank K . Then for all $r \in \{1, \dots, K\}$ we have*

$$d_{\text{L,join}}^r(\mathcal{C}) = d_{\text{H}}^r(\mathcal{C})M_{s-1}.$$

Thus, we can use the properties of the generalized Hamming weights to show the following.

Proposition 30. *Let $\mathcal{C} \subset (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K . Then we have*

1. $d_{\text{L}}(\mathcal{C}) \leq d_{\text{L,join}}^1(\mathcal{C})$.
2. $d_{\text{L,join}}^r(\mathcal{C}) < d_{\text{L,join}}^{r+1}(\mathcal{C})$ for every $1 \leq r < K$.
3. $d_{\text{L,join}}^K(\mathcal{C}) \leq \text{wt}_{\text{L,join}}(\mathcal{C})$.

Proof. The first property follows easily from the definition of the join-Lee support of a vector x . It can be tight, whenever the minimal Lee weight codeword is in the socle, which is not necessary. For the second property we simply use Corollary 29 and the third property also simply follows from the definition of join-Lee support. \square

In fact, we do not recover the exact properties of the generalized Hamming weight codes. We do not have $d_{\text{L}}(\mathcal{C}) = d_{\text{L,join}}^1(\mathcal{C})$ and $\text{wt}_{\text{L,join}}(\mathcal{C}) = d_{\text{L,join}}^K(\mathcal{C})$. This seems to be the price we have to pay in order to drop the absolute homogeneity property and to be able to consider the Lee metric. However, unlike the meet-Lee support we get a nice chain of inequalities:

$$d_{\text{L}}(\mathcal{C}) \leq d_{\text{L,join}}^1(\mathcal{C}) < d_{\text{L,join}}^2(\mathcal{C}) < \dots < d_{\text{L,join}}^K(\mathcal{C}) \leq \text{wt}_{\text{L,join}}(\mathcal{C}).$$

This gives us a new Lee-metric Singleton bound.

Theorem 31. *Let $\mathcal{C} \subset (\mathbb{Z}/p^s\mathbb{Z})^n$ be a (non-degenerate) linear code of rank K . Then we have*

$$d_{\text{L}}(\mathcal{C}) \leq M_{s-1}(n - K + 1) = \left\lfloor \frac{p}{2} \right\rfloor p^{s-1}(n - K + 1).$$

Proof. Using the properties 1.-3. from Proposition 30 we know that

$$d_{\text{L}}(\mathcal{C}) \leq d_{\text{L,join}}^K(\mathcal{C}) - \sum_{i=1}^{K-1} x_i,$$

where

$$x_i = d_{\text{L,join}}^i(\mathcal{C}) - d_{\text{L,join}}^{i-1}(\mathcal{C}).$$

From Corollary 29 we know that

$$x_i = d_{\text{L,join}}^i(\mathcal{C}) - d_{\text{L,join}}^{i-1}(\mathcal{C}) \geq M_{s-1}.$$

We get the claim using that

$$d_{\text{L,join}}^K(\mathcal{C}) = \sum_{i=0}^{s-1} n_i M_{s-1} = n M_{s-1},$$

where we have assumed that the code is non-degenerate. Note that we could have gotten this bound also by directly using

$$d_L(\mathcal{C}) \leq d_{L,\text{join}}^1(\mathcal{C}) = d_H^1(\mathcal{C})M_{s-1} = d_H(\mathcal{C})M_{s-1} \leq (n - K + 1)M_{s-1}.$$

□

This new Singleton bound is clearly much sharper than the previously known Lee-metric Singleton bounds, for example the bound from Theorem 10.

4.1. Density of Optimal Codes with respect to the Join-Lee Support.

Clearly, any code $\mathcal{C} \in (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K attaining this bound can be characterized by the following two properties:

1. The socle $\mathcal{C}_{s-1} = \mathcal{C} \cap \langle p^{s-1} \rangle$ is an MDS code over \mathbb{F}_p .
2. There exists a minimum Lee weight codeword in the socle.

The first property already implies sparsity as n tends to infinity and triviality for $p = 2$. Even the second property is problematic: $d_L(\mathcal{C}_{s-1}) = (n - K + 1)M_{s-1}$, implies that all nonzero entries of a minimal Hamming weight codeword in the socle must be of maximal Lee weight. Using the systematic form of the socle,

$$G_{s-1} = (p^{s-1}\mathbb{I}_K \quad p^{s-1}A),$$

we can immediately see that any row g of G_{s-1} is also of minimal Hamming weight $n - K + 1$. Thus, for g to have only nonzero entries of maximal Lee weight implies $p^{s-1} = M_{s-1}$, which will restrict optimal codes with respect to this bound to $p \in \{2, 3\}$ and any positive integer s . Because of the MDS property over \mathbb{F}_3 , we must have a block length $n \leq 4$.

We can drop the second condition, i.e., there exists a minimal Lee weight codeword in the socle, if we manage to estimate the difference

$$d_{L,\text{join}}^1(\mathcal{C}) - d_L(\mathcal{C}).$$

This task is, however, equally hard as bounding $d_L(\mathcal{C})$ itself.

Knowing hence, that only codes over $\mathbb{Z}/p^s\mathbb{Z}$ for $p = 2, 3$ with length $n = 4$ can attain this bound, the socles of the codes $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ with $p = 3$ and $n \leq 4$ attaining the bound in Theorem 31 are hence generated by a matrix of the form

$$(4.1) \quad \begin{pmatrix} 3^{s-1} & 0 & 0 & A \\ 0 & 3^{s-1} & 0 & B \\ 0 & 0 & 3^{s-1} & C \end{pmatrix},$$

where $A, B, C \in \mathbb{Z}/p^s\mathbb{Z}$ are such that their Lee weight is M_{s-1} .

Example 32. For instance, the code $\mathcal{C} \subseteq (\mathbb{Z}/9\mathbb{Z})^4$ of rank $K = 3$ generated by

$$\begin{pmatrix} 3 & 0 & 0 & 3 \\ 0 & 3 & 0 & 6 \\ 0 & 0 & 3 & 6 \end{pmatrix}.$$

This code has $d_L(\mathcal{C}) = 6$ and thus attains the join-Lee metric Singleton bound as $d_L(\mathcal{C}) = 6 = 3 \cdot (4 - 3 + 1) = M_{s-1}(n - K + 1)$.

Note that for MDS codes, we actually know all r -th generalized Hamming weights: let $\mathcal{C} \subseteq \mathbb{F}_q^n$ be a linear code of dimension k , then

$$d_{\mathbb{H}}^r(\mathcal{C}) = n - k + r.$$

Thus, a natural question that arises is whether the optimal codes with respect to the newly defined Lee-metric Singleton bound have a similar behaviour. That is, we are interested in an expression for the r -th generalized join-Lee weight $d_{\mathbb{L},\text{join}}^r(\mathcal{C})$ for every $r \in \{1, \dots, K\}$.

Proposition 33. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be code of rank K attaining the join-support bound in Theorem 31. Then, for each $r \in \{1, \dots, K\}$, the r -th generalized join-Lee weight is given by*

$$d_{\mathbb{L},\text{join}}^r(\mathcal{C}) = p^{s-1}(n - K + r).$$

Proof. Recall that \mathcal{C} is optimal with respect to the join-support Lee-metric Singleton bound, if its minimum Lee weight code word has the form

$$c_{\min} = (0 \ \dots \ 0 \ p^{s-1} \ 0 \ \dots \ 0 \mid \pm p^{s-1} \ \dots \ \pm p^{s-1}).$$

Thus, it holds $d_{\mathbb{L},\text{join}}^1(\mathcal{C}) = p^{s-1}(n - K + 1)$.

For arbitrary $r \in \{1, \dots, K\}$, the r -dimensional subcodes of \mathcal{C} attaining the r -th generalized join-Lee weight are contained in the socle of the code as well. Hence, they admit a generator matrix G_r which is permutation equivalent to a matrix of the form

$$G_r = (\mathbf{0} \mid p^{s-1}\mathbb{I}_r \mid \pm p^{s-1}\mathbb{J}),$$

where $\mathbf{0}$ is the all-zero matrix of size $r \times (K - r)$ and \mathbb{J} is an all-one matrix of size $r \times (n - K)$. Thus, the desired result follows. \square

4.2. Invariance under Isometry in the Lee Metric.

For the generalized Hamming weights of a linear k -dimensional code $\mathcal{C} \subseteq \mathbb{F}_q^n$, we also know that $d_{\mathbb{H}}^r(\mathcal{C}) = d_{\mathbb{H}}^r(\mathcal{C}')$, for any equivalent code \mathcal{C}' and any $r \in \{1, \dots, k\}$. Also for the generalized join-Lee weights we have the same behaviour.

Proposition 34. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K , then $d_{\mathbb{L},\text{join}}^r(\mathcal{C}) = d_{\mathbb{L},\text{join}}^r(\mathcal{C}')$, for all $r \in \{1, \dots, K\}$ and all \mathcal{C}' which are equivalent to \mathcal{C} , under the Lee-metric isometries.*

Proof. Recall that the Lee-metric isometries only consist of permuting the positions and multiplying any position by 1 or -1 . Thus, all codewords of \mathcal{C}' can be written as $c' = \sigma(c) \star v$, for some permutation σ and $v \in \{1, -1\}^n$, where \star denotes the coordinatewise multiplication and $c \in \mathcal{C}$. Now the claim follows immediately as

$$\begin{aligned} d_{\mathbb{L},\text{join}}^r(\mathcal{C}) &= \min\{|\{\max_{c \in \mathcal{C}}\{\text{wt}_{\mathbb{L}}(c_1)\}, \dots, \max_{c \in \mathcal{C}}\{\text{wt}_{\mathbb{L}}(c_n)\}\}| \mid c \in \mathcal{D} \leq \mathcal{C}, \text{rk}(\mathcal{D}) = r\} \\ &= \min\{|\{\sigma(\max_{c \in \mathcal{C}}\{\text{wt}_{\mathbb{L}}(c_1)\}), \dots, \max_{c \in \mathcal{C}}\{\text{wt}_{\mathbb{L}}(c_n)\}\}| \mid c \in \mathcal{D} \leq \mathcal{C}, \text{rk}(\mathcal{D}) = r\} \\ &= d_{\mathbb{L},\text{join}}^r(\mathcal{C}'). \end{aligned}$$

\square

5. COLUMN SUPPORT FOR THE LEE METRIC

We observe that in order to compute the r -th generalized Hamming weight of a code \mathcal{C} , all we do is considering a generator matrix G and count the number of nonzero columns, i.e., the column weight. In fact, for any r -th generalized Hamming weight one can choose r rows of G which attain the minimal column weight.

For a matrix $A \in \mathcal{R}^{K \times n}$ we will denote by $S_r(A) \in \mathcal{R}^{r \times n}$ all the submatrices of A of size $r \times n$.

Definition 35. Consider a matrix $A = (a_1^\top \cdots a_n^\top) \in \mathcal{R}^{K \times n}$. We define the *column weight*, $\text{wt}_{\mathcal{C}}(A)$, of A by the number of nonzero columns of A , i.e.,

$$\text{wt}_{\mathcal{C}}(A) := |\{i \in \{1, \dots, n\} \mid a_i \neq 0 \in \mathcal{R}^K\}|.$$

The *column support*, $\text{supp}_{\mathcal{C}}(A)$, of A is given by

$$\text{supp}_{\mathcal{C}}(A) := (\max(\text{supp}(a_1)), \dots, \max(\text{supp}(a_n))).$$

Again we have the nice property that $|\text{supp}_{\mathcal{C}}(A)| = \text{wt}_{\mathcal{C}}(A)$. In fact,

$$\text{wt}_{\mathcal{C}}(A) = |\text{supp}_{\mathcal{C}}(A)| = \sum_{i=1}^n \max(\text{supp}(a_i)).$$

Thus, we can define the column support, column weight and the generalized column weights of a code.

Definition 36. Let $\mathcal{C} \subseteq \mathcal{R}^n$ be a linear code of rank K . The *column support* of \mathcal{C} is given by the minimal column support of any generator matrix, i.e.,

$$(5.1) \quad \text{supp}_{\mathcal{C}}(\mathcal{C}) = \min_{G: \langle G \rangle = \mathcal{C}} \text{supp}_{\mathcal{C}}(G).$$

The *column weight* of a code is then given by the size of the column support, i.e.,

$$\text{wt}_{\mathcal{C}}(\mathcal{C}) = |\text{supp}_{\mathcal{C}}(\mathcal{C})|.$$

Finally, the r -th *generalized column weight* of \mathcal{C} is defined as

$$(5.2) \quad d_{\mathcal{C}}^r(\mathcal{C}) = \min\{\text{wt}_{\mathcal{C}}(\mathcal{D}) \mid \mathcal{D} \leq \mathcal{C}, \text{rk}(\mathcal{D}) = r\}.$$

Note that the definition of the r -th generalized column weight of a linear code $\mathcal{C} \subseteq \mathcal{R}^n$ of rank K is equivalent to

$$d_{\mathcal{C}}^r(\mathcal{C}) = \min\{\text{wt}_{\mathcal{C}}(S_r(G)) \mid \text{rk}(\langle S_r(G) \rangle) = r, \langle G \rangle = \mathcal{C}\}.$$

The difficulty of this new definition lies in the choice of the generator matrix instead of the choice of the subcode. This is the only difference to the usual definition of join support and weight.

The difficulty of finding the correct generator matrix to read of the minimal column weights, or the subcode with minimal weight is equivalent.

Let us show the dependency on the choice of generator matrix in the following example.

Example 37. Let us consider $\mathcal{C} \subseteq \mathbb{F}_2^5$ generated by

$$G = \begin{pmatrix} 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{pmatrix}.$$

If we were to compute the column (Hamming) weights of $S_r(G)$, we would get for $S_1(G)$

$$\text{wt}_{\mathcal{C}}((1 \ 0 \ 0 \ 1 \ 1)) = 3.$$

However, this is not the first generalized Hamming weight of the code. There exists a generator matrix G' , such that $S_r(G')$ attains the r -th generalized Hamming weights as column weights, for each $r \in \{1, \dots, k\}$:

$$G' = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \end{pmatrix}.$$

Now we can read of the r -th generalized Hamming weights easily:

$$\begin{aligned} d_{\mathcal{C}}^1 &= \text{wt}_{\mathcal{C}}((1 \ 1 \ 0 \ 0 \ 0)) = 2, \\ d_{\mathcal{C}}^2 &= \text{wt}_{\mathcal{C}}\left(\begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \end{pmatrix}\right) = 3, \\ d_{\mathcal{C}}^3 &= \text{wt}_{\mathcal{C}}\left(\begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \end{pmatrix}\right) = 5. \end{aligned}$$

Thus, the definition is not independent on the choice of generator matrix. Let us now adapt the definitions to the Lee weight.

Definition 38. Consider a matrix $A = (a_1^\top \ \dots \ a_n^\top) \in \mathcal{R}^{K \times n}$. Its *column Lee support* is given by the n -tuple

$$\text{supp}_{\text{L,col}}(A) = (\max(\text{supp}_{\text{L}}(a_1)), \dots, \max(\text{supp}_{\text{L}}(a_n))).$$

The *column Lee weight* of A is given by

$$\text{wt}_{\text{L,col}}(A) = |\text{supp}_{\text{L,col}}(A)| = \sum_{i=1}^n \max(\text{supp}_{\text{L}}(a_i)).$$

Note that this definition asks us to choose in each column the entry of maximal Lee weight.

Example 39. Let us consider the matrix

$$G = \begin{pmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 3 & 3 \end{pmatrix} \in \mathbb{Z}/9\mathbb{Z}^{3 \times 4}.$$

Then, the column Lee support and the column Lee weight of G are given by

$$\text{supp}_{\text{L,col}}(G) = (1, 1, 3, 3) \text{ and } \text{wt}_{\text{L,col}}(G) = 8.$$

We are now able to extend the definitions of column Lee support and column Lee weight to a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K .

Definition 40. Consider a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K . We define its *column Lee support* by the minimal column Lee weight of any generator matrix of \mathcal{C} , i.e.,

$$\text{supp}_{\text{L,col}}(\mathcal{C}) = \min_{G: \langle G \rangle = \mathcal{C}} \text{supp}_{\text{L,col}}(G).$$

The *column Lee weight* of \mathcal{C} is then given by the size of its column Lee support, i.e.,

$$\text{wt}_{\text{L,col}}(\mathcal{C}) = |\text{supp}_{\text{L,col}}(\mathcal{C})|.$$

As in the case for the Hamming metric, also in this case the definition is not independent on the choice of generator matrix. For this, we introduce the following matrix, called *reduced systematic* generator matrix.

Definition 41. Consider a matrix $G \in (\mathbb{Z}/p^s\mathbb{Z})^{K \times n}$ as given in (2.1). We call G to be in *reduced systematic form* if for every entry a of $A_{i,j} \in (\mathbb{Z}/p^{s+1-i}\mathbb{Z})^{k_i \times k_j}$ with $i < j \leq s$ it holds that $\text{wt}_{\text{L}}(a) \leq p^{j-1}$.

We will denote a matrix G in reduced systematic form by G_{rsys} . Let us give an example to clarify Definition 41.

Example 42. Consider $G \in \mathbb{Z}/27\mathbb{Z}^{3 \times 4}$

$$G = \begin{pmatrix} 1 & 14 & 11 & 0 \\ 0 & 9 & 18 & 0 \\ 0 & 0 & 9 & 18 \end{pmatrix}.$$

Note that G is in systematic form as defined in (2.1). By elementary row reduction, i.e., by subtracting suitable multiples of the rows r_j from row r_i with $1 \leq i < j \leq 3$, we obtain a matrix G_{rsys} in reduced systematic form

$$G_{\text{rsys}} = \begin{pmatrix} 1 & 5 & -7 & 0 \\ 0 & 9 & 9 & -18 \\ 0 & 0 & 9 & 18 \end{pmatrix}.$$

By a similar argument used to prove Proposition 3 we observe the following.

Proposition 43. Consider a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K and subtype (k_0, \dots, k_{s-1}) . The code \mathcal{C} is permutation equivalent to a code having a generator matrix in reduced systematic form.

This new systematic form yields a natural upper bound on the column Lee weight of a code \mathcal{C} . For this let us now consider the support subtype outside an information set of size K of the code. Since we can always find a permutation equivalent code, which has an information set in the first K positions, we can assume that we only consider the last $n - K$ columns of a generator matrix in reduced systematic form. In order not to confuse it with the support subtype (n_0, \dots, n_s) of the entire generator matrix, we will denote it by (μ_0, \dots, μ_s) .

Proposition 44. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K and subtype (k_0, \dots, k_{s-1}) and let $(\mu_0, \dots, \mu_{s-1})$ be the support subtypes in the last $n - K$ positions. Then the column Lee weight of \mathcal{C} is upper bounded by

$$\text{wt}_{\text{L,col}}(\mathcal{C}) \leq \sum_{i=0}^{s-1} p^i k_i + \sum_{i=0}^s \mu_i M_i.$$

Proof. By Definition 40 we have

$$\text{wt}_{\text{L,col}}(\mathcal{C}) = \left| \min_{G:\langle G \rangle = \mathcal{C}} \text{supp}_{\text{L,col}}(G) \right|.$$

Furthermore, by Proposition 43, \mathcal{C} admits a generator matrix $G_{\text{rsys}} \in (\mathbb{Z}/p^s\mathbb{Z})^{K \times n}$ in reduced systematic form. Hence, the column Lee weight of G_{rsys} is a natural upper bound to the column Lee weight of the code, i.e.,

$$\text{wt}_{\text{L,col}}(\mathcal{C}) \leq \text{wt}_{\text{L,col}}(G_{\text{rsys}}).$$

Thanks to the form of G_{rsys} , we observe that the maximum Lee weight in the first K columns is given by the entry $(G_{\text{rsys}})_{i,i}$ for $i \in \{1, \dots, K\}$. For the last $n - K$ columns we have to assume the maximal Lee weight. The support subtype (μ_0, \dots, μ_s) in these positions immediately tells us, how many columns are contained in which ideal. Hence, for each column lying in $\langle p^i \rangle$ (where i is maximal for this column) the maximal Lee weight is M_i . This yields the desired result. \square

Let us now introduce the r -th generalized column Lee weights of a code \mathcal{C} .

Definition 45. Given a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K and subtype (k_0, \dots, k_{s-1}) . The r -th generalized column Lee weight of \mathcal{C} is defined as

$$(5.3) \quad d_{\text{L,col}}^r(\mathcal{C}) = \min\{\text{wt}_{\text{L,col}}(\mathcal{D}) \mid \mathcal{D} \leq \mathcal{C}, \text{rk}(\mathcal{D}) = r\}.$$

Similarly to Definition (5.2), the r -th generalized column Lee weight is equivalent to

$$d_{\text{L,col}}^r(\mathcal{C}) = \min\{\text{wt}_{\text{L,col}}(S_r(G)) \mid \text{rk}(\langle S_r(G) \rangle) = r, \langle G \rangle = \mathcal{C}\}.$$

As in the Hamming-metric case, the difficulty lies now in finding a generator matrix attaining the r -th generalized column Lee weights. To visualize this, let us return to our previous example for the Lee-metric support.

Example 46. Let us consider the code $\mathcal{C} \subseteq \mathbb{Z}/9\mathbb{Z}^4$ generated by

$$G = \begin{pmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 3 & 3 \end{pmatrix},$$

which has support subtype $(4, 0, 0)$ and minimum Lee distance 2.

If we compute the minimal column weights of submatrices of G we get

$$\begin{aligned} \text{wt}_{\text{L,col}}((0 \ 1 \ 2 \ 0)) &= 3, \\ \text{wt}_{\text{L,col}}\left(\begin{pmatrix} 0 & 1 & 2 & 0 \\ 0 & 0 & 3 & 3 \end{pmatrix}\right) &= 7, \\ \text{wt}_{\text{L,col}}(G) &= 8. \end{aligned}$$

However, there is a generator matrix of the code which is not in systematic form and which attains smaller column Lee weights:

$$G' = \begin{pmatrix} 8 & 0 & 0 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 8 & 1 & 3 \end{pmatrix}.$$

The r -th generalized Lee weights are then

$$\begin{aligned} d_{\mathbb{L},\text{col}}^1(\mathcal{C}) &= \text{wt}_{\mathbb{L},\text{col}}((8 \ 0 \ 0 \ 1)) = 2 = d_{\mathbb{L}}(\mathcal{C}), \\ d_{\mathbb{L},\text{col}}^2(\mathcal{C}) &= \text{wt}_{\mathbb{L},\text{col}}\left(\begin{pmatrix} 8 & 0 & 0 & 1 \\ 0 & 1 & 8 & 0 \end{pmatrix}\right) = 4, \\ d_{\mathbb{L},\text{col}}^3(\mathcal{C}) &= \text{wt}_{\mathbb{L},\text{col}}\left(\begin{pmatrix} 8 & 0 & 0 & 1 \\ 0 & 1 & 8 & 0 \\ 0 & 0 & 3 & 0 \end{pmatrix}\right) = 6 = \text{wt}_{\mathbb{L},\text{col}}(\mathcal{C}). \end{aligned}$$

Note that both matrices within this example are of reduced systematic form.

Lemma 47. *Let $\mathcal{C} \in (\mathbb{Z}/p^s\mathbb{Z})^n$ be code of rank K . Let $G^{(i)} \in (\mathbb{Z}/p^s\mathbb{Z})^{i \times n}$ of a rank $i \in \{1, \dots, K-1\}$ be a generator matrix of a subcode of \mathcal{C} attaining $d_{\mathbb{L},\text{col}}^i(\mathcal{C})$. Let $c \in \mathcal{C}$ such that $\begin{pmatrix} G^{(i)} \\ c \end{pmatrix}$ is of a generator matrix of a subcode of rank $i+1$. Then it holds,*

$$\text{wt}_{\mathbb{L},\text{col}}\left(\begin{pmatrix} G^{(i)} \\ c \end{pmatrix}\right) > \text{wt}_{\mathbb{L},\text{col}}(G^{(i)}).$$

Proof. Let us define for all columns $j \in \{1, \dots, n\}$ the maximal Lee weight of the j -th column in $G^{(i)}$ as $A_j^{(i)}$. We clearly have $\text{wt}_{\mathbb{L},\text{col}}(G^{(i)}) \leq \text{wt}_{\mathbb{L},\text{col}}\left(\begin{pmatrix} G^{(i)} \\ c \end{pmatrix}\right)$. Thus,

let us assume that $\text{wt}_{\mathbb{L},\text{col}}(G^{(i)}) = \text{wt}_{\mathbb{L},\text{col}}\left(\begin{pmatrix} G^{(i)} \\ c \end{pmatrix}\right)$. Then

$$\sum_{j=1}^n A_j^{(i)} = \sum_{j=1}^n \max\{A_j^{(i)}, \text{wt}_{\mathbb{L}}(c_j)\}$$

and so for all $j \in \{1, \dots, n\}$ we have $\text{wt}_{\mathbb{L}}(c_j) \leq A_j^{(i)}$. However, as $G^{(i)}$ attains $d_{\mathbb{L},\text{col}}^i(\mathcal{C})$, the sum $\sum_{j=1}^n A_j^{(i)}$ is minimal among all rank i subcodes of \mathcal{C} . Hence, there is no index $j \in \{1, \dots, n\}$ for which $\text{wt}_{\mathbb{L}}(c_j) < A_j^{(i)}$ and thus for all j we have $\text{wt}_{\mathbb{L}}(c_j) = A_j^{(i)}$. This implies $c_j = \pm A_j^{(i)}$.

This means that c has in every position the maximal Lee weight over all rows of $G^{(i)}$. Thus, for every row g_ℓ of $G^{(i)}$ with $\ell \in \{1, \dots, i\}$ for which $\text{wt}_{\mathbb{L}}(g_\ell) > \text{wt}_{\mathbb{L}}(c)$, we can add and/or subtract c to decrease its weight. For each row $\ell \in \{1, \dots, i\}$ let us therefore define the sets

$$\begin{aligned} I_\ell^- &= \{j \in \{1, \dots, n\} \mid \text{wt}_{\mathbb{L}}(c_j - g_{\ell j}) < \text{wt}_{\mathbb{L}}(c_j)\}, \\ I_\ell^+ &= \{j \in \{1, \dots, n\} \mid \text{wt}_{\mathbb{L}}(c_j + g_{\ell j}) \leq \text{wt}_{\mathbb{L}}(c_j)\}. \end{aligned}$$

For a fixed row $\ell \in \{1, \dots, i\}$, if

$$\sum_{j \in I_\ell^-} \text{wt}_{\mathbb{L}}(c_j) < \sum_{j \in I_\ell^+} \text{wt}_{\mathbb{L}}(c_j),$$

then we add c to the row g_ℓ . If however,

$$\sum_{j \in I_\ell^+} \text{wt}_{\mathbb{L}}(c_j) \leq \sum_{j \in I_\ell^-} \text{wt}_{\mathbb{L}}(c_j),$$

we subtract c from that row g_ℓ .

We consider now the new row $g'_\ell := c \pm g_\ell$ which has a strictly smaller Lee weight than c . Since the cases are similar, assume that for g_ℓ the first case is true, i.e., $\sum_{j \in I_\ell^-} \text{wt}_L(c_j) < \sum_{j \in I_\ell^+} \text{wt}_L(c_j)$ and thus we add the row c , getting $g'_\ell := c + g_\ell$. Clearly for each position j in I_ℓ^- we at most added a Lee weight of $A_j^{(i)}$, while in each position j in I_ℓ^+ we subtracted a Lee weight of at most $A_j^{(i)}$, thus

$$\begin{aligned} \text{wt}_L(g'_\ell) &= \sum_{j \in I_\ell^+} \text{wt}_L(g_{\ell j} + c_j) + \sum_{j \in I_\ell^-} \text{wt}_L(g_{\ell j} + c_j) \\ &< \sum_{j \in I_\ell^+} \text{wt}_L(g_{\ell j} + c_j) + \sum_{j \in I_\ell^-} A_j^{(i)} + \sum_{j \in I_\ell^-} \text{wt}_L(c_j) \\ &< \sum_{j \in I_\ell^+} \text{wt}_L(c_j) - \sum_{j \in I_\ell^+} A_j^{(i)} + \sum_{j \in I_\ell^+} A_j^{(i)} + \sum_{j \in I_\ell^-} \text{wt}_L(c_j) \\ &= \text{wt}_L(c). \end{aligned}$$

Doing this procedure for every row of the matrix $G^{(i)}$, obtaining the new matrix $G'^{(i)}$ of rank i , we have

$$\text{wt}_{L,\text{col}}(G'^{(i)}) < \text{wt}_{L,\text{col}}(G^{(i)}),$$

since in every row we now reduced the Lee weight, but this is a contradiction to G attaining $d_{L,\text{col}}^i(\mathcal{C})$. \square

Finally, we are able to prove the desired properties for the generalized column Lee weights.

Proposition 48. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K . Then*

1. $d_{L,\text{col}}^1(\mathcal{C}) = d_L(\mathcal{C})$.
2. $d_{L,\text{col}}^r(\mathcal{C}) < d_{L,\text{col}}^{r+1}(\mathcal{C})$ for all $r < K$.
3. $d_{L,\text{col}}^K(\mathcal{C}) = \text{wt}_{L,\text{col}}(\mathcal{C})$.

Proof. For the first property, note that the column Lee weight of a $1 \times n$ matrix is equal to the Lee weight of that n -tuple. Since a minimal Lee-weight codeword c is the rank 1 subcode of \mathcal{C} with the smallest column Lee weight, it attains $\text{wt}_{L,\mathcal{C}}(c) = d_{L,\text{col}}^1(\mathcal{C})$.

The second property follows from Lemma 47. Since then for any matrix $G^{(i+1)} \in (\mathbb{Z}/p^s\mathbb{Z})^{(i+1) \times n}$, which attains $d_{L,\text{col}}^{i+1}(\mathcal{C})$, which we can write as $G^{(i+1)} = \begin{pmatrix} G^{(i)} \\ g' \end{pmatrix}$, we either have that G'_i already attained $d_{L,\text{col}}^i(\mathcal{C})$ and hence

$$d_{L,\text{col}}^i(\mathcal{C}) = \text{wt}_{L,\text{col}}(G^{(i)}) < \text{wt}_{L,\text{col}}(G^{(i+1)}) = d_{L,\text{col}}^{i+1}(\mathcal{C}),$$

or if $G^{(i)}$ did not attain $d_{L,\text{col}}^i(\mathcal{C})$, then

$$d_{L,\text{col}}^i(\mathcal{C}) < \text{wt}_{L,\text{col}}(G^{(i)}) \leq \text{wt}_{L,\text{col}}(G^{(i+1)}).$$

In either case, we get that $d_{L,\text{col}}^i(\mathcal{C}) < d_{L,\text{col}}^{i+1}(\mathcal{C})$.

Lastly, the third property follows immediately from the definition of the column Lee weight of a code \mathcal{C} . \square

The properties in Proposition 48 allow us to deduce a natural Singleton-like bound for the Lee metric.

Theorem 49. *Given a linear code $\mathcal{C} \in (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K . The minimum distance of \mathcal{C} is upper bounded by*

$$d_{\mathbb{L}}(\mathcal{C}) \leq \text{wt}_{\mathbb{L},\text{col}}(\mathcal{C}) - K + 1.$$

Proof. Using the properties given in Proposition 48 we note that

$$(5.4) \quad d_{\mathbb{L}}(\mathcal{C}) = d_{\mathbb{L},\text{col}}^1(\mathcal{C}) \leq d_{\mathbb{L},\text{col}}^K(\mathcal{C}) - \sum_{i=2}^K \left(d_{\mathbb{L},\text{col}}^i(\mathcal{C}) - d_{\mathbb{L},\text{col}}^{i-1}(\mathcal{C}) \right).$$

By the strict inequality between the generalized column Lee weights, we have a difference of at least one, i.e.,

$$d_{\mathbb{L},\text{col}}^i(\mathcal{C}) - d_{\mathbb{L},\text{col}}^{i-1}(\mathcal{C}) \geq 1.$$

Since $d_{\mathbb{L},\text{col}}^K(\mathcal{C}) = \text{wt}_{\mathbb{L},\text{col}}(\mathcal{C})$, the desired bound follows. \square

As for increasing parameters of a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K and subtype (k_0, \dots, k_{s-1}) it becomes harder to compute $\text{wt}_{\mathbb{L},\text{col}}(\mathcal{C})$, applying Proposition 44 we obtain a direct consequence to Theorem 49, which requires no computational effort.

Corollary 50. *Given a linear code $\mathcal{C} \in (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K . The minimum distance of \mathcal{C} is upper bounded by*

$$d_{\mathbb{L}}(\mathcal{C}) \leq \sum_{i=0}^{s-1} p^i k_i + \sum_{i=0}^s \mu_i M_i - K + 1.$$

The bounds given in Theorem 49 and Corollary 50 improve the Singleton bound by Shiromoto [31] and the one by Alderson-Huntemann [1]. In the proof of Theorem 49 we bounded the differences $d_{\mathbb{L},\text{col}}^i(\mathcal{C}) - d_{\mathbb{L},\text{col}}^{i-1}(\mathcal{C})$ by one for every $i = 2, \dots, K$. However, for a relatively small rank K this bound is not very tight. The sum in Equation (5.4) is a telescoping sum, meaning that

$$\sum_{i=2}^K \left(d_{\mathbb{L},\text{col}}^i(\mathcal{C}) - d_{\mathbb{L},\text{col}}^{i-1}(\mathcal{C}) \right) = d_{\mathbb{L},\text{col}}^K(\mathcal{C}) - d_{\mathbb{L},\text{col}}^1(\mathcal{C}) = \text{wt}_{\mathbb{L},\text{col}}(\mathcal{C}) - d_{\mathbb{L}}(\mathcal{C}).$$

Hence, the goal is now to derive a lower bound on the difference $\text{wt}_{\mathbb{L},\text{col}}(\mathcal{C}) - d_{\mathbb{L}}(\mathcal{C})$ allowing us to further tighten the Singleton-like bound.

In the following let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K and subtype (k_0, \dots, k_{s-1}) . Let us introduce the maximal subtype $i \in \{0, \dots, s-1\}$ for which k_i is nonzero, that is

$$\sigma := \max \{ i \in \{0, \dots, s-1\} \mid k_i \neq 0 \}.$$

Proposition 51. *Let p be an odd prime. For a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K and subtype (k_0, \dots, k_{s-1}) and maximal subtype k_σ , we get the following lower bound*

$$\text{wt}_{\mathbb{L},\text{col}}(\mathcal{C}) - d_{\mathbb{L}}(\mathcal{C}) \geq \sum_{i=0}^{\sigma-1} \left(\sum_{j=0}^i k_j \right) \lfloor p/2 \rfloor p^i + (k_\sigma - 1)p^\sigma.$$

Proof. Let us start by focusing on the generalized column Lee weights. Assume that $c_1 \in \mathcal{C}$ is such that $d_{\text{L,col}}^1(\mathcal{C}) = \text{wt}_{\text{L,col}}(\langle c_1 \rangle)$. By Lemma 47, we know that the generalized column Lee weights can be obtained in an iterative fashion. Hence, to find a subcode \mathcal{D}_2 of rank 2 we are looking for a codeword $c_2 \in \mathcal{C}$ such that $\begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$ is of rank 2 and such that it minimizes $\text{wt}_{\text{L,col}}\left(\begin{pmatrix} c_1 \\ c_2 \end{pmatrix}\right)$. We continue with this process until we obtain a matrix

$$G_K := \begin{pmatrix} c_1 \\ \vdots \\ c_K \end{pmatrix}$$

of rank K such that $\text{wt}_{\text{L,col}}(G_K) = \text{wt}_{\text{L,col}}(\mathcal{C}) = d_{\text{L,col}}^K(\mathcal{C})$.

Since the code \mathcal{C} is of subtype $(k_0, \dots, k_\sigma, 0, \dots, 0)$, the rows of the matrix G_K each correspond to one of the σ blocks formed by the systematic form G_{sys} of G_K . To understand the difference of $\text{wt}_{\text{L,col}}(\mathcal{C})$ and the first generalized column Lee weight $d_{\text{L,col}}^1(\mathcal{C})$ we can think of successively removing rows from G_K until we are only left with the minimum weight codeword c_1 . Thinking in the block-wise structure of a generator matrix in systematic form, at some point we will have cancelled k_i rows corresponding to the i -th block of G_{sys} . Hence, the minimal difference subtracted is

$$M_{i-1} - M_i = \lfloor p/2 \rfloor p^{i-1}.$$

Doing this successively for every k_i , with $i \in \{0, \dots, \sigma\}$, gives

$$\sum_{i=0}^{\sigma-1} \binom{i}{j=0} k_j \lfloor p/2 \rfloor p^i.$$

At this point we are left only with a block corresponding to the rows belonging to the maximal subtype k_σ . The minimal difference between the rows of the same block is given by p^σ . Hence, cancelling $(k_\sigma - 1)$ rows yields to a difference of $p^\sigma(k_\sigma - 1)$ and the desired result follows. \square

A natural consequence (combining Propositions 44 and 51) is the next bound on the minimum Lee distance $d_{\text{L}}(\mathcal{C})$ of a code \mathcal{C} of given rank and subtype.

Corollary 52. *Consider a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$, where p is an odd prime. Let \mathcal{C} be of rank K and subtype (k_0, \dots, k_{s-1}) with maximal subtype k_σ and having support subtype $(\mu_0, \dots, \mu_{s-1})$ in the last $n - K$ positions. Then the following upper bound on the minimum Lee distance of \mathcal{C} holds*

$$d_{\text{L}}(\mathcal{C}) \leq \sum_{i=0}^{s-1} p^i k_i + \sum_{i=0}^s \mu_i M_i - \left[\sum_{i=0}^{\sigma-1} \binom{i}{j=0} k_j \lfloor p/2 \rfloor p^i + (k_\sigma - 1)p^\sigma \right].$$

Let us give an example over $\mathbb{Z}/9\mathbb{Z}$.

Example 53. Consider again the code \mathcal{C} generated by

$$G = \begin{pmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 3 & 3 \end{pmatrix},$$

over $\mathbb{Z}/9\mathbb{Z}$. In the last column, the code \mathcal{C} has support subtype $(1, 0, 0)$ and minimum Lee distance 2. Furthermore, we observe that $\sigma = 1$ and support subtype $(1, 0)$. Hence, by Corollary 52

$$d_{\mathbb{L}}(\mathcal{C}) \leq 2 + 3 + 1 \cdot 4 - [2 \cdot 1 + (1 - 1)3] = 7.$$

Similarly to the join-support, examples of codes attaining this bound are codes generated by matrices $G = (p^{s-1}\mathbb{I}_K \quad p^{s-1}A)$ for $A \in (\mathbb{Z}/p^s\mathbb{Z})^{K \times (n-K)}$, where $p = 3$. In fact, for any odd p these codes have a minimum Lee distance $d = p^{s-1}(n - K + 1)$. Furthermore we note that in the last $n - K$ positions we have support subtype $(0, \dots, 0, n - K)$ and $M_{s-1} = \lfloor p/2 \rfloor p^{s-1}$. Hence, inserting these values in the bound given in Corollary 52 gives

$$d_{\mathbb{L}}(\mathcal{C}) \leq p^{s-1}(1 + (n - K) \lfloor p/2 \rfloor).$$

This is equal to $d_{\mathbb{L}}(\mathcal{C})$ exactly if $p = 3$.

For instance, consider again the code $\mathcal{C} \subseteq (\mathbb{Z}/9\mathbb{Z}^4)$ of rank $K = 3$ with generator matrix

$$G = \begin{pmatrix} 3 & 0 & 0 & 3 \\ 0 & 3 & 0 & 6 \\ 0 & 0 & 3 & 6 \end{pmatrix}.$$

This code has minimum Lee distance $d_{\mathbb{L}}(\mathcal{C}) = 6$ and subtype $(k_0, k_1) = (0, 3)$. Hence, we also have $\sigma = 1$. The support subtype in the last $n - K = 1$ positions is $(0, 1, 0)$ and $M_1 = 3$. Computing the bound in Corollary 52 gives then

$$\sum_{i=0}^1 p^i k_i + \sum_{i=0}^2 \mu_i M_i - (k_1 - 1)p = 3 \cdot 3 + 1 \cdot 3 - (3 - 1)3 = 2 \cdot 3 = 6$$

and we conclude that this code is optimal with respect to Lee-metric Singleton bound 52.

5.1. Density of Optimal Codes with respect to the Column-Lee Support.

Let us discuss the density of the codes attaining the bound in Corollary 52. Recall that the bound is derived by $d_{\mathbb{L}}(\mathcal{C}) \leq \text{wt}_{\mathbb{L}, \text{col}}(\mathcal{C}) - (\text{wt}_{\mathbb{L}, \text{col}}(\mathcal{C}) - d_{\mathbb{L}}(\mathcal{C}))$ where we upper bounded the column weight of the code by $\text{wt}_{\mathbb{L}, \text{col}}(\mathcal{C}) \leq \sum_{i=0}^{s-1} p^i k_i + \sum_{i=0}^s \mu_i M_i$. Hence, in order to have codes attaining the bound on the minimum Lee distance, they must attain the bound on the column Lee weight too. That is, their generator matrix G must be in reduced systematic form. Furthermore, the support subtype of the last $n - K$ positions is (μ_0, \dots, μ_s) where in each of the μ_i positions the maximum Lee weight M_i is attained. For instance, a generator matrix may look as follows:

$$G_{\text{rsys}} = \left[\begin{array}{c|c} U & \left[\begin{array}{c|c|c|c} \left[\begin{array}{c} \mu_0 \end{array} \right] & \left[\begin{array}{c} \mu_1 \end{array} \right] & \dots & \left[\begin{array}{c} \mu_s \end{array} \right] \end{array} \right] \end{array} \right]$$

There are two options to attain a Lee weight M_i . Hence, the probability that a generator matrix is of this form is given by the number of such matrices divided by

the number of all matrices, i.e.,

$$\begin{aligned} \prod_{i=0}^{s-1} \left(\frac{2(p^{s-i})^{(k-1)}}{(p^{s-i})^{(k-1)}(p^{s-i} - p^{s-i-1})} \right)^{\mu_i} &= \prod_{i=0}^{s-1} \left(\frac{2}{p^{s-i} - p^{s-i-1}} \right)^{\mu_i} \\ &= 2^{n-K} \prod_{i=0}^{s-1} \left(\frac{1}{p^{s-i}(1 - 1/p)} \right)^{\mu_i} \\ &= 2^{n-K} \prod_{i=0}^{s-1} \left(\frac{p^{i+1}}{p^s(p-1)} \right)^{\mu_i}. \end{aligned}$$

Note that $p^s(p-1) > p^{i+1}$ for every $i \in \{0, \dots, s-1\}$. Hence, the fraction in the product is smaller than 1. Therefore, for $p \rightarrow \infty$ the product tends to 0. The same argument holds if we let s tend to infinity. Similarly, as μ_i depends on n , we note that $\frac{2}{p^{s-i} - p^{s-i-1}} < 1$. This implies that if $n \rightarrow \infty$ the product tends to zero as well. Thus, codes attaining the bound in Corollary 52 are sparse with respect to p , s and n .

Given an optimal code with respect to the Lee-metric Singleton bound 52, one could also ask if the r -th generalized column Lee weights are then also fixed. Since the main problem of the column Lee weight of a code is the computational difficulty, we leave this as an open question.

5.2. Invariance under Isometry in the Lee Metric.

Finally, we ask if the r -th generalized column Lee weights are fixed under isometries.

Proposition 54. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of rank K , then any equivalent code \mathcal{C}' , under the linear Lee-metric isometries is such that*

$$d_{\text{L,col}}^r(\mathcal{C}) = d_{\text{L,col}}^r(\mathcal{C}'),$$

for every $r \in \{1, \dots, K\}$.

Proof. Recall that any generator matrix $G^{(i)}$ of a subcode of rank i of a equivalent \mathcal{C}' can be written as $G^{(i)} = G^{(i)} P \text{diag}(v)$, for some permutation matrix P , $v \in \{1, -1\}^n$ and some generator matrix $G^{(i)}$ of a subcode of rank i of \mathcal{C} . Both, $G^{(i)}$ and $G^{(i)}$ have the same column weight. Now the claim follows immediately as

$$\begin{aligned} d_{\text{L,col}}^r(\mathcal{C}) &= \min\{\text{wt}_{\text{L,col}}(G^{(r)}) \mid \langle G^{(r)} \rangle \leq \mathcal{C}, \text{rk}(\langle G^{(r)} \rangle) = r\} \\ &= \min\{\text{wt}_{\text{L,col}}(G^{(r)} P \text{diag}(v)) \mid \langle G^{(r)} \rangle \leq \mathcal{C}, \text{rk}(\langle G^{(r)} \rangle) = r\} = d_{\text{L,col}}^r(\mathcal{C}'). \end{aligned}$$

□

6. GENERALIZED LEE WEIGHTS FROM FILTRATION

The resulting Lee-metric Singleton bounds in Theorem 31 and Corollary 52 are improving the previously known bounds, however their optimal codes are sparse and the column Lee weight of a code is computationally difficult to compute. We thus ask if fixing the rank of the subcode is the correct direction. In fact, a ring-linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K has very natural subcodes to consider, which are all of rank K .

Definition 55. For each $i \in \{0, \dots, s-1\}$ we define the i -th filtration subcode \mathcal{C}_i of \mathcal{C} as the intersection of \mathcal{C} with the ideal $\langle p^i \rangle$, i.e.,

$$\mathcal{C}_i := \mathcal{C} \cap \langle p^i \rangle.$$

The $(s-1)$ -st filtration \mathcal{C}_{s-1} is commonly known as the *socle* of the code \mathcal{C} .

Note that the filtration subcodes naturally form a chain, namely

$$(6.1) \quad \mathcal{C}_{s-1} \subseteq \mathcal{C}_{s-2} \subseteq \dots \subseteq \mathcal{C}_1 \subseteq \mathcal{C}_0 = \mathcal{C}.$$

We then define a new class of generalized Lee weights, or more concretely generalized Lee distances, coming from filtration subcodes.

Definition 56. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code. For each $r \in \{1, \dots, s\}$ we define the r -th generalized minimum Lee distance of the code \mathcal{C} to be the minimum distance of the filtration subcode \mathcal{C}_{r-1} , that is

$$d_L^r(\mathcal{C}) = d_L(\mathcal{C}_{r-1}).$$

The generalized minimum Lee distances have some natural properties that are summarized in the following.

Proposition 57. Given a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K and subtype (k_0, \dots, k_{s-1}) , let $\sigma := \max\{i \in \{0, \dots, s-1\} \mid k_i \neq 0\}$. Then the generalized minimum Lee distances satisfy

1. $d_L^1(\mathcal{C}) = d_L(\mathcal{C})$,
2. $d_L^r(\mathcal{C}) \leq d_L^{r+1}(\mathcal{C})$ for every $r \in \{1, \dots, s-1\}$,
3. $d_L^r(\mathcal{C}) \leq p^{r-1} + (n-k)M_{r-1}$ for every $r = \sigma+1, \dots, s$.

Proof. The first and second property immediately follow from (6.1).

For the third property we observe that for every $r \in \{\sigma+1, \dots, s\}$, by applying elementary row operations, we can bring a generator matrix G_{r-1} of \mathcal{C}_{r-1} in the form

$$(6.2) \quad G_{r-1} = (p^{r-1}\mathbb{I}_K \quad A),$$

where $A \in (p^{r-1}\mathbb{Z}/p^s\mathbb{Z})^{K \times (n-K)}$. The r -th minimum Lee distance is upper bounded by the Lee weight of any row of G_{r-1} . For each row, the first K positions have a Lee weight of exactly p^{r-1} . In the last $n-K$ positions of each row we assume the maximal Lee weight given by $M_{r-1} := \lfloor p^{s-(r-1)}/2 \rfloor p^{r-1}$ and hence the inequality follows. \square

Due to Property 2. in Proposition 57, we cannot use the usual Singleton-like argument and decrease the weight of the whole code. Instead, we note that any $d_L^r(\mathcal{C})$ is a direct upper bound on the minimum Lee distance. The only question that remains, is how far we have to go down in the filtration to expect the lowest minimum Lee distance, $d_L^1(\mathcal{C})$. In the following we identify several parameters of the code, that are easy to read off from any generator matrix of the code, that indicates which filtration subcode gives an appropriately low upper bound on $d_L(\mathcal{C})$.

The upper bound on the generalized minimum Lee distances in Property 3. of Proposition 57 is relatively loose. This is due to the fact, that we have assumed no knowledge about the matrix A given in (6.2).

As computing the minimum Lee distance of every subcode \mathcal{C}_i is an exhausting task, especially if there is no knowledge about the structure of A , we would like to introduce some more parameters regarding A for the first filtration of \mathcal{C} admitting a generator matrix of the form (6.2). That is the filtration \mathcal{C}_σ with a generator matrix of the form $G_\sigma = (p^\sigma \mathbb{I}_K \ A)$, for some matrix $A \in (p^\sigma \mathbb{Z}/p^s \mathbb{Z})^{K \times (n-k)}$. Let a_{ij} denote the entry of A lying in row i and column j . For each row of A , we determine the maximal power of p appearing and we denote it by

$$\ell_i := \max \left\{ k \in \{\sigma, \dots, s-1\} \mid \exists a_{ij} : \langle a_{ij} \rangle = \langle p^k \rangle, K+1 \leq j \leq n \right\}.$$

Clearly, $\ell_i \geq \sigma$. Let n'_i denote the number of entries of the i -th row of A that live in the ideal $\langle p^{\ell_i} \rangle$, i.e.,

$$n'_{\ell_i} := \left| \left\{ j \in \{K+1, \dots, n\} \mid a_{ij} \in \langle p^{\ell_i} \rangle \right\} \right|.$$

For a given linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s \mathbb{Z})^n$, these parameters help us to understand the evolution of the matrix A in the generator matrices of the filtration subcodes \mathcal{C}_{r-1} , for $r \in \{\sigma+1, \dots, s\}$. In fact, given a generator matrix G_σ of the filtration \mathcal{C}_σ in the form (6.2), the parameters ℓ_i and n'_{ℓ_i} for a row $i \in \{1, \dots, K\}$ allow to understand at which point in the filtration these positions become zero. More precisely, knowing ℓ_i and n'_{ℓ_i} implies that in $\mathcal{C}_{s-\ell_i+\sigma}$ there are n'_{ℓ_i} many zero entries in i -th row of A .

Knowing the number of entries turning into zero in a certain filtration is a huge advantage in bounding the minimum distance of a code. Therefore, we define by $n^{(r-1)}$ the maximal number of zeros we can get in the last $n-K$ positions of a row of a generator matrix of the filtration \mathcal{C}_{r-1} . That is, for every $r \in \{\sigma+1, \dots, s\}$,

$$n^{(r-1)} := \max \left\{ n'_{\ell_i} \mid \ell_i > s-r+\sigma, i \in \{1, \dots, K\} \right\}.$$

If there is no ℓ_i with $\ell_i > s-r+\sigma$, we will set $n^{(r-1)} = 0$. Furthermore, let $\ell^{(r-1)}$ be the corresponding value ℓ_i to $n^{(r-1)}$, i.e.,

$$\ell^{(r-1)} := \max \left\{ \ell_i \mid n'_{\ell_i} = n^{(r-1)}, i \in \{1, \dots, K\} \right\}.$$

We can hence refine the third property in Proposition 57 as follows.

Lemma 58. *Given a linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s \mathbb{Z})^n$ of subtype (k_0, \dots, k_{s-1}) with maximal subtype k_σ . Then, for every $r \in \{\sigma+1, \dots, s\}$, the r -th generalized Lee distance can be upper bounded by*

$$d_{\mathbb{L}}^r(\mathcal{C}) = d_{\mathbb{L}}(\mathcal{C}_{r-1}) \leq p^{r-1} + (n-K-n^{(r-1)})M_{r-1}.$$

Proof. The proof follows in a similar fashion as the proof of Proposition 57 by focusing on the row with the maximal number of zeros in the last $n-K$ columns of \mathcal{C}_{r-1} which is captured in $n^{(r-1)}$. Hence, the remaining $(n-K-n^{(r-1)})$ positions are bounded by the maximal Lee weight in the ideal considering, which is given by M_{r-1} . \square

Example 59. Let us consider a free code $\mathcal{C} \in (\mathbb{Z}/27\mathbb{Z})^5$ spanned by the rows of the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 21 & 6 \\ 0 & 1 & 0 & 10 & 7 \\ 0 & 0 & 1 & 18 & 8 \end{pmatrix} =: (\mathbb{I}_3 \ A).$$

We easily check that

$$\begin{aligned}\ell_1 &= 1 & \text{and} & & n'_{\ell_1} &= 2, \\ \ell_2 &= 0 & \text{and} & & n'_{\ell_2} &= 2, \\ \ell_3 &= 2 & \text{and} & & n'_{\ell_3} &= 1.\end{aligned}$$

Let us now consider the filtration subcodes \mathcal{C}_1 and \mathcal{C}_2 in order to compute the bound given in Proposition 57. Note that in this case $\sigma = 0$ as the code is free. For $\mathcal{C}_\sigma = \mathcal{C}_0 = \mathcal{C}$, the values ℓ_i and n'_i are given above. As $\ell_3 = 2$ and $n'_3 = 1$, at filtration $\mathcal{C}_{3-2+0} = \mathcal{C}_1$ there is one entry equal to zero. Indeed, $\mathcal{C}_1 = \mathcal{C} \cap \langle 3 \rangle$ has a generator matrix of the form

$$\begin{pmatrix} 3 & 0 & 0 & 9 & 18 \\ 0 & 3 & 0 & 3 & 21 \\ 0 & 0 & 3 & 0 & 24 \end{pmatrix},$$

where the last row contains one zero element in the last 2 columns. Note that $n'^{(1)} = n'_{\ell_3} = 1$ and hence, $d_{\mathbb{L}}^2(\mathcal{C}) \leq 3 + (5 - 3 - 1)12 = 15$.

Similarly, at filtration $\mathcal{C}_2 = \mathcal{C} \cap \langle 9 \rangle$ we observe two zero entries in the first row, as

$$\begin{pmatrix} 9 & 0 & 0 & 0 & 0 \\ 0 & 9 & 0 & 9 & 9 \\ 0 & 0 & 9 & 0 & 18 \end{pmatrix}.$$

Here we notice that $n'^{(2)} = n'_{\ell_1} = 2$ and thus $d_{\mathbb{L}}^3(\mathcal{C}) \leq 9 + (5 - 3 - 2)9 = 9$.

By Proposition 57, we know that the r -th generalized Lee distances are in non-decreasing order. Therefore, for any $r \in \{\sigma + 1, \dots, s\}$ the bound in Lemma 58 is a valid upper bound for the minimum Lee distance of a code \mathcal{C} . However, as visible in Example 59, the bounds on the r -th minimum Lee distances do not have to follow the same non-decreasing order. As they all hold as an upper bound to the minimum Lee distance of the code, the following bound is a direct consequence of Lemma 58 by choosing the smallest among the bounds given in the statement.

Corollary 60. *Given a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of subtype (k_0, \dots, k_{s-1}) . For each $r \in \{\sigma + 1, \dots, s\}$ let $\ell \geq 1$ and (ℓ, n') be the pair $(\ell^{(r-1)}, n'^{(r-1)})$ minimizing*

$$p^{s-\ell^{(r-1)}+\sigma} + (n - K - n'^{(r-1)})M_{s-\ell^{(r-1)}+\sigma}.$$

Then the codes minimum distance is bounded by

$$d_{\mathbb{L}}(\mathcal{C}) \leq p^{s-\ell+\sigma} + (n - K - n')M_{s-\ell+\sigma}.$$

As for large rank K this minimum can take a while to compute, we can also derive a slightly weaker bound depending on the maximal value ℓ_i , which is easy to compute.

Corollary 61. *Given a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of subtype (k_0, \dots, k_{s-1}) of maximal subtype k_σ . For each $r \in \{\sigma + 1, \dots, s\}$ let $\ell := \max\{\ell_i \mid i = 1, \dots, K\}$ and define the corresponding value $n' := \max\{n_{\ell_i} \mid \ell_i = \ell, \text{ for } i = 1, \dots, K\}$. Then, the minimum distance is bounded by*

$$d_{\mathbb{L}}(\mathcal{C}) \leq \begin{cases} p^{s-\ell+\sigma} + (n - K - n')M_{s-\ell+\sigma} & \text{if } \ell \geq 1, \\ p^\sigma + (n - K)M_\sigma & \text{else.} \end{cases}$$

In fact, we can identify conditions, leading to four different cases for the bound provided in Corollary 61. For this very last observation, leading to the very last Lee-metric Singleton bound, we first need one last definition. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ be a linear code of maximal subtype k_σ and assume that \mathcal{C}_σ is generated by $(p^\sigma \mathbb{I} \ A)$. Let us denote the entries of A as $a_{i,j}$, for $i \in \{1, \dots, K\}$ and $j \in \{K+1, \dots, n\}$. We define

$$N' = \max\{j \in \{K+1, \dots, n\} \mid p \mid a_{i,j}, i \in \{1, \dots, K\}\}.$$

That is N' is the maximal number of entries in a row of A , which are divisible by p .

Example 62. Let us consider the code over $\mathbb{Z}/27\mathbb{Z}$ generated by

$$G = \begin{pmatrix} 1 & 0 & 3 & 6 \\ 0 & 1 & 18 & 1 \end{pmatrix}.$$

The previous bound from Corollary 61 would take $\ell = 2$ and $n' = 1$, instead $N' = 2$, as in the first row of A we have two entries that are divisible by p . In fact, this indicates the minimum Hamming weight codeword in the socle, in this case 1. Clearly, if N' is large, it is beneficial to go until the socle.

1. Case $\ell = \sigma$ or $n'/2 \leq \frac{p^{s-\ell}-1}{p^{s-\sigma}-1}$. In this case we stay in \mathcal{C}_σ :

$$d_L(\mathcal{C}) \leq p^\sigma + (n - K)M_\sigma$$

2. Case $\ell = s$. In this case we also stay in \mathcal{C}_σ , but observed some zero entries:

$$d_L(\mathcal{C}) \leq p^\sigma + (n - k - n')M_\sigma$$

3. Case $\ell \neq \sigma$ or $\ell \neq s$ and $n'/2 \geq \frac{p^{s-\ell}-1}{p^{s-\sigma}-1}$. In this case we can move to $\mathcal{C}_{s-\ell+\sigma}$:

$$d_L(\mathcal{C}) \leq p^{s-\ell+\sigma} + (n - k - n')M_{s-\ell+\sigma}$$

4. Case: if $n' \leq N' \frac{p^{\ell-\sigma}-p^{\ell-\sigma-1}}{p^{\ell-\sigma}-1} + (n - K - 2) \frac{p^{\ell-\sigma-1}-1}{p^{\ell-\sigma}-1}$. In this case we go to the socle:

$$d_L(\mathcal{C}) \leq p^{s-1} + (n - K - N')M_{s-1}.$$

Note also, that instead of taking the filtration subcodes $\mathcal{C}_i = \mathcal{C} \cap \langle p^i \rangle$, we could have also considered the torsion subcodes.

Definition 63. Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$. For $i \in \{0, \dots, s-1\}$, we call $\tilde{\mathcal{C}}_i = \mathcal{C} \bmod p^{s-i} \subseteq (\mathbb{Z}/p^{s-i}\mathbb{Z})^n$ the i -th torsion code.

We can, however, immediately observe that the i -th torsion code represented as a code over the ambient space is naturally a subcode of the filtration subcode as

$$p^i \tilde{\mathcal{C}}_i \subseteq \mathcal{C}_i \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n,$$

with $\text{rk}(p^i \tilde{\mathcal{C}}_i) = \sum_{j=0}^{i-1} k_j < \text{rk}(\mathcal{C}_i) = K$.

In fact, any generator matrix of $\tilde{\mathcal{C}}_i$ is a truncation of a generator matrix of G , i.e., we cut off the rows belonging to the subtypes k_i, \dots, k_{s-1} .

Thus if we would define the r -th generalized Lee distances as $d_L^r(\mathcal{C}) = d_L(\tilde{\mathcal{C}}_r)$, for $r \in \{0, \dots, s-1\}$ then $d_L(\mathcal{C}) \leq d_L(\mathcal{C}_i) \leq d_L(p^i \tilde{\mathcal{C}}_i)$. Thus, any upper bound on $d_L(p^i \tilde{\mathcal{C}}_i)$ would serve as upper bound on $d_L(\mathcal{C})$, but would be worse than taking directly bounds on the smaller $d_L(\mathcal{C}_i)$.

Finally, we want to note, that the same considerations also apply to the Hamming metric.

Corollary 64. *Given a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of subtype (k_0, \dots, k_{s-1}) of maximal subtype k_σ . For each $r \in \{\sigma + 1, \dots, s\}$ let $\ell := \max \{\ell_i \mid i = 1, \dots, K\}$ and define the corresponding value $n' := \max \{n_{\ell_i} \mid \ell_i = \ell, \text{ for } i = 1, \dots, K\}$. Then, the Hamming minimum distance is bounded by*

$$d_H(\mathcal{C}) \leq \begin{cases} 1 + (n - K - n') & \text{if } \ell \geq 1, \\ 1 + (n - K) & \text{else.} \end{cases}$$

Note, that the Lee-metric version, that is Corollary 61, is not directly implied by the Hamming-metric bound. Such a direct bound would say

$$d_L(\mathcal{C}) \leq \begin{cases} M(1 + (n - K - n')) & \text{if } \ell \geq 1, \\ M(1 + (n - K)) & \text{else.} \end{cases}$$

This is clearly a worse bound than our Lee-metric Singleton bound of Corollary 61.

6.1. Density of Optimal Codes with respect to Filtrations.

One interesting quantity is the number of codes of maximum achievable Lee distance for given parameters. We call a such a code a *maximum Lee distance* (MLD) code. We have already seen that codes attaining the bounds based on the join-support and based on the column support are sparse as p, s and n tend to infinity. In this subsection we discuss the density of MLD codes with respect to the new Lee-metric Singleton bound 61 from the filtration. If nothing else is stated we consider a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K and subtype (k_0, \dots, k_{s-1}) .

Recall that the bound from Corollary 61 is especially tight, if there are many zero positions in a row of a generator matrix of a filtration subcode. Given the rank K of a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ the probability that an entire row of A is zero, where A are the last $n - K$ columns of a generator matrix of a filtration \mathcal{C}_{r-1} with $r \in \{\sigma + 1, \dots, s\}$, is depending on σ , i.e., it depends on whether the code \mathcal{C} is free or not.

For $n \rightarrow \infty$ it is known [4] that

$$\mathbb{P}(\mathcal{C} \text{ is free}) = \begin{cases} 1 & \text{if } R < 1/2, \\ 0 & \text{if } R > 1/2. \end{cases}$$

Hence, in this case we would have to distinguish again the two cases. On the contrary for $p \rightarrow \infty$, it is well-known that the code \mathcal{C} is free with high probability, which implies that $\sigma = 0$. In this case, we have

1. For every $i \in \{1, \dots, K\}$, $\ell_i = 0$. Thus, the bound in Corollary 61 can be reduced to

$$d_L(\mathcal{C}) \leq 1 + (n - K)M,$$

which coincides with the Singleton-like bound provided by [31].

2. There is a $i \in \{1, \dots, K\}$ with $\ell_i \neq 0$. In this case, we can find the pair (ℓ, n') as in Corollary 61 and the minimum Lee distance is bounded by

$$d_L(\mathcal{C}) \leq p^{s-\ell} + (n - K - n')M_{s-\ell}.$$

The following Lemma shows that for $p \rightarrow \infty$ the first case occurs with high probability.

Lemma 65. *For a free linear code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$, as $p \rightarrow \infty$, $\ell = 0$ with high probability.*

Proof. Note that $\mathbb{P}(\ell = 0)$ is the probability that there is no multiple of p contained in the last $n - K$ columns of a generator matrix G of \mathcal{C} in systematic form. More explicitly, it is the probability that all of the entries in the last $n - K$ columns of G are units. That is,

$$\mathbb{P}(\ell = 0) = \left(\frac{(p-1)p^{s-1}}{p^s} \right)^{K(n-K)} = \left(1 - \frac{1}{p} \right)^{K(n-K)}.$$

Hence, letting p grow to infinity and keeping n and K fixed, yields the desired result. \square

This means that, with high probability, MLD codes are sparse as $p \rightarrow \infty$, as codes attaining the bound 10 of Shiromoto are sparse.

Note that, letting s grow to infinity and keeping p fixed, the size of the ring $\mathbb{Z}/p^s\mathbb{Z}$ still grows whereas the probability $\mathbb{P}(\ell = 0)$ is a nonzero constant. This let us suggest, that codes attaining the bound on the minimum distance derived from filtration subcodes might not be sparse.

We start by discussing the case, where the code \mathcal{C} is a free code, hence $\sigma = 0$. Free codes have a generator matrix of the form $(\mathbb{I}_K | A)$, whit $A \in (\mathbb{Z}/p^s\mathbb{Z})^{K \times (n-K)}$. If there is an $0 < \ell < s$ such that $n' = n_{\tilde{\ell}} = n - K$, the filtration $\mathcal{C}_{s-\tilde{\ell}}$ has an entire row equal to zero. This results in having an $(s - \tilde{\ell})$ -th generalized Lee distance of $p^{s-\tilde{\ell}}$ and hence $d_L(\mathcal{C}) \leq p^{s-\tilde{\ell}}$.

Let us investigate on the probability for A having a maximal $0 < \ell_i = \tilde{\ell} < s$ with corresponding $n' = n - K$. This requires that all other rows of A are contained at most in the ideal $\langle p^{\tilde{\ell}} \rangle$. The probability that A is of this form is therefore

$$\begin{aligned} \mathcal{P} &:= \frac{(p^{s-\tilde{\ell}} - p^{s-\tilde{\ell}-1})^{(n-K)} (p^{s-1} - p^{s-\tilde{\ell}-1})^{(K-1)} (p^{s-\tilde{\ell}} - p^{s-\tilde{\ell}-1})^{(n-K-1)(K-1)}}{(p^s)^{(n-K)K}} \\ &= (p^{-\tilde{\ell}} - p^{-\tilde{\ell}-1})^{(n-K)} (p^{s-1} - p^{-\tilde{\ell}-1})^{(K-1)} (p^{-\tilde{\ell}} - p^{-\tilde{\ell}-1})^{(n-K-1)(K-1)} \\ &= \left(\frac{1}{p^{\tilde{\ell}}} - \frac{1}{p^{\tilde{\ell}+1}} \right)^{(n-K-1)K+1} \left(\frac{1}{p} - \frac{1}{p^{\tilde{\ell}+1}} \right)^{(K-1)}. \end{aligned}$$

This probability tends to zero as $n \rightarrow \infty$, and thus MLD codes are sparse with respect to the bound given in Corollary 61 and $n \rightarrow \infty$. However, since \mathcal{P} does not dependent on s , as $s \rightarrow \infty$ and is a nonzero constant, this implies neither sparsity nor density. In any case, we have that with a probability $\mathbb{P}(\mathcal{C} \text{ is free})\mathcal{P}$ the minimum distance of the code is bounded by $d_L(\mathcal{C}) \leq p^{s-\tilde{\ell}}$.

Let us consider now codes that achieve the bound on the minimum Lee distance based on filtration subcodes, i.e., Corollary 61 and check whether this fixes the r -th generalizes Lee distances.

Clearly, if \mathcal{C} has maximal subtype k_σ and attains the bound in Corollary 61, then $d_{\mathbb{L}}(\mathcal{C}) = d_{\mathbb{L}}^1(\mathcal{C}) = \dots = d_{\mathbb{L}}^{\sigma-1}(\mathcal{C}) = d_{\mathbb{L}}(\mathcal{C}_\sigma)$. If $\sigma = s - 1$, or we are in the case 4, i.e., $n' \leq N' \frac{p^{\ell-\sigma} - p^{\ell-\sigma-1}}{p^{\ell-\sigma} - 1} + (n - K - 2) \frac{p^{\ell-\sigma-1} - 1}{p^{\ell-\sigma} - 1}$. In this case we go to the socle, and hence all $d_{\mathbb{L}}^r(\mathcal{C})$ are equal. If we are not in case 4, the behaviour of the filtration subcodes \mathcal{C}_r with $r \geq \sigma$ is more unpredictable.

As already discussed above there are codes with several properties which are attaining the bound in Corollary 61.

One class of codes that we want to consider are those having $n' = n - K$. Assuming that such a code attains the bound, the following result gives us a closed expression for the r -th generalized Lee distances for all r .

Proposition 66. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K , subtype (k_0, \dots, k_s) of maximal subtype k_σ , and tuple $(\ell, n - K)$, such that $d_{\mathbb{L}}(\mathcal{C}) = d_{\mathbb{L}}^{s-\ell+\sigma}(\mathcal{C})$. Then the r -th generalized Lee distance is given by*

$$d_{\mathbb{L}}^r(\mathcal{C}) = \begin{cases} p^{s-\ell+\sigma} & \text{for every } r \leq s - \ell + \sigma, \\ p^r & \text{for every } r > s - \ell + \sigma. \end{cases}$$

Proof. Since $d_{\mathbb{L}}(\mathcal{C}) = d_{\mathbb{L}}^{s-\ell+\sigma}(\mathcal{C})$ and since the r -th generalized Lee distances are increasing in r , we have $d_{\mathbb{L}}^r(\mathcal{C}) = d_{\mathbb{L}}^{s-\ell+\sigma}(\mathcal{C})$ for every $r \leq s - \ell + \sigma$. Hence, the first case is clear. For the second case note that $\mathcal{C}_{s-\ell+\sigma}$ admits a generator matrix containing only zeros in the last $n - K$ columns. These entries remain to be zero for every filtration \mathcal{C}_r with $r > s - \ell + \sigma$. Hence, the minimum distance $d_{\mathbb{L}}(\mathcal{C}_r)$ is always given by p^r . \square

6.2. Invariance under Isometry in the Lee Metric.

Finally, we observe again that the r -th generalized Lee distance for a code $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ coincides with the r -th generalized Lee distance of a code $\mathcal{C}' \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ that is equivalent to \mathcal{C} .

Proposition 67. *Let $\mathcal{C} \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ of rank K and let $\mathcal{C}' \subseteq (\mathbb{Z}/p^s\mathbb{Z})^n$ another code that is equivalent to \mathcal{C} . Then, for every $r \in \{1, \dots, K\}$, we have*

$$d_{\mathbb{L}}^r(\mathcal{C}) = d_{\mathbb{L}}^r(\mathcal{C}').$$

Proof. Let ϕ denote an isometry preserving the Lee distance. Recall that this isometry can only consist of permutations and multiplications by ± 1 . Furthermore, recall that the r -th generalized Lee distance is given by the minimum Lee distance of the r -th filtration subcode \mathcal{C}_r of \mathcal{C} , i.e.,

$$d_{\mathbb{L}}^r(\mathcal{C}) = d_{\mathbb{L}}(\mathcal{C}_r).$$

By the inclusion property of the filtrations, we have $\mathcal{C}_r \subseteq \mathcal{C}$. Note that ϕ additionally preserves this inclusion property, i.e.,

$$\mathcal{C}'_r := \phi(\mathcal{C}_r) \subseteq \phi(\mathcal{C}).$$

Hence, the minimum Lee distances of \mathcal{C}_r and \mathcal{C}'_r coincide. \square

7. COMPARISON OF THE BOUNDS

At this point let us compare the bound of Corollary 61 to the bounds derived from the new puncturing argument (Theorem 14), the join-Lee support (Theorem 31), to the column support (Corollary 52) and to the bounds provided by [31, 1]. We do so by providing first some examples that attain the bound from Corollary 61 and compare it to the other bounds.

Example 68. 1. Let $\mathcal{C} \subseteq (\mathbb{Z}/9\mathbb{Z})^4$ generated by

$$G = \begin{pmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 6 \\ 0 & 0 & 1 & 4 \end{pmatrix}.$$

We quickly observe that this code has a minimum Lee distance $d_L(\mathcal{C}) = 3$. For the last $n - K = 1$ column, we note, that all the entries live in the ideal generated by 1. This means that $\ell = 0$ and $n' = n - K = 1$. Then the bounds are computed as follows.

Filtration:	$d_L(\mathcal{C}) \leq 3$	(Corollary 61)
Join-support:	$d_L(\mathcal{C}) \leq 6$	(Theorem 31)
Column support:	$d_L(\mathcal{C}) \leq 5$	(Corollary 52)
New puncturing:	$d_L(\mathcal{C}) \leq 8$	(Theorem 14)
Shiromoto:	$d_L(\mathcal{C}) \leq 5$	([31])
Alderson - Huntemann:	$d_L(\mathcal{C}) \leq 4$	([1])

2. Let $\mathcal{C} \subseteq (\mathbb{Z}/27\mathbb{Z})^5$ generated by

$$G = \begin{pmatrix} 1 & 10 & 4 & 20 & 9 \\ 0 & 3 & 9 & 18 & 9 \end{pmatrix}.$$

The minimum Lee distance of this code is $d_L(\mathcal{C}) = 9$. For the last $n - K = 3$ columns, we quickly compute $\ell' = 2$ and $n' = 1$. Then the bounds are computed as follows.

Filtration:	$d_L(\mathcal{C}) \leq 9$	(Corollary 60)
Filtration:	$d_L(\mathcal{C}) \leq 9$	(Corollary 61)
Join-support:	$d_L(\mathcal{C}) \leq 36$	(Theorem 31)
Column support:	$d_L(\mathcal{C}) \leq 38$	(Corollary 52)
New puncturing:	$d_L(\mathcal{C}) \leq 48$	(Theorem 14)
Shiromoto:	$d_L(\mathcal{C}) \leq 40$	([31])
Alderson - Huntemann:	not existing	([1])

3. In this example let us consider the code $\mathcal{C} \subseteq (\mathbb{Z}/125\mathbb{Z})^6$ generated by

$$G = \begin{pmatrix} 1 & 0 & 25 & 50 & 75 & 100 \\ 0 & 1 & 2 & 3 & 4 & 5 \end{pmatrix}.$$

This code has minimum distance $d_L(\mathcal{C}) = 5$. Note that the two bounds with respect to the filtration (Corollary 60 and 61) coincide. Hence, we obtain

$$\begin{aligned} \text{Filtration:} \quad & d_L(\mathcal{C}) \leq 5 && \text{(Corollary 60 and 61)} \\ \text{Join-support:} \quad & d_L(\mathcal{C}) \leq 200 && \text{(Theorem 31)} \\ \text{Column support:} \quad & d_L(\mathcal{C}) \leq 247 && \text{(Corollary 52)} \\ \text{New puncturing:} \quad & d_L(\mathcal{C}) \leq 300 && \text{(Theorem 14)} \\ \text{Shiromoto:} \quad & d_L(\mathcal{C}) \leq 249 && \text{([31])} \\ \text{Alderson - Huntemann:} \quad & d_L(\mathcal{C}) \leq 248 && \text{([1])} \end{aligned}$$

We now compare the bounds for different parameters. We will leave out the bound given by the column support, i.e., Corollary 52, as we would need to consider too many different parameters which would not fit in the overview.

(n, K, p^s, σ)	Alderson - Huntemann [1]	Shiromoto [31]	Join-support (Theorem 31)	Filtration (Corollary 61) (ℓ, n')
$(6, 3, 9, 0)$	12	16	12	$(0, 3) : 13$ $(1, 1) : 9$ $(1, 2) : 6$ $(1, 3) : 3$ $(2, 1) : 9$ $(2, 2) : 5$ $(2, 3) : 1$
$(6, 3, 9, 1)$	Not existing	16	12	$(1, \star) : 12$ $(2, 1) : 9$ $(2, 2) : 6$ $(2, 3) : 3$
$(6, 3, 125, 0)$	186	248	200	$(0, 3) : 187$ $(1, 1) : 125$ $(1, 2) : 75$ $(1, 3) : 25$ $(2, 1) : 125$ $(2, 2) : 65$ $(2, 3) : 5$ $(3, 1) : 125$ $(3, 2) : 63$ $(3, 3) : 1$
$(6, 3, 125, 1)$	248 (only for subtype $(0, 3, 0)$)	248	200	$(1, \star) : 185$ $(2, 1) : 125$ $(2, 2) : 75$ $(2, 3) : 2$ $(3, 1) : 125$ $(3, 2) : 65$ $(3, 3) : 5$
$(6, 3, 125, 2)$	310 (only for subtype $(0, 0, 3)$)	248	200	$(2, \star) : 175$ $(3, 1) : 125$

248 (only for sub-type (1, 1, 1))		(3, 2) : 75 (3, 3) : 25
-----------------------------------	--	----------------------------

TABLE 1. Comparison of the bounds on the minimum Lee distance of a code with given parameters.

Let us focus first on a free code, i.e., $\sigma = 0$). Observe, that if the last $n - K$ columns of a generator matrix consist only of nonunits, i.e., $\ell = 0$) the bound by Alderson and Huntemann beats our bounds. However, as soon as $\ell \neq 0$ the new bound based on the minimum distance of filtration subcodes (Corollary 61) always outperforms any other bound. In Table 1 we also observe, that the bound provided by Shiromoto is the loosest.

For nonfree codes, recall that the bound in [1] only works for integer $\mathbb{Z}/p^s\mathbb{Z}$ -dimensions $k > 1$. Furthermore, we note that for a given $\sigma \geq 1$ we always have $\ell \geq \sigma$ and if $\ell = \sigma$ the filtration bound (Corollary 61) is the same for any n' . This is denoted by $n' = \star$ in Table 1. In any of the parameters presented, the bound based on the minimum Lee distance of a filtration subcodes of the code (Corollary 61) outperforms all other bounds.

8. CONCLUSIONS AND FUTURE WORK

In this paper we presented several novel definitions of Lee support and the corresponding generalized Lee weights of subcodes of a fixed rank. These give raise to new Lee-metric Singleton bounds, that beat the previous bound by Shiromoto [31], which follows from a puncturing argument. However, their optimal codes are still sparse for s, n or p going to infinity. This led us to consider different subcodes, namely the filtration subcodes. Bounding their minimum distances gives a sharper Singleton bound, that finally has the desired property; their optimal codes are not sparse for s going to infinity.

The open question remains, whether there exists a bound on the minimum Lee distance of codes, which is such that their optimal codes are dense for s, n or p going to infinity.

ACKNOWLEDGEMENTS

Violetta Weger is supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 899987.

REFERENCES

- [1] Tim L Alderson and Svenja Huntemann. On maximum Lee distance codes. *Journal of Discrete Mathematics*, 2013, 2013.
- [2] Jared Antrobus and Heide Gluesing-Luerssen. Maximal Ferrers diagram codes: constructions and genericity considerations. *IEEE Transactions on Information Theory*, 65(10):6204–6223, 2019.
- [3] Alexander Barg and G David Forney. Random codes: Minimum distances and error exponents. *IEEE Transactions on Information Theory*, 48(9):2568–2573, 2002.

- [4] Eimear Byrne, Anna-Lena Horlemann, Karan Khathuria, and Violetta Weger. Density of free modules over finite chain rings. *Linear Algebra and its Applications*, 651:1–25, 2022.
- [5] Eimear Byrne and Alberto Ravagnani. Partition-balanced families of codes and asymptotic enumeration in coding theory. *Journal of Combinatorial Theory, Series A*, 171:105169, 2020.
- [6] Eimear Byrne and Violetta Weger. Bounds in the lee metric and optimal codes. *Finite Fields and Their Applications*, 87:102151, 2023.
- [7] Sara D Cardell, Marcelo Firer, and Diego Napp. Generalized column distances. *IEEE Transactions on Information Theory*, 66(11):6863–6871, 2020.
- [8] Philippe Delsarte. Bilinear forms over a finite field, with applications to coding theory. *Journal of combinatorial theory, Series A*, 25(3):226–241, 1978.
- [9] Steven T Dougherty. *Algebraic coding theory over finite commutative rings*. Springer, 2017.
- [10] Steven T Dougherty, M Gupta, and Keisuke Shiromoto. On generalized weights for codes over finite rings. *preprint*, 2002.
- [11] Steven T Dougherty and Keisuke Shiromoto. MDR codes over \mathbb{Z}_k . *IEEE Transactions on Information Theory*, 46(1):265–269, 2000.
- [12] Ernest Mukhamedovich Gabidulin. Theory of codes with maximum rank distance. *Problemy peredachi informatsii*, 21(1):3–16, 1985.
- [13] Heide Gluesing-Luerssen. On the sparseness of certain linear MRD codes. *Linear Algebra and its Applications*, 596:145–168, 2020.
- [14] Elisa Gorla and Alberto Ravagnani. Generalized weights of codes over rings and invariants of monomial ideals. *arXiv preprint arXiv:2201.05813*, 2022.
- [15] Elisa Gorla and Flavio Salizzoni. Generalized column distances, 2022.
- [16] Elisa Gorla and Flavio Salizzoni. Generalized weights of convolutional codes. *IEEE Transactions on Information Theory*, 2023.
- [17] Anina Gruica and Alberto Ravagnani. Common complements of linear subspaces and the sparseness of mrd codes. *SIAM Journal on Applied Algebra and Geometry*, 6(2):79–110, 2022.
- [18] A Roger Hammons, P Vijay Kumar, A Robert Calderbank, Neil JA Sloane, and Patrick Solé. The z_4 -linearity of Kerdock, Preparata, Goethals, and related codes. *IEEE Transactions on Information Theory*, 40(2):301–319, 1994.
- [19] Tor Helleseth, Torleiv Kløve, and Johannes Mykkeltveit. The weight distribution of irreducible cyclic codes with block lengths $n_1((q^\ell - 1)/n)$. *Discrete Mathematics*, 18(2):179–211, 1977.
- [20] Anna-Lena Horlemann-Trautmann and Violetta Weger. Information set decoding in the Lee metric with applications to cryptography. *Advances in Mathematics of Communications*, 15(4), 2021.
- [21] Yasuo Komamiya. Application of logical mathematics to information theory. *Proc. 3rd Japan. Nat. Cong. Appl. Math*, 437:3, 1953.
- [22] Jun Kurihara, Ryutaroh Matsumoto, and Tomohiko Uyematsu. Relative generalized rank weight of linear codes and its applications to network coding. *IEEE Transactions on Information Theory*, 61(7):3912–3936, 2015.
- [23] Jun Kurihara, Tomohiko Uyematsu, and Ryutaroh Matsumoto. New parameters of linear codes expressing security performance of universal secure network coding. In *2012 50th Annual Allerton Conference on Communication, Control,*

- and *Computing (Allerton)*, pages 533–540, 2012.
- [24] C Lee. Some properties of nonbinary error-correcting codes. *IRE Transactions on Information Theory*, 4(2):77–82, 1958.
 - [25] Pierre Loidreau. Asymptotic behaviour of codes in rank metric over finite fields. *Designs, codes and cryptography*, 71:105–118, 2014.
 - [26] Alessandro Neri, Anna-Lena Horlemann-Trautmann, Tovohery Randrianarisoa, and Joachim Rosenthal. On the genericity of maximum rank distance and Gabidulin codes. *Designs, Codes and Cryptography*, 86(2):341–363, 2018.
 - [27] Alberto Ravagnani. Generalized weights: an anticode approach. *Journal of Pure and Applied Algebra*, 220(5):1946–1962, 2016.
 - [28] Stefan Ritterhoff, Georg Maringer, Sebastian Bitzer, Violetta Weger, Patrick Karl, Thomas Schamberger, Jonas Schupp, and Antonia Wachter-Zeh. FuLeeca: A Lee-based signature scheme. *Cryptology ePrint Archive*, 2023.
 - [29] Ron M Roth. Maximum-rank array codes and their application to crisscross error correction. *IEEE transactions on Information Theory*, 37(2):328–336, 1991.
 - [30] Beniamino Segre. Curve razionali normali e k -archi negli spazi finiti. *Annali di Matematica Pura ed Applicata*, 39(1):357–379, 1955.
 - [31] Keisuke Shiromoto. Singleton bounds for codes over finite rings. *Journal of Algebraic Combinatorics*, 12(1):95–99, 2000.
 - [32] Richard Singleton. Maximum distance q -nary codes. *IEEE Transactions on Information Theory*, 10(2):116–118, 1964.
 - [33] Violetta Weger, Karan Khathuria, Anna-Lena Horlemann, Massimo Battaglioni, Paolo Santini, and Edoardo Persichetti. On the hardness of the Lee syndrome decoding problem. *Advances in Mathematics of Communications*, pages 0–0, 2022.
 - [34] Victor K Wei. Generalized hamming weights for linear codes. *IEEE Transactions on information theory*, 37(5):1412–1418, 1991.
 - [35] Jay A Wood. The structure of linear codes of constant weight. *Trans. American Math. Soc.*, 354(3):1007–1026, 2001.

¹INSTITUTE OF COMMUNICATION AND NAVIGATION, GERMAN AEROSPACE CENTER, GERMANY

²INSTITUTE OF MATHEMATICS, UNIVERSITY OF ZURICH, SWITZERLAND

Email address: `jessica.bariffi@dlr.de`

³DEPARTMENT OF ELECTRICAL AND COMPUTER ENGINEERING, TECHNICAL UNIVERSITY OF MUNICH, GERMANY

Email address: `violetta.weger@tum.de`