

Renormalization Group Evolution with Scalar Leptoquarks

Sumit Banik^{a,b} Andreas Crivellin^{a,b}

^a*Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

^b*Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland*

E-mail: sumit.banik@psi.ch, andreas.crivellin@psi.ch

ABSTRACT: Leptoquarks are theoretically well-motivated and have received increasing attention in recent years as they can explain several hints for physics beyond the Standard Model. In this article, we calculate the renormalisation group evolution of models with scalar leptoquarks. We compute the anomalous dimensions for all couplings (gauge, Yukawa, Higgs and leptoquarks interactions) of the most general Lagrangian at the two-loop level and the corresponding threshold corrections at one-loop. The most relevant analytic results are presented in the Appendix, while the notebook containing the full expressions can be downloaded at <https://github.com/SumitBanikGit/SLQ-RG>. In our phenomenological analysis, we consider some exemplary cases with focus on gauge and Yukawa coupling unification.

Contents

1	Introduction	1
2	Setup and Conventions	2
3	Renormalization Group Evolution and Phenomenological Analysis	4
4	Conclusions	10
A	Lagrangian terms involving SLQs	10
B	Two-loop β-functions of gauge couplings	13
C	One-Loop β-functions of SM Yukawa couplings	16
D	One-Loop threshold corrections	17

1 Introduction

The Standard Model (SM) of particle physics describes the known fundamental constituents of matter as well as their interactions. While the Higgs particle [1–4], last missing puzzle piece of the SM, was discovered in 2012 at the Large Hadron Collider (LHC) at CERN [5–7], it is clear that the SM cannot be the ultimate theory of Nature: It does not account for the astrophysical observation of Dark Matter nor for the non-vanishing neutrino masses required by neutrino oscillations.

A plethora of possible SM extensions have been proposed in the last decade. In this context, leptoquarks (LQs), i.e. hypothetical new particles which directly couple a quark to a lepton, are very interesting. They were first proposed within Grand Unified Theories (GUTs) [8–11] and composite models [12–15], and squarks can act as LQs within the R-parity violating MSSM (see e.g. Ref. [16] for a review). They were first classified in Ref. [17] into ten possible representations under the SM gauge group, of which five are scalars, and five are vector particles.

In recent years, there has been renewed interest in LQs as they give interesting effects in low energy observables in general [18–24], and could provide solutions to anomalies in semi-leptonic B decays [25–102], the anomalous magnetic moment of the muon (a_μ) [22, 46, 55, 58, 61, 71, 74, 75, 79, 80, 103–125] in particular. The effects of LQs at colliders [103, 126–150], in oblique electroweak parameters as well as Higgs couplings to gauge bosons [151–158], electric dipole moments [159, 160] and proton decay [161, 162] were studied. LQs

were considered as a portal to dark matter [109, 163–170] and as the origin of neutrino masses [58, 61, 77, 119, 125, 171–174]. Furthermore, they have been searched for at the LHC [175–194] resulting in bounds of the order of 1 – 2 TeV (in the absence or suppression of couplings to first generation quarks [23, 142]).

As LQs are possible (light) remnants of a GUT [195], it is interesting to assess their impact on gauge and Yukawa coupling unification. In particular, scalar LQs (SLQs) could be light states of a GUT symmetry-breaking sector [50] and it might even be possible that they are the only new particles (in addition to the SM) up to the GUT scale. In this case, they must alter the (renormalization group evaluation) RGE sufficiently to lead to the required coupling unification [196].

In this article, we will analyze the RGE in models with SLQs ¹. For this, we will calculate the two-loop anomalous dimensions as well as the one-loop threshold corrections (at the LQ scale) and apply these results to the study of gauge and Yukawa coupling unification. For this, the article is structured as follows: In Sec. 2, we present our setup and conventions. In Sec. 3, we derive the β -function, give their analytic expression for some simple cases and examine gauge and Yukawa coupling unification for some specific examples. In Sec. 4, we summarize and discuss the results of this paper. In Appendix A, we give the full SLQ lagrangian, including all five possible SLQs. In Appendix B and C, we give the anomalous dimensions for the gauge (two-loop) and Yukawa couplings (one-loop), respectively. In Appendix D, we collect the one-loop threshold corrections of SM parameters on matching SLQ models with the SM. A notebook with our full results, including Higgs and LQ (self-) interactions etc., is available at [197].

2 Setup and Conventions

SM fields	Q_j	L_j	u_j	d_j	ℓ_j	H
$U(1)_Y, SU(2)_L, SU(3)_c$	$+\frac{1}{6}, \mathbf{2}, \mathbf{3}$	$-\frac{1}{2}, \mathbf{2}, \mathbf{1}$	$+\frac{2}{3}, \mathbf{1}, \mathbf{3}$	$-\frac{1}{3}, \mathbf{1}, \mathbf{3}$	$-1, \mathbf{1}, \mathbf{1}$	$+\frac{1}{2}, \mathbf{2}, \mathbf{1}$
Leptoquark	Φ_1	$\Phi_{\bar{1}}$	Φ_2	$\Phi_{\bar{2}}$	Φ_3	
$U(1)_Y, SU(2)_L, SU(3)_c$	$-\frac{1}{3}, \mathbf{1}, \mathbf{3}$	$-\frac{4}{3}, \mathbf{1}, \mathbf{3}$	$+\frac{7}{6}, \mathbf{2}, \mathbf{3}$	$+\frac{1}{6}, \mathbf{2}, \mathbf{3}$	$-\frac{1}{3}, \mathbf{3}, \mathbf{3}$	

Table 1: Representations of the SM and LQ fields under the $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge group.

The SM fields transform under the $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge group as given in Table 1, where $j = 1, 2, 3$ is a flavour index. The left-handed fermions Q_j and L_j are

¹As for vector LQs a Higgs mechanism is necessary to make the model renormalizable, one can not study the RGEs in a simplified model approach.

doublets under $SU(2)_L$ and decompose into their components as

$$Q_f = \begin{pmatrix} u_{j,L} \\ d_{j,L} \end{pmatrix}, \quad L_f = \begin{pmatrix} \nu_{j,L} \\ \ell_{j,L} \end{pmatrix}, \quad (2.1)$$

while the right-handed fields u_j , d_j and ℓ_j are $SU(2)_L$ singlets which we write as

$$u_j = \begin{pmatrix} u_{j,R} \end{pmatrix}, \quad d_j = \begin{pmatrix} \nu_{j,R} \end{pmatrix}, \quad \ell_j = \begin{pmatrix} \ell_{j,R} \end{pmatrix}. \quad (2.2)$$

The electric charge q is given by the Gell-Mann-Nishijima formula, $q = Y + I_3$ where, I_3 is the third-component of the weak isospin and Y is the hypercharge related to the gauge factor $U(1)_Y$.

The SM Lagrangian is then written as

$$\begin{aligned} \mathcal{L}_{\text{SM}} = & -\frac{1}{4}B^{\mu\nu}B_{\mu\nu} - \frac{1}{4}W^{I,\mu\nu}W_{I,\mu\nu} - \frac{1}{4}G^{\alpha,\mu\nu}G_{\alpha,\mu\nu} \\ & + i(\bar{Q}_f \not{D} Q_f) + i(\bar{L}_f \not{D} L_f) + i\bar{u}_f \not{D} u_f + i\bar{d}_f \not{D} d_f + i\bar{\ell}_f \not{D} \ell_f \\ & + ((D^\mu H)^\dagger D_\mu H) + \mu_H^2 (H^\dagger H)^2 - \lambda_H (H^\dagger H)^2 \\ & - (Y_{f_i f_j}^d (\bar{Q}_{f_i} H) d_{f_j} + Y_{f_i f_j}^\ell (\bar{L}_{f_i} H) \ell_{f_j} + Y_{f_i f_j}^u (\bar{Q}_{f_i} i\sigma_2 H^\dagger) u_{f_j} + \text{h.c.}) \\ & + \mathcal{L}_{\text{gauge-fixing}}, \end{aligned} \quad (2.3)$$

where Y^d , Y^ℓ and Y^u are the Yukawa couplings and $B^{\mu\nu}$ and the field strength tensors defined as,

$$\begin{aligned} B_{\mu\nu} &= \partial_\mu B_\nu - \partial_\nu B_\mu, \\ W_{\mu\nu}^I &= \partial_\mu W_\nu^I - \partial_\nu W_\mu^I + g_2 f^{IJK} W_\mu^J W_\nu^K, \quad I \in \{1, 2, 3\}, \\ G_{\mu\nu}^\alpha &= \partial_\mu G_\nu^\alpha - \partial_\nu G_\mu^\alpha + g_3 f^{\alpha\beta\gamma} G_\mu^\beta G_\nu^\gamma, \quad \alpha \in \{1, \dots, 8\}, \end{aligned} \quad (2.4)$$

corresponding to the gauge groups $U(1)_Y$, $SU(2)_L$ and $SU(3)_c$, respectively. The structure constants of $SU(2)_L$ and $SU(3)_c$ are denoted by f^{IJK} and $f^{\alpha\beta\gamma}$. For better readability, $SU(2)_L$ indices inside parenthesis (\cdots) are contracted such that they form gauge singlets. The covariant derivative for any field ϕ is defined as

$$D_\mu \phi = \partial_\mu \phi - i\tilde{g}_1 Y B_\mu \phi - ig_2 \tau_I W_\mu^I \phi - ig_3 T_\alpha G_\mu^\alpha \phi, \quad (2.5)$$

where Y is the $U(1)_Y$ hypercharge of ϕ . \tilde{g}_1 , g_2 and g_3 are the gauge couplings associated with $U(1)_Y$, $SU(2)_L$ and $SU(3)_c$ gauge groups, respectively. τ_I and T_α are the generators of $SU(2)_L$ and $SU(3)_c$ depending on the representation of ϕ . In the fundamental representation of $SU(2)_L$ and $SU(3)_c$ we have $\tau_I = \frac{\sigma_I}{2}$ and $T_\alpha = \frac{\lambda_\alpha}{2}$, where σ_I are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (2.6)$$

and λ_α are the Gell-Mann matrices.

We now add to the SM the five scalar SLQs which transform under the SM gauge groups as given in Table 1. For later convenience, we decompose the SLQ Lagrangian as

$$\begin{aligned}\mathcal{L}_{\text{LQ}} = & \mathcal{L}_1 + \mathcal{L}_{\bar{1}} + \mathcal{L}_2 + \mathcal{L}_{\bar{2}} + \mathcal{L}_3 + \mathcal{L}_{1\bar{1}} + \mathcal{L}_{12} + \mathcal{L}_{1\bar{2}} + \mathcal{L}_{13} \\ & + \mathcal{L}_{\bar{1}2} + \mathcal{L}_{\bar{1}\bar{2}} + \mathcal{L}_{\bar{1}3} + \mathcal{L}_{2\bar{2}} + \mathcal{L}_{23} + \mathcal{L}_{\bar{2}3} \\ & + \mathcal{L}_{\bar{1}2\bar{2}} + \mathcal{L}_{1\bar{1}2} + \mathcal{L}_{123} + \mathcal{L}_{1\bar{2}3} + \mathcal{L}_{\bar{1}23} \\ & + \mathcal{L}_{\bar{1}2\bar{2}3} + \mathcal{L}_{1\bar{1}2\bar{2}},\end{aligned}\tag{2.7}$$

where, the terms $\mathcal{L}_{ij\dots}$ contain terms involving only the LQs $\{\Phi_i, \Phi_j, \dots\}$. The explicit expressions, as already presented in Ref. [198], for each Lagrangian term are given in Appendix A. Note that for our purpose, it is imperative to have the complete Lagrangian², e.g. terms involving LQ-Higgs interactions, such that the full set of counterterms needed to cancel all generated divergences is available.

3 Renormalization Group Evolution and Phenomenological Analysis

We calculate the complete two-loop β -functions and the one-loop threshold corrections of all couplings of our LQ Lagrangian in the $\overline{\text{MS}}$ -scheme. For the anomalous dimensions, we used the Python package PyR@TE [199].³ The PyR@TE model files and full analytic expressions for the β -functions, as well as the threshold corrections, are available online publicly on Github [197]. The one-loop threshold corrections are presented in Appendix D, which were derived using Matchete [201] and cross-checked for some selected cases with `matchmakereft` [202]. We also checked that including the threshold corrections reduce, as expected, the matching scale dependence. The QCD contribution to some LQ Yukawa couplings was cross-checked with Ref. [203].

We use the following convention throughout this paper for the β -function for any parameter A

$$\beta(A) \equiv \mu \frac{dA}{d\mu} \equiv \frac{1}{(4\pi)^2} \beta^{(1)}(A) + \frac{1}{(4\pi)^4} \beta^{(2)}(A) + \dots\tag{3.1}$$

where μ is the renormalization scale and $\beta^{(n)}(A)$ denotes the contribution at n -loop order and the dots are the higher-order terms. The β -functions form a system of coupled differential equations which we solve numerically, as explained below.

At the one-loop level, we have the following coefficients for the gauge couplings

$$\begin{aligned}\beta^{(1)}(g_1) &= \frac{41}{10} g_1^3 + \left(\frac{n_1}{15} + \frac{16n_{\bar{1}}}{15} + \frac{49n_2}{30} + \frac{n_{\bar{2}}}{30} + \frac{n_3}{5} \right) g_1^3, \\ \beta^{(1)}(g_2) &= -\frac{19}{6} g_2^3 + \left(\frac{n_2}{2} + \frac{n_{\bar{2}}}{2} + 2n_3 \right) g_2^3,\end{aligned}$$

²We disregard B -number violating couplings as they must be small due to constraints from proton decay and therefore will have a small effect on RGEs.

³We cross-checked our results using the *Mathematica* package `RGBeta` [200].




Figure 1: Renormalization group evolution of the gauge couplings ($\alpha_i = g_i^2/4\pi$) in SM and its extensions with the different LQ representations. In order to better visualize the effect of adding LQ, we considered three generations of each LQ of mass 3 TeV. For simplicity, we assumed all Yukawa and Higgs couplings involving LQs to be zero (at the LQ scale).

$$\beta^{(1)}(g_3) = -7g_3^3 + \left(\frac{n_1}{6} + \frac{n_{\tilde{1}}}{6} + \frac{n_2}{3} + \frac{n_{\tilde{2}}}{3} + \frac{n_3}{2}\right)g_3^3, \quad (3.2)$$

where, n_1 , $n_{\tilde{1}}$, n_2 , $n_{\tilde{2}}$ and n_3 denote the number of copies, i.e. generations, of Φ_1 , $\Phi_{\tilde{1}}$, Φ_2 , $\Phi_{\tilde{2}}$ and Φ_3 , respectively. The first term is the well-known SM term, and we have used the $SU(5)$ GUT normalization $g_1 = \sqrt{\frac{5}{3}}\tilde{g}_1$ w.r.t. the values given in Table 1. The one-loop expressions for the β -function for the Yukawa couplings and the corresponding threshold correction are provided in Appendixes C and D, respectively. The two-loop β -function coefficients for the gauge couplings are given in Appendix B.

With these results at hand, we can next study the phenomenology of some selected cases. Here, we focus primarily on the evolution of the gauge and Higgs Yukawa couplings from the electroweak scale to the GUT scale.⁴ As the starting point for the evolution, we take the following initial values of the SM parameters at the top scale, above the top

⁴Note that without a full specification of the GUT theory and its breaking sector, we cannot calculate the one-loop threshold corrections at the GUT scale which would be necessary to obtain a scheme and scale independent result. However, as no large logarithms are involved here, and the gauge coupling is already significantly smaller than the strong coupling at the EW scale, we assume these effects as uncertainties in the calculation.




Figure 2: Left: Two-loop renormalization group evolution of gauge couplings for SM+ Φ_3 . We observe gauge unification at around 10¹⁴ GeV when we set the LQ scale to $m_3 \approx 10^6$ TeV. We vary the initial value of the three diagonal components of Y_3^{LL} between -0.6 to 0.6 (keeping the initial value of all other couplings to zero) to show that their effect on running is minimal. Right: Two-loop renormalization group running of gauge couplings for SM+ $\Phi_2 + \Phi_3$. We observe gauge unification for $m_2 = 3$ TeV and $m_3 = 4 \times 10^3$ TeV.

threshold (i.e. with 6 active flavours) in the $\overline{\text{MS}}$ -scheme [204–238]

$$g_1 = 0.3585 \times \sqrt{\frac{5}{3}}, \quad g_2 = 0.648, \quad g_3 = 1.16, \\ y_t = 0.935, \quad y_b = 0.015, \quad y_\tau = 0.01, \quad \lambda = 0.126. \quad (3.3)$$

We evolve the couplings using the β -functions of the SM up to the LQ scale m_x (with $x = 1, \tilde{1}, 2, \tilde{2}, 3$). Note that here we neglected the tiny Yukawa couplings of the first two generations.

After evolving the couplings to the LQ scale, we include the one-loop threshold corrections determined by comparing the theory with SLQs, to the one without them (i.e. the SM at the LQ scale). This gives a shift to the SM fermion Yukawa couplings depending mainly on the initial values (at the LQ scale) of the LQ Yukawa couplings.⁵ We then run all couplings, the SM ones as well as the ones including LQs, using the β -functions of the full model from the LQ scale to the high scale, for which we take as an upper limit the Planck scale ($\approx 10^{19}$) where gravitational effects would become important.

As a first step, we illustrate the running of the gauge coupling as a function of the scale Q in Fig. 1 for the five different LQ representations for a fixed LQ scale of 3 TeV, which is compatible with direct LHC searches [175–194]. To enhance and highlight the impact of adding the LQs, we considered three generations of the same LQ representation.

We observe that no single generation LQ representation at the TeV scale can lead to gauge coupling unification. One can understand this behaviour of the gauge couplings from

⁵Note that for our SLQ models, the gauge couplings do not receive threshold corrections at one-loop in the $\overline{\text{MS}}$ -scheme.

Eq. (3.2). Since $\beta^{(1)}(g_i) = a_i g_i^3$, where a_i is a constant, $\alpha^{-1} = 4\pi g_i^{-2}$ and the slope of the curve is $\alpha_i^{-1}(\log(\mu))$ is $-8\pi a_i$. We also see that w.r.t. the SM curve (black) in Fig. 1, gauge coupling unification is improved if the slope of α_1^{-1} increases (i.e a_1 decreases) or if the slope of α_2^{-1} and α_3^{-1} decreases (i.e a_2 and a_3 increases). For Φ_1 and $\Phi_{\bar{1}}$, a_2 remains unchanged but a_1 and a_3 increase. Therefore, Φ_1 and $\Phi_{\bar{1}}$ will worsen the unification of gauge couplings. However, for the other three SLQs, both a_2 and a_3 increase, which improves the unification, despite a_1 also increasing. Specifically, $\Phi_{\bar{2}}$ and Φ_3 improve the unification most since the increase in a_1 is much smaller. While the effect for $\Phi_{\bar{2}}$ is too weak, Φ_3 can lead to gauge coupling unification at $\approx 10^{14}$ GeV for an LQ mass of around 10^6 TeV, as shown in Fig. 2 (left).⁶ Furthermore, gauge coupling unification is possible if we added three generations of $\Phi_{\bar{2}}$ as shown in Fig. 1.

Because the LQ Yukawa, Higgs and self-couplings have a minimal effect on the running of gauge couplings, we take their initial value to be zero at the LQ scale. In Fig. 2 (left), we also show the impact of including the LQ Yukawa couplings on SM fermions (which are free parameters). One can see that for the gauge coupling evolution, their effect is suppressed since they only enter at the two-loop level. In fact, we scanned over the initial values of all the three diagonal elements of Y_3^{LL} between -0.6 and 0.6 (while keeping the initial value of other couplings to zero), leading to a slight thickening of the curves.

However, for larger values of the LQ Yukawa couplings non-perturbative values for some couplings, in particular quartic scalar couplings, can be induced. If one requires that this does not happen below a given scale, e.g. the GUT scale, one can set an upper limit on the Yukawa couplings. We show the illustrative example of Φ_3 in Fig. 3 where we plot three different quartic couplings for three large initial values (at $m_3 \approx 10^4$ TeV) of the [3, 3] component of Y_3^{LL} . One can see that the higher the initial value of the LQ Yukawa coupling, the lower the energy at which the quartic couplings reach non-perturbative values. Constraints on certain LQ Yukawa couplings were studied in [240].

We now illustrate the RGE of bottom and tau Yukawa couplings in Fig. 4. While in the SM, bottom-tau Yukawa coupling unification already happens around 10^6 GeV, in LQ models, all representations have a small impact on the RGE of bottom Yukawa. Furthermore, the same is true for the tau one, except for Φ_1 and Φ_2 . For them, the one-loop threshold corrections to the tau Yukawa coupling can be sizable and their running strongly changes (w.r.t. the SM).⁷ The running of the tau Yukawa coupling of SM+ Φ_1 and SM+ Φ_2 are shown in Fig. 4. The tau threshold correction strongly depends on our choice of the initial value of (3, 3) component of Y_1^{RR} , Y_1^{LL} , Y_2^{RL} and Y_2^{LR} which is set

⁶Such a low unification scale would conflict with the limits from proton decay [239], at least for a standard $SU(5)$ GUT.

⁷In this case, also a chirally enhanced effect in the anomalous magnetic moment of the τ lepton is generated [241]. This means that the chirality flips necessary to generate the anomalous magnetic moment does not originate from the rather small tau Yukawa coupling in the SM, but rather from a large coupling of new particles to the Higgs boson [242].




Figure 3: Two-loop renormalization group running of the quartic couplings $Y_3^{(1)}$ (solid), $Y_3^{(3)}$ (dashed) and $Y_3^{(5)}$ (dotted) in SM+ Φ_3 for different initial values at $m_3 \approx 10^4$ TeV of the [3, 3] component of the LQ Yukawa Y_3^{LL} while assuming the other components to be small (as suggested by the B anomalies). We observe that for larger (absolute) initial values of the LQ Yukawa, the quartic couplings become non-perturbative at lower scales.

to 0.105 for dotted lines and the sign of Y_1^{LL} and Y_2^{LR} reversed for the solid lines. The initial value of the remaining LQ couplings was set to zero. In fact, as can be seen in Fig. 4, the trajectories all lie within the quite narrow blue band except for, Φ_1 and Φ_2 . Here a very strong impact on the evaluation of the tau Yukawa is possible due to the chiral enhancement.

We next consider the case, motivated by $SU(5)$ GUTs [50], where SM is extended by Φ_2 and Φ_3 . Here, we find that for $m_2 \approx 3$ TeV and $m_3 \approx 10^{3.5}$ TeV, we can achieve gauge coupling unification at $\approx 10^{13}$ TeV as shown in Fig. 2 (right) as well as Yukawa coupling unification around the same energy scale. However, this scale is again naively in conflict with proton decay. Finally, we consider the case where we extend the SM with (a single representation) all five possible SLQs (see Table 1). In this case, we obtain gauge coupling unification at around 10^{13} TeV for a common LQ scale of 125 TeV as shown in Fig. 5 (top). We also show in Fig. 5 (bottom) that bottom-tau Yukawa coupling unification is possible nearly at the same energy scale for the following initial values of the (3, 3) component of the following LQ Yukawa couplings at the LQ scale

$$Y_1^{RR}_{[3,3]} = 0.1, Y_1^{LL}_{[3,3]} = -0.1, Y_1^{RL}_{[3,3]} = -0.04, Y_1^{LR}_{[3,3]} = 0.1, Y_3^{LL}_{[3,3]} = 0.45, \quad (3.4)$$

while taking the initial value of other LQ couplings to be zero.




Figure 4: Two-loop renormalization group running of bottom and tau Yukawa couplings in SM+ Φ_1 and SM+ Φ_2 for a single-generation SLQ of mass 3 TeV. The dotted line denotes the tau-Yukawa running when we set the initial value of the (3, 3) components of Y_1^{RR} , Y_1^{LL} , Y_2^{RL} and Y_2^{LR} to 0.1. For solid lines, we reverse the sign of the (3, 3) components of Y_1^{LL} and Y_2^{LR} . The blue bands show the variation of running in SM+ Φ_i , for $i = 1, \tilde{1}, 2, \tilde{2}, 3$ for bottom Yukawa and $i = \tilde{1}, \tilde{2}, 3$ for tau Yukawa, while varying the (3, 3) components of LQ Yukawa couplings.




Figure 5: Two-loop renormalization group running of gauge couplings (left), and bottom and tau Yukawa couplings (right) for SM+ $\Phi_1+\Phi_{\tilde{1}}+\Phi_2+\Phi_{\tilde{2}}+\Phi_3$. We observe gauge coupling unification and bottom-tau unification at around the same energy scale $\approx 10^{13}$ GeV when we set the mass of all LQs to ≈ 125 TeV and take the non-zero initial values of LQ couplings given in Eq. (3.4).

4 Conclusions

LQs are well-motivated extensions of the SM. They arise in composite or extra-dimensional setups and, most importantly, are predicted by GUTs. Furthermore, they have been under intensified investigation in the last years as they are potential candidates for describing several tensions between SM measurements and experiments.

In this article, we computed the two-loop renormalization group evolution as well as the one-loop threshold corrections, for all parameters within generic SLQ models. This includes gauge couplings, SM Yukawa couplings, LQ Yukawa couplings (to quarks and leptons), Higgs and LQ (self-)interactions. The appendices collect the full SLQ Lagrangian, two-loop β -functions of the gauge couplings, one-loop threshold corrections and one-loop β -functions for the SM Yukawa couplings. The full analytic expressions, together with the necessary model files, can be obtained from [197].

In our phenomenological analysis, we considered on the case in which one or more SLQs are light remnants of a symmetry-breaking sector of a GUT. In this setup, we focused on the renormalisation group evolution of the gauge and the Yukawa couplings, examining if unification can be achieved. Several simple scenarios were studied: 1) If one adds one of the 5 possible LQ representations to the SM, only Φ_3 (with a mass around 10^6 TeV) can lead to gauge coupling unification at $\approx 10^{15}$ GeV. 2) Extending the SM by all 5 possible LQ representations with a common mass scale ≈ 150 TeV, unification at $\approx 10^{14}$ GeV is achieved. 3) Adding the GUT-motivated LQs Φ_2 and Φ_3 to the SM, with masses of $m_2 = 3$ TeV and $m_3 = 4 \times 10^3$ TeV, respectively, gauge coupling unification occurs at $\approx 10^{13}$ GeV. 4) Concerning the bottom-tau Yukawa unification, which happens at far too low scales in the SM, only the LQs Φ_1 and Φ_2 can lead to a sizable modification, due to possible chiral enhancement. Therefore, by choosing the initial value of LQ Yukawa couplings properly, one can always achieve bottom-tau unification at any desired (high) scale.

Acknowledgements

We would like to thank Lohan Sartore for help in using `PyR@TE` and Anders E. Thomsen for help in using `RGBeta` and `Matchete`. We thank Michael Spira for useful comments on the draft. The work of A.C. is supported by a professorship grant from the Swiss National Science Foundation (No. PP00P21_76884).

A Lagrangian terms involving SLQs

This appendix collects the terms in the SLQ Lagrangian in Eq. (2.7) following the convention in [198]. We begin with the terms \mathcal{L}_1 , $\mathcal{L}_{\bar{1}}$, \mathcal{L}_2 , $\mathcal{L}_{\bar{2}}$ and \mathcal{L}_3 which involve only single LQ interactions that include kinetic terms, mass terms, Yukawa interaction with SM fermions,

quartic interaction with Higgs field and self-quartic interactions.

$$\begin{aligned} \mathcal{L}_1 = & (D_\mu \Phi_1)^\dagger D^\mu \Phi_1 - m_1^2 \Phi_1^\dagger \Phi_1 - Y_1 \left(H^\dagger H \right) \Phi_1^\dagger \Phi_1 + \left[Y_{1,ij}^{RR} \tilde{u}_i^c \ell_j \Phi_1^\dagger + Y_{1,ij}^{LL} \left(\tilde{Q}_i^{c\tau} i\sigma_2 L_j \right) \Phi_1^\dagger \right. \\ & \left. + Y_{1,ij}^{Q,LL} \left(\tilde{Q}_{i,c_1}^{c\tau} i\sigma_2 Q_{j,c_2} \right) \Phi_{1,c_3} \epsilon^{c_1 c_2 c_3} + Y_{1,ij}^{Q,RR} \tilde{u}_{i,c_1}^c d_{j,c_2} \Phi_{1,c_3} \epsilon^{c_1 c_2 c_3} + \text{h.c.} \right] \\ & + \frac{1}{2} Y_1^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_1} \right) \left(\Phi_{1,c_2}^\dagger \Phi_{1,c_2} \right) \end{aligned} \quad (\text{A.1})$$

$$\begin{aligned} \mathcal{L}_{\tilde{1}} = & (D_\mu \Phi_{\tilde{1}})^\dagger D^\mu \Phi_{\tilde{1}} - m_{\tilde{1}}^2 \Phi_{\tilde{1}}^\dagger \Phi_{\tilde{1}} - Y_{\tilde{1}} \left(H^\dagger H \right) \Phi_{\tilde{1}}^\dagger \Phi_{\tilde{1}} + \left[Y_{\tilde{1},ij}^{RR} \tilde{d}_i^c \ell_j \Phi_{\tilde{1}}^\dagger \right. \\ & \left. + Y_{\tilde{1},ij}^{Q,RR} \tilde{u}_{i,c_1}^c u_{j,c_2} \Phi_{\tilde{1},c_3} \epsilon^{c_1 c_2 c_3} + \text{h.c.} \right] + \frac{1}{2} Y_{\tilde{1}}^{(1)} \left(\Phi_{\tilde{1},c_1}^\dagger \Phi_{\tilde{1},c_1} \right) \left(\Phi_{\tilde{1},c_2}^\dagger \Phi_{\tilde{1},c_2} \right) \end{aligned} \quad (\text{A.2})$$

$$\begin{aligned} \mathcal{L}_2 = & (D_\mu \Phi_2)^\dagger D^\mu \Phi_2 - m_2^2 \Phi_2^\dagger \Phi_2 - Y_2 \left(H^\dagger H \right) \Phi_2^\dagger \Phi_2 + \left[Y_{2,ij}^{RL} \left(\Phi_2^\dagger \tilde{u}_i^c i\sigma_2 L_j \right) + Y_{2,ij}^{LR} \left(\tilde{Q}_i \ell_j \Phi_2 \right) \right. \\ & \left. + \text{h.c.} \right] + \frac{1}{2} Y_2^{(1)} \left(\Phi_{2,c_1}^\dagger \Phi_{2,c_1} \right) \left(\Phi_{2,c_2}^\dagger \Phi_{2,c_2} \right) + \frac{1}{2} Y_2^{(3)} \left(\Phi_{2,c_1}^\dagger \Phi_{2,c_2} \right) \left(\Phi_{2,c_2}^\dagger \Phi_{2,c_1} \right) \\ & - Y_{22} \left(H^\dagger i\sigma_2 \Phi_2 \right)^\dagger \left(H^\dagger i\sigma_2 \Phi_2 \right) \end{aligned} \quad (\text{A.3})$$

$$\begin{aligned} \mathcal{L}_{\tilde{2}} = & (D_\mu \Phi_{\tilde{2}})^\dagger D^\mu \Phi_{\tilde{2}} - m_{\tilde{2}}^2 \Phi_{\tilde{2}}^\dagger \Phi_{\tilde{2}} - Y_{\tilde{2}} \left(H^\dagger H \right) \Phi_{\tilde{2}}^\dagger \Phi_{\tilde{2}} + \left[Y_{\tilde{2},ij}^{RL} \left(\Phi_{\tilde{2}}^\dagger \tilde{d}_i^c i\sigma_2 L_j \right) + \text{h.c.} \right] \\ & + \frac{1}{2} Y_{\tilde{2}}^{(1)} \left(\Phi_{\tilde{2},c_1}^\dagger \Phi_{\tilde{2},c_1} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_2} \right) + \frac{1}{2} Y_{\tilde{2}}^{(3)} \left(\Phi_{\tilde{2},c_1}^\dagger \Phi_{\tilde{2},c_2} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_1} \right) \\ & - Y_{\tilde{2}\tilde{2}} \left(H^\dagger i\sigma_2 \Phi_{\tilde{2}} \right)^\dagger \left(H^\dagger i\sigma_2 \Phi_{\tilde{2}} \right) \end{aligned} \quad (\text{A.4})$$

$$\begin{aligned} \mathcal{L}_3 = & (D_\mu \Phi_3)^\dagger D^\mu \Phi_3 - m_3^2 \Phi_3^\dagger \Phi_3 - Y_3 \left(H^\dagger H \right) \Phi_3^\dagger \Phi_3 + \left[Y_{3,ij}^{LL} \left(\tilde{Q}_i^{c\tau} i\sigma_2 (\sigma \cdot \Phi_3)^\dagger L_j \right) \right. \\ & \left. + Y_{3,ij}^{Q,LL} \left(\tilde{Q}_{i,c_1}^{c\tau} i\sigma_2 (\sigma \cdot \Phi_{3,c_3}) Q_{j,c_2} \right) \epsilon^{c_1 c_2 c_3} + \text{h.c.} \right] + \frac{1}{2} Y_3^{(1)} \left(\Phi_{3,c_1}^\dagger \Phi_{3,c_1} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_2} \right) \\ & + \frac{1}{2} Y_3^{(3)} \left(\Phi_{3,c_1}^\dagger \Phi_{3,c_2} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_1} \right) - i Y_{33} \epsilon^{IJK} \left(H^\dagger \sigma^I H \right) \Phi_3^{J\dagger} \Phi_3^K \\ & + \frac{1}{2} Y_3^{(5)} \left(\Phi_{3,c_1}^{I\dagger} \Phi_{3,c_1}^J \Phi_{3,c_2}^{I\dagger} \Phi_{3,c_2}^J \right) \end{aligned} \quad (\text{A.5})$$

where we have omitted, at several places, color indices involving trivial contraction. For example, $Y_1 (H^\dagger H) \Phi_1^\dagger \Phi_1$ in full notation is $Y_1 (H^\dagger H) \Phi_{1,c_1}^\dagger \Phi_{1,c_1}$.

The next set of terms $\mathcal{L}_{1\tilde{1}}$, \mathcal{L}_{12} , $\mathcal{L}_{1\tilde{2}}$, \mathcal{L}_{13} , $\mathcal{L}_{\tilde{1}2}$, $\mathcal{L}_{\tilde{1}\tilde{2}}$, $\mathcal{L}_{\tilde{1}3}$, $\mathcal{L}_{2\tilde{2}}$, \mathcal{L}_{23} and $\mathcal{L}_{\tilde{2}3}$ contain interactions involving precisely two LQs.

$$\mathcal{L}_{1\tilde{1}} = Y_{1\tilde{1}}^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_1} \right) \left(\Phi_{\tilde{1},c_2}^\dagger \Phi_{\tilde{1},c_2} \right) + Y_{1\tilde{1}}^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_2} \right) \left(\Phi_{\tilde{1},c_2}^\dagger \Phi_{\tilde{1},c_1} \right) \quad (\text{A.6})$$

$$\mathcal{L}_{12} = Y_{12}^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_1} \right) \left(\Phi_{2,c_2}^\dagger \Phi_{2,c_2} \right) + Y_{12}'^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_2} \right) \left(\Phi_{2,c_2}^\dagger \Phi_{2,c_1} \right) \quad (\text{A.7})$$

$$\begin{aligned} \mathcal{L}_{1\tilde{2}} = & Y_{1\tilde{2}}^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_1} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_2} \right) + Y_{1\tilde{2}}'^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_2} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_1} \right) \\ & + \left[A_{1\tilde{2}\tilde{2}} \Phi_{1,c_1} \left(\Phi_{\tilde{2},c_2}^\dagger i\sigma_2 \Phi_{\tilde{2},c_3} \right) \epsilon^{c_1 c_2 c_3} - A_{1\tilde{2}} \Phi_1 \left(\Phi_{\tilde{2}}^\dagger H \right) + \text{h.c.} \right] \end{aligned} \quad (\text{A.8})$$

$$\begin{aligned} \mathcal{L}_{13} = & Y_{13}^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_1} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_2} \right) + Y_{13}'^{(1)} \left(\Phi_{1,c_1}^\dagger \Phi_{1,c_2} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_1} \right) + \left[Y_{13} (H^\dagger (\sigma \cdot \Phi_3) H) \Phi_1^\dagger \right. \\ & \left. + \frac{1}{2} Y_{1313} \left(\Phi_{1,c_1}^\dagger \Phi_{3,c_1}^I \Phi_{1,c_2}^\dagger \Phi_{3,c_2}^I \right) + Y_{1333} \left(\Phi_{1,c_1}^\dagger \Phi_{3,c_1}^I \Phi_{3,c_2}^{J\dagger} \Phi_{3,c_2}^K i\epsilon^{IJK} \right) + \text{h.c.} \right] \end{aligned} \quad (\text{A.9})$$

$$\mathcal{L}_{\tilde{1}2} = Y_{\tilde{1}2}^{(1)} \left(\Phi_{\tilde{1},c_1}^\dagger \Phi_{\tilde{1},c_1} \right) \left(\Phi_{2,c_2}^\dagger \Phi_{2,c_2} \right) + Y_{\tilde{1}2}'^{(1)} \left(\Phi_{\tilde{1},c_1}^\dagger \Phi_{\tilde{1},c_2} \right) \left(\Phi_{2,c_2}^\dagger \Phi_{2,c_1} \right) \quad (\text{A.10})$$

$$\mathcal{L}_{\tilde{1}\tilde{2}} = Y_{\tilde{1}\tilde{2}}^{(1)} \left(\Phi_{\tilde{1},c_1}^\dagger \Phi_{\tilde{1},c_1} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_2} \right) + Y_{\tilde{1}\tilde{2}}'^{(1)} \left(\Phi_{\tilde{1},c_1}^\dagger \Phi_{\tilde{1},c_2} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_1} \right) \quad (\text{A.11})$$

$$\begin{aligned} \mathcal{L}_{\tilde{1}3} = & Y_{\tilde{1}3}^{(1)} \left(\Phi_{\tilde{1},c_1}^\dagger \Phi_{\tilde{1},c_1} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_2} \right) + Y_{\tilde{1}3}'^{(1)} \left(\Phi_{\tilde{1},c_1}^\dagger \Phi_{\tilde{1},c_2} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_1} \right) \\ & + \left[Y_{\tilde{1}3} (H^\dagger i\sigma_2 (\sigma \cdot \Phi_3)^\dagger H) \Phi_{\tilde{1}} + \text{h.c.} \right] \end{aligned} \quad (\text{A.12})$$

$$\begin{aligned} \mathcal{L}_{2\tilde{2}} = & Y_{2\tilde{2}}^{(1)} \left(\Phi_{2,c_1}^\dagger \Phi_{2,c_1} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_2} \right) + Y_{2\tilde{2}}'^{(1)} \left(\Phi_{2,c_1}^\dagger \Phi_{2,c_2} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{\tilde{2},c_1} \right) \\ & + Y_{2\tilde{2}}^{(3)} \left(\Phi_{2,c_1}^\dagger \Phi_{\tilde{2},c_1} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{2,c_2} \right) + Y_{2\tilde{2}}'^{(3)} \left(\Phi_{2,c_1}^\dagger \Phi_{\tilde{2},c_2} \right) \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{2,c_1} \right) \\ & + \left[Y_{2\tilde{2}} \left(\Phi_2^\dagger H \right) (H^\dagger i\sigma_2 \Phi_{\tilde{2}}) + \text{h.c.} \right] \end{aligned} \quad (\text{A.13})$$

$$\begin{aligned} \mathcal{L}_{23} = & Y_{23}^{(1)} \left(\Phi_{2,c_1}^\dagger \Phi_{2,c_1} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_2} \right) + Y_{23}'^{(1)} \left(\Phi_{2,c_1}^\dagger \Phi_{2,c_2} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_1} \right) \\ & + Y_{23}^{(3)} \left(\Phi_{2,c_1}^\dagger \sigma^I \Phi_{2,c_1} \right) \left(\Phi_{3,c_2}^{J\dagger} i\epsilon^{IJK} \Phi_{3,c_2}^K \right) + Y_{23}'^{(3)} \left(\Phi_{2,c_1}^\dagger \sigma^I \Phi_{2,c_2} \right) \left(\Phi_{3,c_2}^{J\dagger} i\epsilon^{IJK} \Phi_{3,c_1}^K \right) \\ & + \left[Y_{233} \left(H^\dagger \sigma^I \Phi_{2,c_1} \right) \left(\Phi_{3,c_2}^J i\epsilon^{IJK} \Phi_{3,c_3}^K \right) \epsilon^{c_1 c_2 c_3} + \text{h.c.} \right] \end{aligned} \quad (\text{A.14})$$

$$\mathcal{L}_{\tilde{2}3} = Y_{\tilde{2}3}^{(1)} \left(\Phi_{\tilde{2},c_1}^\dagger \Phi_{\tilde{2},c_1} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_2} \right) + Y_{\tilde{2}3}'^{(1)} \left(\Phi_{\tilde{2},c_1}^\dagger \Phi_{\tilde{2},c_2} \right) \left(\Phi_{3,c_2}^\dagger \Phi_{3,c_1} \right)$$

$$\begin{aligned}
& + Y'_{23}^{(3)} \left(\Phi_{\tilde{2},c_1}^\dagger \sigma^I \Phi_{\tilde{2},c_2} \right) \left(\Phi_{3,c_2}^{J\dagger} i \epsilon^{IJK} \Phi_{3,c_1}^K \right) + Y_{23}^{(3)} \left(\Phi_{\tilde{2},c_1}^\dagger \sigma^I \Phi_{\tilde{2},c_1} \right) \left(\Phi_{3,c_2}^{J\dagger} i \epsilon^{IJK} \Phi_{3,c_2}^K \right) \\
& + \left[Y_{\tilde{2}33} \left(\Phi_{\tilde{2},c_1}^\dagger i \sigma_2 \sigma^I H \right) \left(\Phi_{3,c_2}^J i \epsilon^{IJK} \Phi_{3,c_3}^K \right) \epsilon^{c_1 c_2 c_3} + A_{\tilde{2}3} \left(\Phi_{\tilde{2}}^\dagger (\sigma \cdot \Phi_3) H \right) + \text{h.c.} \right]
\end{aligned} \tag{A.15}$$

The next set of terms $\mathcal{L}_{\tilde{1}2\tilde{2}}$, $\mathcal{L}_{1\tilde{1}2}$, \mathcal{L}_{123} , $\mathcal{L}_{1\tilde{2}3}$ and $\mathcal{L}_{\tilde{1}23}$ contain interactions involving precisely three LQs.

$$\mathcal{L}_{\tilde{1}2\tilde{2}} = A_{\tilde{1}2\tilde{2}} \Phi_{\tilde{1},c_1} \left(\Phi_{2,c_2}^\dagger i \sigma_2 \Phi_{\tilde{2},c_3} \right) \epsilon^{c_1 c_2 c_3} + \text{h.c.} \tag{A.16}$$

$$\mathcal{L}_{1\tilde{1}2} = Y_{1\tilde{1}2} \Phi_{1,c_1} \Phi_{\tilde{1},c_2} \left(\Phi_{2,c_3}^\dagger i \sigma_2 H \right) \epsilon^{c_1 c_2 c_3} + \text{h.c.} \tag{A.17}$$

$$\begin{aligned}
\mathcal{L}_{123} = & Y_{123} \Phi_{1,c_1} \left(H^\dagger (\sigma \cdot \Phi_{3,c_3}) \Phi_{2,c_2} \right) \epsilon^{c_1 c_2 c_3} + Y_{1223} \Phi_{1,c_1}^\dagger \left(\Phi_{2,c_2}^\dagger (\sigma \cdot \Phi_{3,c_2}) \Phi_{2,c_1} \right) \\
& + Y'_{1223} \Phi_{1,c_1}^\dagger \left(\Phi_{2,c_2}^\dagger (\sigma \cdot \Phi_{3,c_1}) \Phi_{2,c_2} \right) + \text{h.c.}
\end{aligned} \tag{A.18}$$

$$\begin{aligned}
\mathcal{L}_{1\tilde{2}3} = & Y_{1\tilde{2}3} \Phi_{1,c_1} \left(\Phi_{\tilde{2},c_2}^\dagger i \sigma_2 (\sigma \cdot \Phi_{3,c_3}) H \right) \epsilon^{c_1 c_2 c_3} + Y_{1\tilde{2}\tilde{2}3} \Phi_{1,c_1}^\dagger \left(\Phi_{\tilde{2},c_2}^\dagger (\sigma \cdot \Phi_{3,c_2}) \Phi_{\tilde{2},c_1} \right) \\
& + Y'_{1\tilde{2}\tilde{2}3} \Phi_{1,c_1}^\dagger \left(\Phi_{\tilde{2},c_2}^\dagger (\sigma \cdot \Phi_{3,c_1}) \Phi_{\tilde{2},c_2} \right) + \text{h.c.}
\end{aligned} \tag{A.19}$$

$$\mathcal{L}_{\tilde{1}23} = Y_{\tilde{1}23} \Phi_{\tilde{1},c_1} \left(\Phi_{2,c_2}^\dagger i \sigma_2 (\sigma \cdot \Phi_{3,c_3}) H \right) \epsilon^{c_1 c_2 c_3} + \text{h.c.} \tag{A.20}$$

Finally, the interaction terms $\mathcal{L}_{\tilde{1}2\tilde{2}3}$ and $\mathcal{L}_{1\tilde{1}2\tilde{2}}$ which involve four different SLQs are as follows,

$$\mathcal{L}_{\tilde{1}2\tilde{2}3} = Y_{\tilde{1}\tilde{2}23} \Phi_{\tilde{1},c_1}^\dagger \left(\Phi_{2,c_2}^\dagger (\sigma \cdot \Phi_{3,c_2}) \Phi_{\tilde{2},c_1} \right) + Y'_{\tilde{1}\tilde{2}23} \Phi_{\tilde{1},c_1}^\dagger \left(\Phi_{2,c_2}^\dagger (\sigma \cdot \Phi_{3,c_1}) \Phi_{\tilde{2},c_2} \right) + \text{h.c.} \tag{A.21}$$

$$\mathcal{L}_{1\tilde{1}2\tilde{2}} = Y_{1\tilde{1}2\tilde{2}} \Phi_{1,c_1}^\dagger \Phi_{\tilde{1},c_1} \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{2,c_2} \right) + Y'_{1\tilde{1}2\tilde{2}} \Phi_{1,c_1}^\dagger \Phi_{\tilde{1},c_2} \left(\Phi_{\tilde{2},c_2}^\dagger \Phi_{2,c_1} \right) + \text{h.c.} \tag{A.22}$$

B Two-loop β -functions of gauge couplings

In this Appendix, we collect the two-loop β -functions of gauge couplings of some of the SLQ models considered in this article.

- SM + Φ_1

$$\begin{aligned}\beta^{(2)}(g_1) = & +\frac{121}{30}g_1^5 + \frac{27}{10}g_1^3g_2^2 + \frac{148}{15}g_1^3g_3^2 - \frac{17}{10}g_1^3\text{Tr}(Y_u^\dagger Y_u) - \frac{1}{2}g_1^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{3}{2}g_1^3\text{Tr}(Y_e^\dagger Y_e) - \frac{13}{5}g_1^3\text{Tr}(Y_1^{RR\dagger} Y_1^{RR}) - g_1^3\text{Tr}(Y_1^{LL\dagger} Y_1^{LL}) \\ & - \frac{4}{5}g_1^3\text{Tr}(Y_1^{Q,LL*} Y_1^{Q,LL}) - 2g_1^3\text{Tr}(Y_1^{Q,RR\dagger} Y_1^{Q,RR})\end{aligned}\quad (\text{B.1})$$

$$\begin{aligned}\beta^{(2)}(g_2) = & +\frac{9}{10}g_1^2g_2^3 + \frac{35}{6}g_2^5 + 12g_2^3g_3^2 - \frac{3}{2}g_2^3\text{Tr}(Y_u^\dagger Y_u) - \frac{3}{2}g_2^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_2^3\text{Tr}(Y_e^\dagger Y_e) - 3g_2^3\text{Tr}(Y_1^{LL\dagger} Y_1^{LL}) - 12g_2^3\text{Tr}(Y_1^{Q,LL*} Y_1^{Q,LL})\end{aligned}\quad (\text{B.2})$$

$$\begin{aligned}\beta^{(2)}(g_3) = & +\frac{37}{30}g_1^2g_3^3 + \frac{9}{2}g_2^2g_3^3 - \frac{67}{3}g_3^5 - 2g_3^3\text{Tr}(Y_u^\dagger Y_u) - 2g_3^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_3^3\text{Tr}(Y_1^{RR\dagger} Y_1^{RR}) - g_3^3\text{Tr}(Y_1^{LL\dagger} Y_1^{LL}) - 8g_3^3\text{Tr}(Y_1^{Q,LL*} Y_1^{Q,LL}) \\ & - 2g_3^3\text{Tr}(Y_1^{Q,RR\dagger} Y_1^{Q,RR})\end{aligned}\quad (\text{B.3})$$

- SM + $\Phi_{\tilde{1}}$

$$\begin{aligned}\beta^{(2)}(g_1) = & +\frac{529}{30}g_1^5 + \frac{27}{10}g_1^3g_2^2 + \frac{388}{15}g_1^3g_3^2 - \frac{17}{10}g_1^3\text{Tr}(Y_u^\dagger Y_u) - \frac{1}{2}g_1^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{3}{2}g_1^3\text{Tr}(Y_e^\dagger Y_e) - 2g_1^3\text{Tr}(Y_{\tilde{1}}^{RR\dagger} Y_{\tilde{1}}^{RR}) + \frac{32}{5}g_1^3\text{Tr}(Y_{\tilde{1}}^{Q,RR*} Y_{\tilde{1}}^{Q,RR})\end{aligned}\quad (\text{B.4})$$

$$\begin{aligned}\beta^{(2)}(g_2) = & +\frac{9}{10}g_1^2g_2^3 + \frac{35}{6}g_2^5 + 12g_2^3g_3^2 - \frac{3}{2}g_2^3\text{Tr}(Y_u^\dagger Y_u) - \frac{3}{2}g_2^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_2^3\text{Tr}(Y_e^\dagger Y_e)\end{aligned}\quad (\text{B.5})$$

$$\begin{aligned}\beta^{(2)}(g_3) = & +\frac{97}{30}g_1^2g_3^3 + \frac{9}{2}g_2^2g_3^3 - \frac{67}{3}g_3^5 - 2g_3^3\text{Tr}(Y_u^\dagger Y_u) - 2g_3^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_3^3\text{Tr}(Y_{\tilde{1}}^{RR\dagger} Y_{\tilde{1}}^{RR}) + 4g_3^3\text{Tr}(Y_{\tilde{1}}^{Q,RR*} Y_{\tilde{1}}^{Q,RR})\end{aligned}\quad (\text{B.6})$$

- SM + Φ_2

$$\begin{aligned}\beta^{(2)}(g_1) = & +\frac{1499}{75}g_1^5 + \frac{87}{5}g_1^3g_2^2 + \frac{524}{15}g_1^3g_3^2 - \frac{17}{10}g_1^3\text{Tr}(Y_u^\dagger Y_u) - \frac{1}{2}g_1^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{3}{2}g_1^3\text{Tr}(Y_e^\dagger Y_e) - \frac{5}{2}g_1^3\text{Tr}(Y_2^{RL\dagger} Y_2^{RL}) - \frac{37}{10}g_1^3\text{Tr}(Y_2^{LR\dagger} Y_2^{LR})\end{aligned}\quad (\text{B.7})$$

$$\begin{aligned}\beta^{(2)}(g_2) = & +\frac{29}{5}g_1^2g_2^3 + \frac{37}{3}g_2^5 + 20g_2^3g_3^2 - \frac{3}{2}g_2^3\text{Tr}(Y_u^\dagger Y_u) - \frac{3}{2}g_2^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_2^3\text{Tr}(Y_e^\dagger Y_e) - \frac{3}{2}g_2^3\text{Tr}(Y_2^{RL\dagger} Y_2^{RL}) - \frac{3}{2}g_2^3\text{Tr}(Y_2^{LR\dagger} Y_2^{LR})\end{aligned}\quad (\text{B.8})$$

$$\begin{aligned}\beta^{(2)}(g_3) = & +\frac{131}{30}g_1^2g_3^3 + \frac{15}{2}g_2^2g_3^3 - \frac{56}{3}g_3^5 - 2g_3^3\text{Tr}(Y_u^\dagger Y_u) - 2g_3^3\text{Tr}(Y_d^\dagger Y_d) \\ & - g_3^3\text{Tr}(Y_2^{RL\dagger} Y_2^{RL}) - g_3^3\text{Tr}(Y_2^{LR\dagger} Y_2^{LR})\end{aligned}\quad (\text{B.9})$$

- SM + $\Phi_{\bar{2}}$

$$\begin{aligned}\beta^{(2)}(g_1) = & + \frac{299}{75}g_1^5 + 3g_1^3g_2^2 + \frac{28}{3}g_1^3g_3^2 - \frac{17}{10}g_1^3\text{Tr}(Y_u^\dagger Y_u) - \frac{1}{2}g_1^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{3}{2}g_1^3\text{Tr}(Y_e^\dagger Y_e) - \frac{13}{10}g_1^3\text{Tr}(Y_{\bar{2}}^{RL\dagger} Y_{\bar{2}}^{RL})\end{aligned}\quad (\text{B.10})$$

$$\begin{aligned}\beta^{(2)}(g_2) = & + g_1^2g_2^3 + \frac{37}{3}g_2^5 + 20g_2^3g_3^2 - \frac{3}{2}g_2^3\text{Tr}(Y_u^\dagger Y_u) - \frac{3}{2}g_2^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_2^3\text{Tr}(Y_e^\dagger Y_e) - \frac{3}{2}g_2^3\text{Tr}(Y_{\bar{2}}^{RL\dagger} Y_{\bar{2}}^{RL})\end{aligned}\quad (\text{B.11})$$

$$\begin{aligned}\beta^{(2)}(g_3) = & + \frac{7}{6}g_1^2g_3^3 + \frac{15}{2}g_2^2g_3^3 - \frac{56}{3}g_3^5 - 2g_3^3\text{Tr}(Y_u^\dagger Y_u) - 2g_3^3\text{Tr}(Y_d^\dagger Y_d) \\ & - g_3^3\text{Tr}(Y_{\bar{2}}^{RL\dagger} Y_{\bar{2}}^{RL})\end{aligned}\quad (\text{B.12})$$

- SM + Φ_3

$$\begin{aligned}\beta^{(2)}(g_1) = & + \frac{207}{50}g_1^5 + \frac{15}{2}g_1^3g_2^2 + 12g_1^3g_3^2 - \frac{17}{10}g_1^3\text{Tr}(Y_u^\dagger Y_u) - \frac{1}{2}g_1^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{3}{2}g_1^3\text{Tr}(Y_e^\dagger Y_e) - 3g_1^3\text{Tr}(Y_3^{LL\dagger} Y_3^{LL}) + \frac{12}{5}g_1^3\text{Tr}(Y_3^{Q,LL*} Y_3^{Q,LL})\end{aligned}\quad (\text{B.13})$$

$$\begin{aligned}\beta^{(2)}(g_2) = & + \frac{5}{2}g_1^2g_2^3 + \frac{371}{6}g_2^5 + 44g_2^3g_3^2 - \frac{3}{2}g_2^3\text{Tr}(Y_u^\dagger Y_u) - \frac{3}{2}g_2^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_2^3\text{Tr}(Y_e^\dagger Y_e) - 9g_2^3\text{Tr}(Y_3^{LL\dagger} Y_3^{LL}) + 36g_2^3\text{Tr}(Y_3^{Q,LL*} Y_3^{Q,LL})\end{aligned}\quad (\text{B.14})$$

$$\begin{aligned}\beta^{(2)}(g_3) = & + \frac{3}{2}g_1^2g_3^3 + \frac{33}{2}g_2^2g_3^3 - 15g_3^5 - 2g_3^3\text{Tr}(Y_u^\dagger Y_u) - 2g_3^3\text{Tr}(Y_d^\dagger Y_d) \\ & - 3g_3^3\text{Tr}(Y_3^{LL\dagger} Y_3^{LL}) + 24g_3^3\text{Tr}(Y_3^{Q,LL*} Y_3^{Q,LL})\end{aligned}\quad (\text{B.15})$$

- SM + $\Phi_2 + \Phi_3$

$$\begin{aligned}\beta^{(2)}(g_1) = & + \frac{1511}{75}g_1^5 + \frac{111}{5}g_1^3g_2^2 + \frac{572}{15}g_1^3g_3^2 - \frac{17}{10}g_1^3\text{Tr}(Y_u^\dagger Y_u) - \frac{1}{2}g_1^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{3}{2}g_1^3\text{Tr}(Y_e^\dagger Y_e) - \frac{5}{2}g_1^3\text{Tr}(Y_2^{RL\dagger} Y_2^{RL}) - \frac{37}{10}g_1^3\text{Tr}(Y_2^{LR\dagger} Y_2^{LR}) \\ & - 3g_1^3\text{Tr}(Y_3^{LL\dagger} Y_3^{LL}) + \frac{12}{5}g_1^3\text{Tr}(Y_3^{Q,LL*} Y_3^{Q,LL})\end{aligned}\quad (\text{B.16})$$

$$\begin{aligned}\beta^{(2)}(g_2) = & + \frac{37}{5}g_1^2g_2^3 + \frac{205}{3}g_2^5 + 52g_2^3g_3^2 - \frac{3}{2}g_2^3\text{Tr}(Y_u^\dagger Y_u) - \frac{3}{2}g_2^3\text{Tr}(Y_d^\dagger Y_d) \\ & - \frac{1}{2}g_2^3\text{Tr}(Y_e^\dagger Y_e) - \frac{3}{2}g_2^3\text{Tr}(Y_2^{RL\dagger} Y_2^{RL}) - \frac{3}{2}g_2^3\text{Tr}(Y_2^{LR\dagger} Y_2^{LR}) \\ & - 9g_2^3\text{Tr}(Y_3^{LL\dagger} Y_3^{LL}) + 36g_2^3\text{Tr}(Y_3^{Q,LL*} Y_3^{Q,LL})\end{aligned}\quad (\text{B.17})$$

$$\begin{aligned}
\beta^{(2)}(g_3) = & + \frac{143}{30} g_1^2 g_3^3 + \frac{39}{2} g_2^2 g_3^3 - \frac{23}{3} g_3^5 - 2 g_3^3 \text{Tr} \left(Y_u^\dagger Y_u \right) - 2 g_3^3 \text{Tr} \left(Y_d^\dagger Y_d \right) \\
& - g_3^3 \text{Tr} \left(Y_2^{RL\dagger} Y_2^{RL} \right) - g_3^3 \text{Tr} \left(Y_2^{LR\dagger} Y_2^{LR} \right) - 3 g_3^3 \text{Tr} \left(Y_3^{LL\dagger} Y_3^{LL} \right) \\
& + 24 g_3^3 \text{Tr} \left(Y_3^{Q,LL*} Y_3^{Q,LL} \right)
\end{aligned} \tag{B.18}$$

C One-Loop β -functions of SM Yukawa couplings

This Appendix presents the one-loop β -functions of the SM Yukawa couplings. We use n_1 , $n_{\tilde{1}}$, n_2 , $n_{\tilde{2}}$ and n_3 to denote the number of generations of Φ_1 , $\Phi_{\tilde{1}}$, Φ_2 , $\Phi_{\tilde{2}}$ and Φ_3 , respectively.

$$\begin{aligned}
\beta^{(1)}(Y_u) = & \frac{3}{2} Y_u Y_u^\dagger Y_u - \frac{3}{2} Y_d Y_d^\dagger Y_u + 3 \text{Tr} \left(Y_u^\dagger Y_u \right) Y_u + 3 \text{Tr} \left(Y_d^\dagger Y_d \right) Y_u + \text{Tr} \left(Y_e^\dagger Y_e \right) Y_u \\
& - \frac{17}{20} g_1^2 Y_u - \frac{9}{4} g_2^2 Y_u - 8 g_3^2 Y_u + n_1 \left(\frac{1}{2} Y_u Y_1^{RR*} Y_1^{RRT} + Y_u Y_1^{Q,RR*} Y_1^{Q,RRT} \right. \\
& \left. + 2 Y_1^{LL*} Y_e^* Y_1^{RRT} + \frac{1}{2} Y_1^{LL*} Y_1^{LLT} Y_u + 8 Y_1^{Q,LL*} Y_d^* Y_1^{Q,RRT} + 4 Y_1^{Q,LL*} Y_1^{Q,LL} Y_u \right) \\
& - n_{\tilde{1}} \left(4 Y_u Y_{\tilde{1}}^{Q,RR*} Y_{\tilde{1}}^{Q,RR} \right) + n_2 \left(Y_u Y_2^{RL} Y_2^{RL\dagger} + 2 Y_2^{LR} Y_e^* Y_2^{RL\dagger} + \frac{1}{2} Y_2^{LR} Y_2^{LR\dagger} Y_u \right) \\
& + n_3 \left(\frac{3}{2} Y_3^{LL*} Y_3^{LLT} Y_u - 12 Y_3^{Q,LL*} Y_3^{Q,LL} Y_u \right)
\end{aligned} \tag{C.1}$$

$$\begin{aligned}
\beta^{(1)}(Y_d) = & \frac{3}{2} Y_d Y_d^\dagger Y_d - \frac{3}{2} Y_u Y_u^\dagger Y_d + 3 \text{Tr} \left(Y_u^\dagger Y_d \right) Y_d + 3 \text{Tr} \left(Y_d^\dagger Y_d \right) Y_d + \text{Tr} \left(Y_e^\dagger Y_e \right) Y_d - \frac{1}{4} g_1^2 Y_d \\
& - \frac{9}{4} g_2^2 Y_d - 8 g_3^2 Y_d + n_1 \left(Y_d Y_1^{Q,RR\dagger} Y_1^{Q,RR} + \frac{1}{2} Y_1^{LL*} Y_1^{LLT} Y_d + 8 Y_1^{Q,LL*} Y_u^* Y_1^{Q,RR} \right. \\
& \left. + 4 Y_1^{Q,LL*} Y_1^{Q,LL} Y_d \right) + n_{\tilde{1}} \left(\frac{1}{2} Y_d Y_{\tilde{1}}^{RR*} Y_{\tilde{1}}^{RRT} \right) + n_2 \left(\frac{1}{2} Y_2^{LR} Y_2^{LR\dagger} Y_d \right) + n_{\tilde{2}} \left(Y_d Y_{\tilde{2}}^{RL} Y_{\tilde{2}}^{RL\dagger} \right) \\
& + n_3 \left(\frac{3}{2} Y_3^{LL*} Y_3^{LLT} Y_d - 12 Y_3^{Q,LL*} Y_3^{Q,LL} Y_d \right)
\end{aligned} \tag{C.2}$$

$$\begin{aligned}
\beta^{(1)}(Y_e) = & \frac{3}{2} Y_e Y_e^\dagger Y_e + 3 \text{Tr} \left(Y_u^\dagger Y_e \right) Y_e + 3 \text{Tr} \left(Y_d^\dagger Y_e \right) Y_e + \text{Tr} \left(Y_e^\dagger Y_e \right) Y_e - \frac{9}{4} g_1^2 Y_e - \frac{9}{4} g_2^2 Y_e \\
& + n_1 \left(+ \frac{3}{2} Y_e Y_1^{RR\dagger} Y_1^{RR} + 6 Y_1^{LL\dagger} Y_u^* Y_1^{RR} + \frac{3}{2} Y_1^{LL\dagger} Y_1^{LL} Y_e \right) + n_{\tilde{1}} \left(\frac{3}{2} Y_e Y_{\tilde{1}}^{RR\dagger} Y_{\tilde{1}}^{RR} \right) \\
& + n_2 \left(3 Y_e Y_2^{LR\dagger} Y_2^{LR} + 6 Y_2^{RL\dagger} Y_u^* Y_2^{LR} + \frac{3}{2} Y_2^{RL\dagger} Y_2^{RL} Y_e \right) + n_{\tilde{2}} \left(\frac{3}{2} Y_{\tilde{2}}^{RL\dagger} Y_{\tilde{2}}^{RL} Y_e \right) \\
& + n_3 \left(\frac{9}{2} Y_3^{LL\dagger} Y_3^{LL} Y_e \right)
\end{aligned} \tag{C.3}$$

D One-Loop threshold corrections

In this Appendix, we collect the one-loop threshold corrections of the SM gauge, Yukawa, and quartic couplings on matching the SM with $\text{SM} + \Phi_1 + \dots + \Phi_3$. To simplify the expressions, we assumed all the LQs to have the same mass m (for unequal mass, see Ref. [197]) and set the dimensionful trilinear couplings to zero. The superscript 0 denotes the original parameters in the SM lagrangian and we use the shorthand notation $L_{xm} = 1 + x \log\left(\frac{\mu^2}{m^2}\right)$.

$$g_1^{(0)} = g_1 - \frac{3g_1^3 \log\left(\frac{\mu}{m^2}\right)}{32\pi^2}, \quad g_2^{(0)} = g_2 - \frac{3g_2^3 \log\left(\frac{\mu}{m^2}\right)}{32\pi^2}, \quad g_3^{(0)} = g_3 - \frac{3g_3^3 \log\left(\frac{\mu}{m^2}\right)}{64\pi^2} \quad (\text{D.1})$$

$$\begin{aligned} \lambda^{(0)} = \lambda & - \frac{3Y_{\tilde{1}3}Y_{\tilde{1}3}^* \log\left(\frac{\mu}{m^2}\right)}{8\pi^2} - \frac{3Y_{\tilde{2}2}Y_{\tilde{2}2}^* \log\left(\frac{\mu}{m^2}\right)}{16\pi^2} - \frac{3Y_{\tilde{1}}^2 \log\left(\frac{\mu}{m^2}\right)}{32\pi^2} - \frac{3Y_{\tilde{2}}^2 \log\left(\frac{\mu}{m^2}\right)}{16\pi^2} \\ & - \frac{3Y_{\tilde{2}2}^2 \log\left(\frac{\mu}{m^2}\right)}{32\pi^2} - \frac{3Y_2Y_{\tilde{2}2} \log\left(\frac{\mu}{m^2}\right)}{16\pi^2} - \frac{3Y_{13}Y_{13}^* \log\left(\frac{\mu}{m^2}\right)}{16\pi^2} - \frac{3Y_1^2 \log\left(\frac{\mu}{m^2}\right)}{32\pi^2} \\ & - \frac{3Y_2^2 \log\left(\frac{\mu}{m^2}\right)}{16\pi^2} - \frac{9Y_3^2 \log\left(\frac{\mu}{m^2}\right)}{32\pi^2} - \frac{3Y_{22}^2 \log\left(\frac{\mu}{m^2}\right)}{32\pi^2} - \frac{3Y_{33}^2 \log\left(\frac{\mu}{m^2}\right)}{16\pi^2} - \frac{3Y_2Y_{22} \log\left(\frac{\mu}{m^2}\right)}{16\pi^2} \end{aligned} \quad (\text{D.2})$$

$$\begin{aligned} Y_u^{(0)} = Y_u & + \frac{Y_u Y_{\tilde{1}}^{Q,RR\dagger} Y_{\tilde{1}}^{Q,RR^T}}{16\pi^2} L_{2m} - \frac{Y_1^{Q,LL^*} Y_d^* Y_1^{Q,RR^T}}{4\pi^2} L_m - \frac{Y_1^{LL^*} Y_e^* Y_1^{RR^T}}{16\pi^2} L_m \\ & - \frac{Y_1^{LL^*} Y_1^{LL^T} Y_u}{128\pi^2} L_{2m} - \frac{3Y_3^{LL^*} Y_3^{LL^T} Y_u}{128\pi^2} L_{2m} - \frac{Y_1^{Q,LL^*} Y_1^{Q,LL^T} Y_u}{16\pi^2} L_{2m} \\ & + \frac{3Y_3^{Q,LL^*} Y_3^{Q,LL} Y_u}{16\pi^2} L_{2m} - \frac{Y_u Y_1^{Q,RR^*} Y_1^{Q,RR^T}}{64\pi^2} L_{2m} - \frac{Y_u Y_1^{RR^*} Y_1^{RR^T}}{128\pi^2} L_{2m} \\ & + \frac{Y_2^{LR} Y_e^\dagger Y_2^{RL^\dagger}}{16\pi^2} L_m - \frac{Y_2^{LR} Y_2^{LR^\dagger} Y_u}{128\pi^2} L_{2m} - \frac{Y_u Y_2^{RL} Y_2^{RL^\dagger}}{64\pi^2} L_{2m} \end{aligned} \quad (\text{D.3})$$

$$\begin{aligned} Y_d^{(0)} = Y_d & - \frac{Y_d Y_{\tilde{1}}^{RR^*} Y_{\tilde{1}}^{RR^T}}{128\pi^2} L_{2m} - \frac{Y_d Y_{\tilde{2}}^{RL} Y_{\tilde{2}}^{RL^\dagger}}{64\pi^2} L_{2m} - \frac{Y_1^{LL^*} Y_1^{LL^T} Y_d}{128\pi^2} L_{2m} \\ & - \frac{3Y_3^{LL^*} Y_3^{LL^T} Y_d}{128\pi^2} L_{2m} - \frac{Y_1^{Q,LL^*} Y_1^{Q,LL^T} Y_d}{16\pi^2} L_{2m} + \frac{3Y_3^{Q,LL^*} Y_3^{Q,LL} Y_d}{16\pi^2} L_{2m} \\ & - \frac{Y_1^{Q,LL^*} Y_u^* Y_1^{Q,RR}}{4\pi^2} L_m - \frac{Y_2^{LR} Y_2^{LR^\dagger} Y_d}{128\pi^2} L_{2m} - \frac{Y_d Y_1^{Q,RR^*} Y_1^{Q,RR}}{64\pi^2} L_{2m} \end{aligned} \quad (\text{D.4})$$

$$Y_e^{(0)} = Y_e - \frac{3Y_{\tilde{2}}^{RL^\dagger} Y_{\tilde{2}}^{RL} Y_e}{128\pi^2} L_{2m} - \frac{3Y_e Y_{\tilde{1}}^{RR^*} Y_{\tilde{1}}^{RR}}{128\pi^2} L_{2m} - \frac{3Y_1^{LL^\dagger} Y_u^* Y_1^{RR}}{16\pi^2} L_m$$

$$\begin{aligned}
& - \frac{3Y_1^{LL\dagger} Y_1^{LL} Y_e}{128\pi^2} L_{2m} - \frac{9Y_3^{LL\dagger} Y_3^{LL} Y_e}{128\pi^2} L_{2m} - \frac{3Y_e Y_2^{LR\dagger} Y_2^{LR}}{64\pi^2} L_{2m} \\
& - \frac{3Y_2^{RL\dagger} Y_2^{RL} Y_e}{128\pi^2} L_{2m} - \frac{3Y_e Y_1^{RR\dagger} Y_1^{RR}}{128\pi^2} L_{2m} + \frac{3Y_2^{RL\dagger} Y_u^\dagger Y_2^{LR}}{16\pi^2} L_m
\end{aligned} \tag{D.5}$$

References

- [1] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, *Phys. Lett.* **12** (1964) 132.
- [2] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, *Phys. Rev. Lett.* **13** (1964) 321.
- [3] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, *Phys. Rev. Lett.* **13** (1964) 508.
- [4] G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, *Global Conservation Laws and Massless Particles*, *Phys. Rev. Lett.* **13** (1964) 585.
- [5] ATLAS collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[1207.7214](#)].
- [6] CMS collaboration, *Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[1207.7235](#)].
- [7] CDF, D0 collaboration, *Evidence for a particle produced in association with weak bosons and decaying to a bottom-antibottom quark pair in Higgs boson searches at the Tevatron*, *Phys. Rev. Lett.* **109** (2012) 071804 [[1207.6436](#)].
- [8] J. C. Pati and A. Salam, *Lepton Number as the Fourth Color*, *Phys. Rev. D* **10** (1974) 275.
- [9] H. Fritzsch and P. Minkowski, *Unified Interactions of Leptons and Hadrons*, *Annals Phys.* **93** (1975) 193.
- [10] H. Georgi and S. L. Glashow, *Unity of All Elementary Particle Forces*, *Phys. Rev. Lett.* **32** (1974) 438.
- [11] H. Georgi, H. R. Quinn and S. Weinberg, *Hierarchy of Interactions in Unified Gauge Theories*, *Phys. Rev. Lett.* **33** (1974) 451.
- [12] J. C. Pati, A. Salam and J. A. Strathdee, *Are Quarks Composite?*, *Phys. Lett. B* **59** (1975) 265.
- [13] J. Wudka, *COMPOSITE LEPTOQUARKS*, *Phys. Lett. B* **167** (1986) 337.
- [14] O. J. P. Eboli and A. V. Olinto, *Composite Leptoquarks in Hadronic Colliders*, *Phys. Rev. D* **38** (1988) 3461.
- [15] B. Gripaios, *Composite Leptoquarks at the LHC*, *JHEP* **02** (2010) 045 [[0910.1789](#)].
- [16] R. Barbier et al., *R-parity violating supersymmetry*, *Phys. Rept.* **420** (2005) 1 [[hep-ph/0406039](#)].
- [17] W. Buchmuller, R. Ruckl and D. Wyler, *Leptoquarks in Lepton - Quark Collisions*, *Phys. Lett. B* **191** (1987) 442.

- [18] O. U. Shanker, *Flavor Violation, Scalar Particles and Leptoquarks*, *Nucl. Phys. B* **206** (1982) 253.
- [19] O. U. Shanker, $\pi\ell$ 2, $K\ell$ 3 and $K^0 - \bar{K}0$ Constraints on Leptoquarks and Supersymmetric Particles, *Nucl. Phys. B* **204** (1982) 375.
- [20] M. Leurer, A Comprehensive study of leptoquark bounds, *Phys. Rev. D* **49** (1994) 333 [[hep-ph/9309266](#)].
- [21] M. Leurer, Bounds on vector leptoquarks, *Phys. Rev. D* **50** (1994) 536 [[hep-ph/9312341](#)].
- [22] S. Davidson, D. C. Bailey and B. A. Campbell, Model independent constraints on leptoquarks from rare processes, *Z. Phys. C* **61** (1994) 613 [[hep-ph/9309310](#)].
- [23] A. Crivellin, M. Hoferichter, M. Kirk, C. A. Manzari and L. Schnell, First-generation new physics in simplified models: from low-energy parity violation to the LHC, *JHEP* **10** (2021) 221 [[2107.13569](#)].
- [24] A. Crivellin, C. Greub, D. Müller and F. Saturnino, Scalar Leptoquarks in Leptonic Processes, *JHEP* **02** (2021) 182 [[2010.06593](#)].
- [25] B. Gripaios, M. Nardecchia and S. A. Renner, Composite leptoquarks and anomalies in B -meson decays, *JHEP* **05** (2015) 006 [[1412.1791](#)].
- [26] R. Alonso, B. Grinstein and J. Martin Camalich, Lepton universality violation and lepton flavor conservation in B -meson decays, *JHEP* **10** (2015) 184 [[1505.05164](#)].
- [27] L. Calibbi, A. Crivellin and T. Ota, Effective Field Theory Approach to $b \rightarrow s\ell\ell^{(\prime)}$, $B \rightarrow K^{(*)}\nu\bar{\nu}$ and $B \rightarrow D^{(*)}\tau\nu$ with Third Generation Couplings, *Phys. Rev. Lett.* **115** (2015) 181801 [[1506.02661](#)].
- [28] S. Fajfer and N. Košnik, Vector leptoquark resolution of R_K and $R_{D^{(*)}}$ puzzles, *Phys. Lett. B* **755** (2016) 270 [[1511.06024](#)].
- [29] B. Bhattacharya, A. Datta, J.-P. Guévin, D. London and R. Watanabe, Simultaneous Explanation of the R_K and $R_{D^{(*)}}$ Puzzles: a Model Analysis, *JHEP* **01** (2017) 015 [[1609.09078](#)].
- [30] D. Buttazzo, A. Greljo, G. Isidori and D. Marzocca, B -physics anomalies: a guide to combined explanations, *JHEP* **11** (2017) 044 [[1706.07808](#)].
- [31] R. Barbieri, G. Isidori, A. Pattori and F. Senia, Anomalies in B -decays and $U(2)$ flavour symmetry, *Eur. Phys. J. C* **76** (2016) 67 [[1512.01560](#)].
- [32] R. Barbieri, C. W. Murphy and F. Senia, B -decay Anomalies in a Composite Leptoquark Model, *Eur. Phys. J. C* **77** (2017) 8 [[1611.04930](#)].
- [33] L. Calibbi, A. Crivellin and T. Li, Model of vector leptoquarks in view of the B -physics anomalies, *Phys. Rev. D* **98** (2018) 115002 [[1709.00692](#)].
- [34] M. Bordone, C. Cornella, J. Fuentes-Martin and G. Isidori, A three-site gauge model for flavor hierarchies and flavor anomalies, *Phys. Lett. B* **779** (2018) 317 [[1712.01368](#)].
- [35] M. Bordone, C. Cornella, J. Fuentes-Martín and G. Isidori, Low-energy signatures of the PS^3 model: from B -physics anomalies to LFV, *JHEP* **10** (2018) 148 [[1805.09328](#)].

- [36] J. Kumar, D. London and R. Watanabe, *Combined Explanations of the $b \rightarrow s\mu^+\mu^-$ and $b \rightarrow c\tau^-\bar{\nu}$ Anomalies: a General Model Analysis*, *Phys. Rev. D* **99** (2019) 015007 [[1806.07403](#)].
- [37] A. Biswas, D. Kumar Ghosh, N. Ghosh, A. Shaw and A. K. Swain, *Collider signature of U_1 Leptoquark and constraints from $b \rightarrow c$ observables*, *J. Phys. G* **47** (2020) 045005 [[1808.04169](#)].
- [38] L. Da Rold and F. Lamagna, *Composite Higgs and leptoquarks from a simple group*, *JHEP* **03** (2019) 135 [[1812.08678](#)].
- [39] A. Crivellin, C. Greub, D. Müller and F. Saturnino, *Importance of Loop Effects in Explaining the Accumulated Evidence for New Physics in B Decays with a Vector Leptoquark*, *Phys. Rev. Lett.* **122** (2019) 011805 [[1807.02068](#)].
- [40] M. Blanke and A. Crivellin, *B Meson Anomalies in a Pati-Salam Model within the Randall-Sundrum Background*, *Phys. Rev. Lett.* **121** (2018) 011801 [[1801.07256](#)].
- [41] J. Heeck and D. Teresi, *Pati-Salam explanations of the B -meson anomalies*, *JHEP* **12** (2018) 103 [[1808.07492](#)].
- [42] I. de Medeiros Varzielas and J. Talbert, *Simplified Models of Flavourful Leptoquarks*, *Eur. Phys. J. C* **79** (2019) 536 [[1901.10484](#)].
- [43] C. Cornella, J. Fuentes-Martin and G. Isidori, *Revisiting the vector leptoquark explanation of the B -physics anomalies*, *JHEP* **07** (2019) 168 [[1903.11517](#)].
- [44] M. Bordone, O. Catà and T. Feldmann, *Effective Theory Approach to New Physics with Flavour: General Framework and a Leptoquark Example*, *JHEP* **01** (2020) 067 [[1910.02641](#)].
- [45] S. Sahoo and R. Mohanta, *Scalar leptoquarks and the rare B meson decays*, *Phys. Rev. D* **91** (2015) 094019 [[1501.05193](#)].
- [46] C.-H. Chen, T. Nomura and H. Okada, *Explanation of $B \rightarrow K^{(*)}\ell^+\ell^-$ and muon $g - 2$, and implications at the LHC*, *Phys. Rev. D* **94** (2016) 115005 [[1607.04857](#)].
- [47] U. K. Dey, D. Kar, M. Mitra, M. Spannowsky and A. C. Vincent, *Searching for Leptoquarks at IceCube and the LHC*, *Phys. Rev. D* **98** (2018) 035014 [[1709.02009](#)].
- [48] D. Bećirević and O. Sumensari, *A leptoquark model to accommodate $R_K^{\text{exp}} < R_K^{\text{SM}}$ and $R_{K^*}^{\text{exp}} < R_{K^*}^{\text{SM}}$* , *JHEP* **08** (2017) 104 [[1704.05835](#)].
- [49] B. Chauhan, B. Kindra and A. Narang, *Discrepancies in simultaneous explanation of flavor anomalies and IceCube PeV events using leptoquarks*, *Phys. Rev. D* **97** (2018) 095007 [[1706.04598](#)].
- [50] D. Bećirević, I. Doršner, S. Fajfer, N. Košnik, D. A. Faroughy and O. Sumensari, *Scalar leptoquarks from grand unified theories to accommodate the B -physics anomalies*, *Phys. Rev. D* **98** (2018) 055003 [[1806.05689](#)].
- [51] O. Popov, M. A. Schmidt and G. White, *R_2 as a single leptoquark solution to $R_{D^{(*)}}$ and $R_{K^{(*)}}$* , *Phys. Rev. D* **100** (2019) 035028 [[1905.06339](#)].
- [52] S. Fajfer, J. F. Kamenik, I. Nisandžić and J. Zupan, *Implications of Lepton Flavor Universality Violations in B Decays*, *Phys. Rev. Lett.* **109** (2012) 161801 [[1206.1872](#)].

- [53] N. G. Deshpande and A. Menon, *Hints of R-parity violation in B decays into $\tau\nu$* , *JHEP* **01** (2013) 025 [[1208.4134](#)].
- [54] M. Freytsis, Z. Ligeti and J. T. Ruderman, *Flavor models for $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **92** (2015) 054018 [[1506.08896](#)].
- [55] M. Bauer and M. Neubert, *Minimal Leptoquark Explanation for the $R_{D^{(*)}}$, R_K , and $(g-2)_\mu$ Anomalies*, *Phys. Rev. Lett.* **116** (2016) 141802 [[1511.01900](#)].
- [56] X.-Q. Li, Y.-D. Yang and X. Zhang, *Revisiting the one leptoquark solution to the $R(D^{(*)})$ anomalies and its phenomenological implications*, *JHEP* **08** (2016) 054 [[1605.09308](#)].
- [57] J. Zhu, H.-M. Gan, R.-M. Wang, Y.-Y. Fan, Q. Chang and Y.-G. Xu, *Probing the R-parity violating supersymmetric effects in the exclusive $b \rightarrow c\ell^-\bar{\nu}_\ell$ decays*, *Phys. Rev. D* **93** (2016) 094023 [[1602.06491](#)].
- [58] O. Popov and G. A. White, *One Leptoquark to unify them? Neutrino masses and unification in the light of $(g-2)_\mu$, $R_{D^{(*)}}$ and R_K anomalies*, *Nucl. Phys. B* **923** (2017) 324 [[1611.04566](#)].
- [59] N. G. Deshpande and X.-G. He, *Consequences of R-parity violating interactions for anomalies in $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$ and $b \rightarrow s\mu^+\mu^-$* , *Eur. Phys. J. C* **77** (2017) 134 [[1608.04817](#)].
- [60] D. Bećirević, N. Košnik, O. Sumensari and R. Zukanovich Funchal, *Palatable Leptoquark Scenarios for Lepton Flavor Violation in Exclusive $b \rightarrow s\ell_1\ell_2$ modes*, *JHEP* **11** (2016) 035 [[1608.07583](#)].
- [61] Y. Cai, J. Gargalionis, M. A. Schmidt and R. R. Volkas, *Reconsidering the One Leptoquark solution: flavor anomalies and neutrino mass*, *JHEP* **10** (2017) 047 [[1704.05849](#)].
- [62] W. Altmannshofer, P. S. Bhupal Dev and A. Soni, *$R_{D^{(*)}}$ anomaly: A possible hint for natural supersymmetry with R-parity violation*, *Phys. Rev. D* **96** (2017) 095010 [[1704.06659](#)].
- [63] S. Kamali, A. Rashed and A. Datta, *New physics in inclusive $B \rightarrow X_c\ell\bar{\nu}$ decay in light of $R(D^{(*)})$ measurements*, *Phys. Rev. D* **97** (2018) 095034 [[1801.08259](#)].
- [64] T. Mandal, S. Mitra and S. Raz, *$R_{D^{(*)}}$ motivated S_1 leptoquark scenarios: Impact of interference on the exclusion limits from LHC data*, *Phys. Rev. D* **99** (2019) 055028 [[1811.03561](#)].
- [65] A. Azatov, D. Bardhan, D. Ghosh, F. Sgarlata and E. Venturini, *Anatomy of $b \rightarrow c\tau\nu$ anomalies*, *JHEP* **11** (2018) 187 [[1805.03209](#)].
- [66] J. Zhu, B. Wei, J.-H. Sheng, R.-M. Wang, Y. Gao and G.-R. Lu, *Probing the R-parity violating supersymmetric effects in $B_c \rightarrow J/\psi\ell^-\bar{\nu}_\ell$, $\eta_c\ell^-\bar{\nu}_\ell$ and $\Lambda_b \rightarrow \Lambda_c\ell^-\bar{\nu}_\ell$ decays*, *Nucl. Phys. B* **934** (2018) 380 [[1801.00917](#)].
- [67] A. Angelescu, D. Bećirević, D. A. Faroughy and O. Sumensari, *Closing the window on single leptoquark solutions to the B-physics anomalies*, *JHEP* **10** (2018) 183 [[1808.08179](#)].
- [68] T. J. Kim, P. Ko, J. Li, J. Park and P. Wu, *Correlation between $R_{D^{(*)}}$ and top quark FCNC decays in leptoquark models*, *JHEP* **07** (2019) 025 [[1812.08484](#)].
- [69] U. Aydemir, T. Mandal and S. Mitra, *Addressing the $\mathbf{R}_{D^{(*)}}$ anomalies with an \mathbf{S}_1 leptoquark from $\mathbf{SO(10)}$ grand unification*, *Phys. Rev. D* **101** (2020) 015011 [[1902.08108](#)].

- [70] H. Yan, Y.-D. Yang and X.-B. Yuan, *Phenomenology of $b \rightarrow c\tau\bar{\nu}$ decays in a scalar leptoquark model*, *Chin. Phys. C* **43** (2019) 083105 [[1905.01795](#)].
- [71] A. Crivellin, D. Müller and T. Ota, *Simultaneous explanation of $R(D^{(*)})$ and $b \rightarrow s\mu^+ \mu^-$: the last scalar leptoquarks standing*, *JHEP* **09** (2017) 040 [[1703.09226](#)].
- [72] D. Marzocca, *Addressing the B-physics anomalies in a fundamental Composite Higgs Model*, *JHEP* **07** (2018) 121 [[1803.10972](#)].
- [73] I. Bigaran, J. Gargalionis and R. R. Volkas, *A near-minimal leptoquark model for reconciling flavour anomalies and generating radiative neutrino masses*, *JHEP* **10** (2019) 106 [[1906.01870](#)].
- [74] A. Crivellin, D. Müller and F. Saturnino, *Flavor Phenomenology of the Leptoquark Singlet-Triplet Model*, *JHEP* **06** (2020) 020 [[1912.04224](#)].
- [75] S. Saad, *Combined explanations of $(g - 2)_\mu$, $R_{D^{(*)}}$, $R_{K^{(*)}}$ anomalies in a two-loop radiative neutrino mass model*, *Phys. Rev. D* **102** (2020) 015019 [[2005.04352](#)].
- [76] P. S. Bhupal Dev, R. Mohanta, S. Patra and S. Sahoo, *Unified explanation of flavor anomalies, radiative neutrino masses, and ANITA anomalous events in a vector leptoquark model*, *Phys. Rev. D* **102** (2020) 095012 [[2004.09464](#)].
- [77] S. Saad and A. Thapa, *Common origin of neutrino masses and $R_{D^{(*)}}$, $R_{K^{(*)}}$ anomalies*, *Phys. Rev. D* **102** (2020) 015014 [[2004.07880](#)].
- [78] W. Altmannshofer, P. S. B. Dev, A. Soni and Y. Sui, *Addressing $R_{D^{(*)}}$, $R_{K^{(*)}}$, muon $g - 2$ and ANITA anomalies in a minimal R-parity violating supersymmetric framework*, *Phys. Rev. D* **102** (2020) 015031 [[2002.12910](#)].
- [79] J. Fuentes-Martín and P. Stangl, *Third-family quark-lepton unification with a fundamental composite Higgs*, *Phys. Lett. B* **811** (2020) 135953 [[2004.11376](#)].
- [80] V. Gherardi, D. Marzocca and E. Venturini, *Low-energy phenomenology of scalar leptoquarks at one-loop accuracy*, *JHEP* **01** (2021) 138 [[2008.09548](#)].
- [81] L. Da Rold and F. Lamagna, *Model for the singlet-triplet leptoquarks*, *Phys. Rev. D* **103** (2021) 115007 [[2011.10061](#)].
- [82] J. Aebischer, G. Isidori, M. Pesut, B. A. Stefanek and F. Wilsch, *Confronting the vector leptoquark hypothesis with new low- and high-energy data*, *Eur. Phys. J. C* **83** (2023) 153 [[2210.13422](#)].
- [83] G. Hiller, D. Loose and K. Schönwald, *Leptoquark Flavor Patterns & B Decay Anomalies*, *JHEP* **12** (2016) 027 [[1609.08895](#)].
- [84] A. Crivellin, D. Müller, A. Signer and Y. Ulrich, *Correlating lepton flavor universality violation in B decays with $\mu \rightarrow e\gamma$ using leptoquarks*, *Phys. Rev. D* **97** (2018) 015019 [[1706.08511](#)].
- [85] A. Crivellin and F. Saturnino, *Explaining the Flavor Anomalies with a Vector Leptoquark (Moriond 2019 update)*, *PoS DIS2019* (2019) 163 [[1906.01222](#)].
- [86] J. Bernigaud, I. de Medeiros Varzielas and J. Talbert, *Finite Family Groups for Fermionic and Leptoquark Mixing Patterns*, *JHEP* **01** (2020) 194 [[1906.11270](#)].

- [87] J. Aebischer, A. Crivellin and C. Greub, *QCD improved matching for semileptonic B decays with leptoquarks*, *Phys. Rev. D* **99** (2019) 055002 [[1811.08907](#)].
- [88] J. Fuentes-Martín, G. Isidori, M. König and N. Selimović, *Vector Leptoquarks Beyond Tree Level*, *Phys. Rev. D* **101** (2020) 035024 [[1910.13474](#)].
- [89] I. de Medeiros Varzielas and G. Hiller, *Clues for flavor from rare lepton and quark decays*, *JHEP* **06** (2015) 072 [[1503.01084](#)].
- [90] S. Sahoo, R. Mohanta and A. K. Giri, *Explaining the R_K and $R_{D^{(*)}}$ anomalies with vector leptoquarks*, *Phys. Rev. D* **95** (2017) 035027 [[1609.04367](#)].
- [91] S. Fajfer, N. Košnik and L. Vale Silva, *Footprints of leptoquarks: from $R_{K^{(*)}}$ to $K \rightarrow \pi\nu\bar{\nu}$* , *Eur. Phys. J. C* **78** (2018) 275 [[1802.00786](#)].
- [92] U. Aydemir, D. Minic, C. Sun and T. Takeuchi, *B -decay anomalies and scalar leptoquarks in unified Pati-Salam models from noncommutative geometry*, *JHEP* **09** (2018) 117 [[1804.05844](#)].
- [93] A. Carvunis, A. Crivellin, D. Guadagnoli and S. Gangal, *The Forward-Backward Asymmetry in $B \rightarrow D^*\ell\nu$: One more hint for Scalar Leptoquarks?*, *Phys. Rev. D* **105** (2022) L031701 [[2106.09610](#)].
- [94] D. Zhang, *Radiative neutrino masses, lepton flavor mixing and muon $g - 2$ in a leptoquark model*, *JHEP* **07** (2021) 069 [[2105.08670](#)].
- [95] D. Marzocca, S. Trifinopoulos and E. Venturini, *From B -meson anomalies to Kaon physics with scalar leptoquarks*, *Eur. Phys. J. C* **82** (2022) 320 [[2106.15630](#)].
- [96] R. Barbieri, C. Cornella and G. Isidori, *Simplified models of vector $SU(4)$ leptoquarks at the TeV*, *Eur. Phys. J. C* **82** (2022) 1161 [[2207.14248](#)].
- [97] C. H. García-Duque, J. M. Cabarcas, J. H. Muñoz, N. Quintero and E. Rojas, *Singlet vector leptoquark model facing recent $LHCb$ and $BABAR$ measurements*, *Nucl. Phys. B* **988** (2023) 116115 [[2209.04753](#)].
- [98] A. Crivellin, B. Fuks and L. Schnell, *Explaining the hints for lepton flavour universality violation with three S_2 leptoquark generations*, *JHEP* **06** (2022) 169 [[2203.10111](#)].
- [99] M. Fernández Navarro and S. F. King, *B -anomalies in a twin Pati-Salam theory of flavour including the 2022 $LHCb$ $R_{K^{(*)}}$ analysis*, *JHEP* **02** (2023) 188 [[2209.00276](#)].
- [100] A. Ismael and S. Khalil, *Resolving R_D and R_{D^*} Anomalies in Adjoint $SU(5)$* , [2301.02226](#).
- [101] S. Iguro and Y. Omura, *A closer look at isodoublet vector leptoquark solution to $R_{D^{(*)}}$ anomaly*, [2306.00052](#).
- [102] J. M. Lizana, J. Matias and B. A. Stefanek, *Explaining the $B_{d,s} \rightarrow K^{(*)}\bar{K}^{(*)}$ non-leptonic puzzle and charged-current B -anomalies via scalar leptoquarks*, [2306.09178](#).
- [103] A. Djouadi, T. Kohler, M. Spira and J. Tutas, *(e b), (e t) TYPE LEPTOQUARKS AT e p COLLIDERS*, *Z. Phys. C* **46** (1990) 679.
- [104] D. Chakraverty, D. Choudhury and A. Datta, *A Nonsupersymmetric resolution of the anomalous muon magnetic moment*, *Phys. Lett. B* **506** (2001) 103 [[hep-ph/0102180](#)].

- [105] K.-m. Cheung, *Muon anomalous magnetic moment and leptoquark solutions*, *Phys. Rev. D* **64** (2001) 033001 [[hep-ph/0102238](#)].
- [106] C. Biggio, M. Bordone, L. Di Luzio and G. Ridolfi, *Massive vectors and loop observables: the $g - 2$ case*, *JHEP* **10** (2016) 002 [[1607.07621](#)].
- [107] G. Couture and H. Konig, *Bounds on second generation scalar leptoquarks from the anomalous magnetic moment of the muon*, *Phys. Rev. D* **53** (1996) 555 [[hep-ph/9507263](#)].
- [108] U. Mahanta, *Implications of BNL measurement of delta $a(\mu)$ on a class of scalar leptoquark interactions*, *Eur. Phys. J. C* **21** (2001) 171 [[hep-ph/0102176](#)].
- [109] F. S. Queiroz, K. Sinha and A. Strumia, *Leptoquarks, Dark Matter, and Anomalous LHC Events*, *Phys. Rev. D* **91** (2015) 035006 [[1409.6301](#)].
- [110] E. Coluccio Leskow, G. D'Ambrosio, A. Crivellin and D. Müller, *$(g - 2)\mu$, lepton flavor violation, and Z decays with leptoquarks: Correlations and future prospects*, *Phys. Rev. D* **95** (2017) 055018 [[1612.06858](#)].
- [111] C.-H. Chen, T. Nomura and H. Okada, *Excesses of muon $g - 2$, $R_{D^{(*)}}$, and R_K in a leptoquark model*, *Phys. Lett. B* **774** (2017) 456 [[1703.03251](#)].
- [112] D. Das, C. Hati, G. Kumar and N. Mahajan, *Towards a unified explanation of $R_{D^{(*)}}$, R_K and $(g - 2)_\mu$ anomalies in a left-right model with leptoquarks*, *Phys. Rev. D* **94** (2016) 055034 [[1605.06313](#)].
- [113] A. Crivellin, M. Hoferichter and P. Schmidt-Wellenburg, *Combined explanations of $(g - 2)_{\mu,e}$ and implications for a large muon EDM*, *Phys. Rev. D* **98** (2018) 113002 [[1807.11484](#)].
- [114] K. Kowalska, E. M. Sessolo and Y. Yamamoto, *Constraints on charmphilic solutions to the muon $g - 2$ with leptoquarks*, *Phys. Rev. D* **99** (2019) 055007 [[1812.06851](#)].
- [115] I. Doršner, S. Fajfer and O. Sumensari, *Muon $g - 2$ and scalar leptoquark mixing*, *JHEP* **06** (2020) 089 [[1910.03877](#)].
- [116] L. Delle Rose, C. Marzo and L. Marzola, *Simplified leptoquark models for precision $l_i \rightarrow l_f \gamma$ experiments: two-loop structure of $O(\alpha_S Y^2)$ corrections*, *Phys. Rev. D* **102** (2020) 115020 [[2005.12389](#)].
- [117] I. Bigaran and R. R. Volkas, *Getting chirality right: Single scalar leptoquark solutions to the $(g - 2)_{e,\mu}$ puzzle*, *Phys. Rev. D* **102** (2020) 075037 [[2002.12544](#)].
- [118] I. Doršner, S. Fajfer and S. Saad, *$\mu \rightarrow e\gamma$ selecting scalar leptoquark solutions for the $(g - 2)_{e,\mu}$ puzzles*, *Phys. Rev. D* **102** (2020) 075007 [[2006.11624](#)].
- [119] K. S. Babu, P. S. B. Dev, S. Jana and A. Thapa, *Unified framework for B-anomalies, muon $g - 2$ and neutrino masses*, *JHEP* **03** (2021) 179 [[2009.01771](#)].
- [120] J. Aebischer, W. Dekens, E. E. Jenkins, A. V. Manohar, D. Sengupta and P. Stoffer, *Effective field theory interpretation of lepton magnetic and electric dipole moments*, *JHEP* **07** (2021) 107 [[2102.08954](#)].
- [121] A. Crivellin, D. Mueller and F. Saturnino, *Correlating $h \rightarrow \mu^+ \mu^-$ to the Anomalous Magnetic Moment of the Muon via Leptoquarks*, *Phys. Rev. Lett.* **127** (2021) 021801 [[2008.02643](#)].

- [122] I. Bigaran, T. Felkl, C. Hagedorn and M. A. Schmidt, *Flavour anomalies meet flavour symmetry*, [2207.06197](#).
- [123] D. Stöckinger and H. Stöckinger-Kim, *On the role of chirality flips for the muon magnetic moment and its relation to the muon mass*, *Front. in Phys.* **10** (2022) 944614.
- [124] S. Parashar, A. Karan, Avnish, P. Bandyopadhyay and K. Ghosh, *Phenomenology of scalar leptoquarks at the LHC in explaining the radiative neutrino masses, muon g-2, and lepton flavor violating observables*, *Phys. Rev. D* **106** (2022) 095040 [[2209.05890](#)].
- [125] S.-L. Chen, W.-w. Jiang and Z.-K. Liu, *Combined explanations of B-physics anomalies, ($g - 2$) _{e,μ} and neutrino masses by scalar leptoquarks*, *Eur. Phys. J. C* **82** (2022) 959 [[2205.15794](#)].
- [126] M. Kramer, T. Plehn, M. Spira and P. M. Zerwas, *Pair production of scalar leptoquarks at the Tevatron*, *Phys. Rev. Lett.* **79** (1997) 341 [[hep-ph/9704322](#)].
- [127] M. Kramer, T. Plehn, M. Spira and P. M. Zerwas, *Pair production of scalar leptoquarks at the CERN LHC*, *Phys. Rev. D* **71** (2005) 057503 [[hep-ph/0411038](#)].
- [128] D. A. Faroughy, A. Greljo and J. F. Kamenik, *Confronting lepton flavor universality violation in B decays with high- p_T tau lepton searches at LHC*, *Phys. Lett. B* **764** (2017) 126 [[1609.07138](#)].
- [129] A. Greljo and D. Marzocca, *High- p_T dilepton tails and flavor physics*, *Eur. Phys. J. C* **77** (2017) 548 [[1704.09015](#)].
- [130] J. Blumlein, E. Boos and A. Kryukov, *Leptoquark pair production in hadronic interactions*, *Z. Phys. C* **76** (1997) 137 [[hep-ph/9610408](#)].
- [131] I. Doršner, S. Fajfer, D. A. Faroughy and N. Košník, *The role of the S_3 GUT leptoquark in flavor universality and collider searches*, *JHEP* **10** (2017) 188 [[1706.07779](#)].
- [132] A. Cerri et al., *Report from Working Group 4: Opportunities in Flavour Physics at the HL-LHC and HE-LHC*, *CERN Yellow Rep. Monogr.* **7** (2019) 867 [[1812.07638](#)].
- [133] P. Bandyopadhyay and R. Mandal, *Revisiting scalar leptoquark at the LHC*, *Eur. Phys. J. C* **78** (2018) 491 [[1801.04253](#)].
- [134] G. Hiller, D. Loose and I. Nišandžić, *Flavorful leptoquarks at hadron colliders*, *Phys. Rev. D* **97** (2018) 075004 [[1801.09399](#)].
- [135] T. Faber, M. Hudec, H. Kolešová, Y. Liu, M. Malinský, W. Porod et al., *Collider phenomenology of a unified leptoquark model*, *Phys. Rev. D* **101** (2020) 095024 [[1812.07592](#)].
- [136] M. Schmaltz and Y.-M. Zhong, *The leptoquark Hunter's guide: large coupling*, *JHEP* **01** (2019) 132 [[1810.10017](#)].
- [137] K. Chandak, T. Mandal and S. Mitra, *Hunting for scalar leptoquarks with boosted tops and light leptons*, *Phys. Rev. D* **100** (2019) 075019 [[1907.11194](#)].
- [138] B. C. Allanach, T. Corbett and M. Madigan, *Sensitivity of Future Hadron Colliders to Leptoquark Pair Production in the Di-Muon Di-Jets Channel*, *Eur. Phys. J. C* **80** (2020) 170 [[1911.04455](#)].

- [139] L. Buonocore, U. Haisch, P. Nason, F. Tramontano and G. Zanderighi, *Lepton-Quark Collisions at the Large Hadron Collider*, *Phys. Rev. Lett.* **125** (2020) 231804 [[2005.06475](#)].
- [140] U. Haisch and G. Polesello, *Resonant third-generation leptoquark signatures at the Large Hadron Collider*, *JHEP* **05** (2021) 057 [[2012.11474](#)].
- [141] C. Borschensky, B. Fuks, A. Kulesza and D. Schwartländer, *Scalar leptoquark pair production at hadron colliders*, *Phys. Rev. D* **101** (2020) 115017 [[2002.08971](#)].
- [142] A. Crivellin, D. Müller and L. Schnell, *Combined constraints on first generation leptoquarks*, *Phys. Rev. D* **103** (2021) 115023 [[2104.06417](#)].
- [143] T. Nomura and H. Okada, *Explanations for anomalies of muon anomalous magnetic dipole moment, $b \rightarrow s\mu\mu^-$, and radiative neutrino masses in a leptoquark model*, *Phys. Rev. D* **104** (2021) 035042 [[2104.03248](#)].
- [144] K. Ban, Y. Jho, Y. Kwon, S. C. Park, S. Park and P.-Y. Tseng, *A comprehensive study of the vector leptoquark with $U(1)$ $B_3 - L_2$ on the B -meson and muon $g - 2$ anomalies*, *PTEP* **2023** (2023) 013B01 [[2104.06656](#)].
- [145] P. Fileviez Perez, C. Murgui and A. D. Plascencia, *Leptoquarks and matter unification: Flavor anomalies and the muon $g - 2$* , *Phys. Rev. D* **104** (2021) 035041 [[2104.11229](#)].
- [146] X. Wang, *Muon $g - 2$ and flavor puzzles in the $U(1)_X$ -gauged leptoquark model*, *JHEP* **08** (2022) 243 [[2108.01279](#)].
- [147] K. Cheung, W.-Y. Keung and P.-Y. Tseng, *Isodoublet vector leptoquark solution to the muon $g - 2$, RK, K^*, RD, D^* , and W -mass anomalies*, *Phys. Rev. D* **106** (2022) 015029 [[2204.05942](#)].
- [148] N. Ghosh, S. K. Rai and T. Samui, *Collider signatures of a scalar leptoquark and vectorlike lepton in light of muon anomaly*, *Phys. Rev. D* **107** (2023) 035028 [[2206.11718](#)].
- [149] C. Borschensky, B. Fuks, A. Jueid and A. Kulesza, *Scalar leptoquarks at the LHC and flavour anomalies: a comparison of pair-production modes at NLO-QCD*, *JHEP* **11** (2022) 006 [[2207.02879](#)].
- [150] I. Doršner, A. Lejlić and S. Saad, *Asymmetric leptoquark pair production at LHC*, *JHEP* **03** (2023) 025 [[2210.11004](#)].
- [151] E. Keith and E. Ma, *S , T , and leptoquarks at HERA*, *Phys. Rev. Lett.* **79** (1997) 4318 [[hep-ph/9707214](#)].
- [152] I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik and N. Košnik, *Physics of leptoquarks in precision experiments and at particle colliders*, *Phys. Rept.* **641** (2016) 1 [[1603.04993](#)].
- [153] A. Bhaskar, D. Das, B. De and S. Mitra, *Enhancing scalar productions with leptoquarks at the LHC*, *Phys. Rev. D* **102** (2020) 035002 [[2002.12571](#)].
- [154] J. Zhang, C.-X. Yue, C.-H. Li and S. Yang, *Constraints on scalar and vector leptoquarks from the LHC Higgs data*, [\[1905.04074\]\(#\)](#).
- [155] V. Gherardi, D. Marzocca and E. Venturini, *Matching scalar leptoquarks to the SMEFT at one loop*, *JHEP* **07** (2020) 225 [[2003.12525](#)].

- [156] A. Crivellin, D. Müller and F. Saturnino, *Leptoquarks in oblique corrections and Higgs signal strength: status and prospects*, *JHEP* **11** (2020) 094 [[2006.10758](#)].
- [157] F. Albergaria and L.avoura, *Oblique corrections from leptoquarks*, [2301.03024](#).
- [158] C. Marzo, *Beyond oblique contributions: Δr and W-mass in models with scalar leptoquarks*, [2212.11174](#).
- [159] W. Dekens, J. de Vries, M. Jung and K. K. Vos, *The phenomenology of electric dipole moments in models of scalar leptoquarks*, *JHEP* **01** (2019) 069 [[1809.09114](#)].
- [160] A. Crivellin and F. Saturnino, *Correlating tauonic B decays with the neutron electric dipole moment via a scalar leptoquark*, *Phys. Rev. D* **100** (2019) 115014 [[1905.08257](#)].
- [161] I. Dorsner, S. Fajfer and N. Kosnik, *Heavy and light scalar leptoquarks in proton decay*, *Phys. Rev. D* **86** (2012) 015013 [[1204.0674](#)].
- [162] I. Doršner, S. Fajfer and O. Sumensari, *Triple-leptoquark interactions for tree- and loop-level proton decays*, *JHEP* **05** (2022) 183 [[2202.08287](#)].
- [163] C. A. Manzari and S. Profumo, *Flavor inspired model for dark matter*, *Phys. Rev. D* **106** (2022) 075025 [[2206.06768](#)].
- [164] R. Mandal, *Fermionic dark matter in leptoquark portal*, *Eur. Phys. J. C* **78** (2018) 726 [[1808.07844](#)].
- [165] G. Bélanger, J. Kumar, D. London and A. Pukhov, *The B anomalies, the U_1 leptoquark and dark matter*, *JHEP* **01** (2023) 041 [[2206.11305](#)].
- [166] B. Belfatto, D. Buttazzo, C. Gross, P. Panci, A. Strumia, N. Vignaroli et al., *Dark Matter abundance via thermal decays and leptoquark mediators*, *JHEP* **06** (2022) 084 [[2111.14808](#)].
- [167] G. Belanger et al., *Leptoquark manoeuvres in the dark: a simultaneous solution of the dark matter problem and the $R_{D^{(*)}}$ anomalies*, *JHEP* **02** (2022) 042 [[2111.08027](#)].
- [168] S. Singirala, S. Sahoo and R. Mohanta, *Light dark matter, rare B decays with missing energy in $L\mu-L\tau$ model with a scalar leptoquark*, *Phys. Rev. D* **105** (2022) 015033 [[2106.03735](#)].
- [169] S. Sahoo, S. Singirala and R. Mohanta, *Dark matter and flavor anomalies in the light of vector-like fermions and scalar leptoquark*, [2112.04382](#).
- [170] A. Mohamadnejad, *Accidental scale-invariant Majorana dark matter in leptoquark-Higgs portals*, *Nucl. Phys. B* **949** (2019) 114793 [[1904.03857](#)].
- [171] U. Mahanta, *Neutrino masses and mixing angles from leptoquark interactions*, *Phys. Rev. D* **62** (2000) 073009 [[hep-ph/9909518](#)].
- [172] F. F. Deppisch, S. Kulkarni, H. Päs and E. Schumacher, *Leptoquark patterns unifying neutrino masses, flavor anomalies, and the diphoton excess*, *Phys. Rev. D* **94** (2016) 013003 [[1603.07672](#)].
- [173] I. Doršner, S. Fajfer and N. Košnik, *Leptoquark mechanism of neutrino masses within the grand unification framework*, *Eur. Phys. J. C* **77** (2017) 417 [[1701.08322](#)].
- [174] K. Cheung, T. Nomura and H. Okada, *A Three-loop Neutrino Model with Leptoquark Triplet Scalars*, *Phys. Lett. B* **768** (2017) 359 [[1701.01080](#)].

- [175] ATLAS collaboration, *Search for leptoquark pair production decaying into $te^- \bar{t}e^+$ or $t\mu^- \bar{t}\mu^+$ in multi-lepton final states in pp collisions at 13 TeV with the ATLAS detector*, [2306.17642](#).
- [176] ATLAS collaboration, *Search for leptoquarks decaying into the $b\tau$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [2305.15962](#).
- [177] ATLAS collaboration, *Search for excited τ -leptons and leptoquarks in the final state with τ -leptons and jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **06** (2023) 199 [[2303.09444](#)].
- [178] ATLAS collaboration, *Search for pair production of third-generation leptoquarks decaying into a bottom quark and a τ -lepton with the ATLAS detector*, [2303.01294](#).
- [179] ATLAS collaboration, *Search for pair-produced scalar and vector leptoquarks decaying into third-generation quarks and first- or second-generation leptons in pp collisions with the ATLAS detector*, *JHEP* **2306** (2023) 188 [[2210.04517](#)].
- [180] ATLAS collaboration, *Search for pair production of third-generation scalar leptoquarks decaying into a top quark and a τ -lepton in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **06** (2021) 179 [[2101.11582](#)].
- [181] ATLAS collaboration, *Search for pair production of scalar leptoquarks decaying into first- or second-generation leptons and top quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 313 [[2010.02098](#)].
- [182] ATLAS collaboration, *Search for pairs of scalar leptoquarks decaying into quarks and electrons or muons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *JHEP* **10** (2020) 112 [[2006.05872](#)].
- [183] ATLAS collaboration, *Searches for third-generation scalar leptoquarks in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *JHEP* **06** (2019) 144 [[1902.08103](#)].
- [184] ATLAS collaboration, *Searches for scalar leptoquarks and differential cross-section measurements in dilepton-dijet events in proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 13$ TeV with the ATLAS experiment*, *Eur. Phys. J. C* **79** (2019) 733 [[1902.00377](#)].
- [185] CMS collaboration, *Searches for additional Higgs bosons and for vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, [2208.02717](#).
- [186] CMS collaboration, *Search for singly and pair-produced leptoquarks coupling to third-generation fermions in proton-proton collisions at $s=13$ TeV*, *Phys. Lett. B* **819** (2021) 136446 [[2012.04178](#)].
- [187] CMS collaboration, *Search for dark matter in events with a leptoquark and missing transverse momentum in proton-proton collisions at 13 TeV*, *Phys. Lett. B* **795** (2019) 76 [[1811.10151](#)].
- [188] CMS collaboration, *Search for pair production of first-generation scalar leptoquarks at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **99** (2019) 052002 [[1811.01197](#)].
- [189] CMS collaboration, *Search for heavy neutrinos and third-generation leptoquarks in hadronic states of two τ leptons and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2019) 170 [[1811.00806](#)].

- [190] CMS collaboration, *Search for leptoquarks coupled to third-generation quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **121** (2018) 241802 [[1809.05558](#)].
- [191] CMS collaboration, *Search for pair production of second-generation leptoquarks at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **99** (2019) 032014 [[1808.05082](#)].
- [192] CMS collaboration, *Search for a singly produced third-generation scalar leptoquark decaying to a τ lepton and a bottom quark in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2018) 115 [[1806.03472](#)].
- [193] CMS collaboration, *Constraints on models of scalar and vector leptoquarks decaying to a quark and a neutrino at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **98** (2018) 032005 [[1805.10228](#)].
- [194] CMS collaboration, *Search for third-generation scalar leptoquarks decaying to a top quark and a τ lepton at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **78** (2018) 707 [[1803.02864](#)].
- [195] A. Preda, G. Senjanovic and M. Zantedeschi, *$SO(10)$: A case for hadron colliders*, *Phys. Lett. B* **838** (2023) 137746 [[2201.02785](#)].
- [196] H. Murayama and T. Yanagida, *A viable $SU(5)$ GUT with light leptoquark bosons*, *Mod. Phys. Lett. A* **7** (1992) 147.
- [197] S. Banik and A. Crivellin, “Renormalization Group Equations of Scalar Leptoquarks.” <https://github.com/SumitBanikGit/SLQ-RG>, 2023.
- [198] A. Crivellin and L. Schnell, *Complete Lagrangian and set of Feynman rules for scalar leptoquarks*, *Comput. Phys. Commun.* **271** (2022) 108188 [[2105.04844](#)].
- [199] L. Sartore and I. Schienbein, *PyR@TE 3*, *Comput. Phys. Commun.* **261** (2021) 107819 [[2007.12700](#)].
- [200] A. E. Thomsen, *Introducing RGBeta: a Mathematica package for the evaluation of renormalization group β -functions*, *Eur. Phys. J. C* **81** (2021) 408 [[2101.08265](#)].
- [201] J. Fuentes-Martín, M. König, J. Pagès, A. E. Thomsen and F. Wilsch, *A Proof of Concept for Matchete: An Automated Tool for Matching Effective Theories*, [2212.04510](#).
- [202] A. Carmona, A. Lazopoulos, P. Olgoso and J. Santiago, *Matchmakereft: automated tree-level and one-loop matching*, *SciPost Phys.* **12** (2022) 198 [[2112.10787](#)].
- [203] T. Plehn, H. Spiesberger, M. Spira and P. M. Zerwas, *Formation and decay of scalar leptoquarks / squarks in e p collisions*, *Z. Phys. C* **74** (1997) 611 [[hep-ph/9703433](#)].
- [204] S. P. Martin and D. G. Robertson, *Standard model parameters in the tadpole-free pure $\overline{\text{MS}}$ scheme*, *Phys. Rev. D* **100** (2019) 073004 [[1907.02500](#)].
- [205] K. G. Chetyrkin, J. H. Kuhn and M. Steinhauser, *RunDec: A Mathematica package for running and decoupling of the strong coupling and quark masses*, *Comput. Phys. Commun.* **133** (2000) 43 [[hep-ph/0004189](#)].
- [206] F. Herren and M. Steinhauser, *Version 3 of RunDec and CRUnDec*, *Comput. Phys. Commun.* **224** (2018) 333 [[1703.03751](#)].
- [207] B. A. Kniehl, A. F. Pikelner and O. L. Veretin, *mr: a C++ library for the matching*

and running of the Standard Model parameters, *Comput. Phys. Commun.* **206** (2016) 84 [[1601.08143](#)].

- [208] G.-y. Huang and S. Zhou, *Precise Values of Running Quark and Lepton Masses in the Standard Model*, *Phys. Rev. D* **103** (2021) 016010 [[2009.04851](#)].
- [209] G. Degrassi, P. Gambino and P. P. Giardino, *The $m_w - m_z$ interdependence in the Standard Model: a new scrutiny*, *JHEP* **05** (2015) 154 [[1411.7040](#)].
- [210] G. Degrassi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori et al., *Higgs mass and vacuum stability in the Standard Model at NNLO*, *JHEP* **08** (2012) 098 [[1205.6497](#)].
- [211] T. P. Cheng, E. Eichten and L.-F. Li, *Higgs Phenomena in Asymptotically Free Gauge Theories*, *Phys. Rev. D* **9** (1974) 2259.
- [212] H. Arason, D. J. Castano, B. Keszthelyi, S. Mikaelian, E. J. Piard, P. Ramond et al., *Renormalization group study of the standard model and its extensions. 1. The Standard model*, *Phys. Rev. D* **46** (1992) 3945.
- [213] C. Ford, D. R. T. Jones, P. W. Stephenson and M. B. Einhorn, *The Effective potential and the renormalization group*, *Nucl. Phys. B* **395** (1993) 17 [[hep-lat/9210033](#)].
- [214] M.-x. Luo and Y. Xiao, *Two loop renormalization group equations in the standard model*, *Phys. Rev. Lett.* **90** (2003) 011601 [[hep-ph/0207271](#)].
- [215] O. V. Tarasov, *Anomalous dimensions of quark masses in the three-loop approximation*, *Phys. Part. Nucl. Lett.* **17** (2020) 109 [[1910.12231](#)].
- [216] L. N. Mihaila, J. Salomon and M. Steinhauser, *Gauge Coupling Beta Functions in the Standard Model to Three Loops*, *Phys. Rev. Lett.* **108** (2012) 151602 [[1201.5868](#)].
- [217] A. V. Bednyakov, A. F. Pikelner and V. N. Velizhanin, *Yukawa coupling beta-functions in the Standard Model at three loops*, *Phys. Lett. B* **722** (2013) 336 [[1212.6829](#)].
- [218] A. V. Bednyakov, A. F. Pikelner and V. N. Velizhanin, *Higgs self-coupling beta-function in the Standard Model at three loops*, *Nucl. Phys. B* **875** (2013) 552 [[1303.4364](#)].
- [219] T. van Ritbergen, J. A. M. Vermaasen and S. A. Larin, *The Four loop beta function in quantum chromodynamics*, *Phys. Lett. B* **400** (1997) 379 [[hep-ph/9701390](#)].
- [220] M. Czakon, *The Four-loop QCD beta-function and anomalous dimensions*, *Nucl. Phys. B* **710** (2005) 485 [[hep-ph/0411261](#)].
- [221] P. A. Baikov, K. G. Chetyrkin and J. H. Kühn, *Five-Loop Running of the QCD coupling constant*, *Phys. Rev. Lett.* **118** (2017) 082002 [[1606.08659](#)].
- [222] T. Luthe, A. Maier, P. Marquard and Y. Schroder, *Complete renormalization of QCD at five loops*, *JHEP* **03** (2017) 020 [[1701.07068](#)].
- [223] F. Herzog, B. Ruijl, T. Ueda, J. A. M. Vermaasen and A. Vogt, *The five-loop beta function of Yang-Mills theory with fermions*, *JHEP* **02** (2017) 090 [[1701.01404](#)].
- [224] F. Jegerlehner and M. Y. Kalmykov, *$O(\alpha, \alpha_s)$ correction to the pole mass of the t quark within the standard model*, *Nucl. Phys. B* **676** (2004) 365 [[hep-ph/0308216](#)].

- [225] P. Marquard, A. V. Smirnov, V. A. Smirnov and M. Steinhauser, *Quark Mass Relations to Four-Loop Order in Perturbative QCD*, *Phys. Rev. Lett.* **114** (2015) 142002 [[1502.01030](#)].
- [226] P. Marquard, A. V. Smirnov, V. A. Smirnov, M. Steinhauser and D. Wellmann, *$\overline{\text{MS}}$ -on-shell quark mass relation up to four loops in QCD and a general $SU(N)$ gauge group*, *Phys. Rev. D* **94** (2016) 074025 [[1606.06754](#)].
- [227] S. P. Martin, *Top-quark pole mass in the tadpole-free $\overline{\text{MS}}$ scheme*, *Phys. Rev. D* **93** (2016) 094017 [[1604.01134](#)].
- [228] N. Gray, D. J. Broadhurst, W. Grafe and K. Schilcher, *Three Loop Relation of Quark (Modified) Ms and Pole Masses*, *Z. Phys. C* **48** (1990) 673.
- [229] S. P. Martin and D. G. Robertson, *Higgs boson mass in the Standard Model at two-loop order and beyond*, *Phys. Rev. D* **90** (2014) 073010 [[1407.4336](#)].
- [230] S. P. Martin, *Z-Boson Pole Mass at Two-Loop Order in the Pure $\overline{\text{MS}}$ Scheme*, *Phys. Rev. D* **92** (2015) 014026 [[1505.04833](#)].
- [231] CMS collaboration, *Measurement of the top quark mass using events with a single reconstructed top quark in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **12** (2021) 161 [[2108.10407](#)].
- [232] CMS collaboration, *Measurement of the top quark mass using proton-proton data at $\sqrt{(s)} = 7$ and 8 TeV*, *Phys. Rev. D* **93** (2016) 072004 [[1509.04044](#)].
- [233] CMS collaboration, *Measurement of the $t\bar{t}$ production cross section, the top quark mass, and the strong coupling constant using dilepton events in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 368 [[1812.10505](#)].
- [234] K. G. Chetyrkin, J. H. Kuhn, A. Maier, P. Maierhofer, P. Marquard, M. Steinhauser et al., *Charm and Bottom Quark Masses: An Update*, *Phys. Rev. D* **80** (2009) 074010 [[0907.2110](#)].
- [235] M. Beneke, A. Maier, J. Piclum and T. Rauh, *The bottom-quark mass from non-relativistic sum rules at NNNLO*, *Nucl. Phys. B* **891** (2015) 42 [[1411.3132](#)].
- [236] S. Narison, *\overline{m}_c and \overline{m}_b from M_{Bc} and improved estimates of f_{Bc} and $f_{Bc(2S)}$* , *Phys. Lett. B* **802** (2020) 135221 [[1906.03614](#)].
- [237] BESIII collaboration, *Precision measurement of the mass of the τ lepton*, *Phys. Rev. D* **90** (2014) 012001 [[1405.1076](#)].
- [238] ARGUS collaboration, *A Measurement of the tau mass*, *Phys. Lett. B* **292** (1992) 221.
- [239] SUPER-KAMIOKANDE collaboration, *Search for proton decay via $p \rightarrow e^+\pi^0$ and $p \rightarrow \mu^+\pi^0$ with an enlarged fiducial volume in Super-Kamiokande I-IV*, *Phys. Rev. D* **102** (2020) 112011 [[2010.16098](#)].
- [240] P. Bandyopadhyay, S. Jangid and A. Karan, *Constraining scalar doublet and triplet leptons with vacuum stability and perturbativity*, *Eur. Phys. J. C* **82** (2022) 516 [[2111.03872](#)].
- [241] A. Crivellin, M. Hoferichter and J. M. Roney, *Toward testing the magnetic moment of the tau at one part per million*, *Phys. Rev. D* **106** (2022) 093007 [[2111.10378](#)].
- [242] A. Crivellin and M. Hoferichter, *The Anomalous Magnetic Moment of the Muon: Beyond the Standard Model via Chiral Enhancement*, 7, 2022, [2207.01912](#).