

Search for Light Dark Photon in the Forward Experiments at the LHC

Yeong Gyun Kim^{a,*}, Kang Young Lee^{b,†}, and Soo-hyeon Nam^{c,‡}

^aDepartment of Science Education, Gwangju National University of Education,
Gwangju 61204, Korea

^bDepartment of Physics Education & Research Institute of Natural Science,
Gyeongsang National University, Jinju 52828, Korea

^cDepartment of Physics, Korea University, Seoul 02841, Korea

Abstract

We investigate detection possibility of light dark photon in the forward experiments at the LHC, such as the SND@LHC and the FASER experiments. We assume that the dark photon mass is smaller than twice of the electron mass. Then the dark photon is long-lived and copiously produced through a neutral pion decay. Such dark photons would easily pass through 100 m of rock in front of the forward experiments and the detector targets, but some portion of them could be converted into an electron-positron pair inside the detector leaving an isolated electromagnetic shower as a new physics signature of the dark photon. Our estimation shows that in the range of kinetic mixing parameter $4 \times 10^{-5} \lesssim \epsilon \lesssim 2 \times 10^{-1}$, more than 10 signal events of the dark photon can be produced assuming 150 fb^{-1} integrated luminosity.

*ygkim@gnue.ac.kr

†kylee.phys@gnu.ac.kr, co-corresponding author

‡glvnsh@gmail.com, corresponding author

Contents

1	Introduction	1
2	A Dark Photon Model	2
3	Dark Photon Production	3
4	Dark Photon Detection	5
5	Conclusion	7

1 Introduction

The ATLAS and CMS detectors at the Large Hadron Collider (LHC) in CERN have been successfully operating and played pivotal roles in discovering Higgs boson in 2012 [1, 2]. While these detectors are well-equipped at large angles relative to the beamline and therefore capable of detecting new physics with high transverse momentum p_T , they have holes in the far-forward direction. This limitation could potentially result in missed opportunities to detect (new) particles produced in the far-forward region.

To address this limitation, the FASER [3] and the SND@LHC [4] experiments were designed to detect high-energy neutrinos and explore new particles that are produced in the far-forward direction at the LHC. These experiments are located 480 meters downstream of the ATLAS interaction point (IP) along the beam collision axis. Charged particles from the IP are deflected by LHC magnets, while the remaining particles are further shielded by the 100m of rock and concrete in front of the detectors. However, neutrinos and feebly interacting new particles would easily pass through this shielding and reach the detectors.

An example of new particles that can be probed in the forward experiments is a dark photon A' , a hypothetical particle which interacts electromagnetically with standard model particles, but with a suppressed interaction strength α' compared to ordinary fine structure constant α . If the dark photon mass $m_{A'}$ is small enough, the main production channel of the dark photons at the LHC is through the decays of mesons, such as neutral pion π^0 . Then, a significant number of dark photons would be produced in the far-forward direction, depending on the interaction strength. If the mass of the dark photon is less than twice the mass of an electron, the dark photon does not decay. With the suppressed interaction strength α' , the majority of the produced dark photons would pass through the 100 m of rock in front of the forward experiments and reach the detector.

When the *light* dark photons pass through detector target material (i.e. tungsten for the forward experiments), they can be converted into a pair of electron and positron, similarly to ordinary energetic photons, but with a much suppressed interaction rate. Therefore the signal signature of the dark photon at the forward experiments would be an isolated electromagnetic (EM) shower, whose shower shape is essentially the same as the one from

ordinary energetic photon. If an ordinary photon enters the tungsten target, the EM shower will start near the detector entrance because the radiation length X_0 of tungsten is just 3.5 mm. On the other hand, if the EM shower of the dark photon occurs in the detector target, it would start at any location within the detector. This is because the ‘effective’ radiation length of the detector material for the dark photon increases by a factor of $1/\epsilon^2 \equiv \alpha/\alpha'$. If $1/\epsilon^2 = 10^4$, for instance, then the effective radiation length of tungsten for the dark photon becomes 35 meters, which is much longer than the actual detector target length (i.e. ~ 30 cm for the SND@LHC detector).

In this paper, we will investigate the detection possibility of the light dark photon in the SND@LHC experiment at the LHC. In section 2, we introduce a dark photon model. We discuss dark photon production and detection in section 3 and 4, respectively. We conclude in section 5.

2 A Dark Photon Model

It might be possible that the standard model (SM) is accompanied by additional gauge structures, such as a new $U(1)_X$ gauge group with a gauge field X . The gauge field X can mix with the SM $U(1)_Y$ gauge field Y through a renormalizable kinetic mixing term $X_{\mu\nu}Y^{\mu\nu}$, where $X_{\mu\nu}$ and $Y_{\mu\nu}$ are the field strengths of X and Y gauge fields, respectively. If the SM particles are all uncharged under the $U(1)_X$, the relevant Lagrangian for the X and Y gauge fields, including a mass term for X , can be written as

$$\mathcal{L} = -\frac{1}{4}X_{\mu\nu}X^{\mu\nu} - \frac{\sigma}{2}X_{\mu\nu}Y^{\mu\nu} - \frac{1}{4}Y_{\mu\nu}Y^{\mu\nu} + j_Y^\mu Y_\mu + \frac{1}{2}m_X^2 X_\mu X^\mu, \quad (1)$$

where j_Y^μ denotes interaction current of gauge field Y .¹

In order to obtain the Lagrangian in the canonical form, we redefine the gauge fields as follows

$$\hat{Y}_\mu = Y_\mu + \sigma X_\mu, \quad \hat{X}_\mu = \sqrt{1 - \sigma^2} X_\mu. \quad (2)$$

Then the Lagrangian is given by

$$\mathcal{L} = -\frac{1}{4}\hat{X}_{\mu\nu}\hat{X}^{\mu\nu} - \frac{1}{4}\hat{Y}_{\mu\nu}\hat{Y}^{\mu\nu} + j_Y^\mu (\hat{Y}_\mu - \frac{\sigma}{\sqrt{1 - \sigma^2}}\hat{X}_\mu) + \frac{1}{2}\frac{m_X^2}{(1 - \sigma^2)}\hat{X}_\mu\hat{X}^\mu. \quad (3)$$

Here we will focus only on the EM interaction in the SM. If we identify the massless gauge boson \hat{Y} as the SM photon field A and the massive gauge boson \hat{X} as the dark photon field A' , the resulting Lagrangian can be written as [7]

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} + \frac{1}{2}m_{A'}^2 A'^2 + j_{em}^\mu (A_\mu + \epsilon A'_\mu), \quad (4)$$

¹The Lagrangian in Eq. (1) seems not manifestly gauge-invariant due to the mass parameter of the $U(1)_X$ gauge boson, m_X . Nonetheless, one can make it gauge-invariant and renormalizable by adopting the Stueckelberg action which has an additional axionic scalar [5]. Then, the topological mass parameter m_X denotes the Stueckelberg coupling of the axionic scalar to the $U(1)_X$ gauge boson, and our Lagrangian corresponds to the case that the Stueckelberg coupling to the $U(1)_Y$ gauge boson is zero in the Stueckelberg extension of the SM with kinetic mixing [6].

where $F_{\mu\nu}$ and $F'_{\mu\nu}$ are the field strengths of the photon A and dark photon A' , respectively, $m_{A'}$ is the mass of the dark photon, and j_{em}^μ is the electromagnetic (EM) current of the SM particles. The dark photon couples to the EM current with a suppressed coupling ϵ compared to the ordinary photon coupling. Therefore, we have two new free parameters $m_{A'}$ and ϵ in this dark photon model, in addition to the SM parameters. The Lagrangian in Eq.(4) is the starting point of our analysis.

3 Dark Photon Production

The LHC at 13 TeV center of mass energy has a total inelastic proton-proton scattering cross section of around 75 mb, as reported by ATLAS and CMS Collaborations [8, 9]. It implies the production of large number of particles that are highly concentrated in the far-forward direction. For example, each hemisphere is expected to have approximately $n_{\pi^0} \approx 2.1 \times 10^{17}$ neutral pions for the 150 fb^{-1} of integrated luminosity. Roughly 11% of all neutral pions in one hemisphere are produced within the pseudo-rapidity range (i.e. $7.2 < \eta < 8.4$) of the SND@LHC experiment. Fig. 1(a) shows the neutral pion production rate in one hemisphere in the (θ_π, p_π) plane, where θ_π and p_π are the meson's angle with respect to the beam axis and momentum, respectively. The neutral pion spectrum was obtained with the Monte Carlo generator EPOS-LHC [10] implemented in the CRMC simulation package [11].


If the dark photon mass is smaller than neutral pion mass, dominant production channel of the dark photons is neutral pion decay, $\pi^0 \rightarrow A'\gamma$. The branching ratio of the decay mode is given by [12]

$$B(\pi^0 \rightarrow A'\gamma) = 2\epsilon^2 \left(1 - \frac{m_{A'}^2}{m_{\pi^0}^2}\right) B(\pi^0 \rightarrow \gamma\gamma), \quad (5)$$


where $B(\pi^0 \rightarrow \gamma\gamma) \simeq 0.99$. The neutral pion is a pseudoscalar, and so (dark) photons are produced isotropically in the meson's rest frames. Fig. 1(b) shows photon production rate in one hemisphere in the $(\theta_\gamma, p_\gamma)$ plane. In the figure, in addition to the pion decay, other production channels of photons such as eta meson decay, proton bremsstrahlung and Drell-Yan process etc. are also included. The number of photons in each hemisphere for 150 fb^{-1} integrated luminosity is $n_\gamma \approx 4.5 \times 10^{17}$. About 10% of photons in one hemisphere are produced in the direction of the SND@LHC detector, i.e. $n_\gamma^{\text{SND}} \approx 4.6 \times 10^{16}$. If the dark photon mass is small enough (i.e. $m_{A'} \sim 0$), the spectrum of dark photons is essentially the same as the one of photons, and the number of dark photons in the direction of the SND@LHC detector scales as

$$n_{A'}^{\text{SND}} = \epsilon^2 \times n_\gamma^{\text{SND}} \approx \epsilon^2 \times 4.6 \times 10^{16} \quad (6)$$


assuming the integrated luminosity of 150 fb^{-1} . Note that around 99% (65%) of the dark photons produced along the SND@LHC direction have energies larger than 10 (100) GeV. We also show simulation results with PYTHIA8 in Fig. 1(c) and 1(d) for the neutral pion and photon, respectively. In this case, the number of photons in the direction of SND@LHC detector is similar to EPOS-LHC prediction, but slightly less, i.e. $n_\gamma^{\text{SND}} \approx 2.6 \times 10^{16}$.




(a) π^0 production by EPOS-LHC



(b) γ production by EPOS-LHC



(c) π^0 production by PYTHIA8



(d) γ production by PYTHIA8

Figure 1: Neutral pion and photon production rates in each hemisphere.

4 Dark Photon Detection

When photons pass through matter, the total number of photons is reduced by the number of interacted photons. The attenuation of photon beam is exponential with respect to travel distance inside absorber,

$$I(x) = I_0 \exp(-\mu x), \quad (7)$$

where I_0 is incident beam intensity, x travel distance. The absorption coefficient μ is given by $\mu = N\sigma$, where N is the density of atoms of the traversed material and σ is the total cross-section per atom. Electron-positron pair production is the most likely photon interaction at high energies. In this case, the absorption coefficient μ is given by $\mu = N\tau_{pair}$ with τ_{pair} , the pair production cross section. For tungsten, the pair production cross section τ_{pair} is about 35 barns/atom and the absorption coefficient is given by

$$\mu = N\tau_{pair} \simeq 2.2/\text{cm} \quad (\text{tungsten}). \quad (8)$$

The absorption coefficient μ is related to the radiation length X_0 of the absorber as follows: $X_0 \simeq 7/9\mu$, which gives $X_0 \simeq 0.35$ cm for the tungsten material.

For light dark photon the pair production cross section is reduced by a factor of ϵ^2 , compared to the photon case. Then the absorption coefficient μ for the dark photon beam is reduced by ϵ^2 accordingly. The photons and dark photons produced in forward direction at the ATLAS IP encounter the 100 m of rock in front of the forward experiment detectors. Therefore, we need to know the absorption coefficient for the rock. The average rock density around CERN is measured to be about 2.5 g/cm³ [13]. The rock can be modeled as a mixture of 41% CaCO₃ and 59% SiO₂ [14]. The radiation lengths are $X_0 \simeq 12.3$ cm for SiO₂ and $X_0 \simeq 8.6$ cm for CaCO₃. From general formula for the radiation length of composite materials, we can obtain the radiation length $X_0 \simeq 10.4$ cm for the rock. In turn, it gives the absorption coefficient μ for the rock,

$$\mu \simeq \frac{7}{9} \times \frac{1}{10.4 \text{ cm}} \simeq 0.075/\text{cm} \quad (\text{rock}). \quad (9)$$

For energetic photons, the attenuation factor $\exp(-\mu x)$ for the 100m of the rock is estimated to be

$$\exp(-0.075/\text{cm} \times 10^4 \text{ cm}) \simeq 0. \quad (10)$$

Therefore, all photons from the ATLAS IP are absorbed by the rock and cannot reach the forward experiment detectors. However, in the case of dark photon, the absorption coefficient μ is reduced by ϵ^2 . For instance, if we assume $\epsilon^2 = 10^{-6}$, the attenuation factor is given by

$$\exp(-\mu x) \simeq \exp(-10^{-6} \times 0.075/\text{cm} \times 10^4 \text{ cm}) \simeq 0.9925 \quad (11)$$

which implies that 99.25% of dark photons just pass through 100 meters of rock and reach the tungsten target of the forward experiment detectors. For the SND@LHC detector, total thickness of the tungsten target is 30 cm. Interaction rates of dark photons inside the tungsten target is given by

$$1 - \exp(-\mu x) = 1 - \exp(-10^{-6} \times 2.2/\text{cm} \times 30 \text{ cm}) \simeq 6.6 \times 10^{-5}. \quad (12)$$




Figure 2: The expected number of events as a function of ϵ , assuming 150 fb^{-1} .

Therefore, the fraction of 6.6×10^{-5} among the dark photons which enter the detector will be detected by the electron-positron pair production inside the detector. For $\epsilon^2 = 10^{-6}$, the number of dark photons which are produced at the ATLAS IP in the direction of the SND@LHC detector is $n_{A'}^{\text{SND}} \simeq 4.6 \times 10^{10}$ for 150 fb^{-1} , as can be seen from the Eq.(6). Then the number of dark photon events inside the detector for 150 fb^{-1} is estimated to be

$$n_{\text{event}} = 4.6 \times 10^{10} \times 0.9925 \times 6.6 \times 10^{-5} \simeq 3.0 \times 10^6. \quad (13)$$

Thus, huge number dark photon events can be obtained with 150 fb^{-1} integrated luminosity, for $\epsilon^2 = 10^{-6}$. For general ϵ values, assuming 150 fb^{-1} , the number of dark photon events in the SND@LHC detector is given by

$$n_{\text{event}} = \epsilon^2 \times 4.6 \times 10^{16} \times \exp(-\epsilon^2 \times 0.075 \times 10^4) \times (1 - \exp(-\epsilon^2 \times 2.2 \times 30)) \quad (14)$$

Fig. 2 shows the number of dark photon events as a function of ϵ . If ϵ is too large, all dark photons will be absorbed by the rock before reaching the detector. On the other hand, if ϵ is too small, all dark photons will just pass through the rock and the detector without leaving any detectable signals in the detector. Therefore, the forward experiments have sensitivity only for a certain range of ϵ . If we require the number of events be larger than 10, the sensitive range of ϵ will be $4 \times 10^{-5} \lesssim \epsilon \lesssim 2 \times 10^{-1}$.

At the detector level, the signature of dark photon signal is an isolated EM shower originated from electron-positron pair production by dark photon interaction with target material. If it were originated from ordinary photon, the isolated EM shower would start at the entrance of the detector because of the small radiation length X_0 . But the starting points of the isolated EM showers originated from dark photons would appear to spread almost equally through detector target, due to the suppressed interaction strength which gives suppressed absorption coefficient μ for dark photon.

A possible dominant background for the dark photon signal would be neutrino-electron scattering $\nu_x e^- \rightarrow \nu_x e^-$, which produces an isolated EM shower. However, it is estimated that the overall background yield is less than one for 150 fb^{-1} integrated luminosity [4]. Furthermore, EM showers originating from (dark) photons are distinguished from those initiated

by electrons with a high efficiency close to 99%. This is mainly because the micrometric accuracy of the nuclear emulsion in the detector is capable of observing the displaced vertex associated with the photon conversion [4]. In this regards, the dark photon signal can be considered as background free.

5 Conclusion

In this work, we investigated detection possibility of light dark photon in the forward experiments at the LHC, especially the SND@LHC. The dark photon interacts with the SM particles through usual EM interaction but with suppressed strength by ϵ and can have an arbitrary mass $m_{A'}$. If dark photon mass is smaller than the neutral pion mass, the dominant production channel of dark photon is a neutral pion decay $\pi^0 \rightarrow \gamma A'$. A copious number of dark photon could be produced from the ATLAS IP, while the exact number of dark photons depends mainly on the new parameter ϵ . About 10% of dark photons in each hemisphere would produce in the pseudo-rapidity range of the SND@LHC experiments. If the dark photon mass is smaller than twice of the electron mass, the dark photons are long-lived. The majority of such dark photons could easily pass through the 100 meters of rock in front of the forward experiments at the LHC, and reach the detector targets of the forward experiments. Although the most of dark photons entering the detector would just pass through the detector, some portion of dark photons could be converted into electron-positron pair leaving an isolated EM shower as a new physics signature of the dark photon. Our estimation shows that in the range of $4 \times 10^{-5} \lesssim \epsilon \lesssim 2 \times 10^{-1}$, more than 10 dark photon events would be produced, assuming 150 fb^{-1} integrated luminosity, and thus it can be detected in the forward experiments at the LHC.

Acknowledgments

This work is supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education under the Grants No. NRF-2021R1F1A1061717 (Y.G.K.) and No. RS-2023-00248860 (S.-h.N.) and also funded by the Ministry of Science and ICT under the Grants No. NRF-2020R1A2C3009918 (S.-h.N.) and No. NRF-2021R1A2C2011003 (K.Y.L.).

References

- [1] **ATLAS** Collaboration, G. Aad *et al.*, “Observation of a New Particle in the Search for the Standard Model Higgs Boson with the ATLAS Detector at the LHC”, *Phys. Lett. B* **716**, 1 (2012).
- [2] **CMS** Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC ”, *Phys. Lett. B* **716**, 30 (2012).
- [3] **FASER** Collaboration, A. Ariga *et al.*, “FASER’s Physics Reach for Long-Lived Particles,” *Phys. Rev. D* **99** (2019) 095011.

- [4] **SND@LHC** Collaboration, C. Ahdida *et al.*, “SND@LHC- Scattering and Neutrino Detector at the LHC”, Tech. Rep. CERN-LHCC-2021-003, LHCCP-016 (CERN, Geneva, 2021).
- [5] E. C. G. Stueckelberg, “Interaction energy in electrodynamics and in the field theory of nuclear forces,” *Helv. Phys. Acta* **11**, 225-244 (1938).
- [6] D. Feldman, Z. Liu and P. Nath, “The Stueckelberg Z-prime Extension with Kinetic Mixing and Milli-Charged Dark Matter From the Hidden Sector,” *Phys. Rev. D* **75**, 115001 (2007).
- [7] M. Fabbrichesi, E. Gabrielli and G. Lanfranchi, “The Dark Photon,” [arXiv:2005.01515](https://arxiv.org/abs/2005.01515) [hep-ph].
- [8] **ATLAS** Collaboration, M. Aaboud *et al.*, “Measurement of the Inelastic Proton-proton Cross Section at $\sqrt{s} = 13$ TeV with the ATLAS Detector at the LHC,” *Phys. Rev. Lett.* **117** (2016) 182002.
- [9] **CMS** Collaboration, H. Van Haevermaet, “Measurement of the inelastic proton-proton cross Section at $\sqrt{s} = 13$ TeV,” *PoS DIS2016* (2016) 198.
- [10] T. Pierog, I. Karpenko, J.M. Katzy, E. Yatsenko, K. Werner, EPOS LHC: test of collective hadronization with data measured at the CERN large hadron collider. *Phys. Rev. C* **92**, 034906 (2015).
- [11] C. Baus, T. Pierog, R. Ulrich, Cosmic Ray Monte Carlo (CRMC), <https://web.iikp.kit.edu/rulrich/crmc.html>.
- [12] B. Batell, M. Pospelov, and A. Ritz, “Exploring Portals to a Hidden Sector Through Fixed Targets,” *Phys. Rev. D* **80**, 095024 (2009).
- [13] E. J. Fern, V. Di Murro, K. Soga, Z. Li, L. Scibile, and J. A. Osborne, “Geotechnical characterisation of a weak sedimentary rock mass at cern, geneva,” *Tunnelling and Underground Space Technology* **77** (2018) 249–260.
- [14] H. Abreu *et al.* [FASER], “First neutrino interaction candidates at the LHC,” *Phys. Rev. D* **104**, no.9, L091101 (2021).