The Forest Filtration of a Graph

Andrés Carnero Bravo

September 6, 2023

Abstract

Given a graph G , we define a filtration of simplicial complexes associated to G , $\mathcal{F}_0(G) \subseteq \mathcal{F}_1(G) \subseteq \cdots \subseteq \mathcal{F}_{\infty}(G)$ where the first complex is the independence complex and the last the complex is formed by the acyclic sets of vertices. We prove some properties of this filtration and we calculate the homotopy type for various families of graphs.

Contents

[Bibliography](#page-33-0) 34

1 Introduction

Given a simple graph G , the sets of edges that induce acyclic graphs form a simplicial complex, the independence complex of the graphic matroid associated to the graph. This complex is pure of dimension $n - k(G) - 1$ and has the homotopy type of the wedge of $T_G(0, 1)$ spheres, where $T_G(x, y)$ is the Tutte polynomial of the graph (see [\[2](#page-33-1)]). If instead of edges we take vertices, we get a complex $\mathcal{F}_{\infty}(G)$ which does not have to be pure and can, for example, have the homotopy type of a wedge of spheres of different dimension, as it does for the complex associated to the k-complete multipartite graph for $k \geq 3$. This complex, defined in [\[13](#page-34-0)] but not studied, is part of a filtration of complexes associated to the graph:

$$
\mathcal{F}_0(G) \subseteq \mathcal{F}_1(G) \subseteq \mathcal{F}_2(G) \subseteq \cdots \subseteq \mathcal{F}_{\infty}(G),
$$

where for each d we take as simplices the set of vertices which induce acyclic graphs of maximum degree at most d. The case $d = 0$ is the independence complex and the case $d = 1$ is also called the 2-independence complex [\[12\]](#page-34-1). To the best of our knowledge, the cases $2 \le d < \infty$ have not been studied before.

2 Preliminaries

We only consider simple graphs, so no loops or multiedges are allowed. Given a graph G , $V(G)$ will be the set of vertices and $E(G)$ the set of edges. For $S \subseteq V(G)$, $G[S]$ is the graph induced by the set S. For a vertex v, $N_G(v) = \{u \in V(G) : uv \in E(G)\}\$ is its open neighborhood, we omit the subindex G if there is no risk of confusion. The degree of a vertex will be denoted $d_G(v) = |N_G(v)|$. The maximum and minimum degrees will be denoted $\Delta(G)$ and $\delta(G)$ respectively. We say G is a forest if it does not have cycles. Given a graph G, its girth $g(G)$ is the length of the smallest cycle in G, if there is no cycle in G we take $g(G) = \infty$.

Given two graphs G and H, there are two graphs over the vertex set $V(G) \times V(H)$:

1. The cartesian product $G \square H$, where $\{(u_1, v_1)(u_2, v_2)\}$ is an edge if $u_1 = u_2$ and $v_1v_2 \in E(H)$ or $u_1u_2 \in E(G)$ and $v_1 = v_2$.

2. The categorical product $G \times H$, where $\{(u_1, v_1)(u_2, v_2)\}\)$ is an edge if $u_1u_2 \in E(G)$ and $v_1v_2 \in E(H)$.

For all the graph definitions not stated here we follow [\[4](#page-33-2)].

A simplicial complex K on the vertex set V is a family of subsets, called simplices, such that if σ is in K, any subset τ is also in K. We say that a simplex δ has dimension $|\delta| - 1$. The q-skeleton of a complex K, denoted sk_qK , is the subcomplex of all simplices with at most $q + 1$ elements.

Given a simplex $\sigma \in K$, its link is the simplicial complex $lk(\sigma) = {\tau \subseteq V : \tau \cap \sigma =$ \emptyset and $\tau \cup \sigma \in K$ and its star is $st(\sigma) = {\tau \in K : \tau \cup \sigma \in K}$. For a vertex we will write $lk(v)$ and $st(v)$ instead of $lk({v})$ or $st({v}).$

We will not distinguish between a complex and its geometric realization. All the homology and cohomology groups will be with integer coefficients.

Theorem 1 (Whitehead's theorem, see [\[7,](#page-34-2) Corollary 4.33]). If X and Y are simply connected CW-complexes and there is a continuous map $f: X \longrightarrow Y$ such that $f_* : H_n(X) \longrightarrow$ $H_n(Y)$ is an isomorphism for each n, then f is an homotopy equivalence.

Given a complex X on n vertices, its Alexander Dual is the complex

$$
X^* = \{ \sigma \subseteq V(X) \colon V(X) - \sigma \notin X \}.
$$

Theorem 2. (see [\[3](#page-33-3)]) Let X be a simplicial complex with n vertex, then

$$
\tilde{H}_i(X) \cong \tilde{H}^{n-i-3}(X^*)
$$

A well know fact we will use is the following result.

Theorem 3. If X is an CW-complex simply connected such that $\tilde{H}_q(X) \cong \mathbb{Z}^a$ for some $q \geqslant 2$ and the rest of the homology groups are trivial, then $X \simeq \bigvee$ a \mathbb{S}^q .

Theorem 4. (see [\[7](#page-34-2)] example $4C.2$) If X is an CW-complex simply connected such that $\tilde{H}_q(X) \cong \mathbb{Z}^a$, $\tilde{H}_{q+1}(X) \cong \mathbb{Z}^b$ for some $q \geqslant 2$ and the rest of the homology groups are trivial, then $X \simeq \bigvee$ a $\mathbb{S}^q\vee \bigvee$ b \mathbb{S}^{q+1} .

Now we give the basic results of homotopy colimits we will need. Taking $\underline{n} = \{1, \ldots, n\}$ and $P_1(\underline{n}) = P(\underline{n}) - {\underline{n}}$, a punctured *n*-cube X consists of:

- a topological space $\mathcal{X}(S)$ for each S in $\mathcal{P}_1(\underline{n})$, and
- a continuous function $f_{S\subseteq T} : \mathcal{X}(S) \longrightarrow \mathcal{X}(T)$ for each $S \subseteq T$,

such that $f_{S\subseteq S} = 1_{\mathcal{X}(S)}$ and for any $R \subseteq S \subseteq T$ the following diagram comutes

A punctured *n*-cube of interest for a given topological space X is the constant punctured cube \mathcal{C}_X , where $\mathcal{C}_X(S) = X$ for any set S and all the functions are 1_X . The colimit of a punctured n-cube is the space

$$
\operatorname{colim}(\mathcal{X}) = \bigsqcup_{S \in \mathcal{P}_1(\underline{n})} \mathcal{X}(S) / \sim
$$

where \sim is the equivalence relation generated by $f_{S \subseteq T_1}(x_S) \sim f_{S \subseteq T_2}(x_S)$ for T_1, T_2 and $S \subseteq T_1, T_2$. From the definition is clear that $\text{colim}(\mathcal{C}_X) \cong X$ for any X.

For any $n \geq 1$ and S in $\mathcal{P}_1(\underline{n})$ we take:

$$
\Delta(S) = \left\{ (t_1, t_2, \dots, t_n) \in \mathbb{R}^n : \sum_{i=1}^n t_i = 1 \text{ and } t_i = 0 \text{ for all } i \in S \right\}
$$

and $d_{S\subseteq T} : \Delta(T) \longrightarrow \Delta(S)$ the corresponding inclusion. Now, for a punctured n-cube X, its homotopy colimit is

$$
\text{hocolim}({\mathcal X})=\bigsqcup_{S\in{\mathcal P}_1(\underline{n})} {\mathcal X}(S)\times \Delta(S)/\sim
$$

where $(x_S, d_{S\subseteq T}(t)) \sim (f_{S\subseteq T}(x_S), t)$. When $n = 2$, we will specify the punctured 2-cube as the diagram

$$
\mathcal{D}: \qquad X \xleftarrow{f} Z \xrightarrow{g} Y
$$

and its homotopy colimit is called the homotopy pushout.

There is a recursive way to compute homotopy colimits of punctured *n*-cubes. Given a punctured *n*-cube X for $n \geq 2$ and defining the punctured $(n - 1)$ -cubes $\mathcal{X}_1(S) = \mathcal{X}(S)$

and $\mathcal{X}_2(S) = \mathcal{X}(S \cup \{n\})$, we have that (Lemma 5.7.6 [\[9\]](#page-34-3))

 $hocolim(\mathcal{X}) \cong hocolim(\mathcal{X}(\underline{n-1}) \longleftarrow hocolim(\mathcal{X}_1) \longrightarrow hocolim(\mathcal{X}_2)).$

If for all $S \subsetneq [n]$ the map

$$
\operatornamewithlimits{colim}_{T \subsetneq S} X_T \longrightarrow X_S
$$

is a cofibration, we say the punctured cube is cofibrant. If we have X_1, \ldots, X_n CWcomplexes such that their intersections are subcomplexes, and we take the punctured cube given by the intersections and the inclusions between them, then the punctured cube is cofibrant and hocolim $(X) \simeq \text{colim}(X)$ (Proposition 5.8.25 [\[9](#page-34-3)]). We will use several times the following known fact about homotopy push-out (see [\[1](#page-33-4)] for a proof).

Lemma 5. Let X, Y, Z be spaces with maps $f: Z \longrightarrow X$ and $g: Z \longrightarrow Y$ such that both maps are null-homotopic. Then

$$
hocolim(\mathcal{S})\simeq X\vee Y\vee\Sigma Z
$$

where

$$
S: \qquad Y \xleftarrow{g} Z \xrightarrow{f} X
$$

Lemma 6. Let X_1, \ldots, X_k simplicial complexes such that the intersection of two or more is contractible or empty, X_i is connected for all i and there is a graph G of size k and a bijection $\gamma: \{1,\ldots,k\} \longrightarrow E(G)$ such that $\bigcap_{i\in S} \gamma(i) \neq \emptyset$ if an only if $\bigcap_{i\in S} X_i \neq \emptyset$ for all non-empty S subset of $\{1, \ldots, k\}$. Then $X = \begin{bmatrix} \end{bmatrix}$ k $i=1$ X_i has the homotopy type of the nerve with the complexes X_i attached to the corresponding point in the nerve.

Proof. By induction on k. For $k = 1, 2$ the result is clear. Assume it is true for any $r \leq k$ and take X_1, \ldots, X_{k+1} simplicial complexes such that the intersection of two or more is contractible or empty, X_i is connected for all i and there is a graph G of size $k + 1$ and a bijection $\gamma: \{1, \ldots, k+1\} \longrightarrow E(G)$ such that $\bigcap_{i \in S} \gamma(i) \neq \emptyset$ if an only if $\bigcap_{i \in S} X_i \neq \emptyset$ for all non-empty S subset of $\{1, \ldots, k + 1\}$. Now, take N the nerve complex of X_1, \ldots, X_{k+1} . For any $i \in \{1, ..., k + 1\}$, $lk(i)$ is:

- (a) Empty if in the corresponding edge both vertices have degree 1.
- (b) Contractible if in the corresponding edge one vertex has degree 1 and the other degree at least 2.

(c) Homotopy equivalent to \mathbb{S}^0 if in the corresponding edge both vertices have degree at least 2.

By the inductive formula for homotopy colimits of punctured cubes, the homotopy colimit of the intersection diagram associated to X_1, \ldots, X_{k+1} is homotopy equivalent to the homotopy pushout of the diagram

 $S: \text{hocolim}(\mathcal{S}_2) \longrightarrow \text{hocolim}(\mathcal{S}_1) \longrightarrow X_{k+1}$,

where S_1 is the homotopy colimit of the intersection diagram associated to $X_1 \cap X_{k+1}, \ldots, X_k \cap$ X_{k+1} , and S_2 is the homotopy colimit of the intersection diagram associated to X_1, \ldots, X_k . Now hocolim $(S_1) \simeq lk(k + 1)$, so we have three possibilities:

- (a) hocolim $(\mathcal{S}_1) = \emptyset$, then hocolim $(\mathcal{S}) \simeq$ hocolim $(\mathcal{S}_2) \sqcup X_{k+1}$
- (b) hocolim $(S_1) \simeq *$, then hocolim $(S) \simeq$ hocolim $(S_2) \vee X_{k+1}$
- (c) hocolim $(S_1) \simeq \mathbb{S}^0$, then hocolim $(S) \simeq$ hocolim $(S_2) \vee \mathbb{S}^1 \vee X_{k+1}$

3 Definition and basic properties

Let G be a graph, we define its *d-forest complex* as the complex

$$
\mathcal{F}_d(G) = \{ \sigma \subseteq V(G) \colon G[\sigma] \text{ is a forest with } \Delta(G[\sigma]) \leq d \};
$$

for $d = \infty$ we take

$$
\mathcal{F}_{\infty}(G) = \{ \sigma \subseteq V(G) \colon G[\sigma] \text{ is a forest} \}.
$$

For $d = 0$, $\mathcal{F}_0(G)$ is the independence complex of G and for $d = 1$ is also called the 2-independence complex of G —the r-independence complex of G has as simplices sets $A \subseteq V(G)$ such that every connected component of $G[A]$ has at most r vertices. Note that if $d + 1 = \min\{r : G \text{ is } K_{1,r}\text{-free}\},\$ then $\mathcal{F}_l(G) = \mathcal{F}_d(G)$ for all $l \geq d$.

Given a graph G let $t_d(G) = \max\{|V(T)|: T \text{ is an induced forest such that } \Delta(T) \leq d\},\$ by definition $t_d(G) = \dim(\mathcal{F}_d(G)) + 1$, therefore knowing the homotopy type of $\mathcal{F}_d(G)$ or its homology groups gives us a lower bound for $t_d(G)$.

 \Box

Theorem 7. For any graph G and all d, the pair $(\mathcal{F}_{d+1}(G), \mathcal{F}_d(G))$ is d-connected.

Proof. For any d, we have that $sk_i\mathcal{F}_d(G) = sk_i\mathcal{F}_{d+1}(G)$ for all $i \le d$ because a forest of order $i + 1$ has maximum degree at most i. Then all the cells in $\mathcal{F}_{d+1}(G) - \mathcal{F}_d(G)$ have dimension greater than d and this implies the result (see [\[7\]](#page-34-2) Corollary 4.12). \Box

By definition the following results are clear.

Proposition 8. For $d \geq 1$,

$$
\mathcal{F}_d(K_n) \simeq \bigvee_{\frac{(n-1)(n-2)}{2}} \mathbb{S}^1.
$$

Proposition 9. For $n \geq 3$ and $d \geq 2$,

$$
\mathcal{F}_d(C_n) \cong \mathbb{S}^{n-2}.
$$

A subset of vertices σ is an independent set if all of its subsets of cardinality 2 are independent. This says that in order to be a simplex of the independence complex, a set of vertices only need have its 1-skeleton contained in the complex. This type of complexes are called *flag complexes*. Now, for $\mathcal{F}_1(G)$, its 1-skeleton is the complete graph of the same order as G , therefore it is not a flag complex in general, because it is not contractible for all graphs. The following result tells us that it has an analogous property but for the 2-skeleton, rather than the 1-skeleton.

Proposition 10. Let σ be a subset of $V(G)$ such that all of its subsets of cardinality 3 are simplices of $\mathcal{F}_1(G)$, then σ is a simplex of $\mathcal{F}_1(G)$.

Proof. If $|\sigma| \leq 3$ the result is clear. Now let $\sigma = \{v_0, v_1, v_2, v_3\}$. Then, for $\tau = \{v_0, v_1, v_2\}$, we have that $G_{\tau} = G[\tau]$ is forest such that $\Delta(G_{\tau}) \leq 1$. Now, v₃ at most can have one neighbor in τ and it must be a vertex of degree 0 in G_{τ} . Therefore $G_{\sigma} = G[\sigma]$ is a graph such that $\Delta(G_{\sigma}) \leq 1$, which implies it is a forest and σ is a simplex of $\mathcal{F}_1(G)$.

Assume the result is true for any subset of at most $k \geq 4$ vertices that has its 2-skeleton in $\mathcal{F}_1(G)$. Let $\sigma = \{v_0, \ldots, v_k\}$ a subset of $k+1$ vertices such that its 2-skeleton is in $\mathcal{F}_1(G)$. By induction hypothesis, $\tau = \{v_0, \ldots, v_{k-1}\}\$ is a simplex of $\mathcal{F}_1(G)$, therefore, taking G_{τ} as before,

$$
G_{\tau} \cong rK_1 \sqcup M_s
$$

with $r, s \geq 0$ and $r + 2s = k + 1$. By hypothesis, v_k can not be adjacent to a vertex in M_s and only can be adjacent to one vertex in rK_1 . So σ induces a graph with maximum degree at most 1 and therefore σ is a simplex of $\mathcal{F}_1(G)$. \Box

This can not be generalized for $\mathcal{F}_d(G)$ with $d \geq 2$ as $\mathcal{F}_d(C_{d+3})$ shows.

If a simplicial complex K is such that $\tilde{H}_q(K) \not\cong 0$, then $f_i(K) \geq f_i(\Delta^{q+1})$ and $f_0(K) =$ $q + 2$ if and only if $K \cong \partial (\Delta^{q+1})$.

Proposition 11. Let G be a graph of order $q + 2$, with $q \ge 1$, then:

- 1. If $\tilde{H}_q(\mathcal{F}_q(G)) \not\cong 0$, then $G \cong K_{1,q+1}$ or $G \cong C_{q+2}$
- 2. If $\tilde{H}_q(\mathcal{F}_\infty(G)) \not\cong 0$, then $G \cong C_{q+2}$

Proof. For $d = q$ or $d = \infty$, we have that $\mathcal{F}_d(G) \cong \partial (\Delta^{q+1})$ and for any proper subset of the vertices S, $\mathcal{F}_d(G[S])$ must be contractible. If $\Delta(G) = q + 1$, then G can not have cycles because $V(G) - \{v\}$ is a simplex for any vertex and $\mathcal{F}_{\infty}(G) \simeq *$. Take v a vertex such that $d_G(v) = q + 1$, then $\mathcal{F}_q(G) = st(v) \cup \mathcal{F}_q(G - v)$ and, because $\tilde{H}_q(\mathcal{F}_q(G - v)) \cong 0$, using the Mayer-Vietoris sequence we have that $\tilde{H}_{q-1}(lk(v)) \not\cong 0$. Therefore $lk(v) \cong \partial(\Delta^q)$. If $q = 1$, then $lk(v)$ is two disjoint vertices from where it follows that $G \cong K_{1,2}$ or $G \cong C_3$. Assume $q \ge 2$, then $N(v)$ must be an independent set and $G \cong K_{1,q+1}$.

Assume $\Delta(G) \leq q$, then G must have a cycle, otherwise $\mathcal{F}_d(G) \simeq *$ for $d = q$ or $d = \infty$. Let $C \leq G$ be an induced cycle. If $V(C) \subsetneq V(G)$, because any proper subset is a simplex, $V(C)$ will be a simplex, but this can not happen. Therefore $G \cong C_{q+2}$. \Box

Proposition 12. If $e \in E(G)$ is bridge, then $\mathcal{F}_{\infty}(G) = \mathcal{F}_{\infty}(G - e)$.

Lemma 13. If $G = G_1 \sqcup G_2$, then for all d,

$$
\mathcal{F}_d(G) = \mathcal{F}_d(G_1) * \mathcal{F}_d(G_2).
$$

Proposition 14. If $G = G_1 \sqcup \cdots \sqcup G_k$, then for $d \ge 0$,

$$
conn(\mathcal{F}_d(G)) \ge 2k - 2 + \sum_{i=1}^k conn(\mathcal{F}_d(G_i)).
$$

 \Box

Proof. This follows from $\mathcal{F}_d(G) = \mathcal{F}_d(G_1) * \cdots * \mathcal{F}_d(G_k)$

Lemma 15. If v is a vertex such that no cycle of G contains it, then $\mathcal{F}_{\infty}(G) \simeq *$.

Proof. Beacause v does not belongs to a cycle, then $\mathcal{F}_{\infty}(G) = \{v\} * \mathcal{F}_{\infty}(G - v)$.

Corollary 16. If $\delta(G) \leq 1$, then $\mathcal{F}_{\infty}(G) \simeq *$.

Corollary 17. If G has a vertex v such that $N_G(v) = \{v_1, v_2\}$, then $\mathcal{F}_{\infty}(G) \simeq \Sigma l k_{\mathcal{F}_{\infty}(G)}(v_i)$ *for* $i = 1, 2$.

Proof. Because $N_G(v) = \{v_1, v_2\}$, then $d_{G-v_i}(v) = 1$ and therefore $\mathcal{F}\infty(G - v_i) \simeq *$. Now $\mathcal{F}_{\infty}(G) \simeq \text{hocolim}(\mathcal{S})$ with $\mathcal{S} \colon \mathcal{F}_{\infty}(G - v_i) \longleftrightarrow lk_{\mathcal{F}(G,\infty)}(v_i) \longrightarrow st_{\mathcal{F}(G,\infty)}(v_i)$, by Lemma [5](#page-4-0) we obtain the result. 口

Proposition 18. Let G be a graph such that $\tilde{H}_q(\mathcal{F}_d(G)) \not\cong 0$ for some d and q, then G has at least $q + 2$ different induced forests of $q + 1$ vertices and maximun degree at most d.

Lemma 19. Let G be a graph that is the union of three graphs G_1, G_2, G_0 such that:

- $G_0 \cong K_3$
- $V(G_0) = \{v, v_1, v_2\}$
- $V(G_1) \cap V(G_0) = \{v_1\}, V(G_2) \cap V(G_0) = \{v_2\}$ and $V(G_1) \cap V(G_2) = \emptyset$

Then, $lk_{\mathcal{F}_{\infty}(G)}(v) \simeq \text{hocolim}(\mathcal{S})$ with S the diagram:

$$
\mathcal{F}_{\infty}(G_1) * \mathcal{F}_{\infty}(G_2 - v_2) \longleftarrow \mathcal{F}_{\infty}(G_1 - v_1) * \mathcal{F}_{\infty}(G_2 - v_2) \longleftarrow \mathcal{F}_{\infty}(G_1 - v_1) * \mathcal{F}_{\infty}(G_2)
$$

Proof. Because

$$
lk_{\mathcal{F}_{\infty}(G)}(v) = (\mathcal{F}_{\infty}(G_1) * \mathcal{F}_{\infty}(G_2 - v_2)) \cup (\mathcal{F}_{\infty}(G_1 - v_1) * \mathcal{F}_{\infty}(G_2))
$$

and

$$
(\mathcal{F}_{\infty}(G_1) * \mathcal{F}_{\infty}(G_2 - v_2)) \cap (\mathcal{F}_{\infty}(G_1 - v_1) * \mathcal{F}_{\infty}(G_2)) = \mathcal{F}_{\infty}(G_1 - v_1) * \mathcal{F}_{\infty}(G_2 - v_2)
$$

we have that

$$
lk_{\mathcal{F}_{\infty}(G)}(v) = \mathrm{colim}(\mathcal{S}) \simeq \mathrm{hocolim}(\mathcal{S})
$$

 \Box

 \Box

Proposition 20. Let G be a graph with n vertices and $g(G) < \infty$, then $\tilde{H}_i(\mathcal{F}_\infty(G)) \cong 0$ for all $i < g(G) - 2$.

Proof. The Alexander Dual has dimension $n - g(G) - 1$, thus $\tilde{H}^k(\mathcal{F}^*_{\infty}(G)) \cong 0$ for all $k > n - g(G) - 1$. By Theorem [2,](#page-2-0) $\tilde{H}_i(\mathcal{F}_\infty(G)) \cong 0$ for all $i < g(G) - 2$. □

In the last proposition we saw that the girth gives us a lower bound for the homological connectivity of $\mathcal{F}_{\infty}(G)$, now we will see that this bound also works for the connectivity, first we show that $g(G) \geq 4$ implies that $\mathcal{F}_{\infty}(G)$ is simply connected, by showing this for $\mathcal{F}_2(G)$.

Proposition 21. Let G be a graph with $g(G) \geq 4$, then $\pi_1(\mathcal{F}_2(G)) \cong 0$.

Proof. We will prove it for connected graphs. We take T a spanning tree of G and take the finitely presented group H_T with set of generators $E(G) \cup E(G^c)$ and with the following relations:

- $uv = 1$ for all the edges of T,
- $(uv)(vw) = uw$ if $\{u, v, w\}$ is a simplex of $\mathcal{F}_2(G)$.

we have that $H_T \cong \pi_1(\mathcal{F}_2(G))$ (see [\[11\]](#page-34-4) Theorem 7.34).

Note that any triple of vertices $\{u, v, w\}$ spans a forest in G because $g(G) \geq 4$, so the 2-skeleton of $\mathcal{F}_2(G)$ contains all possible triangles.

We will show that all generators uv are trivial by induction on the distance $k = d_T(u, v)$. If $k = 1$, this is clear by the first type of relation. Assume uv is trivial if $d_T(u, v) \leq k$. Take uv such that $d_T(u, v) = k + 1$ and take $uw_1w_2 \cdots w_kv$ the uv-path in T. Since $\{u, w_1, v\}$ is a simplex of $\mathcal{F}_2(G)$, the second relation implies $uv = (uw_1)(w_1v) = w_1v$ where we have $d_T(w_1, v) = k.$ \Box

Because $\mathcal{F}_{\infty}(G)$ is always connected, using the last proposition, Propositio[n20](#page-8-0) and the Hurewicz Theorem we have the following result:

Theorem 22. For any graph G, $conn(\mathcal{F}_{\infty}(G)) \geq g(G) - 3$.

4 Homotopy type calculations

In this secction for various families of graphs we will calculate the homotopy type for the forest complexes, for every step in the filtration in most cases; and in some cases, like the cactus graphs, only for the last complex of the filtration.

4.1 Some Graph families

4.1.1 Paths and cycles

The homotopy type of all r-independence complexes of paths was calculated by Salvetti [\[10\]](#page-34-5) using discrete Morse theory. Here we give a different proof for \mathcal{F}_1 using homotopy pushouts, which also shows that the inclusion $\mathcal{F}_1(P_{4r+3}) \hookrightarrow \mathcal{F}_1(P_{4(r+1)})$ is a homotopy equivalence. This will allow us to calculate the homotopy type of $\mathcal{F}_1(C_n)$ avoiding discrete Morse theory, which was the tool used in [\[5\]](#page-34-6).

Proposition 23. [\[10](#page-34-5)]

$$
\mathcal{F}_1(P_n) \simeq \begin{cases} \mathbb{S}^{2r-1} & \text{if } n = 4r \\ * & \text{if } n = 4r+1 \text{ or } n = 4r+2 \\ \mathbb{S}^{2r+1} & \text{if } n = 4r+3 \end{cases}
$$

Proof. For $r = 0$, it is clear that $\mathcal{F}_1(P_1) \simeq * \simeq \mathcal{F}_1(P_2)$. For P_3 , $\mathcal{F}_1(P_3) \simeq K_3$. For $\mathcal{F}_1(P_4)$

$$
lk(v_4) = \mathcal{F}_1(P_2) \cup \{v_3\} * \mathcal{F}_1(P_1) \simeq *
$$

therefore the inclusion $i: \mathcal{F}_1(P_3) \longrightarrow \mathcal{F}_1(P_4)$ is a homotopy equivalence.

Next, we will prove that $\mathcal{F}_1(P_{4r+1}) \simeq \mathcal{F}_1(P_{4r+2}) \simeq *$ for all $r \geq 1$.

Assume that it is true for any $1 \leq r \leq k$. For $\mathcal{F}_1(P_{4(k+1)})$, by induction hypothesis,

$$
lk(v_{4(k+1)}) = \mathcal{F}_1(P_{4k+2}) \cup \{v_{4k+3}\} * \mathcal{F}_1(P_{4k+1}) \simeq *
$$

therefore the inclusion $\mathcal{F}_1(P_{4k+3}) \longrightarrow \mathcal{F}_1(P_{4(k+1)})$ is a homotopy equivalence.

Now, for $\mathcal{F}_1(P_{4(k+1)+1})$ we have

$$
lk(v_{4(k+1)+1}) = \mathcal{F}_1(P_{4k+3}) \cup \{v_{4(k+1)}\} * \mathcal{F}_1(P_{4k+2}).
$$

Setting $X = \mathcal{F}_1(P_{4k+2})$ and $Y = \{v_{4k+3}\}\ast \mathcal{F}_1(P_{4k+2})$, we have by induction hypothesis that

$$
X \cap Y = \mathcal{F}_1(P_{4k+2}) \simeq *
$$

therefore $\mathcal{F}_1(P_{4k+3}) \longrightarrow lk(v_{4(k+1)+1})$ is a homotopy equivalence.

$$
\mathcal{F}_1(P_{4k+3}) \xrightarrow{\simeq} lk(v_{4(k+1)+1}) \xrightarrow{\simeq} \mathcal{F}_1(P_{4(k+1)})
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
st(v_{4(k+1)+1}) \xrightarrow{\simeq} st(v_{4(k+1)+1}) \xrightarrow{\simeq} \mathcal{F}_1(P_{4(k+1)+1})
$$

 $\mathcal{F}_1(P_{4(k+1)+1}) \simeq st(v_{4(k+1)+1}) \simeq *$

For $\mathcal{F}_1(P_{4(k+1)+2})$:

$$
lk(v_{4(k+1)+2}) = \mathcal{F}_1(P_{4(k+1)}) \cup \{v_{4(k+1)+1}\} * \mathcal{F}_1(P_{4k+3});
$$

because $\mathcal{F}_1(P_{4k+3}) \longrightarrow \mathcal{F}_1(P_{4(k+1)})$ is an homotopy equivalence, we have that $lk(v_{4(k+1)+2}) \simeq$ \ast and therefore

$$
\mathcal{F}_1(P_{4(k+1)+2}) \simeq \mathcal{F}_1(P_{4(k+1)+1}) \simeq *.
$$

We have that $\mathcal{F}_1(P_{4(k+1)}) \simeq \mathcal{F}_1(P_{4k+3})$; now for this last complex:

$$
lk(v_{4k+3}) = \mathcal{F}_1(P_{4k+1}) \cup \{v_{4k+2}\} * \mathcal{F}_1(P_{4k}),
$$

where $\mathcal{F}_1(P_{4k+1}) \simeq *$, therefore

$$
lk(v_{4k+3}) \simeq \Sigma \mathcal{F}_1(P_{4k}).
$$

Since $\mathcal{F}_1(P_{4k+2}) \simeq *$, we have that $\mathcal{F}_1(P_{4k+3}) \simeq \Sigma^2 \mathcal{F}_1(P_{4k})$ and

$$
\mathcal{F}_1(P_{4(k+1)}) \simeq \Sigma^2 \mathcal{F}_1(P_{4k}) \simeq \Sigma^2 \mathbb{S}^{2k-1} \simeq \mathbb{S}^{2k+1}.
$$

Doing the exact same argument we can see that $\mathcal{F}_1(P_{4(k+1)+3}) \simeq \Sigma^2 \mathcal{F}_1(P_{4(k+1)})$ and therefore

$$
\mathcal{F}_1(P_{4(k+1)+3}) \simeq \Sigma^2 \mathbb{S}^{2k+1} \simeq \mathbb{S}^{2k+3}.
$$

 \Box

In the proof of the last proposition we saw that the inclusion $\mathcal{F}_1(P_{4k+3}) \longrightarrow \mathcal{F}_1(P_{4(k+1)})$ obtained by erasing the last (or the first) vertex is an homotopy equivalence. We will use this fact in the following corollary.

Corollary 24. $[5]$ $[5]$

$$
\mathcal{F}_1(C_n) \simeq \begin{cases}\n\bigvee_{3} \mathbb{S}^{2r-1} & \text{if } n = 4r \\
\mathbb{S}^{2r-1} & \text{if } n = 4r+1 \\
\mathbb{S}^{2r} & \text{if } n = 4r+2 \\
\mathbb{S}^{2r+1} & \text{if } n = 4r+3\n\end{cases}
$$

Proof. For $n = 3, 4$, the only possible simplices are a vertex or pair of vertices, any set with more vertices will have a 3-path or a cycle. Therefore $\mathcal{F}_1(C_3) \cong K_3$ and $\mathcal{F}_1(C_4) \cong K_4$. For $n = 5$, taking v_1, v_2, v_3, v_4, v_5 the vertices of the cycle with edges $v_i v_{i+1}$, the facets of $\mathcal{F}_1(C_5)$ are $\sigma_i = \{v_i, v_{i+2}, v_{i+3}\}\.$ The edge $v_{i+2}v_{v_i+3}$ only is contained in σ_i , so we can collapse it for all *i*. Therfore $\mathcal{F}_1(C_5) \simeq \mathcal{F}_0(C_5) \cong \mathbb{S}^1$.

Assume $n \geq 6$ and let v_1, \ldots, v_n be the vertices of the cycle. Then $lk(v_n) = K_1 \cup K_2 \cup K_3$ where

$$
K_1 = \mathcal{F}_1(C_n - v_n - v_2 - v_{n-1}) \cong C(\mathcal{F}_1(P_{n-4}))
$$

\n
$$
K_2 = \mathcal{F}_1(C_n - v_n - v_1 - v_{n-2}) \cong C(\mathcal{F}_1(P_{n-4}))
$$

\n
$$
K_3 = \mathcal{F}_1(C_n - v_n - v_1 - v_{n-1}) \cong \mathcal{F}_1(P_{n-3})
$$

Now

$$
K_1 \cap K_2 \cap K_3 = K_1 \cap K_2 = \mathcal{F}_1(C_n - v_n - v_1 - v_2 - v_{n-1} - v_{n-2}) \cong \mathcal{F}_1(P_{n-5})
$$

$$
K_1 \cap K_3 = \mathcal{F}_1(C_n - v_n - v_1 - v_2 - v_{n-1}) \cong \mathcal{F}_1(P_{n-4})
$$

$$
K_2 \cap K_3 = \mathcal{F}_1(C_n - v_n - v_1 - v_{n-1} - v_{n-2}) \cong \mathcal{F}_1(P_{n-4})
$$

$$
K_1 \cup K_2 \cong \Sigma \mathcal{F}_1(P_{n-5})
$$

If $n = 4r, K_1 \cap K_2 \cong \mathcal{F}_1(P_{4(r-2)+3}), K_3 \cong *$ and $K_1 \cap K_3 \cong \mathcal{F}_1(P_{4(r-1)}) \cong K_2 \cap K_3$. By the observation before the corollary, the inclusion $K_1 \cap K_2 \cap K_3 \hookrightarrow K_1 \cap K_3$ is a homotopy equivalence. Therefore $(K_1 \cup K_2) \cap K_3 \simeq K_2 \cap K_3$ and

$$
lk(v_n) \simeq \bigvee_2 \mathbb{S}^{2r-2},
$$

Since

$$
\mathcal{F}_1(C_{4r} - v_n) \simeq \mathbb{S}^{2r-1},
$$

we obtain the result.

If $n = 4r + 1$, $K_1 \cap K_3 \simeq K_2 \cap K_3 \cong \mathcal{F}_1(P_{4(r-1)+1}) \simeq *$ and $K_2 \cup K_3 \simeq K_3$. Because $K_1 \cap K_2 \cap K_3 = K_1 \cap K_2$, we have that

$$
(K_2 \cup K_3) \cap K_1 \simeq K_1 \cap K_3 \simeq *
$$

and

$$
K_1 \cup K_2 \cup K_3 \simeq K_2 \cup K_3 \simeq K_3 \cong \mathcal{F}_1(P_{4(r-1)+2}) \simeq *.
$$

Therefore $\mathcal{F}_1(C_{4r+1}) \simeq \mathcal{F}_1(P_{4r}) \simeq \mathbb{S}^{2r-1}$.

For $n = 4r + 2$ and $n = 4r + 3$, $\mathcal{F}_1(C_n - v_n) \simeq *$, therefore $\mathcal{F}_1(C_n) \simeq \Sigma lk(v_n)$. If $n = 4r + 2$, $K_1 \cap K_2 \cong \mathcal{F}_1(P_{4(r-1)+1}) \cong *$ and $K_1 \cap K_3, K_2 \cap K_3 \cong \mathcal{F}_1(P_{4(r-1)+2}) \cong *$. Then $K_1 \cup K_2 \simeq *$ and $(K_1 \cup K_2) \cap K_3 \simeq *$. From this we have that $lk(v_n) \simeq K_3$, therefore

$$
\mathcal{F}_1(C_{4r+2}) \simeq \Sigma \mathcal{F}_1(P_{4(r-1)+3}) \simeq \mathbb{S}^{2r}.
$$

If $n = 4r + 3$, $K_2 \cap K_3 \cong \mathcal{F}_1(P_{4(r-1)+3})$ and the inclusion $K_2 \cap K_3 \hookrightarrow K_3$ is a homotopy equivalence, therefore $K_2 \cup K_3 \simeq *$. From this $lk(v_n) \simeq \Sigma(K_1 \cap (K_2 \cup K_3)).$ Since $K_1 \cap K_2 \cap K_3 = K_1 \cap K_2$, we have that $K_1 \cap (K_2 \cup K_3) \simeq K_1 \cap K_3$ and

$$
\mathcal{F}_1(C_{4r+3}) \simeq \Sigma^2 \mathcal{F}_1(P_{4(r-1)+3}) \simeq \mathbb{S}^{2r+1}.
$$

Proposition 25.

$$
\mathcal{F}_{\infty}(C_n + e) \cong \mathbb{S}^{n-3}
$$

Proof. Assume the vertices of $G = C_n + e$ are labeled $v, w_1, \ldots, w_r, u, w_{r+1}, \ldots, w_{r+k}$ with $e = vu$ (Figure [1\)](#page-14-1). Because $\mathcal{F}_{\infty}(G - v) \simeq *,$ we have that $\mathcal{F}_{\infty}(G) \simeq \Sigma lk(v)$. Now, $lk(v)$ is formed by the subsets of $V(G - v)$ such that together with v they do not induce a cycle, therefore the facets are

$$
\sigma_0 = [w_1, \ldots, w_r, w_{r+1}, \ldots, w_{r+k}]
$$

and

$$
\sigma_{ij} = [w_1, \ldots, \hat{w}_i, \ldots, w_r, u, w_{r+1}, \ldots, , \hat{w}_{r+j}, \ldots, w_{r+k}]
$$

Figure 1: $C_n + e$

for $1 \leq i \leq r, 1 \leq j \leq k$. If we call K the complex form by σ_0 and its subsets, and L the complex which facets are the simplices σ_{ij} , we get that $lk(v) = K \cup L$ and both of this complexes are contractible, therefore $lk(v) \simeq \Sigma K \cap L$.

Now, taking X the complex with facets $[w_{r+1}, \ldots, \hat{w}_{r+j}, \ldots, w_{r+k}]$ and Y the complex with facets $[w_1, \ldots, \hat{w}_i, \ldots, w_r]$, we have that $K \cap L \cong X * Y$. Because $X \cong \mathbb{S}^{k-2}$ and $Y \cong \mathbb{S}^{r-2}$, we have that $K \cap L \cong \mathbb{S}^{k-2} * \mathbb{S}^{r-2} \cong \mathbb{S}^{r+k-3}$ and, because $r + k = n - 2$, $\mathcal{F}_{\infty}(G) \simeq \mathbb{S}^{n-3}.$ \Box

4.1.2 Double stars

Let $St_{r,s}$ be the *double star* with $V(St_{r,s}) = \{u_0, u_1, \ldots, u_r, v_0, v_1, \ldots, v_s\}$ and $E(St_{r,s}) =$ ${u_iu_0: i > 0} \cup {v_iv_0: i > 0} \cup {u_0v_0}$

Proposition 26.

$$
\mathcal{F}_1(St_{r,s}) \simeq \mathbb{S}^1
$$

and for $2 \leq d < \infty$

$$
\mathcal{F}_d(St_{r,s}) \simeq \bigvee_{\binom{r-1}{d-1}\binom{s-1}{d-1}} \mathbb{S}^{2d-1}
$$

Proof. For $\mathcal{F}_1(St_{r,s})$, the link of u_0 has as facets $\sigma_i = \{u_i, v_1, \ldots, v_s\}$ for all i and $\{v_0\}$, therefore

$$
lk(u_0) \simeq \mathbb{S}^0.
$$

Since $\mathcal{F}_1(St_{r,s} - u_0) \simeq *,$ we have that $\mathcal{F}_1(St_{r,s}) \simeq \Sigma lk(u_0) \simeq \mathbb{S}^1$.

For $d \geq 2$, if $r \leq d - 1$ or $s \leq d - 1$, then $\mathcal{F}_d(St_{r,s}) \simeq *$, because the set $\{u_1, \ldots, u_r\}$ or the set $\{v_1, \ldots, v_s\}$ would be contained in all facets. Assume $r, s \geq d$. The facets of $\mathcal{F}_d(St_{r,s}),$ besides $X = \{u_1, \ldots, u_r, v_1 \ldots, v_s\},$ are of 3 types:

- 1. $\alpha_S = S \cup \{u_0, v_1, \ldots, v_s\}$, where $S \subseteq \{u_1, \ldots, u_r\}$ and $|S| = d$.
- 2. $\beta_S = S \cup \{v_0, u_1, \dots, u_r\}$, where $S \subseteq \{v_1, \dots, v_s\}$ and $|S| = d$.
- 3. $\sigma_{S_1,S_2} = \{u_0, v_0\} \cup S_1 \cup S_2$, where $S_1 \subseteq \{u_1, \ldots, u_r\}$, $S_2 \subseteq \{v_1, \ldots, v_s\}$ and $|S_1| =$ $|S_2| = d - 1.$

Take $\tau = \mathcal{P}(X) - \{\emptyset\}, \alpha$ the complex generated by $\{\alpha_S\}, \beta$ the complex generated by $\{\beta_S\}$ and σ the complex generated by the $\{\sigma_{S_1,S_2}\}\$, $\mathcal{F}_d(St_{r,s}) = \alpha \cup \beta \cup \sigma \cup \tau$. Now, these four complexes are contractible and so are $\alpha \cap \sigma, \beta \cap \sigma, \alpha \cap \tau, \beta \cap \tau$. Also

$$
\alpha \cap \beta \cap \sigma \cap \tau = \alpha \cap \sigma \cap \tau = \beta \cap \sigma \cap \tau = \alpha \cap \beta \cap \sigma = \sigma \cap \tau \cong sk_{d-2}\Delta^{r-1} * sk_{d-2}\Delta^{s-1}
$$

and $\alpha \cap \beta \cap \tau = \alpha \cap \beta$. We compute the homotopy colimit of the punctured 4-cube given by this union using the recursive formula given in the preliminaries. This what the formula gives applied to the top and bottom of the 4-cube:

We find that the complex has the homotopy type of the following homotopy pushout:

$$
S: * \longleftarrow \Sigma(\sigma \cap \tau) \longrightarrow \tau
$$

hocolim $(S) \simeq \Sigma^2(\sigma \cap \tau) \simeq \bigvee_{\binom{r-1}{d-1}\binom{s-1}{d-1}} \mathbb{S}^{2d-1}.$

 \Box

4.1.3 Cactus graphs

For any graph G, we take the block graph $B(G)$ in which the vertices are the blocks of G and the cut-vertices of G, where vB is an edge if v is a vertex of B. If G is connected, then $B(G)$ is a tree.

A graph G is a cactus graph if all of its blocks are isomorphic to a cycle or to K_2 . We will say that a block is saturated if all of its vertices are cut vertices and $sb(G)$ is the number of saturated blocks. A vertex v is saturated if it is shared by two or more saturated blocks, with $sv(G)$ the number of saturated vertices.

Lemma 27. Let G be a cactus graph such that $sb(G) \geq 1$, then there is a saturated block B such that either it does not have saturated vertices, or:

- (i) it has only one saturated vertex v, and
- (ii) the connected component of $B(G) v$ which contains B does not have any other saturated block.

Proof. If there are no saturated vertices, there is nothing to prove. Assume $sv(G) \geq 1$. If there is a saturated block without a saturated vertex, again there is nothing to prove. Assume all saturated blocks have at least one saturated vertex.

Let V_1 be the set of all saturated blocks of G and V_2 the set of all saturated vertices. In the subgraph $T = B(G)[V_1 \cup V_2]$ all the leaves are blocks, because each saturated vertex is in at least two saturated blocks, therefore $d_T(v) \geq 2$ for all the vertices of V_2 . We take $L \subseteq V_1$ the set of all the leaves of T and let (B_1, B_2) be a pair in $L \times L$ such that

$$
d(B_1, B_2) = \max\{d(X, Y): (X, Y) \in L \times L\}
$$

Take v_1 the only saturated vertex in B_1 and v_2 the only saturated vertex in B_2 . We claim that the only B_1B_2 -path in $B(G)$ contains both v_1 and v_2 . If not, then B_1 and B_2 are in different connected components of T and, assuming v_1 is not in the B_1B_2 -path, any leaf B' in the same component of B_1 is such that $d(B', B_2) > d(B_1, B_2)$. Therefore v_1 and v_2 are in the only B_1B_2 -path.

If in $B(G) - v_1$ there are saturated blocks in the same component than B_1 , the distance between these and B_2 is larger that the distance between B_1 and B_2 , which can not happen. Therefore B_1 and v_1 are as wanted. \Box

Lemma 28. Let G be a cactus graph such that all of its blocks are cycles and such that it does not have saturated blocks, then

$$
\mathcal{F}_{\infty}^*(G) \simeq \mathbb{S}^{b(G)-2}.
$$

Proof. Let B_0, \ldots, B_k be the blocks of G. If $k = 0$, then $\mathcal{F}_{\infty}^*(G) = \emptyset = \mathbb{S}^{-1}$. Assume, $k \geq 1$. We take $X_i = V(G) - V(B_i)$ for all i, this are the facets of $\mathcal{F}^*_{\infty}(G)$ and we have that

$$
\bigcap_{i=0}^{k} X_i = \varnothing
$$

$$
\bigcap_{i \in S} X_i \neq \varnothing, \ \ \forall S \subsetneq [k]
$$

Then, its nerve is isomorphic to $\partial \Delta^k \cong \mathbb{S}^{k-1}$. Therefore, $\mathcal{F}^*_{\infty}(G) \simeq \mathbb{S}^{b(G)-1}$.

Lemma 29. Let G be a cactus graph different from K_3 , then $\mathcal{F}_{\infty}(G)$ is simply connected.

 \Box

Proof. If G has only one block and G is not K_3 , G must be a single vertex, K_2 or a cycle with at least 4 vertices, thus $\mathcal{F}_{\infty}(G)$ is contractible or a sphere of dimension at least 2. Assume G has $k \geq 2$ blocks. For each block that is not isomorphic to K_2 we can erase one edge to we obtain T, a spanning tree of G and $\mathcal{F}_{\infty}(G)$. Taking the free group H_T with $E(G) \cup E(G^c)$ as generators and wtih the relations

- $uv = 1$ for all the edges of T
- $(uv)(vw) = uw$ if $\{u, v, w\}$ is a simplex of $\mathcal{F}_{\infty}(G)$

we have that $H_T \cong \pi_1(\mathcal{F}_\infty(G))$ (see [\[11\]](#page-34-4) Theorem 7.34). Take $uv \in E(G) \cup E(G^c) - E(T)$.

If u, v are in the same block, this block must be a cycle. If the cycle has 4 or more vertices, there is a uv-path $uw_1w_2\cdots w_rv$ in T. Now, $\{u, w_1, v\}$, $\{w_1. w_2, v\}$, ..., $\{w_{r-1}, w_r, v\}$ are simplicies of $\mathcal{F}_{\infty}(G)$, then $uv = w_1v = w_2v = \cdots = w_lv = 1$. If the cycle is uvw, because there are $k \geq 2$ blocks, one of the vertices must be a cut vertex:

- If u is a cut vertex, u has a neighbor x in another block such that ux is in T. Then $\{u, v, x\}$ is a simplex of $\mathcal{F}_{\infty}(G)$ and $uv = xv$. Now, $\{v, w, x\}$ and $\{u, w, x\}$ are simplices, thus $xv = xw = uw = 1$. The case in which v is a cut vertex is analogous.
- If w is a cut vertex, w has a neighbor x in another block such that wx is in T. Then $\{u, v, x\}, \{u, w, x\}$ and $\{v, w, x\}$ are simplices. Therefore $xv = vw = 1 = uw = ux$ and $uv = ux = 1$.

If u, v are in different blocks, then there are cut vertices w_1, \ldots, w_r , with $r \geq 1$, such that they are on the only uv-path in T and w_j it is not in the only u w_i -path for any $j > i$, and there are no more cut vertices in the path. Then $\{u, w_1, v\}, \{w_1, w_2, v\}, \ldots, \{w_{r-1}, w_r, v\}$ are simplices and $uv = w_1v = w_2v = \cdots = w_rv = 1$.

Therefore $\pi_1(\mathcal{F}_\infty(G)) \cong H_T \cong 0$.

Corollary 30. Let G a cactus graph such all of its blocks are cycles and does not have saturated blocks, then

 \Box

$$
\mathcal{F}_{\infty}(G) \simeq \mathbb{S}^{n-b(G)-1}.
$$

Proof. If $b(G) = 1$, then G is a cycle and $\mathcal{F}_{\infty}(G) \cong \mathbb{S}^{n-2}$. Assume $b(G) \geq 2$, then, by Lemma [29,](#page-17-0) $\mathcal{F}_{\infty}(G)$ is simply connected and, by Lemma [28,](#page-17-1) $\mathcal{F}_{\infty}^{*}(G) \simeq \mathbb{S}^{b(G)-1}$. Therefore, by Theorem [2,](#page-2-0) $\mathcal{F}_{\infty}(G)$ is a simply connected complex such that its only nontrivial reduced homology group is in dimension $q = n - b(G) - 1$, which is isomorphic to Z. By Theorem [3,](#page-2-1) $\mathcal{F}_{\infty}(G)$ is homotopy equivalent to a sphere of the desired dimension. □

Theorem 31. If G is a cactus graph then $\mathcal{F}_{\infty}(G)$ is either contractible or homotopy equivalent to a sphere of dimension at least $n - b(G) - 1$.

Proof. If $\delta(G) = 1$, then $\mathcal{F}_{\infty}(G) \simeq *$. Assume $\delta(G) = 2$. If there is a cut vertex of degree 2, then $\mathcal{F}_{\infty}(G) \simeq *$. Assume there is no cut vertex of degree 2. If G has a bridge e, then $G - e = G_1 + G_2$ and, by Proposition [12,](#page-7-0) $\mathcal{F}_{\infty}(G) = \mathcal{F}_{\infty}(G_1) * \mathcal{F}_{\infty}(G_2)$. If G has more bridges, then we continue this process until we get that $\mathcal{F}_{\infty}(G) = \mathcal{F}_{\infty}(H_1) * \cdots * \mathcal{F}_{\infty}(H_{r+1}),$ where r is the number of bridges and each H_i is a cactus graph such that every block is a cycle. So, if every $\mathcal{F}_{\infty}(H_i)$ has n_i vertices, is not contractible and is homotopy equivalent to a sphere of dimension at least $n_i - b(H_i) - 1$, $\mathcal{F}_{\infty}(G)$ will be homotopy equivalent to a sphere of dimension at least $n - b(G) + r - 1 > n - b(G) - 1$. Therefore we only need to prove the result for cactus graphs which do not have blocks isomorphic to K_2 .

If G does not have saturated blocks, by Corollary [30,](#page-18-0)

$$
\mathcal{F}_{\infty}(G) \simeq \mathbb{S}^{n-b(G)-1}.
$$

So assume $sb(G) \geq 1$, which implies that $b(G) \geq 4$. Now, we prove the result by induction on $sv(G)$. If $sv(G) = 0$, then take B_0 a saturated block of G and B_1, \ldots, B_k the remaining blocks. Let $X_i = V(G) - V(B_i)$, then X_0, X_1, \ldots, X_k are the facets of $\mathcal{F}^*_{\infty}(G)$. Because B_0 is saturated,

$$
\bigcap_{i=1}^k X_i = \varnothing.
$$

Let $S \subseteq [k] - \{0\}$ such that

$$
\sigma = \bigcap_{i \in S} X_i \neq \varnothing.
$$

Then there is $0 < j \leq k$ such that $j \notin S$ and $V(B_j) \cap \sigma \neq \emptyset$, with B_j a non-saturated block or a saturated block (which can not share vertices with B_0). Then there is a vertex v in $V(B_j)$ such that v is not vertex of the blocks with index in S nor is a vertex of B_0 , therefore $v \in X_0$, $v \in \sigma$ and $X_0 \cap \sigma \neq \emptyset$. From this we get that the nerve is a cone with apex vertex X_0 and $\mathcal{F}_{\infty}^*(G) \simeq *$. Then, by Lemma [29](#page-17-0) and Theorem [2,](#page-2-0) $\mathcal{F}_{\infty}(G)$ is simply connected and all of its reduced homology groups are trivial. Therefore, by Theorem [1,](#page-2-2) $\mathcal{F}_{\infty}(G)$ is contractible. This argument only used that there is an isolated saturated block, a saturated block which does not have saturated vertices; therefore we can assume that there is no isolated saturated block.

Assume the result is true for $sv(G) \leq k$ and let G be a cactus graph with $sv(G) = k + 1$ and with all of its blocks isomorphic to cycles. By Lemma [27](#page-16-1) there is B_0 a saturated block such that only one of its vertices is a saturated vertex, say v , and in the connected component of $B(G) - v$ which contains B_0 there are no more saturated blocks. We call G_1 the subgraph formed by the blocks in this connected component, and G_2 the subgraph induced by the remaining blocks. Then $G = G_1 \cup G_2$ and $G_1 \cap G_2 \cong K_1$. Now

$$
lk_{\mathcal{F}_{\infty}(G)}(v) = lk_{\mathcal{F}_{\infty}(G_1)}(v) * lk_{\mathcal{F}_{\infty}(G_2)}(v)
$$

We will show that $lk_{\mathcal{F}_{\infty}(G_1)}(v) \simeq *$. There are two possibilities:

1. $B_0 \cong C_3$. Then $V(B_0) = \{v, v_1, v_2\}$ and $G_1 = H_1 \cup B_0 \cup H_2$, with $V(H_1) \cap V(B_0) =$ $\{v_1\}, V(H_2) \cap V(B_0) = \{v_2\}$ and $V(H_1) \cap V(H_2) = \emptyset$. Then, by Lemma [19,](#page-8-1) $lk_{\mathcal{F}_{\infty}(G_1)}(v) \simeq \text{hocolim}(\mathcal{S})$ with S the diagram:

$$
\mathcal{F}_{\infty}(H_1) * \mathcal{F}_{\infty}(H_2 - v_2) \longleftrightarrow \mathcal{F}_{\infty}(H_1 - v_1) * \mathcal{F}_{\infty}(H_2 - v_2) \longleftrightarrow \mathcal{F}_{\infty}(H_1 - v_1) * \mathcal{F}_{\infty}(H_2)
$$

By construction, G_1 does not have saturated blocks, then $\delta(H_1 - v_1) = 1$ or it has a cut vertex of degree 2. Therefore $\mathcal{F}_{\infty}(H_1 - v_1) \simeq *$. Analogously, $\mathcal{F}_{\infty}(H_2 - v_2) \simeq *$. From this, we get that hocolim $(\mathcal{S}) \simeq *$.

2. $B_0 \cong C_n$ with $n \ge 4$. Let v_1, v_2 be the neighbors of v in B_0 and take H be the graph obtained from G_1 by erasing v and adding the edge v_1v_2 . Then

$$
lk_{\mathcal{F}_{\infty}(G_1)}(v) = \mathcal{F}_{\infty}(H) \simeq *,
$$

because $\mathcal{F}_{\infty}(H)$ has only one saturated block.

Therefore $lk_{\mathcal{F}_{\infty}(G)}(v) \simeq *$ and $\mathcal{F}_{\infty}(G) \simeq \mathcal{F}_{\infty}(G-v)$. If there is a non-saturated block which contains v, then $\delta(G-v) = 1$ or there is a cut vertex of degree 2, and therefore $\mathcal{F}_{\infty}(G) \simeq *$. Assume that there is no non-saturated block with v among its vertices. Now, in $G - v$, all the remaining edges of the blocks that contain v are bridges, so we can remove them, let H be the graph thus obtained. If $B_0, B_1, \ldots, B_{l-1}$ are the blocks that contain v, with $n_0, n_1, \ldots, n_{l-1}$ their respective orders, then $H = H_1 + \cdots + H_r$ where

$$
r = \sum_{i=0}^{l-1} n_i - 1.
$$

By inductive hypothesis, each $\mathcal{F}_{\infty}(H_i)$ is contractible or is homotopy equivalent to a sphere of dimension at least $|V(H_i)| - b(H_i) - 1$. Then, $\mathcal{F}_{\infty}(H)$ is contractible or it has the homotopy type of a sphere of dimension at least

$$
r-1+\sum_{i=1}^r |V(H_i)|-b(H_i)-1=n-1-(b(G)-l)-1=n-b(G)+l-2>n-b(G)-1.
$$

4.2 Graph operations

4.2.1 Join of graphs

Given two graphs G and H with disjoint vertex sets, we define their join as the graph $G*H$ with $V(G * H) = V(G) \cup V(H)$ and

$$
E(G*H) = E(G) \cup E(H) \cup \{uv: u \in V(G) \text{ and } v \in V(H)\}.
$$

It is well-known that $\mathcal{F}_0(G * H) = \mathcal{F}_0(G) \sqcup \mathcal{F}_0(H)$.

Lemma 32. Let G and H graphs with disjoint vertex sets with orders n_1 and n_2 respectively. Then:

1.
$$
\mathcal{F}_1(G * H) \simeq \mathcal{F}_1(G) \vee \mathcal{F}_1(H) \vee \bigvee_{n_1 n_2 - 1} \mathbb{S}^1
$$

2. If $\mathcal{F}_0(G)$ and $\mathcal{F}_0(H)$ are connected. Then, for all $d \geq 2$

$$
\mathcal{F}_d(G*H) \simeq \left(\bigvee_{n_2-1} \Sigma sk_{d-1} \mathcal{F}_0(G)\right) \vee \left(\bigvee_{n_1-1} \Sigma sk_{d-1} \mathcal{F}_0(H)\right) \vee \left(\bigvee_{(n_1-1)(n_2-1)} \mathbb{S}^2\right) \vee A \vee B
$$

with
$$
A = \mathcal{F}_d(G) \cup C(sk_{d-1}\mathcal{F}_0(G))
$$
 and $B = \mathcal{F}_d(H) \cup C(sk_{d-1}\mathcal{F}_0(H))$

Proof. For $d = 1$,

$$
\mathcal{F}_1(G*H)=\mathcal{F}_1(G)\cup \mathcal{F}_1(H)\cup K_{n_1,n_2}.
$$

Now $\mathcal{F}_1(G) \cap \mathcal{F}_1(H) \cap K_{n_1,n_2} = \mathcal{F}_1(G) \cap \mathcal{F}_1(H) = \emptyset$, therefore $\mathcal{F}_1(G * H)$ is homotopy equivalent to the homotopy pushout of

$$
X \longleftarrow sk_0 \mathcal{F}_1(G) \longrightarrow \mathcal{F}_1(G),
$$

where X is the homotopy pushout of

$$
\mathcal{F}_1(H) \longleftarrow sk_0 \mathcal{F}_1(H) \longrightarrow K_{n_1,n_2}.
$$

Thus

$$
X \simeq \mathcal{F}_1(H) \vee \bigvee_{n_1(n_2-1)} \mathbb{S}^1.
$$

From this the result follows.

For $d \geqslant 2$,

$$
\mathcal{F}_d(G*H) = \mathcal{F}_d(G) \cup \mathcal{F}_d(H) \cup K_1 \cup K_2,
$$

with $K_1 = \bigcup_{u \in V(H)} \{u\} * sk_{d-1} \mathcal{F}_0(G)$ and $K_2 = \bigcup_{u \in V(G)} \{u\} * sk_{d-1} \mathcal{F}_0(H)$. Now:

$$
K_1 \cong \bigvee_{n_2-1} \Sigma sk_{d-1} \mathcal{F}_0(G),
$$

$$
K_2 \cong \bigvee_{n_1-1} \Sigma sk_{d-1} \mathcal{F}_0(H).
$$

Taking $L_1 = \mathcal{F}_d(G)$ and $L_2 = \mathcal{F}_d(H)$, we have that

$$
L_1 \cap L_2 = \emptyset, \ K_1 \cap L_1 = sk_{d-1} \mathcal{F}_0(G), \ K_2 \cap L_2 = sk_{d-1} \mathcal{F}_0(H), \ K_1 \cap K_2 \cong K_{n,m},
$$

$$
L_1 \cap K_1 \cap K_2 = L_1 \cap K_2 \cong \bigvee_{n_1 - 1} \mathbb{S}^0,
$$

$$
L_2 \cap K_2 \cap K_1 = L_2 \cap K_1 \cong \bigvee_{n_2 - 1} \mathbb{S}^0.
$$

Taking $X = K_1 \cup L_1$ and $Y = K_2 \cup L_2$, we have that $\mathcal{F}_d(G * H) = X \cup Y$ and $X \cap Y =$ $(L_1 \cap K_2) \cup (L_2 \cap K_1) \cup (K_1 \cap K_2) = K_1 \cap K_2$. Therefore $\mathcal{F}(G * H, d) \simeq \text{hocolim}(\mathcal{S})$ with

$$
\mathcal{S} \colon X \longrightarrow K_{n,m} \longrightarrow Y
$$

Now, the inclusion $i: K_{n,m} \longrightarrow X$ is really the inclusion $K_{n,m} \longrightarrow K_1$, which is nullhomotopic, and therefore i is null-homotopic. In the same way we see that the inclusion in Y is null-homotopic and that

$$
\mathcal{F}_d(G*H) \simeq X \vee Y \vee \bigvee_{(n_1-1)(n_2-1)} \mathbb{S}^2.
$$

Now, $K_1 \cap L_1 = sk_{d-1}F_0(G)$ and its inclusion in K_1 is null-homotopic, therefore we can compute the homotopy type of X by pasting these two homotopy pushout squares:

$$
sk_{d-1} \mathcal{F}_0(G) \longrightarrow \begin{array}{c} K_1 \\ \downarrow \\ \downarrow \\ L_1 \longrightarrow L_1 \cup C(sk_{d-1} \mathcal{F}_0(G)) \longrightarrow K_1 \vee (L_1 \cup C(sk_{d-1} \mathcal{F}_0(G))) \simeq X \end{array}
$$

Now $L_1 \cup C(sk_{d-1}F_0(G)) = A$. With an similar argument for Y we arrive at the result. \Box

With the last lemma we can construct graphs for which $\mathcal{F}_{\infty}(G)$ is not homotopy equivalent to a wedge of spheres. Let K be a triangulation of the projective plane and let H be the complement graph of the 1-skeleton of the baricentric subdivision, then $\mathcal{F}_0(G) \cong K$ and $G = P_4 * H$ is a graph such that $\mathcal{F}_d(G)$ has torsion for all $d \geq 3$.

Lemma 33. Let G be a graph and take $d \geq 1$, then

$$
\mathcal{F}_d(K_1 * G) \simeq \mathcal{F}_d(G) \cup C(sk_{d-1} \mathcal{F}_0(G))
$$

Proof. The link of the apex vertex is $sk_{d-1}\mathcal{F}_0(G)$, thus the homotopy pushout square

$$
s k_{d-1} \mathcal{F}_0(G) \longrightarrow \begin{matrix} \downarrow \\ \downarrow \\ \mathcal{F}_d(G) \longrightarrow \mathcal{F}_d(G) \cup C(s k_{d-1} \mathcal{F}_0(G)) \end{matrix}
$$

 \Box

computes $\mathcal{F}(K_1 * G, d)$.

Theorem 34. For the complete bipartite graph we have that $\mathcal{F}_0(K_{n,m}) \simeq \mathbb{S}^0$,

$$
\mathcal{F}_1(K_{n,m}) \simeq \bigvee_{nm-1} \mathbb{S}^1,
$$

$$
\mathcal{F}_d(K_{n,m}) \simeq \bigvee_{(n-1)(m-1)} \mathbb{S}^2 \vee \bigvee_{n\binom{m-1}{d}+m\binom{n-1}{d}} \mathbb{S}^d,
$$

for $\infty > d \geqslant 2$ and

$$
\mathcal{F}_{\infty}(K_{n,m}) \simeq \bigvee_{(n-1)(m-1)} \mathbb{S}^2.
$$

Proof. If $d = 0$ is clear. The case $d = 1$ is a particular case of Lemma [32.](#page-21-2) For $d \ge 2$, by Lemma [32](#page-21-2)

$$
\mathcal{F}_d(K_{n,m}) \simeq \left(\bigvee_{m-1} \Sigma sk_{d-1} \mathcal{F}_0(K_n^c)\right) \vee \left(\bigvee_{n-1} \Sigma sk_{d-1} \mathcal{F}_0(K_m^c)\right) \vee \left(\bigvee_{(n-1)(m-1)} \mathbb{S}^2\right) \vee A \vee B
$$

with $A = \mathcal{F}_d(K_n^c) \cup C(sk_{d-1} \mathcal{F}_0(K_n^c))$ and $B = \mathcal{F}_d(K_m^c) \cup C(sk_{d-1} \mathcal{F}_0(K_m^c)).$

Now, for all d, k, r ,

$$
\mathcal{F}_d(K_k^c) \cong \Delta^{k-1}, \; sk_r \mathcal{F}_d(K_k^c) \simeq \bigvee_{\binom{k-1}{r+1}} \mathbb{S}^r;
$$

therefore

$$
A \simeq \bigvee_{\binom{n-1}{d}} \mathbb{S}^d; \ B \simeq \bigvee_{\binom{m-1}{d}} \mathbb{S}^d,
$$

from which we obtain the result.

Corollary 35. Let G_1, G_2, \ldots, G_k be vertex disjoint graphs. For $d \geq 1$, if $\mathcal{F}_d(G_i) \simeq *$ for all i, then

$$
\mathcal{F}_d(G_1 * G_2 * \cdots * G_k) \simeq \bigvee_{\frac{(k-1)(k-2)}{2}} \mathbb{S}^1 \vee \bigvee_{i < j} \mathcal{F}_d(G_i * G_j)
$$

Proof. Let V_i be the vertex set of G_i and take $G = G_1 * G_2 * \cdots * G_k$. If we take vertices from more than two sets of the partition, we will always have a cycle, and therefore each facet of the complex is contained in $V_i \cup V_j$ for some $i \neq j$. Then, taking $X_{ij} = \mathcal{F}_d(G[V_i \cup V_j])$ for $i < j$, we have that $\mathcal{F}_d(G) = \left(\begin{array}{c} \end{array} \right) X_{ij}$ and we can define a bijection $\gamma : \{ij : i < j\} \longrightarrow E(K_k)$ $i \leq j$ such that the hypothesis of Lemma [6](#page-4-1) are achieved. \Box

As an immediate consequence we have the homotopy type for the multipartite graphs Corollary 36. For $d \geq 1$,

$$
\mathcal{F}_d(K_{n_1,\ldots,n_k}) \simeq \bigvee_{\frac{(k-1)(k-2)}{2}} \mathbb{S}^1 \vee \bigvee_{i < j} \mathcal{F}_d(K_{n_i,n_j}).
$$

Theorem 37. [\[8](#page-34-7)]

$$
\mathcal{F}_0(C_n) \simeq \begin{cases} \mathbb{S}^{r-1} \vee \mathbb{S}^{r-1} & \text{if } n = 3r \\ \mathbb{S}^{r-1} & \text{if } n = 3r+1 \\ \mathbb{S}^r & \text{if } n = 3r+2 \end{cases}
$$

Proposition 38. Let W_{n+1} be the wheel on $n + 1$ vertices, then

$$
\mathcal{F}_d(W_{n+1}) \simeq \begin{cases} \mathbb{S}^{3r-2} \vee \mathbb{S}^r \vee \mathbb{S}^r & \text{if } n = 3r \\ \mathbb{S}^{3r-1} \vee \mathbb{S}^r & \text{if } n = 3r+1 \\ \mathbb{S}^{3r} \vee \mathbb{S}^{r+1} & \text{if } n = 3r+2 \end{cases}
$$

 \Box

for $d > \lfloor \frac{n}{2} \rfloor - 1$ and

$$
\mathcal{F}_1(W_{n+1}) \simeq \begin{cases} \sqrt{\mathbb{S}^{2r-1}} \vee \sqrt{\mathbb{S}^1} & \text{if } n = 4r \\ \mathbb{S}^{2r-1} \vee \sqrt{\mathbb{S}^1} & \text{if } n = 4r+1 \\ \mathbb{S}^{2r} \vee \sqrt{\mathbb{S}^1} & \text{if } n = 4r+2 \\ \mathbb{S}^{2r+1} \vee \sqrt{\mathbb{S}^1} & \text{if } n = 4r+3 \\ n-1 & \text{if } n = 4r+3 \end{cases}
$$

Proof. Since $\alpha(C_n) = \lfloor \frac{n}{2} \rfloor$, for $d > \lfloor \frac{n}{2} \rfloor - 1$ we have that $\mathcal{F}_0(C_n) = sk_{d-1}\mathcal{F}_0(C_n)$. By Lemma [33,](#page-23-0)

$$
\mathcal{F}_d(W_{n+1}) \simeq \mathcal{F}_d(C_n) \cup C(\mathcal{F}_0(C_n)).
$$

By Theorem [37,](#page-24-0) the inclusion of the intersection is null-homotopic, therefore

$$
\mathcal{F}_d(W_{n+1}) \simeq \mathcal{F}_\infty(C_n) \vee \Sigma \mathcal{F}_0(C_n)
$$

 \Box

For $d = 1$, $sk_0 \mathcal{F}_0(C_n, 0) = \bigvee_{n=1}^{\infty} \mathbb{S}^0$, the rest of the proof is the same as before.

4.2.2 Graph products

Proposition 39.

$$
\mathcal{F}_{\infty} (P_2 \square P_k) \simeq \begin{cases} \mathbb{S}^{4r-1} & \text{if } k = 3r \\ * & \text{if } k = 3r+1 \\ \mathbb{S}^{4r+2} & \text{if } k = 3r+2. \end{cases}
$$

Proof. By Theorem [22,](#page-9-1) \mathcal{F}_{∞} ($P_2 \square P_k$) is simply connected. We will show that it has at most one non-trivial reduced homology group. The Alexander dual of $\mathcal{F}_{\infty} (P_2 \square P_k)$ has as maximal simplicies the complements of $X_i = \{(i,1), (i+1,1), (i,2), (i+1,2)\}$ for $1 \leq i \leq j$ $k-1$. Taking $U_i = X_i^c$ and U the cover formed by these U_i , we have that

$$
\mathcal{N}(U) \simeq \mathcal{F}_0^*(P_k).
$$

It is standard that [\[8\]](#page-34-7):

$$
\mathcal{F}_0(P_k) \simeq \begin{cases} \mathbb{S}^{r-1} & \text{if } k = 3r \\ * & \text{if } k = 3r + 1 \\ \mathbb{S}^r & \text{if } k = 3r + 2. \end{cases}
$$

Thus, by Theorem [2,](#page-2-0) $\mathcal{N}(U)$ has non-trivial reduced cohomology groups if $k = 3r$ or $k =$ $3r + 2$, in which case the groups are in dimensions are $2(r - 1)$ and $2r - 1$ respectively. Therefore \mathcal{F}_{∞} $(P_2 \square P_k)$ is contractible if $k = 3r + 1$ and

$$
\tilde{H}_q(\mathcal{F}_{\infty}(P_2 \square P_{3r})) \cong \begin{cases} \mathbb{Z} & \text{if } q = 4r - 1 \\ 0 & \text{if } q \neq 4r - 1, \end{cases}
$$

$$
\tilde{H}_q(\mathcal{F}_{\infty}(P_2 \square P_{3r+2})) \cong \begin{cases} \mathbb{Z} & \text{if } q = 4r + 2 \\ 0 & \text{if } q \neq 4r + 2. \end{cases}
$$

By Theorem [3,](#page-2-1) in these cases the complex is homotopy equivalent to a sphere of the desired dimension. □

It is known [\[6](#page-34-8)] that

$$
\mathcal{F}_0(K_n \times K_m) \simeq \bigvee_{(n-1)(m-1)} \mathbb{S}^1.
$$

Now we will show what happens for $d \geq 1$.

Proposition 40.

$$
\mathcal{F}_1(K_n \times K_m) \simeq \bigvee_{\frac{(nm-4)(n-1)(m-1)}{4}} \mathbb{S}^2
$$

Proof. We take $V(K_r) = [r] - \{0\}$ for any r. We proceed by induction on n. For $n = 1$, the result is clear. For $n = 2$ we will prove it by induction on m. For $m = 1, 2$ it is clear and for $m = 3$, $K_2 \times K_3 \cong C_6$. Taking $v_i = (1, i)$ and $u_i = (2, i)$, we have that $lk(v_m) = X \cup Y$, where $Y = \mathcal{F}_1(K_2 \times K_m) - N[v_n]$ and X is the complex with facets $\{u_i, v_i, u_m\}$ for $i \geq m-1$. Then $X \simeq *$, as it is a cone with apex u_m , and $X \cap Y \cong K_{i,m} \simeq *$. Therefore,

$$
lk(v_n) \simeq Y \cong \mathcal{F}_1(K_{1,m-1}) \simeq \bigvee_{m-2} \mathbb{S}^1.
$$

Taking $H = K_2 \times K_m - v_m$, the link of u_m in $\mathcal{F}_1(H)$ has as facets the simplex $\{u_1, \ldots, u_{m-1}\}$ and the edges $\{u_i, v_i\}$ for $i \geq m - 1$, therefore it is contractible and

$$
\mathcal{F}_1(H) \simeq \mathcal{F}_1(H - u_m) \cong \mathcal{F}_1(K_2 \times K_{m-1}) \simeq \bigvee_{\frac{(m-2)(m-3)}{2}} \mathbb{S}^2,
$$

from which the result follows.

Now assume the result is true for $K_r \times K_m$ for all $r \leq n-1$. Take $v_i = (n, i)$, $G_0 = K_n \times K_m$, $G_i = G_{i-1} - v_i$ for $i \geq 1$, $X_{j,k}^i = |\{(j,k),(j,i),(n,k)\}|$ for $k \geq i+1$ and $j \le n - 1, X_{j,k}^i = |\{(j,k),(j,i)\}| \text{ for } k \le i - 1 \text{ and } j \le n - 1,$

$$
X^i = \bigcup_{k \neq i, j \leq n-1} X^i_{j,k}
$$

and $Y^i = \mathcal{F}_1(G_{i-1} - N[v_i])$. Then, taking L_i the link of v_i in $\mathcal{F}_1(G_{i-1})$, we have that

$$
L_i = X^i \cup Y^i.
$$

Now, in X^i , the vertices (j, k) with $j \leq n - 1$ and $k \neq i$ are only in one facet and can be erased, therefore X^i is homotopy equivalent to the subcomplex with maximal facets $\{(j,i),(n,k)\}\$ with $k\geqslant i+1$ and $j\leqslant n-1$, which is isomorphic to $K_{n-1,m-i}$. Because $X^i \cap Y_i$ is isomorphic to this subacomplex, we have that

$$
L_i \simeq Y^i \cong \mathcal{F}_1(K_{n-i,m-1}) \simeq \bigvee_{(m-1)(n-i)-1} \mathbb{S}^1
$$

for $i \leq n-1$. Now, $L_n \simeq Y^n \simeq *$, therefore

$$
\mathcal{F}_1(G_{n-1}) \simeq \mathcal{F}_1(G_n) \cong \mathcal{F}_1(K_{n-1} \times K_m) \simeq \bigvee_{\frac{((n-1)m-4)(n-2)(m-1)}{4}} \mathbb{S}^2.
$$

From this we have that

$$
\mathcal{F}_1(G_0) \simeq \mathcal{F}_1(K_{n-1} \times K_m) \vee \Sigma Y^1 \vee \Sigma Y^2 \vee \cdots \vee \Sigma Y^{n-1}.
$$

Now $\Sigma Y^1 \vee \Sigma Y^2 \vee \cdots \vee \Sigma Y^{n-1}$ is homotopy equivalent to the wedge of

$$
\sum_{i=1}^{m-1} i(n-1) - 1 = \frac{(n-1)m(m-1)}{2} - (m-1)
$$

copies of the 2-sphere. Since

$$
\frac{((n-1)m-4)(n-2)(m-1)}{4} = \sum_{i=1}^{n-2} \frac{im(m-1)}{2} - (m-1),
$$

Figure 2:

we have that $\mathcal{F}_1(K_n \times K_m)$ is homotopy equivalent to the wedge of

$$
\sum_{i=1}^{n-1} \frac{im(m-1)}{2} - (m-1) = \frac{(nm-4)(n-1)(m-1)}{4}
$$

2-spheres.

Lemma 41. For $d \geq 2$, $\mathcal{F}_{d+1}(K_2 \times K_n) \simeq \mathcal{F}_d(K_2 \times K_n)$

Proof. We know that $\mathcal{F}_d(K_2 \times K_n)$ is simply connected for all $d \geq 2$, because $\mathcal{F}_1(K_2 \times K_n)$ is a wedge of 2-spheres. We will show that $H_q(\mathcal{F}_{d+1}(K_2 \times K_n), \mathcal{F}_d(K_2 \times K_n)) \cong 0$ for all q. We know that $H_q(\mathcal{F}_{d+1}(K_2 \times K_n), \mathcal{F}_d(K_2 \times K_n)) \cong 0$ for all $q \leq d$. For $q \geq d+3$, for any q-simplex σ of $\mathcal{F}_{d+1}(K_2 \times K_n)$, we can partition its vertices in two sets V_1, V_2 such that all the vertices in V_i are of the form (i, j) for some j. Next we show that $|V_1| = 0$ or $|V_2| = 0$. If not, we can assume that

$$
|V_1| \le \left\lfloor\frac{d+3}{2}\right\rfloor \le \left\lceil\frac{d+3}{2}\right\rceil \le |V_2|
$$

therefore $|V_2| \geq 3$; there are several cases:

- If $|V_1| = 1$, then $|V_2| \ge d+3$ and the vertex of V_1 has degree at least $d+2$, which can not happen.
- If $|V_1| = 2$, then $|V_2| \ge d + 2$ and there will be at least two vertices of V_2 such their second coordinates are different from those of the vertices of V_1 ; therefore there will be an induced 4-cycle in the vertices of σ , which can not happen.

 \Box

• If $|V_1| \geq 3$, because $|V_2| \geq 3$, there will be an induced 4-cycle or an induced 6-cycle in the vertices of σ , which can not happen.

Therefore $|V_1| = 0$ or $|V_2| = 0$ and σ is a simplex of $\mathcal{F}_d(K_2 \times K_n)$. From this, we have that $H_q(\mathcal{F}_{d+1}(K_2 \times K_n), \mathcal{F}_d(K_2 \times K_n)) \cong 0$ for all $q \geq d+3$.

For $q = d+2$, the only q-simplices of $F_{d+1}(K_2 \times K_n)$ which are not simplices of $\mathcal{F}_d(K_2 \times K_1)$ K_n) are of the form $|V_1| = 1$ and $|V_2| = d + 2$ (or vice versa), where the only vertex of V_1 is adjacent to all but one vertex of V_2 (Figure [2\(a\)\)](#page-28-0). For $q = d + 1$, the only q-simplices of $F_{d+1}(K_2 \times K_n)$ which are not simplices of $\mathcal{F}_d(K_2 \times K_n)$ are of the form $|V_1| = 1$ and $|V_2| = d + 1$ (or vice versa), where the only vertex of V_1 is adjacent to all the vertices of V_2 (Figure [2\(b\)\)](#page-28-1). From all this, we get that there are no relative $d+2$ -cycles and that all of the relative $d + 1$ -cycles are images of some relative $d + 2$ -boundary. Therefore the remaining two relative homology groups are also trivial.

From all this we have that the inclusion $\mathcal{F}_{d+1}(K_2 \times K_n) \longrightarrow \mathcal{F}_d(K_2 \times K_n)$ induces an isomorphism for all homology groups between simply connected complexes, therefore $\mathcal{F}_{d+1}(K_2 \times K_n) \simeq \mathcal{F}_d(K_2 \times K_n).$ \Box

Proposition 42. For $d \geq 2$,

$$
\mathcal{F}_d(K_2 \times K_n) \simeq \bigvee_{\binom{n}{3}} \mathbb{S}^4 \vee \bigvee_{\binom{n-1}{3}} \mathbb{S}^3.
$$

Proof. We only have to prove it for $d = 2$. The result is clear for $n = 1, 2, 3$. Assume $n \ge 4$. Taking $k = \binom{n}{2}$ 3), let X_1, \ldots, X_k be the subcomplexes of $\mathcal{F}_2(K_2 \times K_n)$ corresponding to all the induced 6-cycles. Then $X_i \cong \mathbb{S}^4$. The other facets of $\mathcal{F}_2(K_2 \times K_n)$, besides the ones in some X_i , are $\{1\} \times \underline{n}$ and $\{2\} \times \underline{n}$. Then

$$
\mathcal{F}_2(K_2 \times K_n) = X_1 \cup X_2 \cup \cdots \cup X_k \cup Y_1 \cup Y_2
$$

where $Y_1 = \mathcal{P}(\{1\} \times \underline{n}) - \{\emptyset\}$ and $Y_2 = \mathcal{P}(\{2\} \times \underline{n}) - \{\emptyset\}$. Now we will calculate the homology of $\mathcal{F}_2(K_2 \times K_n)$ using the Mayer-Vietoris spectral sequence. Taking $U =$

Because the nerve of X_1, X_2, \ldots, X_k is isomorphic to the nerve of 2-simplices of $sk_2\Delta^{n-1}$, and U is isomorphic to the suspension of this nerve, we have that the second page is

where $r = \binom{n-1}{3}$. From this we have that $E_{p,q}^{\infty} = E_{p,q}^2$. Therefore

$$
\tilde{H}_q(\mathcal{F}_2(K_2 \times K_n)) \cong \begin{cases} \mathbb{Z}^k & \text{if } q = 4 \\ \mathbb{Z}^r & \text{if } q = 3 \\ 0 & \text{if } q \neq 4,3 \end{cases}
$$

Therefore, because $\mathcal{F}_1(K_2 \times K_n)$ is simply connected, $\mathcal{F}_2(K_2 \times K_n)$ is a simply connected complex which satisfies the hypothesis of Theorem [4,](#page-2-3) from which we see that is has the desired homotopy type.

Theorem 43. For $d \geq 2$,

$$
\mathcal{F}_d(K_n \times K_m) \simeq \bigvee_a \mathbb{S}^4 \vee \bigvee_{b+c} \mathbb{S}^3,
$$

where $a = \binom{m}{2} \binom{n}{3} + \binom{n}{2} \binom{m}{3}$, $b = \binom{m}{2} \binom{n-1}{3} + \binom{n}{2} \binom{m-1}{3}$ and $c = \binom{n-1}{2} \binom{m-1}{2}$.

Proof. In $\mathcal{F}_d(K_n \times K_m)$ the facets have their vertices contained in two rows or two columns, otherwise they will have a cycle. Then, taking the subgraphs

> $H_{i,j} = K_n \times K_m[\{(k,l): l = i \text{ or } l = j\}],$ $G_{i,j} = K_n \times K_m[\{(k,l): k = i \text{ or } k = j\}],$

and the complexes $X_{i,j} = \mathcal{F}_d(H_{i,j}), Y_{i,j} = \mathcal{F}_d(G_{i,j}),$ we have that

$$
\mathcal{F}_d(K_n \times K_m) = \bigcup_{e \in E(K_m)} X_e \cup \bigcup_{e \in E(K_n)} Y_e
$$

From the last Proposition we know that

$$
X_e \simeq \bigvee_{n \atop (3)} s^4 \vee \bigvee_{n-1 \atop (n-1)} s^3
$$

$$
Y_e \simeq \bigvee_{n \atop (3)} s^4 \vee \bigvee_{n-1 \atop (n-1)} s^3
$$

Taking the Mayer-Vietoris spectral sequence, the first page looks like

Where *U* is the nerve of the cover, $a = \binom{n}{2} \binom{m}{3} + \binom{m}{2} \binom{n}{3}$ and $b = \binom{n}{2} \binom{m-1}{3} + \binom{m}{2} \binom{n-1}{3}$. Now, U is isomorphic to the join of the nerve of the X's with the nerve of the Y's, which are homotopy equivalent to K_m and K_n respectively, therefore $\mathcal{U} \simeq \bigvee_c \mathbb{S}^3$ with $c = \binom{n-1}{2} \binom{m-1}{2}$. From all this, we have that the second page of the sequence is

Therefore $E_{p,q}^{\infty} = E_{p,q}^2$ and

$$
\tilde{H}_q(\mathcal{F}_d(K_n \times K_m)) \cong \begin{cases} \mathbb{Z}^a & \text{if } q = 4 \\ \mathbb{Z}^{b+c} & \text{if } q = 3 \\ 0 & \text{if } q \neq 4,3 \end{cases}
$$

As in the proof of the last theorem, we have a simply connected complex which satisfies the hypothesis of Theorem [4.](#page-2-3) □

In [\[1\]](#page-33-4) it was shown that

$$
\mathcal{F}_0(K_2 \times K_m \times K_n) \simeq \bigvee_{\frac{(n-1)(m-1)(nm-2)}{2}} \mathbb{S}^3.
$$

For other $d \geq 1$, because $K_2 \times K_2 \cong K_2 \cup K_2$ we have the following corollary

Corollary 44. For $d \geq 1$,

$$
\mathcal{F}_d(K_2 \times K_2 \times K_n) \simeq \begin{cases} \bigvee_{\substack{(n-2)^2(n-1)^2\\ \text{ odd}}} \mathbb{S}^5 & d=1\\ \bigvee_{\substack{(n-2)^2(n-1)^2\\ \text{ odd}}} \mathbb{S}^8 \vee \bigvee_{\substack{2\binom{n}{3} \binom{n-1}{3}}} \mathbb{S}^7 & d \geq 2 \end{cases}
$$

Question 45. What is the homotopy type of $K_2 \times K_m \times K_n$ for $d \geq 1$?

Acknowledgments. The author wishes to thank Omar Antolín-Camarena for his comments and suggestions which improved this paper.

Bibliography

- [1] Antolín-Camarena, O., and Carnero-Bravo, A. Homotopy type of the independence complex of some categorical products of graphs. https://arxiv.org/abs/2307.12401 (2023).
- [2] Björner, A. Matroid Applications, vol. 40 of Encyclopedia of mathematics and its applications. Cambridge University Press, 2009, ch. The Homology and Shellability of Matroids and Geometric Lattices, pp. 226–283.
- [3] Björner, A., and Tancer, M. Note: Combinatorial alexander duality—a short and elementary proof. Discrete and Computational Geometry 42 (2009), 586–593.
- [4] Chartrand, G., Lesniak, L., and Zhang, P. Graphs and Digraphs. Chapman & Hall/CRC, 2015.
- [5] Deshpande, P., and Singh, A. Higher independence complexes of graphs and their homotopy types. *Journal of the Ramanujan Mathematical Society 36*, 1 (2021), 53–71.
- [6] Goyal, S., Shukla, S., and Singh, A. Homotopy type of independence complexes of certain families of graphs. Contributions to Discrete Mathematics 16, 3 (2021), 74–92.
- [7] HATCHER, A. Algebraic Topology. Cambridge, 2002.
- [8] Kozlov, D. Complexes of directed trees. Journal of Combinatorial Theory, Series A 88 (1999), 112–122.
- [9] Munson, B. A., and Volić, I. Cubical Homotopy Theory, vol. 25 of New mathematical monographs. Cambridge, 2015.
- [10] Paolini, G., and Salvetti, M. Weighted sheaves and homology of artin groups. Algebraic and Geometric Topology 18, 7 (2018), 3943–4000.
- [11] Rotman, J. J. An Introduction to Algebraic Topology. Springer, 1998.
- [12] SALVETTI, M. Combinatorial Methods in Topology and Algebra, vol. 12 of Springer INdAM Series. Springer, 2015, ch. Some Combinatorial Constructions and Relations with Artin Groups, pp. 121–126.
- [13] Zuffi, L. Simplicial complexes from graphs towards graph persistence. Master's thesis, Università di Bologna, 2016.

Andrés Carnero Bravo Instituto de Matemáticas, UNAM, Mexico City, Mexico E-mail address: acarnerobravo@gmail.com