

# Multiscale scanning with nuisance parameters

CLAUDIA KÖNIG

`claudia-juliane.koenig@mathematik.uni-goettingen.de`  
Institute for Mathematical Stochastics, University of Göttingen

AXEL MUNK

`munk@math.uni-goettingen.de`  
Institute for Mathematical Stochastics, University of Göttingen

and  
Felix Bernstein Institute for Mathematical Statistics in the Bioscience,  
University of Göttingen

and  
DFG Cluster of Excellence "Multiscale Bioimaging - From Molecular  
Machines to Networks of Excitable Cells", University of Göttingen Medical  
Center

FRANK WERNER<sup>1</sup>

`frank.werner@uni-wuerzburg.de`  
Julius-Maximilians-Universität Würzburg (JMU), Institute of Mathematics,  
Würzburg, Germany

We develop a multiscale scanning method to find anomalies in a  $d$ -dimensional random field in the presence of nuisance parameters. This covers the common situation that either the baseline-level or additional parameters such as the variance are unknown and have to be estimated from the data. We argue that state of the art approaches to determine asymptotically correct critical values for multiscale scanning statistics will in general fail when such parameters are naively replaced by plug-in estimators. Opposed to this, we suggest to estimate the nuisance parameters on the largest scale and to use (only) smaller scales for multiscale scanning. We prove a uniform invariance principle for the resulting adjusted multiscale statistic (AMS), which is widely applicable and provides a computationally feasible way to simulate asymptotically correct critical values. We illustrate the implications of our theoretical results in a simulation study and in a real data example from super-resolution STED microscopy. This allows us to identify interesting regions inside a specimen in a pre-scan with controlled family-wise error rate.

*Keywords:* scan statistic, invariance principle, limit theorem, multiscale analysis, super-resolution microscopy

*AMS classification numbers:* 60F17, 62H10, 60G50, 62F03.

## 1. Introduction

Scanning a  $d$ -dimensional random field  $Y = (Y_i)_{i \in \{1, \dots, n\}^d}$  of random variables (r.v.'s)  $Y_i$  for anomalies in their distribution is a prominent topic in statistical research which is of great practical use. Applications range from the detection of anomalous clusters in social and biological networks and graphs (Arias-Castro et al., 2011; Chen and Neill, 2014; Sharpnack et al., 2016; Behr et al.,

---

<sup>1</sup>Corresponding author

2020) to detection of structural changes in imaging (Sharpnack, 2018; Caron et al., 2002; Arias-Castro et al., 2006; Kirch et al., 2023), genomics (Reiner-Benaim et al., 2014; Futschik et al., 2014), brain research (Lovato et al., 2021) and cell microscopy (Proksch et al., 2018), to mention a few.

## 1.1. Setup

In this paper, we consider a parametric model of the form

$$Y_i \sim F_{\theta_i, \xi}, \quad i \in I_n^d := \{1, \dots, n\}^d, \quad (1)$$

where the  $Y_i$  are independent, real valued r.v.'s distributed to some unknown  $F_{\theta_i, \xi}$  from a given, known class of distributions  $\mathcal{F}$  indexed in  $\theta$  and  $\xi$ . Here,  $i$  denotes a multi-index, which we assume to index a  $d$ -dimensional grid  $I_n^d = \{1, \dots, n\}^d$ . In what follows,  $\theta_i \in \Theta$  denotes the parameter of interest,  $\xi \in \Xi$  a collection of nuisance parameters, and  $\Theta \subset \mathbb{R}^{d_1}$  and  $\Xi \subset \mathbb{R}^{d_2}$  are the corresponding parameter spaces. Note, that  $\theta_i$  may vary between observations  $Y_i$ , whereas the nuisance parameter  $\xi$  is assumed to be the same for all  $i \in I_n^d$ , but we stress that our analysis can also be transferred to varying  $\xi$ . An exemplary case for model (1) would be a Gaussian field  $Y_i \sim \mathcal{N}(\theta_i, \Sigma)$ , where anomalies  $\theta_i$  occur in the mean as deviation from a ground truth  $\theta_0 \in \Theta$  and the unknown covariance  $\Sigma > 0$  is the nuisance parameter. For the moment, we do not pose structural assumptions on  $F_{\theta, \xi}$ , but require some smoothness and tail properties later on (see Assumption 1 below).

## 1.2. Anomaly detection

Given a baseline or reference intensity  $\theta_0 \in \Theta$  and a region  $R \subset I_n^d$ , the problem to detect an anomaly inside  $R$  can be formulated as the testing problem

$$H_{R,n} : \forall i \in R : \theta_i = \theta_0 \quad (2a)$$

vs.

$$K_{R,n} : \exists i \in R \text{ s.t. } \theta_i \neq \theta_0. \quad (2b)$$

Note, that in practice often  $d_1 = 1$ , i.e.  $\theta_i \in \mathbb{R}$ , and it might be more relevant to consider alternatives of the form  $\theta_i > \theta_0$  or  $\theta_i < \theta_0$ . For the sake of brevity, we consider only two-sided alternatives as in (2b) here, but we emphasize that the one-sided situation can be developed completely analogous (see Remark 3.8 below).

Depending on the structure of  $R$ , the likelihood ratio test (LRT) is known to have favourable properties, see e.g. Lehmann and Romano (2005). Therefore, throughout this paper, we consider the corresponding log LRT statistic

$$T_R(Y, \theta_0, \xi) := \sqrt{2 \log \left( \frac{\sup_{\theta \in \Theta} \prod_{i \in R} f_{\theta, \xi}(Y_i)}{\prod_{i \in R} f_{\theta_0, \xi}(Y_i)} \right)} \quad (3)$$

for (2), where  $f_{\theta, \xi}$  denotes the density or probability distribution function of  $F_{\theta, \xi}$ . Whenever this statistic exceeds a suitably chosen threshold (depending on  $R$  and the prescribed level  $\alpha \in (0, 1)$ ), the corresponding LRT will reject.

## 1.3. Multiscale scanning

However, as position and size of anomalies will in general be unknown, the region of interest  $R$  is not known in advance. Therefore, we aim to *scan* over  $\mathcal{R}_n \subset 2^{I_n^d}$  (with  $2^X$  being the power set of  $X$ ), a set of regions of interest (e.g. the system of all discrete cubes inside  $I_n^d$ ). If the system  $\mathcal{R}_n$  covers a range of spatial scales, then this approach is denoted as *multiscale scanning*, which has a long history in statistics, see e.g. Kulldorff (1997) for an early reference and Arias-Castro et al.

(2006); Walther (2010); Arias-Castro et al. (2011); Neill (2012); Mallik et al. (2013); Chan and Walther (2013); Sharpnack and Arias-Castro (2016); Proksch et al. (2018); König et al. (2020) for more recent works. A particular instance is multiscale change point detection, see e.g. Frick et al. (2014), where the field is of spatial dimension  $d = 1$ , or Chen et al. (2022) for arbitrary dimension  $d$ , and the references given there.

Outgoing from (2), multiscale scanning can be treated as a multiple testing problem

$$H_{R,n} \text{ vs. } K_{R,n} \quad \text{simultaneously over } \mathcal{R}_n. \quad (4)$$

The resulting scanning test then has to combine the local test statistics  $T_R$ ,  $R \in \mathcal{R}_n$ , in (3) in a suitable way, such that the number of false rejections over  $\mathcal{R}_n$  can be controlled. To this end, we aim to bound (asymptotically) the family wise error rate (FWER) by  $\alpha \in (0, 1)$ , i.e. (see e.g. Miller, 1981; Dickhaus, 2014)

$$\sup_{R \in \mathcal{R}_n} \mathbb{P}_{H_{R,n}} [\text{“any false rejection in } R\text{”}] \leq \alpha + o(1) \quad \text{as } n \rightarrow \infty. \quad (5)$$

To combine the local test statistics in a multiscale fashion, it is pertinent to employ a scale calibration  $\tilde{\omega}_n, \omega_n : (0, \infty) \rightarrow (0, \infty)$  to avoid small regions  $R \in \mathcal{R}_n$  from dominating the overall statistic. Here, and in what follows, the cardinality  $|R| \in \mathbb{N}$  of a region  $R \in \mathcal{R}_n$  is denoted as *scale*. This leads to

$$T_n \equiv T_n(Y, \mathcal{R}_n, \theta_0, \xi) := \max_{R \in \mathcal{R}_n} \tilde{\omega}_n(|R|) [T_R(Y, \theta_0, \xi) - \omega_n(|R|)], \quad (6)$$

which we then call a (calibrated) *multiscale test statistic* whenever a range of scales is covered by  $\mathcal{R}_n$ . An appropriate choice of the calibration  $\tilde{\omega}_n$  and  $\omega_n$  is of high importance for the overall method and will determine its detection properties. Different choices of  $\tilde{\omega}_n$  and  $\omega_n$  have been proposed in the literature, mostly to achieve a non-degenerate limiting behaviour of  $T_n$  or to optimize the detection power of the multiple test on all or specific scales. Besides uncalibrated scan statistics used e.g. in Kulldorff (1997); Alm (1997, 1998); Kabluchko and Munk (2008); Neill (2012); Madrid Padilla et al. (2022), we mention Dümbgen and Spokoiny (2001), who proposed a calibration ensuring non-degenerateness of the limiting distribution, and a more recent multivariate analogue by Datta and Sen (2021). Building on this, Dümbgen and Walther (2008) proposed a calibration, which was later shown to be superior over the uncalibrated scan statistic in Chan and Walther (2013), and which is since then widely used in the community, see e.g. Rivera and Walther (2013); Schmidt-Hieber et al. (2013); Frick et al. (2014); König et al. (2020). More recently, other calibrations have been suggested, e.g. leading to Gumbel limits (Sharpnack and Arias-Castro, 2016; Proksch et al., 2018), see also Walther (2020) for a recent review on this topic. In this study, we mostly leave the choice of  $\tilde{\omega}_n$  and  $\omega_n$  to the user up to some mild condition (cf. Assumption 3) on its growth posed below. This choice may e.g. reflect a preference for detecting signals of certain size, which is driven by the application at hand.

The multiscale scanning methodology is now as follows. Given  $Y$ , the values  $T_R(Y, \theta_0, \xi)$ ,  $R \in \mathcal{R}_n$  are computed, and all hypotheses  $H_{R,n}$  such that

$$T_R(Y, \theta_0, \xi) \geq c_{|R|}(\eta) := \frac{\eta}{\tilde{\omega}_n(|R|)} + \omega_n(|R|) \quad (7)$$

are rejected. Here,  $\eta \in \mathbb{R}$  is a global threshold to be determined appropriately. Hence, the scale calibration  $\tilde{\omega}_n, \omega_n$  leads to *scale dependent* local critical values in general, see also Walther (2010); Pein et al. (2017). The global threshold  $\eta$  is now universal, however, and to be chosen such that

$$\mathbb{P}_0 [T_n(Y, \mathcal{R}_n, \theta_0, \xi) > \eta] \leq \alpha + o(1) \quad \text{as } n \rightarrow \infty, \quad (8)$$

for some  $\alpha \in (0, 1)$ . Here and in what follows,  $\mathbb{P}_0$  denotes the law of  $Y$  under the global null hypothesis, i.e.  $\mathbb{P}_0 := \mathbb{P}_{H_{I_n^d}, n}$ . Since controlling the false positives over  $H_{I_n^d}$  implies control over any system of sub-hypotheses  $H_{R,n}$ ,  $R \in \mathcal{R}_n$ , it follows that

$$\sup_{R \in \mathcal{R}_n} \mathbb{P}_{H_{R,n}} [\text{“any false rejection in } R\text{”}] \leq \mathbb{P}_0 [\text{“any false rejection in } I_n^d\text{”}],$$

i.e. (8) ensures (5) to hold true, and hence any of the local rejections is (asymptotically) correct with probability  $\geq 1 - \alpha$ .

This principle to control the FWER underlies (often implicitly) many modern multiscale scanning methods. One central example is to scan an independent Gaussian field  $Y$  for anomalies (elevated means), where  $\theta_0 = \mu_0 = 0$  and  $\xi = \Sigma_0$  is the known covariance. In this case, optimality properties of this method, depending on the chosen calibration  $\tilde{\omega}_n$  and  $\omega_n$ , are known, see e.g. Chan and Walther (2013) for a statement about minimax optimality when using just a single scale, or Sharpnack and Arias-Castro (2016) and König et al. (2020) for an oracle property revealing that scanning over several scales is asymptotically as efficient as scanning over the correct scale only.

#### 1.4. Choosing the global threshold $\eta$

The remaining and potentially difficult question is how to choose the threshold  $\eta$  such that (8) is valid. In principle, this requires knowledge of the distribution of  $T_n$  under  $\mathbb{P}_0$ , which is not easy to obtain, in general. If the distributions  $F_{\theta,\xi} = F_{\mu,\sigma^2}$  are Gaussians with known nuisance parameter  $\xi = \sigma^2$  (variance), and if also the reference intensity  $\mu = \theta_0$  (mean) is known, then it readily turns out that all local test statistics  $T_R$  are constituted by Gaussians again, as

$$T_R(Y, \mu, \sigma) = |R|^{-1/2} \left| \sum_{i \in R} \frac{Y_i - \mu}{\sigma} \right|$$

in this case. Consequently,  $T_n$  can be simulated as the maximum over finitely many (dependent) Gaussian random variables. This way, it is possible to obtain empirical quantiles for  $T_n$  under  $H_0$  and to choose  $\eta$  accordingly.

For many other distributional settings such as Binomial or Poisson distributions, the situation is slightly more delicate. First steps have been taken by Alm (1997, 1998); Aboutkhamseen et al. (2016), who proposed distributional approximations of  $T_n$  in a Poissonian setting under knowledge of a baseline intensity  $\theta_0$ . However, those results are limited to one fixed scale. Another approach is based on tail bounds for  $T_R$  as proven in Walther (2022), which can be combined with a Bonferroni correction to ensure (8) also in a multiscale setting. Complete multiscale approximations of  $T_n$  in (6) have been proposed in case of Bernoulli and Binomial fields by Walther (2010) or in a more general exponential family setting by König et al. (2020) under knowledge of  $\theta_0$  and  $\xi$ . The latter result provides an invariance principle, which allows (given the baseline parameter  $\theta_0$  and the nuisance parameters  $\xi$ ) to approximate the distribution of  $T_n$  by the distribution in the Gaussian case as above, i.e.

$$M_n \equiv M_n(\mathcal{R}_n) := \max_{R \in \mathcal{R}_n} \tilde{\omega}_n(|R|) \left[ |R|^{-1/2} \left| \sum_{i \in R} X_i \right| - \omega_n(|R|) \right] \quad (9)$$

with i.i.d. standard normal r.v.'s  $X_i$ ,  $i \in I_n^d$ . Consequently, it is possible to simulate an appropriate threshold  $\eta$  in (8) in practice as soon as  $\theta_0$  and  $\xi$  are known. The rationale behind approximating  $T_n$  in (6) by  $M_n$  in (9) is, that for sufficiently large scales, a CLT implies that  $T_R \approx |R|^{-1/2} \left| \sum_{i \in R} X_i \right|$  in distribution. However, to ensure that  $M_n$  in fact approximates  $T_n$ , this CLT has to hold true uniformly over all sets in  $\mathcal{R}_n$ , i.e. it is required to couple the process (or at least its maximum)  $(T_R)_{R \in \mathcal{R}_n}$  with  $\left( |R|^{-1/2} \left| \sum_{i \in R} X_i \right| \right)_{R \in \mathcal{R}_n}$  (or the maximum, respectively) and to account for the calibration. Based on classical strong approximation results (Komlós et al., 1975), this approach has also been used in Frick et al. (2014), and more recently been extended in Proksch et al. (2018); König et al. (2020) by using explicit couplings for the suprema of empirical processes developed by Chernozhukov et al. (2014).

## 2. Proposed Method: adjusted multiscale scanning (AMS)

In contrast to existing work, the focus of this study is, however, on the case that both  $\theta_0$  and the nuisance parameter  $\xi$  in (1) are unknown, a typical situation in practice. Note, that this changes the overall situation completely, as the local LRT statistics  $T_R(Y, \theta_0, \xi)$  are no longer accessible. A common remedy which immediately comes to mind is to pre-estimate both  $\theta_0$  and  $\xi$  from  $Y$  globally and then use the corresponding estimators  $\hat{\theta}_0, \hat{\xi}$  as a plug-in in  $T_R$  and consequently in the multiscale extension  $T_n$  in (6). When one aims to *estimate*  $\theta_0$  only, then such an approach is known as pseudo maximum likelihood, and has e.g. been proposed in Gong and Samaniego (1981). Under certain regularity conditions, they prove asymptotic normality of the resulting estimator, but this result is limited to a single scale and cannot be transferred to the case of multiscale testing. A rigorous analysis of a similar approach in case of *testing* is – to the best of our knowledge – missing. In fact, we will show that the distribution of

$$\hat{T}_n := T_n(Y, \mathcal{R}, \hat{\mu}_0, \hat{\sigma}_0)$$

differs also asymptotically from that of  $T_n(Y, \mathcal{R}_n, \theta_0, \xi)$  (and hence also from that of  $M_n(\mathcal{R}_n)$  in (9)) in general, and this is the case even in the most simple Gaussian example. This then raises the question how to adjust the threshold  $\eta$  in (8) properly, an issue which has often been overlooked and has not been discussed in the literature. A notable exception is Walther and Perry (2022), where the tail bounds for  $T_R(Y, \hat{\theta}_0, \hat{\xi})$  in Walther (2022) are combined with a Bonferroni correction to construct corrected values for  $\eta$  in (8). This approach is, however, limited to relatively small systems  $\mathcal{R}_n$  of regions. In this paper, we will close the aforementioned gap and we will show that it is still possible to obtain a uniform invariance principle (see Section 3) for the distribution of  $\hat{T}_n$  as long as the scales in  $\mathcal{R}_n$  are not too small or too large compared to the sample size  $n^d$ . More precisely, under this restriction, we will show that  $\hat{T}_n$  can be approximated in distribution by  $M_n(\mathcal{R}_n)$  as in (9), which can be pre-simulated given  $\mathcal{R}_n$  in advance. From this it becomes apparent that working with thresholds, which do not incorporate the error in estimating  $\theta_0$  and  $\xi$ , does not provide the correct FWER, even asymptotically. The suggested adjusted multiscale scanning (AMS) procedure provides correct thresholds for estimated values of  $\theta_0$  and  $\xi$ .

Let us briefly compare the invariance principle in the paper at hand with previous results, especially those from König et al. (2020). There, an approximation of  $T_n$  in (6) by  $M_n$  in (9) has been proven in the case that  $F_{\theta, \xi}$  is a natural exponential family, both  $\theta_0$  and  $\xi$  are known, and the calibration functions  $\omega_n, \tilde{\omega}_n$  are explicitly chosen as in Dümbgen and Walther (2008), see also Example 3.5 below. In the paper at hand, we extend the above result by dropping all three restrictions. First of all, we allow for distributions  $F_{\theta, \xi}$  beyond the exponential family setting. We will elaborate in Example 3.3 below that e.g. the Gamma distribution can be handled by our theory but not by that from König et al. (2020). However, the major contribution of the paper at hand is the presence of nuisance parameters (including e.g. the reference intensity  $\theta_0$ ), which is of serious importance in practice and serves as the major motivation for our work. This will be illustrated by our data examples in Section 5. Finally, we allow for a wide choice of scale calibrations (see Assumption 3 below) and do not restrict ourselves to one specific choice.

### 2.1. Failure of naive plug-in in the Gaussian model

Let us start by explaining why simply replacing  $\theta_0$  and  $\xi$  by estimators  $\hat{\theta}_0$  and  $\hat{\xi}$  will lead to invalid thresholds  $\eta$  in (8). Therefore consider the homogeneous Gaussian model for  $d = 1$  (see e.g. Frick et al., 2014; Sharpnack and Arias-Castro, 2016) as the most common and best understood example of (1). Here,  $F_{\theta, \xi} = \mathcal{N}(\mu_i, \sigma_0^2)$  parametrized by its mean  $\mu_i \in \mathbb{R}$  and variance  $\sigma_0^2 > 0$ . In this case, it can readily be seen that replacing  $\mu_0$  and / or  $\sigma_0$  by estimators does actually influence the distribution of  $T_n$  and requires corrected critical values (thresholds)  $c_{|R|}(\eta)$  for the local LRT test statistics. To illustrate this, we have depicted the empirical distributions of  $T_n(Y, \mathcal{R}, \mu_0, \sigma_0) = M_n(\mathcal{R})$  as in (9), and a corresponding variant with standard estimators  $\hat{\mu}_0 = \frac{1}{n^d} \sum_{i \in I_n^d} Y_i$  and  $\hat{\sigma}_0^2 = \frac{1}{n^d - 1} \sum_{i \in I_n^d} (X_i - \hat{\mu}_0)^2$  for  $n = 128$  and  $\mathcal{R} = \mathcal{D}_n^d$  being the set of all rectangles in  $2^{I_n^d}$  in

Figure 1. As an exemplary scale calibration, we use the common choice  $\tilde{\omega}_n \equiv 1$  and  $\omega_n(|R|) = \sqrt{2 \log(n^d/|R|)} + 1$ , see Dümbgen and Walther (2008).




Figure 1: QQ (left) and PP-Plots (right) of the empirical distributions of  $M_n(\mathcal{R}) = T_n(Y, \mathcal{R}, 0, 1)$  and the naive approximation  $\hat{T}_n$  in the Gaussian case using all scales 1 – 128, i.e.  $\mathcal{R}_n = \mathcal{D}_n^d$  being the set of all rectangular regions, drawn from 2000 Monte Carlo runs.

It is clearly visible that the distribution of the true scan statistic  $T_n$  differs especially for large quantiles substantially from the one involving any of the estimators. If, e.g., the 80%-quantile of  $M_n$  is used to calibrate  $\hat{T}_n$ , then the corresponding FWER in (5) will be even  $\approx 30\%$ .

The reason for this can already be explained in the easier setting when the true variance  $\sigma_0^2$  is known, i.e. for  $T_n(Y, \mathcal{R}, \hat{\mu}_0, \sigma_0)$  with estimated global baseline  $\hat{\mu}_0$ . In this case, the log LRT statistic  $T_R(Y, \mu_0, \sigma_0)$  compares the local average of  $Y$  on  $R$  with  $\mu_0$ , i.e. in case of unknown  $\mu_0$  with the global average  $\hat{\mu}_0$  on  $I_n^d$ . Consequently, for large regions, the value of  $T_R(Y, \hat{\mu}_0, \sigma_0)$  will be negligible (in particular for  $R = I_n^d$  even constantly be 0), but  $T_R(Y, \mu_0, \sigma_0)$  is not (for  $R = I_n^d$ , this is proportional to a  $\chi_{n^d-1}^2$  distributed random variable). So, as long as the penalization  $\tilde{\omega}_n, \omega_n$  does not hinder these large scales from contributing to  $T_n(Y, \mathcal{R}, \hat{\mu}_0, \sigma_0)$  (and  $T_n(Y, \mathcal{R}, \mu_0, \sigma_0)$ ), the corresponding distributions must be different. On the other hand, a penalization hindering large scales from contributing to  $T_n$  is not desired, as this will decrease the overall detection power of the corresponding LRT tests on such scales. It is well known, see e.g. Dümbgen and Spokoiny (2001), that the penalization chosen here indeed ensures the largest scales to contribute to  $T_n(Y, \mathcal{R}, \mu_0, \sigma_0)$ . Hence, we find that in this case, also the asymptotic distribution (which for this scale penalization is a supremum of a Gaussian process, see Dümbgen and Spokoiny (2001)) should be different.

Note, that the above observation is not restricted to a Gaussian model only, it effectively occurs in any model where the nuisance parameter is estimated and naively plugged into the scan statistic. The largest scales will in general contribute to the distribution-free statistic  $M_n$  in (9), but will under mild assumptions on  $T_R$  (local means) not contribute to  $T_n$  in (6) as argued above.

## 2.2. Adjusted multiscale scanning – AMS

As a consequence, we have to carefully incorporate the influence of estimating global parameters in multiscale scanning in such a way that we allow for a large as possible range of scales. This is the starting point for the subsequent invariance principle in Section 3, which is at the heart of AMS. Afterwards, in Section 5, we illustrate a different scenario - a base signal, which is not known beforehand, has to be incorporated into the multiscale statistic  $T_n$  and hence needs to be estimated from the data.

Whereas in Section 2.1 the failure of the unadjusted multiscale statistic results from estimating a nuisance parameter, in the subsequent example (Section 5) we will face difficulties arising from estimation of the Poisson baseline  $\lambda_0$ . Both issues can be dealt with the main technical result of this paper, a *uniform* invariance principle for estimated base intensity and nuisance parameters.

In the following, we will briefly describe a simplified version of our main result, for the general result see Theorem 3.6 in Section 3. Suppose for simplicity that the estimators  $(\hat{\theta}_n)_{n \in \mathbb{N}}$  and  $(\hat{\xi}_n)_{n \in \mathbb{N}}$

are consistent with the parametric rate  $n^{-d/2}$  (as typically achieved by the MLE). Furthermore let  $\gamma > 0$  an explicitly known exponent depending on  $\omega_n$  and  $\tilde{\omega}_n$ , and let  $(r_n)_n \subset (0, \infty)$  and  $(m_n)_n \subset (0, \infty)$  be sequences such that

$$\log^\gamma(n) = o(r_n) \quad \text{and} \quad m_n = o(n^{-d/2}),$$

as  $n \rightarrow \infty$ . Then, if model (1) satisfies some mild differentiability properties (see Assumption 1 for details), the distribution of  $\hat{T}_n$  can be approximated by that of  $M_n$  in probability as long as  $\mathcal{R}_n$  contains only scales of size between  $r_n$  and  $m_n$ . In other words, we obtain a uniform invariance principle on all not too small (poly-logarithmic in the sample size) and all not too large scales just below the largest ones of order  $n^{-d/2}$ . In view of subsection 2.1, it is obvious that the latter condition is necessary to distinguish the baseline signal from local deviations, hence unavoidable in our context.

If, for a given system of candidate sets  $\mathcal{R}_n$ , we introduce the notation

$$\mathcal{R}_n(c, d) := \{R \in \mathcal{R}_n \mid c \leq |R| \leq d\},$$

then the above result can be formulated as

$$T_n \left( Y, \mathcal{R}_n(r_n, m_n), \hat{\theta}_n, \hat{\xi}_n \right) - M_n(\mathcal{R}_n(r_n, m_n)) \xrightarrow{\mathbb{P}} 0, \quad (10)$$

as  $n \rightarrow \infty$ . The rate of convergence of this approximation will be made precise in Theorem 3.6, which is (up to log factors)  $\mathcal{O}_{\mathbb{P}_0} \left( n^{-d/2} \frac{\sqrt{m_n}}{r_n} + r_n^{1/8} \right)$ .

Our result holds true for many different calibrations, including the previously mentioned one due to Dümbgen and Spokoiny (2001), which ensures that  $M_n$  converges to a non-degenerate continuous limit (and hence, due to (10), so does  $T_n$ ). From our result, similar results can be derived for the scale calibrations by Sharpnack and Arias-Castro (2016) and Proksch et al. (2018), which ensure that  $M_n$  converges to a Gumbel distribution.

With the above result, the AMS method is now carried out as depicted in Algorithm 1, which by construction ensures asymptotic control of the FWER, i.e. (5) holds true.

---

**Algorithm 1** Adjusted multiscale scanning - AMS

---

**Require:** Sample size  $n \in \mathbb{N}$ , dimension  $d \in \mathbb{N}$ , scale system  $\mathcal{R}_n$ , penalization  $\tilde{\omega}_n, \omega_n$ , scale bounds  $r_n < m_n$ , data  $Y$ , FWER level  $\alpha \in (0, 1)$

- 1: Simulate the distribution of  $M_n(\mathcal{R}_n(r_n, m_n))$
- 2: Compute the (empirical)  $(1 - \alpha)$ -quantile  $q_{1-\alpha}$  of  $M_n$ 
  - ▷ Steps 1 and 2 can be precomputed and the results can be stored independent of  $Y$
- 3: Set  $\mathcal{S} \leftarrow \emptyset$
- 4: Compute the estimators  $\hat{\theta}_0, \hat{\xi}$  based on  $Y$  for the baseline intensity  $\theta_0$  and the nuisance parameters  $\xi$
- 5: **for**  $R \in \mathcal{R}_n(r_n, m_n)$  **do**
- 6:     Compute  $T \leftarrow T_R(Y, \hat{\theta}_0, \hat{\xi})$
- 7:     Compute  $c \leftarrow c_{|R|}(q_{1-\alpha})$  as in (7)
- 8:     **if**  $T \geq c$  **then**
- 9:          $\mathcal{S} \leftarrow \mathcal{S} \cup \{R\}$
- 10:     **end if**
- 11: **end for**

**Ensure:**  $\mathcal{S} \subset \mathcal{R}_n$  such that with (asymptotic) probability  $\geq 1 - \alpha$  all  $R \in \mathcal{S}$  contain at least one anomaly

---

In Section 4.1 we will provide details on a computationally efficient implementation of this procedure, which can be carried out by means of fast Fourier transforms as soon as  $T_R$  in (3) is given by a function of the local mean  $\bar{Y}_R := \sum_{i \in R} Y_i$ . This leads to an algorithm for the evaluation of  $M_n$  in (9) and  $(T_R)_{R \in \mathcal{R}_n}$  in  $\mathcal{O}(dNn^d \log(n))$  operations, where  $N$  is the number of scales contained in  $\mathcal{R}_n$ . The corresponding MATLAB<sup>®</sup>-code is available under <https://go.uniwue.de/math-ams>.

### 3. Assumptions and detailed theory

In this section we will now discuss our assumptions and the result (10) in detail. Furthermore, we will examine several examples to demonstrate the validity of our assumptions in realistic scenarios.

To formulate our invariance principle, we will make use of stochastic  $\mathcal{O}$ -notation (see Sect. 2.2 in van der Vaart, 1998): For a sequence  $(X_n)_{n \in \mathbb{N}}$  of real-valued random variables we write  $X_n = \mathcal{O}_{\mathbb{P}}(1)$  iff  $(X_n)_{n \in \mathbb{N}}$  is tight, that is for all  $\varepsilon > 0$  there exists  $M > 0$  such that

$$\sup_{n \in \mathbb{N}} \mathbb{P}[|X_n| > M] < \varepsilon.$$

We write  $X_n = \mathcal{O}_{\mathbb{P}}(a_n)$  with a sequence  $(a_n)_{n \in \mathbb{N}}$  of real numbers, iff  $X_n/a_n = \mathcal{O}_{\mathbb{P}}(1)$ . Note, that in our context this means stochastic convergence for one fixed probability distribution  $\mathbb{P}$  only, and not uniform convergence e.g. over all  $\theta_0 \in \Theta$ . Furthermore, we write  $X_n = o_{\mathbb{P}}(1)$  if  $X_n$  converges stochastically to 0.

Recall, that we abbreviate  $\mathbb{P}_0 = \mathbb{P}_{H_{T_n^d, n}}$ . Effectively, this means that  $\mathbb{P}_0$  depends on  $\theta_0$  and  $\xi$ .

#### 3.1. Assumptions and examples

We start with our assumptions on the distributions  $F_{\theta, \xi}$  in model (1). For notional simplicity we introduce

$$m(\theta, \xi) := \mathbb{E}_{\theta, \xi}[Y] \quad \text{and} \quad v(\theta, \xi) := \mathbb{V}_{\theta, \xi}[Y]. \quad (11)$$

Here and in what follows,  $Y$  denotes a generic random variable with distribution  $Y \sim F_{\theta, \xi}$  according to the parameters mentioned in the subscript.

**Assumption 1** (Model assumptions). *Let  $Y \sim F_{\theta, \xi}$ ,  $(\theta, \xi) \in \Theta \times \Xi \subseteq \mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$  with  $d_1, d_2 \in \mathbb{N}$ . For  $\theta_0 \in \Theta, \xi \in \Xi$  assume that there exists a non-empty open neighbourhood  $U \times V \subset \Theta \times \Xi$  of  $(\theta_0, \xi)$ .*

(a) **Uniformly sub-exponential tails:** *There exist constants  $c_1 > 1, c_2 > 0$  such that*

$$\sup_{(\theta, \xi) \in U \times V} \mathbb{P}_{\theta, \xi}[|Y| > t] \leq c_1 \exp(-c_2 t) \quad \text{for all } t > 0.$$

(b) **Moments exist locally:** *The functions  $m$  and  $v$  in (11) exist and are finite on  $U \times V$  and there exists a constant  $c_v > 0$  such that  $v > c_v$  on  $U \times V$ .*

(c) **Bounded derivatives of  $m$ :** *There exists a constant  $C_m < \infty$  such that  $\|\nabla_{\theta} m\|_2, \|\nabla_{\xi} m\|_2 \leq C_m$  on  $U \times V$ .*

(d) **Bounded and non-vanishing derivatives of  $v$ :** *There exist constants  $0 < c_v < C_v < \infty$  such that  $c_v \leq \|\nabla_{\theta} v\|_2, \|\nabla_{\xi} v\|_2 \leq C_v$  on  $U \times V$ .*

(e) **Taylor approximation:** *There exists a constant  $C_T < \infty$  such that the likelihood ratio statistic  $T_R$  in (3) can be approximated by local means  $\bar{Y}_R = |R|^{-1} \sum_{i \in R} Y_i$  in the sense that*

$$\left| T_R^2(Y, \theta, \xi) - |R| \left( \frac{\bar{Y}_R - m(\theta, \xi)}{\sqrt{v(\theta, \xi)}} \right)^2 \right| \leq C_T |R| \left( \frac{|\bar{Y}_R - m(\theta, \xi)|}{\sqrt{v(\theta, \xi)}} \right)^3 (1 + o_{\mathbb{P}_0}(1)) \quad (12)$$

holds true for all  $R \in \mathcal{R}$  and  $(\theta, \xi) \in U \times V$ .

We briefly comment on these assumptions.

**Remark 3.1.** a) *Assumption 1(a) allows us to control moments of  $Y_i$  and hence of local means  $\bar{Y}_R := \sum_{i \in R} Y_i$  simultaneously over regions  $R \in \mathcal{R}_n$ .*



- b) Assumption 1(b) seems natural, as otherwise no CLT on the small scales can hold true and hence no distributional limit is to be expected.
- c) Assumptions 1(c) and (d) will be used to control the impact caused by replacing  $\theta_0$  and  $\xi$  in (6) by the estimators  $\hat{\theta}_n$  and  $\hat{\xi}_n$ , respectively.
- d) Assumption 1(e) is required to link  $T_R$  to the local means  $\bar{Y}_R$  sufficiently well.

For a better understanding of Assumption 1 let us discuss some examples:

**Example 3.2** (Complete exponential family model). *Suppose that*

$$\{F_{\theta,\xi}\}_{(\theta,\xi)\in\Theta\times\Xi}$$

is a natural exponential family with natural parameter space  $\Theta \times \Xi$ . In this case, each  $Y_i$  has sub-exponential tails (which implies Assumption 1(a)) and the differentiability properties from Assumption 1(c)–(e) hold true (see e.g. Brown, 1986).

This includes both the homogeneous and the heterogeneous Gaussian model as well as the Poisson model, which we employ for our data example in Section 5 on fluorescence microscopy.

**Example 3.3** (Partial exponential family model). *Suppose that for each fixed value of  $\xi \in \Xi$ , the collection  $\{F_{\theta,\xi}\}_{\theta \in \Theta}$  is a natural exponential family. In this case, Assumption 1(a) is again satisfied due to standard theory on exponential families, but the differentiability properties in Assumption 1(c)–(e) have to be verified case by case. As one particular example we consider the Gamma distribution  $\Gamma(\alpha, \beta)$  with shape parameter  $\alpha > 0$ , rate  $\beta > 0$ , and density*

$$f(x) = \frac{1}{\Gamma(\alpha)} \beta^\alpha x^{\alpha-1} \exp(-\beta x).$$

If the shape parameter  $\alpha$  is a nuisance parameter, we obtain  $m(\beta, \alpha) = \alpha/\beta$ ,  $v(\beta, \alpha) = \alpha/\beta^2$  and

$$\frac{\partial}{\partial \beta} m(\beta, \alpha) = -\frac{\alpha}{\beta^2}, \quad \frac{\partial}{\partial \beta} v(\beta, \alpha) = -\frac{2\alpha}{\beta^3}, \quad \frac{\partial}{\partial \alpha} m(\beta, \alpha) = \frac{1}{\beta}, \quad \frac{\partial}{\partial \alpha} v(\beta, \alpha) = \frac{1}{\beta^2},$$

which proves the boundedness conditions in Assumption 1(c) and (d). Furthermore, (12) follows directly from the exponential family structure (see e.g. König et al., 2020).

We also give a counterexample to show the limitations of our Assumptions:

**Example 3.4** (Weibull distribution). *Consider the Weibull distribution with shape parameter  $k > 0$  and rate parameter  $\lambda > 0$ . If  $k \geq 1$ , then all random variables  $Y_i$  will have sub-exponential tails (see e.g. Smith, 1985; Farnum and Booth, 1997), but Assumption 1(e) is not satisfied. Therefore, we compute*

$$\begin{aligned} \log \left( \frac{\sup_{\theta \in \Theta} \prod_{i \in R} f_{(\theta,\xi)}(Y_i)}{\prod_{i \in R} f_{(\theta_0,\xi)}(Y_i)} \right) &= k \log(\lambda_0) - \log \left( \sum_{i \in R} Y_i^k \right) + \frac{\sum_{i \in R} Y_i^k}{\lambda_0^k} - 1 \\ |R| \left( \frac{\bar{Y}_R - m(\theta_0, \xi)}{\sqrt{v(\theta_0, \xi)}} \right)^2 &= |R| \left( \frac{\bar{Y}_R^2 - 2\lambda \Gamma \left(1 + \frac{1}{k}\right) \bar{Y}_R + \lambda^2 \left(\Gamma \left(1 + \frac{1}{k}\right)\right)^2}{\lambda^2 \left[\Gamma \left(1 + \frac{2}{k}\right) - \left(\Gamma \left(1 + \frac{1}{k}\right)\right)^2\right]} \right)^2, \end{aligned}$$

which shows that (12) cannot hold true.

Next we continue with our assumptions on the multiscale scanning procedure described in terms of  $T_n$  in (6). To approximate  $T_n$  by  $M_n$  in (9), we first have to restrict the cardinality of  $\mathcal{R}_n$ .

**Assumption 2** (Polynomial growth of  $\mathcal{R}_n$ ). *There exists constants  $c_1, c_2 > 0$  such that*

$$\#(\mathcal{R}_n) \leq c_1 n^{c_2}. \tag{13}$$

where  $\#$  denotes the number of elements.

As discussed in König et al. (2020), Assumption 2 is rather mild and allows e.g. for  $\mathcal{R}_n$  being the set of all hyper-rectangles, hyper-cubes or half-spaces in  $[0, 1]^d$ .

Our next assumption is on the scale calibration  $\tilde{\omega}_n, \omega_n$  in (6):

**Assumption 3.** *Suppose  $\omega_n, \tilde{\omega}_n : (0, \infty) \rightarrow (0, \infty)$  have the following properties:*

1. *For each  $n \in \mathbb{N}$ ,  $\omega_n$  and  $\tilde{\omega}_n$  are decreasing.*
2. *There exist  $C_\omega > 0$ ,  $\alpha, \tilde{\alpha} > 0$  and  $\beta, \tilde{\beta} \in \mathbb{R}$  such that*

$$\begin{aligned}\omega_n(r) &\leq C_\omega \left( \log \frac{n^d}{r} \right)^\alpha, & |\omega_n'(r)| &\leq C_\omega \left( \log \frac{n^d}{r} \right)^\beta \frac{1}{r}, \\ \tilde{\omega}_n(r) &\leq C_\omega \left( \log \frac{n^d}{r} \right)^{\tilde{\alpha}}, & |\tilde{\omega}_n'(r)| &\leq C_\omega \left( \log \frac{n^d}{r} \right)^{\tilde{\beta}} \frac{1}{r}\end{aligned}$$

for all  $r > 0$  and  $n \in \mathbb{N}$ .

Let us briefly discuss this assumption for common choices considered in the literature:

**Example 3.5.** *a) Dümbgen and Walther (2008); Chan and Walther (2013); Rivera and Walther (2013) consider*

$$\tilde{\omega}_n \equiv 1, \quad \omega_n(|R|) = \sqrt{2\nu(\log(n^d/|R|)) + 1}, \quad (14)$$

where  $\nu \geq 1$  depends on the complexity of the candidate region. These terms fulfil Assumption 3 with  $\alpha = \frac{1}{2}, \beta = -\frac{1}{2}$  and  $\tilde{\alpha} = \tilde{\beta} = 0$ .

*b) Sharpnack and Arias-Castro (2016) consider*

$$\begin{aligned}\tilde{\omega}_n(|R|) &= \sqrt{2(\log(n^d/|R|))}, \\ \omega_n(|R|) &= \sqrt{2(\log(n^d/|R|))} + \frac{(4d-1) \log\left(\sqrt{2(\log(n^d/|R|))}\right) - \log(\sqrt{2\pi})}{\sqrt{2(\log(n^d/|R|))}},\end{aligned} \quad (15)$$

and therefore Assumption 3 is fulfilled with  $\alpha = \tilde{\alpha} = \frac{1}{2}$  and  $\beta = \tilde{\beta} = -\frac{1}{2}$ .

*c) Proksch et al. (2018) consider*

$$\omega_n(|R|) := \tilde{\omega}_n(|R|) := \sqrt{2 \log\left(\frac{Cn^d}{|R|}\right)} + C_d \frac{\log\left(\sqrt{2 \log\left(\frac{Cn^d}{|R|}\right)}\right)}{\sqrt{2 \log\left(\frac{Cn^d}{|R|}\right)}}, \quad (16)$$

with  $C > 1$  and  $C_d$  depending on the dimension and the system of considered scales. These fulfil Assumption 3 with  $\alpha = \tilde{\alpha} = \frac{1}{2}$  and  $\beta = \tilde{\beta} = -\frac{1}{2}$ .

## 3.2. Main result and implications

Now we are in position to state our uniform invariance principle in detail. Therefore let  $(\hat{\theta}_n)_{n \in \mathbb{N}}$  and  $(\hat{\xi})_{n \in \mathbb{N}}$  be sequences of estimators for  $\theta_0$  and  $\xi$  based on  $Y$  such that they are  $s_n$  (e.g.  $s_n = n^{-d/2}$ ) consistent under the null hypothesis, i.e.

$$\|\hat{\theta}_n - \theta_0\|_2 = \mathcal{O}_{\mathbb{P}_0}(s_n^{-1}) \quad \text{and} \quad \|\hat{\xi}_n - \xi\|_2 = \mathcal{O}_{\mathbb{P}_0}(s_n^{-1}), \quad (17)$$

where  $(s_n)_{n \in \mathbb{N}} \subset \mathbb{R}$  is a monotone sequence tending to  $\infty$ . Note, that the MLEs for  $\theta_0$  and  $\xi$  satisfy (17) under Assumption 1 with  $s_n = n^{d/2}$ . Furthermore let

$$\gamma = 12 + 6\tilde{\alpha} + 2 \max\left\{\frac{1}{2}, \alpha, \tilde{\alpha}\right\} + 2 \max\left\{\beta, \tilde{\beta}, 0\right\}$$

with the parameters  $\alpha, \tilde{\alpha}, \beta, \tilde{\beta}$  from Assumption 3 and suppose that

$$\log^\gamma(n) = o(r_n) \text{ as } n \rightarrow \infty, \quad m_n = o(s_n) \text{ as } n \rightarrow \infty. \quad (18)$$

In this situation we can prove the following Gaussian approximation.

**Theorem 3.6** (Gaussian approximation). *Let  $\mathcal{R}_n$  be a set of candidate regions and let  $\omega_n, \tilde{\omega}_n$  be scale calibrations. Grant Assumptions 1, 2 and 3. Furthermore let  $(r_n)_n \subset (0, \infty)$  and  $(m_n)_n \subset (0, \infty)$  be sequences such that (18) holds true and let  $(\hat{\xi}_n)_{n \in \mathbb{N}}$  and  $(\hat{\theta}_n)_{n \in \mathbb{N}}$  are sequences of estimators such that (17) is valid.*

*Then, for a field  $Y = (Y_i)_{i \in I_n^d}$  with i.i.d. random variables  $Y_i \sim F_{\theta_0, \xi}$ , it holds*

$$\left| T_n \left( Y, \mathcal{R}_n(r_n, m_n), \hat{\theta}, \hat{\xi} \right) - M_n(\mathcal{R}_n(r_n, m_n)) \right| = \mathcal{O}_{\mathbb{P}_0} \left( \frac{\log^{\tilde{\alpha}}(n)}{r_n} \sqrt{\frac{m_n}{s_n}} + \left( \frac{\log^\gamma(n)}{r_n} \right)^{\frac{1}{8}} \right), \quad (19)$$

as  $n \rightarrow \infty$ .

**Corollary 3.7.** *Let the assumptions of Theorem 3.6 hold true,  $Y = (Y_i)_{i \in I_n^d}$  as in (1) (independent, but not necessarily identically distributed), and  $\alpha \in [0, 1]$ . If we define*

$$\mathcal{R}_n^* := \left\{ R \in \mathcal{R}_n \mid r_m \leq |R| \leq m_n, T_R \left( Y, \hat{\theta}_n, \hat{\xi}_n \right) \geq \frac{q_{1-\alpha}}{\tilde{\omega}_n(|R|)} + \omega_n(|R|) \right\}$$

with the  $(1 - \alpha)$ -quantile  $q_{1-\alpha}$  of  $M_n$  as in (9), then (5) holds true, i.e.

$$\mathbb{P}[\forall R \in \mathcal{R}_n^* \exists i \in R \text{ s.t. } \theta_i \neq \theta_0] \geq 1 - \alpha + o(1) \quad \text{as } n \rightarrow \infty.$$

**Remark 3.8.** *In the situation that all local alternatives (and tests) are one-sided, i.e.  $\theta_i > \theta_0$ , the above result can readily be generalized by setting*

$$\tilde{T}_R \left( Y, \hat{\theta}_n, \hat{\xi}_n \right) = \begin{cases} T_R \left( Y, \hat{\theta}_n, \hat{\xi}_n \right) & \text{if } \bar{Y}_R > \hat{\theta}_n, \\ 0 & \text{else,} \end{cases}$$

as well as

$$\tilde{T}_n \left( Y, \mathcal{R}_n, \hat{\theta}_n, \hat{\xi}_n \right) := \max_{R \in \mathcal{R}_n} \tilde{\omega}_n(|R|) \left[ \tilde{T}_R \left( Y, \theta_0, \xi \right) - \omega_n(|R|) \right]$$

and

$$\tilde{M}_n(\mathcal{R}_n) := \max_{R \in \mathcal{R}_n, \bar{X}_R > 0} \tilde{\omega}_n(|R|) \left[ |R|^{-1/2} \left| \sum_{i \in R} X_i \right| - \omega_n(|R|) \right].$$

With this notation, it can readily be seen that Theorem 3.6 holds also true for those quantities, i.e. under the same conditions we obtain

$$\left| \tilde{T}_n \left( Y, \mathcal{R}_n(r_n, m_n), \hat{\theta}, \hat{\xi} \right) - \tilde{M}_n(\mathcal{R}_n(r_n, m_n)) \right| = \mathcal{O}_{\mathbb{P}_0} \left( \frac{\log^{\tilde{\alpha}}(n)}{r_n} \sqrt{\frac{m_n}{s_n}} + \left( \frac{\log^\gamma(n)}{r_n} \right)^{\frac{1}{8}} \right)$$

as  $n \rightarrow \infty$ .

**Remark 3.9.** *As a different approach to the invariance principle in Theorem 3.6, it would be particularly elegant to exploit asymptotic equivalence in the Le Cam sense (see e.g. Grama and Nussbaum, 1998; Ray and Schmidt-Hieber, 2018) after a variance stabilizing transformation to a Gaussian model. However, to the best of our knowledge, such results will only be valid for a fixed number of scales. Theorem 3.6 therefore gives rise to interesting questions on how to extend asymptotic equivalence to multiscale frameworks with growing number of tests.*

## 4. Implementation and simulations

In this section, we will first outline details on our implementation and afterwards investigate the behaviour of the adaptive multiscale test in a simulation study.

### 4.1. Implementation

As already mentioned in the introduction, for a generic set  $\mathcal{R}_n$  all local test statistics  $T_R$  in (3) have to be computed individually. To this end, for example, the approximating rectangular system from Walther (2010) can be used to compute local averages on these. Depending on the structure of  $\mathcal{R}_n$ , other efficient computational schemes can be employed. In the following we will focus on the situation that there is a global shape  $B \subset I_n^d$  such that every  $R \in \mathcal{R}_n$  is a rescaled and shifted version of  $B$ . More precisely, for each  $R \in \mathcal{R}_n$  there exist  $t, h \in I_n^d$  with  $t_i + h_i \leq n$  for all  $1 \leq i \leq d$  such that

$$\mathbf{1}_R(x) = \mathbf{1}_B\left(\frac{x-t}{h}\right), \quad x \in I_n^d, \quad (20)$$

where division is meant component-wise and  $\mathbf{1}$  denotes the indicator function. For the set  $\mathcal{R}_n$  of all hyper-rectangles, this is for example the case with  $B = \mathbf{1}_{I_n^d}$ .

If  $R$  obeys (20), we obtain

$$\bar{Y}_R = \frac{1}{|R|} \sum_{i \in R} Y_i = \frac{1}{|R|} \sum_{i \in I_n^d} Y_i \mathbf{1}_B\left(\frac{i-t}{h}\right) = \frac{1}{|R|} \left( Y * \mathbf{1}_B\left(\frac{\cdot}{h}\right) \right)(t),$$

where  $*$  denotes a discrete convolution. Employing the fast Fourier transform (FFT), see Cooley and Tukey (1965), this allows to compute all  $T_R$  with a fixed scale  $h$  by means of three FFTs via

$$(\bar{Y}_R)_h = \frac{1}{h^d |B|} \text{FFT}^{-1} \left( \text{FFT}(Y) \cdot \text{FFT} \left( \mathbf{1} \left( \frac{\cdot}{h} \right) \right) \right).$$

This operation has a computational complexity almost linear in the data, i.e.  $\mathcal{O}(dn^d \log(n))$ , and hence if each  $R \in \mathcal{R}_n$  obeys (20), then  $T_n$  in (6) can be computed within  $\mathcal{O}(dNn^d \log(n))$  where  $N$  is the number of different scales. The same holds true for the evaluation of  $M_n$  in (9). A corresponding implementation in MATLAB<sup>®</sup> is available under <https://go.uniue.de/math-ams>.

### 4.2. Validity of the Gaussian approximation of AMS

We illustrate (10) in homogeneous the Gaussian setting from Subsection 2.1. Figure 2 shows the empirical distribution of  $T_n \left( Y, \mathcal{R}_n(r_n, m_n), \hat{\theta}_n, \hat{\xi}_n \right)$  and  $M_n(\mathcal{R}_n(r_n, m_n))$  with the same  $n$  and  $\tilde{\omega}_n, \omega_n$  as there for  $r_{128} = 4$  and  $m_{128} = 64$ .

We find the distributional fit to be remarkably accurate in all situations, which supports the applicability of the asymptotic result from Theorem 3.6. The distribution of  $M_n$  itself will be shown in Figure 3 for the situation considered in our simulations.

### 4.3. Simulations of level and power

In this section, we will study the finite sample properties of the adjusted multiscale test. We therefore consider two different model situations, namely a heterogeneous Gaussian model and a Poisson model. In both models, we investigate the empirical level and power of the procedure by means of Monte Carlo simulations with  $M = 1000$  runs. As a default setup, we use  $d = 2$ ,  $n = 128$  and  $\mathcal{R}_n$  to be the set of all rectangles in  $I_n^d$  with even side-lengths between 4 and 14 pixels. As penalization functions  $\tilde{\omega}_n$  and  $\omega_n$ , we choose the previously mentioned penalization (14) of Dümbgen and Walther (2008). Empirical quantiles of  $M_n$  in (9) for this situation are




Figure 2: QQ (left) and PP-Plots (right) of the empirical distributions of  $M_n(\mathcal{R}) = T_n(Y, \mathcal{R}, 0, 1)$  and the naive approximation  $\hat{T}_n$  in the Gaussian case under scale restrictions with  $r_{128} = 4$  and  $m_{128} = 64$ , drawn from 2000 Monte Carlo runs. We emphasize that the only difference to Figure 1 is the restriction to scales of sizes 4 – 64.

Table 1: Empirical quantiles  $q_{1-\alpha}$  with different values of  $\alpha$  for the distribution of  $M_n$  in (9) in the case  $n = 128, d = 2$  and  $\mathcal{R}_n$  being the set of all rectangles in  $I_n^d$  with side-lengths between 4 and 14.

$\alpha$	0.2	0.1	0.05	0.025	0.01
$q_{1-\alpha}$	1.2906	1.4677	1.6278	1.7841	1.9768




Figure 3: Empirical density (left) and CDF (right) of  $M_n$  in (9) for  $n = 128$  in  $d = 2$  dimensions with  $\mathcal{R}_n$  being the set of all rectangles in  $I_n^d$  with even side-lengths between 4 and 14 pixels, displayed by a standard kernel density estimator, drawn from 2000 Monte Carlo runs, either calibrated (—) with penalization functions  $\tilde{\omega}_n$  and  $\omega_n$  as in (14), or uncalibrated (- - -).

shown in Table 1, and the empirical density and CDF both for the calibrated and the uncalibrated statistic are displayed in Figure 3.

For power simulations, we consider a  $a^2 = 8 \times 8$  sized anomaly positioned in the centre of the  $128 \times 128$  image. For comparison, we also investigate the corresponding oracle procedure, which arises from evaluating  $T_n(Y, \mathcal{R}_n, \theta_0, \xi)$  with the true values of  $\theta_0$  and  $\xi$ .

To investigate the influence of  $d$  and  $n$ , we additionally perform simulations with  $d = 1, 2$  and  $n = 32, 64, 96, 128$ . Therein we determine the necessary signal strength (mean) of an anomaly of size  $a^d$  ( $a \in \{4, 5, \dots, 13, 14\}$  a fixed number), again positioned in the centre of the domain, such that the empirical power is  $\approx 90\%$ . The corresponding functions  $\theta = \theta(a)$ , where  $\theta$  is either the mean  $\mu$  in the heterogeneous Gaussian model or the parameter  $\lambda$  in the Poisson model, are in fact an upper bound for the (finite sample) separation rate between  $H_0$  and the prescribed alternative. The (asymptotic) separation rate, i.e. the asymptotic detection boundary, provides a sharp notion of the minimal signal strength necessary for detection, which cannot be overcome by any test. Note, that the asymptotic detection boundary for the identification of cuboid signals

with known size  $a$  in Gaussian white noise as considered here has been derived in the literature, see e.g. Chan and Walther (2013); Enikeeva et al. (2018); Kou (2023), as, translated to our notation,

$$\mu \sim \sqrt{\frac{2}{a^d} \log \left( \frac{n^d}{a^d} \right)}. \quad (21)$$

We emphasize that AMS treats a more difficult situation (i.e.  $\theta_0, \xi$  and  $a$  unknown), and will hence not be able to reach this boundary, but as we will show below a certain comparison is still possible.

In the simulations of level and power, we always compare the AMS approach with an oracle approach, where the local LRT statistics  $T_R$  from (3) are evaluated using the true parameters  $\theta_0$  and  $\xi$ .

### 4.3.1. Homogeneous Gaussian model

Let us first consider the case already discussed in the introduction that  $F_{\theta, \xi}$  is a normal distribution  $\mathcal{N}(\mu, \sigma^2)$ , and the variance  $\sigma^2$  is considered as the nuisance parameter and estimated by the sample variance

$$\hat{\sigma}_n^2 := \frac{1}{n^d - 1} \sum_{i \in I_n^d} (Y_i - \bar{Y}_{I_n^d})^2.$$

Then (17) is clearly satisfied under  $\mathbb{P}_0$  with  $s_n = n^{-d/2}$ .

The results of the simulations as described before are depicted in Figure 4. Sub-panels (a) and (b) display the signal and the corresponding data for the default setup ( $d = 2$ ,  $n = 128$ ) with an anomaly of size  $8 \times 8$  pixels and amplitude  $\mu = 0.5$  in the centre, where the variance  $\sigma^2 = 1$ . Sub-panel (c) shows the empirical level for different variances  $\sigma^2$ . We find that the level is kept quite stable over a large range of variances. Sub-panel (d) shows the empirical power for  $\sigma^2 = 1$  against different means  $\mu$ . Notably, there is nearly no power loss of the AMS method compared to the oracle caused by adaptation for an unknown  $\sigma^2$ . Sub-panel (e) is devoted to simulations in dimension  $d = 1$  with  $\sigma^2 = 1$  and different values of  $n \in \{32, 64, 96, 128\}$ . The lines depict that value of  $\mu$ , such that the AMS procedure is able to detect a present anomaly of size  $a$  marked on the  $x$ -axis with amplitude  $\mu$  with a power of  $\approx 90\%$ . We find that a larger anomaly clearly corresponds to a smaller amplitude required for detection. This is in agreement with the asymptotic separation rate in (21), and remarkably, despite the presence of nuisance parameters and the additional difficulty that the length  $a$  is not known, the finite sample shapes of all separation lines in sub-panel (e) are of this form. Furthermore, we find that the influence of the sample size  $n$  on the separation lines is small. Sub-panel (f) shows the same situation in dimension  $d = 2$ . Due to the larger anomaly, which now contains  $a^2$  pixels, the corresponding values for  $\mu$  are smaller. However, the conclusions drawn for the one-dimensional case apply here similarly. Compared with sub-panel (e) we furthermore conclude that the dimension  $d$  does not have a big influence.

### 4.3.2. Poisson model

As a second setup, we consider a Poisson model where  $F_{\theta, \xi}$  is a Poisson distribution  $\text{Poi}(\lambda)$  with unknown background intensity  $\lambda_0$ , which will be estimated by the sample mean, i.e.  $\hat{\lambda}_n := \bar{Y}_{I_n^d}$ . Then (17) is clearly satisfied with  $s_n = n^{-d/2}$ .

The results of the simulations as described before are depicted in Figure 5. Sub-panels (a) and (b) show the signal and the corresponding data for the default setup ( $d = 2$ ,  $n = 128$ ) with an anomaly of size  $8 \times 8$  pixels and amplitude  $\lambda = 1.6$  in the centre, where the background intensity  $\lambda_0 = 1$ . Sub-panel (c) shows the empirical level for different background intensities  $\lambda_0$ . We again find that the level is kept quite stable, even for considerably small values of  $\lambda_0$ . The little bump around  $\lambda_0 = 0.3$  emphasizes that the quantiles are chosen according to the Gaussian approximation in Theorem 3.6. Sub-panel (d) shows the empirical power for  $\lambda_0 = 1$  against different amplitudes




Figure 4: Simulation results in a homogeneous Gaussian model with  $\alpha = 0.1$ . Panels (a) and (b): Signal under the alternative ( $\mu = 0.5$ ) used for this simulation and corresponding data for  $\sigma^2 = 1$ . As the signal is standardized, the standard deviation  $\sigma$  is proportional to the signal-to-noise ratio (SNR). Panel (c): Empirical level for the oracle multiscale scan ( $\sigma^2$  known) and AMS ( $\sigma^2$  unknown, i.e. considered as a nuisance parameter) for different values of  $\sigma$ , smoothed with a moving average filter of width 10. Panel (d): Empirical power versus  $\mu$  for a  $8 \times 8$  anomaly as in (a), smoothed with a moving average filter of width 10. Panels (e) and (f): Value of  $\mu$  required to obtain an empirical power of  $\beta = 0.9$  versus size of a (squared) anomaly with amplitude  $\mu$ .

$\lambda \geq \lambda_0$ . Similarly to the Gaussian case, there is nearly no power loss caused by adaptation for an unknown  $\lambda_0$ . Sub-panel (e) is devoted to simulations in dimension  $d = 1$  with  $\lambda_0 = 1$  and different values of  $n \in \{32, 64, 96, 128\}$ . The lines depict that value of  $\lambda$ , such that the AMS procedure is able to detect a present anomaly of size  $a$  marked on the  $x$ -axis with amplitude  $\lambda$  with a power of  $\approx 90\%$ . We find that a larger anomaly clearly corresponds to a smaller amplitude required for detection, again in agreement with the results on separation rates in the Gaussian case. Furthermore, we find that the influence of the sample size  $n$  on the separation lines is small, despite for the smallest sample size  $n = 32$  where the required value of  $\lambda$  is moderately larger. This might be caused by the worse approximation of  $T_n$  by  $M_n$  in this case. Sub-panel (f) shows the same situation in dimension  $d = 2$ . Due to the larger anomaly, which now contains  $a^2$  pixels, the corresponding values for  $\mu$  are smaller. The conclusions drawn for the one-dimensional case apply here for all considered values of  $n$ . Compared with sub-panel (e) we furthermore conclude that the dimension  $d$  does not have a big influence.

## 5. Data Analysis

A major motivation for our work comes from super-resolution fluorescence microscopy imaging. Fluorescence microscopy is employed to visualize structures of interest (e.g. proteins or protein complexes) in a sample. Therefore, the structures of interest in a sample are labelled biochemically by fluorescent markers, and then scanned spatially along a grid with a diffraction-limited laser spot centred at the current grid point. Whenever a marker is hit by the incoming light it excites with a certain probability, and if so, afterwards light of a different wavelength is emitted and recorded by detectors (cf. Aspelmeier et al., 2015, for details). The later excitation-emission-recording




Figure 5: Simulation results in a Poisson model with  $\alpha = 0.1$ . Panels (a) and (b): Signal under the alternative ( $\lambda_0 = 1$ ,  $\lambda = 1.6$ ) used for this simulation and corresponding data. Panel (c): Empirical level for the oracle multiscale scan ( $\lambda_0$  known) and AMS ( $\lambda_0$  unknown, i.e. considered as a nuisance parameter) for different values of  $\lambda_0$ , smoothed with a moving average filter of width 10. Panel (d): Empirical power versus  $\lambda$  for a  $8 \times 8$  anomaly as in (a), smoothed with a moving average filter of width 10. Panels (e) and (f): Value of  $\lambda$  required to obtain an empirical power of  $\beta = 0.9$  versus size of a (squared) anomaly with amplitude  $\lambda$ .

procedure is repeated several (say  $t \in \mathbb{N}$ ) times, called the measurement time, before the focus is moved to the next grid point.

To obtain a resolution below the Abbe diffraction limit (see Kulaitis et al. (2021) for an explanation in a statistical context), nowadays different techniques are known. Here we will focus on stimulated emission depletion (STED) microscopy, developed by Stefan Hell (see e.g. Hell and Wichmann (1994); Hell (2007) for details), which has been awarded the Nobel price for Chemistry in 2014. It is based on an additional donut-shaped depletion beam, which suppresses fluorescence in the corresponding region and hence reduces effectively the area in which light is emitted. This way, the effective resolution can be decreased dramatically from roughly 200 – 300nm for a standard (confocal) microscope to  $\leq 50$ nm, say.

## 5.1. Mathematical setup

Mathematically, the nature of photon counting in super-resolution microscopy leads to a Poisson field of independent observations (see e.g. Munk et al., 2020). Consequently, we will model the corresponding observations as an independent  $d$ -dimensional (usually  $d = 2$ ) field  $Y$  of Poisson random variables

$$Y_i \sim \text{Poi}(t\lambda_i), \quad i \in I_n^d := \{1, \dots, n\}^d \quad (22)$$

with parameters  $\lambda_i > 0$ ,  $i \in I_n^d$ .

However,  $t$  cannot be chosen arbitrary, as each marker is only able to pass through the cycle of excitation and emission a limited number of times before it bleaches. To avoid masking effects by bleaching, the total illumination time per pixel (called pixel dwell time) is hence set to a (not too large) number  $T > 0$ . It is immediately clear, that a larger value of  $T$  will allow for a more



accurate imaging of the underlying structures of interest. However, even if large areas in the image are mostly empty, these regions have to be scanned and by doing so all markers in the image are hit by light and hence either excited or at least stressed. Therefore, methods which are able to localize interesting structures within a short pre-scan (i.e. with short measurement time  $t \ll T$ ) have achieved great interest recently in many applications and can – in the present context – be seen as an instance of *smart* microscopy, see Strack (2020). Some heuristic methods have been suggested (such as RESCue and others, see Vinçon et al., 2020), which stop measuring at a pixel as soon as a predefined number of photons has been detected, but without any statistical guarantees. Such approaches for a real-time spatial control of the illumination are known as spatially-controlled illumination microscopy (SCIM), see e.g. the review by Krishnaswami et al. (2016). In principle this can be interpreted as scanning each grid point with a substantially smaller pixel dwell time  $t \ll T$  and deciding upon the corresponding measurement  $Y_i$  if the full pixel dwell time  $T$  is applied or if the measurement at the current grid point is stopped. Another approach exploits machine learning to pre-select regions of interest based on shortly measured (i.e. very noisy) data, see e.g. Pandey et al. (2019); Mahecic et al. (2022). All the previously described methods suffer from the fact that they are not able to provide statistical guarantees for correctness of the selected (or more intensively scanned) regions. As we will demonstrate now, AMS does provide a methodology that allows to identify interesting regions with prescribed statistical error.

## 5.2. Used data sets

The data sets to be shown below have been measured in the Institut für Nanophotonik (IFNANO) by René Siegmund especially to apply AMS with different initial scan times  $t$ . The measurement device is a STED super-resolution microscope with an effective resolution of  $\sim 60\text{nm}$ . For each  $t \in \{1\mu\text{s}, 2\mu\text{s}, \dots, 100\mu\text{s}\}$  a data set  $Y_t$  with corresponding dwell time  $t$  (in microseconds  $\mu\text{s}$ ) is recorded. The corresponding value  $t = 100\mu\text{s}$  can be seen as the maximal value  $T$ , which was chosen to avoid bleaching of the fluorescent dyes. In that sense, we will consider  $t = T$  as *ground truth*, since no better measurement is available or can realistically be measured. We aim to achieve a localization of all or most regions of interest by means of AMS based on data with  $t \ll T$ , since this then yields a statistically rigorous method to obtain the spatially-controlled illumination described above with (asymptotically) controlled FWER.

The available raw data image is corrupted by background noise resulting from other contributions such as out-of-focus markers or external photon sources, which also can be modelled as Poissonian random variables, see e.g. Aspelmeier et al. (2015); Munk et al. (2020). Since the background intensity  $\lambda_0 > 0$  varies between experiments, it has to be estimated from the data.

## 5.3. Applying AMS

Applying AMS to the microscopy data in (22) works now as follows. The problem of identifying regions which contain true features can be formulated as detection of regions  $R \subset I_n^d$  such that  $\lambda_i > \lambda_0$  for some  $i \in R$ . However,  $\lambda_0$  is unknown in practice and will hence be estimated from the available data  $Y$  by the global mean  $\hat{\lambda}_n = \bar{Y}_{I_n^d}$ . Below, we have  $n = 400$ ,  $d = 2$ , and we choose  $\mathcal{R}_n$  to be the subset of all rectangles in  $I_n^d$  with scale between  $r_n = 4$  and  $m_n \in \{10, 20\}$ , depending on the investigated structure of interest. The details will be explained below. As we are only interested in active regions, we consider the one-sided testing problem to find regions  $R$  where  $\lambda_i > \lambda_0$ , which can be achieved by restricting the corresponding local rejections  $R$  with  $\bar{Y}_R = \frac{1}{|R|} \sum_{i \in R} Y_i > \hat{\lambda}_n$  (recall Remark 3.8). Furthermore we simulate the threshold  $\eta$  from an asymptotically distribution-free Gaussian approximation  $M_n$  of the test statistic (see (9), with the corresponding adjustment that  $\bar{X}_R > 0$ , cf. also Remark 3.8). We then choose  $\eta$  as the 0.9-quantile  $q_{0.9}^{M_n}$  of this adjusted  $M_n$ . Our main Theorem 3.6 implies that this asymptotically controls the FWER at level  $\alpha$ . The corresponding results will be shown as *significance (fluorescence) maps*, depicting all significant rectangles with colour indicating their size in square-nanometers  $\text{nm}^2$ . If two rectangles overlap, the colour is chosen according to the smaller one, which leads to a map indicating the smallest scale of significance. The major result of this paper implies that the AMS

method ensures the FWER control in (5), and hence each coloured region contains an anomaly with uniform probability  $\geq 0.9$ , i.e. the probability that any of these detections is false is less than 0.1.

## 5.4. Results

### 5.4.1. Data example 1: Single sized crimson beads

As a first data example, we investigate 48nm crimson beads, i.e. carboxylate modified microspheres of 48nm diameter filled with fluorescent markers. Taking into account the microscopes effective resolution of about 60nm, we expect to see circular anomalies of size  $\sim 110$ nm in the data. Therefore we choose  $r_n$  and  $m_n$  such that the smallest box-size is 80nm, and the largest box-size is 400nm. Since each pixel has size  $20\text{nm} \times 20\text{nm}$ , this corresponds to  $r_n = 4$  and  $m_n = 20$ . The measured data for an illumination time of  $t = 5\mu\text{s}$  and  $T = 100\mu\text{s}$  are shown in Figure 6(a) and (d). The corresponding results for AMS are shown in sub-panels (b) and (e). Surprisingly, the test detects most beads based on the  $5\mu\text{s}$  data, even though some of them are barely visible by eye there. Regarding the  $T = 100\mu\text{s}$  data as ground truth, we depict this together with the regions found in (b) in sub-panel (e), revealing that there no false positive detections. Finally, the  $T = 100\mu\text{s}$  data can be used to derive a segmentation into active and inactive regions depicted in (f).




Figure 6: Raw crimson bead data (photon counts, panels (a) and (d)) and corresponding significance maps with level  $\alpha = 0.1$  (smallest significance in  $\text{nm}^2$ , panels (b) and (d)). All significant boxes are plotted with colour indicating their size in  $\text{nm}^2$ . Due to the FWER control, with probability  $\geq 1 - \alpha = 0.9$ , all boxes marked contain anomalies. Using only the smallest scale significance, this leads to the segmentation in (f) showing active regions in red.

### 5.4.2. Data example 2: mixture of differently sized crimson beads

As a second data example, we now consider a mixture of 48nm and 200nm crimson beads. Consequently, the data shown in Figure 7(a) and (d) consists of two structures of different sizes, namely around 48nm and 200nm. Due to the number of markers inside the spheres, the larger structures are significantly brighter than the small ones. We investigate  $t = 15\mu\text{s}$  and  $T = 100\mu\text{s}$ . The AMS procedure is applied with the same parameters as before (i.e. again  $m_n = 20$ , corresponding to a largest box size of 400nm), and the result is shown in sub-panels (b) and (e). It is immediately visible that it detects again nearly all of the small structures already from the measurements taken with  $15\mu\text{s}$  dwell time, which is surprising as some of them are not visible by eye in the data. Once again, regarding the  $T = 100\mu\text{s}$  data as ground truth, we depict this together with the regions found in (b) in sub-panel (e), revealing that there no false positive detections. Finally, the  $T = 100\mu\text{s}$  data can be used to derive a segmentation into active and inactive regions depicted in (f).




Figure 7: Raw crimson bead data (photon counts, logarithmic scale, panels (a) and (d)) and corresponding significance maps with level  $\alpha = 0.1$  (smallest significance in  $\text{nm}^2$ , panels (b) and (e)). All significant boxes are plotted with colour indicating their size in  $\text{nm}^2$ . Due to the FWER control, with probability  $\geq 1 - \alpha = 0.9$ , all boxes marked contain anomalies. Using only the smallest scale significance, this leads to the segmentation in (f) showing active regions in red.

### 5.5. Data example 3: tubulin in neonatal fibroblast

As third data set we finally consider tubulin in neonatal fibroblast. This data set is much more challenging compared to the previous two, since the corresponding structures are filaments and hence, compared to the spheres there, no longer fully contained in single rectangles. To better cover this filamentary structure, which is expected to have width of around 60 – 80nm, we choose

$m_n = 10$  here, i.e. we restrict to boxes with maximal length 200nm. Again, we aim to detect active regions inside the field of view based on noisy  $t = 25\mu\text{s}$  data. It can readily be seen from sub-panel (a) of Figure 8, that the corresponding data is quite noisy, but filamentary structures are clearly visible. Sub-panel (d) of the same figure shows the corresponding  $T = 100\mu\text{s}$  measurement, revealing a folded filamentary structure in the upper right part of the field of view, which was hardly visible by eye in sub-panel (a). With the previously described setup, AMS is able to detect even the folded filamentary structure in the upper left part with a high accuracy, cf. sub-panels (b) and (c). We furthermore find that the method did not yield any false positives despite the relatively high noise-level in the  $t = 25\mu\text{s}$  measurement, as the comparison with the full  $T = 100\mu\text{s}$  ground truth data proves that all significant regions belong to fibroblasts.




Figure 8: Raw tubulin data (photon counts, panels (a) and (d)) and corresponding significance map at level  $\alpha = 0.1$  (smallest significance in  $\text{nm}^2$ , panels (b) and (e)). All significant boxes are plotted with colour indicating their size in  $\text{nm}^2$ . Due to the FWER control, with probability  $\geq 1 - \alpha = 0.9$ , all boxes marked in (b) contain anomalies. Using only the smallest scale significance, this leads to the segmentation in (f) showing active regions in red.

At this point, let us also compare the AMS methodology with direct pixel-wise thresholding. This is e.g. comparable to the techniques used in the previously mentioned SCIM methods (cf. Vinçon et al. (2020); Krishnaswami et al. (2016)). Therefore, we consider simply each pixel, for which the corresponding value  $Y_i$  in (22) exceeds the threshold  $\eta \in \{2, 3\}$ , as an anomaly. Note, that these methods do not lead to any sort of statistical error control. In order to do so, we employ the Benjamini and Hochberg (1995) procedure to obtain a pixel-wise thresholding with controlled false discovery rate (FDR). The results are shown in Figure 9.

Apparently, none of the three methods is able to effectively detect the filamentary structures within the specimen. Even though the hard thresholding with  $\eta = 3$  seems to work best, it still contains several noise pixels in all parts of the image. The FDR-controlled method seems to suffer severely from the inhomogeneity of the image, namely that the right side appears brighter than




Figure 9: Pixel-wise thresholding applied to the raw tubulin data from Figure 8(a). All marked pixels exceeded the corresponding thresholds. Panels (a) and (b) show hard thresholding with no statistical guarantee for  $\eta = 2$  and  $\eta = 3$  respectively, and panel (c) shows the result of the Benjamin-Hochberg procedure with  $\text{FDR} \leq 0.9$ .

the left, and hence the corresponding  $p$ -values of the pixel-wise tests are smaller there. Overall, we can conclude that due to its multiscale structure the AMS procedure clearly outperforms simple pixel-wise thresholding and hence yields a substantial advantage over state-of-the-art SCIM methods.

We also compare the resulting significance maps for different values of  $\alpha$ , see Figure 10. It turns out that the AMS methodology is remarkably stable w.r.t. the significance level  $\alpha$ .




Figure 10: Significance maps from the raw tubulin data in Figure 8(a) for different values of  $\alpha$ .

## 6. Discussion

In this paper, we have developed AMS, a multiscale scanning method allowing for unknown baseline and nuisance parameters. Under a minimal condition on the scales, AMS selects all regions deviating from the global null at guaranteed FWER. To this end we have proven a uniform invariance principle which allows to approximate the multiscale statistic by a distribution-free Gaussian version. This can be used for the computation of (asymptotically) correct critical values. The performance of AMS has been investigated both in simulations and on real data examples from super-resolution microscopy, where it can be seen as an instance of smart microscopy equipped with statistical guarantees. In particular, it outperforms pixel-wise thresholding procedures, including FDR-controlled ones.

We give an FFT based algorithm to perform AMS which roughly scales as  $dNn^d$  where  $N$  is the number of scales contained in  $\mathcal{R}_n$ . This effectively delimits the practical performance of AMS to small dimensions  $d \leq 3$ . However, a combination with ideas from Walther (2010) or using random regions  $R$  might allow to extend AMS to a larger  $d$  in almost linear time. However, it is expected that this comes at the expense of the losing detection power compared to the use of the full grid

$I_n^d$ .

An alternative option to the AMS approach might be based on sample splitting. Therefore, one could separate the spatial observations  $Y_i, i \in I := I_n^d$  into two subsets  $I = I_1 \cup I_2, I_1 \cap I_2 = \emptyset$ , say e.g.  $I_1 = \{1, 3, \dots, n-1\}^d$  and  $I_2 = \{2, 4, \dots, n\}^d$  for even  $n$ , and use  $(Y_i)_{i \in I_1}$  for estimation of  $\theta_0$  and  $\xi$  only. This eases the asymptotic analysis and the statistic  $T_n \left( (Y_i)_{i \in I_2}, \mathcal{R}_n, \hat{\theta}, \hat{\xi} \right)$  will then asymptotically behave exactly like  $T_{n/2} (Y, \mathcal{R}_{n/2}, \theta_0, \xi)$ , and hence previous results on invariance principles might be applicable. However, note that this leads to a severe reduction of the actual resolution of the method (precisely by a factor of 2 due to omitting every second observation) and provides hence a reduced detection power.

A further interesting possible extension of the AMS methodology would be to random fields with spatial dependency. For  $d = 1$  we suggest in the case of simple linear time series error models to follow the approach in Frick et al. (2014), Section 6.1.1 to ease computational effort. Instead of the full likelihood ratio statistic, simply the local LRT statistics for the independent case have to be reweighed by estimates of the total variance. For general dimensions this is more involved. We believe that it might be possible to extend the weak invariance principle for  $m$ -dependent fields in Thm. 2.3 in Kirch et al. (2023) to the multiscale setup in this paper, however, at the expense of a large technical effort.

Finally, as mentioned by a referee, it would be of interest to extend AMS to the situation when no parametric family  $F_{\theta, \xi}$  is assumed. In this case one aims to scan against an (unknown) baseline  $F_0$ , say. Consequently, not only certain nuisance parameters are unknown, rather the entire distribution  $F_0$  and all possible alternatives  $F_i, i \in I_n^d$ . Already for  $d = 1$  this leads to an entirely different situation, as the local LRT statistics have to be replaced by some *non-parametric analogue*, e.g. local median statistics, see Jula Vanegas et al. (2022). How to extend this to general  $d$  is not obvious at all and an interesting task for future research.

## Data availability statement

The data underlying this article, including all software and scripts to reproduce the simulations and data analysis shown in this work are available under <https://go.uniwue.de/math-ams>.

## Acknowledgment

Financial support by the German Research Foundation DFG through subprojects A04 and A07 of CRC 755 and the DFG Cluster of Excellence "MBExC Multiscale Bioimaging - from Molecular Machines to Networks of Excitable Cells" is gratefully acknowledged. We also thank René Siegmund and Alexander Egnér from IFNANO for providing data and helpful discussions. Furthermore, helpful comments and questions raised by the editor, an associate editor and three referees, which led to an improved presentation of the manuscript, are gratefully acknowledged.

## References

- Aboukhamseen, S. M., Soltani, A. R., and Najafi, M. (2016). Modelling cluster detection in spatial scan statistics: formation of a spatial Poisson scanning window and an ADHD case study. *Statist. Probab. Lett.*, 111:26–31.
- Alm, S. E. (1997). On the distributions of scan statistics of a two-dimensional Poisson process. *Adv. in Appl. Probab.*, 29(1):1–18.
- Alm, S. E. (1998). Approximation and simulation of the distributions of scan statistics for Poisson processes in higher dimensions. *Extremes*, 1(1):111–126.
- Arias-Castro, E., Candès, E. J., and Durand, A. (2011). Detection of an anomalous cluster in a network. *Ann. Statist.*, 39(1):278–304.

- Arias-Castro, E., Donoho, D. L., and Huo, X. (2006). Adaptive multiscale detection of filamentary structures in a background of uniform random points. *Ann. Statist.*, 34(1):326–349.
- Aspelmeier, T., Egner, A., and Munk, A. (2015). Modern statistical challenges in high-resolution fluorescence microscopy. *Annu. Rev. Stat. Appl.*, 2:163–202.
- Behr, M., Ansari, M. A., Munk, A., and Holmes, C. (2020). Testing for dependence on tree structures. *Proceedings of the National Academy of Sciences*, 117(18):9787–9792.
- Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. Roy. Statist. Soc. Ser. B*, 57(1):289–300.
- Brown, L. D. (1986). *Fundamentals of statistical exponential families with applications in statistical decision theory*, volume 9 of *Institute of Mathematical Statistics Lecture Notes—Monograph Series*. Institute of Mathematical Statistics, Hayward, CA.
- Caron, Y., Makris, P., and Vincent, N. (2002). A method for detecting artificial objects in natural environments. In *Pattern Recognition, 2002. Proceedings. 16th International Conference on*, volume 1, pages 600–603. IEEE.
- Chan, H. P. and Walther, G. (2013). Detection with the scan and the average likelihood ratio. *Statist. Sinica*, 23(1):409–428.
- Chen, F. and Neill, D. B. (2014). Non-parametric scan statistics for event detection and forecasting in heterogeneous social media graphs. In *Proceedings of the 20th ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 1166–1175. ACM.
- Chen, Y., Wang, T., and Samworth, R. J. (2022). High-dimensional, multiscale online changepoint detection. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 84(1):234–266.
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2014). Gaussian approximation of suprema of empirical processes. *Ann. Statist.*, 42(4):1564–1597.
- Cooley, J. W. and Tukey, J. W. (1965). An algorithm for the machine calculation of complex fourier series. *Mathematics of Computation*, 19:297–301.
- Datta, P. and Sen, B. (2021). Optimal inference with a multidimensional multiscale statistic. *Electron. J. Stat.*, 15(2):5203–5244.
- Dickhaus, T. (2014). *Simultaneous Statistical Inference*. Springer, Heidelberg. With applications in the life sciences.
- Dümbgen, L. and Spokoiny, V. G. (2001). Multiscale testing of qualitative hypotheses. *Ann. Statist.*, 29(1):124–152.
- Dümbgen, L. and Walther, G. (2008). Multiscale inference about a density. *Ann. Statist.*, 36(4):1758–1785.
- Enikeeva, F., Munk, A., and Werner, F. (2018). Bump detection in heterogeneous Gaussian regression. *Bernoulli*, 24(2):1266–1306.
- Farnum, N. R. and Booth, P. (1997). Uniqueness of maximum likelihood estimators of the 2-parameter weibull distribution. *IEEE Trans. Reliab.*, 46(4):523–525.
- Frick, K., Munk, A., and Sieling, H. (2014). Multiscale change point inference. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 76(3):495–580. With 32 discussions by 47 authors and a rejoinder by the authors.
- Futschik, A., Hotz, T., Munk, A., and Sieling, H. (2014). Multiscale DNA partitioning: statistical evidence for segments. *Bioinformatics*, 30(16):2255–2262.



- Gong, G. and Samaniego, F. J. (1981). Pseudomaximum likelihood estimation: theory and applications. *Ann. Statist.*, 9(4):861–869.
- Grama, I. and Nussbaum, M. (1998). Asymptotic equivalence for nonparametric generalized linear models. *Probab. Theory Related Fields*, 111(2):167–214.
- Hell, S. W. (2007). Far-Field Optical Nanoscopy. *Science*, 316(May):1153–1158.
- Hell, S. W. and Wichmann, J. (1994). Breaking the diffraction resolution limit by stimulated emission: stimulated-emission-depletion fluorescence microscopy. *Optics Letters*, 19(11):780–782.
- Jula Vanegas, L., Behr, M., and Munk, A. (2022). Multiscale quantile segmentation. *J. Amer. Statist. Assoc.*, 117(539):1384–1397.
- Kabluchko, Z. (2011). Extremes of the standardized Gaussian noise. *Stochastic Process. Appl.*, 121(3):515–533.
- Kabluchko, Z. and Munk, A. (2008). Exact convergence rate for the maximum of standardized gaussian increments. *Electr. Commun. Probab.*, 13:302–310.
- Kirch, C., Klein, P., and Meyer, M. (2023). Scan statistics for the detection of anomalies in m-dependent random fields with applications to image data.
- Komlós, J., Major, P., and Tusnády, G. (1975). An approximation of partial sums of independent RV’s and the sample DF. I. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 32:111–131.
- König, C., Munk, A., and Werner, F. (2020). Multidimensional multiscale scanning in exponential families: limit theory and statistical consequences. *Ann. Statist.*, 48(2):655–678.
- Kou, J. (2023). Identifying the support of rectangular signals in Gaussian noise. *Comm. Statist. Theory Methods*, 52(10):3262–3289.
- Krishnaswami, V., Van Noorden, C. J. F., Manders, E. M. M., and Hoebe, R. A. (2016). Spatially-controlled illumination microscopy: For prolonged live-cell and live-tissue imaging with extended dynamic range. *Quarterly Reviews of Biophysics*, 49:e19.
- Kulaitis, G., Munk, A., and Werner, F. (2021). What is resolution? A statistical minimax testing perspective on superresolution microscopy. *Ann. Statist.*, 49(4):2292 – 2312.
- Kulldorff, M. (1997). A spatial scan statistic. *Comm. Statist. Theory Methods*, 26(6):1481–1496.
- Lehmann, E. L. and Romano, J. P. (2005). *Testing Statistical Hypotheses*. Springer Texts in Statistics. Springer, New York, third edition.
- Lovato, I., Pini, A., Stamm, A., Taquet, M., and Vantini, S. (2021). Multiscale null hypothesis testing for network-valued data: Analysis of brain networks of patients with autism. *J. R. Stat. Soc. Ser. C Appl. Stat.*, 70(2):372–397.
- Madrid Padilla, O. H., Yu, Y., Wang, D., and Rinaldo, A. (2022). Optimal nonparametric multivariate change point detection and localization. *IEEE Trans. Inform. Theory*, 68(3):1922–1944.
- Mahecic, D., Stepp, W. L., Zhang, C., Griffié, J., Weigert, M., and Manley, S. (2022). Event-driven acquisition for content-enriched microscopy. *Nature Methods*, 19(10):1262–1267.
- Mallik, A., Banerjee, M., and Sen, B. (2013). Asymptotics for  $p$ -value based threshold estimation in regression settings. *Electron. J. Stat.*, 7:2477–2515.
- Miller, Jr., R. G. (1981). *Simultaneous statistical inference*. Springer Series in Statistics. Springer-Verlag, New York-Berlin, second edition.



- Munk, A., Staudt, T., and Werner, F. (2020). *Statistical Foundations of Nanoscale Photonic Imaging*, page 125–143. Springer International Publishing, Cham.
- Neill, D. B. (2012). Fast subset scan for spatial pattern detection. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 74(2):337–360.
- Pandey, G., Westhoff, J. H., Schaefer, F., and Gehrig, J. (2019). A smart imaging workflow for organ-specific screening in a cystic kidney zebrafish disease model. *International Journal of Molecular Sciences*, 20(6).
- Pein, F., Sieling, H., and Munk, A. (2017). Heterogeneous change point inference. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 79(4):1207–1227.
- Proksch, K., Werner, F., and Munk, A. (2018). Multiscale scanning in inverse problems. *Ann. Statist.*, 46(6B):3569–3602.
- Ray, K. and Schmidt-Hieber, J. (2018). The Le Cam distance between density estimation, Poisson processes and Gaussian white noise. *Math. Stat. Learn.*, 1(2):101–170.
- Reiner-Benaim, A., Davis, R. W., and Juneau, K. (2014). Scan statistics analysis for detection of introns in time-course tiling array data. *Statistical applications in genetics and molecular biology*, 13(2):173–190.
- Rivera, C. and Walther, G. (2013). Optimal detection of a jump in the intensity of a Poisson process or in a density with likelihood ratio statistics. *Scand. J. Stat.*, 40(4):752–769.
- Schmidt-Hieber, J., Munk, A., and Dümbgen, L. (2013). Multiscale methods for shape constraints in deconvolution: confidence statements for qualitative features. *Ann. Statist.*, 41(3):1299–1328.
- Sharpnack, J. (2018). Learning patterns for detection with multiscale scan statistics. In Bubeck, S., Perchet, V., and Rigollet, P., editors, *Proceedings of the 31st Conference On Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pages 950–969. PMLR.
- Sharpnack, J. and Arias-Castro, E. (2016). Exact asymptotics for the scan statistic and fast alternatives. *Electron. J. Stat.*, 10(2):2641–2684.
- Sharpnack, J., Rinaldo, A., and Singh, A. (2016). Detecting anomalous activity on networks with the graph fourier scan statistic. *IEEE Transactions on Signal Processing*, 64(2):364–379.
- Smith, R. L. (1985). Maximum likelihood estimation in a class of nonregular cases. *Biometrika*, 72(1):67–90.
- Strack, R. (2020). Smarter microscopes. *Nature Methods*, 17(1):23–23.
- van der Vaart, A. W. (1998). *Asymptotic statistics*, volume 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge.
- Vinçon, B., Geisler, C., and Egner, A. (2020). Pixel hopping enables fast sted nanoscopy at low light dose. *Opt. Express*, 28(4):4516–4528.
- Walther, G. (2010). Optimal and fast detection of spatial clusters with scan statistics. *Ann. Statist.*, 38(2):1010–1033.
- Walther, G. (2020). *Calibrating the Scan Statistic with Size-Dependent Critical Values*, page 1–15. Springer New York, New York, NY.
- Walther, G. (2022). Tail bounds for empirically standardized sums. *Electron. J. Stat.*, 16(1):2406–2431.
- Walther, G. and Perry, A. (2022). Calibrating the scan statistic: finite sample performance versus asymptotics. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 84(5):1608–1639.

## A. Proofs

We start with some helpful abbreviations. For the whole section, let  $X = (X_i)_{i \in I_n^d}$  be a field of i.i.d. standard Gaussians,  $X_i \sim \mathcal{N}(0, 1)$ . As we rely heavily on centred and standardized partial sums, we introduce for  $R \in \mathcal{R}_n$  the quantity

$$Y_R(\theta, \xi) := |R|^{-1} \sum_{i \in R} \left( \frac{Y_i - m(\theta, \xi)}{\sqrt{v(\theta, \xi)}} \right) = \frac{\bar{Y}_R - m(\theta, \xi)}{\sqrt{v(\theta, \xi)}}, \quad (\theta, \xi) \in \Theta \times \Xi$$

with the mean and variance functions  $m$  and  $v$  from Assumption 1. Also recall our abbreviation

$$\bar{X}_R := \frac{1}{|R|} \sum_{i \in R} X_i.$$

### A.1. Preparations

Next we recall a series of helpful results taken from König et al. (2020).

**Lemma A.1** (Coupling). *Let Assumptions 1(a) and 2 hold true, and let  $Y = (Y_i)_{i \in I_n^d}$  be an array of i.i.d. random variables  $Y_i \sim F_{\theta, \xi}$  with  $\theta \in U, \xi \in V$  and let  $(r_n)_{n \in \mathbb{N}} \subset \mathbb{N}$  be a sequence. Then, on the same probability space, there exists an array  $X = (X_i)_{i \in I_n^d}$  of i.i.d. standard Gaussians  $X_i \sim \mathcal{N}(0, 1)$  such that*

$$\mathbb{P} \left[ \left| \max_{R \in \mathcal{R}_n(r_n, n^d)} |R|^{\frac{1}{2}} |Y_R(\theta, \xi)| - \max_{R \in \mathcal{R}_n(r_n, n^d)} |R|^{\frac{1}{2}} |\bar{X}_R| \right| > \delta \right] \leq C \delta^{-3} \left( \frac{\log^{10} n}{r_n} \right)^{1/2}$$

for all  $\delta > 0$  with some universal constant  $C > 0$  independent of  $\theta$  and  $\xi$ .

*Proof.* As the distributions  $F_{\theta, \xi}$  have uniformly sub-exponential tails, this follows directly from Theorem 4.3 in König et al. (2020) in combination with a symmetrization argument as employed in the proof of Theorem 2.5 there.  $\square$

Note, that in terms of  $\mathcal{O}_{\mathbb{P}_0}$  notation, Lemma A.1 yields

$$\max_{R \in \mathcal{R}_n(r_n, n^d)} |R|^{\frac{1}{2}} |Y_R(\theta, \xi)| - \max_{R \in \mathcal{R}_n(r_n, n^d)} |R|^{\frac{1}{2}} |\bar{X}_R| = \mathcal{O}_{\mathbb{P}_0} \left( \left( \frac{\log^{10}(n)}{r_n} \right)^{\frac{1}{6}} \right), \quad (23)$$

uniformly in  $(\theta, \xi) \in U \times V$ .

**Lemma A.2** (Taylor expansion, see Lemma 5.1 in König et al. (2020)). *Let  $Y = (Y_i)_{i \in I_n^d}$  be an array of i.i.d. random variables  $Y_i \sim F_{\theta_0, \xi}$  with fixed  $\theta_0 \in \Theta, \xi \in \Xi$  and let  $(r_n)_{n \in \mathbb{N}} \subset \mathbb{N}$  be a sequence satisfying (18). If Assumptions 1(a) and 2 hold true, then*

$$\max_{R \in \mathcal{R}_n(r_n, n^d)} \left| T_R(Y, \theta_0, \xi) - |R|^{\frac{1}{2}} Y_R(\theta_0, \xi) \right| = \mathcal{O}_{\mathbb{P}_0} \left( \left( \frac{\log^3(n)}{r_n} \right)^{1/4} \right).$$

Note, that the previous results remain true if we replace the upper bound  $n^d$  for the scale size by  $m_n$  with any sequence  $(m_n)_{n \in \mathbb{N}} \subset \mathbb{N}$  satisfying  $r_n \leq m_n$  for all  $n \in \mathbb{N}$ .

**Lemma A.3.** *Let Assumptions 1(a) and 2 hold true, and let  $Y = (Y_i)_{i \in I_n^d}$  be an array of i.i.d. random variables  $Y_i \sim F_{\theta, \xi}$  with fixed  $\theta \in U, \xi \in V$ . If  $(r_n)_{n \in \mathbb{N}} \subset \mathbb{N}$  is a sequence satisfying (18), then we obtain*

$$\max_{R \in \mathcal{R}_n(r_n, m_n)} |R| |Y_R(\theta_0, \xi)|^3 = \mathcal{O}_{\mathbb{P}_0} \left( \left( \frac{\log^3(n)}{r_n} \right)^{\frac{1}{2}} \right)$$

uniformly for  $\theta \in U, \xi \in V$ .

*Proof.* Note, that due to  $\alpha, \tilde{\alpha} > 0$ , (18) implies especially that  $\gamma \geq 4$  and hence  $\log^4(n) = o(r_n)$ .

It is well known (see e.g. Kabluchko (2011)), that for a standard Gaussian array  $X = (X_i)_{i \in I_n^d}$ ,  $X_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$ , one has

$$\mathbb{E} \left[ \max_{R \in \mathcal{R}_n(r_n, m_n)} |R|^{\frac{1}{2}} |\bar{X}_R| \right] \leq C \sqrt{\log(\#\mathcal{R}_n(r_n, m_n))} \quad (24)$$

with some constant  $C > 0$ . Due to Assumption 2 this implies that

$$\frac{1}{\sqrt{\log(n)}} \max_{R \in \mathcal{R}_n(r_n, m_n)} |R|^{\frac{1}{2}} |\bar{X}_R| = \mathcal{O}_{\mathbb{P}_0}(1).$$

Together with (23) this yields

$$\frac{1}{\sqrt{\log(n)}} \max_{R \in \mathcal{R}_n(r_n, m_n)} |R|^{\frac{1}{2}} |Y_R(\theta, \xi)| = \mathcal{O}_{\mathbb{P}_0}(1)$$

uniformly for  $(\theta, \xi) \in U \times V$ , where we used that  $\log^4(n) = o(r_n)$ . Consequently we find

$$\sqrt{\frac{r_n}{\log^3(n)}} \max_{R \in \mathcal{R}_n(r_n, m_n)} |R| |Y_R(\theta_0, \xi)|^3 \leq \sqrt{\frac{1}{\log^3(n)}} \max_{R \in \mathcal{R}_n(r_n, m_n)} |R|^{\frac{3}{2}} |Y_R(\theta_0, \xi)|^3 = \mathcal{O}_{\mathbb{P}_0}(1)$$

uniformly for  $(\theta, \xi) \in U \times V$ , which proves the claim.  $\square$

## A.2. An adjusted Taylor expansion

Now we are in position to derive some preparations for the proof of Theorem 3.6. Therefore we first replace the likelihood ratio statistic  $T_R$  by its Taylor expansion according to Assumption 1(e):

**Lemma A.4.** *Let  $(r_n)_{n \in \mathbb{N}} \subset \mathbb{N}$  and  $(m_n)_{n \in \mathbb{N}} \subset \mathbb{N}$  be sequences tending to  $\infty$  and consider the local statistic  $T_R$  from (3). Suppose that  $Y$  is as in (1) and let Assumptions 1 and 2 be fulfilled. If the estimators  $\hat{\theta}_n$  and  $\hat{\xi}_n$  satisfy (17), then*

$$\max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R(Y, \hat{\theta}_n, \hat{\xi}_n) - |R|^{1/2} |Y_R(\theta_0, \xi)| \right| = \mathcal{O}_{\mathbb{P}_0} \left( \left( \frac{\log^7(n)}{r_n} \right)^{\frac{1}{4}} + \sqrt{\frac{m_n}{s_n}} \right). \quad (25)$$

*Proof.* Let  $U \times V$  be the neighbourhood of  $(\theta_0, \xi)$  as in Assumption 1. By (17), the probability that  $\hat{\theta}_n \in U$  and  $\hat{\xi}_n \in V$  tends to 1, as  $n \rightarrow \infty$ . Hence, it suffices to prove the result conditional on this event. Consequently we can assume  $\hat{\theta}_n \in U$  and  $\hat{\xi}_n \in V$  in the following.

By the triangle inequality and Lemma A.2, we find

$$\begin{aligned} & \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R(Y, \hat{\theta}_n, \hat{\xi}_n) - |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| \right| \\ & \leq \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R(Y, \theta_0, \xi) - |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| \right| \\ & \quad + \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R(Y, \hat{\theta}_n, \hat{\xi}_n) - T_R(Y, \theta_0, \xi) \right| \\ & \leq \sqrt{\max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R^2(Y, \hat{\theta}_n, \hat{\xi}_n) - T_R^2(Y, \theta_0, \xi) \right|} + \mathcal{O}_{\mathbb{P}_0} \left( \left( \frac{\log^3(n)}{r_n} \right)^{\frac{1}{4}} \right), \end{aligned}$$

where we exploited Lemma A.2 and  $|a - b| \leq \sqrt{|a^2 - b^2|}$ . By the triangle inequality and Assumption 1(e) we find furthermore that

$$\begin{aligned}
& \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R^2(Y, \hat{\theta}, \hat{\xi}) - T_R^2(Y, \theta_0, \xi) \right| \\
& \leq \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R^2(Y, \hat{\theta}, \hat{\xi}) - |R| \left| Y_R(\hat{\theta}_n, \hat{\xi}_n) \right|^2 \right| + \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| T_R^2(Y, \theta_0, \xi) - |R| \left| Y_R(\theta_0, \xi) \right|^2 \right| \\
& \quad + \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| |R| \left| Y_R(\hat{\theta}_n, \hat{\xi}_n) \right|^2 - Y_R(\theta_0, \xi)^2 \right| \\
& \leq C_T \max_{R \in \mathcal{R}_n(r_n, m_n)} |R| \left| Y_R(\hat{\theta}_n, \hat{\xi}_n) \right|^3 + C_T \max_{R \in \mathcal{R}_n(r_n, m_n)} |R| \left| Y_R(\theta_0, \xi) \right|^3 \\
& \quad + \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| |R| \left| Y_R(\hat{\theta}_n, \hat{\xi}_n) \right|^2 - Y_R(\theta_0, \xi)^2 \right| \\
& \leq 2C_T \max_{R \in \mathcal{R}_n(r_n, m_n)} |R| \left| Y_R(\theta_0, \xi) \right|^3 + \max_{R \in \mathcal{R}_n(r_n, m_n)} |R| \left| Y_R(\hat{\theta}_n, \hat{\xi}_n) \right|^2 - Y_R(\theta_0, \xi)^2 \Big| \\
& \quad + C_T \max_{R \in \mathcal{R}_n(r_n, m_n)} \left| |R| \left| Y_R(\hat{\theta}_n, \hat{\xi}_n) \right|^3 - Y_R(\theta_0, \xi)^3 \right|,
\end{aligned}$$

where we used  $\left| |a|^3 - |b|^3 \right| = \left| |a^3| - |b^3| \right| \leq |a^3 - b^3|$  for  $a, b \in \mathbb{R}$ . In view of Assumptions 1(a) and 2, the first term in the last display can be bounded using Lemma A.3. The second and third term will be handled using the mean value theorem. Therefore, note that all derivatives of

$$(\theta, \xi) \mapsto Y_R(\theta, \xi)^k, \quad k \in \{2, 3\}$$

are a.s. bounded due to Assumption 1(b)–(d) and the fact that  $\bar{Y}_R$  is a.s. bounded in view of Lemma A.3. This together with (17) implies that

$$\max_{R \in \mathcal{R}_n(r_n, m_n)} |R| \left| Y_R(\hat{\theta}, \hat{\xi})^2 - Y_R(\theta_0, \xi)^2 \right| \leq m_n C \left( \|\hat{\theta}_n - \theta_0\|_2 + \|\hat{\xi}_n - \xi\|_2 \right) = \mathcal{O}_{\mathbb{P}_0} \left( \frac{m_n}{s_n} \right),$$

where  $C$  is some generic constant  $C > 0$ . As  $\max_{R \in \mathcal{R}_n(r_n, m_n)} |R| \left| Y_R(\hat{\theta}_n, \hat{\xi}_n)^3 - Y_R(\theta_0, \xi)^3 \right|$  can be treated similarly, this yields the claim.  $\square$

### A.3. Proof of Theorem 3.6

Now we are ready to prove the Gaussian approximation result from Theorem 3.6.

*Proof of Theorem 3.6.* For notional simplicity, we will throughout this proof drop the index of  $\tilde{\omega}_n$  and  $\omega_n$ . Furthermore we abbreviate  $\tilde{\mathcal{R}}_n := \mathcal{R}_n(r_n, m_n)$ . First of all, we estimate

$$\begin{aligned}
& \left| T_n \left( Y, \tilde{\mathcal{R}}_n, \hat{\theta}, \hat{\xi} \right) - M_n \left( \tilde{\mathcal{R}}_n \right) \right| \\
& = \left| \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( T_R \left( Y, \hat{\theta}_n, \hat{\xi}_n \right) - \omega(|R|) \right) - \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |\bar{X}_R| - \omega(|R|) \right) \right| \\
& \leq \left| \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( T_R \left( Y, \hat{\theta}_n, \hat{\xi}_n \right) - \omega(|R|) \right) - \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \omega(|R|) \right) \right| \\
& \quad + \left| \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \omega(|R|) \right) - \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |\bar{X}_R| - \omega(|R|) \right) \right| \\
& \leq \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \max_{R \in \tilde{\mathcal{R}}_n} \left| T_R \left( Y, \hat{\theta}_n, \hat{\xi}_n \right) - |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| \right| \\
& \quad + \left| \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \omega(|R|) \right) - \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |\bar{X}_R| - \omega(|R|) \right) \right|,
\end{aligned}$$

where we used  $\| \|x\|_\infty - \|y\|_\infty \| \leq \|x - y\|_\infty$ . The first term can be controlled using Assumption 3 and Lemma A.4 by

$$\max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \max_{R \in \tilde{\mathcal{R}}_n} \left| T_R \left( Y, \hat{\theta}_n, \hat{\xi}_n \right) - |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| \right| = \mathcal{O}_{\mathbb{P}_0} \left( \frac{\log^{\tilde{\alpha}}(n)}{r_n} \left( \left( \frac{\log^3(n)}{r_n} \right)^{\frac{1}{4}} + \sqrt{\frac{m_n}{s_n}} \right) \right).$$

So it remains to estimate the second term, i.e. for a suitable sequence  $(c_n)_{n \in \mathbb{N}}$  we want to show that

$$\lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbb{P}_0 \left[ \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \omega(|R|) \right) - \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |\bar{X}_R| - \omega(|R|) \right) \right] > c_n M = 0 \quad (26)$$

This is more subtle, as the coupling in Lemma A.1 only yields a result for the difference of the maxima, and not for the maximum of the differences (as lemma A.4 does). We therefore borrow a slicing technique from Proksch et al. (2018), dividing the set of scales into families on which  $\tilde{\omega}$  and  $\omega$  are almost constant, i.e. into

$$\mathcal{R}_{n,j} := \{R \in \mathcal{R}_n \mid \exp(\epsilon_j) \leq |R| < \exp(\epsilon_{j+1})\}.$$

Here and in what follows we choose

$$\epsilon_1 := \log(r_n), \quad \epsilon_j = \epsilon_1 + \frac{j-1}{J} \log\left(\frac{m_n}{r_n}\right), \quad j = 2, \dots, J.$$

With this definition, for any choice  $J \in \mathbb{N}_{\geq 2}$  we obtain

$$\tilde{\mathcal{R}}_n = \mathcal{R}_n(r_n, m_n) = \{R \in \mathbb{R}_n \mid r_n \leq |R| \leq m_n\} = \bigcup_{j \in J} \mathcal{R}_{n,j}.$$

The parameter  $J \in \mathbb{N}$  will be defined later. On  $\mathcal{R}_{n,j}$  we can approximate  $\tilde{\omega}(|R|)$  and  $\omega(|R|)$  by

$$\tilde{\omega}_{j,n} := \tilde{\omega}(\exp(\epsilon_j)) \quad \text{and} \quad \omega_{j,n} := \omega(\exp(\epsilon_j)),$$

respectively. Then it holds

$$\omega_{j+1,n} \leq \omega(|R|) \leq \omega_{j,n} \quad \text{and} \quad \tilde{\omega}_{j+1,n} \leq \tilde{\omega}(|R|) \leq \tilde{\omega}_{j,n}$$

for all  $R \in \mathcal{R}_{n,j}$ . Now we compute

$$\begin{aligned} & \max_{R \in \mathcal{R}_{n,j}} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \omega(|R|) \right) - \max_{R \in \mathcal{R}_{n,j}} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |\bar{X}_R| - \omega(|R|) \right) \\ & \leq \left( \tilde{\omega}_{j,n} \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \tilde{\omega}_{j+1,n} \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |\bar{X}_R| \right) + (\tilde{\omega}_{j,n} \omega_{j,n} - \tilde{\omega}_{j+1,n} \omega_{j+1,n}) \\ & = \tilde{\omega}_{j,n} \left( \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |\bar{X}_R| \right) + (\tilde{\omega}_{j,n} - \tilde{\omega}_{j+1,n}) \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |\bar{X}_R| \\ & \quad + (\tilde{\omega}_{j,n} \omega_{j,n} - \tilde{\omega}_{j+1,n} \omega_{j+1,n}), \end{aligned}$$

and a similar relation holds true if the roles of  $\bar{X}_R$  and  $Y_R(\theta_0, \xi)$  are interchanged. To bound the differences involving the  $\tilde{\omega}_{j,n}$  terms, we exploit the rough bound

$$\tilde{\omega}_{j,n} = \tilde{\omega}(\exp(\epsilon_j)) \leq C_\omega \left( \log \left( \frac{n^d}{\exp(\epsilon_j)} \right) \right)^{\tilde{\alpha}} \leq C_\omega (\log(n^d))^{\tilde{\alpha}} =: A_n \quad (27)$$

for all  $1 \leq j \leq J$ , and a similar analogue for  $\omega_{j,n}$ . From the mean value theorem we get

$$\begin{aligned}
\tilde{\omega}_{j,n} - \tilde{\omega}_{j+1,n} &= \tilde{\omega}(\exp(\epsilon_{j+1})) - \tilde{\omega}(\exp(\epsilon_j)) \\
&\leq |\exp(\epsilon_j) - \exp(\epsilon_{j+1})| \max_{\zeta \in (\epsilon_j, \epsilon_{j+1})} \tilde{\omega}'(\exp(\zeta)) \\
&\leq C_\omega (\log(n^d))^{\max(\tilde{\beta}, 0)} (\exp(\epsilon_{j+1} - \epsilon_j) - 1) \\
&= C_\omega (\log(n^d))^{\max(\tilde{\beta}, 0)} \left( \exp\left(\frac{1}{J} \log\left(\frac{m_n}{r_n}\right)\right) - 1 \right) \\
&= C_\omega (\log(n^d))^{\max(\tilde{\beta}, 0)} \left( \left(\frac{m_n}{r_n}\right)^{\frac{1}{J}} - 1 \right) =: B_n
\end{aligned}$$

Similar statements hold for  $\omega_{j,n}$  and  $\omega_{j,n} - \omega_{j+1,n}$  and hence using (27) we obtain

$$\begin{aligned}
|\tilde{\omega}_{j,n}\omega_{j,n} - \tilde{\omega}_{j+1,n}\omega_{j+1,n}| &= |\tilde{\omega}_{j,n}(\omega_{j,n} - \omega_{j+1,n}) - \omega_{j+1,n}(\tilde{\omega}_{j,n} - \tilde{\omega}_{j+1,n})| \\
&\leq C_\omega^2 \left( \left(\frac{m_n}{r_n}\right)^{\frac{1}{J}} - 1 \right) \left( (\log(n^d))^{\tilde{\alpha} + \max\{\beta, 0\}} + (\log(n^d))^{\alpha + \max\{\tilde{\beta}, 0\}} \right) =: C_n
\end{aligned}$$

This implies

$$\begin{aligned}
&\left| \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \omega(|R|) \right) - \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |\bar{X}_R| - \omega(|R|) \right) \right| \\
&\leq A_n \max_{1 \leq j \leq J} \left| \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |\bar{X}_R| \right| + B_n \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |\bar{X}_R| + C_n,
\end{aligned}$$

and hence

$$\begin{aligned}
&\mathbb{P}_0 \left[ \left| \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \omega(|R|) \right) - \max_{R \in \tilde{\mathcal{R}}_n} \tilde{\omega}(|R|) \left( |R|^{\frac{1}{2}} |\bar{X}_R| - \omega(|R|) \right) \right| > c_n M \right] \\
&\leq \mathbb{P}_0 \left[ A_n \max_{1 \leq j \leq J} \left| \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |Y_R(\theta_0, \xi)| - \max_{R \in \mathcal{R}_{n,j}} |R|^{\frac{1}{2}} |\bar{X}_R| \right| > \frac{c_n M}{3} \right] \\
&\quad + \mathbb{P}_0 \left[ B_n \max_{R \in \tilde{\mathcal{R}}_n} |R|^{\frac{1}{2}} |\bar{X}_R| > \frac{c_n M}{3} \right] + \mathbb{P}_0 \left[ C_n > \frac{c_n M}{3} \right] \\
&=: \text{I} + \text{II} + \text{III}.
\end{aligned}$$

To bound I, we use Lemma A.1 and the union bound to obtain

$$\text{I} \leq C |J| \frac{3^3 A_n^3}{c_n^3 M^3} \left( \frac{\log^{10}(n)}{r_n} \right)^{\frac{1}{2}}.$$

Here and in what follows,  $C > 0$  is some generic constant, the value of which can change from line to line. For II, we exploit Markov's inequality as well as (24) and find in view of Assumption 2, that

$$\text{II} \leq \frac{\mathbb{E} \left[ \max_{R \in \mathcal{R}_n} |R|^{\frac{1}{2}} |\bar{X}_R| \right]}{c_n M 3^{-1} B_n^{-1}} \leq C \frac{B_n \sqrt{\log(n)}}{c_n M}$$

with some constant  $C > 0$ . This shows that (26) is satisfied as soon as

- (A)  $|J| A_n^3 c_n^{-3} \log^5(n) r_n^{-\frac{1}{2}} = \mathcal{O}(1)$ ,
- (B)  $B_n c_n^{-1} \sqrt{\log(n)} = \mathcal{O}(1)$ , and
- (B)  $C_n = o(c_n)$

hold true. To obtain this, we set

$$J := \lfloor \frac{\log^{\nu+1}(n)}{c_n} \rfloor$$

with some parameter  $\nu > 0$  to be determined. This implies

$$|J| \frac{A_n^3}{c_n^3} \left( \frac{\log^{10}(n)}{r_n} \right)^{\frac{1}{2}} \leq C \frac{\log^{6+3\tilde{\alpha}+\nu}}{c_n^4 \sqrt{r_n}},$$

which proves (B) for

$$c_n := \left( \frac{\log^{12+6\tilde{\alpha}+2\nu}(n)}{r_n} \right)^{\frac{1}{8}}. \quad (28)$$

Next we use  $m_n/r_n \leq n^d$  and use  $\exp(x) - 1 \leq 2x$  for  $x \in [0, 1]$  to find

$$\left( \frac{m_n}{r_n} \right)^{\frac{1}{J}} \leq n^{\frac{d}{J}} = \exp\left( \frac{dc_n}{\log^\nu(n)} \right) \leq 1 + C \frac{c_n}{\log^\nu(n)}$$

for any sufficiently large  $n$  supposed that  $c_n = o(\log^\nu(n))$ . Thus it holds

$$B_n \leq C \frac{c_n}{\log^{\nu - \max\{\tilde{\beta}, 0\}}(n)},$$

and hence (A) is satisfied if  $\nu \geq \max\{\tilde{\beta}, 0\} + \frac{1}{2}$ . It can readily be seen that (C) is now satisfied as soon as we define

$$\nu := \max\left\{ \frac{1}{2}, \alpha, \tilde{\alpha} \right\} + \max\left\{ \beta, \tilde{\beta}, 0 \right\}.$$

Note that this also ensures the required condition  $c_n = o(\log^\nu(n))$  under (18). Hence (26) is satisfied with  $c_n$  as in (28), and thus we have

$$\left| T_n(Y, \tilde{\mathcal{R}}_n, \hat{\theta}, \hat{\xi}) - M_n(\tilde{\mathcal{R}}_n) \right| = \mathcal{O}_{\mathbb{P}_0} \left( \frac{\log^{\tilde{\alpha}}(n)}{r_n} \left( \left( \frac{\log^3(n)}{r_n} \right)^{\frac{1}{4}} + \sqrt{\frac{m_n}{s_n}} \right) + \left( \frac{\log^{12+6\tilde{\alpha}+2\nu}(n)}{r_n} \right)^{\frac{1}{8}} \right),$$

which together with  $\frac{\log^{\tilde{\alpha}+3/44}(n)}{r_n^{5/4}} = \mathcal{O}\left(\frac{\log^{\gamma/8}(n)}{r_n^{1/8}}\right)$  proves the claim.  $\square$