

ON THE DISTRIBUTION OF THE NORM OF PARTITIONS

WALTER BRIDGES AND WILLIAM CRAIG

ABSTRACT. The norm of an integer partition is defined as the product of its parts. This statistic was recently introduced by Schneider in connection to partition zeta functions. In this note, we use the method of moments to study the distribution of the norm under the uniform probability measure on partitions of n as $n \rightarrow \infty$. We use singularity analysis to prove asymptotics for the moments and show as a result that the norm lacks a non-trivial limiting distribution on $[0, \infty)$.

1. INTRODUCTION

A *partition* of a natural number $n \geq 0$ is a non-increasing list $\lambda = (\lambda_1, \dots, \lambda_\ell)$ of non-negative integers such that $\lambda_1 + \lambda_2 + \dots + \lambda_\ell = n$, and we say $\lambda \vdash n$ when λ is a partition of n . In this paper, we focus on the *norm* of a partition $\lambda \vdash n$, defined for $\lambda = (\lambda_1, \dots, \lambda_\ell)$ as the product of its parts, i.e.

$$N(\lambda) := \prod_{j=1}^{\ell} \lambda_j.$$

Interest in this new statistic comes from the work of Schneider and his collaborators [8, 9, 10, 11, 12] to develop a multiplicative theory of partitions that generalizes classical multiplicative number theory. Here, the norm plays a central role in the construction of partition zeta functions [12]. For an overview of recent developments in this area, see [8].

The structure of large partitions is a central theme in the theory of partitions. Celebrated work of Hardy–Ramanujan and Rademacher, using the modularity of the generating function for partitions, yields the well-known asymptotic for $p(n)$, the number of partitions of n :

$$p(n) \sim \frac{1}{4n\sqrt{3}} e^{\pi\sqrt{\frac{2n}{3}}}.$$

Another early result is the Erdős–Lehner Theorem [2, Theorem 1.1], which gives the distribution for the largest part (or equivalently the number of parts) under the uniform measure on partitions of n , as $n \rightarrow \infty$. A modern approach to many statistics for partitions was developed by Fristedt in [4] and was subsequently generalized to a variety of combinatorial structures by Ariata and Tavaré [1]. To study the norm, however, we follow the more classical method of moments.

We first recall the maximum of the norm across partitions of n . It is given as an exercise in [5, pp. 30–31] and [7, p. 5] and is restated in Schneider and Sill’s recent overview of the norm [13, Theorem 11].

Proposition 1.1. *Let $n > 1$. Then we have*

$$M_n := \max\{N(\lambda) : \lambda \vdash n\} = \begin{cases} 3^{\frac{n}{3}} & \text{if } n \equiv 0 \pmod{3}, \\ 4 \cdot 3^{\frac{n-4}{3}} & \text{if } n \equiv 1 \pmod{3}, \\ 2 \cdot 3^{\frac{n-2}{3}} & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

Let E_n denote the expectation operator under the uniform measure on partitions of n . For an integer $\ell \geq 1$, the ℓ -th moment of the norm is

$$E_n(N^\ell) = \frac{\sum_{\lambda \vdash n} N(\lambda)^\ell}{p(n)}.$$

We use standard singularity analysis of the generating function for the ℓ -th moments, to prove the following asymptotic formula. As in Proposition 1.1, the constants that appear to depend on $n \pmod{3}$.

Theorem 1.2. *Fix an integer $\ell \geq 1$. For an integer n , write n_0 for the least residue of $n \pmod{3}$. Then as $n \rightarrow \infty$,*

$$E_n(N^\ell) \sim c_\ell^{(n_0)} \frac{3^{\frac{\ell n}{3}}}{p(n)},$$

where

$$c_\ell^{(n_0)} := \sum_{j=1}^3 e^{-\frac{2\pi i j n_0}{3}} \cdot \frac{1}{3} \prod_{\substack{k \geq 1 \\ k \neq 3}} \left(1 - k^\ell 3^{-\frac{k\ell}{3}} e^{\frac{2\pi i j k}{3}}\right)^{-1}.$$

Remark 1.3. We make the following remarks on Theorem 1.2.

- (1) Following our proof, one could give more terms in the asymptotic expansion above by taking more singularities into account.
- (2) Since the inequality $\text{Var}(E_n) := E_n(N)^2 - E_n(N^2) \geq 0$ fails for sufficiently large n , no rescaling of N can converge to a non-trivial distribution on $[0, \infty)$. This lack of convergence is somewhat similar to certain hook-length statistics for partitions studied by Lang, Wan and Xu [6].
- (3) With a little extra work, one could obtain any finite-length asymptotic expansion for $E_n(N^\ell)$ using singularity analysis as in [3, Chapter VI]. Indeed, it may be possible to use this method to compute moments of a wide range of multiplicative partition statistics. See Schneider's constructions in [10, 12].

Proof of Theorem 1.2. Let $\ell \geq 1$ be an integer. Using straightforward manipulations with generating series, it is clear that

$$F_\ell(q) := 1 + \sum_{n \geq 1} p(n) E_n(N^\ell) q^n = \sum_{\lambda} N(\lambda)^\ell q^{|\lambda|} = \prod_{k \geq 1} \frac{1}{1 - k^\ell q^k},$$

where the sum \sum_{λ} runs over all partitions, regardless of size. As the series $\sum_{k \geq 1} k^\ell q^k$ converges for $|q| < 1$, we see that the product defining F_ℓ converges absolutely for $|q| < 1$,

ℓ	$c_\ell^{(1)}$	$c_\ell^{(2)}$	$c_\ell^{(3)}$
1	97922.939...	97922.904...	97923.267...
2	667.849...	667.848...	668.148...
3	86.275...	86.298...	86.602...
4	26.884...	26.927...	27.247...
5	12.453...	12.513...	12.851...
6	7.157...	7.229...	7.588...
7	4.689...	4.771...	5.152...
8	3.347...	3.435...	3.839...
9	2.535...	2.628...	3.054...
10	2.003...	2.100...	2.548...

TABLE 1. The constants in Theorem 1.2.

so long as q avoids zeros of denominators in the product. Hence, F_ℓ has singularities at

$$q = e^{\frac{2\pi ij}{k}} k^{-\frac{\ell}{k}}, \quad \text{for } 1 \leq j \leq k,$$

and is otherwise holomorphic for $|q| < 1$. Observe that $k \mapsto k^{-\frac{\ell}{k}}$ is an increasing function for $k \geq 3$, and for $k \in \mathbb{N}$ it obtains a minimum at $k = 3$. Thus, $F_\ell(q)$ has a meromorphic continuation to $|q| < 2^{-\frac{1}{2}}$ with simple poles at $q = e^{\frac{2\pi ij}{3}} 3^{-\frac{\ell}{3}}$ for $1 \leq j \leq 3$.

We now apply singularity analysis as described in [3, Chapter VI]. As $q \rightarrow e^{\frac{2\pi ij}{3}} 3^{-\frac{\ell}{3}}$, we have

$$\begin{aligned} F_\ell(q) &\sim \frac{1}{1 - e^{-\frac{2\pi ij}{3}} 3^{\frac{\ell}{3}} q} \cdot \frac{1}{(1 - e^{-\frac{2\pi i(j+1)}{3}} e^{\frac{2\pi ij}{3}})(1 - e^{-\frac{2\pi i(j+2)}{3}} e^{\frac{2\pi ij}{3}})} \prod_{\substack{k \geq 1 \\ k \neq 3}} \left(1 - k^\ell 3^{-\frac{k\ell}{3}} e^{\frac{2\pi ijk}{3}}\right)^{-1} \\ &= \frac{1}{1 - e^{-\frac{2\pi ij}{3}} 3^{\frac{\ell}{3}} q} \cdot \frac{1}{3} \prod_{\substack{k \geq 1 \\ k \neq 3}} \left(1 - k^\ell 3^{-\frac{k\ell}{3}} e^{\frac{2\pi ijk}{3}}\right)^{-1}. \end{aligned}$$

Theorem 1.1 follows immediately from [3, Theorem VI.5]. \square

REFERENCES

- [1] R. Arriata and S. Tavaré, *Independent process approximations for random combinatorial structures*, Adv. Math. **104** (1994), 90–154.
- [2] P. Erdős and J. Lehner, *The distribution of the number of summands in the partitions of a positive integer*, Duke Journal of Mathematics **8** (1941), 335–345.
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*. Cambridge University Press (2009).
- [4] B. Fristedt, *The structure of random large partitions of integers*, Transaction of the American Mathematical Society **337** (1993), 703–735.
- [5] P. Halmos, *Problems for Mathematicians, Young and Old*, Math. Assoc. Amer., 1991.
- [6] H. Lang, H. Wan, N. Xu, *Distributions of hook lengths divisible by 2 or 3*, J. Number Theory, **246** (2023), 227–251.
- [7] D. Newman, *A Problem Seminar*, Springer-Verlag, New York, 1982.

- [8] K. Ono, L. Rolen, R. Schneider, *Explorations in the theory of partition zeta functions*. Exploring the Riemann Zeta Function: 190 years from Riemann's Birth (2017), 223-264.
- [9] K. Ono, R. Schneider, I. Wagner, *Partition-theoretic formulas for arithmetic densities*. Gainesville International Number Theory Conference. Cham: Springer International Publishing, 2016.
- [10] R. Schneider, *Arithmetic of partitions and the q -bracket operator*, Proceedings of the American Mathematical Society **145**, 5, (2017), 1953–1968.
- [11] R. Schneider, *Eulerian series, zeta functions and the arithmetic of partitions*, Ph.D. Thesis, Emory University (2018).
- [12] R. Schneider, *Partition zeta functions*, Research in Number Theory, **2** (2016), 1-17.
- [13] R. Schneider, A.V. Sills, *The product of parts or "norm" of a partition*, Integers: Electronic Journal of Combinatorial Number Theory (2020).