

# IMPROVED $C^{1,1}$ REGULARITY FOR MULTIPLE MEMBRANES PROBLEM

ZHICHAO WANG AND XIN ZHOU

ABSTRACT. We prove the  $C^{1,1}$ -regularity for stationary  $C^{1,\alpha}$  ( $\alpha \in (0, 1)$ ) solutions to the multiple membrane problem. This regularity estimate was essentially used in our recent work on Yau's four minimal spheres conjecture [16].

## 1. INTRODUCTION

The *multiple membranes problem*, firstly studied by Vergara-Caffarelli [14], is to describe the equilibrium position of multiple membranes subject to forces and fixed boundary conditions, and constrained by the condition that they are not allowed to cross each other.

In [15], Vergara-Caffarelli proved  $C^{1,\alpha}$ -regularity for the two membranes that minimize the constant mean curvature functional. The regularity for minimizers of two-membranes problem was improved to be  $C^{1,1}$  by Silvestre [12] in a more general case; in particular, the optimal  $C^{1,1}$ -regularity holds true for prescribing mean curvature functional if the prescribing function does not change sign. The observation was that the height difference of the two membranes is a solution to the obstacle problem and then the classical work due to L. Caffarelli [2] can be applied. We also refer to [3, 4] for the two membranes problem for operators of different forms or fully nonlinear operators.

The problem with more than two membranes is more challenging. For the linear case, Chipot-Vergara-Caffarelli [5] proved the existence of the  $C^{1,\alpha}$  solution. Particularly, for the Laplacian operator, the regularity can be improved to be  $C^{1,1}$  by Savin-Yu [11]. For quasi-linear case, we refer to [1] for the  $p$ -Laplacian operators. In a recent work [10, §11], Sarnataro-Stryker outlined a proof for  $C^{1,1}$ -regularity for minimizers of multiple membranes problem for prescribing mean curvature functionals using [11, 12]; see Remark 1.3(iii) for more discussions.

In this paper, we prove the optimal  $C^{1,1}$ -regularity for all stationary  $C^{1,\alpha}$  solutions to the multiple membranes problem, satisfying a natural bounded first variation assumption; see Remark 1.1. The proof relies only on classical elliptic estimates.

**1.1. Notations and main results.** We consider a general second order elliptic operator of divergence form. For simplicity, we write

$$U_r := B_r(0) \times \mathbb{R} \times B_1(0).$$

Let  $\lambda \in (0, 1)$ ,  $\Lambda > 1$ ,  $\alpha \in (0, 1)$  be constants and  $\mathcal{A}(x, z, \mathbf{p}) \in C^3(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$ ,  $\mathcal{B}(x, z, \mathbf{p}) \in C^2(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n)$  satisfying

$$(1.1) \quad \Lambda |\boldsymbol{\xi}|^2 \geq (\partial_{\mathbf{p}} \mathcal{A})|_{U_2}(\boldsymbol{\xi}, \boldsymbol{\xi}) \geq \lambda |\boldsymbol{\xi}|^2, \quad \forall \boldsymbol{\xi} \in \mathbb{R}^n;$$

$$(1.2) \quad \|\mathcal{A}\|_{C^3(U_2)} + \|\mathcal{B}\|_{C^2(U_2)} \leq \Lambda;$$

$$(1.3) \quad \mathcal{A}(x, 0, 0) = 0, \quad \mathcal{B}(x, 0, 0) = 0.$$

Define an operator  $\mathcal{Q}$  by

$$\mathcal{Q}u := -\operatorname{div} \mathcal{A}(x, u, Du) + \mathcal{B}(x, u, Du).$$

Let  $f^1, \dots, f^n$  and  $g$  be measurable functions. Denote by  $\mathbf{f} = (f^1, \dots, f^n)$ . We say that  $\mathcal{Q}u = -\operatorname{div} \mathbf{f} + g$  in the *weak sense* in  $U \subset \mathbb{R}^n$  if for all  $\phi \in C_c^1(U)$ ,

$$\int_U \mathcal{A}(x, u, Du) D\phi + \mathcal{B}(x, u, Du)\phi - \mathbf{f} \cdot D\phi - g\phi \, dx = 0.$$

We call a collection of  $C^1$  functions  $\{u_j\}_{j=1}^m$  over a smooth domain  $U \subset \mathbb{R}^n$  *ordered graphs* if

$$u_1 \leq u_2 \leq \dots \leq u_m.$$

Let  $\{g_j\}_{j=1}^m$  be a family of measurable function. We say a collection of ordered graphs  $\{u_j\}_{j=1}^m$  is a *weak solution* to the equation

$$(1.4) \quad \sum_{j=1}^m (\mathcal{Q}u_j + g_j) = 0$$

in  $U$ , if for all  $\phi \in C_c^1(U)$ ,

$$\sum_{j=1}^m \int_U \mathcal{A}(x, u_j, Du_j) D\phi + \mathcal{B}(x, u_j, Du_j)\phi + g_j\phi \, dx = 0.$$

We assume that  $\{u_j\}$  satisfy two additional natural conditions:

(I) the *subsystem condition*: for any open set  $U' \subset U$  and  $1 \leq m_1 \leq m_2 \leq m$  so that

$$u_{m_1-1} < u_{m_1} \leq \dots \leq u_{m_2} < u_{m_2+1}, \quad (u_0 := -\infty, u_{m+1} := +\infty), \quad \forall x \in U',$$

we have

$$\sum_{j=m_1}^{m_2} (\mathcal{Q}u_j + g_j) = 0$$

in  $U'$  in the weak sense;

(II) the  $\kappa$ -*condition*: for each  $j = 1, \dots, m$ ,

$$-\kappa \leq \mathcal{Q}u_j \leq \kappa$$

in  $U$  in the weak sense; that is, for all  $\phi \in C_c^1(U)$  with  $\phi \geq 0$ ,

$$-\kappa \int_U \phi \, dx \leq \int_U \mathcal{A}(x, u_j, Du_j) D\phi + \mathcal{B}(x, u_j, Du_j)\phi \, dx \leq \kappa \int_U \phi \, dx.$$

*Remark 1.1.* In Section 1.2 below, we will introduce the connection between this PDE problem and a geometric isotopy energy minimizing problem. The subsystem condition (I) is a direct consequence when the system is only a stationary point for the prescribing mean curvature energy, while the  $\kappa$ -condition (II) is implied by assuming that each graph of  $\{u_j\}$  has bounded  $\kappa$ -first variation. Both conditions are satisfied by energy minimizers, but weak solutions satisfying (I) and (II) form a much more general class than just minimizers. For instance, both conditions are preserved by  $C^{1,\alpha}$  convergence, but the  $C^{1,\alpha}$ -limits are not necessarily isotopy energy minimizers. In particular, the  $C^{1,1}$ -regularity of these  $C^{1,\alpha}$ -limits play a crucial role in our recent work on Yau's four minimal spheres conjecture [16].

We now state our main  $C^{1,1}$ -regularity result.

**Theorem 1.2.** *Let  $\mathcal{Q}$  be an operator satisfying (1.1), (1.2) and (1.3). Let  $\{g_j\}_{j=1}^m$  be a collection of  $C^1$  functions over  $B_1(0)$ . Let  $\{u_j\}_{j=1}^m$  be ordered  $C^{1,\alpha}$  ( $\alpha \in (0,1)$ ) graphs over  $B_1(0) \subset \mathbb{R}^n$  which form a weak solution to (1.4) in  $B_1(0)$  and satisfy the subsystem condition (I) and  $\kappa$ -condition (II) ( $\kappa > 0$ ). Then  $u_j \in C_{loc}^{1,1}(B_1(0))$ . Moreover, if*

$$(1.5) \quad \kappa \leq 1, \quad \|g_j\|_{C^1(B_1(0))} \leq 1, \quad \|u_j\|_{C^{1,\alpha}(B_1(0))} \leq 1,$$

then

$$(1.6) \quad \sum_{j=1}^m \|u_j\|_{C^{1,1}(B_{1/2}(0))} \leq C \sum_{j=1}^m \left( \|u_j\|_{C^{1,\alpha}(B_1(0))} + \|g_j\|_{C^1(B_1(0))} + \kappa \right),$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$ .

*Remark 1.3.* (i) If each term in (1.5) is bounded by  $\Lambda'$ , then (1.6) still holds and the constant  $C$  will also depend on  $\Lambda'$ .

(ii) Our regularity improves Silvestre [12] even for

$$m = 2, \quad \mathcal{A}(x, z, \mathbf{p}) = \mathbf{p} / \sqrt{1 + |\mathbf{p}|^2}, \quad \mathcal{B} = 0,$$

since we don't require that  $g_2 > g_1$ .

(iii) Sarnataro-Stryker [10, Corollary 11.2] outlined a similar estimate for minimizers by virtue of free boundary problems [2] and methods in [11, 12]. Their result applied to arbitrary prescription functions, even though their proof relied on [12] which only holds for prescription functions which do not change sign. In this paper, we give a proof which holds for stationary solutions with arbitrary  $\{g_j\}$  while only using standard elliptic estimates in [6].

**1.2. Background from isotopy minimizing problem.** Fix a smooth open domain  $U \subset \mathbb{R}^n$ , a smooth curvature prescription function  $\bar{h} \in C^\infty(\bar{U} \times [-1, 1])$ , and a smooth Riemannian metric  $g$  in  $\bar{U} \times [-1, 1]$ . We also assume that  $U \times \{0\}$  is a minimal hypersurface in  $(\bar{U} \times [-1, 1], g)$ , so that the second condition of (1.3) is satisfied. Note that all discussion in the paper will be restricted to  $U \times (-1, 1)$ . Let  $\zeta(x, z)$  denote the  $(n+1)$ -dimensional volume element of  $g(x, z)$ . Then the area element w.r.t.  $g$  is a smooth function:

$$F : U \times (-1, 1) \times \mathbb{R}^n \rightarrow [0, \infty),$$

defined as follows: for any  $(x, z, \mathbf{p}) \in U \times (-1, 1) \times \mathbb{R}^n$ , we use  $P_{x,z,\mathbf{p}}$  to denote the  $n$ -dimensional parallelogram in  $\mathbb{R}^{n+1}$  generated by

$$\{e_1, \dots, e_n\}, \quad \text{where } e_i = \partial_{x_i} + \mathbf{p}^i \partial_z.$$

Then,

$$F(x, z, \mathbf{p}) = n\text{-volume of } P_{x,z,\mathbf{p}} \text{ under } g(x, z).$$

Given a  $C^1$ -function  $u : U \rightarrow \mathbb{R}$ , we define the generalized area as the following elliptic functional,

$$\text{Area}(\text{Graph}_u) = \int_U F(x, u, Du) \, dx.$$

The first variation formula of  $\text{Area}(\text{Graph}_u)$  w.r.t. variations  $t \mapsto \text{Graph}_{u+t\phi}$  for a fixed  $\phi \in C_c^1(\Omega)$  is given by

$$(1.7) \quad \delta \text{Area}_u(\phi) = \int_U \partial_{\mathbf{p}} F(x, u(x), Du(x)) \cdot D\phi + \partial_z F(x, u(x), Du(x)) \cdot \phi \, d\mathcal{H}^n(x).$$

Let  $\{u_j\}_{j=1}^m$  be a collection of ordered  $C^{1,\alpha}$ -graph functions over  $U \subset \mathbb{R}^n$ . Let  $\Sigma$  be the varifold induced by the graphs of  $\{u_j\}$  and  $\Omega \in \mathcal{C}(U \times (-1, 1))$  be a Caccioppoli set so that  $\partial\Omega = \sum_{j=1}^m (-1)^{j-1} \llbracket \text{Graph}_{u_j} \rrbracket$  in the sense of currents; see [13]. The  $\mathcal{A}^h$ -functional for the pair  $(\Sigma, \Omega)$ ,  $\mathcal{A}^h(\Sigma, \Omega) = \text{Area}(\Sigma) - \int_{\Omega} \hbar \, d\text{Vol}_g$ , defined in [16, (1.1)] (see also [10, 18]), can be written as

$$\mathcal{A}^h(\Sigma, \Omega) = \text{Area}(\text{Graph}_u) - \int_U \left( \sum_{j=1}^m (-1)^{j-1} \int_{-1}^{u_j(x)} \hbar(x, z) \cdot \zeta(x, z) \, dz \right) dx.$$

For simplicity, we will abuse the notation by writing

$$h(x, z) := \hbar(x, z) \cdot \zeta(x, z) \in C^\infty(\overline{U} \times [-1, 1]).$$

The first variation of  $\mathcal{A}^h(\Sigma, \Omega)$  w.r.t. variations  $t \mapsto \text{Graph}_{u+t\phi}$  is

$$\delta \mathcal{A}_{\Sigma, \Omega}^h(\phi) = \sum_{j=1}^m \int_U \partial_{\mathbf{p}} F(x, u_j, Du_j) \cdot D\phi + \partial_z F(x, u_j, Du_j) \cdot \phi + (-1)^j h(x, u_j) \phi \, dx,$$

for any  $\phi \in C_c^1(U)$ .

We say that  $(\Sigma, \Omega)$  is  $\mathcal{A}^h$ -stationary if for any one parameter family of diffeomorphisms  $\{\varphi_t\}_{-1 \leq t \leq 1}$  of  $U \times (-1, 1)$  satisfying that  $\varphi_t = \text{id}$  in a neighborhood of  $\partial U \times [-1, 1]$ , and  $\varphi_0 = \text{id}$ , then

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{A}^h(\varphi_t(\Sigma), \varphi_t(\Omega)) = 0.$$

Clearly, this implies that for any  $\phi \in C_c^1(U)$ ,

$$(1.8) \quad \sum_{j=1}^m \int_U \partial_{\mathbf{p}} F(x, u_j, Du_j) \cdot D\phi + \partial_z F(x, u_j, Du_j) \cdot \phi + (-1)^j h(x, u_j) \phi \, dx = 0.$$

Moreover, if a subcollection of graphs  $\{u_j\}_{j=m_1}^{m_2}$  do not touch other graphs, we may choose the variation  $\{\varphi_t\}$  to be supported away from other graphs, and then  $\{u_j\}_{j=m_1}^{m_2}$  will satisfy (1.8), which is exactly the subsystem condition (I).

The  $\kappa$ -condition (II) is natural in the sense that, if  $\text{Graph}_{u_j}$  has  $\kappa$ -bounded first variation in  $U$ , then by the first variation formula of the area functional, we have

$$-\kappa \leq -\text{div}_{\mathbf{p}} F(x, u_j, Du_j) + \partial_z F(x, u_j, Du_j) \leq \kappa$$

in the weak sense in  $U$ .

Applying Theorem 1.2 to  $\{u_j\}$ , we obtain the  $C^{1,1}$  regularity as follows.

**Theorem 1.4.** *Fix a constant  $\alpha \in (0, 1)$ . Assume that*

$$(1.9) \quad B_1(0) \times \{0\} \text{ is a minimal hypersurface in } (B_1(0) \times (-1, 1), g).$$

*Let  $\{u_j\}$  be ordered  $\mathcal{A}^h$ -stationary graphs over  $B_1(0)$ . Suppose that for all  $j = 1, \dots, m$ ,  $\text{Graph}_{u_j}$  has  $\kappa$ -bounded first variation. Then  $u_j \in C_{loc}^{1,1}(B_1(0))$ ,  $j = 1, \dots, m$ . Moreover, if*

$$\kappa \leq 1, \quad \|u_j\|_{C^{1,\alpha}(B_1(0))} \leq 1, \quad \|h\|_{C^1(B_1(0) \times [-1, 1])} \leq 1,$$

*then*

$$\sum_{j=1}^m \|u_j\|_{C^{1,1}(B_{1/2}(0))} \leq C \sum_{j=1}^m (\|u_j\|_{C^{1,\alpha}(B_1(0))} + \|h\|_{C^1(B_1(0) \times [-1, 1])} + \kappa),$$

*where  $C$  depends only on  $n, m, \alpha$  and the metric  $g$ .*

*Remark 1.5.* Consider  $\Sigma$  induced by ordered  $C^{1,\alpha}$ -graphs over  $U \subset \mathbb{R}^n$  such that each sheet has bounded first variation. Then by choosing a small enough neighborhood centered at any  $p \in \Sigma$ , and up to a change of coordinates therein, we can always assume (1.9) is satisfied. Indeed, assuming without loss of generality  $p \in U$ , by taking sufficiently small  $\epsilon$ ,  $B_\epsilon^n(p) \times [-\epsilon, \epsilon]$  admits a minimal foliation  $\{\Gamma_t\}$  with  $\partial\Gamma_t$  given by  $\partial B_\epsilon^n(p) \times \{t\}$ ; see [17, Appendix]. In particular, there exists a minimal slice  $\Gamma$  containing  $p$ . By taking sufficiently small  $\epsilon$ ,  $\Gamma$  is sufficiently close to  $B_\epsilon^n(p) \subset \mathbb{R}^n$ . Thus  $\Sigma$  can be written as ordered graphs over  $\Gamma$  with uniformly bounded  $C^{1,\alpha}$ -norm. Then the above theorem can be applied by rescaling  $\epsilon$  to 1.

**1.3. Idea of the proof.** Observe that by the  $\kappa$ -condition, for each  $j = 1, \dots, m$ ,  $Qu_j \in L^\infty$ . Then one can consider the equation of  $u_i - u_j$  and apply the weak Harnack inequality in [6, Theorem 8.17 and Theorem 8.18] to give the bound of  $u_i - u_j$  around the touching set. As a result, the difference of  $u_i$  and  $u_j$  is bounded by  $C\kappa r^2$ , where  $r$  is the distance to the touching set. Such a result together with the subsystem condition will imply the upper bound of the first derivative of  $u_i - u_j$ . These two inequalities will be crucial in the remaining of this paper.

The next step is to prove that  $u_j$  is  $W^{2,2}$ . To do this, we will use an inductive method to prove that the  $L^2$ -integrals of the corresponding difference quotients are uniformly bounded. Suppose that there are  $m$  sheets. Then for any points with density less than  $m$ , one can find a ball so that  $\{u_j\}$  is not connected. Then by the inductive process, one can bound the local  $L^2$ -integral of the difference quotients; see (3.28). On the other hand, for the points with density  $m$ , the first derivative upper bound in the previous paragraph will also give such a small ball with desired bound for the local  $L^2$ -integral of the difference quotients; see (3.29). Then by a covering argument, we will finish the induction and obtain the desired uniform  $L^2$  bound.

After that, we will consider the equation of  $D_k\varphi$ , where  $\varphi$  is the average of  $u_j$ . Thanks to the  $C^1$  estimates of  $u_i - u_j$ , we can adapt the Hölder estimates in [7, Theorem 3.8] to prove that the  $L^2$ -integral of  $D_k\varphi$  over  $B_r(y)$  has an order  $n - \epsilon$  for any  $\epsilon \in (0, 1)$ ; see Lemma 4.2 and Proposition 4.4. Then the argument of Hölder estimates for gradients (see [7, Theorem 3.13]) will be adapted to improve the order estimates of

$$\int_{B_r(y)} |D^2\varphi - (D^2\varphi)_{B_r(y)}|^2 dx.$$

Note that the above two estimates are only valid for the radius  $r$  so that  $\{u_j\}$  is connected in  $B_r(y)$  since we have to use the  $C^1$  estimates of  $u_i - u_j$ . Nevertheless, such an order estimate will be applied to give two delicate estimates (see Lemma 4.6 and Lemma 4.7), which yield a uniform upper bound for

$$\sum_{j=1}^m \frac{1}{|B_r|^2} \int_{B_r(y)} |D^2u_j|^2 dx.$$

This together with the Poincaré inequalities implies the growth estimates of  $u_j$  over a tilt-plane. Such a bound enables us to apply the  $C^{1,\alpha}$  estimates [6, Theorem 8.32] to bound the derivative of  $u_j$  over the tilt-plane, i.e. we obtain the bound of  $|Du_j - (Du_j)_{B_r(y)}|$  for all  $x \in B_r(y)$ . This gives the desired upper bound of  $|Du_j(x) - Du_j(y)|$  by taking  $r = |x - y|$ . Then the main theorem is proved.

**1.4. Outline.** This paper is organized as follows. In Section 2, we introduce some notions and basic inequalities for the coefficients of the equations. Then we will list some variants of elliptic PDEs that will be used in this paper. In Section 3, we give the  $C^1$  norm of the difference of any two sheets. As a consequence, the  $W^{2,2}$  norm of each  $u_j$  is also proved. Section 4 is devoted to

prove the order estimates of the Hessians of  $u_j$ , which imply that  $D^2u_j$  is bounded in the sense of average. Finally, we prove the desired Lipschitz upper bound for  $Du_j$  in Section 5.

**Acknowledgement.** Z. W. would like to thank Professor Jingyi Chen and Professor Ailana Fraser for their support and encouragement. X. Z. is supported by NSF grant DMS-1945178, an Alfred P. Sloan Research Fellowship, and a grant from the Simons Foundation (1026523, XZ).

## 2. PRELIMINARY

In this section, we will derive a number of differential equations to be used later and provide some basic estimates for the coefficients.

**2.1. Some estimates of the coefficients.** We always assume that there exist  $\alpha \in (0, 1)$  and  $\delta_0, \kappa_0, \kappa_1 \in (0, 1]$ , so that for all  $i, j = 1, \dots, m$ , the ordered graphs  $\{u_j\}$  satisfy:

$$(2.1) \quad \sup_{x \in U} |Du_j(x)| + \sup_{x, y \in U} \frac{|Du_j(x) - Du_j(y)|}{|x - y|^\alpha} \leq \delta_0;$$

$$(2.2) \quad |g_j| \leq \kappa_0, \quad |Dg_j| \leq \kappa_1.$$

For simplicity, we denote

$$\kappa^* := \kappa^2 + \kappa_0^2 + \kappa_1^2 + \delta_0^2.$$

**Definition 2.1.** Let  $\mathcal{A}(x, z, \mathbf{p}) \in C^3(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$ ,  $\mathcal{B}(x, z, \mathbf{p}) \in C^2(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n)$  and  $\mathcal{Q}u := -\operatorname{div} \mathcal{A}(x, u, Du) + \mathcal{B}(x, u, Du)$ . For a family of  $C^1$  functions  $\{g_j\}_{j=1}^m$ , we say that  $C^{1, \alpha}$  ordered graphs  $\{u_j\}_{j=1}^m$  over  $U \subset \mathbb{R}^n$  form a solution to  $(\mathcal{Q}, \{g_j\})$  if  $\{u_j\}$  satisfy (1.4) in  $U$  in the weak sense, the subsystem condition (I), and the  $\kappa$ -condition (II).

The following will be used in various blow-up arguments.

**Lemma 2.2.** *Suppose that the ordered collection  $\{u_j\}$  is a solution to  $(\mathcal{Q}, \{g_j\})$  in  $U$ . Let  $y \in \mathbb{R}^n$ ,  $r \in (0, 1]$  and define  $\{v_j\}$  by*

$$v_j(x) = r^{-1}u_j(y + rx).$$

Define a new operator  $\tilde{\mathcal{Q}}$  by

$$\tilde{\mathcal{Q}}v = -\operatorname{div} \tilde{\mathcal{A}}(x, v, Dv) + \tilde{\mathcal{B}}(x, v, Dv),$$

where

$$\tilde{\mathcal{A}}(x, z, \mathbf{p}) := \mathcal{A}(y + rx, rz, \mathbf{p}), \quad \tilde{\mathcal{B}}(x, z, \mathbf{p}) := r\mathcal{B}(y + rx, rz, \mathbf{p}).$$

Define the new functions  $\{\tilde{g}_j\}$  by

$$\tilde{g}_j(x) = rg_j(y + rx).$$

Then  $\{v_j\}$  is a solution to  $(\tilde{\mathcal{Q}}, \{\tilde{g}_j\})$  in  $\tilde{U} := \{r^{-1}(x - y); x \in U\}$ .

*Remark 2.3.* We now describe the changes of  $\delta_0, \kappa_0, \kappa_1$  and  $\kappa$  in the blow-up process. Suppose that in Lemma 2.2, if (2.1) and (2.2) hold for  $\delta_0, \kappa_0, \kappa_1$ . Then we have that

$$\sup_{x \in \tilde{U}} |Dv_j(x)| \leq \delta_0, \quad \sup_{x, x' \in \tilde{U}} \frac{|Dv_j(x) - Dv_j(x')|}{|x - x'|^\alpha} \leq r^\alpha \delta_0;$$

$$|\tilde{g}_j| \leq r\kappa_0, \quad |D\tilde{g}_j| \leq r^2\kappa_1.$$

Moreover, if  $\{u_j\}$  satisfy  $\kappa$ -condition, then  $\{v_j\}$  satisfy  $r\kappa$ -condition. Without loss of generality, we always assume that  $0 < \kappa \leq 1$ .

We now introduce the concept of connected components. Note that each connected component is a solution to  $(\mathcal{Q}, \{g'_j\})$  for the associated  $\{g'_j\} \subset \{g_j\}$  by the subsystem condition.

**Definition 2.4** (Connected component for ordered graphs). Let  $\{u_j\}_{j=1}^m$  be ordered graphs over  $U$ . We say they are connected in  $U$  if and only if for each  $i = 1, \dots, m-1$ , there exists  $x_i \in U$  such that  $u_i(x_i) = u_{i+1}(x_i)$ .

Any ordered graphs  $\{u_j\}$  over  $U$  can be divided into subcollections  $\{u_1, \dots, u_{i_1}\}, \{u_{i_1+1}, \dots, u_{i_2}\}, \dots, \{u_{i_k}, \dots, u_m\}$  so that each collection is connected and the collections are pairwise disjoint; each collection is called a *connected component*.

**Lemma 2.5.** Let  $\{u_j\}_{j=1}^m$  be ordered graphs over  $B_1(0)$ . Given  $t \in (0, 1)$ , then there exists  $r \in [t^m, 1]$  so that  $\{u_j\}$  has the same number of connected components in  $B_r(0)$  and  $B_{tr}(0)$ .

*Proof.* Note that the number of connected components of the system  $\{u_j\}$  in  $B_r(0)$  is decreasing when  $r$  increases. Let  $N(k)$  be the number of connected components of  $\{u_j\}$  in  $B_{r_k}(0)$  for  $r_k := t^k$ . Then the desired result follows from the fact that

$$1 \leq N(1) \leq N(2) \leq \dots \leq N(k) \leq m.$$

□

Let  $u, v$  be two  $C^1$ -functions on  $U$ . When taking the difference of derivatives of  $\mathcal{A}$  and  $\mathcal{B}$ , we will estimate the remainder in terms of the following notations.

$$\begin{aligned} \mathbf{A}(u, v) &= \int_0^1 \partial_{\mathbf{p}} \mathcal{A}(x, tu + (1-t)v, tDu + (1-t)Dv) dt; \\ \mathbf{b}(u, v) &= \int_0^1 \partial_z \mathcal{A}(x, tu + (1-t)v, tDu + (1-t)Dv) dt; \\ \mathbf{c}(u, v) &= \int_0^1 \partial_{\mathbf{p}} \mathcal{B}(x, tu + (1-t)v, tDu + (1-t)Dv) dt; \\ \mathbf{d}(u, v) &= \int_0^1 \partial_z \mathcal{B}(x, tu + (1-t)v, tDu + (1-t)Dv) dt. \end{aligned}$$

Note that by definition, we have

$$\begin{aligned} \mathbf{A}(u, u) &= \partial_{\mathbf{p}} \mathcal{A}(x, u, Du), & \mathbf{b}(u, u) &= \partial_z \mathcal{A}(x, u, Du); \\ \mathbf{c}(u, u) &= \partial_{\mathbf{p}} \mathcal{B}(x, u, Du), & \mathbf{d}(u, u) &= \partial_z \mathcal{B}(x, u, Du). \end{aligned}$$

**Lemma 2.6.** The following formulas follow from a direct computation.

$$\begin{aligned} \mathcal{A}(x, u, Du) - \mathcal{A}(x, v, Dv) &= \mathbf{A}(u, v) \cdot D(u - v) + \mathbf{b}(u, v)(u - v); \\ \mathcal{B}(x, u, Du) - \mathcal{B}(x, v, Dv) &= \mathbf{c}(u, v) \cdot D(u - v) + \mathbf{d}(u, v)(u - v). \end{aligned}$$

The following estimates will be used later.

**Lemma 2.7.** Let  $u_1, u_2, v_1, v_2$  be  $C^1$  functions defined on  $U$ . Then

$$(2.3) \quad \begin{aligned} |\mathbf{A}(u_1, u_2) - \mathbf{A}(v_1, v_2)| + |\mathbf{b}(u_1, u_2) - \mathbf{b}(v_1, v_2)| &\leq \|\mathcal{A}\|_{C^2} (\|u_1 - v_1\|_{C^1} + \|u_2 - v_2\|_{C^1}); \\ |\mathbf{c}(u_1, u_2) - \mathbf{c}(v_1, v_2)| + |\mathbf{d}(u_1, u_2) - \mathbf{d}(v_1, v_2)| &\leq \|\mathcal{B}\|_{C^2} (\|u_1 - v_1\|_{C^1} + \|u_2 - v_2\|_{C^1}). \end{aligned}$$

**2.2. The variants of differential equations.** Since a solution to  $(\mathcal{Q}, \{g_j\})$  is in the form of  $\{u_j\}$ , we can not apply the theory for elliptic PDEs. In this section, we will derive some PDEs satisfied by a single function which can be the average, the derivative, or the difference quotient, etc.

**The equation of the average.** Suppose that  $\{u_j\}$  is a solution to  $(\mathcal{Q}, \{g_j\})$ ; see Definition 2.1. Let

$$\varphi := \frac{1}{m} \sum_{j=1}^m u_j.$$

By Lemma 2.6,

$$\begin{aligned} \mathcal{Q}u_j - \mathcal{Q}\varphi &= -\operatorname{div}\left(\mathbf{A}(u_j, \varphi)D(u_j - \varphi) + \mathbf{b}(u_j, \varphi)(u_j - \varphi)\right) + \\ &\quad + \mathbf{c}(u_j, \varphi)D(u_j - \varphi) + d(u_j, \varphi)(u_j - \varphi). \end{aligned}$$

Observe that

$$m\mathcal{Q}\varphi = \sum_{j=1}^m (\mathcal{Q}\varphi - \mathcal{Q}u_j - g_j).$$

Then  $\varphi$  satisfies the following equation:

$$(2.4) \quad m\mathcal{Q}\varphi = \operatorname{div} \mathbf{f}(\{u_j\}) - g(\{u_j\}, \{g_j\}).$$

Here

$$(2.5) \quad \mathbf{f}(\{u_j\}) := \sum_{j=1}^m \left[ (\mathbf{A}(u_j, \varphi) - \mathbf{A}(\varphi, \varphi))(Du_j - D\varphi) + (\mathbf{b}(u_j, \varphi) - \mathbf{b}(\varphi, \varphi))(u_j - \varphi) \right];$$

$$(2.6) \quad g(\{u_j\}, \{g_j\}) := \sum_{j=1}^m \left[ (\mathbf{c}(u_j, \varphi) - \mathbf{c}(\varphi, \varphi))(Du_j - D\varphi) + (d(u_j, \varphi) - d(\varphi, \varphi))(u_j - \varphi) - g_j \right].$$

Note that in (2.5) and (2.6), the additional terms containing  $\mathbf{A}(\varphi, \varphi)$ ,  $\mathbf{b}(\varphi, \varphi)$ ,  $\mathbf{c}(\varphi, \varphi)$  and  $d(\varphi, \varphi)$  sum to be zero, since  $\sum_{j=1}^m (\varphi - u_j) = 0$ .

**Lemma 2.8.** *Let  $\{u_j\}$  and  $\varphi$  be as above. Let  $\tau > 0$  be a constant so that  $\|u_i - u_j\|_{C^1(B_1(0))} \leq \tau$ . Then*

$$\begin{aligned} |\mathbf{f}(\{u_j\})| &\leq C(m, \Lambda)\tau^2, \quad \sup_{x, y \in B_1(0)} \frac{|\mathbf{f}(\{u_j\})(x) - \mathbf{f}(\{u_j\})(y)|}{|x - y|^\alpha} \leq C(m, \Lambda)\tau; \\ |g(\{u_j\}, \{g_j\})| &\leq C(m, \Lambda)(\tau^2 + \kappa_0). \end{aligned}$$

*Proof.* It follows from Lemma 2.7, equation (2.1) and (2.2).  $\square$

**The equation of differences of two solutions.** Let  $\{u_j\}_{j=1}^m$  and  $\{v_i\}_{i=1}^{m'}$  be two ordered collections of graphs over  $U$ . Suppose that  $\{u_j\}_{j=1}^m$  and  $\{v_i\}_{i=1}^{m'}$  are solutions to  $(\mathcal{Q}, \{g_j\})$  and  $(\mathcal{Q}, \{g'_i\})$ , respectively. Denote by

$$\varphi := \frac{1}{m} \sum_{j=1}^m u_j, \quad \psi := \frac{1}{m'} \sum_{i=1}^{m'} v_i.$$



Then both  $\varphi$  and  $\psi$  satisfy the equation (2.4). We can write them as

$$\begin{aligned} m\mathcal{Q}\varphi &= \operatorname{div} \mathbf{f}(\{u_j\}) - g(\{u_j\}, \{g_j\}); \\ m'\mathcal{Q}\psi &= \operatorname{div} \mathbf{f}(\{v_i\}) - g(\{v_i\}, \{g'_j\}). \end{aligned}$$

Taking the difference and using Lemma 2.6, we obtain the equation for  $\Psi := \varphi - \psi$  as follows:

$$(2.7) \quad \begin{aligned} & -\operatorname{div} \left( \mathbf{A}(\varphi, \psi) D\Psi + \mathbf{b}(\varphi, \psi) \Psi \right) + \mathbf{c}(\varphi, \psi) D\Psi + d(\varphi, \psi) \Psi \\ &= \operatorname{div} \left( \frac{1}{m} \mathbf{f}(\{u_j\}) - \frac{1}{m'} \mathbf{f}(\{v_i\}) \right) - \left( \frac{1}{m} g(\{u_j\}, \{g_j\}) - \frac{1}{m'} g(\{v_i\}, \{g'_j\}) \right). \end{aligned}$$

**The equation of difference quotients.** Given a function  $f$  and a real number  $\tau \neq 0$ , let

$$f^\tau(x) := \frac{f(x + \tau \mathbf{e}) - f(x)}{\tau},$$

where  $\mathbf{e}$  can be any unit vector in  $\mathbb{R}^n$ . For simplicity, denote by

$$\begin{aligned} \mathbf{A}_\tau(u) &= \int_0^1 \partial_{\mathbf{p}} \mathcal{A}(x + t\tau \mathbf{e}, u(x) + t\tau \cdot u^\tau(x), Du(x) + t\tau \cdot Du^\tau(x)) dt; \\ \mathbf{b}_\tau(u) &= \int_0^1 \partial_z \mathcal{A}(x + t\tau \mathbf{e}, u(x) + t\tau \cdot u^\tau(x), Du(x) + t\tau \cdot Du^\tau(x)) dt; \\ \mathbf{c}_\tau(u) &= \int_0^1 \partial_{\mathbf{p}} \mathcal{B}(x + t\tau \mathbf{e}, u(x) + t\tau \cdot u^\tau(x), Du(x) + t\tau \cdot Du^\tau(x)) dt; \\ d_\tau(u) &= \int_0^1 \partial_z \mathcal{B}(x + t\tau \mathbf{e}, u(x) + t\tau \cdot u^\tau(x), Du(x) + t\tau \cdot Du^\tau(x)) dt; \\ \mathfrak{A}_\tau(u) &= \int_0^1 \partial_{x_k} \mathcal{A}(x + t\tau \mathbf{e}, u(x) + t\tau \cdot u^\tau(x), Du(x) + t\tau \cdot Du^\tau(x)) dt; \\ \mathfrak{B}_\tau(u) &= \int_0^1 \partial_{x_k} \mathcal{B}(x + t\tau \mathbf{e}, u(x) + t\tau \cdot u^\tau(x), Du(x) + t\tau \cdot Du^\tau(x)) dt. \end{aligned}$$

where we assume that  $\mathbf{e}$  is the direction of the  $k$ -th coordinate. Then we have that

$$\begin{aligned} \left[ \mathcal{A}(x, u, Du) \right]^\tau &= \mathbf{A}_\tau(u) Du^\tau + \mathbf{b}_\tau(u) u^\tau + \mathfrak{A}_\tau(u), \\ \left[ \mathcal{B}(x, u, Du) \right]^\tau &= \mathbf{c}_\tau(u) Du^\tau + d_\tau(u) u^\tau + \mathfrak{B}_\tau(u). \end{aligned}$$

The following estimates will be used later.

**Lemma 2.9.** *Let  $u, v$  be two  $C^1$  functions. Then we have that for  $x \in U$  with  $B_\tau(x) \subset U$ ,*

$$\begin{aligned} |\mathbf{A}_\tau(u) - \mathbf{A}_\tau(v)| &\leq \|\partial_{\mathbf{p}} \mathcal{A}\|_{C^1} \cdot \|u - v\|_{C^1}; & |\mathbf{b}_\tau(u) - \mathbf{b}_\tau(v)| &\leq \|\partial_z \mathcal{A}\|_{C^1} \cdot \|u - v\|_{C^1}; \\ |\mathbf{c}_\tau(u) - \mathbf{c}_\tau(v)| &\leq \|\partial_{\mathbf{p}} \mathcal{B}\|_{C^1} \cdot \|u - v\|_{C^1}; & |d_\tau(u) - d_\tau(v)| &\leq \|\partial_z \mathcal{B}\|_{C^1} \cdot \|u - v\|_{C^1}; \\ |\mathfrak{A}_\tau(u) - \mathfrak{A}_\tau(v)| &\leq \|\partial_x \mathcal{A}\|_{C^1} \cdot \|u - v\|_{C^1}; & |\mathfrak{B}_\tau(u) - \mathfrak{B}_\tau(v)| &\leq \|\partial_x \mathcal{B}\|_{C^1} \cdot \|u - v\|_{C^1}. \end{aligned}$$

Here we omit the domain  $U \times \mathbb{R} \times B_1(0)$  of the  $C^1$  norms for simplification.

Let  $\{u_j\}$  be ordered graphs over  $U \subset \mathbb{R}^n$  and a solution to  $(\mathcal{Q}, \{g_j\})$ . Then for all  $\phi \in C_c^1(B_1(0))$ , we have

$$\sum_{j=1}^m \int_U \left[ \mathcal{A}(x, u_j, Du_j) \right]^\tau \cdot D\phi + \left[ \mathcal{B}(x, u_j, Du_j) \right]^\tau \cdot \phi + g_j^\tau \phi \, dx = 0.$$

This can be rewritten as

$$\sum_j \left( -\operatorname{div}(\mathbf{A}_\tau(u_j)Du_j^\tau + \mathbf{b}_\tau(u_j)u_j^\tau + \mathfrak{A}_\tau(u_j)) + \mathbf{c}_\tau(u_j)Du_j^\tau + d_\tau(u_j)u_j^\tau + \mathfrak{B}_\tau(u_j) + g_j^\tau \right) = 0,$$

Then we obtain the equation of  $\varphi^\tau$  as

$$(2.8) \quad \sum_{j=1}^m \left[ -\operatorname{div}(\mathbf{A}_\tau(u_j)D\varphi^\tau + \mathbf{b}_\tau(u_j)\varphi^\tau) + \mathbf{c}_\tau(u_j)D\varphi^\tau + d_\tau(u_j)\varphi^\tau \right] = -\operatorname{div}\mathbf{f}_\tau + g_\tau,$$

where

$$(2.9) \quad \mathbf{f}_\tau := \sum_{j=1}^m \left[ (\mathbf{A}_\tau(u_j) - \mathbf{A}_\tau(\varphi))(D\varphi^\tau - Du_j^\tau) + (\mathbf{b}_\tau(u_j) - \mathbf{b}_\tau(\varphi))(\varphi^\tau - u_j^\tau) - \mathfrak{A}_\tau(u_j) \right];$$

$$(2.10) \quad g_\tau := \sum_{j=1}^m \left[ (\mathbf{c}_\tau(u_j) - \mathbf{c}_\tau(\varphi))(D\varphi^\tau - Du_j^\tau) + (d_\tau(u_j) - d_\tau(\varphi))(\varphi^\tau - u_j^\tau) - \mathfrak{B}_\tau(u_j) - g_j^\tau \right].$$

Recall that by (1.3),

$$\int_0^1 \partial_{x_k} \mathcal{A}(x + t\tau\mathbf{e}, 0, 0) dt = 0.$$

By the mean value theorem,

$$(2.11) \quad \begin{aligned} & |\mathfrak{A}_\tau(u_j)| \\ & \leq \int_0^1 \left| \partial_{x_k} \mathcal{A}(x + t\tau\mathbf{e}, u_j(x) + t\tau \cdot u_j^\tau(x), Du_j(x) + t\tau \cdot Du_j^\tau(x)) - \partial_{x_k} \mathcal{A}(x + t\tau\mathbf{e}, 0, 0) \right| dt \\ & \leq \|\partial_z \partial_x \mathcal{A}\|_{C^0(B_1(0))} \left( |u_j(x)| + |u_j(x + \tau\mathbf{e})| \right) + \|\partial_{\mathbf{p}} \partial_x \mathcal{A}\|_{C^0(B_1(0))} \left( |Du_j(x)| + |Du_j(x + \tau\mathbf{e})| \right). \end{aligned}$$

Since  $\mathcal{B}(x, 0, 0) = 0$ , we have  $\mathfrak{B}_\tau(0) = 0$ . Then Lemma 2.9 gives that

$$(2.12) \quad |\mathfrak{B}_\tau(u_j)| \leq \|\partial_x \mathcal{B}\|_{C^1} \|u_j\|_{C^1}.$$

Moreover, we obtain the bound of  $\mathbf{f}_\tau$  and  $g_\tau$  by Lemma 2.9.

**Lemma 2.10.** *With the notions as above, there exists  $C = C(n, m, \Lambda)$  so that for all  $x \in B_{1-\tau}(0)$ ,*

$$\begin{aligned}
|\mathbf{f}_\tau| &\leq C \sum_{j=1}^m \left( \|\partial_{\mathbf{p}} \mathcal{A}\|_{C^1} \|u_j - \varphi\|_{C^1} \cdot |D\varphi^\tau - Du_j^\tau| + \|\partial_z \mathcal{A}\| \cdot \|u_j - \varphi\|_{C^1} \cdot |\varphi^\tau - u_j^\tau| + |\mathfrak{A}_\tau(u_j)| \right) \\
&\leq C \sum_{j=1}^m \left( \|u_j - \varphi\|_{C^1} \cdot |D\varphi^\tau - Du_j^\tau| + \|u_j - \varphi\|_{C^1} \cdot |\varphi^\tau - u_j^\tau| + \|\partial_z \partial_x \mathcal{A}\|_{C^0(B_1(0))} |u_j|_{C^0} + \right. \\
&\quad \left. + \|\partial_{\mathbf{p}} \partial_x \mathcal{A}\|_{C^0(B_1(0))} \cdot |Du_j|_{C^0} \right); \\
|g_\tau| &\leq C \sum_{j=1}^m \left( \|\partial_{\mathbf{p}} \mathcal{B}\|_{C^1} \|u_j - \varphi\|_{C^1} \cdot |D\varphi^\tau - Du_j^\tau| + \|\partial_z \mathcal{B}\|_{C^1} \cdot \|u_j - \varphi\|_{C^1} \cdot |\varphi^\tau - u_j^\tau| + \right. \\
&\quad \left. + |\mathfrak{B}_\tau(u_j)| + |g_j^\tau| \right) \\
&\leq C \sum_{j=1}^m \left( \|u_j - \varphi\|_{C^1} \cdot |D\varphi^\tau - Du_j^\tau| + \|u_j - \varphi\|_{C^1} \cdot |\varphi^\tau - u_j^\tau| + \|\partial_x \mathcal{B}\|_{C^1} \|u_j\|_{C^1} + \right. \\
&\quad \left. + |Dg_j|_{C^0} \right).
\end{aligned}$$

**The equation of the first derivatives.** Let  $\{u_j\}$  be ordered graphs over  $U \subset \mathbb{R}^n$  and a solution to  $(\mathcal{Q}, \{g_j\})$ . If  $\{u_j\} \subset W^{2,2}(U)$ , by letting  $\tau \rightarrow 0$  in (2.8) we have,

$$(2.13) \quad -\operatorname{div} \left( \mathbf{A}^* DD_k \varphi + \mathbf{b}^* D_k \varphi \right) + \mathbf{c}^* DD_k \varphi + d^* D_k \varphi = -\operatorname{div} \left( \hat{\mathbf{f}} - m \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) \right) + \hat{g},$$

where

$$\begin{aligned}
\mathbf{A}^*(x) &:= \sum_{j=1}^m \partial_{\mathbf{p}} \mathcal{A}(x, u_j(x), Du_j(x)); & \mathbf{b}^*(x) &:= \sum_{j=1}^m \partial_z \mathcal{A}(x, u_j(x), Du_j(x)); \\
\mathbf{c}^*(x) &:= \sum_{j=1}^m \partial_{\mathbf{p}} \mathcal{B}(x, u_j(x), Du_j(x)); & d^*(x) &:= \sum_{j=1}^m \partial_z \mathcal{B}(x, u_j(x), Du_j(x));
\end{aligned}$$

and

$$(2.14) \quad \hat{\mathbf{f}} := \sum_{j=1}^m \left[ (\mathbf{A}(u_j, u_j) - \mathbf{A}(\varphi, \varphi))(DD_k \varphi - DD_k u_j) + (\mathbf{b}(u_j, u_j) - \mathbf{b}(\varphi, \varphi))(D_k \varphi - D_k u_j) - (\partial_{x_k} \mathcal{A}(x, u_j, Du_j) - \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi)) \right];$$

$$(2.15) \quad \hat{g} := \sum_{j=1}^m \left[ (\mathbf{c}(u_j, u_j) - \mathbf{c}(\varphi, \varphi))(DD_k \varphi - DD_k u_j) + (d(u_j, u_j) - d(\varphi, \varphi))(D_k \varphi - D_k u_j) - \partial_{x_k} \mathcal{B}(x, u_j, Du_j) - D_k g_j \right].$$

Since  $\mathcal{B}(x, 0, 0) = 0$ , then  $\partial_{x_k} \mathcal{B}(x, 0, 0) = 0$ . By the mean value theorem,

$$(2.16) \quad |\partial_{x_k} \mathcal{B}(x, u_j, Du_j)| \leq \|\partial_{x_k} \mathcal{B}\|_{C^1} \left( |u_j(x)| + |Du_j(x)| \right).$$

**Lemma 2.11.** *Suppose that  $U = B_R(0)$ ,  $R \in (0, 1]$  and  $\{u_j\} \subset W^{2,2}(B_R(0))$ . Then there exists  $C = C(n, m, \Lambda)$  such that*

$$\begin{aligned} \int_{B_R(0)} |\hat{\mathbf{f}}|^2 dx &\leq C \sum_{j=1}^m \int_{B_R(0)} \|u_j - \varphi\|_{C^1(B_R(0))}^2 \left( |DD_k \varphi - DD_k u_j|^2 + |D\varphi - Du_j|^2 + 1 \right) dx; \\ \int_{B_R(0)} |\hat{g}| dx &\leq C \sum_{j=1}^m \int_{B_R(0)} \left( \|u_j - \varphi\|_{C^1(B_R(0))} (|DD_k \varphi - DD_k u_j| + |D\varphi - Du_j|) + \right. \\ &\quad \left. + \|\partial_x \mathcal{B}\|_{C^1} (|u_j(x)| + |Du_j(x)|) + |Dg_j|_{C^0(B_1(0))} \right) dx; \\ \int_{B_R(0)} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) D\phi dx &\leq C \int_{B_R(0)} \left( \|\partial_x \partial_x \mathcal{A}\|_{C^1(U_R)} |\varphi| |\phi| + |D\varphi| |\phi| + |D\varphi - D\varphi(0)| |D\phi| \right. \\ &\quad \left. + (R + |\varphi - \varphi(0)| + |D\varphi - D\varphi(0)|) |D^2 \varphi| |\phi| \right) dx, \end{aligned}$$

for all  $\phi \in C_c(B_R(0))$ .

*Proof.* The first one follows from Lemma 2.7. The same argument together with (2.16) gives the second inequality.

We now prove the third inequality. By a direct computation,

$$\begin{aligned} \operatorname{div} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) &= \sum_{i=1}^n \partial_{x_i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) + \partial_z \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) D\varphi + \\ &\quad + \sum_{i=1}^n \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) DD_i \varphi. \end{aligned}$$

Then we have that

$$\begin{aligned} (2.17) \quad &\int_{B_R(0)} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) \cdot D\phi dx = - \int_{B_R(0)} \operatorname{div} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) \cdot \phi dx \\ &\leq C \int_{B_R(0)} \|\partial_x \partial_x \mathcal{A}\|_{C^1(U_R)} (|\varphi| + |D\varphi|) |\phi| + \|\partial_z \partial_x \mathcal{A}\|_{C^0(U_R)} |D\varphi| |\phi| dx + \\ &\quad + \sum_{i=1}^n \left| \int_{B_R(0)} \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) DD_i \varphi \cdot \phi dx \right|. \end{aligned}$$

Observe that

$$\begin{aligned} \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) DD_i \varphi &= \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(0, \varphi(0), D\varphi(0)) D(D_i \varphi - D_i \varphi(0)) + \\ &\quad + \left( \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) - \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(0, \varphi(0), D\varphi(0)) \right) DD_i \varphi \\ &= \operatorname{div} \left( \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(0, \varphi(0), D\varphi(0)) (D_i \varphi - D_i \varphi(0)) \right) + \\ &\quad + \left( \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) - \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(0, \varphi(0), D\varphi(0)) \right) DD_i \varphi. \end{aligned}$$

This implies that

$$\begin{aligned}
& \left| \int_{B_R(0)} \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) DD_i \varphi \cdot \phi \, dx \right| \\
& \leq \left| \int_{B_R(0)} \left( \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(0, \varphi(0), D\varphi(0)) (D_i \varphi - D_i \varphi(0)) \right) D\phi \, dx \right| + \\
& + \left| \int_{B_R(0)} \left( \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) - \partial_{\mathbf{p}^i} \partial_{x_k} \mathcal{A}(0, \varphi(0), D\varphi(0)) \right) DD_i \varphi \cdot \phi \, dx \right| \\
& \leq C \int_{B_R(0)} |D\varphi - D\varphi(0)| \cdot |D\phi| + (R + |\varphi - \varphi(0)| + |D\varphi - D\varphi(0)|) |D^2 \varphi| |\phi| \, dx.
\end{aligned}$$

This together with (2.17) gives the desired inequality.  $\square$

### 3. THE HEIGHT DIFFERENCE ESTIMATES

In this section, we consider the height difference of two sheets. By the  $\kappa$ -condition, the height difference is a subsolution and a supersolution. Then the weak Harnack inequality gives the growth order around touching points. Then using the inductive methods, we can bound the first derivative of the difference by the distance to the touching sets. Such an estimate is the main result in Section 3.1 and will be crucial whenever we consider a differential equation because all of the remainder terms contain the height differences.

Then in Section 3.2, we first consider the difference quotient of the height difference and then give the  $W^{2,2}$  estimates. Finally, we obtain the uniform  $W^{2,2}$  estimates for each sheet.

**3.1. The first derivatives of the height difference.** Recall that for any  $A \subset \mathbb{R}^n$  and  $x \in \mathbb{R}^n$ ,

$$\text{dist}(x, A) := \inf_{y \in A} \text{dist}(x, y).$$

We define  $\text{dist}(x, \emptyset) = +\infty$ .

**Lemma 3.1.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs over  $B_1(0) \subset \mathbb{R}^n$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (2.1) and (2.2) are satisfied. Then for each  $x_0 \in B_1(0)$  and  $r > 0$  with  $u_{i+1}(x_0) = u_i(x_0)$  for some  $i \in \{1, \dots, m-1\}$  and  $B_{4r}(x_0) \subset B_1(0)$ , we have that*

$$|u_{i+1}(x) - u_i(x)| \leq C \cdot \kappa r^2, \quad \text{for each } x \in B_r(x_0),$$

where  $C$  depends on  $n, \Lambda/\lambda$ .

*Proof.* Recall by the  $\kappa$ -condition in Definition 2.1, for  $j = 1, \dots, m$ ,

$$-\kappa \leq -\text{div} \left( \mathcal{A}(x, u_j, Du_j) \right) + \mathcal{B}(x, u_j, Du_j) \leq \kappa.$$

Let  $w = u_{i+1} - u_i$ . Then by Lemma 2.6,

$$-2\kappa \leq -\text{div} \left( \mathbf{A}(u_{i+1}, u_i) Dw + \mathbf{b}(u_{i+1}, u_i) w \right) + \mathbf{c}(u_{i+1}, u_i) Dw + d(u_{i+1}, u_i) w \leq 2\kappa$$

in the weak sense. Applying weak Harnack inequality [6, Theorem 8.17 and Theorem 8.18] with  $q = n + 1$ , then we have

$$\sup_{x \in B_r(x_0)} w \leq C \left( \inf_{x \in B_r(x_0)} w + r^{2-\frac{2n}{q}} \|2\kappa\|_{L^{q/2}(B_{4r}(x_0))} \right) \leq C \kappa r^2,$$

where  $C = C(n, \Lambda/\lambda)$ . Hence Lemma 3.1 is proved.  $\square$

Now consider a point  $x$  such that  $u_i(x) \neq u_{i+1}(x)$ . Then one can consider the distance  $d$  from  $x$  to the touching set of  $u_i$  and  $u_{i+1}$ . By the choice, in the ball  $B_d(x)$ ,  $\{u_i\}$  has at least two connected components. Then by the subsystem condition (I), the average of each connected components will satisfy a PDE (2.4). Moreover, the difference of the averages will also satisfy a PDE (2.7). Applying the  $C^{1,\alpha}$  interior estimates (Theorem A.1), Lemma 3.1, one can inductively obtain the following estimates of gradients of  $u_i - u_j$ . We remark that such a method for two sheets graphs has been used to study the obstacle problems; see [9, Theorem 3.2].

**Lemma 3.2.** *Let  $\{u_j\}_{j=1}^m$  be ordered  $C^{1,\alpha}$  graphs over  $B_1(0) \subset \mathbb{R}^n$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (2.1) and (2.2) are satisfied. Let  $x_0 \in B_1(0)$  and  $r > 0$  with  $u_i(x_0) = u_{i+1}(x_0)$  for some  $i \in \{1, \dots, m-1\}$  and  $B_{7r}(x_0) \subset B_1(0)$ . Then we have that for each  $x \in B_r(x_0)$ ,*

$$|Du_i(x) - Du_{i+1}(x)| \leq C(n, m, \Lambda, \lambda) \cdot (\kappa + \kappa_0)r.$$

*Proof.* We prove the lemma inductively. So long as  $m = 1$ , then there is nothing to be proved. Suppose that the statements in the lemma hold true for the number of the sheets less than or equal to  $m - 1$ . Now let  $\{u_j\}_{j=1}^m$  be ordered graphs which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Without loss of generality, we assume that  $\{u_j\}_{j=1}^m$  is connected in  $B_1(0)$ .

Now fix  $y \in B_r(x_0)$  and let  $\tilde{r} := \text{dist}(y, \{u_i = u_{i+1}\})$ . Clearly, there exists

$$x_i^* \in \{u_i = u_{i+1}\} \cap B_{2r}(x_0) \text{ with } \tilde{r} = |y - x_i^*|.$$

So long as  $\tilde{r} = 0$ , then  $Du_i(y) = Du_{i+1}(y)$ . We now consider the case of  $\tilde{r} > 0$ . Since  $x_0 \in \{u_i = u_{i+1}\}$ , then it follows that  $\tilde{r} < r$  and  $B_{5\tilde{r}}(x_i^*) \subset B_{7r}(x_0) \subset B_1(0)$ . By the choice of  $\tilde{r}$ , we know that  $u_i < u_{i+1}$  in  $B_{\tilde{r}}(y)$ . By applying Lemma 3.1 in  $B_{5\tilde{r}}(x_i^*)$  for  $u_i$  and  $u_{i+1}$ , we have that

$$(3.1) \quad |u_i(x) - u_{i+1}(x)| \leq C\kappa\tilde{r}^2, \quad \forall x \in B_{\frac{3}{4}\tilde{r}}(x_i^*) \supset B_{\frac{1}{4}\tilde{r}}(y),$$

where  $C = C(n, \Lambda/\lambda)$ . Now we define new functions  $v_1, \dots, v_m : B_1(0) \rightarrow \mathbb{R}$  by

$$v_j(x) := \tilde{r}^{-1}u_j(\tilde{r}x + y).$$

Since  $u_{i+1} > u_i$  in  $y \in B_{\tilde{r}}(y)$ , then  $v_{i+1} > v_i$  in  $B_1(0)$ . Define  $\tilde{\mathcal{Q}}$  by

$$\tilde{\mathcal{Q}}v = -\text{div}\tilde{\mathcal{A}}(x, v, Dv) + \tilde{\mathcal{B}}(x, v, Dv),$$

where

$$\tilde{\mathcal{A}}(x, z, \mathbf{p}) := \mathcal{A}(y + rx, rz, \mathbf{p}), \quad \tilde{\mathcal{B}}(x, z, \mathbf{p}) := r\mathcal{B}(y + rx, rz, \mathbf{p}).$$

Define the new functions  $\{\tilde{g}_j\}$  by

$$\tilde{g}_j(x) = rg_j(y + rx).$$

By Lemma 2.2,  $\{v_j\}$  is a solution to  $(\tilde{\mathcal{Q}}, \{\tilde{g}_j\})$ . Moreover, by the subsystem condition,  $\{v_j\}_{j=i+1}^m$  and  $\{v_j\}_{j=1}^i$  are solutions to  $(\tilde{\mathcal{Q}}, \{\tilde{g}_j\}_{j=i+1}^m)$  and  $(\tilde{\mathcal{Q}}, \{\tilde{g}_j\}_{j=1}^i)$ , respectively.

By Lemma 2.5, there exists  $r_1 \in (9^{-m-1}, 9^{-1})$  such that  $\{v_j\}$  has the same number of connected components in  $B_{9r_1}(0)$  and  $B_{r_1}(0)$ . For simplicity, we assume that  $\{v_j\}_{j=1}^i$  and  $\{v_j\}_{j=i+1}^m$  are connected components in  $B_{r_1}(0)$ . Observe that for  $j = 1, \dots, m$ ,

$$-\kappa\tilde{r} \leq \tilde{\mathcal{Q}}v_j \leq \kappa\tilde{r}.$$

That is,  $\{v_j\}$  satisfies  $\kappa\tilde{r}$ -condition. Since  $\{v_j\}_{j=1}^i$  and  $\{v_j\}_{j=i+1}^m$  are connected in  $B_{r_1}(0)$ , then for any  $j = 1, \dots, i-1$  or  $i+1, \dots, m-1$ , there exists  $x_j^* \in B_{r_1}(0)$  so that  $v_j(x_j^*) = v_{j+1}(x_j^*)$ . Applying Lemma 3.1 in  $B_{8r_1}(x_j^*) \subset B_{9r_1}(0) \subset B_1(0)$ , we have that for all  $x \in B_{2r_1}(x_j^*) \supset B_{r_1}(0)$ ,

$$(3.2) \quad |v_j(x) - v_{j+1}(x)| \leq C(n, \Lambda/\lambda)\kappa\tilde{r}, \quad j = 1, \dots, i-1, i+1, \dots, m-1.$$

On the other hand, by (3.1),

$$|v_i(x) - v_{i+1}(x)| \leq C\kappa\tilde{r}, \quad \forall x \in B_{\frac{1}{4}}(0) \supset B_{r_1}(0).$$

Together with (3.2), we conclude that for all  $x \in B_{r_1}(0)$  and  $j = 1, \dots, m$ ,

$$(3.3) \quad |v_j(x) - v_{j+1}(x)| \leq C(n, \Lambda/\lambda, m)\kappa\tilde{r}.$$

By induction, there exists a constant  $C = C(n, m, \Lambda, \lambda)$  so that for all  $x \in B_{r_1}(0)$ ,

$$(3.4) \quad |Dv_j(x) - Dv_{j+1}(x)| \leq C \cdot (\kappa + \kappa_0)\tilde{r}$$

for  $j = 1, \dots, i-1, i+1, \dots, m$ . Let

$$\varphi := \frac{1}{i} \sum_{j=1}^i v_j, \quad \psi := \frac{1}{m-i} \sum_{j=i+1}^m v_j, \quad \Psi := \varphi - \psi.$$

Then by (2.7),  $\Psi$  satisfies the equation

$$(3.5) \quad -\operatorname{div}(\mathbf{A}(\varphi, \psi)D\Psi + \mathbf{b}(\varphi, \psi)\Psi) + \mathbf{c}(\varphi, \psi)D\Psi + d(\varphi, \psi)\Psi = \operatorname{div} \mathbf{f}^* - g^*,$$

where  $\mathbf{A}, \mathbf{b}, \mathbf{c}$  and  $d$  are defined with respect to  $\tilde{\mathcal{Q}}$ ; and by Lemma 2.8, Lemma 2.2, Equation (3.3) and (3.4),

$$|\mathbf{f}^*| \leq C(n, m, \Lambda, \lambda)(\kappa + \kappa_0)\tilde{r}, \quad \sup_{x, x' \in B_{r_1}(0)} \frac{|\mathbf{f}^*(x) - \mathbf{f}^*(x')|}{|x - x'|^\alpha} \leq C(n, m, \Lambda, \lambda)(\kappa + \kappa_0)\tilde{r};$$

$$|g^*| \leq C(n, m, \Lambda, \lambda)(\kappa + \kappa_0)\tilde{r}.$$

Then the  $C^{1,\alpha}$ -estimates Theorem A.1 give that for  $x \in B_{r_1/2}(0)$ ,

$$|D\Psi(x)| \leq C(n, \alpha)(\|\Psi\|_{C^0(B_{r_1}(0))} + \|g^*\|_{C^0(B_{r_1}(0))} + \|\mathbf{f}^*\|_{C^\alpha(B_{r_1}(0))})$$

$$\leq C(n, m, \Lambda, \lambda)(\kappa + \kappa_0)\tilde{r}.$$

Together with (3.4), we then conclude that

$$|Du_{i+1}(y) - Du_i(y)| = |Dv_{i+1}(0) - Dv_i(0)| \leq C(n, m, \Lambda, \lambda)(\kappa + \kappa_0)\tilde{r}.$$

This completes the induction. By the arbitrariness of  $y$ , we have proved Lemma 3.2.  $\square$

**3.2. Integral estimates of the second derivatives.** Recall that for any function  $f$  and a constant  $\tau \neq 0$ ,

$$f^\tau(x) := \frac{f(x + \tau \mathbf{e}) - f(x)}{\tau},$$

where  $\mathbf{e}$  can be any unit vector in  $\mathbb{R}^n$ .

**Lemma 3.3.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered, connected graphs over  $B_1(0) \subset \mathbb{R}^n$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. Denote by*

$$\eta := \left( \|\partial_z \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_1)} + \|\partial_x \mathcal{B}\|_{C^1(\mathcal{U}_1)} \right) \cdot \sum_{j=1}^m \|u_j\|_{C^0(B_1(0))}.$$

Suppose that there exists a constant  $K > 0$  such that for all  $1 \leq i \leq m$  and  $\tau \in (0, 1/4)$ ,

$$\int_{B_{1-\tau}(0)} |Du_i^\tau - Du_j^\tau|^2 dx \leq K.$$

Then the average  $\varphi = \frac{1}{m} \sum_j u_j$  satisfies

$$\int_{B_{1/2}(0)} |D\varphi^\tau|^2 \leq C(n, m, \Lambda, \lambda) \left( \delta_0^2(K+1) + \eta^2 + \kappa_1 \delta_0 \right).$$

*Proof.* By (2.8), in  $B_{3/4}(0)$ ,

$$(3.6) \quad \sum_{j=1}^m \left[ -\operatorname{div} \left( \mathbf{A}_\tau(u_j) D\varphi^\tau + \mathbf{b}_\tau(u_j) \varphi^\tau \right) + \mathbf{c}_\tau(u_j) D\varphi^\tau + d_\tau(u_j) \varphi^\tau \right] = -\operatorname{div} \mathbf{f}_\tau + g_\tau,$$

where  $\mathbf{f}_\tau$  and  $g_\tau$  are given by (2.9) and (2.10). By (1.1) and (2.2), we have

$$(3.7) \quad \lambda \leq \mathbf{A}_\tau \leq \Lambda, \quad |\mathbf{b}_\tau| + |\mathbf{c}_\tau| + |d_\tau| \leq 3\Lambda.$$

Moreover, by (2.1),

$$(3.8) \quad |u_j^\tau| \leq \sup |Du_j| \leq \delta_0.$$

Since  $\{u_j\}$  is connected, then for all  $j = 1, \dots, m-1$ , there exists  $x_j \in B_1(0)$  such that  $u_j(x_j) = u_{j+1}(x_j)$ , which yields that for all  $x \in B_1(0)$ ,

$$\begin{aligned} |u_j(x) - u_{j+1}(x)| &\leq |u_j(x) - u_j(x_j)| + |u_{j+1}(x_j) - u_{j+1}(x)| \\ &\leq 2\delta_0|x - x_j| \leq 4\delta_0. \end{aligned}$$

By (2.1),

$$|Du_j(x) - Du_{j+1}(x)| \leq 2\delta_0.$$

By the arbitrariness of  $j$ , we conclude that

$$\|u_j - \varphi\|_{C^1(B_1(0))} \leq C(m)\delta_0.$$

This together with Lemma 2.10 implies

$$(3.9) \quad |\mathbf{f}_\tau| \leq C(n, m, \Lambda) \sum_{j=1}^m \left( \delta_0 |D\varphi^\tau - Du_j^\tau| + \eta + \delta_0 \right);$$

$$(3.10) \quad |g_\tau| \leq C \sum_{j=1}^m \left( \delta_0 |D\varphi^\tau - Du_j^\tau| + \delta_0 + \eta + \kappa_1 \right);$$

Now let  $\phi = \varphi^\tau \cdot \zeta^2$ , where  $\zeta \in C_c^1(B_{3/4}(0))$  is a cut-off function so that

$$(3.11) \quad |D\zeta| \leq C; \quad 0 \leq \zeta \leq 1; \quad \zeta \equiv 1 \text{ on } B_{1/2}(0).$$



Then by taking  $\phi$  as a test function in (3.6) and using (3.7), we have

$$\begin{aligned} \int_{B_{3/4}(0)} |D\varphi^\tau|^2 \zeta^2 \, dx &\leq C \int_{B_{3/4}(0)} \left( |D\varphi^\tau| |\varphi^\tau| |D\zeta^2| + |\varphi^\tau| |D(\varphi^\tau \zeta^2)| + |D\varphi^\tau| |\varphi^\tau| \zeta^2 + |\varphi^\tau|^2 \zeta^2 + \right. \\ &\quad \left. + |\mathbf{f}_\tau| (|D\varphi^\tau| \zeta^2 + |\varphi^\tau| |D\zeta^2|) + |g_\tau| |\varphi^\tau| \zeta^2 \right) \, dx \\ &\leq \frac{1}{2} \int_{B_{3/4}(0)} |D\varphi^\tau|^2 \zeta^2 \, dx + C \int_{B_{3/4}(0)} \left( |\varphi^\tau|^2 |D\zeta|^2 + |\varphi^\tau|^2 \zeta^2 + |\mathbf{f}_\tau|^2 \zeta^2 \right. \\ &\quad \left. + |g_\tau| |\varphi^\tau| \zeta^2 \right) \, dx. \end{aligned}$$

where  $C = C(m, \Lambda, \lambda)$ . Plugging (3.8), (3.9), (3.10) and (3.11) into the last inequality, we obtain

$$\begin{aligned} \int_{B_{3/4}(0)} |D\varphi^\tau|^2 \zeta^2 \, dx &\leq C \left( \sum_{j=1}^m \int_{B_{3/4}(0)} \delta_0^2 + \delta_0^2 |D\varphi^\tau - Du_j^\tau|^2 \, dx + \eta^2 + \kappa_1 \delta_0 \right) \\ &\leq C \left( \delta_0^2 (K+1) + \eta^2 + \kappa_1 \delta_0 \right), \end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda)$ . This finishes the proof of Lemma 3.3.  $\square$

Now we are ready to prove the hessian estimates for height differences. We will use the inductive method on the number of sheets  $m$ . For the point  $x$  which has density less than  $m$ , we will take  $r_1$  to be the largest radius so that  $\{u_j\}$  is not connected in  $B_{r_1}(x)$ . Then by induction, we can prove the integral estimates over  $B_{c(m)r_1}(x)$ , where  $c(m) \in (0, 1)$  is a constant depending only on  $m$ ; Lemma 3.3 will be used in this part. Moreover, for the touching points, we will use Lemma 3.2 to find such a small radius with desired integral bound. Finally, we will finish the induction by taking a covering argument.

**Lemma 3.4.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs over  $B_1(0) \subset \mathbb{R}^n$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. Denote by*

$$\eta := \left( \|\partial_z \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_1)} + \|\partial_x \mathcal{B}\|_{C^1(\mathcal{U}_1)} \right) \cdot \sum_{j=1}^m \|u_j\|_{C^0(B_1(0))}.$$

*If in addition that  $\{u_j\}$  is connected in  $B_{1/16}(0)$ , then for all  $\tau \in (0, 1/32)$  and  $j = 1, \dots, m-1$ ,*

$$\int_{B_{1/32}(0)} |Du_j^\tau - Du_{j+1}^\tau|^2 \, dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right),$$

*where  $C = C(n, m, \Lambda, \lambda)$ . This gives that  $u_i - u_j \in W^{2,2}(B_1(0))$  and*

$$\int_{B_{1/32}(0)} |D^2(u_i - u_j)|^2 \, dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right).$$

*As a corollary, if  $\{u_j\}$  is connected in  $B_s(0)$  and  $s \leq 1/32$ , then*

$$\int_{B_s(0)} |D^2(u_i - u_j)|^2 \, dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right) s^n.$$

*Proof.* We are going to prove the lemma by induction. So long as  $m = 1$ , then there is nothing to be proved. Suppose that the lemma is true when the number of the sheets is less than or

equal to  $m - 1$ . Now let  $\{u_j\}_{j=1}^m$  be ordered graphs which is a solution to  $(\mathcal{Q}, \{g_j\})$ . We further assume that  $\{u_j\}$  is connected in  $B_{1/16}(0)$ . For any  $x \in B_1(0)$ , let

$$r_1(x) := \sup \{r' < 1 - |x|; \{u_j\} \text{ is not connected in } B_{r'}(x)\}.$$

Then  $r_1(x_0) > 0$  if  $u_1(x_0) \neq u_m(x_0)$  for  $x_0 \in B_1(0)$ .

*Step 1: Decomposing  $\{u_j\}$  into connected components in small balls.*

Note that by Lemma 2.5, there exists  $r(x) \in (16^{-m}r_1(x), r_1(x))$  so that  $\{u_j\}$  has same number of connected components in  $B_{r(x)}(x)$  and  $B_{r(x)/16}(x)$ . Since  $\{u_j\}$  is connected in  $B_{1/16}(0)$ , then for each  $i = 1, \dots, m - 1$ , there exists  $x_i \in B_{1/16}(0)$  with  $u_i(x_i) = u_{i+1}(x_i)$ . Clearly,  $B_{1/8}(0) \subset B_{3/16}(x_i)$  and  $B_{3/4}(x_i) \subset B_1(0)$ . In particular, for all  $x \in B_{1/16}(0)$ ,

$$\text{dist}(x, \{u_i = u_{i+1}\}) \leq \frac{1}{8} < 1 - |x|.$$

Then by Lemma 3.1 and Lemma 3.2, there exists  $C = C(n, m, \Lambda, \lambda)$  such that for all  $x \in B_{1/16}(0)$ ,

$$\begin{aligned} |u_i(x) - u_{i+1}(x)| &\leq C \cdot \kappa [\text{dist}(x, \{u_i = u_{i+1}\})]^2 \leq C\kappa(r(x))^2; \\ |Du_i(x) - Du_{i+1}(x)| &\leq C \cdot (\kappa + \kappa_0) \text{dist}(y, \{u_i = u_{i+1}\}) \leq C(\kappa + \kappa_0)r(x). \end{aligned}$$

By the arbitrariness of  $i$ , we conclude that for all  $x \in B_{1/16}(0)$  and  $i, j \in \{1, \dots, m\}$ ,

$$(3.12) \quad |u_i(x) - u_j(x)| \leq C\kappa(r(x))^2; \quad |Du_i(x) - Du_j(x)| \leq C(\kappa + \kappa_0)r(x).$$

Now we fix  $y \in B_{1/16}(0)$  and use  $r$  to denote  $r(y)$  when there is no ambiguity. For any  $x \in B_1(0)$ , define

$$v_j(x) := r^{-1}u_j(rx + y), \quad \tilde{g}_j(x) = rg_j(rx + y).$$

Then by the choice of  $r_1$  and  $r$ ,  $\{v_j\}$  is not connected in  $B_1(0)$ . By relabeling, we have connected components

$$\{v_{1,1}, \dots, v_{1,i_1}\}, \quad \{v_{2,1}, \dots, v_{2,i_2}\}, \quad \dots, \quad \{v_{m,1}, \dots, v_{m,i_m}\},$$

and the associated functions

$$\{\tilde{g}_{1,1}, \dots, \tilde{g}_{1,i_1}\}, \quad \{\tilde{g}_{2,1}, \dots, \tilde{g}_{2,i_2}\}, \quad \dots, \quad \{\tilde{g}_{m,1}, \dots, \tilde{g}_{m,i_m}\}.$$

We assume that the order is preserved, i.e.

$$v_{1,1} \leq \dots \leq v_{1,i_1} < v_{2,1} \leq \dots < v_{m,1} \leq \dots \leq v_{m,i_m}.$$

Then by (3.12), for all  $j, k \in \{1, \dots, m\}$  and  $1 \leq \ell \leq i_j, 1 \leq \ell' \leq i_k$ ,

$$(3.13) \quad |v_{j,\ell}(x) - v_{k,\ell'}(x)| \leq C\kappa r(y), \quad |Dv_{j,\ell}(x) - Dv_{k,\ell'}(x)| \leq C \cdot (\kappa + \kappa_0)r(y).$$

Define

$$\varphi_j := \frac{1}{i_j} \sum_{\ell=1}^{i_j} v_{j,\ell}.$$

By the subsystem condition in Definition 2.1 and Lemma 2.2, for each  $j = 1, \dots, m$ ,  $\{v_{j,\ell}\}_\ell$  is a solution to  $(\tilde{\mathcal{Q}}, \{\tilde{g}_{j,\ell}\}_\ell)$ , where

$$\tilde{\mathcal{Q}}v = -\text{div } \tilde{\mathcal{A}}(x, v, Dv) + \tilde{\mathcal{B}}(x, v, Dv),$$

and

$$\tilde{\mathcal{A}}(x, z, \mathbf{p}) := \mathcal{A}(y + rx, rz, \mathbf{p}), \quad \tilde{\mathcal{B}}(x, z, \mathbf{p}) := r\mathcal{B}(y + rx, rz, \mathbf{p}).$$

Clearly, by (1.1),

$$(3.14) \quad \lambda \leq \partial_{\mathbf{p}} \widetilde{\mathcal{A}} \leq \Lambda, \quad \|\widetilde{\mathcal{A}}\|_{C^2} + \|\widetilde{\mathcal{B}}\|_{C^2} \leq \Lambda.$$

Moreover, by the definition of  $\widetilde{\mathcal{A}}$  and  $\widetilde{\mathcal{B}}$ ,

$$(3.15) \quad \begin{aligned} \|\partial_x \widetilde{\mathcal{A}}\|_{C^1} + \|\partial_z \widetilde{\mathcal{A}}\|_{C^1} + \|\widetilde{\mathcal{B}}\|_{C^2} &\leq C(n, \Lambda) \cdot r, \\ \|\partial_z \partial_x \widetilde{\mathcal{A}}\|_{C^0} + \|\partial_x \widetilde{\mathcal{B}}\|_{C^1} + \|\partial_z \widetilde{\mathcal{B}}\|_{C^1} &\leq C(n, \Lambda) \cdot r^2. \end{aligned}$$

Then for any  $\tau \in (0, 1/4)$ ,  $\varphi_j^\tau$  satisfies the equation in (2.8) (w.r.t.  $(\widetilde{\mathcal{Q}}, \{\widetilde{g}_{j,\ell}\}_\ell)$ ) in  $B_{1-\tau}(0)$  in the weak sense,

$$-\operatorname{div} \left( \mathbf{A}_\tau^j D\varphi_j^\tau + \mathbf{b}_\tau^j \varphi_j^\tau \right) + \mathbf{c}_\tau^j D\varphi_j^\tau + d_\tau^j \varphi_j^\tau = -\operatorname{div} \mathbf{f}_\tau^j + g_\tau^j,$$

where

$$\begin{aligned} \mathbf{A}_\tau^j &:= \sum_{\ell=1}^{i_j} \mathbf{A}_\tau(v_{j,\ell}); \quad \mathbf{b}_\tau^j := \sum_{\ell=1}^{i_j} \mathbf{b}_\tau(v_{j,\ell}); \quad \mathbf{c}_\tau^j := \sum_{\ell=1}^{i_j} \mathbf{c}_\tau(v_{j,\ell}); \quad d_\tau^j = \sum_{\ell=1}^m d_\tau(v_{j,\ell}); \\ \mathbf{f}_\tau^j &:= \sum_{\ell=1}^{i_j} \left( (\mathbf{A}_\tau(v_{j,\ell}) - \mathbf{A}_\tau(\varphi_j)) (D\varphi_j^\tau - Dv_{j,\ell}^\tau) + (\mathbf{b}_\tau(v_{j,\ell}) - \mathbf{b}_\tau(\varphi_j)) (\varphi_j^\tau - v_{j,\ell}^\tau) - \mathfrak{A}_\tau(v_{j,\ell}) \right); \\ g_\tau^j &= \sum_{\ell=1}^{i_j} \left( (\mathbf{c}_\tau(v_{j,\ell}) - \mathbf{c}_\tau(\varphi_j)) (D\varphi_j^\tau - Dv_{j,\ell}^\tau) + (d_\tau(v_{j,\ell}) - d_\tau(\varphi_j)) (\varphi_j^\tau - v_{j,\ell}^\tau) - \mathfrak{B}_\tau(v_{j,\ell}) - \widetilde{g}_{j,\ell}^\tau \right). \end{aligned}$$

Lemma 2.10 gives the bound of  $\mathbf{f}_\tau^j$  and  $g_\tau^j$ :

$$\begin{aligned} \left| \mathbf{f}_\tau^j + \sum_{\ell=1}^{i_j} \mathfrak{A}_\tau(v_{j,\ell}) \right| &\leq C \sum_{\ell} \left( \|v_{j,\ell} - \varphi_j\|_{C^1} \cdot |D\varphi_j^\tau - Dv_{j,\ell}^\tau| + \|v_{j,\ell} - \varphi_j\|_{C^1} \cdot |\varphi_j^\tau - v_{j,\ell}^\tau| \right); \\ \left| g_\tau^j + \sum_{\ell=1}^{i_j} \mathfrak{B}_\tau(v_{j,\ell}) \right| &\leq C \sum_{\ell} \left( \|v_{j,\ell} - \varphi_j\|_{C^1} \cdot |D\varphi_j^\tau - Dv_{j,\ell}^\tau| + \|v_{j,\ell} - \varphi_j\|_{C^1} \cdot |\varphi_j^\tau - v_{j,\ell}^\tau| + \right. \\ &\quad \left. + |D\widetilde{g}_{j,\ell}^\tau|_{C^0} \right); \end{aligned}$$

This together with (3.13) and (3.15) gives that

$$(3.16) \quad \left| \mathbf{f}_\tau^j + \sum_{\ell=1}^{i_j} \mathfrak{A}_\tau(v_{j,\ell}) \right| \leq C \left( (\kappa + \kappa_0) r \sum_{\ell=1}^{i_j} |D\varphi_j^\tau - Dv_{j,\ell}^\tau| + (\kappa + \kappa_0)^2 r^2 \right).$$

$$(3.17) \quad \left| g_\tau^j + \sum_{\ell=1}^{i_j} \mathfrak{B}_\tau(v_{j,\ell}) \right| \leq C \left( (\kappa + \kappa_0) r \sum_{\ell=1}^{i_j} |D\varphi_j^\tau - Dv_{j,\ell}^\tau| + (\kappa + \kappa_0)^2 r^2 + \kappa_1 r^2 \right).$$

*Step 2: Estimating the height of two connected components.*

For  $j, k \in \{1, \dots, \mathbf{m}\}$ , letting  $\Psi := \varphi_j - \varphi_k$ , we obtain the equation for  $\Psi^\tau$  in  $B_{1-\tau}(0)$  as follows:

$$(3.18) \quad -\operatorname{div}(\overline{\mathbf{A}} D\Psi^\tau + \overline{\mathbf{b}} \Psi^\tau) + \overline{\mathbf{c}} D\Psi^\tau + \overline{d} \Psi^\tau = -\operatorname{div} \overline{\mathbf{f}} + \overline{g},$$

where

$$\begin{aligned}\bar{\mathbf{A}} &:= \frac{1}{i_j} \mathbf{A}_\tau^j; & \bar{\mathbf{b}} &:= \frac{1}{i_j} \mathbf{b}_\tau^j; & \bar{\mathbf{c}} &:= \frac{1}{i_j} \mathbf{c}_\tau^j; & d &:= \frac{1}{i_j} d_\tau^j; \\ \bar{\mathbf{f}} &:= -\left(\frac{1}{i_j} \mathbf{A}_\tau^j - \frac{1}{i_k} \mathbf{A}_\tau^k\right) D\varphi_k^\tau - \left(\frac{1}{i_j} \mathbf{b}_\tau^j - \frac{1}{i_k} \mathbf{b}_\tau^k\right) \varphi_k^\tau + \left(\frac{1}{i_j} \mathbf{f}_\tau^j - \frac{1}{i_k} \mathbf{f}_\tau^k\right); \\ \bar{g} &:= -\left(\frac{1}{i_j} \mathbf{c}_\tau^j - \frac{1}{i_k} \mathbf{c}_\tau^k\right) D\varphi_k^\tau - \left(\frac{1}{i_j} d_\tau^j - \frac{1}{i_k} d_\tau^k\right) \varphi_k^\tau + \left(\frac{1}{i_j} g_\tau^j - \frac{1}{i_k} g_\tau^k\right).\end{aligned}$$

By Lemma 2.9 and (3.13), (3.14), (3.15)

$$(3.19) \quad \left| \frac{1}{i_j} \mathbf{A}_\tau^j - \frac{1}{i_k} \mathbf{A}_\tau^k \right| \leq C \sup_{\ell, \ell'} |\mathbf{A}_\tau(v_{j,\ell}) - \mathbf{A}_\tau(v_{k,\ell'})| \leq C(n, m, \Lambda, \lambda)(\kappa + \kappa_0)r.$$

A similar argument gives that

$$(3.20) \quad \left| \frac{1}{i_j} \mathbf{b}_\tau^j - \frac{1}{i_k} \mathbf{b}_\tau^k \right| + \left| \frac{1}{i_j} \mathbf{c}_\tau^j - \frac{1}{i_k} \mathbf{c}_\tau^k \right| + \left| \frac{1}{i_j} d_\tau^j - \frac{1}{i_k} d_\tau^k \right| \leq C(n, m, \Lambda, \lambda)(\kappa + \kappa_0)r^2;$$

$$(3.21) \quad \left| \frac{1}{i_j} \sum_{\ell=1}^{i_j} \mathfrak{A}_\tau(v_{j,\ell}) - \frac{1}{i_k} \sum_{\ell=1}^{i_k} \mathfrak{A}_\tau(v_{k,\ell}) \right| \leq \|\partial_x \widetilde{\mathcal{A}}\|_{C^1} \|v_{j,\ell} - v_{k,\ell'}\|_{C^1} \leq C(\kappa + \kappa_0)r^2;$$

$$(3.22) \quad \left| \frac{1}{i_j} \sum_{\ell=1}^{i_j} \mathfrak{B}_\tau(v_{j,\ell}) - \frac{1}{i_k} \sum_{\ell=1}^{i_k} \mathfrak{B}_\tau(v_{k,\ell}) \right| \leq \|\partial_x \widetilde{\mathcal{B}}\|_{C^1} \|v_{j,\ell} - v_{k,\ell'}\|_{C^1} \leq C(\kappa + \kappa_0)r^3.$$

Obviously, (3.21) combining with (3.16) gives that

$$\begin{aligned}(3.23) \quad & \left| \frac{1}{i_j} \mathbf{f}_\tau^j - \frac{1}{i_k} \mathbf{f}_\tau^k \right| \\ & \leq \frac{1}{i_j} \left| \mathbf{f}_\tau^j + \sum_{\ell=1}^{i_j} \mathfrak{A}_\tau(v_{j,\ell}) \right| + \left| \frac{1}{i_j} \sum_{\ell=1}^{i_j} \mathfrak{A}_\tau(v_{j,\ell}) - \frac{1}{i_k} \sum_{\ell=1}^{i_k} \mathfrak{A}_\tau(v_{k,\ell}) \right| + \frac{1}{i_k} \left| \mathbf{f}_\tau^k + \sum_{\ell=1}^{i_k} \mathfrak{A}_\tau(v_{k,\ell}) \right| \\ & \leq C(\kappa + \kappa_0)r \sum_{\ell=1}^{i_j} |D\varphi_j^\tau - Dv_{j,\ell}^\tau| + C(\kappa + \kappa_0)r^2 + C(\kappa + \kappa_0)r \sum_{\ell=1}^{i_k} |D\varphi_k^\tau - Dv_{k,\ell}^\tau|.\end{aligned}$$

Similarly, (3.22) and (3.17) imply

$$\begin{aligned}(3.24) \quad & \left| \frac{1}{i_j} g_\tau^j - \frac{1}{i_k} g_\tau^k \right| \\ & \leq \frac{1}{i_j} \left| g_\tau^j + \sum_{\ell=1}^{i_j} \mathfrak{B}_\tau(v_{j,\ell}) \right| + \left| \frac{1}{i_j} \sum_{\ell=1}^{i_j} \mathfrak{B}_\tau(v_{j,\ell}) - \frac{1}{i_k} \sum_{\ell=1}^{i_k} \mathfrak{B}_\tau(v_{k,\ell}) \right| + \frac{1}{i_k} \left| g_\tau^k + \sum_{\ell=1}^{i_k} \mathfrak{B}_\tau(v_{k,\ell}) \right| \\ & \leq C(\kappa + \kappa_0)r \sum_{\ell=1}^{i_j} |D\varphi_j^\tau - Dv_{j,\ell}^\tau| + C(\kappa + \kappa_0 + \kappa_1)r^2 + C(\kappa + \kappa_0)r \sum_{\ell=1}^{i_k} |D\varphi_k^\tau - Dv_{k,\ell}^\tau|.\end{aligned}$$

By (3.19), (3.20), (3.23) and (3.24), we conclude that

$$(3.25) \quad \begin{aligned} |\bar{\mathbf{f}}| + |\bar{g}| &\leq C \cdot (\kappa + \kappa_0)r(|D\varphi_k^\tau| + |\varphi_k^\tau|) + C(\kappa + \kappa_0)r \sum_{\ell=1}^{i_j} |D\varphi_j^\tau - Dv_{j,\ell}^\tau| + \\ &+ C(\kappa + \kappa_0)r \sum_{\ell=1}^{i_k} |D\varphi_k^\tau - Dv_{k,\ell}^\tau| + C(\kappa + \kappa_0 + \kappa_1)r^2, \end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda)$ .

Now let  $\zeta \in C_c^1(B_{1/64}(0))$  be a cut-off function so that

$$0 \leq \zeta \leq 1; \quad |D\zeta| \leq C; \quad \zeta \equiv 1 \text{ for all } x \in B_{1/128}(0).$$

By taking  $\phi = \Psi^\tau \zeta^2$  as a test function in (3.18) and using the ellipticity of  $\bar{\mathbf{A}}$  and the bound of  $\bar{\mathbf{b}}, \bar{\mathbf{c}}, \bar{\mathbf{d}}$ , we obtain that

$$\begin{aligned} \int_{B_{3/4}(0)} |D\Psi^\tau|^2 \zeta^2 dx &\leq C \int_{B_{3/4}(0)} \left( |\Psi^\tau|^2 |D\zeta|^2 + |D\Psi^\tau| |\Psi^\tau| \zeta^2 + |\Psi^\tau|^2 \zeta^2 + \right. \\ &\quad \left. + |\bar{\mathbf{f}}| (|D\Psi^\tau| \zeta^2 + |D\zeta|^2 |\Psi^\tau|) + |\bar{g}| \zeta^2 |\Psi^\tau| \right) dx \\ &\leq \frac{1}{2} \int_{B_{3/4}(0)} |D\Psi^\tau|^2 \zeta^2 dx + C \int_{B_{3/4}(0)} (\kappa + \kappa_0)^2 r^2 + |\bar{\mathbf{f}}|^2 \zeta^2 + |\bar{g}|^2 \zeta^2 dx, \end{aligned}$$

where we used  $|\Psi^\tau| \leq C \cdot (\kappa + \kappa_0)r$  following from (3.13) and the mean value theorem. After simplification, it becomes

$$(3.26) \quad \int_{B_{3/4}(0)} |D\Psi^\tau|^2 \zeta^2 dx \leq C \int_{B_{3/4}(0)} (\kappa + \kappa_0)^2 r^2 + |\bar{\mathbf{f}}|^2 \zeta^2 + |\bar{g}|^2 \zeta^2 dx.$$

Now we estimate the right hand side of this equation. Observe that

$$\begin{aligned} &\left( \|\partial_z \partial_x \tilde{\mathcal{A}}\|_{C^0(B_1(0))} + \|\partial_x \tilde{\mathcal{B}}\|_{C^1(B_1(0))} \right) \cdot \sum_{\ell=1}^{i_j} \|v_{j,\ell}\|_{C^0(B_1(0))} \\ &\leq \left( r^2 \|\partial_z \partial_x \mathcal{A}\|_{C^0(B_1(0))} + r^2 \|\partial_x \mathcal{B}\|_{C^1(B_1(0))} \right) \cdot \sum_{\ell=1}^m \|u_\ell\|_{C^0(B_1(0))} \cdot r^{-1} \leq \eta r. \end{aligned}$$

Thus by induction,

$$(3.27) \quad \int_{B_{1/32}(0)} |Dv_{j,\ell}^\tau - D\varphi_j^\tau|^2 dx \leq C \left( (1 + \eta^2 \cdot r^2)(\kappa r + \kappa_0 r)^2 + \kappa_1^2 r^2 \right) \leq C(1 + \eta^2)r^2.$$

This together with Lemma 3.3 (by taking  $K, \delta_0, \eta, \kappa_1$  to be  $C(1 + \eta^2)r^2, \delta_0, \eta r, \kappa_1 r^2$  therein) gives that

$$\begin{aligned} \int_{B_{1/64}(0)} |D\varphi_j^\tau|^2 dx &\leq C \left( \delta_0^2 (1 + \eta^2 r^2) + \eta^2 r^2 + \kappa_1 \delta_0 r^2 \right) \\ &\leq C(1 + \eta^2), \end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda)$  and we used the fact that  $\delta_0, \kappa, \kappa_1, r \leq 1$ . Plugging this into (3.25) and noting that  $|\varphi_k^\tau| \leq \sup |D\varphi_k| \leq \delta_0 < 1$ , we have

$$\begin{aligned} \int_{B_{3/4}(0)} |\bar{\mathbf{f}}|^2 \zeta^2 + |\bar{\mathbf{g}}|^2 \zeta^2 dx &\leq C \int_{B_{1/64}(0)} \left( (\kappa + \kappa_0)^2 r^2 (|D\varphi_k^\tau|^2 \zeta^2 + \sum_{\ell=1}^{i_j} |D\varphi_j^\tau - Dv_{j,\ell}^\tau|^2) \right. \\ &\quad \left. + (\kappa + \kappa_0)^2 r^2 \sum_{\ell=1}^{i_k} |D\varphi_k^\tau - Dv_{k,\ell}^\tau|^2 + (\kappa + \kappa_0 + \kappa_1)^2 r^2 \right) dx \\ &\leq C \left( (1 + \eta^2) (\kappa + \kappa_0)^2 r^2 + \kappa_1^2 r^2 \right). \end{aligned}$$

Plugging this into (3.26), we then have that

$$\int_{B_{1/128}(0)} |D\Psi^\tau|^2 dx \leq C \left( (1 + \eta^2) (\kappa + \kappa_0)^2 r^2 + \kappa_1^2 r^2 \right),$$

which combining with (3.27) yields that

$$\int_{B_{1/128}(0)} |Dv_{j,\ell}^\tau - Dv_{k,\ell'}^\tau|^2 dx \leq C \left( (1 + \eta^2) (\kappa + \kappa_0)^2 r^2 + \kappa_1^2 r^2 \right).$$

Note that  $v_j$  is from scaling  $u_j$ . Hence we conclude that

$$(3.28) \quad \int_{B_{\hat{r}(y)}(y)} |Du_j^\tau - Du_k^\tau|^2 dx \leq C \left( (1 + \eta^2) (\kappa + \kappa_0)^2 + \kappa_1^2 \right) \hat{r}^n, \quad \hat{r}(y) := r(y)/128.$$

*Step 3: Completing the proof by a covering argument.*

So far, we have defined  $\hat{r}(y)$  for all  $y \in B_{1/16}(0) \setminus \{u_1 = u_m\}$  so that (3.28) is satisfied. Now we define  $\hat{r}(y) = \tau$  for  $y \in B_{1/16}(0) \cap \{u_1 = u_m\}$ . In this case,  $Du_i(y) = Du_j(y)$ . Then by (3.12), for  $\tau \in (0, 1/32)$  and  $x \in B_\tau(0)$ ,

$$\begin{aligned} |Du_i^\tau(x) - Du_j^\tau(x)| &\leq \frac{|Du_i(x + \tau e) - Du_j(x + \tau e)|}{\tau} + \frac{|Du_i(x) - Du_j(x)|}{\tau} \\ &\leq C \cdot (\kappa + \kappa_0) \tau / \tau = C(n, m, \Lambda, \lambda) \cdot (\kappa + \kappa_0). \end{aligned}$$

This implies

$$(3.29) \quad \int_{B_{\hat{r}(y)}(y)} |Du_i^\tau(x) - Du_j^\tau(x)|^2 dx \leq C \cdot (\kappa + \kappa_0)^2 \hat{r}^n.$$

Now that  $\{B_{\hat{r}(y)}(y); y \in B_{1/32}(0)\}$  forms a covering of  $B_{1/32}(0)$ . Hence one can find a finite cover  $\{B_{\hat{r}(y_k)}(y_k); k = 1, \dots, N\}$  so that  $B_{\hat{r}(y_k)/2}(y_k)$  does not intersect  $B_{\hat{r}(y_i)/2}(y_i)$  for  $k \neq i$ . Then by (3.28) and (3.29),

$$\begin{aligned} \int_{B_{1/32}(0)} |Du_i^\tau - Du_j^\tau|^2 dx &\leq \sum_{k=1}^N \int_{B_{\hat{r}(y_k)}(y_k)} |Du_i^\tau - Du_j^\tau|^2 dx \\ &\leq C \sum_{k=1}^N \left( (1 + \eta^2) (\kappa + \kappa_0)^2 + \kappa_1^2 \right) [\hat{r}(y_k)]^n \\ &\leq C \cdot \left( (1 + \eta^2) (\kappa + \kappa_0)^2 + \kappa_1^2 \right). \end{aligned}$$

This completes the proof Lemma 3.4.  $\square$

So far, we have proved the hessian estimates of the height difference. This combining with Lemma 3.3 will give the hessian estimates for the average, which yields the integral bound of  $D^2u_j$ .

Recall that

$$\boldsymbol{\kappa}^* := \kappa^2 + \kappa_0^2 + \kappa_1^2 + \delta_0^2,$$

and

$$\eta := \left( \|\partial_z \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_1)} + \|\partial_x \mathcal{B}\|_{C^1(\mathcal{U}_1)} \right) \cdot \sum_{j=1}^m \|u_j\|_{C^0(B_1(0))}.$$

**Proposition 3.5.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs over  $B_1(0) \subset \mathbb{R}^n$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. Then*

$$\sum_{j=1}^m \int_{B_{3/4}(0)} |D^2u_j|^2 dx \leq C(n, m, \Lambda, \lambda)(\boldsymbol{\kappa}^* + \eta^2).$$

*Proof.* For every  $y \in B_{3/4}(0)$ , it follows that  $B_{1/4}(y) \subset B_1(0)$ . By Lemma 2.5, there exists  $r \in [16^{-m}, 1]$  so that  $\{u_j\}$  has the same number of connected components in  $B_{r/4}(y)$  and  $B_{r/64}(y)$ . Denote by  $\{v_j\}_{j=1}^m$  be one of the connected components. Note that  $\kappa, \kappa_0, \kappa_1 \leq 1$ . Then by Lemma 3.4 in  $B_{r/4}(y)$  for  $\{v_j\}$ , we have that for  $i, j = 1, \dots, m$ ,

$$(3.30) \quad \int_{B_{r/128}(y)} |D^2v_i - D^2v_j|^2 dx \leq C\left((1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2\right) \leq C(\boldsymbol{\kappa}^* + \eta^2).$$

Then applying Lemma 3.3 in  $B_{r/128}(y)$ , we obtain

$$\int_{B_{r/256}(y)} |D^2\varphi|^2 dx \leq C\left(\delta_0^2 \cdot C(\boldsymbol{\kappa}^* + \eta^2 + 1) + \eta^2 + \kappa_1\delta_0\right) \leq C(\boldsymbol{\kappa}^* + \eta^2).$$

where  $C = C(n, m, \Lambda, \lambda)$  and  $\varphi = \frac{1}{m} \sum_{j=1}^m v_j$ . This together with (3.30) gives that for  $j = 1, \dots, m$ ,

$$\int_{B_{r/256}(y)} |D^2v_j|^2 dx \leq C(\boldsymbol{\kappa}^* + \eta^2).$$

In particular, letting  $r_m := 16^{-m-2}$ , by the arbitrariness of  $v_j$ , we conclude that for  $j = 1, \dots, m$ ,

$$\int_{B_{r_m}(y)} |D^2u_j|^2 dx \leq C(\boldsymbol{\kappa}^* + \eta^2).$$

By taking a finite cover of  $B_{3/4}(0)$  with small balls with radius  $r_m$ , then Proposition 3.5 is proved.  $\square$

#### 4. ORDER ESTIMATES OF THE HESSIAN

This section is devoted to prove a monotonicity formula for the average of  $|D^2u_j|^2$  over small balls, which implies that  $\int_{B_r(y)} |D^2u_j|^2 dx$  has order  $n$ . Since we have proved that each  $u_j$  is  $W^{2,2}$ , then we are able to consider the divergence form equation for  $Du_j$ ,  $j = 1, \dots, m$ . We will follow the strategy in proving Hölder regularity for gradients of solutions of divergence form, where one should approximate the  $D_k u$  ( $k = 1, \dots, n$ ) by a harmonic function; see [7, §3.4]. The main difficulty here is that the connectedness of  $\{u_j\}$  may be not preserved and then there are no height difference estimates in Section 3.

To overcome it, we will use an inductive method. Note that at the radius where  $\{u_j\}$  is not connected, the height difference estimates are still valid. By the subsystem condition, each connected component will satisfy a partial differential equation. Furthermore, the difference of the average of connected component will satisfy a weak solution. This together with the inequality from the induction will give the desired inequality.

In Section 4.1, We will first prove the order  $n - \epsilon$  for any  $\epsilon > 0$ . The idea is to approximate the average by a harmonic function which will help us to give the order of the average. Combining with the height difference estimates, we will obtain the order for each  $u_j$ .

In Section 4.2, we will prove the improved order estimates for the graph of  $u_j$  over a tilt-plane. Note that the inductive method can not be proceed since the optimal order is  $n$  instead of  $n + \alpha$  in the minimal surface equation. Here we will consider the order of the average and then use the inductive method to estimate

$$\sigma^*(r) := \sum_{j=1}^m \int_{B_r(0)} |D^2 u_j - (D^2 u_j)_{B_r(0)}|^2 dx.$$

Finally, in Section 4.3, we will give a monotonicity formula of  $r^{-n} \sigma^*(r)$ .

In this section, we will use the following classical results. Note that here we only require the inequalities for  $r > r_0$ .

**Lemma 4.1** ([7, Lemma 3.4; 8, Lemma 14.4.16]). *Let  $\gamma > 1$ ,  $R_0 > 0$ ,  $\beta \in (0, 1/2)$ ,  $\mu > 2\beta$ ,  $\kappa > 0$  and  $\sigma(r)$  be a nonnegative, monotonically increasing function on  $[0, R_0]$  satisfying*

$$\sigma(r) \leq \gamma \left( \left( \frac{r}{R} \right)^\mu + \delta \right) \sigma(R) + \kappa R^{\mu-2\beta}$$

for all  $0 < r_0 < r \leq R \leq R_0$  with  $\delta < \delta_1(\gamma, \mu, \beta) = (2\gamma)^{-\frac{\mu}{\beta}}$ . We then have

$$\sigma(r) \leq \gamma_1 \left( \frac{r}{R} \right)^{\mu-\beta} \sigma(R) + \gamma_2 \kappa r^{\mu-2\beta}$$

with  $\gamma_1, \gamma_2$  depending on  $\gamma, \mu, \beta$ .

*Proof.* Let

$$\tau := (2\gamma)^{-\frac{1}{\beta}}.$$

Then  $\delta < \delta_1 = \tau^\mu$ . By assumption,

$$\sigma(\tau R) \leq 2\gamma \tau^\mu \sigma(R) + \kappa R^{\mu-2\beta} = \tau^{\mu-\beta} \sigma(R) + \kappa R^{\mu-2\beta}.$$

Since  $r < R$ , then there exists an integer  $k \geq 0$  so that

$$\tau^{k+1} R < r \leq \tau^k R.$$

Without loss of generality, we assume that  $k \geq 1$ . Then we have that

$$\begin{aligned} \sigma(r) &\leq \sigma(\tau^k R) \leq \tau^{(\mu-\beta)k} \sigma(R) + \sum_{j=0}^{k-1} \kappa (\tau^{k-1-j} R)^{\mu-2\beta} \tau^{(\mu-\beta)j} \\ &\leq \tau^{\beta-\mu} \cdot \left( \frac{r}{R} \right)^{\mu-\beta} \sigma(R) + \kappa \tau^{(k-1)(\mu-2\beta)} R^{\mu-2\beta} (1 - \tau^\beta)^{-1} \\ &\leq (2\gamma)^{\frac{\mu-\beta}{\beta}} \left( \frac{r}{R} \right)^{\mu-\beta} \sigma(R) + \kappa \cdot (1 - (2\gamma)^{-1})^{-1} \cdot (2\gamma)^{\frac{2(\mu-2\beta)}{\beta}} \cdot r^{\mu-2\beta}. \end{aligned}$$



Then the lemma follows by taking

$$\gamma_1 = (2\gamma)^{\frac{\mu-\beta}{\beta}}; \quad \gamma_2 = (1 - (2\gamma)^{-1})^{-1} \cdot (2\gamma)^{\frac{2(\mu-2\beta)}{\beta}}.$$

□

**4.1. A primary order estimate.** Let  $\{u_j\}$  be  $C^{1,\alpha}$  ordered graphs on  $B_1(0)$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that  $\{u_j\}$  is connected in  $B_s(0)$  for some  $s \in (0, 1/16]$ . Then by Lemma 3.1 and Lemma 3.2, there exists  $C = C(n, m, \Lambda, \lambda)$  for  $x \in B_{1/16}(0) \setminus B_s(0)$ ,

$$(4.1) \quad |u_i(x) - u_j(x)| \leq C \cdot \kappa |x|^2, \quad |Du_i(x) - Du_j(x)| \leq C \cdot (\kappa + \kappa_0) |x|.$$

We now derive a local integral estimate for the average of  $\{u_j\}$ . This process is similar to the Hölder estimates [7, Theorem 3.8] by approximating with a harmonic function. However, we have more error terms on the right hand side. To overcome it, we will write the equation of  $D_k \varphi$  as a form of (4.17) and use (4.1) to control all of the error terms.

Recall that

$$\kappa^* := \kappa^2 + \kappa_0^2 + \kappa_1^2 + \delta_0^2,$$

and

$$\eta := \left( \|\partial_z \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_1)} + \|\partial_x \mathcal{B}\|_{C^1(\mathcal{U}_1)} \right) \cdot \sum_{j=1}^m \|u_j\|_{C^0(B_1(0))}.$$

For simplicity, let

$$\epsilon := \min \left\{ \frac{1}{2}\alpha, \frac{1}{2}(1 - \alpha) \right\}.$$

**Lemma 4.2.** *Let  $\{u_j\}$  be  $C^{1,\alpha}$  ordered graphs on  $B_1(0)$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. If  $\{u_j\}_{j=1}^m$  is connected in  $B_s(0)$ , with  $s \leq r < R$ , then we have that*

$$\int_{B_r(0)} |D^2 \varphi|^2 dx \leq C \left( \frac{r}{R} \right)^{n-\epsilon} \int_{B_R(0)} |D^2 \varphi|^2 dx + C(\kappa^* + \eta^2) r^{n-2\epsilon},$$

where  $\varphi = \frac{1}{m} \sum_{j=1}^m u_j$  and  $C = C(n, m, \Lambda, \lambda, \alpha)$ .

*Proof of Lemma 4.2.* We first assume that  $r \leq R \leq 1/64$ . By Remark 3.5,  $u_j \in W_{loc}^{2,2}(B_1(0))$ . Then by (2.13),

$$(4.2) \quad -\operatorname{div} \left( \mathbf{A}^* D D_k \varphi + \mathbf{b}^* D_k \varphi \right) + \mathbf{c}^* D D_k \varphi + d^* D_k \varphi = -\operatorname{div} \left( \hat{\mathbf{f}} - m \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) \right) + \hat{g},$$

where  $\hat{\mathbf{f}}$  and  $\hat{g}$  are defined in (2.14) and (2.15); and

$$\begin{aligned} \mathbf{A}^*(x) &:= \sum_{j=1}^m \partial_{\mathbf{p}} \mathcal{A}(x, u_j(x), Du_j(x)); & \mathbf{b}^*(x) &:= \sum_{j=1}^m \partial_z \mathcal{A}(x, u_j(x), Du_j(x)); \\ \mathbf{c}^*(x) &:= \sum_{j=1}^m \partial_{\mathbf{p}} \mathcal{B}(x, u_j(x), Du_j(x)); & d^*(x) &:= \sum_{j=1}^m \partial_z \mathcal{B}(x, u_j(x), Du_j(x)). \end{aligned}$$

We divide the proof into four steps.

**Step I:** *Estimating the coefficients in (4.2).*

Observe that by (2.1), for all  $x, y \in B_R(0)$ ,

$$(4.3) \quad |u_j(x) - u_j(y)| \leq 2\delta_0 R, \quad |Du_j(x) - Du_j(y)| \leq 2\delta_0 R^\alpha.$$

By (1.2) and the definition of  $\mathbf{A}^*, \mathbf{b}^*, \mathbf{c}^*, d^*$ ,

$$(4.4) \quad |\mathbf{A}^*| + |\mathbf{b}^*| + |\mathbf{c}^*| + |d^*| \leq C(m, \Lambda).$$

By the definition of  $\mathbf{A}^*$ , we have that

$$|\mathbf{A}^*(x) - \mathbf{A}^*(y)| \leq \sum_{j=1}^m \left( \|\partial_x \partial_{\mathbf{p}} \mathcal{A}\|_{C^0} |x - y| + \|\partial_z \partial_{\mathbf{p}} \mathcal{A}\|_{C^0} |u_j(x) - u_j(y)| + \|\partial_{\mathbf{p}} \partial_{\mathbf{p}} \mathcal{A}\|_{C^0} |Du_j(x) - Du_j(y)| \right).$$

Together with (4.3), we have that

$$(4.5) \quad \sup_{x, y \in B_R(0)} |\mathbf{A}^*(x) - \mathbf{A}^*(y)| \leq C(m, \Lambda)(R + \delta_0 R + \delta_0 R^\alpha) \leq C(m, \Lambda)(R + \delta_0 R^\alpha).$$

A similar argument gives that

$$(4.6) \quad \sup_{x, y \in B_R(0)} \left( |\mathbf{b}^*(x) - \mathbf{b}^*(y)| + |\mathbf{c}^*(x) - \mathbf{c}^*(y)| + |d^*(x) - d^*(y)| \right) \leq C(m, \Lambda)(R + \delta_0 R^\alpha).$$

On the other hand, plugging (4.3) and (4.1) into Lemma 2.11, we have that

$$(4.7) \quad \int_{B_R(0)} |\hat{\mathbf{f}}|^2 dx \leq C \sum_{j=1}^m \int_{B_R(0)} (\kappa + \kappa_0)^2 R^2 \left( |DD_k \varphi - DD_k u_j|^2 + 1 \right) dx;$$

$$(4.8) \quad \int_{B_R(0)} |\hat{g}| dx \leq C \sum_{j=1}^m \int_{B_R(0)} (\kappa + \kappa_0) R \left( |DD_k \varphi - DD_k u_j| + (\kappa + \kappa_0) R \right) + \eta + \delta_0 + \kappa_1 dx.$$

Recall that by Lemma 3.4, for  $1 \leq i \leq j \leq m$ ,

$$\int_{B_R(0)} |D^2 u_i - D^2 u_j|^2 dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right) R^n,$$

which implies that

$$\int_{B_R(0)} |D^2 \varphi - D^2 u_j|^2 dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right) R^n.$$

Plugging the above two inequalities into (4.7) and (4.8), we have

$$(4.9) \quad \int_{B_R(0)} |\hat{\mathbf{f}}|^2 dx \leq C(1 + \eta^2) \kappa^* R^{n+2},$$

$$(4.10) \quad \int_{B_R(0)} |\hat{g}| dx \leq C(\sqrt{\kappa^*} + \eta \kappa^*) R^n,$$

where  $C = C(n, m, \Lambda, \lambda)$  and we used that  $\kappa, \kappa_0, \kappa_1, \delta_0 \leq 1$ .

By the definition of  $\varphi$  and (4.3), for  $x, y \in B_R(0)$ ,

$$(4.11) \quad |D_k \varphi| \leq \delta_0, \quad |D_k \varphi(x) - D_k \varphi(y)| \leq \delta_0 (2R)^\alpha.$$

Plugging it into Lemma 2.11, we obtain that for all  $\phi \in C_c^1(B_R(0))$ ,

$$(4.12) \quad \left| \int_{B_R(0)} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) D\phi dx \right| \leq C \int_{B_R(0)} (\eta + \delta_0) |\phi| + \delta_0 R^\alpha |D\phi| + R^\alpha |D^2 \varphi| |\phi| dx.$$

□

**Step II:** *Constructing a harmonic function to approximate  $D_k \varphi$ .*

Let  $f$  be the weak solution of

$$(4.13) \quad -\operatorname{div}\left(\mathbf{A}^*(0)Df\right) = 0, \quad f|_{\partial B_R(0)} = D_k\varphi - D_k\varphi(0).$$

Let

$$\phi := D_k\varphi - D_k\varphi(0) - f.$$

Obviously,  $\phi \in W_0^{1,2}(B_R(0))$ . Then we claim that

$$(4.14) \quad \int_{B_R(0)} |Df|^2 dx \leq C(m, \lambda, \Lambda) \int_{B_R(0)} |DD_k\varphi|^2 dx.$$

Indeed, by taking  $\phi$  as a test function in (4.13), we have

$$\begin{aligned} \int_{B_R(0)} |Df|^2 dx &\leq C(\lambda) \int_{B_R(0)} |\mathbf{A}^*(0)| |DD_k\varphi| |Df| dx \\ &\leq C(\lambda) \int_{B_R(0)} |\mathbf{A}^*(0)|^2 |DD_k\varphi|^2 dx + \frac{1}{2} \int_{B_R(0)} |Df|^2 dx, \end{aligned}$$

which implies

$$(4.15) \quad \int_{B_R(0)} |Df|^2 dx \leq C(\lambda) \int_{B_R(0)} |\mathbf{A}^*(0)|^2 |DD_k\varphi|^2 dx.$$

Plugging (4.4) into (4.15), we then obtain (4.14). By (4.11) and the maximum principle for  $f$ ,

$$|f(x)| \leq \delta_0(2R)^\alpha.$$

It follows that

$$(4.16) \quad |\phi| \leq |D_k\varphi - D_k\varphi(0)| + |f| \leq 2^{1+\alpha}\delta_0 R^\alpha.$$

□

**Step III:** Bounding  $L^2$  norm of  $\phi$  by  $\varphi$ .

Note that

$$\operatorname{div}(\mathbf{b}^*(0)D_k\varphi(0)) = 0, \quad \operatorname{div}(\mathbf{c}^*(0)D_k\varphi(0)) = 0.$$

This together with the definition of  $f$  in (4.13) and (4.2) gives that

$$(4.17) \quad \begin{aligned} &-\operatorname{div}\left(\mathbf{A}^*D(D_k\varphi - f) + \mathbf{b}^*(D_k\varphi - D_k\varphi(0)) - \mathbf{c}^*(0)(D_k\varphi - D_k\varphi(0))\right) + d^*D_k\varphi \\ &= -\operatorname{div}\left((\mathbf{A}^*(0) - \mathbf{A}^*)Df + (\mathbf{b}^*(0) - \mathbf{b}^*)D_k\varphi(0) + \hat{\mathbf{f}} - m\partial_{x_k}\mathcal{A}(x, \varphi, D\varphi)\right) \\ &\quad - (\mathbf{c}^* - \mathbf{c}^*(0))DD_k\varphi + \hat{g}. \end{aligned}$$

Then by taking  $\phi$  as a test function in (4.17) and using (4.4), (4.5), (4.6), (4.11), (4.12) and (4.16), we obtain

$$\begin{aligned}
& \int_{B_R(0)} |D\phi|^2 dx \\
& \leq C(n, m, \Lambda, \lambda) \int_{B_R(0)} \left( (|\mathbf{b}^*| + |\mathbf{c}^*(0)|) |D_k\varphi - D_k\varphi(0)| |D\phi| + |d^*| |D_k\varphi| |\phi| + \right. \\
& \quad + |\mathbf{A}^*(0) - \mathbf{A}^*| |Df| |D\phi| + |\mathbf{b}^*(0) - \mathbf{b}^*| |D_k\varphi(0)| |D\phi| + |\mathbf{c}^* - \mathbf{c}^*(0)| |DD_k\varphi| |\phi| + \\
& \quad \left. + |\hat{\mathbf{f}}| |D\phi| + |\hat{g}| |\phi| \right) dx + m \left| \int_{B_R(0)} \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi) D\phi dx \right| \\
& \leq C \int_{B_R(0)} \left( \delta_0 R^\alpha |D\phi| + \delta_0 \cdot \delta_0 R^\alpha + (R + \delta_0 R^\alpha) |Df| |D\phi| + (R + \delta_0 R^\alpha) \cdot \delta_0 |D\phi| + \right. \\
& \quad \left. + (R + \delta_0 R^\alpha) \delta_0 R^\alpha |DD_k\varphi| + |\hat{\mathbf{f}}| |D\phi| + |\hat{g}| \delta_0 R^\alpha + (\delta_0 + \eta) \delta_0 R^\alpha + R^\alpha |D^2\varphi| \delta_0 R^\alpha \right) dx \\
& \leq \frac{1}{4} \int_{B_R(0)} |D\phi|^2 dx + C \int_{B_R(0)} \left( \delta_0^2 R^\alpha + R^{2\alpha} (|Df|^2 + |D^2\varphi|^2) + |\hat{\mathbf{f}}|^2 + |\hat{g}| \delta_0 R^\alpha + \delta_0 \eta R^\alpha \right) dx.
\end{aligned}$$

This together with (4.14) implies

$$(4.18) \quad \int_{B_R(0)} |D\phi|^2 dx \leq C \int_{B_R(0)} \left( (\delta_0^2 + \delta_0 \eta) R^\alpha + R^{2\alpha} |D^2\varphi|^2 + |\hat{\mathbf{f}}|^2 + |\hat{g}| \delta_0 R^\alpha \right) dx.$$

Plugging (4.9) and (4.10) into (4.18), we conclude that

$$(4.19) \quad \int_{B_R(0)} |DD_k\varphi - Df|^2 dx \leq C \int_{B_R(0)} R^{2\alpha} |D^2\varphi|^2 dx + C(\kappa^* + \eta^2) R^{n+\alpha}.$$

By harmonicity of  $f$  [7, Lemma 3.10; 8, Lemma 14.4.5], there exists a constant  $A(n)$  such that

$$\int_{B_r(0)} |Df|^2 dx \leq A(n) \left( \frac{r}{R} \right)^n \int_{B_R(0)} |Df|^2 dx.$$

This together with (4.19) and (4.14) gives

$$\begin{aligned}
\int_{B_r(0)} |DD_k\varphi|^2 & \leq \int_{B_r(0)} 2|Df|^2 + 2|DD_k\varphi - Df|^2 dx \\
& \leq A' \left( \left( \frac{r}{R} \right)^n + R^{2\alpha} \right) \int_{B_R(0)} |D^2\varphi|^2 dx + C(\kappa^* + \eta^2) R^{n+\alpha},
\end{aligned}$$

where  $A' = A'(n, m, \Lambda, \lambda, \alpha)$  and we now fix such a constant.  $\square$

**Step IV:** *Proving the proposition case by case.*

By the arbitrariness of  $k$ , we conclude that for all  $r < R \leq 1/64$ ,

$$(4.20) \quad \int_{B_r(0)} |D^2\varphi|^2 \leq nA' \left( \left( \frac{r}{R} \right)^n + R^{2\alpha} \right) \int_{B_R(0)} |D^2\varphi|^2 dx + C(\kappa^* + \eta^2) R^{n+\alpha}.$$

Then Lemma 4.1 applies to give a positive constant  $R_0 = R_0(n, A', \alpha) < 1/64$  with

$$R_0^{2\alpha} < (2nA)^{-\frac{n}{\epsilon}}$$

and the following possibilities:

- If  $s < r \leq R \leq R_0$ , then by Lemma 4.1,

$$\int_{B_r(0)} |D^2\varphi|^2 dx \leq C \left(\frac{r}{R}\right)^{n-\epsilon} \int_{B_R(0)} |D^2\varphi|^2 dx + C(\kappa^* + \eta^2)r^{n-2\epsilon},$$

where  $C$  is depending only on  $n, m, \Lambda, \lambda, \alpha$ .

- If  $r > R_0$ , then the inequality is trivial because  $R_0$  depends only on  $n, m, \Lambda, \lambda, \alpha$  and  $R/r \leq 1/R_0$ . Indeed, it follows that

$$\int_{B_r(0)} |D^2\varphi|^2 dx \leq C \cdot R_0^{-n} \left(\frac{r}{R}\right)^{n-\epsilon} \int_{B_R(0)} |D^2\varphi|^2 dx.$$

- If  $r < R_0 < R$ , then by the first case,

$$\begin{aligned} \int_{B_r(0)} |D^2\varphi|^2 dx &\leq C \left(\frac{r}{R_0}\right)^{n-\epsilon} \int_{B_{R_0}(0)} |D^2\varphi|^2 dx + C(\kappa^* + \eta^2)r^{n-2\epsilon} \\ &\leq CR_0^{-n+\epsilon} \left(\frac{r}{R}\right)^{n-\epsilon} \int_{B_R(0)} |D^2\varphi|^2 dx + C(\kappa^* + \eta^2)r^{n-2\epsilon}. \end{aligned}$$

This finishes the proof of Lemma 4.2. □

*Remark 4.3.* To prove Lemma 4.2, one can replace (4.19) by a weaker inequality, e.g.

$$\int_{B_R(0)} |DD_k\varphi - Df|^2 dx \leq C \int_{B_R(0)} R^\alpha |D^2\varphi|^2 dx + C(\kappa^* + \eta^2)R^n,$$

which can be derived by a much easier way without (4.17). However, the order on the right hand side of (4.19) will be essentially used in Lemma 4.5.

Combining Lemma 4.2 and Lemma 3.4, we obtain the primary order estimates for all  $r < R \leq t$  even if  $\{u_j\}$  is disconnected in  $B_R(0)$ .

**Proposition 4.4.** *Let  $t \leq 1$ . Let  $\{u_j\}$  be  $C^{1,\alpha}$  ordered graphs on  $B_t(0)$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. There exists a constant  $C = C(n, m, \Lambda, \lambda, \alpha)$  so that for all  $r < R \leq t$ ,*

$$\sum_{j=1}^m \int_{B_r(0)} |D^2u_j|^2 \leq C \left(\frac{r}{R}\right)^{n-\epsilon} \sum_{j=1}^m \int_{B_R(0)} |D^2u_j|^2 dx + C(\kappa^* + \eta^2)r^{n-2\epsilon}.$$

*Proof.* So long as  $r \geq t/16$ , then the inequality is trivial. Now we assume that  $r \leq t/16$ . We are going to prove the lemma inductively on  $m$ . When  $m = 1$ , then the inequality follows from Lemma 4.2. Suppose that the lemma is true for the number of sheets less than or equal to  $m - 1$ . Let  $\{u_j\}_{j=1}^m$  be a solution to  $(\mathcal{Q}, \{g_j\})$  and is connected in  $B_t(0)$ . Now set

$$s := \inf\{r > 0; \{u_j\}_{j=1}^m \text{ is connected in } B_r(0)\}.$$

*Case 1:*  $r < R \leq s$ . Then  $\{u_j\}_{j=1}^m$  can be divided into connected components in  $B_s(0)$ . By induction, each connected component satisfies the inequality in the proposition. Then the desired result follows by taking the sum.

*Case 2:*  $s \leq r < R$ . Then by Lemma 4.2,

$$\int_{B_r(0)} |D^2\varphi|^2 dx \leq C \left(\frac{r}{R}\right)^{n-\epsilon} \int_{B_R(0)} |D^2\varphi|^2 dx + C(\kappa^* + \eta^2)r^{n-2\epsilon},$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$ . Since  $r \leq t/16$ , then Lemma 3.4 applies to give that for  $1 \leq i, j \leq m$ ,

$$\int_{B_r(0)} |D^2 u_i - D^2 u_j|^2 dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right) r^n \leq C(\kappa^* + \eta^2) r^n.$$

Combining the above two inequalities, we obtain

$$\sum_{j=1}^m \int_{B_r(0)} |D^2 u_j|^2 dx \leq C \left( \frac{r}{R} \right)^{n-\epsilon} \int_{B_R(0)} \sum_{j=1}^m |D^2 u_j|^2 dx + C(\kappa^* + \eta^2) r^{n-2\epsilon},$$

which is the desired inequality.

*Case 3:*  $r \leq s \leq R$ . By Case 1, there exists  $C = C(n, m, \Lambda, \lambda, \alpha)$  such that

$$\sum_{j=1}^m \int_{B_r(0)} |D^2 u_j|^2 dx \leq C \left( \frac{r}{s} \right)^{n-\epsilon} \int_{B_s(0)} \sum_{j=1}^m |D^2 u_j|^2 dx + C(\kappa^* + \eta^2) r^{n-2\epsilon}.$$

Then by Case 2,

$$\sum_{j=1}^m \int_{B_s(0)} |D^2 u_j|^2 dx \leq C \left( \frac{s}{R} \right)^{n-\epsilon} \int_{B_R(0)} \sum_{j=1}^m |D^2 u_j|^2 dx + C(\kappa^* + \eta^2) s^{n-2\epsilon}.$$

The above two inequalities give that

$$\sum_{j=1}^m \int_{B_r(0)} |D^2 u_j|^2 dx \leq C \left( \frac{r}{R} \right)^{n-\epsilon} \int_{B_R(0)} \sum_{j=1}^m |D^2 u_j|^2 dx + C(\kappa^* + \eta^2) r^{n-2\epsilon}.$$

This completes the proof of Proposition 4.4.  $\square$

**4.2. Improved order estimates.** Let  $u \in W^{1,2}(B_1(0))$ . Then the function defined by

$$X(\in \mathbb{R}^n) \mapsto \int_{B_1(0)} |Du - X|^2 dx$$

is convex and has a minimum at  $X = (D^2 u)_{B_1(0)}$ ; that is

$$(4.21) \quad \int_{B_1(0)} |Du - (Du)_{B_1(0)}|^2 dx \leq \int_{B_1(0)} |Du - X|^2 dx, \quad \forall X \in \mathbb{R}^n.$$

We now adapt the argument of Hölder estimates for gradients in [7, Theorem 3.13] to give the improved order estimates.

**Lemma 4.5.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs on  $B_t(0)$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. Suppose that  $\{u_j\}$  is connected in  $B_s(0)$  ( $s < t$ ) with*

$$(4.22) \quad \sum_{j=1}^m \int_{B_t} |D^2 u_j|^2 dx \leq \mathcal{K} t^{n-\alpha}$$

for some constant  $\mathcal{K}$ . Then for  $s \leq r < R \leq t$ ,

$$\begin{aligned} \int_{B_r(0)} |DD_k \varphi - (DD_k \varphi)_{B_r(0)}|^2 dx &\leq C \left( \frac{r}{R} \right)^{n+1} \int_{B_R(0)} |DD_k \varphi - (DD_k \varphi)_{B_R(0)}|^2 dx + \\ &\quad + C(\mathcal{K} + \kappa^* + \eta^2) r^{n+\alpha}. \end{aligned}$$

where  $C$  is a constant depending only on  $n, m, \Lambda, \lambda, \alpha$ .

*Proof.* So long as  $r \geq t/16$ , then  $R/r \leq 16$  and by (4.21),

$$\begin{aligned} \int_{B_r(0)} |D^2\varphi - (D^2\varphi)_{B_r(0)}|^2 dx &\leq \int_{B_r(0)} |D^2\varphi - (D^2\varphi)_{B_R(0)}|^2 dx \\ &\leq 16^{n+1} \left(\frac{r}{R}\right)^{n+1} \int_{B_R(0)} |D^2\varphi - (D^2\varphi)_{B_R(0)}|^2 dx. \end{aligned}$$

Now we assume that  $s < t/16$  so that Lemma 3.4 applies. By Proposition 4.4, there exists a constant  $\mathcal{B} = \mathcal{B}(n, m, \Lambda, \lambda, \alpha)$  so that for  $r < t$ ,

$$\sum_{j=1}^m \int_{B_r(0)} |D^2 u_j|^2 dx \leq \mathcal{B} \left(\frac{r}{t}\right)^{n-\epsilon} \sum_{j=1}^m \int_{B_t(0)} |D^2 u_j|^2 dx + \mathcal{B}(\kappa^* + \eta^2) r^{n-2\epsilon}.$$

Together with (4.22) and  $2\epsilon \leq \alpha$ , we conclude that

$$(4.23) \quad \sum_{j=1}^m \int_{B_r(0)} |D^2 u_j|^2 dx \leq \mathcal{B}(\mathcal{K} + \kappa^* + \eta^2) r^{n-\alpha}.$$

Recall that by (4.2),  $\varphi$  satisfies the equation

$$(4.24) \quad -\operatorname{div}\left(\mathbf{A}^* D D_k \varphi + \mathbf{b}^* D_k \varphi\right) + \mathbf{c}^* D D_k \varphi + d^* D_k \varphi = -\operatorname{div}\left(\hat{\mathbf{f}} - m \partial_{x_k} \mathcal{A}(x, \varphi, D\varphi)\right) + \hat{g}.$$

We now fix  $1 \leq k \leq n$ . For any  $R < t$ , let  $f$  be the weak solution of

$$\operatorname{div}\left(\mathbf{A}^*(0) D f\right) = 0, \quad f|_{\partial B_R(0)} = D_k \varphi - D_k \varphi(0).$$

Note that by (4.21),

$$(4.25) \quad \int_{B_R(0)} |D f - (D f)_{B_R(0)}|^2 dx \leq \int_{B_R(0)} |D f - (D D_k \varphi)_{B_R(0)}|^2 dx.$$

By harmonicity of  $f$  [7, Lemma 3.10; 8, Lemma 14.4.5],

$$(4.26) \quad \int_{B_r(0)} |D f - (D f)_{B_r(0)}|^2 dx \leq A(n) \left(\frac{r}{R}\right)^{n+2} \int_{B_R(0)} |D f - (D f)_{B_R(0)}|^2 dx.$$

Then we claim that

$$(4.27) \quad \int_{B_R(0)} |D f - (D D_k \varphi)_{B_R(0)}|^2 dx \leq C(m, \Lambda, \lambda) \int_{B_R(0)} |D D_k \varphi - (D D_k \varphi)_{B_R(0)}|^2 dx.$$

Indeed, since  $\operatorname{div} \mathbf{A}^*(0) (D D_k \varphi)_{B_R(0)} = 0$ , then by the definition of  $f$ ,

$$\operatorname{div}\left(\mathbf{A}^*(0) (D f - (D D_k \varphi)_{B_R(0)})\right) = 0.$$

By taking  $\phi := D_k\varphi - D_k\varphi(0) - f$  as a test function, we obtain

$$\begin{aligned}
& \int_{B_R(0)} |Df - (DD_k\varphi)_{B_R(0)}|^2 dx \\
& \leq C(m, \lambda) \int_{B_R(0)} \mathbf{A}^*(0) \left( Df - (DD_k\varphi)_{B_R(0)} \right) \left( Df - (DD_k\varphi)_{B_R(0)} \right) dx \\
& = C(m, \lambda) \int_{B_R(0)} \mathbf{A}^*(0) \left( Df - (DD_k\varphi)_{B_R(0)} \right) \left( DD_k\varphi - (DD_k\varphi)_{B_R(0)} \right) dx \\
& \leq C(m, \Lambda, \lambda) \int_{B_R(0)} |DD_k\varphi - (DD_k\varphi)_{B_R(0)}|^2 dx + \frac{1}{2} \int_{B_R(0)} |Df - (DD_k\varphi)_{B_R(0)}|^2 dx,
\end{aligned}$$

where we used (4.4) in the last inequality. Then (4.27) is proved. Combining (4.25) and (4.27), we obtain

$$(4.28) \quad \int_{B_R(0)} |Df - (Df)_{B_R(0)}|^2 dx \leq C(m, \Lambda, \lambda) \int_{B_R(0)} |DD_k\varphi - (DD_k\varphi)_{B_R(0)}|^2 dx.$$

Observe that

$$(4.29) \quad \int_{B_r(0)} |(Df)_{B_r(0)} - (DD_k\varphi)_{B_r(0)}|^2 dx \leq \int_{B_r(0)} |Df - DD_k\varphi|^2 dx.$$

Then we have

$$\begin{aligned}
& \int_{B_r(0)} |DD_k\varphi - (DD_k\varphi)_{B_r(0)}|^2 dx \\
& \leq 3 \int_{B_r(0)} |DD_k\varphi - Df|^2 + |Df - (Df)_{B_r(0)}|^2 + |(Df)_{B_r(0)} - (DD_k\varphi)_{B_r(0)}|^2 dx \\
& \leq 6 \int_{B_r(0)} |DD_k\varphi - Df|^2 dx + 3 \int_{B_r(0)} |Df - (Df)_{B_r(0)}|^2 dx.
\end{aligned}$$

Plugging (4.19) and (4.26) in it, we obtain

$$\begin{aligned}
\int_{B_r(0)} |DD_k\varphi - (DD_k\varphi)_{B_r(0)}|^2 dx & \leq C \int_{B_R(0)} \left( R^{2\alpha} |D^2\varphi|^2 + (\kappa^* + \eta^2) R^\alpha \right) dx + \\
& \quad + 3A(n) \left( \frac{r}{R} \right)^{n+2} \int_{B_R(0)} |Df - (Df)_{B_R(0)}|^2 dx,
\end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda)$ . This together with (4.23) and (4.28) gives that

$$\begin{aligned}
\int_{B_r(0)} |DD_k\varphi - (DD_k\varphi)_{B_r(0)}|^2 dx & \leq 3A(n) \left( \frac{r}{R} \right)^{n+2} \int_{B_R(0)} |DD_k\varphi - (DD_k\varphi)_{B_R(0)}|^2 dx \\
& \quad + C(\mathcal{K} + \kappa^* + \eta^2) R^{n+\alpha},
\end{aligned}$$

where we used the fact that  $\delta_0, \alpha \leq 1$ . Then by Lemma 4.1, there exists  $C = C(n, A, \alpha)$  so that for any  $r < R \leq t$ ,

$$\begin{aligned}
\int_{B_r(0)} |DD_k\varphi - (DD_k\varphi)_{B_r(0)}|^2 dx & \leq C \left( \frac{r}{R} \right)^{n+1} \int_{B_R(0)} |DD_k\varphi - (DD_k\varphi)_{B_R(0)}|^2 dx + \\
& \quad + C(\mathcal{K} + \kappa^* + \eta^2) r^{n+\alpha}.
\end{aligned}$$

This completes the proof.  $\square$



So far, we have proved the improved order estimates over some tilt-plane for any given  $r$  so that  $\{u_j\}$  is connected. The following gives that the tilt-plane will be close to each other in the order of  $\alpha$ .

**Lemma 4.6.** *Let  $\{u_j\}_{j=1}^m$ ,  $(\mathcal{Q}, \{g_j\})$ ,  $\mathcal{K}$ ,  $t > 0$  be the same as in Lemma 4.5. In particular, we assume that  $\{u_j\}$  is connected in  $B_s(0)$ . Let  $C_1$  be the constant in Lemma 4.5. Then for  $s \leq r \leq R \leq t$ ,*

$$|(D^2\varphi)_{B_R(0)} - (D^2\varphi)_{B_r(0)}|^2 \leq C_2 \left( \frac{R}{t^{n+1}} \int_{B_t(0)} |D^2\varphi - (D^2\varphi)_{B_t(0)}|^2 dx + (\mathcal{K} + \kappa^* + \eta^2)R^\alpha \right),$$

where  $C_2 = C_2(n, \alpha, C_1)$ .

*Proof.* For  $r \in [s, t]$ , let

$$\sigma(r) := \int_{B_r(0)} |D^2\varphi - (D^2\varphi)_{B_r(0)}|^2 dx.$$

Then by Lemma 4.5,

$$\sigma(r) \leq C_1 \left( \frac{r}{t} \right)^{n+1} \sigma(t) + C_1 (\mathcal{K} + \kappa^* + \eta^2) r^{n+\alpha}, \quad C_1 = C_1(n, m, \Lambda, \lambda, \alpha).$$

For any  $s \leq r < R \leq t$ , we then have

$$\begin{aligned} |(D^2\varphi)_{B_R(0)} - (D^2\varphi)_{B_r(0)}|^2 &\leq \frac{2}{|B_r|} \int_{B_r(0)} \left( |D^2\varphi - (D^2\varphi)_{B_R(0)}|^2 + |D^2\varphi - (D^2\varphi)_{B_r(0)}|^2 \right) \\ &\leq \frac{2}{|B_r|} (\sigma(R) + \sigma(r)) \\ &\leq \frac{4C_1}{|B_r|} \left[ \left( \frac{R}{t} \right)^{n+1} \sigma(t) + (\mathcal{K} + \kappa^* + \eta^2) R^{n+\alpha} \right]. \end{aligned}$$

In particular, for  $s \leq \hat{R}_i = 2^{-i}R$ ,

$$\left| (D^2\varphi)_{B_{\hat{R}_i}(0)} - (D^2\varphi)_{B_{\hat{R}_{i+1}}(0)} \right|^2 \leq \frac{2^{n+2}C_1}{|B_1|} \left( \frac{\hat{R}_i}{t^{n+1}} \cdot \sigma(t) + (\mathcal{K} + \kappa^* + \eta^2) \hat{R}_i^\alpha \right).$$

It then implies that

$$\begin{aligned} |(D^2\varphi)_{B_R(0)} - (D^2\varphi)_{B_{\hat{R}_k}(0)}|^2 &\leq \left( \sum_{i=0}^{k-1} \left| (D^2\varphi)_{B_{\hat{R}_i}(0)} - (D^2\varphi)_{B_{\hat{R}_{i+1}}(0)} \right| \right)^2 \\ &\leq \frac{2^{n+2}C_1}{|B_1|} \left[ \sum_{i=0}^{k-1} \left( \sqrt{\frac{\sigma(t)}{t^{n+1}}} \cdot \hat{R}_i^{\frac{1}{2}} + \sqrt{\mathcal{K} + \kappa^* + \eta^2} \hat{R}_i^{\frac{\alpha}{2}} \right) \right]^2 \\ &\leq \frac{2^{n+2}C_1}{|B_1|} \left[ \frac{1}{1 - 2^{-\frac{\alpha}{2}}} \left( \sqrt{\frac{\sigma(t)}{t^{n+1}}} \cdot R^{\frac{1}{2}} + \sqrt{\mathcal{K} + \kappa^* + \eta^2} R^{\frac{\alpha}{2}} \right) \right]^2 \\ &\leq \frac{2^{n+3}C_1}{|B_1|} \cdot (1 - 2^{-\frac{\alpha}{2}})^{-2} \left( \frac{\sigma(t)}{t^{n+1}} \cdot R + (\mathcal{K} + \kappa^* + \eta^2) R^\alpha \right). \end{aligned}$$

Suppose that  $\hat{R}_{k+1} \leq r < \hat{R}_k$ . Then

$$\begin{aligned} |(D^2\varphi)_{B_R(0)} - (D^2\varphi)_{B_r(0)}|^2 &\leq 2|(D^2\varphi)_{B_R(0)} - (D^2\varphi)_{B_{\hat{R}_k}(0)}|^2 + 2|(D^2\varphi)_{B_{\hat{R}_k}(0)} - (D^2\varphi)_{B_r(0)}|^2 \\ &\leq \frac{2^{n+4}C_1}{|B_1|} \cdot (1 - 2^{-\frac{\alpha}{2}})^{-2} \left( \frac{\sigma(t)}{t^{n+1}} \cdot R + (\mathcal{K} + \kappa^* + \eta^2)R^\alpha \right) + \\ &\quad + \frac{8C_1}{|B_r|} \left[ \left( \frac{\hat{R}_k}{t} \right)^{n+1} \sigma(t) + (\mathcal{K} + \kappa^* + \eta^2)\hat{R}_k^{n+\alpha} \right] \\ &\leq \frac{2^{n+5}C_1}{|B_1|} \cdot (1 - 2^{-\frac{\alpha}{2}})^{-2} \left( \frac{\sigma(t)}{t^{n+1}} \cdot R + (\mathcal{K} + \kappa^* + \eta^2)R^\alpha \right). \end{aligned}$$

Then the lemma follows by taking

$$C_2 = \frac{2^{n+5}C_1}{|B_1|} \cdot (1 - 2^{-\frac{\alpha}{2}})^{-2}.$$

□

Recall that

$$\sigma^*(r) := \sum_{j=1}^m \int_{B_r(0)} |D^2u_j - (D^2u_j)_{B_r(0)}|^2 dx.$$

Lemma 4.5 combining with Lemma 3.4 implies the order estimates of  $\sigma^*(r)$ .

**Lemma 4.7.** *Let  $\{u_j\}_{j=1}^m$ ,  $(\mathcal{Q}, \{g_j\})$ ,  $\mathcal{K}$ ,  $t > 0$ ,  $\kappa^*$  be the same as in Lemma 4.5. Then there exists a constant  $C$  depending only on  $n, m, \Lambda, \lambda, \alpha$ , so that for all  $r \leq t$ ,*

$$\sigma^*(r) \leq C \left( \frac{r}{t} \right)^{n+1} \sigma^*(t) + C(\mathcal{K} + \kappa^* + \eta^2)r^n.$$

*Proof.* So long as  $r \geq t/16$ , then  $t/r \leq 16$  and by (4.21),

$$\sigma^*(r) \leq \sum_{j=1}^m \int_{B_r(0)} |D^2u_j - (D^2u_j)_{B_t(0)}|^2 dx \leq 16^{n+1} \left( \frac{r}{t} \right)^{n+1} \sigma^*(t).$$

This is the desired inequality. We now assume that  $r \leq t/16$  and then Lemma 3.4 can be applied.

We prove the lemma inductively. When  $m = 1$ , then the desired inequality follows from Lemma 4.5 by taking  $R = t$ .

Now we assume that the lemma is true when the number of the sheets is less than or equal to  $m - 1$ . We also assume that  $\{u_j\}$  is connected in  $B_t(0)$ ; otherwise, the desired inequality follows from induction. Let

$$s := \inf\{r > 0; \{u_j\} \text{ is connected in } B_r(0)\}.$$

By Lemma 4.5, there exists  $C = C(n, m, \Lambda, \lambda, \alpha)$  such that for  $s < r \leq t$ ,

$$\begin{aligned} &\int_{B_r(0)} |DD_k\varphi - (DD_k\varphi)_{B_r(0)}|^2 dx \\ &\leq C \left( \frac{r}{t} \right)^{n+1} \int_{B_t(0)} |DD_k\varphi - (DD_k\varphi)_{B_t(0)}|^2 dx + C(\mathcal{K} + \kappa^* + \eta^2)r^{n+\alpha} \\ &\leq C \left( \frac{r}{t} \right)^{n+1} \sigma^*(t) + C(\mathcal{K} + \kappa^* + \eta^2)r^{n+\alpha}. \end{aligned}$$

Recall that by Lemma 3.4, for  $s < r \leq t$ ,

$$\begin{aligned}
(4.30) \quad & \int_{B_r(0)} |D^2 u_i - D^2 u_j - (D^2 u_i)_{B_r(0)} + (D^2 u_j)_{B_r(0)}|^2 dx \\
& \leq 2 \int_{B_r(0)} |D^2 u_i - D^2 u_j|^2 dx + 2|B_r| \cdot |(D^2 u_i - D^2 u_j)_{B_r(0)}|^2 \\
& \leq 4 \int_{B_r(0)} |D^2 u_i - D^2 u_j|^2 dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right) r^n.
\end{aligned}$$

Thus we obtain that for  $s < r \leq t$ ,

$$\begin{aligned}
& \int_{B_r(0)} |D^2 u_i - (D^2 u_i)_{B_r(0)}|^2 dx \\
& \leq \int_{B_r(0)} \sum_{j=1}^m \left( |D^2 u_i - D^2 u_j - (D^2 u_i)_{B_r(0)} + (D^2 u_j)_{B_r(0)}|^2 \right) + m |D^2 \varphi - (D^2 \varphi)_{B_r(0)}|^2 dx \\
& \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right) r^n + C \left( \frac{r}{t} \right)^{n+1} \sigma^*(t) + C(\mathcal{K} + \kappa^* + \eta^2) r^{n+\alpha} \\
& \leq C \left( \frac{r}{t} \right)^{n+1} \sigma^*(t) + C(\mathcal{K} + \kappa^* + \eta^2) r^n,
\end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$ . This implies the desired inequality.

It remains to consider the case of  $r < s$ . Observe that we have proved that

$$(4.31) \quad \sigma^*(s) \leq C \left( \frac{s}{t} \right)^{n+1} \sigma^*(t) + C(\mathcal{K} + \kappa^* + \eta^2) s^n.$$

By Proposition 4.4, there exists a constant  $\mathcal{B} = \mathcal{B}(n, m, \Lambda, \lambda, \alpha)$  so that

$$\sum_{j=1}^m \int_{B_s(0)} |D^2 u_j|^2 dx \leq \mathcal{B} \left( \frac{s}{t} \right)^{n-\epsilon} \int_{B_t(0)} \sum_{j=1}^m |D^2 u_j|^2 dx + \mathcal{B}(\kappa^* + \eta^2) s^{n-2\epsilon}.$$

Together with (4.22) and  $2\epsilon \leq \alpha$ , we obtain that

$$(4.32) \quad \sum_{j=1}^m \int_{B_s(0)} |D^2 u_j|^2 dx \leq \mathcal{B}(\mathcal{K} + \kappa^* + \eta^2) s^{n-\alpha}.$$

Let  $\mathcal{K}' := \mathcal{B}(\mathcal{K} + \kappa^* + \eta^2)$ . Note that in  $B_s(0)$ ,  $\{u_j\}_{j=1}^m$  can be divided into two disjoint ordered graphs  $\{u_j\}_{j=1}^i$  and  $\{u_j\}_{j=i+1}^m$ . Thus by induction using (4.32) in place of (4.22), we have for all  $r < s$ ,

$$\begin{aligned}
& \sum_{j=1}^i \int_{B_r(0)} |D^2 u_j - (D^2 u_j)_{B_r(0)}|^2 dx \leq C \left( \frac{r}{s} \right)^{n+1} \sum_{j=1}^i \int_{B_s(0)} |D^2 u_j - (D^2 u_j)_{B_s(0)}|^2 dx + \\
& \quad + C(\mathcal{K}' + \kappa^* + \eta^2) r^n; \\
& \sum_{j=i+1}^m \int_{B_r(0)} |D^2 u_j - (D^2 u_j)_{B_r(0)}|^2 dx \leq C \left( \frac{r}{s} \right)^{n+1} \sum_{j=i+1}^m \int_{B_s(0)} |D^2 u_j - (D^2 u_j)_{B_s(0)}|^2 dx + \\
& \quad + C(\mathcal{K}' + \kappa^* + \eta^2) r^n.
\end{aligned}$$

The taking the sum of the two inequalities above, we obtain

$$\sigma^*(r) \leq C \left(\frac{r}{s}\right)^{n+1} \sigma^*(s) + C(\mathcal{K} + \kappa^* + \eta^2)r^n,$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$ . Plugging (4.31) into it, we obtain that

$$\begin{aligned} \sigma^*(r) &\leq C \left(\frac{r}{s}\right)^{n+1} \left[ \left(\frac{s}{t}\right)^{n+1} \sigma^*(t) + C(\mathcal{K} + \kappa^* + \eta^2)s^n \right] + C(\mathcal{K} + \kappa^* + \eta^2)r^n \\ &\leq C \left(\frac{r}{t}\right)^{n+1} \sigma^*(t) + C(\mathcal{K} + \kappa^* + \eta^2)r^n. \end{aligned}$$

This finishes the proof of Lemma 4.7.  $\square$

**4.3. Monotonicity of the the second derivatives.** Now we are ready to prove a monotonicity formula for the average of  $L^2$ -integral of  $D^2u_j$ . When  $m = 1$ , the desired result follows from Lemma 4.6 and Lemma 4.7. For  $m \geq 2$ , we will use Lemma 3.4 and run an inductive method.

**Proposition 4.8.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs on  $B_t(0)$  ( $t \leq 1$ ) which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. For simplicity, denote by*

$$\bar{\sigma}(r) := \sum_{j=1}^m \frac{1}{|B_r|} \int_{B_r(0)} |D^2u_j|^2 dx, \quad \forall r \leq t.$$

Then there exists a constant  $C = C(n, m, \Lambda, \lambda, \alpha)$  so that

$$\bar{\sigma}(r) \leq C(\bar{\sigma}(t) + \kappa^* + \eta^2).$$

*Proof.* We first note that

$$(4.33) \quad \sum_{j=1}^m |(D^2u_j)_{B_t(0)}|^2 \leq \sum_{j=1}^m \frac{1}{|B_t|} \int_{B_t(0)} |D^2u_j|^2 dx = \bar{\sigma}(t).$$

By (4.21), we have that

$$(4.34) \quad \frac{1}{|B_r|} \cdot \sigma^*(r) = \sum_{j=1}^m \frac{1}{|B_r|} \int_{B_r(0)} |D^2u_j - (D^2u_j)_{B_r(0)}|^2 dx \leq \bar{\sigma}(r).$$

We prove the proposition inductively. When  $m = 1$ , we have by (4.33),

$$\begin{aligned} \int_{B_r(0)} |D^2u|^2 dx &\leq 3 \int_{B_r(0)} |D^2u - (D^2u)_{B_r(0)}|^2 + |(D^2u)_{B_r(0)} - (D^2u)_{B_t(0)}|^2 + |(D^2u)_{B_t(0)}|^2 dx \\ &\leq 3\sigma^*(r) + 3|B_r| \cdot |(D^2u)_{B_r(0)} - (D^2u)_{B_t(0)}|^2 + 3|B_r|\bar{\sigma}(t). \end{aligned}$$

Applying Lemma 4.6 and Lemma 4.7,

$$\begin{aligned} \int_{B_r(0)} |D^2u|^2 dx &\leq C \left(\frac{r}{t}\right)^{n+1} \sigma^*(t) + C(\bar{\sigma}(t) + \kappa^* + \eta^2)r^n + \\ &\quad + C|B_r| \left( \frac{t}{t^{n+1}} \sigma^*(t) + (\bar{\sigma}(t) + \kappa^* + \eta^2)t^\alpha \right) + C\bar{\sigma}(t) \cdot r^n. \end{aligned}$$

This together with (4.34) implies that

$$\int_{B_r(0)} |D^2u|^2 dx \leq C(\bar{\sigma}(t) + \kappa^* + \eta^2)r^n.$$

Then the desired inequality follows immediately.

Now suppose that the lemma is true when the number of sheets is less than or equal to  $m - 1$ . We also assume that  $\{u_j\}$  is connected in  $B_t(0)$ ; otherwise, the desired inequality follows from induction. Let

$$s := \inf\{r > 0; \{u_j\} \text{ is connected in } B_r(0)\}.$$

*Case 1:*  $s \leq r \leq t$ . Without loss of generality, we assume that  $r < t/32$  so that Lemma 3.4 applies. Then by Lemma 4.6, for  $s \leq r \leq t$ ,

$$(4.35) \quad \begin{aligned} & |(D^2\varphi)_{B_t(0)} - (D^2\varphi)_{B_r(0)}|^2 \\ & \leq C \left( \frac{t}{t^{n+1}} \int_{B_t(0)} |D^2\varphi - (D^2\varphi)_{B_t(0)}|^2 dx + (\bar{\sigma}(t) + \kappa^* + \eta^2)t^\alpha \right) \\ & \leq C\bar{\sigma}(t) + C(\kappa^* + \eta^2)t^\alpha, \end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$  and we used (4.34) in the last inequality. On the other hand, by Lemma 4.7 and (4.34), there exists a constant  $C$  depending only on  $n, m, \Lambda, \lambda, \alpha$  so that for all  $r \leq t$ ,

$$\begin{aligned} \int_{B_r(0)} |D^2\varphi - (D^2\varphi)_{B_r(0)}|^2 dx & \leq C \left( \frac{r}{t} \right)^{n+1} \sigma^*(t) + C(\bar{\sigma}(t) + \kappa^* + \eta^2)r^n \\ & \leq C(\bar{\sigma}(t) + \kappa^* + \eta^2)r^n. \end{aligned}$$

This together with (4.35) and (4.33) implies that

$$\begin{aligned} \int_{B_r(0)} |D^2\varphi|^2 dx & \leq 3 \int_{B_r(0)} |D^2\varphi - (D^2\varphi)_{B_r(0)}|^2 + |(D^2\varphi)_{B_r(0)} - (D^2\varphi)_{B_t(0)}|^2 + |(D^2\varphi)_{B_t(0)}|^2 dx \\ & \leq C(\bar{\sigma}(t) + \kappa^* + \eta^2)r^n. \end{aligned}$$

Recall that by Lemma 3.4, there exists  $C = C(n, m, \Lambda, \lambda)$  so that

$$\int_{B_r(0)} |D^2u_i - D^2u_j|^2 dx \leq C \cdot \left( (1 + \eta^2)(\kappa + \kappa_0)^2 + \kappa_1^2 \right) r^n \leq C \cdot (\kappa^* + \eta^2)r^n.$$

We then conclude that

$$\begin{aligned} \int_{B_r(0)} |D^2u_i|^2 dx & \leq 2 \int_{B_r(0)} |D^2\varphi - D^2u_i|^2 + |D^2\varphi|^2 dx \\ & \leq C(m) \sum_{j=1}^m \int_{B_r(0)} |D^2u_j - D^2u_i|^2 + |D^2\varphi|^2 dx \\ & \leq C(\bar{\sigma}(t) + \kappa^* + \eta^2)r^n, \end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$ . This is the desired inequality.

*Case 2:*  $r < s$ . By the definition of  $s$ ,  $\{u_j\}_{j=1}^m$  can be divided into two ordered graphs  $\{u_j\}_{j=1}^i$  and  $\{u_j\}_{j=i+1}^m$  in  $B_s(0)$ . Thus by induction, there exists  $C = C(n, m, \Lambda, \lambda, \alpha)$  so that

$$\begin{aligned} \sum_{j=1}^i \frac{1}{|B_r|} \int_{B_r(0)} |D^2u_j|^2 dx & \leq C \left( \sum_{j=1}^i \frac{1}{|B_s|} \int_{B_s(0)} |D^2u_j|^2 dx + \kappa^* + \eta^2 \right); \\ \sum_{j=i+1}^m \frac{1}{|B_r|} \int_{B_r(0)} |D^2u_j|^2 dx & \leq C \left( \sum_{j=i+1}^m \frac{1}{|B_s|} \int_{B_s(0)} |D^2u_j|^2 dx + \kappa^* + \eta^2 \right). \end{aligned}$$

Combining the two inequalities together, we conclude that

$$(4.36) \quad \bar{\sigma}(r) \leq C(\bar{\sigma}(s) + \kappa^* + \eta^2).$$

Observe that by Case 1,

$$\bar{\sigma}(s) \leq C(\bar{\sigma}(t) + \kappa^* + \eta^2).$$

Plugging this into (4.36), then the desired inequality follows immediately. This finishes the proof of Proposition 4.8.  $\square$

## 5. LIPSCHITZ UPPER BOUND FOR THE FIRST DERIVATIVES

In this section, we will give the upper bound of the Lipschitz norm of the first derivatives of ordered graphs in Theorem 1.2.

Note that we have proved the uniform upper bound for the  $\bar{\sigma}(r)$  for all small  $r$ . This together with Poincaré inequality (Lemma 5.1) gives that the graphs  $\{u_j\}$  will be close to a plane  $P$  in the sense of  $L^2$ ; see Lemma 5.2. We will then consider the differential equation of the difference to the plane  $P$ . Applying the  $C^{1,\alpha}$  estimates to the divergence form, we will then obtain the upper bound of the derivative of such new functions by  $\bar{\sigma}(r)$  and  $\kappa^*$ . We remark that Lemma 3.1 and Lemma 3.2 are used to bound the remainder terms in this process. To finish the proof, we will apply Proposition 4.8 and Proposition 3.5 to bound  $\bar{\sigma}(r)$  by  $\kappa^*$  and  $\eta$ ; see Theorem 5.5 for more details.

Recall that

$$\kappa^* := \kappa^2 + \kappa_0^2 + \kappa_1^2 + \delta_0^2,$$

and

$$\eta := \left( \|\partial_z \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_1)} + \|\partial_x \mathcal{B}\|_{C^1(\mathcal{U}_1)} \right) \cdot \sum_{j=1}^m \|u_j\|_{C^0(B_1(0))}.$$

**5.1. The first derivative upper bound.** Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs on  $B_t(0)$  ( $t \leq 1$ ) which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Recall that

$$\bar{\sigma}(r) := \sum_{j=1}^m \frac{1}{|B_r|} \int_{B_r(0)} |D^2 u_j|^2 dx.$$

The following estimates are well-known and we provide it here for the readers' convenience.

**Lemma 5.1** (Poincaré inequalities [6, Estimate (7.45)]). *For any convex domain  $\Omega$  and  $u \in W^{1,p}(\Omega)$ ,  $1 \leq p < +\infty$ ,*

$$\|u - u_{B_r}\|_{L^p(\Omega)} \leq r^{1-n} d^n \|Du\|_{L^p(\Omega)}, \quad d = \text{diam } \Omega.$$

This yields the following inequalities.

**Lemma 5.2.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha} \cap W^{2,2}$  ordered graphs on  $B_t(0)$  ( $t \leq 1$ ). Then for  $r \leq t$ ,*

$$\begin{aligned} \sum_{j=1}^m \frac{1}{|B_r|} \int_{B_r(0)} |Du_j(x) - (Du_j)_{B_r(0)}|^2 dx &\leq 2^{2n} \bar{\sigma}(r) r^2; \\ \sum_{j=1}^m \frac{1}{|B_r|} \int_{B_r(0)} |u_j(x) - (u_j)_{B_r(0)} - (Du_j)_{B_r(0)} \cdot x|^2 dx &\leq 2^{4n} \bar{\sigma}(r) r^4. \end{aligned}$$

Now we are going to use the equation to bound oscillations. We will use the following notions. Let  $\{u_j\}$  be ordered graphs on  $B_r(0)$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Recall that by (2.4),

$$(5.1) \quad m\mathcal{Q}\varphi = \operatorname{div} \mathbf{f}(\{u_j\}) - g(\{u_j\}, \{g_j\})$$

in  $B_r(0)$  in the weak sense, where  $\mathbf{f}(\{u_j\})$  and  $g(\{u_j\}, \{g_j\})$  are defined in (2.5) and (2.6), respectively. Let

$$\varphi_r = (\varphi)_{B_r(0)} + (D\varphi)_{B_r(0)} \cdot x.$$

Note that

$$\begin{aligned} \mathcal{Q}\varphi_r &:= -\operatorname{div} \mathcal{A}(x, \varphi_r, D\varphi_r) + \mathcal{B}(x, \varphi_r, D\varphi_r) \\ &= -\operatorname{tr}(\partial_x \mathcal{A}(x, \varphi_r, D\varphi_r)) - \partial_z \mathcal{A}(x, \varphi_r, D\varphi_r) \cdot (D\varphi)_{B_r(0)} + \mathcal{B}(x, \varphi_r, D\varphi_r). \end{aligned}$$

**Lemma 5.3.** *There exists a constant  $C = C(n, m, \Lambda, \lambda)$  such that for all  $x \in B_r(0)$ ,*

$$(5.2) \quad -C(\delta_0 + \eta) \leq \mathcal{Q}\varphi_r \leq C(\delta_0 + \eta),$$

*Proof.* Since  $\mathcal{A}(x, 0, 0) = 0$ , then

$$\begin{aligned} |\partial_x \mathcal{A}(x, \varphi_r, D\varphi_r)| &\leq \|\partial_z \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_r)} |\varphi_r| + \|\partial_p \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_r)} |D\varphi_r| \\ &\leq \|\partial_z \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_r)} \left( |\varphi|_{C^0(B_r(0))} + r |D\varphi|_{C^0(B_r(0))} \right) + \|\partial_p \partial_x \mathcal{A}\|_{C^0(\mathcal{U}_r)} |D\varphi|_{C^0(B_r(0))} \\ &\leq C(\eta + \delta_0). \end{aligned}$$

Similarly,

$$|\mathcal{B}(x, \varphi_r, D\varphi_r)| \leq C(\eta + \delta_0).$$

On the other hand,

$$|\partial_z \mathcal{A}(x, \varphi_r, D\varphi_r) \cdot (D\varphi)_{B_r(0)}| \leq C\delta_0.$$

Hence the lemma is proved.  $\square$

For simplicity, we denote

$$\begin{aligned} \Psi(x) &:= \varphi(x) - (\varphi)_{B_r(0)} - (D\varphi)_{B_r(0)} \cdot x; \\ \Psi_j(x) &:= u_j(x) - (u_j)_{B_r(0)} - (Du_j)_{B_r(0)} \cdot x. \end{aligned}$$

By (5.1) and Lemma 2.6, we obtain

$$(5.3) \quad \begin{aligned} \operatorname{div}(\mathbf{A}(\varphi, \varphi_r)D\Psi + \mathbf{b}(\varphi, \varphi_r)\Psi) + \mathbf{c}(\varphi, \varphi_r)D\Psi + d(\varphi, \varphi_r)\Psi \\ = \operatorname{div} \mathbf{f}(\{u_j\}) - g(\{u_j\}, \{g_j\}) - m\mathcal{Q}\varphi_r, \end{aligned}$$

where  $\mathbf{f}, g$  are the same in (5.1).

The following result gives that the oscillation of  $\Psi_j$  has the second order. Plugging this into the interior  $C^{1,\alpha}$  estimates gives the oscillation of  $Du_j$  in small balls. We remark that the interior estimates can not be applied to  $Du_j$  because  $u_j$  itself does not satisfy a weak partial differential equation.

**Lemma 5.4.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs on  $B_1(0)$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. There exists a constant  $C = C(n, m, \Lambda, \lambda, \alpha)$  so that for each  $1 \leq j \leq m$ ,*

$$\sup_{x \in B_{r/2}(0)} |u_j(x) - (u_j)_{B_r(0)} - (Du_j)_{B_r(0)} \cdot x| \leq C(\sqrt{\overline{\sigma}(r)} + \kappa + \kappa_0 + \delta_0 + \eta)r^2;$$

$$\sup_{x \in B_{r/4}(0)} |Du_j(x) - (Du_j)_{B_r(0)}| \leq C(\sqrt{\overline{\sigma}(r)} + \kappa + \kappa_0 + \delta_0 + \eta)r.$$

*Proof.* We divide the proof into two cases.

*Case I:*  $r \geq 1/16^{m+1}$ . Then by (2.1),

$$\begin{aligned} \sup_{x \in B_{r/2}(0)} |u_j(x) - (u_j)_{B_r(0)} - (Du_j)_{B_r(0)} \cdot x| &\leq C\delta_0 \leq C \cdot 16^{2m+2}r^2; \\ \sup_{x \in B_{r/4}(0)} |Du_j(x) - (Du_j)_{B_r(0)}| &\leq C\delta_0 \leq C \cdot 16^{2m+2}r^2. \end{aligned}$$

These are the desired inequalities.

*Case II:*  $r \leq 1/16^{m+1}$ . Applying Lemma 2.5, one can find  $R \in [r, 16^m r]$  such that  $\{u_j\}$  has the same connected components in  $B_{16R}(0)$  and  $B_R(0)$ . Choose one connected component and still denote it by  $\{u_j\}$ . By Lemma 3.1 and Lemma 3.2, there exists a constant  $C = C(n, m, \Lambda, \lambda)$  so that for all  $x \in B_R(0) \supset B_r(0)$ ,

$$(5.4) \quad |u_i - u_j| \leq C\kappa r^2; \quad |Du_i - Du_j| \leq C(\kappa + \kappa_0)r.$$

Let  $\mathbf{f}$  and  $g$  be the notions in (5.3). Then by Lemma 2.8 and (5.2),

$$(5.5) \quad |\mathbf{f}(\{u_j\})| \leq C(m, \Lambda)(\kappa + \kappa_0)^2 r^2;$$

$$(5.6) \quad \sup_{x, y \in B_r(0)} \frac{|\mathbf{f}(\{u_j\})(x) - \mathbf{f}(\{u_j\})(y)|}{|x - y|^\alpha} \leq C(m, \Lambda)(\kappa + \kappa_0)r;$$

$$(5.7) \quad \tilde{g} := |g(\{u_j\}, \{g_j\})| + m|\mathcal{Q}\varphi_r| \leq C(m, \Lambda)((\kappa + \kappa_0)r + \kappa_0 + \delta_0 + \eta).$$

Now we fix a constant  $q > n$  (e.g.  $q = n + 1$ ). By a direct computation,

$$\int_{B_r(0)} \|\mathbf{f}\|^q dx \leq C(\kappa + \kappa_0)^{2q} r^{n+2q}.$$

Thus by the local estimates for weak solutions [6, Theorem 8.17] with  $u(x) = \Psi(x)$ ,  $p = 2$  and  $2R = r$ , then there exists  $C = C(n, q)$  so that

$$\sup_{x \in B_{r/2}(0)} |\Psi(x)| \leq C(r^{-\frac{n}{2}} \|\Psi(x)\|_{L^2(B_r(0))} + r^{1-\frac{n}{q}} \|\mathbf{f}\|_{L^q(B_r(0))} + r^{2-\frac{2n}{q}} \|\tilde{g}\|_{L^{q/2}(B_r(0))}).$$

By Lemma 5.2 and the estimates for  $\mathbf{f}$  and  $\tilde{g}$  in (5.5) and (5.7), we hence conclude that

$$\begin{aligned} \sup_{x \in B_{r/2}(0)} |\Psi(x)| &\leq C \left( r^{-\frac{n}{2}} \cdot \sqrt{\sigma(r)} \cdot r^{2+\frac{n}{2}} + r^{1-\frac{n}{q}} (\kappa + \kappa_0)^2 r^{\frac{n}{q}+2} + \right. \\ (5.8) \quad &\quad \left. + r^{2-\frac{2n}{q}} ((\kappa + \kappa_0)r + \kappa_0 + \delta_0 + \eta) \cdot r^{\frac{2n}{q}} \right) \\ &\leq C \left( \sqrt{\sigma(r)} + \kappa + \kappa_0 + \delta_0 + \eta \right) r^2. \end{aligned}$$

On the other hand, by (5.4),

$$\begin{aligned} |\Psi_i - \Psi_j| &\leq |u_i - u_j| - |(u_i)_{B_r(0)} - (u_j)_{B_r(0)}| + |(Du_i)_{B_r(0)} - (Du_j)_{B_r(0)}| \cdot r \\ &\leq C(\kappa + \kappa_0)r^2, \end{aligned}$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$ . This together with (5.8) gives that for  $x \in B_{r/2}(0)$ ,

$$|\Psi_j| \leq |\Psi_j - \Psi| + |\Psi| \leq \frac{1}{m} \sum_{i=1}^m |\Psi_j - \Psi_i| + |\Psi| \leq C \left( \sqrt{\sigma(r)} + \kappa + \kappa_0 + \delta_0 + \eta \right) r^2.$$

This proves the first inequality in the lemma.



It remains to estimate the derivative of  $\Psi$ . Indeed, by the  $C^{1,\alpha}$  estimates Theorem A.1,

$$\begin{aligned} \sup_{x \in B_{r/4}(0)} |D\Psi| \leq C(n, m, \Lambda) & \left( \sup_{x \in B_{r/2}(0)} r^{-1} |\Psi(x)| + r^\alpha \sup_{x, x' \in B_{r/2}(0)} \frac{|\mathbf{f}(x) - \mathbf{f}(x')|}{|x - x'|^\alpha} + \right. \\ & \left. + \sup_{x \in B_{r/2}(0)} |\mathbf{f}(x)| + \sup_{x \in B_{r/2}(0)} r |\tilde{g}| \right). \end{aligned}$$

Plugging (5.8), (5.5), (5.6), and (5.7) into it, we conclude that

$$(5.9) \quad \begin{aligned} \sup_{x \in B_{r/4}(0)} |D\Psi| & \leq C \left( \sqrt{\bar{\sigma}(r)} + \kappa + \kappa_0 + \delta_0 + \eta \right) r + (\kappa + \kappa_0) r^{1+\alpha} + (\kappa + \kappa_0)^2 r^2 \\ & \leq C \left( \sqrt{\bar{\sigma}(r)} + \kappa + \kappa_0 + \delta_0 + \eta \right) r. \end{aligned}$$

On the other hand, (5.4) gives that for  $x \in B_r(0)$ ,

$$|D\Psi_j - D\Psi| = |Du_j - (Du_j)_{B_r(0)} - D\varphi - (D\varphi)_{B_r(0)}| \leq C(\kappa + \kappa_0)r.$$

Combining with (5.9), we conclude that for all  $x \in B_{r/4}(0)$ ,

$$|D\Psi_j| \leq C \left( \sqrt{\bar{\sigma}(r)} + \kappa + \kappa_0 + \delta_0 + \eta \right) r.$$

Hence the lemma is proved.  $\square$

**5.2. Lipschitz estimates of the first derivatives.** In this section, we will prove the Lipschitz upper bound for ordered graphs by using Lemma 5.4. The key step is to apply Proposition 4.8 and Proposition 3.5 to bound  $\bar{\sigma}(r)$  by  $\kappa^*$  and  $\eta$ .

**Theorem 5.5.** *Let  $\{u_j\}_{j=1}^m$  be  $C^{1,\alpha}$  ordered graphs on  $B_1(0)$  which is a solution to  $(\mathcal{Q}, \{g_j\})$ . Suppose that (1.1), (1.2), (1.3), (2.1) and (2.2) are satisfied. Then there exists  $C = C(n, m, \Lambda, \lambda, \alpha)$  so that for  $j = 1, \dots, m$ ,*

$$\sup_{x, y \in B_{1/2}(0)} \frac{|Du_j(x) - Du_j(y)|}{|x - y|} \leq C(\kappa + \kappa_0 + \kappa_1 + \delta_0 + \eta).$$

*Proof.* Let  $x_1, x_2 \in B_{1/2}(0)$  and  $r := |x_1 - x_2|$ . So long as  $r \geq 1/32$ , then by (2.1),

$$\frac{|Du_j(x_1) - Du_j(x_2)|}{|x_1 - x_2|} \leq \delta_0 \cdot |x_1 - x_2|^{\alpha-1} \leq 32\delta_0.$$

This is the desired inequality. Now we assume that  $r < 1/32$ . By Proposition 3.5, there exists  $C = C(n, m, \Lambda, \lambda)$  so that

$$(5.10) \quad \sum_{j=1}^m \int_{B_{3/4}(0)} |D^2 u_j|^2 dx \leq C(\kappa^* + \eta^2).$$

For each  $y \in B_{1/2}(0)$  and  $t \in (0, 1/2]$ , we define

$$\bar{\sigma}(t; y) := \sum_{j=1}^m \frac{1}{|B_t|} \int_{B_t(y)} |D^2 u_j|^2 dx.$$

Then (5.10) implies that for all  $y \in B_{1/2}(0)$ ,

$$\bar{\sigma}(1/4; y) \leq C(\kappa^* + \eta^2).$$

Applying Proposition 4.8 in  $B_{1/4}(y)$ , we obtain that for all  $t \leq 1/4$ ,

$$(5.11) \quad \bar{\sigma}(t; y) \leq C(\bar{\sigma}(1/4; y) + \kappa^* + \eta^2) \leq C(\kappa^* + \eta^2),$$

where  $C = C(n, m, \Lambda, \lambda, \alpha)$ . Observe that  $r \leq 1/32$  and  $B_{4r}(x_1) \subset B_{1/4}(x_1)$ . Then by Lemma 5.4 in  $B_{1/4}(x_1)$ , there exists a constant  $C = C(n, m, \Lambda, \lambda, \alpha)$  so that

$$\begin{aligned} |Du_j(x_1) - (Du_j)_{B_{4r}(x_1)}| &\leq C(\sqrt{\bar{\sigma}(4r; x_1)} + \kappa + \kappa_0 + \delta_0 + \eta)r; \\ |Du_j(x_2) - (Du_j)_{B_{4r}(x_1)}| &\leq C(\sqrt{\bar{\sigma}(4r; x_1)} + \kappa + \kappa_0 + \delta_0 + \eta)r. \end{aligned}$$

This together with (5.11) (by taking  $y = x_1$  therein) implies

$$\begin{aligned} |Du_j(x_1) - Du_j(x_2)| &\leq |Du_j(x_1) - (Du_j)_{B_{4r}(x_1)}| + |Du_j(x_2) - (Du_j)_{B_{4r}(x_1)}| \\ &\leq C(\sqrt{\bar{\sigma}(4r; x_1)} + \kappa + \kappa_0 + \delta_0 + \eta)r \\ &\leq C(\sqrt{\kappa^* + \eta^2}) \cdot |x_1 - x_2|, \end{aligned}$$

which yields

$$|Du_j(x_1) - Du_j(x_2)| \leq C(\sqrt{\kappa^*} + \eta) \cdot |x_1 - x_2|.$$

This completes the proof of Theorem 5.5.  $\square$

#### APPENDIX A. ESTIMATES OF SECOND ORDER PARTIAL DIFFERENTIAL EQUATIONS

We first consider the operators  $L$  of the form

$$Lu = D_i(a^{ij}(x)D_j u + b^i u) + c^i(x)D_i u + d(x)u,$$

where coefficients  $a^{ij}, b^i, c^i, d$  ( $i, j = 1, \dots, n$ ) are assumed to be measurable functions on a domain  $\Omega \subset \mathbb{R}^n$ . Let  $f^i, g, i = 1, \dots, n$  be locally integrable functions on  $\Omega$ . We shall assume that  $L$  is strictly elliptic in  $\Omega$ ; that is, there exists a positive number  $\lambda$  such that

$$(A.1) \quad \sum_{i,j} a^{ij} \xi_i \xi_j \geq \lambda |\xi|^2, \quad \forall x \in \Omega, \xi \in \mathbb{R}^n.$$

We also assume that the coefficients  $a^{ij}, b^i \in C^\alpha(\bar{\Omega})$ ,  $c^i, d, g \in L^\infty(\Omega)$ , and  $f^i \in C^\alpha(\bar{\Omega})$ . Suppose

$$(A.2) \quad \max_{1 \leq i, j \leq n} \left\{ |a^{ij}, b^i|_{C^\alpha(\bar{\Omega})}, |c^i, d|_{L^\infty(\bar{\Omega})} \right\} \leq K.$$

We present some estimates of the solution to the equation

$$(A.3) \quad Lu = D_i f^i + g, \quad \text{weakly in } B_r(0).$$

For the convenience, we denote  $\mathbf{f} = (f^1, \dots, f^n)$ .

**Theorem A.1** ([6, Theorem 8.32]). *Let  $u \in C^{1,\alpha}(\bar{B}_1(0))$  be a weak solution to (A.3) in  $B_r(0)$  for some  $r \in (0, 1]$ . Suppose that  $L$  satisfies the assumption (A.1) and (A.2) for  $\Omega = B_1(0)$ . Then we have*

$$|Du|_{C^0(B_{r/2}(0))} \leq C \left( r^{-1} |u|_{C^0(B_r(0))} + r |g|_{C^0(B_r(0))} + |f|_{C^0(B_r(0))} + r^\alpha \sup_{x,y \in B_r(0)} \frac{|\mathbf{f}(x) - \mathbf{f}(y)|}{|x - y|^\alpha} \right)$$

for  $C = C(n, \lambda, K)$ , where  $\lambda$  is given by (A.1), and  $K$  by (A.2).

*Proof.* Define

$$\tilde{a}^{ij}(x) := a^{ij}(rx), \quad \tilde{b}^i(x) = rb^i(rx); \quad \tilde{c}^i(x) = rc^i(rx), \quad \tilde{d}(x) = r^2d(rx).$$

Then by (A.1) and (A.2),

$$\sum_{i,j} \tilde{a}^{ij}(x) \xi_i \xi_j \geq \lambda |\xi|^2, \quad \forall x \in B_1(0), \xi \in \mathbb{R}^n;$$

$$\max_{1 \leq i, j \leq n} \left\{ |\tilde{a}^{ij}, \tilde{b}^i|_{C^\alpha(\overline{B}_1(0))}, |\tilde{c}^i, \tilde{d}|_{L^\infty(\overline{B}_1(0))} \right\} \leq Kr^\alpha \leq K.$$

Let

$$\tilde{u}(x) := r^{-1}\tilde{u}(rx); \quad \tilde{\mathbf{f}}^i(x) = (\tilde{f}^1, \dots, \tilde{f}^n) := \mathbf{f}^i(rx); \quad \tilde{g}(x) := rg(rx).$$

Then  $\tilde{u}$  satisfies the equation

$$D_i(\tilde{a}^{ij}(x)D_j\tilde{u} + \tilde{b}^i\tilde{u}) + \tilde{c}^i(x)D_i\tilde{u} + d(x)\tilde{u} = D_i\tilde{f}^i + \tilde{g}.$$

Applying [6, Theorem 8.32] to  $\tilde{u}$  in  $B_1(0)$ , we obtain that

$$|\tilde{u}|_{C^{1,\alpha}(B_{1/2}(0))} \leq C \left( |\tilde{u}|_{C^0(B_1(0))} + |\tilde{g}|_{C^0(B_1(0))} + |\tilde{\mathbf{f}}|_{C^0(B_1(0))} + \sup_{x,y \in B_1(0)} \frac{|\tilde{\mathbf{f}}(x) - \tilde{\mathbf{f}}(y)|}{|x - y|^\alpha} \right)$$

$$= C \left( r^{-1}|u|_{C^0(B_r(0))} + r|g|_{C^0(B_r(0))} + |\mathbf{f}|_{C^0(B_r(0))} + r^\alpha \sup_{x,y \in B_r(0)} \frac{|\mathbf{f}(x) - \mathbf{f}(y)|}{|x - y|^\alpha} \right).$$

for  $C = C(n, \lambda, K)$ . This finishes the proof of Theorem A.1.  $\square$

## REFERENCES

- [1] Assis Azevedo, José-Francisco Rodrigues, and Lisa Santos, *The  $N$ -membranes problem for quasilinear degenerate systems*, *Interfaces Free Bound.* **7** (2005), no. 3, 319–337. MR2171135
- [2] Luis Caffarelli, *The obstacle problem revisited*, *J. Fourier Anal. Appl.* **4** (1998), no. 4-5, 383–402. MR1658612
- [3] Luis Caffarelli, Daniela De Silva, and Ovidiu Savin, *The two membranes problem for different operators*, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **34** (2017), no. 4, 899–932. MR3661864
- [4] Luis Caffarelli, Luis Duque, and Hernán Vivas, *The two membranes problem for fully nonlinear operators*, *Discrete Contin. Dyn. Syst.* **38** (2018), no. 12, 6015–6027. MR3917798
- [5] Michel Chipot and Giorgio Vergara-Caffarelli, *The  $N$ -membranes problem*, *Appl. Math. Optim.* **13** (1985), no. 3, 231–249. MR806627
- [6] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, *Classics in Mathematics*, Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition. MR1814364
- [7] Qing Han and Fanghua Lin, *Elliptic partial differential equations*, Second, *Courant Lecture Notes in Mathematics*, vol. 1, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2011. MR2777537
- [8] Jürgen Jost, *Partial differential equations*, Third, *Graduate Texts in Mathematics*, vol. 214, Springer, New York, 2013. MR3012036
- [9] Fang-Hua Lin, *Regularity for a class of parametric obstacle problems (integrand, integral current, prescribed mean curvature, minimal surface system)*, ProQuest LLC, Ann Arbor, MI, 1985. Thesis (Ph.D.)—University of Minnesota. MR2634480
- [10] Lorenzo Sarnataro and Douglas Stryker, *Optimal regularity for minimizers of the prescribed mean curvature functional over isotopies* (2023), available at arXiv:math/2304.02722.
- [11] Ovidiu Savin and Hui Yu, *On the multiple membranes problem*, *J. Funct. Anal.* **277** (2019), no. 6, 1581–1602. MR3985514
- [12] Luis Silvestre, *The two membranes problem*, *Comm. Partial Differential Equations* **30** (2005), no. 1-3, 245–257. MR2131053

- [13] Leon Simon, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983. MR756417 (87a:49001)
- [14] Giorgio Vergara Caffarelli, *Regolarità di un problema di disequazioni variazionali relativo a due membrane*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat. (8) **50** (1971), 659–662. MR304835
- [15] ———, *Variational inequalities for two surfaces of constant mean curvature*, Arch. Rational Mech. Anal. **56** (1974/75), 334–347. MR355721
- [16] Zhichao Wang and Xin Zhou, *Existence of four minimal spheres in  $S^3$  with a bumpy metric* (2023), available at arXiv:math/2305.08755.
- [17] Brian White, *Curvature estimates and compactness theorems in 3-manifolds for surfaces that are stationary for parametric elliptic functionals*, Invent. Math. **88** (1987), no. 2, 243–256. MR880951
- [18] Xin Zhou and Jonathan J. Zhu, *Existence of hypersurfaces with prescribed mean curvature I—generic min-max*, Camb. J. Math. **8** (2020), no. 2, 311–362. MR4091027

SHANGHAI CENTER FOR MATHEMATICAL SCIENCE, 2005 SONGHU ROAD, FUDAN UNIVERSITY, SHANGHAI, 200438, CHINA

*Email address:* zhichao@fudan.edu.cn

DEPARTMENT OF MATHEMATICS, 531 MALOTT HALL, CORNELL UNIVERSITY, ITHACA, NY 14853, USA

*Email address:* xinzhou@cornell.edu