# SYMPLECTIC TABLEAUX AND QUANTUM SYMMETRIC PAIRS

HIDEYA WATANABE

ABSTRACT. We provide a new branching rule from the general linear group  $GL_{2n}(\mathbb{C})$  to the symplectic group  $Sp_{2n}(\mathbb{C})$  by establishing a simple algorithm which gives rise to a bijection from the set of semistandard tableaux of a fixed shape to a disjoint union of several copies of sets of symplectic tableaux of various shapes. The algorithm arises from representation theory of a quantum symmetric pair of type  $AII_{2n-1}$ , which is a q-analogue of the classical symmetric pair ( $\mathfrak{gl}_{2n}(\mathbb{C}), \mathfrak{sp}_{2n}(\mathbb{C})$ ).

## 1. INTRODUCTION

1.1. Branching rules. Let G be a group and  $\hat{G}$  a complete set of representatives of the equivalence classes of certain irreducible G-modules. Let H be a subgroup of G. It is a fundamental problem to determine how a given irreducible G-module  $V \in \hat{G}$  decomposes into irreducible H-submodules (if it does):

$$V \simeq \bigoplus_{W \in \hat{H}} W^{m_{V,W}}.$$

An explicit description of the multiplicities  $m_{V,W}$  is called a *branching rule*.

The problem of finding branching rules for certain pairs (G, H) of classical groups (the general/special linear groups  $GL_m(\mathbb{C})$ ,  $SL_m(\mathbb{C})$ , symplectic groups  $Sp_{2n}(\mathbb{C})$ , and (special) orthogonal groups  $O_m(\mathbb{C})$ ,  $SO_m(\mathbb{C})$ ) has been studied for a long time, and several (partial) answers have been obtained (see [HTW05] and references therein).

In the present paper, we focus on the irreducible polynomial representations for the pair  $(G, H) = (GL_{2n}(\mathbb{C}), Sp_{2n}(\mathbb{C}))$ . The equivalence classes of irreducible polynomial representations of  $GL_{2n}(\mathbb{C})$  (resp.,  $Sp_{2n}(\mathbb{C})$ ) are parametrized by the set  $\operatorname{Par}_{\leq 2n}$  (resp.,  $\operatorname{Par}_{\leq n}$ ) of partitions of length at most 2n (resp., n). For each  $\lambda \in \operatorname{Par}_{\leq 2n}$  and  $\nu \in \operatorname{Par}_{\leq n}$ , let  $m_{\lambda,\nu}$  denote the corresponding multiplicity.

Littlewood [Lit40] provided a partial branching rule. Namely, he determined the multiplicities  $m_{\lambda,\nu}$  for all  $\lambda, \nu \in \operatorname{Par}_{< n}$ , but not for all  $\lambda \in \operatorname{Par}_{< 2n}$ .

Sundaram [Sun90] gave a complete branching rule. The key ingredients for her theorem are King's symplectic tableaux ([Kin76]), Berele's insertion scheme for  $Sp_{2n}(\mathbb{C})$  ([Ber86]), and Sundaram's algorithm. In her branching rule, the multiplicities are determined by counting certain tableaux, which we call symplectic Littlewood-Richardson tableaux.

Naito and Sagaki [NaSa05] proposed a conjectural branching rule in terms of Littelmann paths. The conjecture was proved by Schumann and Torres [ScTo18].

1.2. **Results.** In the present paper, we introduce a simple algorithm which gives rise to a bijection

$$\mathrm{LR}^{\mathrm{AII}}:\mathrm{SST}_{2n}(\lambda)\to\bigsqcup_{\substack{\nu\in\mathrm{Par}_{\leq n}\\\nu\subset\lambda}}Sp\mathrm{T}_{2n}(\nu)\times\mathrm{Rec}_{2n}(\lambda/\nu)$$

Date: August 4, 2023.

<sup>2020</sup> Mathematics Subject Classification. Primary 05E10; Secondary 17B10, 17B37.

#### HIDEYA WATANABE

which sends a semistandard Young tableau of shape  $\lambda \in \operatorname{Par}_{\leq 2n}$  with entries in  $[1, 2n] := \{1, \ldots, 2n\}$  to a pair consisting of a symplectic tableau of some shape  $\nu \in \operatorname{Par}_{\leq n}$  with entries in [1, 2n] and a tableau, called a *recording tableau*, of skew shape  $\lambda/\nu$ . As a byproduct, it turns out that the multiplicity  $m_{\lambda,\nu}$  coincides with the number  $|\operatorname{Rec}_{2n}(\lambda/\nu)|$  of recording tableaux of shape  $\lambda/\nu$ . Therefore, our algorithm provides a new branching rule for  $(GL_{2n}(\mathbb{C}), Sp_{2n}(\mathbb{C}))$ . Moreover, it has deep representation theoretical information as we will see in the next subsection. We call the bijection the *Littlewood-Richardson map* since it can be regarded as a generalization of the branching rule, known as the Littlewood-Richardson rule, for the pair  $(GL_m(\mathbb{C}) \times GL_m(\mathbb{C}), GL_m(\mathbb{C}))$ .

Let us briefly explain our algorithm. Given  $T \in SST_{2n}(\lambda)$ , let  $\mathbf{a} = (a_1, \ldots, a_l)$  denote the first column of T (read from top to bottom), and S the other part. For the column  $\mathbf{a}$ , define a new column red( $\mathbf{a}$ ) to be the one obtained from  $\mathbf{a}$  by removing the entries in the set rem( $\mathbf{a}$ ), which is defined by the following recursive formula:

$$\operatorname{rem}(\mathbf{a}) := \begin{cases} \emptyset & \text{if } l \leq 1, \\ \operatorname{rem}(a_1, \dots, a_{l-2}) \sqcup \{a_{l-1}, a_l\} & \text{if } l \geq 2, \ a_l \in 2\mathbb{Z}, \ a_{l-1} = a_l - 1, \text{ and} \\ a_l < 2l - |\operatorname{rem}(a_1, \dots, a_{l-2})| - 1, \\ \operatorname{rem}(a_1, \dots, a_{l-1}) & \text{otherwise.} \end{cases}$$

Then, define a new tableau  $\operatorname{suc}(T)$  to be the product  $\operatorname{red}(\mathbf{a}) * S$  (in the plactic monoid). Set  $P^0 := T$ ,  $\nu^0 := \lambda$ , and  $Q^0$  to be the unique tableau of shape  $\lambda/\lambda$ . For each  $k \ge 0$ , set  $P^{k+1} := \operatorname{suc}(P^k)$ ,  $\nu^{k+1}$  to be the shape of  $P^{k+1}$ , and  $Q^{k+1}$  to be the tableau of shape

 $\lambda/\nu^{k+1}$  such that

$$Q^{k+1}(i,j) = \begin{cases} Q^k(i,j) & \text{if } (i,j) \notin D(\nu^k), \\ k+1 & \text{if } (i,j) \in D(\nu^k), \end{cases}$$

where  $D(\nu^k)$  denotes the Young diagram of the partition  $\nu^k$ . It turns out that this procedure eventually terminates; there exists a unique integer  $k_0 \ge 0$  such that

$$P^{k} = P^{k_{0}}, \ \nu^{k} = \nu^{k_{0}}, \ \text{and} \ Q^{k} = Q^{k_{0}} \ \text{ for all } k \ge k_{0}$$

Set

$$P^{AII}(T) := P^{k_0}, \quad Q^{AII}(T) := Q^{k_0}$$

Now, the output  $LR^{AII}(T)$  of the algorithm is the pair  $(P^{AII}(T), Q^{AII}(T))$ :

$$\mathrm{LR}^{\mathrm{AII}}(T) = (P^{\mathrm{AII}}(T), Q^{\mathrm{AII}}(T)).$$

**Example 1.2.1.** Let n = 3,  $\lambda = (4, 3, 2, 2, 1)$ , and consider the semistandard tableau T of shape  $\lambda$  given by

T =	1	1	<b>2</b>	4
	2	2	3	
	4	4		
	5	6		
	6			

Then, we have

$$\operatorname{rem}(1, 2, 4, 5, 6) = \{1, 2, 5, 6\},\$$

and

$$P^{1} = \underbrace{4}_{4} * \underbrace{124}_{23}_{4} = \underbrace{124}_{23}_{44}, \quad Q^{1} = \underbrace{1}_{1}_{1}$$

The next step is computed as follows: We have

$$\operatorname{rem}(1, 2, 4, 6) = \{1, 2\},\$$

and

$$P^{2} = \frac{4}{6} * \frac{2}{4} = \frac{2}{4} + \frac{4}{3}, \quad Q^{2} = \frac{1}{2}$$

Let us proceed to the next step: We have

$$\operatorname{rem}(2, 3, 4, 6) = \{3, 4\},\$$

and

$$P^{3} = \frac{2}{6} * \frac{4}{4} = \frac{2}{6}, \quad Q^{3} = \frac{1}{2}$$

Since  $rem(2,6) = \emptyset$ , the algorithm now terminates. Hence, we finally obtain

$$\mathrm{LR}^{A\mathrm{II}}(T) = \begin{pmatrix} 2 & 4 & 4 \\ \hline 6 & & 1 \\ \hline & & 3 & 2 \\ & & 3 & 1 \\ \hline & & 1 \end{pmatrix}.$$

1.3. Quantum symmetric pairs. Although the author believes that the bijectivity of the Littlewood-Richardson map can be proved in the realm of combinatorics, we prove it via representation theory of a quantum symmetric pair of type  $AII_{2n-1}$ , from which the algorithm to compute the Littlewood-Richardson map arises.

A quantum symmetric pair is a quantum analogue of a classical symmetric pair (e.g.,  $(\mathfrak{gl}_{2n}(\mathbb{C}), \mathfrak{sp}_{2n}(\mathbb{C}))$ , where  $\mathfrak{gl}_{2n}(\mathbb{C})$  (resp.,  $\mathfrak{sp}_{2n}(\mathbb{C})$ ) denotes the general linear Lie algebra (resp., the symplectic Lie algebra)). It consists of a Drinfeld-Jimbo quantum group U and a Letzter *i*quantum group U<sup>*i*</sup>. We refer the reader unfamiliar with quantum symmetric pair to a survey paper [Wan21]. In the present paper, we consider only a quantum symmetric pair of type  $AII_{2n-1}$ ; the U and U<sup>*i*</sup> are quantum analogues of the universal enveloping algebras of  $\mathfrak{gl}_{2n}(\mathbb{C})$  and  $\mathfrak{sp}_{2n}(\mathbb{C})$ , respectively.

For each  $\lambda \in \operatorname{Par}_{\leq 2n}$ , there exists a finite-dimensional irreducible U-module  $V(\lambda)$  with a distinguished basis of the form  $\{b_T \mid T \in \operatorname{SST}_{2n}(\lambda)\}$ , called the canonical basis ([Lus93]). Similarly, for each  $\nu \in \operatorname{Par}_{\leq n}$ , there exists a finite-dimensional irreducible U<sup>i</sup>-module  $V^i(\nu)$  ([Mol06], [Wat21]). In the present paper, we prove that the U<sup>i</sup>-module  $V^i(\nu)$  admits a distinguished basis of the form  $\{b_T^i \mid T \in \operatorname{SpT}_{2n}(\nu)\}$ .

Then, for each  $\lambda \in Par_{\leq 2n}$ , by composing several  $\mathbf{U}^i$ -module homomorphisms, we construct a  $\mathbf{U}^i$ -module isomorphism

$$\mathrm{LR}^{\mathrm{AII}}: V(\lambda) \to \bigoplus_{\nu \in \mathrm{Par}_{\leq n}} (V^{\imath}(\nu) \otimes \mathbb{Q}(q) \mathrm{Rec}_{2n}(\lambda/\nu)),$$

which we call the quantum Littlewood-Richardson map. Here, each  $x \in \mathbf{U}^i$  acts on each summand of the right-hand side as  $x \otimes id$ . The isomorphism is a q-analogue of the Littlewood-Richardson map on  $\mathrm{SST}_{2n}(\lambda)$  in the following sense: For each  $T \in \mathrm{SST}_{2n}(\lambda)$ , we have  $\mathrm{LR}^{\mathrm{AII}}(b_T) \equiv b^i_{P^{\mathrm{AII}}(T)} \otimes Q^{\mathrm{AII}}(T)$  modulo  $q^{-1}$  (in a suitable sense). This fact implies the bijectivity of the Littlewood-Richardson map.

#### HIDEYA WATANABE

Therefore, the Littlewood-Richardson map tells us not only the multiplicities  $m_{\lambda,\nu}$  but also how the irreducible U-module  $V(\lambda)$  decomposes into irreducible U<sup>*i*</sup>-submodules at  $q = \infty$ . Hence, this result must be closely related to the theory of crystal bases for quantum symmetric pairs (*cf.* [Wat21+]).

1.4. **Organization.** The paper is organized as follows. In Section 2, we collect terminology and basic results concerning partitions and tableaux which are necessary to formulate our main algorithm. Then, we state our main theorem in Section 3. It consists of the bijectivity of the Littlewood-Richardson map and an explicit description of the recording tableaux. The rest of the paper is devoted to proving the theorem. In Section 4, we factor the reduction map ( $\mathbf{a} \mapsto \operatorname{red}(\mathbf{a})$ ) into small pieces so that we can prove its injectivity. After reviewing representation theory of  $\mathfrak{gl}_{2n}(\mathbb{C})$ ,  $\mathfrak{sp}_{2n}(\mathbb{C})$ ,  $\mathbf{U}$ , and  $\mathbf{U}^i$  in Sections 5 and 6, we prove the surjectivity of the Littlewood-Richardson map via an investigation into the quantum Littlewood-Richardson map in Section 7. We finally complete the proof of our main theorem in Section 8 by relating the recording tableaux to the symplectic Littlewood-Richardson tableaux.

Acknowledgements. This work was supported by JSPS KAKENHI Grant Number JP22KJ2603.

Notation. Throughout the paper, we fix positive integers m and n.

Given two integers a and b, let [a, b] denote the integer interval:

$$[a,b] := \{c \in \mathbb{Z} \mid a \le c \le b\}.$$

# 2. Preliminaries from combinatorics

In this section, we collect terminology and basic results concerning partitions and tableaux which are necessary to formulate our main algorithm in the next section.

## 2.1. Partitions. A *partition* is a non-increasing sequence

$$\lambda = (\lambda_1, \ldots, \lambda_l)$$

of positive integers. We often regard a non-increasing sequence of nonnegative integers as a partition by ignoring the zero's. Each  $\lambda_i$  is referred to as a *part* of  $\lambda$ . It is convenient to set  $\lambda_i := 0$  for i > l. The sum  $\sum_{i=1}^{l} \lambda_i$  of parts is called the *size* of  $\lambda$ , and is denoted by  $|\lambda|$ . The number l of parts of  $\lambda$  is called the *length* of  $\lambda$ , and is denoted by  $\ell(\lambda)$ . We regard the empty sequence () as a unique partition of length 0. Let Par denote the set of all partitions.

For each  $l \in \mathbb{Z}_{\geq 0}$ , let  $\operatorname{Par}_{\leq l}$  denote the set of all partitions of length at most l.

For each  $l \in \mathbb{Z}_{\geq 0}$ , let  $\varpi_l$  denote the partition of length l whose parts are all 1:

(2.1) 
$$\varpi_l = (1^l) = (\underbrace{1, \dots, 1}_l)$$

Until the end of this subsection, let us fix a partition  $\lambda$ . The Young diagram of shape  $\lambda$  is the set

$$D(\lambda) := \{ (i, j) \mid 1 \le i \le \ell(\lambda) \text{ and } 1 \le j \le \lambda_i \}.$$

As usual, we visualize it by a collection of boxes; e.g.,

$$D(4,3,2,2,1) =$$

Given a partition  $\lambda$  and a number  $j \in [1, \lambda_1]$ , set

(2.2) 
$$\operatorname{col}_{j}(\lambda) := \sharp \{ i \in [1, \ell(\lambda)] \mid \lambda_{i} \geq j \}.$$

For example, if  $\lambda = (4, 3, 2, 2, 1)$ , then

$$(\operatorname{col}_1(\lambda), \operatorname{col}_2(\lambda), \operatorname{col}_3(\lambda), \operatorname{col}_4(\lambda)) = (5, 4, 2, 1).$$

**Definition 2.1.1.** We say that the partition  $\lambda$  has even columns if  $col_i(\lambda)$  is even for all  $j \in [1, \lambda_1].$ 

Given a partition  $\mu \in \text{Par}$ , we write  $\mu \subseteq \lambda$  to indicate that  $D(\mu) \subseteq D(\lambda)$ , or equivalently,

$$\mu_i \leq \lambda_i \text{ for all } 1 \leq i \leq \ell(\lambda).$$

For each  $\mu \in \text{Par}$  with  $\mu \subseteq \lambda$ , set

$$|\lambda/\mu| := |\lambda| - |\mu|$$
 and  $D(\lambda/\mu) := D(\lambda) \setminus D(\mu)$ .

**Definition 2.1.2.** Let  $\mu \in \text{Par such that } \mu \subseteq \lambda$ . We say that  $\lambda/\mu$  is a *vertical strip*, and write  $\mu \subseteq_{\operatorname{vert}} \lambda$  if

$$\mu_i \geq \lambda_i - 1$$
 for all  $i \in [1, \ell(\lambda)]$ 



The following is immediate from the definition of vertical strips.

**Lemma 2.1.3.** Let  $\lambda'$  denote the partition  $(\lambda_1 - 1, \ldots, \lambda_{\ell(\lambda)} - 1)$ , and  $\mu$  be a partition such that  $\lambda' \subseteq_{vert} \mu$ . Then, we have  $\mu \subseteq_{vert} \lambda$  if and only if  $\ell(\mu) \leq \ell(\lambda)$ .

2.2. Semistandard tableaux. In this subsection, we fix a partition  $\lambda$ .

A Young tableau, or simply a tableau, of shape  $\lambda$  is a map

$$T: D(\lambda) \to \mathbb{Z}_{>0}$$

The partition  $\lambda$  is called the *shape* of T, and denoted by sh(T). As usual, we visualize a tableau by filling the boxes of  $D(\lambda)$  with positive integers; e.g., the following is a tableau of shape (4, 3, 2, 2, 1).

(2.3)



Let T be a tableau of shape  $\lambda$ . Given a positive integer k, let T[k] denote the number of occurrences of the entry k in T:

(2.4) 
$$T[k] := \sharp\{(i,j) \in D(\lambda) \mid T(i,j) = k\}.$$

For each  $j \in [1, \lambda_1]$ , set

$$w_j^{\text{col}}(T) := (T(\text{col}_j(\lambda), j), \dots, T(2, j), T(1, j))$$

(see (2.2) for the definition of  $\operatorname{col}_j$ ). It is the sequence of entries in the *j*-th column of T read from bottom to top. The *column word* of T is the sequence  $w^{\operatorname{col}}(T)$  of entries obtained by concatenating the  $w_j^{\operatorname{col}}(T)$ 's:

(2.5) 
$$w^{\operatorname{col}}(T) := w_1^{\operatorname{col}}(T) \circ \cdots \circ w_{\lambda_1}^{\operatorname{col}}(T).$$

For example, if T is the tableau in (2.3), then

$$w^{\text{col}}(T) = (6, 5, 4, 2, 1, 6, 4, 2, 1, 3, 2, 4).$$

The tableau T is said to be *semistandard* if the entries increase weakly from left to right along the rows, and strictly from top to bottom along the columns. Namely,

 $T(i,j) \leq T(i,j+1) \text{ and } T(i,j) < T(i+1,j) \text{ ~for all } (i,j) \in D(\lambda),$ 

where we set  $T(i', j') := \infty$  if  $(i', j') \notin D(\lambda)$ . For example, the tableau in (2.3) is semistandard. Let  $SST_m(\lambda)$  denote the set of all semistandard tableaux of shape  $\lambda$  with entries in [1, m].

The generating function

(2.6) 
$$s_{\lambda}(x_1, \dots, x_m) := \sum_{T \in SST_m(\lambda)} \mathbf{x}^{\operatorname{wt}(T)} \in \mathbb{Z}[x_1, \dots, x_m]$$

is called the *Schur function*, where

(2.7)  $\operatorname{wt}(T) := (T[1], \dots, T[m]), \quad \mathbf{x}^{(a_1, \dots, a_m)} := x_1^{a_1} \cdots x_m^{a_m}.$ 

The Schur functions are symmetric polynomials:

$$s_{\lambda}(x_{\sigma(1)},\ldots,x_{\sigma(m)})=s_{\lambda}(x_1,\ldots,x_m)$$

for all permutation  $\sigma$  on [1, m].

For  $\mu \in \text{Par with } \mu \subseteq \lambda$ , a *tableau* of shape  $\lambda/\mu$  is a map

$$D(\lambda/\mu) \to \mathbb{Z}_{>0}.$$

Let  $\operatorname{Tab}(\lambda/\mu)$  denote the set of all tableaux of shape  $\lambda/\mu$ . The notion of semistandard tableaux of shape  $\lambda/\mu$  is defined in the obvious way. Let  $\operatorname{SST}_m(\lambda/\mu)$  denote the set of all semistandard tableaux of shape  $\lambda/\mu$  with entries in [1, m]. For example, the following is a semistandard tableau of shape (4, 3, 2, 2, 1)/(2, 2, 1):



2.3. Plactic monoid. The set of all semistandard tableaux with entries in [1, m] forms a monoid, called the *plactic monoid* (*cf.* [Ful97, Sections 1.1 and A.2]). For the reader's convenience, we recall here its definition. In order to describe the monoid structure, we need to introduce the *column insertion algorithm*, which receives a pair (w, T) of a positive integer w and a semistandard tableau T as an input, and returns a new semistandard tableau  $w \to T$  as an output as follows. Set  $\lambda := \operatorname{sh}(T)$ , and

$$w_0 := w, \quad T_0 := T.$$

For each j > 0, given a pair  $(w_{j-1}, T_{j-1})$ , set

(2.8) 
$$r_j := \min\{r \in [1, \operatorname{col}_j(\lambda) + 1] \mid T(r, j) \ge w_{j-1}\} \\ w_j := T(r_j, j),$$

where we set  $T(i', j') = \infty$  if  $(i', j') \notin D(\lambda)$  (see (2.2) for the definition of  $\operatorname{col}_j$ ). Also set  $\lambda^j$  to be the partition such that

$$D(\lambda^j) = \begin{cases} D(\lambda) & \text{if } r_j \le \operatorname{col}_j(\lambda), \\ D(\lambda) \sqcup \{(r_j, j)\} & \text{if } r_j = \operatorname{col}_j(\lambda) + 1, \end{cases}$$

and  $T_j$  to be the semistandard tableau of shape  $\lambda^j$  such that

$$T_{j}(i',j') = \begin{cases} w_{j-1} & \text{if } (i',j') = (r_{j},j), \\ T_{j-1}(i',j') & \text{if } (i',j') \neq (r_{j},j), \end{cases} \text{ for all } (i',j') \in D(\lambda^{j}).$$

Let  $s \ge 1$  denote the minimal integer such that  $r_j = \operatorname{col}_j(\lambda) + 1$ . Then, the semistandard tableau  $T_s$  is the one  $w \to T$ .

The sequence

$$\operatorname{br}(w,T) := (r_1,\ldots,r_s)$$

is called the *bumping route*.

**Example 2.3.1.** Let w = 7 and

$$T = \underbrace{\begin{array}{c}1 & 2 & 2 & 3\\3 & 4 & 5\\4 & 5 & 6\\6 & 6 & 9\\7 & 7 & 10\\8 & 8\\10\end{array}}_{W \to T}$$
$$w \to T = \underbrace{\begin{array}{c}1 & 2 & 2 & 3\\3 & 4 & 5 & 9\\3 & 4 & 5 & 6\\4 & 5 & 6\end{array}}_{M \times T}, \quad \operatorname{br}(w, T) = (5, 5, 4, 2).$$

Then, we have

The following proposition can be straightforwardly deduced from the definitions.

**Proposition 2.3.2.** Let  $w, T, r_1, \ldots, r_s$  be as above. Set  $S := w \to T$  and  $\mu := \operatorname{sh}(S)$ . Then, the following hold.

- (1)  $r_1 \geq \cdots \geq r_s$
- (2)  $w \le T(r_1, 1) \le T(r_2, 2) \le \dots \le T(r_{s-1}, s-1) < T(r_s, s) = \infty.$

(3)  $\lambda \subset \mu$ . Moreover,  $D(\mu/\lambda) = \{(r_s, s)\}.$ 

(4) For each  $(i, j) \in D(\mu)$ , we have

$$S(i,j) = \begin{cases} w & \text{if } (i,j) = (r_1,1), \\ T(r_{j-1},j-1) & \text{if } j \in [2,s] \text{ and } i = r_j, \\ T(i,j) & \text{otherwise.} \end{cases}$$

Given two semistandard tableaux S and T, their product S \* T in the plactic monoid is given by

$$S * T := w_1 \to (\dots \to (w_r \to T) \dots)$$

where  $(w_1, \ldots, w_r) = w^{\text{col}}(S)$  (see (2.5) for the definition of  $w^{\text{col}}$ ).

Regarding bumping routes of successive insertions, the following is known.

**Proposition 2.3.3** ([Ful97, Exercise 3 in Section A.2]). Let  $\lambda \in \operatorname{Par}_{\leq m}$ ,  $T \in \operatorname{SST}_m(\lambda)$ , and  $w, w' \in [1, m]$  be such that w < w'. Let us write

$$\operatorname{br}(w,T) = (r_1, \dots, r_s), \quad \operatorname{br}(w', (w \to T)) = (r'_1, \dots, r'_{s'}).$$

Then, we have  $s' \leq s$  and  $r'_j > r_j$  for all  $j \in [1, s']$ .

The following is known as (a combinatorial version of) Pieri's formula.

**Proposition 2.3.4.** Let  $\lambda \in \operatorname{Par}_{\leq m}$  and  $k \in [0, m]$ . The assignment  $(S, T) \to S * T$  gives rise to a bijection

$$*: \mathrm{SST}_m(\varpi_k) \times \mathrm{SST}_m(\lambda) \to \bigsqcup_{\substack{\mu \in \mathrm{Par}_{\leq m} \\ \lambda \subseteq \mu \text{ and } |\mu/\lambda| = k}} \mathrm{SST}_m(\mu).$$

2.4. Symplectic tableaux. In this subsection, we fix a partition  $\lambda \in Par_{\leq 2n}$ .

**Definition 2.4.1** ([Kin76, Section 4]). A semistandard tableau  $T \in SST_{2n}(\lambda)$  is said to be *symplectic* if

$$T(k,1) \ge 2k-1$$
 for all  $k \in [1,\ell(\lambda)]$ .

Let  $SpT_{2n}(\lambda)$  denote the set of all symplectic tableaux of shape  $\lambda$ .

**Proposition 2.4.2.** If  $SpT_{2n}(\lambda) \neq \emptyset$ , then  $\ell(\lambda) \leq n$ .

*Proof.* Assume contrary that  $\ell(\lambda) > n$ . Let  $T \in SpT_{2n}(\lambda)$ . Then, we have

$$T(n+1,1) \ge 2(n+1) - 1 = 2n + 1 > 2n.$$

This contradicts that the entries of T are in [1, 2n]. Thus, the assertion follows.

For each  $\nu \in \operatorname{Par}_{\leq n}$ , the generating function

(2.9) 
$$s_{\nu}^{Sp}(y_1, \dots, y_n) := \sum_{T \in \operatorname{SpT}_{2n}(\nu)} \mathbf{y}^{\operatorname{wt}^{Sp}(T)} \in \mathbb{Z}[y_1^{\pm 1}, \dots, y_n^{\pm 1}]$$

is called the symplectic Schur function, where

(2.10) 
$$\operatorname{wt}^{Sp}(T) := (T[1] - T[2], T[3] - T[4], \dots, T[2n-1] - T[2n]).$$

The symplectic Schur functions are linearly independent.

**Lemma 2.4.3.** Let  $T \in SST_{2n}(\lambda)$ . If T is not symplectic, then there exists a unique  $i \in [2, 2n]$  such that

$$T(i, 1) < 2i - 1$$
 and  $T(k, 1) \ge 2k - 1$  for all  $k \in [1, i - 1]$ .

Moreover, we have

$$T(i, 1) = 2i - 2$$
 and  $T(i - 1, 1) = 2i - 3$ .

Proof. Since T is not symplectic, there exists  $i \in [1, \ell(\lambda)]$  such that T(i, 1) < 2i - 1 (note that the number *i* cannot be 1 since T(1, 1) is always greater than or equal to  $1 (= 2 \cdot 1 - 1)$ ). We may take the minimal *i* among such integers. Then, the first assertion is clear.

Now, we have

$$2i - 3 = 2(i - 1) - 1 \le T(i - 1, 1) < T(i, 1) < 2i - 1.$$

This implies the second assertion.

2.5. Reduction map. For each  $l \in [0, m]$ , a tableau in  $SST_m(\varpi_l)$  (see (2.1) for the definition of  $\varpi_l$ ) can be represented by an increasing sequence  $(a_1, \ldots, a_l)$  of integers in [1, m]. We often regard such sequences **a**, **b**, etc. as sets, and consider their cardinalities  $|\mathbf{a}|$ , disjoint unions  $\mathbf{a} \sqcup \mathbf{b}$ , set differences  $\mathbf{a} \setminus \mathbf{b}$ , and so on.

In this subsection, we fix  $l \in [0, 2n]$  and  $\mathbf{a} = (a_1, \ldots, a_l) \in SST_{2n}(\varpi_l)$ .

**Definition 2.5.1.** The set of *removable entries* of  $\mathbf{a}$  is the subset rem( $\mathbf{a}$ ) of  $\mathbf{a}$  defined as follows.

(1) If  $l \le 1$ , then rem $(\mathbf{a}) = \emptyset$ . (2) If l > 1, then  $\operatorname{rem}(\mathbf{a}) = \begin{cases} \operatorname{rem}(a_1, \dots, a_{l-2}) \sqcup \{a_{l-1}, a_l\} & \text{if } a_l \in 2\mathbb{Z}, \ a_{l-1} = a_l - 1, \text{ and} \\ a_l < 2l - |\operatorname{rem}(a_1, \dots, a_{l-2})| - 1, \\ \operatorname{rem}(a_1, \dots, a_{l-1}) & \text{otherwise.} \end{cases}$ 

**Definition 2.5.2.** The reduction map on  $SST_{2n}(\varpi_l)$  is the map

red : 
$$\operatorname{SST}_{2n}(\varpi_l) \to \bigsqcup_{k=0}^{2n} \operatorname{SST}_{2n}(\varpi_k); \ \mathbf{a} \mapsto \mathbf{a} \setminus \operatorname{rem}(\mathbf{a}).$$

For example, if we take  $\mathbf{a} = (1, 3, 4, 5, 6, 7, 11, 12, 13, 14)$ , then we have

 $\operatorname{rem}(\mathbf{a}) = \{3, 4, 5, 6, 13, 14\}, \quad \operatorname{red}(\mathbf{a}) = (1, 7, 11, 12).$ 

The following is immediate from the definition.

# Proposition 2.5.3.

- (1) If  $a_l \in \text{rem}(\mathbf{a})$ , then  $a_l \in 2\mathbb{Z}$ .
- (2) If  $a_l \notin \operatorname{rem}(\mathbf{a})$ , then  $\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \ldots, a_{l-1})$ .
- (3) If  $a_l \in \text{rem}(\mathbf{a})$ , then  $\text{rem}(a_1, \ldots, a_{l-1}) = \text{rem}(a_1, \ldots, a_{l-2})$ .
- (4)  $\operatorname{rem}(a_1,\ldots,a_k) \subseteq \operatorname{rem}(\mathbf{a})$  for all  $k \in [0,l]$ .

2.6. Successor map. In this subsection, we fix  $\lambda \in \operatorname{Par}_{\leq 2n}$ , and set  $l := \ell(\lambda)$ . Let  $\lambda'$  denote the partition  $(\lambda_1 - 1, \ldots, \lambda_l - 1)$ .

Let us define a map

(2.11) 
$$d: \mathrm{SST}_{2n}(\lambda) \to \mathrm{SST}_{2n}(\varpi_l) \times \mathrm{SST}_{2n}(\lambda')$$

as follows. For each  $T \in SST_{2n}(\lambda)$ , the image d(T) is the pair  $(\mathbf{a}, T')$  consisting of the first column  $\mathbf{a}$  of T:

 $\mathbf{a} = (T(1,1), T(2,1), \dots, T(l,1)),$ 

and the other part T':

$$T'(i,j) = T(i,j+1)$$
 for all  $(i,j) \in D(\lambda')$ .

**Definition 2.6.1.** The successor map is the composite

suc := \* 
$$\circ$$
 (red, id)  $\circ d$  : SST<sub>2n</sub>( $\lambda$ )  $\rightarrow \bigsqcup_{\mu \in \operatorname{Par}_{\leq 2n}} \operatorname{SST}_{2n}(\mu)$ ,

where \* denotes the multiplication map of the plactic monoid.

## Example 2.6.2. Let



Then, we have



**Lemma 2.6.3.** Let  $T \in SST_{2n}(\lambda)$ . Set  $d(T) = (\mathbf{a}, T')$ , and write  $\mathbf{a} = (a_1, \ldots, a_l)$  and red $(\mathbf{a}) = \mathbf{b} = (b_1, \ldots, b_k)$ . Set  $S^0 := T'$ ,  $S^t := b_t \to S^{t-1}$ , and  $\mu^t := \operatorname{sh}(S^t)$ . Let us write  $\operatorname{br}(b_t, S^{t-1}) = (r_{t,1}, \ldots, r_{t,s_t})$ . Let  $r_{t,0}$  be such that  $a_{r_{t,0}} = b_t$ . Then, the following hold for all  $t \in [1, k]$ :

- (1)  $D(\mu^t / \lambda') = \{(r_{u,s_u}, s_u) \mid u \in [1, t]\}.$ (2)  $S^t(i, j) = \begin{cases} T(r_{u,j-1}, j) & \text{if } u \in [1, t], \ j \in [1, s_u], \ and \ i = r_{u,j}, \\ T(i, j+1) & \text{otherwise.} \end{cases}$

- (3)  $s_1 \ge s_2 \ge \cdots \ge s_k$ . (4)  $r_{t,j} > r_{t-1,j} > \cdots > r_{1,j}$  for all  $j \in [0, s_t]$ . (5)  $r_{t,j} = \min\{r \in [r_{t-1,j}+1, \operatorname{col}_j(\mu^{t-1})+1] \mid T(r, j+1) \ge T(r_{t,j-1}, j)\}$  for all  $j \in [1, s_t]$ , where we set  $r_{0,j} = 0$ . (6)  $r_{t,0} \ge r_{t,1} \ge \cdots \ge r_{t,s_t}$ .

*Proof.* The first and second assertions can be deduced from Proposition 2.3.2 (3) and (4)by induction on t. The third and fourth assertions follow from Proposition 2.3.3.

Next, let us prove the fifth assertion. For each  $j \in [0, s_t]$ , set

$$w_{t,j} := \begin{cases} b_t & \text{if } j = 0, \\ S^{t-1}(r_{t,j}, j) & \text{if } j > 0. \end{cases}$$

Then, by equation (2.8), we have

(2.12) 
$$r_{t,j} = \min\{r \in [1, \operatorname{col}_j(\mu^{t-1}) + 1] \mid S^{t-1}(r, j) \ge w_{t,j-1}\}.$$

This, together with the second and fourth assertions, implies our claim.

Finally, let us prove the last assertion by induction on t; we understand that  $s_0 = 0$  so that the assertion for t = 0 is clear. We have

$$r_{t,1} \geq \cdots \geq r_{t,s_t}$$

by Proposition 2.3.2 (1). Hence, we only need to show that  $r_{t,0} \ge r_{t,1}$ . By equation (2.12), we have

$$r_{t,1} \le \operatorname{col}_1(\mu^{t-1}) + 1$$

Hence, there is nothing to prove if  $r_{t,0} > \operatorname{col}_1(\mu^{t-1})$ . Therefore, suppose that  $r_{t,0} \leq \operatorname{col}_1(\mu^{t-1})$ . Then, we have

$$T(r_{t,0}, 1) \le T(r_{t,0}, 2).$$

Since  $a_{r_{t,0}} = b_t > b_{t-1} = a_{r_{t-1,0}}$ , we see that  $r_{t,0} > r_{t-1,0}$ . By our induction hypothesis, it holds that  $r_{t-1,0} \ge r_{t-1,1}$ . Summarizing above, we have

$$r_{t-1,1} < r_{t,0} \le \operatorname{col}_1(\mu^{t-1}) \text{ and } T(r_{t,0},2) \ge T(r_{t,0},1)$$

Then, the fifth assertion implies that  $r_{t,1} \leq r_{t,0}$ , as desired.

**Lemma 2.6.4.** Let  $T \in SST_{2n}(\lambda)$  and set  $\mu := \operatorname{sh}(\operatorname{suc}(T))$ . Then, we have  $\mu \subseteq_{vert} \lambda$ . Moreover, the equality holds if and only if  $\operatorname{suc}(T) = T$ .

*Proof.* We use the notation in Lemma 2.6.3. Note that  $S^k = suc(T)$  and  $\mu^k = \mu$ . By Proposition 2.3.4, we see that

$$\lambda' \subseteq_{\operatorname{vert}} \mu \text{ and } |\mu/\lambda'| = |\operatorname{red}(\mathbf{a})|.$$

Therefore, in order to prove that  $\mu \subseteq_{\text{vert}} \lambda$ , we only need to show that  $\ell(\mu) \leq l$  (see Lemma 2.1.3).

For each  $t \in [0, k]$ , by Lemma 2.6.3 (1) we have

(2.13) 
$$\ell(\mu) = \ell(\lambda') + \sharp\{t \in [1,k] \mid s_t = 1\}$$

Let  $u \in [1, k]$  be such that  $r_{u,0} \leq \ell(\lambda')$ . Then, we have

$$r_{u,1} \le r_{u,0} \le \ell(\lambda') \le \ell(\mu^{u-1}) = \operatorname{col}_1(\mu^{u-1}),$$

where the first inequality follows from Lemma 2.6.3 (6). This implies that  $s_u > 1$ . Hence, we see that

 $\sharp\{t \in [1,k] \mid s_u = 1\} \le k - \ell(\lambda').$ 

This, together with equation (2.13), implies the first assertion.

Next, let us prove the second assertion. Suppose that  $\mu = \lambda$ . In this case, we must have  $red(\mathbf{a}) = \mathbf{a}$  since

$$0 = |\lambda/\mu| = |\lambda/\lambda'| - |\mu/\lambda'| = l - |\operatorname{red}(\mathbf{a})| = |\operatorname{rem}(\mathbf{a})|.$$

On the other hand, it is clear that

$$\mathbf{a} * T' = T.$$

Therefore, we obtain suc(T) = T, as desired. The opposite direction is trivial.

# 3. Main result

3.1. Statement. Let  $\lambda \in \operatorname{Par}_{\leq 2n}$  and  $T \in \operatorname{SST}_{2n}(\lambda)$ . For each  $k \geq 0$ , define a semistandard tableau  $P^k$ , a partition  $\nu^{\overline{k}}$ , and a tableau  $Q^k$  inductively as follows. First, set

$$P^0 := T, \quad \nu^0 := \lambda,$$

and  $Q^0$  to be the unique tableau of shape  $\lambda/\lambda$ . For  $k \ge 0$ , set

$$P^{k+1} := \operatorname{suc}(P^k), \quad \nu^{k+1} := \operatorname{sh}(P^{k+1}).$$

and  $Q^{k+1}$  to be the tableau of shape  $\lambda/\nu^{k+1}$  given by

$$Q^{k+1}(i,j) := \begin{cases} Q^k(i,j) & \text{ if } (i,j) \notin D(\nu^k), \\ k+1 & \text{ if } (i,j) \in D(\nu^k). \end{cases}$$

Note that by Lemma 2.6.4, there exists a unique  $k_0 \ge 0$  such that

$$\lambda = \nu^0 \underset{\text{vert}}{\supset} \nu^1 \underset{\text{vert}}{\supset} \nu^2 \underset{\text{vert}}{\supset} \cdots \underset{\text{vert}}{\supset} \nu^{k_0}$$

and

$$P^{k} = P^{k_{0}}, \ \nu^{k} = \nu^{k_{0}}, \ Q^{k} = Q^{k_{0}} \text{ for all } k \ge k_{0}$$

**Definition 3.1.1.** Let  $T, k_0$  be as above. Define two tableaux  $P^{AII}(T)$  and  $Q^{AII}(T)$  to be  $P^{k_0}$  and  $Q^{k_0}$ , respectively.

See Example 1.2.1 for an example of this algorithm. The following is immediate from the definition.

**Lemma 3.1.2.** Let  $\lambda, T, \nu^k, k_0$  be as above. Set  $\nu := \nu^{k_0}$ .

(1) For each  $(i, j) \in D(\lambda/\nu)$ , we have

$$Q^{\text{AII}}(T)(i,j) = \min\{k \in [1,k_0] \mid (i,j) \notin D(\nu^k)\}.$$

(2) For each  $k \ge 0$ , we have

$$D(\nu^{k}) = D(\nu) \sqcup \{(i, j) \in D(\lambda/\nu) \mid Q^{AII}(T)(i, j) > k\}.$$

**Definition 3.1.3.** Let  $\lambda \in \operatorname{Par}_{\leq 2n}$  and  $\nu \in \operatorname{Par}_{\leq n}$  be such that  $\nu \subseteq \lambda$ . A tableau Q of shape  $\lambda/\nu$  is said to be a *recording tableau* if there exists  $T \in \operatorname{SST}_{2n}(\lambda)$  such that  $Q^{\operatorname{AII}}(T) = Q$ . Let  $\operatorname{Rec}_{2n}(\lambda/\nu)$  denote the set of recording tableaux of shape  $\lambda/\nu$ .

Let  $\lambda \in \operatorname{Par}_{\leq 2n}$  and  $\nu \in \operatorname{Par}_{\leq n}$  be such that  $\nu \subseteq \lambda$ . Let  $\operatorname{Rec}_{2n}(\lambda/\nu)$  denote the set of tableaux Q of shape  $\lambda/\nu$  satisfying the following:

(R1) The entries of Q strictly decrease along the rows from left to right.

- (R2) The entries of Q weakly decrease along the columns from top to bottom.
- (R3) For each k > 0, the number Q[k] (see (2.4) for the definition) is even.
- (R4) For each k > 0, it holds that

$$Q[k] \ge 2(\ell(\nu^{k-1}) - n),$$

where  $\nu^{k-1}$  is the partition such that

$$D(\nu^{k-1}) = D(\nu) \sqcup \{(i,j) \in D(\lambda/\nu) \mid Q(i,j) \ge k\}.$$

(R5) For each r, k > 0, let  $Q_{\leq r}[k]$  denote the number of occurrences of k in Q in the r-th row or above. Then, the following inequality holds:

$$Q_{\leq r}[k+1] \leq Q_{\leq r}[k].$$

Now, we are ready to state the main result in this paper.

Theorem 3.1.4. Let  $\lambda \in \operatorname{Par}_{<2n}$ .

(1) The assignment  $T \mapsto (P^{AII}(T), Q^{AII}(T))$  gives rise to a bijection

$$\mathrm{LR}^{\mathrm{AII}} : \mathrm{SST}_{2n}(\lambda) \to \bigsqcup_{\substack{\nu \in \mathrm{Par}_{\leq n} \\ \nu \subseteq \lambda}} (Sp\mathrm{T}_{2n}(\nu) \times \mathrm{Rec}_{2n}(\lambda/\nu)).$$

(2) For each  $\nu \in \operatorname{Par}_{\leq n}$  such that  $\nu \subseteq \lambda$ , we have

$$\operatorname{Rec}_{2n}(\lambda/\nu) = \operatorname{Rec}_{2n}(\lambda/\nu).$$

**Definition 3.1.5.** We call the map  $LR^{AII}$  in Theorem 3.1.4 the *Littlewood-Richardson* map.

The rest of this paper is devoted to proving the theorem. Since the proof is involved, we give an outline in the next subsection.

3.2. Outline of the proof. First, we reformulate the reduction map as a composite of certain maps (Corollary 4.4.2 (1)). As a result, we see that the reduction map on  $SST_{2n}(\varpi_l)$  is injective (Corollary 4.4.2 (2)).

Next, by studying some properties of the reduction map, we deduce that the tableau  $P^{AII}(T)$  is symplectic (Proposition 7.1.1). Also, we prove the injectivity of the Littlewood-Richardson map by using the successor map (Proposition 7.1.3).

To prove the surjectivity of the Littlewood-Richardson map, we use representation theory of a quantum symmetric pair  $(\mathbf{U}, \mathbf{U}^i)$  of type  $AII_{2n-1}$ . It is known that for each  $\lambda \in Par_{\leq 2n}$ , there exists an irreducible **U**-module  $V(\lambda)$  with basis of the form  $\{b_T \mid T \in$  $SST_{2n}(\lambda)\}$ . Similarly, for each  $\nu \in Par_{\leq n}$ , there exists an irreducible  $\mathbf{U}^i$ -module  $V^i(\nu)$ . We show that it admits a distinguished basis of the form  $\{b_T^i \mid T \in SpT_{2n}(\nu)\}$  (Proposition 6.4.1). Then, we lift the Littlewood-Richardson map to a  $\mathbf{U}^i$ -module homomorphism

$$V(\lambda) \to \bigoplus_{\substack{\nu \in \operatorname{Par}_{\leq n} \\ \nu \subseteq \lambda}} (V^{\iota}(\nu) \otimes \mathbb{Q}(q) \operatorname{Rec}_{2n}(\lambda/\nu))$$

which maps  $b_T$  to  $b_{P^{AII}(T)} \otimes Q^{AII}(T)$  modulo  $q^{-1}$  for all  $T \in SST_{2n}(\lambda)$  (Proposition 6.5.2). Here, each  $x \in \mathbf{U}^i$  acts on each summand of the right-hand side as  $x \otimes id$ . The complete reducibility of the right-hand side implies that this homomorphism is surjective (Theorem 7.2.1). Hence, we conclude that the Littlewood-Richardson map is surjective (Corollary 7.2.2).

Finally, by studying the successor map in detail, we verify that  $\operatorname{Rec}_{2n}(\lambda/\nu) \subseteq \operatorname{Rec}_{2n}(\lambda/\nu)$ (Lemma 8.3.1). On the other hand, we show that there is an injective map from  $\operatorname{Rec}_{2n}(\lambda/\nu)$ to the set  $\operatorname{LRT}_{2n}^{Sp}(\lambda/\nu)$  of symplectic Littlewood-Richardson tableaux (Lemma 8.3.2). Then, we conclude that  $\operatorname{Rec}_{2n}(\lambda/\nu) = \operatorname{Rec}_{2n}(\lambda/\nu)$  (Theorem 8.3.3) by proving that both  $|\operatorname{Rec}_{2n}(\lambda/\nu)|$  and  $|\operatorname{LRT}_{2n}^{Sp}(\lambda/\nu)|$  are equal to the multiplicity  $m_{\lambda,\nu}$  by Sundaram's branching rule (Theorem 8.2.3) and the bijectivity of the Littlewood-Richardson map.

#### 4. Factorization of the reduction map

The aim of this section is to prove Corollary 4.4.2, which describes the reduction map as a composite of several maps. The other part is devoted to preparing the proof.

4.1. Combinatorial *R*-matrices. Recall from (2.1) the partitions  $\varpi_l$  for  $l \ge 0$ .

**Definition 4.1.1** (cf. [NaYa97, Rule 3.10]). Let  $k, l \in [0, m]$ . The combinatorial *R*-matrix is the map

 $R = R_{k,l} : \mathrm{SST}_m(\varpi_k) \times \mathrm{SST}_m(\varpi_l) \to \mathrm{SST}_m(\varpi_l) \times \mathrm{SST}_m(\varpi_k)$ 

defined as follows. Let  $\mathbf{a} = (a_1, \ldots, a_k) \in SST_m(\varpi_k)$  and  $\mathbf{b} = (b_1, \ldots, b_l) \in SST_m(\varpi_l)$ .

#### HIDEYA WATANABE

(1) When  $k \leq l$ . For each  $r \in [1, k]$ , define  $i_r \in [1, l]$  inductively as follows. Set  $i_1$  to be the minimum  $i \in [1, l]$  such that  $b_i \geq a_1$ ; when such i does not exist, we set  $i_1 := 1$ . Suppose that  $r \geq 2$  and we have determined  $i_1, \ldots, i_{r-1}$ . Set  $i_r$  to be the minimum  $i \in [1, l] \setminus \{i_1, \ldots, i_{r-1}\}$  such that  $b_i \geq a_r$ ; when such i does not exist, we set  $i_r := \min([1, l] \setminus \{i_1, \ldots, i_{r-1}\})$ . Then, we set

$$R(\mathbf{a}, \mathbf{b}) := (\mathbf{a} \sqcup \mathbf{b}'', \mathbf{b}'),$$

where

$$\mathbf{b}' := (b_{i_1}, \ldots, b_{i_k}), \quad \mathbf{b}'' := \mathbf{b} \setminus \mathbf{b}'.$$

(2) When  $k \ge l$ . For each  $r \in [1, l]$ , define  $i_r \in [1, k]$  inductively as follows. Set  $i_1$  to be the maximum  $i \in [1, k]$  such that  $a_i \le b_1$ ; when such i does not exist, we set  $i_1 := k$ . Suppose that  $r \ge 2$  and we have determined  $i_1, \ldots, i_{r-1}$ . Set  $i_r$  to be the maximum  $i \in [1, k] \setminus \{i_1, \ldots, i_{r-1}\}$  such that  $a_i \le b_r$ ; when such i does not exist, we set  $i_r := \max([1, k] \setminus \{i_1, \ldots, i_{r-1}\})$ . Then, we set

$$R(\mathbf{a}, \mathbf{b}) := (\mathbf{a}', \mathbf{b} \sqcup \mathbf{a}''),$$

where

$$\mathbf{a}' := (a_{i_1}, \ldots, a_{i_l}), \quad \mathbf{a}'' := \mathbf{a} \setminus \mathbf{a}'$$

**Proposition 4.1.2** ([NaYa97, Proposition 3.21]). The map  $R_{k,l}$  is a bijection with inverse  $R_{l,k}$ .

Given an integer  $a \in [1, m]$ , set

(4.1) 
$$a^{\vee} := [1,m] \setminus \{a\} \in \mathrm{SST}_m(\varpi_{m-1}).$$

Lemma 4.1.3. We have

$$R(m, m^{\vee}) = (1^{\vee}, 1).$$

*Proof.* With the same notation as Definition 4.1.1(1), we see that

 $i_1 = 1.$ 

Hence, the assertion follows.

**Lemma 4.1.4.** Let  $t \in [1, m - 1]$ . Then, we have

$$R(t^{\vee}, (1, \dots, t)) = ((1, \dots, t - 1, m), m^{\vee}).$$

*Proof.* With the same notation as Definition 4.1.1(2), we see inductively that

$$i_1 = 1, i_2 = 2, \dots, i_{t-1} = t - 1$$
, and  $i_t = m - 1$ .

Hence, the assertion follows.

**Lemma 4.1.5.** Let  $\mathbf{a} = (a_1, \ldots, a_k) \in \text{SST}_m(\varpi_k)$  and  $\mathbf{b} = (b_1, \ldots, b_l) \in \text{SST}_m(\varpi_l)$  with  $k \leq l$ . Suppose that for each  $r \in [1, k]$ , there exists  $j_r \in [1, l]$  such that  $a_r = b_{j_r}$ . Then, we have

$$R(\mathbf{a},\mathbf{b}) = (\mathbf{b},\mathbf{a})$$

and

$$R(\mathbf{b}, \mathbf{a}) = (\mathbf{a}, \mathbf{b}).$$

*Proof.* The first assertion is clear from Definition 4.1.1 (1). The second assertion follows from the first one and Proposition 4.1.2.  $\Box$ 

14

**Lemma 4.1.6.** Let  $\mathbf{a} = (a_1, \ldots, a_k) \in SST_m(\varpi_k)$ , and  $r, s \in [1, m]$  be such that  $a_k + 1 < r \leq s$ . Then, we have

$$R(s^{\vee}, \mathbf{a} \sqcup [r, s]) = (\mathbf{a} \sqcup [r - 1, s - 1], (r - 1)^{\vee})$$

and

$$R(\mathbf{a} \sqcup [r-1, s-1], (r-1)^{\vee}) = (s^{\vee}, \mathbf{a} \sqcup [r, s])$$

*Proof.* With the same notation as Definition 4.1.1(2), we see inductively that

 $i_1 = a_1, \dots, i_k = a_k, i_{k+1} = r, i_{k+2} = r+1, \dots, i_{k+s-r} = s-1$ , and  $i_{k+s-r+1} = r-1$ .

Hence, the first assertion follows. Now, the second assertion follows from Proposition 4.1.2.  $\hfill \Box$ 

4.2. Some properties of removable entries. In this subsection, we fix  $al \in [0, 2n]$  and  $\mathbf{a} = (a_1, \ldots, a_l) \in SST_{2n}(\varpi_l)$ .

**Lemma 4.2.1.** Let  $i \in [1, l]$  be such that  $a_j \notin \operatorname{rem}(\mathbf{a})$  for all  $j \in [i, l]$ . Then, we have  $\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \ldots, a_{i-1}).$ 

*Proof.* The assertion is deduced by iterative applications of Proposition 2.5.3 (2).  $\Box$ 

**Proposition 4.2.2.** For each  $i \in [1, l]$ , we have  $a_i \in rem(\mathbf{a})$  if and only if one of the following hold:

- (1)  $a_i \notin 2\mathbb{Z}, i < l, a_{i+1} = a_i + 1, and a_i < 2i |\operatorname{rem}(a_1, \dots, a_{i-1})|.$
- (2)  $a_i \in 2\mathbb{Z}, i > 1, a_{i-1} = a_i 1, and a_i < 2i |\operatorname{rem}(a_1, \dots, a_{i-2})| 1.$

*Proof.* We prove the assertion by induction on l. If  $l \leq 1$ , then the assertion is clear since we have  $\operatorname{rem}(\mathbf{a}) = \emptyset$  and neither i < l nor i > 1 for all  $i \in [1, l]$ . Hence, assume that l > 1 and the assertion holds for  $0, 1, \ldots, l-1$ .

First, suppose that

(4.2) 
$$a_l \in 2\mathbb{Z}, \ a_{l-1} = a_l - 1, \ \text{and} \ a_l < 2l - |\operatorname{rem}(a_1, \dots, a_{l-2})| - 1$$

By Definition 2.5.1, we have

(4.3) 
$$\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \dots, a_{l-2}) \sqcup \{a_{l-1}, a_l\}$$

When  $i \leq l-2$ , equation (4.3) implies that  $a_i \in \text{rem}(\mathbf{a})$  if and only if  $a_i \in \text{rem}(a_1, \ldots, a_{l-2})$ . Hence, the assertion follows from our induction hypothesis.

When  $i \ge l-1$ , equation (4.3) implies that  $a_i \in \text{rem}(\mathbf{a})$ . On the other hand, condition (4.2) implies the first (resp., second) condition in the statement when i = l-1 (resp., i = l). Therefore, the assertion follows in this case.

Next, suppose that condition (4.2) fails. By Definition 2.5.1, it holds that

$$\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \ldots, a_{l-1})$$

Then, the assertion follows from our induction hypothesis.

For each  $a \in \mathbb{Z}$ , set

(4.4) 
$$s(a) := \begin{cases} a+1 & \text{if } a \notin 2\mathbb{Z}, \\ a-1 & \text{if } a \in 2\mathbb{Z} \end{cases}$$

**Lemma 4.2.3.** Let  $a \in [1, 2n]$ . Then, we have  $a \in \text{rem}(\mathbf{a})$  if and only if  $s(a) \in \text{rem}(\mathbf{a})$ . Consequently, we have  $|\text{rem}(\mathbf{a})| \in 2\mathbb{Z}$ .

*Proof.* The assertion follows from Proposition 4.2.2.

#### HIDEYA WATANABE

**Lemma 4.2.4.** Suppose that  $a_l \in 2\mathbb{Z}$  and that there exists  $r \in [0, l-1]$  such that  $a_{l-r} = a_l - r \in \text{rem}(\mathbf{a})$ . Then, we have  $a_{l-t} \in \text{rem}(\mathbf{a})$  for all  $t \in [0, r]$ . Moreover, if  $r \notin 2\mathbb{Z}$ , then it holds that

$$\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \dots, a_{l-r-1}) \sqcup [a_l - r, a_l].$$

*Proof.* Since  $a_{l-r} = a_l - r$  and  $a_{l-r} < a_{l-r+1} < \cdots < a_l$ , we have  $a_{l-t} = a_l - t$  for all  $t \in [0, r]$ .

Let us prove for each  $t \in [1, r]$  that if  $a_{l-t} \in \text{rem}(\mathbf{a})$ , then  $a_{l-t+1} \in \text{rem}(\mathbf{a})$ . It is clear that this claim implies the first assertion.

First, suppose that  $t \notin 2\mathbb{Z}$ . Then we have

$$a_{l-t+1} = a_{l-t} + 1 = s(a_{l-t}) \in rem(\mathbf{a})$$

by Lemma 4.2.3.

Next, suppose that  $t \in 2\mathbb{Z}$ . By Proposition 4.2.2, we have  $a_{l-t-1} = a_{l-t} - 1$  and

 $a_{l-t} < 2(l-t) - |\operatorname{rem}(a_1, \dots, a_{l-t-2})| - 1.$ 

Then, by Definition 2.5.1, we obtain

(4.5) 
$$|\operatorname{rem}(a_1, \dots, a_{l-t})| = |\operatorname{rem}(a_1, \dots, a_{l-t-2})| + 2$$

On the other hand, since  $t \in [1, r] \cap 2\mathbb{Z}$ , we see that l - t + 1 < l. Hence, we have  $a_{l-t+2} = a_{l-t+1} + 1$  and

$$a_{l-t+1} = a_{l-t} + 1 < 2(l-t) - |\operatorname{rem}(a_1, \dots, a_{l-t-2})| = 2(l-t+1) - |\operatorname{rem}(a_1, \dots, a_{l-t})|,$$

by equation (4.5). Then, Proposition 4.2.2 implies

$$a_{l-t+1} \in \operatorname{rem}(\mathbf{a}),$$

as desired.

So far, we have proved the first assertion. The second assertion follows from the following equality, which is obtained from Definition 2.5.1 under our hypothesis that  $r \notin 2\mathbb{Z}$ :

$$\operatorname{rem}(a_1, \dots, a_{l-r+2k+1}) = \operatorname{rem}(a_1, \dots, a_{l-r+2k-1}) \sqcup \{a_{l-r+2k}, a_{l-r+2k+1}\}$$

for all  $k \in [0, (r-1)/2]$ . Thus, we complete the proof.

**Lemma 4.2.5.** Suppose that  $a_l \in 2\mathbb{Z}$  and that there exists  $r \in [1, l-1]$  such that  $a_{l-r} = a_l - r \notin 2\mathbb{Z}$ . Then, we have

$$\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \dots, a_{l-r-1}) \sqcup [a_l - t, a_l],$$

where  $t \in [0, r]$  is the maximal odd integer such that

$$a_l - t < 2(l - t) - |\operatorname{rem}(a_1, \dots, a_{l-r-1})|;$$

when such t does not exist, we set t := -1.

*Proof.* Let  $t' \in [0, r]$  denote the maximal odd integer such that  $a_{l-t'} \in \text{rem}(\mathbf{a})$ ; when such t' does not exist, we set t' := -1. Then, by Lemma 4.2.4, we have

$$\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \dots, a_{l-t'-1}) \sqcup [a_l - t', a_l]$$

Moreover, since  $a_{l-r}, a_{l-r+1}, \ldots, a_{l-t'-1} \notin \text{rem}(\mathbf{a})$  by the definition of t', Lemma 4.2.1 implies that

(4.6) 
$$\operatorname{rem}(a_1, \dots, a_{l-t'-1}) = \operatorname{rem}(a_1, \dots, a_{l-r-1}).$$

Hence, in order to complete the proof, we only need to show that t' = t.

First, suppose that  $t' \neq -1$ . Then, by Proposition 4.2.2 and equation (4.6), we have

$$a_l - t' = a_{l-t'} < 2(l - t') - |\operatorname{rem}(a_1, \dots, a_{l-t'-1})| = 2(l - t') - |\operatorname{rem}(a_1, \dots, a_{l-r-1})|$$

This implies that  $t' \leq t$ . In particular,  $t \neq -1$ . Then, the definition of t and equation (4.6) imply that

$$a_{l-t} = a_l - t < 2(l-t) - |\operatorname{rem}(a_1, \dots, a_{l-r-1})| = 2(l-t) - |\operatorname{rem}(a_1, \dots, a_{l-t-1})|.$$

Hence, we obtain  $t \leq t'$ . Therefore, we conclude that t' = t, as desired in this case (when  $t' \neq -1$ ).

Next, suppose that t' = -1. We only need to show that there exists no odd integer  $t \in [0, r]$  such that

$$a_l - t < 2(l - t) - |\operatorname{rem}(a_1, \dots, a_{l-r-1})|.$$

Assume contrary. Then, by Proposition 4.2.2, we see that  $a_{l-t} \in \text{rem}(\mathbf{a})$ . This contradicts that t' = -1. Therefore, we obtain t' = t, as desired.

**Lemma 4.2.6.** Let  $i \in [1, l]$  be such that  $a_i \notin \text{rem}(\mathbf{a})$ . Then, we have

$$\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \dots, a_{i-1}) \sqcup ((a_{i+1}, \dots, a_l) \cap \operatorname{rem}(\mathbf{a})).$$

*Proof.* By Proposition 2.5.3 (4), the right-hand side is contained in the left-hand side. Hence, we only need to show the opposite containment.

Let  $j \in [1, l]$  be such that  $a_j \in \text{rem}(\mathbf{a})$ . By our assumption, it must hold that  $j \neq i$ .

First, suppose that j < i-1. In this case, we have  $a_j \in \text{rem}(a_1, \ldots, a_{i-1})$  by Proposition 4.2.2.

Next, suppose that j = i - 1. In this case, we have  $a_j \in 2\mathbb{Z}$ ; otherwise, it holds that  $a_i = a_{j+1} = a_j + 1$  by Proposition 4.2.2 and  $a_j + 1 = s(a_j) \in \text{rem}(\mathbf{a})$  by Lemma 4.2.3, which contradicts that  $a_i \notin \text{rem}(\mathbf{a})$ . Then, we obtain  $a_j \in \text{rem}(a_1, \ldots, a_{i-1})$  by Proposition 4.2.2.

Finally, suppose that j > i. In this case, it is clear that

$$a_j \in (a_{i+1}, \ldots, a_l) \cap \operatorname{rem}(\mathbf{a})$$

Thus, we complete the proof.

Proposition 4.2.7. We have

 $0 \le l - |\operatorname{rem}(\mathbf{a})| \le \min(l, 2n - l).$ 

In particular, it holds that

$$|\operatorname{rem}(\mathbf{a})| \ge 2(l-n).$$

*Proof.* Set  $r := |\operatorname{rem}(\mathbf{a})|$ . Since  $\operatorname{rem}(\mathbf{a})$  is a subset of  $\{a_1, \ldots, a_l\}$ , we have

$$0 \le l - r \le l.$$

Hence, we only need to show that  $l - r \leq 2n - l$ , or equivalently,

$$r \ge 2(l-n).$$

We prove this inequality by induction on l. When  $l \leq 1$ , the claim is trivial since we have  $\operatorname{rem}(\mathbf{a}) = \emptyset$  and  $n \geq 1$ .

Assume that l > 1 and our claim is true for  $0, 1, \ldots, l - 1$ . Set

$$\mathbf{a}' := (a_1, \dots, a_{l-1}) \text{ and } r' := |\operatorname{rem}(\mathbf{a}')|.$$

Then, we have  $r \ge r'$ . Note that  $a_{l-1} \le 2n - 1$ .

Suppose first that  $a_{l-1} < 2n - 1$ . Then, we have

 $\mathbf{a}' \in \mathrm{SST}_{2(n-1)}(\varpi_{l-1}).$ 

By our induction hypothesis, we obtain

$$r \ge r' \ge 2((l-1) - (n-1)) = 2(l-n),$$

as desired.

Next, suppose that  $a_{l-1} = 2n - 1$  and  $a_{l-1} \in \text{rem}(\mathbf{a})$ . In this case, Lemma 4.2.3 and Definition 2.5.1 imply that  $a_l = 2n \in \text{rem}(\mathbf{a})$  and

$$\operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1, \dots, a_{l-2}) \sqcup \{a_{l-1}, a_l\},$$

respectively. Set

$$r'' := |\operatorname{rem}(a_1, \ldots, a_{l-2})|.$$

Then, by the same argument as in the previous case, we obtain

$$r = r'' + 2 \ge 2((l-2) - (n-1)) + 2 = 2(l-n),$$

as desired.

Finally, suppose that  $a_{l-1} = 2n - 1$  and  $a_{l-1} \notin \text{rem}(\mathbf{a})$ . In this case, Proposition 4.2.2 implies

$$2n - 1 = a_{l-1} \ge 2(l-1) - r''$$

Furthermore, since  $r'' \in 2\mathbb{Z}$  by Lemma 4.2.3, it follows that

$$2n - 2 \ge 2(l - 1) - r'',$$

equivalently,

 $r'' \ge 2(l-n).$ 

Now, the assertion follows from the fact that  $r \ge r''$ .

Corollary 4.2.8. Let  $i \in [1, l]$ .

- (1) If  $a_i$  is odd, then  $|\operatorname{rem}(a_1, \ldots, a_{i-1})| \ge 2i a_i 1$ .
- (2) If  $a_i$  is even, then  $|\operatorname{rem}(a_1,\ldots,a_i)| \ge 2i a_i$ .

*Proof.* Observe that  $(a_1, \ldots, a_{i-1}) \in SST_{a_i-1}(\varpi_{i-1})$  in the first case, while  $(a_1, \ldots, a_i) \in SST_{a_i}(\varpi_i)$  in the second case. Then, the assertion follows from Proposition 4.2.7.

4.3. Some properties of the reduction map. In this subsection, we fix  $l \in [0, 2n]$  and  $\mathbf{a} = (a_1, \ldots, a_l) \in SST_{2n}(\varpi_l)$ .

**Lemma 4.3.1.** Let  $l \in [1, 2n]$  be an even integer. Then, we have

$$\operatorname{red}(1,2,\ldots,l)=()$$

*Proof.* By Corollary 4.2.8 (2), we have

 $|\operatorname{rem}(1, 2, \dots, l)| \ge l.$ 

This implies that

 $rem(1, 2, \ldots, l) = [1, l].$ 

Therefore, the assertion follows.

**Lemma 4.3.2.** Let 
$$i \in [1, l]$$
 be such that  $a_j \notin \operatorname{rem}(\mathbf{a})$  for all  $j \in [i, l]$ . Then, we have  
  $\operatorname{red}(\mathbf{a}) = \operatorname{red}(a_1, \ldots, a_{i-1}) \sqcup (a_i, \ldots, a_l).$ 

*Proof.* The assertion follows from Lemma 4.2.1.

18

**Lemma 4.3.3.** Suppose that  $a_l \in 2\mathbb{Z}$  and that there exists  $r \in [0, l-1]$  such that  $r \notin 2\mathbb{Z}$  and  $a_{l-r} = a_l - r \in \text{rem}(\mathbf{a})$ . Then, we have

$$\operatorname{red}(\mathbf{a}) = \operatorname{red}(a_1, \dots, a_{l-r-1})$$

*Proof.* The assertion follows from Lemma 4.2.4.

**Lemma 4.3.4.** Suppose that  $a_l \in 2\mathbb{Z}$  and that there exists  $r \in [1, l-1]$  such that  $a_{l-r} = a_l - r \notin 2\mathbb{Z}$ . Then, we have

$$red(\mathbf{a}) = red(a_1, \dots, a_{l-r-1}) \sqcup [a_l - r, a_l - t - 1],$$

where  $t \in [0, r]$  is the maximal odd integer such that

$$a_l - t < 2(l - t) - |\operatorname{rem}(a_1, \dots, a_{l-r-1})|;$$

when such t does not exist, we set t := -1.

*Proof.* The assertion follows from Lemma 4.2.5.

**Lemma 4.3.5.** Let us write  $red(\mathbf{a}) = (a_{i_1}, a_{i_2}, \ldots, a_{i_k})$  for some  $k \in [0, l]$  and  $1 \le i_1 < i_2 < \cdots < i_k \le l$ . Then, for each  $t \in [1, k]$ , we have

$$i_t = t + |\operatorname{rem}(a_1, a_2, \dots, a_{i_t-1})|$$

*Proof.* Clearly, we have

$$i_t = t + |(a_1, \ldots, a_{i_t}) \cap \operatorname{rem}(\mathbf{a})|$$

Since  $a_{i_t} \notin \text{rem}(\mathbf{a})$ , Lemma 4.2.6 implies

$$(a_1,\ldots,a_{i_t})\cap \operatorname{rem}(\mathbf{a}) = \operatorname{rem}(a_1,\ldots,a_{i_t-1})$$

Hence, the assertion follows.

**Proposition 4.3.6.** The tableau red(**a**) is symplectic. Consequently, the assignment  $\mathbf{a} \mapsto \operatorname{red}(\mathbf{a})$  gives rise to a map

red : 
$$\operatorname{SST}_{2n}(\varpi_l) \to \bigsqcup_{\substack{0 \le k \le \min(l, 2n-l)\\ l-k \in 2\mathbb{Z}}} Sp \operatorname{T}_{2n}(\varpi_k)$$

for each  $l \in [0, 2n]$ .

*Proof.* Let us write  $\operatorname{red}(\mathbf{a}) = (a_{i_1}, \ldots, a_{i_k})$  for some  $k \leq l$  and  $1 \leq i_1 < \cdots < i_k \leq l$ . Assume contrary that  $\operatorname{red}(\mathbf{a})$  is not symplectic. Then, by Lemma 2.4.3, there exists  $r \in [2, k]$  such that

$$a_{i_r} = 2r - 2, \ a_{i_{r-1}} = 2r - 3,$$

and

$$a_{i_s} \ge 2s - 1$$
 for all  $s \in [1, r - 2]$ .

In particular, we have

$$i_{r-1} = i_r - 1.$$

By Lemma 4.3.5, it holds that

$$i_{r-1} = (r-1) + |\operatorname{rem}(a_1, \dots, a_{i_r-2})|$$

Therefore, we have

$$2i_{r-1} - |\operatorname{rem}(a_1, \dots, a_{i_r-2})| = 2(r-1) + |\operatorname{rem}(a_1, \dots, a_{i_r-2})| \ge 2r - 2$$

So far, we have shown that  $a_{i_r}$  is even,  $a_{i_r-1} = a_{i_r} - 1$ , and

$$a_{i_r} = 2r - 2 < 2i_r - |\operatorname{rem}(a_1, \dots, a_{i_r-2})| - 1.$$

19

Then, Proposition 4.2.2 implies that  $a_{i_r} \in \text{rem}(\mathbf{a})$ . However, this contradicts that  $a_{i_r}$  is an entry of red( $\mathbf{a}$ ). Thus, we complete the proof.

**Proposition 4.3.7.** We have  $red(\mathbf{a}) = \mathbf{a}$  if and only if  $\mathbf{a}$  is symplectic.

*Proof.* First, suppose that  $red(\mathbf{a}) = \mathbf{a}$ . Then, Proposition 4.3.6 implies that the tableaux  $\mathbf{a}$  is symplectic.

Next, suppose that **a** is symplectic. Assume contrary that  $red(\mathbf{a}) \neq \mathbf{a}$ . Then, by Proposition 4.2.2, there exists  $i \in [2, l]$  such that  $a_i < 2i - 1$ . This contradicts that **a** is symplectic. Thus, the assertion follows.

**Corollary 4.3.8.** Let  $\nu \in \operatorname{Par}_{\leq n}$  and  $T \in \operatorname{SST}_{2n}(\nu)$ . Then, we have  $\operatorname{suc}(T) = T$  if and only if T is symplectic.

*Proof.* Let us write  $d(T) = (\mathbf{a}, T')$ . By Lemma 2.6.4 and its proof, we have  $\operatorname{suc}(T) = T$  if and only if  $\operatorname{red}(\mathbf{a}) = \mathbf{a}$ . The latter is equivalent to that  $\mathbf{a}$  is symplectic, which is then equivalent to that T is symplectic. Thus, the assertion follows.

4.4. Factorization of the reduction map. Recall the combinatorial *R*-matrices, the map s, and the map  $\cdot^{\vee}$  from Definition 4.1.1, (4.4), and (4.1), respectively.

**Proposition 4.4.1.** Let  $l \in [2, 2n]$ ,  $\mathbf{a} = (a_1, \dots, a_l) \in SST_{2n}(\varpi_l)$ . Let us write  $R(s(a_l)^{\vee}, (a_1, \dots, a_{l-1})) = (\mathbf{c}, \mathbf{d})$ for some  $(\mathbf{c}, \mathbf{d}) \in SST_{2n}(\varpi_{l-1}) \times SST_{2n}(\varpi_{2n-1})$ , and

$$R(\operatorname{red}(\mathbf{c}), \mathbf{d}) = (s(b_k)^{\vee}, \mathbf{b}')$$

for some  $b_k \in \mathbb{Z}$  and  $\mathbf{b}' = (b_1, \ldots, b_{k-1}) \in SST_{2n}(\varpi_{k-1})$ , where  $k := |\operatorname{red}(\mathbf{c})| + 1$ . Then, we have  $b_k > b_{k-1}$  and

$$\mathbf{b} := (b_1, \dots, b_k) = \begin{cases} (1, 2) & \text{if } a_l = l \in 2\mathbb{Z}, \\ \operatorname{red}(\mathbf{a}) & \text{otherwise.} \end{cases}$$

*Proof.* Set  $\mathbf{a}' := (a_1, \ldots, a_{l-1}).$ 

First, suppose that  $a_l = l \in 2\mathbb{Z}$ . Then, we have

$$(s(a_l)^{\vee}, (1, \dots, l-1)) = ((l-1)^{\vee}, \mathbf{a}')$$

$$\stackrel{R}{\mapsto} ((1, \dots, l-2, 2n), (2n)^{\vee}) = (\mathbf{c}, \mathbf{d})$$

$$\stackrel{(\text{red,id})}{\mapsto} ((2n), (2n)^{\vee}) = (\text{red}(\mathbf{c}), \mathbf{d})$$

$$\stackrel{R}{\mapsto} (1^{\vee}, (1)) = (s(b_k)^{\vee}, \mathbf{b}').$$

Here, we used Lemmas 4.1.4, 4.3.2, 4.3.1, and 4.1.3. As a result, we obtain  $\mathbf{b} = (1, 2)$ , as desired.

Next, suppose that  $a_l \notin 2\mathbb{Z}$ . Then, we have

$$(s(a_l)^{\vee}, \mathbf{a}') = ((a_l + 1)^{\vee}, \mathbf{a}')$$

$$\stackrel{R}{\mapsto} (\mathbf{a}', (a_l + 1)^{\vee}) = (\mathbf{c}, \mathbf{d})$$

$$\stackrel{(\text{red}, \text{id})}{\mapsto} (\text{red}(\mathbf{a}'), (a_l + 1)^{\vee}) = (\text{red}(\mathbf{c}), \mathbf{d})$$

$$\stackrel{R}{\mapsto} ((a_l + 1)^{\vee}, \text{red}(\mathbf{a}')) = (s(b_k)^{\vee}, \mathbf{b}').$$

Here, we used Lemma 4.1.5. This implies that  $\mathbf{b}' = \operatorname{red}(\mathbf{a}')$  and  $b_k = s(a_l + 1) = a_l$ . Hence, we obtain

$$b_k = a_l > a_{l-1} \ge b_{k-1}$$

since  $\mathbf{b}' = \operatorname{red}(\mathbf{a}')$  is a subsequence of  $\mathbf{a}' = (a_1, \ldots, a_{l-1})$ .

On the other hand, since  $a_l + 1 \notin \text{rem}(\mathbf{a})$ , we have  $a_l = s(a_l + 1) \notin \text{rem}(\mathbf{a})$  by Lemma 4.2.3. Then, Lemma 4.3.2 implies

$$\operatorname{red}(\mathbf{a}) = \operatorname{red}(\mathbf{a}') \sqcup \{a_l\} = \mathbf{b},$$

as desired.

Finally, suppose that  $a_l \in 2\mathbb{Z}$  and  $a_l \neq l$ . Set

$$r := \max\{i \in [0, l-1] \mid a_{l-i} = a_l - i\}, \quad \mathbf{a}'' := (a_1, \dots, a_{l-r-1}).$$

Note that r < l - 1. Then, we have  $\mathbf{a} = \mathbf{a}'' \sqcup [a_l - r, a_l], \ \mathbf{a}' = \mathbf{a}'' \sqcup [a_l - r, a_l - 1], s(a_l) = a_l - 1$ , and

$$(\mathbf{c}, \mathbf{d}) = R((a_l - 1)^{\vee}, \mathbf{a}') = ((\mathbf{a}'' \sqcup [a_l - r - 1, a_l - 2]), (a_l - r - 1)^{\vee}).$$

The last equality follows from Lemma 4.1.6.

(1) When  $r \in 2\mathbb{Z}$ . Since  $a_l - r - 1 \notin 2\mathbb{Z}$  and  $a_l - 2 \in 2\mathbb{Z}$ , we can apply Lemma 4.3.4 to obtain

$$\operatorname{red}(\mathbf{c}) = \operatorname{red}(\mathbf{a}'') \sqcup [a_l - r - 1, a_l - t - 3],$$

where t is the maximal odd integer in [0, r-1] such that

 $(a_l - 2) - t < 2((l - 1) - t) - |\operatorname{rem}(\mathbf{a}'')|;$ 

when such t does not exist, we set t := -1. Then, by Lemma 4.1.6 again, we have

 $R(\operatorname{red}(\mathbf{c}), \mathbf{d}) = ((a_l - t - 2)^{\vee}, \operatorname{red}(\mathbf{a}'') \sqcup [a_l - r, a_l - t - 2]).$ 

This implies that  $\mathbf{b}' = \operatorname{red}(\mathbf{a}'') \sqcup [a_l - r, a_l - t - 2], b_k = s(a_l - t - 2) = a_l - t - 1,$ and

$$\mathbf{b} = \operatorname{red}(\mathbf{a}'') \sqcup [a_l - r, a_l - t - 1].$$

On the other hand, we have

$$\mathbf{a} = \mathbf{a}'' \sqcup [a_l - r, a_l] = (\mathbf{a}'' \sqcup (a_l - r)) \sqcup [a_l - r + 1, a_l].$$

By Lemma 4.3.4, we have

$$\operatorname{red}(\mathbf{a}) = \operatorname{red}(\mathbf{a}'' \sqcup (a_l - r)) \sqcup [a_l - r + 1, a_l - t' - 1],$$

where t' is the maximal odd integer in [0, r-1] such that

$$a_l - t' < 2(l - t') - |\operatorname{rem}(\mathbf{a}'' \sqcup (a_l - r))|;$$

when such t' does not exist, we set t' := -1. Since  $s(a_l - r) = a_l - r - 1 \notin \mathbf{a}''$ , we have

 $\operatorname{rem}(\mathbf{a}'' \sqcup (a_l - r)) = \operatorname{rem}(\mathbf{a}'') \text{ and } \operatorname{red}(\mathbf{a}'' \sqcup (a_l - r)) = \operatorname{red}(\mathbf{a}'') \sqcup (a_l - r).$ 

Therefore, we obtain

$$\operatorname{red}(\mathbf{a}) = \operatorname{red}(\mathbf{a}'') \sqcup [a_l - r, a_l - t' - 1]$$

Now, one can straightforwardly verify that t' = t. Hence, we conclude that

$$\operatorname{red}(\mathbf{a}) = \operatorname{red}(\mathbf{a}'') \sqcup [a_l - r, a_l - t - 1] = \mathbf{b}_l$$

as desired.

(2) When  $r \notin 2\mathbb{Z}$  and  $a_l - r - 1 \notin \text{rem}(\mathbf{c})$ . This case can be proved as in the same way as the previous case.

(3) When  $r \notin 2\mathbb{Z}$  and  $a_l - r - 1 \in \text{rem}(\mathbf{c})$ . Since  $a_l - r - 2 = s(a_l - r - 1) \in \text{rem}(\mathbf{c})$ , we see that

$$a_{l-r-1} = a_l - r - 2.$$

 $\operatorname{Set}$ 

$$\mathbf{a}''' := (a_1, \ldots, a_{l-r-2}).$$

Then, we have

$$\mathbf{c} = \mathbf{a}''' \sqcup [a_l - r - 2, a_l - 2]$$

with  $a_l - r - 2 \in \text{rem}(\mathbf{c})$ . By Lemma 4.3.3, we obtain

$$red(\mathbf{c}) = red(\mathbf{a}''')$$

Then, Lemma 4.1.5 implies

$$R(\operatorname{red}(\mathbf{c}), \mathbf{d}) = (\mathbf{d}, \operatorname{red}(\mathbf{c})),$$

and consequently,

$$\mathbf{b} = \operatorname{red}(\mathbf{a}''') \sqcup (a_l - r - 2).$$

On the other hand, we have

(

$$\mathbf{a} = (\mathbf{a}^{\prime\prime\prime} \sqcup (a_l - r - 2)) \sqcup [a_l - r, a_l].$$

Since  $a_l - r - 2 \notin 2\mathbb{Z}$ , Proposition 2.5.3 (2) implies that

$$\operatorname{rem}(\mathbf{a}''' \sqcup (a_l - r - 2)) = \operatorname{rem}(\mathbf{a}''').$$

On the other hand, the fact that  $a_l - r - 2 \in \text{rem}(\mathbf{c})$ , together with Proposition 4.2.2, implies

$$a_l - r - 2 < 2(l - r - 1) - |\operatorname{rem}(\mathbf{a}''')|.$$

These two equalities and Proposition 4.2.2 ensure that

$$a_l - r \in \operatorname{rem}(\mathbf{a}).$$

Then, Lemma 4.3.3 implies

$$\operatorname{red}(\mathbf{a}) = \operatorname{red}(\mathbf{a}''') \sqcup (a_l - r - 2) = \mathbf{b},$$

as desired.

In order to state the main result in this section, let us introduce four maps:

• 
$$\bigvee : \mathrm{SST}_{2n}(\varpi_l) \to \mathrm{SST}_{2n}(\varpi_1) \times \mathrm{SST}_{2n}(\varpi_{l-1}); \ (a_1, \ldots, a_l) \mapsto ((a_l), (a_1, \ldots, a_{l-1})).$$

•  $K : \mathrm{SST}_{2n}(\varpi_1) \to \mathrm{SST}_{2n}(\varpi_{2n-1}); \ (a) \mapsto s(a)^{\vee}.$ 

$$\bigwedge : \mathrm{SST}_{2n}(\varpi_1) \times \mathrm{SST}_{2n}(\varpi_k) \to \mathrm{SST}_{2n}(\varpi_{k+1}) \sqcup \{0\};$$
$$((a), (a_1, \dots, a_k)) \mapsto \begin{cases} (a_1, \dots, a_k, a) & \text{if } a > a_k \\ 0 & \text{if } a \le a_k \end{cases}$$

1

where 0 is a formal symbol.

• 
$$\pi : \mathrm{SST}_{2n}(\varpi_{k'}) \to \bigsqcup_{k''=0}^{2n} \mathrm{SST}_{2n}(\varpi_{k''}); \ \mathbf{a} \mapsto \begin{cases} () & \text{if } \mathbf{a} = (1,2), \\ \mathbf{a} & \text{if } \mathbf{a} \neq (1,2). \end{cases}$$

Corollary 4.4.2. Let  $l \in [0, 2n]$ .

(1) If  $l \geq 2$ , then the composite

$$\pi \circ \bigwedge \circ (K^{-1}, \mathrm{id}) \circ R \circ (\mathrm{red}, \mathrm{id}) \circ R \circ (K, \mathrm{id}) \circ \bigvee$$

is well defined on  $SST_{2n}(\varpi_l)$  and coincides with the reduction map.

(2) The reduction map on  $SST_{2n}(\varpi_l)$  is injective.

*Proof.* The first assertion follows from Proposition 4.4.1 and Lemma 4.3.1. The second assertion for  $l \leq 1$  is trivial, and for  $l \geq 2$  follows from the first one since each factor in the composite is injective (on a suitable domain).

**Corollary 4.4.3.** Let  $\lambda \in \operatorname{Par}_{\leq 2n}$ . Then, the successor map in injective on  $\operatorname{SST}_{2n}(\lambda)$ .

*Proof.* The assertion follows from the injectivity of the map d (see (2.11) for the definition), the reduction map (Corollary 4.4.2 (2)), and the Pieri's formula (Proposition 2.3.4).

#### 5. Preliminaries from Lie Algebras

In this section, we briefly review representation theory of  $\mathfrak{gl}_{2n}(\mathbb{C})$  and  $\mathfrak{sp}_{2n}(\mathbb{C})$ .

5.1. General linear algebras. Let  $\mathfrak{g}$  denote the general linear algebra  $\mathfrak{gl}_{2n}(\mathbb{C})$ , that is, the complex Lie algebra  $\mathbb{C}$  consisting of all  $2n \times 2n$  complex matrices.

For each  $i, j \in [1, 2n]$ , let  $E_{i,j}$  denote the matrix unit with entry 1 at (i, j) position. For each  $i \in [1, 2n]$ , set

$$d_i := E_{i,i}.$$

Also, for each  $i \in [1, 2n - 1]$ , set

$$h_i := d_i - d_{i+1}$$

Let M be a finite-dimensional  $\mathfrak{g}$ -module. Then, it decomposes into its weight spaces:

$$M = \bigoplus_{\mathbf{w} = (w_1, \dots, w_{2n}) \in \mathbb{Z}^{2n}} M_{\mathbf{w}}$$

where

$$M_{\mathbf{w}} := \{ m \in M \mid d_i m = w_i m \text{ for all } i \in [1, 2n] \}.$$

The character of M is the Laurent polynomial given by

$$\operatorname{ch}_{\mathfrak{g}} M = \sum_{\mathbf{w} \in \mathbb{Z}^{2n}} (\dim M_{\mathbf{w}}) x_1^{w_1} \cdots x_{2n}^{w_{2n}} \in \mathbb{Z}[x_1^{\pm 1}, \dots, x_{2n}^{\pm 1}].$$

For each  $\lambda \in \operatorname{Par}_{\leq 2n}$ , there exists a unique, up to isomorphism, finite-dimensional irreducible  $\mathfrak{g}$ -module  $V^{\mathfrak{g}}(\lambda)$  such that

$$\operatorname{ch}_{\mathfrak{g}} V^{\mathfrak{g}}(\lambda) = s_{\lambda}(x_1, \dots, x_{2n}),$$

where the right-hand side denotes the Schur function (see (2.6) for the definition).

5.2. Symplectic Lie algebras. Consider the 2*n*-dimensional complex vector space  $\mathbb{C}^{2n}$  with a standard basis  $\{e_1, \ldots, e_{2n}\}$ . Let  $\langle, \rangle$  denote the skew-symmetric bilinear form on  $\mathbb{C}^{2n}$  given by

$$\langle e_i, e_j \rangle = \begin{cases} 1 & \text{if } j - i = n, \\ -1 & \text{if } i - j = n, \\ 0 & \text{otherwise.} \end{cases}$$

Let  $\mathfrak{s}$  denote the symplectic Lie algebra  $\mathfrak{sp}_{2n}(\mathbb{C})$ , that is, the subalgebra of  $\mathfrak{g}$  consisting of  $X \in \mathfrak{g}$  such that

$${}^{t}XJ_n + J_nX = O_{2n},$$

where  $O_{2n}$  denote the zero matrix of size 2n,

$$J_n := \begin{pmatrix} O_n & I_n \\ -I_n & O_n \end{pmatrix}$$

the matrix representation of the skew-symmetric bilinear form  $\langle,\rangle$  with respect to the standard basis with  $I_n$  the identity matrix of size n.

For each  $i \in [1, n]$ , set

$$h'_i := \begin{cases} h_i - h_{n+i} & \text{if } i < n, \\ d_n - d_{2n} & \text{if } i = n. \end{cases}$$

Then, we have  $h'_i \in \mathfrak{s}$ .

Let M be a finite-dimensional  $\mathfrak{s}$ -module. Then, it decomposes into its weight spaces:

$$M = \bigoplus_{\mathbf{z}=(z_1,\dots,z_n)\in\mathbb{Z}^n} M_{\mathbf{z}},$$

where

$$M_{\mathbf{z}} := \{ m \in M \mid h'_i m = (z_i - z_{i+1})m \text{ for all } i \in [1, n-1] \text{ and } h'_n m = z_n m \}$$

The character of M is the Laurent polynomial given by

$$\operatorname{ch}_{\mathfrak{s}} M = \sum_{\mathbf{z} \in \mathbb{Z}^n} (\dim M_{\mathbf{z}}) y_1^{z_1} \cdots y_n^{z_n} \in \mathbb{Z}[y_1^{\pm 1}, \dots, y_n^{\pm 1}].$$

For each  $\nu \in \operatorname{Par}_{\leq n}$ , there exists a unique finite-dimensional irreducible  $\mathfrak{s}$ -module  $V^{\mathfrak{s}}(\nu)$  such that

$$\operatorname{ch}_{\mathfrak{s}} V^{\mathfrak{s}}(\nu) = s_{\nu}^{Sp}(y_1, \dots, y_n)$$

where the right-hand side denotes the symplectic Schur function (see (2.9) for the definition).

Let  $M = \bigoplus_{\mathbf{w} \in \mathbb{Z}^{2n}} M_{\mathbf{w}}$  be a finite-dimensional  $\mathfrak{g}$ -module. For each  $m \in M_{\mathbf{w}}$ , we have

$$h'_{i}m = \begin{cases} (w_{i} - w_{i+1} - w_{n+i} + w_{n+i+1})m & \text{if } i < n, \\ (w_{n} - w_{2n})m & \text{if } i = n, \end{cases} \text{ for all } i \in [1, n].$$

Therefore, as an  $\mathfrak{s}$ -module, the M decomposes as

$$M = \bigoplus_{\mathbf{z} \in \mathbb{Z}^n} M_{\mathbf{z}}, \quad M_{\mathbf{z}} = \bigoplus_{\mathbf{w} \in (\operatorname{res}^{\mathfrak{s}})^{-1}(\mathbf{z})} M_{\mathbf{w}},$$

where

$$\operatorname{res}^{\mathfrak{s}}: \mathbb{Z}^{2n} \to \mathbb{Z}^{n}; \ (w_{1}, \dots, w_{2n}) \mapsto (w_{1} - w_{n+1}, w_{2} - w_{n+2}, \dots, w_{n} - w_{2n}).$$

This observation implies that

$$\operatorname{ch}_{\mathfrak{s}} M = \operatorname{res}^{\mathfrak{s}}(\operatorname{ch}_{\mathfrak{a}} M),$$

where

$$\operatorname{res}^{\mathfrak{s}}: \mathbb{Z}[x_1^{\pm 1}, \dots, x_{2n}^{\pm 1}] \to \mathbb{Z}[y_1^{\pm 1}, \dots, y_n^{\pm 1}]$$

denotes the ring homomorphism such that

$$\operatorname{res}^{\mathfrak{s}}(x_i) = \begin{cases} y_i & \text{if } i \leq n, \\ y_{i-n}^{-1} & \text{if } i > n. \end{cases}$$

In particular, for each  $\lambda \in \operatorname{Par}_{\leq 2n}$ , we obtain

(5.1) 
$$\operatorname{ch}_{\mathfrak{s}} V^{\mathfrak{g}}(\lambda) = s_{\lambda}(y_1, \dots, y_n, y_1^{-1}, \dots, y_n^{-1}).$$

On the other hand, as an  $\mathfrak{s}$ -module,  $V^{\mathfrak{g}}(\lambda)$  decomposes into a direct sum of several copies of  $V^{\mathfrak{s}}(\nu)$  for various  $\nu \in \operatorname{Par}_{\leq n}$ :

(5.2) 
$$V^{\mathfrak{g}}(\lambda) \simeq \bigoplus_{\nu \in \operatorname{Par}_{\leq n}} V^{\mathfrak{s}}(\nu)^{m_{\lambda,\nu}} \text{ for some } m_{\lambda,\nu} \ge 0.$$

An explicit descriptions of the multiplicities  $m_{\lambda,\nu}$  is called a *branching rule*.

5.3. A Non-standard realization of the symplectic algebra. Let  $\mathfrak{k}$  denote the Lie subalgebra of  $\mathfrak{g}$  generated by

 $\{E_{2i-1,2i}, E_{2i,2i-1} \mid i \in [1,n]\} \sqcup \{E_{2i+1,2i} + E_{2i-1,2i+2} \mid i \in [1,n-1]\}.$ 

Then, there exists an isomorphism

$$f:\mathfrak{s}\to\mathfrak{k}$$

of Lie algebras such that

$$f(h'_i) = \begin{cases} (-1)^{i-1}(h_{2i-1} + h_{2i+1}) & \text{if } i < n, \\ (-1)^{n-1}h_{2n-1} & \text{if } i = n, \end{cases}$$

(cf. [Wat21, Section 4.3]).

The notions of weight  $\mathfrak{k}$ -modules and their characters are defined in the same way as those for  $\mathfrak{s}$  by replacing  $h'_i$  with  $f(h'_i)$ .

For each  $\nu \in \operatorname{Par}_{\leq n}$ , the irreducible  $\mathfrak{s}$ -module  $V^{\mathfrak{s}}(\nu)$  is equipped with a  $\mathfrak{k}$ -module structure via the isomorphism f. Let  $V^{\mathfrak{k}}(\nu)$  denote this  $\mathfrak{k}$ -module. Clearly, we have

$$\operatorname{ch}_{\mathfrak{k}} V^{\mathfrak{k}}(\nu) = \operatorname{ch}_{\mathfrak{s}} V^{\mathfrak{s}}(\nu) = s_{\nu}^{Sp}(y_1, \dots, y_n).$$

Let M be a  $\mathfrak{g}$ -module,  $\mathbf{w} = (w_1, \ldots, w_{2n}) \in \mathbb{Z}^{2n}$ , and  $m \in M_{\mathbf{w}}$ . Then, for each  $i \in [1, n]$ , we have

$$f(h'_i)m = \begin{cases} (-1)^{i-1}(w_{2i-1} - w_{2i} + w_{2i+1} - w_{2i+2})m & \text{if } i < n, \\ (-1)^{n-1}(w_{2n-1} - w_{2n})m & \text{if } i = n. \end{cases}$$

Therefore, defining a map

$$\operatorname{res}^{\mathfrak{k}}:\mathbb{Z}^{2n}\to\mathbb{Z}^n$$

by

$$\operatorname{res}^{\mathfrak{e}}(w_1,\ldots,w_{2n})=(w_1-w_2,-w_3+w_4,w_5-w_6,\ldots,(-1)^{n-1}(w_{2n-1}-w_{2n})),$$

we obtain the following weight space decomposition of M as a  $\mathfrak{k}$ -module:

$$M = \bigoplus_{\mathbf{z} \in \mathbb{Z}^n} M_{\mathbf{z}}, \quad M_{\mathbf{z}} = \bigoplus_{\mathbf{w} \in (\operatorname{res}^{\mathfrak{k}})^{-1}(\mathbf{z})} M_{\mathbf{w}}$$

Consequently, we have

(5.3) 
$$\operatorname{ch}_{\mathfrak{k}} M = \operatorname{res}^{\mathfrak{e}}(\operatorname{ch}_{\mathfrak{g}} M),$$

where

$$\operatorname{res}^{\mathfrak{k}}: \mathbb{Z}[x_1^{\pm 1}, \dots, x_{2n}^{\pm 1}] \to \mathbb{Z}[y_1^{\pm 1}, \dots, y_n^{\pm 1}]$$

denotes the ring homomorphism given by

$$\operatorname{res}^{\mathfrak{k}}(x_{2i-1}) = \begin{cases} y_i & \text{if } i \text{ is odd,} \\ y_i^{-1} & \text{if } i \text{ is even,} \end{cases} \operatorname{res}^{\mathfrak{k}}(x_{2i}) = \begin{cases} y_i^{-1} & \text{if } i \text{ is odd,} \\ y_i & \text{if } i \text{ is even,} \end{cases} \text{ for all } i \in [1, n].$$

Proposition 5.3.1. We have

$$\operatorname{ch}_{\mathfrak{k}} V^{\mathfrak{g}}(\lambda) = \sum_{\nu \in \operatorname{Par}_{\leq n}} m_{\lambda,\nu} s_{\nu}^{Sp}(y_1, \dots, y_n).$$

*Proof.* Using the fact that the Schur functions are symmetric, we compute as follows:

$$\operatorname{ch}_{\mathfrak{k}} V^{\mathfrak{g}}(\lambda) \stackrel{(5.3)}{=} \operatorname{res}^{\mathfrak{k}}(\operatorname{ch}_{\mathfrak{g}} V^{\mathfrak{g}}(\lambda))$$

$$= s_{\lambda}(y_{1}, y_{1}^{-1}, y_{2}^{-1}, y_{2}, \dots, y_{n}^{(-1)^{n-1}}, y_{n}^{(-1)^{n}})$$

$$= s_{\lambda}(y_{1}, \dots, y_{n}, y_{1}^{-1}, \dots, y_{n}^{-1})$$

$$\stackrel{(5.1)}{=} \operatorname{ch}_{\mathfrak{s}} V^{\mathfrak{g}}(\lambda)$$

$$\stackrel{(5.2)}{=} \sum_{\nu \in \operatorname{Par}_{\leq n}} m_{\lambda,\nu} s_{\nu}^{Sp}(y_{1}, \dots, y_{n}).$$

Thus, the assertion follows.

## 6. Preliminaries from quantum symmetric pairs

In this section, we briefly review representation theory of the quantum group **U** of  $\mathfrak{gl}_{2n}(\mathbb{C})$  and an *i*quantum group  $\mathbf{U}^i$  of  $\mathfrak{sp}_{2n}(\mathbb{C})$ .

6.1. Quantum group of type A. Let U denote the quantum group  $U_q(\mathfrak{gl}_{2n})$ , that is, the unital associative algebra over  $\mathbb{Q}(q)$  with generators

$$E_i, F_i, D_k^{\pm 1}$$
 for  $i \in [1, 2n - 1]$  and  $k \in [1, 2n]$ 

subject to the following relations:

$$\begin{split} D_k D_k^{-1} &= D_k^{-1} D_k = 1, \\ D_k D_l &= D_l D_k, \\ D_k E_i &= q^{\delta_{k,i} - \delta_{k,i+1}} E_i D_k, \quad D_k F_i = q^{-\delta_{k,i} + \delta_{k,i+1}} F_i D_k, \\ E_i F_j - F_j E_i &= \delta_{i,j} \frac{K_i - K_i^{-1}}{q - q^{-1}}, \\ E_i E_j &= E_j E_i, \quad F_i F_j = F_j F_i, \\ E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0, \quad F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 = 0, \end{split}$$

where

$$K_i := D_i D_{i+1}^{-1}$$

A U-module M is said to be a *weight module* if it admits a decomposition

$$M = \bigoplus_{\mathbf{w} \in \mathbb{Z}^{2n}} M_{\mathbf{w}}$$

as a vector space such that

$$M_{\mathbf{w}} = \{ m \in M \mid D_i m = q^{w_i} m \text{ for all } i \in [1, 2n] \}$$

The character of a finite-dimensional weight **U**-module M is the Laurent polynomial  $\operatorname{ch} M \in \mathbb{Z}[x_1^{\pm 1}, \ldots, x_{2n}^{\pm 1}]$  given by

$$\operatorname{ch} M = \sum_{\mathbf{w} \in \mathbb{Z}^{2n}} (\dim M_{\mathbf{w}}) \mathbf{x}^{\mathbf{w}}.$$

26

By [Lus93, 19.1.1], there exists a unique anti-algebra automorphism  $\rho$  on U such that

$$\rho(E_i) = qK_iF_i, \quad \rho(F_i) = qK_i^{-1}E_i, \quad \rho(K_i) = K_i \text{ for all } i \in I$$

A symmetric bilinear form (,) on a U-module M is said to be contragredient if

$$(xm_1, m_2) = (m_1, \rho(x)m_2)$$
 for all  $x \in \mathbf{U}, m_1, m_2 \in M$ 

A basis B of such M is said to be *almost orthonormal* if

$$(b_1, b_2) \in \delta_{b_1, b_2} + q^{-1} \mathbb{Q}[\![q^{-1}]\!]$$
 for all  $b_1, b_2 \in B$ .

For each  $m_1, m_2 \in M$ , we write  $m_1 \equiv m_2$  to indicate that

$$m_1 - m_2 \in q^{-1} \mathbb{Q}[\![q^{-1}]\!] B.$$

Given two **U**-modules M and N with almost orthonormal bases  $B_M$  and  $B_N$  respectively, we see that the tensor product  $B_M \otimes B_N := \{b_1 \otimes b_2 \mid (b_1, b_2) \in B_M \times B_N\}$  forms an almost orthonormal basis of  $M \otimes N$  with respect to the natural bilinear form.

For each  $\lambda \in \operatorname{Par}_{\leq 2n}$ , there exists a unique, up to isomorphism, finite-dimensional irreducible U-module  $V(\lambda)$  such that

$$\operatorname{ch} V(\lambda) = s_{\lambda}(x_1, \dots, x_{2n}).$$

The weight space  $V(\lambda)_{\lambda}$  is one-dimensional. Fix a nonzero vector  $v_{\lambda} \in V(\lambda)_{\lambda}$ . Then, there exists a unique contragredient form (,) on  $V(\lambda)$  such that  $(v_{\lambda}, v_{\lambda}) = 1$ .

There exists a distinguished basis  $\mathbf{B}(\lambda)$ , called the *canonical basis*, of  $V(\lambda)$ . It is of the form

$$\mathbf{B}(\lambda) = \{ b_T \mid T \in \mathrm{SST}_{2n}(\lambda) \},\$$

and the vector  $b_T$  is a weight vector of weight wt(T) (see (2.7) for the definition):

$$b_T \in V(\lambda)_{\mathrm{wt}(T)}.$$

The canonical basis forms an almost orthonormal basis.

### Example 6.1.1.

(1) The U-module structure of  $V(\varpi_1)$  is as follows:

$$E_i b_{(a)} = \delta_{a,i+1} b_{(i)},$$
  

$$F_i b_{(a)} = \delta_{a,i} b_{(i+1)},$$
  

$$D_k b_{(a)} = q^{\delta_{a,k}} b_{(a)}.$$

(2) The U-module structure of  $V(\varpi_{2n-1})$  is as follows (recall the map  $\cdot^{\vee}$  from (4.1)):

$$\begin{split} E_i b_{a^{\vee}} &= \delta_{a,i} b_{(i+1)^{\vee}}, \\ F_i b_{a^{\vee}} &= \delta_{a,i+1} b_{i^{\vee}}, \\ D_k b_{a^{\vee}} &= q^{1-\delta_{a,k}} b_{a^{\vee}}. \end{split}$$

# **Proposition 6.1.2** (*cf.* [Kwo09]).

(1) Let  $l \in [1, 2n]$ . Then, there exists an injective U-module homomorphism

$$\bigvee: V(\varpi_l) \to V(\varpi_1) \otimes V(\varpi_{l-1})$$

such that

$$\bigvee (b_{(a_1,\ldots,a_l)}) \equiv b_{(a_l)} \otimes b_{(a_1,\ldots,a_{l-1})}$$

for all  $(a_1, \ldots, a_l) \in SST_{2n}(\varpi_l)$ .

(2) Let  $k' \in [0, 2n - 1]$ . Then, there exists a U-module homomorphism

 $\bigwedge: V(\varpi_1) \otimes V(\varpi_{k'}) \to V(\varpi_{k'+1})$ 

such that

$$\bigwedge (b_{(a)} \otimes b_{(a_1,...,a_{k'})}) \equiv \begin{cases} b_{(a_1,...,a_{k'},a)} & \text{if } a > a_{k'}, \\ 0 & \text{if } a \le a_{k'}. \end{cases}$$

for all  $a \in [1, 2n]$  and  $(a_1, \ldots, a_{k'}) \in SST_{2n}(\varpi_{k'})$ . (3) Let  $k, l \in [0, 2n]$ . Then, there exists a U-module isomorphism

$$R = R_{k,l} : V(\varpi_k) \otimes V(\varpi_l) \to V(\varpi_l) \otimes V(\varpi_k)$$

such that

$$R(b_{\mathbf{a}} \otimes b_{\mathbf{b}}) \equiv b_{\mathbf{c}} \otimes b_{\mathbf{d}}$$
  
for all  $\mathbf{a} \in SST_{2n}(\varpi_k)$  and  $\mathbf{b} \in SST_{2n}(\varpi_l)$ , where  
 $(\mathbf{c}, \mathbf{d}) := R(\mathbf{a}, \mathbf{b}).$ 

(4) Let 
$$l \in [0, 2n]$$
 and  $\mu \in Par_{\leq 2n}$ . Then, there exists a U-module isomorphism

$$*: V(\varpi_l) \otimes V(\mu) \to \bigoplus_{\substack{\lambda \in \operatorname{Par}_{\leq 2n} \\ \mu \subseteq \lambda \\ vert}} V(\lambda)$$

such that

$$*(b_S \otimes b_T) \equiv b_{S*T}$$

for all  $(S,T) \in SST_{2n}(\varpi_l) \times SST_{2n}(\mu)$ .

(5) Let  $\lambda \in \operatorname{Par}_{\leq 2n}$ , and set  $l := \ell(\lambda)$ . Let  $\lambda'$  denote the partition  $(\lambda_1 - 1, \dots, \lambda_l - 1)$ . Then, there exists a U-module homomorphism

$$d: V(\lambda) \to V(\varpi_l) \otimes V(\lambda')$$

such that

$$d(b_T) \equiv b_{\mathbf{a}} \otimes b_{T'}$$

for all  $T \in SST_{2n}(\lambda)$ , where

$$(\mathbf{a}, T') := d(T).$$

6.2. Quantum symmetric pair of type AII. For each  $i \in [1, n]$ , set

$$B_{2i} := F_{2i} - q[E_{2i-1}, [E_{2i+1}, E_{2i}]_{q^{-1}}]_{q^{-1}} K_{2i}^{-1} \in \mathbf{U}.$$

Here,  $[,]_{q^{-1}}$  denotes the q-commutator given by

$$[x, y]_{q^{-1}} := xy - q^{-1}yx.$$

Let  $\mathbf{U}^i$  denote the subalgebra of  $\mathbf{U}$  generated by

$$\{E_{2i-1}, F_{2i-1}, K_{2i-1}^{\pm 1} \mid i \in [1, n]\} \sqcup \{B_{2i} \mid i \in [1, n-1]\}.$$

The pair  $(\mathbf{U}, \mathbf{U}^i)$  forms a quantum symmetric pair of type  $AII_{2n-1}$ .

A U<sup>i</sup>-module M is said to be a *weight module* if it admits a decomposition

$$M = \bigoplus_{\mathbf{z} \in \mathbb{Z}^n} M_{\mathbf{z}}$$

as a vector space such that

$$M_{\mathbf{z}} = \{ m \in M \mid K_{2i-1}m = q^{z_{2i-1}-z_{2i}}m \text{ for all } i \in [1,n] \}.$$

The character of a finite-dimensional weight  $\mathbf{U}^i$ -module M is the Laurent polynomial  $\operatorname{ch}_i M \in \mathbb{Z}[y_1^{\pm 1}, \ldots, y_n^{\pm 1}]$  given by

$$\operatorname{ch}_{i} M = \sum_{\mathbf{z} \in \mathbb{Z}^{n}} (\dim M_{\mathbf{z}}) \mathbf{y}^{\mathbf{z}}.$$

**Proposition 6.2.1.** Let M be a finite-dimensional weight U-module. Then, we have  $\operatorname{ch}_{i} M = \operatorname{res}^{\mathfrak{k}}(\operatorname{ch} M).$ 

For each  $\nu \in \operatorname{Par}_{\leq n}$ , there exists, up to isomorphism, a unique finite-dimensional irreducible  $\mathbf{U}^{i}$ -module  $V^{i}(\nu)$  such that

$$\operatorname{ch}_{i} V^{i}(\nu) = s_{\nu}^{Sp}(y_{1}, \dots, y_{n})$$

(cf. [Wat21, Proposition 3.3.9 and Corollary 4.3.2]).

The anti-automorphism  $\rho$  on U restricts to an anti-automorphism on U<sup>*i*</sup>. The notions of contragredient inner product, almost orthogonal basis, and binary relation  $\equiv$  are defined for U<sup>*i*</sup>-modules as in the same way as U-modules.

Let M be a **U**-module equipped with a contragredient inner product. When we regard the **U**-module M as a **U**<sup>*i*</sup>-module by restriction, the inner product is still contragredient. In particular, the irreducible **U**-module  $V(\lambda)$ , regarded as a **U**<sup>*i*</sup>-module, admits a contragredient inner product. Hence, it is completely reducible.

**Proposition 6.2.2.** Let  $\lambda \in \text{Par}_{\leq 2n}$ . Then, the multiplicity of  $V^i(\nu)$  in  $V(\lambda)$  coincides with  $m_{\lambda,\nu}$ .

Proof. Let us write

$$V(\lambda) \simeq \bigoplus_{\nu \in \operatorname{Par}_{< n}} V^{i}(\nu)^{m'_{\lambda,\nu}}$$

for some  $m'_{\lambda,\nu} \geq 0$ . Then, we have

$$\sum_{\nu \in \operatorname{Par}_{<_n}} m'_{\lambda,\nu} s_{\nu}^{Sp}(y_1, \dots, y_n) = \operatorname{ch}_{\iota} V(\lambda) = \operatorname{res}^{\mathfrak{k}}(s_{\lambda}(x_1, \dots, x_{2n})) = \operatorname{ch}_{\mathfrak{k}} V^{\mathfrak{g}}(\lambda),$$

where the second equality follows from Proposition 6.2.1. Now, the assertion follows from Proposition 5.3.1 and the linearly independence of the symplectic Schur functions.  $\Box$ 

**Lemma 6.2.3.** There exists a  $U^i$ -module isomorphism

$$K: V(\varpi_1) \to V(\varpi_{2n-1})$$

such that

$$K(b_{(a)}) = b_{s(a)^{\vee}}$$
 for all  $a \in [1, 2n]$ .

*Proof.* Clearly, the linear map K defined as above is an isomorphism. Using Example 6.1.1, one can straightforwardly verify that

$$K(xb_{(a)}) = xb_{s(a)^{\vee}}$$
 for all  $x \in \mathbf{U}^i$  and  $a \in [1, 2n]$ 

Hence, the map K is a  $\mathbf{U}^{i}$ -module isomorphism.

**Lemma 6.2.4.** There exists a  $U^i$ -module homomorphism

$$\pi: V(\varpi_2) \to V(\varpi_2) \oplus V(\varpi_0)$$

such that

$$\pi(b_{(a_1,a_2)}) \equiv \begin{cases} b_{()} & \text{if } (a_1,a_2) = (1,2), \\ b_{(a_1,a_2)} & \text{if } (a_1,a_2) \neq (1,2). \end{cases}$$

for all  $1 \le a_1 < a_2 \le 2n$ .

*Proof.* The assertion follows from [Wat23+, Theorem 4.3.1]. Alternatively, one can verify that the vector  $w_0 := b_{(1,2)} - q^{-2}b_{(3,4)} \in V(\varpi_2)$  spans the U<sup>*i*</sup>-submodule isomorphic to  $V(\varpi_0)$  (cf. [Wat23, Lemma 4.1.4]).

## 6.3. Reduction map and successor map.

**Definition 6.3.1.** The *reduction map* is the  $U^i$ -module homomorphism

$$\operatorname{red} = \operatorname{red}_{l} : V(\varpi_{l}) \to \bigoplus_{\substack{0 \le k \le \min(l, 2n-l)\\ l-k \in 2\mathbb{Z}}} V(\varpi_{k})$$

defined inductively as follows.

- (1) If  $l \leq 1$ , then the reduction map is the identity map.
- (2) If l > 1, then the reduction map is the composite

$$\pi \circ \bigwedge \circ (K^{-1} \otimes \mathrm{id}) \circ R \circ (\mathrm{red}_{l-1} \otimes \mathrm{id}) \circ R \circ (K \otimes \mathrm{id}) \circ \bigvee$$

**Proposition 6.3.2.** For each  $\mathbf{a} \in SST_{2n}(\varpi_l)$ , we have

$$\operatorname{red}(b_{\mathbf{a}}) \equiv b_{\operatorname{red}(\mathbf{a})}$$

*Proof.* The assertion follows from Corollary 4.4.2 (1), Proposition 6.1.2, and Lemmas 6.2.3 and 6.2.4.  $\Box$ 

**Definition 6.3.3.** Let  $\lambda = (\lambda_1, \ldots, \lambda_l) \in \operatorname{Par}_{\leq 2n}$ . Set  $\lambda' := (\lambda_1 - 1, \ldots, \lambda_l - 1)$ . The successor map is the U<sup>*i*</sup>-module homomorphism

$$\operatorname{suc}: V(\lambda) \to \bigoplus_{\substack{\lambda' \subseteq \mu \\ \operatorname{vert}}} V(\mu)$$

defined to be the composite

suc := 
$$* \circ (\operatorname{red} \otimes \operatorname{id}) \circ d$$
.

**Proposition 6.3.4.** Let  $\lambda \in Par_{\leq 2n}$  and  $T \in SST_{2n}(\lambda)$ . Then, we have

$$\operatorname{suc}(b_T) \equiv b_{\operatorname{suc}(T)}.$$

 $\square$ 

*Proof.* The assertion follows from Propositions 6.1.2 and 6.3.2.

# 6.4. An orthonormal basis of $V^{i}(\nu)$ .

**Proposition 6.4.1.** Let  $\nu \in \operatorname{Par}_{\leq n}$ . Then, there exists a basis  $\{b_T^i \mid T \in SpT_{2n}(\nu)\}$  of  $V^i(\nu)$  and a  $\mathbf{U}^i$ -module homomorphism

$$p_{\nu}: V(\nu) \to V^{i}(\nu)$$

such that

$$p_{\nu}(b_T) \equiv \begin{cases} b_T^i & \text{if } T \in Sp \mathcal{T}_{2n}(\nu), \\ 0 & \text{if } T \notin Sp \mathcal{T}_{2n}(\nu), \end{cases}$$

for all  $T \in SST_{2n}(\nu)$ .

*Proof.* Let  $p_0: V(\nu) \to V(\nu)$  and  $p_1: V(\nu) \to \bigoplus_{\xi \subset \nu \\ \text{vert}} V(\xi)$  denote the composite of the successor map suc:  $V(\nu) \to \bigoplus_{\xi} V(\xi)$  followed by the projections, respectively. Then, by Corollary 4.3.8, we have

$$p_0(b_T) \equiv \begin{cases} b_T & \text{if } T \in Sp \mathcal{T}_{2n}(\nu), \\ 0 & \text{if } T \notin Sp \mathcal{T}_{2n}(\nu), \end{cases} \quad p_1(b_T) \equiv \begin{cases} 0 & \text{if } T \in Sp \mathcal{T}_{2n}(\nu), \\ b_{\text{suc}(T)} & \text{if } T \notin Sp \mathcal{T}_{2n}(\nu), \end{cases}$$

This implies that  $p_0(V(\nu))$  and  $p_1(V(\nu))$  contain linearly independent subsets  $\{p_0(b_T) \mid T \in SpT_{2n}(\nu)\}$  and  $\{p_1(b_T) \mid T \notin SpT_{2n}(\nu)\}$ , respectively. Since the successor map on  $SST_{2n}(\nu)$  is injective by Corollary 4.4.3, we see that they are bases of the two spaces.

Now, we compute as

$$\operatorname{ch}_{\iota} p_0(V(\nu)) = \sum_{T \in SpT_{2n}(\nu)} \mathbf{y}^{\operatorname{wt}^{Sp}(T)} = \operatorname{ch}_{\iota} V^{\iota}(\nu)$$

This implies  $p_0(\nu) \simeq V^i(\nu)$ . Thus, we complete the proof.

# 6.5. Quantum Littlewood-Richardson map.

**Definition 6.5.1.** The quantum Littlewood-Richardson map is the  $\mathbf{U}^i$ -module homomorphism

$$\mathrm{LR}^{\mathrm{AII}}: V(\lambda) \to \bigoplus_{\substack{\nu \in \mathrm{Par}_{\leq n} \\ \nu \subseteq \lambda}} V^{i}(\nu) \otimes \mathbb{Q}(q) \mathrm{Rec}_{2n}(\lambda/\nu)$$

defined to be the composite of the  $\mathbf{U}^i$ -module homomorphism

$$V(\lambda) \to \bigoplus_{\substack{\nu \in \operatorname{Par}_{\leq n} \\ \nu \subseteq \lambda}} V(\nu) \otimes \mathbb{Q}(q) \operatorname{SST}_{2n}(\lambda)$$

which sends  $b_T$  to  $\operatorname{suc}^k(b_T) \otimes T$  for each  $T \in \operatorname{SST}_{2n}(\lambda)$ , where k > 0 is such that  $\operatorname{suc}^k(S) = P^{\operatorname{AII}}(S)$  for all  $S \in \operatorname{SST}_{2n}(\lambda)$ , and the sum of  $\mathbf{U}^i$ -homomorphisms of the form

$$V(\nu) \otimes \mathbb{Q}(q) \mathrm{SST}_{2n}(\lambda) \to V^{i}(\nu) \otimes \mathbb{Q}(q) \mathrm{Rec}_{2n}(\lambda/\nu)$$

which sends  $b_S \otimes T$  to  $\delta_{S,P^{AII}(T)} p_{\nu}(b_{P^{AII}(T)}) \otimes Q^{AII}(T)$  for each  $S \in SST_{2n}(\nu)$  and  $T \in SST_{2n}(\lambda)$ .

The following is immediate from the definition.

**Proposition 6.5.2.** Let  $T \in SST_{2n}(\lambda)$ . Then, we have

$$\operatorname{LR}^{\operatorname{AII}}(b_T) \equiv b_{P^{\operatorname{AII}}(T)}^i \otimes Q^{\operatorname{AII}}(T).$$

### 7. Surjectivity and codomain of the Littlewood-Richardson map

In this section, we complete the proof of the first assertion of Theorem 3.1.4.

## 7.1. Injectivity and codomain of the Littlewood-Richardson map.

**Proposition 7.1.1.** Let  $\lambda \in \operatorname{Par}_{\leq 2n}$  and  $T \in \operatorname{SST}_{2n}(\lambda)$ . Then, the tableau  $P^{\operatorname{AII}}(T)$  is symplectic. Consequently, we have

$$\operatorname{LR}^{\operatorname{AII}}(T) \in SpT_{2n}(\nu) \times \operatorname{Rec}_{2n}(\lambda/\nu)$$

for some  $\nu \in \operatorname{Par}_{\leq n}$  such that  $\nu \subseteq \lambda$ .

*Proof.* Since  $suc(P^{AII}(T)) = P^{AII}(T)$ , it is symplectic by Corollary 4.3.8.

Let  $\lambda, \mu \in \operatorname{Par}_{\leq 2n}$  be such that  $\mu \subseteq \lambda$ . Define a map

$$\widetilde{\operatorname{suc}} : \operatorname{SST}_{2n}(\mu) \times \operatorname{Tab}(\lambda/\mu) \to \bigsqcup_{\substack{\mu' \in \operatorname{Par}_{\leq 2n} \\ \mu' \subseteq \mu \\ \psi' \subset \mu \\ \operatorname{vert}}} \operatorname{SST}_{2n}(\mu') \times \operatorname{Tab}(\lambda/\mu')$$

as follows. Let  $(S, R) \in SST_{2n}(\mu) \times Tab(\lambda/\mu)$ . Set

$$k := \max\{R(i,j) \mid (i,j) \in D(\lambda/\mu)\},\$$

and  $\mu' := \operatorname{sh}(\operatorname{suc}(S))$ . Then,

$$\widetilde{\operatorname{suc}}(S, R) = (\operatorname{suc}(S), R'),$$

where  $R' \in \operatorname{Tab}(\lambda/\mu')$  is such that

$$R'(i,j) = \begin{cases} R(i,j) & \text{if } (i,j) \notin D(\mu), \\ k+1 & \text{if } (i,j) \in D(\mu), \end{cases} \text{ for all } (i,j) \in D(\lambda/\mu').$$

**Lemma 7.1.2.** Let  $\lambda, \mu \in \operatorname{Par}_{\leq 2n}$  be such that  $\mu \subseteq \lambda$ . Then, the map  $\widetilde{suc}$  on  $\operatorname{SST}_{2n}(\mu) \times \operatorname{Tab}(\lambda/\mu)$  is injective.

*Proof.* Let  $(S_1, R_1), (S_2, R_2) \in SST_{2n}(\mu) \times Tab(\lambda/\mu)$  be such that

$$(\operatorname{suc}(S_1), R'_1) := \operatorname{suc}(S_1, R_1) = \operatorname{suc}(S_2, R_2) = (\operatorname{suc}(S_2), R'_2).$$

Then, for all  $(i, j) \in D(\lambda/\mu)$ , we have

$$R_1(i,j) = R'_1(i,j) = R'_2(i,j) = R_2(i,j).$$

This implies that

$$R_1 = R_2$$

Next, we show that  $S_1 = S_2$ . Let us write

$$d(S_i) = (\mathbf{a}_i, S'_i)$$
 for each  $i = 1, 2$ .

Then, we have

$$\operatorname{red}(\mathbf{a}_1) * S_1' = \operatorname{suc}(S_1) = \operatorname{suc}(S_2) = \operatorname{red}(\mathbf{a}_2) * S_2'$$

Since  $|\operatorname{sh}(S'_1)| = |\operatorname{sh}(S'_2)|$ , we must have

$$|\operatorname{red}(\mathbf{a}_1)| = |\operatorname{red}(\mathbf{a}_2)|.$$

Then, by Proposition 2.3.4, we obtain

$$(\operatorname{red}(\mathbf{a}_1), S_1') = (\operatorname{red}(\mathbf{a}_2), S_2')$$

Now, Corollary 4.4.2 (2) implies

$$\mathbf{a}_1 = \mathbf{a}_2$$

Hence, we deduce

$$S_1 = \mathbf{a}_1 * S_1' = \mathbf{a}_2 * S_2' = S_2,$$

as desired.

**Proposition 7.1.3.** Let  $\lambda \in \operatorname{Par}_{\leq 2n}$ . Then, the Littlewood-Richardson map on  $\operatorname{SST}_{2n}(\lambda)$  is injective.

32

*Proof.* Let  $T_1, T_2 \in SST_{2n}(\lambda)$  be such that  $LR^{AII}(T_1) = LR^{AII}(T_2)$ . Let us write

 $LR^{AII}(T_1) = (P, Q),$ 

and set  $\nu := \operatorname{sh}(P)$ .

First, note that

$$LR^{AII}(T_1) = \widetilde{\operatorname{suc}}^{k_0}(T_1, Q^0)$$

for a sufficiently large  $k_0 > 0$ , where  $Q^0$  denotes the unique tableau of shape  $\lambda/\lambda$ . For each  $k \in [0, k_0]$ , let  $\nu^k$  denote the partition whose Young diagram is

$$D(\nu^k) = D(\nu) \sqcup \{(i,j) \in D(\lambda/\nu) \mid Q(i,j) > k\}$$

Then, we see that

$$\operatorname{sh}(\operatorname{suc}^k(T_1)) = \nu^k \text{ for all } k \in [0, k_0].$$

Since  $LR^{AII}(T_2) = LR^{AII}(T_1) = (P, Q)$ , we have also

$$\operatorname{sh}(\operatorname{suc}^k(T_2)) = \nu^k \text{ for all } k \in [0, k_0].$$

Applying Lemma 7.1.2, we deduce that

$$\operatorname{suc}^{k}(T_{1}) = \operatorname{suc}^{k}(T_{2})$$
 for all  $k \in [0, k_{0}]$ 

by descending induction on k. In particular, we obtain

$$T_1 = \operatorname{suc}^0(T_1) = \operatorname{suc}^0(T_2) = T_2$$

as desired.

## 7.2. Surjectivity of the Littlewood-Richardson map.

**Theorem 7.2.1.** The quantum Littlewood-Richardson map on  $V(\lambda)$  is a  $\mathbf{U}^i$ -module isomorphism.

*Proof.* Let  $Q \in \operatorname{Rec}_{2n}(\lambda/\nu)$ . Then, there exists  $T \in \operatorname{SST}_{2n}(\lambda)$  such that  $Q^{AII}(T) = Q$ . By Proposition 6.5.2, we have

$$\operatorname{LR}^{\operatorname{AII}}(b_T) \equiv b_{P^{\operatorname{AII}}(T)} \otimes Q \neq 0.$$

Since the summand  $V^{i}(\nu) \otimes \mathbb{Q}(q)Q$  is an irreducible  $\mathbf{U}^{i}$ -module, we see that it is contained in the image of LR<sup>AII</sup>. Since the recording tableau Q is arbitrary, the LR<sup>AII</sup> on  $V(\lambda)$  is surjective.

The injectivity follows from that of the Littlewood-Richardson map on  $SST_{2n}(\lambda)$  and Proposition 6.5.2

**Corollary 7.2.2.** The Littlewood-Richardson map on  $SST_{2n}(\lambda)$  is bijective.

## 8. Characterization of the recording tableaux

In this section, we prove the second assertion of Theorem 3.1.4.

33

#### HIDEYA WATANABE

8.1. Some properties of the successor map. Let  $\lambda = (\lambda_1, \ldots, \lambda_l) \in \operatorname{Par}_{\leq 2n}$  and  $T \in \operatorname{SST}_{2n}(\lambda)$ . Let us write  $d(T) = (\mathbf{a}, S)$ ,  $\mathbf{a} = (a_1, \ldots, a_l)$ , and  $\operatorname{red}(\mathbf{a}) = \mathbf{b} = (b_1, \ldots, b_k)$ . For each  $t \in [0, k]$ , set

$$S_t := \begin{cases} S & \text{if } t = 0, \\ b_t \to S_{t-1} & \text{if } t > 0, \end{cases}$$

and  $\lambda^t := \operatorname{sh}(S_t)$ . Let us write

$$br(b_t, S_{t-1}) = (r_{t,1}, r_{t,2}, \dots, r_{t,s_t}).$$

Also, let  $r_{t,0} \in [1, l]$  be such that

$$a_{r_{t,0}} = b_t.$$

Set

$$br_{\leq t} := \{ (r_{u,j}, j) \mid u \in [1, t], \ j \in [1, s_u] \}.$$

Set  $T' := \operatorname{suc}(T)$ ,  $\mu := \operatorname{sh}(T')$ , and  $l' := \ell(\mu)$ .

Define  $\mathbf{a}' = (a'_1, \dots, a'_{l'}), S', \mathbf{b}' = (b'_1, \dots, b'_{k'}), S'_{t'}, \mu^{t'}, (r'_{t',1}, \dots, r'_{t',s'_{t'}})$  for  $t' \in [1, k']$  in the same way as before.

**Lemma 8.1.1.** Let  $t \in [1, k]$  be such that  $r_{t,0} = r_{t,1}$  and  $a'_{r_{t,1}+1} = a'_{r_{t,1}} + 1$ . Then, we have  $a_{r_{t,0}+1} = a_{r_{t,0}} + 1$ .

*Proof.* By Lemma 2.6.3 (2), we have

$$a'_{r_{t,1}} = T'(r_{t,1}, 1) = T(r_{t,0}, 1) = a_{r_{t,0}}$$

First, suppose that  $(r_{t,1} + 1, 1) \notin br_{\leq k}$ . Then, we have

$$T'(r_{t,1}+1,1) = T(r_{t,1}+1,2).$$

Hence, we deduce that

$$a'_{r_{t,1}+1} = T'(r_{t,1}+1,1) = T(r_{t,1}+1,2) \ge T(r_{t,1}+1,1) = a_{r_{t,1}+1}$$

This implies that

$$a_{r_{t,0}+1} = a_{r_{t,1}+1} \le a'_{r_{t,1}+1} = a'_{r_{t,1}} + 1 = a_{r_{t,0}} + 1.$$

Therefore, the assertion follows.

Next, suppose that  $(r_{t,1} + 1, 1) \in br_{\leq k}$ . Let us write  $r_{t,1} + 1 = r_{u,1}$  for some  $u \in [1, k]$ . Then, we have

$$T'(r_{t,1}+1,1) = T'(r_{u,1},1) = T(r_{u,0},1).$$

This implies that

$$a'_{r_{t,1}+1} = T'(r_{t,1}+1,1) = T(r_{u,0},1) = a_{r_{u,0}}$$

Since we have

$$r_{1,1} < r_{2,1} < \dots < r_{k,1}$$

by Lemma 2.6.3 (4), it must hold that u > t. Noting that  $a_{r_{u,0}} > a_{r_{t,0}}$ , we obtain as before that

$$a_{r_{u,0}} = a_{r_{t,0}} + 1.$$

This implies that  $r_{u,0} = r_{t,0} + 1$  since  $(a_1, \ldots, a_k)$  is a strictly increasing sequence. Hence, the assertion follows.

**Lemma 8.1.2.** Let  $t \in [0, k]$ . Then, we have

$$\sharp\{t' \in [1, k'] \mid r'_{t', 0} \le r_{t, 1}\} \ge t,$$

where we set  $r_{0,1} := 0$ . In particular, it holds that  $k' \ge k$ .

*Proof.* For each  $t \in [0, k]$ , set

$$N(t) := \sharp \{ t' \in [1, k'] \mid r'_{t', 0} \le r_{t, 1} \}.$$

Note that we have

$$N(t) = \sharp\{i \in [1, r_{t,1}] \mid a'_i \notin \operatorname{rem}(\mathbf{a}')\} = r_{t,1} - |(a'_1, \dots, a'_{r_{t,1}}) \cap \operatorname{rem}(\mathbf{a}')|$$

Let us prove the assertion by induction on t. When t = 0, the assertion is trivial. Assume that t > 0 and the assertion holds for  $0, 1, \ldots, t - 1$ .

First, suppose that  $a'_{r_{t,1}} \notin \operatorname{rem}(\mathbf{a}')$ . Then, we have

$$N(t) \ge N(t-1) + 1.$$

Hence, we deduce from our induction hypothesis that

$$N(t) \ge N(t-1) + 1 \ge (t-1) + 1 = t,$$

as desired.

Next, suppose that  $a'_{r_{t,1}} \in \text{rem}(\mathbf{a}')$  and  $a'_{r_{t,1}} \notin 2\mathbb{Z}$ . Then, by Proposition 4.2.2, we must have  $r_{t,1} < l, a'_{r_{t,1}+1} = a'_{r_{t,1}} + 1$ , and

$$a'_{r_{t,1}} < 2r_{t,1} - |\operatorname{rem}(a'_1, \dots, a'_{r_{t,1}-1})|.$$

Also, by Corollary 4.2.8, we have

$$|\operatorname{rem}(a'_1,\ldots,a'_{r_{t,1}-1})| \ge 2r_{t,1} - a'_{r_{t,1}} - 1.$$

Hence, we obtain

$$|\operatorname{rem}(a'_1,\ldots,a'_{r_{t,1}-1})| = 2r_{t,1} - a'_{r_{t,1}} - 1.$$

Therefore,

$$N(t) = r_{t,1} - (|\operatorname{rem}(a'_1, \dots, a'_{r_{t,1}-1})| + 1) = a'_{r_{t,1}} - r_{t,1}.$$

On the other hand, by Corollary 4.2.8 again, we have

$$|\operatorname{rem}(a_1,\ldots,a_{r_{t,0}-1})| \ge 2r_{t,0} - a_{r_{t,0}} - 1.$$

Also, since  $a_{r_{t,0}} \notin \text{rem}(\mathbf{a})$ , it holds that

$$r_{t,0} - |\operatorname{rem}(a_1, \dots, a_{r_{t,0}-1})| = t.$$

Hence,

$$a_{r_{t,0}} - r_{t,0} \ge t - 1.$$

Combining above, we obtain

$$N(t) = a'_{r_{t,1}} - r_{t,0} + (r_{t,0} - r_{t,1}) \ge t + (r_{t,0} - r_{t,1}) - 1$$

Therefore, the assertion follows when  $r_{t,0} > r_{t,1}$ . Hence, we only need to consider the case when  $r_{t,0} = r_{t,1}$ . In this case, Lemma 8.1.1 implies

$$a_{r_{t,0}+1} = a_{r_{t,0}} + 1.$$

Since  $a_{r_{t,0}} \notin \text{rem}(\mathbf{a})$ , we have

$$a_{r_{t,0}} \ge 2r_{t,0} - |\operatorname{rem}(a_1, \dots, a_{r_{t,0}-1})| = r_{t,0} + t.$$

Now, we deduce

$$N(t) = a'_{r_{t,1}} - r_{t,1} = a_{r_{t,0}} - r_{t,0} \ge t,$$

as desired.

Finally, suppose that  $a'_{r_{t,1}} \in \text{rem}(\mathbf{a}')$  and  $a'_{r_{t,1}} \in 2\mathbb{Z}$ . In this case, one can deduce the assertion in a similar way to the previous case.

Thus, we complete the proof.

#### HIDEYA WATANABE

**Lemma 8.1.3.** Let  $t \in [0, k]$ . Then, we have  $s'_t \ge s_t$  and  $r'_{t,j-1} \le r_{t,j}$  for all  $j \in [1, s_t]$ , where we set  $s_0 = s'_0 = 0$ .

*Proof.* We proceed by induction on t. The case when t = 0 is trivial. Hence, assume that t > 0 and the assertion holds for  $0, 1, \ldots, t - 1$ .

We prove the assertion by induction on j. By lemma 8.1.2 and the fact that  $r'_{1,0} < r'_{2,0} < \cdots < r'_{k',0}$ , we obtain

$$r_{t,0}' \le r_{t,1}.$$

Now, assume that j > 1 and we have  $s'_t, s_t \ge j-1$  and  $r'_{t,j-2} \le r_{t,j-1}$ . When  $s_t = j-1$ , there is nothing to prove. Hence, assume that  $s_t \ge j$ .

By Lemma 2.6.3, we have

$$r'_{t,j-1} = \min\{r' \in [1, \operatorname{col}_{j-1}(\mu^{t-1}) + 1] \mid T'(r', j) \ge T'(r'_{t,j-2}, j-1)\}.$$

Note that our induction hypothesis implies that

$$r'_{t-1,j-1} \le r_{t-1,j}$$

The right-hand side is less than  $r_{t,j}$  by Lemma 2.6.3 (4). Also we have

$$r_{t,j} \le \operatorname{col}_j(\lambda^{t-1}) + 1$$

and

$$\operatorname{col}_{j}(\lambda^{t-1}) \le \operatorname{col}_{j}(\lambda^{k}) = \operatorname{col}_{j}(\mu^{0}) \le \operatorname{col}_{j-1}(\mu^{0}) \le \operatorname{col}_{j-1}(\mu^{t-1})$$

Observe that

$$\begin{aligned} T'(r_{t,j},j) &= T(r_{t,j-1},j), \\ T'(r'_{t,j-2},j-1) &= \begin{cases} T(r_{u,j-2},j-1) & \text{if } r'_{t,j-2} = r_{u,j-1} \text{ for some } u \in [1,k], \\ T(r'_{t,j-2},j) & \text{otherwise,} \end{cases} \end{aligned}$$

and

$$T(r_{u,j-2}, j-1) \le T(r_{u,j-1}, j) \le T(r_{t,j-1}, j),$$
  
$$T(r'_{t,j-2}, j) \le T(r_{t,j-1}, j).$$

By above, we obtain  $r'_{t,j-1} \leq r_{t,j}$  and  $s'_t \geq j$ , as desired.

**Proposition 8.1.4.** Let  $r \in [0, l]$ . Then, we have

*Proof.* For each  $t \in [1, k]$ , we have  $s'_t \ge s_t$  and  $r'_{t,s_t-1} \le r_{t,s_t}$ . Hence, if  $r_{t,s_t} \le r$ , then

$$r'_{t,s'_t} \le r'_{t,s_t-1} \le r_{t,s_t} \le r.$$

Thus, the assertion follows.

### 8.2. Symplectic Littlewood-Richardson tableaux.

**Definition 8.2.1.** Let  $\lambda \in \operatorname{Par}_{\leq 2n}$  and  $\nu \in \operatorname{Par}_{\leq n}$ . A tableau  $T \in \operatorname{Tab}(\lambda/\nu)$  is said to be a symplectic Littlewood-Richardson tableau if it satisfies the following.

- (1)  $T \in SST_{2n}(\lambda/\nu)$ .
- (2) Let  $(w_1, \ldots, w_N)$  denote the column-word  $w^{\text{col}}(T)$  of T. Then, the reversed word  $(w_N, \ldots, w_1)$  is a lattice permutation: For each  $r \in [1, N]$  and  $k \in [1, 2n 1]$ , the number of occurrences of k in the subsequence  $(w_N, \ldots, w_r)$  is greater than or equal to that of k + 1.

- (3) The sequence wt(T) = (T[1], T[2],...,T[2n]) is a partition which has even columns (see Definition 2.1.1).
- (4) If T(i,j) = 2k + 1 for some  $(i,j) \in D(\lambda/\nu)$  and  $k \in \mathbb{Z}_{\geq 0}$ , then we have  $i \leq n + k$ .

Let  $LRT_{2n}^{Sp}(\lambda/\nu)$  denote the set of all symplectic Littlewood-Richardson tableaux of shape  $\lambda/\nu$ .

**Remark 8.2.2.** Let  $T \in SST_{2n}(\lambda/\nu)$  be such that the sequence wt(T) is a partition which has even columns. Then, we have  $T \in LRT_{2n}^{Sp}(\lambda/\nu)$  if and only if the reversed column-word fits  $\lambda/\nu$  n-symplectically in the sense of [Sun90, Definition 3.9].

The symplectic Littlewood-Richardson tableaux provide us a branching rule:

**Theorem 8.2.3** (cf. [Sun90, Corollary 3.12]). Let  $\lambda \in \operatorname{Par}_{\leq 2n}$  and  $\nu \in \operatorname{Par}_{\leq n}$  be such that  $\nu \subseteq \lambda$ . Then, we have

$$m_{\lambda,\nu} = |\mathrm{LRT}_{2n}^{Sp}(\lambda/\nu)|.$$

8.3. Characterization of the recording tableaux. Let  $\lambda \in \operatorname{Par}_{\leq 2n}$ ,  $\nu \in \operatorname{Par}_{\leq n}$  be such that  $\nu \subseteq \lambda$ .

Lemma 8.3.1. We have

$$\operatorname{Rec}_{2n}(\lambda/\nu) \subseteq \operatorname{Rec}_{2n}(\lambda/\nu).$$

*Proof.* Let  $Q \in \operatorname{Rec}_{2n}(\lambda/\nu)$ . We only need to show that  $Q \in \operatorname{Rec}_{2n}(\lambda/\nu)$ , that is, to verify that the tableau Q satisfies conditions (R1)–(R5) in Section 3.

By the definition of recording tableaux, there exists  $T \in SST_{2n}(\lambda)$  such that  $Q^{AII}(T) = Q$ . For each  $k \in \mathbb{Z}_{\geq 0}$ , set

$$P^{k} := \begin{cases} T & \text{if } k = 0, \\ \operatorname{suc}(P^{k-1}) & \text{if } k > 0, \end{cases} \quad \nu^{k} := \operatorname{sh}(P^{k}).$$

Let us write

$$d(P^k) = (\mathbf{a}^k, S^k), \ \mathbf{a}^k = (a_1^k, \dots, a_{l_k}^k), \ \operatorname{red}(\mathbf{a}^k) = \mathbf{b}^k = (b_1^k, \dots, b_{l'_k}^k),$$

and

$$\operatorname{br}(b_t^k, b_{t-1}^k \to (\dots \to (b_1^k \to S^{k-1}))) = (r_{t,1}^k, r_{t,2}^k, \dots, r_{t,s_t^k}^k)$$

First, let us verify conditions (R1) and (R2). Let  $(i, j) \in D(\lambda/\nu)$  and write Q(i, j) = k. Suppose that  $(i, j + 1) \in D(\lambda/\nu)$  (resp.,  $(i + 1, j) \in D(\lambda/\nu)$ ). Then, since  $\nu^k \subseteq \nu^{k-1}$ , we must have  $(i, j + 1) \notin D(\nu^{k-1})$  (resp.,  $(i + 1, j) \notin D(\nu^k)$ ). This, together with Lemma 3.1.2, implies that  $Q(i, j + 1) \leq k - 1$  (resp.,  $Q(i + 1, j) \leq k$ ), as desired.

Next, let us verify conditions (R3) and (R4). Let k > 0. Then, we have

$$Q[k] = |\nu^{k-1}/\nu^k|$$

by Lemma 3.1.2. Since  $P^k = \operatorname{suc}(P^{k-1}) = \operatorname{red}(\mathbf{a}^{k-1}) * S^{k-1}$ , it holds that

$$\nu^{k-1}/\nu^k| = |\operatorname{rem}(\mathbf{a}^{k-1})|.$$

The right-hand side is even by Lemma 4.2.3, and is greater than or equal to  $2(\ell(\nu^{k-1}) - n)$  by Proposition 4.2.7.

Finally, let us verify condition (R5). Let r, k > 0. Then, we have

$$Q_{\leq r}[k] = \sharp\{i \in [1, r] \mid Q(i, j) = k \text{ for some } j\}$$
$$= |[1, r] \setminus \{r_{t, s_t^{k-1}}^{k-1} \mid t \in [1, l'_{k-1}]\}|$$
$$= r - \sharp\{t \in [1, l'_{k-1}] \mid r_{t, s_t^{k-1}}^{k-1} \leq r\}.$$

Hence, we deduce that

$$Q_{\leq r}[k] - Q_{\leq r}[k+1] = \{t' \in [1, l'_k] \mid r^k_{t', s^k_{t'}} \leq r\} - \{t \in [1, l'_{k-1}] \mid r^{k-1}_{t, s^{k-1}_t} \leq r\}.$$

The right-hand side is nonnegative by Proposition 8.1.4.

Thus, we complete the proof.

Lemma 8.3.2. Consider the map

$$\operatorname{Rec}_{2n}(\lambda/\nu) \to \operatorname{Tab}(\lambda/\nu)$$

which sends each  $R \in \operatorname{Rec}_{2n}(\lambda/\nu)$  to the tableaux of shape  $\lambda/\nu$  whose (i, j) entry is equal to  $R_{\leq i}[R(i, j)]$ ; the number of occurrences of R(i, j) in R in the *i*-th row or above. Then, it is injective and its image is contained in  $\operatorname{LRT}_{2n}^{Sp}(\lambda/\nu)$ .

*Proof.* Let  $R \in \operatorname{Rec}_{2n}(\lambda/\nu)$ , and  $S \in \operatorname{Tab}(\lambda/\nu)$  be the image of R under the map above. We only need to show that the tableau S satisfies the conditions (1)–(4) in Definition 8.2.1.

First, let us verify condition (1). By condition (R1), each k' > 0 appears as an entry of R at most once in each row. In particular, we have

$$R[k'] \le \ell(\lambda) \le 2n$$
 for all  $k' > 0$ .

This implies that the entries of S are in [1, 2n].

Let  $(i, j) \in D(\lambda/\nu)$  and set k := R(i, j). Suppose that  $(i, j + 1) \in D(\lambda/\nu)$  (resp.,  $(i+1, j) \in D(\lambda/\nu)$ ), and set k' := R(i, j+1) < k (resp.,  $k'' := R(i+1, j) \le k$ ). Here, we used condition (R1) (resp., (R2)). Then, we have

$$S(i, j+1) = R_{\leq i}[k'] \geq R_{\leq i}[k] = S(i, j),$$

(resp.,

$$S(i+1,j) = R_{\leq i+1}[k''] = R_{\leq i}[r''] + 1 \ge R_{\leq i}[r] + 1 = S(i,j) + 1,$$

where the inequality follows from condition (R5). This implies that S is semistandard.

Second, let us verify condition (2). Let us write

$$w_{\rm col}(S) = (w_1, \dots, w_N)$$
 and  $w_{\rm col}(R) = (w'_1, \dots, w'_N).$ 

Let  $r \in [1, N]$  and  $k \in [1, 2n]$ . Then, we have

$$\sharp\{t \in [r, N] \mid w_t = k\} = \sharp\{k' > 0 \mid \sharp\{t' \in [r, N] \mid w_{t'}' = k'\} \ge k\}.$$

The right-hand side decreases as k increases. Hence, the assertion follows.

Next, let us verify condition (3). By condition (R5), we have

$$R[1] \ge R[2] \ge \cdots$$

Then, we can consider the tableau whose *j*-th column consists of exactly R[j] *j*'s. Clearly, its shape is wt(S). Noting that  $R[j] \in 2\mathbb{Z}$  by condition (R3), we see that the partition wt(S) has even columns.

Finally, let us verify condition (4). Let  $(i, j) \in D(\lambda/\nu)$  and suppose that S(i, j) = 2k+1 for some  $k \ge 0$ . When  $i \le n$ , there is nothing to prove. Hence, assume that i > n.

Set r := R(i, j). Then, we have  $R_{\leq i}[r] = 2k + 1$ . Let  $\mu$  be the partition whose Young diagram is given by

$$D(\mu) = D(\nu) \sqcup \{(i,j) \in D(\lambda/\nu) \mid R(i,j) \ge r\}.$$

We have  $R_{\leq i}[r] \geq 2(i-n)$ . Otherwise, we obtain

$$R[r] = R_{\leq \ell(\mu)}[r] \leq R_{\leq i}[r] + (\ell(\mu) - i) < 2(i - n) + (\ell(\mu) - i) \leq 2(\ell(\mu) - i)$$

which contradicts condition (R4).

Now, we have

$$2k + 1 = R_{\le i}[r] \ge 2(i - n).$$

This implies that  $2k \ge 2(i-n)$ , and then the assertion follows.

Thus, we complete the proof.

**Theorem 8.3.3.** We have 
$$\operatorname{Rec}_{2n}(\lambda/\nu) = \operatorname{Rec}_{2n}(\lambda/\nu)$$
.

*Proof.* By Lemmas 8.3.1 and 8.3.2, we have

$$\operatorname{Rec}_{2n}(\lambda/\nu) \subseteq \operatorname{Rec}_{2n}(\lambda/\nu)$$

and

(8.1)

$$|\operatorname{Rec}_{2n}(\lambda/\nu)| \le |\widetilde{\operatorname{Rec}}_{2n}(\lambda/\nu)| \le |\operatorname{LRT}_{2n}^{Sp}(\lambda/\nu)|$$

Hence, we only need to show that

$$|\operatorname{Rec}_{2n}(\lambda/\nu)| = |\operatorname{LRT}_{2n}^{Sp}(\lambda/\nu)|.$$

Recall from Theorem 8.2.3 that

$$|\operatorname{LRT}_{2n}^{Sp}(\lambda/\nu)| = m_{\lambda,\nu}.$$

On the other hand, by Theorem 7.2.1 and Proposition 6.2.2, we obtain

$$|\operatorname{Rec}_{2n}(\lambda/\nu)| = m_{\lambda,\nu}.$$

Therefore, equation (8.1) holds.

Thus, we complete the proof.

## References

- [Ber86] A. Berele, A Schensted-type correspondence for the symplectic group, J. Combin. Theory Ser. A 43 (1986), no. 2, 320–328.
- [Ful97] W. Fulton, Young Tableaux. With Applications to Representation Theory and Geometry, London Mathematical Society Student Texts, 35. Cambridge University Press, Cambridge, 1997. x+260 pp.
- [HTW05] R. E. Howe, E. Tan, and J. F. Willenbring, Stable branching rules for classical symmetric pairs, Trans. Amer. Math. Soc. 357 (2005), no. 4, 1601–1626.
- [Kin76] R. C. King, Weight multiplicities for the classical groups, Group theoretical methods in physics (Fourth Internat. Colloq., Nijmegen, 1975), pp. 490–499. Lecture Notes in Phys., Vol. 50, Springer, Berlin, 1976.
- [Kwo09] J. Kwon, Crystal graphs and the combinatorics of Young tableaux, Handbook of algebra. Vol. 6, 473–504, Handb. Algebr., 6, Elsevier/North-Holland, Amsterdam, 2009.
- [Lit40] D. E. Littlewood, The Theory of Group Characters and Matrix Representations of Groups, Reprint of the second (1950) edition. AMS Chelsea Publishing, Providence, RI, 2006. viii+314 pp.
- [Lus93] G. Lusztig, Introduction to Quantum Groups, Reprint of the 1994 edition. Modern Birkhäuser Classics. Birkhäuser/Springer, New York, 2010. xiv+346 pp.
- [Mol06] A. I. Molev, Representations of the twisted quantized enveloping algebra of type  $C_n$ , Mosc. Math. J. 6 (2006), no. 3, 531–551, 588.

#### HIDEYA WATANABE

- [NaSa05] S. Naito and D. Sagaki, An approach to the branching rule from  $\mathfrak{sl}_{2n}(\mathbb{C})$  to  $\mathfrak{sp}_{2n}(\mathbb{C})$  via Littelmann's path model, J. Algebra 286 (2005), no. 1, 187–212.
- [NaYa97] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, Selecta Math. (N.S.) 3 (1997), no. 4, 547–599.
- [ScTo18] B. Schumann and J. Torres, A non-Levi branching rule in terms of Littelmann paths, Proc. Lond. Math. Soc. (3) 117 (2018), no. 5, 1077–1100.
- [Sun90] S. Sundaram, Tableaux in the representation theory of the classical Lie groups, Invariant theory and tableaux (Minneapolis, MN, 1988), 191–225, IMA Vol. Math. Appl., 19, Springer, New York, 1990.
- [Wan21] W. Wang, Quantum symmetric pairs, arXiv:2112.10911.
- [Wat21] H. Watanabe, Classical weight modules over *i*quantum groups, J. Algebra 578 (2021), 241–302.
- [Wat21+] H. Watanabe, Crystal bases of modified *i*quantum groups of certain quasi-split types, to appear in Algebr. Represent. Theory.
- [Wat23] H. Watanabe, Stability of *i*canonical bases of irreducible finite type of real rank one, Represent. Theory 27 (2023), 1–29.
- [Wat23+] H. Watanabe, Stability of *i*canonical bases of locally finite type, arXiv:2306.12199.

(H. WATANABE) OSAKA CENTRAL ADVANCED MATHEMATICAL INSTITUTE, OSAKA METROPOLITAN UNIVERSITY, OSAKA, 558-8585, JAPAN

Email address: watanabehideya@gmail.com