A DEFORMATION OF BOREL EQUIVARIANT HOMOTOPY

GABRIEL ANGELINI-KNOLL, MARK BEHRENS, EVA BELMONT, AND HANA JIA KONG

ABSTRACT. We describe a deformation of the ∞ -category of Borel *G*-spectra for a finite group *G*. This provides a new presentation of the *a*-complete real Artin–Tate motivic stable homotopy category when $G = C_2$ and gives a new interpretation of the *a*-completed C_2 -effective slice spectral sequence. As a new computational tool, we present a modified Adams–Novikov spectral sequence which computes the RO(G)-graded Mackey functor valued homotopy of Borel *G*-spectra.

Contents

1. Introduction	2
1.1. Overview	2
1.2. The main constructions	3
1.3. Computing the MANSS	4
1.4. Future work	6
1.5. Organization of the paper	6
1.6. Conventions and notation	7
1.7. Acknowledgements	8
2. Filtered objects and deformations	8
2.1. Filtered objects	8
2.2. Tower functors	11
2.3. Décalage	13
2.4. Deformations	13
3. A tale of two spectral sequences	14
3.1. The C_2 -effective slice spectral sequence	15
3.2. The modified <i>E</i> -based Adams spectral sequence	16
3.3. The E -MASS as a generalized Adams spectral sequence	18
3.4. The MANSS and the C_2 -effective slice spectral sequence	20
4. Computing with the MANSS	22
4.1. The E_2 -term of the MANSS	23
4.2. The case of $MU_{\mathbb{R}}$ -projective spectra	25
4.3. The case of a trivial C_2 -action	29
4.4. The C-motivic MANSS and the algebraic AHSS	31
5. Applications	38
5.1. Example: ko_{C_2} at $p = 2$	38
5.2. Example: $tmf(2)$ at $p = 3$	42
References	48

Date: September 29, 2023.

1. INTRODUCTION

1.1. **Overview.** Motivic homotopy theory, developed by Voevodsky and Morel (eg. [MV99]), was famously used as a tool to attack the Milnor conjecture and the Bloch–Kato conjecture, as well as other deep problems in number theory and algebraic geometry. It became useful in stable homotopy theory for a different reason: the "Betti realization" functor from the \mathbb{C} -motivic homotopy category to the stable homotopy category makes computations of stable homotopy groups of spheres more accessible [IWX23]. The \mathbb{R} -motivic homotopy category is also equipped with a Betti realization functor to the C_2 -equivariant stable homotopy category; work in progress by Guillou and Isaksen uses this to compute C_2 -equivariant stable homotopy groups. It is natural to ask whether there is a related category that supports an analogous Betti realization functor to G-equivariant spectra for other finite groups G. However, we do not expect such a comparison category to come from motivic homotopy theory; the C_2 case works because \mathbb{C} is an extension of \mathbb{R} with Galois group C_2 , and the Artin–Schreier theorem implies that e and C_2 are the only finite groups that can occur as absolute Galois groups.

Instead, we turn to another perspective by building the desired category as a deformation of the equivariant category. Work of Gheorghe–Wang–Xu [GWX21] (see also [GIKR22], [Pst23]) provided a new characterization of the category of (*p*-completed, cellular) \mathbb{C} -motivic spectra as a *deformation* (see Definition 2.4.2) of (*p*-completed) ordinary spectra. The Tate twist element τ (see [HKO11, Lem. 23]) provides a deformation parameter, and Betti realization can be described as inverting τ . Gheorghe–Isaksen–Krause–Ricka [GIKR22] provided a model for this deformation using the category of modules over a certain algebra in filtered spectra coming from the Adams–Novikov filtration.

The filtered spectra model can be further generalized: given a well-behaved filtration functor (see Definition 2.2.1) in some appropriate category \mathcal{C} , the deformation perspective provides an appropriate new category \mathcal{D} equipped with a comparison functor $\mathcal{D} \to \mathcal{C}$, analogous to the Betti realization functor. The construction comes with a "deformation parameter" τ and the Betti realization functor is given by inverting τ . Using this perspective, Burklund– Hahn–Senger [BHS20] showed that a variant of the cellular *p*-complete \mathbb{R} -motivic category (the Artin–Tate category) arises from a filtration functor on C_2 -equivariant spectra. However, this is not amenable to generalization to an arbitrary finite group G, because this filtration hinges on the existence of the C_2 -spectrum $MU_{\mathbb{R}}$ known as Real complex cobordism. No analogue of $MU_{\mathbb{R}}$ for other finite groups G has been found despite over a decade of searching (see [HHR11, HSW23]).

In this paper, we overcome this difficulty in the setting of Borel G-spectra for a finite group G. We give an alternate construction of the filtration that doesn't involve $MU_{\mathbb{R}}$ and can be generalized to other groups. Our paper gives two interpretations of this filtration:

- as a generalized Adams filtration in Borel G-spectra with an appropriate base (Definition 2.4.6), or
- as a modified Adams–Novikov filtration which mixes the (non-equivariant) Adams– Novikov filtration and the homotopy fixed points filtration (see Section 3.2).

We show these are equivalent (Proposition 3.3.3) and show that the $G = C_2$ case of our construction recovers the *a*-completion of the motivic category studied in [BHS20] (Theorem 3.4.9). The second construction produces a spectral sequence, which we call the *modified Adams-Novikov spectral sequence* (MANSS). We demonstrate that it is amenable to computations in Section 5.

In the case of $G = C_2$, our construction provides a new description of a completed form of the C_2 -effective slice spectral sequence introduced by the fourth author in [Kon20]. The C_2 -effective slice spectral sequence is defined using Betti realization and Voevodsky's effective slice filtration on the \mathbb{R} -motivic homotopy category [Voe02]. It agrees with the Hill–Hopkins– Ravenel slice spectral sequence [HHR16] for Real orientable C_2 -spectra, and in the non-Real orientable case it is more tractable than the slice spectral sequence. Burklund–Hahn–Senger [BHS20] have shown that the C_2 -effective slice spectral sequence agrees with the décalage (a degree-shifting operation) of the $MU_{\mathbb{R}}$ -based Adams–Novikov spectral sequence, and hence (after completion) with the décalage of the MANSS. Moreover, for other groups G, the MANSS can be viewed as a Borel G-equivariant analogue of the C_2 -effective spectral sequence up to décalage.

1.2. The main constructions. We will use the presentations expressed in the language of filtered spectra by [GIKR22] and [BHS20]. A filtered spectrum $X_{\bullet} \in \mathcal{C}^{\mathbb{Z}}$ is simply a \mathbb{Z} -indexed sequence of objects X_i and maps

$$\cdots \leftarrow X_{-2} \leftarrow X_{-1} \leftarrow X_0 \leftarrow X_1 \leftarrow X_2 \leftarrow \cdots$$

in a category C. We regard this sequence as a decreasing filtration of $X_{-\infty} \coloneqq \operatorname{colim}_i X_{-i}$. There is a symmetric monoidal structure on filtered objects given by *Day convolution* (2.1.4).

Given a filtered object X_{\bullet} and a Postnikov truncation functor $\tau_{\geq \bullet}$, one can apply a degreeshifting operation called *décalage* (see §2.3) to form another filtered object $\text{Déc}^{\tau}(X)_{\bullet}$. The central idea, as introduced in [GIKR22], is to study the category of modules over Déc(X):

(*)
$$\operatorname{Mod}(\mathcal{C}^{\mathbb{Z}},\operatorname{Déc}^{\tau}(X)).$$

The filtrations relevant to this paper are

- (1) the Adams-Novikov filtration $\Gamma_{MU_p} = \overline{MU_p}^{\otimes \bullet} \otimes X$ of a spectrum X (where $\overline{(-)}$ denotes the fiber of the unit map);
- (2) the Real Adams–Novikov filtration $\Gamma_{\mathbb{R}} = \overline{MU_{\mathbb{R},2}}^{\otimes \bullet} \otimes X$ of a C_2 -spectrum X;
- (3) the MU[G]-based Adams filtration $\Gamma_{MU_p[G]} = \overline{MU_p \wedge G_+}^{\otimes \bullet} \otimes X$ of a (Borel) G-spectrum X;
- (4) the modified Adams–Novikov filtration, defined to be the Day convolution in Borel G-spectra of
 - (a) the homotopy fixed points filtration of a Borel G-spectrum $X^h = F(EG_+, X)$, given by filtering EG by skeleta, and
 - (b) the (nonequivariant) Adams–Novikov filtration of the sphere.

Filtrations (1) and (2) pertain to the motivating results; (3) and (4) are our constructions.

Theorem 1.2.1 ([GIKR22]). Applying the construction (*) to the Adams–Novikov filtration with τ as the Postnikov filtration recovers the cellular \mathbb{C} -motivic category $\mathrm{SH}^{\mathrm{cell}}(\mathbb{C})_{i2}$.

Here and henceforth, we use $(-)_{ip}$ to denote the category of modules over the *p*-complete unit.

Theorem 1.2.2 ([BHS20]). Applying the construction (*) to the Real Adams–Novikov filtration with τ as even slice décalage (Notation 2.3.2) gives the Artin–Tate category $\text{SH}(\mathbb{R})_{i2}^{AT}$, generated by bigraded suspensions $\Sigma^{i,j} \text{Spec}(k)_+$, where $k \in \{\mathbb{R}, \mathbb{C}\}$.

Note that $\text{Spec}(\mathbb{R})_+$ is the sphere spectrum, so this is analogous to the cellular category but with one more generator.

Our main construction is the following.

Definition (Definition 2.4.6). Define the category of *complete artificial motivic spectra* $\widehat{\operatorname{SH}}_{G,p}^{\operatorname{art}}$ to be the construction (*) applied to the MU[G]-based Adams filtration, with τ as the even Postnikov filtration.

Our main results are the following three theorems, which justify the definition above by relating it to the construction by [BHS20] in the case $G = C_2$. The third theorem says that that (3) has the same spectral sequence as (4).

Theorem (Theorem 3.4.7). For $X \in \text{Sp}_{C_2,i^2}$, there is an equivalence

$$\operatorname{D\acute{e}c}^{\operatorname{ev.post}}\Gamma_{MU_2[C_2]} \simeq (\Gamma_{\mathbb{R}})'$$

of filtered Borel C_2 -spectra, where $(-)^h = F((EC_2)_+, -)$ denotes Borel completion and Déc^{ev.post} denotes the even Postnikov décalage.

Theorem (Theorem 3.4.9). There is an equivalence

$$\widehat{\operatorname{SH}}_{C_2,2}^{\operatorname{art}} \simeq (\operatorname{SH}(\mathbb{R})_{i2}^{\operatorname{AT}})_a^{\wedge}$$

where the completion of $\text{SH}(\mathbb{R})_{i2}^{\text{AT}}$ is with respect to the motivic lift *a* of the equivariant Euler class a_{σ} defined in [BHS20, Eq. (1)].

Theorem (Theorems 3.4.2). For $G = C_2$, the spectral sequence associated to the modified Adams–Novikov filtration agrees with the $MU_p[C_2]$ -based Adams spectral sequence from the E_2 -term onwards. In particular, the construction (*) applied to the modified Adams–Novikov filtration also recovers $(SH(\mathbb{R})_{i2}^{AT})_a^{\wedge}$.

While the comparison theorems use $\Gamma_{MU_2[C_2]}$, we are interested in the modified Adams– Novikov filtration because it is amenable to computation.

1.3. Computing the MANSS. We study the modified Adams–Novikov spectral sequence (MANSS), the spectral sequence associated to the modified Adams–Novikov filtration as discussed in the previous section. Computing this spectral sequence is relevant because, in many cases such as ours, computing the homotopy groups of a construction of the form (*) boils down to computing the spectral sequence associated to the filtration in question; see [BHS22, Theorem A.1].

The MANSS is an RO(G)-graded spectral sequence which is a kind of mixture of the (non-equivariant) Adams–Novikov spectral sequence (ANSS) and the homotopy fixed point spectral sequence (HFPSS), and as such is very amenable to computations.

We introduce the following tools for computing with the MANSS:

(1) In Section 4.1, we introduce a pair of double complex spectral sequences which converge to the E_2 -term of the *p*-primary MANSS:

$${}_{I}\underline{\underline{E}}_{2}^{*,*,\star}(G/H) = \operatorname{Ext}_{(MU_{p})_{*}MU_{p}}^{*,*}(MU_{*}, H^{*}(H; (MU_{p})_{*}\Sigma^{-\star}X)) \Longrightarrow \overset{\operatorname{MANSS}}{\longrightarrow} \underline{\underline{E}}_{2}^{*,*}(X)(G/H),$$
$${}_{II}\underline{\underline{E}}_{2}^{*,*,\star}(G/H) = H^{*}(H; \operatorname{Ext}_{(MU_{p})_{*}MU_{p}}^{*,*}((MU_{p})_{*}, (MU_{p})_{*}\Sigma^{-\star}X)) \Longrightarrow \overset{\operatorname{MANSS}}{\longrightarrow} \underline{\underline{E}}_{2}^{*,*}(X)(G/H).$$

- (2) We show these double complex spectral sequences collapse in the case where X is the Borel completion of a genuine C_2 -spectrum which is $MU_{\mathbb{R}}$ -projective (i.e. its $MU_{\mathbb{R}}$ -homology is free on a set of generators in degrees which are multiples of ρ). We give a complete computation of the E_2 -term of the 2-primary MANSS for such spectra in terms of the ANSS E_2 -term of the underlying non-equivariant spectrum X^e , in a manner that mirrors the computation of the E_2 -term of the C_2 -effective slice spectral sequence for $MU_{\mathbb{R}}$ -projective spectra [BHS20, Thm. 5.6].
- (3) In Section 4.3 we show that these double complex spectral sequences also collapse in the case where X is a Borel C_2 -spectrum with trivial C_2 -action and torsion-free MU-homology. In this case, the MANSS converges to

$$\pi_{i+j\sigma}^{C_2} F((EC_2)_+, X) \cong \pi_i(F(P_j^\infty, X)),$$

and we give a complete computation of the MANSS E_2 -term of such spectra in terms of the ANSS E_2 -term of X, which is different from (2), where the generators are now parameterized by the cells of the associated stunted projective spectra.

For X as in (3) with MU-homology concentrated in even degrees, we introduce a complex motivic lift of the MANSS (the *motivic* MANSS)

$${}^{\mathrm{MANSS}}_{\mathrm{mot}} E^{*,\star,*}_r(\nu_{\mathbb{C}} X)$$

in which X is replaced with the motivic lift $\nu_{\mathbb{C}}(X) \in SH(\mathbb{C})_{i2}$ and the stunted projective spaces P_j^{∞} are replaced with the motivic lifts $\nu_{\mathbb{C}}(P_j^{\infty})$. We have

$${}^{\text{MANSS}}_{\text{mot}} E_r^{*,\star}(X) \cong {}^{\text{MANSS}}_{\text{mot}} E_r^{*,\star,*}(\nu_{\mathbb{C}} X)[\tau^{-1}].$$

We prove the following theorem.

Theorem (Theorem 4.4.16). The motivic MANSS

$${}^{\mathrm{MANSS}}_{\mathrm{mot}} E^{*,*+j\sigma,*}_{r}(\nu_{\mathbb{C}}(X) \otimes C\tau)$$

is isomorphic (after reindexing) to the accelerated algebraic Atiyah–Hirzebruch spectral sequence (AHSS)

$$\underset{\text{acc,alg}}{^{\text{AHSS}}} E_1^{k,s,t}(X^{P_j^{\infty}}) = \text{Ext}_{MU_*MU}^{s,t}(MU_*, MU_*(X^{P_j^{2k}/P_j^{2k-2}})) \implies \lim_m \text{Ext}_{MU_*MU}^{s,t}(MU_*, MU_*(X^{P_j^{2m}}))$$

obtained from filtering P_i^{∞} by its even skeleta.¹

The significance of this theorem is that the differentials in the algebraic Atiyah–Hirzebruch spectral sequence are computable in terms of the attaching maps of P_j^{∞} . We can then employ the zig-zag of spectral sequences

$$(1.3.1)$$

$$\stackrel{\text{MANSS}}{\text{mot}} E_{2r+1}^{*,*+j\sigma,*}(\nu_{\mathbb{C}}(X) \otimes C\tau) \longleftarrow \stackrel{\text{MANSS}}{\text{mot}} E_{2r+1}^{*,*+j\sigma,*}(\nu_{\mathbb{C}}(X)) \longrightarrow \stackrel{\text{MANSS}}{\text{mot}} E_{2r+1}^{*,*+j\sigma,*}(\nu_{\mathbb{C}}(X))[\tau^{-1}]$$

$$\parallel$$

$$\stackrel{\text{AHSS}}{\underset{\text{acc,alg}}{\text{AHSS}}} E_{r}^{*,*,*}(X_{j}^{P^{\infty}})$$

$$\stackrel{\text{MANSS}}{\underset{\text{MANSS}}{\text{MANSS}}} E_{2r+1}^{*,*+j\sigma,*}(X)$$

to lift differentials from the algebraic AHSS to the MANSS.

For any G and X we observe that the restriction map

$$\pi_V^G(X^h) \to \pi_{|V|}(X)$$

induces a map of spectral sequences

(1.3.2)
$$^{\text{MANSS}}E_r^{*,\star}(X) \to {}^{\text{ANSS}}E_r^{*,|\star|}(X)$$

from the MANSS to the Adams–Novikov spectral sequence of the underlying non-equivariant spectrum X (Corollary 4.1.5). This gives another means of determining MANSS differentials.

¹Because the filtration is by even skeleta instead of skeleta, we refer to this spectral sequence as the *accelerated algebraic Atiyah–Hirzebruch spectral sequence*.

1.3.1. *Computational examples.* We end this paper with some explicit computations of the MANSS.

- In the case of $G = C_2$, p = 2, and $X = ko_{C_2}$, we show that all of the differentials and multiplicative structure in the MANSS for ko_{C_2} can be deduced from (1.3.1) and (1.3.2). This recovers the main generating differentials in the C_2 -effective slice spectral sequence determined by the fourth author in [Kon20].
- In the case of $G = C_3$, p = 3, and X = tmf(2), the MANSS agrees with the $RO(C_3)$ graded homotopy fixed point spectral sequence for tmf(2). Following [HHR11, HSW23], we extend the $RO(C_3)$ -grading to an $RO^{\succ}(C_3)$ -grading which includes the "spoke sphere" S^{\succ} . Wilson computed the slice spectral sequence for tmf(2) in [Wil17]. We record an observation of Senger, that the μ_3 -orientation of tmf(2) [HSW23], the Hill-Hopkins-Ravenel norm, and known differentials in the integer graded homotopy fixed point spectral sequence (which is closely related to the decent spectral sequence for tmf₍₃₎) completely determine the structure of the $RO^{\succ}(C_3)$ -graded MANSS.

1.4. **Future work.** In future work, we plan to investigate how to extend our deformation $\widehat{\operatorname{SH}}_{G,p}^{\operatorname{art}}$ of $\operatorname{Sp}_{ip}^{BG}$ to a deformation $\operatorname{SH}_{G,p}^{\operatorname{art}}$ of $\operatorname{Sp}_{G,p}^{\operatorname{def}}$. Let $\mathbb{S}_{C_p,p}^{\operatorname{def}}$ denote $\operatorname{D\acute{e}c}^{\operatorname{ev.post}}(MU_p[G]^{\otimes \bullet+1})$. In the case of $G = C_p$, the idea would be to define the desired filtered C_p -spectrum $\Gamma_{C_p} \mathbb{S}_p$ as a pullback

where a_{λ} is the Euler class of the 2-dimensional representation of C_p given by rotation by $2\pi/p$. This would require the construction of a suitable filtration $\Gamma_{C_p}^{\Phi} a_{\lambda}^{-1} \mathbb{S}_p$ of the geometrically local sphere $a_{\lambda}^{-1} \mathbb{S}_p$ which maps to the localized filtration $a_{\lambda}^{-1} \mathbb{S}_{C_p,p}^{\text{def}}$.

In the case of $G = C_2$, this amounts to understanding the *a*-localization of the \mathbb{R} -motivic sphere in the Artin–Tate category. Unfortunately, this is the one object that the authors of [BHS20] flag as very mysterious, so the proposed program will encounter significant challenges. Nevertheless, the completion of this program would yield a notable application. The "Geffective slice spectral sequence"

$$E_r^{\star,\star}(\Gamma_G X)$$

associated to the filtered object $\Gamma_G X$ would potentially allow for much of the computational payoffs of the hypothetical odd primary analog BP_{μ_p} of $BP_{\mathbb{R}}$ discussed in [HHR11], while circumventing the need to construct BP_{μ_p} itself. Note that Hill–Hopkins–Ravenel [HHR11, p.1-5] have suggested that if such a spectrum BP_{μ_p} were known to exist, the its norm

$$N_{C_p}^{C_{p^2}}BP_{\mu_p}$$

could be used to resolve the 3-primary Kervaire invariant problem.

1.5. Organization of the paper. In Section 2, we introduce the foundations of filtered spectra which will be used throughout the paper. We end this section with our definition of complete artificial motivic spectra.

In Section 3, we recall the construction of the C_2 -effective slice spectral sequence. We then define, for E an arbitrary ring spectrum, the E-modified Adams spectral sequence. We prove that this modified Adams spectral sequence is actually the Adams spectral sequence based on the Borel G-spectrum E[G]. The modified Adams–Novikov spectral sequence (MANSS) is the instance of this spectral sequence where $E = MU_p$. We finish this section by proving that the Borel completion of the C_2 -effective slice filtration is the even Postnikov décalage of the MANSS filtration.

In Section 4, we introduce our main computational tools. We define the double complex spectral sequences, and compute their E_2 -terms. We then consider for $G = C_2$ the special case when X is the Borel completion of an $MU_{\mathbb{R}}$ -projective C_2 -spectrum (see Definition 3.1.3), and the special case when X has a trivial C_2 -action. We define the motivic MANSS, and show that mod τ , the motivic MANSS is isomorphic to the algebraic Atiyah–Hirzebruch spectral sequence.

In Section 5, we apply our methods to completely compute the MANSS in the case of the C_2 -spectrum ko_{C_2} , and the C_3 -spectrum tmf(2).

1.6. Conventions and notation. Throughout, we will use the theory of ∞ -categories (quasicategories) as developed by [Joy02, Lur09, Lur17]. Since the nerve functor from 1-categories to ∞ -categories is fully faithful, we will not include this functor in the notation. We will also refer to ∞ -categories simply as categories from this point onwards. Throughout, G will denote a finite group. We use the following notations:

\mathbb{Z}	(\mathbb{Z}, \leq) as a partially ordered set
\mathbb{Z}^{δ}	\mathbb{Z} as a discrete category
$\mathcal{C}^{\mathrm{Fil}}$	$\operatorname{Fun}(\mathbb{Z}^{\operatorname{op}},\mathcal{C})$
$\mathcal{C}^{\mathrm{Gr}}$	$\operatorname{Fun}(\mathbb{Z}^{\delta},\mathcal{C})$
Pr^{L}	the category of presentable categories and left adjoints
$\mathrm{Pr}^{L,\mathrm{st}}$	the category of presentable stable categories and left adjoints
$\operatorname{Sp}, \operatorname{Sp}_G, \operatorname{Sp}^{BG}$	the category of spectra, genuine G -spectra, Borel G -spectra
$\mathrm{SH}(k)$	Morel–Voevodsky ∞ -category of motivic spectra over k [MV99]
\mathcal{C}_{ip}	∞ -category of modules over the <i>p</i> -complete unit in \mathcal{C}
\otimes	symmetric monoidal product in the category at hand
$\operatorname{Ext}^{s,t}((MU_p)_*X)$	$\operatorname{Ext}_{(MU_p)_*MU_p}^{s,t}((MU_p)_*,(MU_p)_*X)$

Recall that a presentably symmetric monoidal, stable category is an object in the category $\operatorname{CAlg}(\operatorname{Pr}^{L,\operatorname{st}})$. We follow [BHS20, p.16] and use the following conventions: Given $\mathcal{C} \in \operatorname{CAlg}(\operatorname{Pr}^{L,\operatorname{st}})$, let \mathcal{C}_p denote the category of *p*-complete objects in \mathcal{C} . For $\mathcal{D} \in \{\mathcal{C}, \mathcal{C}^{\operatorname{Fil}}, \mathcal{C}^{\operatorname{Gr}}\}$ and $\mathcal{C} \in \{\operatorname{Sp}, \operatorname{Sp}_G, \operatorname{Sp}^{BG}\}$, let X_p denote the *p*-completion of $X \in \mathcal{D}$ and let $\mathcal{D}_{ip} = \operatorname{Mod}(\mathcal{D}; \mathbb{1}_p)$ denote the category of modules over $\mathbb{1}_p$. When working in one of these categories \mathcal{D}_{ip} , we will write \otimes for the symmetric monoidal structure $\otimes_{\mathbb{1}_p}$. We use this convention in Definition 2.4.6, Proposition 2.4.7 and throughout Section 3.1, Section 3.4, Section 4, and Section 5. Note that we still use the symbol \otimes_R to mean the (underived) tensor product in the category of modules over a discrete ring and we use \otimes to mean the usual tensor product of (graded) abelian groups in these sections. We expect that the meaning is clear from the context. We also write X_E for Bousfield localization of a spectrum X at a spectrum E.

We write $\operatorname{map}_{\mathcal{C}}$ for the mapping spectrum of a stable category \mathcal{C} or simply map when the context is clear. Similarly, we write $\operatorname{Map}_{\mathcal{C}}$ for the mapping space of a category \mathcal{C} and Map when the category \mathcal{C} is clear from context. If \mathcal{C} is \mathcal{D} -linear for some $\mathcal{D} \in \operatorname{CAlg}(\operatorname{Pr}^{L,\operatorname{st}})$ we write $\operatorname{map}_{\mathcal{C}}^{\mathcal{D}}(-,-): \mathcal{C}^{\operatorname{op}} \times \mathcal{C} \to \mathcal{D}$ for the mapping object in \mathcal{D} of the category \mathcal{C} or simply $\operatorname{map}^{\mathcal{D}}$ when \mathcal{C} is clear from the context.

For $X \in \text{Sp}_G$, and $V \in RO(G)$, we write

$$\pi_{V}^{H}(X) \coloneqq [\mathbb{S}^{V}, X]^{H} = \pi_{0}((\Sigma^{-V}X)^{H})$$
7

for the RO(G)-graded H-equivariant homotopy groups of X. As H-varies, these form a Mackey functor, which we denote

$$\underline{\pi}_V(X).$$

For $X \in \operatorname{Sp}^{BG}$, we will similarly define

$$\pi_V^H(X^h) \coloneqq \pi_0((\Sigma^{-V}X)^{hH}),$$

and we denote the associated Mackey functor

$$\underline{\pi}_V(X^n).$$

We shall write $\{E_r^{f,n}\}$ to denote a spectral sequence valued in abelian groups, and $\{\underline{E}_r^{f,n}\}$ to denote a spectral sequence valued in *G*-Mackey functors. Given a spectral sequence $\{\underline{E}_r^{f,n}\}$ valued in Mackey functors, we will write $\{E_r^{f,n}\}$ (removing the underline) to denote the value of the spectral sequence of Mackey functors at G/G:

$$E_r^{f,n} \coloneqq \underline{E}_r^{f,n}(G/G).$$

1.7. Acknowledgements. This project was begun in May 2021 at an American Institute for Mathematics workshop Equivariant techniques in stable homotopy theory and finalized during a SQuaRE at the American Institute for Mathematics. The authors thank AIM for providing a supportive and mathematically rich environment. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 1010342555. The second author was partially supported by the NSF grant DMS-2005476, and he completed part of this project as a visiting scholar at Northwestern University. The third author was supported by the NSF grant DMS-2204357. The fourth author was supported by the NSF grant DMS-1926686. The authors thank Andrew Senger, for generously sharing his computations of the $RO^{\succ}(C_p)$ -graded homotopy fixed-point spectral sequence of $E_{k(p-1)}$. The authors also benefited from conversations with Benjamin Antieau, Robert Burklund, Jeremy Hahn, Mike Hill, Max Johnson, and Noah Riggenbach.

2. Filtered objects and deformations

Throughout this section, let C be a presentably symmetric monoidal, stable category and assume C is generated by a set S of compact dualizable objects. In this situation, we say that the set S rigidly generates C.

2.1. Filtered objects. Here we summarize results about filtered and graded objects in C, denoted C^{Fil} and C^{Gr} (see Section 1.6). By our running assumptions on C, we know that $C^{\text{Fil}}, C^{\text{Gr}} \in \text{CAlg}(\text{Pr}^{L,\text{st}})$ where we equip each of these categories with the Day convolution symmetric monoidal structure [Day71, Gla16]. There is an associated graded functor

$$\operatorname{Gr}^*: \mathcal{C}^{\operatorname{Fil}} \to \mathcal{C}^{\operatorname{Gr}},$$

which is symmetric monoidal [GP18, Theorem 1.13]. Given a filtered spectrum A_{\bullet} we define $A(-i)_{\bullet}$ such that $A(-i)_{j} = A_{j+i}$. Letting $\mathbb{1}_{\mathcal{C}}$ denote the unit of the symmetric monoidal product on \mathcal{C} , we write $\Sigma^{s} \mathbb{1}_{\mathcal{C}}(-t)$ for the *s*-th suspension of the symmetric monoidal unit shifted by *t* and note that the homotopy groups in the category of graded objects in \mathcal{C} are bigraded in the sense that for $A \in \mathcal{C}^{\text{Gr}}$,

$$\pi_{s,t}A \coloneqq \pi_0 \operatorname{map}(\Sigma^s \mathbb{1}_{\mathcal{C}}(-t), A).$$

If \mathcal{C} is Sp_G -linear, we have a functor

$$\operatorname{map}^{\operatorname{Sp}_G}(-,-) = \mathcal{C}^{\operatorname{op}} \times \mathcal{C} \to \operatorname{Sp}_G$$

and \mathcal{C}^{Fil} and \mathcal{C}^{Gr} are also Sp_G -linear. Note that since \mathcal{C} is tensored over Sp_G we can can tensor with representation spheres to produce $\Sigma^V X \coloneqq \mathbb{S}^V \otimes X$ for any $X \in \mathcal{C}$. We can therefore consider the $RO(G) \times \mathbb{Z}$ -graded Mackey functor valued homotopy of $A \in \mathcal{C}^{\text{Fil}}$ with value

$$\underline{\pi}_{V,t}A \coloneqq \underline{\pi}_0 \operatorname{map}^{\operatorname{Sp}_G}(\Sigma^V \mathbb{1}_{\mathcal{C}}(-t), A)$$

at $(V, t) \in RO(G) \times \mathbb{Z}$.

Construction 2.1.1. Given an object $I \in C^{\text{Fil}}$, the spectral sequence associated to the filtered object I is the spectral sequence that comes from applying $\pi_*(-) \coloneqq \pi_* \operatorname{map}(\mathbb{1}_{\mathcal{C}}, -)$ to the tower of cofiber sequences



where $I_n/I_{n+1} := \operatorname{cofib}(I_{n+1} \to I_n)$. If $I_{\infty} = 0$ then we say I is complete and the associated conditionally convergent spectral sequence has signature

$$E_1^{f,n}(I_{\bullet}) = \pi_n(\operatorname{Gr}^f I) \implies \pi_n I_{-\infty}$$

where $I_{-\infty}$ is the colimit. The differentials take the form

$$d_r: E_r^{f,n} \to E_r^{f+r,n-1}.$$

We will also use a variant of this construction for Sp_G -linear categories. Suppose \mathcal{C} is Sp_G -linear and $I \in \mathcal{C}^{\text{Fil}}$. When $I_{\infty} = 0$ then we say I is complete and there is a conditionally convergent spectral sequence of Mackey functors with signature

$$\underline{E}_1^{f,V} = \underline{\pi}_V(\operatorname{Gr}^f I) \implies \underline{\pi}_V I_{-\infty}.$$

The differentials take the form

$$d_r: \underline{E}_r^{f,V} \to \underline{E}_r^{f+r,V-1}.$$

One source of filtered objects comes from cosimplicial objects. Given a cosimplicial object

$$X^{\bullet}: \Delta \to \mathcal{C}$$

we get a filtered object $\{fil_n X^{\bullet}\}$ by taking

(2.1.2)
$$\operatorname{fil}_n X^{\bullet} \coloneqq \operatorname{fiber}(\operatorname{Tot} X^{\bullet} \to \operatorname{Tot}_{n-1} X^{\bullet})$$

(where by convention, we define $\operatorname{Tot}_k X^{\bullet}$ to be trivial if k is negative). The associated spectral sequence is then the Bousfield–Kan spectral sequence, which we will denote:

$$E_r^{s,n}(X^{\bullet}) \coloneqq E_r^{s,n}(\operatorname{fil}_{\bullet} X^{\bullet}).$$

Remark 2.1.3. Our indexing convention differs from the standard indexing convention for the Bousfield–Kan spectral sequence. To illustrate this, suppose that E is a flat homotopy commutative ring spectrum, and that X is a spectrum. Then (with our indexing convention) the E-based Adams Spectral Sequence (E-ASS) for X is the spectral sequence associated to the cosimplicial object $E^{\otimes \bullet +1} \otimes X$ with

$$E-ASS E_2^{s,n} := E_2^{s,n} (E^{\otimes \bullet +1} \otimes X) \cong \operatorname{Ext}_{E_*E}^{s,s+n} (E_*, E_*X) \implies \pi_n(X_E^{\wedge})$$

where $X_E^{\wedge} := \operatorname{Tot} E^{\otimes \bullet + 1} \otimes X$ is the *E*-nilpotent completion of *X*.

Suppose I and J are filtered objects. Let $I\otimes J$ denote the Day convolution [Gla16]; concretely, this is

(2.1.4)
$$(I \otimes J)_n = \operatorname{colim}_{i+j \ge n} I_i \otimes J_j$$

on objects.

Lemma 2.1.5. Let d_1^I and d_1^J , respectively, denote the d_1 differentials in Construction 2.1.1 associated to the filtered objects I, J. Then we have

(2.1.6)
$$\operatorname{Gr}^{n}(I \otimes J) = \bigoplus_{i+j=n} \operatorname{Gr}^{i} I \otimes \operatorname{Gr}^{j} J$$

and whenever

$$x \otimes y \in \operatorname{im}\left(\pi_s(\operatorname{Gr}^i I) \otimes \pi_s(\operatorname{Gr}^j J) \longrightarrow \bigoplus_{i+j=n} \pi_s(\operatorname{Gr}^i I \otimes \operatorname{Gr}^j J)\right)$$

we have

$$d_1^{I \otimes J}(x \otimes y) = d_1^I(x) \otimes y + (-1)^{|x|} x \otimes d_1^J(y).$$

Proof. Equation (2.1.6) follows from the fact that the functor

$$\operatorname{Gr}^* \colon \mathcal{C}^{\operatorname{Fil}} \to \mathcal{C}^{\operatorname{Gr}}$$

is symmetric monoidal [GP18, Theorem 1.13] (see also [AKS18, Theorem 3.3.10]). For the statement about $d_1^{I\otimes J}$, we first claim that if $z \in \pi_s(\operatorname{Gr}^i I \otimes \operatorname{Gr}^j J)$ then

(2.1.7)
$$d_1(z) \in \pi_s(\operatorname{Gr}^{i+1} I \otimes \operatorname{Gr}^j J) \oplus \pi_s(\operatorname{Gr}^i I \otimes \operatorname{Gr}^{j+1} J).$$

To see this, let $P_{i,j}$ denote the pushout in the diagram

There is a map of cofiber sequences

for each pair of integers i, j such that i + j = n (see [AKS18, Lemma 3.3.1]). The diagram (2.1.8) shows that the image of $P_{i,j}$ is in the image of the inclusion

$$(\operatorname{Gr}^{i+1}I \otimes \operatorname{Gr}^{j}J) \oplus (\operatorname{Gr}^{i}I \otimes \operatorname{Gr}^{j+1}J) \to \operatorname{Gr}^{n+1}(I \otimes J)$$

proving (2.1.7). Moreover, the diagram

commutes (see [May01, Lemma 4.7]) and so, writing \tilde{d}_1^K for the map of spectra inducing the map d_1^K , we have

$$\tilde{d}_1^{I\otimes J} = \tilde{d}_1^I \otimes \operatorname{id} + \operatorname{id} \otimes \tilde{d}_1^J.$$

Finally, naturality of the map

$$\pi_{s}(\operatorname{Gr}^{i} I) \otimes \pi_{s}(\operatorname{Gr}^{j} J) \longrightarrow \bigoplus_{i+j=n} \pi_{s}(\operatorname{Gr}^{i} I \otimes \operatorname{Gr}^{j} J)$$

gives rise to the desired statement, where the sign comes from the natural isomorphism

$$\Sigma^{-1} \mathrm{Gr}^{i} I \otimes \mathrm{Gr}^{j} J \cong \mathrm{Gr}^{i} I \otimes \Sigma^{-1} \mathrm{Gr}^{j}.$$

Lemma 2.1.9. Suppose $\lim_{s} I_s \otimes J_i = 0$ and $\lim_{s} I_i \otimes J_s = 0$ for each $i, j \in \mathbb{Z}$. Then $\lim_{s} (I \otimes J)_s = 0$.

Proof. Write $\operatorname{colim}_{i+j\geq s} I_i \otimes J_j$ as the coequalizer of the diagram

$$\oplus_{i+j=s+1}I_i\otimes J_j \xrightarrow{\longrightarrow} \oplus_{i+j}I_i\otimes J_j$$

where one map is induced by the structure maps of I and the other is induced by the structure maps of J. Consequently, the limit $\lim_{s \to j} \operatorname{colim}_{i+j\geq s} I_i \otimes J_j$ can be identified as the coequalizer of the diagram

$$(\lim_{s} \oplus_{i+j=s+1} I_i \otimes J_j) \xrightarrow{\longrightarrow} (\lim_{s} \oplus_{i+j=s} I_i \otimes J_j)$$

where one map is induced by the structure maps of I and the other map is induced by the structure maps of J. It therefore suffices to show that $\lim_{s} \bigoplus_{i+j=s} I_i \otimes J_j = 0$. Now we note that this limit is given by the limit

$$\dots \xrightarrow{\longrightarrow} \oplus_{i+j=s} I_i \otimes J_j \xrightarrow{\longrightarrow} \oplus_{i+j=s} I_i \otimes J_j \xrightarrow{\longrightarrow} \dots$$

where the top maps are induced by the structure maps of I and the bottom maps are induced by the structure maps of J. By examination, the limit of the top maps is zero and the limit of the bottom maps are zero. Consequently, the limit of the differences of the two maps is also zero.

2.2. Tower functors. Fix a finite group G and a category $\mathcal{C} \in CAlg(Pr^{L,st})$.

Definition 2.2.1 ([BHS20, p. 71]). A tower functor in C is a lax symmetric monoidal functor

$$T: \mathcal{C} \to \mathcal{C}^{\mathrm{Fil}}$$

Next we recall two examples of tower functors that come from different families of truncations of Sp_G , namely those arising from the regular slice filtration and the Postnikov filtration.

Definition 2.2.2 ([Ull13]). Let $\operatorname{Sp}_{G,\geq n}^{\operatorname{slice}}$ denote the coreflective subcategory of Sp_G generated by the set $\{G_+ \wedge_H S^{\rho_H k}\}$ where *H* is a subgroup of *G*, ρ_H is the regular representation of *H*, and $|\rho_H| k \geq n$. The associated functor

$$P_{\geq \bullet} \colon \operatorname{Sp}_{C_2} \to \operatorname{Sp}_{C_2}^{\operatorname{Fil}}$$

is a tower functor (see [BHS20, Construction 4.2]). Similarly, $P_{\geq 2\bullet}$ is a tower functor called the even slice tower.

Definition 2.2.3 ([HY18]). Let $\operatorname{Sp}_{G,\geq n}^{\operatorname{post}}$ denote the localizing subcategory generated by $G_+ \wedge_H \operatorname{\mathbb{S}}^k$ for H a subgroup of G and $k \geq n$. The associated functor

$$\tau_{\geq \bullet} \colon \operatorname{Sp}_{C_2} \to \operatorname{Sp}_{C_2}^{\operatorname{Fil}}$$

is a tower functor. Similarly, $\tau_{\geq 2^{\bullet}}$ is a tower functor called the even Postnikov tower. This follows because it is a t-structure that is compatible with the symmetric monoidal structure

(the full subcategory $\text{Sp}_{G,\geq 0}^{\text{post}}$ clearly contains the symmetric monoidal unit it is closed under the symmetric monoidal product (see [Lur17, Remark 2.2.1.2]).

The following result will be useful later.

Lemma 2.2.4. There is an inclusion of subcategories $\operatorname{Sp}_{C_2,\geq n}^{\operatorname{post}} \subseteq \operatorname{Sp}_{C_2,\geq n}^{\operatorname{slice}}$. Consequently, there is a natural transformation of tower functors

$$\tau_{\geq \bullet} \implies P_{\geq \bullet}$$

After postcomposing with the functor $(-)^e \colon \operatorname{Sp}_{C_2}^{\operatorname{Fil}} \to \operatorname{Sp}^{\operatorname{Fil}}$, this natural transformation is an equivalence.

Proof. We check the generators of $\operatorname{Sp}_{C_2,\geq n}^{\operatorname{post}}$ are in $\operatorname{Sp}_{C_2,\geq n}^{\operatorname{slice}}$. By definition, we have $C_{2+} \otimes \mathbb{S}^k \in \operatorname{Sp}_{C_2,\geq n}^{\operatorname{slice}}$, for all $k \geq n$. It suffices to show $\mathbb{S}^k \in \operatorname{Sp}_{C_2,\geq n}^{\operatorname{slice}}$ for all $k \geq n$. Since a localizing subcategory is closed under suspension, it suffices to show $\mathbb{S}^n \in \operatorname{Sp}_{C_2,\geq n}^{\operatorname{slice}}$. For n = 1, this is implied by the cofiber sequence

$$C_{2+} \wedge S^1 \to S^1 \to S^{\rho}.$$

The case for larger n follows from the multiplicative relation [Ull13, Cor. 4.2].

The underlying equivalence follows from the fact that both filtrations forget to the classical Postnikov filtration on underlying spectra. \Box

Remark 2.2.5. Recall that there is a map $a: \mathbb{S}^{-\sigma} \to \mathbb{S}$ in Sp_{C_2} called the Euler class with the property that completion at a

$$(-)_a^{\wedge} \colon \operatorname{Sp}_{C_2} \longrightarrow \operatorname{Sp}_{C_2}$$

agrees with the cofree functor $F(EC_{2+}, -)$ sending a C_2 -spectrum X to its underlying Borel C_2 -spectrum. We also note that we can identify the essential image of the functor $(-)_a^{\wedge}$ with the category of Borel C_2 -spectra Sp^{BC₂}.

Lemma 2.2.6. Let $f: X \to Y$ be a map in Sp_{C_2} . If f is an underlying equivalence, then f induces an equivalence between the *a*-completions $f_a^{\wedge}: X_a^{\wedge} \to Y_a^{\wedge}$.

Proof. Since a-completion is equivalent to the cofree construction

$$F(EC_{2+},-): \operatorname{Sp}_{C_2} \to \operatorname{Sp}_{C_2},$$

it preserves weak equivalences (see, e.g. [HM17, §2.2.1]). The result follows since the essential image of the functor $F(EC_{2+}, -) = (-)_a^{\wedge}$ is Sp^{BC_2} whose weak equivalences are detected by the underlying spectra.

We therefore have the following corollary of the last two lemmas.

Proposition 2.2.7. There is a natural equivalence of tower functors

$$(\tau_{\geq \bullet}(-))_a^{\wedge} \xrightarrow{\simeq} (P_{\geq \bullet}(-))_a^{\wedge} \colon \operatorname{Sp}_{C_2} \to (\operatorname{Sp}^{BC_2})^{\operatorname{Fil}}$$

Proof. By Lemma 2.2.4 and Lemma 2.2.6 it suffices to check that the functor $(-)_a^{\wedge}$: Sp^{Fil}_{C₂} → Sp^{Fil}_{C₂} is lax symmetric monoidal. Since $(-)_a^{\wedge} = F(EC_{2+}, -)$: Sp^{Fil}_{C₂} → (Sp^{BC₂})^{Fil} is right adjoint to inclusion, it suffices to show that the inclusion functor is symmetric monoidal, which is evident.

2.3. Décalage. We described (2.1.2) how to associate a filtered object $\operatorname{fil}_{\bullet}(X^{\bullet})$ to a cosimplicial object X^{\bullet} . We now describe how a tower functor allows us to associate a different filtered object (the "*T*-décalage" filtration).

Definition 2.3.1. Given a tower functor $T: \mathcal{C} \to \mathcal{C}^{\text{Fil}}$, and a cosimplicial object $X^{\bullet} \in \mathcal{C}^{\Delta}$, the *T*-décalage of X^{\bullet} defined to be the filtered object given by

$$\operatorname{D\acute{e}c}^T(X^{\bullet}) \coloneqq \operatorname{Tot} T(X^{\bullet}).$$

For example, if $T = \tau_{\geq \bullet}$, then $\operatorname{D\acute{e}c}^T(X^{\bullet})_w = \operatorname{Tot}_i \tau_{\geq w} X^i$.

Notation 2.3.2. We refer to the $P_{\geq \bullet}$ -décalage as the *slice décalage* (Déc^{slice}) and the $\tau_{\geq \bullet}$ -décalage as the *Postnikov décalage* (Déc^{post}). We refer to the $P_{\geq 2\bullet}$ -décalage as the *even slice décalage* (Déc^{ev.slice}) and the $\tau_{\geq 2\bullet}$ -décalage as the *even Postnikov décalage* (Déc^{ev.post}).

Remark 2.3.3. The Postnikov décalage described above corresponds to "classical décalage" on the filtered object $\operatorname{fil}_{\bullet} X^{\bullet}$ via the Beilinson *t*-structure. See [GIKR22, Rmk. 3.7] for a discussion of this point.

Postnikov décalage has the effect of shifting the pages of the associated spectral sequences.

Theorem 2.3.4 (Hedenlund [Hed20, Thm. II.3.2]). Suppose X^{\bullet} is a cosimplicial spectrum. There is a page-shifting isomorphism of spectral sequences

$$E_{r+1}^{s,n}(X^{\bullet}) \cong E_r^{s+n,n}(\operatorname{D\acute{e}c}^{\operatorname{post}}(X^{\bullet})).$$

Finally, suppose that X^{\bullet} is a cosimplicial spectrum with $\pi_* X^s$ concentrated in even degrees. Then $E_r^{s,n}(X^{\bullet})$ is concentrated in degrees where s + n is even, the associated graded of the Postnikov décalage of X^{\bullet} is concentrated in even degrees, and we therefore have

(2.3.5)
$$E_{2r+1}^{s,n}(X^{\bullet}) \cong E_{2r}^{s+n,n}(\operatorname{D\acute{e}c}^{\operatorname{post}}(X^{\bullet})) \cong E_r^{\frac{s+n}{2},n}(\operatorname{D\acute{e}c}^{\operatorname{ev.post}}(X^{\bullet})).$$

2.4. **Deformations.** Here we present our deformation of Borel *G*-equivariant stable homotopy theory. We will see in Section 3.4 that this gives a different presentation of the *a*-complete Artin–Tate real motivic stable homotopy category. We will also demonstrate in Section 4 that this perspective is useful for computing RO(G)-graded homotopy groups of Borel *G*-spectra. First, we need to define a deformation of an object in CAlg($Pr^{L,st}$).

Notation 2.4.1. Given a \mathcal{C}^{Fil} -algebra \mathcal{D} in CAlg($\Pr^{L,\text{st}}$), we write $\mathbb{1}_{\mathcal{D}}(i) \coloneqq \mathbb{1}_{\mathcal{D}} \otimes \mathbb{1}_{\mathcal{C}}(i)$. Definition 2.4.2 ([BHS20, Definition C.14]). A *deformation* of \mathcal{C} is a diagram



in $\operatorname{CAlg}(\operatorname{Pr}^{L,\operatorname{st}})$ such that

(1) there is a map

$$i: \mathbb{Z} \to \ker \pi_0 \operatorname{Pic}(\operatorname{Re})$$

of abelian groups sending n to $\mathbb{1}_{\mathcal{C}^{def}}(n)$

(2) the map

$$\operatorname{Map}_{\mathcal{C}^{\operatorname{def}}}(\mathbb{1}_{\mathcal{C}^{\operatorname{def}}}(i),\mathbb{1}_{\mathcal{C}^{\operatorname{def}}}(j)) \to \operatorname{Map}(\mathbb{1}_{\mathcal{C}},\mathbb{1}_{\mathcal{C}})$$

induced by Re is an equivalence for all integers $i \leq j$.

(3) the category \mathcal{C}^{def} is rigidly generated by a set $\{c(X_{\alpha})(i)\}$ where \mathcal{C} is rigidly generated by the set $\{X_{\alpha}\}$.

Finally, we define our deformation of Borel G-spectra. Though this definition is currently unmotivated, our project in the next two sections will be to motivate this definition by showing that it recovers a suitable completion of the Artin–Tate \mathbb{R} -motivic stable homotopy category (see Section 3.4) and that it is useful for computations (see Section 4). First, we define an object in CAlg(Sp^{Fil}_G). This will involve the E-based Amitsur complex.

$$(2.4.3) E \Longrightarrow E^{\otimes 2} \Longrightarrow \dots$$

which is the standard cosimplicial spectrum whose totalization is the *E*-nilpotent completion \mathbb{S}_{E}^{\wedge} and associated Bousfield–Kan spectral sequence can be identified with the *E*-based Adams spectral sequence.

Remark 2.4.4. Given an associative monoid $E \in \text{Sp}^{BG}$, then the functor $\text{Déc}^{\text{post}}(-\otimes E^{\otimes \bullet +1})$ given by the Postnikov décalage of the functor $-\otimes E^{\otimes \bullet +1}$ is again a tower functor (see [BHS20, Construction C.9]).

Notation 2.4.5. Given an associative monoid E in G-spectra, we write $E[G] = E \otimes G_+$ for the group ring, which is again an associative monoid in G-spectra.

Definition 2.4.6. Let $\mathbb{S}_{G,p}^{\text{def}} \coloneqq \text{Déc}^{\text{ev.post}}(MU_p[G]^{\otimes \bullet +1})$ be the even Postnikov décalage of the $MU_p[G]$ -based Amitsur complex (2.4.3) in the category Sp_{ip}^{BG} . Define the category of complete artificial motivic spectra to be

$$\widehat{\operatorname{SH}}_{G,p}^{\operatorname{art}} \coloneqq \operatorname{Mod}((\operatorname{Sp}_{ip}^{BG})^{\operatorname{Fil}}; \mathbb{S}_{G,p}^{\operatorname{def}}).$$

Proposition 2.4.7. The category $\widehat{SH}_{G,p}^{\operatorname{art}}$ is a deformation of $\operatorname{Sp}_{ip}^{BG}$.

Proof. This follows directly from [BHS20, Proposition C.2] and Remark 2.4.4 using $F = D\acute{e}c^{\text{ev.post}}(-\otimes MU_p[G]^{\otimes \bullet +1})$. The fact that $\mathbb{S}_{G,p}^{\text{def}}$ is a commutative monoid in $(\operatorname{Sp}_{ip}^{BG})^{\text{Fil}}$ follows from the proof of [PP21, Theorem 5.64], which generalizes verbatim to $\operatorname{Sp}_{ip}^{BG}$.

3. A TALE OF TWO SPECTRAL SEQUENCES

Work of the fourth author [Kon20] demonstrated the utility of the C_2 -effective slice spectral sequence for computing C_2 -equivariant homotopy groups. In this section, we give a new perspective on the C_2 -effective slice spectral sequence. Specifically, we construct the modified Adams–Novikov spectral sequence (MANSS), which is useful for computing the homotopy groups of Borel G-spectra for any finite group G. In the case of $G = C_2$, we compare this to the C_2 -effective slice spectral sequence.

One advantage of the modified Adams–Novikov spectral sequence is that the differentials can be computed from the classical Adams–Novikov spectral sequence for the underlying spectrum and the equivariant attaching maps for EG. When X is a Borel G-spectrum with trivial action the differentials in the MANSS can often be deduced from the attaching maps of BG together with the differentials in the Adams–Novikov spectral sequence for the underlying non-equivariant spectrum X^e . We illustrate this phenomenon in Section 5. 3.1. The C_2 -effective slice spectral sequence. Recall that we have the C_2 -equivariant Betti realization functor from the \mathbb{R} -motivic stable homotopy category:

$$\operatorname{Be}_{C_2}$$
: $\operatorname{SH}(\mathbb{R})_{i2} \to \operatorname{Sp}_{C_2,i2}$.

The C_2 -effective slice spectral sequence [Kon20] is the spectral sequence associated to the filtration we now define. Suppose $X = \text{Be}_{C_2}(\overline{X})$. There there is an effective slice filtration

$$\cdots \to \operatorname{fil}_{f+1}^{\operatorname{eff}} \overline{X} \to \operatorname{fil}_f^{\operatorname{eff}} \overline{X} \to \ldots$$

in \mathbb{R} -motivic homotopy theory [Voe02] with filtration quotients

$$s_f(\overline{X}) = \operatorname{fil}_f^{\operatorname{eff}} \overline{X} / \operatorname{fil}_{f+1}^{\operatorname{eff}} \overline{X}$$

We call the C_2 -Betti realization of this filtration the C_2 -effective slice filtration of X associated to \overline{X} . The C_2 -effective slice filtration gives a functor

$$i_* : \mathrm{SH}(\mathbb{R})_{i2} \to \mathrm{Sp}_{C_2,i2}^{\mathrm{Fil}}$$
$$\overline{X} \mapsto \mathrm{Be}_{C_2}(\mathrm{fil}_{\bullet}^{\mathrm{eff}}X)$$

Slice décalage of the $MU_{\mathbb{R},2}$ -based Amitsur complex gives another functor valued in filtered C_2 -spectra:

$$\Gamma_{\mathbb{R}} : \operatorname{Sp}_{C_{2},i2} \to \operatorname{Sp}_{C_{2},i2}^{\operatorname{Fil}} \\ X \mapsto \operatorname{D\acute{e}c}^{\operatorname{ev.slice}} \left(X \otimes MU_{\mathbb{R},2}^{\otimes \bullet +1} \right).$$

The main theorem of [BHS20] is the following.

Theorem 3.1.1 (Galois reconstruction). The functor

is an equivalence, and there is an equivalence

$$i_* \mathbb{S}_{\mathbb{R},2} \simeq \Gamma_{\mathbb{R}} \mathbb{S}_{C_2,2}.$$

It follows that when restricted to $\operatorname{Sp}_{C_2,i2}$, the functor $\Gamma_{\mathbb{R}}$ actually takes values in

$$\operatorname{Mod}(\operatorname{Sp}_{C_2,i2}^{\operatorname{Fil}}; i_* \mathbb{S}_{\mathbb{R},2})$$

and the composite

$$\nu_{\mathbb{R}} \coloneqq i_*^{-1} \circ \Gamma_{\mathbb{R}} : \operatorname{Sp}_{C_2, i2} \to \operatorname{SH}(\mathbb{R})_{i2}^{\operatorname{AT}}$$

is a section of the functor Be_{C_2} . Given $X \in \operatorname{Sp}_{C_2,i2}$, the C_2 -effective slice filtration of X relative to $\nu_{\mathbb{R}}(X)$ is the filtered spectrum $\Gamma_{\mathbb{R}}X$. Therefore we shall refer to the filtration $\Gamma_{\mathbb{R}}X$ of X simply as the C_2 -effective slice filtration of X (without reference to the motivic spectrum $\nu_{\mathbb{R}}(X)$ the C_2 -effective slice filtration is taken with respect to).

The C_2 -effective slice spectral sequence for X associated to \overline{X} is the $RO(C_2) \times \mathbb{Z}$ -graded spectral sequence associated to the filtered C_2 -spectrum $i_*\overline{X}$ and it has signature

$$C_2$$
-ESSS $\underline{E}_1^{f,V} = \underline{\pi}_V \operatorname{Be}_{C_2}(s_f(\overline{X})) \implies \underline{\pi}_V X.$

In the case where $\overline{X} = \nu_{\mathbb{R}}(X)$, we shall simply refer to this spectral sequence as the C₂-effective slice spectral sequence of X (C₂-ESSS), and the E₁-term can be identified with

$$\underline{\pi}_{\star} \mathrm{Gr}^{*}(\Gamma_{\mathbb{R}}(X)).$$

One of the most attractive aspects of the C_2 -ESSS is the computability of its E_2 -term for the class of $MU_{\mathbb{R}}$ -projective spectra. Recall the following definition from [BHS20].

Definition 3.1.3. A spectrum $X \in \operatorname{Sp}_{C_2,i2}$ is $MU_{\mathbb{R}}$ -projective if $MU_{\mathbb{R},2} \otimes X$ is a retract of a C_2 -spectrum of the form $\bigoplus_i \Sigma^{n_i \rho} MU_{\mathbb{R},2}$.

For example, the (2-completion of) the C_2 -Betti realization of any regular localized quotient of MGL_2 in the sense of [Hea19, Definition 3.4] is $MU_{\mathbb{R}}$ -projective by [Hea19, §4]. The following theorem explains how the E_2 -term of the C_2 -ESSS for an $MU_{\mathbb{R}}$ -projective $X \in \text{Sp}_{i2}$ surprisingly only depends on the Adams–Novikov E_2 -term for the underlying non-equivariant spectrum $X^e \in \text{Sp}_{i2}$.

Theorem 3.1.4 (Burklund–Hahn–Senger [BHS20, Thm. 5.6]). Suppose that $X \in \text{Sp}_{C_2,i2}$ is $MU_{\mathbb{R}}$ -projective. Then the E_1 -term of its C_2 -ESSS is given by

$${}^{C_2 \text{-ESSS}}\underline{E}_1^{f,V}(X) = \bigoplus_s \text{Ext}^{s,2f}((MU_2)_*(X^e) \otimes \underline{\pi}_{V-f\rho+s}H\underline{\mathbb{Z}}_2).$$

3.2. The modified *E*-based Adams spectral sequence. Given $X \in \text{Sp}^{BG}$, and an associative monoid *E* in Sp^{BG} we can form the *E*-based Adams spectral sequence of *X*. Let \overline{E} denote the fiber of the unit map $\overline{E} = \text{fib}(\mathbb{S} \to E)$. We can then construct the standard *E*-based Adams resolution

of X. We view this as a filtered spectrum

$$\Gamma_E X \coloneqq \{\cdots \to \overline{E}^{\otimes 2} \otimes X \to \overline{E} \otimes X \to X = X = \dots\} \in (\operatorname{Sp}^{BG})^{\operatorname{Fil}}$$

Let

$$X_E^{\wedge} \coloneqq \operatorname{Tot}(E^{\bullet+1} \otimes X)$$

be the totalization of the *E*-based Amitsur complex tensored with X (2.4.3). If the map $X \to X_E^{\wedge}$ is an equivalence, we have

$$\lim_{\circ} \Gamma_E(X)_s = 0.$$

Of course, we always have

$$\operatorname{colim} \Gamma_E(X)_s = X.$$

Next, we will set up the homotopy fixed point spectral sequence so that it also conditionally converges to the colimit. Define

$$EG^{(k)} \coloneqq |\mathrm{sk}_k B_{\bullet}(G, G, *)|$$

where $B_{\bullet}(G, G, *)$ is the usual two-sided bar construction whose geometric realization is EG. Let $EG^{(-1)} = \emptyset$ by convention. The homotopy fixed point spectral sequence of $X \in Sp^{BG}$ is the spectral sequence associated to the tower

$$F(EG_+^{(0)}, X) \leftarrow F(EG_+^{(1)}, X) \leftarrow F(EG_+^{(2)}, X) \leftarrow \cdots$$

whose inverse limit in Sp^{BG} is X.

We will explicitly describe the decreasing filtration of X which gives rise to this tower. Write $\widetilde{E}G^{(k)}$ for the cofiber of the collapse map

$$EG_+^{(k)} \to S^0.$$

Then there is a cofiber sequence

$$F(\widetilde{E}G^{(s)}, X) \to X \to F(EG^{(s)}_+, X).$$

It follows that the homotopy fixed points spectral sequence computing $\underline{\pi}_{\star}(X)$ arises from the following decreasing filtration of X:

Here, the filtration quotients in this diagram are computed using the diagram of cofiber sequences

which identifies $\Sigma^{-1} \widetilde{E} G^{(s)} / \widetilde{E} G^{(s-1)}$ with $E G_+^{(s)} / E G_+^{(s-1)}$ for $s \ge 0$. We regard this construction as a functor

$$\Gamma^h \colon \mathbb{Z}^{\mathrm{op}} \to \mathrm{Sp}^{BG}$$

Specifically, set

$$\Gamma^{h}(X)_{s} \coloneqq \begin{cases} X & \text{for } s \leq 0\\ F(\widetilde{E}G^{(s-1)}_{+}, X) & \text{for } s > 0. \end{cases}$$

This has been arranged so that $\lim_{s} \Gamma^{h}(X)_{s} = 0$ and $\operatorname{colim}_{s} \Gamma^{h}(X)_{s} = X$. To see this, note that since the map

$$EG_+ \longrightarrow S^0$$

is a non-equivariant equivalence, the space $\widetilde{E}G$ is non-equivariantly contractible.

Now we mix these two filtrations in the following way. By tensoring the Adams resolution for S with X, we produce the *E*-Adams resolution for X in Sp^{BG} . If we also filter X by the filtration $\Gamma^h(X)$ as above then we produce an object

$$X_{\bullet,\blacksquare} \in \operatorname{Fun}(\mathbb{Z}^{\operatorname{op}} \times \mathbb{Z}^{\operatorname{op}}, \operatorname{Sp}^{BG})$$

defined on an object (i, j) by

(3.2.3)
$$X_{i,j} = \begin{cases} \overline{E}^{\otimes i} \otimes X & \text{for } i > 0, j \le 0 \\ X & \text{for } i \le 0, j \le 0 \\ \overline{E}^{\otimes i} \otimes F(\widetilde{E}G_{+}^{(j-1)}, X) & \text{for } i > 0, j > 0 \\ F(\widetilde{E}G_{+}^{(j-1)}, X) & \text{for } i \le 0, j > 0. \end{cases}$$

Note that $X_{\bullet,\bullet}$ is simply the exterior tensor product $\Gamma_E(\mathbb{S})_{\bullet}\overline{\otimes} \Gamma^h(X)_{\bullet}$.

We then consider the Day convolution

$$\Gamma_E^h(X)_{\bullet} \coloneqq \Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(X)_{\bullet}$$

By definition, $\Gamma_E^h(X)_{\bullet}$ is the left Kan extension in the diagram

and it is explicitly defined on objects by

$$\Gamma_E^h(X)_s = (\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(X)_{\bullet})_s = \operatorname{colim}_{a+b \ge s} \Gamma_E(\mathbb{S})_a \otimes \Gamma^h(X)_b.$$

Unpacking this, we compute that

(3.2.4)
$$\Gamma_E^h(X)_s = \begin{cases} X & \text{if } s \le 0\\ \underset{a+b\ge s}{\operatorname{colim}} \overline{E}^{\otimes a} \otimes F(\widetilde{E}G^{(b-1)}_+, X) & \text{if } s > 0. \end{cases}$$

Let \mathbb{N}^{op} denote the full subcategory of \mathbb{Z}^{op} generated by the nonnegative integers and let \mathbb{N}^{δ} denote the category whose objects are nonnegative integers and whose only morphisms are the identity maps. By Lemma 2.1.5 we can identify

(3.2.5)
$$\operatorname{cofib}(\Gamma_E^h(X)_{s+1} \to \Gamma_E^h(X)_s) = \bigoplus_{a+b=s} E \otimes \overline{E}^{\otimes a} \otimes F(EG_+^{(b)}/EG_+^{(b-1)}, X).$$

Notation 3.2.6. For $\star \in RO(G)$ we write $\star = * + \diamond$ where * is the virtual dimension of \star (representing a sum of trivial representations), and \diamond is a virtual representation of virtual dimension 0.

Using this notation, we have a $RO(G) \times \mathbb{Z}$ -graded spectral sequence with signature

$$\underline{E}^{-\mathrm{MASS}}\underline{E}_{1}^{s,n+\diamond}(X) = \bigoplus_{a+b=s} \underline{\pi}_{n+\diamond}(E \otimes \overline{E}^{\otimes a} \otimes F(EG_{+}^{(b)}/EG_{+}^{(b-1)},X)) \implies \underline{\pi}_{n+\diamond}(X_{E}^{h})$$

where

We shall refer to this spectral sequence as the modified *E*-based Adams spectral sequence (*E*-MASS) for *X*. The fact that the modified *E*-based Adams spectral sequence conditionally convergences whenever the *E*-based Adams spectral sequence converges follows from Lemma 2.1.9: The fact $\lim_{a} \left(\overline{E}^{\otimes a} \otimes F(\tilde{E}G^{(b)}, X) \right) = 0$ follows from the fact that $\tilde{E}G^{(b)}$ is dualizable and the *E*-based Adams spectral sequence for *X* converges. The fact that $\lim_{b} \left(\overline{E}^{\otimes a} \otimes F(\tilde{E}G^{(b)}, X) \right) = 0$ follows because \overline{E} has trivial action and $\widetilde{E}G$ is contractible in the category of Borel *G*-spectra.

3.3. The *E*-MASS as a generalized Adams spectral sequence. The following lemma will be useful for identifying certain filtration quotients.

Lemma 3.3.1. There is an equivalence

$$F(EG_{+}^{(k)}/EG_{+}^{(k-1)},\mathbb{S}) \simeq \mathbb{S}[G] \otimes \overline{\mathbb{S}[G]}^{\otimes k}$$

where $\overline{\mathbb{S}[G]} := \operatorname{fiber}(\mathbb{S} \to \mathbb{S}[G]).$

Proof. By definition, $EG^{(k)} \approx |\mathrm{sk}_k B_{\bullet}(G, G, *)|$. Consequently, there is an equivalence $F(EG_+^{(k)}, \mathbb{S}) \approx \mathrm{Tot}_k(F(B_{\bullet}(G, G, *)_+, \mathbb{S})).$

By [MNN17, Proposition 2.14], there is therefore an equivalence

$$F(EG_{+}^{(k)}, \mathbb{S}) \simeq \operatorname{cofib}(\overline{\mathbb{S}[G]}^{\otimes k+1} \to \mathbb{S})$$

The lemma then follows from the diagram of fiber sequences



Remark 3.3.2. When $G = C_2$, we have $\overline{\mathbb{S}[C_2]} = \mathbb{S}^{-\sigma} := F(\mathbb{S}^{\sigma}, \mathbb{S})$. To see this, note that there is a fiber sequence

$$\mathbb{S}^{-\sigma} \xrightarrow{a} \mathbb{S}^{0} \to Ca = \mathbb{S}[C_{2}]$$

so $\overline{\mathbb{S}[C_2]} = \mathbb{S}^{-\sigma}$, and the map $\overline{\mathbb{S}[C_2]}^{\otimes k} = \mathbb{S}^{-k\sigma} \to \mathbb{S}$ is simply a^k . Consequently, by Lemma 3.3.1 we have equivalences

$$EC_{2+}^{(k)}/EC_{2+}^{(k-1)} \simeq \Sigma^{k\sigma}Ca$$

for each $k \geq 0$.

Now we prove the main result from this section.

Proposition 3.3.3 (The *E*-MASS is a generalized Adams spectral sequence). The modified *E*-based Adams spectral sequence for a finite group *G* agrees with the E[G]-based Adams spectral sequence from the E_2 -term onwards.

Proof. To prove this result, it suffices to prove that the filtered spectrum that we use to construct the modified E-based Adams spectral sequence is in fact an E[G]-based Adams resolution.

Since $\widetilde{EG}^{(s)}$ is finite, the modified *E*-based Adams spectral sequence for *X* is the spectral sequence associated to the Day convolution

$$\Gamma_E^h(X)_{\bullet} \simeq \Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X.$$

(see Lemma 3.3.1). We therefore wish to show that

$$\cdots \to (\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X)_1 \to (\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X)_0 = X$$

is an E[G]-injective resolution of X in the sense of [Mil81]. We need to check that

- (1) there is an equivalence $\operatorname{Gr}^{n}(\Gamma_{E}(\mathbb{S})_{\bullet} \otimes \Gamma^{h}(\mathbb{S})_{\bullet} \otimes X) \simeq E[G] \otimes M$ for some spectrum M and consequently, $\operatorname{Gr}^{n}(\Gamma_{E}(\mathbb{S})_{\bullet} \otimes \Gamma^{h}(\mathbb{S})_{\bullet} \otimes X)$ is E[G]-injective;
- (2) the map

$$(\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X)_n \to \operatorname{Gr}^n(\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X)$$

induces a monomorphism after applying $E[G]_*$ for all $n \ge 0$ (note that the case n = 0 is clear because $(\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X)_1 = \overline{E[G]}$).

The first part, follows from (3.2.5) since

$$(3.3.4) \qquad \bigoplus_{a+b=s} E \otimes \overline{E}^{\otimes a} \otimes \mathbb{S}[G] \otimes \overline{\mathbb{S}[G]}^{\otimes b} \otimes X \simeq E[G] \otimes (\bigoplus_{a+b=s} \overline{E}^{\otimes a} \otimes \overline{\mathbb{S}[G]}^{\otimes b}) \otimes X$$

for each $s \ge 0$.

To prove the second part, it suffices to show that the induced map

$$E[G]_*((\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X)_{n+1}) \to E[G]_*((\Gamma_E(\mathbb{S})_{\bullet} \otimes \Gamma^h(\mathbb{S})_{\bullet} \otimes X)_n)$$

is the zero map. To prove this, we first observe that the source and target can be identified with $E[G]_*$ applied a finite colimit. It therefore suffices to show that the maps

$$E[G]_*(\overline{E}^{\otimes i} \otimes \overline{\mathbb{S}[G]}^{\otimes j}) \to E[G]_*(\overline{E}^{\otimes i} \otimes \overline{\mathbb{S}[G]}^{\otimes j-1})$$

and

$$E[G]_*(\overline{E}^{\otimes i} \otimes \overline{\mathbb{S}[G]}^{\otimes j}) \to E[G]_*(\overline{E}^{\otimes i-1} \otimes \overline{\mathbb{S}[G]}^{\otimes j})$$

are the zero maps, but this is the case because $E^{\otimes i} \otimes \mathbb{S}[G]^{\otimes \bullet}$ is an $\mathbb{S}[G]$ -Adams resolution for each i and $E^{\otimes \bullet} \otimes \mathbb{S}[G]^{\otimes j}$ is an E-Adams resolution for each j.

Corollary 3.3.5. The *E*-MASS for $G = C_2$ agrees with the *E*/*a*-based Adams spectral sequence from the E_2 -term onwards.

Proof. This follows from the observation that $Ca \simeq \mathbb{S}[C_2]$ and Proposition 3.3.3.

Corollary 3.3.6. If $X \in \text{Sp}_G$ is a ring spectrum, the *E*-MASS for X is a spectral sequence of algebras, and in general the *E*-MASS is a spectral sequence of modules over the *E*-MASS for S.

3.4. The MANSS and the C_2 -effective slice spectral sequence. We now introduce the main tool that we use this paper.

Definition 3.4.1. In the special case of Section (3.3) where $E = MU_p$,² we shall refer to the MU_p -MASS as the *p*-primary modified Adams–Novikov spectral sequence (MANSS)

$$\underline{MANSS} \underline{E}_{1}^{s,n+\diamond}(X) = \bigoplus_{a+b=s} \underline{\pi}_{n+\diamond}(MU_{p} \otimes \overline{MU}_{p}^{\otimes a} \otimes F(EG_{+}^{(b)}/EG_{+}^{(b-1)},X)) \implies \underline{\pi}_{\star}(X_{MU_{p}}^{h})$$

where $X_{MU_p}^h$ is defined as in (3.2.7).

The following corollaries follow from the evident generalization of Proposition 3.3.3 to the category Sp_{ip}^{BG} .

Corollary 3.4.2 (MANSS is MU_p/a -based ASS). The *p*-primary MANSS for $G = C_2$ agrees with the MU_p/a -based Adams spectral sequence from the E_2 -term onwards.

Proof. This follows from 3.3.5 by taking $E = MU_p$.

Corollary 3.4.3. If $X \in \text{Sp}_{G,ip}$ is a ring spectrum, the *p*-primary MANSS for X is a spectral sequence of algebras, and in general the *p*-primary MANSS is a spectral sequence of modules over the *p*-primary MANSS for \mathbb{S}_p .

Proof. This follows from the fact that $E[G] \simeq F(\Sigma^{\infty}G_+, E)$.

The following corollary of Proposition 3.3.3 begins to explain the motivation behind Definition 2.4.6.

²Here we work in the category $\operatorname{Sp}_{ip}^{BG}$ so that all instances of \otimes should be interpreted as $\otimes_{\operatorname{Sp}}$

Corollary 3.4.4. In the category $Sp_{G,ip}^{\text{Fil}}$ there is an equivalence

$$\mathbb{S}_{G,p}^{\operatorname{def}} \coloneqq \operatorname{D\acute{e}c}^{\operatorname{ev.post}}(MU_p[G]^{\otimes \bullet + 1}) \simeq \operatorname{D\acute{e}c}^{\operatorname{ev.post}}\Gamma_{MU_p}^h \mathbb{S}_p.$$

We already showed that the 2-primary MANSS for the group $G = C_2$ agrees with the MU_2/a -based Adams spectral sequence in Corollary 3.4.2. Therefore to compare the MANSS with the C_2 -ESSS it suffices to compare the MU_2/a -based Adams spectral sequence to the C_2 -ESSS. Since the MANSS is a spectral sequence in Borel G-spectra, it corresponds to an a-complete p-complete filtered object in the C_2 -equivariant category under the equivalence

$$(\operatorname{Sp}_{C_2,ip})^{\wedge}_a \simeq \operatorname{Sp}_{ip}^{BG}.$$

For the rest of this section we will work in the category $\text{Sp}_{C_2,ip}$, and objects of Sp_{ip}^{BG} will be regarded as *a*-complete objects of $\text{Sp}_{C_2,ip}$.

Lemma 3.4.5. Let $X \in \text{Sp}_{C_2,i2}$. For any $i \ge 1$, there is a natural equivalence

(3.4.6)
$$(\tau_{\geq n}X) \otimes (Ca)^{\otimes i} \to \tau_{\geq n}(X \otimes (Ca)^{\otimes i}).$$

Proof. Since $\tau_{\geq n}$ is closed under the operation $-\otimes Ca$, the existence of a map follows from the fact that $\tau_{\geq n}$ is part of the data of a *t*-structure on Sp^{BC2}.

For any C_2 -spectrum Z, we have

$$\pi_k^e(\tau_{\ge n}(Z \otimes Ca)) = \begin{cases} \pi_k^e(Z \otimes Ca), \ k \ge n \\ 0, \ \text{otherwise} \end{cases}, \text{ and } \underline{\pi}_k^{C_2}(\tau_{\ge n}(Z \otimes Ca)) = \begin{cases} \pi_k^e(Z), \ k \ge n \\ 0, \ \text{otherwise.} \end{cases}$$

By induction, we compute the right hand side of (3.4.6)

$$\underline{\pi}_k\Big(\tau_{\geq n}(X\otimes(Ca)^{\otimes i})\Big) = \begin{cases} \underline{\pi}_k((\tau_{\geq n}X)\otimes(Ca)^{\otimes i}), \ k \geq n = \begin{cases} \pi_k^e(X\otimes(Ca)^{\otimes i-1}), \ \text{at } C_2/C_2, \\ \pi_k^e(X\otimes(Ca)^{\otimes i}), \ \text{at } C_2/e, \end{cases} \\ 0, \ \text{otherwise.} \end{cases}$$

The Mackey functor valued homotopy of the left hand side equals

$$\underline{\pi}_{k}\Big((\tau_{\geq n}X)\otimes(Ca)^{\otimes i}\Big) = \begin{cases} \pi_{k}^{e}\Big((\tau_{\geq n}X\otimes(Ca)^{\otimes i-1}\Big), \text{ for } C_{2}/C_{2} = \begin{cases} \pi_{k}^{e}(X\otimes(Ca)^{\otimes i-1}), \ k \geq n\\ 0, \text{ otherwise}, \end{cases}\\ \pi_{k}^{e}\Big((\tau_{\geq n}X)\otimes(Ca)^{\otimes i}\Big), \text{ for } C_{2}/e = \begin{cases} \pi_{k}^{e}(X\otimes(Ca)^{\otimes i}), \ k \geq n\\ 0, \text{ otherwise}. \end{cases}\end{cases}$$

Since Mackey functor valued homotopy detects weak equivalences and these identifications are compatible with the Mackey functor structure maps, the result follows. \Box

Theorem 3.4.7 (Comparison of C_2 -ESSS and MU_2/a -based ASS). Let $X \in \text{Sp}_{C_2,i2}$. There is a equivalence of filtered objects in $\text{Sp}_{C_2,i2}$

$$\left(\operatorname{D\acute{e}c}^{\operatorname{ev.slice}}(MU_{\mathbb{R},2}^{\otimes \bullet+1} \otimes X)\right)_{a}^{\wedge} \simeq \operatorname{D\acute{e}c}^{\operatorname{ev.post}}\left(\left((MU_{2}/a)^{\otimes \bullet+1} \otimes X\right)\right).$$

Proof. This follows by the composite of natural equivalences

$$\left(\lim_{i} P_{\geq 2n} (MU_{\mathbb{R},2}^{\otimes i+1} \otimes X)\right)_{a}^{\wedge} \stackrel{\text{commute limits}}{\simeq} \lim_{i} \left(\left(P_{\geq 2n} (MU_{\mathbb{R},2}^{\otimes i+1} \otimes X)\right)_{a}^{\wedge} \right) \right)^{\wedge} \\ \stackrel{\text{Lem.2.2.4+Lem.2.2.6}}{\simeq} \lim_{i} \left(\left(\tau_{\geq 2n} (MU_{\mathbb{R},2}^{\otimes i+1} \otimes X) \otimes Ca^{\otimes i+1} \right) \right)^{\wedge} \\ \stackrel{\text{def}}{\simeq} \lim_{j} \lim_{i} \left(\tau_{\geq 2n} (MU_{\mathbb{R},2}^{\otimes i+1} \otimes X) \otimes Ca^{\otimes j+1} \right) \\ \stackrel{\text{diagonal}}{\simeq} \lim_{i} \left(\tau_{\geq 2n} (MU_{\mathbb{R},2}^{\otimes i+1} \otimes X) \otimes Ca^{\otimes i+1} \right) \\ \stackrel{\text{Lem.3.4.5}}{\simeq} \lim_{i} \left(\tau_{\geq 2n} ((MU_{\mathbb{R},2}/a)^{\otimes i+1} \otimes X) \right) \\ \stackrel{\text{shearing}}{\simeq} \lim_{i} \left(\tau_{\geq 2n} ((MU_{\mathbb{R},2}/a)^{\otimes i+1} \otimes X) \right).$$

Here the shearing isomorphism is the equivalence

$$MU_{\mathbb{R},2}/a \simeq MU_2/a$$

which follows from the fact that MU_2 and $MU_{\mathbb{R},2}$ have equivalent underlying ring spectra. The proof is then complete by Corollary 3.4.2.

Associated to the *a*-completion of the C_2 -effective slice filtration is the *a*-completed C_2 effective slice spectral sequence

$${}^{C_2\text{-ESSS}}\underline{E}_1^{f,n+\diamond}(X)_a^{\wedge} = \underline{\pi}_{n+\diamond}(\operatorname{Gr}^f\Gamma_{\mathbb{R}}X) \implies \underline{\pi}_{n+\diamond}(X_{MU}^h).$$

The following corollary follows from (2.3.5) (note that the hypothesis that $MU_*(X^e)$ is even implies that the MANSS has no nontrivial d_r differentials for r even).

Corollary 3.4.8. Suppose that $(MU_2)_*(X^e)$ is concentrated in even degrees. Then for $r \ge 1$, there is a (page shifting) isomorphism of spectral sequences

$$\underline{E}_{2r+1}^{s,n+\diamond}(X) \cong \underline{E}_r^{n+s}(X)^{\wedge}_{r-s}.$$

We can now give the first evidence for why the category of artificial complete motivic spectra is a useful deformation of Borel G-spectra.

Theorem 3.4.9 (MANSS Galois reconstruction). There are equivalences

$$\widehat{\mathrm{SH}}_{C_2,2}^{\mathrm{art}} \coloneqq \mathrm{Mod}((\mathrm{Sp}_{i2}^{BC_2})^{\mathrm{Fil}}; \mathbb{S}_{G,2}^{\mathrm{def}}) \simeq \mathrm{Mod}((\mathrm{Sp}_{ip}^{BG})^{\mathrm{Fil}}; \mathrm{D\acute{e}c}^{\mathrm{ev.post}}\Gamma^h_{MU_2} \mathbb{S}_2) \simeq (\mathrm{SH}(\mathbb{R})_{i2}^{\mathrm{AT}})_a^{\wedge}.$$

Proof. This follows from Proposition 2.2.7, Theorem 3.4.7, and Theorem 3.1.1.

4. Computing with the MANSS

We compute the E_2 -term of (*p*-primary) MANSS (see Definition 3.4.1) using double complex spectral sequences in Section 4.1. We then specialize to the case $G = C_2$, p = 2, $G/H = C_2/C_2$ and write

$$^{\text{MANSS}}E_r(X) := {}^{\text{MANSS}}\underline{E}_r(X)(C_2/C_2)$$

We express the E_2 -term using $A^{s,2t} := \operatorname{Ext}^{s,2t}((MU_p)_*X^e)$ in two cases:

(1) when X is $MU_{\mathbb{R}}$ -projective (Section 4.2), and

(2) when X has trivial C_2 -action and $(MU_2)_* X^e$ is torsion free (Section 4.3).

In the first case, we show the MANSS relates to the ANSS and the C_2 -effective SS; in the second case, by studying a motivic version of MANSS, we relate it to the Atiyah–Hirzebruch spectral sequence.

Assume $x \in A^{s,2t}$ (see Notation 4.2.3). In this section, we introduce the following notations.

G. Angelini-Knoll, M. Behrens, E. Belmont, and H. J. Kong

Notation	Degree	Definition	From
\overline{x}	$(t, t\rho - s)$	Notation 4.2.9	C_2 -effective SS
\overline{ux}	$(t, t\rho - s + 1 - \sigma)$	Notation 4.2.9	C_2 -effective SS
$\langle x \rangle$	$(s, t\rho - s)$	Notation 4.2.3	double complex SS
$\langle ux \rangle$	$(s, t\rho - s)$	Notation 4.2.3	double complex SS
[x]	(s, 2t - s)	Notation 4.3.6	compare AHSS
[ux]	$(s, 2t - s + 1 - \sigma, t)$	Notation 4.3.6	compare AHSS
[x]	(s, 2t - s, t, t)	Proposition 4.4.9	motivic MANSS
[ux]	$(s, 2t - s + 1 - \sigma, t)$	Proposition 4.4.9	motivic MANSS

4.1. The E_2 -term of the MANSS. Let X be an object of $\operatorname{Sp}_{ip}^{BG}$. Recall from Section 3.2 that the E_1 -term of the *p*-primary MANSS is

$$\overset{\text{MANSS}}{\underline{E}}_{1}^{f,V}(X) = \bigoplus_{a+b=f} \underline{\pi}_{V} Z_{a,b}$$

where

$$Z_{a,b} = F(EG^{(a)}/EG^{(a-1)}, X) \otimes MU_p \otimes \overline{MU}_p^{\otimes b}.$$

In other words, the E_1 -term ($\underline{E}_1^{\star,*}(X), d_1^{\text{MANSS}}$) is the total complex of the double complex $\bigoplus_{a,b} \underline{\pi}_{\star}(Z_{a,b})$ whose horizontal differential $d_1^{\text{ANSS}}: \underline{\pi}_{\star}Z_{a,b} \to \underline{\pi}_{\star}Z_{a,b+1}$ is the Adams–Novikov d_1 and vertical differential $d_1^{\text{hfpss}}: \underline{\pi}_{\star} Z_{a,b} \to \underline{\pi}_{\star} Z_{a+1,b}$ is the homotopy fixed point spectral sequence d_1 . We wish to compute the cohomology $E_2^{\star,*}$ of the total complex. There are two double complex spectral sequences

(4.1.1)
$${}_{I}\underline{E}_{2}^{a,b,\star}(X) = H^{b}(H^{a}(\bigoplus_{i,j} \underline{\pi}_{\star}(Z_{i,j}), d_{1}^{hfpss}), d_{1}^{ANSS}) \Longrightarrow {}^{MANSS}\underline{E}_{2}^{a+b,\star}(X)$$

(4.1.2)
$$_{II}\underline{E}_{2}^{a,b,\star}(X) = H^{a}(H^{b}(\bigoplus_{i,j}\underline{\pi}_{\star}(Z_{i,j}), d_{1}^{\text{ANSS}}), d_{1}^{\text{hfpss}}) \Longrightarrow \overset{\text{MANSS}}{\underline{E}}_{2}^{a+b,\star}(X)$$

which come from filtering the double complex by i and j, respectively [McC01, §2.4]. We use these spectral sequences to describe the E_2 -term of the MANSS.

Proposition 4.1.3. There is an identification

Ν

$$\underline{E}_{2}^{a,b,n+\diamond}(X)(G/H) = \operatorname{Ext}^{b,a+b+n}(H^{a}(H;(MU_{p})_{*}(\mathbb{S}^{-i_{H}^{*}\diamond}\otimes X)))$$

In particular, the value of the E_2 -term of the MANSS at G/e is

$$\underline{AANSS}\underline{E}_{2}^{b,V}(X)(G/e) = \operatorname{Ext}^{b,b+|V|}((MU_{p})_{*}(X^{e})).$$

Proof. Since $EG^{(i)}/EG^{(i-1)}$ is dualizable, $(MU_p)_*\overline{MU_p}$ is flat over $(MU_p)_*$, and MU_p and MU_p have trivial G-action, we have:

$$\underline{\pi}_{*+\diamond}(Z_{i,j}) \cong \underline{\pi}_{*}(F(EG^{(i)}/EG^{(i-1)}, \mathbb{S}^{\diamond} \otimes X) \otimes MU_{p} \otimes \overline{MU_{p}}^{\otimes j})$$

$$(4.1.4) \cong \underline{\pi}_{*}(F(EG^{(i)}/EG^{(i-1)}, \mathbb{S}^{\diamond} \otimes X \otimes MU_{p})) \otimes_{(MU_{p})_{*}} (MU_{p})_{*} \overline{MU_{p}})^{\otimes_{(MU_{p})_{*}j}}.$$

At the G-set G/H, this is $(MU_p)_* \overline{MU_p}^{\otimes j}$ tensored with the E_2 -term of a homotopy fixed points spectral sequence, and so we have

$$H^{a}(\underline{\pi}_{*+\diamond}(Z_{\bullet,j})(G/H), d_{1}^{\mathrm{hfpss}}) \cong H^{a}(H; MU_{*}(\mathbb{S}^{-i_{H}^{*}\diamond} \otimes X)) \otimes_{(MU_{p})_{*}} (MU_{p})_{*} \overline{MU_{p}^{\otimes_{(MU_{p})_{*}j}}}.$$

This is isomorphic to the cobar complex

$$C^{j,*}_{(MU_p)_*MU_p}(H^a(H; (MU_p)_*(\mathbb{S}^{-i_H^*} \otimes X)))$$
23

for the $(MU_p)_*MU_p$ -comodule

$$H^{a}(H;(MU_{p})_{*}(\mathbb{S}^{-i_{H}^{*}\diamond}\otimes X))$$

and we obtain $I\underline{E}_2^{a,b,*+\diamond}(G/H)$ by applying d_1^{ANSS} . In the H = e case, the double complex spectral sequence collapses at $I\underline{E}_2$ because it is concentrated in degree a = 0.

Corollary 4.1.5. The restriction to the trivial subgroup $e \leq G$ gives a map of spectral sequences from the value of the MANSS at G/H to the classical Adams–Novikov spectral sequence. The induced map on E_2 -terms is

$$\underline{HANSS} \underline{E}_{2}^{b,V}(X)(G/H) \longrightarrow \operatorname{Ext}_{(MU_{p})_{*}MU_{p}}^{b,b+|V|}((MU_{p})_{*}(X^{e})).$$

Proposition 4.1.6. There is an identification

$$H \underline{E}_{2}^{a,b,n+\diamond}(X)(G/H) \cong H^{a}(H; \operatorname{Ext}^{b,a+b+n}((MU_{p})_{*}(\mathbb{S}^{-\diamond} \otimes X))).$$

Moreover, when X^e is MU_p -nilpotent complete and the E_2 -term of the Adams–Novikov spectral sequence for X^e is concentrated on the zero line, then this spectral sequence collapses and we can identify the E_2 -term of the MANSS with

$$\underline{E}_2^{a,n+\diamond}(G/H) = H^a(H;\pi_n(\Sigma^{-\diamond}X)).$$

Proof. Starting with (4.1.4), we have

$$II \underline{E}_{1}^{a,b,n+\diamond}(X)(G/H) \cong \operatorname{Ext}^{b,a+b+n}(\underline{\pi}_{*}(F(EG^{(a)}/EG^{(a-1)}, \mathbb{S}^{-\diamond} \otimes X \otimes MU_{p})(G/H)))$$

$$\cong (\operatorname{Ext}^{b,a+b+n}(\pi_{*}(F(EG^{(a)}/EG^{(a-1)}, \mathbb{S}^{-\diamond} \otimes X \otimes MU_{p})^{H})))$$

$$\cong (\operatorname{Ext}^{b,a+b+n}(\pi_{*}((\mathbb{S}[G] \otimes \overline{\mathbb{S}[G]}^{\otimes a} \otimes \mathbb{S}^{-\diamond} \otimes X \otimes MU_{p}))))^{H}$$

$$\cong (\operatorname{Ext}^{b,a+b+n}((MU_{p})_{*}((\mathbb{S}[G] \otimes \overline{\mathbb{S}[G]}^{\otimes a}) \otimes_{(MU_{p})_{*}} \pi_{*}(\mathbb{S}^{-\diamond} \otimes X \otimes MU_{p}))))^{H}$$

$$\cong (\operatorname{Ext}^{b,a+b+n}(H\mathbb{Z}_{*}(\mathbb{S}[G] \otimes \overline{\mathbb{S}[G]}^{\otimes a}) \otimes (MU_{p})_{*}(\mathbb{S}^{-\diamond} \otimes X)))^{H}$$

$$\cong (H\mathbb{Z}_{*}(\mathbb{S}[G] \otimes \overline{\mathbb{S}[G]}^{\otimes a}) \otimes \operatorname{Ext}^{b,a+b+n}((MU_{p})_{*}(\mathbb{S}^{-\diamond} \otimes X))))^{H}$$

where we use the description in Lemma 3.3.1 of the dual of the fact that $F(EG^{(a)}/EG^{(a-1)}, \mathbb{S}) \simeq \mathbb{S}[G] \otimes \overline{\mathbb{S}[G]}^{\otimes a}$. This is precisely the cellular chain complex computing the cohomology of BH with local coefficients in the $\mathbb{Z}[H]$ -module given by the Ext group. Thus we have

$$_{II}\underline{E}_{2}^{a,b,n+\diamond}(G/H) \cong H^{a}(H; \operatorname{Ext}^{b,a+b+n}((MU_{p})_{*}(\mathbb{S}^{-\diamond} \otimes X))).$$

Corollary 4.1.7. The MANSS is independent of a choice of Adams–Novikov resolution. The MANSS is also independent of the choice of S[G]-Adams resolution.

Proof. Given an Adams–Novikov resolution L_{\bullet} of X^e and a map of Adams–Novikov resolutions $\overline{MU}_p^{\bullet} \otimes X^e \to L_{\bullet}$, Proposition 4.1.6 produces a map of double complex spectral sequences that is an isomorphism on the E_2 -terms and satisfies the conditions of [Boa99, Theorem 8.3]. The second statement follows from the same argument using Proposition 4.1.3 and noting that tower (3.2.1) may be regarded as a S[G]–Adams resolution of X. □

The reader will notice that the E_2 -terms in Propositions 4.1.3 and 4.1.6 compare to the E_2 -term of the homotopy fixed point spectral sequence. The following proposition gives a relationship between the *E*-MASS and the homotopy fixed point spectral sequence.

Proposition 4.1.8. For a *G*-spectrum *X*, let ${}^{hfpss}\underline{E}_r$ denote the E_r -term of the homotopy fixed points spectral sequence, which is obtained by applying $\underline{\pi}_{\star}$ to the filtration (3.2.1). There is a map of spectral sequences given on E_r -terms by

$$\underline{E}_{r}^{s,V}(X) \xrightarrow{E-\text{MASS}} \underline{E}_{r}^{s,V}(X)$$

Proof. For each $a, b \in \mathbb{Z}$ satisfying $a + b = n \ge 0$, there is a canonical map

$$F(EG_+ \otimes \widetilde{E}G^{(a)}, X) \otimes \overline{E}^b \longrightarrow \operatorname{colim}_{a+b=n} F(EG_+ \otimes \widetilde{E}G^{(a)}, X) \otimes \overline{E}^{\otimes b}.$$

Fixing b = 0, this gives a map of filtered spectra producing the map of spectral sequences. \Box

4.2. The case of $MU_{\mathbb{R}}$ -projective spectra. We now focus on the case of $G = C_2$ and p = 2, and let MANSS = (X) = MANSS = (X)(G + G)

$$^{\text{IANSS}}E_r(X) := {}^{\text{MANSS}}\underline{E}_r(X)(C_2/C_2).$$

If $X \in \text{Sp}_{C_2,i2}$ is $MU_{\mathbb{R}}$ -projective (Definition 3.1.3), then Theorem 3.1.4 gives a complete computation of the E_1 -term of the C_2 -ESSS. However, this does not immediately carry over to a description of the E_1 -term of the *a*-completed C_2 -ESSS, which by Corollary 3.4.8 is the E_2 -term of the MANSS. In this section, we will instead argue that the second double complex spectral sequence

$$_{II}E_2^{a,b,n-j+j\sigma}(X) = H^a(C_2; \operatorname{Ext}^{b,b+n}((MU_2)_*(\mathbb{S}^{j-j\sigma} \otimes X))) \implies \operatorname{MANSS} E_2^{a+b,n-j+j\sigma}(X)$$

for such X collapses. This method of computing the E_2 -term of the MANSS has the added benefit of giving explicit meaning to the generators.

It is noteworthy that the property of being $MU_{\mathbb{R}}$ -projective is not preserved by *a*-completion. Thus we are in the unusual situation of computing the E_2 -term of the MANSS, an inherently Borel equivariant construction, by assuming the Borel equivariant spectrum has a genuine equivariant lift which is $MU_{\mathbb{R}}$ -projective.

Remark 4.2.1. A similar analysis to that performed in this section can be used to prove the other double complex spectral sequence ${}_{I}E_{r}^{*,*,*}(X)$ also collapses at E_{2} for $MU_{\mathbb{R}}$ -projective X, but the resulting presentation of the E_{2} -term of the MANSS will be less useful for us.

We first identify the E_2 -term

$${}_{II}E_2^{a,b,n-j+j\sigma}(X) = H^a(C_2; \operatorname{Ext}^{b,b+n}((MU_2)_*(\mathbb{S}^{j-j\sigma} \otimes X)))$$

of the second double complex spectral sequence in the $MU_{\mathbb{R}}$ -projective case. The first task is to identify the action of C_2 .

Lemma 4.2.2. Suppose $X \in \text{Sp}_{C_{2},i2}$ is $MU_{\mathbb{R}}$ -projective. The action of the generator of C_{2} on

$$\operatorname{Ext}^{s,2t}((MU_2)_*(\mathbb{S}^{j-j\sigma}\otimes X^e))$$

is multiplication by $(-1)^{t+j}$.

Proof. This follows directly from the $MU_{\mathbb{R}}$ -projective condition. To see this, note that C_2 acts on the underlying spectrum MU_2 of $MU_{\mathbb{R},2}$ by complex conjugation. Therefore, the generator of C_2 acts on $\pi_{2f}(MU_{\mathbb{R},2}^e \otimes \mathbb{S}^{j-j\sigma})$ by $(-1)^{t+j}$ (see [HHR16, Rmk. 5.14]).

Notation 4.2.3. For an abelian group A, we let

$$A[2] := \{ x \in A : 2x = 0 \}.$$

Let $X \in \text{Sp}_{C_2,i2}$ be $MU_{\mathbb{R}}$ -projective, and define

$$A^{s,2t} = \operatorname{Ext}^{s,2t}((MU_2)_*X^e).$$

Then by Lemma 4.2.2 we have

(4.2.4)
$$A^{s,2t} \cong H^0(C_2; \operatorname{Ext}^{s,2t}((MU_2)_*(\mathbb{S}^{t-t\sigma} \otimes X^e))) = {}_{II}E_2^{0,s,t\rho-s}(X).$$

Given $x \in A^{s,2t}$, we denote its image under (4.2.4) by $\langle x \rangle$. By Lemma 4.2.2, there is also an isomorphism

(4.2.5)
$$A^{s,2t}[2] \cong H^0(C_2; \operatorname{Ext}^{s,2t}((MU_2)_*(\mathbb{S}^{(t+1)-(t+1)\sigma} \otimes X^e)))$$
$$= {}_{II}E_2^{0,s,t\rho-s+1-\sigma}(X).$$

Given $x \in A^{s,2t}[2]$, we denote its image under (4.2.5) by $\langle ux \rangle$.

Recall (see [HK01]) that there is a subring (the "positive cone")

$$\mathbb{Z}[u^2, a]/(2a) \subset \pi_{\star}^{C_2} H \underline{\mathbb{Z}}.$$

with $|a| = -\sigma$ and $|u| = 1 - \sigma$. The images of the elements a and u^2 in $\pi_{\star} H \underline{\mathbb{Z}}_a^{\wedge}$ induce isomorphisms

$$\pi_{*+\diamond}^{C_2} H \underline{\mathbb{Z}}_a^{\wedge} \cong H^{-*}(\mathbb{S}_{hC_2}^{\diamond}) \cong H^{-*}(C_2; H^0(\mathbb{S}^{\diamond})) \cong \mathbb{Z}[u^{\pm 2}, a]/(2a).$$

There is an isomorphism

$$\mathbb{Z}_{2}[u^{2}, a]/(2a) \cong H^{*}(C_{2}; \operatorname{Ext}^{0,0}_{(MU_{2})_{*}MU_{2}}((MU_{2})_{*}, (MU_{2})_{*}(\mathbb{S}_{2}^{-\diamond}))) = {}_{II}E_{2}^{*,0,\diamond}(\mathbb{S}_{2}).$$

The elements u^2 and a detect the corresponding elements in

$$\pi_{*+\diamond}^{C_2} H \underline{\mathbb{Z}}_2 \cong \operatorname{Ext}^{0,0}((MU_2)_* \otimes \pi_{*+\diamond}^{C_2} H \underline{\mathbb{Z}}) = {}^{C_2 - \operatorname{ESSS}} E_1^{0,*+\diamond}(\mathbb{S}_2)_a^{\wedge} \cong {}^{\operatorname{MANSS}} E_3^{-*,*+\diamond}(\mathbb{S}_2)$$

in the second double complex spectral sequence. Therefore these elements are permanent cycles in the double complex spectral sequence, and the second double complex spectral sequence is a spectral sequence of modules over $\mathbb{Z}[u^{\pm 2}, a]/(2a)$. The following proposition is an elementary immediate consequence of Lemma 4.2.2 using the well-known group cohomology of C_2 with arbitrary coefficients.

Proposition 4.2.6. Let $X \in \text{Sp}_{C_2,i2}$ be $MU_{\mathbb{R}}$ -projective. As before, define

$$A^{s,2t} = \operatorname{Ext}^{s,2t}((MU_2)_*X^e).$$

Then under the isomorphisms (4.2.4) and (4.2.5) we have

$${}_{II}E_2^{*,*,\star}(X) \cong \left(A^{*,*} \otimes \mathbb{Z}[u^{\pm 2},a]/(2a)\right) \oplus \left(A^{*,*}[2] \otimes \mathbb{F}_2[u^{\pm 2},a]\right).$$

Corollary 4.2.7. For $MU_{\mathbb{R}}$ -projective X, every element of $_{II}E_2^{*,*,*}(X)$ is of the form $u^{2i}a^j\langle x\rangle$ or $u^{2i}a^j\langle ux\rangle$ for $i \in \mathbb{Z}, j \ge 0$, and $x \in A^{*,*}$ (respectively $x \in A^{*,*}[2]$).

Our strategy for proving that the second double complex spectral sequence collapses will be to produce elements in ${}^{\text{MANSS}}E_2^{*,\star}(X)$ which the generators in Corollary 4.2.7 converge to. To do this, we will require the following "universal coefficient theorem".

Lemma 4.2.8. For any torsion-free $(MU_2)_*MU_2$ -comodule N, there is a non-canonically split short exact sequence

$$0 \longrightarrow \operatorname{Ext}^{*,*}(N) \otimes \underbrace{\mathbb{F}_2}_{\alpha} \xrightarrow{p} \operatorname{Ext}^{*,*}(N/2) \xrightarrow{\iota}_{\beta} \operatorname{Tor}_1^{\mathbb{Z}}(\operatorname{Ext}^{*+1,*}(N), \mathbb{F}_2) \longrightarrow 0.$$

If N is an algebra in the category of comodules, then the short exact sequence is a short exact sequence of modules over $\operatorname{Ext}^{*,*}(N)$, and the splitting is a splitting in the category of $\operatorname{Ext}^{*,*}(N)$ -modules.

Proof. The short exact sequence is easily deduced from the long exact sequence in Ext coming from the short exact sequence

$$0 \to N \xrightarrow{\cdot 2} N \to N/2 \to 0.$$

The splitting map p comes from choosing a splitting p' of the short exact sequence of free abelian groups

$$0 \longrightarrow Z^{*,*} \xrightarrow{\downarrow \cdots \qquad } C^{*,*}(N) \xrightarrow{d} B^{*+1,*} \longrightarrow 0$$

where $C^{*,*}(N)$ is the cobar complex for N, and $Z^{*,*}$ and $B^{*,*}$ are the subcomplexes of cocycles and coboundaries, respectively. The composite

$$p'': C^{*,*}(N) \to Z^{*,*} \to \operatorname{Ext}^{*,*}(N)$$

is then verified to be a map of cochain complexes, where the target is given zero differential. The splitting map p is then obtained by applying $H^{*,*}(-)$ to the map of cochain complexes

$$C^{*,*}(N) \otimes \mathbb{F}_2 \xrightarrow{p'' \otimes 1} \operatorname{Ext}^{*,*}(N) \otimes \mathbb{F}_2.$$

Now assume that N is a comodule algebra. Then the maps α and β are easily checked to be maps of modules over $\operatorname{Ext}^{*,*}(N)$. Because the module structure on $\operatorname{Ext}^{*,*}(N) \otimes \mathbb{F}_2$ and $\operatorname{Ext}^{*,*}(N/2)$ come from the maps of rings

$$\operatorname{Ext}^{*,*}(N) \to \operatorname{Ext}^{*,*}(N) \otimes \mathbb{F}_2 \xrightarrow{\alpha} \operatorname{Ext}^{*,*}(N/2),$$

the fact that p is a map of $\operatorname{Ext}^{*,*}(N)$ -modules follows from the commutativity of the following diagram

Notation 4.2.9. Suppose that $X \in \text{Sp}_{C_2,i2}$ is $MU_{\mathbb{R}}$ -projective, and choose an element $x \in \text{Ext}^{s,2t}((MU_2)_*X^e)$. Let \overline{x} denote the image of this element under the map

(4.2.10)
$$\operatorname{Ext}^{s,2t}((MU_2)_*X^e \otimes \pi_0^{C_2}H\underline{\mathbb{Z}}) \to {}^{C_2\operatorname{-ESSS}}E_1^{t,t\rho-s}(X)$$

induced by Theorem 3.1.4. By abuse of notation, we will also use \overline{x} to denote the image of this element under the map

(4.2.11)
$$C_2\text{-ESSS} E_1^{t,t\rho-s}(X) \to C_2\text{-ESSS} E_1^{t,t\rho-s}(X)_a^{\wedge} \cong {}^{\text{MANSS}} E_2^{s,t\rho-s}(X).$$

Now choose a splitting ι of the short exact sequence

(4.2.12)
$$0 \longrightarrow A^{s-1,2t} \otimes \mathbb{F}_2 \longrightarrow \operatorname{Ext}^{s-1,2t}((MU_2)_*(X^e)/2) \longrightarrow \operatorname{Tor}_1^{\mathbb{Z}}(A^{s,2t},\mathbb{F}_2) \longrightarrow 0$$

as in Lemma 4.2.8 (note $(MU_2)_*X^e$ is torsion-free by the $MU_{\mathbb{R}}$ -projective hypothesis). For

$$x \in A^{s,2t}[2] = \operatorname{Tor}_{1}^{\mathbb{Z}}(A^{s,2t}, \mathbb{F}_{2}),$$

27

let \overline{ux} denote the image of $\iota(x)$ under the maps $\operatorname{Ext}^{s-1,2t}((MU_2)_*X^e \otimes \mathbb{F}_2\{a\}) = \operatorname{Ext}^{s-1,2t}((MU_2)_*X^e \otimes \pi_{-\sigma}H\underline{\mathbb{Z}}) \to {}^{C_2 - ESSS}E_1^{t,t\rho-s+1-\sigma}(X)$ and $C_2 - \operatorname{ESSS}E_1^{t,t\rho-s+1-\sigma}(X) \to {}^{C_2 - \operatorname{ESSS}}E_1^{t,t\rho-s+1-\sigma}(X)_a^{\wedge} \cong {}^{\operatorname{MANSS}}E_2^{s,t\rho-s+1-\sigma}(X).$

 E_1 $(X) \rightarrow E_1$ $(X)_a \stackrel{\text{\tiny def}}{=} E_2$ (X). **Remark 4.2.13.** Given $x \in \text{Ext}^{s,2t}((MU_2)_*X^e)$ of order 2, the element \overline{ux} defined above is not unique, as it depends on a choice of life. However, the chort exect accuracy (4,2,12) implies

not unique, as it depends on a choice of lift. However, the short exact sequence (4.2.12) implies that any two lifts \overline{ux} and \overline{ux}' differ by an element of the form $a\overline{y}$ for $y \in \operatorname{Ext}^{s-1,2t}((MU_2)_*X^e)$.

Remark 4.2.14. If X is a ring spectrum, the maps (4.2.10) and (4.2.11) are maps of rings, and therefore we have

$$\overline{x} \cdot \overline{y} = \overline{xy}$$

Moreover the map ι in (4.2.12) is a map of modules over $A^{*,*}$ by Lemma 4.2.8. It follows that we have

$$\overline{x} \cdot \overline{uy} = \overline{uxy}.$$

Proposition 4.2.15. Let $X \in \operatorname{Sp}_{i2}^{BC_2}$ be $MU_{\mathbb{R}}$ -projective, and $A^{*,*}$ as above. For each $x \in A^{*,*}$, the element \overline{x} is detected in the second double complex spectral sequence by the element $\langle x \rangle$, and for each $x \in A^{*,*}[2]$, the element \overline{ux} is detected by $\langle ux \rangle$. Therefore, the double complex spectral sequence collapses, and every element of $\operatorname{MANSS} E_2^{*,*}(X)$ is of the form $u^{2j}a^i\overline{x}$ or $u^{2j}a^i\overline{ux}$ for $j \in \mathbb{Z}$, $i \geq 0$, and x as above.

Proof. Using the fact that there is a C_2 -equivariant isomorphism $(MU_2)_*((X \otimes Ca)^e) \cong (MU_2)_*(X^e[C_2])$, the E_2 -term of the second double complex spectral sequence for $X \otimes Ca$ is given by

$$H^{\ell}(C_2; \operatorname{Ext}^{*,*}((MU_2)_*(\mathbb{S}^{-\diamond} \otimes (X \otimes Ca)^e))) \cong \begin{cases} A^{*,*} \otimes \mathbb{Z}[u^{\pm}], & \ell = 0, \\ 0, & \ell > 0. \end{cases}$$

Here we use the powers of u as a placeholder for the fact that the computation is independent of \diamond , but this also reflects the module structure over $\mathbb{Z}[u^{\pm 2}, a]/(2a)$ (where a acts as zero). Therefore, the second double complex spectral sequence for $X \otimes Ca$ collapses. Consider the map of second double complex spectral sequences

induced by the map

 $X \to X \otimes Ca.$

Under this map, for each element $x \in A^{s,2t}$, the elements $\langle x \rangle$ and \overline{x} map to $u^{-t}x$, and for each element $x \in A^{s,2t}[2]$, the elements $\langle ux \rangle$ and \overline{ux} map to $u^{-t+1}x$. Note that by Remark 4.2.13, this is independent of the choice of \overline{ux} , because a acts trivially on the second double complex spectral sequence for $X \otimes Ca$. In particular, the map

$$_{II}E_2^{0,*,\star}(X) \to _{II}E_2^{0,*,\star}(X \otimes Ca)$$

is an injection. We may therefore deduce that the elements $\langle x \rangle$ detect \overline{x} and the elements $\langle ux \rangle$ detect \overline{ux} in the double complex spectral sequence for X. It follows that the elements $\langle x \rangle$ and $\langle ux \rangle$ in $_{II}E_2^{*,*,*}(X)$ are all permanent cycles in the second double complex spectral sequence, and thus, by Corollary 4.2.7, the double complex spectral sequence $\{_{II}E_r^{*,*,*}(X)\}$ collapses at the E_2 -term.

4.3. The case of a trivial C_2 -action. In this section we continue to assume that $G = C_2$ and p = 2. Instead of assuming that $X \in \operatorname{Sp}_{i2}^{BC_2}$ is $MU_{\mathbb{R}}$ -projective, we simply assume that Xis a Borel C_2 -spectrum with $(MU_2)_*X$ torsion-free, but we make the additional assumption that the action of C_2 on X is *trivial*.

The next lemma says that the C_2/C_2 level of the MANSS, which arises from applying $\pi_{*,*}^{C_2}$ to the filtered Borel C_2 -spectrum $\Gamma_{MU_2}^h$, can also be viewed as a collection of non-equivariant spectral sequences, one for each weight j. In Section 4.4, we will define a \mathbb{C} -motivic analogue of the MANSS which generalizes this C_2/C_2 level version.

Lemma 4.3.1. Suppose $X \in \operatorname{Sp}_{i2}^{BC_2}$ has trivial action. Fix $j \in \mathbb{Z}$ and let

(4.3.2)
$$\Gamma_{MU_2}^{hC_2}(X)[j]_s \coloneqq \begin{cases} X^{P_j^{\infty}} & \text{if } s \le 0\\ \underset{a+b\ge s}{\text{colim}} \left(\overline{MU_2}^{\otimes a} \otimes X\right)^{P_{j+b}^{\infty}} & \text{if } s > 0. \end{cases}$$

The spectral sequence associated to this filtered object agrees with the C_2/C_2 -level of the MANSS starting in the E_2 -term.

This agrees on the level of filtrations with the C_2 -fixed points of a variant of the MANSS which uses a different filtration of EC_2 , namely the filtration of $S^{\infty\sigma} \simeq EC_2$ coming from $S^{k\sigma}$ rather than the skeletal filtration of the bar complex model of EC_2 ; this is reflected in Lemma 4.3.1. Nonetheless, as in Proposition 3.3.3, the resulting spectral sequences agree starting on the E_2 -term.

Remark 4.3.3. The reader may wonder why the tensor with \overline{MU}_2 happens inside the parentheses in (4.3.2), when this tensoring happens outside the parentheses in (3.2.4). However, the tensor in (3.2.4) is taken in the category of Borel *G*-spectra, and therefore homotopy fixed points are taken after taking the tensor.

Next we identify the associated graded of (4.3.2) to give a new description of the C_2/C_2 level of the MANSS E_1 term (see Definition 3.4.1).

Lemma 4.3.4. There is an isomorphism

$${}^{\text{MANSS}}E_1^{s,2t-s+j\sigma}(X) \cong \bigoplus_{a+b=s} C^b_{\text{cell}}(\Sigma^{-j}P_j^{\infty};\mathbb{Z}) \otimes C^{a,2t+j}_{(MU_2)_*MU_2}((MU_2)_*X).$$

where $C^{*,*}_{(MU_2)_*MU_2}$ denotes the reduced cobar complex and C^*_{cell} denotes the cellular cochain complex.

Thus, the MANSS E_1 -term is the total complex of the double complex

$$C^{\bullet}_{\operatorname{cell}}(\Sigma^{-j}P_j^{\infty};\mathbb{Z}) \otimes C^{\bullet,2*+j}_{(MU_2)_*MU_2}((MU_2)_*X)).$$

Intuitively, the MANSS is a combination of the ANSS for X and the Atiyah–Hirzebruch spectral sequence (AHSS) for $X^{P_j^{\infty}}$. The goal of this subsection and the next is to make this intuition precise.

We have already related the MANSS to the ANSS in Corollary 4.1.5. In this subsection we will show that the first double complex spectral sequence always collapses in the case at hand, giving us a computation of the E_2 -term of the MANSS which has an Atiyah–Hirzebruch flavor to it. In the next subsection we will then add some additional hypotheses on X which will allow us to relate higher differentials in the MANSS to differentials in the ANSS for X and the algebraic AHSS for Ext^{*,*}($(MU_2)_*(X^{P_j^{\infty}})$).

As in the previous subsection, for brevity we adopt the notation

$$A^{s,2t} \coloneqq \operatorname{Ext}^{s,2t}((MU_2)_*X^e).$$
29

Because both the cobar complex and the cellular chain complex are torsion-free, we have a non-canonically split Künneth short exact sequence (4.3.5)

$$0 \to \bigoplus_{a+b=s} A^{a,2t+j} \otimes H^b(\Sigma^{-j}P_j^{\infty}) \xrightarrow{\alpha} \operatorname{MANSS} E_2^{s,2t-s+j\sigma} \xrightarrow{\beta} \bigoplus_{a+b=s+1} \operatorname{Tor}_1^{\mathbb{Z}}(A^{a,2t+j}, H^b(\Sigma^{-j}P_j^{\infty})) \to 0.$$

Notation 4.3.6. Let $1 \in H^0(P_0^{\infty})$ denote the unit. Given $x \in A^{s,2t}$, we let

$$[x] \in {}^{\mathrm{MANSS}} E_2^{s, 2t-s}(X)$$

denote the image of $x \otimes 1$ under the first map in (4.3.5). This gives a map

(4.3.7)
$$A^{s,2t} \hookrightarrow {}^{\text{MANSS}}E_2^{s,2t-s}(X).$$

Furthermore, there is an isomorphism

$$\operatorname{Tor}_{1}^{\mathbb{Z}}(A^{s,2t}, H^{1}(\Sigma^{1}P_{-1}^{\infty})) \cong A^{s,2t}[2].$$

By choosing a splitting of the second map of (4.3.5), we get a map

(4.3.8)
$$A^{s,2t}[2] \hookrightarrow {}^{\text{MANSS}} E_2^{s,2t-s+1-\sigma}(X).$$

If $x \in A^{s,2t}[2]$, then we will denote the image of x under these maps by [ux].

Recall from the previous subsection that ${}^{\text{MANSS}}E_2^{*,\star}(X)$ is a module over $\pi_{\star}^{C_2}H\underline{\mathbb{Z}}_a^{\wedge} = \mathbb{Z}[a, u^{\pm 2}]/(2a)$. By [BMMS86, Theorem V.2.4] we have

$$\pi_{i+j\sigma}^{C_2}(H\underline{\mathbb{Z}}_a^{\wedge}) \cong H^{-i}(P_j^{\infty}),$$

and under this isomorphism the element $a^k u^{2l} \in \pi_{2l-(k+2l)\sigma}^{C_2}(H\underline{\mathbb{Z}}_a^{\wedge})$ corresponds to a generator of $H^{-2l}(P_{-k-2l}^{\infty})$, and the first map of (4.3.5) is a map of $\pi_{\star}^{C_2}(H\underline{\mathbb{Z}}_a^{\wedge})$ -modules.

Remark 4.3.9. Let x be an element in $A^{s,2t}[2]$. Analogous to the definition of \overline{ux} (4.2.9), the choice of the element [ux] may not be unique, but may depend on the choice of splitting of (4.3.8). However different lifts [ux] coming from two choices of splitting differ by an element of the form

$$\sum_{i} \left(\frac{a^2}{u^2}\right)^i a[y_i]$$

for $y_i \in A^{s-1-2i,2t}$.

Remark 4.3.10. If X is a ring spectrum, the map α in (4.3.5) is a map of rings, and therefore we have

$$[x] \cdot [y] = [xy].$$

Moreover the map β in (4.3.5) is a map of modules over $A^{*,*}$. It follows from an argument analogous to that of Lemma 4.2.8 that any splitting of (4.3.5) is a splitting of modules over $A^{*,*}$, and therefore given $x \in A^{s,2t}$ and $y \in A^{s',2t'}[2]$, we have

$$[x] \cdot [uy] = [uxy].$$

The following proposition is an immediate consequence of the Künneth theorem (4.3.5).

Proposition 4.3.11. For $X \in \text{Sp}_{i2}^{BG}$ with trivial action and MU_*X -torsion-free, under the maps (4.3.7) and (4.3.8) there is an isomorphism

$${}^{\text{MANSS}}E_2^{*,\star}(X) \cong \left(A^{*,*} \otimes \mathbb{Z}[u^{\pm 2}, a]/(2a)\right) \oplus \left(A^{*,*}[2] \otimes \mathbb{F}_2[u^{\pm 2}, a]\right).$$

$$30$$

Corollary 4.3.12. For X as in Proposition 4.3.11, the first double complex spectral sequence

$${}_{I}E_{2}^{a,b,2t-a-b-j+j\sigma}(X) = \operatorname{Ext}^{a,2t}((MU_{2})_{*}X \otimes H^{b+j}(P_{j}^{\infty})) \implies \operatorname{MANSS}E_{2}^{a+b,2t-a-b-j+j\sigma}(X)$$

collapses.

Proof. Assume P_j^{∞} is given a finite type cellular structure. Let $B^k \subseteq Z^k \subseteq C_{cell}^k(P_j^{\infty})$ denote the subgroups of coboundaries and cocycles of the cellular cochain complex. Then we have a resolution of free abelian groups

$$0 \to B^k \to Z^k \to H^k(P_j^\infty) \to 0.$$

Since $(MU_2)_*(X)$ is torsion free, we have a short exact sequence of $(MU_2)_*(MU_2)$ -comodules

$$0 \to (MU_2)_* X \otimes B^k \to (MU_2)_* X \otimes Z^k \to (MU_2)_* \otimes H^k(P_j^{\infty}) \to 0$$

and therefore an exact sequence on Ext:

$$A^{a,2t} \otimes B^k \xrightarrow{(1)} A^{a,2t} \otimes Z^k \to \operatorname{Ext}^{a,2t}((MU_2)_*X \otimes H^k(P_j^\infty)) \to A^{a+1,2t} \otimes B^k \xrightarrow{(2)} A^{a+1,2t} \otimes Z^k.$$

It follows that we have a short exact sequence ("universal coefficient theorem") (4.3.13)

$$0 \to A^{a,2t} \otimes H^k(P_j^{\infty}) \to \operatorname{Ext}^{a,2t}((MU_2)_*X \otimes H^k(P_j^{\infty})) \to \operatorname{Tor}_1^{\mathbb{Z}}(A^{a+1,2t}, H^k(P_j^{\infty})) \to 0$$

involving the cokernel of (1) and the kernel of (2). Comparing the description of $_{I}E_{2}$ in (4.3.13) with the description of $^{\text{MANSS}}E_{2}$ in (4.3.5), we see that there can be no differentials in the first double complex spectral sequence.

Remark 4.3.14. A similar argument, using the classical Universal Coefficient Theorem for cohomology, shows that the second double complex spectral sequence also collapses.

4.4. The C-motivic MANSS and the algebraic AHSS. We continue with the assumption in the last subsection that $X \in \operatorname{Sp}_{i2}^{BC_2}$ is a Borel spectrum with trivial C_2 action, and $(MU_2)_*X$ is torsion-free. We will now proceed with our goal of relating differentials in the MANSS to those in the Atiyah–Hirzebruch spectral sequence (AHSS). Our strategy will be to use complex motivic homotopy as a means of intrinsically encoding the Adams–Novikov aspect of the MANSS. For this purpose we will add the following assumption for the rest of the section.

Assumption 4.4.1. The graded group $(MU_2)_*X$ is concentrated in even degrees.

Let

$$\nu_{\mathbb{C}}: \mathrm{Sp}_{i2} \to \mathrm{SH}(\mathbb{C})_{i2}$$

be the section of Betti realization Be : $\mathrm{SH}(\mathbb{C})_{i2} \to \mathrm{Sp}_{i2}$ constructed in [GIKR22]. Letting $\tau \in \pi_{0,-1}^{\mathbb{C}} \mathbb{S}_2$ denote the canonical generator, we let $C\tau$ denote the cofiber of τ , and define for $Z \in \mathrm{SH}(\mathbb{C})$

$$Z/\tau \coloneqq Z \otimes C\tau.$$

We shall define a complex motivic lift $\operatorname{MaNSS}_{mot} E_r^{*,\star,*}(\nu_{\mathbb{C}}X)$ of the MANSS (the "motivic MANSS"). There will then be a zig-zag

$$^{\text{MANSS}}E_r^{*,\star}(X) \cong {}^{\text{MANSS}}_{\text{mot}}E_r^{*,\star,*}(\nu_{\mathbb{C}}X[\tau^{-1}]) \leftarrow {}^{\text{MANSS}}_{\text{mot}}E_r^{*,\star,*}(\nu_{\mathbb{C}}X) \to {}^{\text{MANSS}}_{\text{mot}}E_r^{*,\star,*}(\nu_{\mathbb{C}}X/\tau)$$

and we will show that the spectral sequence $\underset{\text{mot}}{\overset{\text{MANSS}}{\text{mot}}} E_r^{*,*+j\sigma,*}(\nu_{\mathbb{C}}X/\tau)$ is a variant of the algebraic AHSS for $\text{Ext}^{*,*}((MU_2)_*(X^{P_j^{\infty}}))$.

Because of our hypothesis that MU_*X is even, we have

(4.4.2)
$$\pi_{n,t}^{\mathbb{C}}(\nu_{\mathbb{C}}X/\tau) \cong \operatorname{Ext}^{2t-n,2t}((MU_{2})_{*}X) = A^{2t-n,2t}$$

and the τ -Bockstein spectral sequence for $\nu_{\mathbb{C}} X$ is equivalent to the ANSS for X. Specifically, we have isomorphisms

$$(MU_2)_{2t}(X) \cong (MGL_2)_{2t,t}(\nu_{\mathbb{C}}X)$$

and with respect to these isomorphisms we have an isomorphism

$$(MGL_2)_{*,*}(\nu_{\mathbb{C}}X) \cong (MU_2)_*(X)[\tau]$$

The motivic ANSS (the MGL_2 -based ASS in $SH(\mathbb{C})_{i2}$) for $\nu_{\mathbb{C}}X$ takes the form

$${}^{\mathrm{ANSS}}_{\mathrm{mot}} E_2^{s,n,w}(\nu_{\mathbb{C}} X) \Longrightarrow \pi_{n-s,w}^{\mathbb{C}} \nu_{\mathbb{C}} X$$

We have

$${}^{\text{ANSS}}_{\text{mot}} E_2^{s,2w-s,w}(\nu_{\mathbb{C}} X) \cong A^{s,2w}$$

and we denote the image of an element $x \in A^{s,2w}$ under the above isomorphism by

$$\tilde{x} \in {}^{\text{ANSS}}_{mot} E_2^{s, 2w-s, w}(\nu_{\mathbb{C}} X).$$

In the case where $X = \mathbb{S}_2$ the element $\tau \in \pi_{0,-1}^{\mathbb{C}} \mathbb{S}_2$ lifts to an element which we also denote

$$\tau \in {}^{\text{ANSS}}_{\text{mot}} E_2^{-2,0,-1}(\mathbb{S}_2)$$

This gives $\operatorname{MNSS}_{\mathrm{mot}} E_2^{*,*,*}(\nu_{\mathbb{C}} X)$ the structure of a $\mathbb{Z}[\tau]$ -module, and we have

$${}^{\text{ANSS}}_{\text{mot}} E_2^{*,*,*}(\nu_{\mathbb{C}} X) \cong A^{*,*}[\tau].$$

A differential

$$d_{2r+1}(x) = y$$

in the ANSS for X corresponds to a differential

$$d_{2r+1}(\tilde{x}) = \tau^r \tilde{y}$$

in the motivic ANSS for $\nu_{\mathbb{C}} X$.

Define

$$\nu_{\mathbb{C}}(X, P_j^{\infty}) \coloneqq \lim_{m} \nu_{\mathbb{C}}(X^{P_j^{2m}}).$$

We take the limit outside the $\nu_{\mathbb{C}}$ because $\nu_{\mathbb{C}}$ does not commute with limits in general. We will be interested in the complex motivic homotopy groups

$$\pi_{i,w}^{\mathbb{C}}\nu_{\mathbb{C}}(X,P_j^{\infty})$$

as lifts of the equivariant homotopy groups

$$\pi_{i+j\sigma}^{C_2}\boldsymbol{X}^h\cong\pi_i(\boldsymbol{X}^{P_j^\infty}).$$

Remark 4.4.3. It may be the case that there is an equivalence

$$\nu_{\mathbb{C}}(X,P_{j}^{\infty})\simeq \left(\nu_{\mathbb{C}}X\right)^{L_{j}^{\infty}}$$

where the spectra L_j^{∞} are the stunted versions of $B\mu_2$ studied by [GR23], [Qui19], but we are unable to determine this.

Lemma 4.4.4. The maps $P_j^{2m} \to P_{j+1}^{2m}$ induce cofiber sequences

$$\nu_{\mathbb{C}}(X^{P_{j+1}^{2m}}) \to \nu_{\mathbb{C}}(X^{P_{j}^{2m}}) \to \Sigma^{-j,-\lceil j/2 \rceil} \nu_{\mathbb{C}} X$$
$$\nu_{\mathbb{C}}(X, P_{j+1}^{\infty}) \to \nu_{\mathbb{C}}(X, P_{j}^{\infty}) \to \Sigma^{-j,-\lceil j/2 \rceil} \nu_{\mathbb{C}} X.$$
$$32$$

Proof. Following, for example, [Beh06, Sec. 8], using the torsion-free hypothesis on $(MU_2)_*X$, we have isomorphisms of $(MU_2)_*$ -modules

(4.4.5)
$$(MU_{2})_{*}(X^{P_{2k-1}^{2m}}) \cong (MU_{2})_{*}(X) \otimes_{(MU_{2})_{*}} \frac{x^{k}((MU_{2})_{*}[x]/(x^{m}))}{\langle x^{i-1}[2](x) : i \geq k \rangle},$$
$$(MU_{2})_{*}(X^{P_{2k}^{2m}}) \cong (MU_{2})_{*}(X) \otimes_{(MU_{2})_{*}} \frac{x^{k}((MU_{2})_{*}[x]/(x^{m}))}{\langle x^{i-1}[2](x) : i \geq k+1 \rangle}$$

where |x| = -2 and $[2](x) \in (MU_2)_*[[x]]$ is the 2-series of the universal formal group law. Applying $\nu_{\mathbb{C}}$ to the cofiber sequences

$$X^{P_{2j+1}^{2m}} \to X^{P_{2j}^{2m}} \to \Sigma^{-2j} X$$
$$\Sigma^{-2j} X \to X^{P_{2j}^{2m}} \to X^{P_{2j-1}^{2m}}$$

the result then follows from [GIKR22, Lem. 3.13, Prop. 3.18].

Definition 4.4.6. Consider the following motivic lift $\Gamma_{MGL_2}^{h\mu_2}(\nu_{\mathbb{C}}X)[j] \in SH(\mathbb{C})^{Fil}$ of the MANSS filtration:

$$\Gamma_{MGL_2}^{h\mu_2}(\nu_{\mathbb{C}}X)[j]_s \coloneqq \begin{cases} \nu_{\mathbb{C}}(X, P_j^{\infty}) & \text{if } s \le 0\\ \underset{a+b\ge s}{\text{colim}} \overline{MGL_2}^{\otimes a} \widehat{\otimes} \nu_{\mathbb{C}}(X, P_{j+b}^{\infty}) & \text{if } s > 0. \end{cases}$$

where the $\widehat{\otimes}$ notation is defined by

$$Z\widehat{\otimes}\nu_{\mathbb{C}}(X,P_{\ell}^{\infty}) \coloneqq \lim_{m} Z \otimes \nu_{\mathbb{C}}(X^{P_{\ell}^{2m}})$$

for $Z \in SH(\mathbb{C})_{i2}$. We define the *motivic MANSS* to be the spectral sequence associated to this filtration.

Calculating the associated graded of this object gives a motivic analogue of Lemma 4.3.4: $\underset{\text{mot}}{^{\text{MANSS}}}E_1^{s,2t-s+j\sigma,*}(\nu_{\mathbb{C}}X) \cong \bigoplus_{a+b=s} C_{\text{cell,mot}}^{b+j,*}(\nu_{\mathbb{C}}P_j^{\infty}) \otimes C_{(MGL_2)_{*,*}MGL_2}^{a,2t+j,*}((MGL_2)_{*,*}\nu_{\mathbb{C}}X).$

where we are able to commute the limit in $\widehat{\otimes}$ past the colimit since the former is sequential and the latter can be taken to be finite. Non-finiteness issues involved in dualizing $\nu_{\mathbb{C}}(\mathbb{S}, P_j^{\infty})$ to $\nu_{\mathbb{C}}(P_i^{\infty})$ are explained in [GIKR22, Lemma 3.24]. Analogous arguments to (4.3.5) hold in the motivic case and we obtain the following.

Lemma 4.4.7. There is a (non-canonically split) short exact sequence

$$(4.4.8) \quad 0 \to \bigoplus_{a+b=s} A^{a,2t} \otimes_{\mathbb{Z}_2} H^{b+j,w}(\nu_{\mathbb{C}} P_j^{\infty}) \to \underset{\text{mot}}{\overset{\text{MANSS}}{\text{mot}}} E_2^{s,2t-s-j+j\sigma,t-w}(\nu_{\mathbb{C}} X)$$
$$\to \bigoplus_{a+b=s+1} \operatorname{Tor}_1^{\mathbb{Z}_2}(A^{a,2t}, H^{b+j,w}(\nu_{\mathbb{C}} P_j^{\infty})) \to 0.$$

The same argument as Corollary 4.3.12 gives the following.

Proposition 4.4.9. There is an isomorphism

(4.4.10)
$$\underset{\text{mot}}{^{\text{MANSS}}} E_2^{s,2t-s-j+j\sigma,t-w}(\nu_{\mathbb{C}}X) \cong \bigoplus_{a+b=s} \text{Ext}^{a,2t}((MU_2)_*X \otimes_{\mathbb{Z}_2} H^{b+j,w}(\nu_{\mathbb{C}}P_j^{\infty})).$$

In particular, we have

(4.4.11)
$$\overset{\text{MANSS}}{\text{mot}} E_2^{*,*,*}(\nu_{\mathbb{C}} X) \cong \overset{\text{MANSS}}{33} E_2^{*,*}(X) \otimes \mathbb{Z}[\tau].$$

All elements of the E_2 -term of the motivic MANSS may be represented by elements of the form

$$au^m a^k u^{2l}[x] \text{ or } au^m a^k u^{2l}[uy]$$

for $k,m\geq 0,\,l\in\mathbb{Z},\,x\in A^{s,2t},\,y\in A^{s,2t}[2],$ and

$$[x] \in \underset{\text{mot}}{^{\text{MANSS}}E_2^{s,2t-s,t}}(\nu_{\mathbb{C}}X), \qquad [uy] \in \underset{\text{mot}}{^{\text{MANSS}}E_2^{s,2t-s+1-\sigma,t}}(\nu_{\mathbb{C}}X), \\ \tau \in \underset{\text{mot}}{^{\text{MANSS}}E_2^{-2,0,-1}}(\nu_{\mathbb{C}}X), \qquad a \in \underset{\text{mot}}{^{\text{MANSS}}E_2^{1,-\sigma,0}}(\nu_{\mathbb{C}}X), \\ u^2 \in \underset{\text{mot}}{^{\text{MANSS}}E_2^{0,2-2\sigma,1}}(\nu_{\mathbb{C}}X).$$

Betti realization induces an equivalence [DI10]

(4.4.12)
$$\operatorname{SH}(\mathbb{C})_{i2}^{\mathrm{T}}[\tau^{-1}] \simeq \operatorname{Sp}_{i2}$$

where $\text{SH}(\mathbb{C})_{i2}^{\text{T}}$ denotes the subcategory of $\text{SH}(\mathbb{C})_{i2}$ of cellular objects. The following is an immediate consequence of (4.4.11).

Proposition 4.4.13. The τ -local motivic MANSS with E_2 -term $\underset{\text{mot}}{\overset{\text{MANSS}}{\text{mot}}} E_2^{*,\star,*}(\nu_{\mathbb{C}}X)[\tau^{-1}]$ is isomorphic to the 2-primary MANSS with E_2 -term $\overset{\text{MANSS}}{\overset{\text{MANSS}}{\text{mot}}} E_r^{*,\star}(X)[\tau,\tau^{-1}]$.

Proof. The canonical map induces an isomorphism on E_2 -terms.

We will now endeavor to identify the spectral sequence we obtain by smashing the motivic MANSS filtration with $C\tau$. Define

$$\operatorname{Ext}^{*,*}((\widehat{MU}_2)_*(X^{P_j^{\infty}})) \coloneqq \lim_m \operatorname{Ext}^{*,*}((MU_2)_*(X^{P_j^{2m}}))$$

so we have

$$\pi_{n,t}^{\mathbb{C}}\nu_{\mathbb{C}}(X,P_{j}^{\infty}) = \operatorname{Ext}^{t-n,t}((\widehat{MU}_{2})_{*}(X^{P_{j}^{\infty}})).$$

By (4.4.5), there are short exact sequences

(4.4.14)
$$0 \to (MU_2)_*(X^{P_{2k+1}^{2m}}) \to (MU_2)_*(X^{P_{2k}^{2m}}) \to (MU_2)_*(\Sigma^{-2k}X) \to 0$$

$$(4.4.15) \qquad 0 \to (MU_2)_* (\Sigma^{-2k} X) \xrightarrow{\cdot x^{k-1} [2](x)} (MU_2)_* (X^{P_{2k}^{2m}}) \to (MU_2)_* (X^{P_{2k-1}^{2m}}) \to 0$$

Applying Ext(-) produces long exact sequences. We observe that each of the terms in these long exact sequences as well as each of the kernels and cokernels of the maps in the long exact sequences, are Mittag-Leffler sequences in m and consequently applying the functor $\lim_m \text{Ext}(-)$ also produces a long exact sequence. These long exact sequences splice together to give us an *algebraic Atiyah-Hirzebruch spectral sequence (AHSS)* (compare with [Beh06, Sec. 8]):

$$\underset{\text{alg}}{^{\text{AHSS}}}E_1^{l,s,t}(X,P_j^{\infty}) \implies \text{Ext}^{s,t}((\widehat{MU}_2)_*(X^{P_j^{\infty}}))$$

with

$${}^{\text{AHSS}}_{\text{alg}} E_1^{l,s,t}(X, P_j^{\infty}) = \begin{cases} \text{Ext}^{s,t}((MU_2)_*(\Sigma^{-2k}X)), & l = 2k \ge j, \\ \text{Ext}^{s+1,t}((MU_2)_*(\Sigma^{-2k}X)), & l = 2k-1 \ge j, \\ 0, & l < j. \end{cases}$$

The differentials in the algebraic AHSS are determined by the effects of the attaching maps of cells in P_i^{∞} on Ext($(MU_2)_*X$).

There is a variant of the algebraic AHSS which involves taking 2 cells at a time. Specifically, applying $\lim_{m \to \infty} \operatorname{Ext}(-)$ to the decreasing filtration

$$(MU_2)_*(X^{P_j^{2m}}) \leftarrow \cdots \leftarrow (MU_2)_*(X^{P_{2k-1}^{2m}}) \leftarrow (MU_2)_*(X^{P_{2k+1}^{2m}}) \leftarrow \cdots$$

34

of $(MU_2)_*(X^{P_j^{2m}})$ when j is odd and the decreasing filtration

$$(MU_2)_*(X^{P_j^{2m}}) \leftarrow (MU_2)_*(X^{P_{j+1}^{2m}}) \leftarrow \dots \leftarrow (MU_2)_*(X^{P_{2k-1}^{2m}}) \leftarrow (MU_2)_*(X^{P_{2k+1}^{2m}}) \leftarrow \dots$$

when j is even gives rise to a variant of the algebraic AHSS which we will called the *accelerated* algebraic AHSS:

$$\stackrel{\text{AHSS}}{\underset{\text{alg,acc}}{\text{AHSS}}} E_1^{k,s,t}(X, P_j^{\infty}) \implies \text{Ext}^{s,t}((\widehat{MU}_2)_*(X^{P_j^{\infty}}))$$

with

$${}^{\text{AHSS}}_{\text{alg,acc}} E_1^{k,s,t}(X, P_j^{\infty}) = \begin{cases} \text{Ext}^{s,t}(MU_*(\Sigma^{-2k}X \otimes M(2))), & k > j/2, \\ \text{Ext}^{s,t}(MU_*(\Sigma^{-j}X)), & j \text{ even and } k = j/2, \\ 0, & \text{otherwise.} \end{cases}$$

Note that the same considerations as before apply so that applying \lim_m to the long exact sequence produces another long exact sequence.

We define the motivic MANSS mod τ to be the spectral sequence with signature

$$\underset{\text{mot}}{^{\text{MANSS}}} E_1^{s,i+j\sigma,w}(\nu_{\mathbb{C}}X/\tau) \implies \pi_{i,w}^{\mathbb{C}}(\nu_{\mathbb{C}}(X,P_j^{\infty})/\tau)$$

associated to the filtered motivic spectrum $\Gamma^h_{MGL_2}(\nu_{\mathbb{C}}X)[j]/\tau$. Since we have

$$H^{b+j,w}(\nu_{\mathbb{C}}((P_j^{\infty})_2)/\tau) = \begin{cases} \mathbb{Z}_2, & b = 0, j \text{ even}, w = \frac{j}{2}, \\ \mathbb{Z}/2, & b > 0, b+j \text{ even}, w = \frac{b+j}{2}, \\ 0 & \text{otherwise}, \end{cases}$$

by (4.4.10) we have

$$\underset{\text{mot}}{\text{MANSS}} E_2^{s,i+j\sigma,w}(\nu_{\mathbb{C}}X/\tau) = \begin{cases} \text{Ext}^{2w-i,b}((MU_2)_*X \otimes_{\mathbb{Z}_2} H^{b-2w,\frac{b}{2}-w}(\nu_{\mathbb{C}}((P_j^{\infty})_2)/\tau)) & b=i+s+j \text{ even}, \\ 0 & \text{otherwise.} \end{cases}$$

The following proposition gives an isomorphism between the motivic MANSS mod τ and the accelerated algebraic AHSS. In particular, it implies that mod τ , the differentials in the motivic MANSS are given by the effects of attaching maps in P_j^{∞} on Ext.

Theorem 4.4.16. The isomorphisms

$$\operatorname{Ext}^{2w-i,i+s+j}((MU_{2})_{*}X \otimes_{\mathbb{Z}_{2}} H^{i+s+j-2w,\frac{i+s+j}{2}-w}(\nu_{\mathbb{C}}P_{j}^{\infty}/\tau)) \\ \cong \begin{cases} \operatorname{Ext}^{2w-i,2w}((MU_{2})_{*}(\Sigma^{-j}X)), & i+s-2w=0, j \text{ even}, \\ \operatorname{Ext}^{2w-i,2w}((MU_{2})_{*}(X \otimes \Sigma^{2w-i-s-j}M(2))), & i+s-2w>0, \\ 0, & \text{otherwise} \end{cases}$$

induce an isomorphism of spectral sequences

Proof. The fundamental observation is that if $Y \in \text{Sp}_{i2}$ and $(MU_2)_*Y$ even, then

(4.4.17)
$$\pi_{i,w}^{\mathbb{C}}(\overline{MGL_{2}}^{\otimes a} \otimes \nu_{\mathbb{C}}(Y)/\tau) = \begin{cases} \operatorname{Ext}^{2w-i,2w}((MU_{2})_{*}Y), & i < 2w - a_{i}\\ Z^{2w-i,2w}((MU_{2})_{*}Y), & i = 2w - a_{i}\\ 0, & i > 2w - s \end{cases}$$

where $Z^{2w-i,2w}((MU_2)_*Y) \subseteq C^{2w-i,2w}_{(MU_2)_*MU_2}((MU_2)_*Y)$ denotes the subgroup of cocycles in the normalized cobar complex. This is proven by induction on *a* using the cofiber sequences

$$\overline{MGL_2^{\otimes a+1}} \otimes \nu_{\mathbb{C}}(X)/\tau \to \overline{MGL_2^{\otimes a}} \otimes \nu_{\mathbb{C}}(X)/\tau \to MGL_2 \otimes \overline{MGL_2^{\otimes a}} \otimes \nu_{\mathbb{C}}(X)/\tau.$$

We will prove the theorem by directly comparing the exact couple of the accelerated Atiyah–Hirzebruch spectral sequence starting from the E_2 -term to the exact couple of the motivic MANSS, run at double speed, starting from the E_1 -term. To this end, we will first fix integers j and s and define a spectral sequence to compute

$$\pi_{i,w}^{\mathbb{C}}\Gamma_{MGL_2}^{h\mu_2}(\nu_{\mathbb{C}}X)[j]_s/\tau.$$

We define a tower of spectra

where

$$Y_{\ell} \coloneqq \operatorname{colim}_{\substack{a+b \ge s \\ a \le \ell}} (\overline{MGL_2}^{\otimes a} \widehat{\otimes} \nu_{\mathbb{C}}(X, P_{j+b}^{\infty})) / \tau$$

and

$$S_f = \begin{cases} \operatorname{cof}(Y_{f+1} \to Y_f), & \text{if } 0 \le f \le s - 1, \\ Y_s, & \text{if } f = s, \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

Observe that we have

$$\pi_{i,w}^{\mathbb{C}} S_f = \begin{cases} (MGL_2 \otimes \overline{MGL_2}^{\otimes f}) \widehat{\otimes} \nu_{\mathbb{C}}(X, P_{j+s-f}^{\infty}) / \tau & \text{if } 1 \le f \le s \text{ and} \\ \overline{MGL_2}^{\otimes s} \widehat{\otimes} \nu_{\mathbb{C}}(X, P_j^{\infty}) & \text{if } f = 0. \end{cases}$$

so when $1 \leq f \leq s$ we have

(4.4.18)
$$\pi_{i,w}^{\mathbb{C}}S_f = \begin{cases} \widehat{C}_{(MU_2)_*MU_2}^{f,2w}(\widehat{MU_2})_*(X^{P_{j+s-f}^{\infty}})), & i = 2w - f, \\ 0, & \text{otherwise.} \end{cases}$$

Applying $\pi_{i,w}^{\mathbb{C}}$, we obtain a conditionally convergent spectral sequence with signature

(4.4.19)
$$E_1^{i,f,w} = \pi_{i,w}S_f \implies Y_0,$$

with

$$E_{1}^{i,f,w} = \begin{cases} \widehat{C}_{(MU_{2})_{*}MU_{2}}^{f,2w}((\widehat{MU}_{2})_{*}(X^{P_{j+s-f}^{\infty}})), & i = 2w - f, 0 \le f < s \\ Z^{s,2w}((\widehat{MU}_{2})_{*}(X^{P_{j}^{\infty}})), & i = 2w - s, f = s \\ \text{Ext}^{2w-i,2w}((\widehat{MU}_{2})_{*}(X^{P_{j}^{\infty}})), & i < 2w - s, f = s \\ 0, & \text{otherwise} \end{cases}$$

by (4.4.18) and (4.4.17). Note that in each stem *i*, there is at most one non trivial bidegree (at filtration f = 2w - i when $2w - s \le i \le 2w$, or f = s for i < 2w - s). Therefore, we will have no extensions and the spectral sequence is strongly convergent. For degree reasons, the only possibly non-trivial differentials are

$$(4.4.20) d_1: \widehat{C}^{f,2w}_{(MU_2)_*MU_2}((\widehat{MU}_2)_*(X^{P^{\infty}_{j+s-f}})) \to \widehat{C}^{f+1,2w}_{(MU_2)_*MU_2}((\widehat{MU}_2)_*(X^{P^{\infty}_{j+s-f-1}})),$$

and

(4.4.21)
$$d_1: \widehat{C}^{s-1,2w}_{(MU_2)_*MU_2}((\widehat{MU}_2)_*(X^{P_{j+1}^{\infty}})) \to Z^{s,2w}((\widehat{MU}_2)_*(X^{P_j^{\infty}})).$$

Now, assume that s + i + j is odd. Then j + s + i - 2w + 1 is even so the map

$$\widehat{C}^{*,*}_{(MU_2)_*MU_2}((\widehat{MU}_2)_*X^{P^{\infty}_{j+s+i-2w+1}}) \twoheadrightarrow \widehat{C}^{*,*}_{(MU_2)_*MU_2}((\widehat{MU}_2)_*X^{P^{\infty}_{j+s+i-2w}})$$

is a surjection by (4.4.15), and the map

$$\widehat{C}^{*,*}_{(MU_2)_*MU_2}((\widehat{MU}_2)_*X^{P^{\infty}_{j+s+i-2w}}) \hookrightarrow \widehat{C}^{*,*}_{(MU_2)_*MU_2}((\widehat{MU}_2)_*X^{P^{\infty}_{j+s+i-2w-1}})$$

is an injection by (4.4.14). We compute (4.4.20) using the commutative diagrams

$$\pi_{i,w}^{\mathbb{C}} S_{2w-i} = \widehat{C}_{(MU_2)_*MU_2}^{2w-i,2w} ((\widehat{MU}_2)_* X^{P_{j+s+i-2w}^{\infty}}) \xrightarrow{\alpha_{2w-i+1}} \pi_{i-1,w}^{\mathbb{C}} Y_{2w-i+1} \xrightarrow{d_1} \widehat{C}_{(MU_2)_*MU_2}^{2w-i+1,2w} ((\widehat{MU}_2)_* X^{P_{j+s+i-2w}^{\infty}}) \xrightarrow{\alpha_{2w-i+1}} \pi_{i-1,w}^{\mathbb{C}} S_{2w-i+1} = \widehat{C}_{(MU_2)_*MU_2}^{2w-i+1,2w} ((\widehat{MU}_2)_* X^{P_{j+s+i-2w-1}^{\infty}})$$

and

We can compute the differential (4.4.21) in a similar way. We therefore have

(4.4.22)
$$\pi_{i,w}^{\mathbb{C}}\Gamma_{MGL_2}^{h\mu_2}(\nu_{\mathbb{C}}X)[j]_s/\tau \cong \operatorname{Ext}^{2w-i,2w}((\widehat{MU}_2)_*X^{P_{i+s-2w+j}^{\infty}})$$

for i + s + j odd. By construction, these isomorphisms are compatible in the sense that

is a commutative diagram. Note that since

$${}^{\text{MANSS}}_{\text{mot}} E_2^{s,i+j\sigma,w} (\nu_{\mathbb{C}} X / \tau)$$

is concentrated in degrees with i + j + s even, it only has non-trivial d_r -differentials for r odd, and there are isomorphisms

$${}^{\text{MANSS}}_{\text{mot}} E^{s,i+j\sigma,w}_{2r}(\nu_{\mathbb{C}}X/\tau) \cong {}^{\text{MANSS}}_{\text{mot}} E^{s,i+j\sigma,w}_{2r+1}(\nu_{\mathbb{C}}X/\tau).$$

To prove the theorem, we must show that under the isomorphism

$${}^{\text{MANSS}}_{\text{mot}} E_2^{s,i+j\sigma,w}(\nu_{\mathbb{C}} X/\tau) \cong {}^{\text{AHSS}}_{\text{acc,alg}} E_1^{\frac{i+j+s}{2}-w,2w-i,2w}(X,P_j^{\infty})$$

the d_{2r+1} differentials in the motivic MANSS for $\nu_{\mathbb{C}} X / \tau$ coincide with the d_r differentials in the accelerated algebraic AHSS for $X^{P_j^{\infty}}$. This follows from the following commutative

diagram



where i + j + s is even, $k \coloneqq \frac{i+j+s-2w}{2}$, $a \coloneqq 2w - i + 1$, we write

$$\Gamma_s = \Gamma_{MGL_2}^{n\mu_2} (\nu_{\mathbb{C}} X / \tau) [j]_s$$

and $Z_1^{*,*,*} \subseteq \frac{\text{MANSS}}{\text{mot}} E_1^{*,*,*}(\nu_{\mathbb{C}} X / \tau)$ denotes the subgroup of d_1 -cycles.

5. Applications

We end with some computations of the MANSS to illustrate our perspective.

5.1. Example: ko_{C_2} at p = 2. Using the (2-primary) MANSS (see Definition 3.4.1), we revisit the computation of the $RO(C_2)$ -graded homotopy groups of the C_2 -homotopy fixed points of the 2-complete C_2 -equivariant connective real K-theory spectrum $\pi_{**}^{C_2}(ko_{C_2})_{2,a}^{\wedge}$. By [BHS20, §10], this is the same information as the Artin–Tate homotopy groups of the a-complete very effective Hermitian K-theory $\pi_{***}^{\text{AT}}(kq_{2,a}^{\wedge})$. The MANSS for $(ko_{C_2})_{2,a}^{\wedge}$ has signature

$$^{\text{MANSS}}E_2^{s,i+j\sigma}((ko_{C_2})_{2,a}^{\wedge}) \implies \pi_{i+j\sigma}^{C_2}((ko_{C_2})_{2,a}^{\wedge}).$$

Recall that the underlying spectrum of $(ko_{C_2})_{2,a}^{\wedge}$ is the 2-complete connective real K-theory spectrum ko_2 , whose Adams–Novikov E_2 -term is of the form

^{ANSS}
$$E_2^{*,*}(ko_2) = \mathbb{Z}_2[h_1, v_1^2]/2h_1$$

where the generators have degree $|h_1| = (1, 2)$ and $|v_1^2| = (0, 4)$. Here we use Adams grading convention.

There are two avenues to computing the E_2 -term of the MANSS. On one hand, we have the following lemma.

Lemma 5.1.1. The spectrum $(ko_{C_2})_2^{\wedge}$ is $MU_{\mathbb{R}}$ -projective.

Proof. Smashing the "Wood cofiber sequence"

$$\Sigma^{\sigma}(ko_{C_2})_2^{\wedge} \xrightarrow{\eta_2^{\wedge}} (ko_{C_2})_2^{\wedge} \to k\mathbb{R}_2$$

of [GHIR20, Prop. 10.13] with $MU_{\mathbb{R},2}$, and using the fact that the Hurewicz image of $[\eta] \in \pi_{\sigma}^{C_2}(S)$ in $\pi_{\sigma}MU_{\mathbb{R}}$ is trivial, we deduce that $MU_{\mathbb{R},2} \otimes ko_{C_2}$ is a summand of $MU_{\mathbb{R},2} \otimes k\mathbb{R}$. The result follows from the fact that $k\mathbb{R}$ is $MU_{\mathbb{R}}$ -projective [Hea19, Sec. 4].

It follows from Proposition 4.2.15 and Remark 4.2.14 that we have

$$^{\text{MANSS}}E_2^{*,\star}((ko_{C_2})_{2,a}^{\wedge}) \cong \mathbb{Z}_2[u^{\pm 2}, a, \overline{h_1}, \overline{v_1^2}]/(2a, 2\overline{h_1}) \oplus \mathbb{F}_2[u^{\pm 2}, a, \overline{h_1}, \overline{v_1^2}]\{\overline{uh_1}\}$$

Here, labelling degrees as (filtration, $RO(C_2)$ -stem), we have

$$|u^{2}| = (0, 2 - 2\sigma), \qquad |a| = (1, -\sigma),$$

$$|\overline{h_{1}}| = (1, \sigma), \qquad |\overline{v_{1}}| = (0, 2\rho),$$

$$|\overline{uh_{1}}| = (1, 1).$$

Note that while in general the generator $\overline{uh_1}$ depends on a choice of a lift, by Remark 4.2.13, this lift is unique in this case because $\operatorname{Ext}^{0,2}(MU_*ko) = 0$.

However, as the underlying spectrum ko_2^{\wedge} of $(ko_{C_2})_{2,a}^{\wedge}$ has a trivial C_2 -action, Proposition 4.3.11 gives a *different* presentation of the MANSS E_2 -term:

^{MANSS}
$$E_2^{*,\star}((ko_{C_2})_{2,a}^{\wedge}) \cong \mathbb{Z}_2[u^{\pm 2}, a, [h_1], [v_1^2]]/(2a, 2[h_1]) \oplus \mathbb{F}_2[u^{\pm 2}, a, [h_1], [v_1^2]]\{[uh_1]\}.$$

with

$$\begin{split} |[h_1]| &= (1,1), \\ |[uh_1]| &= (1,2-\sigma). \end{split}$$

Note that analogous to the case of $\overline{uh_1}$, the generator $[uh_1]$ has the potential to depend on a choice of a lift, but by Remark 4.3.9, because $\operatorname{Ext}^{0,2}((MU_2)_*ko_2) = 0$, this lift is also unique in this case.

Finally, Proposition 4.4.9 and Remark 4.3.10 gives the motivic MANSS E_2 -term: ${}^{\text{MANSS}}_{\text{mot}}E_2^{*,\star,*}((\nu_{\mathbb{C}}((ko)_2)) \cong \mathbb{Z}_2[u^{\pm 2}, a, [h_1], [v_1^2], \tau]/(2a, 2[h_1]) \oplus \mathbb{F}_2[u^{\pm 2}, a, [h_1], [v_1^2], \tau] \{[uh_1]\}.$ Recording tridegrees as (filtration, $RO(C_2)$ -stem, motivic weight), we have

$$\begin{aligned} |u^2| &= (0, 2 - 2\sigma, 1), & |a| &= (1, -\sigma, 0), \\ |[h_1]| &= (1, 1, 1), & |[v_1^2]| &= (0, 4, 2), \\ |[uh_1]| &= (1, 2 - \sigma, 1), & |\tau| &= (-2, 0, -1). \end{aligned}$$

Theorem 4.4.16 implies that mod τ the motivic AHSS can be identified with the accelerated algebraic AHSS for $\text{Ext}((\widehat{MU}_2)_*ko_2^{P_j^{\infty}})$. Under this correspondence, the class $a^k u^{2l}[u^{\varepsilon}x]$ corresponds to the class

$$x[-2l-\varepsilon] \in \operatorname{AHSS}_{\operatorname{alg}} E_1^{*,*,*}(ko_2, P_{-k-2l-\varepsilon}^{\infty})$$

which represents "x on the $(-2l - \varepsilon)$ -cell of $P_{-k-2l-\varepsilon}^{\infty}$ ".

The following theorem explains the philosophy that in the case of a trivial action, MANSS differentials can be derived from ANSS differentials and algebraic AHSS differentials.

Theorem 5.1.2. In the MANSS for $(ko_{C_2})_{2,a}^{\wedge}$, the d_3 -differentials are determined by

(1) $d_3(a) = 0,$ (2) $d_3([v_1^2]) = [h_1^3],$ (3) $d_3(u^2) = a^2[h_1],$ (4) $d_3([uh_1]) = \frac{a^2}{u^2}[uh_1^2].$

For dimensional reasons we have $E_4 = E_{\infty}$.

Proof. To prove (1), note that the element a is always a permanent cycle in the MANSS due to the fact that it comes from $\pi_{-\sigma}^{C_2} \mathbb{S}^0$.

To prove (2), note that for dimensional reasons, we have

$$d_3([v_1^2]) = \alpha[h_1^3] + \beta \frac{a^2}{u^2} [v_1^2 h_1]$$

for $\alpha, \beta \in \mathbb{F}_2$. Consider the map from the MANSS to the ANSS from Corollary 4.1.5. In the Adams–Novikov spectral sequence, we have $d_3(v_1^2) = h_1^3$. Since $[v_1^2]$ maps to v_1^2 and $\alpha[h_1^3] + \beta \frac{a^2}{u^2}[v_1^2 h_1]$ maps to αh_1^2 , we deduce that $\alpha = 1$. In the motivic MANSS for $\nu_{\mathbb{C}}(ko_2)$ we therefore have

$$d_3([v_1^2]) = \alpha \tau[h_1^3] + \beta \frac{a^2}{u^2} [v_1^2 h_1]$$
$$= \tau[h_1^3] + \beta \frac{a^2}{u^2} [v_1^2 h_1].$$

Therefore in the motivic MANSS for $\nu_{\mathbb{C}}(ko_2)/\tau$ we have

$$d_3([v_1^2]) = \beta \frac{a^2}{u^2} [v_1^2 h_1].$$

Since the 0-cell splits off of P_0^{∞} , we have

$$d_1([v_1^2]) = 0$$

in the accelerated algebraic AHSS for $Ext((\widehat{MU}_2)_*ko_2^{P_0^{\infty}})$. By Theorem 4.4.16 we deduce $\beta = 0.$

To prove (3), note that for dimensional reasons, in the MANSS we have

$$d_3(u^2) = \alpha a^2[h_1]$$

for $\alpha \in \mathbb{F}_2$. This corresponds to a differential

$$d_3(u^2) = \alpha a^2[h_1]$$

in the motivic MANSS for $\nu_{\mathbb{C}}(ko_2)$, and hence a differential

$$d_3(u^2) = \alpha a^2[h_1]$$

in the motivic MANSS for $\nu_{\mathbb{C}}(ko_2)/\tau$. However, since the 0-cell attaches to the -2-cell of P_{-2}^{∞} by an η -attaching map, we have

$$d_1(u^2) = a^2[h_1]$$

in the accelerated algebraic AHSS for $Ext((\widehat{MU}_2)_* P_{-2}^{\infty})$. It follows that $\alpha = 1$. (Note (3) could also have been deduced from Proposition 4.1.8.)

To prove (4), note that for dimensional reasons, in the MANSS we have

$$d_3([uh_1]) = \alpha \frac{a^2}{u^2} [uh_1^2]$$

for $\alpha \in \mathbb{F}_2$. This corresponds to a differential

$$d_3([uh_1]) = \alpha \frac{a^2}{u^2} [uh_1^2]$$
40

in the motivic MANSS for $\nu_{\mathbb{C}}(ko_2)$, and hence a differential

$$d_3([uh_1]) = \alpha \frac{a^2}{u^2} [uh_1^2]$$

in the motivic MANSS for $\nu_{\mathbb{C}}(ko_2)/\tau$. However, since the 1-cell attaches to the -1-cell of P_{-1}^{∞} by an η -attaching map, we have

$$d_1([uh_1]) = \frac{a^2}{u^2}[uh_1^2]$$

in the accelerated algebraic AHSS for $Ext((\widehat{MU}_2)_* P_{-1}^{\infty})$. It follows that $\alpha = 1$.

The element $[uh_1]$ is not a product in the E_2 -term of the MANSS for $(ko_{C_2})_{2,a}^{\wedge}$ because u does not exist on the E_2 -term. The following proposition is therefore needed to completely identify the multiplicative structure in ${}^{\text{MANSS}}E((ko_{C_2})_{2,a}^{\wedge})$.

Proposition 5.1.3. In $^{\text{MANSS}}E_2((ko_{C_2})_{2,a}^{\wedge})$ there is a multiplicative relation

$$[uh_1]^2 = u^2[h_1^2] + a^2[v_1^2].$$

Proof. For degree reasons, we know that

$$[uh_1]^2 = \alpha u^2 [h_1^2] + \beta a^2 [v_1^2]$$

for some $\alpha, \beta \in \mathbb{F}_2$. By the Leibniz rule, we have

$$0 = d_3([uh_1]^2) = d_3(\alpha u^2[h_1^2] + \beta a^2[v_1^2])$$
$$= \alpha a^2[h_1^3] + \beta a^2[h_1^3].$$

Therefore, we have $\alpha = \beta$. Note that under the map of spectral sequences from the MANSS to the ANSS given by Corollary 4.1.5, we have $[uh_1]$ maps to h_1 . Since $h_1^2 \neq 0$, we conclude that $\alpha = \beta = 1$.

When $X \in \text{Sp}_{C_2,i2}$ is $MU_{\mathbb{R}}$ -projective, X^e has a trivial C_2 -action, and $x \in \text{Ext}^{s,2t}((MU_2)_*X^e)$, then we have two different MANSS classes:

$$\overline{x} \in {}^{\text{MANSS}} E^{s,t-s+t\sigma}(X_a^{\wedge}), \text{ and}$$
$$[x] \in {}^{\text{MANSS}} E_2^{s,2t-s}(X_a^{\wedge}).$$

By Lemma 4.2.2, we either have t = 2k or 2x = 0. Write $t = 2k + \varepsilon$. By considering the map to the ANSS (Corollary 4.1.5), we have

$$[x] \equiv u^{2k} \overline{u^{\varepsilon} x} \pmod{a}.$$

For example, when $X = (ko_{C_2})_{2,a}^{\wedge}$, we deduce that

$$[v_1^2] = u^2 \overline{v_1^2},$$

$$[h_1] = \overline{uh_1}, \text{ and}$$

$$[uh_1] = u^2 \overline{h_1}$$

for dimensional reasons. In particular, squaring the last of these equations and using Proposition 5.1.3 gives

$$u^{4}\overline{h}_{1}^{2} = u^{2}[h_{1}^{2}] + a^{2}[v_{1}^{2}].$$

Dividing by u^4 , we get

(5.1.4)
$$\overline{h_1^2} = u^{-2} [h_1^2] + u^{-4} a^2 [v_1^2].$$

which gives an example of an $x \in \text{Ext}$ where $[x] \neq u^t \overline{x}$. More generally we have

$$\overline{h_1^{2k}} = (u^{-2}[h_1^2] + u^{-4}a^2[v_1^2])^k.$$

Analogous to the case of $[uh_1]$, the element $\overline{uh_1}$ is not a product in the E_2 -term of the MANSS for $(ko_{C_2})_{2,a}^{\wedge}$. We deduce the following exotic multiplicative relation, which was originally proven by the fourth author in [Kon20] in the C_2 -ESSS.

Proposition 5.1.5. In $^{\text{MANSS}}E((ko_{C_2})_{2,a}^{\wedge})$ there is a multiplicative relation

$$\overline{uh_1}^2 = u^2 \overline{h_1^2} + a^2 \overline{v_1^2}$$

Proof. Solving (5.1.4) for $[h_1^2]$, we have

$$[h_1]^2 = u^2 \overline{h_1^2} + u^{-2} a^2 [v_1^2]$$
$$= u^2 \overline{h_1^2} + a^2 \overline{v_1^2}.$$

The result follows from the fact that $[h_1] = \overline{uh_1}$.

We deduce the following differentials in the C_2 -ESSS naming convention, which correspond to the generating differentials in the C_2 -ESSS for ko_{C_2} as computed by the fourth author in [Kon20]. These differentials are implied by the differentials computed in Theorem 5.1.2 and the multiplicative relation from Proposition 5.1.5 so we only mention them in order to provide a dictionary between the computations above and the computations in [Kon20].

Corollary 5.1.6. In the MANSS for $(ko_{C_2})_{2,a}^{\wedge}$, we have

$$d_3(u^2) = a^2 \overline{uh_1}$$
, and $d_3(\overline{v}_1^2) = \overline{uh_1^3}$.

Proof. The first differential follows immediately from Theorem 5.1.2(1). The second is computed from Theorem 5.1.2 and Proposition 5.1.3. On the one hand, we have

$$d_3(u^2 \overline{v_1^2}) = d_3([v_1^2]) = (\overline{uh_1})^3 \stackrel{\text{Thm}.5.1.2}{=} [h_1]^3 = \overline{uh_1} \cdot (u^2 \overline{h_1^2} + a^2 \overline{v_1^2}).$$

However, the Leibniz rule gives

$$d_3(u^2\overline{v_1^2}) = a^2\overline{uh_1}\cdot\overline{v_1^2} + u^2d_3(\overline{v_1^2}).$$

We deduce that

$$d_3(\overline{v_1^2}) = \overline{uh_1} \cdot \overline{h_1^2} = \overline{uh_1^3}.$$

5.2. Example: tmf(2) at p = 3. We now consider the 3-complete spectrum $tmf(2)_3$ of topological modular forms for $\Gamma(2)$ studied in depth in [HSW23]. Recall that a $\Gamma(2)$ -structure on an elliptic curve C is a choice of isomorphism

$$\mathbb{F}_2 \oplus \mathbb{F}_2 \xrightarrow{\cong} C[2]$$

where C[2] denotes the 2-torsion subgroup-scheme of C. Associated to the Deligne-Mumford compactification of the moduli stack of elliptic curves with level 2 structure is a non-connective commutative ring spectrum Tmf(2) [Sto12, §5.3]. The group $GL_2(\mathbb{F}_2) \cong \Sigma_3$ acts transitively on the set of $\Gamma(2)$ -structures, and this gives Tmf(2) the structure of a commutative Borel Σ_3 spectrum, which we may regard as a genuine commutative Σ_3 -spectrum. Following [HSW23, Notation 1.7], we define tmf(2) \in Sp $_{\Sigma_3}$ to be the Σ_3 -equivariant connective cover of Tmf(2). The periodic version of tmf(2)₃ is given by

$$\mathrm{TMF}(2)_3 \coloneqq \mathrm{tmf}(2)_3[\Delta^{-1}] \simeq (\mathrm{tmf}(2)_{E(2)})_3.$$

By [Sto12, Theorem 6.1], we have

$$\text{TMF}_3 \simeq \text{TMF}(2)_3^{h\Sigma_3}.$$

Recall that there is a decomposition $\Sigma_3 \cong C_3 \rtimes C_2$ where C_2 acts nontrivially on C_3 . The HFPSS for the C_2 -action on the parametrized C_3 -homotopy fixed points $\text{TMF}(2)_3^{h_{C_2}C_3}$ (see [GM95] [QS21, 1.6. Definition]) degenerates to give an isomorphism

$$\pi_* \mathrm{TMF}_3 \cong \pi_* (\mathrm{TMF}(2)_3^{h_{C_2}C_3})^{h_{C_2}}$$

We remark that K(2)-locally, there is a maximal finite subgroup $G_{12} = C_3 \rtimes C_4$ of the second Morava stabilizer group where C_4 acts nontrivially on C_3 . We have

$$\Sigma_3 = G_{12}/C_2' = C_3 \rtimes (C_4/C_2')$$

where C'_2 is the subgroup of C_4 of order 2. There is an equivalence

$$\mathrm{TMF}(2)_{K(2)} \simeq E_2^{hC_2' \times \mathrm{Gal}(\mathbb{F}_9/\mathbb{F}_3)}$$

where E_2 is the Morava *E*-theory spectrum associated to the formal group of the unique supersingular elliptic curve over \mathbb{F}_9 . Thus we have (see [Beh20, Proposition 1.6.12])

$$\mathrm{TMF}_{K(2)} \simeq E_2^{h(G_{12} \rtimes \mathrm{Gal})} \simeq (E_2^{hC_2' \rtimes \mathrm{Gal}})^{h\Sigma_3}$$

With respect to the decomposition $\Sigma_3 = C_3 \rtimes C_2$, let γ be a choice of generator of the subgroup C_3 and μ be the generator of the subgroup C_2 . We have [Sto12, Sec. 7]

$$\pi_* \operatorname{tmf}(2)_3 \cong \mathbb{Z}_3[e_1, e_2].$$

The Σ_3 -action is encoded by

$$\gamma(e_1) = e_2, \ \gamma(e_2) = -e_1 - e_2$$

 $\mu(e_1) = -e_2, \ \mu(e_2) = -e_1.$

Define the element $\Delta^{1/2} \in \pi_{12} \operatorname{tmf}(2)_3$ to be $\Lambda^{1/2} := 4$

$$\Delta^{1/2} \coloneqq 4e_1e_2(e_1 + e_2)$$

so we have $(\Delta^{1/2})^2 = \Delta \in \pi_{24} \text{tmf}(2)_3$ [Sto12, Eq. (16)]. Note that

$$\Delta^{1/2} = -4e_1 \cdot \gamma(e_1) \cdot \gamma^2(e_1).$$

We will now compute the $(C_3$ -equivariant 3-primary) MANSS for tmf(2)^h₃ where

$$\operatorname{cmf}(2)_3^n \coloneqq F(EC_{3+}, \operatorname{tmf}(2)_3)$$

is the Borel completion of $tmf(2)_3$.

Remark 5.2.1. We warn the reader that the MANSS computation for $\text{tmf}(2)_3$ in this subsection is not particularly novel or deep. We shall see that the MANSS in this case is isomorphic to the $RO(C_3)$ -graded HFPSS. This is expected given the fact that E_2 is μ_3 -oriented in the sense of [HSW23] and the MANSS should agree with the homotopy completion of the slice spectral sequence in this case.

Indeed, the behaviour of the $RO(C_3)$ -graded HFPSS spectral sequence of E_2 is well-known to experts, and is essentially described in [HHR11], where it is related to a conjectural description of the odd primary slice spectral sequence. This conjectural description is verified in [Wil17]. A detailed account of the behavior of the $RO(C_p)$ -graded HFPSS for the Morava E-theory spectra $E_{n(p-1)}$ is the subject of a forthcoming paper of Andrew Senger. We are grateful to Senger for both generously sharing his results, and allowing us to use the n = 1case of his computations to illustrate this odd primary example of the MANSS. Write $RO(C_3) = \mathbb{Z}\{1, \lambda\}$, where λ is the standard 2-dimensional representation of C_3 given by rotation by 120 degrees. Note that the real regular representation $\rho = \mathbb{R}[C_3]$ of C_3 is given by $\rho = 1 + \lambda$. Following [HHR11, Wil17, HSW23], we shall find that the computation will be cleaner via the introduction of a spoke-enhanced representation grading. Let

$$S^{\succ} \subset S^{\lambda}$$

denote the unreduced suspension of the 3rd roots of unity in λ (so that S^{\succ} is the 1-skeleton of the usual C_3 -CW complex structure of S^{λ}). Since S^{λ} is the C_3 -analog of the C_2 -representation sphere $S^{2\sigma}$, in some sense S^{\succ} plays the role of " $S^{\lambda/2}$ ", as it is the C_3 analog of S^{σ} . For example, there is a cofiber sequence

$$(5.2.2) (C_3)_+ \to S^0 \to S^{\succ}.$$

However, we avoid taking this analogy too literally, as

$$S^{\succ} \otimes S^{\succ} \simeq S^{\lambda} \vee (C_3)_+ \otimes S^2 \neq S^{\lambda}.$$

We define $S^{\rightarrow} := D(S^{\succ})$, but note that S^{\succ} is not invertible. We do have

$$S^{\lambda \to \succ} \simeq S^{\flat}$$

which implies that

$$S^{\rightarrow} \simeq S^{\succ} \otimes S^{-\lambda}.$$

We therefore define

$$RO^{\succ}(C_3) = \frac{RO(C_3) \oplus \mathbb{Z}\{\succ\}}{(2\succ = \lambda)}$$

and we then have

$$-(\succ) = \succ - \lambda \in RO^{\succ}(C_3)$$

Given $i + j\lambda + \varepsilon \succ$ (with $\varepsilon \in \{0, 1\}$), we define for $X \in \operatorname{Sp}_{C_3}$,

$$\pi_{i+j\lambda+\varepsilon}^{C_3}(X) \coloneqq \left[S^{i+j\lambda} \otimes (S^{\succ})^{\varepsilon}, X\right]^{C_3}.$$

In this subsection, we will use $\pi_{\bigstar}^{C_3}$ to denote $RO^{\succ}(C_3)$ -graded homotopy groups. If $R \in \operatorname{Sp}_{C_3}$ is a ring spectrum, then $\pi_{\bigstar}^{C_3}R$ is a module over $\pi_{\bigstar}^{C_3}R$. Let

$$a_{\succ} \colon S^{-\succ} \to S^0$$

denote the Spanier-Whitehead dual of the second map of (5.2.2). The λ -suspension of this map is the canonical inclusion

$$a_{\succ} \colon S^{\succ} \hookrightarrow S^{\lambda}.$$

If we abusively also use a_{\succ} to denote the Spanier-Whitehead dual of this map

$$a_{\succ} \colon S^{-\lambda} \to S^{-\flat}$$

then it makes sense to form a_{\succ}^2 , and we have

$$a_{\succ}^2 = a_{\lambda}.$$

For any $X \in \operatorname{Sp}_{C_3}$, the $RO^{\succ}(C_3)$ -graded homotopy groups $\pi_{\bigstar}^{C_3}$ are a module over $\mathbb{Z}[a_{\succ}]$.

Our goal is to compute the $RO^{\succ}(C_3)$ -graded MANSS with signature

$${}^{\text{MANSS}}E^{s,\bigstar}(\text{tmf}(2)_3^h) \implies \pi_{\bigstar}^{C_3}\text{tmf}(2)_3^h.$$

Consider the second double complex spectral sequence

$${}_{II}E^{a,b,\bigstar}(\operatorname{tmf}(2)_3^h) = H^a(C_3; \operatorname{Ext}^{b,a+b}((MU_3)_*(\operatorname{tmf}(2)_3 \otimes S^{-\bigstar}))) \implies {}^{\operatorname{MANSS}}E^{a+b,\bigstar}(\operatorname{tmf}(2)_3^h) = 44$$

Since $\pi_* \operatorname{tmf}(2)_3$ is concentrated in even degrees, it is complex orientable, and therefore its ANSS collapses with $\operatorname{Ext}^{b,*}((MU_3)_*\operatorname{tmf}(2)_3) = 0$ for b > 0. In particular, the second double complex spectral sequence collapses to give an isomorphism

(5.2.3)
$$^{\text{MANSS}}E_2^{s,\bigstar}(\operatorname{tmf}(2)_3^h) \cong H^s(C_3; \pi_s(\operatorname{tmf}(2)_3 \otimes S^{-\bigstar})).$$

It follows from Proposition 4.1.8 that the MANSS is isomorphic to the HFPSS. We first identify the E_2 -term of this spectral sequence.

Proposition 5.2.4. There are preferred generators c_4 and c_6 and an additive isomorphism

$$(5.2.5) \quad ^{\text{MANSS}} E^{*, \bigstar} (\operatorname{tmf}(2)_{3}^{h}) \cong \\ \frac{\pi_{\bigstar}^{C_{3}} H \mathbb{Z}_{3}^{h} [c_{4}, c_{6}, \Delta^{1/2}] / (a_{\succ} c_{4}, a_{\succ} c_{6}, c_{4}^{3} - c_{6}^{2} - 12^{3} (\Delta^{1/2})^{2}) \{ \overline{v}_{1}, \overline{v}_{1}', u_{\succ} c_{4}, u_{\succ}' c_{4}, u_{\succ} c_{6}, u_{\succ}' c_{6} \}}{(c_{4} \cdot \overline{v}_{1}, c_{6} \cdot \overline{v}_{1}, a_{\succ} \cdot \overline{v}_{1}', c_{6} \cdot \overline{v}_{1}', a_{\succ} \cdot u_{\succ} c_{4}, a_{\succ} \cdot u_{\succ}' c_{4}, a_{\succ} \cdot u_{\succ} c_{6}, a_{\succ} \cdot u_{\succ}' c_{6}, c_{6} \cdot u_{\succ} c_{6})}$$

Furthermore the inclusion

$$\pi_{\bigstar}^{C_3} H \mathbb{Z}_3^h[c_4, c_6, \Delta^{1/2}] / (a_{\succ} c_4, a_{\succ} c_6, c_4^3 - c_6^2 - 12^3 (\Delta^{1/2})^2) \hookrightarrow {}^{\text{MANSS}} E^{*,\bigstar}(\operatorname{tmf}(2)_3^h)$$

is an inclusion of rings, and the isomorphism (5.2.5) is an isomorphism of modules over this subring.

Proof. Our first task will be to compute $\pi_* \operatorname{tmf}(2)_3$ as a C_3 -module. Let \mathbb{Z}^{\succ} denote the kernel of the augmentation $\mathbb{Z}[C_3] \to \mathbb{Z}$. Note that we have an isomorphism of C_3 -modules:

$$H_1(S^{\succ};\mathbb{Z})\cong\mathbb{Z}^{\succ}$$

As a $\mathbb{Z}[C_3]$ -module we have

$$\pi_{0} \operatorname{tmf}(2)_{3} = \mathbb{Z}_{3}\{1\}$$

$$\pi_{4} \operatorname{tmf}(2)_{3} = \mathbb{Z}_{3}^{\succ}\{e_{1}\}$$

$$\pi_{8} \operatorname{tmf}(2)_{3} = \mathbb{Z}_{3}[C_{3}]\{e_{1}^{2}\}$$

$$\pi_{12} \operatorname{tmf}(2)_{3} = \mathbb{Z}_{3}\{\Delta^{1/2}\} \oplus \mathbb{Z}_{3}[C_{3}]\{e_{1}^{2}e_{2}\}$$

$$\pi_{16} \operatorname{tmf}(2)_{3} = \mathbb{Z}_{3}^{\succ}\{\Delta^{1/2}e_{1}\} \oplus \mathbb{Z}_{3}[C_{3}]\{e_{1}^{4}\}$$

$$\pi_{20} \operatorname{tmf}(2)_{3} = \mathbb{Z}_{3}[C_{3}]\{\Delta^{1/2}e_{1}^{2}, e_{1}^{4}e_{2}\}$$

$$\vdots$$

In general, we have an isomorphism of C_3 -modules

(5.2.6)
$$\pi_* \operatorname{tmf}(2)_3 \cong \mathbb{Z}_3[\Delta^{1/2}] \otimes \left(\mathbb{Z}\{1\} \oplus \mathbb{Z}^{\succ}\{e_1\} \oplus \mathbb{Z}[C_3]\{e_1^{2i}e_2^{\varepsilon} : i > 0, \varepsilon \in \{0, 1\}\}\right).$$

Note that we have (using the analog of [BMMS86, Theorem V.2.4])

$$\pi_{i+j\lambda+\varepsilon}^{C_3} H \mathbb{Z}_3^h \cong H^{-i}((EC_3)_+ \otimes_{C_3} S^{j\lambda+\varepsilon}; \mathbb{Z}_3)$$
$$\cong H^{-i-2j-\varepsilon}(C_3; \mathbb{Z}_3^{\varepsilon}).$$

It follows from elementary group cohomology computations that we have

(5.2.7)
$$\pi_{\bigstar}^{C_3} H \mathbb{Z}_3^h \cong \mathbb{Z}_3[u_{\lambda}^{\pm}, a_{\succ}]/(3a_{\succ})$$

with

$$|u_{\lambda}| = 2 - \lambda$$
, and $|a_{\succ}| = - \succ$.

Observe that we have

$$^{\text{MANSS}}E^{0,\lambda+\succ+1}(\text{tmf}(2)_3) = H^0(C_3;\pi_0(\text{tmf}(2)_3 \otimes S^{-\lambda-\succ-1}))$$
$$\cong H^0(C_3;\mathbb{Z}_3^{\succ} \otimes \mathbb{Z}^{\succ})$$
$$\cong H^0(C_3;\mathbb{Z}_3) \oplus H^0(C_3;\mathbb{Z}_3[C_3])$$
$$\cong \mathbb{Z}_3 \oplus \mathbb{Z}_3.$$

We let \overline{v}_1 denote a generator of the first summand, and \overline{v}'_1 denote a generator of the second summand. We let c_4 be a generator of

$$^{\text{MANSS}} E^{0,8}(\text{tmf}(2)_3^h) = H^0(C_3; \pi_8 \text{tmf}_0(2)_3)$$

= $H^0(C_3; \mathbb{Z}_3[C_3]\{e_1^2\})$
= $\mathbb{Z}_3,$

and let $u_{\succ}c_4$ and $u'_{\succ}c_4$ denote the generators of

$$^{\text{MANSS}}E^{0,7+\succ}(\text{tmf}(2)_3^h) = H^0(C_3; \pi_7 \text{tmf}_0(2)_3 \otimes S^{-\succ})$$
$$= H^0(C_3; \mathbb{Z}_3[C_3]\{e_1^2\} \otimes \mathbb{Z}^{\succ})$$
$$= \mathbb{Z}_3 \oplus \mathbb{Z}_3.$$

Similarly we take c_6 to be a generator of the second summand of

^{MANSS}
$$E^{0,12}(\operatorname{tmf}(2)_3^h) = H^0(C_3; \pi_{12} \operatorname{tmf}_0(2)_3)$$

= $H^0(C_3; \mathbb{Z}_3\{\Delta^{1/2}\} \oplus \mathbb{Z}_3[C_3]\{e_1^2e_2\})$
= $\mathbb{Z}_3 \oplus \mathbb{Z}_3$

where the first summand is generated by $\Delta^{1/2}$. We let $u_{\succ}c_6$ and $u'_{\succ}c_6$ be the generators of the first and second summands of

$$^{\text{MANSS}}E^{0,11+\succ}(\text{tmf}(2)_3^h) = H^0(C_3; \pi_{11}\text{tmf}_0(2)_3 \otimes S^{-\succ})$$
$$= H^0(C_3; \mathbb{Z}_3\{\Delta^{1/2}\} \otimes \mathbb{Z}^{\succ} \oplus \mathbb{Z}_3[C_3]\{e_1^2e_2\} \otimes \mathbb{Z}^{\succ})$$
$$= \mathbb{Z}_3 \oplus \mathbb{Z}_3.$$

If we take c_4 and c_6 to be the generators coming from the image of

$$\pi_* \operatorname{tmf}_3 \to \pi_* \operatorname{tmf}(2)_3$$

then the result follows from (5.2.3), (5.2.6), and (5.2.7).

The μ_3 -orientation of tmf(2) in [HSW23, Theorem 1.9.] gives rise to a C_3 -equivariant map

$$\overline{v}_1: S^{\lambda + \succ + 1} \to \operatorname{tmf}(2)_3$$

which is an isomorphism on π_4^e (after 3-completion). Since tmf(2)₃ is a C_3 -commutative ring spectrum, the norm-restriction adjunction [HHR16, Prop. 2.27] gives rise to a map

$$N_e^{C_3} \operatorname{Res}_e^{C_3} \operatorname{tmf}(2)_3 \to \operatorname{tmf}(2)_3.$$

Therefore, we may norm the element $e_1 \in \pi_4^e \operatorname{tmf}(2)_3$ to get an equivariant enhancement of $\Delta^{1/2}$:

$$\overline{\Delta}^{1/2} \coloneqq -4N^{C_3}(e_1) \in \pi^{C_3}_{4\rho} \operatorname{tmf}(2)_3.$$

Proposition 5.2.8. In the MANSS for $tmf(2)_3$ there are differentials

$$d_{5}(u_{\lambda}^{8}(\overline{\Delta}^{1/2})^{2}) \doteq a_{\succ}^{5}u_{\lambda}^{7}(\overline{\Delta}^{1/2})^{2},$$
$$d_{5}(u_{\lambda}) \doteq a_{\succ}^{5}\overline{v}_{1},$$
$$d_{9}(a_{\succ}u_{\lambda}^{17}\overline{v}_{1}(\overline{\Delta}^{1/2})^{4}) \doteq a_{\succ}^{10}u_{\lambda}^{15}(\overline{\Delta}^{1/2})^{5}$$
$$d_{9}(u_{\lambda}^{2}\overline{v}_{1}) \doteq a_{\succ}^{9}\overline{\Delta}^{1/2}.$$

These generate all of the differentials in the MANSS under the Leibniz rule and in particular we find that

$$u_{\lambda}^{i}u_{\succ}^{\varepsilon}c_{4}, u_{\succ}^{\prime}c_{4}, u_{\lambda}^{i}u_{\succ}^{\varepsilon}c_{6}, u_{\succ}^{\prime}c_{6}, u_{\lambda}\overline{v}_{1}, \overline{v}_{1}^{\prime}, u_{\lambda}^{3}$$

are permanent cycles. It follows that $(\Delta^{1/2})^3 = u_{\lambda}^{12} (\overline{\Delta}^{1/2})^3$ is a permanent cycle, and the MANSS is therefore $(\Delta^{1/2})^3$ -periodic.

Proof. The HFPSS/MANSS for $tmf(2)_3$ maps to the HFPSS for $TMF(2)_3$ converging to the homotopy groups of

$$\mathrm{TMF}(2)_3^{hC_3} \simeq \mathrm{TMF}_3 \vee \Sigma^{36} \mathrm{TMF}_3.$$

The structure of the HFPSS for TMF(2)₃ in integer degrees is well known. It is a shifted direct sum of two copies of the HFPSS which converges to π_* TMF₃ [Bau08]. The E_2 -term is given by

$${}^{HFPSS}E_2^{s,t}(\text{TMF}(2)_3) \cong H^s(C_3; \pi_t \text{TMF}(2)_3) \cong \frac{\mathbb{Z}_3[c_4, c_6, \Delta^{\pm 1/2}, \beta] \otimes E[\alpha]}{(c_4^3 - c_6^2 - 12^3 (\Delta^{1/2})^2, 3\alpha, 3\beta, \alpha c_4, \alpha c_6, \beta c_4, \beta c_6)}$$

and under the map from the MANSS for $tmf(2)_3$ to the HFPSS for $TMF_0(2)$ we have

$$a_{\succ}u_{\lambda}\overline{v}_{1} \mapsto \alpha$$
, and $a_{\succ}^{2}u_{\lambda}^{3}\overline{\Delta}^{1/2} \mapsto \beta$.

In the HFPSS for $\text{TMF}(2)_3$ there are differentials

$$d_5(\Delta) \doteq \alpha \beta^2$$
, and $d_9(\Delta^2 \alpha) \doteq \beta^5$.

Here, we use \doteq to indicate both sides of the equation are equal up multiplication by an element of \mathbb{Z}_3^{\times} . This implies that in the MANSS for tmf(2)₃, we have differentials

$$d_5(u_{\lambda}^8(\overline{\Delta}^{1/2})^2) \doteq a_{\succ}^5 u_{\lambda}^7(\overline{\Delta}^{1/2})^2,$$

$$d_9(a_{\succ} u_{\lambda}^{17} \overline{v}_1(\overline{\Delta}^{1/2})^4) \doteq a_{\succ}^{10} u_{\lambda}^{15}(\overline{\Delta}^{1/2})^5.$$

The element a_{\succ} is a permanent cycle because it is in the Hurewicz image, and the elements \overline{v}_1 and $\overline{\Delta}^{1/2}$ are permanent cycles because we constructed elements of $\pi_{\bigstar}^{C_3} \operatorname{tmf}(2)_3^h$ which these converge to. It follows that we have differentials

$$d_5(u_{\lambda}) \doteq a_{\succ}^5 \overline{v}_1$$
, and $d_9(u_{\lambda}^2 \overline{v}_1) \doteq a_{\succ}^9 \overline{\Delta}^{1/2}$.

By dimension considerations, we observe that these differentials generate all of the differentials in the MANSS under the Leibniz rule and the result follows. \Box

A deformation of Borel equivariant homotopy

References

- [AKS18] Gabe Angelini-Knoll and Andrew Salch. A May-type spectral sequence for higher topological Hochschild homology. *Algebr. Geom. Topol.*, 18(5):2593–2660, 2018.
- [Bau08] Tilman Bauer. Computation of the homotopy of the spectrum tmf. In Groups, homotopy and configuration spaces, volume 13 of Geom. Topol. Monogr., pages 11–40. Geom. Topol. Publ., Coventry, 2008.
- [Beh06] Mark Behrens. Root invariants in the Adams spectral sequence. Trans. Amer. Math. Soc., 358(10):4279–4341, 2006.
- [Beh20] Mark Behrens. Topological modular and automorphic forms. In *Handbook of homotopy theory*, CRC Press/Chapman Hall Handb. Math. Ser., pages 221–261. CRC Press, Boca Raton, FL, 2020.
- [BHS20] Robert Burklund, Jeremy Hahn, and Andrew Senger. Galois reconstruction of Artin-Tate ℝ-motivic spectra, 2020. Preprint: arXiv:2010.10325.
- [BHS22] Robert Burklund, Jeremy Hahn, and Andrew Senger. On the boundaries of highly connected, almost closed manifolds, 2022. Preprint: 1910.14116.
- [BMMS86] R. R. Bruner, J. P. May, J. E. McClure, and M. Steinberger. H_{∞} ring spectra and their applications, volume 1176 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1986.
- [Boa99] J. Michael Boardman. Conditionally convergent spectral sequences. In Homotopy invariant algebraic structures. A conference in honor of J. Michael Boardman. AMS special session on homotopy theory, Baltimore, MD, USA, January 7–10, 1998, pages 49–84. Providence, RI: American Mathematical Society, 1999.
- [Day71] Brian Day. Construction of biclosed categories. Bull. Aust. Math. Soc., 5:139–140, 1971.
- [DI10] Daniel Dugger and Daniel C. Isaksen. The motivic Adams spectral sequence. *Geom. Topol.*, 14(2):967–1014, 2010.
- [GHIR20] Bertrand J. Guillou, Michael A. Hill, Daniel C. Isaksen, and Douglas Conner Ravenel. The cohomology of C_2 -equivariant $\mathcal{A}(1)$ and the homotopy of ko_{C2}. Tunis. J. Math., 2(3):567–632, 2020.
- [GIKR22] Bogdan Gheorghe, Daniel C. Isaksen, Achim Krause, and Nicolas Ricka. C-motivic modular forms. J. Eur. Math. Soc. (JEMS), 24(10):3597–3628, 2022.
- [Gla16] Saul Glasman. Day convolution for ∞ -categories. Math. Res. Lett., 23(5):1369–1385, 2016.
- [GM95] J. P. C. Greenlees and J. P. May. Generalized Tate cohomology, volume 543 of Mem. Am. Math. Soc. Providence, RI: American Mathematical Society (AMS), 1995.
- [GP18] Owen Gwilliam and Dmitri Pavlov. Enhancing the filtered derived category. J. Pure Appl. Algebra, 222(11):3621–3674, 2018.
- [GR23] Thomas Gregersen and John Rognes. On the motivic Segal conjecture, 2023. Preprint: arXiv:2302.12659.
- [GWX21] Bogdan Gheorghe, Guozhen Wang, and Zhouli Xu. The special fiber of the motivic deformation of the stable homotopy category is algebraic. *Acta Math.*, 226(2):319–407, 2021.
- [Hea19] Drew Heard. On equivariant and motivic slices. Algebr. Geom. Topol., 19(7):3641–3681, 2019.
- [Hed20] Alice Hedenlund. *Multiplicative Tate Spectral Sequences*. PhD thesis, University of Oslo, 2020.
- [HHR11] Michael A. Hill, Michael J. Hopkins, and Douglas C. Ravenel. On the 3-primary Arf-Kervaire invariant problem, 2011. https://people.math.rochester.edu/faculty/doug/mypapers/odd.pdf.
- [HHR16] Michael A Hill, Michael J Hopkins, and Douglas C Ravenel. On the nonexistence of elements of Kervaire invariant one. Annals of Mathematics, pages 1–262, 2016.
- [HK01] Po Hu and Igor Kriz. Real-oriented homotopy theory and an analogue of the Adams-Novikov spectral sequence. *Topology*, 40(2):317–399, 2001.
- [HKO11] Po Hu, Igor Kriz, and Kyle Ormsby. Remarks on motivic homotopy theory over algebraically closed fields. J. K-Theory, 7(1):55–89, 2011.
- [HM17] Michael A. Hill and Lennart Meier. The C_2 -spectrum Tmf₁(3) and its invertible modules. Algebr. Geom. Topol., 17(4):1953–2011, 2017.
- [HSW23] Jeremy Hahn, Andrew Senger, and Dylan Wilson. Odd primary analogs of real orientations. Geom. Topol., 27(1):87–129, 2023.
- [HY18] Michael Hill and Carolyn Yarnall. A new formulation of the equivariant slice filtration with applications to C_p -slices. Proceedings of the American Mathematical Society, 146(8):3605–3614, 2018.
- [IWX23] Daniel C. Isaksen, Guozhen Wang, and Zhouli Xu. Stable homotopy groups of spheres: from dimension 0 to 90. Publ. Math. Inst. Hautes Études Sci., 137:107–243, 2023.
- [Joy02] A. Joyal. Quasi-categories and Kan complexes. J. Pure Appl. Algebra, 175(1-3):207–222, 2002.

- [Kon20] Hana Jia Kong. The C_2 -effective spectral sequence for C_2 -equivariant connective real K-theory, 2020. Preprint: arXiv:2004.00806.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Lur17] Jacob Lurie. Higher algebra. https://www.math.ias.edu/~lurie/papers/HA.pdf, 2017. Electronic book.
- [May01] J. P. May. The additivity of traces in triangulated categories. Adv. Math., 163(1):34–73, 2001.
- [McC01] John McCleary. A user's guide to spectral sequences, volume 58 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, second edition, 2001.
- [Mil81] Haynes R. Miller. On relations between Adams spectral sequences, with an application to the stable homotopy of a Moore space. J. Pure Appl. Algebra, 20:287–312, 1981.
- [MNN17] Akhil Mathew, Niko Naumann, and Justin Noel. Nilpotence and descent in equivariant stable homotopy theory. *Adv. Math.*, 305:994–1084, 2017.
- [MV99] Fabien Morel and Vladimir Voevodsky. A¹-homotopy theory of schemes. Inst. Hautes Études Sci. Publ. Math., (90):45−143, 1999.
- [PP21] Piotr Pstragowski and Irakli Patchkoria. Adams spectral sequences and Franke's algebraicity conjecture, 2021. Preprint: arXiv:2110.03669.
- [Pst23] Piotr Pstragowski. Synthetic spectra and the cellular motivic category. *Invent. Math.*, 232(2):553–681, 2023.
- [QS21] J. D. Quigley and Jay Shah. On the parametrized Tate construction, 2021. Preprint: arXiv:2110.07707.
- [Qui19] J. D. Quigley. The motivic Mahowald invariant. Algebr. Geom. Topol., 19(5):2485–2534, 2019.
- [Sto12] Vesna Stojanoska. Duality for topological modular forms. Doc. Math., 17:271–311, 2012.
- [Ull13] John Richard Ullman. On the regular slice spectral sequence. PhD thesis, Massachusetts Institute of Technology, 2013.
- [Voe02] Vladimir Voevodsky. Open problems in the motivic stable homotopy theory. I. In Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998), volume 3 of Int. Press Lect. Ser., pages 3–34. Int. Press, Somerville, MA, 2002.
- [Wil17] Dylan Wilson. Equivariant, Parameterized, and Chromatic Homotopy Theory. Phd thesis, Northwestern University, Evanston, IL, June 2017.

Département de Mathématiques, Institut Galilée, Université Paris 13, 99 av. J.B. Clément, FR-93430 Villetaneuse, France

Email address: angelini-knoll@math.univ-paris13.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN, USA *Email address:* mbehren1@nd.edu

DEPARTMENT OF MATHEMATICS, APPLIED MATHEMATICS, AND STATISTICS, CASE WESTERN RESERVE UNIVERSITY, CLEVELAND, OH, USA

 $Email \ address: \verb"eva.belmont@case.edu"$

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA, USA *Email address*: hana.jia.kong@gmail.com