

Model-independent reconstruction of the primordial curvature power spectrum from PTA data

Zhu Yi,^a Zhi-Qiang You,^{b,*} and You Wu^{c,d,*}

^aAdvanced Institute of Natural Sciences, Beijing Normal University, Zhuhai 519087, China

^bHenan Academy of Sciences, Zhengzhou 450046, Henan, China

^cCollege of Mathematics and Physics, Hunan University of Arts and Science, Changde, 415000, China

^dDepartment of Physics and Synergistic Innovation Center for Quantum Effects and Applications, Hunan Normal University, Changsha, Hunan 410081, China

E-mail: yz@bnu.edu.cn, you_zhiqiang@whu.edu.cn, youwuphy@gmail.com

Abstract. Recently released data from pulsar timing array (PTA) collaborations provide strong evidence for a stochastic signal consistent with a gravitational-wave background, potentially originating from scalar-induced gravitational waves (SIGWs). However, in order to determine whether the SIGWs with a specific power spectrum of curvature perturbations can account for the PTA signal, one needs to estimate the energy density of the SIGWs, which can be computationally expensive. In this paper, we use a model-independent approach to reconstruct the primordial curvature power spectrum using a free spectrum cross over from 10^1 Mpc^{-1} to 10^{20} Mpc^{-1} with NANOGrav 15-year data set. Our results can simplify the task of assessing whether a given primordial curvature power spectrum can adequately explain the observed PTA signal without calculating the energy density of SIGWs.

*Corresponding author.

Contents

1	Introduction	1
2	The scalar-induced gravitational waves	2
3	The model-independent parameterization	4
4	Methodology and Results	6
5	Conclusion	7

1 Introduction

The detection of a common-spectrum process demonstrating the Hellings-Downs angular correlation feature inherent in gravitational waves (GWs) has been reported by the North American Nanohertz Observatory for Gravitational Waves (NANOGrav) [1, 2], Parkers Pulsar Timing Array (PPTA) [3, 4], European Pulsar Timing Array (EPTA) [5, 6], and Chinese Pulsar Timing Array (CPTA) [7], recently. Assuming the source to be an ensemble of binary supermassive black hole inspirals and adopting a fiducial $f^{-2/3}$ characteristic-strain spectrum, its estimated strain amplitude is approximately $\sim 10^{-15}$ at a reference frequency of 1 yr^{-1} [1, 4, 6, 7]. However, the precise source of this intriguing signal, whether attributed to supermassive black hole binaries (SMBHBs) or other cosmological phenomena, remains a subject of active investigation [8–29]. Among the various hypotheses, a particularly promising candidate is the scalar-induced gravitational waves (SIGWs); and this paper focuses on investigating the scenario where the observed signal arises from SIGWs. For other physical processes contributing to the PTA band, please see [30–43].

SIGWs, accompanied by the formation of primordial black holes (PBHs), originate from primordial curvature perturbations generated during the inflationary epoch [44–87]. The requirement for significant SIGWs is a power spectrum amplitude for primordial curvature perturbations, approximately $\mathcal{A}_\zeta \sim \mathcal{O}(0.01)$, which is seven orders of magnitude larger than constraints from cosmic microwave background (CMB) anisotropy observations, $\mathcal{A}_\zeta = 2.1 \times 10^{-9}$ [88]. This rapid enhancement in the primordial curvature power spectrum can be accomplished through inflation models with a transitional ultra-slow-roll phase [89–124]. Detectable not only by PTAs but also by space-based GW detectors such as LISA [125, 126], Taiji [127], TianQin [128, 129], and DECIGO [130], SIGWs provide invaluable insights into the physics of the early Universe. For instance, through the observational data of SIGWs, constraints on primordial curvature perturbations at small scales can be obtained [131, 132].

In exploring the possibility of SIGWs as an explanation for the PTA signal, the primordial curvature power spectrum is commonly characterized through special models, such as the δ -function, lognormal form, or broken power law parameterizations

[8, 18]. By employing these parameterizations and using the PTA data, one can derive constraints on the primordial curvature power spectrum at small scales. However, a drawback of this approach lies in its dependency on the specific parameterizations of the curvature power spectrum. To overcome this limitation, in this paper, we adopt a linear interpolation parameterization with a large number of bins. The linear interpolation parameterization is versatile enough to depict various types of primordial curvature power spectra, as long as the number of bins is sufficiently large. By employing linear interpolation parameterization, researchers can analyze and interpret data in a more general way, allowing for the exploration of a broader range of possibilities and reducing the influence of preconceived assumptions about the underlying functional form of the power spectrum. This near-model-independent methodology can provide more robust constraints on the primordial curvature power spectrum. By this parameterization, we reconstruct the primordial curvature power spectrum, and the amplitude is displayed in a free-spectrum framework in this paper.

The organization of this paper is as follows. Section 2 gives the energy density of the SIGWs, and section 3 introduces the model-independent parameterization of the primordial curvature power spectrum. In Section 4, we present our results, and the conclusions are drawn in Section 5.

2 The scalar-induced gravitational waves

If the scalar perturbations seeded from the primordial curvature perturbations generated during inflation are large enough, the secondary order of the linear scalar perturbations can act as a source to induce GWs. In this section, we give the energy density of SIGWs during radiation domination in detail. The perturbed metric in the Newtonian gauge can be expressed as

$$ds^2 = a^2(\eta) \left[-(1 + 2\Phi)d\eta^2 + \left\{ (1 - 2\Phi)\delta_{ij} + \frac{1}{2}h_{ij} \right\} dx^i dx^j \right], \quad (2.1)$$

where Φ represents the Bardeen potential, and h_{ij} is the tensor perturbation following the transverse-traceless gauge condition, $\partial^i h_{ij} = h_i^i = 0$. In the Fourier space, the tensor perturbation can be represented as

$$h_{ij}(\mathbf{x}, \eta) = \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} e^{i\mathbf{k}\cdot\mathbf{x}} [h_{\mathbf{k}}^+(\eta) e_{ij}^+(\mathbf{k}) + h_{\mathbf{k}}^\times(\eta) e_{ij}^\times(\mathbf{k})]. \quad (2.2)$$

Here, $e_{ij}^+(\mathbf{k})$ and $e_{ij}^\times(\mathbf{k})$ represent the plus and cross-polarization tensors,

$$\begin{aligned} e_{ij}^+(\mathbf{k}) &= \frac{1}{\sqrt{2}} [e_i(\mathbf{k}) e_j(\mathbf{k}) - \bar{e}_i(\mathbf{k}) \bar{e}_j(\mathbf{k})], \\ e_{ij}^\times(\mathbf{k}) &= \frac{1}{\sqrt{2}} [e_i(\mathbf{k}) \bar{e}_j(\mathbf{k}) + \bar{e}_i(\mathbf{k}) e_j(\mathbf{k})], \end{aligned} \quad (2.3)$$

and $e_i(\mathbf{k})$ and $\bar{e}_i(\mathbf{k})$ are orthonormal basis vectors orthogonal to the wave vector \mathbf{k} , $\mathbf{e} \cdot \bar{\mathbf{e}} = \mathbf{e} \cdot \mathbf{k} = \bar{\mathbf{e}} \cdot \mathbf{k} = 0$.

Disregarding anisotropic stress, the equation of motion for tensor perturbations in Fourier space, for either polarization, takes the form

$$h_{\mathbf{k}}''(\eta) + 2\mathcal{H}h_{\mathbf{k}}'(\eta) + k^2 h_{\mathbf{k}}(\eta) = 4S_{\mathbf{k}}(\eta), \quad (2.4)$$

where η is the conformal time, $d\eta = dt/a$, a prime denotes the derivative with respect to the conformal time, $\mathcal{H} = a'/a$ represents the conformal Hubble parameter, and the source term is given by

$$S_{\mathbf{k}} = \int \frac{d^3\tilde{\mathbf{k}}}{(2\pi)^{3/2}} e_{ij}(\mathbf{k}) \tilde{k}^i \tilde{k}^j \left[2\Phi_{\tilde{\mathbf{k}}}\Phi_{\mathbf{k}-\tilde{\mathbf{k}}} + \frac{1}{\mathcal{H}^2} (\Phi_{\tilde{\mathbf{k}}} + \mathcal{H}\Phi_{\tilde{\mathbf{k}}}) (\Phi'_{\mathbf{k}-\tilde{\mathbf{k}}} + \mathcal{H}\Phi_{\mathbf{k}-\tilde{\mathbf{k}}}) \right], \quad (2.5)$$

where $\Phi_{\mathbf{k}}$ is the Bardeen potential in the Fourier space. By using the primordial curvature perturbations $\zeta_{\mathbf{k}}$ generated in the inflation, the Bardeen potential can be expressed as

$$\Phi_{\mathbf{k}} = \frac{3 + 3w}{5 + 3w} T(k, \eta) \zeta_{\mathbf{k}}, \quad (2.6)$$

where $T(k, \eta)$ is the transfer function,

$$T(k, \eta) = 3 \left[\frac{\sin(k\eta/\sqrt{3}) - (k\eta/\sqrt{3}) \cos(k\eta/\sqrt{3})}{(k\eta/\sqrt{3})^3} \right]. \quad (2.7)$$

The solution for $h_{\mathbf{k}}(\eta)$ in equation (2.4) can be obtained through the Green's function method,

$$h_{\mathbf{k}}(\eta) = \frac{4}{a(\eta)} \int_0^\eta d\bar{\eta} G_{\mathbf{k}}(\eta, \bar{\eta}) a(\bar{\eta}) S_{\mathbf{k}}(\bar{\eta}), \quad (2.8)$$

where the Green function is

$$G_{\mathbf{k}}(\eta, \bar{\eta}) = \frac{\sin[k(\eta - \bar{\eta})]}{k}. \quad (2.9)$$

The definition of the power spectrum of SIGWs is

$$\langle h_{\mathbf{k}}(\eta) h_{\mathbf{p}}(\eta) \rangle = \delta^3(\mathbf{k} + \mathbf{p}) \frac{2\pi^2}{k^3} \mathcal{P}_h(k, \eta). \quad (2.10)$$

Combining Eqs. (2.5)-(2.10), we get [46, 47, 52, 133–135]

$$\mathcal{P}_h(k, \eta) = 4 \int_0^\infty dv \int_{|1-v|}^{1+v} du \left[\frac{4v^2 - (1 - u^2 + v^2)^2}{4uv} \right]^2 I_{RD}^2(u, v, x) \mathcal{P}_\zeta(kv) \mathcal{P}_\zeta(ku), \quad (2.11)$$

where \mathcal{P}_ζ is the primordial curvature power spectrum with the definition,

$$\langle \zeta_{\mathbf{k}} \zeta_{\mathbf{p}} \rangle = \delta^3(\mathbf{k} + \mathbf{p}) \frac{2\pi^2}{k^3} \mathcal{P}_\zeta. \quad (2.12)$$

Here, $u = |\mathbf{k} - \tilde{\mathbf{k}}|/k$, $v = |\tilde{\mathbf{k}}|/k$, $x = k\eta$, and

$$I_{\text{RD}}(u, v, x) = \int_0^x dy y \sin(x-y) \{3T(uy)T(vy) + y[T(vy)uT'(uy) + vT'(vy)T(uy)] + y^2 uvT'(uy)T'(vy)\}, \quad (2.13)$$

where the subscript ‘‘RD’’ denotes the radiation-dominated era with the equation of state satisfying $w = 1/3$. The energy density of Gravitational waves is

$$\Omega_{\text{GW}}(k, \eta) = \frac{1}{24} \left(\frac{k}{aH} \right)^2 \overline{\mathcal{P}_h(k, \eta)}. \quad (2.14)$$

Combining equation (2.11) and definition (2.14), we obtain [133, 135]

$$\Omega_{\text{GW}}(k, \eta) = \frac{1}{6} \left(\frac{k}{aH} \right)^2 \int_0^\infty dv \int_{|1-v|}^{1+v} du \left[\frac{4v^2 - (1 - u^2 + v^2)^2}{4uv} \right]^2 \overline{I_{\text{RD}}^2(u, v, x)} \mathcal{P}_\zeta(kv) \mathcal{P}_\zeta(ku), \quad (2.15)$$

where $\overline{I_{\text{RD}}^2}$ is the oscillation time average of I_{RD}^2 . At late time, $k\eta \gg 1$, i.e., $x \rightarrow \infty$, the time average of $I_{\text{RD}}^2(u, v, x \rightarrow \infty)$ is [52]

$$\begin{aligned} \overline{I_{\text{RD}}^2(v, u, x \rightarrow \infty)} &= \frac{1}{2x^2} \left(\frac{3(u^2 + v^2 - 3)}{4u^3v^3} \right)^2 \left\{ \pi^2 (u^2 + v^2 - 3)^2 \Theta(v + u - \sqrt{3}) \right. \\ &\quad \left. + \left(-4uv + (u^2 + v^2 - 3) \ln \left| \frac{3 - (u+v)^2}{3 - (u-v)^2} \right| \right)^2 \right\}. \end{aligned} \quad (2.16)$$

The energy density evolution of gravitational waves follows the same way as that of radiation. Using this property, we can easily determine the current energy density of gravitational waves

$$\Omega_{\text{GW}}(k, \eta_0) = \frac{\Omega_{r,0} \Omega_{\text{GW}}(k, \eta)}{\Omega_r(\eta)}, \quad (2.17)$$

where $\Omega_{r,0}$ represents the current energy density of radiation, and $\Omega_r(\eta) = 1$ during the generation of SIGWs.

3 The model-independent parameterization

In investigating the issue of SIGWs, the primordial curvature power spectra are often characterized using specific forms, where the most frequently employed parameterizations are the δ -function, lognormal form, and broken power law parameterizations. Consequently, the outcomes of such investigations are dependent on these specific parameterizations. Hence, the exploration of a model-independent or, at the very least, near-model-independent approach is indeed valuable. Following this principle, we parameterize the primordial curvature power spectrum using a linear interpolation form

with a total of n bins. The linear interpolation parameterization is robust to depict different kinds of primordial curvature power spectra, as long as the number of bins is large enough. While the more bins imply a larger amount of computation, we need to strike a balance between these two contradictions. In the following, we give the precise details of the linear interpolation parameterization of the primordial curvature power spectrum.

On the larger scales where $k \leq k_{\text{large}} = 1 \text{ Mpc}^{-1}$, the primordial curvature power spectrum is already well-constrained by CMB observational data. The primordial curvature power spectrum can be effectively characterized using a power-law form,

$$\mathcal{P}_\zeta(k) = A_{\text{CMB}} \left(\frac{k}{k_*} \right)^{n_s - 1}, \quad (3.1)$$

where, from the CMB observational data, the amplitude is $A_{\text{CMB}} = 2.1 \times 10^{-9}$, the scalar tilt is $n_s = 0.965$, and the pivot scale is $k_* = 0.05 \text{ Mpc}^{-1}$ [88].

For the scales where $k_{\text{large}} < k \leq k_{\text{end}} = 10^{20} \text{ Mpc}^{-1}$, we divide the primordial curvature power spectrum into three distinct regions: $k_{\text{large}} < k < k_{\text{pta1}}$, $k_{\text{pta1}} \leq k \leq k_{\text{pta2}}$, and $k_{\text{pta2}} < k \leq k_{\text{end}}$. We set the scale points k_{pta} as follows

$$\log_{10}(k_{\text{pta1}}/\text{Mpc}^{-1}) = 5, \quad \log_{10}(k_{\text{pta2}}/\text{Mpc}^{-1}) = 10, \quad (3.2)$$

ensuring that the region between these scale points adequately covers the PTA scales. The smallest scale k_{end} is determined by the e -folds of the inflation, e -folds $\approx \ln(k_{\text{end}}/k_*) = 50$. To implement linear interpolation, selecting appropriate data points is essential. For the scales within $k_{\text{large}} < k < k_{\text{pta1}}$, we set the scale points k_i as $\log_{10}(k_{i+1}/\text{Mpc}^{-1}) = \log_{10}(k_i/\text{Mpc}^{-1}) + 1$, beginning with $\log_{10}(k_1/\text{Mpc}^{-1}) = 2$. For the scales within $k_{\text{pta1}} \leq k \leq k_{\text{pta2}}$, we set the scale points k_i as $\log_{10}(k_{i+1}/\text{Mpc}^{-1}) = \log_{10}(k_i/\text{Mpc}^{-1}) + 1/4$ with the initial point $\log_{10}(k_1/\text{Mpc}^{-1}) = \log_{10}(k_{\text{pta1}}/\text{Mpc}^{-1})$. Due to their relevance to PTA scales, additional scale points are chosen for this range. Finally, for scales within $k_{\text{pta2}} < k \leq k_{\text{end}}$, we set the scale points k_i as $\log_{10}(k_{i+1}/\text{Mpc}^{-1}) = \log_{10}(k_i/\text{Mpc}^{-1}) + 1$ with the initial point $\log_{10}(k_1/\text{Mpc}^{-1}) = \log_{10}(k_{\text{pta2}}/\text{Mpc}^{-1}) + 1$. These scale points k_i are displayed in Figure 1.

Using these chosen scale points k_i , for scales where $k_{\text{large}} < k \leq k_{\text{end}} = 10^{20} \text{ Mpc}^{-1}$, the primordial curvature power spectrum can be represented through the linear interpolation form,

$$\log_{10} \mathcal{P}_\zeta(k) = \log_{10} A_i + \frac{\log_{10} A_{i+1} - \log_{10} A_i}{\log_{10} k_{i+1} - \log_{10} k_i} \times (\log_{10} k - \log_{10} k_i), \quad k_i < k \leq k_{i+1}. \quad (3.3)$$

In conclusion, the primordial curvature power spectrum is expressed as

$$\log_{10} \mathcal{P}_\zeta(k) = \begin{cases} \log_{10} A_{\text{CMB}} + (n_s - 1) \log_{10} (k/k_*), & k \leq k_{\text{large}}, \\ \log_{10} A_i + \frac{\log_{10} A_{i+1} - \log_{10} A_i}{\log_{10} k_{i+1} - \log_{10} k_i} \times (\log_{10} k - \log_{10} k_i), & k_i < k \leq k_{i+1}. \end{cases} \quad (3.4)$$

4 Methodology and Results

In this section, we use a Bayesian analysis on the NANOGrav 15-year data set to reconstruct the amplitude of the primordial curvature power spectrum described by Eq. (3.4). We utilize the 14 frequency bins of the NANOGrav 15-year data set [1, 8] to fit the posterior distributions of these amplitudes A_i . The analysis was carried out using the `Bilby` code [136], employing the `dynesty` algorithm for nested sampling [137] with 1000 live points (`nlive` = 1000). The log-likelihood function was obtained by evaluating the energy density of SIGWs at the 14 specific frequency bins. Subsequently, we computed the sum of the logarithms of the probability density functions from 14 independent kernel density estimates [138]. Consequently, the likelihood function can be expressed as

$$\ln \mathcal{L}(\Theta) = \sum_{i=1}^{14} \ln \mathcal{L}_i(\Omega_{\text{GW}}(f_i, \Theta)), \quad (4.1)$$

where Θ represents the collection of parameters in the parameterization given by Eq. (3.4), and the relation of frequency and wave number is

$$f = \frac{k}{2\pi} \simeq 1.6 \text{ nHz} \left(\frac{k}{10^6 \text{ Mpc}^{-1}} \right). \quad (4.2)$$

For scales within the range of $2 \leq \log_{10}(k_i/\text{Mpc}^{-1}) \leq 5$, the priors for the amplitude $\log_{10} A_i$ employ a uniform distribution denoted as $U[-10, \log_{10}(A_c)]$, where A_c is the lower limit of the μ -distortion or BBN constraints at the corresponding scale. For other scales, the prior for each $\log_{10} A_i$ is $U[-10, 0]$, and posterior distributions are illustrated in Figure 1.

Using the linear interpolated model (3.4), the constraints on the continuous primordial curvature power spectrum are actually on the bin amplitude parameters A_i . The violins in Figure 1 illustrate the posteriors of the amplitudes of the primordial curvature power spectrum A_i . Although the full scales information of the primordial curvature power spectrum is contained in SIGWs as demonstrated by the integration of Eq. (2.15), the NANOGrav 15-year data set can only provide constraints on the primordial curvature power spectrum within a specific range of scales, approximately $k/\text{Mpc}^{-1} \sim [10^6, 10^9]$. Outside of this scale range, there are few constraints on the amplitudes of the primordial curvature power spectrum. For the scales $k/\text{Mpc}^{-1} > 13$, the posteriors of the amplitudes A_i are similar with $k/\text{Mpc}^{-1} = 13$ (almost no constraints on amplitudes), so we don't display them in the Figure 1. In Figure 1, the light green shaded region is excluded by the Planck 2018 data [88], and the narrow allowed region was derived using Planck TT, TE, EE+lowE+lensing+BK15 data and 12 knots for the cubic spline interpolation [88]. Because of the high precision measurements of CMB anisotropies by the Planck mission, the constraints on the primordial curvature power spectrum at large scales are quite strong. The orange and gray regions are the constraints from the effect on the ratio between neutron and proton during the Big Bang nucleosynthesis (BBN) [139] and μ -distortion of CMB [140], respectively.

To assess the quality of our model-independent reconstruction, we compare them with model-dependent scenarios, as illustrated in Figure 1. The red line and black




Figure 1. The constraints on the primordial curvature power spectrum. The violins are the allowed regions to account for the NANOGrav 15-year data set in a free-spectrum way. The light green shaded region is excluded by the CMB observations [88]. The orange and gray regions are the constraints from the effect on the ratio between neutron and proton during the Big Bang nucleosynthesis (BBN) [139] and μ -distortion of CMB [140], respectively. The red and black dashed lines are the primordial curvature power spectrum with lognormal (LN) and broken-power-law forms (BPL).

dashed line represent the primordial curvature power spectrum using the lognormal form and the broken power-law form, respectively. These two parameterizations utilize the best-fitted values that account for the NANOGrav 15-year data set via SIGWs [18]. The model-independent reconstruction is consistent with the model-dependent results. Consequently, using this model-independent reconstruction, we can directly assess whether a given primordial curvature power spectrum can explain the NANOGrav 15-yrs signal via SIGWs, without the need for calculating the corresponding SIGWs explicitly. In Figure 1, only two bins can be well constrained, while others are weakly constrained, making it difficult to distinguish different parameterizations of the primordial curvature power spectrum. This is consistent with the previous results that the NANOGrav 15-year data set can not effectively differentiate between various parameterizations of the primordial curvature power spectrum, as discussed in Ref. [8, 18].

5 Conclusion

The signal detected by NANOGrav, PPTA, EPTA, and CPTA collaborations recently can be explained by SIGWs. However, when investigating the plausibility of

SIGWs as an explanation for the PTA signals, the primordial curvature power spectrum often needs to be parameterized in specific forms. As a result, the conclusions drawn depend on the special models. In this paper, we parameterize the primordial curvature power spectrum by using a linear interpolation method. This parameterization is near-model-independent for it can depict a wide array of possible primordial curvature power spectra, as long as the number of bins is large enough. By adopting this linear interpolation parameterization and treating the NANOGrav signal as SIGWs, we utilize Bayesian analysis to obtain the model-independent reconstruction of the primordial curvature power spectrum that could explain the observed PTA signal. The model-independent reconstruction of the primordial curvature power spectrum is displayed through a free-spectrum framework that consists of 34 bins. The model-independent reconstructions are consistent with results obtained when using specific forms of the primordial curvature power spectrum, such as the lognormal and broken power-law forms, to fit the NANOGrav 15-year data set. Therefore, we can now just compare primordial curvature power spectra with the model-independent reconstruction to easily assess their compatibility with the PTA signals, without taking time to calculate the energy density of SIGWs. This streamlines the assessment process and simplifies the task of determining whether a given primordial curvature power spectrum can adequately explain the observed PTA signals.

Acknowledgments

ZY is supported by the National Natural Science Foundation of China under Grant No. 12205015 and the supporting fund for young researcher of Beijing Normal University under Grant No. 28719/310432102. ZQY is supported by the National Natural Science Foundation of China under Grant No. 12305059.

References

- [1] NANOGrav collaboration, *The NANOGrav 15 yr Data Set: Evidence for a Gravitational-wave Background*, *Astrophys. J. Lett.* **951** (2023) L8 [[2306.16213](#)].
- [2] NANOGrav collaboration, *The NANOGrav 15 yr Data Set: Observations and Timing of 68 Millisecond Pulsars*, *Astrophys. J. Lett.* **951** (2023) L9 [[2306.16217](#)].
- [3] A. Zic et al., *The Parkes Pulsar Timing Array Third Data Release*, [2306.16230](#).
- [4] D.J. Reardon et al., *Search for an Isotropic Gravitational-wave Background with the Parkes Pulsar Timing Array*, *Astrophys. J. Lett.* **951** (2023) L6 [[2306.16215](#)].
- [5] EPTA collaboration, *The second data release from the European Pulsar Timing Array - I. The dataset and timing analysis*, *Astron. Astrophys.* **678** (2023) A48 [[2306.16224](#)].
- [6] EPTA, INPTA: collaboration, *The second data release from the European Pulsar Timing Array - III. Search for gravitational wave signals*, *Astron. Astrophys.* **678** (2023) A50 [[2306.16214](#)].

- [7] H. Xu et al., *Searching for the Nano-Hertz Stochastic Gravitational Wave Background with the Chinese Pulsar Timing Array Data Release I*, *Res. Astron. Astrophys.* **23** (2023) 075024 [2306.16216].
- [8] NANOGrav collaboration, *The NANOGrav 15 yr Data Set: Search for Signals from New Physics*, *Astrophys. J. Lett.* **951** (2023) L11 [2306.16219].
- [9] EPTA collaboration, *The second data release from the European Pulsar Timing Array: V. Implications for massive black holes, dark matter and the early Universe*, 2306.16227.
- [10] Z. Yi, Q. Gao, Y. Gong, Y. Wang and F. Zhang, *Scalar induced gravitational waves in light of Pulsar Timing Array data*, *Sci. China Phys. Mech. Astron.* **66** (2023) 120404 [2307.02467].
- [11] G. Franciolini, A. Iovino, Junior., V. Vaskonen and H. Veermae, *Recent Gravitational Wave Observation by Pulsar Timing Arrays and Primordial Black Holes: The Importance of Non-Gaussianities*, *Phys. Rev. Lett.* **131** (2023) 201401 [2306.17149].
- [12] L. Liu, Z.-C. Chen and Q.-G. Huang, *Implications for the non-Gaussianity of curvature perturbation from pulsar timing arrays*, 2307.01102.
- [13] S. Vagnozzi, *Inflationary interpretation of the stochastic gravitational wave background signal detected by pulsar timing array experiments*, *JHEAp* **39** (2023) 81 [2306.16912].
- [14] Y.-F. Cai, X.-C. He, X.-H. Ma, S.-F. Yan and G.-W. Yuan, *Limits on scalar-induced gravitational waves from the stochastic background by pulsar timing array observations*, 2306.17822.
- [15] Y.-C. Bi, Y.-M. Wu, Z.-C. Chen and Q.-G. Huang, *Implications for the supermassive black hole binaries from the NANOGrav 15-year data set*, *Sci. China Phys. Mech. Astron.* **66** (2023) 120402 [2307.00722].
- [16] Y.-M. Wu, Z.-C. Chen and Q.-G. Huang, *Cosmological Interpretation for the Stochastic Signal in Pulsar Timing Arrays*, 2307.03141.
- [17] G. Franciolini, D. Racco and F. Rompineve, *Footprints of the QCD Crossover on Cosmological Gravitational Waves at Pulsar Timing Arrays*, 2306.17136.
- [18] Z.-Q. You, Z. Yi and Y. Wu, *Constraints on primordial curvature power spectrum with pulsar timing arrays*, *JCAP* **11** (2023) 065 [2307.04419].
- [19] J.-H. Jin, Z.-C. Chen, Z. Yi, Z.-Q. You, L. Liu and Y. Wu, *Confronting sound speed resonance with pulsar timing arrays*, *JCAP* **09** (2023) 016 [2307.08687].
- [20] L. Liu, Z.-C. Chen and Q.-G. Huang, *Probing the equation of state of the early Universe with pulsar timing arrays*, *JCAP* **11** (2023) 071 [2307.14911].
- [21] H. An, B. Su, H. Tai, L.-T. Wang and C. Yang, *Phase transition during inflation and the gravitational wave signal at pulsar timing arrays*, 2308.00070.
- [22] Z. Zhang, C. Cai, Y.-H. Su, S. Wang, Z.-H. Yu and H.-H. Zhang, *Nano-Hertz gravitational waves from collapsing domain walls associated with freeze-in dark matter in light of pulsar timing array observations*, *Phys. Rev. D* **108** (2023) 095037 [2307.11495].

- [23] B. Das, N. Jaman and M. Sami, *Gravitational wave background from quintessential inflation and NANOGrav data*, *Phys. Rev. D* **108** (2023) 103510 [[2307.12913](#)].
- [24] S. Balaji, G. Domènech and G. Franciolini, *Scalar-induced gravitational wave interpretation of PTA data: the role of scalar fluctuation propagation speed*, *JCAP* **10** (2023) 041 [[2307.08552](#)].
- [25] X.K. Du, M.X. Huang, F. Wang and Y.K. Zhang, *Did the nHZ Gravitational Waves Signatures Observed By NANOGrav Indicate Multiple Sector SUSY Breaking?*, [2307.02938](#).
- [26] V.K. Oikonomou, *Flat energy spectrum of primordial gravitational waves versus peaks and the NANOGrav 2023 observation*, *Phys. Rev. D* **108** (2023) 043516 [[2306.17351](#)].
- [27] Z. Yi, Z.-Q. You, Y. Wu, Z.-C. Chen and L. Liu, *Exploring the NANOGrav Signal and Planet-mass Primordial Black Holes through Higgs Inflation*, [2308.14688](#).
- [28] Z.-C. Chen, Q.-G. Huang, C. Liu, L. Liu, X.-J. Liu, Y. Wu et al., *Prospects for Taiji to detect a gravitational-wave background from cosmic strings*, [2310.00411](#).
- [29] L. Liu, Y. Wu and Z.-C. Chen, *Simultaneously probing the sound speed and equation of state of the early Universe with pulsar timing arrays*, [2310.16500](#).
- [30] X.-J. Zhu, W. Cui and E. Thrane, *The minimum and maximum gravitational-wave background from supermassive binary black holes*, *Mon. Not. Roy. Astron. Soc.* **482** (2019) 2588 [[1806.02346](#)].
- [31] J. Li, Z.-C. Chen and Q.-G. Huang, *Measuring the tilt of primordial gravitational-wave power spectrum from observations*, *Sci. China Phys. Mech. Astron.* **62** (2019) 110421 [[1907.09794](#)].
- [32] Z.-C. Chen, C. Yuan and Q.-G. Huang, *Non-tensorial gravitational wave background in NANOGrav 12.5-year data set*, *Sci. China Phys. Mech. Astron.* **64** (2021) 120412 [[2101.06869](#)].
- [33] Y.-M. Wu, Z.-C. Chen and Q.-G. Huang, *Constraining the Polarization of Gravitational Waves with the Parkes Pulsar Timing Array Second Data Release*, *Astrophys. J.* **925** (2022) 37 [[2108.10518](#)].
- [34] Z.-C. Chen, Y.-M. Wu and Q.-G. Huang, *Searching for isotropic stochastic gravitational-wave background in the international pulsar timing array second data release*, *Commun. Theor. Phys.* **74** (2022) 105402 [[2109.00296](#)].
- [35] Z.-C. Chen, Y.-M. Wu and Q.-G. Huang, *Search for the Gravitational-wave Background from Cosmic Strings with the Parkes Pulsar Timing Array Second Data Release*, *Astrophys. J.* **936** (2022) 20 [[2205.07194](#)].
- [36] PPTA collaboration, *Constraining ultralight vector dark matter with the Parkes Pulsar Timing Array second data release*, *Phys. Rev. D* **106** (2022) L081101 [[2210.03880](#)].
- [37] IPTA collaboration, *Searching for continuous Gravitational Waves in the second data release of the International Pulsar Timing Array*, *Mon. Not. Roy. Astron. Soc.* **521** (2023) 5077 [[2303.10767](#)].
- [38] Y.-M. Wu, Z.-C. Chen and Q.-G. Huang, *Search for stochastic gravitational-wave*

- background from massive gravity in the NANOGrav 12.5-year dataset, *Phys. Rev. D* **107** (2023) 042003 [[2302.00229](#)].
- [39] Y.-M. Wu, Z.-C. Chen and Q.-G. Huang, *Pulsar timing residual induced by ultralight tensor dark matter*, *JCAP* **09** (2023) 021 [[2305.08091](#)].
- [40] INTERNATIONAL PULSAR TIMING ARRAY collaboration, *Comparing recent PTA results on the nanohertz stochastic gravitational wave background*, [2309.00693](#).
- [41] Y.-M. Wu, Z.-C. Chen, Y.-C. Bi and Q.-G. Huang, *Constraining the Graviton Mass with the NANOGrav 15-Year Data Set*, [2310.07469](#).
- [42] Y.-C. Bi, Y.-M. Wu, Z.-C. Chen and Q.-G. Huang, *Constraints on the velocity of gravitational waves from NANOGrav 15-year data set*, [2310.08366](#).
- [43] Z.-C. Chen, Y.-M. Wu, Y.-C. Bi and Q.-G. Huang, *Search for Non-Tensorial Gravitational-Wave Backgrounds in the NANOGrav 15-Year Data Set*, [2310.11238](#).
- [44] S. Hawking, *Gravitationally collapsed objects of very low mass*, *Mon. Not. Roy. Astron. Soc.* **152** (1971) 75.
- [45] B.J. Carr and S.W. Hawking, *Black holes in the early Universe*, *Mon. Not. Roy. Astron. Soc.* **168** (1974) 399.
- [46] K.N. Ananda, C. Clarkson and D. Wands, *The Cosmological gravitational wave background from primordial density perturbations*, *Phys. Rev. D* **75** (2007) 123518 [[gr-qc/0612013](#)].
- [47] D. Baumann, P.J. Steinhardt, K. Takahashi and K. Ichiki, *Gravitational Wave Spectrum Induced by Primordial Scalar Perturbations*, *Phys. Rev. D* **76** (2007) 084019 [[hep-th/0703290](#)].
- [48] R. Saito and J. Yokoyama, *Gravitational wave background as a probe of the primordial black hole abundance*, *Phys. Rev. Lett.* **102** (2009) 161101 [[0812.4339](#)].
- [49] L. Alabidi, K. Kohri, M. Sasaki and Y. Sendouda, *Observable Spectra of Induced Gravitational Waves from Inflation*, *JCAP* **09** (2012) 017 [[1203.4663](#)].
- [50] M. Sasaki, T. Suyama, T. Tanaka and S. Yokoyama, *Primordial black holes—perspectives in gravitational wave astronomy*, *Class. Quant. Grav.* **35** (2018) 063001 [[1801.05235](#)].
- [51] T. Nakama, J. Silk and M. Kamionkowski, *Stochastic gravitational waves associated with the formation of primordial black holes*, *Phys. Rev. D* **95** (2017) 043511 [[1612.06264](#)].
- [52] K. Kohri and T. Terada, *Semianalytic calculation of gravitational wave spectrum nonlinearly induced from primordial curvature perturbations*, *Phys. Rev. D* **97** (2018) 123532 [[1804.08577](#)].
- [53] H. Di and Y. Gong, *Primordial black holes and second order gravitational waves from ultra-slow-roll inflation*, *JCAP* **07** (2018) 007 [[1707.09578](#)].
- [54] S.-L. Cheng, W. Lee and K.-W. Ng, *Primordial black holes and associated gravitational waves in axion monodromy inflation*, *JCAP* **07** (2018) 001 [[1801.09050](#)].

- [55] R.-G. Cai, S. Pi, S.-J. Wang and X.-Y. Yang, *Resonant multiple peaks in the induced gravitational waves*, *JCAP* **05** (2019) 013 [[1901.10152](#)].
- [56] R.-g. Cai, S. Pi and M. Sasaki, *Gravitational Waves Induced by non-Gaussian Scalar Perturbations*, *Phys. Rev. Lett.* **122** (2019) 201101 [[1810.11000](#)].
- [57] R.-G. Cai, S. Pi, S.-J. Wang and X.-Y. Yang, *Pulsar Timing Array Constraints on the Induced Gravitational Waves*, *JCAP* **10** (2019) 059 [[1907.06372](#)].
- [58] R.-G. Cai, Z.-K. Guo, J. Liu, L. Liu and X.-Y. Yang, *Primordial black holes and gravitational waves from parametric amplification of curvature perturbations*, *JCAP* **06** (2020) 013 [[1912.10437](#)].
- [59] Y. Wu, *Merger history of primordial black-hole binaries*, *Phys. Rev. D* **101** (2020) 083008 [[2001.03833](#)].
- [60] R.-G. Cai, Y.-C. Ding, X.-Y. Yang and Y.-F. Zhou, *Constraints on a mixed model of dark matter particles and primordial black holes from the galactic 511 keV line*, *JCAP* **03** (2021) 057 [[2007.11804](#)].
- [61] S. Pi and M. Sasaki, *Gravitational Waves Induced by Scalar Perturbations with a Lognormal Peak*, *JCAP* **09** (2020) 037 [[2005.12306](#)].
- [62] G. Domènech, S. Pi and M. Sasaki, *Induced gravitational waves as a probe of thermal history of the universe*, *JCAP* **08** (2020) 017 [[2005.12314](#)].
- [63] L. Liu, Z.-K. Guo and R.-G. Cai, *Effects of the surrounding primordial black holes on the merger rate of primordial black hole binaries*, *Phys. Rev. D* **99** (2019) 063523 [[1812.05376](#)].
- [64] L. Liu, Z.-K. Guo and R.-G. Cai, *Effects of the merger history on the merger rate density of primordial black hole binaries*, *Eur. Phys. J. C* **79** (2019) 717 [[1901.07672](#)].
- [65] L. Liu, Z.-K. Guo, R.-G. Cai and S.P. Kim, *Merger rate distribution of primordial black hole binaries with electric charges*, *Phys. Rev. D* **102** (2020) 043508 [[2001.02984](#)].
- [66] L. Liu, X.-Y. Yang, Z.-K. Guo and R.-G. Cai, *Testing primordial black hole and measuring the Hubble constant with multiband gravitational-wave observations*, *JCAP* **01** (2023) 006 [[2112.05473](#)].
- [67] L. Liu, Z.-Q. You, Y. Wu and Z.-C. Chen, *Constraining the merger history of primordial-black-hole binaries from GWTC-3*, *Phys. Rev. D* **107** (2023) 063035 [[2210.16094](#)].
- [68] Z.-C. Chen, C. Yuan and Q.-G. Huang, *Pulsar Timing Array Constraints on Primordial Black Holes with NANOGrav 11-Year Dataset*, *Phys. Rev. Lett.* **124** (2020) 251101 [[1910.12239](#)].
- [69] C. Yuan, Z.-C. Chen and Q.-G. Huang, *Scalar induced gravitational waves in different gauges*, *Phys. Rev. D* **101** (2020) 063018 [[1912.00885](#)].
- [70] C. Yuan, Z.-C. Chen and Q.-G. Huang, *Log-dependent slope of scalar induced gravitational waves in the infrared regions*, *Phys. Rev. D* **101** (2020) 043019 [[1910.09099](#)].

- [71] C. Yuan, Z.-C. Chen and Q.-G. Huang, *Probing primordial–black-hole dark matter with scalar induced gravitational waves*, *Phys. Rev. D* **100** (2019) 081301 [[1906.11549](#)].
- [72] B. Carr, K. Kohri, Y. Sendouda and J. Yokoyama, *Constraints on primordial black holes*, *Rept. Prog. Phys.* **84** (2021) 116902 [[2002.12778](#)].
- [73] L. Liu, O. Christiansen, Z.-K. Guo, R.-G. Cai and S.P. Kim, *Gravitational and electromagnetic radiation from binary black holes with electric and magnetic charges: Circular orbits on a cone*, *Phys. Rev. D* **102** (2020) 103520 [[2008.02326](#)].
- [74] L. Liu, O. Christiansen, W.-H. Ruan, Z.-K. Guo, R.-G. Cai and S.P. Kim, *Gravitational and electromagnetic radiation from binary black holes with electric and magnetic charges: elliptical orbits on a cone*, *Eur. Phys. J. C* **81** (2021) 1048 [[2011.13586](#)].
- [75] T. Papanikolaou, C. Tzerefos, S. Basilakos and E.N. Saridakis, *Scalar induced gravitational waves from primordial black hole Poisson fluctuations in $f(R)$ gravity*, *JCAP* **10** (2022) 013 [[2112.15059](#)].
- [76] T. Papanikolaou, C. Tzerefos, S. Basilakos and E.N. Saridakis, *No constraints for $f(T)$ gravity from gravitational waves induced from primordial black hole fluctuations*, *Eur. Phys. J. C* **83** (2023) 31 [[2205.06094](#)].
- [77] A. Chakraborty, P.K. Chanda, K.L. Pandey and S. Das, *Formation and Abundance of Late-forming Primordial Black Holes as Dark Matter*, *Astrophys. J.* **932** (2022) 119 [[2204.09628](#)].
- [78] Z.-C. Chen and Q.-G. Huang, *Merger Rate Distribution of Primordial-Black-Hole Binaries*, *Astrophys. J.* **864** (2018) 61 [[1801.10327](#)].
- [79] Z.-C. Chen, F. Huang and Q.-G. Huang, *Stochastic Gravitational-wave Background from Binary Black Holes and Binary Neutron Stars and Implications for LISA*, *Astrophys. J.* **871** (2019) 97 [[1809.10360](#)].
- [80] Z.-C. Chen and Q.-G. Huang, *Distinguishing Primordial Black Holes from Astrophysical Black Holes by Einstein Telescope and Cosmic Explorer*, *JCAP* **08** (2020) 039 [[1904.02396](#)].
- [81] Z.-C. Chen, C. Yuan and Q.-G. Huang, *Confronting the primordial black hole scenario with the gravitational-wave events detected by LIGO-Virgo*, *Phys. Lett. B* **829** (2022) 137040 [[2108.11740](#)].
- [82] Z.-C. Chen, S.-S. Du, Q.-G. Huang and Z.-Q. You, *Constraints on primordial-black-hole population and cosmic expansion history from GWTC-3*, *JCAP* **03** (2023) 024 [[2205.11278](#)].
- [83] L. Liu and S.P. Kim, *Merger rate of charged black holes from the two-body dynamical capture*, *JCAP* **03** (2022) 059 [[2201.02581](#)].
- [84] L.-M. Zheng, Z. Li, Z.-C. Chen, H. Zhou and Z.-H. Zhu, *Towards a reliable reconstruction of the power spectrum of primordial curvature perturbation on small scales from GWTC-3*, *Phys. Lett. B* **838** (2023) 137720 [[2212.05516](#)].
- [85] Z.-C. Chen, S.P. Kim and L. Liu, *Gravitational and electromagnetic radiation from*

- binary black holes with electric and magnetic charges: hyperbolic orbits on a cone*, *Commun. Theor. Phys.* **75** (2023) 065401 [[2210.15564](#)].
- [86] D.-S. Meng, C. Yuan and Q.-G. Huang, *Primordial black holes generated by the non-minimal spectator field*, *Sci. China Phys. Mech. Astron.* **66** (2023) 280411 [[2212.03577](#)].
- [87] S. Garcia-Saenz, Y. Lu and Z. Shuai, *Scalar-Induced Gravitational Waves from Ghost Inflation and Parity Violation*, [2306.09052](#).
- [88] PLANCK collaboration, *Planck 2018 results. X. Constraints on inflation*, *Astron. Astrophys.* **641** (2020) A10 [[1807.06211](#)].
- [89] J. Martin, H. Motohashi and T. Suyama, *Ultra Slow-Roll Inflation and the non-Gaussianity Consistency Relation*, *Phys. Rev. D* **87** (2013) 023514 [[1211.0083](#)].
- [90] H. Motohashi, A.A. Starobinsky and J. Yokoyama, *Inflation with a constant rate of roll*, *JCAP* **09** (2015) 018 [[1411.5021](#)].
- [91] Z. Yi and Y. Gong, *On the constant-roll inflation*, *JCAP* **03** (2018) 052 [[1712.07478](#)].
- [92] J. Garcia-Bellido and E. Ruiz Morales, *Primordial black holes from single field models of inflation*, *Phys. Dark Univ.* **18** (2017) 47 [[1702.03901](#)].
- [93] C. Germani and T. Prokopec, *On primordial black holes from an inflection point*, *Phys. Dark Univ.* **18** (2017) 6 [[1706.04226](#)].
- [94] H. Motohashi and W. Hu, *Primordial Black Holes and Slow-Roll Violation*, *Phys. Rev. D* **96** (2017) 063503 [[1706.06784](#)].
- [95] J.M. Ezquiaga, J. Garcia-Bellido and E. Ruiz Morales, *Primordial Black Hole production in Critical Higgs Inflation*, *Phys. Lett. B* **776** (2018) 345 [[1705.04861](#)].
- [96] H. Di and Y. Gong, *Primordial black holes and second order gravitational waves from ultra-slow-roll inflation*, *JCAP* **07** (2018) 007 [[1707.09578](#)].
- [97] G. Ballesteros, J. Beltran Jimenez and M. Pieroni, *Black hole formation from a general quadratic action for inflationary primordial fluctuations*, *JCAP* **06** (2019) 016 [[1811.03065](#)].
- [98] I. Dalianis, A. Kehagias and G. Tringas, *Primordial black holes from α -attractors*, *JCAP* **01** (2019) 037 [[1805.09483](#)].
- [99] F. Bezrukov, M. Pauly and J. Rubio, *On the robustness of the primordial power spectrum in renormalized Higgs inflation*, *JCAP* **02** (2018) 040 [[1706.05007](#)].
- [100] K. Kannike, L. Marzola, M. Raidal and H. Veermäe, *Single Field Double Inflation and Primordial Black Holes*, *JCAP* **09** (2017) 020 [[1705.06225](#)].
- [101] J. Lin, Q. Gao, Y. Gong, Y. Lu, C. Zhang and F. Zhang, *Primordial black holes and secondary gravitational waves from k and G inflation*, *Phys. Rev. D* **101** (2020) 103515 [[2001.05909](#)].
- [102] J. Lin, S. Gao, Y. Gong, Y. Lu, Z. Wang and F. Zhang, *Primordial black holes and scalar induced gravitational waves from Higgs inflation with noncanonical kinetic term*, *Phys. Rev. D* **107** (2023) 043517 [[2111.01362](#)].
- [103] Q. Gao, Y. Gong and Z. Yi, *Primordial black holes and secondary gravitational waves from natural inflation*, *Nucl. Phys. B* **969** (2021) 115480 [[2012.03856](#)].

- [104] Q. Gao, *Primordial black holes and secondary gravitational waves from chaotic inflation*, *Sci. China Phys. Mech. Astron.* **64** (2021) 280411 [[2102.07369](#)].
- [105] Z. Yi, Y. Gong, B. Wang and Z.-h. Zhu, *Primordial black holes and secondary gravitational waves from the Higgs field*, *Phys. Rev. D* **103** (2021) 063535 [[2007.09957](#)].
- [106] Z. Yi, Q. Gao, Y. Gong and Z.-h. Zhu, *Primordial black holes and scalar-induced secondary gravitational waves from inflationary models with a noncanonical kinetic term*, *Phys. Rev. D* **103** (2021) 063534 [[2011.10606](#)].
- [107] Z. Yi and Z.-H. Zhu, *NANOGrav signal and LIGO-Virgo primordial black holes from the Higgs field*, *JCAP* **05** (2022) 046 [[2105.01943](#)].
- [108] Z. Yi, *Primordial black holes and scalar-induced gravitational waves from the generalized Brans-Dicke theory*, *JCAP* **03** (2023) 048 [[2206.01039](#)].
- [109] F. Zhang, Y. Gong, J. Lin, Y. Lu and Z. Yi, *Primordial non-Gaussianity from G-inflation*, *JCAP* **04** (2021) 045 [[2012.06960](#)].
- [110] S. Pi, Y.-l. Zhang, Q.-G. Huang and M. Sasaki, *Scalaron from R^2 -gravity as a heavy field*, *JCAP* **05** (2018) 042 [[1712.09896](#)].
- [111] A.Y. Kamenshchik, A. Tronconi, T. Vardanyan and G. Venturi, *Non-Canonical Inflation and Primordial Black Holes Production*, *Phys. Lett. B* **791** (2019) 201 [[1812.02547](#)].
- [112] C. Fu, P. Wu and H. Yu, *Primordial Black Holes from Inflation with Nonminimal Derivative Coupling*, *Phys. Rev. D* **100** (2019) 063532 [[1907.05042](#)].
- [113] C. Fu, P. Wu and H. Yu, *Scalar induced gravitational waves in inflation with gravitationally enhanced friction*, *Phys. Rev. D* **101** (2020) 023529 [[1912.05927](#)].
- [114] I. Dalianis, S. Karydas and E. Papantonopoulos, *Generalized Non-Minimal Derivative Coupling: Application to Inflation and Primordial Black Hole Production*, *JCAP* **06** (2020) 040 [[1910.00622](#)].
- [115] A. Gundhi and C.F. Steinwachs, *Scalaron–Higgs inflation reloaded: Higgs-dependent scalaron mass and primordial black hole dark matter*, *Eur. Phys. J. C* **81** (2021) 460 [[2011.09485](#)].
- [116] D.Y. Cheong, S.M. Lee and S.C. Park, *Primordial black holes in Higgs- R^2 inflation as the whole of dark matter*, *JCAP* **01** (2021) 032 [[1912.12032](#)].
- [117] F. Zhang, *Primordial black holes and scalar induced gravitational waves from the E model with a Gauss-Bonnet term*, *Phys. Rev. D* **105** (2022) 063539 [[2112.10516](#)].
- [118] F. Zhang, J. Lin and Y. Lu, *Double-peaked inflation model: Scalar induced gravitational waves and primordial-black-hole suppression from primordial non-Gaussianity*, *Phys. Rev. D* **104** (2021) 063515 [[2106.10792](#)].
- [119] S. Kawai and J. Kim, *Primordial black holes from Gauss-Bonnet-corrected single field inflation*, *Phys. Rev. D* **104** (2021) 083545 [[2108.01340](#)].
- [120] R.-G. Cai, C. Chen and C. Fu, *Primordial black holes and stochastic gravitational wave background from inflation with a noncanonical spectator field*, *Phys. Rev. D* **104** (2021) 083537 [[2108.03422](#)].

- [121] P. Chen, S. Koh and G. Tumurtushaa, *Primordial black holes and induced gravitational waves from inflation in the Horndeski theory of gravity*, [2107.08638](#).
- [122] R. Zheng, J. Shi and T. Qiu, *On primordial black holes and secondary gravitational waves generated from inflation with solo/multi-bumpy potential **, *Chin. Phys. C* **46** (2022) 045103 [[2106.04303](#)].
- [123] A. Karam, N. Koivunen, E. Tomberg, V. Vaskonen and H. Veermäe, *Anatomy of single-field inflationary models for primordial black holes*, *JCAP* **03** (2023) 013 [[2205.13540](#)].
- [124] A. Ashoorioon, A. Rostami and J.T. Firouzjaee, *EFT compatible PBHs: effective spawning of the seeds for primordial black holes during inflation*, *JHEP* **07** (2021) 087 [[1912.13326](#)].
- [125] K. Danzmann, *LISA: An ESA cornerstone mission for a gravitational wave observatory*, *Class. Quant. Grav.* **14** (1997) 1399.
- [126] LISA collaboration, *Laser Interferometer Space Antenna*, [1702.00786](#).
- [127] W.-R. Hu and Y.-L. Wu, *The Taiji Program in Space for gravitational wave physics and the nature of gravity*, *Natl. Sci. Rev.* **4** (2017) 685.
- [128] TIANQIN collaboration, *TianQin: a space-borne gravitational wave detector*, *Class. Quant. Grav.* **33** (2016) 035010 [[1512.02076](#)].
- [129] Y. Gong, J. Luo and B. Wang, *Concepts and status of Chinese space gravitational wave detection projects*, *Nature Astron.* **5** (2021) 881 [[2109.07442](#)].
- [130] S. Kawamura et al., *The Japanese space gravitational wave antenna: DECIGO*, *Class. Quant. Grav.* **28** (2011) 094011.
- [131] K. Inomata and T. Nakama, *Gravitational waves induced by scalar perturbations as probes of the small-scale primordial spectrum*, *Phys. Rev. D* **99** (2019) 043511 [[1812.00674](#)].
- [132] Z. Yi and Q. Fei, *Constraints on primordial curvature spectrum from primordial black holes and scalar-induced gravitational waves*, *Eur. Phys. J. C* **83** (2023) 82 [[2210.03641](#)].
- [133] Y. Lu, Y. Gong, Z. Yi and F. Zhang, *Constraints on primordial curvature perturbations from primordial black hole dark matter and secondary gravitational waves*, *JCAP* **12** (2019) 031 [[1907.11896](#)].
- [134] K. Inomata, M. Kawasaki, K. Mukaida, Y. Tada and T.T. Yanagida, *Inflationary primordial black holes for the LIGO gravitational wave events and pulsar timing array experiments*, *Phys. Rev. D* **95** (2017) 123510 [[1611.06130](#)].
- [135] J.R. Espinosa, D. Racco and A. Riotto, *A Cosmological Signature of the SM Higgs Instability: Gravitational Waves*, *JCAP* **09** (2018) 012 [[1804.07732](#)].
- [136] G. Ashton et al., *BILBY: A user-friendly Bayesian inference library for gravitational-wave astronomy*, *Astrophys. J. Suppl.* **241** (2019) 27 [[1811.02042](#)].
- [137] J. Skilling, *Nested Sampling*, *AIP Conf. Proc.* **735** (2004) 395.
- [138] C.J. Moore and A. Vecchio, *Ultra-low-frequency gravitational waves from cosmological and astrophysical processes*, *Nature Astron.* **5** (2021) 1268 [[2104.15130](#)].

- [139] K. Inomata, M. Kawasaki and Y. Tada, *Revisiting constraints on small scale perturbations from big-bang nucleosynthesis*, *Phys. Rev. D* **94** (2016) 043527 [[1605.04646](#)].
- [140] D.J. Fixsen, E.S. Cheng, J.M. Gales, J.C. Mather, R.A. Shafer and E.L. Wright, *The Cosmic Microwave Background spectrum from the full COBE FIRAS data set*, *Astrophys. J.* **473** (1996) 576 [[astro-ph/9605054](#)].