SEQUENCES WITH IDENTICAL AUTOCORRELATION SPECTRA

DANIEL J. KATZ, ADEEBUR RAHMAN, AND MICHAEL J WARD

ABSTRACT. Aperiodic autocorrelation measures the similarity between a finite-length sequence of complex numbers and translates of itself. Autocorrelation is important in communications, remote sensing, and scientific instrumentation. The autocorrelation function reports the aperiodic autocorrelation at every possible translation. Knowing the autocorrelation function of a sequence is equivalent to knowing the magnitude of its Fourier transform. Resolving the lack of phase information is called the phase problem. We say that two sequences are isospectral to mean that they have the same aperiodic autocorrelation function. Sequences used in technological applications often have restrictions on their terms: they are not arbitrary complex numbers, but come from an alphabet that may reside in a proper subring of the complex field or may come from a finite set of values. For example, binary sequences involve terms equal to only +1 and -1. In this paper, we investigate the necessary and sufficient conditions for two sequences to be isospectral, where we take their alphabet into consideration. There are trivial forms of isospectrality arising from modifications that predictably preserve the autocorrelation, for example, negating sequences or both conjugating their terms and writing them in reverse order. By an exhaustive search of binary sequences up to length 34, we find that nontrivial isospectrality among binary sequences does occur, but is rare. We say that a positive integer n is barren to mean that there are no nontrivially isospectral binary sequences of length n. For integers $n \leq 34$, we found that the barren ones are 1-8, 10, 11, 13, 14, 19, 22, 23, 26, and 29. We prove that any multiple of a non-barren number is also not barren, and pose an open question as to whether there are finitely or infinitely many barren numbers.

1. INTRODUCTION

In many physical measurements of wave phenomena, detectors are unable to discern phases. This loss of phase information is called the phase problem, a terminology that arose in x-ray crystallography, where diffraction

Date: 14 August 2023.

This paper is based upon work supported in part by the National Science Foundation under Grants CCF-1815487 and CCF-2206454. Adeebur Rahman was supported by the Ramanujan Research Scholarship from the Department of Mathematics, California State University, Northridge. Michael J Ward was supported by the Efrem Ostrow Scholarship from the Department of Mathematics, California State University, Northridge.

patterns give the magnitude of the Fourier transform of the electron density without the phase information [BE22]. Knowing the magnitude of the Fourier transform is the same as knowing the autocorrelation of the electron density, which in general is not sufficient information to recover the electron density itself. This paper concerns itself with the aperiodic one-dimensional discrete problem of phases, that is, the extent to which one can deduce a sequence from its aperiodic autocorrelation. See [BE22] on the the practical importance of the aperiodic problem of phases.

Because we consider aperiodic autocorrelation, a sequence is any doubly infinite sequence $f = (\ldots, f_{-1}, f_0, f_1, f_2, \ldots)$ of complex numbers such that only finitely many of the terms are nonzero. We identify this sequence with $f(z) = \sum_{j \in \mathbb{Z}} f_j z^j \in \mathbb{C}[z, z^{-1}]$, where $\mathbb{C}[z, z^{-1}]$ is the ring of Laurent polynomials with complex coefficients. The support of f, written $\operatorname{supp}(f)$, is the set $\{j \in \mathbb{Z} : f_j \neq 0\}$. A segment is a set of consecutive integers. The length of a sequence f, written $\operatorname{len}(f)$, is the cardinality of the smallest segment that contains $\operatorname{supp}(f)$. A contiguous sequence f is a sequence where $\operatorname{supp}(f)$ is a segment. A unimodular sequence is a contiguous sequence fwhere $|f_j| = 1$ for every $j \in \operatorname{supp}(f)$. For any positive integer m, an m-ary sequence is a unimodular sequence where f_j is an mth root of unity for every $j \in \operatorname{supp}(f)$; when m = 2, we have a binary sequence, where $f_j \in \{1, -1\}$ for every $j \in \operatorname{supp}(f)$.

For a sequence $f = (\dots, f_{-1}, f_0, f_1, f_2, \dots)$ and an integer s, the aperiodic autocorrelation of f at shift s is

$$C_f(s) = \sum_{j \in \mathbb{Z}} f_{j+s} \overline{f_j}.$$

Note that the finite support of f guarantees that $C_f(s) \neq 0$ for only finitely many $s \in \mathbb{Z}$. The Laurent polynomial interpretation of sequences provides a convenient formalism for calculating autocorrelation. To use this formalism, for $f(z) = \sum_{j \in \mathbb{Z}} f_j z^j \in \mathbb{C}[z, z^{-1}]$ we define the *conjugate of* f(z), written $\overline{f(z)}$, to be $\sum_{j \in \mathbb{Z}} \overline{f_j} z^{-j}$. In this case one readily shows that

$$f(z)\overline{f(z)} = \sum_{s \in \mathbb{Z}} C_f(s) z^s.$$

We call this last Laurent polynomial the *autocorrelation spectrum of* f because it organizes every autocorrelation value $C_f(s)$ as the coefficient of z^s , so that one can read off the autocorrelation value at any shift.

We are interested in the extent to which the autocorrelation spectrum $\sum_{s\in\mathbb{Z}} C_f(s)z^s$ determines the sequence f from which it is derived. We say that two sequences f and g are *isospectral* to mean that they have identical autocorrelation spectra, that is, $C_f(s) = C_g(s)$ for every $s \in \mathbb{Z}$, or equivalently, $\sum_{s\in\mathbb{Z}} C_f(s)z^s = \sum_{s\in\mathbb{Z}} C_g(s)z^s$, or equivalently $f(z)\overline{f(z)} = g(z)\overline{g(z)}$. Isospectrality is an equivalence relation. Note that since $\mathbb{C}[z, z^{-1}]$ is an integral domain, no nonzero sequence is isospectral to the zero sequence.

Note that the conjugation map $f(z) \mapsto \overline{f(z)}$ is an automorphism of $\mathbb{C}[z, z^{-1}]$. A subring R of \mathbb{C} that is closed under complex conjugation is said to be *conjugate-closed*. If R is a such a subring, then conjugation restricts to an automorphism of $R[z, z^{-1}]$. Therefore, if R is a conjugate-closed subring of \mathbb{C} , then our conjugation map $f(z) \mapsto \overline{f(z)}$ maps 0, units, irreducibles, and reducible elements of $R[z, z^{-1}]$ respectively to 0, units, irreducibles, and reducible elements of $R[z, z^{-1}]$.

If R is a subring of \mathbb{C} , then R^* denotes the multiplicative group of units of R and R^{\circledast} denotes the set of all unimodular units in R, which is a subgroup of R^* . If R is conjugate-closed, then every unimodular element u of R is a unit of R because $u\overline{u} = 1$. If R is a conjugate-closed subring of \mathbb{C} , then we say that two sequences f and g are R-similar and write $f \sim_R g$ to mean that $f(z) = uz^j g(z)$ for some $u \in R^{\circledast}$ and $j \in \mathbb{Z}$. Note that R-similarity is an equivalence relation whose equivalence classes are called R-similarity classes and are notated

(1)
$$[f]_R = \{ uz^j f : u \in R^{\circledast}, j \in \mathbb{Z} \}.$$

The *R*-similarity relation in $R[z, z^{-1}]$ is a finer equivalence relation than the relation of being associate. For a conjugate-closed subring of \mathbb{C} , the *conjugate* of an *R*-similarity class is the set of all conjugates of the elements in the class; this set is itself an *R*-similarity class.

If u is a unimodular element of \mathbb{C} , then a *u*-palindrome is a sequence f(z) such that $\overline{f(z)} = uz^j f(z)$ for some $j \in \mathbb{Z}$. A palindrome is just a 1-palindrome. Note that the zero sequence is a *u*-palindrome for every unimodular u, but a nonzero sequence cannot be both a *u*-palindrome and a *v*-palindrome for distinct u and v. If R is a conjugate-closed subring of \mathbb{C} , then an *R*-palindrome is any *u*-palindrome for some $u \in R^{\circledast}$. In other words, an *R*-palindrome is a sequence that is *R*-similar to its own conjugate. Thus, an *R*-similarity class either consists entirely of *R*-palindromic sequences (in which case the class is its own conjugate, so we call it self-conjugate or palindromic) or it has no *R*-palindromic sequences (in which case our class is distinct from its conjugate). Therefore, non-self-conjugate *R*-similarity classes come in conjugate pairs.

If R is a conjugate-closed subring of \mathbb{C} , then R-similar sequences are isospectral, and we should note that a sequence f(z) is isospectral to $\overline{f(z)}$ (as well as any sequence R-similar to $\overline{f(z)}$). If R is a conjugate-closed subring of \mathbb{C} , then we say that two sequences in $f(z), g(z) \in R[z, z^{-1}]$ are trivially isospectral over R to mean that they are either R-similar to each other or one is R-similar to the conjugate of the other. Trivial isospectrality over R is an equivalence relation that refines isospectrality and is refined by R-similarity. The trivial isospectrality class of some f(z) over R is the union of the R-similarity class of f(z) and the R-similarity class of $\overline{f(z)}$. If f(z) is an R-palindrome, these two classes are the same, so the trivial isospectrality class of f(z) is just the (self-conjugate) R-similarity class of f(z), but if f(z) is not an R-palindrome, then the trivial isospectrality class of f(z) is a disjoint union of two (non-self-conjugate) *R*-similarity classes (that of f(z) and that of $\overline{f(z)}$).

Throughout this paper $\mathbb{N} = \{0, 1, 2...\}$. The following theorem tells us which *R*-similarity classes contain sequences isospectral to a given sequence for certain subrings *R* of \mathbb{C} .

Theorem 1.1. Let R be a conjugate-closed subring of \mathbb{C} that is a unique factorization domain. Let f be a sequence whose terms lie in R. If f = 0, then the only sequence isospectral to f is itself. If $f \neq 0$, then suppose that

(2)
$$f(z) = u f_1(z)^{a_1} \cdots f_m(z)^{a_m} g_1(z)^{b_1} \cdots g_n(z)^{b_n} \overline{g_1(z)}^{c_1} \cdots \overline{g_n(z)}^{c_n}$$

is a factorization of f(z) into nonassociate irreducibles $f_1(z), \ldots, f_n(z), g_1(z), \ldots, g_n(z), \overline{g_1(z)}, \ldots, \overline{g_n(z)}$ and unit u of $R[z, z^{-1}]$ where $f_1(z), \ldots, f_m(z)$ are R-palindromic and $g_1(z), \ldots, g_n(z)$ are not, and we have $a = (a_1, \ldots, a_m) \in \mathbb{N}^m$ and $b = (b_1, \ldots, b_n), c = (c_1, \ldots, c_n) \in \mathbb{N}^n$. Then the set of all sequences in $R[z, z^{-1}]$ that are isospectral to f(z) is

(3)
$$\bigcup_{\substack{b',c'\in\mathbb{N}^n\\b'+c'=b+c}} \left[uf_1(z)^{a_1}\cdots f_m(z)^{a_m}g_1(z)^{b'_1}\cdots g_n(z)^{b'_n}\overline{g_1(z)}^{c'_1}\cdots \overline{g_n(z)}^{c'_n} \right]_R,$$

which we have written as a disjoint union of R-similarity classes. There are $\prod_{j=1}^{n} (b_j + c_j + 1)$ classes in this disjoint union. If $b_j + c_j$ is even for every $j \in \{1, 2, ..., n\}$, then precisely one of these classes is self-conjugate, namely the one with b' = c' = (b + c)/2; otherwise there are no self-conjugate classes in the disjoint union. The rest of the classes occur in conjugate pairs. Therefore, the isospectrality class of f is the disjoint union of $\left[\left(\prod_{j=1}^{n} (b_j + c_j + 1)\right)/2\right]$ trivial isospectrality classes. The sequence f is nontrivially isospectral to some other sequence in $R[z, z^{-1}]$ if and only if $\prod_{j=1}^{n} (b_j + c_j + 1) \geq 3$.

Remark 1.2. When $R = \mathbb{C}$ in Theorem 1.1, the polynomial f(z) factors into linear factors, and then one recapitulates the results described in Theorem 2.4 of [BP15], which obtains results already shown in [Fej16]. If f(z)represents a sequence of length $\ell \geq 2$, then there are $\ell - 1$ linear factors in (2) and then the number of nontrivial isospectrality classes in the isospectrality class of f(z) is at most $2^{\ell-2}$, as observed in [BP15, Cor. 2.6]. The maximum of $2^{\ell-2}$ is achieved if and only if either (i) $m = 0, n = \ell - 1$, and $\{b_j, c_j\} = \{0, 1\}$ for every $j \in \{1, \ldots, \ell - 1\}$ or (ii) $\ell = 3$ and n = 1 and $b_1 + c_1 = 2$.

Remark 1.3. Beinert and Plonka [BP15, Remark 2.7] also consider what happens when $R = \mathbb{R}$, the real field, in the situation outlined in Theorem 1.1, and point out that a real sequence f(z) of length ℓ can have $2^{\ell-2}$ nontrivial isospectrality classes in its isospectrality class only if all its roots are real (i.e., if and only if f(z) splits in $\mathbb{R}[x]$).

Theorem 1.1 limits the circumstances under which R-palindromes may be isospectral to each other.

Corollary 1.4. Let R be a conjugate-closed subring of \mathbb{C} that is a unique factorization domain. If f and g are R-palindromes that are isospectral, then they must be R-similar.

Furthermore, we prove more technical results that show that certain kinds of *R*-palindromes may not be isospectral to each other. If *R* is a conjugateclosed ring, let R^{*2} be the subgroup of squares in R^* , that is, $R^{*2} = \{r^2 : r \in R^*\}$, and let $R^{\otimes 2}$ be the subgroup of R^{*2} consisting of unimodular elements, that is, $R^{\otimes 2} = \{r^2 : r \in R^{\otimes}\}$.

Corollary 1.5. Let R be a conjugate-closed subring of \mathbb{C} that is a unique factorization domain. Let $u, v \in R^{\circledast}$ and let $f(z) \in R[z, z^{-1}]$ be a u-palindrome and $g \in R[z, z^{-1}]$ be a v-palindrome that is isospectral to f. Then either f = g = 0 or else $v/u \in R^{\circledast 2}$.

When $R = \mathbb{Z}$, the only units in R are ± 1 , so the only R-palindromes are palindromes (i.e., 1-palindromes) and *antipalindromes*, which are (-1)-palindromes, so one consequence of Corollary 1.5 is the following.

Corollary 1.6. It is not possible for a palindrome in $\mathbb{Z}[z, z^{-1}]$ to be isospectral to an antipalindrome in $\mathbb{Z}[z, z^{-1}]$ unless both the sequences are 0.

We restrict the relation of isospectrality to binary sequences: an equivalence class of this relation is called a *binary isospectrality class*. We also restrict the notion of trivial isospectrality over \mathbb{Z} to binary sequences so that f(z) and g(z) are trivially isospectral if and only if $f(z) = uz^j g(z)$ or $f(z) = uz^j g(z)$ for some $u \in \{-1, 1\}$ and $j \in \mathbb{Z}$, and an equivalence class of this relation is called a *trivial binary isospectrality class*. Trivial binary isospectrality refines binary isospectrality, so every binary isospectrality class is a disjoint union of trivial binary isospectrality classes. The *volume* of a binary isospectrality class equals the number of trivial binary isospectrality classes in this union, and a binary isospectrality class of volume greater than one is called *nontrivial*. We used a computer to find all nontrivial binary isospectrality classes for binary sequences of lengths 1 through 34. In Table 1, we indicate how many nontrivial binary isospectrality classes there are of each volume. We represent the distribution of volumes of nontrivial isospectrality classes in a compact notation $n_1[v_1] + n_2[v_2] + \cdots + n_t[v_t]$, which means that there are n_i classes of volume v_i for each $i \in \{1, 2, \ldots, t\}$. If an entry for a particular sequence length is blank, it means that there are no nontrivial isospectrality classes for binary sequences of that length. We did not discover any nontrivial binary isospectrality class C of odd volume: by Theorem 1.1 and Corollary 1.6 such a C would contain a one trivial binary isospectrality class that would consist either entirely of palindromes or entirely of antipal indromes, and all other trivial isospectrality classes in Cwould occur in conjugate pairs.

sequence	frequency [volume]	sequence	frequency [volume]
length	of nontrivial classes	length	of nontrivial classes
1		18	42 [2]
2		19	
3		20	44 [2]
4		21	67 [2]
5		22	
6		23	
7		24	422 [2]
8		25	36[2]
9	$1 \ [2]$	26	
10		27	348 [2] + 1 [4]
11		28	$180 \ [2]$
12	8 [2]	29	
13		30	1214 [2]
14		31	26 [2]
15	14 [2]	32	1136 [2]
16	12 [2]	33	1105 [2]
17	$1 \ [2]$	34	30 [2]

TABLE 1. Nontrivial binary isospectrality classes

A positive integer n is said to be *barren* if there are no nontrivially isospectral binary sequences of length n. Table 1 shows that the numbers from 1 to 8, along with 10, 11, 13, 14, 19, 22, 23, 26, and 29 are barren. The following result, which is proved using a concatenation construction, explains why many numbers are not barren.

Proposition 1.7. Let m, n be positive integers such that m|n. If m is not barren, then n is not barren.

Perusal of Table 1 shows that barren numbers seem to become more sparse as the length increases. This leads to the following open question.

Open Problem 1.8. Are there finitely or infinitely many barren numbers?

The phenomenon of isospectrality is significant for the study of Golay complementary pairs. A Golay complementary pair (or just Golay pair or complementary pair) is a pair (f,g) of sequences such that for every nonzero shift s, the sum of the aperiodic autocorrelation of f at shift s is the opposite of the aperiodic autocorrelation of g at shift s. In our Laurent polynomial interpretation, this is equivalent to saying that $f(z)\overline{f(z)} + g(z)\overline{g(z)}$ is a constant. A Golay sequence is a sequence that occurs as one of the two sequences in a Golay complementary pair. If (f,g) is a complementary pair, then (f, h) is a complementary pair for every h that is isospectral to g. Golay defined complementary pairs in [Gol51, p. 469], and found a construction [Gol61, pp. 85–86] of binary complementary pairs (i.e., complementary pairs where both sequences in the pair are binary) that yields $n!2^{n+2}$ pairs consisting of sequences of length 2^n for each positive integer n. To the authors' knowledge, these are the only Golay pairs consisting of binary sequences of length 2^n known to date. See [DJ99, Theorem 3] for an easier way of conceptualizing Golay's construction via multivariable Boolean functions, and see [KM22, Example 37] for a simpler way of indexing variables in Davis and Jedwab's construction. The $n!2^{n+2}$ known complementary pairs of length 2^n involve $n!2^n$ distinct binary sequences that the make up $n!2^{n-2}$ trivial \mathbb{Z} -isospectrality classes of size 4. If we find any binary sequence that is nontrivially isospectral to one of these $n!2^n$ known binary Golay sequences, then we will have discovered a new binary Golay pair.

The rest of this paper is organized as follows. In Section 2, we prove Theorem 1.1, our main result. In Section 3, we prove its Corollaries 1.4, 1.5, and 1.6. In Section 4, we prove Proposition 1.7.

2. Proof of Theorem 1.1

We restate Theorem 1.1 and prove it here.

Theorem 2.1. Let R be a subring of \mathbb{C} that is closed under conjugation and is a unique factorization domain. Let f be a sequence whose terms lie in R. If f = 0, then the only sequence isospectral to f is itself. If $f \neq 0$, then suppose that

$$f(z) = u f_1(z)^{a_1} \cdots f_m(z)^{a_m} g_1(z)^{b_1} \cdots g_n(z)^{b_n} \overline{g_1(z)}^{c_1} \cdots \overline{g_n(z)}^{c_n}$$

is a factorization of f(z) into nonassociate irreducibles $f_1(z), \ldots, f_n(z), g_1(z), \ldots, g_n(z), \overline{g_1(z)}, \ldots, \overline{g_n(z)}$ and unit u of $R[z, z^{-1}]$ where $f_1(z), \ldots, f_m(z)$ are R-palindromic and $g_1(z), \ldots, g_n(z)$ are not, and we have $a = (a_1, \ldots, a_m) \in \mathbb{N}^m$ and $b = (b_1, \ldots, b_n), c = (c_1, \ldots, c_n) \in \mathbb{N}^n$. Then the set of all sequences in $R[z, z^{-1}]$ that are isospectral to f(z) is

$$\bigcup_{\substack{b',c'\in\mathbb{N}^n\\b'+c'=b+c}} \left[uf_1(z)^{a_1}\cdots f_m(z)^{a_m}g_1(z)^{b'_1}\cdots g_n(z)^{b'_n}\overline{g_1(z)}^{c'_1}\cdots \overline{g_n(z)}^{c'_n} \right]_R,$$

which we have written as a disjoint union of R-similarity classes. There are $\prod_{j=1}^{n} (b_j + c_j + 1)$ classes in this disjoint union. If $b_j + c_j$ is even for every $j \in \{1, 2, ..., n\}$, then precisely one of these classes is self-conjugate, namely the one with b' = c' = (b + c)/2; otherwise there are no self-conjugate classes in the disjoint union. The rest of the classes occur in conjugate pairs. Therefore, the isospectrality class of f is the disjoint union of $\left[\left(\prod_{j=1}^{n} (b_j + c_j + 1)\right)/2\right]$ trivial isospectrality classes. The sequence f is nontrivially isospectral to some other sequence in $R[z, z^{-1}]$ if and only if $\prod_{j=1}^{n} (b_j + c_j + 1) \geq 3$.

Proof. Let h(z) be a sequence in $R[z, z^{-1}]$. Then h(z) is isospectral to f(z) if and only if $h(z)\overline{h(z)} = f(z)\overline{f(z)}$. Therefore, if h(z) is isospectral to f(z), then the unique factorization of h(z) can only contain the irreducibles in $f(z)\overline{f(z)}$. Thus, in searching for the sequences isospectral to f(z) we can confine ourselves to sequences h(z) that can be written as

$$h(z) = v f_1(z)^{a'_1} \cdots f_m(z)^{a'_m} g_1(z)^{b'_1} \cdots g_n(z)^{b'_n} \overline{g_1(z)}^{c'_1} \cdots \overline{g_n(z)}^{c'_n},$$

for some unit $v \in R[z, z^{-1}]$ and $a' = (a'_1, \ldots, a'_m) \in \mathbb{N}^m$ and $b' = (b'_1, \ldots, b'_n)$, $c' = (c'_1, \ldots, c'_n) \in \mathbb{N}^n$. For such a sequence h(z), the product $h(z)\overline{h(z)}$ has a unique factorization with $2a'_i$ factors of each f_i and $b'_j + c'_j$ factors of each g_j . Since $f(z)\overline{f(z)}$ has $2a_i$ factors of each f_i and $b_j + c_j$ factors of each g_j , then h(z) cannot be isospectral to f(z) unless it is of the form h(z) = vH(z)where

$$H(z) = f_1(z)^{a_1} \cdots f_m(z)^{a_m} g_1(z)^{b'_1} \cdots g_n(z)^{b'_n} \overline{g_1(z)}^{c'_1} \cdots \overline{g_n(z)}^{c'_n}$$

with v a unit in $R[z, z^{-1}]$ and b' + c' = b + c. Let $F(z) = u^{-1}f(z)$, so f(z) = uF(z), so that $F(z)\overline{F(z)} = H(z)\overline{H(z)}$ Thus, $f(z)\overline{f(z)} = h(z)\overline{h(z)}$ if and only if $u\overline{u} = v\overline{v}$, i.e., if and only if $(v/u)\overline{(v/u)} = 1$, which happens if and only if $v/u = xz^k$ for some unimodular x and some $k \in \mathbb{Z}$. This is true if and only if $h(z) = vH(z) = xz^k uH(z)$, i.e., if and only if h(z) is R-similar to uH(z), and this is true if and only if h(z) is in the union specified in the statement of this theorem. This union is disjoint because the representatives that we show for the R-similarity classes in the union are all nonassociates of each other.

The number of *R*-similarity classes in our disjoint union equals the number of pairs $(b', c') \in \mathbb{N}^n \times \mathbb{N}^n$ such that b' + c' = b + c. This last constraint forces $b' \in \prod_{j=1}^n \{0, 1, \ldots, b_j + c_j\}$, and for each such b', there is a unique $c' = b + c - b' \in \mathbb{N}^n$ such that b' + c' = b + c, so we have precisely

(4)
$$\prod_{j=1}^{n} |\{0, 1, \dots, b_j + c_j\}| = \prod_{j=1}^{n} (b_j + c_j + 1) \text{ classes}$$

The conjugate of a representative

$$r(z) = u f_1(z)^{a_1} \cdots f_m(z)^{a_m} g_1(z)^{b'_1} \cdots g_n(z)^{b'_n} \overline{g_1(z)}^{c'_1} \cdots \overline{g_n(z)}^{c'_r}$$

of one of the R-similarity classes in our disjoint union is

$$\overline{r(z)} = \overline{u}wz^k f_1(z)^{a_1} \cdots f_m(z)^{a_m} g_1(z)^{c'_1} \cdots g_n(z)^{c'_n} \overline{g_1(z)}^{b'_1} \cdots \overline{g_n(z)}^{b'_n}$$

for some unimodular $w \in R$ and $k \in \mathbb{Z}$ because each of $f_1(z), \ldots, f_m(z)$ is *R*-palindromic. Since *u* is a unit in $R[z, z^{-1}]$, so is \overline{u} and then $\overline{u}/u = xz^j$ for some unimodular $x \in R$ and some $j \in \mathbb{Z}$. Therefore, $\overline{r(z)}$ is *R*-similar to

$$s(z) = u f_1(z)^{a_1} \cdots f_m(z)^{a_m} g_1(z)^{c'_1} \cdots g_n(z)^{c'_n} \overline{g_1(z)}^{b'_1} \cdots \overline{g_n(z)}^{b'_n}$$

so that the conjugate of $[r(z)]_R$ is $[s(z)]_R$. Thus, $[r(z)]_R$ is self-conjugate if and only if b' = c'. This can be true of only one class (the one with b' = c' = (b+c)/2 if $b_j + c_j$ is even for every j) or none at all (otherwise), hence the rest on the R-similarity classes occur in conjugate pairs. Recall from the Introduction that each trivial isospectrality class is either the disjoint union of two non-self-conjugate R-similarity classes or equal to a single self-conjugate isospectrality class. Let N be the number of R-similarity classes (counted by (4)) in our disjoint union (3). If N is even, then there are no self-conjugate R-similarity classes, so the number of trivial isospectrality classes in our disjoint union is $N/2 = \lceil N/2 \rceil$. If N is odd, then there is exactly one self-conjugate R-similarity class, so the number of trivial isospectrality class of f is the disjoint union of $\left[\left(\prod_{j=1}^n (b_j + c_j + 1)\right)/2\right]$ trivial isospectrality classes. Now if there are at least 3 classes in the disjoint union, then at least one

Now if there are at least 3 classes in the disjoint union, then at least one of them is not $[f]_R$ or $[\overline{f}]_R$, therefore f is non-trivially isospectral to some sequence.

Conversely, if f is non-trivially isospectral to some sequence g, then $[f]_R$ and $[g]_R$ are two distinct R-similarity classes in the disjoint union. At most one of these classes can be self-conjugate, and since the rest come in conjugate pairs, there must be at least three classes in the disjoint union. \Box

3. Palindromic classes

Since Theorem 1.1 only allows for at most one palindromic R-similarity class within an isospectrality class, the following corollary, which was stated as Corollary 1.4 in the Introduction, is immediate.

Corollary 3.1. Let R be a conjugate-closed subring of \mathbb{C} that is a unique factorization domain. If f and g are R-palindromes that are isospectral, then they must be R-similar.

Proof. If f = 0, then it is isospectral to g if and only if g = 0, in which case f and g are clearly R-similar. Assume that $f \neq 0$ henceforth. By Theorem 1.1 there is at most one palindromic class isospectral to f in the disjoint union (3) of R-similarity classes containing all the sequences in $R[z, z^{-1}]$ that are isospectral to f. Since f and g are both R-palindromes, they must be in this one R-similarity class, so f and g are R-similar.

Before we prove Corollary 1.5, we prove a preliminary result on R-similar R-palindromes.

Lemma 3.2. Let R be a conjugate-closed subring of \mathbb{C} that is a unique factorization domain. Let $u, v \in R^{\circledast}$ and let $f(z) \in R[z, z^{-1}]$ be a u-palindrome and $g \in R[z, z^{-1}]$ be a v-palindrome that is R-similar to f. Then either f = g = 0 or else $v/u \in R^{\circledast 2}$.

Proof. We know that $\overline{f(z)} = uz^i f(z)$ and $\overline{g(z)} = vz^j g(z)$ for some $i, j \in \mathbb{Z}$. Since f(z) and g(z) are *R*-similar, we have $f(z) = wz^k g(z)$ for some $w \in R^{\circledast}$ and $k \in \mathbb{Z}$. This proves that f = 0 if and only if g = 0, so assume both are nonzero from now on. Then

$$\begin{split} f(z) &= u^{-1} z^{-i} f(z) \\ &= u^{-1} z^{-i} \overline{w z^k g(z)} \\ &= u^{-1} z^{-i} w^{-1} z^{-k} \overline{g(z)} \\ &= u^{-1} z^{-i} w^{-1} z^{-k} v z^j g(z) \\ &= u^{-1} z^{-i} w^{-1} z^{-k} v z^j w^{-1} z^{-k} f(z) \\ &= (v/u) w^{-2} z^{-i+j-2k} f(z). \end{split}$$

Since $R[z, z^{-1}]$ is an integral domain, this forces $v/u = w^2 \in R^{\otimes 2}$.

Now we can prove Corollary 1.5 readily.

Corollary 3.3. Let R be a conjugate-closed subring of \mathbb{C} that is a unique factorization domain. Let $u, v \in R^{\circledast}$ and let $f(z) \in R[z, z^{-1}]$ be a u-palindrome and $g \in R[z, z^{-1}]$ be a v-palindrome that is isospectral to f. Then either f = g = 0 or else $v/u \in R^{\circledast 2}$.

Proof. Corollary 3.1 shows that f and g must be R-similar, so then the conclusion follows from Lemma 3.2.

Since 1 is the only square unit in \mathbb{Z} , we have the following consequence, recorded in the Introduction as Corollary 1.6.

Corollary 3.4. It is not possible for a palindrome in $\mathbb{Z}[z, z^{-1}]$ to be isospectral to an antipalindrome in $\mathbb{Z}[z, z^{-1}]$ unless both of the sequences are 0.

Proof. When $R = \mathbb{Z}$, we have $R^{\circledast} = \{1, -1\}$ and so $R^{\circledast 2} = \{1\}$, so by Corollary 3.3, if a *u*-palindrome *f* is isospectral to a *v*-palindrome *g* then either f = g = 0 or u = v.

4. BARREN INTEGERS

If $a = a_0 + a_1 z + \dots + a_{\ell-1} z^{\ell-1}$ is a binary sequence of length ℓ and $b = b_0 + b_1 z + \dots + b_{m-1} z^{m-1}$ is a binary sequence of length m, then the concatenation of a with b is the sequence $a(z) + z^{\ell}b(z)$. Notice that if $a(z) = a_0 + \dots + a_{\ell-1} z^{\ell-1}$ is a binary sequence of length ℓ , then the concatenation of a with a is $a(z) + z^{\ell}a(z) = (1 + z^{\ell})a(z)$. Our goal for this section is to prove Proposition 1.7, which we do using a concatenation construction. The following result shows how concatenation of sequences works in our Laurent polynomial formalism; it is readily proved by induction, so we record it without proof.

Lemma 4.1. Let $a(z) = a_0 + a_1 z + \cdots + a_{\ell-1} z^{\ell-1}$ be a sequence of length ℓ and let m be a positive integer. The sequence obtained by concatenating mcopies of a together is $(1 + z^{\ell} + z^{2\ell} + \cdots + z^{(m-1)\ell})a(z)$.

Now we are ready to restate and prove 1.7.

Proposition 4.2. Let m, n be positive integers such that m|n. If m is not barren, then n is not barren.

Proof. If m is not barren, then there exist nontrivially isospectral binary sequences f(z) and g(z) of length m. Since m|n, we have $n = m\ell$ for some integer ℓ . Since f(z) and g(z) are isospectral, we know $f(z)\overline{f(z)} = g(z)\overline{g(z)}$. We can concatenate ℓ copies of f(z) together to make F(z) and ℓ copies of g(z) together to make G(z). Let $h(z) = 1 + z^m + z^{2m} + \cdots z^{(\ell-1)m}$. Then by Lemma 4.1, F(z) = h(z)f(z), and G(z) = h(z)g(z).

$$F(z)\overline{F(z)} = h(z)f(z)\overline{h(z)f(z)}$$
$$= h(z)\overline{h(z)}f(z)\overline{f(z)}$$
$$= h(z)\overline{h(z)}g(z)\overline{g(z)}$$
$$= G(z)\overline{G(z)},$$

so F(z) and G(z) are isospectral sequences of length n. Now we need to show they are nontrivially isospectral.

Suppose that F(z) and G(z) are Z-similar to show contradiction. Then $F(z) = \pm z^j G(z)$ for some $j \in \mathbb{Z}$. Therefore, $h(z)f(z) = \pm z^j G(z) = \pm z^j h(z)g(z)$. Since $\mathbb{Z}[z, z^{-1}]$ is an integral domain, we get $f(z) = \pm z^j g(z)$. This would mean that f(z) and g(z) are Z-similar, but f(z) and g(z) are nontrivially isospectral, so this cannot happen. Therefore, F(z) and G(z) are not Z-similar.

Suppose that F(z) is \mathbb{Z} -similar to $\overline{G(z)}$ to show contradiction. That is $F(z) = \pm z^k \overline{G(z)}$ for some $k \in \mathbb{Z}$. Then

$$h(z)f(z) = \pm z^k \overline{G(z)} = \pm \overline{h(z)g(z)} = \pm z^{(1-\ell)m} h(z)\overline{g(z)}$$

Since $\mathbb{Z}[z, z^{-1}]$ is an integral domain, we have that $f(z) = \pm z^{(1-\ell)m}\overline{g(z)}$. This would mean that f(z) is \mathbb{Z} -similar to $\overline{g(z)}$, but this cannot happen because f(z) and g(z) are nontrivially isospectral. Therefore, F(z) and $\overline{G(z)}$ are not \mathbb{Z} -similar.

Since F(z) is *R*-similar to neither G(z) nor $\overline{G(z)}$, we see that F(z) and G(z) are nontrivially isospectral sequences of length *n*. Hence *n* is not barren.

References

- [BE22] Tamir Bendory and Dan Edidin. Algebraic theory of phase retrieval. Notices Amer. Math. Soc., 69(9):1487–1495, 2022.
- [BP15] Robert Beinert and Gerlind Plonka. Ambiguities in one-dimensional discrete phase retrieval from Fourier magnitudes. J. Fourier Anal. Appl., 21(6):1169–1198, 2015.
- [DJ99] James A. Davis and Jonathan Jedwab. Peak-to-mean power control in OFDM, Golay complementary sequences, and Reed-Muller codes. *IEEE Trans. Inform. Theory*, 45(7):2397–2417, 1999.
- [Fej16] Leopold Fejér. Über trigonometrische Polynome. J. Reine Angew. Math., 146:53– 82, 1916.

- [Gol51] Marcel J. E. Golay. Static multislit spectrometry and its application to the panoramic display of infrared spectra. J. Opt. Soc. Am., 41(7):468–472, Jul 1951.
- [Gol61] Marcel J. E. Golay. Complementary series. IRE Trans., IT-7:82–87, 1961.
- [KM22] Daniel J. Katz and Eli Moore. Sequence pairs with lowest combined autocorrelation and crosscorrelation. IEEE Trans. Inform. Theory, 68(12):8251–8267, 2022.

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, NORTHRIDGE, UNITED STATES

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, NORTHRIDGE, UNITED STATES

Department of Mathematics, California State University, Northridge, United States