

FINITE GROUPS, SMOOTH INVARIANTS AND ISOLATED QUOTIENT SINGULARITIES

AMIRAM BRAUN

Dept. of Mathematics, University of Haifa, Haifa, Israel 31905

abraun@math.haifa.ac.il

ABSTRACT. Let $G \subset SL(V)$ be a finite group, $p = \text{char } F$ and $\dim_F V$ is finite. Let $S(V)$ be the symmetric algebra of V , $S(V)^G$ the subring of G -invariants, and V^* the dual space of V .

Theorem A. $S(V)^G$ is a polynomial ring if and only if:

- (1) G is generated by transvections and,
- (2) $S(V)^{G_U}$ is a polynomial ring for each subspace $U \subset V^*$ with $\dim_F U = 1$, where $G_U = \{g \in G \mid g(u) = u, \forall u \in U\}$.

As a consequence we obtain the following classification result. \hat{A} stands for the completion of A at its unique graded maximal ideal.

Theorem B. Suppose F is algebraically closed and $S(V)^G$ is an isolated singularity. Then $\widehat{S(V)^G} \cong \widehat{S(W)^H}$, where $\dim_F W = \dim_F V$, $(|H|, p) = 1$ and (H, W) is a mod p reduction (in the sense of Brauer) of a member in the Zassenhaus-Vincent-Wolf list of complex isolated quotient singularities (e.g. [26])

Mathematics Subject Classification: Primary: 13A50, 14B05, 14J17, 14L30; Secondary: 14L24.

1. Introduction

Let $G \subset GL(V)$ be a finite group, F a field and $\dim_F V$ is finite. Let $S(V)$ be the symmetric algebra of V and $S(V)^G := \{x \in S(V) \mid g(x) = x, \forall g \in G\}$. The basic question we address here is:

Question. When is $S(V)^G$ a polynomial ring:

This was completely settled in the non-modular case as follows.

Date: August 31, 2023.

Theorem. (*Shephard-Todd-Chevalley-Serre*, [1, Sec. 7.2]). *Suppose $(|G|, p) = 1$, where $p = \text{char } F$ or $p = 0$. Then $S(V)^G$ is a polynomial ring if and only if G is generated by pseudo-reflections.*

In the modular case, that is if $p \mid |G|$, the above question is still open. Special cases are described next.

Theorem. (*H. Nakajima*, [17]). *Suppose G is a p -group and $F = F_p$, the prime field. Then $S(V)^G$ is a polynomial ring if and only if V has a basis $\{x_1, \dots, x_n\}$ such that*

- (1) $Fx_1 + \dots + Fx_i$ is a G -module for $i = 1, \dots, n$;
- (2) $|\text{orbit}(x_1)| \cdots |\text{orbit}(x_n)| = |G|$.

Theorem. (*Kemper-Malle*, [12], [8, p. 119]). *Suppose V is an irreducible G -module. Then $S(V)^G$ is a polynomial ring, if and only if:*

- (1) G is generated by pseudo-reflections, and;
- (2) $S(V)^{G_U}$ is a polynomial ring, for each subspace $U \subset V^*$, the dual space of V , where $G_U = \{g \in G \mid g(u) = u, \forall u \in U\}$ (*Steinberg condition*).

For an arbitrary V we have:

Theorem. (*Serre* [19], [5, V, §6, Ex. 8]). *Suppose $S(V)^G$ is a polynomial ring. Then (1) and (2) hold.*

The necessity of condition (2) was firstly shown in [21] in case $F = \mathbb{C}$. So Kemper-Malle's theorem implies that Serre's necessary conditions are also sufficient, if V is also irreducible. This suggests that the irreducibility assumption on V can be removed.

Conjecture C. *$S(V)^G$ is a polynomial ring if and only if:*

- (1) G is generated by pseudo-reflections, and;
- (2) $S(V)^{G_U}$ is a polynomial ring for each subspace $U \subset U^*$ with $\dim_F U = 1$.

The shift to $\dim_F U = 1$ is a simple consequence of the above theorem by Serre. A proof of Conjecture C, in case $\dim_F V = 3$, is given by D. A. Stepanov [23] and Shchigolev-Stepanov [20].

An analog result for p -groups was recently established in:

Theorem. (*A. Braun [2]*). *Suppose G is a finite p -group and $\dim_F V \geq 4$. Then $S(V)^G$ is a polynomial ring if and only if:*

- (1) $S(V)^G$ is a Cohen-Macaulay, and;
- (2) $S(V)^{G_U}$ is a polynomial ring, for each $U \subset V^*$ with $\dim_F U = 2$.

Our main objective here is to affirm the previous conjecture if $G \subset SL(V)$. We prove the following.

Theorem A. *Suppose $G \subset SL(V)$ is a finite group. Then $S(V)^G$ is a polynomial ring if and only if:*

- (1) G is generated by transvections, and;
- (2) $S(V)^{G_U}$ is a polynomial ring for each subspace $U \subset V^*$ with $\dim_F U = 1$.

Recall that if $G \subset SL(V)$, the only possible pseudo-reflection $g \in G$ is a transvection, that is $\text{rank}(g - 1) = 1$ and $(g - 1)^2(V) = 0$. In particular $g^p = 1$, where $p = \text{char } F$.

Actually condition (1) can be weakened in Theorem A to:

- (1') G is generated by its p -sylov subgroups.

Indeed let $g \in G$, $g^{p^e} = 1$, be one of the generators in (1'). Then $(g - 1)^{p^e} = 0$ on V^* , implying that $\ker_{V^*}(g - 1) \neq 0$, and hence $\exists U \subset V^*$, $\dim_F U = 1$, with $g \in G_U$. By (2) G_U is generated by pseudo-reflections which are necessarily transvections, since $G \subset SL(V)$. So each such g can be replaced by these transvection generators.

The proof of Theorem A is achieved by combining the following two separate theorems.

Theorem 1. *Suppose conditions (1) and (2) of Theorem A hold. Let $W \subset V$ be a proper G -submodule. Set $H := \{g \in G \mid (g - 1)(V) \subseteq W\}$, $m = \dim_F W$. Then:*

- (1) $(WS(V))^H = z_1 S(V)^H + \cdots + z_m S(V)^H$, where $z_1, \dots, z_m \in S(V)^G$ and are part of a minimal homogenous generating set of the polynomial ring $S(V)^H$;
- (2) $S(V)^G / (z_1, \dots, z_m) \cong [S(V)^H / (WS(V))^H]^{G/H}$ and is an isolated singularity as well as Cohen-Macaulay.

Theorem 2. *Let $G \subset SL(U)$ be a finite group generated by transvections, where U is an irreducible G -module and F is algebraically closed. Let $A \subset S(U)$ be a graded subring having the following properties:*

- (1) $A = F[x_1, \dots, x_d]$ is a polynomial subring, $\dim A = d = \dim_F U$, where $\{x_1, \dots, x_d\}$ are homogenous generators;
- (2) G acts faithfully by graded automorphisms on A .

Then $\deg x_1 = \dots = \deg x_d$ and $M := Fx_1 + \dots + Fx_d$ is an irreducible G -module.

Consequently the action of G on $A = S(M)$ is obtained from the linear irreducible action of G on M .

The proof of Theorem 2 relies on finite group classification results, listing all the possible pairs (G, U) . One then goes through the list and verifies, case by case, the validity of the statements. This rather lengthy procedure is dealt with in Section 3.

The proof of Theorem 1 is motivated and partially relies on the practice of changing polynomial ring generators, as initiated in [2]. The relevance of such a result is suggested by [17, Lemma 2.13].

The proof of Theorem A is then established by choosing $U = V/W$ where W is a maximal G -submodule of V . Theorem 2 implies that the action of G/H on the polynomial ring $A := S(V)^H/(WS(V))^H$ is a linear action. Since $A^{G/H}$ is an isolated singularity, it follows that [12] can be applied to conclude that $A^{G/H}$ is a polynomial ring. However this last step also requires that G/H is generated by pseudo-reflections on $M = Fx_1 + \dots + Fx_d$, (actually on M^*), where $A = F[x_1, \dots, x_d]$ ($d = \dim_F U$, $\deg x_1 = \dots = \deg x_d$). To establish this we crucially use the facts that $G \subseteq SL(V)$ as well as $G/H \subseteq SL(U)$ are generated by transvections (on V , respectively U), and consequently the (abstract) group G/H is generated by its p -sylov subgroups.

The lack of this last fact is a serious obstacle in removing the assumption $G \subseteq SL(V)$ in Theorem A, and thus in proving Conjecture C in full generality.

Theorem B states that a modular isolated quotient singularity, if $G \subset SL(V)$, is isomorphic, after completion to a non-modular one. This in turn can be obtained as the completion of a mod p reduction of a complex isolated quotient singularity. The list of all complex isolated quotient singularities is due to Zassenhaus-Vincent-Wolf, and can be seen e.g. in [26]. They feature in differential geometry as the solutions to the "Clifford-Klein spherical space form problem".

The key ingredients in the proof of Theorem B are that $S(V)^{T(G)}$ is a polynomial ring, where $T(G) := \langle g | g \text{ is a transvection on } V \rangle$, and that $(|G/T(G)|, p) = 1$. Both are

consequences of Theorem A. Now this implies by H. Cartan's theorem that:

$$\widehat{S(V)^G} = (S(V)^{\widehat{T(G)}})^{G/T(G)} \cong S(\widehat{W})^{G/T(G)},$$

where $W := Fy_1 + \cdots + Fy_n$, $F[[y_1, \dots, y_n]] = S(\widehat{V})^{\widehat{T(G)}}$, and $G/T(G)$ acts linearly on W . Also one may assume that W is void of fixed points for each $h \in G/T(G)$, $h \neq 1$. This last fact is a key property in the classification of complex isolated quotient singularities. The shift, from complex classification results to those over algebraically closed F with $\text{char } F = p > 0$, is possible since $(|G/T(G)|, p) = 1$, and this is well known to finite group theorists (and others). It is sketched (e.g.) in [22, Thm. 3.13]. All of this is detailed in Section 4.

The following is obtained as a consequence of Theorem A (without appealing to Theorem B) and a result of S. Kovacs [14].

Proposition. *Let $G \subset SL(V)$ be a finite group, $\dim_F V$ is finite and F is perfect. Suppose $S(V)^G$ is an isolated singularity. Then:*

- (1) $S(V)^G$ is a rational singularity, and
- (2) $S(V)^G$ has a non-commutative crepant resolution, provided it is Gorenstein.

For the definition of rational singularity in prime characteristic we refer to [14, Def. 1.3].

2. Theorem A

Throughout this section we shall use the following notations:

$W \subset V$ is a G -submodule, $H := \{g \in G \mid (g-1)(V) \subseteq W\}$. Clearly H is a normal subgroup of G .

Lemma 2.1. *Let $y = a_n v^n + \cdots + a_1 v + a_0 \in (WS(V))^H$, be a homogenous element with $a_i \in WS(V')$, where $W \subseteq V' \subset V$ is a codimension 1 subspace of V , and $v \in V$, with $Fv + V' = V$. Then:*

$$\left[\sum_{s=0}^{i-1} \phi(a_{n-s}) \delta(v)^{i-s} \binom{n-s}{i-s} \right] + \delta(a_{n-i}) = 0, \text{ where } \phi \in H \text{ and } \delta := \phi - 1.$$

Proof: $0 = \delta(y) = \delta(a_n v^n + \cdots + a_1 v + a_0)$. We shall compute the coefficient of v^{n-i} in the last expression. Recall that $\delta(v^j) = \sum_{r=0}^{j-1} v^r \delta(v)^{j-r} \binom{j}{r}$ (e.g [2, p.7]). $\delta(a_j v^j) = \delta(a_j) v^j + \phi(a_j) \delta(v^j) = \delta(a_j) v^j + \phi(a_j) (\sum_{r=0}^{j-1} v^r \delta(v)^{j-r} \binom{j}{r})$.

So for $j = n, n-1, \dots, n-i+1$, v^{n-i} will occur with coefficient $\phi(a_j)\delta(v)^{j-(n-i)}\binom{j}{n-i}$ (taking $r = n-i$).

For $j = n-i$, the coefficient of v^{n-i} is $\delta(a_{n-i})$. Changing indices by taking $j = n-s$, we get that the coefficient of v^{n-i} in the above expression is:

$$A_{n-i} := \sum_{s=0}^{i-1} \phi(a_{n-s})\delta(v)^{i-s}\binom{n-s}{i-s} + \delta(a_{n-i}), \text{ (using } \binom{n-s}{n-i} = \binom{n-s}{i-s}\text{)}.$$

Now $0 = \sum_{i=0}^n A_{n-i}v^{n-i}$. Also $\delta(V) \subseteq W$ implies that $\phi(a_{n-s})\delta(v)^{i-s}, \delta(a_{n-i}) \in WS(V')$. Consequently $A_{n-i} \in WS(V')$ and the above algebraic equation of v over $S(V')$ implies $A_{n-i} = 0, i = 0, \dots, n$. \square

Lemma 2.2. *Keeping the notation of Lemma 2.1, we have for $k < n$:*

$$\sum_{i=0}^k \binom{n-i}{k-i} a_{n-i} v^{k-i} \in (WS(V))^H.$$

Proof: Let $\phi \in H$ and $\delta := \phi - 1$. Using $\delta(xy) = \delta(x)y + \phi(x)\delta(y)$, we get:

$$\begin{aligned} & \delta\left[\sum_{i=0}^k \binom{n-i}{k-i} a_{n-i} v^{k-i}\right] \\ &= \sum_{i=0}^k \binom{n-i}{k-i} \delta(a_{n-i})v^{k-i} + \sum_{i=0}^{k-1} \phi(a_{n-i}) \left(\sum_{j=1}^{k-i} v^{k-i-j} \delta(v)^j \binom{k-i}{j}\right) \\ &= \sum_{t=0}^k v^{k-t} \left\{ \sum_{s=0}^{t-1} \binom{n-s}{k-s} \phi(a_{n-s})\delta(v)^{t-s} \binom{k-s}{t-s} + \binom{n-t}{k-t} \delta(a_{n-t}) \right\} \end{aligned}$$

(using the convention $\sum^{-1} = 0$). But $\binom{n-s}{k-s} \binom{k-s}{t-s} = \binom{n-s}{n-t} \binom{n-t}{k-t} = \binom{n-s}{t-s} \binom{n-t}{k-t}$, implying that the coefficient of v^{k-t} is:

$$\begin{aligned} & \left\{ \left[\sum_{s=0}^{t-1} \phi(a_{n-s})\delta(v)^{t-s} \binom{n-s}{t-s} \right] + \delta(a_{n-t}) \right\} \binom{n-t}{k-t} \\ &= A_{n-t} \binom{n-t}{k-t} = 0, \end{aligned}$$

where the last equality is by Lemma 2.1. \square

Proposition 2.3. *Let $y = a_n v^n + \dots + a_1 v + a_0 \in (W(S(V)))^H$, be an homogenous element, where $W \subseteq V' \subset V$, $Fv + V' = V$, and $a_i \in WS(V')$, $i = 0, \dots, n$. Then $y = b_n v^n + \dots + b_1 v + a_0$, where $b_i \in (WS(V))^H$, $i = 1, \dots, n$, and $\deg b_i = \deg a_i < \deg y$, $i = 1, \dots, n$.*

Proof: By Lemma 2.2 (taking $k = n-1$), we get

$$b_1 = \binom{n}{n-1} a_n v^{n-1} + \binom{n-1}{n-2} a_{n-1} v^{n-2} + \dots + \binom{2}{1} a_2 v + a_1 \in (WS(V))^H.$$

Hence:

$$y = [-\binom{n}{n-1} + 1]a_nv^n + [-\binom{n-1}{n-2} + 1]a_{n-1}v^{n-1} + \cdots + [-\binom{2}{1} + 1]a_2v^2 + b_1v + a_0.$$

By using Lemma 2.2 (taking $k = n - 2$), we replace $[-\binom{2}{1} + 1]a_2$ by:

$$b_2 := [-\binom{2}{1} + 1]\left\{\binom{n}{n-2}a_nv^{n-2} + \binom{n-1}{n-3}a_{n-1}v^{n-3} + \cdots + \binom{3}{1}a_3v + a_2\right\} \in (WS(V))^H$$

and

$$y = [1 - \binom{n}{n-1} - (-\binom{2}{1} + 1)\binom{n}{n-2}]a_nv^n + \cdots + [1 - \binom{3}{2} - (-\binom{2}{1} + 1)\binom{3}{1}]a_3v^3 + b_2v^2 + b_1v + a_0.$$

One continues in this way, constructing b_3, \dots, b_n . \square

Note 2.4.

- (1) b_1, \dots, b_n depend on the elements a_1, \dots, a_n and hence on the choices of V', v .
- (2) Proposition 2.3 is a key result, enabling the replacement of each member of a minimal homogenous generating set of $(WS(V))^H$ by one in $S(V)^G$, as claimed in Theorem 1. The element y is repeatedly rewritten in accordance to the pseudo-reflection used ψ .

We shall also need the following simple observation.

Lemma 2.5. *Let $\psi \in G - H$ be a pseudo-reflection. Then $W \subseteq \ker(\psi - 1)$.*

Proof: $\psi \notin H$ and hence $(\psi - 1)(V) \not\subseteq W$. If $W \not\subseteq \ker(\psi - 1)$ then $(\psi - 1)(W) \neq 0$, and then $(\psi - 1)(V) = (\psi - 1)(W)$ (since $\text{rank}(\psi - 1) = 1$). But $(\psi - 1)(W) \subseteq W$ (since W is a G -submodule), leading to a contradiction. \square

We shall need the following version of [2, Prop. 2.13].

Proposition 2.6. *Let $A = \bigoplus_n A_n$ be an \mathbb{N} -graded polynomial ring over a field $F = A_0$ and $\mathfrak{G} \subset \text{Aut}_{\text{gr}}(A)$ a finite subgroup of graded automorphisms. Let $x \in A^{\mathfrak{G}}$ be an homogenous element. Set $\mathfrak{H} := \ker(\mathfrak{G} \rightarrow \mathfrak{G}|_{A/xA})$, $\mathfrak{m} := A_+$, the irrelevant maximal ideal of A and $\mathfrak{n} := A_+^{\mathfrak{G}}$. Suppose:*

- (1) $x \in \mathfrak{m} - \mathfrak{m}^2$;
- (2) $A_{\mathfrak{n}}^{\mathfrak{G}}$ satisfies Serre's S_{m+1} and R_m conditions with $m \geq 2$.

Then $A^{\mathfrak{G}}/(x) = (A^{\mathfrak{H}}/xA^{\mathfrak{H}})^{\mathfrak{G}/\mathfrak{H}}$, and $[A^{\mathfrak{G}}/(x)]_{\mathfrak{n}/(x)}$ satisfies S_m and R_{m-1} .

Note. In [2, Prop 2.13] the following extra assumption is required: " $\mathfrak{G}/\mathfrak{H}$ acts faithfully on $A^{\mathfrak{H}}/xA^{\mathfrak{H}}$ ". This is obsolete in view of [17, Lemma 2.8], by taking $\mathfrak{p} := xA$, which is \mathfrak{G} -stable and \mathfrak{H} is by definition the inertia subgroup of \mathfrak{p} .

Actually we shall need the stronger version of Proposition 2.6 where $A_{\mathfrak{n}}^{\mathfrak{G}}, [A^{\mathfrak{G}}/(x)]_{\mathfrak{n}/(x)}$ are replaced by $A^{\mathfrak{G}}, A^{\mathfrak{G}}/(x)$ (respectively). For that we use the following:

Lemma 2.7. *Let $B = \bigoplus_{n \geq 0} B_n$ be a commutative Noetherian graded ring with $B_0 = F$, a field. Let $\mathfrak{n} := \bigoplus_{n > 0} B_n$. Then $B_{\mathfrak{n}}$ satisfies R_m, S_m , if and only if B satisfies R_m, S_m (respectively).*

Proof: Let $\mathfrak{p} \in \text{spec } R$ with height $= m$, $\mathfrak{p}^* :=$ the largest graded ideal in \mathfrak{p} . Then by [4, Lemma 1.5.6(a)] \mathfrak{p}^* is a prime ideal and $\text{height}(\mathfrak{p}) = 1 + \text{height}(\mathfrak{p}^*)$, if \mathfrak{p} is not graded [4, Thm. 1.5.8(b)]. Consequently if \mathfrak{p} is not graded then $\text{height}(\mathfrak{p}^*) = m - 1$. But $\mathfrak{p}^* \subset \mathfrak{n}$ implying, if $B_{\mathfrak{n}}$ satisfies R_m , that $B_{\mathfrak{p}^*}$ is regular. This implies by [4, Ex. 2.24(a)] that $B_{\mathfrak{p}}$ is regular. Hence B satisfies R_m . The converse implication is trivial.

Similarly by [4, Thm. 1.5.9] $\text{depth } B_{\mathfrak{p}} = \text{depth } B_{\mathfrak{p}^*} + 1$, if \mathfrak{p} is not graded. So if $B_{\mathfrak{n}}$ satisfies Serre's S_m condition then $\text{depth } B_{\mathfrak{p}^*} \geq \min(m, \text{height } \mathfrak{p}^*)$. Hence if \mathfrak{p} is not graded $\text{depth } B_{\mathfrak{p}} = 1 + \text{depth } B_{\mathfrak{p}^*} \geq \min(1 + \text{height } \mathfrak{p}^*) = \min(m, \text{height } \mathfrak{p})$. \square

Using similar arguments we have:

Lemma 2.8. *Let $B = \bigoplus_{n \geq 0} B_n$ be a graded Noetherian commutative ring with $B_0 = F$, a field and $d = \dim B$. Suppose B satisfies R_{d-1} . Then B is an isolated singularity.*

Proof: Let $\mathfrak{n} := \bigoplus_{n \geq 1} B_n$, the unique graded maximal ideal of B . Let \mathfrak{p} be a maximal ideal of B which is not graded. Let \mathfrak{p}^* be the largest graded ideal in \mathfrak{p} . As in Lemma 2.7 $\text{height}(\mathfrak{p}) = 1 + \text{height}(\mathfrak{p}^*)$. Hence $\text{height}(\mathfrak{p}^*) \leq d - 1$ and therefore $B_{\mathfrak{p}^*}$ is regular implying by [4, Ex. 2.24(a)] that $B_{\mathfrak{p}}$ is regular. \square

Corollary 2.9. *Suppose F is perfect. Then the following are equivalent:*

- (1) $S(V)^G$ is an isolated singularity;
- (2) $S(V)^{G_U}$ is a polynomial ring for each $U \subset V^*$, $\dim_F U = 1$.

Proof: By Lemma 2.8 item (1) is equivalent to: $S(V)^G$ satisfies R_{d-1} , where $d = \dim_F V = \dim S(V)^G$. The rest follows from [2, Lemma 2.4]. \square

The following is a consequence of Proposition 2.6 and Lemma 2.7. Actually it is part of the proof of Theorem 1.

Lemma 2.10. *Suppose $H := \{g \in G \mid (g-1)(V) \subseteq W\}$ and $(z_1, \dots, z_k, y_{k+1}, \dots, y_m) = (WS(V))^H$, where $\{z_1, \dots, z_k, y_{k+1}, \dots, y_m\}$ is part of a minimal homogenous generating set of the polynomial ring $S(V)^H$. Assume also that:*

- (1) $\{z_1, \dots, z_k\} \subseteq S(V)^G$;
- (2) $S(V)^G$ satisfies Serre's R_{d-1} condition ($d = \dim_F V$) and is Cohen-Macaulay.

Then $S(V)^G/(z_1, \dots, z_k) \cong (S(V)^H/z_1S(V)^H + \dots + z_kS(V)^H)^{G/H}$, and it satisfies Serre's R_{d-1-k} condition as well as being Cohen-Macaulay.

Proof: This is done by induction. By assumption $z_1, \dots, z_k \in \mathfrak{m} - \mathfrak{m}^2$ where $\mathfrak{m} := S(V)_+^H$, the irrelevant maximal ideal of $S(V)^H$. Suppose we have by induction that

$$S(V)^G/(z_1, \dots, z_{i-1}) \cong (S(V)^H/z_1S(V)^H + \dots + z_{i-1}S(V)^H)^{G/H}$$

satisfies $R_{d-1-(i-1)}$ -condition and is Cohen-Macaulay.

Set $I := z_1S(V)^H + \dots + z_{i-1}S(V)^H$, $J := z_1S(V)^H + \dots + z_iS(V)^H$, $A := S(V)^H/I$, $\bar{\mathfrak{m}} := \mathfrak{m}/I$, $\mathfrak{G} := G/H$, $x := \bar{z}_i = z_i + (z_1, \dots, z_{i-1}) \in S(V)^G/(z_1, \dots, z_{i-1})$.

$x \in \bar{\mathfrak{m}} - \bar{\mathfrak{m}}^2$, since \bar{z}_i is part of the minimal generating set of the polynomial ring A .

Since $\mathfrak{G} = G/H$ acts faithfully on $S(V)^H/(WS(V))^H$ by [17, Lemma 2.8] and $J \subset (WS(V))^H$, it follows that \mathfrak{G} acts faithfully on $S(V)^H/J = A/xA$. Hence $\mathfrak{H} := \ker(\mathfrak{G} \rightarrow \mathfrak{G}|_{A/xA}) = 1$. Therefore by Proposition 2.6 $A^{\mathfrak{G}}/(x) \cong (A/xA)^{\mathfrak{G}}$. Hence:

$$(2.1) \quad (S(V)^H/I)^{G/H} / \bar{z}_i(S(V)^H/I)^{G/H} \cong (S(V)^H/J)^{G/H}.$$

By the inductive assumption, the left hand side of (2.1) is isomorphic to:

$$[S(V)^G/(z_1, \dots, z_{i-1})] / \bar{z}_i[S(V)^G/(z_1, \dots, z_{i-1})] = S(V)^G/(z_1, \dots, z_i).$$

In conclusion $S(V)^G/(z_1, \dots, z_i) \cong [S(V)^H/z_1S(V)^H + \dots + z_iS(V)^H]^{G/H}$. Also by Proposition 2.6 and Lemma 2.7 $[S(V)^G/(z_1, \dots, z_i)]$ satisfies R_{d-1-i} and S_{d-i} . Since $\dim S(V)^G/(z_1, \dots, z_i) = d - i$, it follows that it is Cohen-Macaulay. \square

The following result is classical. The earliest reference I could find is [6], but it only deals with the local analog. Similarly [4, Prop. 2.2.4], [18, Cor. 1.5] merely handle the local case. So here is a direct proof.

Lemma 2.11. *Let $A = F[x_1, \dots, x_n]$ be a graded polynomial ring, where x_1, \dots, x_n are homogenous elements. Suppose A/\mathfrak{n} is a graded polynomial ring, where $\mathfrak{n} \subseteq \mathfrak{m} := (x_1, \dots, x_n)$*

is a graded ideal. Then $\mathfrak{n} = y_{s+1}A + \cdots + y_nA$, $\{y_{s+1}, \dots, y_n\}$ is part of a minimal homogenous generating set of A , and $s = \dim A/\mathfrak{n}$

Proof: We clearly may assume that $s < n$,

Set $\bar{\mathfrak{m}} := \mathfrak{m}/\mathfrak{n} = (A/\mathfrak{n})_+$, $\bar{A} := A/\mathfrak{n}$. Then $s = \dim_{\bar{A}/\bar{\mathfrak{m}}} \bar{\mathfrak{m}}/\bar{\mathfrak{m}}^2 = \dim_{A/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = \dim_{A/\mathfrak{m}} \mathfrak{m}/(\mathfrak{m}^2 + \mathfrak{n})$. Hence $\{\tilde{x}_1, \dots, \tilde{x}_n\}$, the images of $\{x_1, \dots, x_n\}$ in $\mathfrak{m}/(\mathfrak{m}^2 + \mathfrak{n})$ are linearly dependent.

We may assume that $\{\tilde{x}_1, \dots, \tilde{x}_s\}$ are linearly independent over A/\mathfrak{m} . Also all the linear dependencies are taking place in the homogenous components of $\mathfrak{m}/(\mathfrak{m}^2 + \mathfrak{n})$. Hence for each $i > s$ we have $I_i \subset \{1, \dots, s\}$ such that $\tilde{x}_i = \sum_{j \in I_i} \alpha_{ij} \tilde{x}_j$, $\alpha_{ij} \in A$, $\deg \tilde{x}_i = \deg \tilde{x}_j$, $j \in I_i$. So since $A/\mathfrak{m} = F$ we can take $\alpha_{ij} \in F$. Therefore $\forall i > s$, $m_i := x_i - \sum_{j \in I_i} \alpha_{ij} x_j \in \mathfrak{m}^2 + \mathfrak{n}$. But since $\{x_1, \dots, x_n\}$ are linearly independent $\pmod{\mathfrak{m}^2}$, it follows that $m_i \notin \mathfrak{m}^2$, $\forall i > s$.

Clearly $\{x_1, \dots, x_s, m_{s+1}, \dots, m_n\}$ is a minimal generating set of $F[x_1, \dots, x_n]$. Let $y_i \in \mathfrak{n}$ be a homogenous element with $y_i - m_i \in \mathfrak{m}^2$, $i = s+1, \dots, n$. Then $A = F[x_1, \dots, x_n] = F[x_1, \dots, x_s, m_{s+1}, \dots, m_n] = F[x_1, \dots, x_s, y_{s+1}, \dots, y_n]$. So $\{y_{s+1}, \dots, y_n\}$ is part of a minimal generating set of A implying that $(y_{s+1}) \subset (y_{s+1}, y_{s+2}) \subset \cdots \subset (y_{s+1}, \dots, y_n)$ is a proper chain of prime ideals in A . Hence $\text{height}(y_{s+1}, \dots, y_n) = n - s$. But $(y_{s+1}, \dots, y_n) \subseteq \mathfrak{n}$ and $\text{height}(\mathfrak{n}) = n - \dim A/\mathfrak{n} = n - s$, so $(y_{s+1}, \dots, y_n) = \mathfrak{n}$. \square

Corollary 2.12. *Keeping the notation of this section, and assuming that $S(V)^H$ is a polynomial ring, where $H := \{g \in G \mid (g-1)(V) \subseteq W\}$. Then $(WS(V))^H = y_1 S(V)^H + \cdots + y_m S(V)^H$, where $m = \dim_F W$ and $\{y_1, \dots, y_m\}$ is a part of a minimal homogenous generating set of $S(V)^H$.*

Proof: Recall that $(WS(V))^H = WS(V) \cap S(V)^H$. Then by [17, Lemma 2.11] (the start of the proof of item (2), using $G(V, W) = H$), $S(V)^H / (WS(V))^H$ is a polynomial ring. Consequently by Lemma 2.11, with $A := S(V)^H$, $\mathfrak{n} := (WS(V))^H$, the result holds. The last equality follows since $\text{height}((WS(V))^H) = \text{height}(WS(V)) = \dim_F W$. \square

We shall now prove Theorem 1 (of the introduction).

Theorem 2.13. *Suppose $\dim_F V = d$, $G \subset SL(V)$ is generated by transvections and $S(V)^{G_U}$ is a polynomial ring $\forall U \subset V^*$, $\dim_F U = 1$. Let $W \subset V$ be a G -submodule and $H := \{g \in G \mid (g-1)(V) \subseteq W\}$. Then:*

- (1) $(WS(V))^H = z_1 S(V)^H + \cdots + z_m S(V)^H$, where $z_i \in S(V)^G$, $i = 1, \dots, m$ and $m = \dim_F W$,

- (2) $\{z_1, \dots, z_m\}$ is part of a minimal homogenous generating set of $S(V)^H$;
- (3) $S(V)^G/(z_1, \dots, z_m) \cong [S(V)^H/(WS(V))^H]^{G/H}$ and it satisfies Serre's condition R_{d-1-m} as well as being Cohen-Macaulay.

Proof: It follows from [2, Lemma 2.4] that $S(V)^G$ satisfies R_{d-1} and also $S(V)^{G_U}$ is a polynomial ring for each $U \subset V^*$, $\dim_F U \geq 1$. Consequently, since $H = G_{W^\perp}$ [2, Lemma 2.1], it follows that $S(V)^H$ is a polynomial (where $W^\perp = \{f \in V^* | f(W) = 0\}$). Also by [11, Thm. 3.1] it follows that $S(V)^{G_U}$ being a Cohen-Macaulay ring, for each $U \subset V^*$, $\dim_F U \geq 1$ implies that $S(V)^G$ is Cohen-Macaulay.

By Corollary 2.12 $(WS(V))^H = y_1 S(V)^H + \dots + y_m S(V)^H$, $m = \dim_F W = \text{height}((WS(V))^H)$, and $\{y_1, \dots, y_m\}$ are part of a minimal homogenous generating set of $S(V)^H$.

We order $\{y_1, \dots, y_m\}$, according to their degree's $\deg y_1 \leq \deg y_2 \leq \dots \leq \deg y_m$.

Observe that if $y_1 \in S(W)$, then $y_1 \in S(W) \cap S(V)^H = S(W)^H$. Let ψ be a pseudo-reflection with $\psi \notin H$. Then by Lemma 2.5 $W \subseteq \ker(\psi - 1)$, hence since $G = \langle H, \psi \mid \psi \text{ is a pseudo-reflection, } \psi \notin H \rangle$, we get that $y_1 \in S(W)^G \subseteq S(V)^G$.

We shall next show that $y_1 \in S(W)^H$. We assume that $y_1 \notin S(W)$. Let $W \subseteq V' \subset V$, with $\dim_F V' = \dim_F V - 1$, and $v \in V - V'$. H acts trivially on V/W implying that V' is an H -module. Suppose $y_1 \notin WS(V')$. Since $y_1 \in WS(V)$, then $y_1 = \sum_{i=0}^s a_i v^i$, $a_i \in WS(V')$, $i = 0, \dots, s$, $a_s \neq 0$, $s > 0$. It follows that $\deg y_1 = \deg a_i + i$, $i = 0, \dots, s$, and $\deg y_1 = \deg a_s + s > \deg a_s$.

Let $g \in H$ be arbitrary and $\delta = g - 1$. Then:

$$(2.2) \quad 0 = \delta(y_1) = \sum_{i=0}^s \delta(a_i) v^i + \sum_{i=0}^s a_i \delta(v^i) + \sum_{i=0}^s \delta(a_i) \delta(v^i).$$

Since $\delta(v^i) = i v^{i-1} \delta(v) + \sum_{j=0}^{i-2} \binom{i}{j} v^j \delta(v)^{i-j}$, equation (2.2) takes the form $0 = \delta(a_s) v^s + \sum_{i=0}^{s-1} x_i v^i$, $x_i \in S(V')$. Since $a_s \in WS(V')$, then $\delta(a_s) \in WS(V')$, so if $\delta(a_s) \neq 0$ we reach an algebraic dependence of v over $S(V')$. Therefore $\delta(a_s) = 0$, $\forall g \in H$. So $a_s \in (WS(V'))^H \subseteq (WS(V))^H$ and $\deg(a_s) < \deg y_1$, $\forall i$, an absurd. So $y_1 \in WS(V')$, Continuing in this way with $V'' \subset V'$, $\dim_F V'' = \dim_F V' - 1$, $W \subseteq V''$, $v \in V' - V''$, we reach $y_1 \in S(W)^H$, as claimed.

We shall now consider the following properties:

- (1') $(WS(V))^H = (z_1, \dots, z_k, y_{k+1}, \dots, y_m)$ with $\deg z_i = \deg y_i$, $i = 1, \dots, k$, and $\{z_1, \dots, z_k\} \subseteq S(V)^G$;

- (2') $\{z_1, \dots, z_k, y_{k+1}, \dots, y_m\}$ is a part of a minimal homogenous generating set of $S(V)^H$,
and;
(3') $S(V)^G/(z_1, \dots, z_k) \cong (S(V)^H/z_1S(V)^H + \dots + z_kS(V)^H)^{G/H}$, and it satisfies Serre's
 R_{d-1-k} -condition, as well as being Cohen-Macaulay.

It follows from Lemma 2.10, Corollary 2.12 and the previous reasoning that the choice $z_1 = y_1$ satisfies (1'), (2') and (3').

So we assume that (1'), (2') and (3') hold for k . Set $y := y_{k+1}$, so $\deg y \geq \deg z_i$, $i = 1, \dots, k$. We shall now proceed to show that we can replace y by $z \in S(V)^G$ and have in place all the requirements of (1'), (2') and (3').

Let $\psi \in G - H$, where ψ is a transvection. Set $V' := \ker(\psi - 1)$, $v \in V - V'$ with $Fv + V' = V$. By Lemma 2.5 $W \subseteq V'$. Also by Proposition 2.3 we have $y = b_nv^n + \dots + b_1v + a_0$, $b_i \in (WS(V))^H$, $\deg b_i = \deg a_i$. $i = 1, \dots, n$, $a_0 \in WS(V')$ (as well as $a_i \in WS(V')$, $i = 1, \dots, n$). Also $\deg b_i = \deg a_i < \deg y$, implying by (1') that $b_i \in z_1S(V)^H + \dots + z_kS(V)^H$. Since $z_i \in S(V)^G$, $i = 1, \dots, k$, it follows that $\psi(b_i), \delta(b_i) \in z_1S(V) + \dots + z_kS(V)$ where $\delta := \psi - 1$. Also $\delta(a_0) = 0$, since $\delta(V') = 0$. Consequently:

$$\delta(y) = \sum_{i=1}^n \delta(b_iv^i) + \delta(a_0) = \sum_{i=1}^n (\delta(b_i)v^i + \psi(b_i)\delta(v^i)) \in z_1S(V) + \dots + z_kS(V).$$

Also $S(V)^H$ is G -stable since H is normal in G . So $\delta(y) \in S(V)^H \cap (z_1S(V) + \dots + z_kS(V)) = S(V)^H \cap (z_1S(V)^H + \dots + z_kS(V)^H)S(V)$. Since $S(V)^H$ is a polynomial ring and $S(V)$ is Cohen-Macaulay (being a polynomial ring), it follows from [4, Prop. 1.5.15 and 2.2.11] that $S(V)$ is projective and hence free over $S(V)^H$. Hence by [15, Thm. 7.5(ii)] $S(V)^H \cap (z_1S(V)^H + \dots + z_kS(V)^H)S(V) = z_1S(V)^H + \dots + z_kS(V)^H$. Consequently $\delta(y) \in z_1S(V)^H + \dots + z_kS(V)^H$, for each such ψ . Observe that there is no reference here to V' any longer. Hence: $\bar{y} := y + z_1S(V)^H + \dots + z_kS(V)^H$ satisfies:

$$\bar{y} \in [S(V)^H/z_1S(V)^H + \dots + z_kS(V)^H]^{G/H} = S(V)^G/(z_1, \dots, z_k).$$

Therefore there exists $z \in S(V)^G$, $\deg z = \deg y$ with $z - y \in z_1S(V)^H + \dots + z_kS(V)^H$. Clearly $z_i, y \in (WS(V))^H$, $i = 1, \dots, k$, imply $z \in (WS(V))^H \cap S(V)^G$. Also $z_1S(V)^H + \dots + z_kS(V)^H + yS(V)^H = z_1S(V)^H + \dots + z_kS(V)^H + zS(V)^H$. Moreover if $z - y \neq 0$, then $\deg(z - y) = \deg y \geq \deg z_i$, $i = 1, \dots, k$, so $z - y = \sum_{i=1}^k z_i c_i$, $c_i \in S(V)^H$ and c_i when expressed in the homogenous generators of $S(V)^H$, the ones which appear must have degree smaller than $\deg y$, so y does not appear in the expression of c_i in these generators. So z

can replace y in the minimal homogenous generating set of $S(V)^H$. All in all we verified with $z_{k+1} := z \in S(V)^G$, that:

- (1') $(WS(V))^H = z_1 S(V)^H + \cdots + z_{k+1} S(V)^H + y_{k+2} S(V)^H + \cdots + y_m S(V)^H$;
- (2') $\{z_1, \dots, z_k, z_{k+1}, y_{k+2}, \dots, y_m\}$ is part of a minimal homogenous generating set of $S(V)^H$;

Finally (3') follows from Lemma 2.10.

The theorem now follows by taking $k = m$. \square

The proof of Theorem A

We are given that $G \subset SL(V)$ is generated by transvections on V (or V^*) and $S(V)^{G_U}$ is a polynomial ring for each $U \subset V^*$ with $\dim_F U = 1$. We want to show that $S(V)^G$ is a polynomial ring.

As in the proof of [2, Thm. B, p.20] we may assume that F is algebraically closed, as well as $S(V)^G$ satisfies R_{d-1} , where $d = \dim_F V$ and $S(V)^G$ is Cohen-Macaulay.

Let $W \subset V$ be a maximal G -submodule, that is $0 \neq V/W$ is a simple G -module. Set $H := \{g \in G \mid (g-1)(V) \subseteq W\}$. Hence by [2, Lemma 2.1] $H = G_{W^\perp}$, where $W^\perp := \{f \in V^* \mid f(W) = 0\}$, implying that $S(V)^H$ is a polynomial ring.

By Theorem 2.13 (= Theorem 1) $(WS(V))^H = z_1 S(V)^H + \cdots + z_m S(V)^H$, with $\{z_1, \dots, z_m\}$ are in $S(V)^G$, $m = \dim_F W = \text{height}(WS(V))^H$. Moreover z_1, \dots, z_m are part of a minimal homogenous generating set of $S(V)^H$, and $S(V)^G / (z_1, \dots, z_m) \cong [(S(V)^H / (WS(V))^H)]^{G/H}$, satisfying R_{d-1-m} and the Cohen-Macaulay property.

As in Corollary 2.12 $A := S(V)^H / (WS(V))^H$ is a polynomial ring with $\dim A = d - m = \dim_F V/W$. By [17, Lemma 2.8] G/H acts faithfully on A , this action is the restriction of the faithful (linear and irreducible) action of G/H on V/W and on $S(V/W)$. By [2, Lemma 2.6] G/H is generated by the transvections $\{\sigma_1, \dots, \sigma_t\}$ on V/W (and on $(V/W)^*$).

Let $A = F[x_1, \dots, x_{d-m}]$, where x_i is homogenous, $i = 1, \dots, d - m$. By Theorem 3.1 (=Theorem 2, which is proved in section 3) $\deg x_1 = \cdots = \deg x_{d-m}$ and $M = Fx_1 + \cdots + Fx_{d-m}$ is an irreducible G/H -module. Consequently $A = S(M)$ and the action of G/H on A is induced from a linear faithful action of G/H on M . Since $S(M)^{G/H} = A^{G/H}$ satisfies R_{d-1-m} and F is algebraically closed, it follows from [2, Lemma 2.4] that $S(M)^{(G/H)_U}$ is a polynomial ring for each subspace $U \subset M^*$ with $\dim_F U = 1$.

We have that $\sigma_i^p = 1$, $i = 1, \dots, t$. So $\sigma_i^p = 1$ also holds with σ_i , regarded now as an automorphism on $A = S(M)$ (although it may fail to act as a pseudo-reflection on M^*). However $(\sigma_i - 1)^p = 0$ on M^* , showing that $\ker(\sigma_i - 1) \neq 0$ (on M^*), hence $\exists U_i \subset M^*$, $\dim_F U_i = 1$ with $\sigma_i \in (G/H)_{U_i}$, $i = 1, \dots, t$. Since $S(M)^{(G/H)_{U_i}}$ is a polynomial ring it follows that $(G/H)_{U_i}$ is generated by the pseudo-reflections on M^* , $\forall i$, implying that G/H is generated by pseudo-reflections on M^* .

To conclude, G/H acting faithfully, linearly and irreducibly on M and hence on M^* , is generated by pseudo-reflections on M^* and $S(M)^{(G/H)_{U_i}}$ is a polynomial ring $\forall U \subset M^*$ with $\dim_F U = 1$. Invoking [12, Main theorem] we get that $A^{G/H} \cong S(V)^G/(z_1, \dots, z_m)$ is a polynomial ring in $d - m$ generators. Hence $S(V)^G$ is generated by $m + (d - m) = d$ elements, showing that it is a polynomial ring. \square

Theorem 1 and Theorem 2 imply the following necessary conditions.

Proposition 2.14. *Let $G \subset SL(V)$, $p = \text{char } F$ and $S(V)^G$ is a polynomial ring. Let $W \subset V$ be a maximal G -submodule. Set $H = \{g \in G \mid (g - 1)(V) \subseteq W\}$. Then one of the following holds:*

- (1) $\dim_F V/W = 1$;
- (2) $\dim_F V/W = 2$, $G/H \cong SL_2(q)$, $p|q$, or $G/H = SL_2(5)$, $p = 3$;
- (3) $n := \dim_F V/W \geq 3$, $G/H \cong SL_n(q)$, $p|q$;
- (4) $p = 2$, $G/H \cong D \rtimes S_n$, where $n := \dim_F V/W > 1$, $D = \{(\xi^{a_1}, \dots, \xi^{a_n}) \mid \sum_{i=1}^n a_i \equiv 0(m)\}$, where ξ is a primitive m -th root of unity.

Proof: By Theorems 1 and 2, $S(M)^{G/H}$ is a polynomial ring, where $A := S(V)^H/(WS(V))^H = F[x_1, \dots, x_n] = S(M)$, $M := Fx_1 + \dots + Fx_n$, $\deg x_1 = \dots = \deg x_n$ and M is a faithful irreducible G/H -module. So we are in the setup analyzed by [12], so the above possibilities are the only ones admitting a polynomial ring of invariants. \square

3. Theorem 2

In this section we shall prove the following (Theorem 2 of the introduction).

Theorem 3.1. *Let F be an algebraically closed field with $\text{char } F = p > 0$. Let $G \subset GL(U)$ be a finite group generated by transvections, where U is an irreducible finite dimensional G -module. Let $A \subset S(U)$ be a graded polynomial subring, having the following properties:*

- (1) $A = F[x_1, \dots, x_n]$ is a polynomial subring, $\dim A = n = \dim_F U$, where $\{x_1, \dots, x_n\}$ are homogenous generators;
- (2) G acts faithfully by graded automorphisms on A .

Then $\deg x_1 = \dots = \deg x_n$ and $Fx_1 + \dots + Fx_n$ is an irreducible G -module.

Remark 3.2. The possibilities for G as defined above are naturally divided into two cases:

- (i) G acts primitively on U , or
- (ii) G acts imprimitively on U .

In case (i) the possibilities for G are listed (following [10, Thm. 2], [27]) in [12, Thm. 1.5] and are as follows:

- (a) $G = SL_n(q)$, $Sp_n(q)$, $SU_n(q)$, with $(n, q) \neq (3, 2)$, $p|q$;
- (b) $n \geq 4$ is even, $p = 2$, $G = SO_n^\pm(q)$ with q even;
- (c) $n \geq 6$ is even, $p = 2$, $G = S_{n+1}$ or $G = S_{n+2}$;
- (d) $(n, p) = (2, 3)$, $G = SL_2(5)$;
- (e) $(n, p) = (3, 2)$, $G = 3 \cdot A_6$,
- (f) $(n, p) = (6, 2)$, $G = 3 \cdot U_4(3) \cdot 2$.

In case (ii) [27, 1.8, 1.9] imply that $p = 2$ and G is a monomial subgroup.

We shall deal with cases (i) and (ii) separately.

It is of interest to remark that the assumption $p \geq 3$ simplifies the proof considerably as follows:

- if case (i) holds, we merely need to handle cases (a) and (d);
- Case (ii) does not occur.

The proof of Theorem 2 in case (i)

We assume that G acts primitively on U .

Suppose we have, by negation, $\deg x_1 = \dots = \deg_{x_{r_1}} = a_1 < \deg x_{r_1+1} = \dots = \deg x_{r_2} = a_2 < \dots < \deg x_{r_{k-1}+1} = \dots = \deg x_{r_k} = a_k$, $r_k = n$.

Set $M_1 := Fx_1 + \dots + Fx_{r_1}$, $M_i := (B_i + Fx_{r_{i-1}+1} + \dots + Fx_{r_i})/B_i$, where $B_i = F[x_1, \dots, x_{r_{i-1}}]_{a_i}$, the homogenous subspace of degree a_i , $i = 2, \dots, k$. Clearly M_i is a G -module, with $\dim_F M_i < n$ for each i . In fact $\dim_F M_i = r_i - r_{i-1}$, where $r_0 = 0$, and $\sum_{i=1}^k \dim_F M_i = n$.

Let $0 = X_{0i} \subset X_{1i} \subset \cdots \subset X_{ji} \subset \cdots \subset X_{li} = M_i$, be a G -submodule decomposition series of $M_i \forall i$. That is $X_{ji}/X_{j-1,i}$ is an irreducible G -module $\forall i, j$. Therefore $\dim_F X_{ji}/X_{j-1,i} < n$, $\forall i, j$.

In each of the possible cases for G , we find, in Corollaries 3.5, 3.7, 3.9, 3.13 and Lemma 3.11, a subgroup $L \subseteq G$ with the following properties:

- (I) $X_{ji}/X_{j-1,i}$ is a trivial L -module, $\forall i, j$;
- (II) L is not a p -group.

Consequently $\exists t$ such that $(g-1)^{p^t}(M_i) = 0$, $i = 1, \dots, k$, $\forall g \in L$. Therefore $\exists s$, such that $(g-1)^{p^s}(x_j) = 0$, $\forall j = 1, \dots, n$, $\forall g \in L$. This shows by assumption (4) (the faithful action of G on A) that L is a p -group, a contradiction. Consequently $\deg x_1 = \cdots = \deg x_n$.

A similar argument shows that $M_1 = Fx_1 + \cdots + Fx_n$ is an irreducible G -module.

This ends the proof of case (i), in the subcases (a), (b), (c) and (f).

Case (d) is handled in Lemma 3.14 and case (e) in Corollary 3.16. \square

We recall (e.g [13, Definition (5.3.1)]) the following.

Definition 3.3. Let G be a finite group and F a field. We define:

$$R_F(G) = \min\{m \mid \exists \text{ injective homomorphism } \varphi : G \rightarrow PGL_m(F)\},$$

$$R_p(G) = \min\{R_F(G) \mid F \text{ is a field, } \text{char } F = p\},$$

$$R_{p'}(G) = \min\{R_s(G) \mid s \text{ prime, } s \neq p\}.$$

Lemma 3.4. *Let F be an algebraically closed field, with $\text{char } F = p > 0$. Let H be a finite group with $H/Z(H)$ being simple non-abelian. Assume that $(|Z(H)|, p) = 1$ and p divides $|H|$. Let M be a non-trivial irreducible FH -module. Then one of the following holds:*

- (1) $\dim_F M \geq R_p(H/Z(H))$;
- (2) M is a trivial H_p -module, where $H_p = \langle P \mid P \text{ is a } p\text{-sylov subgroup of } H \rangle$.

Proof: Set $\varphi : H \rightarrow GL(M)$, the natural homomorphism and $N := \ker \varphi$.

Suppose firstly that $N \not\subseteq Z(H)$. Then $NZ(H)/Z(H)$ is a non-trivial normal subgroup of $H/Z(H)$, so $NZ(H) = H$. Since by assumption $N \neq H$, it follows that $1 \neq H/N = NZ(H)/N \cong Z(H)/N \cap Z(H)$, which is by assumption, a group of order prime to p . Therefore $H_p N/N = N/N$ and $H_p \subseteq N$ as claimed.

Suppose now that $N \subseteq Z(H)$. Assume firstly that $\varphi(H) \subseteq Z(GL(M)) = F1_M$. H_p is generated by elements of p -power order $\{g_1, \dots, g_r\}$. Hence $\varphi(g_i)^{p^{e_i}} = 1_M$. But $\varphi(g_i) = \lambda_i 1_M$, $\lambda_i \in F$, $i = 1, \dots, r$. Hence $\lambda_i^{p^{e_i}} = 1$, and $\lambda_i = 1$, $i = 1, \dots, r$. Hence $\varphi(g_i) = 1_M$, $i = 1, \dots, r$ and $H_p \subseteq N$. as claimed.

Finally assume that $\varphi(H) \not\subseteq Z(GL(M))$. By Schur's Lemma $\varphi(Z(H)) \subseteq F1_M = Z(GL(M))$. Therefore the induced map $\bar{\varphi} : H/Z(H) \rightarrow GL(M)/Z(GL(M)) = PGL(M)$ is a non-trivial injective homomorphism. Consequently $\dim_F M \geq R_p(H/Z(H))$. \square

Corollary 3.5. *Let $F = \bar{F}$ and $\text{char } F = p$. Let $H = SL_n(q)$, $Sp_n(q)$ or $SU_n(q)$, where $p|q$ and $(n, p) \neq (3, 2)$. Let M be an irreducible FH -module. Then either $\dim_F M \geq n$ or M is a trivial $H_p = H$ -module.*

Proof: We have (respectively) $|Z(H)| = (n, q - 1)$, $(2, q - 1)$ or $(n, q + 1)$. Hence $(p, |Z(H)|) = 1$. So all requirements of Lemma 3.4 are in place. Moreover H is generated by elements of p -order (they are transvections in the natural representation of H). Consequently $H = H_p$. Also $R_p(H/Z(H)) = n$, where the last equality is by [13, Tab. 5.4.C]. \square

Definition 3.6. The complex reflection Mitchell group G_{34} has the structure $G_{34} = 6.PSU_4(3).2$ (Wikipedia). In fact $Z(G_{34}) = \langle h \rangle$, where $h = \text{diag}(\xi, \xi, \xi, \xi, \xi, \xi)$, $\xi^6 = 1$, $\xi \in \mathbb{C}$ (written in its 6-dimensional natural representation). $W_2(G_{34})$ denotes the mod 2 reduction of G_{34} . That is $G_{34}/\langle \text{diag}(-1, -1, -1, -1, -1, -1) \rangle = W_2(G_{34})$. If \bar{h} is the image of h in $W_2(G_{34})$, then $\bar{h} = \text{diag}(\bar{\xi}, \bar{\xi}, \bar{\xi}, \bar{\xi}, \bar{\xi}, \bar{\xi})$, $\bar{\xi}^3 = 1$, $\bar{\xi} \in F(\text{char } F = 2)$. Moreover $Z(W_2(G_{34})) = \langle \bar{h} \rangle$ and $W_2(G_{34})$ has the structure $W_2(G_{34}) = 3.U_4(3).2$.

Let $\pi : W_2(G_{34}) \rightarrow W_2(G_{34})/\langle \bar{h} \rangle$ be the natural projection. Set $H := \pi^{-1}(U_4(3))$. We clearly have $H/\langle \bar{h} \rangle \cong U_4(3)$, a simple group, implying that $Z(H) = \langle \bar{h} \rangle = Z(W_2(G_{34}))$. In particular $(|Z(H)|, 2) = 1$, but $2||H|$, and H_2 is not a 2-group.

Corollary 3.7. *Let $G = W_2(G_{34})$ and $H \subset G$ as in Definition 3.6. Let M be a non-trivial irreducible FH -module, where $F = \bar{F}$ and $\text{char } F = 2$. Then either $\dim_F M \geq 6$ or M is a trivial H_2 -module.*

Proof: By Definition 3.6 all the requirements of Lemma 3.4 are valid for H and $H/Z(H) \cong U_4(3)$. Now by [13, Thm 5.3.9, Tab. 5.3.A] we have $R_2(U_4(3)) \geq R_{3'}(U_4(3)) = 6$. \square

Definition 3.8. Let $G = SO_n^+(q)$, $n \geq 6$ is even, q is even, $\text{char } F = 2$, or $G = SO_n^-(q)$, $n \geq 4$ is even, q is even, $\text{char } F = 2$. Set $H := \Omega_n^\pm(q)$ (with the above restriction on n).

Then we have : $H/Z(H) = P\Omega_n^\pm(q)$ is a simple group (e.g. [13, Tab. 5.1.A]). Moreover $Z(H) \subseteq M(H/Z(H))$, the Schur multiplier of $H/Z(H)$, and $|M(H/Z(H))| = (4, q^{\frac{n}{2}} - 1), (4, q^{\frac{n}{2}} + 1)$ (respectively).

Corollary 3.9. *Let G, H be as in Definition 3.8. Let M be a non-trivial irreducible FH -module, where $\text{char } F = 2, F = \bar{F}$. Then either $\dim_F M \geq n$ or M is a trivial H_2 -module.*

Proof: By Definition 3.8 all the requirements of Lemma 3.4 are valid for H and $H/Z(H) \cong P\Omega_n^\pm(q)$ is a simple group. Now by [13, Prop. 5.4.13] we have $R_2(P\Omega_n^\pm(q)) = n$. \square

The missing cases: $SO_4^+(q)$ where q is even, will be treated next. One difficulty is that $\Omega_4^+(q)$ is not a simple group.

Note 3.10. : The case of $SO_4^+(q)$, q is even.

Using [13, (2.5.11), (2.5.12), Section 2.5, Description 4, Prop. 2.9.1(iv), Prop. 2.9.3] and the notations there, we have: $I = O_4^+(q) = SO_4^+(q) = S, \Omega = \Omega_4^+(q), S = \Omega \langle g \rangle$, where g is any reflection.

In fact $\Omega.2 = S$, where in case of $q > 2$, Ω is the unique subgroup of index 2. Actually $\Omega = S'$. In this case we also have that: $\Omega_4^+(q) = L_1L_2, L_i \triangleleft \Omega_4^+(q), i = 1, 2, L_1 \cap L_2 = 1, L_i \cong SL_2(q), i = 1, 2$ (and the latter is a simple group since $q > 2$). Moreover $gL_1g^{-1} = L_2, gL_2g^{-1} = L_1$.

In case of $q = 2$, Ω is not the only index 2 subgroup of S , but we still have $\Omega_4^+(2) = L_1L_2, L_i \triangleleft \Omega_4^+(2), i = 1, 2, L_1 \cap L_2 = 1, L_i \cong SL_2(2) \cong S_3, i = 1, 2$. Again $gL_1g^{-1} = L_2, gL_2g^{-1} = L_1$. Let $N_i :=$ the unique normal subgroup of $L_i, i = 1, 2$. So N_i is cyclic of order 3, $i = 1, 2, |N_1N_2| = 9$, and $gN_1g^{-1} = N_2, gN_2g^{-1} = N_1$.

Lemma 3.11. *Let $F = \bar{F}$, $\text{char } F = 2$ and $S = SO_4^+(q)$, where q is even. Let M be a non-trivial irreducible FS -module. Then either:*

- (1) $\dim_F M \geq 4$, or
- (2) M is a trivial Ω -module in case $q > 2$, or a trivial N_1N_2 -module in case $q = 2$.

Proof: Suppose by negation that $\dim_F M \leq 3$. Assume firstly that M is a faithful S -module. Recall that S is generated by elements of order 2. Since $\dim_F M \leq 3$ and $\text{char } F = 2$, it follows that each such element is a transvection on M .

If S acts primitively on M , the pair (M, S) should appear in the list of [12, Thm. 1.5], but it does not. If S acts imprimitively on M , this implies that $S \cong D \rtimes S_3$ (if $\dim_F M = 3$)

and $D \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$, m is odd. If $\dim_F M = 2$, $S \cong D \rtimes S_2$, where $D \cong \mathbb{Z}/m\mathbb{Z}$, m is odd. Both possibilities are inconsistent with the structure of $SO_4^+(q)$, q even. Therefore $1 \neq N := \ker(S \rightarrow SL(M))$.

If $N \cap \Omega = 1$, then $N \times \Omega \approx S$, so $|N| = 2$. Hence N is central in S in contradiction to [13, Tab. 2.1.4]. Therefore $1 \neq K := N \cap \Omega$.

If $K \cap L_1 = 1$, then K commutes with L_1 (elementwise), so $K \subseteq L_2$. Therefore $N_2 \subseteq K$ (if $q = 2$), or $L_2 = K$ (if $q > 2$). Consequently $N_1 = gN_2g^{-1} \subseteq gKg^{-1} \subseteq gNg^{-1} = N \Rightarrow N_1 \subseteq N \cap \Omega = K$, if $q = 2$. So $N_1N_2 \subset N$ in this case. Or if $q > 2$ then $L_1 = gL_2g^{-1} \subseteq gKg^{-1} \subseteq gNg^{-1} = N$, so $L_1 \subseteq N \cap \Omega = K$, so $\Omega = L_1L_2 \subseteq N$ in this case. A similar argument applies if $K \cap L_1 \neq 1$. \square

In order to deal with case (c), that is $n \geq 6$ and is even, $p = 2$, $G = S_{n+1}$ or S_{n+2} , we need the following result of A. Wagner.

Theorem 3.12. [24] *S_m , with $m > 6$ have a unique faithful modular 2 representation of least degree, this degree being $m - 1$, or $m - 2$ according as m is odd or even.*

Corollary 3.13. *Let G be as in case (c). Suppose M is a non-trivial irreducible FG -module. Then either:*

- (1) $\dim_F M \geq n$, or
- (2) M is a trivial A_{n+1} -module, if $G = S_{n+1}$ (or a trivial A_{n+2} -module, if $G = S_{n+2}$).

Proof: Suppose $\dim_F M < n$. Then, by Thm. 3.12, M is an unfaithful G -module. Let $N := \ker(G \rightarrow GL(M))$. Then N is normal in G and the result follows. \square

Lemma 3.14. *Let $(n, p) = (2, 3)$, $G = SL_2(5)$ be as in case (d). Then $M_1 = Fx_1 + Fx_2$ is an irreducible G -module and $\deg x_1 = \deg x_2$, where the notations are as in the proof of Thm. 3.1.*

Proof: If $a_1 = \deg x_1 < \deg x_2 = a_2$, then $M_1 = Fx_1$, $B_2 = F[x_1]_{a_2}$, $M_2 = B_2 + Fx_2/B_2$. So M_1, M_2 are trivial G -modules (since G is generated by elements of order 3). Therefore $(g - 1)(M_i) = 0$, $i = 1, 2$, implying that $(g - 1)^2(x_j) = 0$, $j = 1, 2 \forall g \in G$. Consequently $(g^3 - 1)(x_j) = (g - 1)^3(x_j) = 0$, $j = 1, 2, \forall g \in G$. This shows, by assumption (4), that $g^3 = 1, \forall g \in G$, so G is a 3-groups, an obvious absurd. So $M_1 = Fx_1 + Fx_2$, and a similar argument ensures that M_1 is irreducible G -module. This settles Thm 3.1 in case (d). \square

Lemma 3.15. *Let $G = 3.A_6$, $(n, p) = (3, 2)$ be as in case (e). Let M be a non-trivial FG -module with $\dim_F M = 2$, where $\text{char } F = 2$. Then M is a reducible G -module.*

Proof: Assume by negation that M is irreducible. Suppose firstly that M is unfaithful. Let $N := \ker(G \rightarrow GL(M))$. Let $\langle h \rangle$ be the normal subgroup of G satisfying $|\langle h \rangle| = 3$ and $G/\langle h \rangle \cong A_6$. If $N \cap \langle h \rangle = 1$, then $N \cong N\langle h \rangle / \langle h \rangle$ is a non-trivial normal subgroup of $G/\langle h \rangle \cong A_6$. Hence $N\langle h \rangle = G$ and $N \cong A_6$. But G is generated by order 2 elements, so the same holds for $G/N \cong \langle h \rangle$, in contradiction to $|\langle h \rangle| = 3$. Therefore $\langle h \rangle \subseteq N$.

If $\langle h \rangle = N$, then M is a faithful $G/\langle h \rangle$ -module. But $G/\langle h \rangle \cong A_6$. So by [13, Prop. 5.3.7(ii)] $R_2(A_6) = 3$, a contradiction.

If $\langle h \rangle \subset N$ (properly), then $N/\langle h \rangle$ is a non-trivial normal subgroup of $G/\langle h \rangle \cong A_6$, implying that $N = G$, so M is a trivial G -module, violating the assumptions.

All in all M is a 2-dimensional faithful G -module. Recall that G is generated by elements of order 2. Since $\dim_F M = 2$ and $\text{char } F = 2$, each such element is a transvection on M . If G acts primitively on M , then the pair (M, G) would have appeared in the list of [12, Thm. 1.5], but it does not.

So we must conclude that G acts imprimitively on M . Therefore (since $G \subset SL(M)$) we get that $G \cong D \rtimes S_2$, where D is abelian, so G is solvable, a contradiction. \square

Corollary 3.16. *Let $G = 3.A_6$, $(n, p) = (3, 2)$ be as in case (e). Then Theorem 3.1 holds.*

Proof: We assume by negation that $\dim_F M_i \leq 2$, $\forall i$. Let $\{X_{ji}\}$ be the decomposition series of M_i for each i . Then by Lemma 3.15 $X_{ji}/X_{j-1,i}$ is a trivial 1-dimensional G -module $\forall i$. Consequently $(g-1)^{2^s}(M_i) = 0$, $\forall i, \forall g \in G$, implying that $(g^{2^t} - 1)(x_j) = (g-1)^{2^t}(x_j) = 0$, $j = 1, 2, 3$, $\forall g \in G$. Therefore by assumption (4) $g^{2^t} = 1$, $\forall g \in G$, and G is a 2-group, an obvious contradiction. Hence $\deg x_1 = \deg x_2 = \deg x_3$ and $M_1 = Fx_1 + Fx_2 + Fx_3$ is an irreducible G -module. \square

The proof of Theorem 2 in case (ii)

So $G \subset GL(U)$ is a irreducible imprimitive group, generated by transvections on U , $\dim_F U = n$, F is algebraically closed. So $G \subset SL(U)$. By [27, 1.8, 1.9] G is a monomial subgroup and $\text{char } F = 2$. Therefore $G = D \rtimes S_n$, where $D = \{\text{diag}(\xi^{a_1}, \dots, \xi^{a_n}) \mid \sum_{i=1}^n a_i \equiv 0 \pmod{m}\}$, and $|\langle \xi \rangle| = m$, $\xi \in F$, is a primitive m -th root of unity. Therefore m is an odd integer.

Note 3.17. (1) $G \neq S_n$, since U is an irreducible G -module.

(2) $D \cong \langle \xi \rangle \times \cdots \times \langle \xi \rangle$, ($(n-1)$ times), so $|D| = m^{n-1}$.

Lemma 3.18. Let M be an FG -module with $\dim_F M = n-i$. Then there exists a subgroup $K \subseteq D$ having the following properties:

(1) $|K| \geq m^i$;

(2) M is a trivial K -module.

Proof: Let $\varphi : G \rightarrow GL(M)$ be the natural homomorphism. G is generated by order 2 elements, implying that $\varphi(G) \subseteq SL(M)$.

Now $g^m = 1$, $\forall g \in D$, and $(m, 2) = 1$, imply that $\varphi(D)$ is a commutative group of semi-simple transformations on M , and is therefore simultaneously diagonalizable (F is algebraically closed). Therefore after a base change of M we have:

$$\varphi(D) \subseteq \{\text{diag}(\lambda_1, \dots, \lambda_{n-i}) \mid \lambda_j \in F, j = 1, \dots, n-i\}.$$

So $\det(\varphi(D)) = 1$ implies that $\prod_{j=1}^{n-i} \lambda_j = 1$. Also $g^m = 1$ implies that $\lambda_j^m = 1$, so $\lambda_j = \xi^{a_j}$, where $\xi \in F$ is the m -th primitive root of unity, $j = 1, \dots, n-i$.

Consequently:

$$\varphi(D) \subseteq \{\text{diag}(\xi^{a_1}, \dots, \xi^{a_{n-i}}) \mid \sum_{j=1}^{n-i} a_j \equiv 0 \pmod{m}\} \cong \langle \xi \rangle \times \cdots \times \langle \xi \rangle, (n-i-1)\text{-times}.$$

Hence $|\varphi(D)| \leq m^{n-i-1}$. Therefore $|K| \geq m^i$, where $K := \ker(\varphi|_D : D \rightarrow GL(M))$. \square

Corollary 3.19. Suppose one of the following holds:

(1) $k = 2$, that is $\dim_F M_1 + \dim_F M_2 = n$;

(2) $\dim_F M_i = n-l$, for some i , $\dim M_j = 1$, $\forall j \neq i$.

Then M_s is a trivial K -module $\forall s$, where $K \subseteq D$ is a subgroup with $K \neq 1$.

Proof: Set $K_i := \ker(\varphi_i : D \rightarrow GL(M_i)) \forall i$. If (1) holds then by Lemma 3.18 $|K_i| \geq m^{n-\dim_F M_i}$, $i = 1, 2$. Hence $|K_1||K_2| \geq m^{2n-n} = m^n > m^{n-1} = |D|$. Therefore $K := K_1 \cap K_2 \neq 1$, will do.

Suppose (2) holds. G is generated by order 2 elements, so M_j is a trivial G -module $\forall j \neq i$. Now by Lemma 3.18 $|K_i| \geq m^l$, so $K := K_i$ acts trivially on $M_s \forall s$. \square

The proof of Theorem 2, $G = D \rtimes S_n$, n is even, $n > 6$

Recall that S_n is a subgroup of G . As before, we consider the sequence of G -modules M_i , $i = 1, \dots, k$. If $\dim_F M_i \leq n - 3$ then by Theorem 3.12, M_i is an unfaithful S_n -module. Consequently M_i is a trivial A_n -module $\forall i$. This implies as before that $(g - 1)^{2^s}(x_j) = 0$, $\forall g \in A_n$, $j = 1, \dots, n$, for some fixed s . Therefore by the faithful action assumption of G on $A = F[x_1, \dots, x_n]$, we have $g^{2^s} = 1 \forall g \in A_n$. So A_n is a 2-group, an obvious contradiction.

If $\dim_F M_i = n - 2$, and $\dim_F M_j = 2$, for some j . Then by Corollary 3.19(1) $\{M_i, M_j\}$ are trivial K -modules for some $K \subseteq D$, $K \neq 1$.

If $\dim_F M_i = n - 2$, $\dim_F M_j = \dim_F M_l = 1$, where $j \neq l$. Then by Corollary 3.19(2) $\{M_i, M_j, M_l\}$ are trivial K -modules, for some $K \subseteq D$, $K \neq 1$.

If $\dim_F M_i = n - 1$, $\dim_F M_j = 1$, then again by Corollary 3.19(1) $\{M_i, M_j\}$ are trivial K -modules for some $K \subseteq D$, $K \neq 1$.

So in all of these cases $(g - 1)^{2^i}(x_j) = 0$, $j = 1, \dots, n$, $\forall g \in K$. So K is a 2-group, in contradiction to $|K|$ being odd. So $\deg x_1 = \dots = \deg x_n$, and $M_1 = Fx_1 + \dots + Fx_n$ is irreducible G -module. \square

The case of $G = D \rtimes S_4$ (that is $n = 4$)

By negation, we need to consider the following possibilities:

- (1) $\dim_F M_i = 3$, $\dim_F M_j = 1$;
- (2) $\dim_F M_i = 2$, $\dim_F M_j = 2$, $i \neq j$;
- (3) $\dim_F M_i = 2$, $\dim_F M_j = \dim_F M_l = 1$, $j \neq l$;
- (4) $\dim_F M_i = 1$, $i = 1, 2, 3, 4$.

In all cases we are led by Corollary 3.19 to the existence of $K \subseteq D$, $K \neq 1$ and M_l is a trivial K -module $\forall l$. This implies as before that K is a 2-group, a contradiction.

The case of $G = D \rtimes S_2$ (that is $n=2$)

We either have $\dim_F M_1 = 2$ and so $\deg x_1 = \deg x_2$, or $\dim_F M_1 = \dim_F M_2 = 1$ and we argue as before. \square

The case of $G = D \rtimes S_6$ (that is $n=6$)

By negation, we need to consider the following possibilities:

- (1) $\dim_F M_i = 5$, $\dim_F M_j = 1$;
- (2) $\dim_F M_i = 4$, $\dim_F M_j = 2$;
- (3) $\dim_F M_i = 4$, $\dim_F M_j = \dim_F M_r = 1$, $j \neq r$;

- (4) $\dim_F M_i = 3, \dim_F M_j = 3, i \neq j$;
- (5) $\dim_F M_i = 3, \dim_F M_j = 2, \dim_F M_r = 1$;
- (6) $\dim_F M_i = 3, \dim_F M_j = \dim_F M_r = \dim_F M_l = 1$, for 3 distinct indices j, r, l ;
- (7) $\dim_F M_i = \dim_F M_j = \dim_F M_r = 2$, for 3 distinct indices i, j, r ;
- (8) $\dim_F M_i = \dim_F M_j = 2, i \neq j, \dim_F M_r = \dim_F M_l = 1, r \neq l$;
- (9) $\dim_F M_i = 2, \dim_F M_j = \dim_F M_t = \dim_F M_l = \dim_F M_r = 1$, for 4 distinct indices j, t, l, r ;
- (10) $\dim_F M_i = 1, \forall i = 1, \dots, 6$.

The only cases which are not directly handled by Corollary 3.19 are (5), (7) and (8).

In case (5) we have by Lemma 3.18, that $|K_i| \geq m^3, |K_j| \geq m^4$, hence since $|D| = m^5$, $|K_i \cap K_j| = \frac{|K_i||K_j|}{|K_i K_j|} \geq \frac{|K_i||K_j|}{|D|} \geq \frac{m^7}{m^5} = m^2$. Since M_r is a trivial G -module, it follows that $\{M_i, M_j, M_r\}$ are trivial $K_i \cap K_j$ -modules. So the rest of the argument follows as before.

In case (7) $|K_i| \geq m^4, |K_j| \geq m^4, |K_r| \geq m^4$. But $|K_i K_j||K_i \cap K_j| = |K_i||K_j|$, so $|K_i \cap K_j| = \frac{|K_i||K_j|}{|K_i K_j|} \geq \frac{|K_i||K_j|}{|D|} \geq \frac{m^8}{m^5} = m^3$. Consequently $|(K_i \cap K_j) \cap K_r| = \frac{|K_i \cap K_j||K_r|}{|(K_i \cap K_j)K_r|} \geq \frac{m^3 \cdot m^4}{|D|} = m^2$. Therefore $\{M_i, M_j, M_r\}$ are trivial $(K_i \cap K_j \cap K_r)$ -modules. So the rest follows as before.

In case (8), $|K_i| \geq m^4, |K_j| \geq m^4$, so as above $|K_i \cap K_j| \geq m^3$. Now $\{M_r, M_l\}$ are trivial G -modules. Hence $\{M_i, M_j, M_r, M_l\}$ are trivial $K_i \cap K_j$ -modules, so we argue as before. \square

The proof of Theorem 2, $G = D \rtimes S_n$, n is odd, $n \geq 6$

As before we consider the sequence of G -modules $M_i, i = 1, \dots, k$.

If $\dim_F M_i \leq n - 2, \forall i$, then by Theorem 3.12, M_i is an unfaithful S_n -module $\forall i$. This implies that M_i is a trivial A_n -module for each i . Consequently $(g-1)^{2^s}(x_j) = 0, j = 1, \dots, n, \forall g \in A_n$, implying that A_n is a 2-group, an obvious contradiction.

If $\dim_F M_i = n - 1, \dim_F M_j = 1$, then by Corollary 3.9 $\{M_i, M_j\}$ are trivial K -modules, for $1 \neq K \subset D$, a subgroup of D . Consequently $(g-1)^{2^s}(x_j) = 0, j = 1, \dots, n, \forall g \in K$, implying that K is a 2-group, a contradiction.

All in all, $\deg x_1 = \dots = \deg x_n, M_1 = Fx_1 + \dots + Fx_n$ is an irreducible G -module. \square

The case of $G = D \rtimes S_5$ (that is $n = 5$)

By negation, we need to consider the following possibilities:

- (1) $\dim_F M_i = 4, \dim_F M_j = 1$;

- (2) $\dim_F M_i = 3, \dim_F M_j = 2$;
- (3) $\dim_F M_i = 3, \dim_F M_j = 1 = \dim_F M_l = 1, j \neq l$;
- (4) $\dim_F M_i = \dim_F M_j = 2, i \neq j, \dim_F M_l = 1$;
- (5) $\dim_F M_i = 2, \dim_F M_j = 1, \forall j \neq i$;
- (6) $\dim_F M_i = 1, i = 1, \dots, 5$.

The only case not covered by Corollary 3.19 is (4). Here we get, by Lemma 3.18, $|K_i| \geq m^3, |K_j| \geq m^3$, hence $|K_i \cap K_j| = \frac{|K_i||K_j|}{|K_i K_j|} \geq \frac{m^3 m^3}{|D|} = \frac{m^6}{m^4} = m^2$. Since G acts trivially on M_l , it follows that $\{M_i, M_j, M_l\}$ are trivial $K_i \cap K_j$ -modules. Therefore $(g-1)^4(x_s) = 0, s = 1, \dots, 5, \forall g \in K_i \cap K_j$. Hence $K_i \cap K_j$ is a 2-group. A contradiction. The conclusion follows as before. \square

The case of $G = D \rtimes S_3$, (that is $n=3$)

By negation, we have to consider the following possibilities:

- (1) $\dim_F M_i = 2, \dim_F M_j = 1$;
- (2) $\dim_F M_i = 1, i = 1, 2, 3$.

Both cases are handled by Corollary 3.19, leading to a contradiction. \square

4. Isolated quotient singularities in prime characteristic

We assume throughout the present section that $G \subset SL(V)$ is a finite group, $\dim_F V$ is finite, F is a field with $\text{char } F = p > 0$.

We recall the following.

Theorem 4.1. [26, Theorem 6.1.11], [22, Theorem 6.3.1]. *There exists a list of $\{(W, H)\}$, where $\dim_{\mathbb{C}} W$ is finite, $H \subset GL(W)$ is a finite group, such that each isolated quotient singularities over \mathbb{C} is isomorphic to one of $S(W)^H$.*

We shall refer to the above list as the Zassenhaus-Vincent-Wolf list.

The main result in this section is the following.

Theorem B. *Suppose $S(V)^G$ is an isolated singularity and F is algebraically closed. Then $\widehat{S(V)^G} \cong \widehat{S(W)^H}$, where $(p, |H|) = 1, \dim_F W = \dim_F V$ and (W, H) is a mod p reduction of a direct sum of members in the Zassenhaus-Vincent-Wolf list.*

Proposition 4.2. *Suppose $G \subset SL(V)$, F is perfect, $\text{char } F = p > 0$ and $S(V)^G$ is an isolated singularity. Then:*

- (1) $S(V)^{T(G)}$ is a polynomial ring;
- (2) $T(G) = \langle P | P \text{ is a } p\text{-sylov subgroup of } G \rangle$, and consequently $(|G/T(G)|, p) = 1$.

Here $T(G)$ is the subgroup generated by all transvections on V (or V^*).

Proof: Let $0 \neq f \in V^*$ and $U = Ff$. By Corollary 2.9 $S(V)^{G_U}$ is a polynomial ring, hence G_U is generated by transvections implying $G_U \subseteq T(G)$. Hence $G_U = T(G)_U$. Therefore by Theorem A $S(V)^{T(G)}$ is a polynomial ring.

Moreover, let P be a p -sylov subgroup of G , then by a classical result $\exists f \in V^*$, $f \neq 0$, with $P \subseteq G_U$, $U := Ff$. Since $G_U = T(G)_U$, it follows that $P \subseteq T(G)$. Hence $\langle P | P \text{ is a } p\text{-sylov subgroup of } G \rangle \subseteq T(G)$.

The converse inclusion is trivial since every transvection is of order p . So (2) is verified. \square

Proposition 4.3. *Let $G \subset SL(V)$ be a finite group and F is perfect. Suppose $S(V)^G$ is an isolated singularity. Then:*

- (1) $S(V)^G$ has rational singularity;
- (2) $S(V)^G$ has a non-commutative crepant resolution if it is Gorenstein.

Proof: By Proposition 4.2 $S(V)^T$ is a polynomial ring and $(|G/T(G)|, p) = 1$. Therefore $S(V)^G = (S(V)^T)^{G/T(G)}$ is a direct summand of $S(V)^T$ (using $\frac{1}{r} \sum_{h \in G/T(G)} h$, as the Reynolds operator, where $r = |G/T(G)|$). Consequently by [9, Thm. 2.1] $S(V)^G$ is F -regular. In particular it is F -rational. Consequently by [14, Cor. 1.10] $S(V)^G$ has rational singularity. This settles item (1).

We now show that the skew group ring $S(V)^{T(G)} * G/T(G)$ is a non-commutative crepant resolution (if $S(V)^G$ is Gorenstein). We firstly recall that the extension $S(V)^G \subset S(V)^{T(G)}$ is unramified in codimension 1 [19, Lemma p.05]. Consequently as in [3, Cor. 2.24(2)], using $S(V)^G = (S(V)^{T(G)})^{G/T(G)}$, we have $S(V)^{T(G)} * G/T(G) \cong \text{End}_{S(V)^G} S(V)^{T(G)}$. Since $(|G/T(G)|, p) = 1$ it follows that $gl. \dim[S(V)^{T(G)} * G/T(G)]$ is finite [16, Thm. 5.6]. Now being a Cohen-Macaulay $S(V)^G$ -module (since $S(V)^T$ is such), it follows that $S(V)^{T(G)} * G/T(G)$ is a homologically homogenous PI ring. The reflexivity of $S(V)^T$ as a $S(V)^G$ -module follows from the normality of $S(V)^T$. This settles item (2). \square

The next property will be needed in the sequel.

Definition 4.4. We say that V is void of fixed points if $V^g = 0$, $\forall g \in G$, $g \neq 1$.

Remark 4.5. V is void of fixed points if and only if V^* is void of fixed points. This is so since if $V^g \neq 0$, for some $g \in G$, then $(V^*)^{g^{-1}} \neq 0$. To verify this, observe that if A is the matrix of g on V with respect to a basis $\{v_1, \dots, v_n\}$ of V , then A^t is the matrix of g^{-1} on V^* with respect to the dual basis $\{f_1, \dots, f_n\}$. Then $v := \sum_{i=1}^n \xi_i v_i \in V^g$ implies $\sum_{i=1}^n \xi_i f_i \in (V^*)^{g^{-1}}$.

The following result appears in [22, Lemma 2.4] with the additional assumption $(|G|, p) = 1$.

Lemma 4.6. *Suppose F is perfect and G is free from pseudo-reflections. Then the following are equivalent:*

- (1) $S(V)^G$ is an isolated singularity;
- (2) V is void of fixed points.

Proof: By Corollary 2.9, (1) is equivalent to: $S(V)^{G_U}$ is a polynomial ring for each $U \subset V^*$, $\dim_F U = 1$. Suppose (1) holds. If $V^g \neq 0$, for some $g \in G$, $g \neq 1$, then by Remark 4.5 $(V^*)^{g^{-1}} \neq 0$, hence $\exists U \subset V^*$, $\dim_F U = 1$ with $g^{-1} \in G_U$, so $G_U \neq 1$. Now $S(V)^{G_U}$ is a polynomial ring, so G_U is generated by pseudo-reflection, in contradiction to the assumption. Conversely if (2) holds on V , then (2) holds on V^* . Hence $G_U = 1$, for each $U \subset V^*$, implying that $S(V)^{G_U} = S(V)$ is a polynomial ring, hence (1) holds. \square

The following is well known.

Lemma 4.7. *Let (A, \mathfrak{m}) be a local ring which is the localization of a finitely generated algebra over a field. Then the following are equivalent:*

- (1) A satisfies R_i (respectively S_i);
- (2) \hat{A} , the \mathfrak{m} -adic completion, satisfies R_i (respectively S_i).

Consequently A is an isolated singularity (respectively normal) if and only if \hat{A} is such.

Proof: The implication (2) \Rightarrow (1) is valid for every Noetherian local ring as follows from [15, Thm. 23.9(i)].

To prove the converse direction, recall that A is a G -ring [15, Cor.p.259]. Hence the map $A \rightarrow \hat{A}$ is regular [15, p.256]. Consequently for every prime $\mathfrak{p} \in \text{spec } A$ the fiber ring $(\hat{A})_{\mathfrak{p}}/\mathfrak{p}(\hat{A})_{\mathfrak{p}}$ is geometrically regular. Hence $(\hat{A})_{\mathfrak{p}}/\mathfrak{p}(\hat{A})_{\mathfrak{p}}$ satisfies R_i and S_i . Hence by [15, Thm. 23.9(ii),(iii)] \hat{A} satisfies R_i (respectively S_i). \square

We shall need the following version of H. Cartan's theorem [22, Lemma 2.3]:

Theorem 4.8. *Let $F[[x_1, \dots, x_n]]$ be a power series ring and $H \subseteq \text{Aut } F[[x_1, \dots, x_n]]$ is a finite group with $(|H|, p) = 1$. Suppose $\mathfrak{m} := (x_1, \dots, x_n)$ is H -stable. Then:*

- (1) *There are parameters $\{y_1, \dots, y_n\}$ such that $F[[x_1, \dots, x_n]] = F[[y_1, \dots, y_n]]$ and H acts linearly on $W := Fy_1 + \dots + Fy_n$;*
- (2) *$F[[x_1, \dots, x_n]]^H \cong \widehat{S(W)^H}$.*

Proof: For (1) we refer to [22, Lemma 2.3]. Now the inclusion $\widehat{S(W)^H} = F[\widehat{y_1, \dots, y_n}]^H \subseteq F[[y_1, \dots, y_n]]^H = F[[x_1, \dots, x_n]]^H$ is clear. Note that by Galois theory $[Q(F[[x_1, \dots, x_n]]) : Q(F[[x_1, \dots, x_n]]^H)] = |H|$. So we only need to affirm that $[Q(\widehat{S(W)}) : Q(\widehat{S(W)^H})] \leq |H|$, since this implies that $Q(\widehat{S(W)^H}) = Q(F[[x_1, \dots, x_n]]^H) = Q(F[[x_1, \dots, x_n]]^H)$, and then the normality of $\widehat{S(W)^H}$ (which follows from the normality of $S(W)^H$) implies that $\widehat{S(W)^H} = F[[x_1, \dots, x_n]]^H$.

Set $\mathfrak{n} = S(W)_+^H$, $\mathfrak{m} = S(W)_+$. Then \mathfrak{m} is the unique maximal above \mathfrak{n} , hence $\widehat{S(W)} = \lim_{\leftarrow i} S(W)/\mathfrak{n}^i S(W)$ implying that $\widehat{S(W)} = S(W) \otimes_{S(W)^H} \widehat{S(W)^H}$. Hence $[Q(\widehat{S(W)}) : Q(\widehat{S(W)^H})] \leq [Q(S(W)) : Q(S(W)^H)] = |H|$. \square

In order to translate the Zassenhaus-Vincent-Wolf list from objects over \mathbb{C} to ones over an algebraically closed field F , with $\text{char } F = p > 0$, we need the following result (well known to finite group theorists). This is also sketched in [22, Theorem 3.13]

Proposition 4.9. *Let G be a finite groups with $(|G|, p) = 1$ and F is algebraically closed. Then:*

- (1) *There exists an isomorphism:*

$$\psi : \{ \text{Irreducible } \mathbb{C}G\text{-modules} \} \rightarrow \{ \text{Irreducible } FG\text{-modules} \};$$

- (2) *$M^g = 0, \forall 1 \neq g \in G$ if and only if $\psi(M)^g = 0, \forall 1 \neq g \in G$.*

Proof: Observe firstly that by [7, Thm. 83.5], since $(|G|, p) = 1$,

$$\begin{aligned} & \text{The number of non-isomorphic irreducible } FG\text{-modules} \\ &= \text{The number of } p\text{-regular conjugacy classes} \\ &= \text{The number of conjugacy classes} \\ &= \text{The number of non-isomorphic irreducible } \mathbb{C}G\text{-modules.} \end{aligned}$$

So the cardinality of both sets appearing in (1) is the same.

Let $K := \mathbb{Q}(\xi)$, ξ is a primitive m -th root of unity, where m is a prime number, $p \neq m \geq \exp(G)$. Then by Brauer's theorem [7, Thm. (41.1)], K is a splitting for G . So $M = M_0 \otimes_K \mathbb{C}$, $M_0 \subset M$, and M_0 is an irreducible KG -module. Let $R := \mathbb{Z}[\xi]$, so R is the ring of integers of K [25, Thm. 2.6]. Let P be a prime ideal in R , with $P \cap \mathbb{Z} = p\mathbb{Z}$. Then there exists a $R_P G$ -module N , $N \subset M_0$, with $N \otimes_{R_P} K = M_0$ (e.g. [7, Thm. (75.2)]).

Now N is an indecomposable $R_P G$ -module. Otherwise $M_0 = N \otimes_{R_P} K$ would have been decomposable as KG -module. Let $\bar{N} := N/PN$. Set $\bar{K} := R_P/P_P = \mathbb{F}_p(\bar{\xi}) \subset F$, $\bar{\xi}$ a primitive m -th root of the unity in F (we identify $\langle \bar{\xi} \rangle$ with the group of m -th roots of unity in F). Then by [7, Cor. (83.7)] \bar{K} is a splitting field for G .

$\bar{K}G$ is a semi-simple \bar{K} -algebra since $(p, |G|) = 1$, hence \bar{N} is a projective $\bar{K}G$ -module. Consequently by [7, Thm. (77.1)] N is a projective $R_P G$ -module, implying that it is a direct summand of a free $R_P G$ -module.

We shall next show that \bar{N} is an irreducible $\bar{K}G$ -module. Let $\hat{R}_P :=$ the P -adic completion of R_P . Since N is an indecomposable $R_P G$ -module, it follows from the proof of [7, Thm. (76.29)] that $\hat{R}_P N$ is an indecomposable $\hat{R}_P G$ -module. Also it is easily seen that $\hat{R}_P N$ is a projective $\hat{R}_P G$ -module, implying that it is a direct summand of a free $\hat{R}_P G$ -module.

Let $\bar{K}G = (\bar{K}G)\epsilon_1 \oplus \cdots \oplus (\bar{K}G)\epsilon_m$ be a decomposition of $\bar{K}G$ into a direct sum of irreducible $\bar{K}G$ modules, where $\{\epsilon_1, \dots, \epsilon_m\}$ are orthogonal idempotents, with $\bar{1} = \epsilon_1 + \cdots + \epsilon_m$. By [7, Thm. (77.11)] there are orthogonal idempotents $\{e_1, \dots, e_m\}$ in $\hat{R}_P G$, with $1 = e_1 + \cdots + e_m$ and $\bar{e}_i = \epsilon_i$, $i = 1, \dots, m$. Therefore $\hat{R}_P G = (\hat{R}_P G)e_1 \oplus \cdots \oplus (\hat{R}_P G)e_m$, and $\overline{(\hat{R}_P G)e_i} := (\hat{R}_P G)e_i/P(\hat{R}_P G)e_i = (\bar{K}G)\epsilon_i$, $i = 1, \dots, m$. Hence $(\hat{R}_P G)e_i$ is an indecomposable projective $\hat{R}_P G$, $\forall i = 1, \dots, m$.

Since $\hat{R}_P N$ is an indecomposable projective $\hat{R}_P G$ -module, which is a direct summand of a free $\hat{R}_P G$ -module, it follows from the Krull-Schmidt theorem for $\hat{R}_P G$ -modules [7, Thm. (76.26)] that $\hat{R}_P N \cong (\hat{R}_P G)e_i$, for some i . Therefore:

$$\bar{N} = N/PN = \hat{R}_P N/P\hat{R}_P N \cong \overline{(\hat{R}_P G)e_i} = (\bar{K}G)\epsilon_i, \text{ is an irreducible } \bar{K}G\text{-module.}$$

Moreover since \bar{K} is a splitting field, it follows that \bar{N} is an absolutely irreducible $\bar{K}G$ -module. So we define: $\psi(M) := \bar{N} \otimes_{\bar{K}} F$. That this map is well defined (e.g. independent of the choice in P) follows from [7, Cor. (82.2)].

We finally need to verify that ψ is 1 – 1 (the onto-ness will follow from the beginning paragraph).

Let T, U be two irreducible representations of KG with characters α, β (respectively). Let $\bar{T} = \bar{U}$ denote the induced $\bar{K}G$ -representations, and τ their common character..

Let χ be the Brauer character associated to τ , e.g. [7, p.589]. Then $\overline{\chi(x)} = \tau(x)$, $\forall x \in G$ ($(p, |G|) = 1$, so every $x \in G$ is p -regular). On the other hand, since α, β are coming from $R_p G$ -representations, it follows by [7, p.589] that $\chi = \alpha$, $\chi = \beta$. Therefore by [7, (30.12), (30.14), (30.15)] T and U are equivalent. This settles item (1).

To prove (2), suppose that $\psi(M)^g \neq 0$, then $\psi(M) = \bar{N} \otimes_K F$ implies that $\bar{N}^g \neq 0$. Let $0 \neq \bar{x} \in \bar{N}^g$, and $x \in N$ is some preimage of \bar{x} in N . Then $z := \frac{1}{r} \sum_{i=0}^{r-1} g^i(x) \in N$, where $r = |g|$. Moreover $g(z) = z$. So $z \in N^G \subseteq M^g$.

We only need to affirm that $z \neq 0$. This follows from $\bar{z} = \overline{\frac{1}{r} \sum_{i=0}^{r-1} g^i(x)} = \frac{1}{r} \sum_{i=0}^{r-1} g^i(\bar{x}) = \bar{x} \neq 0$.

The reverse implication is easier. If $M^g \neq 0$, then $M_0^g \neq 0$. So pick $0 \neq x \in M_0^g$. We can find $x_1 \in N^g$ with $x_1 \notin P_P N$ (say $x = \pi^i x_1$, $x_1 \notin P_P N$, $(\pi) = P_P$). Then $\bar{x}_1 \in \bar{N}^g \subseteq \psi(M)^g$. This settles (2).□

The proof of Theorem B

By Proposition 4.2 $S(V)^{T(G)}$ is a polynomial ring and $(|G/T(G)|, p) = 1$. Set $H := G/T(G)$. By Cartan's theorem (Theorem 4.8) $S(\widehat{V})^{T(G)} = F[[y_1, \dots, y_n]]$ and $G/T(G)$ acts linearly on $W := Fy_1 + \dots + Fy_n$. Consequently $\widehat{S(V)}^G \cong (S(\widehat{V})^{T(G)})^{G/T(G)} = F[[y_1, \dots, y_n]]^{G/T(G)} \cong \widehat{S(W)}^H$. By Lemma 4.7 the isolated quotient singularity property of $S(V)^G$ is translated to $\widehat{S(W)}^H$ and then to $S(W)^H$.

We may assume that H is free from pseudo-reflections on W . Indeed, if the subgroup of H generated by pseudo-reflections $P(H) \neq 1$, then since $(|H|, p) = 1$, $S(W)^{P(H)}$ is a polynomial ring, and again by Cartan's theorem $\widehat{S(V)}^G \cong \widehat{S(W)}^H \cong (S(\widehat{W})^{P(H)})^{H/P(H)} = F[[z_1, \dots, z_n]]^{H/P(H)}$ and on $W_1 := Fz_1 + \dots + Fz_n$, $H_1 := H/P(H)$ acts linearly. Since $|H_1| < |H|$, this process must terminate.

Consequently by Lemma 4.6 W is void of fixed points, that is $W^g = 0$, $\forall g \in H$, $g \neq 1$.

Now since $(|H|, p) = 1$, W is a completely irreducible FH -module, so $W = W_1 \oplus \dots \oplus W_r$, where W_i is an irreducible FH -module and $W_i^g = 0$, $\forall g \in H$, $g \neq 1$, $i = 1, \dots, r$.

Consequently by Proposition 4.9 there exists a $\mathbb{C}H$ -module $M := M_1 \oplus \dots \oplus M_r$, where M_i is an irreducible $\mathbb{C}H$ -module with $M_i^g = 0$, $\forall g \in H$, $g \neq 1$, such that $\psi(M_i) = W_i$, $i = 1, \dots, r$. Now M_i , $i = 1, \dots, r$ is a member of the Zassenhaus-Vincent-Wolf list (with respect to H) since $M_i^g = 0 \forall g \in H$, $g \neq 1$, [26, Thm. 5.3.1, 6.1.11, 6.3.1, 6.3.6].□

We shall now present modular isolated quotient singularity examples in arbitrary dimension (≥ 3). They are always Cohen-Macaulay, but not necessarily Gorenstein.

Example 4.10. Let $V = Fv + Fw_n \cdots + Fw_1$, be a $n + 1$ -dimensional F -vectorspace, with $n \geq 2$, and $a \in F$ is a primitive l -th root of unity. We also require $(n, l) = 1$.

Let $g := \text{diag}(a, \dots, a, a^{-n}, a)$, $h := 1_{n+1} + e_{1, n+1}$.

The action on the basis elements is given by:

$$g(v) = av, g(w_i) = \begin{cases} aw_i, & i \neq 2; \\ a^{-n}w_2, & i = 2. \end{cases}, h(w_i) = w_i, i = 1, \dots, n, h(v) = v + w_1.$$

It is easy to verify that h is a transvection, $g, h \in SL(V)$ and $gh = hg$. Hence $G := \langle g, h \rangle$ is a commutative finite group, $|G| = pl$ and $G = \{g^i h^j \mid 0 \leq i \leq l - 1, 0 \leq j \leq p - 1\}$. Set $H := \langle h \rangle$.

We next verify that $S(V)^G$ is an isolated singularity. For that we show that $S(V)^{G_U}$ is a polynomial ring for each $U \subset V^*$ with $\dim_F U = 1$. Now by [2, Lemma 2.1] $x \in G_U$ if and only if $(x - 1)(V) \subseteq U^\perp$, where $U^\perp = \{y \in V \mid f(y) = 0\}$, $U = Ff$. Clearly $\dim_F U^\perp = \dim_F V - 1 = n$.

Case 1: $w_r \notin U^\perp$, for some $r \in \{1, \dots, n\}$.

Let $x = g^i h^j \in G_U$, $0 \leq i \leq l - 1$, $0 \leq j \leq p - 1$, then $(x - 1)(V) \subseteq U^\perp$, hence:

$$(x - 1)(w_r) = (g^i h^j)(w_r) - w_r = g^i(w_r) - w_r = \begin{cases} (a^i - 1)w_r, & \text{if } r \neq 2; \\ (a^{-in} - 1)w_2, & \text{if } r = 2. \end{cases}.$$

Since $(l, n) = 1$, it follows that $i = 0$ and so $x = h^j \in H$ and $G_U \subseteq H$. Therefore either $G_U = 1$ or $G_U = H$ and in both cases $S(V)^{G_U}$ is a polynomial ring.

Case 2: $U^\perp = Fw_n + \cdots + Fw_1$.

So $v \notin U^\perp$. Similarly if $x = g^i h^j \in G_U$, then $(x - 1)(v) \in U^\perp$. But $(x - 1)(v) = (g^i h^j)(v) - v = g^i(v + jw_1) - v = a^i v + ja^i w_1 - v = (a^i - 1)v + ja^i w_1$. But $w_1 \in U^\perp$, hence $(a^i - 1)v \in U^\perp$, forcing $i = 0$ and $x \in H$. Hence $G_U \subseteq H$ and again $G_U = 1$ or $G_U = H$ and in both cases $S(V)^{G_U}$ is a polynomial ring.

All isolated quotient singularities are Cohen-Macaulay as follows from [11, Thm. 3.1].

Now $S(V)^H = F[v^p - w_1^{p-1}v, w_n, \dots, w_1] = S(M)$, where $M := F(v^p - w_1^{p-1}v) + Fw_n + \cdots + Fw_1$. We compute the action of g^i on M :

$$g^i(v^p - w_1^{p-1}v) = (g^i(v))^p - (g^i(w_1))^{p-1}g^i(v) = (a^i v)^p - (a^i w_1)^{p-1}a^i v = a^{ip}(v^p - w_1^{p-1}v),$$

$$g^i(w_r) = \begin{cases} a^i w_r, & r \neq 2; \\ a^{-in} w_2, & r = 2. \end{cases} .$$

Hence $S(V)^G = S(M)^{\langle g \rangle}$ is Gorenstein if and only if $\det_M g^i = 1$, $\forall i$, if and only if $\deg_M g = 1$, that is $a^{p-1} = 1$, so $a \in \mathbb{F}_p$. So if $a \in F - \{0\}$, then $S(V)^G$ is Gorenstein if and only if $l = p - 1$.

5. Acknowledgements

Thanks are due to Yuval Ginosar, Martin Liebeck and Gunter Malle for their help in navigating through the turbulent waters of finite group theory.

REFERENCES

- [1] D. J. Benson, *Polynomial Invariants of Finite Groups*, LMS Lecture Notes Series, Vol. 190, Cambridge University Press, Cambridge, 1993.
- [2] A. Braun, *p-Groups and the polynomial ring of invariants question*, *Advances in Mathematics* 397 (2022).
- [3] A. Braun, *On the Gorenstein property for modular invariants*, *J. Algebra* 345 (2011) 81–99.
- [4] W. Bruns, J. Herzog *Cohen-Macaulay rings*, Cambridge Univ. Press, Cambridge (1998).
- [5] N. Bourbaki, *Groupes et algèbres de Lie*, IV, V et VI, Herman, Paris (1968).
- [6] C. Chevalley, *On the theory of local rings*, *Ann. Math.* 44 (1943), 690–708.
- [7] C. W. Curtis, I. Reiner *Representation Theory of Finite Groups and Associative Algebras*, Wiley Classics Library Edition, 1988.
- [8] H. Derksen, G. Kemper, *Computational Invariant Theory*, 2nd Edition, Springer-Verlag, 2015.
- [9] H. Hochster, C. Huneke *Tight closure and strong F-regularity*, *Soc. Math. de France, Memoire n° 38*, 1989, 119–133.
- [10] W. Kantor, *Subgroups of classical groups generated by long root elements*, *Trans. Amer. Math. Soc.* 248 (1979), 69–96.
- [11] G. Kemper, *Loci in quotients by finite groups, pointwise stabilizers and the Buchsbaum property*, *J. reine angew. Math.* 547 (2002), 347–379.
- [12] G. Kemper, G. Malle, *The finite irreducible groups with polynomial ring of invariants*, *Transformation Groups* 2 (1997), 57–89.
- [13] P. B. Kleidman, M. W. Liebeck *The subgroup structure of the finite classical groups*, Cambridge Univ. Press (1990).
- [14] S. J. Kovacs, *Rational singularities*, arXiv: 1703.02269v13 [math. AG], <https://doi.org/10.48550/arXiv.1703.02269>.
- [15] H. Matsumura, *Commutative Ring Theory*, Cambridge Univ. Press, Cambridge, (1986).
- [16] J. C. McConnell, J. C. Robson *Noncommutative Noetherian Rings*, Graduate Studies in Mathematics, vol. 30, A.M.S., Providence, Rhode Island, 2001.
- [17] H. Nakajima, *Regular Rings of Invariants of Unipotent Groups*, *J. Algebra.* 85, 253–286 (1983).
- [18] J. D. Sally, *Number of generators of ideals in local rings*, Marcel Dekker, Inc, New-York-Based, 1978.
- [19] J. -P. Serre, *Groupes finis d’automorphismes d’anneaux locaux réguliers*, *Colloque d’Algèbre ENSJF*, Paris, 1967, pp. 8-01–8-11.
- [20] Shchigolev, Vladimir, Stepanov, Dmitry *Three dimensionanl isolated quotient singularities in even characteristic*, *Glasg. Math. J.* 60 (2018), no. 2, 435–445.
- [21] R. Steinberg, *Differential equations invariant under finite reflection groups*, *Trans. Amer. Math. Soc.* 112 (1964), 392–400.
- [22] D. A. Stepanov, *Gorenstein Isolated Quotient Singularities Over \mathbb{C}* , *Proc. Edinburgh Math. Soc.* (2014) 57, 811–830.
- [23] D. A. Stepanov, *Three dimensional isolated quotient singularities in odd characteristic*, *Math. Sb.* 207 (2016), no. 6, 113–128.
- [24] A. Wagner, *The faithful linear representation of least degree of S_n and A_n over a field of characteristic 2*, *Math. Zeit.* 151 (1976), no. 2, 127–137.

- [25] Washington, Lawrence C. *Introduction to cyclotomic fields*, Grad. Texts in Math., 83, Springer-Verlag, New-York, 2nd edition, 1997.
- [26] J. A. Wolf, *Spaces of Constant Curvature*, AMS Chelsea Publishing, Providence, Rhode Island, 2010.
- [27] A. E. Zalesskii and V. N. Serežkin, *Linear groups generated by transvections*, Izv. Akad. Nauk SSSR, Ser. Mat. 40 (1976), no. 1 26–49 (in Russian). English translation: A. E. Zalesskii and V. N. Serežkin, *Linear groups generated by transvections*, Math. USSR Izv. 10 (1976), 25–46.