

The $\mathcal{N} = 2, 4$ Supersymmetric Linear $W_\infty[\lambda]$ Algebras for Generic λ Parameter

Changhyun Ahn[†], and Man Hea Kim^{*}

[†] Department of Physics, Kyungpook National University, Taegu 41566, Korea

^{*} Department of Physics Education, Sunchon National University, Sunchon 57922, Korea

ahn@knu.ac.kr, manhea.kim10000@gmail.com

Abstract

The four different kinds of currents are given by the multiple (β, γ) and (b, c) ghost systems with a multiple product of derivatives. We determine their complete algebra where the structure constants depend on the deformation parameter λ appearing in the conformal weights of above fields nontrivially and depend on the generic spins h_1 and h_2 appearing on the left hand sides in the (anti)commutators. By taking the linear combinations of these currents, the $\mathcal{N} = 4$ supersymmetric linear $W_\infty[\lambda]$ algebra (and its $\mathcal{N} = 4$ superspace description) for generic λ is obtained explicitly. Moreover, we determine the $\mathcal{N} = 2$ supersymmetric linear $W_\infty[\lambda]$ algebra for arbitrary λ . As a by product, the λ deformed bosonic $W_{1+\infty}[\lambda] \times W_{1+\infty}[\lambda + \frac{1}{2}]$ subalgebra (a generalization of Pope, Romans and Shen's work in 1990) is obtained. The first factor is realized by (b, c) fermionic fields while the second factor is realized by (β, γ) bosonic fields. The degrees of the polynomials in λ for the structure constants are given by $(h_1 + h_2 - 2)$. Each $w_{1+\infty}$ algebra from the celestial holography is reproduced by taking the vanishing limit of other deformation parameter q at $\lambda = 0$ with the contractions of the currents.

Contents

1	Introduction	1
2	The $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$ algebra	4
2.1	The bifundamental currents	4
2.2	The OPEs or (anti)commutators between the bifundamentals	6
2.3	The $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$ algebra	10
2.4	The $\mathcal{N} = 4$ supersymmetric OPE in $\mathcal{N} = 4$ superspace	15
3	The $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda]$ algebra	16
3.1	The singlet currents and the adjoint currents	16
3.2	The OPEs or (anti)commutators between the singlets and the adjoints	16
4	Conclusions and outlook	20
A	The remaining OPEs or (anti)commutators between the bifundamentals	22
B	The remaining $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$ algebra	28
C	The basic component OPEs for $\mathcal{N} = 4$ supersymmetric OPE	39
D	The $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda]$ algebra	47

1 Introduction

The simplest infinitely generated W -algebra, which has been named as w_∞ algebra [1], contains generators of spin (or weight) $2, 3, 4, \dots$. The operator product expansion (OPE) between these generators has simple form in the sense that there are only the second and the first order poles. The first order pole is given by the derivative of the second order pole with different relative coefficients. Once the spins of the generators on the left hand side are given by h_1 and h_2 , then the spin appearing in the second order pole on the right hand side is equal to $(h_1 + h_2 - 2)$. In terms of commutator, the mode dependent structure constant is given by $[(h_2 - 1)m - (h_1 - 1)n]$ where m is the mode for the generator of spin h_1 while n is the mode for the generator of spin h_2 . We can easily observe that this structure constant is odd under the exchange of $h_1 \leftrightarrow h_2$ and $m \leftrightarrow n$. This extra minus sign is consistent with the antisymmetry on the left hand side of the commutator under this exchange.

By considering all the possible terms on the right hand side of the commutator as well as the central term, the W_∞ algebra [2, 3] has been constructed. On the right hand side of the commutator or OPE, the generators of the spin $(h_1 + h_2 - 2)$, the spin $(h_1 + h_2 - 4)$, \dots , the spin 2 for even $(h_1 + h_2)$ or the spin 3 for odd $(h_1 + h_2)$ appear. The structure constants and the central term are determined by using the Jacobi identities. By rescaling the generators by spin dependent power of a parameter q , the commutator has q dependence explicitly. If we take q to zero limit, then the generator of spin $(h_1 + h_2 - 2)$ together with the central term survive [4]. Also the structure constant becomes the one in previous paragraph. Therefore, the centrally extended w_∞ algebra is reproduced under this process. By including the spin 1 generator in the commutator or OPE, the $W_{1+\infty}$ algebra [5] is obtained. The corresponding structure constants and the central term are similar to the ones for W_∞ algebra. Moreover, the generator of the spin 2 for even $(h_1 + h_2)$ or the spin 1 for odd $(h_1 + h_2)$ on the right hand side appears.

By considering the complex fermion realization of $W_{1+\infty}$ algebra and complex boson realization of W_∞ algebra, the $\mathcal{N} = 2$ supersymmetric W_∞ algebra [6] where the bosonic sector is $W_{1+\infty} \times W_\infty$ is found. By construction, there is no spin 1 generator from the complex bosons. The two supersymmetry currents can be constructed from the bilinears between the above complex bosons and complex fermions with possible derivatives. In particular, the right hand side of anticommutator between the two supersymmetry currents contains the bosonic currents of $W_{1+\infty}$ and W_∞ as well as the central term. All the structure constants and the central terms are determined by Jacobi identities.

In [7], the $U(K)$ matrix generalization of W_∞ algebra which is denoted by W_∞^K algebra is obtained. From the K values of complex bosons, the generators of this algebra can be written in terms of bilinear of these complex bosons contracted by $(K^2 - 1)$ generators of $SU(K)$ and $K \times K$ identity matrix with possible derivatives. Of course, when the two generators in the commutator are represented by this identity matrix, then this leads to the one in [3]. Moreover, in [8], the similar analysis (in the notation of $W_{1+\infty}^L$) for L values of complex fermions has been done and the generators are described by the bilinear of these complex fermions contracted by $(L^2 - 1)$ generators of $SU(L)$ and the identity matrix which is $L \times L$ matrix together with various derivatives. Along the line of [6], the $W_\infty^{K,L}$ algebra [9] has been obtained and the bosonic sector is given by $W_{1+\infty}^L \times W_\infty^K$. Instead of having the singlets and the adjoints of the $SU(K)$ and $SU(L)$, the double index notation in [8] leads to the fact that there are no antisymmetric f symbol or symmetric d symbol in the commutators. Under the condition of $K = L$, the $\mathcal{N} = 2$ supersymmetry arises.

In [10, 11], it is known that the $\mathcal{N} = 2$ supersymmetric linear $W_\infty[\lambda]$ algebra is isomorphic

to the $\mathcal{N} = 2$ supersymmetric linear $W_\infty[\frac{1}{2} - \lambda]$ algebra. There exists a deformation parameter λ such that the $\mathcal{N} = 2$ supersymmetric $W_\infty[\lambda]$ (which is equivalent to $W_\infty^{1,1}[\lambda]$ in later notation) algebra can be truncated on the various subalgebras for the particular values of λ . At $\lambda = 0$, the two fermions are identified with each other and one of the bosons is identified with the derivative of other boson. Then the algebra reduces to the W_∞ algebra. At $\lambda = \frac{1}{2}$, the two bosons are identified with each other and one of the fermions is identified with the derivative of other fermion. This implies the W_∞ algebra. As a subalgebra, the bosonic algebra contains $W_\infty[\lambda]$ algebra and $W_\infty[\lambda + \frac{1}{2}]$ algebra.

In [12], the multiple numbers of complex fermions and complex bosons are introduced to find the exact correspondence between the $\mathcal{N} = (2, 2)$ SYK model [13, 14, 15] and the $\mathcal{N} = 2$ (β, γ) and (b, c) systems [10, 11]. Because there are multiple numbers of chiral multiplets (or Fermi multiplets) in the former, the multiple numbers of (β, γ) and (b, c) fields are crucial. See also the previous work of [16]. Note that the fundamental OPEs between these fields have the first order poles, compared to the ones in [6]. The generators of spin 1 consisting of the bilinear of β field and γ field with contraction and spin $\frac{1}{2}$ consisting of β field and c field with contraction occur on the right hand sides of the (anti)commutators. At $\lambda = 0$, the previous result in [17] is reproduced. The bosonic subalgebra of the $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda = 0]$ algebra is given by $W_{1+\infty}^K[\lambda = 0]$ generated by fermions (b, c) and $W_\infty^K[\lambda + \frac{1}{2} = \frac{1}{2}]$ generated by bosons (β, γ) . On the other hand, at $\lambda = \frac{1}{2}$, the bosonic subalgebra of the $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda = \frac{1}{2}]$ algebra is given by $W_\infty^K[\lambda = \frac{1}{2}]$ generated by fermions and $W_{1+\infty}^K[\lambda + \frac{1}{2} = 1]$ generated by bosons.

Recently, as noted in [18], the structure constants obtained in [12] hold for some constraints on the spins for the left hand sides of the algebras with nonzero λ : for general h_1 with the constrained h_2 . In [19], some OPEs between the $\mathcal{N} = 4$ multiplets of low spins h_1 and h_2 are found explicitly for nonzero λ . Moreover, the results in [18] satisfy with some constraints on the spins also¹.

In this paper, we would like to construct the OPEs or (anti)commutators between the bifundamentals under the $SU(2) \times SU(2)$ for *nonzero* deformation parameter λ with *generic* h_1 and h_2 , by reconsidering [12, 18]. Once these are determined explicitly, then the complete OPEs between the $\mathcal{N} = 4$ multiplets, related to the $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$

¹At $\lambda = 0$, all the (anti)commutators are true. At nonzero λ , under the condition of $h_1 = h_2$, all the (anti)commutators which do not contain the spin 1 or the spin $\frac{1}{2}$ currents on the left hand side are valid. If they contain one of these on the left hand sides, the structure constants depend on the λ and cannot be written as the known structure constants in [18]. For $h_1 \neq h_2$ at nonzero λ , in general, the right hand sides of the (anti)commutators contain the extra structures which cannot be written in terms of the structure constants of [18] and depend on the λ . According to the results of present paper, all these constraints are gone and we have the explicit (anti)commutators with complete structure constants for any λ, h_1 and h_2 .

algebra, are obtained and the full algebra between the singlets and the adjoints in the $SU(K)$, related to the $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda]$ algebra, are determined completely.

In section 2, the complete algebra between the bifundamentals under the $SU(2) \times SU(2)$ is obtained for any h_1, h_2 and deformation parameter λ . Then the $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$ algebra is determined and its $\mathcal{N} = 4$ superspace description is presented.

In section 3, the complete $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda]$ algebra between the singlets and the adjoints under the $SU(K)$ is found.

In section 4, the main results of this paper are summarized.

In Appendices, all the details which do not appear in previous sections are given ².

2 The $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$ algebra

2.1 The bifundamental currents

The operator product expansions of the (β, γ) and (b, c) systems are given by [16]

$$\gamma^{i,\bar{a}}(z) \beta^{\bar{j},b}(w) = \frac{1}{(z-w)} \delta^{i\bar{j}} \delta^{\bar{a}b} + \dots, \quad c^{i,\bar{a}}(z) b^{\bar{j},b}(w) = \frac{1}{(z-w)} \delta^{i\bar{j}} \delta^{\bar{a}b} + \dots. \quad (2.1)$$

There exist fundamental indices a, b and antifundamental indices \bar{a}, \bar{b} of $SU(2)$. Similarly there are fundamental indices i, j and antifundamental indices \bar{i}, \bar{j} of $SU(N)$. The (β, γ) fields are bosonic operators while the (b, c) fields are fermionic operators. Note that the conformal weights of (β, γ) fields are given by $(\lambda, 1 - \lambda)$ while the conformal weights of (b, c) fields are given by $(\frac{1}{2} + \lambda, \frac{1}{2} - \lambda)$, under the stress energy tensor (2.4) ³.

The $SU(N)$ singlet currents can be determined by summing over the (anti)fundamental indices of $SU(N)$ ⁴ and are given by [19, 18] from the work of [10, 11]

$$W_{F,h}^{\lambda,\bar{a}b}(z) = (-4q)^{h-2} \sum_{i=0}^{h-1} a^i(h, \lambda + \frac{1}{2}) \partial_z^{h-i-1} ((\partial_z^i b^{\bar{l}\bar{b}}) \delta_{\bar{l}\bar{l}} c^{l\bar{a}})(z),$$

² In the abstract, we use a simplified notation for $W_\infty[\lambda]$ algebra. For the $\mathcal{N} = 4$ supersymmetric case, the $W_\infty[\lambda]$ stands for $W_\infty^{2,2}[\lambda]$ and for the $\mathcal{N} = 2$ supersymmetric case, the $W_\infty[\lambda]$ stands for $W_\infty^{K,K}[\lambda]$. The bosonic subalgebras stand for $W_{1+\infty}^K[\lambda] \times W_{1+\infty}^K[\lambda + \frac{1}{2}]$. Maybe the $W_{1+\infty}[\lambda]$ is better than the $W_\infty[\lambda]$ appearing in the title but we keep the notation of [10, 11].

³ Under the $\lambda \rightarrow \lambda + \frac{1}{2}$, the conformal weights for the former become the ones for the latter and under the $\lambda \rightarrow \lambda - \frac{1}{2}$, they for the latter become the ones for the former. These transformations will provide the symmetries in the bosonic subalgebras later.

⁴The free field theory construction we consider appears in the realization of $\mathcal{N} = 4$ coset $\frac{SU(N+2)}{SU(N)}$ model [20]. Then the currents we consider, which are bilinear in the free fields, should be invariant under the denominator $SU(N)$. In the subsections 2.3 and 2.4, the free field realization associated with this $\mathcal{N} = 4$ coset model, under the identification of the λ with the corresponding parameter appearing in the $\mathcal{N} = 4$ coset model [19], can be found.

$$\begin{aligned}
W_{B,h}^{\lambda,\bar{a}\bar{b}}(z) &= (-4q)^{h-2} \sum_{i=0}^{h-1} a^i(h, \lambda) \partial_z^{h-i-1} ((\partial_z^i \beta^{\bar{l}\bar{b}}) \delta_{\bar{l}\bar{l}} \gamma^{l\bar{a}})(z), \\
Q_{h+\frac{1}{2}}^{\lambda,\bar{a}\bar{b}}(z) &= \sqrt{2} (-4q)^{h-1} \sum_{i=0}^{h-1} \beta^i(h+1, \lambda) \partial_z^{h-i-1} ((\partial_z^i b^{\bar{l}\bar{b}}) \delta_{\bar{l}\bar{l}} \gamma^{l\bar{a}})(z), \\
\bar{Q}_{h+\frac{1}{2}}^{\lambda,a\bar{b}}(z) &= \sqrt{2} (-4q)^{h-1} \sum_{i=0}^h \alpha^i(h+1, \lambda) \partial_z^{h-i} ((\partial_z^i \beta^{\bar{l}a}) \delta_{\bar{l}\bar{l}} c^{\bar{l}\bar{b}})(z).
\end{aligned} \tag{2.2}$$

By counting the number of (anti)fundamental indices, there exist four components labeled by $(\bar{a}\bar{b}) = (11, 12, 21, 22)$ in each current. Then there are eight bosonic currents, $W_{F,h}^{\lambda,\bar{a}\bar{b}}$ and $W_{B,h}^{\lambda,\bar{a}\bar{b}}$, for the weight $h = 1, 2, \dots$ and eight fermionic currents, $Q_{h+\frac{1}{2}}^{\lambda,\bar{a}\bar{b}}$ and $\bar{Q}_{h+\frac{1}{2}}^{\lambda,\bar{a}\bar{b}}$, for the weight $h + \frac{1}{2} = \frac{3}{2}, \frac{5}{2}, \dots$ as well as four fermionic currents $\bar{Q}_{\frac{1}{2}}^{\lambda,a\bar{b}}$ of the weight $\frac{1}{2}$ in (2.2), which is rescaled by $\frac{1}{2}$, compared to the one in [18]. Note that four fermionic currents $Q_{\frac{1}{2}}^{\lambda,\bar{a}\bar{b}}$ of the weight $\frac{1}{2}$ are identically zero.

The relative coefficients appearing in (2.2) depend on the conformal weight h and deformation parameter λ explicitly. They are given by the binomial coefficients denoted by parentheses and the rising Pochhammer symbols where $(a)_n \equiv a(a+1)\cdots(a+n-1)$ and n is a nonnegative integer [3] as follows [10, 11]:

$$\begin{aligned}
a^i(h, \lambda) &\equiv \binom{h-1}{i} \frac{(-2\lambda - h + 2)_{h-1-i}}{(h+i)_{h-1-i}}, \quad 0 \leq i \leq h-1, \\
\alpha^i(h, \lambda) &\equiv \binom{h-1}{i} \frac{(-2\lambda - h + 2)_{h-1-i}}{(h+i-1)_{h-1-i}}, \quad 0 \leq i \leq h-1, \\
\beta^i(h, \lambda) &\equiv \binom{h-2}{i} \frac{(-2\lambda - h + 2)_{h-2-i}}{(h+i)_{h-2-i}}, \quad 0 \leq i \leq h-2.
\end{aligned} \tag{2.3}$$

The λ dependence appears in the numerators of (2.3). The degrees of the polynomials in λ are given by $(h-1-i)$, $(h-1-i)$ and $(h-2-i)$ respectively. The corresponding ones in (2.2) are given by $(h-1)$, $(h-1)$, $(h-1)$ and h respectively.

The stress energy tensor of conformal weight 2 is given by [10, 11, 19, 18]

$$L = W_{B,2}^{\lambda,\bar{a}a} + W_{F,2}^{\lambda,\bar{a}a}, \tag{2.4}$$

and the corresponding central charge is

$$c_{cen} = 6N(1-4\lambda), \tag{2.5}$$

which depends on the deformation parameter λ explicitly. The above bosonic and fermionic currents in (2.2) are quasiprimary operators under the stress energy tensor (2.4)⁵.

⁵ The rank $(q_{syk} + 1)$ of the interaction of the two dimensional $\mathcal{N} = (2, 2)$ SYK model corresponds to $(q_{syk} + 1) = \frac{1}{2\lambda}$.

2.2 The OPEs or (anti)commutators between the bifundamentals

Let us calculate the OPEs between the currents in (2.2). As an intermediate step, we obtain the following OPE, from (2.1), between the first currents without any derivatives⁶

$$\begin{aligned}
((\partial_z^{i_1} b^{\bar{l}b}) \delta_{\bar{l}l} c^{l\bar{a}})(z) ((\partial_w^{i_2} b^{\bar{l}d}) \delta_{\bar{l}l} c^{l\bar{c}})(w) &= \delta^{\bar{a}d} \delta^{\bar{c}b} (-1)^{i_2+1} \frac{N i_1! j_2!}{(i_1 + i_2 + 1)!} \partial_z^{i_1+i_2+1} \left[\frac{1}{(z-w)} \right] \\
&\quad + \delta^{\bar{a}d} \sum_{r=0}^{i_2} (-1)^r \binom{i_2}{r} \partial_z^r \left[\frac{((\partial_w^{i_1+i_2-r} b^{\bar{l}b}) \delta_{\bar{l}l} c^{l\bar{c}})(w)}{(z-w)} \right] \\
&\quad - \delta^{\bar{c}b} \sum_{r=0}^{i_1} (-1)^r \binom{i_1}{r} \partial_w^r \left[\frac{((\partial_w^{i_1+i_2-r} b^{\bar{l}d}) \delta_{\bar{l}l} c^{l\bar{a}})(w)}{(z-w)} \right] \\
&\quad + \dots
\end{aligned} \tag{2.6}$$

The central term in (2.6) comes from the double contractions of this OPE. Note that the right hand side contains the quadratic terms in (b, c) fields with various derivatives.

Then it is straightforward to calculate the following OPE

$$\begin{aligned}
&\partial_z^{h_1-i_1-1} ((\partial_z^{i_1} b^{\bar{l}b}) \delta_{\bar{l}l} c^{l\bar{a}})(z) \partial_w^{h_2-i_2-1} ((\partial_w^{i_2} b^{\bar{l}d}) \delta_{\bar{l}l} c^{l\bar{c}})(w) \\
&= \delta^{\bar{a}d} \delta^{\bar{c}b} (-1)^{h_2} \frac{N i_1! i_2! (i_1 + i_2 + 2)_{h_1+h_2-i_1-i_2-2}}{(h_1 + h_2 - 1)!} \partial_z^{h_1+h_2-1} \left[\frac{1}{(z-w)} \right] \\
&\quad + \delta^{\bar{a}d} \sum_{r=0}^{i_2} (-1)^r \binom{i_2}{r} \partial_z^{h_1-i_1-1+r} \partial_w^{h_2-i_2-1} \left[\frac{((\partial_w^{i_1+i_2-r} b^{\bar{l}b}) \delta_{\bar{l}l} c^{l\bar{c}})(w)}{(z-w)} \right] \\
&\quad - \delta^{\bar{c}b} \sum_{r=0}^{i_1} (-1)^r \binom{i_1}{r} \partial_z^{h_1-i_1-1} \partial_w^{h_2-i_2-1+r} \left[\frac{((\partial_w^{i_1+i_2-r} b^{\bar{l}d}) \delta_{\bar{l}l} c^{l\bar{a}})(w)}{(z-w)} \right] + \dots
\end{aligned} \tag{2.7}$$

Let us consider the numerator appearing inside the bracket of the second line of (2.7). We expect that we should write down it in terms of the derivative of $W_{F,h+1}^{\lambda,\bar{a}\bar{b}}$. The number of derivative of b field is taken as s and the conformal weight for the derivative term is $(s+1)$. Then we can think of all possible terms having the weight $(s+1)$ from the currents, $W_{F,1}^{\lambda,\bar{a}\bar{b}}$ with s derivatives, \dots , $W_{F,s}^{\lambda,\bar{a}\bar{b}}$ with one derivative. We introduce the undetermined coefficients appearing in these derivative terms. They depend on the λ parameter and the number of derivative of b field s . Obviously, the derivatives of c field should vanish and all the derivatives of b field have the numerical values 1. By varying the s starting from 0 to 10, the above undetermined coefficients can be fixed and the s dependence can be determined explicitly. It

⁶The Thielemans package [21] is used together with a mathematica [22] all the times in this paper.

turns out that the following nontrivial identity satisfies

$$((\partial_z^s b^{\bar{l}b}) \delta_{\bar{l}l} c^{l\bar{a}})(z) = \sum_{h=0}^s (-1)^{s+1} 4^{-h} q^{1-h} Y_{W_F}(s-h, s+1, \lambda) \partial_z^{s-h} W_{F,h+1}^{\lambda, \bar{a}b}(z), \quad (2.8)$$

where the λ dependent coefficient is given by

$$Y_{W_F}(h, s, \lambda) \equiv \frac{2(-2\lambda - s + 1)_h [s-1]_{h-1}}{h! [2s-h-1]_{h-1}} = \frac{2(-2\lambda - s + 1)_h (1-s)_{h-1}}{h! (1-2s+h)_{h-1}}. \quad (2.9)$$

The degree of this polynomial in λ is h and the falling Pochhammer symbol $[a]_n \equiv a(a-1)\cdots(a-n+1)$ is used. Moreover, the identity $[a]_n = (-1)^n (-a)_n$ is used.

The contribution from $((\partial^{i_1+i_2-r} b^{\bar{l}b}) \delta_{\bar{l}l} c^{l\bar{c}})$, by using (2.9), can be written as

$$\begin{aligned} & \sum_{h_3=1}^{i_1+i_2-r+1} (-1)^{i_1+i_2-r+1} 4^{1-h_3} q^{2-h_3} \frac{2(-i_1-i_2+r-2\lambda)_{i_1+i_2-r-h_3+1} (-i_1-i_2+r)_{i_1+i_2-r-h_3}}{(i_1+i_2-r-h_3+1)! (-i_1-i_2+r-h_3)_{i_1+i_2-h_3-r}} \\ & \times \partial_w^{i_1+i_2-r+1-h_3} W_{F,h_3}^{\lambda, \bar{c}b}(w), \end{aligned} \quad (2.10)$$

by identifying $s = i_1 + i_2 - r$, $h = h_3 - 1$ and $\bar{a} = \bar{c}$. After multiplying all the factors into (2.10), we obtain

$$\begin{aligned} & \delta^{\bar{a}d} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_2} \sum_{h_3=1}^{h_1+h_2-1} (-1)^{h_1+h_2+i_1+i_2+1} 4^{h_1+h_2-h_3-3} q^{h_1+h_2-h_3-2} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \\ & \times \frac{2(-i_1-i_2+r-2\lambda)_{i_1+i_2-r-h_3+1} (-i_1-i_2+r)_{i_1+i_2-r-h_3}}{(i_1+i_2-r-h_3+1)! (-i_1-i_2+r-h_3)_{i_1+i_2-h_3-r}} \binom{i_2}{r} \\ & \times \partial_z^{h_1-i_1-1+r} \partial_w^{h_2-i_2-1} \left[\frac{\partial_w^{i_1+i_2-r-h_3+1} W_{F,h_3}^{\lambda, \bar{c}b}(w)}{(z-w)} \right], \end{aligned} \quad (2.11)$$

where the upper bound for h_3 , (i_1+i_2-r+1) , is changed into (h_1+h_2-1) because the values of $h_3 \geq (i_1+i_2-r+2)$ do not contribute due to the factorial function $(i_1+i_2-r-h_3+1)!$ in the denominator. This changing of upper bound in (2.11) is crucial to the later discussion.

By introducing a new variable $r' \equiv i_1+i_2-r$ and removing a prime notation, the following relation satisfies

$$\begin{aligned} & \delta^{\bar{a}d} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \sum_{h_3=1}^{h_1+h_2-1} (-1)^{h_1+h_2+i_1+i_2+1} 4^{h_1+h_2-h_3-3} q^{h_1+h_2-h_3-2} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \\ & \times \frac{2 \prod_{x=0}^{r-h_3} (-r-2\lambda+x) (-r)_{-h_3+r}}{(1-h_3+r)! (-h_3-r)_{-h_3+r}} \binom{i_2}{i_1+i_2-r} \\ & \times \partial_z^{h_1+i_2-r-1} \partial_w^{h_2-i_2-1} \left[\frac{\partial_w^{r-h_3+1} W_{F,h_3}^{\lambda, \bar{c}b}(w)}{(z-w)} \right], \end{aligned} \quad (2.12)$$

where we intentionally add the zero value which is equal to the summation of r from $r = 0$ to $r = i_1 - 1$ from the above binomial. Moreover, we can move the derivative appearing inside of the bracket (2.12) by using the relation

$$\begin{aligned} \left[\frac{\partial_w^{r-h_3+1} W_{F,h_3}^{\lambda,\bar{c}\bar{b}}(w)}{(z-w)} \right] &= \sum_{k'=0}^{r-h_3+1} \binom{r-h_3+1}{r-h_3+1-k'} \partial_z^{k'} \partial_w^{r-h_3+1-k'} \left[\frac{W_{F,h_3}^{\lambda,\bar{c}\bar{b}}(w)}{(z-w)} \right] \\ &= \sum_{k=h_2-i_2-1}^{h_2-h_3-i_2+r} \binom{r-h_3+1}{1+k-h_2+i_2} \partial_z^{-k+h_2-h_3-i_2+r} \partial_w^{1+k-h_2+i_2} \left[\frac{W_{F,h_3}^{\lambda,\bar{c}\bar{b}}(w)}{(z-w)} \right], \end{aligned} \quad (2.13)$$

where the new variable $k \equiv -k' + h_2 - h_3 - i_2 + r$ which is related to the upper bound of (2.13) is introduced.

By substituting (2.13) into (2.12), the following result satisfies

$$\begin{aligned} \delta^{\bar{a}\bar{d}} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} (-1)^{h_1+h_2+i_1+i_2+1} 4^{h_1+h_2-h_3-3} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \\ \times q^{h_1+h_2-h_3-2} \frac{2 \prod_{x=0}^{r-h_3} (-r-2\lambda+x) (-r)_{-h_3+r}}{(1-h_3+r)! (-h_3-r)_{-h_3+r}} \binom{i_2}{i_1+i_2-r} \binom{r-h_3+1}{1+k-h_2+i_2} \\ \times \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k \left[\frac{W_{F,h_3}^{\lambda,\bar{c}\bar{b}}(w)}{(z-w)} \right]. \end{aligned} \quad (2.14)$$

Note that the upper bound and the lower bound for k in (2.14) are changed but the extra terms do not contribute because of the constraint on the k appearing in the binomial.

By using the relations $(-x)_n = (-1)^n (x-n+1)_n$ and $(a)_n = \frac{(a+n-1)!}{(a-1)!}$, the following relation satisfies

$$\frac{2(-r)_{-h_3+r}}{(1-h_3+r)! (-h_3-r)_{-h_3+r}} = 4 \binom{r}{h_3-1} \frac{(2h_3-1)!}{(h_3+r)!}. \quad (2.15)$$

Therefore, the contribution from the second term of (2.7) with the correct factors, together with (2.15), can be described by

$$\begin{aligned} \delta^{\bar{a}\bar{d}} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! S_{F,R}^{h_1,h_2,h_3,k}(\lambda) \\ \times \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k \left[\frac{W_{F,h_3}^{\lambda,\bar{c}\bar{b}}(w)}{(z-w)} \right], \end{aligned} \quad (2.16)$$

where the λ dependent coefficients in (2.16) are given by

$$S_{F,R}^{h_1,h_2,h_3,k}(\lambda) \equiv \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \right]$$

$$\times \binom{r}{h_3-1} \binom{i_2}{i_1+i_2-r} \binom{1-h_3+r}{1-h_2+i_2+k} \prod_{j=0}^{r-h_3} (-r-2\lambda+j) \Big]. \quad (2.17)$$

We can do similar analysis by taking the third term of (2.7)⁷.

Then the final OPE between $W_{F,h_1}^{\lambda,\bar{a}b}(z)$ and $W_{F,h_2}^{\lambda,\bar{c}d}(w)$, by considering two other contributions, is summarized by

$$\begin{aligned} W_{F,h_1}^{\lambda,\bar{a}b}(z) W_{F,h_2}^{\lambda,\bar{c}d}(w) &= \delta^{\bar{a}d} \delta^{\bar{c}b} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} (-1)^{h_1} 4^{h_1+h_2-4} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \frac{N i_1! i_2!}{(i_1 + i_2 + 1)!} \\ &\quad \times q^{h_1+h_2-4} \partial_z^{h_1+h_2-1} \left[\frac{1}{(z-w)} \right] \\ &\quad + \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3 - 1)! \\ &\quad \times \left(\delta^{\bar{a}d} S_{F,R}^{h_1,h_2,h_3,k}(\lambda) \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k \left[\frac{W_{F,h_3}^{\lambda,\bar{c}b}(w)}{(z-w)} \right] \right. \\ &\quad \left. - \delta^{\bar{c}b} S_{F,L}^{h_1,h_2,h_3,k}(\lambda) \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \left[\frac{W_{F,h_3}^{\lambda,\bar{a}d}(w)}{(z-w)} \right] \right) + \dots, \end{aligned} \quad (2.18)$$

where other type of λ dependent coefficients is⁸

$$\begin{aligned} S_{F,L}^{h_1,h_2,h_3,k}(\lambda) &\equiv \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \right. \\ &\quad \times \left. \binom{r}{h_3-1} \binom{i_1}{i_1+i_2-r} \binom{1-h_3+r}{1-h_1+i_1+k} \prod_{j=0}^{r-h_3} (-r-2\lambda+j) \right]. \end{aligned} \quad (2.19)$$

The highest power of $\frac{1}{(z-w)}$ in the field dependent terms of (2.18) is given by $(h_1 + h_2 - h_3)$. Then the total conformal spin of the right hand side of this term is given by $(h_1 + h_2)$. On the other hands, the corresponding lowest power of $\frac{1}{(z-w)}$ is 1 when $k = h_1 + h_2 - h_3 - 1$ for $\delta^{\bar{a}d}$ and $k = 0$ for $\delta^{\bar{c}b}$. The number of derivative with respect to w is given by $(h_1 + h_2 - h_3 - 1)$. Note that the number of powers in the derivatives appearing in the last two lines (2.18) arises in symmetric way. Compared to the one in (2.17), the difference in (2.19) appears in some elements of binomials. Or according to the exchange of $i_1 \leftrightarrow i_2$ and $h_1 \leftrightarrow h_2$, the structure constant (2.17) becomes the structure constant (2.19) or vice versa. The free indices in (2.18) are given by h_1 and h_2 while i_1, i_2, h_3 and k are dummy variables.

⁷If the upper bound is less than the lower bound in the product notation appearing in (2.17), we treat them to be 1 according to the “empty” product.

⁸There is no meaning for the assignments in the R and L appearing in (2.17) and (2.19).

It is straightforward to determine the commutator corresponding to (2.18) by using the standard method of conformal field theory. According to the analysis of [3], the derivative terms, ∂_z^h and $\partial_w^{h'}$ can be replaced by $(-1)^h [m + h - 1]_h$ and $(-1)^{h'} [n + h' - 1]_{h'}$ respectively. It turns out that ⁹

$$\begin{aligned} \left[(W_{F,h_1}^{\lambda,\bar{a}b})_m, (W_{F,h_2}^{\lambda,\bar{c}d})_n \right] &= \delta^{\bar{a}d} \delta^{\bar{c}b} q^{h_1+h_2-4} c_{W_F}^{h_1,h_2}(\lambda) [m + h_1 - 1]_{h_1+h_2-1} \delta_{m+n} \\ &\quad + \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3 - 1)! \\ &\quad \times \left[\delta^{\bar{a}d} S_{F,R}^{h_1,h_2,h_3,k}(\lambda) [m + h_1 - 1]_{h_1+h_2-h_3-k-1} [n + h_2 - 1]_k (W_{F,h_3}^{\lambda,\bar{c}b})_{m+n} \right. \\ &\quad \left. - \delta^{\bar{c}b} S_{F,L}^{h_1,h_2,h_3,k}(\lambda) [m + h_1 - 1]_k [n + h_2 - 1]_{h_1+h_2-h_3-k-1} (W_{F,h_3}^{\lambda,\bar{a}d})_{m+n} \right], \end{aligned} \quad (2.20)$$

where mode independent central term appearing in (2.20) is given by

$$c_{W_F}^{h_1,h_2}(\lambda) = N \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} (-1)^{h_2-1} 4^{h_1+h_2-4} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \frac{i_1! i_2!}{(i_1 + i_2 + 1)!}. \quad (2.21)$$

The OPE (2.18) or commutator (2.20) is one of the main result of this paper. In Appendix A we will present other remaining OPEs or (anti)commutators. We have found the complete algebra between the currents in (2.2) for generic deformation parameter λ for any conformal weights h_1 and h_2 where $h_1, h_2 = 1, 2, 3, \dots$. At $\lambda = 0$, obviously, the previous results [18] are reproduced.

2.3 The $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$ algebra

The $\mathcal{N} = 4$ multiplet can be described by [23, 24, 25] the following $\mathcal{N} = 4$ super field

$$\Phi^{(h)}(Z) = \Phi_0^{(h)}(z) + \theta^i \Phi_{\frac{1}{2}}^{(h),i}(z) + \theta^{4-i,j} \Phi_1^{(h),ij}(z) + \theta^{4-i} \Phi_{\frac{3}{2}}^{(h),i}(z) + \theta^{4-0} \Phi_2^{(h)}(z), \quad (2.22)$$

where $SO(4)$ indices $i, j = 1, 2, \dots, 4$. The conventions for the $\mathcal{N} = 4$ superspace coordinates and the left covariant spinor derivatives are taken from [23, 25]. The Grassmann coordinate θ^i has a weight $-\frac{1}{2}$ ¹⁰.

⁹Let us emphasize that inside of the summations there are various Pochhammer symbols. By using the mathematica [22], for example, the summation over k will lead to some complicated combinations of special functions with h_1, h_2, h_3 and modes m, n dependences. If we substitute the particular values for h_1, h_2 and h_3 , then still the resulting expression is again complicated and is not written in terms of polynomials of m and n . Therefore, we introduce a function which depends on the above variables h_1, h_2 and h_3 and use a Table and Do commands inside a mathematica. Then we obtain the expected polynomials of m, n for h_1, h_2 and h_3 .

¹⁰The coordinates of $\mathcal{N} = 4$ superspace can be written in terms of (Z, \bar{Z}) where $Z = (z, \theta^i)$, $\bar{Z} = (\bar{z}, \bar{\theta}^i)$ and the $SO(4)$ -vector index i is given by $i = 1, \dots, 4$. The left covariant spinor derivative is defined by

The lowest component of (2.22) having a weight h is described by [18]

$$\Phi_0^{(h)} = \frac{q^{2-h}(-4)^{h-2}}{(2h-1)} \left[-(h-2\lambda) W_{F,h}^{\lambda,\bar{a}a} + (h-1+2\lambda) W_{B,h}^{\lambda,\bar{a}a} \right], \quad (2.23)$$

where the bosonic currents are given by (2.2) and there are summation over $SU(2)$ indices.

It is known that the next lowest components in (2.22) can be determined by using the four supersymmetry currents of $\mathcal{N} = 4$ stress energy tensor and the above lowest component (2.23). By computing the first order poles of these OPEs, we obtain [18] with an extra minus sign

$$\begin{aligned} \Phi_{\frac{1}{2}}^{(h),1} &= -q^{1-h} (-4)^{h-3} \left[-\frac{1}{2} (Q_{h+\frac{1}{2}}^{\lambda,11} + i\sqrt{2} Q_{h+\frac{1}{2}}^{\lambda,12} + 2i\sqrt{2} Q_{h+\frac{1}{2}}^{\lambda,21} - 2 Q_{h+\frac{1}{2}}^{\lambda,22} \right. \\ &\quad \left. + 2 \bar{Q}_{h+\frac{1}{2}}^{\lambda,11} + 2i\sqrt{2} \bar{Q}_{h+\frac{1}{2}}^{\lambda,12} + i\sqrt{2} \bar{Q}_{h+\frac{1}{2}}^{\lambda,21} - \bar{Q}_{h+\frac{1}{2}}^{\lambda,22}) \right], \\ \Phi_{\frac{1}{2}}^{(h),2} &= -q^{1-h} (-4)^{h-3} \left[\frac{i}{2} (Q_{h+\frac{1}{2}}^{\lambda,11} + 2i\sqrt{2} Q_{h+\frac{1}{2}}^{\lambda,21} - 2 Q_{\frac{3}{2}}^{\lambda,22} + 2 \bar{Q}_{h+\frac{1}{2}}^{\lambda,11} + 2i\sqrt{2} \bar{Q}_{h+\frac{1}{2}}^{\lambda,12} - \bar{Q}_{h+\frac{1}{2}}^{\lambda,22}) \right], \\ \Phi_{\frac{1}{2}}^{(h),3} &= -q^{1-h} (-4)^{h-3} \left[\frac{i}{2} (Q_{h+\frac{1}{2}}^{\lambda,11} + i\sqrt{2} Q_{h+\frac{1}{2}}^{\lambda,12} - 2 Q_{h+\frac{1}{2}}^{\lambda,22} + 2 \bar{Q}_{h+\frac{1}{2}}^{\lambda,11} + i\sqrt{2} \bar{Q}_{h+\frac{1}{2}}^{\lambda,21} - \bar{Q}_{h+\frac{1}{2}}^{\lambda,22}) \right], \\ \Phi_{\frac{1}{2}}^{(h),4} &= -q^{1-h} (-4)^{h-3} \left[\frac{1}{2} Q_{h+\frac{1}{2}}^{\lambda,11} + Q_{h+\frac{1}{2}}^{\lambda,22} - \bar{Q}_{h+\frac{1}{2}}^{\lambda,11} - \frac{1}{2} \bar{Q}_{h+\frac{1}{2}}^{\lambda,22} \right], \end{aligned} \quad (2.24)$$

where the fermionic currents are given in (2.2).

The following six components of (2.22) can be determined by calculating the OPEs between the previous four supersymmetry currents and the components appearing in (2.24) and reading off the first order poles [18] with an extra minus sign

$$\begin{aligned} \Phi_1^{(h),12} &= q^{1-h} (-4)^{h-3} \left[2i W_{B,h+1}^{\lambda,11} - \sqrt{2} W_{B,h+1}^{\lambda,12} - 2i W_{B,h+1}^{\lambda,22} \right. \\ &\quad \left. + 2i W_{F,h+1}^{\lambda,11} - 2\sqrt{2} W_{F,h+1}^{\lambda,12} - 2i W_{F,h+1}^{\lambda,22} \right], \\ \Phi_1^{(h),13} &= q^{1-h} (-4)^{h-3} \left[-2i W_{B,h+1}^{\lambda,11} + 4\sqrt{2} W_{B,h+1}^{\lambda,21} + 2i W_{B,h+1}^{\lambda,22} \right. \\ &\quad \left. - 2i W_{F,h+1}^{\lambda,11} + 2\sqrt{2} W_{F,h+1}^{\lambda,21} + 2i W_{F,h+1}^{\lambda,22} \right], \end{aligned}$$

$D^i = \theta^i \frac{\partial}{\partial z} + \frac{\partial}{\partial \theta^i}$ with the anticommutators $\{D^i, D^j\} = 2\delta^{ij} \frac{\partial}{\partial z}$. The notation θ^{4-0} stands for $\theta^1 \theta^2 \theta^3 \theta^4$. The complement $4-i$ is defined such that $\theta^1 \theta^2 \theta^3 \theta^4 = \theta^{4-i} \theta^i$ [23, 25].

$$\begin{aligned}
\Phi_1^{(h),14} &= q^{1-h} (-4)^{h-3} \left[2 W_{B,h+1}^{\lambda,11} + i\sqrt{2} W_{B,h+1}^{\lambda,12} + 4i\sqrt{2} W_{B,h+1}^{\lambda,21} - 2 W_{B,h+1}^{\lambda,22} \right. \\
&\quad \left. - 2 W_{F,h+1}^{\lambda,11} - 2i\sqrt{2} W_{F,h+1}^{\lambda,12} - 2i\sqrt{2} W_{F,h+1}^{\lambda,21} + 2 W_{F,h+1}^{\lambda,22} \right], \\
\Phi_1^{(h),23} &= q^{1-h} (-4)^{h-3} \left[- 2 W_{B,h+1}^{\lambda,11} - i\sqrt{2} W_{B,h+1}^{\lambda,12} - 4i\sqrt{2} W_{B,h+1}^{\lambda,21} + 2 W_{B,h+1}^{\lambda,22} \right. \\
&\quad \left. - 2 W_{F,h+1}^{\lambda,11} - 2i\sqrt{2} W_{F,h+1}^{\lambda,12} - 2i\sqrt{2} W_{F,h+1}^{\lambda,21} + 2 W_{F,h+1}^{\lambda,22} \right], \\
\Phi_1^{(h),24} &= q^{1-h} (-4)^{h-3} \left[- 2i W_{B,h+1}^{\lambda,11} + 4\sqrt{2} W_{B,h+1}^{\lambda,21} + 2i W_{B,h+1}^{\lambda,22} \right. \\
&\quad \left. + 2i W_{F,h+1}^{\lambda,11} - 2\sqrt{2} W_{F,h+1}^{\lambda,21} - 2i W_{F,h+1}^{\lambda,22} \right], \\
\Phi_1^{(h),34} &= q^{1-h} (-4)^{h-3} \left[- 2i W_{B,h+1}^{\lambda,11} + \sqrt{2} W_{B,h+1}^{\lambda,12} + 2i W_{B,h+1}^{\lambda,22} \right. \\
&\quad \left. + 2i W_{F,h+1}^{\lambda,11} - 2\sqrt{2} W_{F,h+1}^{\lambda,21} - 2i W_{F,h+1}^{\lambda,22} \right], \tag{2.25}
\end{aligned}$$

where the bosonic currents are given by (2.2).

The next components [18] with an extra minus sign can be determined by calculating the OPEs between the supersymmetry currents and the previous components (2.25)

$$\begin{aligned}
\tilde{\Phi}_{\frac{3}{2}}^{(h),1} &\equiv \Phi_{\frac{3}{2}}^{(h),1} - \frac{1}{(2h+1)} (1-4\lambda) \partial \Phi_{\frac{1}{2}}^{(h),1} \\
&= q^{-h} (-4)^{h-3} \left[- \frac{1}{2} (Q_{h+\frac{3}{2}}^{\lambda,11} + i\sqrt{2} Q_{h+\frac{3}{2}}^{\lambda,12} + 2i\sqrt{2} Q_{h+\frac{3}{2}}^{\lambda,21} - 2 Q_{h+\frac{3}{2}}^{\lambda,22} \right. \\
&\quad \left. - 2 \bar{Q}_{h+\frac{3}{2}}^{\lambda,11} - 2i\sqrt{2} \bar{Q}_{h+\frac{3}{2}}^{\lambda,12} - i\sqrt{2} \bar{Q}_{h+\frac{3}{2}}^{\lambda,21} + \bar{Q}_{h+\frac{3}{2}}^{\lambda,22}) \right], \\
\tilde{\Phi}_{\frac{3}{2}}^{(h),2} &\equiv \Phi_{\frac{3}{2}}^{(h),2} - \frac{1}{(2h+1)} (1-4\lambda) \partial \Phi_{\frac{1}{2}}^{(h),2} \\
&= q^{-h} (-4)^{h-3} \left[\frac{i}{2} (Q_{h+\frac{3}{2}}^{\lambda,11} + 2i\sqrt{2} Q_{h+\frac{3}{2}}^{\lambda,21} - 2 Q_{h+\frac{3}{2}}^{\lambda,22} - 2 \bar{Q}_{h+\frac{3}{2}}^{\lambda,11} - 2i\sqrt{2} \bar{Q}_{h+\frac{3}{2}}^{\lambda,12} + \bar{Q}_{h+\frac{3}{2}}^{\lambda,22}) \right], \\
\tilde{\Phi}_{\frac{3}{2}}^{(h),3} &\equiv \Phi_{\frac{3}{2}}^{(h),3} - \frac{1}{(2h+1)} (1-4\lambda) \partial \Phi_{\frac{1}{2}}^{(h),3} \\
&= q^{-h} (-4)^{h-3} \left[\frac{i}{2} (Q_{h+\frac{3}{2}}^{\lambda,11} + i\sqrt{2} Q_{h+\frac{3}{2}}^{\lambda,12} - 2 Q_{h+\frac{3}{2}}^{\lambda,22} - 2 \bar{Q}_{h+\frac{3}{2}}^{\lambda,11} - i\sqrt{2} \bar{Q}_{h+\frac{3}{2}}^{\lambda,21} + \bar{Q}_{h+\frac{3}{2}}^{\lambda,22}) \right],
\end{aligned}$$

$$\begin{aligned}\tilde{\Phi}_{\frac{3}{2}}^{(h),4} &\equiv \Phi_{\frac{3}{2}}^{(h),4} - \frac{1}{(2h+1)} (1-4\lambda) \partial \Phi_{\frac{1}{2}}^{(h),4} \\ &= q^{-h} (-4)^{h-3} \left[\frac{1}{2} (Q_{h+\frac{3}{2}}^{\lambda,11} + 2 Q_{h+\frac{3}{2}}^{\lambda,22} + 2 \bar{Q}_{h+\frac{3}{2}}^{\lambda,11} + \bar{Q}_{h+\frac{3}{2}}^{\lambda,22}) \right],\end{aligned}\quad (2.26)$$

which are quasiprimary under the stress energy tensor (2.4). The eight fermionic currents for fixed h in (2.2) are distributed into (2.24) and (2.26) where h is replaced by $(h-1)$.

The last component [18] with an extra minus sign of (2.22) obtained by the OPE between the supersymmetry currents and the components in (2.26), which is quasiprimary under the stress energy tensor (2.4) is

$$\tilde{\Phi}_2^{(h)} \equiv \Phi_2^{(h)} - \frac{1}{(2h+1)} (1-4\lambda) \partial^2 \Phi_0^{(h)} = -q^{-h} (-4)^{h-3} \left[2 (W_{B,h+2}^{\lambda,\bar{a}a} + W_{F,h+2}^{\lambda,\bar{a}a}) \right]. \quad (2.27)$$

Again, the bosonic currents are given by (2.2). The eight bosonic currents for fixed h in (2.2) are distributed into (2.23), (2.25) where h is replaced by $(h-1)$ and (2.27) where h is replaced by $(h-2)$.

It is straightforward, from (2.20) and (A.6) or (2.18) and (A.3), to calculate all the (anti)commutators or OPEs between the $\mathcal{N}=4$ multiplet explicitly. For example, the commutator between the lowest components is described by ¹¹

$$\begin{aligned}& \left[(\Phi_0^{(h_1)})_m, (\Phi_0^{(h_2)})_n \right] = C_{0,0}^{h_1,h_2}(\lambda) [m+h_1-1]_{h_1+h_2-1} \delta_{m+n} \\ &+ \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} \frac{(-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-4}}{(2h_1-1)(2h_2-1)} (2h_3-1)! \\ & \times \left(\left[(h_1-2\lambda)(h_2-2\lambda) S_{F,R}^{h_1,h_2,h_3,k} - (h_1-1+2\lambda)(h_2-1+2\lambda) S_{B,R}^{h_1,h_2,h_3,k} \right] \right. \\ & \times [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \\ & - \left. \left[(h_1-2\lambda)(h_2-2\lambda) S_{F,L}^{h_1,h_2,h_3,k} - (h_1-1+2\lambda)(h_2-1+2\lambda) S_{B,L}^{h_1,h_2,h_3,k} \right] \right. \\ & \times [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \Bigg) (\Phi_0^{(h_3)})_{m+n} \\ &+ \sum_{h_3=-1}^{h_1+h_2-3} \sum_{k=0}^{h_1+h_2-h_3-3} \frac{(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-5}}{2(2h_1-1)(2h_2-1)} (2h_3+2)! \\ & \times \left(\left[(h_1-2\lambda)(h_2-2\lambda)(h_3+1+2\lambda) S_{F,R}^{h_1,h_2,h_3+2,k} \right] \right)\end{aligned}$$

¹¹For simplicity, we do not include q dependence explicitly. Or we consider $q=1$ case.

$$\begin{aligned}
& + (h_1 - 1 + 2\lambda)(h_2 - 1 + 2\lambda)(h_3 + 2 - 2\lambda) S_{B,R}^{h_1,h_2,h_3+2,k} \Big] \\
& \times [m + h_1 - 1]_{h_1+h_2-h_3-k-3} [n + h_2 - 1]_k \\
& - \left[(h_1 - 2\lambda)(h_2 - 2\lambda)(h_3 + 1 + 2\lambda) S_{F,L}^{h_1,h_2,h_3+2,k} \right. \\
& + (h_1 - 1 + 2\lambda)(h_2 - 1 + 2\lambda)(h_3 + 2 - 2\lambda) S_{B,L}^{h_1,h_2,h_3+2,k} \Big] \\
& \times [m + h_1 - 1]_k [n + h_2 - 1]_{h_1+h_2-h_3-k-3} \Bigg) (\tilde{\Phi}_2^{(h_3)})_{m+n}, \tag{2.28}
\end{aligned}$$

where the mode independent central term is given by ¹²

$$\begin{aligned}
C_{0,0}^{h_1,h_2}(\lambda) = & \frac{2N(-1)^{h_1+1} 4^{2(h_1+h_2-4)}}{(2h_1-1)(2h_2-1)} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \frac{i_1! i_2!}{(i_1+i_2+1)!} \\
& \times \left[(h_1 - 2\lambda)(h_2 - 2\lambda) a^{i_1}(h_1, \lambda + \tfrac{1}{2}) a^{i_2}(h_2, \lambda + \tfrac{1}{2}) \right. \\
& \left. - (h_1 - 1 + 2\lambda)(h_2 - 1 + 2\lambda) a^{i_1}(h_1, \lambda) a^{i_2}(h_2, \lambda) \right]. \tag{2.29}
\end{aligned}$$

We can easily check that if we interchange between $h_1 \leftrightarrow h_2$ and $m \leftrightarrow n$, the right hand side of (2.28) becomes the original one with an extra minus sign, by using the definitions in (A.4) together with (A.5). The combination of $(h_1 + h_2)$ appears and the h_1 dependent factor and its h_2 dependent factor occur simultaneously. The corresponding OPE for (2.28) can be found in (C.1). From (2.23) and (2.27), the commutators between the first component and the last component consist of the ones between the bosonic currents in (2.2). This implies that the bosonic subalgebras of the $\mathcal{N} = 4$ supersymmetric linear $W_\infty[\lambda]$ algebra arises from the linear combination of (2.23) and (2.27).

The remaining 14 (anti)commutators are presented in (B.1). The complete algebra between the $\mathcal{N} = 4$ multiplet in (2.22) for generic deformation parameter λ for any conformal weights h_1 and h_2 where $h_1, h_2 = 1, 2, 3, \dots$ is summarized by (2.28) and (B.1).

¹² We obtain the corresponding OPE as follows. The central term has the additional sign factor $(-1)^{h_1+h_2-1}$ from the falling Pochhammer symbol. Moreover, the derivative term $\partial_z^{h_1+h_2-1} \left[\frac{1}{(z-w)} \right]$ should be multiplied. For the two falling Pochhammer symbols in (2.28), we should transform as $(-1)^{h_1+h_2-h_3-1} \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k$ where the power of minus sign comes from the upper bound of k in the summation and the corresponding operator appears as $\frac{\Phi_0^{(h_3)}(w)}{(z-w)}$ on the right hand side. Other terms can be done similarly.

2.4 The $\mathcal{N} = 4$ supersymmetric OPE in $\mathcal{N} = 4$ superspace

By using the $\mathcal{N} = 4$ supersymmetry, the above algebra (2.28) and (B.1) can be written in terms of the following $\mathcal{N} = 4$ super OPE¹³

$$\begin{aligned}
\Phi^{(h_1)}(Z_1) \Phi^{(h_2)}(Z_2) &= \frac{1}{z_{12}^{h_1+h_2}} C_{0,0}^{h_1,h_2}(\lambda) (h_1 + h_2 - 1)! \\
&+ \frac{\theta^{4-0}}{z_{12}^{h_1+h_2+2}} \left[(-1)^{h_1+h_2} C_{0,2}^{h_2,h_1}(\lambda) + \frac{(1-4\lambda)}{(2h_1+1)} C_{0,0}^{h_1,h_2}(\lambda) \right] (h_1 + h_2 + 1)! + \sum_{p=1}^{h_1+h_2-1} \frac{1}{z_{12}^p} \\
&\times \left[\sum_{h_3=1}^{h_1+h_2-p} f_1^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p} \Phi^{(h_3)} - \frac{1}{256} f_2^{h_1,h_2,1,p}(\lambda) \partial^{h_1+h_2-p} \mathbf{J} \right. \\
&+ \sum_{h_3=2}^{h_1+h_2-p} f_2^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p} \left(D^{4-0} - \frac{(1-4\lambda)}{(2h_3-3)} \partial^2 \right) \Phi^{(h_3-2)} \Big] (Z_2) \\
&+ \sum_{p=1}^{h_1+h_2} \frac{\theta^i}{z_{12}^p} \left[\sum_{h_3=0}^{h_1+h_2-p} f_3^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p} D^i \right. \\
&+ \sum_{h_3=0}^{h_1+h_2-p-1} f_4^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p-1} \left(D^{4-i} + \frac{(1-4\lambda)}{(2h_3+1)} \partial D^i \right) \Big] \Phi^{(h_3)}(Z_2) \\
&+ \sum_{p=1}^{h_1+h_2} \frac{\theta^{4-ij}}{z_{12}^p} \left[\sum_{h_3=0}^{h_1+h_2-p} f_5^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p} D^{4-ij} \right. \\
&+ \sum_{h_3=0}^{h_1+h_2-p} f_6^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p} D^{ij} \Big] \Phi^{(h_3)}(Z_2) \\
&+ \sum_{p=1}^{h_1+h_2+1} \frac{\theta^{4-i}}{z_{12}^p} \left[\sum_{h_3=0}^{h_1+h_2-p} f_7^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p} \left(D^{4-i} + \frac{(1-4\lambda)}{(2h_3+1)} \partial D^i \right) \right. \\
&+ \sum_{h_3=0}^{h_1+h_2-p+1} f_8^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p+1} \partial D^i \Big] \Phi^{(h_3)}(Z_2) \\
&+ \sum_{p=1}^{h_1+h_2+1} \frac{\theta^{4-0}}{z_{12}^p} \left[\sum_{h_3=2}^{h_1+h_2-p+2} f_9^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p+2} \left(D^{4-0} - \frac{(1-4\lambda)}{(2h_3-3)} \partial^2 \right) \Phi^{(h_3-2)} \right. \\
&- \frac{1}{256} f_9^{h_1,h_2,1,p}(\lambda) \partial^{h_1+h_2-p+2} \mathbf{J} + \sum_{h_3=1}^{h_1+h_2-p+2} f_{10}^{h_1,h_2,h_3,p}(\lambda) \partial^{h_1+h_2-h_3-p+2} \Phi^{(h_3)} \Big] (Z_2)
\end{aligned}$$

¹³ The $\mathcal{N} = 4$ super OPE between the stress energy tensor is

$$\mathbf{J}(Z_1) \mathbf{J}(Z_2) = -\frac{\theta_{12}^{4-0}}{z_{12}^2} N + \frac{\theta_{12}^{4-i}}{z_{12}} D^i \mathbf{J}(Z_2) + \frac{\theta_{12}^{4-0}}{z_{12}} 2 \partial \mathbf{J}(Z_2) + \dots$$

The $\mathcal{N} = 4$ super OPE between the $\mathcal{N} = 4$ stress energy tensor and the $\mathcal{N} = 4$ multiplet is presented in (C.3).

$$+ \dots \quad (2.30)$$

The central term comes from the two contributions in (C.1). Depending on the number of Grassmann coordinates, each component OPE in (C.1) is located at the right place. The conformal weight of $(h_1 + h_2)$ on the left hand side is satisfied also on the right hand side. Note that there is a relation $\Phi^{(0)} \equiv \frac{1}{64} \mathbf{J}$. The \mathbf{J} dependent terms arise on the right hand side of (2.30). The various structure constants are given by (B.2) and (C.2)¹⁴. The upper bound for h_3 summation comes from the vanishing of the derivatives. Similarly the upper bound for p summation comes from the fact that the conformal weight is given by $(h_1 + h_2)$. There are $(1 - 4\lambda)$ factors in (2.30) where the previous central term (2.5) associated with the stress energy tensor vanishes at $\lambda = \frac{1}{4}$. The result of (2.30) is a generalization of [19] where the $h_1 + h_2 \leq 4$ cases are obtained. At $\lambda = 0$, the previous results in [18] are reproduced.

Therefore, the above expression (2.30) is equivalent to the component results in (2.28) and (B.1).

3 The $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda]$ algebra

3.1 The singlet currents and the adjoint currents

By multiplying the generators of $SU(K)$ $t_{a\bar{b}}^{\hat{A}}$ and $\delta_{a\bar{b}}$ into (2.2)¹⁵, we construct the following singlets and adjoint currents [17]

$$\begin{aligned} W_{F,h} &\equiv W_{F,h}^{\bar{a}b} \delta_{b\bar{a}}, & W_{F,h}^{\hat{A}} &\equiv W_{F,h}^{\bar{a}b} t_{b\bar{a}}^{\hat{A}}, & W_{B,h} &\equiv W_{B,h}^{\bar{a}b} \delta_{b\bar{a}}, & W_{B,h}^{\hat{A}} &\equiv W_{B,h}^{\bar{a}b} t_{b\bar{a}}^{\hat{A}}, \\ Q_{h+\frac{1}{2}} &\equiv Q_{h+\frac{1}{2}}^{\bar{a}b} \delta_{b\bar{a}}, & Q_{h+\frac{1}{2}}^{\hat{A}} &\equiv Q_{h+\frac{1}{2}}^{\bar{a}b} t_{b\bar{a}}^{\hat{A}}, & \bar{Q}_{h+\frac{1}{2}} &\equiv \bar{Q}_{h+\frac{1}{2}}^{a\bar{b}} \delta_{a\bar{b}}, & \bar{Q}_{h+\frac{1}{2}}^{\hat{A}} &\equiv \bar{Q}_{h+\frac{1}{2}}^{a\bar{b}} t_{a\bar{b}}^{\hat{A}}, \end{aligned} \quad (3.1)$$

where $\hat{A} = 1, 2, \dots, (K^2 - 1)$. It is straightforward to calculate the OPEs between the singlets and the adjoints by using the OPEs appearing in section 2.

3.2 The OPEs or (anti)commutators between the singlets and the adjoints

The OPE between the singlets, from (3.1) and (2.18), can be summarized by

$$W_{F,h_1}^\lambda(z) W_{F,h_2}^\lambda(w) = K c_{W_F}^{h_1, h_2}(\lambda) q^{h_1+h_2-4} (-1)^{h_1+h_2+1} \partial_z^{h_1+h_2-1} \left[\frac{1}{(z-w)} \right]$$

¹⁴If the upper bound is less than the lower bound in the summation notation appearing in (2.30), we treat them to be 0 according to the “empty” sum.

¹⁵In this section, the previous $SU(2)$ is replaced by $SU(K)$.

$$\begin{aligned}
& + \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
& \times \left(S_{F,R}^{h_1,h_2,h_3,k} \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k - S_{F,L}^{h_1,h_2,h_3,k} \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \right) \left[\frac{W_{F,h_3}^\lambda(w)}{(z-w)} \right] + \dots \quad (3.2)
\end{aligned}$$

Due to the contractions between the group indices, the field dependent term appears as a common factor. The relations (2.21), (2.17) and (2.19) are assumed.

With one adjoint index, the following OPE satisfies

$$\begin{aligned}
W_{F,h_1}^\lambda(z) W_{F,h_2}^{\lambda,\hat{A}}(w) & = \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
& \times \left(S_{F,R}^{h_1,h_2,h_3,k} \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k - S_{F,L}^{h_1,h_2,h_3,k} \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \right) \left[\frac{W_{F,h_3}^{\lambda,\hat{A}}(w)}{(z-w)} \right] + \dots \quad (3.3)
\end{aligned}$$

There is no central term and the expression (3.3) is the same as the one without a central term in (3.2) except the adjoint index appearing on the current.

With two adjoint indices, we obtain

$$\begin{aligned}
W_{F,h_1}^{\lambda,\hat{A}}(z) W_{F,h_2}^{\lambda,\hat{B}}(w) & = \delta^{\hat{A}\hat{B}} c_{W_F}^{h_1,h_2}(\lambda) q^{h_1+h_2-4} (-1)^{h_1+h_2+1} \partial_z^{h_1+h_2-1} \left[\frac{1}{(z-w)} \right] \\
& + \frac{1}{2} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
& \times \left(\left(S_{F,R}^{h_1,h_2,h_3,k} \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k - S_{F,L}^{h_1,h_2,h_3,k} \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \right) \right. \\
& \times \left[\frac{(\frac{2}{K} \delta^{\hat{A}\hat{B}} W_{F,h_3}^\lambda + d^{\hat{A}\hat{B}\hat{C}} W_{F,h_3}^{\lambda,\hat{C}})(w)}{(z-w)} \right] \\
& + \left(S_{F,R}^{h_1,h_2,h_3,k} \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k + S_{F,L}^{h_1,h_2,h_3,k} \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \right) \\
& \left. \times \left[\frac{i f^{\hat{A}\hat{B}\hat{C}} W_{F,h_3}^{\lambda,\hat{C}}(w)}{(z-w)} \right] \right) + \dots \quad (3.4)
\end{aligned}$$

On the right hand side of (3.4), there are singlets and adjoints. Because we are dealing with $K \geq 3$, there exists a nontrivial symmetric d symbol.

We present some OPEs between the singlets where $h_1 + h_2 \leq 6$ corresponding to (3.2) as follows:

$$W_{F,1}^\lambda(z) W_{F,1}^\lambda(w) = \frac{1}{(z-w)^2} \frac{N K}{16 q^2} + \dots ,$$

$$\begin{aligned}
W_{F,1}^\lambda(z) W_{F,2}^\lambda(w) &= \frac{1}{(z-w)^3} \frac{NK\lambda}{2q} + \frac{1}{(z-w)^2} W_{F,1}^\lambda(w) + \dots, \\
W_{F,1}^\lambda(z) W_{F,3}^\lambda(w) &= \frac{1}{(z-w)^4} 2NK\lambda^2 + \frac{1}{(z-w)^3} 8q\lambda W_{F,1}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^2} \left[2W_{F,2}^\lambda - 4q\lambda \partial_w W_{F,1}^\lambda \right] (w) + \dots, \\
W_{F,2}^\lambda(z) W_{F,2}^\lambda(w) &= \frac{1}{(z-w)^4} \frac{NK(1-12\lambda^2)}{2} + \frac{1}{(z-w)^2} 2W_{F,2}^\lambda(w) \\
&\quad + \frac{1}{(z-w)} \partial_w W_{F,2}^\lambda(w) + \dots, \\
W_{F,2}^\lambda(z) W_{F,3}^\lambda(w) &= \frac{1}{(z-w)^5} 8qNK\lambda(1-4\lambda^2) \\
&\quad + \frac{1}{(z-w)^4} 16q^2(1-4\lambda^2) W_{F,1}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^2} 3W_{F,3}^\lambda(w) + \frac{1}{(z-w)} \partial_w W_{F,3}^\lambda(w) + \dots, \\
W_{F,3}^\lambda(z) W_{F,3}^\lambda(w) &= \frac{1}{(z-w)^6} \frac{32q^2NK}{3}(1-15\lambda^2+20\lambda^4) \\
&\quad + \frac{1}{(z-w)^4} 64q^2(1-\lambda^2) W_{F,2}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^3} 32q^2(1-\lambda^2) \partial_w W_{F,2}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^2} \left[4W_{F,4}^\lambda + \frac{48q^2}{5}(1-\lambda^2) \partial_w^2 W_{F,2}^\lambda \right] (w) \\
&\quad + \frac{1}{(z-w)} \left[2\partial W_{F,4}^\lambda + \frac{32q^2}{15}(1-\lambda^2) \partial_w^3 W_{F,2}^\lambda \right] (w) + \dots. \tag{3.5}
\end{aligned}$$

Compared to the ones in [4], the above OPEs (3.5) becomes the corresponding expressions appearing in [4] with some typo (in the last expression of (3.38) in [4], the numerical value $\frac{1}{6}$ should be 1) when we take $\lambda \rightarrow 0$ limit with $q = \frac{1}{4}$. The NK corresponds to their central term. In other words, the $W_{1+\infty}^K[\lambda = 0]$ algebra is realized. We also add the third and the last ones in (3.5). We present other OPEs in Appendix D corresponding to (3.3) and (3.4)¹⁶.

¹⁶ At $\lambda = \pm\frac{1}{2}$, we can easily see that the last three OPEs in (3.5) and the next OPE below will lead to the ones in (3.21) of [4] which are some OPEs in the W_∞ algebra. For convenience, we present the following OPE

$$\begin{aligned}
W_{F,2}^\lambda(z) W_{F,4}^\lambda(w) &= \frac{1}{(z-w)^6} 32q^2NK\lambda^2(1-4\lambda^2) + \frac{1}{(z-w)^5} 128q^3\lambda(1-4\lambda^2) W_{F,1}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^4} \left[\frac{48q^2}{5}(7-12\lambda^2) W_{F,2}^\lambda - 64q^3\lambda(1-4\lambda^2) \partial_w W_{F,1}^\lambda \right] (w) + \frac{1}{(z-w)^2} 4W_{F,4}^\lambda(w)
\end{aligned}$$

In (D.1) where the three commutators (3.2), (3.3) and (3.4) are also added for convenience, the complete (anti)commutators for the $\mathcal{N} = 2$ supersymmetric $W_\infty^{K,K}[\lambda]$ algebra are presented.

In particular, at $\lambda = 0$, the above three OPEs, (3.2), (3.3) and (3.4), or the first three commutators appearing in (D.1), lead to the one in [8]. In this sense, these for nonzero λ are the λ deformed $W_{1+\infty}$ algebra (or $\hat{SU}(K)_{k=N}$ algebra) which is denoted as $W_{1+\infty}^K[\lambda]$ algebra in the footnote 2 of the introduction. By taking the contractions of the currents with vanishing q limit for $\lambda = 0$, as noted by [26], the $w_{1+\infty}$ algebra (which is the extension of w_∞ algebra [1]) is reproduced from (3.2). In the OPE language, there are only second and first order poles in (3.5) under this limit by ignoring the central terms¹⁷. For $h_1 = 1$, there exists only second order pole.

We can analyze the similar OPEs or (anti)commutators corresponding to the bosonic subalgebra coming from the (β, γ) fields. By comparing (2.20) with the first relation of (A.6) together with (A.5), the three OPEs corresponding to (3.2), (3.3) and (3.4) can be obtained by changing $\lambda \rightarrow \lambda - \frac{1}{2}$ in the structure constants and the central terms¹⁸. There exists an extra minus sign in the central term. These for nonzero λ are the λ deformed $W_{1+\infty}$ algebra which is denoted as $W_{1+\infty}^K[\lambda + \frac{1}{2}]$ algebra in the footnote 2 of the introduction.

Therefore, the bosonic subalgebra of $\mathcal{N} = 2$ supersymmetric linear $W_\infty[\lambda]$ algebra is described by $W_{1+\infty}^K[\lambda] \times W_{1+\infty}^K[\lambda + \frac{1}{2}]$. Note that the transformations described in the footnote 3 can be applied here. The point is that the (b, c) fields realization is applied in the first factor while the (β, γ) fields realization is applied in the second factor. We realize that the $W_{1+\infty}$ algebra is realized by both fermionic and bosonic constructions. The deformation parameter λ appears in our construction and plays the role in each bosonic subalgebra with a little different form. The first one is described by λ while the second one is described by $(\lambda + \frac{1}{2})$. Therefore, we have found a matrix generalization of [10, 11] with explicit forms for the structure constants

$$+ \frac{1}{(z-w)} \partial_w W_{F,4}^\lambda(w) + \dots$$

In this OPE, at $\lambda = \pm \frac{1}{2}$, the spin 1 current does not appear due to the $(1 - 4\lambda^2)$ factor. This also happens for the OPE between the spin 2 and the spin 3 of (3.5). We can check that this OPE reduces to the one in [4] at $\lambda = 0$ with $q = \frac{1}{4}$. Therefore, in our notation, the $W_\infty^K[\lambda = \pm \frac{1}{2}]$ algebra is obtained. We have checked that the structure constant in (D.5) satisfies $P_F^{h_1, h_2, 1}(m, n, \pm \frac{1}{2}) = 0$ for $h_1, h_2 > 1$. This implies that there is a decoupling of $W_{F,1}^\lambda(w)$ at $\lambda = \pm \frac{1}{2}$ in the full OPEs.

¹⁷Inside Thielemans package [21], the command `SetOPEOptions[OPEMethod, ClassicalOPEs]`; provides no double contractions and this leads to the vanishing central terms in the OPEs.

¹⁸We observe that at $\lambda = \frac{1}{2}$, the corresponding $W_{1+\infty}^K[\lambda + \frac{1}{2} = 1]$ algebra can be reproduced. Similarly, the $W_\infty^K[\lambda + \frac{1}{2} = \frac{1}{2}]$ algebra is obtained at $\lambda = 0$. As in the footnote 16, the structure constant in (D.5) where $S_{F,R}^{h_1, h_2, h_3, k}(\lambda)$ is replaced by $S_{B,R}^{h_1, h_2, h_3, k}(\lambda)$ (and $S_{F,L}^{h_1, h_2, h_3, k}(\lambda)$ is replaced by $S_{B,L}^{h_1, h_2, h_3, k}(\lambda)$) satisfies $P_B^{h_1, h_2, 1}(m, n, 0) = 0$ for $h_1, h_2 > 1$. There is a decoupling of $W_{B,1}^\lambda(w)$ at $\lambda = 0$ in the full OPEs.

4 Conclusions and outlook

In this paper, we have found the OPEs or (anti)commutators between the bifundamentals in (2.18) and (A.3) (or (2.20) and (A.6)). In deriving these, it was crucial to prove (2.8) and (A.1). Based on these results, the $\mathcal{N} = 4$ supersymmetric $W_\infty^{2,2}[\lambda]$ algebra is obtained by (2.28) and (B.1) (or (2.30)). Finally, the $\mathcal{N} = 2$ supersymmetric $W_\infty^{K,K}[\lambda]$ algebra is determined by (D.1). In arriving at these two algebras, the above OPEs or (anti)commutators between the bifundamentals are heavily used. One of the main results in this paper, as mentioned in the abstract, is a generalization of $W_{1+\infty}$ algebra found by [5] by the presence of the deformation parameter λ . Originally, this parameter appears in the conformal weights in the (β, γ) and (b, c) ghost systems (See also the footnote 3) and plays the role of the rank of the interaction in the two dimensional $\mathcal{N} = (2, 2)$ SYK model. See also the footnote 5.

It is an open problem to describe the present results in the $\mathcal{N} = 2$ superspace starting from the commuting B field and an anticommuting C field [10, 11]. It is known that the

¹⁹ As done in [11] where the λ is replaced by $(\frac{1}{2} - \lambda)$, for the transformation (with multiple numbers of each free field) of

$$c^{l\bar{a}} \mapsto \epsilon_1 \delta^l_{\bar{l}} \delta^{\bar{a}}_b \beta^{\bar{l}b}, \quad \beta^{\bar{l}a} \mapsto -i \epsilon_2 \delta^{\bar{l}}_l \delta^a_{\bar{b}} c^{l\bar{b}} \quad \gamma^{l\bar{a}} \mapsto -\epsilon_1 \delta^l_{\bar{l}} \delta^{\bar{a}}_b b^{\bar{l}b}, \quad b^{\bar{l}a} \mapsto i \epsilon_2 \delta^{\bar{l}}_l \delta^a_{\bar{b}} \gamma^{l\bar{b}}, \quad (3.6)$$

where ϵ_1 and ϵ_2 are real anticommuting parameters, the corresponding currents are transformed as follows

$$\begin{aligned} W_{F,h}^{\lambda,\bar{a}b}(z) &\mapsto (-1)^h i(\epsilon_1 \epsilon_2) W_{B,h}^{\frac{1}{2}-\lambda,b\bar{a}}(z), & W_{B,h}^{\lambda,\bar{a}b}(z) &\mapsto (-1)^h i(\epsilon_1 \epsilon_2) W_{F,h}^{\frac{1}{2}-\lambda,b\bar{a}}(z), \\ Q_{h+\frac{1}{2}}^{\lambda,\bar{a}b}(z) &\mapsto (-1)^{h+1} i(\epsilon_1 \epsilon_2) Q_{h+\frac{1}{2}}^{\frac{1}{2}-\lambda,b\bar{a}}(z), & \bar{Q}_{h+\frac{1}{2}}^{\lambda,a\bar{b}}(z) &\mapsto (-1)^{h+1} i(\epsilon_1 \epsilon_2) \bar{Q}_{h+\frac{1}{2}}^{\frac{1}{2}-\lambda,\bar{b}a}(z). \end{aligned} \quad (3.7)$$

Note that the group indices are reversed after transformation. We can make the above transformations (3.7) into the previous OPE (2.18) and it turns out that we obtain the singular terms of the OPE of $(-1)^{h_1} W_{B,h_1}^{\frac{1}{2}-\lambda,b\bar{a}}(z) (-1)^{h_2+1} Q_{h_2+\frac{1}{2}}^{\frac{1}{2}-\lambda,d\bar{c}}(w)$. On the other hand, we can extract this OPE from the first equation of (A.3). The singular terms are exactly the same as the previous one and the central term has opposite sign due to the different statistics in (3.6). Similarly, the singular terms for the OPEs $(-1)^{h_1} W_{B,h_1}^{\frac{1}{2}-\lambda,b\bar{a}}(z) (-1)^{h_2+1} Q_{h_2+\frac{1}{2}}^{\frac{1}{2}-\lambda,d\bar{c}}(w)$, $(-1)^{h_1} W_{B,h_1}^{\frac{1}{2}-\lambda,b\bar{a}}(z) (-1)^{h_2+1} \bar{Q}_{h_2+\frac{1}{2}}^{\frac{1}{2}-\lambda,\bar{d}c}(w)$ and $(-1)^{h_1+1} Q_{h_1+\frac{1}{2}}^{\frac{1}{2}-\lambda,\bar{a}b}(z) (-1)^{h_2+1} \bar{Q}_{h_1+\frac{1}{2}}^{\frac{1}{2}-\lambda,c\bar{d}}(w)$ can be determined from the second, the third, and the last in (A.3) by using the transformations (3.7) respectively. Or we can extract them from the fourth, the fifth and the last OPEs in (A.3). The singular terms are the same. For the last one, the central term has opposite sign as before (3.6). This analysis implies that the first, the fourth and the fifth of (A.3) are redundant because they can be determined from (2.18), the second and the third of (A.3) up to the central term. Recall that the enveloping algebra of $OSp(1, 2)$ in an associative algebra between the wedge generators is factored over the ideal in [11]. The quadratic Casimir operator of $OSp(1, 2)$ is given by $\lambda(\lambda - \frac{1}{2})$ which is invariant under the above transformation $\lambda \rightarrow \frac{1}{2} - \lambda$. The linear combination described by the Casimir is proportional to the identity operator. The multiplication properties of this algebra depends on λ via the Casimir operator value $\lambda(\lambda - \frac{1}{2})$. We thank the referee for raising this point.

bosonic W_∞ algebra with $\lambda = 0$ can be obtained from the bosonic $W_{1+\infty}$ algebra with $\lambda = 0$ by decoupling the spin 1 current. It is an open problem to observe whether that kind of truncation for the nonzero λ , by introducing the new currents of spins from the old currents of spins via (non)linear combinations, appears or not. It would be interesting to find how the large (N, k) 't Hooft like limit in the $\mathcal{N} = 4$ coset model appears in the present result for any h_1 and h_2 [19].

Recently, the wedge subalgebra of $w_{1+\infty}$ algebra [1] has been identified with the symmetries on the celestial sphere [27, 28] in the context of celestial conformal field theory [29]. See also relevant works in [30, 31, 32, 33, 34, 35, 36, 37]²⁰. The $\mathcal{N} = 1$ supersymmetric $w_{1+\infty}$ algebra plays the important role in the corresponding $\mathcal{N} = 1$ supersymmetric Einstein-Yang-Mills theory [26, 17]. It would be interesting to find whether the results of this paper provide the corresponding celestial conformal field theories or not, by taking the proper limits on the two parameters q and λ . More explicitly, can we see the mode dependent structure constants in the (anti)commutators from the celestial conformal field theories?

Acknowledgments

CA thanks M. Pate for general discussions on the celestial holography. This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (No. 2023R1A2C1003750).

²⁰Moreover, in [38], the chiral algebra [39] from the $\mathcal{N} = 2$ gauge theories in four dimensions is studied in the context of twisted holography. See also the relevant work in [40]. We thank the referee for pointing this out. In particular, the small $\mathcal{N} = 4$ superconformal algebra [41, 42] is reproduced in [39]. On the other hand, when $\lambda = 0$, the (A.1) of [19] (which is the component results corresponding to the expression in the footnote 13) contains the equations of (2.1), (2.2) and (2.3) of [43], as a subalgebra, which describe the above small $\mathcal{N} = 4$ superconformal algebra. In their construction [39], the conformal weights for (β, γ) and (b, c) are given by $\frac{1}{2}, \frac{1}{2}, 1$ and 0 respectively. This implies the case of $\lambda = \frac{1}{2}$. Then by taking the symmetry between λ and $(\frac{1}{2} - \lambda)$ described in the footnote 19, we arrive at the previous $\lambda = 0$ case. It would be interesting to see whether their construction provides the extension of small $\mathcal{N} = 4$ superconformal algebra or not.

A The remaining OPEs or (anti)commutators between the bifundamentals

The other three identities in the context of (2.8) can be summarized by

$$\begin{aligned} ((\partial_z^s \beta^{\bar{l}\bar{b}}) \delta_{\bar{l}\bar{l}} \gamma^{l\bar{a}})(z) &= \sum_{h=0}^s (-1)^{s+1} 4^{-h} q^{1-h} Y_{W_B}(s-h, s+1, \lambda) \partial_z^{s-h} W_{B,h+1}^{\lambda, \bar{a}b}(z), \\ ((\partial_z^s b^{\bar{l}\bar{b}}) \delta_{\bar{l}\bar{l}} \gamma^{l\bar{a}})(z) &= \sum_{h=0}^s (-1)^s 2^{-\frac{1}{2}} 4^{-h} q^{-h} Y_Q(s-h, s+1, \lambda) \partial_z^{s-h} Q_{h+\frac{3}{2}}^{\lambda, \bar{a}b}(z), \\ ((\partial_z^s \beta^{\bar{l}\bar{b}}) \delta_{\bar{l}\bar{l}} c^{l\bar{a}})(z) &= \sum_{h=0}^s (-1)^{s+1} 2^{\frac{3}{2}} 4^{-h} q^{1-h} Y_{\bar{Q}}(s-h, s+1, \lambda) \partial_z^{s-h} \bar{Q}_{h+\frac{1}{2}}^{\lambda, b\bar{a}}(z). \end{aligned} \quad (\text{A.1})$$

The left hand sides of (A.1) appear in the last three bifundamentals (2.2) without derivatives. Here the coefficients appearing on the right hand sides of (A.1) are

$$\begin{aligned} Y_{W_B}(h, s, \lambda) &\equiv \frac{2(-2\lambda - s + 2)_h [s-1]_{h-1}}{h! [2s-h-1]_{h-1}}, \\ Y_Q(h, s, \lambda) &\equiv \frac{(-2\lambda - s + 1)_h [s-1]_h}{h! [2s-h]_h}, \\ Y_{\bar{Q}}(h, s, \lambda) &\equiv \frac{(-2\lambda - s + 2)_h [s-1]_h}{h! [2s-h-2]_h}. \end{aligned} \quad (\text{A.2})$$

In the similar expressions to (2.6), the identities in (A.1) enable us to express the right hand sides of the corresponding OPEs by using the original bifundamentals (2.2) with possible derivatives.

The remaining OPEs between the bifundamentals, by taking the similar analysis done in the subsection 2.2 together with (A.1) and (A.2), are

$$\begin{aligned} W_{B,h_1}^{\lambda, \bar{a}b}(z) W_{B,h_2}^{\lambda, \bar{c}d}(w) &= \delta^{\bar{a}d} \delta^{\bar{c}b} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} (-1)^{h_1+1} (4q)^{h_1+h_2-4} a^{i_1}(h_1, \lambda) a^{i_2}(h_2, \lambda) \frac{N i_1! i_2!}{(i_1+i_2+1)!} \\ &\quad \times \partial_z^{h_1+h_2-1} \left[\frac{1}{(z-w)} \right] \\ &+ \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\ &\quad \times \left(\delta^{\bar{a}d} S_{B,R}^{h_1, h_2, h_3, k}(\lambda) \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k \left[\frac{W_{B,h_3}^{\lambda, \bar{c}b}(w)}{(z-w)} \right] \right. \\ &\quad \left. - \delta^{\bar{c}b} S_{B,L}^{h_1, h_2, h_3, k}(\lambda) \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \left[\frac{W_{B,h_3}^{\lambda, \bar{a}d}(w)}{(z-w)} \right] \right) + \dots, \end{aligned}$$

$$\begin{aligned}
W_{F,h_1}^{\lambda,\bar{a}b}(z) Q_{h_2+\frac{1}{2}}^{\lambda,\bar{c}d}(w) &= \delta^{\bar{a}d} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
&\quad \times T_F^{h_1,h_2,h_3,k}(\lambda) \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k \left[\frac{Q_{h_3+\frac{1}{2}}^{\lambda,\bar{c}b}(w)}{(z-w)} \right] + \dots, \\
W_{F,h_1}^{\lambda,\bar{a}b}(z) \bar{Q}_{h_2+\frac{1}{2}}^{\lambda,c\bar{d}}(w) &= \delta^{\bar{d}\bar{b}} \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
&\quad \times \bar{T}_F^{h_1,h_2,h_3,k}(\lambda) \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \left[\frac{\bar{Q}_{h_3+\frac{1}{2}}^{\lambda,c\bar{a}}(w)}{(z-w)} \right] + \dots, \\
W_{B,h_1}^{\lambda,\bar{a}b}(z) Q_{h_2+\frac{1}{2}}^{\lambda,\bar{c}d}(w) &= \delta^{\bar{c}\bar{b}} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
&\quad \times T_B^{h_1,h_2,h_3,k}(\lambda) \partial_z^k \partial_w^{h_1+h_2-h_3-k-1} \left[\frac{Q_{h_3+\frac{1}{2}}^{\lambda,\bar{a}d}(w)}{(z-w)} \right] + \dots, \\
W_{B,h_1}^{\lambda,\bar{a}b}(z) \bar{Q}_{h_2+\frac{1}{2}}^{\lambda,c\bar{d}}(w) &= \delta^{\bar{a}c} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
&\quad \times \bar{T}_B^{h_1,h_2,h_3,k}(\lambda) \partial_z^{h_1+h_2-h_3-k-1} \partial_w^k \left[\frac{\bar{Q}_{h_3+\frac{1}{2}}^{\lambda,b\bar{d}}(w)}{(z-w)} \right] + \dots, \\
Q_{h_1+\frac{1}{2}}^{\lambda,\bar{a}b}(z) \bar{Q}_{h_2+\frac{1}{2}}^{\lambda,c\bar{d}}(w) &= \delta^{\bar{a}c} \delta^{\bar{d}b} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2} 2(-1)^{h_1+1} (4q)^{h_1+h_2-2} \beta^{i_1}(h_1+1, \lambda) \alpha^{i_2}(h_2+1, \lambda) \\
&\quad \times \frac{N i_1! i_2!}{(i_1+i_2+1)!} \partial_z^{h_1+h_2} \left[\frac{1}{(z-w)} \right] \\
&\quad + \sum_{h_3=1}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} 2(-1)^{h_1+h_2+1} (4q)^{h_1+h_2-h_3} (2h_3-1)! \\
&\quad \times \left(\delta^{\bar{a}c} U_F^{h_1,h_2,h_3,k}(\lambda) \partial_z^{h_1+h_2-h_3-k} \partial_w^k \left[\frac{W_{F,h_3}^{\lambda,\bar{d}b}(w)}{(z-w)} \right] \right. \\
&\quad \left. + \delta^{\bar{d}b} U_B^{h_1,h_2,h_3,k}(\lambda) \partial_z^k \partial_w^{h_1+h_2-h_3-k} \left[\frac{W_{B,h_3}^{\lambda,\bar{a}c}(w)}{(z-w)} \right] \right) + \dots. \tag{A.3}
\end{aligned}$$

We observe that the powers of derivatives in the second(fourth) OPE appear in the third(fifth) OPE under the exchange of $z \leftrightarrow w$. Compared to the ones for $\lambda = 0$, the algebra given by (2.18) and (A.3) has the summation over the dummy variable k appearing in the structure constants and derivatives. In general, the highest order poles occur when the bosonic currents have the spin 1 or the fermionic currents have the spin $\frac{3}{2}$ (or $\frac{1}{2}$) and the lowest order pole is given by the first order pole. Between them, all the other structures of poles can be read off

from the explicit structure constants and the derivatives.

The structure constants appearing in (A.3) as well as the ones in section 2 are given by

$$\begin{aligned}
S_{F,R}^{h_1,h_2,h_3,k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_2}{i_1+i_2-r} \binom{1-h_3+r}{1-h_2+i_2+k} \prod_{j=0}^{r-h_3} (-r-2\lambda+j) \Big], \\
S_{F,L}^{h_1,h_2,h_3,k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda + \frac{1}{2}) a^{i_2}(h_2, \lambda + \frac{1}{2}) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_1}{i_1+i_2-r} \binom{1-h_3+r}{1-h_1+i_1+k} \prod_{j=0}^{r-h_3} (-r-2\lambda+j) \Big], \\
S_{B,R}^{h_1,h_2,h_3,k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda) a^{i_2}(h_2, \lambda) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_2}{i_1+i_2-r} \binom{1-h_3+r}{1-h_2+i_2+k} \prod_{j=0}^{r-h_3} (1-r-2\lambda+j) \Big], \\
S_{B,L}^{h_1,h_2,h_3,k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda) a^{i_2}(h_2, \lambda) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_1}{i_1+i_2-r} \binom{1-h_3+r}{1-h_1+i_1+k} \prod_{j=0}^{r-h_3} (1-r-2\lambda+j) \Big], \\
T_F^{h_1,h_2,h_3,k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r+1)!} a^{i_1}(h_1, \lambda + \frac{1}{2}) \beta^{i_2}(h_2+1, \lambda) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_2}{i_1+i_2-r} \binom{1-h_3+r}{1-h_2+i_2+k} \prod_{j=0}^{r-h_3} (-r-2\lambda+j) \Big], \\
\bar{T}_F^{h_1,h_2,h_3,k}(h_1, \lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda + \frac{1}{2}) \alpha^{i_2}(h_2+1, \lambda) \right. \\
&\quad \times \binom{r}{h_3} \binom{i_1}{i_1+i_2-r} \binom{-h_3+r}{1-h_1+i_1+k} \prod_{j=0}^{r-h_3-1} (1-r-2\lambda+j) \Big], \\
T_B^{h_1,h_2,h_3,k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r+1)!} a^{i_1}(h_1, \lambda) \beta^{i_2}(h_2+1, \lambda) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_1}{i_1+i_2-r} \binom{1-h_3+r}{1-h_1+i_1+k} \prod_{j=0}^{r-h_3} (-r-2\lambda+j) \Big],
\end{aligned}$$

$$\begin{aligned}
\bar{T}_B^{h_1, h_2, h_3, k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} a^{i_1}(h_1, \lambda) \alpha^{i_2}(h_2+1, \lambda) \right. \\
&\quad \times \binom{r}{h_3} \binom{i_2}{i_1+i_2-r} \binom{-h_3+r}{-h_2+i_2+k} \prod_{j=0}^{r-h_3-1} (1-r-2\lambda+j) \Big], \\
U_F^{h_1, h_2, h_3, k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} \beta^{i_1}(h_1+1, \lambda) \alpha^{i_2}(h_2+1, \lambda) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_2}{i_1+i_2-r} \binom{1-h_3+r}{-h_2+i_2+k} \prod_{j=0}^{r-h_3} (-r-2\lambda+j) \Big], \\
U_B^{h_1, h_2, h_3, k}(\lambda) &= \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2} \sum_{r=0}^{i_1+i_2} \left[\frac{(-1)^{i_1+i_2}}{(h_3+r)!} \beta^{i_1}(h_1+1, \lambda) \alpha^{i_2}(h_2+1, \lambda) \right. \\
&\quad \times \binom{r}{h_3-1} \binom{i_1}{i_1+i_2-r} \binom{1-h_3+r}{1-h_1+i_1+k} \prod_{j=0}^{r-h_3} (1-r-2\lambda+j) \Big]. \tag{A.4}
\end{aligned}$$

The k dependence appears in one of the binomials and the dummy variables i_1, i_2 and r appear in various places. The λ dependence appears in the numerators.

Note that there are some relations, from the symmetries of bosonic subalgebra,

$$S_{B,R}^{h_1, h_2, h_3, k}(\lambda) = S_{F,R}^{h_1, h_2, h_3, k}(\lambda - \tfrac{1}{2}), \quad S_{B,L}^{h_1, h_2, h_3, k}(\lambda) = S_{F,L}^{h_1, h_2, h_3, k}(\lambda - \tfrac{1}{2}). \tag{A.5}$$

This can be understood by changing the conformal weights $(\tfrac{1}{2} + \lambda, \tfrac{1}{2} - \lambda)$ for (b, c) fields into $(\tfrac{1}{2} + \lambda - \tfrac{1}{2}, \tfrac{1}{2} - \lambda + \tfrac{1}{2})$ which are the ones for (β, γ) fields. See also the footnote 3. Then the OPE or commutator between the currents $W_{B,h}^{\lambda, \bar{a}b}$ can be obtained also from the one between the currents $W_{F,h}^{\lambda, \bar{a}b}$ with the relation (A.5) up to a central term ²¹.

The (anti)commutators corresponding to (A.3), by following the description around (2.20), can be described by

$$\begin{aligned}
[(W_{B,h_1}^{\lambda, \bar{a}b})_m, (W_{B,h_2}^{\lambda, \bar{c}d})_n] &= \delta^{\bar{a}d} \delta^{\bar{c}b} c_{W_B}^{h_1, h_2}(\lambda) q^{h_1+h_2-4} [m+h_1-1]_{h_1+h_2-1} \delta_{m+n} \\
&\quad + \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
&\quad \times \left[\delta^{\bar{a}d} S_{B,R}^{h_1, h_2, h_3, k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k (W_{B,h_3}^{\lambda, \bar{c}b})_{m+n} \right. \\
&\quad \left. - \delta^{\bar{c}b} S_{B,L}^{h_1, h_2, h_3, k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} (W_{B,h_3}^{\lambda, \bar{a}d})_{m+n} \right],
\end{aligned}$$

²¹Sometimes we do not write down the structure constants with argument λ for the space of lines. They do depend on the λ except the relations in (A.8).

$$\begin{aligned}
& \left[(W_{F,h_1}^{\lambda,\bar{a}b})_m, (Q_{h_2+\frac{1}{2}}^{\lambda,\bar{c}d})_r \right] = \delta^{\bar{a}d} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times T_F^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k (Q_{h_3+\frac{1}{2}}^{\lambda,\bar{c}b})_{m+r}, \\
& \left[(W_{F,h_1}^{\lambda,\bar{a}b})_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\bar{c}\bar{d}})_r \right] = \delta^{\bar{d}\bar{b}} \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_F^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} (\bar{Q}_{h_3+\frac{1}{2}}^{\lambda,c\bar{a}})_{m+r}, \\
& \left[(W_{B,h_1}^{\lambda,\bar{a}b})_m, (Q_{h_2+\frac{1}{2}}^{\lambda,\bar{c}d})_r \right] = \delta^{\bar{c}\bar{b}} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times T_B^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} (Q_{h_3+\frac{1}{2}}^{\lambda,\bar{a}d}(w))_{m+r}, \\
& \left[(W_{B,h_1}^{\lambda,\bar{a}b})_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\bar{c}\bar{d}})_r \right] = \delta^{\bar{a}\bar{c}} \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_B^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k (\bar{Q}_{h_3+\frac{1}{2}}^{\lambda,b\bar{d}})_{m+r}, \\
& \left\{ (Q_{h_1+\frac{1}{2}}^{\lambda,\bar{a}b})_r, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\bar{c}\bar{d}})_s \right\} = \delta^{\bar{a}c} \delta^{\bar{d}b} q^{h_1+h_2-2} c_{Q\bar{Q}}^{h_1,h_2}(\lambda) [r+h_1-\frac{1}{2}]_{h_1+h_2} \delta_{r+s} \\
& \quad + \sum_{h_3=1}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} 2(-1)^{h_3+1} (4q)^{h_1+h_2-h_3} (2h_3-1)! \\
& \quad \times \left[\delta^{\bar{a}c} U_F^{h_1,h_2,h_3,k} [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k} [s+h_2-\frac{1}{2}]_k (W_{F,h_3}^{\lambda,\bar{d}b})_{r+s} \right. \\
& \quad \left. + \delta^{\bar{d}b} U_B^{h_1,h_2,h_3,k} [r+h_1-\frac{1}{2}]_k [s+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k} (W_{B,h_3}^{\lambda,\bar{a}c})_{r+s} \right]. \quad (\text{A.6})
\end{aligned}$$

The λ dependences for the structure constants are ignored to save the space but they do depend on the λ as in (A.4).

The central terms appearing in (A.6) are given by

$$\begin{aligned}
c_{W_B}^{h_1,h_2}(\lambda) & \equiv N \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2-1} (-1)^{h_2} 4^{h_1+h_2-4} a^{i_1}(h_1, \lambda) a^{i_2}(h_2, \lambda) \frac{i_1! i_2!}{(i_1+i_2+1)!}, \\
c_{Q\bar{Q}}^{h_1,h_2}(\lambda) & \equiv N \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2} 2(-1)^{h_2+1} 4^{h_1+h_2-2} \beta^{i_1}(h_1+1, \lambda) \alpha^{i_2}(h_2+1, \lambda) \frac{i_1! i_2!}{(i_1+i_2+1)!}.
\end{aligned} \quad (\text{A.7})$$

Then the complete algebra between the bifundamentals is given by either (2.20) and (A.6) or (2.18) and (A.3). The central terms are characterized by (2.21) and (A.7) while mode independent structure constants are given by (A.4).

There are nontrivial relations between the structure constants (A.4) at $\lambda = 0$ and the ones

in [18] as follows:

$$\begin{aligned}
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2} 4^{h_1+h_2-h_3-2} (2h_3-1)! S_{F,R}^{h_1,h_2,h_3,k}(0) [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \\
&= \frac{1}{2} p_F^{h_1,h_2,h_1+h_2-h_3-2}(m,n), \\
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} 4^{h_1+h_2-h_3-2} (2h_3-1)! S_{F,L}^{h_1,h_2,h_3,k}(0) [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \\
&= \frac{1}{2} p_F^{h_1,h_2,h_1+h_2-h_3-2}(m,n), \\
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2} 4^{h_1+h_2-h_3-2} (2h_3-1)! S_{B,R}^{h_1,h_2,h_3,k}(0) [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \\
&= \frac{1}{2} p_B^{h_1,h_2,h_1+h_2-h_3-2}(m,n), \\
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} 4^{h_1+h_2-h_3-2} (2h_3-1)! S_{B,L}^{h_1,h_2,h_3,k}(0) [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \\
&= \frac{1}{2} p_B^{h_1,h_2,h_1+h_2-h_3-2}(m,n), \\
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} 4^{h_1+h_2-h_3-2} (2h_3)! T_F^{h_1,h_2,h_3,k}(0) [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k \\
&= q_F^{h_1,h_2+\frac{1}{2},h_1+h_2-h_3-2}(m,r), \\
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2} 4^{h_1+h_2-h_3-2} (2h_3)! \bar{T}_F^{h_1,h_2,h_3,k}(0) [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} \\
&= q_F^{h_1,h_2+\frac{1}{2},h_1+h_2-h_3-2}(m,r), \\
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} 4^{h_1+h_2-h_3-2} (2h_3)! T_B^{h_1,h_2,h_3,k}(0) [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} \\
&= q_B^{h_1,h_2+\frac{1}{2},h_1+h_2-h_3-2}(m,r), \\
& \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} 4^{h_1+h_2-h_3-2} (2h_3)! \bar{T}_B^{h_1,h_2,h_3,k}(0) [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k \\
&= q_B^{h_1,h_2+\frac{1}{2},h_1+h_2-h_3-2}(m,r), \\
& \sum_{k=0}^{h_1+h_2-h_3} (-1)^{h_3+1} 4^{h_1+h_2-h_3} (2h_3-1)! U_F^{h_1,h_2,h_3,k}(0) [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k} [s+h_2-\frac{1}{2}]_k \\
&= \frac{1}{2} o_F^{h_1+\frac{1}{2},h_2+\frac{1}{2},h_1+h_2-h_3}(r,s),
\end{aligned}$$

$$\begin{aligned} & \sum_{k=0}^{h_1+h_2-h_3} (-1)^{h_3+1} 4^{h_1+h_2-h_3} (2h_3-1)! U_B^{h_1,h_2,h_3,k}(0) [r+h_1-\frac{1}{2}]_k [s+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k} \\ & = \frac{1}{2} o_B^{h_1+\frac{1}{2}, h_2+\frac{1}{2}, h_1+h_2-h_3}(r, s). \end{aligned} \quad (\text{A.8})$$

We obtain these relations by using (2.20) and (A.6) at $\lambda = 0$. The six structure constants appearing in (A.8), $p_F^{h_1,h_2,h_3}(m,n)$, $p_B^{h_1,h_2,h_3}(m,n)$, $q_F^{h_1,h_2,h_3}(m,r)$, $q_B^{h_1,h_2,h_3}(m,r)$, $o_F^{h_1,h_2,h_3}(r,s)$ and $o_B^{h_1,h_2,h_3}(r,s)$ are given in [18] explicitly. That is, each structure constant appearing on the left hand side has the dummy variable k and for generic h_1, h_2 and h_3 , only after this is summed over k together with mode and k dependent factors, the previous structure constant appearing on the right hand side is reproduced.

B The remaining $\mathcal{N} = 4$ supersymmetric linear $W_\infty^{2,2}[\lambda]$ algebra

The total number of (anti)commutators between the $\mathcal{N} = 4$ multiplet is given by 15. One of them is given by (2.28). The 14 remaining (anti)commutators are given by

$$\begin{aligned} & \left[(\Phi_0^{(h_1)})_m, (\Phi_{\frac{1}{2}}^{(h_2),i})_r \right] = \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} \frac{2(-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-5}}{(2h_1-1)} (2h_3)! \\ & \times \left(\left[(h_1-2\lambda) T_F^{h_1,h_2,h_3,k} + (h_1-1+2\lambda) \bar{T}_B^{h_1,h_2,h_3,k} \right] \right. \\ & \times [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k \\ & + \left[(h_1-2\lambda) \bar{T}_F^{h_1,h_2,h_3,k} + (h_1-1+2\lambda) T_B^{h_1,h_2,h_3,k} \right] \\ & \times [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} \Bigg) (\Phi_{\frac{1}{2}}^{(h_3),i})_{m+r} \\ & + \sum_{h_3=0}^{h_1+h_2-2} \sum_{k=0}^{h_1+h_2-h_3-2} \frac{2(-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-6}}{(2h_1-1)} (2h_3+2)! \\ & \times \left(\left[(h_1-2\lambda) T_F^{h_1,h_2,h_3+1,k} - (h_1-1+2\lambda) \bar{T}_B^{h_1,h_2,h_3+1,k} \right] \right. \\ & \times [m+h_1-1]_{h_1+h_2-h_3-k-2} [r+h_2-\frac{1}{2}]_k \\ & - \left[(h_1-2\lambda) \bar{T}_F^{h_1,h_2,h_3+1,k} - (h_1-1+2\lambda) T_B^{h_1,h_2,h_3+1,k} \right] \\ & \times [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-2} \Bigg) (\tilde{\Phi}_{\frac{3}{2}}^{(h_3),i})_{m+r}, \end{aligned}$$

$$\begin{aligned}
& \left[(\Phi_0^{(h_1)})_m, (\Phi_1^{(h_2),ij})_n \right] = \sum_{h_3=0}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3-1} \frac{2(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-5}}{(2h_1-1)} (2h_3+1)! \\
& \times \left(\left[(h_1-2\lambda) S_{F,R}^{h_1,h_2+1,h_3+1,k} - (h_1-1+2\lambda) S_{B,R}^{h_1,h_2+1,h_3+1,k} \right] \right. \\
& \times [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2]_k \\
& - \left. \left[(h_1-2\lambda) S_{F,L}^{h_1,h_2+1,h_3+1,k} - (h_1-1+2\lambda) S_{B,L}^{h_1,h_2+1,h_3+1,k} \right] \right. \\
& \times [m+h_1-1]_k [n+h_2]_{h_1+h_2-h_3-k-1} \left. \right) (\Phi_1^{(h_3),ij})_{m+n} \\
& + \sum_{h_3=0}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3-1} \frac{2(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-5}}{(2h_1-1)} (2h_3+1)! \\
& \times \left(\left[(h_1-2\lambda) S_{F,R}^{h_1,h_2+1,h_3+1,k} + (h_1-1+2\lambda) S_{B,R}^{h_1,h_2+1,h_3+1,k} \right] \right. \\
& \times [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2]_k \\
& - \left. \left[(h_1-2\lambda) S_{F,L}^{h_1,h_2+1,h_3+1,k} + (h_1-1+2\lambda) S_{B,L}^{h_1,h_2+1,h_3+1,k} \right] \right. \\
& \times [m+h_1-1]_k [n+h_2]_{h_1+h_2-h_3-k-1} \left. \right) (\tilde{\Phi}_1^{(h_3),ij})_{m+n}, \\
& \left[(\Phi_0^{(h_1)})_m, (\tilde{\Phi}_{\frac{3}{2}}^{(h_2),i})_r \right] = \sum_{h_3=0}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} \frac{2(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-4}}{(2h_1-1)} (2h_3)! \\
& \times \left(\left[(h_1-2\lambda) T_F^{h_1,h_2+1,h_3,k} - (h_1-1+2\lambda) \bar{T}_B^{h_1,h_2+1,h_3,k} \right] \right. \\
& \times [m+h_1-1]_{h_1+h_2-h_3-k} [r+h_2+\frac{1}{2}]_k \\
& - \left. \left[(h_1-2\lambda) \bar{T}_F^{h_1,h_2+1,h_3,k} - (h_1-1+2\lambda) T_B^{h_1,h_2+1,h_3,k} \right] \right. \\
& \times [m+h_1-1]_k [r+h_2+\frac{1}{2}]_{h_1+h_2-h_3-k} \left. \right) (\Phi_{\frac{1}{2}}^{(h_3),i})_{m+r} \\
& + \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} \frac{2(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-5}}{(2h_1-1)} (2h_3+2)! \\
& \times \left(\left[(h_1-2\lambda) T_F^{h_1,h_2+1,h_3+1,k} + (h_1-1+2\lambda) \bar{T}_B^{h_1,h_2+1,h_3+1,k} \right] \right. \\
& \times [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2+\frac{1}{2}]_k
\end{aligned}$$

$$\begin{aligned}
& + \left[(h_1 - 2\lambda) \bar{T}_F^{h_1, h_2+1, h_3+1, k} + (h_1 - 1 + 2\lambda) T_B^{h_1, h_2+1, h_3+1, k} \right] \\
& \times [m + h_1 - 1]_k [r + h_2 + \frac{1}{2}]_{h_1+h_2-h_3-k-1} \Bigg) (\tilde{\Phi}_{\frac{3}{2}}^{(h_3), i})_{m+r}, \\
& \left[(\Phi_0^{(h_1)})_m, (\tilde{\Phi}_2^{(h_2)})_n \right] = C_{0,2}^{h_1, h_2}(\lambda) [m + h_1 - 1]_{h_1+h_2+1} \delta_{m+n} \\
& + \sum_{h_3=1}^{h_1+h_2+1} \sum_{k=0}^{h_1+h_2-h_3+1} \frac{2(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-3}}{(2h_1-1)} (2h_3-1)! \\
& \times \left(\left[(h_1 - 2\lambda) S_{F,R}^{h_1, h_2+2, h_3, k} + (h_1 - 1 + 2\lambda) S_{B,R}^{h_1, h_2+2, h_3, k} \right] \right. \\
& \times [m + h_1 - 1]_{h_1+h_2-h_3-k+1} [n + h_2 + 1]_k \\
& \left. - \left[(h_1 - 2\lambda) S_{F,L}^{h_1, h_2+2, h_3, k} + (h_1 - 1 + 2\lambda) S_{B,L}^{h_1, h_2+2, h_3, k} \right] \right. \\
& \times [m + h_1 - 1]_k [n + h_2 + 1]_{h_1+h_2-h_3-k+1} \Bigg) (\Phi_0^{(h_3)})_{m+n} \\
& + \sum_{h_3=-1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} \frac{(-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-4}}{(2h_1-1)} (2h_3+2)! \\
& \times \left(\left[(h_1 - 2\lambda)(h_3 + 1 + 2\lambda) S_{F,R}^{h_1, h_2+2, h_3+2, k} \right. \right. \\
& \left. \left. - (h_1 - 1 + 2\lambda)(h_3 + 2 - 2\lambda) S_{B,R}^{h_1, h_2+2, h_3+2, k} \right] \right. \\
& \times [m + h_1 - 1]_{h_1+h_2-h_3-k-1} [n + h_2 + 1]_k \\
& \left. - \left[(h_1 - 2\lambda)(h_3 + 1 + 2\lambda) S_{F,L}^{h_1, h_2+2, h_3+2, k} \right. \right. \\
& \left. \left. - (h_1 - 1 + 2\lambda)(h_3 + 2 - 2\lambda) S_{B,L}^{h_1, h_2+2, h_3+2, k} \right] \right. \\
& \times [m + h_1 - 1]_k [n + h_2 + 1]_{h_1+h_2-h_3-k-1} \Bigg) (\tilde{\Phi}_2^{(h_3)})_{m+n}, \\
& \left\{ (\Phi_{\frac{1}{2}}^{(h_1), i})_r, (\Phi_{\frac{1}{2}}^{(h_2), j})_s \right\} = \delta^{ij} C_{\frac{1}{2}, \frac{1}{2}}^{h_1, h_2}(\lambda) [r + h_1 - \frac{1}{2}]_{h_1+h_2} \delta_{r+s} \\
& + \delta^{ij} \sum_{h_3=1}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-4} (2h_3-1)! \\
& \times \left(\left(U_B^{h_2, h_1, h_3, k} - U_F^{h_1, h_2, h_3, k} \right) [r + h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k} [s + h_2 - \frac{1}{2}]_k \right)
\end{aligned}$$

$$\begin{aligned}
& + \left(U_B^{h_1, h_2, h_3, k} - U_F^{h_2, h_1, h_3, k} \right) [r + h_1 - \frac{1}{2}]_k [s + h_2 - \frac{1}{2}]_{h_1+h_2-h_3-k} \left(\Phi_0^{(h_3)} \right)_{r+s} \\
& + \delta^{ij} \sum_{h_3=-1}^{h_1+h_2-2} \sum_{k=0}^{h_1+h_2-h_3-2} 2(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-6} (2h_3+2)! \\
& \times \left(\left((h_3+1+2\lambda) U_F^{h_1, h_2, h_3+2, k} + (h_3+2-2\lambda) U_B^{h_2, h_1, h_3+2, k} \right) \right. \\
& \times [r + h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k-2} [s + h_2 - \frac{1}{2}]_k \\
& + \left((h_3+1+2\lambda) U_F^{h_2, h_1, h_3+2, k} + (h_3+2-2\lambda) U_B^{h_1, h_2, h_3+2, k} \right) \\
& \times [r + h_1 - \frac{1}{2}]_k [s + h_2 - \frac{1}{2}]_{h_1+h_2-h_3-k-2} \left. \right) \left(\tilde{\Phi}_2^{(h_3)} \right)_{r+s} \\
& + \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-5} (2h_3+1)! \left[\right. \\
& \times \left(\left(U_F^{h_1, h_2, h_3+1, k} + U_B^{h_2, h_1, h_3+1, k} \right) [r + h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k-1} [s + h_2 - \frac{1}{2}]_k \right. \\
& - \left. \left(U_F^{h_2, h_1, h_3+1, k} + U_B^{h_1, h_2, h_3+1, k} \right) [r + h_1 - \frac{1}{2}]_k [s + h_2 - \frac{1}{2}]_{h_1+h_2-h_3-k-1} \right) \left(\Phi_1^{(h_3), ij} \right)_{r+s} \\
& + \left(\left(U_F^{h_1, h_2, h_3+1, k} - U_B^{h_2, h_1, h_3+1, k} \right) [r + h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k-1} [s + h_2 - \frac{1}{2}]_k \right. \\
& - \left. \left(U_F^{h_2, h_1, h_3+1, k} - U_B^{h_1, h_2, h_3+1, k} \right) [r + h_1 - \frac{1}{2}]_k [s + h_2 - \frac{1}{2}]_{h_1+h_2-h_3-k-1} \right) \left(\tilde{\Phi}_1^{(h_3), ij} \right)_{r+s} \left. \right], \\
& \left[(\Phi_{\frac{1}{2}}^{(h_1), i})_r, (\Phi_1^{(h_2), jk})_m \right] = \delta^{ij} \sum_{h_3=0}^{h_1+h_2} \sum_{t=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-4} (2h_3)! \\
& \times \left(\left(T_F^{h_2+1, h_1, h_3, t} - \bar{T}_B^{h_2+1, h_1, h_3, t} \right) [r + h_1 - \frac{1}{2}]_t [m + h_2]_{h_1+h_2-h_3-t} \right. \\
& + \left. \left(T_B^{h_2+1, h_1, h_3, t} - \bar{T}_F^{h_2+1, h_1, h_3, t} \right) [r + h_1 - \frac{1}{2}]_{h_1+h_2-h_3-t} [m + h_2]_t \right) \left(\Phi_{\frac{1}{2}}^{(h_3), k} \right)_{r+m} \\
& + \delta^{ij} \sum_{h_3=0}^{h_1+h_2-1} \sum_{t=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-5} (2h_3+2)! \\
& \times \left(\left(T_F^{h_2+1, h_1, h_3+1, t} + \bar{T}_B^{h_2+1, h_1, h_3+1, t} \right) [r + h_1 - \frac{1}{2}]_t [m + h_2]_{h_1+h_2-h_3-t-1} \right.
\end{aligned}$$

$$\begin{aligned}
& + \left(T_B^{h_2+1, h_1, h_3+1, t} + \bar{T}_F^{h_2+1, h_1, h_3+1, t} \right) [r+h_1 - \frac{1}{2}]_{h_1+h_2-h_3-t-1} [m+h_2]_t \left(\tilde{\Phi}_{\frac{3}{2}}^{(h_3), k} \right)_{r+m} \\
& - \delta^{ik} \left[j \leftrightarrow k \right] \\
& + \epsilon^{ijkl} \sum_{h_3=0}^{h_1+h_2} \sum_{t=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-4} (2h_3)! \\
& \times \left(\left(T_F^{h_2+1, h_1, h_3, t} + \bar{T}_B^{h_2+1, h_1, h_3, t} \right) [r+h_1 - \frac{1}{2}]_t [m+h_2]_{h_1+h_2-h_3-t} \right. \\
& \left. - \left(T_B^{h_2+1, h_1, h_3, t} + \bar{T}_F^{h_2+1, h_1, h_3, t} \right) [r+h_1 - \frac{1}{2}]_{h_1+h_2-h_3-t} [m+h_2]_t \right) \left(\Phi_{\frac{1}{2}}^{(h_3), l} \right)_{r+m} \\
& + \epsilon^{ijkl} \sum_{h_3=0}^{h_1+h_2-1} \sum_{t=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-5} (2h_3+2)! \\
& \times \left(\left(T_F^{h_2+1, h_1, h_3+1, t} - \bar{T}_B^{h_2+1, h_1, h_3+1, t} \right) [r+h_1 - \frac{1}{2}]_t [m+h_2]_{h_1+h_2-h_3-t-1} \right. \\
& \left. - \left(T_B^{h_2+1, h_1, h_3+1, t} - \bar{T}_F^{h_2+1, h_1, h_3+1, t} \right) [r+h_1 - \frac{1}{2}]_{h_1+h_2-h_3-t-1} [m+h_2]_t \right) \left(\tilde{\Phi}_{\frac{3}{2}}^{(h_3), l} \right)_{r+m}, \\
& \left\{ (\Phi_{\frac{1}{2}}^{(h_1), i})_r, (\tilde{\Phi}_{\frac{3}{2}}^{(h_2), j})_s \right\} = \delta^{ij} C_{\frac{1}{2}, \frac{3}{2}}^{h_1, h_2}(\lambda) [r+h_1 - \frac{1}{2}]_{h_1+h_2+1} \delta_{r+s} \\
& + \delta^{ij} \sum_{h_3=1}^{h_1+h_2+1} \sum_{k=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-3} (2h_3-1)! \\
& \times \left[\left(U_F^{h_1, h_2+1, h_3, k} + U_B^{h_2+1, h_1, h_3, k} \right) [r+h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k+1} [s+h_2 + \frac{1}{2}]_k \right. \\
& \left. - \left(U_B^{h_1, h_2+1, h_3, k} + U_F^{h_2+1, h_1, h_3, k} \right) [r+h_1 - \frac{1}{2}]_k [s+h_2 + \frac{1}{2}]_{h_1+h_2-h_3-k+1} \right] \left(\Phi_0^{(h_3)} \right)_{r+s} \\
& + \delta^{ij} \sum_{h_3=-1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} 2(-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-5} (2h_3+2)! \\
& \times \left[\left((h_3+1+2\lambda) U_F^{h_1, h_2+1, h_3+2, k} - (h_3+2-2\lambda) U_B^{h_2+1, h_1, h_3+2, k} \right) \right. \\
& \times [r+h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k-1} [s+h_2 + \frac{1}{2}]_k \\
& \left. + \left((h_3+2-2\lambda) U_B^{h_1, h_2+1, h_3+2, k} - (h_3+1+2\lambda) U_F^{h_2+1, h_1, h_3+2, k} \right) \right. \\
& \times [r+h_1 - \frac{1}{2}]_k [s+h_2 + \frac{1}{2}]_{h_1+h_2-h_3-k-1} \left. \right] \left(\tilde{\Phi}_2^{(h_3)} \right)_{r+s}
\end{aligned}$$

$$\begin{aligned}
& + \sum_{h_3=0}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-4} (2h_3+1)! \left[\right. \\
& \times \left(\left(U_F^{h_1, h_2+1, h_3+1, k} - U_B^{h_2+1, h_1, h_3+1, k} \right) [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k} [s+h_2+\frac{1}{2}]_k \right. \\
& + \left(U_F^{h_2+1, h_1, h_3+1, k} - U_B^{h_1, h_2+1, h_3+1, k} \right) [r+h_1-\frac{1}{2}]_k [s+h_2+\frac{1}{2}]_{h_1+h_2-h_3-k} \left. \right) (\Phi_1^{(h_3),ij})_{r+s} \\
& + \left(\left(U_F^{h_1, h_2+1, h_3+1, k} + U_B^{h_2+1, h_1, h_3+1, k} \right) [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k} [s+h_2+\frac{1}{2}]_k \right. \\
& + \left. \left. \left(U_F^{h_2+1, h_1, h_3+1, k} + U_B^{h_1, h_2+1, h_3+1, k} \right) [r+h_1-\frac{1}{2}]_k [s+h_2+\frac{1}{2}]_{h_1+h_2-h_3-k} \right) (\tilde{\Phi}_1^{(h_3),ij})_{r+s} \right], \\
& \left[(\Phi_{\frac{1}{2}}^{(h_1),i})_r, (\tilde{\Phi}_2^{(h_2)})_m \right] = \sum_{h_3=0}^{h_1+h_2+1} \sum_{k=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-3} (2h_3)! \\
& \times \left[\left(T_F^{h_2+2, h_1, h_3, k} - \bar{T}_B^{h_2+2, h_1, h_3, k} \right) [r+h_1-\frac{1}{2}]_k [m+h_2+1]_{h_1+h_2-h_3-k+1} \right. \\
& + \left. \left(\bar{T}_F^{h_2+2, h_1, h_3, k} - T_B^{h_2+2, h_1, h_3, k} \right) [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k+1} [m+h_2+1]_k \right] (\Phi_{\frac{1}{2}}^{(h_3),i})_{r+m} \\
& + \sum_{h_3=0}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-4} (2h_3+2)! \\
& \times \left[\left(T_F^{h_2+2, h_1, h_3+1, k} + \bar{T}_B^{h_2+2, h_1, h_3+1, k} \right) [r+h_1-\frac{1}{2}]_k [m+h_2+1]_{h_1+h_2-h_3-k} \right. \\
& - \left. \left(\bar{T}_F^{h_2+2, h_1, h_3+1, k} + T_B^{h_2+2, h_1, h_3+1, k} \right) [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k} [m+h_2+1]_k \right] (\tilde{\Phi}_{\frac{3}{2}}^{(h_3),i})_{r+m}, \\
& \left[(\Phi_1^{(h_1),ij})_m, (\Phi_1^{(h_2),kl})_n \right] = \left(\delta^{ik} \delta^{jl} - \delta^{il} \delta^{jk} \right) C_{1,1,-}^{h_1,h_2}(\lambda) [m+h_1]_{h_1+h_2+1} \delta_{m+n} \\
& + \epsilon^{ijkl} C_{1,1,+}^{h_1,h_2}(\lambda) [m+h_1]_{h_1+h_2+1} \delta_{m+n} \\
& + \left(\delta^{ik} \delta^{jl} - \delta^{il} \delta^{jk} \right) \sum_{h_3=1}^{h_1+h_2+1} \sum_{t=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-3} (2h_3-1)! \\
& \times \left[\left(S_{F,R}^{h_1+1, h_2+1, h_3, t} - S_{B,R}^{h_1+1, h_2+1, h_3, t} \right) [m+h_1]_{h_1+h_2-h_3-t+1} [n+h_2, t] \right. \\
& - \left. \left(S_{F,L}^{h_1+1, h_2+1, h_3, t} - S_{B,L}^{h_1+1, h_2+1, h_3, t} \right) [m+h_1]_t [n+h_2]_{h_1+h_2-h_3-t+1} \right] (\Phi_0^{(h_3)})_{m+n}
\end{aligned}$$

$$\begin{aligned}
& + \left(\delta^{ik} \delta^{jl} - \delta^{il} \delta^{jk} \right) \sum_{h_3=-1}^{h_1+h_2-1} \sum_{t=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} 2^{-1} 4^{2(h_1+h_2-h_3)-4} (2h_3+2)! \\
& \times \left[\left((h_3+1+2\lambda) S_{F,R}^{h_1+1,h_2+1,h_3+2,t} + (h_3+2-2\lambda) S_{B,R}^{h_1+1,h_2+1,h_3+2,t} \right) \right. \\
& \times [m+h_1]_{h_1+h_2-h_3-t-1} [n+h_2]_t \\
& - \left((h_3+1+2\lambda) S_{F,L}^{h_1+1,h_2+1,h_3+2,t} + (h_3+2-2\lambda) S_{B,L}^{h_1+1,h_2+1,h_3+2,t} \right) \\
& \times [m+h_1]_t [n+h_2]_{h_1+h_2-h_3-t-1} \Bigg] (\tilde{\Phi}_2^{(h_3)})_{m+n} \\
& + \epsilon^{ijkl} \sum_{h_3=0}^{h_1+h_2+1} \sum_{t=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-3} (2h_3-1)! \\
& \times \left[\left(S_{F,R}^{h_1+1,h_2+1,h_3,t} + S_{B,R}^{h_1+1,h_2+1,h_3,t} \right) [m+h_1]_{h_1+h_2-h_3-t+1} [n+h_2]_t \right. \\
& - \left(S_{F,L}^{h_1+1,h_2+1,h_3,t} + S_{B,L}^{h_1+1,h_2+1,h_3,t} \right) [m+h_1]_t [n+h_2]_{h_1+h_2-h_3-t+1} \Bigg] (\Phi_0^{(h_3)})_{m+n} \\
& + \epsilon^{ijkl} \sum_{h_3=-1}^{h_1+h_2-1} \sum_{t=0}^{h_1+h_2-h_3-1} (-1)^{h_1+h_2+1} 2^{-1} 4^{2(h_1+h_2-h_3)-4} (2h_3+2)! \\
& \times \left[\left((h_3+1+2\lambda) S_{F,R}^{h_1+1,h_2+1,h_3+2,t} - (h_3+2-2\lambda) S_{B,R}^{h_1+1,h_2+1,h_3+2,t} \right) \right. \\
& \times [m+h_1]_{h_1+h_2-h_3-t-1} [n+h_2]_t \\
& - \left((h_3+1+2\lambda) S_{F,L}^{h_1+1,h_2+1,h_3+2,t} - (h_3+2-2\lambda) S_{B,L}^{h_1+1,h_2+1,h_3+2,t} \right) \\
& \times [m+h_1]_t [n+h_2]_{h_1+h_2-h_3-t-1} \Bigg] (\tilde{\Phi}_2^{(h_3)})_{m+n} \\
& + \delta^{ik} \sum_{h_3=0}^{h_1+h_2} \sum_{t=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-4} (2h_3+1)! \Bigg[\\
& \times \left(\left(S_{F,R}^{h_1+1,h_2+1,h_3+1,t} + S_{B,R}^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_{h_1+h_2-h_3-t} [n+h_2]_t \right. \\
& + \left(S_{F,L}^{h_1+1,h_2+1,h_3+1,t} + S_{B,L}^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_t [n+h_2]_{h_1+h_2-h_3-t} \Bigg) (\Phi_1^{(h_3),jl})_{m+n} \\
& + \left(\left(S_{F,R}^{h_1+1,h_2+1,h_3+1,t} - S_{B,R}^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_{h_1+h_2-h_3-t} [n+h_2]_t \right.
\end{aligned}$$

$$\begin{aligned}
& + \left(S_{F,L}^{h_1+1,h_2+1,h_3+1,t} - S_{B,L}^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_t [n+h_2]_{h_1+h_2-h_3-t} \left(\tilde{\Phi}_1^{(h_3),jl} \right)_{m+n} \Bigg] \\
& - \delta^{il} \left[j \leftrightarrow k \right] - \delta^{jk} \left[i \leftrightarrow j \right] + \delta^{jl} \left[i \leftrightarrow j, k \leftrightarrow l \right], \\
& \left[(\Phi_1^{(h_1),ij})_m, (\tilde{\Phi}_{\frac{3}{2}}^{(h_2),k})_r \right] = \delta^{ik} \sum_{h_3=0}^{h_1+h_2+1} \sum_{t=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2+1-h_3)-3} (2h_3)! \\
& \times \left[\left(T_F^{h_1+1,h_2+1,h_3,t} + \bar{T}_B^{h_1+1,h_2+1,h_3,t} \right) [m+h_1]_{h_1+h_2-h_3-t+1} [r+h_2+\frac{1}{2}]_t \right. \\
& \left. + \left(T_B^{h_1+1,h_2+1,h_3,t} + \bar{T}_F^{h_1+1,h_2+1,h_3,t} \right) [m+h_1]_t [r+h_2+\frac{1}{2}]_{h_1+h_2-h_3-t+1} \right] (\Phi_{\frac{1}{2}}^{(h_3),j})_{m+r} \\
& + \delta^{ik} \sum_{h_3=0}^{h_1+h_2} \sum_{t=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2+1-h_3)-4} (2h_3+2)! \\
& \times \left[\left(T_F^{h_1+1,h_2+1,h_3+1,t} - \bar{T}_B^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_{h_1+h_2-h_3-t} [r+h_2+\frac{1}{2}]_t \right. \\
& \left. + \left(T_B^{h_1+1,h_2+1,h_3+1,t} - \bar{T}_F^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_t [r+h_2+\frac{1}{2}]_{h_1+h_2-h_3-t} \right] (\tilde{\Phi}_{\frac{3}{2}}^{(h_3),j})_{m+r} \\
& - \delta^{jk} \left[i \leftrightarrow j \right] \\
& + \epsilon^{ijkl} \sum_{h_3=0}^{h_1+h_2+1} \sum_{t=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2+1-h_3)-3} (2h_3)! \\
& \times \left[\left(T_F^{h_1+1,h_2+1,h_3,t} - \bar{T}_B^{h_1+1,h_2+1,h_3,t} \right) [m+h_1]_{h_1+h_2-h_3-t+1} [r+h_2+\frac{1}{2}]_t \right. \\
& \left. - \left(T_B^{h_1+1,h_2+1,h_3,t} - \bar{T}_F^{h_1+1,h_2+1,h_3,t} \right) [m+h_1]_t [r+h_2+\frac{1}{2}]_{h_1+h_2-h_3-t+1} \right] (\Phi_{\frac{1}{2}}^{(h_3),l})_{m+r} \\
& + \epsilon^{ijkl} \sum_{h_3=0}^{h_1+h_2} \sum_{t=0}^{h_1+h_2-h_3} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2+1-h_3)-4} (2h_3+2)! \\
& \times \left[\left(T_F^{h_1+1,h_2+1,h_3+1,t} + \bar{T}_B^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_{h_1+h_2-h_3-t} [r+h_2+\frac{1}{2}]_t \right. \\
& \left. - \left(T_B^{h_1+1,h_2+1,h_3+1,t} + \bar{T}_F^{h_1+1,h_2+1,h_3+1,t} \right) [m+h_1]_t [r+h_2+\frac{1}{2}]_{h_1+h_2-h_3-t} \right] (\tilde{\Phi}_{\frac{3}{2}}^{(h_3),l})_{m+r},
\end{aligned}$$

$$\begin{aligned}
& \left[(\Phi_1^{(h_1),ij})_m, (\tilde{\Phi}_2^{(h_2)})_n \right] = \sum_{h_3=0}^{h_1+h_2+1} \sum_{k=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-3} (2h_3+1)! \\
& \times \left[\left(\left(S_{F,R}^{h_1+1,h_2+2,h_3+1,k} + S_{B,R}^{h_1+1,h_2+2,h_3+1,k} \right) [m+h_1]_{h_1+h_2-h_3-k+1} [n+h_2+1]_k \right. \right. \\
& - \left(S_{F,L}^{h_1+1,h_2+2,h_3+1,k} + S_{B,L}^{h_1+1,h_2+2,h_3+1,k} \right) [m+h_1]_k [n+h_2+1]_{h_1+h_2-h_3-k+1} \left. \right) (\Phi_1^{(h_3),ij})_{m+n} \\
& + \left(\left(S_{F,R}^{h_1+1,h_2+2,h_3+1,k} - S_{B,R}^{h_1+1,h_2+2,h_3+1,k} \right) [m+h_1]_{h_1+h_2-h_3-k+1} [n+h_2+1]_k \right. \\
& - \left. \left. \left(S_{F,L}^{h_1+1,h_2+2,h_3+1,k} - S_{B,L}^{h_1+1,h_2+2,h_3+1,k} \right) [m+h_1]_k [n+h_2+1]_{h_1+h_2-h_3-k+1} \right) (\tilde{\Phi}_1^{(h_3),ij})_{m+n} \right], \\
& \left\{ (\tilde{\Phi}_{\frac{3}{2}}^{(h_2),i})_r, (\tilde{\Phi}_{\frac{3}{2}}^{(h_2),j})_s \right\} = \delta^{ij} C_{\frac{3}{2}, \frac{3}{2}}^{h_1, h_2}(\lambda) [r+h_1+\frac{1}{2}]_{h_1+h_2+2} \delta_{r+s} \\
& + \delta^{ij} \sum_{h_3=1}^{h_1+h_2+2} \sum_{k=0}^{h_1+h_2-h_3+2} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-2} (2h_3-1)! \\
& \times \left[\left(U_F^{h_1+1,h_2+1,h_3,k} - U_B^{h_2+1,h_1+1,h_3,k} \right) [r+h_1+\frac{1}{2}]_{h_1+h_2-h_3-k+2} [s+h_2+\frac{1}{2}]_k \right. \\
& + \left. \left(U_F^{h_2+1,h_1+1,h_3,k} - U_B^{h_1+1,h_2+1,h_3,k} \right) [r+h_1+\frac{1}{2}]_k [s+h_2+\frac{1}{2}]_{h_1+h_2-h_3-k+2} \right] (\Phi_0^{(h_3)})_{r+s} \\
& + \delta^{ij} \sum_{h_3=-1}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} 2(-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-4} (2h_3+2)! \\
& \times \left[\left((h_3+1+2\lambda) U_F^{h_1+1,h_2+1,h_3+2,k} + (h_3+2-2\lambda) U_B^{h_2+1,h_1+1,h_3+2,k} \right) \right. \\
& \times [r+h_1+\frac{1}{2}]_{h_1+h_2-h_3-k} [s+h_2+\frac{1}{2}]_k \\
& + \left. \left((h_3+1+2\lambda) U_F^{h_2+1,h_1+1,h_3+2,k} + (h_3+2-2\lambda) U_B^{h_1+1,h_2+1,h_3+2,k} \right) \right. \\
& \times [r+h_1+\frac{1}{2}]_k [s+h_2+\frac{1}{2}]_{h_1+h_2-h_3-k} \left. \right] (\tilde{\Phi}_2^{(h_3)})_{r+s} \\
& + \sum_{h_3=0}^{h_1+h_2+1} \sum_{k=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-3} (2h_3+1)! \left[\right. \\
& \times \left(\left(U_F^{h_1+1,h_2+1,h_3+1,k} + U_B^{h_2+1,h_1+1,h_3+1,k} \right) [r+h_1+\frac{1}{2}]_{h_1+h_2-h_3-k+1} [s+h_2+\frac{1}{2}]_k \right. \\
& - \left. \left. \left(U_F^{h_2+1,h_1+1,h_3+1,k} + U_B^{h_1+1,h_2+1,h_3+1,k} \right) [r+h_1+\frac{1}{2}]_k [s+h_2+\frac{1}{2}]_{h_1+h_2-h_3-k+1} \right) (\Phi_1^{(h_3),ij})_{r+s} \right]
\end{aligned}$$

$$\begin{aligned}
& + \left(\left(U_F^{h_1+1, h_2+1, h_3+1, k} - U_B^{h_2+1, h_1+1, h_3+1, k} \right) [r+h_1+\frac{1}{2}]_{h_1+h_2-h_3-k+1} [s+h_2+\frac{1}{2}]_k \right. \\
& \quad \left. - \left(U_F^{h_2+1, h_1+1, h_3+1, k} - U_B^{h_1+1, h_2+1, h_3+1, k} \right) [r+h_1+\frac{1}{2}]_k [s+h_2+\frac{1}{2}]_{h_1+h_2-h_3-k+1} \right) (\tilde{\Phi}_1^{(h_3), ij})_{r+s}, \\
& \left[(\tilde{\Phi}_{\frac{3}{2}}^{(h_2), i})_r, (\tilde{\Phi}_2^{(h_2)})_m \right] = \sum_{h_3=0}^{h_1+h_2+2} \sum_{k=0}^{h_1+h_2-h_3+2} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-2} (2h_3)! \\
& \times \left[\left(T_F^{h_2+2, h_1+1, h_3, k} + \bar{T}_B^{h_2+2, h_1+1, h_3, k} \right) [r+h_1+\frac{1}{2}]_k [m+h_2+1]_{h_1+h_2-h_3-k+2} \right. \\
& \quad \left. - \left(\bar{T}_F^{h_2+2, h_1+1, h_3, k} + T_B^{h_2+2, h_1+1, h_3, k} \right) [r+h_1+\frac{1}{2}]_{h_1+h_2-h_3-k+2} [m+h_2+1]_k \right] (\Phi_{\frac{1}{2}}^{(h_3), i})_{r+m} \\
& + \sum_{h_3=-1}^{h_1+h_2+1} \sum_{k=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2} 4^{2(h_1+h_2-h_3)-3} (2h_3+2)! \\
& \times \left[\left(T_F^{h_2+2, h_1+1, h_3+1, k} - \bar{T}_B^{h_2+2, h_1+1, h_3+1, k} \right) [r+h_1+\frac{1}{2}]_k [m+h_2+1]_{h_1+h_2-h_3-k+1} \right. \\
& \quad \left. + \left(\bar{T}_F^{h_2+2, h_1+1, h_3+1, k} - T_B^{h_2+2, h_1+1, h_3+1, k} \right) [r+h_1+\frac{1}{2}]_{h_1+h_2-h_3-k+1} [m+h_2+1]_k \right] (\tilde{\Phi}_{\frac{3}{2}}^{(h_3), i})_{r+m}, \\
& \left[(\tilde{\Phi}_2^{(h_1)})_m, (\tilde{\Phi}_2^{(h_2)})_n \right] = C_{2,2}^{h_1, h_2}(\lambda) [m+h_1+1]_{h_1+h_2+3} \delta_{m+n} \\
& + \sum_{h_3=1}^{h_1+h_2+3} \sum_{k=0}^{h_1+h_2-h_3+3} (-1)^{h_1+h_2+1} 4^{2(h_1+h_2-h_3)-1} (2h_3-1)! \\
& \times \left[\left(S_{F,R}^{h_1+2, h_2+2, h_3, k} - S_{B,R}^{h_1+2, h_2+2, h_3, k} \right) [m+h_1+1]_{h_1+h_2-h_3-k+3} [n+h_2+1]_k \right. \\
& \quad \left. - \left(S_{F,L}^{h_1+2, h_2+2, h_3, k} - S_{B,L}^{h_1+2, h_2+2, h_3, k} \right) [m+h_1+1]_k [n+h_2+1]_{h_1+h_2-h_3-k+3} \right] (\Phi_0^{(h_3)})_{m+n} \\
& + \sum_{h_3=-1}^{h_1+h_2+1} \sum_{k=0}^{h_1+h_2-h_3+1} (-1)^{h_1+h_2} 2^{-1} 4^{2(h_1+h_2-h_3)-2} (2h_3+2)! \\
& \times \left[\left((h_3+1+2\lambda) S_{F,R}^{h_1+2, h_2+2, h_3+2, k} + (h_3+2-2\lambda) S_{B,R}^{h_1+2, h_2+2, h_3+2, k} \right) \right. \\
& \quad \left. \times [m+h_1+1]_{h_1+h_2-h_3-k+1} [n+h_2+1]_k \right. \\
& \quad \left. - \left((h_3+1+2\lambda) S_{F,L}^{h_1+2, h_2+2, h_3+2, k} + (h_3+2-2\lambda) S_{B,L}^{h_1+2, h_2+2, h_3+2, k} \right) \right]
\end{aligned}$$

$$\times [m + h_1 + 1]_k [n + h_2 + 1]_{h_1 + h_2 - h_3 - k + 1} \left[(\tilde{\Phi}_2^{(h_3)})_{m+n} \right] \quad (B.1)$$

Note that we introduce the fields $\tilde{\Phi}_1^{(h),ij} \equiv \frac{1}{2}\epsilon^{ijkl}\Phi_1^{(h),kl}$. In the sixth commutator, we use a simplified notation $[j \leftrightarrow k]$ which stands for the substitution $j \leftrightarrow k$ in the previous expressions. We do not write down the arguments for the structure constants appearing in (A.4) in (B.1)²² and the structure constants are given by (A.4).

The λ dependent coefficients appearing in (B.1) are

$$\begin{aligned} C_{0,2}^{h_1,h_2}(\lambda) &\equiv \frac{N(-1)^{h_1} 4^{2(h_1+h_2)-6}}{(2h_1-1)} \sum_{i_1=0}^{h_1-1} \sum_{i_2=0}^{h_2+1} \frac{i_1! i_2!}{(i_1+i_2+1)!} \\ &\quad \times \left[(h_1 - 2\lambda)a^{i_1}(h_1, \lambda + \frac{1}{2})a^{i_2}(h_2 + 2, \lambda + \frac{1}{2}) \right. \\ &\quad \left. + (h_1 - 1 + 2\lambda)a^{i_1}(h_1, \lambda)a^{i_2}(h_2 + 2, \lambda) \right], \\ C_{\frac{1}{2},\frac{1}{2}}^{h_1,h_2}(\lambda) &\equiv 2N(-1)^{h_1} 4^{2(h_1+h_2)-4} \sum_{i_1=0}^{h_1} \sum_{i_2=0}^{h_2} \frac{i_1! i_2!}{(i_1+i_2+1)!} \\ &\quad \times \left[\beta^{i_1}(h_1 + 1, \lambda)\alpha^{i_2}(h_2 + 1, \lambda) + \alpha^{i_1}(h_1 + 1, \lambda)\beta^{i_2}(h_2 + 1, \lambda) \right], \\ C_{\frac{1}{2},\frac{3}{2}}^{h_1,h_2}(\lambda) &\equiv 2N(-1)^{h_1+1} 4^{2(h_1+h_2)-7} \sum_{i_1=0}^{h_1} \sum_{i_2=0}^{h_2+1} \frac{i_1! i_2!}{(i_1+i_2+1)!} \\ &\quad \times \left(\beta^{i_1}(h_1 + 1, \lambda)\alpha^{i_2}(h_2 + 2, \lambda) - \alpha^{i_1}(h_1 + 1, \lambda)\beta^{i_2}(h_2 + 2, \lambda) \right), \\ C_{1,1,\pm}^{h_1,h_2}(\lambda) &\equiv 2N(-1)^{h_1+1} 4^{2(h_1+h_2)-7} \sum_{i_1=0}^{h_1} \sum_{i_2=0}^{h_2} \frac{i_1! i_2!}{(i_1+i_2+1)!} \\ &\quad \times \left[a^{i_1}(h_1 + 1, \lambda + \frac{1}{2})a^{i_2}(h_2 + 1, \lambda + \frac{1}{2}) \pm a^{i_1}(h_1 + 1, \lambda)a^{i_2}(h_2 + 1, \lambda) \right], \\ C_{\frac{3}{2},\frac{3}{2}}^{h_1,h_2}(\lambda) &\equiv 2N(-1)^{h_1} 4^{2(h_1+h_2)-6} \sum_{i_1=0}^{h_1+1} \sum_{i_2=0}^{h_2+1} \frac{i_1! i_2!}{(i_1+i_2+1)!} \\ &\quad \times \left[\beta^{i_1}(h_1 + 2, \lambda)\alpha^{i_2}(h_2 + 2, \lambda) + \alpha^{i_1}(h_1 + 2, \lambda)\beta^{i_2}(h_2 + 2, \lambda) \right], \end{aligned}$$

²²The corresponding OPEs can be obtained from the analysis of the footnote 12.

$$C_{2,2}^{h_1,h_2}(\lambda) \equiv 2 N (-1)^{h_1+1} 4^{2(h_1+h_2)-5} \sum_{i_1=0}^{h_1+1} \sum_{i_2=0}^{h_2+1} \frac{i_1! i_2!}{(i_1+i_2+1)!} \\ \times \left[a^{i_1}(h_1+2, \lambda + \frac{1}{2}) a^{i_2}(h_2+2, \lambda + \frac{1}{2}) - a^{i_1}(h_1+2, \lambda) a^{i_2}(h_2+2, \lambda) \right], \quad (\text{B.2})$$

where the previous relations (2.3) can be applied. Note that the first and the third in (B.2) have nonzero values although the components of the $\mathcal{N} = 4$ multiplet on the left hand side are different from each other because of (2.26) and (2.27).

C The basic component OPEs for $\mathcal{N} = 4$ supersymmetric OPE

The fundamental five OPEs for the $\mathcal{N} = 4$ superspace description (2.30) are enough to write the component OPEs in terms of $\mathcal{N} = 4$ super OPE because other corresponding OPEs to (B.1) can be obtained by taking super derivatives and putting the fermionic coordinates to zero in $\mathcal{N} = 4$ superspace²³. They are given by

$$\Phi_0^{(h_1)}(z) \Phi_0^{(h_2)}(w) = \frac{1}{(z-w)^{h_1+h_2}} C_{0,0}^{h_1,h_2}(\lambda) (h_1+h_2-1)! \\ + \sum_{p=1}^{h_1+h_2-1} \sum_{h_3=1}^{h_1+h_2-p} \frac{1}{(z-w)^p} \left[f_1^{h_1,h_2,h_3,p}(\lambda) \partial_w^{h_1+h_2-h_3-p} \Phi_0^{(h_3)} \right. \\ \left. + f_2^{h_1,h_2,h_3,p}(\lambda) \partial_w^{h_1+h_2-h_3-p} \left(\Phi_2^{(h_3-2)} - \frac{(1-4\lambda)}{(2h_3-3)} \partial_w^2 \Phi_0^{(h_3-2)} \right) \right] (w) + \dots, \\ \Phi_{\frac{1}{2}}^{(h_1),i}(z) \Phi_0^{(h_2)}(w) = \sum_{p=1}^{h_1+h_2} \frac{1}{(z-w)^p} \left[\sum_{h_3=0}^{h_1+h_2-p} f_3^{h_1,h_2,h_3,p}(\lambda) \partial_w^{h_1+h_2-h_3-p} \Phi_{\frac{1}{2}}^{(h_3),i} \right. \\ \left. - \sum_{h_3=0}^{h_1+h_2-p-1} f_4^{h_1,h_2,h_3,p}(\lambda) \partial_w^{h_1+h_2-h_3-p-1} \left(\Phi_{\frac{3}{2}}^{(h_3),i} - \frac{(1-4\lambda)}{(2h_3+1)} \partial_w \Phi_{\frac{1}{2}}^{(h_3),i} \right) \right] (w) \\ + \dots, \\ \Phi_1^{(h_1),ij}(z) \Phi_0^{(h_2)}(w) = \sum_{p=1}^{h_1+h_2} \sum_{h_3=0}^{h_1+h_2-p} \frac{1}{(z-w)^p} \left[-f_5^{h_1,h_2,h_3,p}(\lambda) \partial_w^{h_1+h_2-h_3-p} \Phi_1^{(h_3),ij} \right.$$

²³ The exact relations between the components and its superfields in (2.22) can be summarized by [23, 25]

$$\Phi_0^{(h)} \leftrightarrow \Phi^{(h)}, \quad \Phi_{\frac{1}{2}}^{(h),i} \leftrightarrow D^i \Phi^{(h)}, \quad \Phi_1^{(h),ij} \leftrightarrow -\frac{1}{2!} \varepsilon^{ijkl} D^k D^l \Phi^{(h)}, \\ \Phi_{\frac{3}{2}}^{(h),i} \leftrightarrow \frac{1}{3!} \varepsilon^{ijkl} D^j D^k D^l \Phi^{(h)}, \quad \Phi_2^{(h)} \leftrightarrow \frac{1}{4!} \varepsilon^{ijkl} D^i D^j D^k D^l \Phi^{(h)}.$$

$$\begin{aligned}
& - f_6^{h_1, h_2, h_3, p}(\lambda) \partial_w^{h_1+h_2-h_3-p} \tilde{\Phi}_1^{(h_3), ij} \Big] (w) + \dots , \\
\Phi_{\frac{3}{2}}^{(h_1), i}(z) \Phi_0^{(h_2)}(w) &= \sum_{p=1}^{h_1+h_2+1} \frac{1}{(z-w)^p} \left[- \sum_{h_3=0}^{h_1+h_2-p} f_7^{h_1, h_2, h_3, p}(\lambda) \partial_w^{h_1+h_2-h_3-p} \Phi_{\frac{3}{2}}^{(h_3), i} \right. \\
& + \left. \sum_{h_3=0}^{h_1+h_2-p+1} f_8^{h_1, h_2, h_3, p}(\lambda) \partial_w^{h_1+h_2-h_3-p+1} \Phi_{\frac{1}{2}}^{(h_3), i} \right] (w) + \dots , \\
\Phi_2^{(h_1)}(z) \Phi_0^{(h_2)}(w) &= \frac{(h_1+h_2+1)!}{(z-w)^{h_1+h_2+2}} \left[(-1)^{h_1+h_2} C_{0,2}^{h_2, h_1}(\lambda) + \frac{(1-4\lambda)}{(2h_1+1)} C_{0,0}^{h_1, h_2}(\lambda) \right] \\
& + \sum_{p=1}^{h_1+h_2+1} \sum_{h_3=1}^{h_1+h_2-p+2} \frac{1}{(z-w)^p} \left[f_9^{h_1, h_2, h_3, p}(\lambda) \partial_w^{h_1+h_2-h_3-p+2} \right. \\
& \times \left. \left(\Phi_2^{(h_3-2)} - \frac{(1-4\lambda)}{(2h_3-3)} \partial_w^2 \Phi_0^{(h_3-2)} \right) \right. \\
& + \left. f_{10}^{h_1, h_2, h_3, p}(\lambda) \partial_w^{h_1+h_2-h_3-p+2} \Phi_0^{(h_3)} \right] (w) + \dots . \tag{C.1}
\end{aligned}$$

For fixed $\Phi_0^{(h_2)}(w)$, all the five components with z dependence are acted in (C.1) and some of the coefficients in (2.29) and (B.2) are used. Based on (C.1) together with the footnote 23, the $\mathcal{N}=4$ super OPE (2.30) can be determined completely.

The λ dependent structure constants appearing in (C.1) are given by

$$\begin{aligned}
f_1^{h_1, h_2, h_3, p}(\lambda) &\equiv \sum_{k=0}^{h_1+h_2-h_3-1} \frac{4^{2(h_1+h_2-h_3)-4} (2h_3-1)!(p-1)!}{(2h_1-1)(2h_2-1)} \\
&\times \left[(-1)^{h_1+h_2+k+1} \binom{k}{h_1+h_2-h_3-p} ((h_1-2\lambda)(h_2-2\lambda) S_{F,R}^{h_1, h_2, h_3, k} \right. \\
&- (h_1-1+2\lambda)(h_2-1+2\lambda) S_{B,R}^{h_1, h_2, h_3, k}) \\
&+ (-1)^{h_3+k+1} \binom{h_1+h_2-h_3-k-1}{h_1+h_2-h_3-p} ((h_1-2\lambda)(h_2-2\lambda) S_{F,L}^{h_1, h_2, h_3, k} \right. \\
&- (h_1-1+2\lambda)(h_2-1+2\lambda) S_{B,L}^{h_1, h_2, h_3, k}) \Big], \\
f_2^{h_1, h_2, h_3, p}(\lambda) &\equiv \sum_{k=0}^{h_1+h_2-h_3-1} \frac{4^{2(h_1+h_2-h_3)-1} (2h_3-2)!(p-1)!}{2(2h_1-1)(2h_2-1)} \left[(-1)^{h_1+h_2+k} \right. \\
&\times \binom{k}{h_1+h_2-h_3-p} ((h_1-2\lambda)(h_2-2\lambda)(h_3-1+2\lambda) S_{F,R}^{h_1, h_2, h_3, k} \\
&+ (h_1-1+2\lambda)(h_2-1+2\lambda)(h_3-2\lambda) S_{B,R}^{h_1, h_2, h_3, k}) \Big]
\end{aligned}$$

$$\begin{aligned}
& + (-1)^{h_3+k} \binom{h_1+h_2-h_3-k-1}{h_1+h_2-h_3-p} \left((h_1-2\lambda)(h_2-2\lambda)(h_3-1+2\lambda) \right. \\
& \quad \times S_{F,L}^{h_1,h_2,h_3,k} + (h_1-1+2\lambda)(h_2-1+2\lambda)(h_3-2\lambda) S_{B,L}^{h_1,h_2,h_3,k} \Big) \Big], \\
f_3^{h_1,h_2,h_3,p}(\lambda) & \equiv \sum_{k=0}^{h_1+h_2-h_3-1} \frac{2 \times 4^{2(h_1+h_2-h_3)-5} (2h_3)!(p-1)!}{(2h_2-1)} \\
& \quad \times \left[(-1)^{h_3+k+1} \binom{h_1+h_2-h_3-k-1}{h_1+h_2-h_3-p} \right. \\
& \quad \times \left((h_2-2\lambda) T_F^{h_2,h_1,h_3,k} + (h_2-1+2\lambda) \bar{T}_B^{h_2,h_1,h_3,k} \right) \\
& \quad + (-1)^{h_1+h_2+k} \binom{k}{h_1+h_2-h_3-p} \\
& \quad \times \left. \left((h_2-2\lambda) \bar{T}_{F,L}^{h_2,h_1,h_3,k} + (h_2-1+2\lambda) T_B^{h_2,h_1,h_3,k} \right) \right], \\
f_4^{h_1,h_2,h_3,p}(\lambda) & \equiv \sum_{k=0}^{h_1+h_2-h_3-2} \frac{2 \times 4^{2(h_1+h_2-h_3)-6} (2h_3+2)!(p-1)!}{(2h_2-1)} \\
& \quad \times \left[(-1)^{h_3+k+1} \binom{h_1+h_2-h_3-k-2}{h_1+h_2-h_3-p-1} \right. \\
& \quad \times \left((h_2-2\lambda) T_F^{h_2,h_1,h_3+1,k} - (h_2-1+2\lambda) \bar{T}_B^{h_2,h_1,h_3+1,k} \right) \\
& \quad + (-1)^{h_1+h_2+k} \binom{k}{h_1+h_2-h_3-p-1} \\
& \quad \times \left. \left((h_2-2\lambda) \bar{T}_{F,L}^{h_2,h_1,h_3+1,k} - (h_2-1+2\lambda) T_B^{h_2,h_1,h_3+1,k} \right) \right], \\
f_5^{h_1,h_2,h_3,p}(\lambda) & \equiv \sum_{k=0}^{h_1+h_2-h_3-1} \frac{2 \times 4^{2(h_1+h_2-h_3)-5} (2h_3+1)!(p-1)!}{(2h_2-1)} \\
& \quad \times \left[(-1)^{h_3+k} \binom{h_1+h_2-h_3-k-1}{h_1+h_2-h_3-p} \right. \\
& \quad \times \left(-(h_2-2\lambda) S_{F,R}^{h_2,h_1+1,h_3+1,k} + (h_2-1+2\lambda) S_{B,R}^{h_2,h_1+1,h_3+1,k} \right) \\
& \quad + (-1)^{h_1+h_2+k} \binom{k}{h_1+h_2-h_3-p} \\
& \quad \times \left. \left(-(h_2-2\lambda) S_{F,L}^{h_2,h_1+1,h_3+1,k} + (h_2-1+2\lambda) S_{B,L}^{h_2,h_1+1,h_3+1,k} \right) \right],
\end{aligned}$$

$$\begin{aligned}
f_6^{h_1, h_2, h_3, p}(\lambda) &\equiv \sum_{k=0}^{h_1+h_2-h_3-1} \frac{2 \times 4^{2(h_1+h_2-h_3)-5}(2h_3+1)!(p-1)!}{(2h_2-1)} \\
&\quad \times \left[(-1)^{h_3+k} \binom{h_1+h_2-h_3-k-1}{h_1+h_2-h_3-p} \right. \\
&\quad \times \left(-(h_2-2\lambda)S_{F,R}^{h_2,h_1+1,h_3+1,k} - (h_2-1+2\lambda)S_{B,R}^{h_2,h_1+1,h_3+1,k} \right) \\
&\quad + (-1)^{h_1+h_2+k} \binom{k}{h_1+h_2-h_3-p} \\
&\quad \left. \times \left(-(h_2-2\lambda)S_{F,L}^{h_2,h_1+1,h_3+1,k} - (h_2-1+2\lambda)S_{B,L}^{h_2,h_1+1,h_3+1,k} \right) \right], \\
f_7^{h_1, h_2, h_3, p}(\lambda) &\equiv -f_3^{h_1+1, h_2, h_3+1, p}(\lambda) + \sum_{k=0}^{h_1+h_2-h_3-1} \frac{4^{2(h_1+h_2-h_3)-5}(2h_3+2)!(p-1)!(1-4\lambda)}{2(2h_1+1)(2h_2-1)} \\
&\quad \times \left[(-1)^{h_3+k} \binom{h_1+h_2-h_3-k-2}{h_1+h_2-h_3-p} \right. \\
&\quad \times \left((h_2-2\lambda)T_F^{h_2,h_1,h_3+1,k} - (h_2-1+2\lambda)\bar{T}_B^{h_2,h_1,h_3+1,k} \right) \\
&\quad + (-1)^{h_1+h_2+k+1} \binom{k}{h_1+h_2-h_3-p} \\
&\quad \left. \times \left((h_2-2\lambda)\bar{T}_F^{h_2,h_1,h_3+1,k} - (h_2-1+2\lambda)T_B^{h_2,h_1,h_3+1,k} \right) \right], \\
f_8^{h_1, h_2, h_3, p}(\lambda) &\equiv -4^{-2} f_4^{h_1+1, h_2, h_3-1, p}(\lambda) \\
&\quad + \sum_{k=0}^{h_1+h_2-h_3} \frac{4^{2(h_1+h_2-h_3)-4}(2h_3)!(p-1)!(1-4\lambda)}{2(2h_1+1)(2h_2-1)} \\
&\quad \times \left[(-1)^{h_3+k} \binom{h_1+h_2-h_3-k-1}{h_1+h_2-h_3-p+1} \right. \\
&\quad \times \left((h_2-2\lambda)T_F^{h_2,h_1,h_3,k} + (h_2-1+2\lambda)\bar{T}_B^{h_2,h_1,h_3,k} \right) \\
&\quad + (-1)^{h_1+h_2+k+1} \binom{k}{h_1+h_2-h_3-p+1} \\
&\quad \left. \times \left((h_2-2\lambda)\bar{T}_F^{h_2,h_1,h_3,k} + (h_2-1+2\lambda)T_B^{h_2,h_1,h_3,k} \right) \right], \\
f_9^{h_1, h_2, h_3, p}(\lambda) &\equiv \sum_{k=0}^{h_1+h_2-h_3+1} \frac{4^{2(h_1+h_2-h_3)}(2h_3-2)!(p-1)!}{(2h_2-1)} \left[\right. \\
&\quad \times \left. \left((-1)^{h_3+k+1} \binom{h_1+h_2-h_3-k+1}{h_1+h_2-h_3-p+2} \right) \right]
\end{aligned}$$

$$\begin{aligned}
& \times \left((h_2 - 2\lambda)(h_3 - 1 + 2\lambda)S_{F,R}^{h_2,h_1+2,h_3,k} \right. \\
& - (h_2 - 1 + 2\lambda)(h_3 - 2\lambda)S_{B,R}^{h_2,h_1+2,h_3,k} \Big) \\
& + (-1)^{h_1+h_2+k+1} \binom{k}{h_1 + h_2 - h_3 - p + 2} \\
& \times \left((h_2 - 2\lambda)(h_3 - 1 + 2\lambda)S_{F,L}^{h_2,h_1+2,h_3,k} \right. \\
& - (h_2 - 1 + 2\lambda)(h_3 - 2\lambda)S_{B,L}^{h_2,h_1+2,h_3,k} \Big) \\
& + \frac{(1 - 4\lambda)}{2 \times 4(2h_1 + 1)(2h_1 - 1)} \left((-1)^{h_1+h_2+k} \binom{k}{h_1 + h_2 - h_3 - p + 2} \right. \\
& \times \left((h_1 - 2\lambda)(h_2 - 2\lambda)(h_3 - 1 + 2\lambda)S_{F,R}^{h_1,h_2,h_3,k} \right. \\
& + (h_1 - 1 + 2\lambda)(h_2 - 1 + 2\lambda)(h_3 - 2\lambda)S_{B,R}^{h_1,h_2,h_3,k} \Big) \\
& + (-1)^{h_3+k} \binom{h_1 + h_2 - h_3 - k - 1}{h_1 + h_2 - h_3 - p + 2} \\
& \times \left((h_1 - 2\lambda)(h_2 - 2\lambda)(h_3 - 1 + 2\lambda)S_{F,L}^{h_1,h_2,h_3,k} \right. \\
& + (h_1 - 1 + 2\lambda)(h_2 - 1 + 2\lambda)(h_3 - 2\lambda)S_{B,L}^{h_1,h_2,h_3,k} \Big) \Big) \Big], \\
f_{10}^{h_1,h_2,h_3,p}(\lambda) & \equiv \sum_{k=0}^{h_1+h_2-h_3+1} \frac{4^{2(h_1+h_2-h_3)-3}(2h_3-1)!(p-1)!}{(2h_2-1)} \left[\right. \\
& \times 2 \left((-1)^{h_3+k} \binom{h_1 + h_2 - h_3 - k + 1}{h_1 + h_2 - h_3 - p + 2} \right. \\
& \times \left((h_2 - 2\lambda)S_{F,R}^{h_2,h_1+2,h_3,k} + (h_2 - 1 + 2\lambda)S_{B,R}^{h_2,h_1+2,h_3,k} \right) \\
& + (-1)^{h_1+h_2+k} \binom{k}{h_1 + h_2 - h_3 - p + 2} \\
& \times \left((h_2 - 2\lambda)S_{F,L}^{h_2,h_1+2,h_3,k} + (h_2 - 1 + 2\lambda)S_{B,L}^{h_2,h_1+2,h_3,k} \right) \Big) \\
& + \frac{(1 - 4\lambda)}{4(2h_1 + 1)(2h_1 - 1)} \left((-1)^{h_1+h_2+k+1} \binom{k}{h_1 + h_2 - h_3 - p + 2} \right. \\
& \times \left((h_1 - 2\lambda)(h_2 - 2\lambda)S_{F,R}^{h_1,h_2,h_3,k} \right. \\
& - (h_1 - 1 + 2\lambda)(h_2 - 1 + 2\lambda)S_{B,R}^{h_1,h_2,h_3,k} \Big) \\
& - (-1)^{h_3+k} \binom{h_1 + h_2 - h_3 - k - 1}{h_1 + h_2 - h_3 - p + 2} \Big) \Big]
\end{aligned}$$

$$\begin{aligned} & \times \left((h_1 - 2\lambda)(h_2 - 2\lambda) S_{F,L}^{h_1, h_2, h_3, k} \right. \\ & \left. - (h_1 - 1 + 2\lambda)(h_2 - 1 + 2\lambda) S_{B,L}^{h_1, h_2, h_3, k} \right) \Bigg). \end{aligned} \quad (\text{C.2})$$

Again, the arguments λ in (A.4) are ignored in (C.2) for convenience. The summation over k is put to these structure constants with other k dependent factors.

The $\mathcal{N} = 4$ super OPE between the $\mathcal{N} = 4$ stress energy tensor and the $\mathcal{N} = 4$ multiplet can be summarized by

$$\begin{aligned} \mathbf{J}(Z_1) \Phi^{(h)}(Z_2) &= \frac{1}{z_{12}^h} C_1^h(\lambda) + \frac{\theta_{12}^{4-0}}{z_{12}^{h+2}} C_2^h(\lambda) \\ &+ \sum_{p=1}^h \frac{1}{z_{12}^p} \left[\sum_{h_3=1}^{h-p} g_1^{h, h_3, p}(\lambda) \partial^{h-h_3-p} \Phi^{(h_3)} + \sum_{h_3=2}^{h-p} g_2^{h, h_3, p}(\lambda) \partial^{h-h_3-p} D^{4-0} \Phi^{(h_3-2)} \right. \\ &\quad \left. + v_1^{h, p}(\lambda) \partial^{h-p} \mathbf{J} \right] (Z_2) \\ &+ \sum_{p=1}^h \frac{\theta_{12}^i}{z_{12}^p} \left[\sum_{h_3=0}^{h-p} g_3^{h, h_3, p}(\lambda) \partial^{h-h_3-p} D^i \Phi^{(h_3)} + \sum_{h_3=0}^{h-p-1} g_4^{h, h_3, p}(\lambda) \partial^{h-h_3-p-1} D^{4-i} \Phi^{(h_3)} \right] (Z_2) \\ &+ \sum_{p=1}^h \sum_{h_3=0}^{h-p} \frac{\theta_{12}^{4-ij}}{z_{12}^p} \left[g_5^{h, h_3, p}(\lambda) D^{4-ij} + g_6^{h, h_3, p}(\lambda) D^{ij} \right] \partial^{h-h_3-p} \Phi^{(h_3)} (Z_2) \\ &+ \sum_{p=1}^{h+1} \frac{\theta_{12}^{4-i}}{z_{12}^p} \left[\sum_{h_3=0}^{h-p} g_7^{h, h_3, p}(\lambda) \partial^{h-h_3-p} D^{4-i} \Phi^{(h_3)} + \sum_{h_3=0}^{h-p+1} g_8^{h, h_3, p}(\lambda) \partial^{h-h_3-p+1} D^i \Phi^{(h_3)} \right] (Z_2) \\ &+ \sum_{p=1}^{h+1} \frac{\theta_{12}^{4-0}}{z_{12}^p} \left[\sum_{h_3=2}^{h-p+2} g_9^{h, h_3, p}(\lambda) \partial^{h-h_3-p+2} D^{4-0} \Phi^{(h_3-2)} + \sum_{h_3=1}^{h-p+2} g_{10}^{h, h_3, p}(\lambda) \partial^{h-h_3-p+2} \Phi^{(h_3)} \right. \\ &\quad \left. + v_2^{h, p}(\lambda) \partial^{h-p+2} \mathbf{J} \right] (Z_2) + \dots \end{aligned} \quad (\text{C.3})$$

All the terms in (C.3) should have the spin h . On the right hand side, there exists the $\mathcal{N} = 4$ stress energy tensor \mathbf{J} .

The central terms and the structure constants appearing in (C.3) are given by

$$\begin{aligned} C_1^h(\lambda) &\equiv -\frac{N 4^{2(h-2)} (h-1)!}{(2h-1)} \sum_{i=0}^{h-1} \frac{i!}{(i+1)!} \left[(h-2\lambda) a^i(h, \lambda + \frac{1}{2}) + (h-1+2\lambda) a^i(h, \lambda) \right], \\ C_2^h(\lambda) &\equiv -\frac{N 4^{2(h-2)} (h+1)!}{(2h-1)} \sum_{i=0}^{h-1} \frac{1}{(i+1)(i+2)} \left[(h-2\lambda)(2+3i) a^i(h, \lambda + \frac{1}{2}) \right. \\ &\quad \left. + (h-1+2\lambda)(2+3i) a^i(h, \lambda) \right]. \end{aligned}$$

$$\begin{aligned}
& + (h - 1 + 2\lambda)(i - 2)a^i(h, \lambda) \Big] , \\
g_1^{h,h_3,p}(\lambda) & \equiv \frac{256}{p} \left[f_{10}^{-1,h,h_3,p+1}(\lambda) - \frac{(1 - 4\lambda)}{(2h_3 + 1)} f_9^{-1,h,h_3+2,p+1}(\lambda) \right] , \\
g_2^{h,h_3,p}(\lambda) & \equiv \frac{256}{p} f_9^{-1,h,h_3,p+1}(\lambda) , \\
g_3^{h,h_3,p}(\lambda) & \equiv 128 \left[f_3^{0,h,h_3,p}(\lambda) + \frac{(1 - 4\lambda)}{(2h_3 + 1)} f_4^{0,h,h_3,p}(\lambda) \right] , \\
g_4^{h,h_3,p}(\lambda) & \equiv 128 f_4^{0,h,h_3,p}(\lambda) , \\
g_5^{h,h_3,p}(\lambda) & \equiv 64 f_5^{0,h,h_3,p}(\lambda) , \\
g_6^{h,h_3,p}(\lambda) & \equiv 64 f_6^{0,h,h_3,p}(\lambda) , \\
g_7^{h,h_3,p}(\lambda) & \equiv \frac{4^{2(h-h_3)-1}}{2(2h-1)} (2h_3+2)!(p-1)! \sum_{k=0}^{h-h_3-1} \left[(-1)^{h_3+k+1} \binom{h-h_3-k-1}{h-h_3-p} \right. \\
& \quad \times \left((h-2\lambda) T_F^{h,1,h_3+1,k}(\lambda) + (h-1+2\lambda) \bar{T}_B^{h,1,h_3+1,k}(\lambda) \right) \\
& \quad + (-1)^{h_2+k} \binom{k}{h-h_3-p} \left((h-1+2\lambda) T_B^{h,1,h_3+1,k}(\lambda) + (h-2\lambda) \bar{T}_F^{h,1,h_3+1,k}(\lambda) \right) \\
& \quad - \frac{(1-4\lambda)}{2} \left((-1)^{h+k} \binom{k}{h-h_3-p} (h-2\lambda) \bar{T}_F^{h,0,h_3+1,k}(\lambda) \right. \\
& \quad \left. \left. + (-1)^{h_3+k} \binom{h-h_3-k-2}{h-h_3-p} (h-1+2\lambda) \bar{T}_B^{h,0,h_3+1,k}(\lambda) \right) \right] , \\
g_8^{h,h_3,p}(\lambda) & \equiv -\frac{4^{2(h-h_3)}}{2(2h-1)} (2h_3)!(p-1)! \sum_{k=0}^{h-h_3} \left[(-1)^{h_3+k} \binom{h-h_3-k}{h-h_3-p+1} \right. \\
& \quad \times \left((h-2\lambda) T_F^{h,1,h_3,k}(\lambda) - (h-1+2\lambda) \bar{T}_B^{h,1,h_3,k}(\lambda) \right) \\
& \quad + (-1)^{h_2+k} \binom{k}{h-h_3-p+1} \left((h-1+2\lambda) T_B^{h,1,h_3,k}(\lambda) - (h-2\lambda) \bar{T}_F^{h,1,h_3,k}(\lambda) \right) \\
& \quad + \frac{(1-4\lambda)}{2} \left((-1)^{h+k} \binom{k}{h-h_3-p+1} (h-2\lambda) \bar{T}_F^{h,0,h_3,k}(\lambda) \right. \\
& \quad \left. \left. - (-1)^{h_3+k} \binom{h-h_3-k-1}{h-h_3-p+1} (h-1+2\lambda) \bar{T}_B^{h,0,h_3,k}(\lambda) \right) \right] + \frac{(1-4\lambda)}{(2h_3+1)} g_7^{h,h_3,p}(\lambda) , \\
g_9^{h,h_3,p}(\lambda) & \equiv \frac{4^{2(h-h_3)+3}}{(2h-1)} (2h_3-2)!(p-1)! \sum_{k=0}^{h-h_3+1} \left[(-1)^{h_2+k+1} \binom{k}{h-h_3-p+2} \right]
\end{aligned}$$

$$\begin{aligned}
& \times \left((h - 2\lambda)(h_3 - 1 + 2\lambda) S_{F,R}^{2,h,h_3,k}(\lambda) - (h - 1 + 2\lambda)(h_3 - 2\lambda) S_{B,R}^{2,h,h_3,k}(\lambda) \right) \\
& - (-1)^{h_3+k} \binom{h - h_3 - k + 1}{h - h_3 - p + 2} \left((h - 2\lambda)(h_3 - 1 + 2\lambda) S_{F,L}^{2,h,h_3,k}(\lambda) \right. \\
& - (h - 1 + 2\lambda)(h_3 - 2\lambda) S_{B,L}^{2,h,h_3,k}(\lambda) \Big) \\
& + \frac{(1 - 4\lambda)}{4} (-1)^{h+k} \binom{k}{h - h_3 - p + 2} \left((h - 2\lambda)(h_3 - 1 + 2\lambda) S_{F,R}^{1,h,h_3,k}(\lambda) \right. \\
& - (h - 1 + 2\lambda)(h_3 - 2\lambda) S_{B,R}^{1,h,h_3,k}(\lambda) \Big) \\
& - \frac{(1 - 4\lambda)}{4} (-1)^{h_3+k} \binom{h - h_3 - k}{h - h_3 - p + 2} \left((h - 2\lambda)(h_3 - 1 + 2\lambda) S_{F,L}^{1,h,h_3,k}(\lambda) \right. \\
& - (h - 1 + 2\lambda)(h_3 - 2\lambda) S_{B,L}^{1,h,h_3,k}(\lambda) \Big) \Big], \\
g_{10}^{h,h_3,p}(\lambda) & \equiv \frac{4^{2(h-h_3)+1}}{2(2h-1)} (2h_3-1)!(p-1)! \sum_{k=0}^{h-h_3+1} \left[(-1)^{h_2+k} \binom{k}{h - h_3 - p + 2} \right. \\
& \times \left((h - 2\lambda) S_{F,R}^{2,h,h_3,k}(\lambda) + (h - 1 + 2\lambda) S_{B,R}^{2,h,h_3,k}(\lambda) \right) \\
& + (-1)^{h_3+k} \binom{h - h_3 - k + 1}{h - h_3 - p + 2} \left((h - 2\lambda) S_{F,L}^{2,h,h_3,k}(\lambda) + (h - 1 + 2\lambda) S_{B,L}^{2,h,h_3,k}(\lambda) \right) \\
& - \frac{(1 - 4\lambda)}{4} (-1)^{h+k} \binom{k}{h - h_3 - p + 2} \left((h - 2\lambda) S_{F,R}^{1,h,h_3,k}(\lambda) \right. \\
& + (h - 1 + 2\lambda) S_{B,R}^{1,h,h_3,k}(\lambda) \Big) \\
& + \frac{(1 - 4\lambda)}{4} (-1)^{h_3+k} \binom{h - h_3 - k}{h - h_3 - p + 2} \left((h - 2\lambda) S_{F,L}^{1,h,h_3,k}(\lambda) \right. \\
& + (h - 1 + 2\lambda) S_{B,L}^{1,h,h_3,k}(\lambda) \Big) \Big] - \frac{(1 - 4\lambda)}{(2h_3 + 1)} g_9^{h,h_3+2,p}(\lambda), \\
v_1^{h,p}(\lambda) & \equiv -\frac{1}{p} \left[f_9^{-1,h,1,p+1}(\lambda) + 4(1 - 4\lambda) f_9^{-1,h,2,p+1}(\lambda) \right], \\
v_2^{h,p}(\lambda) & \equiv -\frac{1}{256} \left[g_9^{h,1,p}(\lambda) + 4(1 - 4\lambda) g_9^{h,2,p}(\lambda) \right], \tag{C.4}
\end{aligned}$$

where the previous relations (A.4) and (C.2) are used in (C.4).

D The $\mathcal{N} = 2$ supersymmetric linear $W_\infty^{K,K}[\lambda]$ algebra

The whole 26 (anti)commutators are summarized by

$$\begin{aligned}
& [(W_{F,h_1}^\lambda)_m, (W_{F,h_2}^\lambda)_n] = K c_{W_F}^{h_1,h_2}(\lambda) q^{h_1+h_2-4} [m+h_1-1]_{h_1+h_2-1} \delta_{m+n} \\
& + \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
& \times \left(S_{F,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \right. \\
& \left. - S_{F,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \right) (W_{F,h_3}^\lambda)_{m+n}, \\
& [(W_{F,h_1}^\lambda)_m, (W_{F,h_2}^{\lambda,\hat{A}})_n] = \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
& \times \left(S_{F,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \right. \\
& \left. - S_{F,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \right) (W_{F,h_3}^{\lambda,\hat{A}})_{m+n}, \\
& [(W_{F,h_1}^{\lambda,\hat{A}})_m, (W_{F,h_2}^{\lambda,\hat{B}})_n] = \delta^{\hat{A}\hat{B}} c_{W_F}^{h_1,h_2}(\lambda) q^{h_1+h_2-4} [m+h_1-1]_{h_1+h_2-1} \delta_{m+n} \\
& + \frac{1}{2} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
& \times \left[\left(S_{F,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \right. \right. \\
& \left. \left. - S_{F,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \right) \right. \\
& \times \left(\frac{2}{K} \delta^{\hat{A}\hat{B}} (W_{F,h_3}^\lambda)_{m+n} + d^{\hat{A}\hat{B}\hat{C}} (W_{F,h_3}^{\lambda,\hat{C}})_{m+n} \right) \\
& + \left(S_{F,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \right. \\
& \left. + S_{F,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \right) \\
& \left. \times i f^{\hat{A}\hat{B}\hat{C}} (W_{F,h_3}^{\lambda,\hat{C}})_{m+n} \right], \\
& [(W_{B,h_1}^\lambda)_m, (W_{B,h_2}^\lambda)_n] = K c_{W_B}^{h_1,h_2}(\lambda) q^{h_1+h_2-4} [m+h_1-1]_{h_1+h_2-1} \delta_{m+n} \\
& + \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3-1)!
\end{aligned}$$

$$\begin{aligned}
& \times \left(\begin{array}{l} S_{B,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \\ - S_{B,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \end{array} \right) (W_{B,h_3}^\lambda)_{m+n}, \\
\left[(W_{B,h_1}^\lambda)_m, (W_{B,h_2}^{\lambda,\hat{A}})_n \right] &= \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
&\quad \times \left(\begin{array}{l} S_{B,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \\ - S_{B,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \end{array} \right) (W_{B,h_3}^{\lambda,\hat{A}})_{m+n}, \\
\left[(W_{B,h_1}^{\lambda,\hat{A}})_m, (W_{B,h_2}^{\lambda,\hat{B}})_n \right] &= \delta^{\hat{A}\hat{B}} c_{W_B}^{h_1,h_2}(\lambda) q^{h_1+h_2-4} [m+h_1-1]_{h_1+h_2-1} \delta_{m+n} \\
&\quad + \frac{1}{2} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3-1)! \\
&\quad \times \left[\left(\begin{array}{l} S_{B,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \\ - S_{B,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \end{array} \right) \right. \\
&\quad \times \left(\frac{2}{K} \delta^{\hat{A}\hat{B}} (W_{B,h_3}^\lambda)_{m+n} + d^{\hat{A}\hat{B}\hat{C}} (W_{B,h_3}^{\lambda,\hat{C}})_{m+n} \right) \\
&\quad + \left(\begin{array}{l} S_{B,R}^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [n+h_2-1]_k \\ + S_{B,L}^{h_1,h_2,h_3,k} [m+h_1-1]_k [n+h_2-1]_{h_1+h_2-h_3-k-1} \end{array} \right) \\
&\quad \times i f^{\hat{A}\hat{B}\hat{C}} (W_{B,h_3}^{\lambda,\hat{C}})_{m+n} \Big], \\
\left[(W_{F,h_1}^\lambda)_m, (Q_{h_2+\frac{1}{2}}^\lambda)_r \right] &= \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
&\quad \times T_F^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k (Q_{h_3+\frac{1}{2}}^\lambda)_{m+r}, \\
\left[(W_{F,h_1}^\lambda)_m, (Q_{h_2+\frac{1}{2}}^{\lambda,\hat{A}})_r \right] &= \left[(W_{F,h_1}^{\lambda,\hat{A}})_m, (Q_{h_2+\frac{1}{2}}^\lambda)_r \right] \\
&= \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
&\quad \times T_F^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k (Q_{h_3+\frac{1}{2}}^{\lambda,\hat{A}})_{m+r},
\end{aligned}$$

$$\begin{aligned}
& \left[(W_{F,h_1}^{\lambda,\hat{A}})_m, (Q_{h_2+\frac{1}{2}}^{\lambda,\hat{B}})_r \right] = \frac{1}{2} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times T_F^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k \\
& \quad \times \left(\frac{2}{K} \delta^{\hat{A}\hat{B}} (Q_{h_3+\frac{1}{2}}^{\lambda})_{m+r} + (if^{\hat{A}\hat{B}\hat{C}} + d^{\hat{A}\hat{B}\hat{C}}) (Q_{h_3+\frac{1}{2}}^{\lambda,\hat{C}})_{m+r} \right), \\
& \left[(W_{F,h_1}^{\lambda})_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda})_r \right] = \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_F^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} (\bar{Q}_{h_3+\frac{1}{2}}^{\lambda})_{m+r}, \\
& \left[(W_{F,h_1}^{\lambda})_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\hat{A}})_r \right] = \left[(W_{F,h_1}^{\lambda,\hat{A}})_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda})_r \right] \\
& \quad = \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_F^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} (\bar{Q}_{h_3+\frac{1}{2}}^{\lambda,\hat{A}})_{m+r}, \\
& \left[(W_{F,h_1}^{\lambda,\hat{A}})_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\hat{B}})_r \right] = \frac{1}{2} \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_F^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} \\
& \quad \times \left(\frac{2}{K} \delta^{\hat{A}\hat{B}} (Q_{h_3+\frac{1}{2}}^{\lambda})_{m+r} - (if^{\hat{A}\hat{B}\hat{C}} - d^{\hat{A}\hat{B}\hat{C}}) (Q_{h_3+\frac{1}{2}}^{\lambda,\hat{C}})_{m+r} \right), \\
& \left[(W_{B,h_1}^{\lambda})_m, (Q_{h_2+\frac{1}{2}}^{\lambda})_r \right] = \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times T_B^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} (Q_{h_3+\frac{1}{2}}^{\lambda})_{m+r}, \\
& \left[(W_{B,h_1}^{\lambda})_m, (Q_{F,h_2+\frac{1}{2}}^{\lambda,\hat{A}})_r \right] = \left[(W_{B,h_1}^{\lambda,\hat{A}})_m, (Q_{h_2+\frac{1}{2}}^{\lambda})_r \right] \\
& \quad = \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times T_B^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} (Q_{h_3+\frac{1}{2}}^{\lambda,\hat{A}})_{m+r}, \\
& \left[(W_{B,h_1}^{\lambda,\hat{A}})_m, (Q_{h_2+\frac{1}{2}}^{\lambda,\hat{B}})_r \right] = \frac{1}{2} \sum_{h_3=1}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times T_B^{h_1,h_2,h_3,k} [m+h_1-1]_k [r+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k-1} \\
& \quad \times \left(\frac{2}{K} \delta^{\hat{A}\hat{B}} (Q_{h_3+\frac{1}{2}}^{\lambda})_{m+r} - (if^{\hat{A}\hat{B}\hat{C}} - d^{\hat{A}\hat{B}\hat{C}}) (Q_{h_3+\frac{1}{2}}^{\lambda,\hat{C}})_{m+r} \right),
\end{aligned}$$

$$\begin{aligned}
& \left[(W_{B,h_1}^\lambda)_m, (\bar{Q}_{h_2+\frac{1}{2}}^\lambda)_r \right] = \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_B^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k (\bar{Q}_{h_3+\frac{1}{2}}^\lambda)_{m+r}, \\
& \left[(W_{B,h_1}^\lambda)_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\hat{A}})_r \right] = \left[(W_{B,h_1}^{\lambda,\hat{A}})_m, (\bar{Q}_{h_2+\frac{1}{2}}^\lambda)_r \right] \\
& = \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_B^{h_1,h_2,h_3,k} [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k (\bar{Q}_{h_3+\frac{1}{2}}^{\lambda,\hat{A}})_{m+r}, \\
& \left[(W_{B,h_1}^{\lambda,\hat{A}})_m, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\hat{B}})_r \right] = \sum_{h_3=0}^{h_1+h_2-1} \sum_{k=0}^{h_1+h_2-h_3-1} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3-2} (2h_3)! \\
& \quad \times \bar{T}_B^{h_1,h_2,h_3,k}(\lambda) [m+h_1-1]_{h_1+h_2-h_3-k-1} [r+h_2-\frac{1}{2}]_k \\
& \quad \cdot \times \left(\frac{2}{K} \delta^{\hat{A}\hat{B}} (Q_{h_3+\frac{1}{2}}^\lambda)_{m+r} + (if^{\hat{A}\hat{B}\hat{C}} + d^{\hat{A}\hat{B}\hat{C}}) (Q_{h_3+\frac{1}{2}}^{\lambda,\hat{C}})_{m+r} \right), \\
& \left\{ (Q_{h_1+\frac{1}{2}}^\lambda)_r, (\bar{Q}_{h_2+\frac{1}{2}}^\lambda)_s \right\} = K c_{Q\bar{Q}}^{h_1,h_2}(\lambda) q^{h_1+h_2-2} [r+h_1-\frac{1}{2}]_{h_1+h_2} \delta_{r+s} \\
& \quad + \sum_{h_3=1}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} 2 (-1)^{h_3+1} (4q)^{h_1+h_2-h_3} (2h_3-1)! \\
& \quad \times \left(U_F^{h_1,h_2,h_3,k} [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k} [s+h_2-\frac{1}{2}]_k (W_{F,h_3}^\lambda)_{r+s} \right. \\
& \quad \left. + U_B^{h_1,h_2,h_3,k} [r+h_1-\frac{1}{2}]_k [s+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k} (W_{B,h_3}^\lambda)_{r+s} \right), \\
& \left\{ (Q_{h_1+\frac{1}{2}}^{\lambda,\hat{A}})_r, (\bar{Q}_{h_2+\frac{1}{2}}^\lambda)_s \right\} = \left\{ (Q_{h_2+\frac{1}{2}}^\lambda)_r, (\bar{Q}_{h_1+\frac{1}{2}}^{\lambda,\hat{A}})_s \right\} \\
& = \sum_{h_3=1}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} 2 (-1)^{h_3+1} (4q)^{h_1+h_2-h_3} (2h_3-1)! \\
& \quad \times \left(U_F^{h_1,h_2,h_3,k} [r+h_1-\frac{1}{2}]_{h_1+h_2-h_3-k} [s+h_2-\frac{1}{2}]_k (W_{F,h_3}^{\lambda,\hat{A}})_{r+s} \right. \\
& \quad \left. + U_B^{h_1,h_2,h_3,k} [r+h_1-\frac{1}{2}]_k [s+h_2-\frac{1}{2}]_{h_1+h_2-h_3-k} (W_{B,h_3}^{\lambda,\hat{A}})_{r+s} \right), \\
& \left\{ (Q_{h_1+\frac{1}{2}}^{\lambda,\hat{A}})_r, (\bar{Q}_{h_2+\frac{1}{2}}^{\lambda,\hat{B}})_s \right\} = \delta^{\hat{A}\hat{B}} c_{Q\bar{Q}}^{h_1,h_2}(\lambda) q^{h_1+h_2-2} [r+h_1-\frac{1}{2}]_{h_1+h_2} \delta_{r+s} \\
& \quad + \sum_{h_3=1}^{h_1+h_2} \sum_{k=0}^{h_1+h_2-h_3} (-1)^{h_3+1} (4q)^{h_1+h_2-h_3} (2h_3-1)!
\end{aligned}$$

$$\begin{aligned}
& \times \left[\left(U_F^{h_1, h_2, h_3, k} [r + h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k} [s + h_2 - \frac{1}{2}]_k \right. \right. \\
& + U_B^{h_1, h_2, h_3, k} [r + h_1 - \frac{1}{2}]_k [s + h_2 - \frac{1}{2}]_{h_1+h_2-h_3-k} \Big) \\
& \times \left(\frac{2}{K} \delta^{\hat{A}\hat{B}} (W_{F,h_3}^\lambda)_{r+s} + d^{\hat{A}\hat{B}\hat{C}} (W_{F,h_3}^{\lambda,\hat{C}})_{r+s} \right) \\
& + \left(U_F^{h_1, h_2, h_3, k} [r + h_1 - \frac{1}{2}]_{h_1+h_2-h_3-k} [s + h_2 - \frac{1}{2}]_k \right. \\
& - U_B^{h_1, h_2, h_3, k} [r + h_1 - \frac{1}{2}]_k [s + h_2 - \frac{1}{2}]_{h_1+h_2-h_3-k} \Big) \\
& \left. \left. \times i f^{\hat{A}\hat{B}\hat{C}} (W_{B,h_3}^{\lambda,\hat{C}})_{r+s} \right] . \tag{D.1}
\end{aligned}$$

The central terms are given by (2.21) and (A.7). For $K \geq 3$, there are nonzero d symbols. In (D.1), we do not write down the λ dependences appearing in (A.4) explicitly in order to save the space ²⁴ and again, the structure constants are given by (A.4).

We continue to present the description in (3.5) for other cases having the adjoint indices and we obtain the following results for several h_1 and h_2

$$\begin{aligned}
W_{F,1}^\lambda(z) W_{F,1}^{\lambda,\hat{A}}(w) &= 0 + \dots , \\
W_{F,1}^\lambda(z) W_{F,2}^{\lambda,\hat{A}}(w) &= \frac{1}{(z-w)^2} W_{F,1}^{\lambda,\hat{A}}(w) + \dots , \\
W_{F,1}^\lambda(z) W_{F,3}^{\lambda,\hat{A}}(w) &= \frac{1}{(z-w)^3} 8 q \lambda W_{F,1}^{\lambda,\hat{A}}(w) + \frac{1}{(z-w)^2} \left[2 W_{F,2}^{\lambda,\hat{A}} - 4 q \lambda \partial_w W_{F,1}^{\lambda,\hat{A}} \right](w) + \dots , \\
W_{F,2}^\lambda(z) W_{F,2}^{\lambda,\hat{A}}(w) &= \frac{1}{(z-w)^2} 2 W_{F,2}^{\lambda,\hat{A}}(w) + \frac{1}{(z-w)} \partial_w W_{F,2}^{\lambda,\hat{A}}(w) + \dots , \\
W_{F,2}^\lambda(z) W_{F,3}^{\lambda,\hat{A}}(w) &= \frac{1}{(z-w)^4} 16 q^2 (1-2\lambda)(1+2\lambda) W_{F,1}^{\lambda,\hat{A}}(w) + \frac{1}{(z-w)^2} 3 W_{F,3}^{\lambda,\hat{A}}(w) \\
&+ \frac{1}{(z-w)} \partial_w W_{F,3}^{\lambda,\hat{A}}(w) + \dots , \\
W_{F,3}^\lambda(z) W_{F,3}^{\lambda,\hat{A}}(w) &= \frac{1}{(z-w)^4} 64 q^2 (1-\lambda)(1+\lambda) W_{F,2}^{\lambda,\hat{A}}(w) \\
&+ \frac{1}{(z-w)^3} 32 q^2 (1-\lambda)(1+\lambda) \partial_w W_{F,2}^{\lambda,\hat{A}}(w) \\
&+ \frac{1}{(z-w)^2} \left[4 W_{F,4}^{\lambda,\hat{A}} + \frac{48 q^2}{5} (1-\lambda)(1+\lambda) \partial_w^2 W_{F,2}^{\lambda,\hat{A}} \right](w)
\end{aligned}$$

²⁴The corresponding OPEs can be obtained from the description of the footnote 12.

$$\begin{aligned}
& + \frac{1}{(z-w)} \left[2 \partial_w W_{F,4}^{\lambda,\hat{A}} + \frac{32 q^2}{15} (1-\lambda)(1+\lambda) \partial_w^3 W_{F,2}^{\lambda,\hat{A}} \right] (w) + \cdots , \\
W_{F,1}^{\lambda,\hat{A}}(z) W_{F,1}^{\lambda,\hat{B}}(w) & = \frac{1}{(z-w)^2} \delta^{\hat{A}\hat{B}} \frac{N}{16 q^2} - \frac{1}{(z-w)} \frac{i}{4q} f^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}}(w) + \cdots , \\
W_{F,1}^{\lambda,\hat{A}}(z) W_{F,2}^{\lambda,\hat{B}}(w) & = \frac{1}{(z-w)^3} \delta^{\hat{A}\hat{B}} \frac{N \lambda}{2q} \\
& + \frac{1}{(z-w)^2} \left[\delta^{\hat{A}\hat{B}} \frac{1}{K} W_{F,1}^{\lambda} + \frac{1}{2} d^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}} - i \lambda f^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}} \right] (w) \\
& - \frac{1}{(z-w)} \frac{i}{4q} f^{\hat{A}\hat{B}\hat{C}} W_{F,2}^{\lambda,\hat{C}}(w) + \cdots , \\
W_{F,1}^{\lambda,\hat{A}}(z) W_{F,3}^{\lambda,\hat{B}}(w) & = \frac{1}{(z-w)^4} 2 N \lambda^2 \delta^{\hat{A}\hat{B}} \\
& + \frac{1}{(z-w)^3} \left[\delta^{\hat{A}\hat{B}} \frac{8q\lambda}{K} W_{F,1}^{\lambda} + 4q\lambda d^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}} \right. \\
& \left. - \frac{4q}{3}(1+2\lambda^2) i f^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}} \right] (w) \\
& + \frac{1}{(z-w)^2} \left[\delta^{\hat{A}\hat{B}} \left(\frac{2}{K} W_{F,2}^{\lambda} - q \frac{4\lambda}{K} \partial_w W_{F,1}^{\lambda} \right) + d^{\hat{A}\hat{B}\hat{C}} \left(W_{F,2}^{\lambda,\hat{C}} - 2q\lambda \partial_w W_{F,1}^{\lambda,\hat{C}} \right) \right. \\
& \left. - i f^{\hat{A}\hat{B}\hat{C}} \left(\lambda W_{F,2}^{\lambda,\hat{C}} - \frac{2q}{3}(1+2\lambda^2) \partial_w W_{F,1}^{\lambda,\hat{C}} \right) \right] (w) \\
& - \frac{1}{(z-w)} \frac{i}{4q} f^{\hat{A}\hat{B}\hat{C}} W_{F,3}^{\lambda,\hat{C}}(w) + \cdots , \\
W_{F,2}^{\lambda,\hat{A}}(z) W_{F,2}^{\lambda,\hat{B}}(w) & = \frac{1}{(z-w)^4} \frac{N(1-12\lambda^2)}{2} \delta^{\hat{A}\hat{B}} \\
& - \frac{1}{(z-w)^3} 2q(1-2\lambda)(1+2\lambda) i f^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}}(w) \\
& + \frac{1}{(z-w)^2} \left[\delta^{\hat{A}\hat{B}} \frac{2}{K} W_{F,2}^{\lambda} + d^{\hat{A}\hat{B}\hat{C}} W_{F,2}^{\lambda,\hat{C}} \right. \\
& \left. - q(1-2\lambda)(1+2\lambda) i f^{\hat{A}\hat{B}\hat{C}} \partial_w W_{F,1}^{\lambda,\hat{C}} \right] (w) \\
& + \frac{1}{(z-w)} \left[\delta^{\hat{A}\hat{B}} \frac{1}{K} \partial_w W_{F,2}^{\lambda} + \frac{1}{2} d^{\hat{A}\hat{B}\hat{C}} \partial_w W_{F,2}^{\lambda,\hat{C}} \right. \\
& \left. - i f^{\hat{A}\hat{B}\hat{C}} \left(\frac{1}{4q} W_{F,3}^{\lambda,\hat{C}} + \frac{q}{3}(1-2\lambda)(1+2\lambda) \partial_w^2 W_{F,1}^{\lambda,\hat{C}} \right) \right] (w) + \cdots ,
\end{aligned}$$

$$\begin{aligned}
W_{F,2}^{\lambda,\hat{A}}(z) W_{F,3}^{\lambda,\hat{B}}(w) &= \frac{1}{(z-w)^5} \delta^{\hat{A}\hat{B}} 8 q N \lambda(1-2\lambda)(1+2\lambda) \\
&\quad + \frac{1}{(z-w)^4} \left[\delta^{\hat{A}\hat{B}} \frac{16 q^2 (1-2\lambda)(1+2\lambda)}{K} W_{F,1}^\lambda \right. \\
&\quad + 8 q^2 (1-2\lambda)(1+2\lambda) d^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}} \\
&\quad \left. - 8 q^2 \lambda(1-2\lambda)(1+2\lambda) i f^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}} \right] (w) \\
&\quad - \frac{1}{(z-w)^3} \frac{16 q}{3} (1-\lambda)(1+\lambda) i f^{\hat{A}\hat{B}\hat{C}} W_{F,2}^{\lambda,\hat{C}} (w) \\
&\quad + \frac{1}{(z-w)^2} \left[\delta^{\hat{A}\hat{B}} \frac{3}{K} W_{F,3}^\lambda + \frac{3}{2} d^{\hat{A}\hat{B}\hat{C}} W_{F,3}^{\lambda,\hat{C}} \right. \\
&\quad \left. - \frac{4 q}{3} (1-\lambda)(1+\lambda) i f^{\hat{A}\hat{B}\hat{C}} \partial_w W_{F,2}^{\lambda,\hat{C}} \right] (w) \\
&\quad + \frac{1}{(z-w)} \left[\delta^{\hat{A}\hat{B}} \frac{1}{K} \partial_w W_{F,3}^\lambda + \frac{1}{2} d^{\hat{A}\hat{B}\hat{C}} \partial_w W_{F,3}^{\lambda,\hat{C}} \right. \\
&\quad \left. - i f^{\hat{A}\hat{B}\hat{C}} \left(\frac{1}{4 q} W_{F,4}^{\lambda,\hat{C}} + \frac{4 q}{15} (1-\lambda)(1+\lambda) \partial_w^2 W_{F,2}^{\lambda,\hat{C}} \right) \right] (w) + \dots ,
\end{aligned}$$

$$\begin{aligned}
W_{F,3}^{\lambda,\hat{A}}(z) W_{F,3}^{\lambda,\hat{B}}(w) &= \frac{1}{(z-w)^6} \frac{32 q^2 N (1-15\lambda^2+20\lambda^4)}{3} \delta^{\hat{A}\hat{B}} \\
&\quad - \frac{1}{(z-w)^5} \frac{128 q^3}{3} (1-\lambda)(1+\lambda)(1-2\lambda)(1+2\lambda) i f^{\hat{A}\hat{B}\hat{C}} W_{F,1}^{\lambda,\hat{C}} (w) \\
&\quad + \frac{1}{(z-w)^4} \left[\delta^{\hat{A}\hat{B}} \frac{64 q^2 (1-\lambda)(1+\lambda)}{K} W_{F,2}^\lambda + 32 q^2 (1-\lambda)(1+\lambda) d^{\hat{A}\hat{B}\hat{C}} W_{F,2}^{\lambda,\hat{C}} \right. \\
&\quad \left. - \frac{64 q^3}{3} (1-\lambda)(1+\lambda)(1-2\lambda)(1+2\lambda) i f^{\hat{A}\hat{B}\hat{C}} \partial_w W_{F,1}^{\lambda,\hat{C}} \right] (w) \\
&\quad + \frac{1}{(z-w)^3} \left[\delta^{\hat{A}\hat{B}} \frac{32 q^2 (1-\lambda)(1+\lambda)}{K} \partial_w W_{F,2}^\lambda \right. \\
&\quad \left. + 16 q^2 (1-\lambda)(1+\lambda) d^{\hat{A}\hat{B}\hat{C}} \partial_w W_{F,2}^{\lambda,\hat{C}} - i f^{\hat{A}\hat{B}\hat{C}} \left(\frac{8 q}{3} (2-\lambda)(2+\lambda) W_{F,3}^{\lambda,\hat{C}} \right. \right. \\
&\quad \left. \left. + \frac{64 q^3}{3} (1-\lambda)(1+\lambda)(1-2\lambda)(1+2\lambda) \partial_w^2 W_{F,1}^{\lambda,\hat{C}} \right) \right] (w) \\
&\quad + \frac{1}{(z-w)^2} \left[\delta^{\hat{A}\hat{B}} \left(\frac{4}{K} W_{F,4}^\lambda + \frac{48 q^2 (1-\lambda)(1+\lambda)}{5 K} \partial_w^2 W_{F,2}^\lambda \right) \right]
\end{aligned}$$

$$\begin{aligned}
& + d^{\hat{A}\hat{B}\hat{C}} \left(2 W_{F,4}^{\lambda,\hat{C}} + \frac{24}{5} (1-\lambda)(1+\lambda) \partial_w^2 W_{F,2}^{\lambda,\hat{C}} \right) \\
& - i f^{\hat{A}\hat{B}\hat{C}} \left(\frac{4}{3} q (2-\lambda)(2+\lambda) \partial_w W_{F,3}^{\lambda,\hat{C}} \right. \\
& \left. + \frac{16}{9} q^3 (1-\lambda)(1+\lambda)(1-2\lambda)(1+2\lambda) \partial_w^3 W_{F,1}^{\lambda,\hat{C}} \right) \Big] (w) \\
& + \frac{1}{(z-w)} \left[\delta^{\hat{A}\hat{B}} \left(\frac{2}{K} \partial_w W_{F,4}^\lambda + \frac{32 q^2 (1-\lambda)(1+\lambda)}{15 K} \partial_w^3 W_{F,2}^\lambda \right) \right. \\
& + d^{\hat{A}\hat{B}\hat{C}} \left(\partial_w W_{F,4}^{\lambda,\hat{C}} + \frac{16}{15} q^2 (1-\lambda)(1+\lambda) \partial_w^3 W_{F,2}^{\lambda,\hat{C}} \right) \\
& - i f^{\hat{A}\hat{B}\hat{C}} \left(\frac{1}{4q} W_{F,5}^{\lambda,\hat{C}} + \frac{8}{21} q (2-\lambda)(2+\lambda) \partial_w^2 W_{F,3}^{\lambda,\hat{C}} \right. \\
& \left. + \frac{16}{45} q^3 (1-\lambda)(1+\lambda)(1-2\lambda)(1+2\lambda) \partial_w^4 W_{F,1}^{\lambda,\hat{C}} \right] (w) + \dots . \quad (\text{D.2})
\end{aligned}$$

At $\lambda = 0$ with $N = 1$, the six OPEs (3.5) and (D.2) can be seen from the previous work [8]. By redefining the fields having the adjoint index with the extra factor q and ignoring the central terms, we observe that there are the second and first order poles in the first half of (D.2) (for $h_1 = 1$, there is only the second order pole) while there exists the first order pole in the last half of (D.2) under the $q \rightarrow 0$ limit with $\lambda = 0$ [26].

We present some structure constants we are using above for each h_1, h_2, h_3 and k , from (2.17), as follows:

$$\begin{aligned}
S_{F,R}^{1,1,1,0}(\lambda) &= 1, & S_{F,R}^{1,2,1,0}(\lambda) &= \frac{1}{2}(2\lambda-1), & S_{F,R}^{1,2,1,1}(\lambda) &= 0, \\
S_{F,R}^{1,2,2,0}(\lambda) &= -\frac{1}{6}, & S_{F,R}^{1,3,1,0}(\lambda) &= \frac{1}{3}(\lambda-1)(2\lambda-1), & S_{F,R}^{1,3,1,1}(\lambda) &= \frac{1}{6}(\lambda-1)(2\lambda-1), \\
S_{F,R}^{1,3,1,2}(\lambda) &= 0, & S_{F,R}^{1,3,2,0}(\lambda) &= -\frac{1}{6}(\lambda-1), & S_{F,R}^{1,3,2,1}(\lambda) &= 0, \\
S_{F,R}^{1,3,3,0}(\lambda) &= \frac{1}{120}, & S_{F,R}^{2,1,1,0}(\lambda) &= 0, & S_{F,R}^{2,1,1,1}(\lambda) &= \frac{1}{2}(2\lambda+1), \\
S_{F,R}^{2,1,2,0}(\lambda) &= -\frac{1}{6}, & S_{F,R}^{2,2,1,0}(\lambda) &= -\frac{1}{12}(2\lambda-1)(2\lambda+1), & S_{F,R}^{2,2,1,1}(\lambda) &= \frac{1}{12}(2\lambda-1)(2\lambda+1), \\
S_{F,R}^{2,2,1,2}(\lambda) &= -\frac{1}{12}(2\lambda-1)(2\lambda+1), & S_{F,R}^{2,2,2,0}(\lambda) &= \frac{1}{12}, & S_{F,R}^{2,2,2,1}(\lambda) &= -\frac{1}{12}, \\
S_{F,R}^{2,2,3,0}(\lambda) &= \frac{1}{120}, & S_{F,R}^{2,3,1,0}(\lambda) &= -\frac{1}{12}(\lambda-1)(2\lambda-1)(2\lambda+1), & S_{F,R}^{2,3,1,1}(\lambda) &= 0, \\
S_{F,R}^{2,3,1,2}(\lambda) &= 0, & S_{F,R}^{2,3,1,3}(\lambda) &= 0, & S_{F,R}^{2,3,2,0}(\lambda) &= \frac{1}{15}(\lambda-1)(\lambda+1), \\
S_{F,R}^{2,3,2,1}(\lambda) &= -\frac{1}{30}(\lambda-1)(\lambda+1), & S_{F,R}^{2,3,2,2}(\lambda) &= \frac{1}{90}(\lambda-1)(\lambda+1), & S_{F,R}^{2,3,3,0}(\lambda) &= -\frac{1}{120},
\end{aligned}$$

$$\begin{aligned}
S_{F,R}^{2,3,3,1}(\lambda) &= \frac{1}{240}, & S_{F,R}^{2,3,4,0}(\lambda) &= -\frac{1}{5040}, & S_{F,R}^{3,1,1,0}(\lambda) &= 0, \\
S_{F,R}^{3,1,1,1}(\lambda) &= \frac{1}{6}(\lambda+1)(2\lambda+1), & S_{F,R}^{3,1,1,2}(\lambda) &= \frac{1}{3}(\lambda+1)(2\lambda+1), & S_{F,R}^{3,1,2,0}(\lambda) &= 0, \\
S_{F,R}^{3,1,2,1}(\lambda) &= -\frac{1}{6}(\lambda+1), & S_{F,R}^{3,1,3,0}(\lambda) &= \frac{1}{120}, & S_{F,R}^{3,2,1,0}(\lambda) &= 0, \\
S_{F,R}^{3,2,1,1}(\lambda) &= 0, & S_{F,R}^{3,2,1,2}(\lambda) &= 0, & S_{F,R}^{3,2,1,3}(\lambda) &= -\frac{1}{12}(\lambda+1)(2\lambda-1)(2\lambda+1), \\
S_{F,R}^{3,2,2,0}(\lambda) &= \frac{1}{90}(\lambda-1)(\lambda+1), & S_{F,R}^{3,2,2,1}(\lambda) &= -\frac{1}{30}(\lambda-1)(\lambda+1), \\
S_{F,R}^{3,2,2,2}(\lambda) &= \frac{1}{15}(\lambda-1)(\lambda+1), & S_{F,R}^{3,2,3,0}(\lambda) &= -\frac{1}{240}, & S_{F,R}^{3,2,3,1}(\lambda) &= \frac{1}{120}, \\
S_{F,R}^{3,2,4,0}(\lambda) &= -\frac{1}{5040}, & S_{F,R}^{3,3,1,0}(\lambda) &= \frac{1}{180}(\lambda-1)(\lambda+1)(2\lambda-1)(2\lambda+1), \\
S_{F,R}^{3,3,1,1}(\lambda) &= -\frac{1}{180}(\lambda-1)(\lambda+1)(2\lambda-1)(2\lambda+1), \\
S_{F,R}^{3,3,1,2}(\lambda) &= \frac{1}{180}(\lambda-1)(\lambda+1)(2\lambda-1)(2\lambda+1), \\
S_{F,R}^{3,3,1,3}(\lambda) &= -\frac{1}{180}(\lambda-1)(\lambda+1)(2\lambda-1)(2\lambda+1), \\
S_{F,R}^{3,3,1,4}(\lambda) &= \frac{1}{180}(\lambda-1)(\lambda+1)(2\lambda-1)(2\lambda+1), \\
S_{F,R}^{3,3,2,0}(\lambda) &= -\frac{1}{90}(\lambda-1)(\lambda+1), & S_{F,R}^{3,3,2,1}(\lambda) &= \frac{1}{60}(\lambda-1)(\lambda+1), \\
S_{F,R}^{3,3,2,2}(\lambda) &= -\frac{1}{60}(\lambda-1)(\lambda+1), & S_{F,R}^{3,3,2,3}(\lambda) &= \frac{1}{90}(\lambda-1)(\lambda+1), \\
S_{F,R}^{3,3,3,0}(\lambda) &= -\frac{1}{1260}(\lambda-2)(\lambda+2), & S_{F,R}^{3,3,3,1}(\lambda) &= \frac{1}{840}(\lambda-2)(\lambda+2), \\
S_{F,R}^{3,3,3,2}(\lambda) &= -\frac{1}{1260}(\lambda-2)(\lambda+2), & S_{F,R}^{3,3,4,0}(\lambda) &= \frac{1}{5040}, & S_{F,R}^{3,3,4,1}(\lambda) &= -\frac{1}{5040}, \\
S_{F,R}^{3,3,5,0}(\lambda) &= \frac{1}{362880}, & S_{F,L}^{h_2,h_1,h_3,k}(\lambda) &= S_{F,R}^{h_1,h_2,h_3,k}(\lambda),
\end{aligned} \tag{D.3}$$

where the corresponding $S_{F,L}^{h_2,h_1,h_3,k}(\lambda)$ can be obtained by using the last relation of (D.3). Note that the λ dependences above do not have a λ factor and therefore, they do not vanish at $\lambda = 0$. Some of the structure constants do not depend on the λ . For other structure constants, the similar analysis can be done. It is rather nontrivial to observe the previous relations (A.8) because these structure constants at $\lambda = 0$ have their nontrivial numerical values. We have checked that all the λ dependent structure constants for nonzero λ have the factor λ explicitly after subtracted by those evaluated at $\lambda = 0$.

By introducing a new variable

$$h_3 = h_1 + h_2 - h - 2, \tag{D.4}$$

we can sum over the variable k for the first commutator in (D.1) as follows:

$$\begin{aligned} P_F^{h_1, h_2, h}(m, n, \lambda) &\equiv \sum_{k=0}^{h+1} (-1)^{h_1+h_2-h-2} 4^h (2(h_1 + h_2 - h) - 5)! \\ &\times \left(S_{F,R}^{h_1, h_2, h_1+h_2-h-2, k}(\lambda) [m+h_1-1]_{h-k+1} [n+h_2-1]_k \right. \\ &\quad \left. - S_{F,L}^{h_1, h_2, h_1+h_2-h-2, k}(\lambda) [m+h_1-1]_k [n+h_2-1]_{h-k+1} \right). \end{aligned} \quad (\text{D.5})$$

Then we obtain the following structure constants by using (D.3), (D.4) and (D.5),

$$\begin{aligned} P_F^{1,1,-1}(m, n, \lambda) &= 0, & P_F^{1,2,-1}(m, n, \lambda) &= 0, & P_F^{1,2,0}(m, n, \lambda) &= m, \\ P_F^{1,3,-1}(m, n, \lambda) &= 0, & P_F^{1,3,0}(m, n, \lambda) &= 2m, & P_F^{1,3,1}(m, n, \lambda) &= 4m(2m+n)\lambda, \\ P_F^{2,1,-1}(m, n, \lambda) &= 0, & P_F^{2,1,0}(m, n, \lambda) &= -n, & P_F^{2,2,-1}(m, n, \lambda) &= 0, \\ P_F^{2,2,0}(m, n, \lambda) &= (m-n), & P_F^{2,2,1}(m, n, \lambda) &= 0, & P_F^{2,3,-1}(m, n, \lambda) &= 0, \\ P_F^{2,3,0}(m, n, \lambda) &= (2m-n), & P_F^{2,3,1}(m, n, \lambda) &= 0, \\ P_F^{2,3,2}(m, n, \lambda) &= -\frac{8}{3}(m-1)m(m+1)(2\lambda-1)(2\lambda+1), & P_F^{3,1,-1}(m, n, \lambda) &= 0, \\ P_F^{3,1,0}(m, n, \lambda) &= -2n, & P_F^{3,1,1}(m, n, \lambda) &= -4n(m+2n)\lambda, \\ P_F^{3,2,-1}(m, n, \lambda) &= 0, & P_F^{3,2,0}(m, n, \lambda) &= (m-2n), & P_F^{3,2,1}(m, n, \lambda) &= 0, \\ P_F^{3,2,2}(m, n, \lambda) &= \frac{8}{3}(n-1)n(n+1)(2\lambda-1)(2\lambda+1), & P_F^{3,3,-1}(m, n, \lambda) &= 0, \\ P_F^{3,3,0}(m, n, \lambda) &= 2(m-n), & P_F^{3,3,1}(m, n, \lambda) &= 0, \\ P_F^{3,3,2}(m, n, \lambda) &= -\frac{16}{15}(m-n)(2m^2-mn+2n^2-8)(\lambda-1)(\lambda+1), \\ P_F^{3,3,3}(m, n, \lambda) &= 0. \end{aligned} \quad (\text{D.6})$$

On the other hand, we write down the previous structure constants appearing in [12] as follows:

$$\begin{aligned} p_F^{1,2,0}(m, n, \lambda) &= m, & p_F^{1,3,0}(m, n, \lambda) &= 2m, & p_F^{2,1,0}(m, n, \lambda) &= -n, \\ p_F^{2,2,0}(m, n, \lambda) &= (m-n), & p_F^{2,3,0}(m, n, \lambda) &= (2m-n), \\ p_F^{2,3,2}(m, n, \lambda) &= -\frac{8}{3}(m-1)m(m+1)(2\lambda-1)(2\lambda+1), \\ p_F^{3,1,0}(m, n, \lambda) &= -2n, & p_F^{3,2,0}(m, n, \lambda) &= (m-2n), \\ p_F^{3,2,2}(m, n, \lambda) &= \frac{8}{3}(n-1)n(n+1)(2\lambda-1)(2\lambda+1), \\ p_F^{3,3,0}(m, n, \lambda) &= 2(m-n), \\ p_F^{3,3,2}(m, n, \lambda) &= -\frac{16}{15}(m-n)(2m^2-mn+2n^2-8)(\lambda-1)(\lambda+1). \end{aligned} \quad (\text{D.7})$$

Comparing (D.6) with (D.7), the structure constants $P_F^{1,3,1}(m, n, \lambda)$ and $P_F^{3,1,1}(m, n, \lambda)$ do not appear in (D.7). This indicates that the structure constants in [44] are not appropriate for the present work.

For convenience, we present the following OPEs corresponding to (3.5)

$$\begin{aligned}
W_{B,1}^\lambda(z) W_{B,1}^\lambda(w) &= -\frac{1}{(z-w)^2} \frac{N K}{16 q^2} + \dots, \\
W_{B,1}^\lambda(z) W_{B,2}^\lambda(w) &= -\frac{1}{(z-w)^3} \frac{N K (2\lambda-1)}{4 q} + \frac{1}{(z-w)^2} W_{B,1}^\lambda(w) + \dots, \\
W_{B,1}^\lambda(z) W_{B,3}^\lambda(w) &= -\frac{1}{(z-w)^4} \frac{N K (2\lambda-1)^2}{2} + \frac{1}{(z-w)^3} 4q (2\lambda-1) W_{B,1}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^2} \left[2 W_{B,2}^\lambda - 2q (2\lambda-1) \partial_w W_{B,1}^\lambda \right] (w) + \dots, \\
W_{B,2}^\lambda(z) W_{B,2}^\lambda(w) &= \frac{1}{(z-w)^4} N K (1 - 6\lambda + 6\lambda^2) + \frac{1}{(z-w)^2} 2 W_{B,2}^\lambda(w) + \frac{1}{(z-w)} \partial_w W_{B,2}^\lambda(w) \\
&\quad + \dots, \\
W_{B,2}^\lambda(z) W_{B,3}^\lambda(w) &= \frac{1}{(z-w)^5} 16q N K (\lambda-1)\lambda(2\lambda-1) + \frac{1}{(z-w)^4} 64q^2 \lambda(\lambda-1) W_{B,1}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^2} 3 W_{B,3}^\lambda(w) + \frac{1}{(z-w)} \partial_w W_{B,3}^\lambda(w) + \dots, \\
W_{B,2}^\lambda(z) W_{B,4}^\lambda(w) &= \frac{1}{(z-w)^6} 32q^2 N K \lambda(\lambda-1)(2\lambda-1)^2 \\
&\quad - \frac{1}{(z-w)^5} 256q^3 \lambda(\lambda-1)(2\lambda-1) W_{B,1}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^4} \left[-\frac{192q^2}{5} (3\lambda^2 - 3\lambda - 1) W_{B,2}^\lambda \right. \\
&\quad \left. + 128q^3 \lambda(\lambda-1)(2\lambda-1) \partial_w W_{B,1}^\lambda \right] (w) \\
&\quad + \frac{1}{(z-w)^2} 4 W_{B,4}^\lambda(w) + \frac{1}{(z-w)} \partial_w W_{B,4}^\lambda(w) + \dots, \\
W_{B,3}^\lambda(z) W_{B,3}^\lambda(w) &= -\frac{1}{(z-w)^6} \frac{16q^2 N K}{3} (-3 + 10\lambda + 30\lambda^2 - 80\lambda^3 + 40\lambda^4) \\
&\quad - \frac{1}{(z-w)^4} 16q^2 (2\lambda-3)(2\lambda+1) W_{B,2}^\lambda(w) \\
&\quad - \frac{1}{(z-w)^3} 8q^2 (2\lambda-3)(2\lambda+1) \partial_w W_{B,2}^\lambda(w) \\
&\quad + \frac{1}{(z-w)^2} \left[4 W_{B,4}^\lambda - \frac{12q^2}{5} (2\lambda-3)(2\lambda+1) \partial_w^2 W_{B,2}^\lambda \right] (w)
\end{aligned}$$

$$+ \frac{1}{(z-w)} \left[2 \partial_w W_{B,4}^\lambda - \frac{8q^2}{15} (2\lambda - 3)(2\lambda + 1) \partial_w^3 W_{B,2}^\lambda \right] (w) + \dots \quad (\text{D.8})$$

Note that the structure constants appearing in (D.8) can be reproduced by changing $\lambda \rightarrow \lambda - \frac{1}{2}$ in (3.5) with opposite signs for the central terms.

References

- [1] I. Bakas, “The Large n Limit of Extended Conformal Symmetries,” Phys. Lett. B **228**, 57 (1989) doi:10.1016/0370-2693(89)90525-X
- [2] C. N. Pope, L. J. Romans and X. Shen, “The Complete Structure of W(Infinity),” Phys. Lett. B **236**, 173-178 (1990) doi:10.1016/0370-2693(90)90822-N
- [3] C. N. Pope, L. J. Romans and X. Shen, “W(infinity) and the Racah-wigner Algebra,” Nucl. Phys. B **339**, 191-221 (1990) doi:10.1016/0550-3213(90)90539-P
- [4] C. N. Pope, “Lectures on W algebras and W gravity,” [arXiv:hep-th/9112076 [hep-th]].
- [5] C. N. Pope, L. J. Romans and X. Shen, “A New Higher Spin Algebra and the Lone Star Product,” Phys. Lett. B **242**, 401-406 (1990) doi:10.1016/0370-2693(90)91782-7
- [6] E. Bergshoeff, C. N. Pope, L. J. Romans, E. Sezgin and X. Shen, “The Super W(infinity) Algebra,” Phys. Lett. B **245**, 447-452 (1990) doi:10.1016/0370-2693(90)90672-S
- [7] I. Bakas and E. Kiritsis, “Grassmannian Coset Models and Unitary Representations of W(infinity),” Mod. Phys. Lett. A **5**, 2039-2050 (1990) doi:10.1142/S0217732390002328
- [8] S. Odake and T. Sano, “W(1) + infinity and superW(infinity) algebras with SU(N) symmetry,” Phys. Lett. B **258**, 369-374 (1991) doi:10.1016/0370-2693(91)91101-Z
- [9] S. Odake, “Unitary representations of W(infinity) algebras,” Int. J. Mod. Phys. A **7**, 6339-6356 (1992) doi:10.1142/S0217751X9200288X [arXiv:hep-th/9111058 [hep-th]].
- [10] E. Bergshoeff, M. A. Vasiliev and B. de Wit, “The SuperW(infinity) (lambda) algebra,” Phys. Lett. B **256**, 199-205 (1991) doi:10.1016/0370-2693(91)90673-E
- [11] E. Bergshoeff, B. de Wit and M. A. Vasiliev, “The Structure of the superW(infinity) (lambda) algebra,” Nucl. Phys. B **366**, 315-346 (1991) doi:10.1016/0550-3213(91)90005-I

- [12] C. Ahn, “The $\mathcal{N} = 2$ supersymmetric $w_{1+\infty}$ symmetry in the two-dimensional SYK models,” JHEP **05**, 115 (2022) doi:10.1007/JHEP05(2022)115 [arXiv:2203.03105 [hep-th]].
- [13] J. Murugan, D. Stanford and E. Witten, “More on Supersymmetric and 2d Analogs of the SYK Model,” JHEP **08**, 146 (2017) doi:10.1007/JHEP08(2017)146 [arXiv:1706.05362 [hep-th]].
- [14] K. Bulycheva, “ $\mathcal{N} = 2$ SYK model in the superspace formalism,” JHEP **04**, 036 (2018) doi:10.1007/JHEP04(2018)036 [arXiv:1801.09006 [hep-th]].
- [15] C. Ahn and C. Peng, “Chiral Algebras of Two-Dimensional SYK Models,” JHEP **07**, 092 (2019) doi:10.1007/JHEP07(2019)092 [arXiv:1812.05106 [hep-th]].
- [16] T. Creutzig, Y. Hikida and P. B. Ronne, “Extended higher spin holography and Grassmannian models,” JHEP **11**, 038 (2013) doi:10.1007/JHEP11(2013)038 [arXiv:1306.0466 [hep-th]].
- [17] C. Ahn, “A deformed supersymmetric $w_{1+\infty}$ symmetry in the celestial conformal field theory,” Eur. Phys. J. C **82**, no.7, 630 (2022) doi:10.1140/epjc/s10052-022-10582-9 [arXiv:2202.02949 [hep-th]].
- [18] C. Ahn, “The structure of the $\mathcal{N} = 4$ supersymmetric linear $W_\infty[\lambda]$ algebra,” Eur. Phys. J. C **83**, no.7, 615 (2023) doi:10.1140/epjc/s10052-023-11752-z [arXiv:2208.07000 [hep-th]].
- [19] C. Ahn, “N=4 supersymmetric linear $W_\infty[\lambda]$ algebra,” Phys. Rev. D **106**, no.2, 026008 (2022) doi:10.1103/PhysRevD.106.026008 [arXiv:2205.04024 [hep-th]].
- [20] L. Eberhardt, M. R. Gaberdiel and I. Rienacker, “Higher spin algebras and large $\mathcal{N} = 4$ holography,” JHEP **03**, 097 (2018) doi:10.1007/JHEP03(2018)097 [arXiv:1801.00806 [hep-th]].
- [21] K. Thielemans, “A Mathematica package for computing operator product expansions,” Int. J. Mod. Phys. C **2**, 787-798 (1991) doi:10.1142/S0129183191001001
- [22] Wolfram Research, Inc., Mathematica, Version 13.0.0, Champaign, IL (2021).
- [23] K. Schoutens, “O(n) Extended Superconformal Field Theory in Superspace,” Nucl. Phys. B **295**, 634-652 (1988) doi:10.1016/0550-3213(88)90539-1

- [24] M. Beccaria, C. Candu and M. R. Gaberdiel, “The large $N = 4$ superconformal W_∞ algebra,” JHEP **06**, 117 (2014) doi:10.1007/JHEP06(2014)117 [arXiv:1404.1694 [hep-th]].
- [25] C. Ahn and M. H. Kim, “The operator product expansion between the 16 lowest higher spin currents in the $\mathcal{N} = 4$ superspace,” Eur. Phys. J. C **76**, no.7, 389 (2016) doi:10.1140/epjc/s10052-016-4234-2 [arXiv:1509.01908 [hep-th]].
- [26] C. Ahn, “Towards a supersymmetric $w_{1+\infty}$ symmetry in the celestial conformal field theory,” Phys. Rev. D **105**, no.8, 086028 (2022) doi:10.1103/PhysRevD.105.086028 [arXiv:2111.04268 [hep-th]].
- [27] A. Strominger, “ $w(1+\infty)$ and the Celestial Sphere,” [arXiv:2105.14346 [hep-th]].
- [28] A. Strominger, “ $w_{1+\infty}$ Algebra and the Celestial Sphere: Infinite Towers of Soft Graviton, Photon, and Gluon Symmetries,” Phys. Rev. Lett. **127**, no.22, 221601 (2021) doi:10.1103/PhysRevLett.127.221601
- [29] S. Pasterski, M. Pate and A. M. Raclariu, “Celestial Holography,” [arXiv:2111.11392 [hep-th]].
- [30] A. Fotopoulos, S. Stieberger, T. R. Taylor and B. Zhu, “Extended Super BMS Algebra of Celestial CFT,” JHEP **09**, 198 (2020) doi:10.1007/JHEP09(2020)198 [arXiv:2007.03785 [hep-th]].
- [31] A. Guevara, E. Himwich, M. Pate and A. Strominger, “Holographic symmetry algebras for gauge theory and gravity,” JHEP **11**, 152 (2021) doi:10.1007/JHEP11(2021)152 [arXiv:2103.03961 [hep-th]].
- [32] E. Himwich, M. Pate and K. Singh, “Celestial operator product expansions and $w_{1+\infty}$ symmetry for all spins,” JHEP **01**, 080 (2022) doi:10.1007/JHEP01(2022)080 [arXiv:2108.07763 [hep-th]].
- [33] H. Jiang, “Holographic chiral algebra: supersymmetry, infinite Ward identities, and EFTs,” JHEP **01**, 113 (2022) doi:10.1007/JHEP01(2022)113 [arXiv:2108.08799 [hep-th]].
- [34] J. Mago, L. Ren, A. Y. Srikant and A. Volovich, “Deformed $w_{1+\infty}$ Algebras in the Celestial CFT,” SIGMA **19**, 044 (2023) doi:10.3842/SIGMA.2023.044 [arXiv:2111.11356 [hep-th]].

- [35] W. Bu, S. Heuveline and D. Skinner, “Moyal deformations, $W_{1+\infty}$ and celestial holography,” JHEP **12**, 011 (2022) doi:10.1007/JHEP12(2022)011 [arXiv:2208.13750 [hep-th]].
- [36] W. Melton, S. A. Narayanan and A. Strominger, “Deforming soft algebras for gauge theory,” JHEP **03**, 233 (2023) doi:10.1007/JHEP03(2023)233 [arXiv:2212.08643 [hep-th]].
- [37] P. Drozdov and T. Kimura, “Structure of deformed $w_{1+\infty}$ symmetry and topological generalization in Celestial CFT,” [arXiv:2306.11693 [math-ph]].
- [38] K. Costello, N. M. Paquette and A. Sharma, “Burns space and holography,” JHEP **10**, 174 (2023) doi:10.1007/JHEP10(2023)174 [arXiv:2306.00940 [hep-th]].
- [39] C. Beem, M. Lemos, P. Liendo, W. Peelaers, L. Rastelli and B. C. van Rees, “Infinite Chiral Symmetry in Four Dimensions,” Commun. Math. Phys. **336**, no.3, 1359-1433 (2015) doi:10.1007/s00220-014-2272-x [arXiv:1312.5344 [hep-th]].
- [40] R. Bittleston, S. Heuveline and D. Skinner, “The celestial chiral algebra of self-dual gravity on Eguchi-Hanson space,” JHEP **09**, 008 (2023) doi:10.1007/JHEP09(2023)008 [arXiv:2305.09451 [hep-th]].
- [41] M. Ademollo, L. Brink, A. D’Adda, R. D’Auria, E. Napolitano, S. Sciuto, E. Del Giudice, P. Di Vecchia, S. Ferrara and F. Gliozzi, *et al.* “Supersymmetric Strings and Color Confinement,” Phys. Lett. B **62**, 105-110 (1976) doi:10.1016/0370-2693(76)90061-7
- [42] M. Ademollo, L. Brink, A. D’Adda, R. D’Auria, E. Napolitano, S. Sciuto, E. Del Giudice, P. Di Vecchia, S. Ferrara and F. Gliozzi, *et al.* “Dual String with U(1) Color Symmetry,” Nucl. Phys. B **111**, 77-110 (1976) doi:10.1016/0550-3213(76)90483-1
- [43] C. Ahn, M. R. Gaberdiel and M. H. Kim, “The small $\mathcal{N} = 4$ superconformal \mathcal{W}_∞ algebra,” J. Phys. A **53**, no.39, 395401 (2020) doi:10.1088/1751-8121/aba5ba [arXiv:2004.07439 [hep-th]].
- [44] C. Ahn and M. H. Kim, “The $\mathcal{N} = 4$ higher spin algebra for generic μ parameter,” JHEP **02**, 123 (2021) doi:10.1007/JHEP02(2021)123 [arXiv:2009.04852 [hep-th]].