Magnetic quadrupole moment of 175 Lu and parity-violating polarization degree of levels in 175 LuOH⁺

Igor Kurchavov,^{1,*} Daniel Maison,^{1,†} Leonid Skripnikov,^{1,2,‡} Matt Grau,^{3,§} and Alexander Petrov^{1,2,¶}

¹Petersburg Nuclear Physics Institute named by B.P. Konstantinov of National

Research Center "Kurchatov Institute" (NRC "Kurchatov Institute" - PNPI),

1 Orlova roscha mcr., Gatchina, 188300 Leningrad region, Russia

²Saint Petersburg State University, 7/9 Universitetskaya nab., St. Petersburg, 199034 Russia

³Department of Physics, Old Dominion University, Norfolk, VA 23529

(Dated: Received: date / Revised version: date)

The calculation of the parity-violating polarizations in the external electric field, which are associated with the electron electric dipole moment (eEDM) and magnetic quadrupole moment (MQM) of the ¹⁷⁵Lu nucleus, as well as the determination of the rovibrational structure for the ¹⁷⁵LuOH⁺ cation, is performed. Beyond the bending of the molecule, the slight effect of the stretching of the distance between Lu and OH is taken into account. This study is required for the preparation of the experiment and for the extraction of the eEDM and MQM values of ¹⁷⁵Lu from future measurements.

INTRODUCTION

The pursuit of understanding the fundamental symmetries governing the laws of physics has been a central focus in the field of theoretical and experimental physics. Among these symmetries, the invariance with respect to charge conjugation (\mathcal{C}), spatial parity (\mathcal{P}), and time reversion (\mathcal{T}) has long been considered crucial. However, in the latter half of the 20th century, experimental evidence emerged, confirming the violation of both \mathcal{P} and the combined \mathcal{CP} symmetries in weak interactions. Nowadays, the violation of \mathcal{CP} symmetry holds immense interest in the fields of cosmology and astrophysics, as it is considered one of the three fundamental conditions for baryogenesis [1]. As a result, the exploration of novel manifestations of symmetry violation has emerged as a prominent research area in contemporary theoretical and experimental physics [2].

One of the approaches used to explore these phenomena is by studying the electron electric dipole moments (eEDM) and nuclear magnetic quadrupole moments (MQM), which serve as highly sensitive probes for testing the boundaries of the Standard Model of electroweak interactions and its extensions [3–8]. The search for both *e*EDMand MQM has been gaining momentum. Recently the JILA group has obtained a new constraint on the eEDM, $|d_e| < 4.1 \times 10^{-30} \text{ e} \cdot \text{cm}$ (90% confidence) [9], using the ¹⁸⁰Hf¹⁹F⁺ ions trapped in a rotating electric field. It further improves the latest ACME collaboration result obtained in 2018, $|d_e| \lesssim 1.1 \cdot 10^{-29} \ e \cdot cm$ [10], by a factor of 2.4 and the first result $|d_{\rm e}| \lesssim 1.3 \times 10^{-28}$ on the ${}^{180}\text{Hf}{}^{19}\text{F}{}^+$ ions [11] by a factor of about 32. Planned experiments aiming to measure MQM with the use of molecules offer a pathway to investigate \mathcal{P}, \mathcal{T} odd nuclear forces, quark chromo-EDMs, and other CPviolating quark interactions [12].

In this paper, we focus on the 175 Lu MQM in the LuOH⁺ molecular ion, specifically in the ground rota-

tional level of the first excited bending vibrational mode. The choice of LuOH⁺ is motivated by its electronic structure similarities to YbOH, while demonstrating even higher sensitivity to nuclear \mathcal{CP} -violation effects due to the significant electric quadrupole moment of ¹⁷⁵Lu [13].

In a polar molecule with a heavy element, the \mathcal{T}, \mathcal{P} -violating energy shifts induced by *e*EDM and MQM are

$$\Delta E_{\mathcal{P},\mathcal{T}} = P_e E_{\text{eff}} d_e + P_M W_M M,\tag{1}$$

where d_e is the value of the eEDM, M is the value of the MQM, factors $E_{\rm eff}$ and W_M are determined by the electronic structure of the molecule, P_e and P_M are the corresponding \mathcal{P}, \mathcal{T} -odd polarization coefficients. To extract M and d_e from the measured energy shift $\Delta E_{\mathcal{P},\mathcal{T}}$, one needs to know E_{eff} , W_M , P_e and P_M values. The value of W_M was calculated in Ref. [13], E_{eff} was calculated in Ref. [14]. An important task is to distinguish between the two sources of symmetry violation. This can be achieved by using different molecules or different electronic states of the same molecule. However, as can be seen from Eq. (1), it is also possible to utilize two or more sublevels of the same electronic state with different P_e/P_M ratios. Therefore, an accurate method for estimating these ratios is necessary. In Ref. [14], we calculated the value of P_e for ¹⁷⁵LuOH⁺. In Ref. [15], a method for calculating P_M was developed and applied to the ¹⁷³YbOH molecule. The main objective of this study is to apply this method to calculate P_M for ¹⁷⁵LuOH⁺. Furthermore, more precise data for P_e , hyperfine structure, and spectroscopic constants are obtained.

METHODS

Following Ref. [16], we present the Hamiltonian in the molecular reference frame as follows:

$$\ddot{\mathbf{H}} = \ddot{\mathbf{H}}_{\text{mol}} + \ddot{\mathbf{H}}_{\text{hfs}} + \ddot{\mathbf{H}}_{\text{ext}}.$$
 (2)

Two approaches for the molecular Hamiltonian $\hat{\mathbf{H}}_{mol}$ are used:

$$\hat{\mathbf{H}}_{\rm mol}^{\rm I} = \frac{(\hat{\mathbf{J}} - \hat{\mathbf{J}}^{e-v})^2}{2\mu R_e^2} + \frac{(\hat{\mathbf{J}}^v)^2}{2\mu_{OH}r^2} + V(R_e, \theta) \qquad (3)$$

and

$$\hat{\mathbf{H}}_{\rm mol}^{\rm II} = -\frac{\hbar^2}{2\mu} \frac{\partial^2}{\partial R^2} + \frac{(\hat{\mathbf{J}} - \hat{\mathbf{J}}^{e-v})^2}{2\mu R^2} + \frac{(\hat{\mathbf{J}}^v)^2}{2\mu_{OH}r^2} + V(R,\theta),\tag{4}$$

where μ is the reduced mass of the Lu-OH system, μ_{OH} is the reduced mass of the OH, $\hat{\mathbf{J}}$ is the total electronic, vibrational, and rotational angular momentum, $\hat{\mathbf{J}}^{e-v} =$ $\mathbf{J}^e + \mathbf{J}^v$ is the electronic-vibrational momentum, \mathbf{J}^e , is the electronic momentum, $\hat{\mathbf{J}}^{v}$ is the vibrational momentum, R is the distance between Lu and the center mass of OH, $R_e = 1.930$ Å is the corresponding equilibrium value for R, r = 0.954 Å is OH bond length, θ is the angle between OH and the axis (z axis of the molecular frame) directed from Lu to the OH center of mass and $V(R,\theta)$ is the potential energy surface obtained in the electronic structure calculations. The condition $\theta = 0$ corresponds to the linear configuration where the O atom is between Lu and H ones. R, r and θ are the so called Jacobi coordinates, see Fig. 1.



FIG. 1: Jacobi coordinates for the LuOH⁺ cation.

Using the Hamiltonian $\hat{H}^{\rm I}_{\rm mol},$ we neglect the influence of the stretching ν_1 (associated with R) and OH ligand ν_3 (associated with r) vibrational modes. However, we still consider the bending modes ν_2 (associated with θ) with fixed R and r. In this approach, the spectroscopic constants and P_e coefficient in Ref. [14] were calculated. Using the Hamiltonian \hat{H}_{mol}^{II} , we additionally take into account the influence of the stretching mode. In this paper, we recalculated the spectroscopic constants (and obtained a new one for the stretching mode) and the P_e value using \hat{H}_{mol}^{II} . We also calculated the P_M polarization in both approaches.

The Hamiltonian for the hyperfine interaction of elec-

trons with Lu and H nuclei reads

$$\hat{\mathbf{H}}_{\rm hfs} = -g_{\rm H} \mathbf{I}^{\rm H} \cdot \sum_{a} \left(\frac{\boldsymbol{\alpha}_{2a} \times \boldsymbol{r}_{2a}}{r_{2a}^3} \right) + \\ -g_{\rm Lu} \mu_N \mathbf{I}^{\rm Lu} \cdot \sum_{a} \left(\frac{\boldsymbol{\alpha}_a \times \boldsymbol{r}_{1a}}{r_{1a}^3} \right) \quad (5)$$
$$\cdot e^2 \sum_{q} (-1)^q \hat{Q}_q^2 (\mathbf{I}^{\rm Lu}) \sum_{a} \sqrt{\frac{2\pi}{5}} \frac{Y_{2q}(\theta_{1a}, \phi_{1a})}{r_{1a}^3}$$

where g_{Lu} and g_{H} are the g-factors of the lutetium and hydrogen nuclei, α_a are the Dirac matrices for the *a*-th electron, r_{1a} and r_{2a} are their radius-vectors in the coordinate system centered on the Lu and H nuclei, $\hat{Q}_q^2(\mathbf{I}^{\mathrm{Lu}})$ is the quadrupole moment operator for ¹⁷⁵Lu nucleus, $I^{\text{Lu}} = 7/2$, $I^{\text{H}} = 1/2$ are nuclear spins for ¹⁷⁵Lu and ¹H. The Stark Hamiltonian

$$\dot{\mathbf{H}}_{\text{ext}} = -\mathbf{D} \cdot \mathbf{E} \tag{6}$$

describes the interaction of the molecule with the external electric field, and **D** is the dipole moment operator.

Wavefunctions, rovibrational energies, and hyperfine structure were obtained by numerically diagonalizing the Hamiltonian (2) over the basis set of electronicrotational-vibrational-nuclear spins wavefunctions.

$$\Psi_{\Omega m\omega} P_{lm}(\theta) \Theta^J_{M_J,\omega}(\alpha,\beta) U^{\rm H}_{M_I^{\rm H}} U^{\rm Lu}_{M_I^{\rm Lu}}.$$
 (7)

or

$$\Psi_{\Omega m\omega}\chi_{\nu_1}(R)P_{lm}(\theta)\Theta^J_{M_J,\omega}(\alpha,\beta)U^{\rm H}_{M_I^{\rm H}}U^{\rm Lu}_{M_I^{\rm Lu}} \qquad (8)$$

for \hat{H}^{I}_{mol} or \hat{H}^{II}_{mol} respectively. Here $\Theta^{J}_{M_{J},\omega}(\alpha,\beta) =$ $\sqrt{(2J+1)/4\pi}D^J_{M_J,\omega}(\alpha,\beta,\gamma=0)$ is the rotational wavefunction, $D^J_{M_I,\omega}$ is Wigner function, α,β correspond to azimuthal and polar angles of the molecular z axis (directed from Lu to the center of mass of OH group), $U_{M_{\rm r}^{\rm H}}^{\rm H}$ and $U_{M_{ru}^{Lu}}^{Lu}$ are the hydrogen and lute tium nuclear spin wavefunctions, M_J is the projection of the molecular (electronic-rotational-vibrational) angular momentum $\hat{\mathbf{J}}$ on the lab axis, ω is the projection of the same momentum on z axis of the molecular frame, $M_I^{\rm H}$ and $M_I^{\rm Lu}$ are the projections of the nuclear angular momenta of hydrogen and lutetium on the lab axis, $P_{lm}(\theta)$ is the associated Legendre polynomial, l is the vibration angular momentum and m is its projection on the molecular axis, Ω is the projection of the total electronic angular momentum on the molecular axis z for linear configuration. $\Psi_{\Omega m\omega}$ is the electronic wavefunction (see Ref. [16] for details).

In this calculation functions with $\omega - m = \Omega = \pm 1/2$, l = 0 - 30 and $m = 0, \pm 1, \pm 2, J = 1/2, 3/2, 5/2$ were included in the basis sets (7) and (8). The ground vibrational state $\nu_2 = 0$ corresponds to m = 0, the first excited bending mode $\nu_2 = 1$ to $m = \pm 1$, the second excited bending mode has states with $m = 0, \pm 2$ etc. A common designation ν_2^m for the bending vibrational levels will be used below. No momenta is associated with ν_1 quantum number.

Provided that the *electronic-vibrational* matrix elements are known, the matrix elements of $\hat{\mathbf{H}}$ between states in the basis set (8) can be calculated with help of the angular momentum algebra [16, 17] mostly in the same way as for the diatomic molecules [18]. The corresponding matrix elements are taken from Ref. [14]. To calculate the \mathcal{T}, \mathcal{P} -odd shifts the average value of corresponding Hamiltonians [19]

$$H_{\rm MQM} = -\frac{M}{2I^{\rm Lu}(2I^{\rm Lu}-1)}T_{ik}\frac{3}{2r^5}\epsilon_{jli}\alpha_j r_l r_k, \quad (9)$$

where ϵ_{jli} is the unit antisymmetric tensor, α is the vector of Dirac matrices,

$$T_{ik} = I_i^{\text{Lu}} I_k^{\text{Lu}} + I_k^{\text{Lu}} I_i^{\text{Lu}} - \frac{2}{3} \delta_{i,k} I^{\text{Lu}} (I^{\text{Lu}} + 1), \qquad (10)$$

$$H_{\rm EDM} = 2d_e \begin{pmatrix} 0 & 0\\ 0 & \boldsymbol{\sigma} \boldsymbol{E} \end{pmatrix} , \qquad (11)$$

E is the inner molecular electric field, and σ are the Pauli matrices. [15, 16] were evaluated.

For the calculation of the potential energy surface, we employed two models to account for electron correlation effects: coupled cluster with single and double excitation amplitudes (CCSD) and coupled cluster with single, double, and perturbative triple excitation amplitudes (CCSD(T)). These calculations were performed within the Dirac-Coulomb Hamiltonian framework. For Lu, we utilized the uncontracted Dyall's AE3Z basis set [20], while for F, we employed the aug-cc-PVTZ-DK basis set [21, 22]. In these calculations, the correlation treatment excluded the 1s..3d electrons, and a virtual orbital energy cutoff of 70 Hartree was set.

We obtained the CCSD and CCSD(T) potential surface $V(R,\theta)$ on the grid of coordinates (R_i,θ_k) ,

$$\{R_i\} = 1.771, 1.877, 1.930, 1.983, 2.089 \mathring{A}$$
(12)

$$\{\theta_k\} = 0^{\circ}, 5^{\circ}, 10^{\circ}, 15^{\circ}, 20^{\circ}, 25^{\circ}, 55^{\circ}, 90^{\circ}, 122^{\circ}, 155^{\circ}, 180^{\circ}$$
(13)

The potentials were approximated by polynomial $\sum_{n=0}^{7} \sum_{m=0}^{4} C_{nm} \theta^n R^m$. Since the linear configuration has minimum energy, the coefficients $C_{nm} = 0$ for n = 1.

RESULTS AND DISCUSSIONS

In Fig. 2 and Table I, we present the calculated potential energy surface and spectroscopic properties. It can be see that the results for the CCSD and CCSD(T)

models are in close agreement with each other. Incorporating non-iterative triple cluster amplitudes leads to a decrease in vibrational energies by $4-8 \text{ cm}^{-1}$. The use of Hamiltonian \hat{H}_{mol}^{II} instead of \hat{H}_{mol}^{I} has a slightly greater impact on the vibrational energies. The final l-doubling value for $\nu_2 = 1$ of 24.5 MHz is approximately 1 MHz higher than the value obtained with a frozen R variable.

In Fig. 3 the calculated polarizations P_e , P_M for the selected levels of the lowest N = 1 rotational state of the first excited $\nu_2 = 1$ bending vibrational mode with frozen R approximation as a functions of the external electric field are given. The selected 14 levels, numbered 43–56 (see Tables II and III), are those which were chosen in Ref. [14] for the eEDM search. The corresponding energies (order of 31 850 MHz) are given in Fig. 3 panel (a) of Ref. [14]. Using only levels numbered 43–56 is not enough, however, if nonzero MQM of ¹⁷⁵Lu nucleus is assumed. The reason is that the ratio $P_e/P_M \sim -10.5$ is about the same for all these levels that makes it impossible to distinguish eEDM and MQM contributions. Similarly to P_{e} [14] there are levels with close values of P_M . These states differ by only projection $M_I^{\rm H} = \pm 1/2$ which almost does not influence P_e and P_M .

The numerical data for P_e , P_M and hyperfine energies for all N = 1 levels for E = 50 and E = 100 V/cm are given in Tables II and III respectively. To assess the influence of the stretching mode calculations with Hamiltonians \hat{H}^{I}_{mol} and \hat{H}^{II}_{mol} were performed. One can see that accounting for stretching mode leads to a decreasing of notable P_e, P_M values up to about 5% for E = 50 and 4% for E = 100. This is explained by increasing l-doubling value describing the energy difference between levels of opposite parity at zero electric field when stretching mode is taken into account.

Since MQM induced energy shift is proportional to P_M

TABLE I: Calculated vibrational energy levels (cm^{-1}) rotational constants B (cm⁻¹), and l-doubling (MHz) for the excitation modes of stretching $\nu_1 = 0 - 1$ and bending $\nu_2 = 0 - 2$ quanta of ¹⁷⁵LuOH⁺. Ligand mode ν_3 quanta is zero in calculations.

Parameter	CCSD(T) (R-frozen)	CCSD	CCSD(T)
$ \nu_1 = 0, \nu_2 = 0 $	0	0	0
$\nu_1 = 0, \nu_2 = 1$	442	438	434
$\nu_1 = 1, \nu_2 = 0$		750	745
$\nu_1 = 0, \nu_2 = 2^0$	871	864	856
$\nu_1 = 0, \nu_2 = 2^2$	898	887	879
$B(\nu_1 = 0, \nu_2 = 0)$	0.2879	0.2874	0.2868
$B(\nu_1 = 0, \nu_2 = 1)$	0.2881	0.2869	0.2863
$B(\nu_1 = 1, \nu_2 = 0)$		0.2862	0.2855
$B(\nu_1 = 0, \nu_2 = 2^0)$	0.2883	0.2870	0.2864
$B(\nu_1 = 0, \nu_2 = 2^2)$	0.2882	0.2919	0.2912
<i>l</i> -doubling $(\nu_1 = 0, \nu_2 = 1)$	23.5	24.4	24.5
<i>l</i> -doubling $(\nu_1 = 0, \nu_2 = 2^2)$	0.005	0.012	0.012



FIG. 2: Potential surface $V(\theta, R)$ for CCSD(T) calculations with polynomial interpolation.

for the MQM searches the levels with large ${\cal P}_M$ values are preferred. Besides, to distinguish *e*EDM and MQM contributions, the levels with different P_e/P_M ratios have to be used. As an example of the proposed *e*EDM contribution exclusion scheme, let us consider the first and fifty third levels for E = 50 V/cm. For the first level we have $\delta E^1 = -0.3740 E_{\text{eff}} d_e + 0.0794 W_M M$; for the fifty third level $\delta E^{53} = -0.3524 E_{\text{eff}} d_e + 0.0333 W_M M$. Then the combination $\delta E^1 - 1.0613 \delta E^{53} = 0.044 W_M M$ is independent of eEDM and can be used for MQM extraction. Similarly, for the electric field E = 100 V/cm, for example, one can choose levels numbered 1 (with $\delta E^1 =$ $-0.4715 E_{\text{eff}} d_e + 0.1001 W_M M$ and a ratio $P_e / P_M \approx 4.71$) and 43 (with $\delta E^{43} = 0.5294 E_{\text{eff}} d_e - 0.0502 W_M M$ and the ratio equal to 10.55). Then the obtained combination is $1.1229\delta E^1 + \delta E^{43} = 0.0622 W_M M$. We note also that our choice of the levels is only an example. On the base of Tables II, III and using formula

$$MW_M = \frac{1}{\Delta} \left(P_e^x \delta E^y - P_e^y \delta E^x \right),$$

$$\Delta = P_e^x P_M^y - P_e^y P_M^x$$
(14)

one can choose alternative appropriate levels for the MQM search. Here x and y are numbers of chosen levels. Similarly the *e*EDM contribution can be determined.

Electric field E = 100 V/cm provides almost saturated values for P_e and P_M . As it was shown in Letter [23] that in the l - doubling structure, the P value tends to reach half of the maximum value for molecules with Hund's case b. Calculations showed that all levels have polarizations $P_e < 0.58$ and $P_M < 0.12$.



FIG. 3: Calculated polarizations $P_e/7$ (solid) and P_M (dashed) for the selected levels (see text for details) for the different values of M_F of the lowest N = 1rotational level of the first excited the $\nu_2 = 1$ bending vibrational mode of ¹⁷⁵LuOH+ as functions of the external electric field.

To access these energy levels in order to perform the *e*EDM and MQM precision measurement, trapped LuOH⁺ ions that are initially prepared in the ground rovibrational state by either optical pumping or quantum logic spectroscopy. From this point either of the states in the pair necessary to distinguish the *e*EDM from MQM can be populated by driving a Raman transition using a pair far-detuned infrared lasers with a difference frequency tuned to the energy of the desired state. As the pairs states with opposite projection of $M_I^{\rm H} = \pm 1/2$ are separated by several MHz they will be straightforward to resolve experimentally.

CONCLUSION

We calculated spectroscopic constants for the lowest vibrational levels, parity-violating polarizations P_e and P_M associated with *e*EDM and MQM of ¹⁷⁵Lu energy shifts in the first excited bending mode for the ¹⁷⁵LuOH⁺ cation. The levels suitable for the MQM search are determined.

Electronic structure calculations have been carried out using computing resources of the federal collective usage center Complex for Simulation and Data Processing for Mega-science Facilities at National Research Centre "Kurchatov Institute", http://ckp.nrcki.ru/.

Electronic correlation calculation of the poten-

tial energy surface was supported by the Russian Science Foundation (Grant No. 19-72-10019-P (https://rscf.ru/en/project/22-72-41010/). Dirac-Hartree-Fock calculations were supported by Foundation for the Advancement of Theoretical Physics and Mathematics "BASIS" grant according to the Research Project No. 21-1-2-47-1.

- * kurchavov_ip@pnpi.nrcki.ru
- † daniel.majson@gmail.com, maison_de@pnpi.nrcki.ru
- ‡ skripnikov_lv@pnpi.nrcki.ru,leonidos239@gmail.com
- § mgrau@odu.edu
- ¶ petrov_an@pnpi.nrcki.ru
- [1] A. D. Sakharov, JETP lett. 5, 24 (1967).
- [2] M. S. Safronova, D. Budker, D. DeMille, D. F. J. Kimball, A. Derevianko, and C. W. Clark, Rev. Mod. Phys. 90, 025008 (2018).
- [3] R. Alarcon, J. Alexander, V. Anastassopoulos, T. Aoki, R. Baartman, S. Baeßler, L. Bartoszek, D. H. Beck, F. Bedeschi, and R. B. et al., (2022), arXiv:2203.08103 [hep-ph].
- [4] T. FUKUYAMA, International Journal of Modern Physics A 27, 1230015 (2012).
- [5] M. Pospelov and A. Ritz, Phys. Rev. D 89, 056006 (2014), arXiv:1311.5537 [hep-ph].
- [6] Y. Yamaguchi and N. Yamanaka, Phys. Rev. Lett. 125, 241802 (2020), arXiv:2003.08195 [hep-ph].
- [7] Y. Yamaguchi and N. Yamanaka, Phys. Rev. D 103, 013001 (2021), arXiv:2006.00281 [hep-ph].
- [8] V. Flambaum, D. DeMille, and M. Kozlov, Physical review letters 113, 103003 (2014).
- [9] T. S. Roussy, L. Caldwell, T. Wright, W. B. Cairncross, Y. Shagam, K. B. Ng, N. Schlossberger, S. Y. Park, A. Wang, J. Ye, and E. A. Cornell, Science **381**, 46 (2023), https://www.science.org/doi/pdf/10.1126/science.adg4084.
- [10] V. Andreev, D. Ang, D. DeMille, J. Doyle, G. Gabrielse, J. Haefner, N. Hutzler, Z. Lasner, C. Meisenhelder, B. O'Leary, et al., Nature 562, 355 (2018).
- [11] W. B. Cairncross, D. N. Gresh, M. Grau, K. C. Cossel, T. S. Roussy, Y. Ni, Y. Zhou, J. Ye, and E. A. Cornell, Phys. Rev. Lett. **119**, 153001 (2017).
- [12] L. V. Skripnikov, A. N. Petrov, A. V. Titov, and V. V. Flambaum, Phys. Rev. Lett. **113**, 263006 (2014).
- [13] D. E. Maison, L. V. Skripnikov, V. V. Flambaum, and M. Grau, J. Comp. Phys. **153**, 224302 (2020).
- [14] D. E. Maison, L. V. Skripnikov, G. Penyazkov, M. Grau, and A. N. Petrov, Phys. Rev. A 106, 062827 (2022).
- [15] I. Kurchavov and A. Petrov, Phys. Rev. A 106, 062806 (2022).
- [16] A. Petrov and A. Zakharova, Phys. Rev. A 105, L050801 (2022).
- [17] L. D. Landau and E. M. Lifshitz, *Quantum mechanics*, 3rd ed. (Pergamon, Oxford, 1977).
- [18] A. N. Petrov, Phys. Rev. A 83, 024502 (2011).
- [19] M. G. Kozlov, V. I. Fomichev, Yu. Yu. Dmitriev, L. N. Labzovsky, and A. V. Titov, **20**, 4939 (1987).
- [20] A. S. P. Gomes, K. G. Dyall, and L. Visscher, Theor. Chim. Acta 127, 369 (2010).

- [21] T. H. Dunning, Jr, J. Comp. Phys. 90, 1007 (1989).
- [22] R. A. Kendall, T. H. Dunning, Jr, and R. J. Harrison, J. Comp. Phys. 96, 6796 (1992).
- [23] A. Petrov and A. Zakharova, Phys. Rev. A 105, L050801 (2022).

TABLE II: The calculated energies (in MHz), polarizations for the different projections of the total angular momentum M_F of the lowest N = 1 rotational level of the first excited v = 1 bending vibrational mode of ¹⁷⁵LuOH⁺ for the value of the external electric field E = 50 V/cm. Levels are numbered by the increasing energy. Zero energy level corresponds to the lowest energy of N = 1 states at zero electric field. Calculation with frozen R variable are marked by (f).

#	M_{F}	En(f)	En	$P_{e}(\mathbf{f})$	P_{e}	$P_M(f)$	P_M	#	M_{F}	En(f)	En	$P_{e}(\mathbf{f})$	P_{e}	$P_M(f)$	P_M
1	2.5	-5.5	-5.3	-0.3812	-0.3740	0.0809	0.0794	49	0.5	31851.8	31849.4	0.0014	0.0013	-0.0001	-0.0001
$\frac{1}{2}$	1.5	-5.3	-5.1	-0.3811	-0.3738	0.0809	0.0794	50	0.5	31873.7	31872.1	-0.0013	-0.0012	0.0001	0.0001
3	1.5	-1.7	-1.7	-0.2450	-0.2374	0.0520	0.0504	51	0.5	31874.6	31872.9	-0.2151	-0.2070	0.0204	0.0195
4	0.5	-1.6	-1.6	-0.2436	-0.2359	0.0517	0.0501	52	1.5	31874.6	31872.9	-0.2164	-0.2082	0.0206	0.0197
5	0.5	-0.2	-0.2	-0.0013	-0.0013	0.0003	0.0003	53	1.5	31876.9	31875.2	-0.3631	-0.3524	0.0345	0.0333
6	0.5	23.3	24.3	0.0011	0.0011	-0.0002	-0.0002	54	2.5	31877.0	31875.2	-0.3631	-0.3524	0.0345	0.0333
7	1.5	24.9	25.8	0.2448	0.2372	-0.0520	-0.0503	55	2.5	31880.1	31878.3	-0.4457	-0.4355	0.0423	0.0411
8	0.5	25.0	25.9	0.2436	0.2360	-0.0517	-0.0501	56	3.5	31880.3	31878.4	-0.4457	-0.4355	0.0423	0.0411
9	2.5	28.7	29.5	0.3808	0.3736	-0.0808	-0.0793	57	4.5	32043.2	32042.7	0.3530	0.3435	-0.0477	-0.0475
10	1.5	28.9	29.7	0.3807	0.3734	-0.0808	-0.0793	58	5.5	32043.3	32042.9	0.3528	0.3433	-0.0477	-0.0475
11	4.5	118.7	119.3	-0.3374	-0.3333	0.0703	0.0694	59	3.5	32045.9	32045.4	0.3266	0.3165	-0.0442	-0.0437
12	3.5	118.9	119.5	-0.3374	-0.3333	0.0703	0.0694	60	4.5	32046.1	32045.5	0.3265	0.3164	-0.0442	-0.0437
13	3.5	122.2	122.8	-0.3040	-0.2987	0.0633	0.0622	61	2.5	32048.4	32047.8	0.2853	0.2750	-0.0386	-0.0380
14	2.5	122.4	122.9	-0.3039	-0.2986	0.0633	0.0622	62	3.5	32048.5	32048.0	0.2853	0.2750	-0.0386	-0.0380
15	2.5	125.3	125.8	-0.2452	-0.2391	0.0511	0.0498	63	1.5	32050.5	32049.9	0.2203	0.2108	-0.0298	-0.0291
16	1.5	125.4	125.9	-0.2449	-0.2387	0.0510	0.0497	64	2.5	32050.6	32050.0	0.2205	0.2111	-0.0298	-0.0292
17	1.5	127.5	127.9	-0.1446	-0.1396	0.0301	0.0291	65	0.5	32052.0	32051.3	0.1202	0.1141	-0.0162	-0.0158
18	0.5	127.5	127.9	-0.1419	-0.1368	0.0296	0.0285	66	1.5	32052.0	32051.4	0.1241	0.1179	-0.0168	-0.0163
19	0.5	128.4	128.7	-0.0024	-0.0025	0.0005	0.0005	67	0.5	32052.5	32051.8	0.0036	0.0036	-0.0005	-0.0005
20	0.5	151.7	153.1	0.0020	0.0021	-0.0004	-0.0004	68	0.5	32075.7	32075.9	-0.0035	-0.0035	0.0005	0.0005
21	1.5	152.6	153.9	0.1444	0.1394	-0.0301	-0.0290	69	0.5	32076.3	32076.5	-0.1200	-0.1139	0.0162	0.0158
22	0.5	152.6	154.0	0.1422	0.1371	-0.0296	-0.0286	70	1.5	32076.3	32076.5	-0.1238	-0.1177	0.0168	0.0163
23	2.5	154.8	156.1	0.2450	0.2388	-0.0510	-0.0497	71	1.5	32077.8	32077.9	-0.2198	-0.2103	0.0297	0.0291
24	1.5	154.9	156.2	0.2447	0.2385	-0.0510	-0.0497	72	2.5	32077.8	32078.0	-0.2201	-0.2106	0.0298	0.0291
25	3.5	158.0	159.2	0.3036	0.2984	-0.0632	-0.0621	73	2.5	32080.0	32080.1	-0.2847	-0.2743	0.0385	0.0379
26	2.5	158.1	159.3	0.3036	0.2983	-0.0632	-0.0621	74	3.5	32080.1	32080.2	-0.2847	-0.2743	0.0385	0.0379
27	4.5	161.7	162.8	0.3369	0.3329	-0.0702	-0.0693	75	3.5	32082.8	32082.8	-0.3257	-0.3156	0.0441	0.0437
28	3.5	161.9	163.0	0.3369	0.3328	-0.0702	-0.0693	76	4.5	32082.9	32083.0	-0.3256	-0.3155	0.0441	0.0437
29	2.5	688.8	689.0	0.0456	0.0440	-0.0091	-0.0088	77	4.5	32085.9	32085.9	-0.3520	-0.3424	0.0477	0.0474
30	3.5	688.8	689.1	0.0457	0.0440	-0.0091	-0.0088	78	5.5	32086.0	32086.1	-0.3518	-0.3422	0.0477	0.0474
31	1.5	689.4	689.6	0.0312	0.0300	-0.0062	-0.0060	79	4.5	32319.7	32315.5	0.0083	0.0120	0.0108	0.0112
32	2.5	689.4	689.7	0.0314	0.0302	-0.0063	-0.0061	80	3.5	32319.9	32315.6	0.0082	0.0119	0.0107	0.0111
33	0.5	689.7	690.0	0.0145	0.0138	-0.0029	-0.0028	81	3.5	32320.3	32316.1	0.0063	0.0092	0.0083	0.0086
34	1.5	689.8	690.0	0.0162	0.0156	-0.0032	-0.0031	82	2.5	32320.4	32316.2	0.0062	0.0089	0.0081	0.0084
35	0.5	689.9	690.1	0.0016	0.0015	-0.0003	-0.0003	83	2.5	32320.8	32316.5	0.0044	0.0064	0.0058	0.0060
36	0.5	713.4	714.6	-0.0063	-0.0063	0.0013	0.0013	84	1.5	32320.8	32316.5	0.0039	0.0057	0.0052	0.0054
37	1.5	713.5	714.7	-0.0195	-0.0192	0.0039	0.0038	85	0.5	32321.0	32316.7	0.0014	0.0020	0.0018	0.0019
38	0.5	713.5	714.7	-0.0099	-0.0093	0.0020	0.0019	86	1.5	32321.0	32316.8	0.0025	0.0036	0.0032	0.0034
39	1.5	713.6	714.8	-0.0285	-0.0270	0.0057	0.0054	87	0.5	32321.2	32316.9	0.0007	0.0011	0.0010	0.0010
40	2.5	713.6	714.8	-0.0331	-0.0322	0.0066	0.0064	88	4.5	32342.1	32338.3	-0.0069	-0.0105	-0.0107	-0.0111
41	2.5	713.7	714.9	-0.0450	-0.0431	0.0090	0.0086	89	3.5	32342.1	32338.3	-0.0056	-0.0085	-0.0086	-0.0089
42	3.5	713.8	715.0	-0.0463	-0.0446	0.0092	0.0089	90	2.5	32342.1	32338.3	-0.0041	-0.0062	-0.0062	-0.0065
43	2.5	31845.7	31843.5	0.4474	0.4374	-0.0426	-0.0414	91	1.5	32342.1	32338.3	-0.0025	-0.0038	-0.0038	-0.0039
44	3.5	31845.8	31843.6	0.4473	0.4373	-0.0426	-0.0414	92	0.5	32342.1	32338.3	-0.0008	-0.0013	-0.0013	-0.0013
45	1.5	31848.7	31846.4	0.3642	0.3536	-0.0347	-0.0335	93	3.5	32342.3	32338.5	-0.0066	-0.0101	-0.0103	-0.0107
46	2.5	31848.8	31846.5	0.3642	0.3536	-0.0347	-0.0335	94	2.5	32342.3	32338.5	-0.0049	-0.0074	-0.0075	-0.0078
47	0.5	31850.9	31848.6	0.2156	0.2075	-0.0205	-0.0196	95	1.5	32342.3	32338.5	-0.0030	-0.0045	-0.0046	-0.0047
48	1.5	31851.0	31848.7	0.2170	0.2089	-0.0206	-0.0198	96	0.5	32342.3	32338.5	-0.0010	-0.0015	-0.0015	-0.0016

TABLE III: The calculated energies (in MHz), polarizations for the different projections of the total angular momentum M_F of the lowest N = 1 rotational level of the first excited v = 1 bending vibrational mode of ¹⁷⁵LuOH⁺ for the value of the external electric field E = 100 V/cm. Levels are numbered by the increasing energy. Zero energy level corresponds to the lowest energy of N = 1 states at zero electric field. Calculation with frozen R variable are marked by (f).

	11	E (f)	Ð	D (C)	л		ת		11	\mathbf{E} (f)	F	D (C)	л	D (f)	
#	MF	En(f)	En	$\frac{P_e(I)}{P_e(I)}$	Pe	$P_M(I)$	P_M	#	MF	En(I)	En	$P_e(I)$	P_e	$P_M(f)$	P_M
1	2.5	-16.2	-15.9	-0.4750	-0.4715	0.1009	0.1001	49	0.5	31850.8	31848.4	-0.0001	-0.0001	0.0000	0.0000
2	1.5	-16.0	-15.8	-0.4750	-0.4714	0.1009	0.1001	50	0.5	31872.6	31870.9	0.0002	0.0002	0.0000	0.0000
3	1.5	-6.2	-6.0	-0.3811	-0.3739	0.0809	0.0794	51	0.5	31875.8	31874.0	-0.3626	-0.3519	0.0345	0.0333
4	0.5	-6.1	-5.9	-0.3809	-0.3736	0.0809	0.0793	52	1.5	31875.8	31874.1	-0.3625	-0.3517	0.0345	0.0332
5	0.5	-0.9	-0.9	-0.0004	-0.0004	0.0001	0.0001	53	1.5	31882.8	31880.9	-0.4904	-0.4813	0.0466	0.0454
6	0.5	22.5	23.5	0.0004	0.0004	-0.0001	-0.0001	54	2.5	31882.8	31881.0	-0.4903	-0.4812	0.0466	0.0454
7	1.5	27.9	28.8	0.3807	0.3734	-0.0808	-0.0792	55	2.5	31891.1	31889.2	-0.5331	-0.5256	0.0506	0.0496
8	0.5	28.0	28.8	0.3804	0.3731	-0.0807	-0.0792	56	3.5	31891.2	31889.3	-0.5330	-0.5255	0.0506	0.0495
9	2.5	38.6	39.3	0.4741	0.4706	-0.1006	-0.0999	57	4.5	32026.8	32026.5	0.4008	0.3931	-0.0542	-0.0543
10	1.5	38.7	39.4	0.4741	0.4706	-0.1006	-0.0998	58	5.5	32027.0	32026.6	0.4006	0.3929	-0.0542	-0.0543
11	4.5	102.3	103.0	-0.3830	-0.3816	0.0798	0.0795	59	3.5	32033.0	32032.5	0.3896	0.3814	-0.0526	-0.0526
12	3.5	102.5	103.2	-0.3830	-0.3815	0.0798	0.0795	60	4.5	32033.1	32032.7	0.3894	0.3812	-0.0526	-0.0526
13	3.5	110.4	111.0	-0.3691	-0.3668	0.0769	0.0764	61	2.5	32039.0	32038.5	0.3692	0.3603	-0.0498	-0.0496
14	2.5	110.5	111.2	-0.3691	-0.3667	0.0769	0.0764	62	3.5	32039.1	32038.6	0.3691	0.3601	-0.0498	-0.0496
15	2.5	118.1	118.7	-0.3370	-0.3329	0.0702	0.0693	63	1.5	32044.6	32044.1	0.3258	0.3159	-0.0439	-0.0435
16	1.5	118.2	118.8	-0.3369	-0.3329	0.0702	0.0693	64	2.5	32044.7	32044.2	0.3257	0.3157	-0.0439	-0.0435
17	1.5	124.6	125.1	-0.2450	-0.2388	0.0510	0.0497	65	0.5	32049.2	32048.6	0.2201	0.2107	-0.0297	-0.0290
18	0.5	124.7	125.2	-0.2446	-0.2384	0.0509	0.0497	66	1.5	32049.2	32048.6	0.2203	0.2109	-0.0297	-0.0290
19	0.5	127.7	128.1	-0.0004	-0.0004	0.0001	0.0001	67	0.5	32051.2	32050.5	0.0002	0.0002	0.0000	0.0000
20	0.5	151.0	152.4	0.0003	0.0004	-0.0001	-0.0001	68	0.5	32074.1	32074.3	-0.0002	-0.0002	0.0000	0.0000
21	1.5	154.1	155.4	0.2448	0.2386	-0.0510	-0.0497	69	0.5	32076.2	32076.4	-0.2195	-0.2101	0.0296	0.0289
22	0.5	154.2	155.4	0.2444	0.2382	-0.0509	-0.0496	70	1.5	32076.3	32076.4	-0.2197	-0.2102	0.0296	0.0290
23	2.5	161.0	162.1	0.3365	0.3324	-0.0701	-0.0692	71	1.5	32081.2	32081.3	-0.3246	-0.3146	0.0438	0.0434
24	1.5	161.1	162.2	0.3364	0.3324	-0.0701	-0.0692	72	2.5	32081.3	32081.3	-0.3245	-0.3145	0.0438	0.0433
25	3.5	169.2	170.3	0.3684	0.3660	-0.0767	-0.0762	73	2.5	32087.6	32087.6	-0.3676	-0.3585	0.0496	0.0495
26	2.5	169.4	170.4	0.3683	0.3660	-0.0767	-0.0762	74	3.5	32087.7	32087.7	-0.3674	-0.3584	0.0496	0.0495
27	4.5	178.0	179.1	0.3821	0.3806	-0.0796	-0.0793	75	3.5	32094.7	32094.6	-0.3876	-0.3793	0.0524	0.0524
28	3.5	178.2	179.2	0.3821	0.3806	-0.0796	-0.0793	76	4.5	32094.8	32094.7	-0.3874	-0.3792	0.0524	0.0524
29	2.5	687.2	687.5	0.0794	0.0772	-0.0159	-0.0155	77	4.5	32102.1	32102.1	-0.3988	-0.3909	0.0540	0.0542
30	3.5	687.3	687.6	0.0794	0.0772	-0.0159	-0.0155	78	5.5	32102.3	32102.2	-0.3985	-0.3907	0.0541	0.0542
31	1.5	689.4	689.7	0.0585	0.0565	-0.0117	-0.0113	79	4.5	32318.7	32314.5	0.0150	0.0217	0.0190	0.0198
32	2.5	689.5	689.8	0.0585	0.0566	-0.0117	-0.0113	80	3.5	32318.9	32314.6	0.0150	0.0216	0.0190	0.0198
33	0.5	690.9	691.1	0.0311	0.0299	-0.0062	-0.0060	81	3.5	32320.9	32316.6	0.0119	0.0172	0.0154	0.0160
34	1.5	690.9	691.1	0.0315	0.0303	-0.0063	-0.0061	82	2.5	32321.0	32316.7	0.0118	0.0171	0.0153	0.0159
35	0.5	691.4	691.6	0.0003	0.0003	-0.0001	-0.0001	83	2.5	32322.4	32318.2	0.0083	0.0121	0.0109	0.0113
36	0.5	714.8	716.0	-0.0032	-0.0035	0.0006	0.0007	84	1.5	32322.5	32318.3	0.0082	0.0120	0.0108	0.0112
37	0.5	715.0	716.2	-0.0285	-0.0270	0.0057	0.0054	85	1.5	32323.4	32319.2	0.0043	0.0063	0.0057	0.0059
38	1.5	715.0	716.2	-0.0322	-0.0311	0.0064	0.0062	86	0.5	32323.4	32319.2	0.0040	0.0057	0.0052	0.0054
39	1.5	715.3	716.5	-0.0588	-0.0568	0.0001	0.0002	87	0.5	32323.8	32319.5	0.0003	0.0005	0.0002	0.0001
40	2.5	715.4	716.5	-0.0593	-0.0574	0.0118	0.0110	88	4.5	32344 1	32340.3	-0.0126	-0.0190	-0.0187	-0.0195
10	$2.0 \\ 2.5$	715.9	717.0	-0.0804	-0.0783	0.0110	0.0111	89	3.5	32344 3	32340.4	-0.0120	-0.0190	-0.0107	-0.0184
42	$\frac{2.0}{3.5}$	715.9	717.1	-0.0805	-0.0784	0.0161	0.0156	90	3.5	32344 4	32340.6	-0.0120	-0.0100	-0.0161	-0.0164
12	25	31833 7	31831.6	0.5366	0.5204	-0.0512	-0.0502	01	25	32344 4	32340.6	-0.00110	-0.0100	-0.0101	-0.0132
4Δ	$\frac{2.0}{3.5}$	31833.8	31831.8	0.5365	0.5294	-0.0512	-0.0502	02	$\frac{2.0}{1.5}$	32344 4	323/0.6	-0.0054	-0.0133	-0.0127	-0.00102
45	15	318/11 9	31830 0	0.0000	0.0290	-0.0470	-0.0450	03	0.5	32344.5	323/0.6	-0.0018	-0.0001	-0.0077	-0.0027
-10 /16	2.5	318/11 2	31830 1	0.4920	0.4009	-0.0470	-0.0459	93	25	32344.0	32340.0	_0 00010	-0.0125	-0.0120	-0.0135
40 17	05	318/17 7	318/15 /	0.4921	0.4000	-0.03/7	-0.0409	05	15	32344.0	32340.7	-0.0061	-0.0100	-0.0101	_0 0080
-±1 18	15	31847 7	31845.4	0.3030	0.0002	0.0347	-0.0332	90 06	1.0 0 5	32344.0	32340.0 333/0 9	0.0001	0.0009	0.0000	0.0009
40	1.0	31041.1	51645.5	0.5057	0.5551	-0.0347	-0.0399	90	0.0	JZJ44. (52540.8	-0.0021	-0.0031	-0.0050	-0.0031