

Enumeration of saturated and unsaturated substituted N-heterocycles

Stefan Schuster* and Tatjana Malycheva

Dept. of Bioinformatics, Matthias Schleiden Institute, Friedrich Schiller University of Jena, Ernst-Abbe-Pl. 2, 07743 Jena, Germany

*Corresponding author, e-mail: stefan.schu@uni-jena.de

Abstract

Mathematical and computational approaches in chemistry and biochemistry fill a gap in respect to the analysis of the physicochemical features of compounds and their functionality and provide an overview of known as well as yet unknown, but hypothetically possible structures. Nitrogen containing heterocycles such as aziridine, azetidine and pyrrolidine bear a high potential in pharmacology, biotechnology and synthetic biology. Here, we present a mathematical enumeration procedure for all possible azaheterocycles with at least one substituent depending on the number of atoms in the ring, in the sense of saturated and unsaturated congeners. One subgroup belonging to that substance class is constituted by ring-shaped amino acids with a secondary amino group such as proline. A recursion formula is derived, which results in a modified Lucas number series. Moreover, an explicit formula for determining the number of such substances based on the Golden Ratio is given and a second one, based on binomial coefficients, is newly derived. This enumeration is a helpful tool for construction or complementation of virtual compound databases and for computer-assisted chemical synthesis route planning.

Introduction

In view of many applications, it is instrumental to comprehend the combinatorial multitude of chemical substances of a given class. For example, in metabolomics, in the study of prebiotic or biotic evolution and in synthetic organic chemistry, it is important to know the maximum number of substances that can be expected. To enumerate theoretically possible structures, mathematical methods are helpful tools (Henze et al. 1931, Balaban et al. 1988, Haranczyk et al. 2010, Böcker et al. 2014, Fichtner et al., 2017). Enumeration is instrumental in constructing or complementing virtual compound databases and for computer-assisted chemical synthesis route planning. Combinatorial analysis, similar to enumeration studies on alcohols (Henze et al. 1931) and hydrocarbons (Balaban et al. 1988) was carried out on branched and unbranched aliphatic amino acids without any rings (Grützmann et al. 2011, Fichtner et al. 2017). There is a plethora of non-proteinogenic amino acids, occurring, for example, in toxic peptides (Arp et al. 2008, Kroteń et al. 2010, Zahradníčková et al 2022). For instance, jessenipeptin involves dehydro-2-aminobutyric acid, D-allo-threonine, L-

diaminobutyric acid and D-proline (Arp et al. 2008). The latter belongs to the substance class under study here. Many other examples are reviewed in Fichtner et al. (2017). It can be expected that many more substances are synthesized in living organisms than have been discovered so far.

The aim of the present paper is to derive a method for calculating the number of nitrogen-containing heterocycles containing at least one substituent connected to the ring by a single bond. We allow for varying numbers of double bonds in the ring. The enumeration is performed in dependence on ring size, while the substituent remains unchanged. We focus on molecules in which the substituent is connected at a position adjacent to the nitrogen (2-position or α -position). One subgroup belonging to that substance class is constituted by ring-shaped α -amino acids with a secondary amino group such as azetidine-2-carboxylic acid, proline, pipercolic acid, and baikiain (Zahradníčková et al 2022). However, the same enumeration procedure can be applied to azaheterocycles in which the substituent is located at another position as long as this does not lead to an additional plane of symmetry.

Since the number of hydrogen atoms is not the same in all structures of a given size if the number of double bonds varies, our approach is about counting congeners rather than isomers. Methods for enumerating congeners are scarce so far (Haranczyk et al., 2010; Schuster et al., 2017).

Throughout this paper, we exclude adjacent (allenic) double bonds, for two reasons. First, the bond angle would then amount to about 180° . That would be very difficult to realise in (closed) rings unless another two adjacent double bonds or a triple bond are situated on the opposite side of the ring. Second, allenic double bonds and triple bonds only occur very rarely in biomolecules (some occurrences are reviewed in Fichtner et al. 2017).

Several of the substances under consideration are rather unstable and, thus, short-lived. The question arises what the minimum lifetime of a molecule should be in order to be considered as a separate substance. Here, we use a very general approach in that we count separately, a priori, all conceivable structures of the type defined above with one exception: resonance structures (also known as Kekulé structures) (Lukovits et al. 2003, Hosoya 2005) of rings with fully conjugated double bonds. That is, resonance structures in a given ring are considered equivalent because the π electrons are delocalised. However, for the sake of clarity in the mathematical presentation, we will first consider them separately and only later modify the calculation so as to combine them.

Mathematical enumeration

Many substances involving one or more (non-allenic) double bonds can exist as cis- and trans-isomers depending on whether the single bonds are situated on the same or on different sides, respectively. Here we do not consider such isomers to be different. In fact, in three- to seven-membered rings, only the cis-isomer can occur for steric reasons (Neuenschwander et al. 2011). Moreover, we do not distinguish stereoisomers (such as L-proline and D-proline).

The smallest meaningful number for a ring is $n = 3$. Then, there are four possible ring structures, notably one involving single bonds only and three with one double bond each (Fig. 1). An example is provided by 2H-azirine-2-carboxylic acid, which is part of the

antibacterial agent azirinomycin produced by *Streptomyces aureus* (Stapley et al. 1971). As azirinomycin is also toxic to humans, it cannot be used in human medicine.




Figure 1. Three-membered N-heterocycles with one substituent R (-COOH; -CHO; -NH₂ etc.). Names refer to the unsubstituted rings. The left and right 2H-azirines are substituted at the 2 and 3 positions, respectively.

For $n = 4$, there are six possible ring structures, notably one involving single bonds only, four with one double bond each and one with two non-adjacent double bonds each (Fig. 2). For these two double bonds, two locations are feasible: in the representation shown in Fig. 2, they could be located in the vertical or horizontal direction. These resonance structures are equivalent and, thus, are finally counted only once. An example of a substance with a four-membered ring is azetidone-2-carboxylic acid, which is produced as a defence chemical against herbivory in many plants such as the lily of the valley (Fowden 1963).




Figure 2. Four-membered N-heterocycles with one substituent R (-COOH; -CHO; -NH₂ etc.). Names refer to the unsubstituted rings. The left and right 1-azetidines are substituted at the 2 and 4 positions, respectively, and so are the left and right 2-azetidines. While both Kekulé structures of azete are shown, they are finally counted only once.

Counting resonance structures separately

As mentioned above, it is convenient to first count resonance structures separately and later modify the method so as to consider them in the way usual in chemistry. We denote the

numbers of these (partially hypothetical) structures by y_n . As mentioned above, the initial values read:

$$y_3 = 4, y_4 = 7. \quad (1)$$

Now we derive a recursion formula for calculating the number series y_k . Assume we know all y_k from $k = 3$ (see Fig. 1) up to $k = n$ and wish to calculate y_{n+1} . We consider the bond between the C_α and C_β atoms and insert an additional carbon between these two atoms (see Fig. 3). In this way, we generate an $(n+1)$ -membered ring.

If the original bond under consideration is a single bond, we make two single bonds out of it. If it is a double bond, we keep that type next to the C_α atom and make the second new bond a single bond. This is located between the new and original C_β atoms, with the latter being the new C_γ atom. The question arises how we can add a double bond between these two carbon atoms. As we do not know the type of bond between the original C_β and C_γ atoms (if $n > 3$) or between the original C_β and N atoms (if $n = 3$) and exclude allenic bonds, we cannot simply add such a bond. Instead, we start from the $(n-1)$ -membered ring and insert two additional carbons between the original C_α and C_β atoms. We do that in such a way that we keep the type of bond next to the C_α atom. If it is a single bond, we will insert a double bond between the new C_β and C_γ and a single bond between the new C_γ and C_δ (or N). If it is a double bond, we will insert a single bond between the new C_β and C_γ and a double bond between the new C_γ and C_δ (or N).




Figure 3. Illustration of the recursive enumeration method. Solid dots, variable chain length. R, substituent. Red bar (or green bar) together with black bar: double bond that is maintained in the transition to ring size $n+1$. Blue bar together with black bar: double bond that is inserted in the extension provided that the bond between C_α and C_β was a single bond. Further explanations, see text.

Combining the two procedures (starting at $n-1$ and n), we arrive at the recursion equation

$$y_{n+1} = y_n + y_{n-1} \quad (2)$$

with the initial conditions (1).

There is no overlap between the molecules thus generated because the y_n molecules generated by starting from length n have a single bond between the new C β and C γ , while the y_{n-1} molecules generated by starting from length $n-1$ have a double bond at that position or, alternatively, the pattern double-single-double, which does not occur in the transition from n to $n+1$. Moreover, all possibilities of extending the molecules according to the defined rules are covered.

Eqs. (1) and (2) define the series of Lucas numbers (Koshy 2001, Taskara et al. 2010, Azarian 2012, Fowler et al. 2014). The series reads

$$y_n = 1, 3, 4, 7, 11, 18, 29, 47, 76, 123, 199, 322, 521, 843, \dots, \quad (3)$$

where we included to $y_1 = 1$ and $y_2 = 3$. It is called after French mathematician François Édouard Anatole Lucas. In the Online Encyclopedia of Integer Sequences (www.oeis.org), the Lucas series is, together with some of its properties, given under no. A000204. It is worth noting that the recursion formula (2) with the alternative initial conditions $x_1 = 1$, $x_2 = 1$ leads to the famous Fibonacci numbers (Koshy 2001).

There is a related problem in mathematics. In graph theory, a matching of a graph is a set of edges (arcs) without common vertices (Diestel 2000). That edge set corresponds to the double bonds in biomolecules because the latter must not be adjacent. The total number of matchings is the Hosoya index or topological index, which leads to the Fibonacci series in the case of linear graphs (Hosoya 1973, 1988, Schuster et al. 2017). It can also be computed for monocyclic graphs and then leads to the Lucas series (Hosoya 1973). The topological index has been used also in chemistry, notably for coding the complicated structures of cyclic and branched hydrocarbons and for predicting some of their thermodynamic quantities such as the boiling point (Hosoya 1973). Hosoya (1988, 2005) used that quantity to extend the famous Hückel rule.

The topological index is sometimes written as the sum of the numbers $p(G,k)$ of matchings including $k = 0, 1, 2$ etc. edges (with G denoting the graph) (Koshy, 2001). In the application under study here, these are the numbers of substances with 0, 1, 2 etc. double bonds. To our knowledge, we are the first to have noticed the relationship to double bonds and, thus, the use of the topological index for enumerating congeners with varying numbers of double bonds (Schuster et al, 2017; Fichtner et al., 2017).

As pointed out by Hosoya (1973), ring-shaped molecules can formally be considered even from $n=1$ on. A monogon and a digon may be defined, respectively, as a point graph and a graph with two points joined by two lines. We can formally define that there is only one structure of a one-membered ring. There are three matchings for the digon: no edge, one of the edges or the other one. While we started with the ring sizes of three and four and, accordingly with y_3 and y_4 , we can go backwards in the recursion to $y_2 = 3$ and $y_1 = 1$.

Eq. (2) is a recursion equation in that the numbers for certain ring sizes are calculated from the numbers of smaller rings. It would be helpful to derive an explicit formula, by which the number y_n can directly be calculated from the ring size, n . This can be achieved by an exponential ansatz $y_n = \lambda^n$. Substituting this into Eq. (2) leads to the quadratic equation $\lambda^2 - \lambda - 1 = 0$. The two solutions of that equation are then used as the bases of the linear combination of two exponential functions (Koshy, 2001):

$$y_n = \left(\frac{1+\sqrt{5}}{2}\right)^n + \left(\frac{1-\sqrt{5}}{2}\right)^n \quad (4)$$

The coefficients in this linear combination are both equal to one; they have been determined from the initial values $y_1 = 1$, $y_2 = 3$. The quantity $(1 + \sqrt{5})/2$ is the famous Golden Ratio, 1.618... (Koshy, 2001). As the absolute value of the second basis, $|1 - \sqrt{5}|/2$, is between zero and one, the influence of the second term in the explicit formula gets smaller and smaller as n increases. Already from $n = 2$ on, we can simplify the formula to

$$y_n = \text{round}\left(1 + \frac{\sqrt{5}}{2}\right)^n \quad (5)$$

This implies that the limit of the ratio of two consecutive numbers is the Golden Ratio.

Alternative formula based on binomial coefficients

Besides the explicit formula (5) for the Fibonacci numbers, there is an alternative formula in the form of a sum of binomial coefficients, which was derived by Lucas in 1876 (cf. Koshy, 2001; Azarian, 2012). For the case of fatty acids and unbranched aliphatic amino acids, that formula has an instructive interpretation in terms of double bonds (Schuster et al., 2017): each binomial coefficient $\binom{n-k}{k}$ quantifies the number of structures involving exactly k double bonds (corresponding to the k -combinations in mathematics). It involves $n-k$ rather than n because double bonds must not be adjacent to each other. Inspired by that reasoning, we have derived an analogous formula for Lucas numbers:

$$y_n = \sum_{k=0}^r \binom{n+1-k}{k} - \sum_{k=0}^s \binom{n-3-k}{k} \quad (6)$$

where $r = \text{floor}[(n+1)/2]$ and $s = \text{floor}[(n-3)/2]$. The floor function denotes the largest integer that is less than or equal to its argument.

Formula (6) can be explained as follows. We cut the ring at the N atom and thus transform it to an acyclic (straight) molecule. Formally, the N atom now occurs twice (Fig. 4). The first sum in Eq. (6) gives the number of congeners in straight molecules (Schuster et al. 2017). The formula there slightly differs because in fatty acids, no double bond is allowed next to the carboxyl group.

Now we have to take into account that in the cyclic molecule, there cannot be two adjacent double bonds next to the N atom. Therefore, we have to subtract all congeners that have both a double bond at the upper end and another one at the lower end of the straight molecule. Since at the second and second last positions, only single bonds are allowed then,

we need to subtract the possibilities in the central part, which involves four bond positions less (in the red ellipse in Fig. 4). Thus, the sum is running up to $n-3-k$ rather than $n+1-k$.



Figure 4. Schematic representation of the transformation of the ring molecule into a straight-chain aliphatic molecule by a cut at the heteroatom, to explain the derivation of formula (6). Alternatively, between the N atom (here shown twice) and any of the two adjacent C atoms, there might be a single bond. However, to explain the second term in Eq. (6), we consider the case where there are double bonds at both positions.

Combining resonance structures

Now, we return to taking into account that rings with fully conjugated double bond systems have the property that the double bonds are delocalised. Thus, it cannot be said whether these bonds are located at odd-numbered (1, 3, ...) or even-numbered positions. This applies not only to aromatic rings (e.g. picolinic acid) but also to non-aromatic rings. Even if the latter rings subsist in non-planar conformations (e.g. boat conformation), these are rather short-lived and undergo transitions to other conformations. Every now and then, such transitions lead to a transient planar conformation, which enables a delocalisation of electrons. Therefore, the numbers of congeners of all rings with even-numbered ring sizes are by one lower than given by the Lucas series.

Let x_n denote the number of theoretically possible, ring-shaped congeners involving n carbons, without distinguishing resonance structures. Including formally the numbers x_1 and x_2 , we obtain the series:

$$x_n = 1, 2, 4, 6, 11, 17, 29, 46, 76, 122, 199, 321, 521, 842, \dots \quad (7)$$

Also this series is given in the Online Encyclopedia of Integer Sequences (www.oeis.org), notably under number A014217.

The monogon can only involve either the C α atom or a nitrogen but not both. Thus, it does not correspond to a meaningful amino acid. In the digon, the C α atom and nitrogen can only be linked by one bond, either a single or a double bond. In the context of amino acids, these two possibilities correspond to glycine and dehydroglycine (iminoacetic acid). Interestingly, the second number in series (7), which is 2, correctly describes the number of structures for $n = 2$. This is because the two hypothetical situations where the double bond is on one or the other side are equivalent anyway.

Also the number series x_n can be calculated explicitly, notably by a formula given in www.oeis.org (number A014217):

$$x_n = \text{floor}\left\{\left(1 + \frac{\sqrt{5}}{2}\right)^n\right\} \quad (8)$$

For example, $(1.618\dots)^4 = 6.854\dots$, which gives 6. To understand formula (8), one can start from eq. (4) and take into account that the basis of the second power term therein is $-0.618\dots$, which is the negative reciprocal of the Golden Ratio. The powers $[-0.618\dots]^n$ are numbers between -1 and 0 for odd n and between 0 and 1 for even n . Thus, y_n is equal to $\text{floor}\{[(1 + \sqrt{5})/2]^n\}$ for odd n and $\text{floor}\{[(1 + \sqrt{5})/2]^n\} + 1$ for even n . Together with the definition of x_n in terms of y_n , this reasoning leads to eq. (8).

The alternative formula (6) can be modified as follows to give the number series (7):

$$x = \sum_{k=0}^r \binom{n+1-k}{k} - \sum_{k=0}^s \binom{n-3-k}{k} - \frac{1+(-1)^n}{2} \quad (9)$$

For every even n , the additional term equals -1 , while for every odd n , it equals zero.

Discussion

Here, we have presented a method for calculating the number of all possible azaheterocycles as a function of ring size, with the ~~additional~~ flexibility that various numbers of double bonds are allowed. We have excluded adjacent double bonds because they occur very rarely in ring-shaped molecules. We have given both a (relatively simple) recursion equation and two explicit formulas. The resulting number series is closely related to the Lucas series known from number theory (Hosoya 1973, Koshy 2001, Taskara et al. 2010, Azarian 2012) in that every second number is equal to a Lucas number while the numbers in between are by one lower than in the Lucas series (www.oeis.org, accession no. number A014217).

In addition to an explicit formula for the Lucas series, which is based on the Golden Ratio (Koshy, 2001), we have derived a second formula, based on binomial coefficients, Eq. (6). To the best of our knowledge, this is a novel result. While Taskara et al. (2010) and Azarian (2012) dealt with the relationships between Lucas numbers and binomial coefficients, they have not given Eq. (6). Equation (iii) in Theorem 2.1 given by Azarian (2012) reads

$$\frac{1}{5}[y_{n+5} - y_{n+1}] = \sum_{k=0}^q \binom{n+2-k}{k} \quad (10)$$

with $q = \text{floor}[(n+2)/2]$. Eq. (10) seems to be related to Eq. (6). Note that the terms $n+1-k$ and $n-3-k$ in Eq. (6) differ by 4, and so do the subscripts in Eq. (10). It is worthwhile checking in future studies whether one of these two equations can be derived from the other one.

As mentioned in the Introduction, acyclic aliphatic amino acids have been enumerated earlier (Grützmann et al., 2011; Fichtner et al., 2017). For the unbranched substances including single and possibly double bonds, the famous Fibonacci series is obtained (Fichtner et al., 2017). Importantly, the ratio of two consecutive numbers tends to the Golden Ratio for the Fibonacci series, the Lucas series and for the number series (7) derived here.

The presented method for enumeration bears manifold potential applications. In metabolomics, it is of interest to know how many different substances can be expected, and so it is in synthetic biology. In the latter field, this helps to assess how long it would take, approximately, to synthesize a complete library of them. Analogous considerations are

relevant in patent applications. In the study of evolution, it is useful to analyse possible alternatives to the biomolecules found in extant living organisms.

Enumeration by the same method is also applicable to cyclic azacycloalkane or azacycloalkene analogs with more than one substituent. An example is azirinomycin, which is 3-methyl-2H-azirine-2-carboxylic acid (Stapley et al. 1971). It is worth mentioning that rings without any substituent obey different enumeration rules because they show a higher symmetry. The same is true for some molecules with several identical substituents if an additional symmetry plane occurs.

Other applications of Lucas numbers in mathematical chemistry have been found earlier. In addition to the above-mentioned applications in chemistry (Hosoya 1973, 1988, 2005), that series is also useful to describe the numbers of Kekulé structures in certain classes of polycyclic aromatic compounds. For $[n]$ circulenes, the number 2 has to be added to every second Lucas number (Bergan et al., 1987; Cyvin et al., 1991). The series itself can be used for a class of aromatic compounds for which Fowler et al. (2014) suggested the term lucacenes.

Three- and four-membered heterocyclic rings are strained due to their small bond angles and thus, most of them are highly reactive and unstable (Behre et al., 2012, Banert et al. 2013). Moreover, 1H-azirines are much less stable than their tautomers 2H-azirines because they are, in addition, anti-aromatic and are, thus, very unlikely to be found in natural products (Zavoruev et al. 1990, Banert et al. 2013). It is a difficult decision what the minimum lifetime of a molecule should be so that it is considered as a separate substance. We leave this decision to further studies and have here taken into account, a priori, all conceivable structures of the type under study with one exception: different resonance/Kekulé structures of aromatic compounds.

The term enumeration has two meanings. In the sense used in the present paper, it refers to calculating the number of possible structures. In contrast, constructive enumeration requires that all structures are listed successively in silico, for example, by a graph-theoretical approach (Lukovits, 2003). For small and moderate ring sizes, this can be done relatively easily, as we have done in Figs. 1 and 2. For larger rings, it will be of interest to establish a suitable code for systematic, constructive enumeration in the future. Implementation into virtual compound libraries, which are used in computer-assisted drug discovery, testing of hypotheses and experiment optimization, are potential promising applications (Faulon et al. 2005).

Acknowledgment: The authors thank Severin Sasso (Leipzig) and Hans-Dieter Arndt (Jena) for stimulating discussions. Financial support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) within the SFB 1127 ChemBioSys, grant no. 239748522, is gratefully acknowledged.

Literature

- Arp, J., S. Götze, R. Mukherji, D.J. Mattern, M. García-Altare, M. Klapper, et al. (2018). Synergistic activity of cosecreted natural products from amoebae-associated bacteria. *Proceedings National Academy Sciences U.S.A.* 115(15): 3758-3763.

- Azarian, M.K. (2012). Identities involving Lucas or Fibonacci and Lucas numbers as binomial sums. *International Journal of Contemporary Mathematical Sciences*, Vol. 7, 2012, no. 45, 2221 - 2227
- Balaban, A. T., J. W. Kennedy and L. Quintas (1988). The number of alkanes having n carbons and a longest chain of length d : an application of a theorem of Pólya. *Journal of Chemical Education* 65(4): 304.
- Banert, K., S. Bochmann, M. Hagedorn and F. Richter (2013). Stable but chimeric antiaromatic 1H-azirines? A threefold reinvestigation. *Tetrahedron Letters* 54(46): 6185-6188.
- Behre, J., R. Voigt, I. Althöfer and S. Schuster (2012). On the evolutionary significance of the size and planarity of the proline ring. *Naturwissenschaften* 99(10): 789-799.
- Bergan, J.L., Cyvin, B.N. and Cyvin S.J. (1987). The Fibonacci numbers, and Kekulé structures of some corona-condensed benzenoids (corannulenes). *Acta Chimica Hungarica* 124: 299-314.
- Böcker, S. and S. Wagner (2014). Counting glycans revisited. *Journal of Mathematical biology* 69(4): 799-816.
- Cyvin, S.J., E. Brendsdal, J. Brunvoll and M. Skaret (1991). Corannulene as a member of circulenes: its topological properties and molecular vibrations. *Journal of Molecular Structure* 247: 119-127.
- Diestel, R. (2000). An accessibility theorem for infinite graph minors. *Journal of Graph Theory* 35(4): 273-277.
- Faulon, J. L., D. P. Visco Jr and D. Roe (2005). Enumerating molecules. *Reviews in computational Chemistry* 21: 209-286.
- Fichtner, M., K. Voigt and S. Schuster (2017). The tip and hidden part of the iceberg: Proteinogenic and non-proteinogenic aliphatic amino acids. *Biochimica et Biophysica Acta (BBA)-General Subjects* 1861(1): 3258-3269.
- Fowden, L. (1963). Amino-acid analogues and the growth of seedlings. *Journal of Experimental Botany* 14(3): 387-398.
- Fowler, P. W., S. Cotton, D. Jenkinson, W. Myrvold and W. H. Bird (2014). Equiaromatic benzenoids: Arbitrarily large sets of isomers with equal ring currents. *Chemical Physics Letters* 597: 30-35.
- Grützmann, K., S. Böcker and S. Schuster (2011). Combinatorics of aliphatic amino acids. *Naturwissenschaften* 98(1): 79-86.
- Haranczyk, M., T. Puzyn and E.G. Ng (2010). On enumeration of congeners of common persistent organic pollutants. *Environmental Pollution* 158(8): 2786-2789.
- Henze, H. R. and C. M. Blair (1931). The number of structurally isomeric alcohols of the methanol series. *Journal of the American chemical society* 53(8): 3042-3046.
- Hosoya, H. (1973). Topological index and Fibonacci numbers with relation to chemistry. *Fibonacci Quart* 11(3): 255-266.
- Hosoya, H. (1988). "On some counting polynomials in chemistry." *Discrete applied mathematics* 19(1-3): 239-257.
- Hosoya, H. (2005). Aromaticity index can predict and explain the stability of polycyclic conjugated hydrocarbons. *Monatshefte für Chemie/Chemical Monthly* 136: 1037-1054.

- Koshy, T. (2001). *Fibonacci and Lucas Numbers with Applications*, Volume 2, John Wiley & Sons.
- Kroteń, M.A., M. Bartoszewicz and I. Swiecicka (2010). Cereulide and valinomycin, two important natural dodecadeptide with ionophoretic activities, *Polish Journal of Microbiology* 59: 3–10.
- Lukovits, I., A. Graovac, E. Kálmán, G. Kaptay, P. Nagy, S. Nikolić, J. Sytchev and N. Trinajstić (2003). Nanotubes: Number of Kekulé structures and aromaticity. *Journal of chemical information and computer sciences* 43(2): 609-614.
- Neuenschwander, U. and I. Hermans (2011). The conformations of cyclooctene: consequences for epoxidation chemistry. *The Journal of Organic Chemistry* 76(24): 10236-10240.
- Schuster, S., M. Fichtner and S. Sasso (2017). Use of Fibonacci numbers in lipidomics—Enumerating various classes of fatty acids. *Scientific reports* 7(1): 1-9.
- Stapley, E.O., D. Hendlin, M. Jackson, A.K. Miller, S. Hernandez and J.M. Mata (1971). Azirinomycin. I Microbial production and biological characteristics. *The Journal of Antibiotics* 24(1): 42-47.
- Taskara, N., K. Uslu and H.H. Gulec (2010). On the properties of Lucas numbers with binomial coefficients. *Applied Mathematics Letters* 23: 68-72.
- Zahradníčková, H., S. Opekar, L. Římnáčová, P. Šimek and M. Moos (2022). Chiral secondary amino acids, their importance, and methods of analysis. *Amino Acids* 54(5): 687-719.
- Zavoruev, S. and R.-I. Rakauskas (1990). Thermodynamic stability and reactivity of the antiaromatic heterocycle 1H-azirine. *Theoretical and Experimental Chemistry* 25(4): 445-450.