

ON THE SYMPLECTIC STRUCTURE OVER THE MODULI SPACE OF PROJECTIVE STRUCTURES ON A SURFACE

INDRANIL BISWAS

ABSTRACT. The moduli space of projective structures on a compact oriented surface Σ has a holomorphic symplectic structure, which is constructed by pulling back, using the monodromy map, the Atiyah–Bott–Goldman symplectic form on the character variety $\text{Hom}(\pi_1(\Sigma), \text{PSL}(2, \mathbb{C}))/\text{PSL}(2, \mathbb{C})$. We produce another construction of this symplectic form. This construction shows that the symplectic form on the moduli space is actually algebraic. Note that the monodromy map is only holomorphic and *not* algebraic, so the first construction does not give algebraicity of the pulled back form.

1. INTRODUCTION

Let Σ be a compact oriented C^∞ surface of genus g . The corresponding Teichmüller space $\text{Teich}(\Sigma)$ is the quotient of $\text{Com}(\Sigma)$, the space of C^∞ complex structures on Σ compatible with its orientation, by the natural action of the group $\text{Diff}^0(\Sigma)$ given by the connected component, containing the identity element, of the group $\text{Diff}^+(\Sigma)$ defined by the orientation preserving diffeomorphisms of Σ . The mapping class group $\text{Mod}(\Sigma) := \text{Diff}^+(\Sigma)/\text{Diff}^0(\Sigma)$ acts on $\text{Teich}(\Sigma)$, and the quotient $\mathcal{M}_g := \text{Teich}(\Sigma)/\text{Mod}(\Sigma)$ is the moduli space of smooth complex projective curves of genus g .

A projective structure on Σ is defined by giving a coordinate atlas, where each coordinate function has target \mathbb{CP}^1 , such that all the transition functions are Möbius transformations (same as holomorphic automorphisms of \mathbb{CP}^1). Let $\text{Proj}(\Sigma)$ denote the space of C^∞ projective structures on Σ compatible with its orientation. A projective structure on Σ produces a flat $\text{PSL}(2, \mathbb{C})$ -bundle on Σ together with a (non-flat) reduction of structure group to a Borel subgroup of $\text{PSL}(2, \mathbb{C})$. Sending a projective structure to the monodromy homomorphism of the corresponding flat $\text{PSL}(2, \mathbb{C})$ -bundle, we obtain a holomorphic map from $\mathcal{P}(\Sigma) := \text{Proj}(\Sigma)/\text{Diff}^0(\Sigma)$ to the $\text{PSL}(2, \mathbb{C})$ -character variety for Σ . This map is a local biholomorphism, and hence the pullback of the Atiyah–Bott–Goldman symplectic form on the character variety gives a holomorphic symplectic form on $\mathcal{P}(\Sigma)$. The symplectic form on the character variety is preserved by the action of $\text{Mod}(\Sigma)$, and also the monodromy map is $\text{Mod}(\Sigma)$ -equivariant. Therefore, the above holomorphic symplectic form on $\mathcal{P}(\Sigma)$ descends to a holomorphic symplectic form Θ_g on $\mathcal{P}(\Sigma)_g := \mathcal{P}(\Sigma)/\text{Mod}(\Sigma)$.

Now, $\mathcal{P}(\Sigma)_g$ is an algebraic orbifold; in fact, it is an algebraic torsor over \mathcal{M}_g for the cotangent bundle $\Omega_{\mathcal{M}_g}$. The above monodromy map is *not* algebraic because the degree of the map is infinite.

2010 *Mathematics Subject Classification.* 14H15, 32G15, 14H60.

Key words and phrases. Projective structure, character variety, symplectic form, Teichmüller space.

Our aim here is to give an alternative construction of the symplectic form Θ_g on $\mathcal{P}(\Sigma)_g$. Our construction shows that the form Θ_g is actually algebraic.

2. A PARAMETER SPACE FOR PROJECTIVE STRUCTURES

2.1. Projective structure. A holomorphic coordinate chart on a connected Riemann surface X is a pair (U, ϕ) , where $U \subset X$ is an open subset and $\phi : U \rightarrow \mathbb{CP}^1$ is a holomorphic embedding. A holomorphic coordinate atlas on X is a collection of coordinate charts $\{(U_i, \phi_i)\}_{i \in I}$ such that $X = \bigcup_{i \in I} U_i$. A projective structure on X is given by a holomorphic coordinate atlas $\{(U_i, \varphi_i)\}_{i \in I}$ such that for every $i, j \in I \times I$ and every nonempty connected component $V_c \subset U_i \cap U_j$, there is an element $A_{j,i}^c \in \text{Aut}(\mathbb{CP}^1) = \text{PSL}(2, \mathbb{C})$ whose restriction to $\phi_i(V_c)$ coincides with $(\phi_j \circ \phi_i^{-1})|_{\phi_i(V_c)}$. Two holomorphic coordinate atlases $\{(U_i, \phi_i)\}_{i \in I}$ and $\{(U_i, \phi_i)\}_{i \in I'}$ satisfying the above condition are called *equivalent* if their union $\{(U_i, \phi_i)\}_{i \in I \cup I'}$ also satisfies the above condition. A *projective structure* on X is an equivalence class of holomorphic coordinate atlases satisfying the above condition (see [Gu]).

Fix the Borel subgroup

$$B := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}(2, \mathbb{C}) \mid b = 0 \right\} \subset \text{PSL}(2, \mathbb{C}). \quad (2.1)$$

Let $P \rightarrow X$ be a holomorphic principal $\text{PSL}(2, \mathbb{C})$ -bundle on X equipped with a holomorphic connection D . Let $P_B \subset P$ be a holomorphic reduction of structure group, to the subgroup B in (2.1), given by a holomorphic section

$$\sigma : X \rightarrow P/B \quad (2.2)$$

of the natural projection $f : P/B \rightarrow X$. The connection D on P induces a connection on the fiber bundle f , which, in turn, decomposes the tangent bundle $T(P/B)$ as $T(P/B) = T_f \oplus \mathcal{H}$, where $T_f \subset T(P/B)$ is the relative tangent bundle for the projection f and $\mathcal{H} \subset T(P/B)$ is the horizontal subbundle for the connection on P/B . Let

$$\tilde{\sigma} : TX \rightarrow \sigma^* T_f \quad (2.3)$$

be the composition of homomorphisms

$$TX \xrightarrow{d\sigma} \sigma^* T(P/B) = \sigma^* T_f \oplus \sigma^* \mathcal{H} \rightarrow \sigma^* T_f$$

constructed using the above decomposition of $T(P/B)$, where $d\sigma$ is the differential of σ in (2.2).

A projective structure on X is a triple (P, P_B, D) as above such that the homomorphism $\tilde{\sigma}$ in (2.3) is an isomorphism [Gu].

The holomorphic cotangent bundle of X will be denoted by K_X . The extensions of \mathcal{O}_X by K_X are parametrized by

$$H^1(X, \text{Hom}(\mathcal{O}_X, K_X)) = H^1(X, K_X) = \mathbb{C}.$$

Let

$$0 \rightarrow K_X \rightarrow V \rightarrow \mathcal{O}_X \rightarrow 0 \quad (2.4)$$

be the extension corresponding to $1 \in H^1(X, \text{Hom}(\mathcal{O}_X, K_X)) = \mathbb{C}$.

Let

$$P^0 \longrightarrow X \tag{2.5}$$

be the principal $\text{PSL}(2, \mathbb{C})$ -bundle given by $\mathbb{P}(V)$; the fiber of P^0 over any $x \in X$ is the space of all holomorphic isomorphisms $\mathbb{C}\mathbb{P}^1 \longrightarrow P_x^0$. The line subbundle $K_X \hookrightarrow V$ in (2.4) produces a reduction of structure group

$$P_B^0 \subset P^0 \tag{2.6}$$

to the Borel subgroup $B \subset \text{PSL}(2, \mathbb{C})$ in (2.1). The fiber of P_B^0 over any $x \in X$ is the space of all isomorphisms $\mathbb{C}\mathbb{P}^1 \longrightarrow P_x^0$ that take the point $\mathbb{C} \cdot e_2 \in \mathbb{C}\mathbb{P}^1$ to the point of P_x^0 given by the line $(K_X)_x \subset V_x$, where $\{e_1, e_2\}$ is the standard basis of \mathbb{C}^2 .

For any projective structure (P, P_B, D) on X , the pair (P, P_B) coincides with (P^0, P_B^0) [Gu]. Furthermore, for every connection D on P^0 , the triple (P^0, P_B^0, D) is actually a projective structure on X . From these it follows that the space of all projective structures on X is an affine space modeled on the vector space $H^0(X, K_X^{\otimes 2})$.

2.2. A symplectic form. Fix a compact oriented C^∞ surface Σ of genus $g \geq 2$. Denote by $\text{Diff}^+(\Sigma)$ the group of all orientation preserving diffeomorphisms of Σ . Let $\text{Diff}^0(\Sigma) \subset \text{Diff}^+(\Sigma)$ be the connected component containing the identity element. Then the quotient

$$\text{Mod}(\Sigma) := \pi_0(\text{Diff}^+(\Sigma)) = \text{Diff}^+(\Sigma)/\text{Diff}^0(\Sigma)$$

is known as the mapping class group. Let $\text{Com}(\Sigma)$ denote the space of C^∞ complex structures on Σ compatible with its orientation. The quotient $\text{Teich}(\Sigma) := \text{Com}(\Sigma)/\text{Diff}^0(\Sigma)$ for the natural action of $\text{Diff}^0(\Sigma)$ on $\text{Com}(\Sigma)$ is the Teichmüller space for Σ . The group $\text{Mod}(\Sigma)$ acts property discontinuously on $\text{Teich}(\Sigma)$. The corresponding orbifold

$$\mathcal{M}_g = \text{Teich}(\Sigma)/\text{Mod}(\Sigma)$$

is the moduli space of smooth complex projective curves of genus g .

Let $\text{Proj}(\Sigma)$ denote the space of all projective structures on the C^∞ surface Σ compatible with its orientation. The group $\text{Diff}^+(\Sigma)$ has a natural action on it. Consider the quotient

$$\mathcal{P}(\Sigma) := \text{Proj}(\Sigma)/\text{Diff}^0(\Sigma).$$

The natural map $\text{Proj}(\Sigma) \longrightarrow \text{Com}(\Sigma)$ produces a projection

$$\varpi : \mathcal{P}(\Sigma) \longrightarrow \text{Teich}(\Sigma) \tag{2.7}$$

The action of the group $\text{Diff}^+(\Sigma)$ on $\text{Proj}(\Sigma)$ produces an action of $\text{Mod}(\Sigma)$ acts on $\mathcal{P}(\Sigma)$. Let

$$\mathcal{P}_g := \mathcal{P}(\Sigma)/\text{Mod}(\Sigma) \tag{2.8}$$

be the corresponding orbifold. The map ϖ in (2.7) produces a map

$$\varpi' : \mathcal{P}_g = \mathcal{P}(\Sigma)/\text{Mod}(\Sigma) \longrightarrow \text{Teich}(\Sigma)/\text{Mod}(\Sigma) = \mathcal{M}_g. \tag{2.9}$$

There is a universal family $\mathcal{X}_{\text{Teich}(\Sigma)}/\text{Teich}(\Sigma)$ of Riemann surfaces on $\text{Teich}(\Sigma)$. The fiber of the map ϖ in (2.7) over any $t \in \text{Teich}(\Sigma)$ is an affine space for the space all

quadratic differentials on the Riemann surface $(\mathcal{X}_{\text{Teich}(\Sigma)})_t$. In fact, $\mathcal{P}(\Sigma)$ is a holomorphic torsor over $\text{Teich}(\Sigma)$ for the holomorphic cotangent bundle $\Omega_{\text{Teich}(\Sigma)}$ of $\text{Teich}(\Sigma)$.

We note that \mathcal{P}_g has a natural algebraic structure such that the map ϖ' in (2.9) is algebraic. In fact, \mathcal{P}_g is an algebraic torsor over \mathcal{M}_g for the cotangent bundle $\Omega_{\mathcal{M}_g}$ of \mathcal{M}_g .

Given $x \in \Sigma$, the fundamental group $\pi_1(\Sigma, x)$ has $2g$ generators that are subject to a single relation. So the space of all homomorphisms from $\pi_1(\Sigma, x)$ to $\text{PSL}(2, \mathbb{C})$ that lift to a homomorphism from $\pi_1(\Sigma, x)$ to $\text{SL}(2, \mathbb{C})$ constitute a Zariski closed subset of $\text{PSL}(2, \mathbb{C})^{2g}$. This Zariski closed subset is evidently affine. The action of $\text{PSL}(2, \mathbb{C})$ by conjugation on this affine space of homomorphisms is also in the affine category. The homomorphisms that are irreducible in the sense that their image in $\text{PSL}(2, \mathbb{C})$ is Zariski dense, and lift to a homomorphism from $\pi_1(\Sigma, x)$ to $\text{SL}(2, \mathbb{C})$, make up an open-dense subset $\mathcal{R}'(\Sigma)$. It has the property that $\text{PSL}(2, \mathbb{C})$ acts freely on it. In fact, we can form this orbit variety in the affine category. This affine variety is called the *character variety* of Σ and is denoted by $\mathcal{R}(\Sigma)$. This $\mathcal{R}(\Sigma)$ is smooth of dimension $6(g-1)$. Since the inner automorphisms of $\pi_1(\Sigma, x)$ are killed by this procedure, the variety $\mathcal{R}(\Sigma)$ is actually independent of the choice of our base point $x \in \Sigma$ and (thus) comes with an action of the mapping class group $\text{Mod}(\Sigma)$ (which acts on $\pi_1(\Sigma, x)$ by outer automorphisms). We can interpret $\mathcal{R}(\Sigma)$ as the moduli space of topologically trivial flat principal $\text{PSL}(2, \mathbb{C})$ -bundles on Σ with irreducible monodromy.

The character variety $\mathcal{R}(\Sigma)$ also comes with a closed nondegenerate 2-form (to which we refer simply as an *algebraic symplectic structure*), [AB], [Go], that we shall denote by

$$\Theta_\Sigma \in H^0(\mathcal{R}(\Sigma), \bigwedge^2 T^*\mathcal{R}(\Sigma)). \quad (2.10)$$

It is natural, and therefore it is preserved by the action of $\text{Mod}(\Sigma)$. In [BL] an alternative construction of Θ_Σ is given.

Since a projective structure on Σ determines a principal $\text{PSL}(2, \mathbb{C})$ -bundle with a flat connection, we have a natural, $\text{Mod}(\Sigma)$ -equivariant monodromy map

$$\Psi : \mathcal{P}(\Sigma) \longrightarrow \mathcal{R}(\Sigma). \quad (2.11)$$

This map Ψ is known to be a local (holomorphic) isomorphism [Hu, p. 272, Theorem], [He], but its fibers are not finite (this follows from [Ba, p. 141, Theorem 6.2] (see [Ba, p. 120]) and [GKM]). The pullback of the form in (2.10)

$$\Psi^*\Theta_\Sigma \in H^0(\mathcal{P}(\Sigma), \bigwedge^2 T^*\mathcal{P}(\Sigma))$$

is a holomorphic symplectic form on $\mathcal{P}(\Sigma)$. Since Θ_Σ is preserved by the action of $\text{Mod}(\Sigma)$, and Ψ is $\text{Mod}(\Sigma)$ -equivariant, it follows that $\Psi^*\Theta_\Sigma$ descends to a holomorphic symplectic form on \mathcal{P}_g (see (2.8))

$$\Theta_g \in H^0(\mathcal{P}_g, \bigwedge^2 T^*\mathcal{P}_g). \quad (2.12)$$

3. SYMPLECTIC FORM ON MODULI OF PROJECTIVE STRUCTURES

Recall that \mathcal{P}_g has an algebraic structure. We will construct an algebraic 2-form on \mathcal{P}_g . For that we need to describe the tangent bundle $T\mathcal{P}_g$.

3.1. Infinitesimal deformations of projective structures. Let X be a smooth complex projective curve X of genus $g \geq 2$, and let δ be a projective structure on X . Then there is a holomorphic differential operator of order three

$$\Delta_\delta : TX \longrightarrow K_X^{\otimes 2}; \quad (3.1)$$

we will briefly recall its construction (see [Hu, pp. 264–265]). Since the space of projective structures on an open subset $U \subset X$ is an affine space for $H^0(U, K_U^{\otimes 2})$, given $\chi \in H^0(U, TU)$, and any coordinate function ϕ on U compatible with δ , the Lie derivative $L_\chi \phi$ of ϕ with respect to χ is an element of $H^0(U, K_U^{\otimes 2})$. This element

$$L_\chi \phi \in H^0(U, K_U^{\otimes 2}) \quad (3.2)$$

is independent of the choice of ϕ . The differential operator Δ_δ in (3.1) sends χ to $L_\chi \phi$.

We will describe an alternative construction of Δ_δ . The natural isomorphism of $\text{Aut}(\mathbb{CP}^1)$ with $\text{PSL}(2, \mathbb{C})$ gives an isomorphism

$$\mathfrak{sl}(2, \mathbb{C}) \xrightarrow{\sim} H^0(\mathbb{CP}^1, T\mathbb{CP}^1), \quad (3.3)$$

where $\mathfrak{sl}(2, \mathbb{C})$ is the Lie algebra of $\text{PSL}(2, \mathbb{C})$, that takes the Lie algebra operation on $\mathfrak{sl}(2, \mathbb{C})$ to the Lie bracket operation of vector fields. For any $n \geq 0$, let

$$\eta_n : \mathbb{CP}^1 \times H^0(\mathbb{CP}^1, T\mathbb{CP}^1) \longrightarrow J^n(T\mathbb{CP}^1) \quad (3.4)$$

be the natural restriction map of global sections, where $\mathbb{CP}^1 \times H^0(\mathbb{CP}^1, T\mathbb{CP}^1)$ is the trivial vector bundle on \mathbb{CP}^1 with fiber $H^0(\mathbb{CP}^1, T\mathbb{CP}^1)$. The map η_2 is an isomorphism. So the composition of homomorphisms

$$\eta_3 \circ (\eta_2)^{-1} : J^2(T\mathbb{CP}^1) \longrightarrow J^3(T\mathbb{CP}^1)$$

has the following property: Consider the natural jet sequence

$$0 \longrightarrow K_{\mathbb{CP}^1}^{\otimes 2} \xrightarrow{\alpha} J^3(T\mathbb{CP}^1) \xrightarrow{\beta} J^2(T\mathbb{CP}^1) \longrightarrow 0. \quad (3.5)$$

We have $\beta \circ (\eta_3 \circ (\eta_2)^{-1}) = \text{Id}_{J^2(T\mathbb{CP}^1)}$. We note that $\eta_3 \circ (\eta_2)^{-1}$ is equivariant for the actions of $\text{PSL}(2, \mathbb{C})$ on $J^3(T\mathbb{CP}^1)$ and $J^2(T\mathbb{CP}^1)$ given by the action of $\text{PSL}(2, \mathbb{C})$ on \mathbb{CP}^1 .

Let

$$\Delta_{\mathbb{P}^1} : J^3(T\mathbb{CP}^1) \longrightarrow K_{\mathbb{CP}^1}^{\otimes 2} \quad (3.6)$$

be the unique homomorphism such that $\text{kernel}(\Delta_{\mathbb{P}^1}) = \eta_3((\eta_2)^{-1}(J^2(T\mathbb{CP}^1)))$ and

$$\Delta_{\mathbb{P}^1} \circ \alpha = \text{Id}_{K_{\mathbb{CP}^1}^{\otimes 2}}, \quad (3.7)$$

where α is the homomorphism in (3.5). So $\Delta_{\mathbb{P}^1} \circ \alpha$ is a holomorphic differential operator of order three from $T\mathbb{CP}^1$ to $K_{\mathbb{CP}^1}^{\otimes 2}$ which is invariant under the action of $\text{PSL}(2, \mathbb{C})$. The symbol of $\Delta_{\mathbb{P}^1} \circ \alpha$ is a holomorphic section of $\text{Hom}(T\mathbb{CP}^1, K_{\mathbb{CP}^1}^{\otimes 2}) \otimes (T\mathbb{CP}^1)^{\otimes 3} = \mathcal{O}_{\mathbb{CP}^1}$.

From (3.7) it follows immediately that the symbol of $\Delta_{\mathbb{P}^1} \circ \alpha$ is the constant function 1 on $\mathbb{C}\mathbb{P}^1$.

Consequently, given any holomorphic coordinate chart (U, ϕ) on X compatible with the projective structure δ , we have a differential operator $TU \rightarrow K_U^{\otimes 2}$ which is simply the transport of $\Delta_{\mathbb{P}^1}$ using the biholomorphism ϕ . Since $\Delta_{\mathbb{P}^1}$ is $\mathrm{PSL}(2, \mathbb{C})$ -invariant, these locally defined differential operators on X patch together compatibly to produce a global third order differential operator $TX \rightarrow K_X^{\otimes 2}$. This differential operator coincides with Δ_δ in (3.1). Indeed, this follows immediately by comparing their expressions in terms of holomorphic coordinate functions on X compatible with the projective structure δ .

So, $\Delta_\delta : J^3(X) \rightarrow K_X^{\otimes 2}$ gives a splitting of the jet sequence

$$0 \rightarrow K_X^{\otimes 2} \rightarrow J^3(TX) \rightarrow J^2(TX) \rightarrow 0 \quad (3.8)$$

on X . From this it follows immediately that the symbol of the differential operator Δ_δ in (3.1) is the constant function 1 (see also [Hu, p. 273]). Consequently, the sheaf of solutions of Δ_δ is a complex local system on X ; let

$$\Lambda_\delta \rightarrow X \quad (3.9)$$

be the local system given by the sheaf of solutions of Δ_δ (the notation of [Hu] is being used). Since Δ_δ is given by a splitting of (3.8), the holomorphic vector bundle underlying the local system Λ_δ is identified with $J^2(TX)$.

Consider the holomorphic principal $\mathrm{PSL}(2, \mathbb{C})$ -bundle P^0 in (2.5). Let D be the holomorphic connection on P^0 corresponding to the projective structure δ . The connection on the adjoint bundle $\mathrm{ad}(P^0)$ (for P^0) induced by D will be denoted by \tilde{D} . The complex local system on X given by the sheaf of flat sections of $\mathrm{ad}(P^0)$, for the connection \tilde{D} , will be denoted by

$$\underline{\mathrm{ad}}(P^0). \quad (3.10)$$

Lemma 3.1. *The two complex local systems Λ_δ (constructed in (3.9)) and $\underline{\mathrm{ad}}(P^0)$ (in (3.10)) on X are canonically identified.*

Proof. The kernel of the projection $\Delta_{\mathbb{P}^1}$ in (3.6) is $J^2(T\mathbb{C}\mathbb{P}^1)$, because it is given by a splitting of the exact sequence in (3.5). In particular, the vector bundle corresponding to the local system defined by the sheaf of solutions of $\Delta_{\mathbb{P}^1}$ is $J^2(T\mathbb{C}\mathbb{P}^1)$. Recall that η_2 identifies $J^2(T\mathbb{C}\mathbb{P}^1)$ with $\mathbb{C}\mathbb{P}^1 \times H^0(\mathbb{C}\mathbb{P}^1, T\mathbb{C}\mathbb{P}^1) = \mathbb{C}\mathbb{P}^1 \times \mathfrak{sl}(2, \mathbb{C})$ (see (3.3)). So the kernel of the projection $\Delta_{\mathbb{P}^1}$ in (3.6) is identified with $\mathbb{C}\mathbb{P}^1 \times \mathfrak{sl}(2, \mathbb{C})$. This identification is evidently $\mathrm{PSL}(2, \mathbb{C})$ -equivariant; since $\Delta_{\mathbb{P}^1}$ is $\mathrm{PSL}(2, \mathbb{C})$ -invariant, the kernel of $\Delta_{\mathbb{P}^1}$ is equipped with an action of $\mathrm{PSL}(2, \mathbb{C})$. This action of $\mathrm{PSL}(2, \mathbb{C})$ on $\mathrm{kernel}(\Delta_{\mathbb{P}^1}) = \mathbb{C}\mathbb{P}^1 \times \mathfrak{sl}(2, \mathbb{C})$ coincides with the diagonal action of $\mathrm{PSL}(2, \mathbb{C})$.

Let (U, ϕ) be any coordinate chart on X compatible with the projective structure δ . Using ϕ , the kernel of $\Delta_{\mathbb{P}^1}$ produces a constant subsheaf of TU . Similarly, the constant sheaf $\mathbb{C}\mathbb{P}^1 \times \mathfrak{sl}(2, \mathbb{C})$ produces a constant sheaf on U with stalk $\mathfrak{sl}(2, \mathbb{C})$, using ϕ . The above isomorphism of the kernel of $\Delta_{\mathbb{P}^1}$ with $\mathbb{C}\mathbb{P}^1 \times \mathfrak{sl}(2, \mathbb{C})$ produces an isomorphism of these two local systems on U .

The local system Λ_δ on X is constructed by patching together these locally defined constant subsheaves of TU , while the local system $\underline{\text{ad}}(P^0)$ is constructed by patching together these locally defined constant sheaves with stalk $\mathfrak{sl}(2, \mathbb{C})$; the patchings are done using the transition functions for the coordinate charts compatible with δ . Now from the above observation that the kernel $\Delta_{\mathbb{P}^1}$ is $\text{PSL}(2, \mathbb{C})$ -equivariantly identified with $\mathbb{C}\mathbb{P}^1 \times \mathfrak{sl}(2, \mathbb{C})$ it follows immediately that these locally defined isomorphisms on X produce a global isomorphism between Λ_δ and $\underline{\text{ad}}(P^0)$. \square

Recall that the vector bundles on X underlying the local systems Λ_δ and $\underline{\text{ad}}(P^0)$ are $J^2(TX)$ and $\text{ad}(P^0)$ respectively. Therefore, Lemma 3.1 produces an isomorphism of $J^2(TX)$ with $\text{ad}(P^0)$; to clarify, this isomorphism depends on the projective structure δ . Using this isomorphism between $J^2(TX)$ and $\text{ad}(P^0)$, the bilinear trace form on $\mathfrak{sl}(2, \mathbb{C})$ produces a fiberwise nondegenerate symmetric pairing

$$f_\delta : J^2(TX) \otimes J^2(TX) \longrightarrow \mathcal{O}_X \quad (3.11)$$

Let

$$\mathcal{B}_\bullet : \mathcal{B}_0 := TX \xrightarrow{\Delta_\delta} \mathcal{B}_1 := K_X^{\otimes 2} \quad (3.12)$$

be the two-term complex on X , where Δ_δ is the differential operator in (3.1); to clarify, the sheaf \mathcal{B}_i in (3.12) is at the i -th position of the complex.

The space of infinitesimal deformations of the pair (X, δ) are parametrized by the hypercohomology $\mathbb{H}^1(\mathcal{B}_\bullet)$ [Hu, p. 266]. From the short exact sequence of sheaves

$$0 \longrightarrow \Lambda_\delta \longrightarrow TX \xrightarrow{\Delta_\delta} K_X^{\otimes 2} \longrightarrow 0$$

(see (3.9)) we know that

$$\mathbb{H}^1(\mathcal{B}_\bullet) = H^1(X, \Lambda_\delta). \quad (3.13)$$

Consider the short exact sequence of complexes

$$\mathcal{B}_\bullet : \begin{array}{ccc} 0 & & 0 \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & K_X^{\otimes 2} \\ \downarrow & & \downarrow \text{id} \\ TX & \xrightarrow{\Delta_\delta} & K_X^{\otimes 2} \\ \downarrow \text{id} & & \downarrow \\ TX & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0 & & 0 \end{array}$$

on X . It produces a short exact sequence of hypercohomologies

$$\begin{aligned} 0 = H^0(X, TX) &\longrightarrow H^0(X, K_X^{\otimes 2}) \xrightarrow{\alpha_1} \mathbb{H}^1(\mathcal{B}_\bullet) \\ &\xrightarrow{\alpha_2} H^1(X, TX) \longrightarrow H^1(X, K_X^{\otimes 2}) = 0. \end{aligned} \quad (3.14)$$

The homomorphism α_1 in (3.14) corresponds to moving the projective structure keeping the Riemann surface X fixed [Hu, p. 267 (iii)], and the homomorphism α_2 corresponds to the forgetful map that sends an infinitesimal deformation of (X, δ) to the infinitesimal deformation of X corresponding to it obtained by simply forgetting the projective structure [Hu, p. 266 (ii)].

3.2. A two-form on the moduli space of projective structures. As before, X is a smooth complex projective curve of genus $g \geq 2$, and δ is a projective structure on X . Consider the differential operator Δ_δ in (3.1). We will show that it satisfies a certain identity.

The natural contraction homomorphism

$$TX \otimes K_X^{\otimes 2} \longrightarrow K_X \quad (3.15)$$

will be denoted by “ \cdot ”. Let

$$p_0 : J^3(TX) \longrightarrow TX \quad (3.16)$$

be the natural projection.

The \mathcal{O}_X -linear homomorphism $\Delta_\delta : J^3(TX) \longrightarrow K_X^{\otimes 2}$ in (3.1) produces an \mathcal{O}_X -linear homomorphism

$$H_1 : \text{Sym}^2(J^3(TX)) \longrightarrow K_X \quad (3.17)$$

that sends any $(v \otimes w) + (w \otimes v) \in J^3(TX)_x \otimes J^3(TX)_x$, $x \in X$, to

$$p_0(v) \cdot \Delta_\delta(w) + p_0(w) \cdot \Delta_\delta(v) \in (K_X)_x,$$

where p_0 is the projection in (3.16), and “ \cdot ” is the pairing in (3.15).

We will construct another homomorphism H_2 from $\text{Sym}^2(J^3(TX) \otimes J^3(TX))$ to K_X . For that we first recall that given vector bundles V and W on X , and $i \geq 0$, there are natural homomorphisms

$$f_1 : J^{i+1}(V) \longrightarrow J^1(J^i(V)) \quad \text{and} \quad f_2 : J^i(V) \otimes J^i(W) \longrightarrow J^i(V \otimes W). \quad (3.18)$$

Now we have the following composition of homomorphisms

$$\begin{aligned} \text{Sym}^2(J^3(TX)) &\hookrightarrow J^3(TX) \otimes J^3(TX) \xrightarrow{f_1 \otimes f_1} J^1(J^2(TX)) \otimes J^1(J^2(TX)) \\ &\xrightarrow{f_2} J^1(J^2(TX) \otimes J^2(TX)) \xrightarrow{J^1(f_\delta)} J^1(\mathcal{O}_X) \xrightarrow{d} K_X \end{aligned} \quad (3.19)$$

(see (3.18) for $f_1 \otimes f_1$ and (3.11) for $J^1(f_\delta)$); note that the de Rham differential $d : \mathcal{O}_X \longrightarrow K_X$, being a first order differential operator, is given by an \mathcal{O}_X -linear homomorphism $J^1(\mathcal{O}_X) \longrightarrow K_X$. Let

$$H_2 : \text{Sym}^2(J^3(TX)) \longrightarrow K_X \quad (3.20)$$

be the homomorphism given by the composition of homomorphisms in (3.19).

Lemma 3.2. *The homomorphism H_1 in (3.17) coincides with the homomorphism H_2 in (3.20).*

Proof. In view of the explicit description of Δ_δ in (3.2), this is a rather straightforward computation. We will give an alternative proof of it using representations of $\mathrm{PSL}(2, \mathbb{C})$.

It suffices to prove that

- (1) $H_1 = H_2$ for $X = \mathbb{CP}^1$, and
- (2) the identification $H_1 = H_2$ on \mathbb{CP}^1 is equivariant for the action of $\mathrm{PSL}(2, \mathbb{C})$ on \mathbb{CP}^1 .

Indeed, in that case using coordinate charts on X , compatible with the projective structure δ , we may transport the identification $H_1 = H_2$ (for \mathbb{CP}^1) to open subsets of X ; the equivariance condition in (2) then ensures that these locally defined identifications over X patch together compatibly to give a global identification $H_1 = H_2$ over X .

Since the homomorphism $\eta_3 \circ (\eta_2)^{-1} : J^2(T\mathbb{CP}^1) \rightarrow J^3(T\mathbb{CP}^1)$ (see (3.4)) gives a $\mathrm{PSL}(2, \mathbb{C})$ -equivariant splitting of the short exact sequence in (3.5), we have

$$J^3(T\mathbb{CP}^1) = J^2(T\mathbb{CP}^1) \oplus K_{\mathbb{CP}^1}^{\otimes 2}.$$

This implies that

$$\mathrm{Sym}^2(J^3(T\mathbb{CP}^1)) = \mathrm{Sym}^2(J^2(T\mathbb{CP}^1)) \oplus K_{\mathbb{CP}^1}^{\otimes 4} \oplus (K_{\mathbb{CP}^1}^{\otimes 2} \otimes J^2(T\mathbb{CP}^1)), \quad (3.21)$$

and this isomorphism commutes with the actions of $\mathrm{PSL}(2, \mathbb{C})$. The vector bundle $\mathrm{Sym}^2(J^2(T\mathbb{CP}^1))$ is trivial, because $J^2(T\mathbb{CP}^1)$ is trivialized by η_2 in (3.4). Therefore,

$$H^0(\mathbb{CP}^1, \mathrm{Hom}(\mathrm{Sym}^2(J^2(T\mathbb{CP}^1)), K_{\mathbb{CP}^1})) = 0 \quad (3.22)$$

as the degree of $K_{\mathbb{CP}^1}$ is negative. Also, we have

$$H^0(\mathbb{CP}^1, \mathrm{Hom}(K_{\mathbb{CP}^1}^{\otimes 4}, K_{\mathbb{CP}^1})) = H^0(\mathbb{CP}^1, (T\mathbb{CP}^1)^{\otimes 3}).$$

This implies that

$$H^0(\mathbb{CP}^1, \mathrm{Hom}(K_{\mathbb{CP}^1}^{\otimes 4}, K_{\mathbb{CP}^1})) = \mathrm{Sym}^6(\mathbb{C}^2)$$

as a $\mathrm{PSL}(2, \mathbb{C})$ -module. Now taking invariants for the action of $\mathrm{PSL}(2, \mathbb{C})$,

$$H^0(\mathbb{CP}^1, \mathrm{Hom}(K_{\mathbb{CP}^1}^{\otimes 4}, K_{\mathbb{CP}^1}))^{\mathrm{PSL}(2, \mathbb{C})} = 0. \quad (3.23)$$

Next, we have

$$\begin{aligned} & H^0(\mathbb{CP}^1, \mathrm{Hom}(K_{\mathbb{CP}^1}^{\otimes 2} \otimes J^2(T\mathbb{CP}^1), K_{\mathbb{CP}^1})) \\ &= H^0(X, J^2(T\mathbb{CP}^1))^* \otimes H^0(X, \mathrm{Hom}(K_{\mathbb{CP}^1}^{\otimes 2}, K_{\mathbb{CP}^1})) \end{aligned}$$

because $J^2(T\mathbb{CP}^1)$ is the trivial bundle $\mathbb{CP}^1 \times H^0(X, J^2(T\mathbb{CP}^1))$ (trivialized by η_2 in (3.4)). Consequently, we have

$$\begin{aligned} & H^0(\mathbb{CP}^1, \mathrm{Hom}(K_{\mathbb{CP}^1}^{\otimes 2} \otimes J^2(T\mathbb{CP}^1), K_{\mathbb{CP}^1})) = H^0(X, J^2(T\mathbb{CP}^1))^* \otimes H^0(X, T\mathbb{CP}^1) \\ &= \mathfrak{sl}(2, \mathbb{C})^* \otimes \mathfrak{sl}(2, \mathbb{C}) = \mathfrak{sl}(2, \mathbb{C})^* \otimes \mathfrak{sl}(2, \mathbb{C})^* \end{aligned}$$

as a $\mathrm{PSL}(2, \mathbb{C})$ -module. Now taking invariants for the action of $\mathrm{PSL}(2, \mathbb{C})$,

$$H^0(\mathbb{CP}^1, \mathrm{Hom}(K_{\mathbb{CP}^1}^{\otimes 2} \otimes J^2(T\mathbb{CP}^1), K_{\mathbb{CP}^1}))^{\mathrm{PSL}(2, \mathbb{C})} = \mathbb{C}. \quad (3.24)$$

Note that $1 \in (\mathfrak{sl}(2, \mathbb{C})^* \otimes \mathfrak{sl}(2, \mathbb{C})^*)^{\mathrm{PSL}(2, \mathbb{C})}$ is given by the map $\mathfrak{sl}(2, \mathbb{C}) \otimes \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathbb{C}$ defined by $A \otimes B \rightarrow \mathrm{trace}(AB)$.

In view of (3.22), (3.23) and (3.24), from (3.21) we conclude that

$$H^0(\mathbb{C}\mathbb{P}^1, \text{Hom}(\text{Sym}^2(J^3(T\mathbb{C}\mathbb{P}^1)), K_X))^{\text{PSL}(2, \mathbb{C})} = \mathbb{C}.$$

In other words, up to multiplication by a scalar, there is a unique nonzero $\text{PSL}(2, \mathbb{C})$ -equivariant homomorphism from $\text{Sym}^2(J^3(T\mathbb{C}\mathbb{P}^1))$ to K_X . Hence the homomorphisms H_1 and H_2 coincide up to multiplication by a scalar when $X = \mathbb{C}\mathbb{P}^1$. It is now easy to see that $H_1 = H_2$ when $X = \mathbb{C}\mathbb{P}^1$. \square

Take two distinct point $a, b \in X$. Let $U_a := X \setminus \{b\}$ and $U_b := X \setminus \{a\}$ be the open subsets; define $U := U_a \cap U_b$. The 1-cocycles for \mathcal{B}_\bullet consists of

$$(\theta, \omega_a, \omega_b) \in H^0(U, TU) \oplus H^0(U_a, K_{U_a}^{\otimes 2}) \oplus H^0(U_b, K_{U_b}^{\otimes 2})$$

such that $\omega_a - \omega_b = \Delta_\delta(\theta)$, where Δ_δ is the differential operator in (3.1). A cocycle $(\theta, \omega_a, \omega_b)$ is a 1-coboundary if there is an element

$$(\theta_a, \theta_b) \in H^0(U_b, TU_a) \oplus H^0(U_b, TU_b)$$

such that $\theta = \theta_a - \theta_b$, $\omega_a = \Delta_\delta(\theta_a)$ and $\omega_b = \Delta_\delta(\theta_b)$. The hypercohomology $\mathbb{H}^1(\mathcal{B}_\bullet)$ (see (3.12)) is identified with the quotient of 1-cocycles by the 1-coboundaries.

The residue of any $\alpha \in H^0(U, K_U)$ at the point $a \in X$ will be denoted by $\text{Res}_a(\alpha)$.

Take two 1-cocycles $\mathbf{c}^1 := (\theta^1, \omega_a^1, \omega_b^1)$ and $\mathbf{c}^2 := (\theta^2, \omega_a^2, \omega_b^2)$ for \mathcal{B}_\bullet in (3.12). So we have

$$\theta^1 \cdot \omega_b^2, \theta^2 \cdot \omega_a^1 \in H^0(U, K_U).$$

Define

$$\widehat{\Theta}'(\mathbf{c}^1, \mathbf{c}^2) := \text{Res}_a(\theta^1 \cdot \omega_b^2 + \theta^2 \cdot \omega_a^1) \in \mathbb{C}, \quad (3.25)$$

where “ \cdot ” is the natural pairing $TX \otimes K_X^{\otimes 2} \rightarrow K_X$.

Theorem 3.3. *If \mathbf{c}^1 in (3.25) is a 1-coboundary, then $\widehat{\Theta}'(\mathbf{c}^1, \mathbf{c}^2) = 0$.*

Proof. Since $\mathbf{c}^1 := (\theta^1, \omega_a^1, \omega_b^1)$ is a 1-coboundary, there is

$$(\theta_a, \theta_b) \in H^0(U_b, TU_a) \oplus H^0(U_b, TU_b)$$

such that $\theta^1 = \theta_a - \theta_b$, $\omega_a^1 = \Delta_\delta(\theta_a)$ and $\omega_b^1 = \Delta_\delta(\theta_b)$. So $\theta^1 \cdot \omega_b^2 + \theta^2 \cdot \omega_a^1$ in (3.25) has the expression

$$\begin{aligned} \theta^1 \cdot \omega_b^2 + \theta^2 \cdot \omega_a^1 &= (\theta_a - \theta_b) \cdot \omega_b^2 + \theta^2 \cdot \Delta_\delta(\theta_a) = \theta_a \cdot \omega_b^2 + \theta^2 \cdot \Delta_\delta(\theta_a) - \theta_b \cdot \omega_b^2 \\ &= \theta_a \cdot (\omega_a^2 + \Delta_\delta(\theta^2)) + \theta^2 \cdot \Delta_\delta(\theta_a) - \theta_b \cdot \omega_b^2 = \theta_a \cdot \omega_a^2 - \theta_b \cdot \omega_b^2 + H_1(\theta_a, \theta^2), \end{aligned} \quad (3.26)$$

where H_1 is the homomorphism in (3.17). Since the residues of $\theta_a \cdot \omega_a^2$ and $\theta_b \cdot \omega_b^2$ vanish at the point a (and at b),

$$\widehat{\Theta}'(\mathbf{c}^1, \mathbf{c}^2) := \text{Res}_a(H_1(\theta_a, \theta^2)) \quad (3.27)$$

(see (3.25) and (3.26)). From Lemma 3.2,

$$\text{Res}_a(H_1(\theta_a, \theta^2)) = \text{Res}_a(H_2(\theta_a, \theta^2)); \quad (3.28)$$

on the other hand, $H_2(\theta_a, \theta^2)$ is exact on $U := U_a \cap U_b$ (the final homomorphism in (3.19) is the de Rham differential). Hence we have

$$\text{Res}_a(H_2(\theta_a, \theta^2)) = 0.$$

Now from (3.27) we know that $\widehat{\Theta}'(\mathbf{c}^1, \mathbf{c}^2) = 0$. □

Theorem 3.3 has the following immediate consequence.

Corollary 3.4. *The pairing $\widehat{\Theta}'$ in (3.25) produces an alternating form*

$$\widehat{\Theta}(\delta) : \bigwedge^2 \mathbb{H}^1(\mathcal{B}_\bullet) \longrightarrow \mathbb{C}.$$

Note that $\widehat{\Theta}$ in Corollary 3.4 produces a 2-form on the moduli space \mathcal{P}_g in (2.8). From its construction it is evident that $\widehat{\Theta}$ is an algebraic form on \mathcal{P}_g .

From the construction of the 2-form $\widehat{\Theta}$ on \mathcal{P}_g it is straightforward to deduce that it coincides with the symplectic form Θ_g in (2.12).

REFERENCES

- [AB] M. F. Atiyah and R. Bott, The Yang-Mills equations over Riemann surfaces, *Philos. Trans. Roy. Soc. London* **308** (1983), 523–615.
- [Ba] S. Baba, A Schottky decomposition theorem for complex projective structures, *Geom. Topol.* **14** (2010), 117–151.
- [BL] I. Biswas and E. Looijenga, Projective structures on Riemann surfaces, *Pure Appl. Math. Quart.* (to appear).
- [GKM] D. Gallo, M. Kapovich and A. Marden, The monodromy groups of Schwarzian equations on closed Riemann surfaces, *Ann. of Math.* **151** (2000) 625–704.
- [Go] W. M. Goldman, The symplectic nature of fundamental groups of surfaces, *Adv. in Math.* **54** (1984), 200–225.
- [Gu] R. C. Gunning, *Lectures on Riemann Surfaces*, Mathematical Notes 2, Princeton University Press, Princeton, NJ, 1966.
- [He] D. A. Hejhal, Monodromy groups and linearly polymorphic functions, *Acta Math.* **135** (1975), 1–55.
- [Hu] J. H. Hubbard, The monodromy of projective structures, *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, pp. 257–275, Ann. of Math. Stud., 97, Princeton Univ. Press, Princeton, N.J., 1981.

MATHEMATICS DEPARTMENT, SHIV NADAR UNIVERSITY, NH91, TEHSIL DADRI, GREATER NOIDA,
UTTAR PRADESH 201314, INDIA

Email address: indranil.biswas@snu.edu.in, indranil29@gmail.com