A Study of the Long-Term Behavior of Hybrid Systems with Symmetries via Reduction and the Frobenius-Perron Operator*

Maria Oprea[†], Aden Shaw[‡], Robi Huq[§], Kaito Iwasaki[¶], Dora Kassabova[∥], and William Clark[#]

Abstract. Hybrid dynamical systems are systems which undergo both continuous and discrete transitions. As typical in dynamical analysis, an essential goal is to study the long-term behavior of these systems. In this work, we present two different novel approaches for studying these systems. The first approach is based on constructing an analog of the Frobenius-Perron (transport) operator for hybrid systems. Rather than tracking the evolution of a single trajectory, this operator encodes the asymptotic nature of an ensemble of trajectories. The second approach presented applies to an important subclass of hybrid systems, mechanical impact systems. We develop an analog of Lie-Poisson(-Suslov) reduction for left-invariant impact systems on Lie groups. In addition to the Hamiltonian (and constraints) being left-invariant, the impact surface must also be a right coset of a normal subgroup. This procedure allows a reduction from a 2n-dimensional system to an (n + 1)-dimensional one. We conclude the paper by presenting numerical results on a diverse array of applications.

Key words. Hybrid Systems, Hybrid Reduction, Transfer Operator

MSC codes. 34A38, 34D05, 37C30, 37C83, 70F25

1. Introduction. A characteristic pursuit in dynamical systems is to study and understand their long-term behavior. One approach to understand a system's asymptotic properties is through a probabilistic lens. If the dynamics preserve a probability measure, then the celebrated Poincaré recurrence and ergodic theorems can be applied. These results are well-studied in the cases where the evolution is either discrete (an iterated map) or continuous (a flow induced by a differential equation) [20]; however, many real world phenomena are described by systems that fail to be exclusively discrete nor continuous [15]. Consider the pedagogical example of a bouncing ball: as it flies through the air, its motion is continuous (ballistic motion is described by a differential equation), but when it hits the ground there is an instantaneous change in momentum (by applying the so-called impact map), see Figure 1. We call such systems *hybrid systems*—dynamical systems whose evolution are subject to *both* continuous and discrete laws [5].

Although hybrid systems are ubiquitous in real life applications, their rigorous study is made difficult due to combination of continuous and discrete dynamics, and is consequently less developed in the literature. As such, the overarching goal of this work is to extend preexisting theory to hybrid systems to (a) study such systems through a probabilistic lens and (b) leverage

Funding: This work was funded by the NSF grant DMS-1645643 and AFOSR Award No. MURI FA9550-32-1-0400.

[†]Center of Applied Mathematics, Cornell University, Ithaca, NY (mao237@cornell.edu).

[‡]Department of Mathematics, Rose-Hulman Institute of Technology, Terre Haute, IN (shawap@rose-hulman.edu).

[§]Department of Mathematics, University of Minnesota Twin Cities, Minneapolis, MN (huq00011@umn.edu).

[¶]Department of Mathematics, University of Michigan, Ann Arbor, MI (kaitoi@umich.edu).

^{||}Department of Mathematics, University of Washington, Seattle, WA (dmk285@uw.edu).

[#]Department of Mathematics, Ohio University, Athens, OH (clarkw3@ohio.edu)

the geometry of such systems in order to reduce the difficulty/dimension of the problem.



(a) Bouncing ball in $T^*\mathbb{R}$ (b) Bouncing Ball with coefficient of elasticity of c = 0.5.

Figure 1: Plots of the motion of the bouncing ball. Left: Phase plot on $T^*\mathbb{R}$ with the momentum and position of the ball. Right: Trajectory of the ball, which tracks the height x(t) of the ball. The guard is given by $S = \{(x, p) \in T^*\mathbb{R} \cong \mathbb{R}^2 : x = 0, p < 0\}.$

Consider the continuous-time dynamical system $\dot{x} = X(x)$, where X is a vector-field on the ambient manifold M. Let S be a distinguished codimension one (embedded) submanifold

called the *guard*; this encodes location of the discrete transitions. Finally, we introduce the *reset* map, $\Delta \colon S \to M$, which dictates the discrete transition. Such a system will be called *hybrid* and its dynamics will be governed by the following rule

(1.1)
$$\begin{cases} \dot{x} = X(x), & x \notin \mathcal{S}, \\ x^+ = \Delta(x^-), & x \in \mathcal{S}. \end{cases}$$

This system exhibits continuous behavior away from the guard and discrete when the state enters the guard. If the guard is in its image, i.e., $\Delta(S) \cap S \neq \emptyset$, then multiple resets can immediately occur—a phenomenon called *beating*. To



Figure 2: Schematic drawing of a hybrid system.

avoid this, we will assume that the guard is disjoint from its image under the reset map. More technical details for the hybrid system governed by (1.1) will be discussed in Section 2.1.

The Frobenius-Perron and Koopman operators are adjoint linear transfer operators that describe the evolution of scalar observables. Linear operator theory provides a framework for studying non-linear dynamical systems by obtaining infinite-dimensional linear representations of the system, which can be used for the spectral analysis of nonlinear dynamical systems [6, 27]. This allows for an elegant study of the asymptotic behaviour by looking at successive iterates of the Koopman and Frobenius-Perron operators, and their proprieties in the limit of infinitely many iterations.

Despite the fact that Koopman's original ideas were published nearly a century ago, work on the subject was not popularized until fairly recently, due to a lack of efficient algorithms and computing power capable of calculating the Koopman operator. This theory is well-defined for both discrete- and continuous-time systems as right composition. However, current methods have shown that even when taking ino advantage the linearity of the Koopman operator, there are limitations when handling higher-dimensional systems [31]. This leads us into our second approach—geometric dimension reduction.

The theory of reduction plays an important role in classical mechanics. Lie-Poisson reduction concerns the case when the configuration space is a Lie group, G, and reduces the dynamics from the symplectic manifold T^*G to the Poisson manifold \mathfrak{g}^* effictively decreasing the dimension of the system from 2n to n [26, Chapter 13]. Discrete analogs have been proposed [25], where the Lie group is finite dimensional, and the Lagrangian is discrete. Hybrid systems pose a challenge due to the fact that the equations of motion come from a continuous Hamiltonian, whereas the discrete impacts are provided by an arbitrary map which might not come from a discrete Lagrangian as in [25]. Therefore, neither the continuous nor the discrete reduction can be directly applied.

It is possible to leverage the symmetries of the continuous part of a hybrid system, but in order to do this, it is necessary to document the locations and effects of the discrete impacts. This is why, for hybrid systems, reduction from 2n to n is not achievable. In this paper we propose a method (see 3.22) that allows reduction to n + 1 dimensions, where the 'extra' dimension is used to record the locations of the discrete impacts on the guard.

The goal of this work is to extend and combine these theories such that they can be used on more complex hybrid systems. For instance, the state space of a robotic device is often high dimensional, especially when one has to take into account a large number of components that work together to fulfill the functionality of the device. By reducing the dimension and studying the long-term behaviour of this system one could understand robustness and stability.

This manuscript is organized as follows: Section 2 contains the necessary preliminaries of hybrid systems (2.1), the classical Frobenius-Perron and Koopman operators (2.2), and reduction (2.3). Section 3 contains our results, specifically the hybrid Frobenius-Perron operator (3.1) and hybrid Lie-Poisson reduction (3.2). Section 4 contains numerical results for four examples: the bouncing ball (4.1), the Chaplygin sleigh (4.2), matrix groups (4.3), and an SIR type disease model (4.4). Conclusions and future work are presented in Section 5. Detailed proofs for technical lemmas can be found in the Appendix.

2. Preliminaries. This section is devoted to defining common terminology related to our results and is broken up into three subsections: hybrid systems, Frobenius-Perron and Koopman operators, and reduction. An understanding of hybrid systems is necessary for understanding both of the main results, so hybrid systems are covered first, and each of the remaining subsections correspond to a main result.

2.1. Hybrid Dynamical Systems. The purpose of this section is to define the types of systems we are working with. First, we will focus on hybrid dynamical systems, which are continuous dynamical systems which exhibit occasional discrete transitions. The state space will always be a smooth manifold M. The continuous dynamics will be given in terms of a vector field, and the jumps happen when the trajectory of the system hits the guard denoted by \mathcal{S} [18]. Following hybrid systems, we will refine our discussion to impact systems which are a subset of hybrid systems.

Definition 2.1 (Hybrid dynamical system). A hybrid dynamical system \mathcal{H} , abbreviated by HDS, is 4-tuple $\mathcal{H} = (M, \mathcal{S}, \Delta, X)$ with the following properties

- (H.1) M is a finite-dimensional smooth manifold,
- (H.2) $\mathcal{S} \subset M$ is an embedded smooth manifold where codim $\mathcal{S} = 1$,
- (H.3) $\Delta: \mathcal{S} \to M$ is a smooth map whose image is an embedded submanifold,
- (H.4) $X: M \to TM$ is a smooth vector field, and
- (H.5) $\mathcal{S} \cap \Delta(\mathcal{S}) = \emptyset$ and $\operatorname{codim}(\overline{\mathcal{S}} \cap \Delta(\mathcal{S})) \ge 2$.

s

As stated in the introduction, the submanifold S is called the quard and the map Δ is the reset.

Away from the guard the trajectories of a hybrid dynamical system are integral curves of the vector field X, whereas, on the guard the system is equivalent to a discrete dynamical system with map Δ . Putting the continuous and the discrete parts of the trajectory together generates the hybrid flow.

Definition 2.2 (Hybrid Flow). Let $\mathcal{H} = (M, \mathcal{S}, \Delta, X)$ be an HDS. The map $\varphi^{\mathcal{H}} : \operatorname{dom}(\varphi^{\mathcal{H}}) \subset$ $\mathbb{R} \times M \to M$ is the hybrid flow if

$$\frac{d}{dt}\varphi^{\mathcal{H}}(t,x) = X_{\varphi^{\mathcal{H}}(t,x)}, \qquad \varphi^{\mathcal{H}}(t,x) \notin \mathcal{S};$$
$$\lim_{s \to t^{+}} \varphi^{\mathcal{H}}(s,x) = \Delta\left(\lim_{s \to t^{-}} \varphi^{\mathcal{H}}(s,x)\right), \quad \varphi^{\mathcal{H}}(t,x) \in \mathcal{S}.$$

Hybrid systems can fail to be complete in two qualitatively different ways: finite time blow ups of the continuous trajectory, and the Zeno phenomenon. The latter is special to hybrid systems, and it happens when a trajectory undergoes infinitely many jumps in a finite amount of time.

Zeno trajectories pose a difficult challenge. In order to compute the hybrid flow as time approaches Zeno one needs to be able to perform infinitely accurate event detection. From a theoretical perspective, what happens after the time becomes greater than t_{∞} is still unclear [2]. It is generally difficult to determine whether or not a state will have a Zeno trajectory or not. However, for reasonable HDSs which possess an invariant volume-form, the set of points which are Zeno form a null set [9]. As volume-forms are smooth objects, the HDS must satisfy some amount of smoothness conditions.

Definition 2.3 (Quasi-smooth dependence property).

Let $\mathcal{H} = (M, \mathcal{S}, \Delta, X)$ be a HDS with flow $\varphi^{\mathcal{H}}$. \mathcal{H} has the quasi-smooth dependence property if for every $x \in M \setminus \mathcal{S}$ and $t \in \mathbb{R}$ such that $\varphi^{\mathcal{H}}(t, x) \notin \mathcal{S}$, there exists an open neighborhood $x \in U$ such that $U \cap \mathcal{S} = \emptyset$ and the map $\varphi^{\mathcal{H}}(t, \cdot) : U \to M$ is smooth. Furthermore, if \mathcal{H} is a HDS with the quasi-smooth dependence property, then \mathcal{H} is a smooth HDS.

In this work we will assume that all the systems have the quasi-smooth dependence propriety (as well as completeness, unless otherwise stated); this is not a strong assumption cf. [8]. This assumption allows for a meaningful study of how volumes preserve across applications of the reset map, which is crucial when defining the Frobenius-Perron operator. For a volume form $\mu \in \Omega^n(M)$ to be be invariant under the hybrid flow within a smooth HDS, μ must be invariant under both the continuous-time dynamics and the reset. Invariance under the continuous dynamics follows when $\mathcal{L}_X \mu = 0$ (equivalently, $\operatorname{div}_{\mu}(X) = 0$), where \mathcal{L} denotes the Lie derivative. Invariance across resets occurs when

$$\Delta^* i_X \mu = \iota_{\mathcal{S}}^* i_X \mu,$$

where $\iota_{\mathcal{S}} \colon \mathcal{S} \hookrightarrow M$ is the inclusion and i_X is the interior product [9]. This prompts the following definition.

Definition 2.4 (Hybrid Jacobian of Δ). Let $\mathcal{H} = (M, \mathcal{S}, \Delta, X)$ be a smooth HDS and $\mu \in \Omega^n(M)$ a volume-form. The unique function $\mathcal{J}^X_{\mu}(\Delta) \in C^{\infty}(\mathcal{S})$ such that

(2.1)
$$\Delta^* i_X \mu = \mathcal{J}^X_\mu(\Delta) \cdot \iota^*_{\mathcal{S}} i_X \mu,$$

is called the hybrid Jacobian of Δ .

Proposition 2.5 (Theorem 9 in [9]). Let $\mathcal{H} = (M, \mathcal{S}, \Delta, X)$ be a smooth HDS and $\mu \in \Omega^n(M)$ a volume-form. Then μ is invariant under the flow of \mathcal{H} if $\mathcal{L}_X \mu = 0$ and its hybrid Jacobian $\mathcal{J}^X_\mu(\Delta)$ is 1.

Example 1 (Filippov Systems). Consider the plane $M = \mathbb{R}^2$, with the guard given by

$$\mathcal{S} = \left\{ (x, y) \in \mathbb{R}^2 : xy = 0 \right\},\$$

and the identity reset map of $\Delta(x, y) = \mathrm{Id}_{\mathbb{R}^2}(x, y) = (x, y)$. Furthermore, consider the volume form $\mu = dx \wedge dy$ and the following discontinuous vector field.

$$X_{(x,y)} = \begin{cases} \left(\frac{y}{\alpha} - x\right) \frac{\partial}{\partial x} - (\alpha x + y) \frac{\partial}{\partial y} & \text{for } xy > 0;\\ (\alpha y - x) \frac{\partial}{\partial x} - \left(y + \frac{x}{\alpha}\right) \frac{\partial}{\partial y} & \text{for } xy < 0. \end{cases}$$

To compute the hybrid Jacobian, we notice that Δ is the identity so we must compute $\iota_{\mathcal{S}}^* i_X \mu$ on both sides of the guard

$$i_X dx \wedge dy = \begin{cases} \left(\frac{y}{\alpha} - x\right) dy + (\alpha x + y) dx & \text{for } xy > 0; \\ (\alpha y - x) dy + \left(y + \frac{x}{a}\right) dx & \text{for } xy < 0. \end{cases}$$

Substituting either x = 0 or y = 0 shows that the hybrid Jacobian is

$$\mathcal{J}^X_\mu(\mathrm{Id}_{\mathbb{R}^2}) = \alpha^2.$$

As a consequence of Theorem 3.12 below, an invariant density will exist when $\alpha = \exp(\pi)$. In

this case, if (x, y) lies within the first quadrant, then an invariant density is given by

$$\rho(x,y) = e^{-2\tau}, \quad \tan \tau = \frac{\alpha x}{y}.$$

In the case that the points do not lie in the first quadrant, ρ can be extended through the following map

$$\tilde{\rho}(x,y) = \begin{cases} \rho(x,y), & x > 0, \, y > 0; \\ \rho(-y,x), & x > 0, \, y < 0; \\ \rho(-x,-y), & x < 0, \, y < 0; \\ \rho(y,-x), & x < 0, \, y > 0, \end{cases}$$



Figure 3: This is the invariant density for this example. The jump discontinuity between quadrants is precisely the hybrid Jacobian: $\exp(\pi)$.

which is displayed in Figure 3.

As will be seen in Section 3.1, the hybrid Jacobian will be central in the study of the hybrid Frobenius-Perron operator.

2.1.1. Mechanical Impact Systems. A physically motivated class of hybrid systems are mechanical impact systems. In these systems, the continuous dynamics obey Hamiltonian/Lagrangian mechanics while the reset map encodes the momentum/velocity jump at a physical impact. Throughout this work, we will be primarily in the Hamiltonian setting, but the results can be translated to the Lagrangian framework with little difficulty.

Let Q be an n dimensional smooth manifold representing the configuration space of the system of interest. To each point $q \in Q$, the possible momenta will be elements of the cotangent space $p \in T_q^*Q$. The union of all possible momenta attached to all possible points is called the cotangent bundle or the phase space, and is denoted by T^*Q . In a similar fashion, the possible velocities will be tangent vectors, $v \in T_qQ$, and the collection of all positions and velocities will be the tangent bundle TQ.

The cotangent bundle is equipped with the canonical symplectic form which we call $\omega \in \Omega^2(T^*Q)$, and in local coordinates is given by $\omega = dq^i \wedge dp_i$. For a given Hamiltonian function $H: T^*Q \to \mathbb{R}$, Hamilton's equations of motion are described by

where i_X denotes the interior product, $i_X \omega(Y) = \omega(X, Y)$ for any vector field Y, dH is the differential of H, and X is the resulting Hamiltonian vector field. There exist various levels of regularity for Hamilton functions; throughout this work, we will assume that all Hamiltonians are of mechanical type.

Definition 2.6 (Natural Hamiltonians). A Hamiltonian $H: T^*Q \to \mathbb{R}$ is called natural/mechanical if there exists a Riemannian metric \coprod on Q and a function $V: Q \to \mathbb{R}$ such that

$$H(q,p) = \frac{1}{2}(\mathbf{U}^{-1})_q(p,p) + V(q),$$

where \coprod^{-1} is the induced co-metric.

For a natural Hamiltonian we can explicitly write the write the Hamiltonian vector field from (2.2) and obtain the following coordinate representation of Hamilton's equations of motion:

(2.3)
$$\dot{q}^{k} = \frac{\partial H}{\partial p_{k}} = (\Pi^{-1})^{kj} p_{j};$$
$$\dot{p}_{k} = -\frac{\partial H}{\partial q^{k}} = -\frac{1}{2} \frac{\partial (\Pi^{-1})^{ij}}{\partial q^{k}} p_{i} p_{j} - \frac{\partial V}{\partial q^{k}}.$$

The metric underlying a mechanical Hamiltonian has the additional feature of being able to isomorphically transform vectors to covectors.

Definition 2.7 (Musical Isomorphisms). Given a Riemannian manifold (Q, \amalg) the musical isomorphisms are the maps

$$b: \mathfrak{X}(Q) \to \Omega^{1}(Q) : X \mapsto \amalg(X, \cdot), \\ \sharp: \Omega^{1}(Q) \to \mathfrak{X}(Q) : \alpha \mapsto \amalg^{-1}(\alpha, \cdot),$$

which are defined so that $\sharp = \flat^{-1}$. In coordinates the musical isomorphisms can be written as

 $(X^{\flat})_i = \coprod_{ij} X^j, \quad and \quad (\alpha^{\sharp})^i = (\amalg^{-1})^{ij} \alpha_j.$

Remark 2.8. If the Hamiltonian is natural then one can isomorphically switch between the Hamiltonian and Lagrangian formalism using the fiber derivative $\mathbb{F}L: TQ \to T^*Q$, where $L: TQ \to \mathbb{R}$ is the Lagrangian, and the inverse fiber derivative $\mathbb{F}H: T^*Q \to TQ$. In the mechanical case, the fiber derivatives agree with the musical isomorphisms.

Impacts induce instantaneous changes in the momenta in T^*Q and are triggered by the positions in Q. We assume that the impact surface Σ is a smooth codimension 1 submanifold of the configuration space Q, which can locally be written as a regular level set of a smooth function $s: Q \to \mathbb{R}$. Away from Σ , the dynamics are integral curves of the Hamiltonian vector field generated by H. We put these pieces together in the following definition:

Definition 2.9 (Impact system). An impact system is a tuple $\mathcal{I} = (Q, H, \Sigma)$ such that

- (I.1) Q is a finite-dimensional smooth manifold,
- (I.2) $H: T^*Q \to \mathbb{R}$ is a natural Hamiltonian,
- (I.3) $\Sigma \subset Q$ is an orientable embedded smooth manifold where $\operatorname{codim} \Sigma = 1$.

The three pieces of information in Definition 2.9 are enough to completely specify a hybrid dynamical system $(M, X, \mathcal{S}, \widetilde{\Delta})$ through the following process.

- (h.1) $M = T^*Q$ is the phase space containing positions and momenta (q, p).
- (h.4) X is given by X_H , the Hamiltonian vector field generated by H through equation (2.2).

(h.2) The guard is the set of all outward pointing momenta attached to positions in Σ , so

$$S = \{(q, p) \in T^*Q|_{\Sigma} : \pi_Q^* ds(X_H) > 0\},\$$

where $\pi_Q: T^*Q \to Q$ is the canonical projection.

(h.3) The impact map is identity in the first n components $\Delta(q, p) = (q, \Delta(p))$. In the last n components, $\Delta(p)$ can be computed from the Weierstrass-Erdmann corner conditions [14, Section 15]. Let p^+ and p^- denote be the values of the momenta before and after the impact, respectively, with H^+ and H^- their corresponding values of the Hamiltonian. Then the corner conditions are given by

(2.4)
$$H^- = H^+ \quad \text{and} \quad p^+ - p^- = \varepsilon \, ds.$$

Remark 2.10. Throughout this paper, the impact surface $\Sigma \subset Q$ will be independent of time, but in general, the local defining function s is an element of $C^{\infty}(\mathbb{R} \times Q)$, which locally defines $\Sigma_t \subset \mathbb{R} \times Q$. Then, the left side of (2.4) becomes $H^- - H^+ = \varepsilon \partial s / \partial t$, which can be physically intuited as the impact surface adding to/taking away from the energy of the system.

We can rephrase Equation (2.4) in a coordinate-free manner with

(2.5)
$$(\mathrm{Id} \times \Delta)^* \vartheta_H = \iota^* \vartheta_H \quad \text{and} \quad \vartheta_H = p_i dx^i - H dt$$

where $\iota: \mathbb{R} \times Q \hookrightarrow \mathbb{R} \times T^*Q$ is the inclusion map and ϑ_H is called the action form. Note that (2.5) works for both time-dependent and time-independent impact surfaces, but will be used only for the latter.

In the case where the Hamiltonian in (2.4) is natural, there exists a unique solution [9, 10] given by

(2.6)
$$\tilde{\Delta}(q,p) = (q, R(q)p), \quad \text{where} \quad R(q)p = p - 2\frac{\mathrm{II}^{-1}(ds,p)}{\mathrm{II}^{-1}(ds,ds)}ds.$$

2.1.2. Constraints. Constraints are ubiquitous in mechanical systems and dictate either allowable positions or velocities. There are two distinguished classes of constraints: integral or holonomic and non-integrable/nonholonomic. While holonomic systems maintain their Hamiltonian structure, nonholonomic systems are *not* Hamiltonian as they do not come from variational principles. Fortunately, holonomic systems can be viewed as a special case of nonholonomic systems and the theory for the latter still describes the former. As a result, we will treat all constraints as nonholonomic and all constraints will be *linear* in the velocities. For a more in-depth treatment of nonholonomic mechanics see [4, Chapter 2].

To obtain the nonholonomic equations of motion, one must use the Lagrange-d'Alembert principle [26, Chapter 7]. In the Hamiltonian formalism these give raise to the constraint Hamiltonian vector field.

Definition 2.11 (Constrained Hamiltonian vector field). Let (Q, ω, H) be a natural Hamiltonian system and $\mathcal{D} \subset TQ$ be a regular distribution. Let $\eta^{\alpha} \in \Omega^{1}(Q)$ be a collection of k 1-forms which locally describe \mathcal{D} via annihilation. Then the nonholonomic Hamiltonian vector field $X_{H}^{\mathcal{D}}$, is the unique vector field such that

$$i_{X_H^{\mathcal{D}}}\omega|_{\mathcal{D}^*} = dH|_{\mathcal{D}^*} + \lambda_\alpha \pi_Q^* \eta^\alpha|_{\mathcal{D}^*},$$

where λ_{α} are multipliers to enforce the constraints,

$$\eta^{\alpha} \big(\mathbb{F}H(q, p) \big) = 0, \quad 1 \le \alpha \le k,$$

and $\mathbb{F}H(\mathcal{D}^*) = \mathcal{D}$ is the induced (co)distribution on T^*Q , i.e., $\mathcal{D}^* = \{(q, p) \in T^*Q : p = g(\dot{q}, \cdot) \text{ for some } (q, \dot{q}) \in \mathcal{D}\}.$

Example 2 (The Chaplygin sleigh). The Chaplygin sleigh is a pedagogical example of a (nonholonomically) constrained system. The configuration space of the Chaplygin sleigh is SE_2 , which tracks the x and y position of the sleigh along with the direction θ its facing. The Lagrangian of this system is

$$L = \frac{1}{2} \left(m(\dot{x}^2 + \dot{y}^2) + (I + ma^2)\dot{\theta}^2 - 2ma\dot{\theta}\left(\dot{x}\sin\theta - \dot{y}\cos\theta\right) \right)$$

where m is the mass of the sleigh, a is the distance between the center of mass and the front of the sleigh, I is the moment of inertia about the center of mass. To disallow for sliding in transverse directions, the sleigh is given the knife edge constraint,

(2.7)
$$\dot{x}\sin\theta - \dot{y}\cos\theta = 0$$
, or $\eta = \sin(\theta) \, dx - \cos(\theta) \, dy$.

Given the constraint (2.7), the corresponding distribution is

$$\mathcal{D} = \left\{ (x, y, \theta, \dot{x}, \dot{y}, \dot{\theta}) \in TQ : \dot{x} \sin \theta - \dot{y} \cos \theta = 0 \right\},\$$

which shows the relation between the allowable configurations and the constraints. More information can be found within [4, Chapter 1.7].

We now combine the constraint continuous dynamics with the discrete dynamics given by impacts. The object that stores all the information about the constrained dynamics and the discrete transitions is called a constrained impact system, and is defined below.

Definition 2.12 (Constrained impact system). A constrained impact system is a 4-tuple $\mathcal{I} = (Q, H, \Sigma, \mathcal{D})$ such that

- (I.1) Q is a finite-dimensional smooth manifold,
- (I.2) $H: T^*Q \to \mathbb{R}$ is a natural Hamiltonian,
- (I.3) $\mathcal{D} \subset TQ$ is a regular distribution,
- (I.4) $\Sigma \subset M$ is an embedded smooth manifold where $\operatorname{codim} \Sigma = 1$.

Example 2 (Continued). In addition to the to the previous properties of the Chaplygin sleigh, we now define an impact surface for the system

$$s: SE_2 \to \mathbb{R} : (x, y, \theta) \mapsto (\theta - \theta_0)(\theta + \theta_0).$$

Given this map, $\Sigma = s^{-1}(\{0\}) = \{(x, y, \theta) \in SE_2 : (\theta - \theta_0)(\theta + \theta_0) = 0\}$. With the impact surface, the Chaplygin sleigh becomes a constrained impact system in contrast to previously being a constrained mechanical system.

Next, the corresponding reset map for this system is given by $\Delta(v, \omega, \theta) = (v, -\omega, \theta)$, which represents the sleigh automatically changing direction when the critical angle θ_0 is reached. This example will be further continued in Section 4.2.

2.2. The Frobenius-Perron and Koopman Operators. When studying the long time behaviour of dynamical systems through the probabilistic lens, one is interested in the evolution of densities instead of single points. Consider a large number of initial points $x_1, \ldots, x_n \in X$, sampled randomly from some probability density function f_0 , and assume that the dynamics are given by a 1-parameter group action $\phi_t \colon X \to X$. Each of these points are going to move around under the dynamics to points $x_1(t), \ldots, x_n(t)$ after some time t. If n is large and $t \to \infty$, computing, for each individual trajectory, its value at time t becomes very expensive. Now, assume that there exists some other probability density f_t such that $x_1(t), \ldots, x_n(t)$ are roughly obtained by sampling n points from this distribution. Then in order to analyze the long term behavior of the system it is sufficient to look at how f_t evolves over time, and in particular, at $\lim_{t\to\infty} f_t$. The Frobenius-Perron operator dictates the evolution of f_t .

In the following, we provide the formal definitions of the Frobenius-Perron operator for discrete and continuous time systems as well as its infinitesimal generators [21].

Definition 2.13 (Discrete Frobenius-Perron Operator). Let $\varphi \colon M \to M$ be a nonsingular measurable transformation on a measure space (M, \mathcal{A}, μ) . The (discrete) Frobenius-Perron operator corresponding to φ is the unique linear operator $P \colon L^1(M, \mathcal{A}, \mu) \to L^1(M, \mathcal{A}, \mu)$ defined by

(2.8)
$$\int_E Pf(x) \, d\mu = \int_{\varphi^{-1}(E)} f(x) \, d\mu, \quad \text{for all } E \in \mathcal{A}$$

Remark 2.14 (Measure-theoretic considerations). Throughout this paper, all sigma algebras are assumed to be the Borel algebra generated by the topology on the manifold (Q or M, depending on the context), so measure spaces will be denoted by doubles (M, μ) rather than triples (M, \mathcal{A}, μ) . Furthermore, all sets are assumed to be measurable unless stated otherwise. In addition, we interchangeably use our volume-forms as measures. Formally, $\mu(E) \coloneqq \int_E \mu$ for $\mu \in \Omega^n(M)$ and a measurable set $E \subseteq M$.

Given an initial probability distribution $f \in L^1$, the Frobenius-Perron operator gives an evolution of f by the transformation φ . Furthermore, for any $f \in L^1$, if φ is an *invertible* non-singular transformation, then

$$Pf(x) = f(\varphi^{-1}(x))\mathcal{J}^{-1}(x),$$

where \mathcal{J}^{-1} is the determinant of the inverse of the Jacobian matrix. If φ fails to be invertible,

(2.9)
$$Pf(x) = \sum_{y \in \varphi^{-1}(\{x\})} f(y) \mathcal{J}^{-1}(y)$$

The discrete Frobenius-Perron operator has a natural continuous-time system analog.

Definition 2.15 (Continuous Frobenius-Perron Operator). Let $\varphi: M \times \mathbb{R} \to M$ be a flow and denote $\varphi_t(x) = \varphi(x, t)$. Suppose that the flow is nonsingular and measurable on the measure space (M, μ) . For $E \subseteq M$, the (continuous) Frobenius-Perron operator $P_t: L^1(M, \mu) \to L^1(M, \mu)$ for each $t \in \mathbb{R}$ corresponding to φ_t is defined by

$$\int_{E} P_t f(x) \, d\mu = \int_{\varphi_{-t}(E)} f(x) \, d\mu \qquad \text{for all } E \subseteq M$$

Remark 2.16. The Frobenius-Perron and Koopman operators are adjoints of one another. Letting $K_t: L^{\infty}(M,\mu) \to L^{\infty}(M,\mu)$ be the Koopman operator, which is defined as

$$K_t g(x) = g(\varphi_t(x))$$

and $\langle \cdot, \cdot \rangle \colon L^1 \times L^\infty \to \mathbb{R}$ denote the duality pairing (see [21, Page 212] for duality pairings in this context) between L^1 and L^∞ , then $\langle P_t f, g \rangle = \langle f, K_t g \rangle$.

2.2.1. Infinitesimal Generators. Computing the Frobenius-Perron operator directly from the definition requires knowledge of the flow φ_t which is the solution to the differential equation $\dot{x} = X(x)$. This is often infeasible, especially if the analytical expression for φ_t is not known. This problem can be sidestepped by considering the infinitesimal generator of the Frobenius-Perron operator, which allows us to describe the evolution of an initial probability density function as the solution to a linear partial differential equation.

Definition 2.17 (Infinitesimal Generator of Frobenius-Perron Operator). Let $\{P_t : t \in \mathbb{R}\}$ be a family of Frobenius-Perron operators corresponding to a the flow $\{\varphi_t\}$ arising from the differential equation $\dot{x} = X(x)$. The infinitesimal generator for the Frobenius-Perron operator, $A: \operatorname{dom}(A) \subset L^1(M, \mu) \to L^1(M, \mu)$ is defined as

(2.10)
$$Af = \lim_{\tau \to 0} \frac{P_{t+\tau}f - P_tf}{\tau}.$$

We now state a theorem which allows for the computation of both A and P_t , assuming that f is differentiable.

Theorem 2.18 ([21, Theorem 7.6.11]). Let $X \in \mathfrak{X}(M)$ be a smooth vector field inducing the flow φ_t . Suppose that the measure space (M, μ) has a smooth measure, i.e., $\mu \in \Omega^n(M)$ is a differentiable volume-form. Then the infinitesimal generator A for the Frobenius-Perron operator is given by

(2.11)
$$Af = -df(X) - f \cdot \operatorname{div}_{\mu} X.$$

Moreover, if $u(t, x) = P_t f(x)$, then u solves the partial differential equation

(2.12)
$$\frac{\partial u}{\partial t} + du(X) = -u \cdot \operatorname{div}_{\mu}(X), \qquad u(0,x) = f(x)$$

where $\operatorname{div}_{\mu}(X)$ is the divergence of the vector field X with respect to the volume form μ .

Proof. Note that (2.12) holds as $AP_t = P_t A$, so it remains to show (2.11). Let $f \in L^1$ be an initial probability distribution function, then

$$\int_E (P_t f)\mu = \int_{\varphi_{-t}(E)} f\mu = \int_E (\varphi_{-t})^* f \cdot (\varphi_{-t})^* \mu,$$

which provides $(P_t f)\mu = (\varphi_{-t})^* f \cdot (\varphi_{-t})^* \mu$. Differentiating both sides with respect to t and applying the product rule yields

$$(Af)\mu = -df(X)\mu - f \cdot \operatorname{div}_{\mu}(X)\mu.$$

This immediately implies (2.11) as $\mu \neq 0$.

2.3. Reduction. Numerical computations of the Frobenius-Perron operator become costly with increasing dimension. This is why reduction techniques should be employed to make numerical simulations more tractable. At its core, the theory of reduction tries to deal with the following question: How can we reduce the dimensionality of a system, if there is access to additional information? There are different approaches to answering this question, depending on the type of insight available. In this work, we deal with systems that exhibit symmetry. Hybrid systems with symmetry are present in a myriad of examples, ranging from multi-legged locomotion [16, 34], image recognition and robotic perception [23] to climate dynamics [12].

Mathematically, symmetries are expressed as actions of a Lie group on the state space Q, which is assumed to be a manifold. From here, depending on what kind of structure is available on the configuration manifold there are different reduction techniques that can be applied. For instance, if the manifold is the cotangent bundle equipped with a symplectic form then we are in the realm of cotangent reduction [19]. On the other hand, if we are dealing with a Poisson manifold, then techniques of Poisson reduction can be applied. A particular case that fits in both of these scenarios is Lie-Poisson reduction, which deals with situations when the state space is a Lie group, acting on itself by left or right translation [17]. For example, this is the case for systems whose coordinates are Cartesian ($Q = \mathbb{R}^n$), polar ($Q = \mathbb{S}^1$, SO_n, etc.), or a combination ($Q = SE_n$). Classical Lie-Poisson reduction is capable of reducing the dimension from 2n down to n.

Recent efforts have been made towards creating a theory of cotangent reduction for hybrid systems [1]. Other versions of reduction have been carried out for impact systems, e.g. [3, 11, 13] and the references therein; however, there is still need for an analysis of Lie-Poisson reduction for hybrid systems. As will be discussed in Section 3.2.1 for hybrid systems on Lie groups, the best reduction we can attain is from 2n to n + 1 dimensions.

2.3.1. Lie-Poisson Reduction. In the following we present the continuous version of Lie Poisson reduction. Building upon this prerequisite, we will develop hybrid Lie Poisson reduction in Section 3.1. Let G be a Lie group and denote by \mathfrak{g} and \mathfrak{g}^* its Lie algebra and dual, respectively.



Figure 4: Schematic drawing of continuous Lie-Poisson reduction.

Consider the action of G on T^*G defined by the lift of left translations: $g \cdot \alpha = \ell_{g^{-1}}^* \alpha$. The momentum map of this action and the momentum map for the lift of right translations are

$$J_L \colon T^*G \to \mathfrak{g}^* : \alpha_g \mapsto r_g^* \alpha_g,$$
$$J_R \colon T^*G \to \mathfrak{g}^* : \alpha_g \mapsto \ell_g^* \alpha_g,$$

respectively. The cotangent bundle is equipped with the canonical symplectic form $\omega = dg \wedge dp_g$, which gives raise to the usual Poisson bracket

$$\{\cdot, \cdot\} \colon C^{\infty}(T^*G) \times C^{\infty}(T^*G) \to C^{\infty}(T^*G)$$
$$\colon (f, k) \mapsto \omega(X_f, X_k),$$

where X_f and X_k are the Hamiltonian vector fields of f and k, respectively. We assume that the functions in $C^{\infty}(T^*G)$ are left invariant. Then, using the right momentum map, one can project the dynamics on T^*G to \mathfrak{g}^* without losing essential information about the full trajectories (Figure 4).

The quotient manifold T^*G/G is isomorphic to the dual of the Lie algebra \mathfrak{g}^* [17, Chapter 13.3] and can be showed to be a Poisson manifold with Poisson bracket given by [24]

$$\{f,k\}(\mu) = \langle \mu, [df,dk] \rangle, \quad \text{for } f,k \in C^{\infty}(\mathfrak{g}^*).$$

where the bracket [df, dk] denotes the Lie bracket in \mathfrak{g} . Note that the differentials dk and df can indeed be considered elements of the Lie algebra, due to the identification of \mathfrak{g}^{**} with \mathfrak{g} .

Now, assume that the dynamics on the full cotangent bundle is given by a left invariant Hamiltonian $H: T^*G \to \mathbb{R}$. Using the available Poisson structure, it is possible to define a reduced dynamical system on \mathfrak{g}^* . This is summarized in the following theorem.

Theorem 2.19. Let $H: T^*G \to \mathbb{R}$ be a left invariant Hamiltonian, so $H(q, p_q) = H(gq, \ell_g^*p_q)$ for all $g \in G$ and $(q, p_q) \in T^*G$. Then, the restriction of H to \mathfrak{g}^* , denoted by $h = H|_{\mathfrak{g}^*}$, satisfies

$$H = h \circ J_R$$
 if and only if $H(q, p_q) = h(J_R(q, p_q)),$

and the flow Φ_t of H on T^*G is related to the flow ϕ_t of h on \mathfrak{g}^* by

$$J_R \circ \Phi_t = \phi_t \circ J_R$$

Moreover, the reduced equations of motion on g^* can be written as

(2.13)
$$\frac{d}{dt}\phi(t)(\mu_0) = \dot{\mu} = \operatorname{ad}_{dh}^* \mu,$$

where $\operatorname{ad}_{dh}^*: \mathfrak{g}^* \to \mathfrak{g}^*$ and $\operatorname{ad}_{dh}^* \mu(\xi) = \mu([dh, \xi])$ for all $\xi \in \mathfrak{g}^*$.

Proof. See Theorem 13.4.1 in [17].

Example 3. Consider the action of SO₃ on itself. The Lie algebra of SO₃ is $\mathfrak{so}_3 \cong \mathbb{R}^3$ with Lie bracket $[u, v] = u \times v$ and its dual $\mathfrak{so}_3^* \cong \mathbb{R}^3$ consists of row vectors p. The right momentum map is $J_R(A, p) = pA$, and the reduced Poisson bracket on \mathfrak{so}_3 is: $\{f, k\}(p) = p(df \times dk) = p^T \cdot (df \times dk)$. The reduced equations of motion (2.13) become $\dot{\mu}^T = \operatorname{ad}_{dh}^* \mu = \mu^T \times dh$ or $\dot{\mu} = \mu \times dh$ if μ is viewed here as a column vector.

In Section 3.2.1, we will extend the reduction presented above to hybrid systems.

3. Results. Our results are twofold. First, we define the hybrid Frobenius-Perron operator, which is the analogous operator to the continuous case, extended for when the underlying dynamics is hybrid. Then, we extend Lie-Poisson reduction to hybrid systems whose state space is a Lie group.

3.1. Hybrid Frobenius-Perron and Koopman operators. In this section, we derive the infinitesimal generator for the Frobenius-Perron operator of hybrid systems. As hybrid systems are a mixture of both discrete- and continuous-time systems, the infinitesimal generator will be a combination of both (2.9) and (2.12).

Definition 3.1. Let $\varphi_t^{\mathcal{H}} \colon M \to M$ be a hybrid flow on the differentiable manifold M. Assume there exists a differentiable volume form $\mu \in \Omega^n(M)$ such that (M,μ) is a measure space and that the flow $\varphi_t^{\mathcal{H}}$ is complete. Then, the hybrid Frobenius-Perron operator corresponding to $\varphi_t^{\mathcal{H}}$ is the unique linear operator $P_t^{\mathcal{H}} \colon L^1(M,\mu) \to L^1(M,\mu)$ implicitly defined by

(3.1)
$$\int_{E} P_{t}^{\mathcal{H}} f \, d\mu = \int_{\varphi_{-t}^{\mathcal{H}}(E)} f \, d\mu \qquad \text{for all } E \subseteq M.$$

When $P_t^{\mathcal{H}}$ is continuous, the hybrid infinitesimal generator is defined to be

(3.2)
$$A^{\mathcal{H}}f = \lim_{\Delta t \to 0} \frac{P_{t+\Delta t}^{\mathcal{H}}f - P_{t}^{\mathcal{H}}f}{\Delta t}.$$

As in the continuous case, our goal is to find an explicit formula for the infinitesimal generator. Moreover, as in Theorem 2.18 we want to find a partial differential equation for $u(t,x) \coloneqq P_t^{\mathcal{H}} f(x)$.

Remark 3.2. Before we proceed, we will further examine (3.2). Assume E is a sufficiently small ball about some point $x \in M \setminus S$, and that $\varphi_{-t}^{\mathcal{H}}(E) \cap S = \emptyset$ for some time $t \in \mathbb{R}$. Then, within E, the dynamics of point x follows the continuous part of $\varphi_t^{\mathcal{H}}$, which is smooth by the quasi-smooth dependence propriety. Since we are taking the limit as $\Delta t \to 0$, we can assume that $\varphi_s^{\mathcal{H}}(E) \cap S = \emptyset$ for $-t - \Delta t \leq s \leq -t$. Hence, the hybrid infinitesimal generator should be the same as in the continuous case.

Now consider what happens when $E \cap S \neq \emptyset$. At a first glance, it is tempting to define the transfer operator as the one corresponding to the discrete reset map Δ ; however, the guard S as well as its tangent space is n-1 dimensional, whereas the image of the reset map is M is n dimensional. This is an issue since both sides of (3.1) integrate μ , which is a volume form on M. In order to deal with this mismatch in dimensionality, we will extend the differential of Δ , so that it acts on vectors in the tangent space of M restricted to base points in S. We will do this by taking advantage of the fact that any vector in the tangent space at $x \in S$ can be decomposed into its component along the flow v_x , and $v_s \in T_x S$ as in Figure 5.

In the following, we will make the decomposition from the previous remark more precise. Definition 3.3 (Extended differential). Let $\mathcal{H} = (M, \mathcal{S}, \Delta, X)$ be a smooth HDS that satisfies propriety 2.3 and let $x \in \mathcal{S}$. The augmented differential of the reset map is the linear map

$$\Delta_*^X \colon T_x M \to T_{\Delta(x)} M : v \mapsto \begin{cases} \Delta_* v & \text{if } v \in T_x \mathcal{S}; \\ c X_{\Delta(x)} & \text{if } v = c X_x \in \operatorname{Span}(X_x). \end{cases}$$

When defining the discrete part of the Frobenius-Perron operator, we will use the Jacobian of the augmented differential rather than the Jacobian of the reset map. This fixes the dimensionality issue, and is indeed the correct procedure as will become apparent in the proof of Theorem 3.11 below.

The remainder of this section is dedicated to finding a hybrid analog of formula (2.9). The result is the summarized in Theorem 3.12, and is the main contribution of this section. As in the discrete case, the goal is to perform a change of coordinates in the integral (3.1) so as



Figure 5: Decomposition of vectors in the tangent space of the ambient manifold M, with base point on the guard.

to obtain an explicit formula for the Frobenius-Perron operator. Due to the dimensionality issues presented in Remark 3.2, this is not straightforward, and will be carefully analysed in the following.

We begin by giving a precise definition of the determinant for general manifolds. The determinant will be important later on, when extending the change of variables formula from the discrete case (2.9).

Definition 3.4 (Form determinant [32]). Let M and N be n-dimensional smooth manifolds with volume-forms $\mu \in \Omega^n(M)$ and $\eta \in \Omega^n(N)$. Let $F: TM \to TN$ be a smooth map which is linear in the fibers. Then the determinant of F with respect to μ and η is defined to be the unique $\mathcal{C}^{\infty}(M)$ function such that

$$\det_{\mu \to \eta}(F) \cdot \mu = F^* \eta$$

Moreover, if $F: TM \to TM$ then we will write $\det_{\mu \to \mu}(F)$ as $\det_{\mu}(F)$.

Remark 3.5. If $f: M \to N$ is a smooth map, $\det_{\mu \to \eta}(f_*)$ is the usual Jacobian determinant.

Remark 3.6. The existence and uniqueness of the determinant comes from the fact that both μ and $F^*\eta$ are top-forms on M. The determinant does not depend on the vectors that μ and $F^*\eta$ act on. In particular,

$$\det_{\mu \to \eta}(F) \cdot \mu(v^1, \dots, v^n) = F^* \eta(v^1, \dots, v^n) \quad \text{for all } v^1, \dots, v^n \in T_x M.$$

When $F = f_*$, there is an elegant formula for the determinant of the inverse transformation. Lemma 3.7. For a diffeomorphism $f: M \to N$ and volume-forms $\mu \in \Omega^n(M)$ and $\eta \in \Omega^n(N)$,

$$\det_{\mu \to \eta}(f_*^{-1}) = \frac{1}{\det_{\eta \to \mu}(f_*) \circ f^{-1}}.$$

Proof. See Appendix B.1.

We now turn to the case where $f: S \to M$ is the reset map Δ . This is precisely where the augmented differential becomes important since it offers a way to relate the determinant of the reset map with respect to the induced measures on S and $\Delta(S)$ to the determinant of a linear map from TM to TM with respect to just μ . We formalize this through the following lemma.

Lemma 3.8 (Transverse property). Let $\Delta^X_*: TM|_{\mathcal{S}} \to TM|_{\Delta(\mathcal{S})}$ be the augmented differential and let $\iota_{\mathcal{S}}: \mathcal{S} \hookrightarrow M$ and $\iota_{\Delta(\mathcal{S})}: \Delta(\mathcal{S}) \hookrightarrow M$ denote the respective inclusion maps. If X is transverse to \mathcal{S} , then

$$\det_{\mu \to \mu} \Delta_*^X = \det_{\alpha \to \beta} \Delta_*$$

where $\alpha = \iota_{\mathcal{S}}^* i_X \mu$, $\beta = \iota_{\Delta(\mathcal{S})}^* i_X \mu$. *Proof.* See Appendix A.1.

Keeping the same notation as in Lemma 3.8, the definition of the determinant gives us

$$\det_{\alpha \to \beta} (\Delta_*) \cdot \alpha = \Delta^* \beta$$

and plugging in the expressions for α and β , we see that

(3.3)
$$\det_{\alpha \to \beta} (\Delta_*) \cdot \iota_S^* i_X \mu = \Delta^* \iota_{\Delta(S)}^* i_X \mu$$

which is precisely the defining equation for the hybrid Jacobian $\mathcal{J}^X_{\mu}(\Delta)$. Thus, by the uniqueness of the determinant we have

$$\det_{\alpha \to \beta} \Delta_* = \mathcal{J}^X_{\mu}(\Delta), \quad \text{and finally}, \quad \det_{\mu \to \mu} \Delta^X_* = \mathcal{J}^X_{\mu}(\Delta).$$

In other words, the hybrid Jacobian is equal to the determinant of the augmented differential, as stated precisely in the following lemma.

Lemma 3.9 (Relation between determinant and Jacobian). The determinant of the augmented differential is the hybrid Jacobian, i.e.,

$$\det_{\iota \to \mu} \Delta^X_* = \mathcal{J}^X_\mu(\Delta).$$

Combining lemmas 3.9 and 3.7, we obtain a formula for how the measure changes under the coordinate transformations $x \mapsto \Delta(x)$.

Lemma 3.10. Suppose that $\Delta: S \to \Delta(S)$ is invertible and the linear transformation Δ_*^X is non-singular. Then,

$$\mu_{\Delta^{-1}(x)}\Big((\Delta_*^X)^{-1} v^1, \dots, (\Delta_*^X)^{-1} v^n \Big) = \frac{1}{(\mathcal{J}^X_{\mu}(\Delta) \circ \Delta^{-1})(x)} \cdot \mu_x(v^1, \dots, v^n),$$

for all $v^1, \ldots, v^n \in T_x M|_{\mathcal{S}}$.

Proof. By the definition of the form determinant,

(3.4)
$$\mu_{\Delta^{-1}(x)}\left(\left(\Delta_{*}^{X}\right)^{-1}v^{1},\ldots,\left(\Delta_{*}^{X}\right)^{-1}v^{n}\right) = \det_{\mu}\left(\left(\Delta_{*}^{X}\right)^{-1}\right) \cdot \mu_{x}(v^{1},\ldots,v^{n})$$

Utilizing Lemma 3.7 to rewrite (3.4), we obtain

(3.5)
$$\mu_{\Delta^{-1}(x)}\left(\left(\Delta_{*}^{X}\right)^{-1}v^{1},\ldots,\left(\Delta_{*}^{X}\right)^{-1}v^{n}\right) = \frac{1}{\left(\det_{\mu}(\Delta_{*}^{X})\circ\Delta^{-1}\right)(x)}\cdot\mu_{x}(v^{1},\ldots,v^{n}),$$

and Lemma 3.9 gives us the desired result.

We are now ready to state the main theorem.

Theorem 3.11 (Hybrid infinitesimal generator for invertible Δ). Let \mathcal{H} be a smooth hybrid dynamical system with an invertible reset map Δ and dynamics given by (1.1). Let $\mu \in \Omega^n(M)$ be a reference volume-form and suppose that $\mathcal{J}^X_{\mu}(\Delta) \neq 0$. Then, the hybrid transfer operator $u(t,x) \coloneqq P_t^{\mathcal{H}} f(x)$ satisfies the following

(3.6)
$$\begin{cases} \frac{\partial u}{\partial t} + du(X) = -u \cdot \operatorname{div}_{\mu}(X) & \text{for } x \notin \Delta(\mathcal{S}); \\ u(t^{+}, x) = \frac{u(t^{-}, \Delta^{-1}(x))}{\left(\mathcal{J}_{\mu}^{X}(\Delta) \circ \Delta^{-1}\right)(x)} & \text{for } x \in \Delta(\mathcal{S}). \end{cases}$$

Proof. We will first derive the continuous case of (3.6); let $x \notin \Delta(S)$. To do so, perform a change of coordinates in Equation (3.1) to obtain

(3.7)
$$\int_{E} P_{t}^{\mathcal{H}} f(x) d\mu = \int_{\varphi_{-t}^{\mathcal{H}}(E)} f(x) d\mu = \int_{E} \left(f \circ \varphi_{-t}^{\mathcal{H}} \right) (x) \left(\varphi_{-t}^{\mathcal{H}} \right)^{*} \mu,$$

which holds for any measurable $E \subseteq M$. Pointwise, we have

(3.8)
$$P_t^{\mathcal{H}} f(x) \mu = \left(f \circ \varphi_{-t}^{\mathcal{H}} \right) (x) \left(\varphi_{-t}^{\mathcal{H}} \right)^* \mu$$

Since $x \notin \Delta(\mathcal{S})$, the quasi-smooth dependence propriety guarantees the existence of a small interval $(-\varepsilon, \varepsilon)$ such that $\varphi_{-\tau}^{\mathcal{H}}(x) \notin \Delta(\mathcal{S})$ for all $\tau \in (-\varepsilon, \varepsilon)$, which, in turn, implies the map $\tau \mapsto \varphi_{-\tau}^{\mathcal{H}}$ is smooth in a neighborhood of x. We can then differentiate (3.8) with respect to t to obtain

$$\frac{\partial u}{\partial t}\mu = -du(X)\mu - u \cdot \operatorname{div}_{\mu}(X)\mu.$$

Finally, dividing out by $\mu \neq 0$ yields the $x \notin \Delta(\mathcal{S})$ case of (3.6).

Now, suppose that $x \in \Delta(\mathcal{S})$ at t = 0, i.e., the starting point at x is right after an impact has occurred. Yet again using the quasi-smooth dependence property, let $\varepsilon \in \mathbb{R}^+$ be such that $\varphi_{-t}^{\mathcal{H}}(x) \notin \mathcal{S}$ for all $t \in (0, \varepsilon)$ so that $\varphi_{-t}^{\mathcal{H}}(x) = \varphi_{-t}(x)$ and $\lim_{t \searrow 0} \varphi_{-t}^{\mathcal{H}}(x) = \Delta^{-1}(x)$. The goal will be to use equation (3.8) and take the limit as $t \searrow 0$.

First, we analyze the behavior of the pushforward of vectors tangent to S by the flow $\varphi_{-t}^{\mathcal{H}}$, and we claim that

(3.9)
$$\lim_{t \searrow 0} (\varphi_{-t}^{\mathcal{H}})_* v = (\Delta_*^X)^{-1} v, \quad \text{for all } v \in T_x M.$$



(a) Pushforward of the flow on vectors in the tangent space of the impact surface. For a curve $\gamma(s) \in \Delta(S)$ that passes through x, the curve $\Delta^{-1}(\gamma(s)) \in S$ passes through $y = \Delta^{-1}(x)$. Since $\lim_{t \searrow 0} (\varphi_{-t}^{\mathcal{H}}) = \Delta^{-1}\gamma$, in the limit the pushforward v under the flow is the tangent vector at y of $\Delta^{-1}(\gamma)$ depicted in red.



(b) The effect of the pushforward of the flow on vectors which are tangent to the flow. The representative curve of X_x is the flow itself $\varphi_s^{\mathcal{H}}$ depicted in green. By the semigroup propriety, when we compose it with $\varphi_{-t}^{\mathcal{H}}$ we get the same flow, but with a time shift of -t depicted in orange. Differentiating and taking the limit as $t \searrow 0$ gives $X_{\Delta^{-1}(x)}$.

Figure 6: Schematic drawing of the two cases in the proof of Theorem 3.11.

To show this, we consider two cases based on the decomposition $T_x M = T_x \Delta(\mathcal{S}) \oplus \text{Span}(X_x)$. First, let $v \in T_x \Delta(\mathcal{S})$. We use an argument similar to that of Theorem 4.2 in [9], which is depicted in Figure 6a.

$$\lim_{t \to 0} (\varphi_{-t}^{\mathcal{H}})_* v = \Delta_*^{-1} v \qquad \text{Since } -t < 0 \text{ for all } t \in (0, \varepsilon)$$
$$= (\Delta_*^X)^{-1} (v) \qquad \text{By the definition of } (\Delta_*^X)^{-1}$$

Now, let $v = cX_x \in \text{Span}(X_x)$ as illustrated in Figure 6b. By the semigroup propriety of $\varphi_{-t}^{\mathcal{H}}$ and the definition of $(\Delta_*^X)^{-1}$, we have that

$$\lim_{t \searrow 0} (\varphi_{-t}^{\mathcal{H}})_* cX_x = cX_{\Delta^{-1}(x)} = (\Delta_*^X)^{-1} (cX_x)$$

Next, we take the limit of Equation (3.8) as $t \searrow 0$ and get

(3.10)
$$u(0^+, x)\mu = \lim_{t \searrow 0} u(t, x)\mu = \lim_{t \searrow 0} P_t^{\mathcal{H}} f(x)\mu = \lim_{t \searrow 0} \left(f \circ \varphi_{-t}^{\mathcal{H}} \right) (x) \left(\varphi_{-t}^{\mathcal{H}} \right)^* \mu.$$

As in the continuous case, we want to cancel μ_x from both sides. To do so, Let $v^1, \ldots, v^n \in$

 $T_x M$ be arbitrary. Plugging v^1, \ldots, v^n into Equation (3.10), we see that

$$\begin{aligned} u(0^+, x)\mu_x(v^1, \dots, v^n) &= \lim_{t \searrow 0} \left(f \circ \varphi_{-t}^{\mathcal{H}}(x) \right) \cdot \left(\varphi_{-t}^{\mathcal{H}} \right)^* \mu_x(v^1, \dots, v^n) & \text{By Equation (3.10)} \\ &= \lim_{t \searrow 0} \left(f \circ \varphi_{-t}^{\mathcal{H}}(x) \right) \cdot \mu_{\varphi_{-t}^{\mathcal{H}}(x)} \left((\varphi_{-t}^{\mathcal{H}})_* v^1, \dots, (\varphi_{-t}^{\mathcal{H}})_* v^n \right) & \text{By the definition of the pushforward} \\ &= f(\Delta^{-1}(x)) \cdot \mu_{\Delta^{-1}(x)} \left((\Delta_*^X)^{-1}(v^1), \dots, (\Delta_*^X)^{-1}(v^n) \right) & \text{By Equation (3.9)} \\ &= \frac{f(\Delta^{-1}(x))}{\left(\mathcal{J}_{\mu}^X(\Delta) \right) (\Delta^{-1}(x))} \mu_x(v^1, \dots, v^n) & \text{By Lemma 3.10} \end{aligned}$$

$$(3.11) \qquad = \frac{u(0^-, \Delta^{-1}(x))}{\left(\mathcal{J}_{\mu}^X(\Delta) \circ \Delta^{-1} \right)(x)} \mu_x(v^1, \dots, v^n) & \text{By the definition of } u \end{aligned}$$

Since v^1, \ldots, v^n are arbitrary and $\mu_x \neq 0$, we have obtained the discrete case of (3.6) for t = 0.

Now, consider $\varphi_{-t}^{\mathcal{H}}(x) \in \Delta(\mathcal{S})$ for $t \neq 0$ and without loss of generality, consider t > 0 since the t < 0 case follows from applying the same steps in the opposite direction.

$$\begin{aligned} u(t^+, x) &= P_t^{\mathcal{H}} u(0^+, x) & \text{By definition of } u \\ &= P_t^{\mathcal{H}} \frac{u(0^-, \Delta^{-1}(x))}{\left(\mathcal{J}_{\mu}^X(\Delta) \circ \Delta^{-1}\right)(x)} & \text{By Equation (3.11)} \\ &= \frac{1}{\left(\mathcal{J}_{\mu}^X(\Delta) \circ \Delta^{-1}\right)(x)} P_t^{\mathcal{H}} u(0^-, \Delta^{-1}(x)) & \text{Since } \left(\mathcal{J}_{\mu}^X(\Delta) \circ \Delta^{-1}\right)(x) \in \mathbb{R} \\ &= \frac{1}{\left(\mathcal{J}_{\mu}^X(\Delta) \circ \Delta^{-1}\right)(x)} u(t^-, \Delta^{-1}(x)) & \text{Since } t > 0 \end{aligned}$$

which concludes the proof.

In general, the reset map need not be invertible. Assuming that Δ^{-1} has finitely many preimages, Theorem 3.11 still holds. The precise statement is as follows.

Theorem 3.12 (General hybrid infinitesimal generator). Let \mathcal{H} be a smooth hybrid dynamical system with dynamics given by (1.1) and suppose $\Delta^{-1}(\{x\})$ is finite. Additionally, let $\mu \in \Omega^n(M)$ be a reference volume-form and suppose that $\mathcal{J}^X_{\mu}(\Delta) \neq 0$. Then, the hybrid transfer operator $u(t,x) \coloneqq P_t^{\mathcal{H}} f(x)$ satisfies the following

(3.12)
$$\begin{cases} \frac{\partial u}{\partial t} + du(X) = -u \cdot \operatorname{div}_{\mu}(X) & \text{for } x \notin \Delta(\mathcal{S}); \\ u(t^+, x) = \sum_{y \in \Delta^{-1}(\{x\})} \frac{1}{(\mathcal{J}^X_{\mu}(\Delta))(y)} u(t^-, y) & \text{for } x \in \Delta(\mathcal{S}). \end{cases}$$

Proof. As the continuous case of Theorem 3.12 is identical to that of 3.11, it remains to show

the $x \in \Delta(S)$ case of (3.12). Doing so, let $\Delta^{-1}(\{x\}) = \{y_1, \ldots, y_n\}$ and define $E \subset M$ to be a half-neighborhood about $x \in \Delta(S)$ such that

$$\lim_{\tau \searrow t} \varphi_{-\tau}^{\mathcal{H}}(E) = \bigcup_{i=1}^{n} E_i,$$

where each E_i is a half-neighborhood about y_i and $\{E_i\}$ is pairwise disjoint as in Figure 7. Let $\varphi_t^i = \varphi_t^{\mathcal{H}}|_{E_i}$ denote the hybrid flow restricted to each ball E_i so that each φ_t^i is invertible. Now, we will apply the proof of 3.11 to each φ_i^i and

the multiple preimages. Generalizing



apply the proof of 3.11 to each φ_t^i and Figure 7: Diagrammatic construction of E_i and its then combine the results to account for corresponding φ_{-t}^i when n = 3.

the change of coordinates in (3.7), becomes a sum over the φ_{-t}^{i} 's as follows

$$\int_{E} P_{t}^{\mathcal{H}} f(x) \, d\mu = \sum_{i=1}^{n} \int_{\varphi_{-t}^{i}(E)} f(x) \, d\mu = \sum_{i=1}^{n} \int_{E} \left(f \circ \varphi_{-t}^{i} \right) (x) \left(\varphi_{-t}^{i} \right)^{*} \mu_{t}$$

which by similar reasoning, implies that

(3.13)
$$P_t^{\mathcal{H}} f(x)\mu = \sum_{i=1}^n (f \circ \varphi_{-t}^i)(x)(\varphi_{-t}^i)^*\mu.$$

Following the same steps that imply Equation (3.11) gives us

(3.14)
$$\lim_{t \searrow 0} \left(f \circ \varphi_{-t}^i \right)(x) \left(\varphi_{-t}^i \right)^* \mu_x(v^1, \dots, v^n) = \frac{f(y_i)}{\left(\mathcal{J}^X_\mu(\Delta) \right)(y_i)} \mu_x(v^1, \dots, v^n).$$

since Δ is invertible when restricted to E_i . Summing (3.14) over each $y_i \in \Delta^{-1}(\{x\})$ and equating with the left-hand side of (3.13) gives

$$u(t^+, x) = \sum_{i=1}^n \frac{1}{\left(\mathcal{J}^X_\mu(\Delta)\right)(y_i)} u(t^-, y_i) = \sum_{y \in \Delta^{-1}(\{x\})} \frac{1}{\left(\mathcal{J}^X_\mu(\Delta)\right)(y)} u(t^-, y),$$

which is precisely the discrete part of Equation (3.12).

Remark 3.13 ($\Delta^{-1}(\{x\})$) with infinite cardinality). Consider the smooth HDS

$$(M, \mathcal{S}, \Delta, X) = (\mathbb{R}, \mathbb{Z} \setminus \{0\}, n \mapsto 0, \partial/\partial x),$$

which implies $\Delta(S) = \{0\}$. This is an HDS that does not undergo beating or any sort of Zeno properties, but $\Delta^{-1}(\{0\}) = \mathbb{Z} \setminus \{0\}$, which is infinite. This implies that we necessarily cannot

assume that the preimage of a point in $\Delta(\mathcal{S})$ has finite cardinality. Suppose $\mu = dx$. Then,

$$\Delta^* i_X \mu = \Delta^* dx(X) = \Delta^* 1 = 1 \circ \Delta = 1$$
$$\iota_S^* i_X \mu = \iota_S^* dx(X) = 1 \circ \iota_S = 1$$

which implies that $\mathcal{J}^X_{\mu}(\Delta) = 1$. Then,

(3.15)
$$u(0^+, 0) = \sum_{y \in \Delta^{-1}(\{0\})} \frac{1}{\mathcal{J}^X_\mu(\Delta)(y)} u(0^-, y) = \sum_{y \in \mathbb{Z} \setminus \{0\}} u(0^-, y)$$

which can blowup for certain choices of f.

3.2. Hybrid Reduction. The primary goal of this work is to understand and compute the Frobenius-Perron operator for mechanical impact systems which results in solving (3.6). Unfortunately, the dimension in many examples makes direct numerical study difficult. The goal of this section is to extend the idea of Lie-Poisson reduction (Theorems 2.19 and its nonholonomic counterpart) to impact systems. Other versions of reduction have been carried out for impact systems, e.g. [3, 11, 13] and the references therein. Classical Lie-Poisson reduction is capable of reducing the dimension from 2n down to n. When impacts are considered, it will be shown that the dimension can only be reduced to n + 1.

3.2.1. Hybrid Lie-Poisson reduction. Lie-Poisson reduction is possible when the Hamiltonian is left-invariant as the dynamics can be translated to the identity element. The natural extension to impact systems is for the impact surface, $\Sigma \subset G$, to be left-invariant as well. As will be shown below, this is true precisely when Σ is the *right* coset of a normal subgroup.

Before we proceed, we provide a summary of the notation used in this section.

Term	Notation	Element(s)
Lie Group	G	g,g_0
Lie Subgroup	K	$k, ilde{k}$
Lie Algebra	g	ξ
Lie Algebra of K	Ŕ	$\delta k,\widetilde{\delta k}$
Dual Lie Algebra	\mathfrak{g}^*	ζ
Impact Surface	$\Sigma = Kg_0$	kg_0
Metric on G	Ц	N/A

Table 1: Notation for Section 3.2.1

Throughout this section we consider mechanical impact systems as defined in 2.9, which, by definition, means that the natural Hamiltonian can be written as

$$H(q,p) = \frac{1}{2} \mathbf{L}^{-1}(p,p) + V(q)$$

Remark 3.14. We can go between G's Lie algebra \mathfrak{g} and its dual \mathfrak{g}^* using the musical isomorphisms. In particular $\xi := \zeta^{\sharp} = (\Pi^{-1})^{ij} \zeta_j$.

Assume now that we are in the setting of Lie-Poisson reduction, i.e., H is left invariant and the state space is a Lie group G. Then the full equations of motion for the mechanical impact system (G, H, Σ) are given by the following Proposition.

Proposition 3.15. Let $H: T^*G \to \mathbb{R}$ be a natural, left-invariant Hamiltonian and $h: \mathfrak{g}^* \to \mathbb{R}$ be its restriction to the identity. For an impact surface, $\Sigma \subset G$, let $s: G \to \mathbb{R}$ be such that zero is a regular value and (locally) $\Sigma = s^{-1}(\{0\})$. Then, between impacts, the usual Lie-Poisson equations hold

(3.16)
$$\begin{cases} \dot{\zeta} = \mathrm{ad}_{dh}^* \zeta; \\ \dot{g} = (\ell_g)_* \zeta^{\sharp}, \qquad \text{for } s(g) \neq 0, \end{cases}$$

and upon impact, the left-translation of the corner conditions (2.4) becomes

(3.17)
$$\begin{cases} \zeta \mapsto \zeta - 2 \frac{\left(\ell_g\right)^* ds_g\left(\zeta^{\sharp}\right)}{\ell_g^* ds_g\left(\left(\ell_{g^{-1}}\right)_* ds_g^{\sharp}\right)} \left(\ell_g\right)^* ds_g; & \text{for } s(g) = 0, \\ g \mapsto g, \end{cases}$$

where $\sharp: \Omega^1(G) \to \mathfrak{X}(G)$ is the musical isomorphism, as defined in 2.7.

Proof. For an arbitrary manifold Q with natural Hamiltonian and non-moving impact surface, the variational impact equations are given by [9]:

(3.18)
$$\begin{cases} p^{+} = p^{-} - \frac{2p^{-}(\nabla s_{q})}{ds_{q}(\nabla s_{q})} ds_{q}; \\ q^{+} = q^{-}, \end{cases} \text{ for } q \in \Sigma, \end{cases}$$

where $\nabla s_q = (ds_q)^{\sharp}$. By [17], the system will follow the continuous Lie-Poisson equations of motion as follows when away from impacts.

$$\begin{cases} \dot{\zeta} = \mathrm{ad}_{dh}^* \zeta; \\ \dot{g} = (\ell_{g^{-1}})_* \zeta^{\sharp} \end{cases}$$

In order to get the impact equations on the dual of the Lie algebra we need to left translate (3.18) to the origin. For $g \in G$,

$$p^{+} = p^{-} - \frac{2p^{-}(\nabla s_{g})}{ds_{g}(\nabla s_{g})}ds_{g}$$
 Starting with (3.18)
$$\iff (\ell_{g})^{*}p^{+} = (\ell_{g})^{*}p^{-} - (\ell_{g})^{*}\frac{2p^{-}(\nabla s_{g})}{ds_{g}(\nabla s_{g})}ds_{g}$$
 After left translating
$$\iff \zeta^{+} = \zeta^{-} - \frac{2\zeta^{-}((\ell_{g^{-1}})_{*}\nabla s_{g})}{\ell_{g}^{*}ds_{g}((\ell_{g^{-1}})_{*}\nabla s_{g})}(\ell_{g})^{*}ds_{g}$$
 By the definition of ζ

where $\zeta^+ = (\ell_q)^* p^+$ and $\zeta^- = (\ell_q)^* p^-$ are elements of \mathfrak{g}^* . The last step is to note that

$\zeta(\ell_{g^{-1}})_* \nabla s_g = g^{-1} \cdot \zeta_i(\Pi^{-1}) i j (ds_g)_j$	Expanding in local coordinate representaion
$= (ds_g)_j g^{-1} \cdot (\mathrm{II}^{-1})^{ij} \zeta_i$	Since $(ds_g)_j \in \mathbb{R}$ for each j
$= ds_g (\ell_{g^{-1}})_* \zeta^\sharp$	Contracting indices
$=\ell_{q^{-1}}^* ds_g(\zeta^\sharp)$	By the definition of the pullback

The goal now is to provide an analog of continuous Lie-Poisson reduction 2.19 for the hybrid equations of motion presented above 3.15. This is not straightforward due to the reasons presented in the following:

Remark 3.16. The power of the Lie-Poisson equations (2.13) is that it decouples the momentum from the position dynamics. This fails to be the case for the impact case for two main reasons: 1) the impact map for ζ in (3.17) depends on g as well as ζ , and 2) switching between (3.16) and (3.17) depends on g (i.e. on the position on the impact surface). As will be shown, by requiring the impact set to be left-invariant, the first issue can be resolved. The second issue requires more structure i.e. that of a right coset of a normal subgroup.

The overarching idea is the following: we want to be able to detect when the impact occurs without having to keep track of the entire n dimensional trajectory in G. We try to do so in the least number of dimensions possible. One crucial piece of information is *whether* the impact is happening or not. Hence, the question to be asked is: what propriety does Σ need to have so that *where* the impact happens does not affect the reduced hybrid equations of motion? From the following definitions and lemmas it will become apparent that Σ needs to be a right coset.

Definition 3.17 (Impact *u*-stabilizer). Given some $u \in \Sigma \subseteq G$, define the impact u- stabilizer to be the set



$$G_u(\Sigma) = \{g \in G : gu \in \Sigma\}.$$
 Figure 8: Depiction of Σ elements remaining in Σ after multiplication by $G_u(\Sigma)$ elements.

In other words $G_u(\Sigma)$, consists of all group elements that, when acting on u, keep Σ invariant. A special property holds when Σ is a right coset.

Lemma 3.18. Let K be a subgroup of G, let $\Sigma = Kg_0$ for some $g_0 \in G$, and define $G_u(\Sigma)$ as in Definition 3.17. Then, $G_u(\Sigma) = K$.

Proof. We will show $G_u(\Sigma) \subseteq K$ and then $K \subseteq G_u(\Sigma)$. Let $g \in G_u(\Sigma)$ be arbitrary so $gu \in \Sigma$ by definition. Since $\Sigma = Kg_0$, there exists a $k \in K$ such that $gu = kg_0$, which implies $g = kg_0u^{-1}$. Since $u \in \Sigma = Kg_0$, we know that $u = \tilde{k}g_0$ for some $\tilde{k} \in K$. Substituting this in, $g = kg_0u^{-1} = kg_0(\tilde{k}g_0)^{-1} = kg_0g_0^{-1}\tilde{k}^{-1} = k\tilde{k}^{-1}$. Since $K \leq G$, we know $g = k\tilde{k}^{-1} \in K$, so $G_u(\Sigma) \subseteq K$. Now, let $k \in K$ be arbitrary. Using similar arguments as in the first

part, $ku = k\tilde{k}g_0$ for some $\tilde{k} \in K$, and since $K \leq G$, we know that $k\tilde{k} \in K$, which implies $ku \in Kg_0 = \Sigma$. Thus, $G_u(\Sigma) = K$.

In order to reduce the dimension to n+1, it should not matter whether the impact happens at u or at another point in $G_u(\Sigma)$. The previous lemma implies that for right cosets $G_u(\Sigma) = K$ for any u. Assuming that points in $G_u(\Sigma)$ are equivalent for the reset map, this implies that for right cosets, the only thing we would need to keep track of is the direction normal to K. This is indeed one dimensional if K is codimension 1. Now we need to see when the equivalence assumption actually holds. For this we define the tangent preserving propriety.

Definition 3.19 (Tangent preserving). Let Σ be a submanifold of G (not necessarily a subgroup). Then, Σ is said to be (left) tangent preserving if for $u \in \Sigma$ and $g \in G_u(\Sigma)$, then

$$(\ell_q)_* T_u \Sigma = T_{qu} \Sigma$$



Figure 9: Geometric depiction of the tangent preserving propriety, where Σ has the tangent preserving propriety and Σ' does not, so $(\ell_g)_* v \in T\Sigma$ and $(\ell_g)_* v \notin T\Sigma'$.

Lemma 3.20. If $\Sigma = Kg_0$ for a subgroup $K \leq G$, then Σ is (left) tangent preserving.

Proof. Let $g \in G_u = K$ be arbitrary. By Lemma 3.18, we have $g\Sigma = gKg_0 = Kg_0 = \Sigma$, so $\ell_g \colon \Sigma \to \Sigma$ is a diffeomorphism, which implies that $(\ell_g)_*T_u\Sigma = T_{gu}\Sigma$. Thus, Σ is tangent preserving.

The tangent preserving propriety tells us that we can left translate a vector along Σ , and still stay in the tangent space of Σ . See Figure 9 for a schematic drawing.

Lemma 3.21. Assume, as before, that $K \leq G$ is a subgroup and let $\Sigma = Kg_0$ for some $g_0 \in G$. If there exists a smooth function $s \in C^{\infty}(G)$ such that $s^{-1}(\{0\}) = \Sigma$, then for all $u \in S$, $g \in G_u(\Sigma)$, there exists an $\alpha_{u,g} \in \mathbb{R}$ such that $\alpha_{u,g} ds_u = (\ell_g)^* ds_{ug}$.

Proof. Since $\Sigma = Kg_0$, we use Lemma 3.19 states that $(\ell_g)_*T_u\Sigma = T_{gu}\Sigma$ for any $g \in K$. By the definition of $s \in C^{\infty}(G)$, we know $T_u\Sigma = \ker ds_u$ for all $u \in \Sigma$. Then applying the tangent preserving property,

$$\ker ds_u = \left(\ell_{g^{-1}}\right)_* \ker ds_{gu} = \ker((\ell_g)^* ds_{gu}).$$

As $T_u\Sigma = \ker ds_u = \ker((\ell_g)^* ds_{gu}) \subset T_uG$ is a codimension 1 subspace, it follows that ds_u and $(\ell_g)^* ds_{gu}$ are linearly dependent. Since $T_u\Sigma = \ker ds_u = \ker((\ell_g)^* ds_{gu})$ is an codimension 1 vector field, it follows that the range of both ds_u and ds_{gu} is one dimensional and thus ds_u and ds_{gu} must differ by a constant i.e. there exists an $\alpha_{u,g} \in \mathbb{R}$ such that $\alpha_{u,g} ds_u = ds_{ug}$.

Hence, if the tangent preserving property is fulfilled, the only information that we lose by translating vectors along Σ is contained within a constant.



Figure 10: Depiction of independence of local section as guaranteed by 3.22. The lift of $q = [g_0]$ in the case of $\mathbb{S}^1 \subset \mathbb{T}^2$ is the circle depicted in red. The tangent vectors to any section project down to the same \dot{q} in the reduced space $\mathbb{S}^1 \setminus \mathbb{T}^2$.

With the help of the previous lemmas, we can address the issues in Remark 3.16 in the theorem below.

Theorem 3.22 (Impact Lie-Poisson Reduction). Let G be a Lie group, $g_0 \in G$ be arbitrary, and $\Sigma = Kg_0 \subseteq G$ be a right coset of codimension 1 for a closed, normal, codimension 1 Lie subgroup K of G. Denote the natural projection map by $\pi: G \to K \setminus G$, and the Lie algebra of K by \mathfrak{K} .

Let $H: T^*G \to \mathbb{R}$ be a natural left-invariant Hamiltonian and let $h = H|_{\mathfrak{g}^*}$ be its restriction to the identity. Suppose (g(t), p(t)) follows the hybrid flow $\varphi_t^{\mathcal{H}}$ and let $\zeta(t) = (\ell_{g(t)})^*p(t)$. Let $\sigma: K \setminus G \to G$ be a local section, let $q \in K \setminus G$, and let $\Delta \zeta \in \operatorname{Ann}(\mathfrak{K})$ be such that $h(\zeta) = h(\zeta + \Delta \zeta)$. Then, the equations of motion can be written as

(3.19)
$$\begin{cases} \dot{\zeta} = \operatorname{ad}_{dh}^* \zeta, \\ \dot{q} = d\pi_{\sigma(q)} (\ell_{\sigma(q)})_* \zeta^{\sharp}, \\ \end{cases} \quad q \notin \pi(\Sigma),$$

(3.20)
$$\begin{cases} \zeta \mapsto \zeta + \Delta \zeta, \\ q \mapsto q, \end{cases} \qquad q \in \pi(\Sigma)$$

Remark 3.23. In classical Lie-Poisson reduction, the space can be reduced from T^*G to \mathfrak{g}^* . In the hybrid case, the reduction stops at $\mathfrak{g}^* \times (K \setminus G)$. This extra term is used to determine whether or not an impact occurs. *Proof.* What needs to be shown is that 1) the dynamics on $q \in K \setminus G$ are well-defined, i.e. the choice of section does not matter and 2) the reduced hybrid equations (3.19) and (3.20) do not depend on $g \in G$.

We begin with the first part. There are two components: the discrete and the continuous equations. For the continuous part, the equations of motion are given as in Proposition 3.15

$$\begin{cases} \dot{\zeta} = \operatorname{ad}_{dh}^* \zeta; \\ \dot{g} = (\ell_g)_* \zeta^{\sharp}, & \text{if } g \notin \Sigma. \end{cases}$$

As $K \leq G$ is a subgroup that is closed in the topology of G, there exists a unique manifold structure such that $\pi: G \to K \setminus G$ is a smooth surjective submersion [33]. Therefore, for a smooth curve $g(t) \in G$, we have that $q(t) \coloneqq \pi(g(t)) \in K \setminus G$ is a smooth curve of equivalence classes. Additionally, as $K \leq G$ has codimension 1, the quotient manifold is a 1-dimensional manifold.

For a curve in $g(t) \in G$ satisfying $\dot{g} = (\ell_g)_* \zeta^{\sharp}$, the projected curve, $q = \pi(g)$, satisfies

(3.21)
$$\frac{d}{dt}q(t) = \frac{d}{dt}\pi\left(g(t)\right) = d\pi_{g(t)}\left(\frac{d}{dt}g(t)\right) = d\pi_{g(t)}\left(\ell_{g(t)}\right)_*\zeta^{\sharp}.$$

For all $k \in K$ and $g \in G$, $\pi(kg) = \pi(g)$, so differentiating $\pi(g)$ yields

$$\left(\ell_k\right)^* d\pi_{kg} = d\pi_g$$

Let $\sigma: K \setminus G \to G$ be a smooth (local) section. Choosing a local section is equivalent to choosing a representative $g_0(t)$ of the equivalence class q(t). Any curve $g(t) \in G$ can be written as $g(t) = k(t)\sigma(\pi(g(t)))$ for some curve $k(t) \in K$ as in Figure 11. Calling $g_0(t) \coloneqq \sigma(\pi(g(t)))$, the dynamics on $q = \pi(g)$ in equation (3.21) are

$$\dot{q} = d\pi_{k(t)g_0(t)} \left(\ell_{k(t)g_0(t)}\right)_* \zeta^{\sharp} = \left(\ell_{k(t)}\right)^* d\pi_{k(t)g_0(t)} \left(\ell_{g_0(t)}\right)_* \zeta^{\sharp} = d\pi_{g_0(t)} \left(\ell_{g_0(t)}\right)_* \zeta^{\sharp}.$$

Note that the previous equation does not depend on g(t) anymore, but only on q(t) and on the choice of local section σ . Now, suppose we choose a different section $\tilde{g}_0(t) = \tilde{\sigma}(\pi(g(t)))$. Then, there exists $\tilde{k}(t) \in K$ such that $\tilde{g}_0(t) = \tilde{k}(t)g_0(t)$ (see Figure 10). This is because both g_0 and \tilde{g}_0 are lifts of q(t). Comparing these two different representatives yields

$$d\pi_{\tilde{g}_{0}(t)} \left(\ell_{\tilde{g}_{0}(t)}\right)_{*} \zeta^{\sharp} = d\pi_{\tilde{k}(t)g_{0}(t)} \left(\ell_{\tilde{k}(t)g_{0}(t)}\right)_{*} \zeta^{\sharp}$$
$$= \left(\ell_{\tilde{k}(t)}\right)^{*} d\pi_{g_{0}(t)} \left(\ell_{g_{0}(t)}\right)_{*} \zeta^{\sharp}$$
$$= d\pi_{g_{0}(t)} \left(\ell_{g_{0}(t)}\right)_{*} \zeta^{\sharp}.$$

where the last equality comes from the fact that $\pi(kg) = \pi(g)$ for all $k \in K$ and $g \in G$. Therefore, the equations of motion are independent on the choice of section.

In order to find the impact map we use the corner conditions (2.4) and adapt them for the case of Lie groups.



Figure 11: Schematic drawing of Impact Lie-Poisson reduction, where g(t) represents the true dynamics of the system, which differs from $g_0(t)$ by some subgroup element k(t) for every t. Since Σ projects down to one point $[g_0] \in K \setminus G$, it is sufficient to keep track of q(t) in order to figure out when impacts happen.

Since the impact surface is time independent $\delta t = \mathbb{R}$ and we get the energy conservation condition: $H^+ = H^-$. In particular, $H^+|_{\mathfrak{g}} = H^-|_{\mathfrak{g}}$ and $h^+|_{\mathfrak{g}} = h^-|_{\mathfrak{g}}$. For the momentum equation, we left translate to the identity

$$((\ell_g)^*(p^+ - p^-))((\ell_{g^{-1}})\delta g) = 0 \implies (\zeta^+ - \zeta^-) (\ell_{g^{-1}})_* \delta g = 0$$

where $\zeta^{\pm} := \ell_a^* p^{\pm}$.

From the corner conditions we know that δg has to lie in $T_g \Sigma$ and since Σ is a right coset $\delta g = (r_{g_0})_* \dot{k}$ for $\dot{k} \in T_k K$. Therefore,

$$(\ell_{g^{-1}})_*\delta g = (\ell_{g_0^{-1}})_*(\ell_{k^{-1}})_*(r_{g_0})_*\dot{k} = (\mathrm{Ad}_{g_0^{-1}})_*(\ell_{k^{-1}})_*\dot{k}.$$

Since K is a subgroup $(\ell_{k^{-1}})_*\dot{k} = \widetilde{\delta k} \in \mathfrak{K}$. Additionally, K is normal, so its lie algebra is closed under the adjoint, i.e., $\operatorname{Ad}_g \xi \in \mathfrak{K}$ for any $g \in G$ and $\xi \in \mathfrak{K}$. In particular this holds for $g = g_0$ and for $\xi = \widetilde{\delta k}$ so $(\operatorname{Ad}_{g_0^{-1}})(\ell_{k^{-1}})_*\dot{k} := \delta k \in \mathfrak{K}$. Since δg can be any vector in $T_g \Sigma$, δk spans \mathfrak{K} .

For the q variable $g^+ = g^-$ implies that $\pi(g^+) = \pi(g^-)$, so $q^+ = q^-$. Hence we obtain the impact map

$$\begin{cases} \zeta^+ = \zeta^- + \Delta \zeta \\ q^+ = q^- \end{cases} \quad \text{for all } \Delta \zeta \in \operatorname{Ann}(\mathfrak{K}) \text{ such that } h(\zeta^+) = h(\zeta^-) \end{cases}$$

The impact equation on $K \setminus G$ are identity, independent of the local section chosen for the continuous part. Moreover, the trajectory in q(t) is continuous, as shown in Figure 4. Impacts occur when $g(t)g_0(t)^{-1} \in K$ i.e. when the projection q(t) passes through the equivalence class $[g_0]$. This process is also independent on g(t).

Remark 3.24. If the Hamiltonian is natural, ζ^{\sharp} is defined such that $\coprod(\zeta^{\sharp}, \cdot) = \zeta$. However, ζ^{\sharp} is well-defined even if the Hamiltonian is not natural as long as the fiber derivative is invertible. The formula is given by $\zeta^{\sharp} := \mathbb{F}L^{-1}(\zeta)$.

Remark 3.25. The proof requires the existence of a *local* section, and the choice of the section $\sigma: K \setminus G \to G$ does not matter. As such, no global section needs to exist.

In the following we give an explicit expression for $\Delta \zeta$.

Corollary 3.26. If $H: T^*G \to \mathbb{R}$ is a natural left invariant Hamiltonian of the form $H(g, p) = \frac{1}{2} \prod^{-1}(p, p)$ where \prod is a Riemannian metric, and if K is a connected, closed, and codimension 1 subgroup of G, then there exists a smooth function $s: G \to \mathbb{R}$ such that $\Sigma = s^{-1}(\{0\})$ and $\Delta \zeta$ is given by

(3.22)
$$\Delta \zeta = -\frac{2ds_{g_0}((\ell_{g_0^{-1}})_*(\zeta^{-})^{\sharp})}{(\ell_{g_0}^*)ds_{g_0}((\ell_{g_0^{-1}})_*\nabla s_{g_0})}(\ell_{g_0})^*ds_{g_0}.$$

Proof. Since K is closed and connected, $K \setminus G$ is a 1-dimensional connected manifold, and $\pi: G \to K \setminus G$ is a surjective submersion, and hence, differentiable. Thus $K \setminus G$ must be isomorphic to either \mathbb{R} or \mathbb{S}^1 . Without loss of generality, assume $K \setminus G \cong \mathbb{R}$ and let $j: K \setminus G \to \mathbb{R}$ be the isomorphism. Define s to be

$$s: G \to \mathbb{R}: g \mapsto j(\pi(g)) - j(\pi(g_0))$$

Note that $g \in \Sigma$ if and only if $[g] = [g_0]$, which implies that $s(g) = j(\pi(g)) - j(\pi(g_0)) = 0$. Moreover, since both j and π are differentiable, s is also differentiable.

From Proposition 3.15 we get

$$\Delta \zeta = -\frac{2\zeta^{-}((\ell_{g^{-1}})_* \nabla s_g)}{ds_g(\nabla s_g)} (\ell_g)^* ds_g.$$

This formula is already close to the desired result, but it depends on g. Our goal is to use the right coset property in order to eliminate the g dependency.

From Lemma 3.21 there exists an $\alpha_{u,g} \in \mathbb{R}$ such that $\ell_g^* ds_{ug} = \alpha_{u,g} ds_u$. Let $g = kg_0 \in \Sigma$, $u = k \in G_{g_0}$ and $g = g_0 \in \Sigma$. Then, $\ell_k^* ds_{kg_0} = \alpha_{k,g_0} ds_g$ and similarly $(\ell_{k^{-1}})_* \nabla s_{kg_0} = \alpha_{k,g_0}^{-1} \nabla s_{g_0}$. Moreover, $\ell_{kg_0}^* = (\ell_k \circ \ell_{g_0})^* = \ell_{g_0}^* \ell_k^*$, and analogously $(\ell_{(kg_0)^{-1}})_* = (\ell_{g_0}^{-1})_* (\ell_{k^{-1}})_*$. Putting these two observations together into the formula for $\Delta \zeta$ we obtain

$$\Delta \zeta = -\frac{2\zeta^{-}((\ell_{g_{0}^{-1}})_{*}\alpha_{k,g_{0}}^{-1}\nabla s_{g})}{\ell_{g_{0}}^{*}\alpha_{k,g_{0}}ds_{g_{0}}(\alpha_{k,g_{0}}^{-1}(\ell_{g_{0}^{-1}})_{*}\nabla s_{g_{0}})}(\ell_{g_{0}})^{*}\alpha_{k,g_{0}}ds_{g_{0}}.$$

Since α_{k,g_0} is a nonzero constant, we can pull it out of the pullback and pushforward, and since it appears in both the numerator and the denominator, we are left with

$$\Delta \zeta = -\frac{2\zeta^{-}((\ell_{g_{0}^{-1}})_{*}\nabla s_{g_{0}})}{\ell_{g_{0}}^{*}ds_{g_{0}}((\ell_{g_{0}^{-1}})_{*}\nabla s_{g_{0}})}(\ell_{g_{0}})^{*}ds_{g_{0}}.$$

The final result follows from the same derivation as the end of the proof of Proposition 3.15.

Remark 3.27. The normality of the subgroup K is crucial for Hybrid Lie-Poisson reduction. If dropped, $\Delta \zeta$ might not be independent of g, and hence, the reset equations cannot be reduced. This is illustrated in the following counterexample: Consider the Lie group G = Aff(1) and the subgroup K of G, which are the following sets closed under matrix multiplication

$$G = \left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} : a \in \mathbb{R} \setminus \{0\} \text{ and } b \in \mathbb{R} \right\} \text{ and } K = \left\{ \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} : a \in \mathbb{R} \right\}.$$

K a codimension 1 closed subgroup of G; however, K is not closed under the adjoint, and hence, not normal. Next, choose $g_0 = \begin{pmatrix} a_0 & b_0 \\ 0 & 1 \end{pmatrix}$ so then for $g \in \Sigma$ and $\delta g \in T_g \Sigma$, we have

$$\delta g = \begin{pmatrix} \dot{\alpha} a_0 & \dot{\alpha} b_0 \\ 0 & 0 \end{pmatrix} \quad \text{for } \begin{pmatrix} \dot{\alpha} & 0 \\ 0 & 0 \end{pmatrix} \in \mathfrak{K}.$$

However,

$$(\ell_{g^{-1}})_*\delta g = \frac{\dot{\alpha}}{\alpha} \begin{pmatrix} 1 & b_0/a_0 \\ 0 & 0 \end{pmatrix} \notin \mathfrak{K}.$$

The dual of the Lie algebra \mathfrak{g} consists of vectors $P = (p_a, p_b)$, and the dual pairing $P(A) = p_a \dot{a} + p_b \dot{b}$ is the usual dot product between P and (\dot{a}, \dot{b}) . Left translation of P is $\ell_C^* P = (cp_a, cp_b)$ where $C = \begin{pmatrix} c & d \\ 0 & 1 \end{pmatrix}$. With $\zeta = \ell_A P$, the left translated corner conditions become

$$\left(a(p_a^+ - p_a^-) \quad a(p_b^+ - p_b^-) \right) \begin{pmatrix} \dot{\alpha}/\alpha \\ \dot{\alpha}b_0/\alpha a_0 \end{pmatrix} = a(p_a^+ - p_a^-) \frac{\dot{\alpha}}{\alpha} + \frac{\dot{\alpha}b_0}{\alpha a_0} a(p_b^+ - p_b^-).$$

Now use the fact that $a = \alpha a_0$ and that the above has to be 0 for any α due to the corner conditions to obtain

$$a_0(p_a^+ - p_a^-) + b_0(p_b^+ - p_b^-) = 0$$

This equation does not depend on α anymore which is good, but it also implies that $(p_a^+ - p_a^-)/(p_b^+ - p_b^-) = -b_0/a_0$. With $p_a^+ - p_a^- = \beta$, $\Delta \zeta = (\beta, -b_0 a \beta/a_0)$ which depends on a. So the reset map is not independent of the group element.

4. Applications. The theory developed above enables us to speed up the computation of the Frobenius-Perron operator for hybrid systems. First, solving the partial differential equation (3.6) is faster than computing the Frobenius-Perron operator using the definition (3.1). For the latter one would need to sample a large number of initial points from the given distribution, and compute their trajectories for a long period of time. This endeavour suffers greatly under curse of dimensionality. Second, by doing impact reduction on the hybrid system at hand we can reduce the dimensionality, which further decreases the computation memory and time.

In the following we illustrate the theory, and its applicability on a number of diverse examples. First, we consider the classical examples of the bouncing ball and the nonholonomic Chaplygin sleigh. Next, we provide analytical results in the case where the configuration space is $\operatorname{GL}_n(\mathbb{R})$ and the impact surface is the normal subgroup $\operatorname{SL}_n(\mathbb{R})$. Finally, we employ our method to analyse an SIR model of an epidemic, with event-based human intervention. The numerical simulations are done using a semi-Lagrangian discretization scheme. At each grid point a characteristic is run backward for some fixed Δt , and the value of the solution is updated by using the nearest grid point to the foot of the characteristic. Although this method is quite crude, the simulations are much faster than the ones that use the definition based approach to computing the Frobenius-Perron operator (see baseline in [9]). All simulations are performed in matlab on a MacBook Pro (10 cores and 32 GB memory). The code can be found at https://github.com/Mary199810/Hybrid-transfer-operators.git.

4.1. The bouncing ball. The canonical pedagogical example is the bouncing ball with elastic impacts. The equations of motion in the Hamiltonian framework are

$$\dot{x} = \frac{p}{m}, \quad \dot{p} = -mg,$$

impacts occur when the ball strikes the ground (x = 0), and the reset map is $\Delta(0, p) = (0, -p)$. Thus when the ball hits the ground, the sign momentum is reversed since no dissipation occurs during impact. The hybrid continuity equation, (3.6), is

(4.1)
$$\begin{cases} \frac{\partial u}{\partial t} + \frac{p}{m} \frac{\partial u}{\partial x} - mg \frac{\partial u}{\partial p} = 0, & \text{for } x \neq 0; \\ u(t^+, 0, p) = u(t^-, 0, -p), & \text{otherwise.} \end{cases}$$

Consider an initial distribution $\rho(x, p) = e^{-(x-1)^2 - p^2}$ and let m = 1 and g = 1. The results of the simulations are shown in Figure 12. The simulation time was 252.213084 seconds.

4.2. The Chaplygin sleigh with angle impacts. The Chaplygin sleigh was introduced in the work of Chaplygin [7], and is one of the best studied examples of nonholonomic systems. It describes the motion of a rigid body in the plane supported on a knife edge. The body can move freely without friction in the direction parallel to the edge, and it can rotate around it, but it cannot move perpendicularly. The Chaplygin sleigh model is useful for describing the motion of a car, in particular when discussing parallel parking [28], for motion planning problems of skaters in the plane [30], and for more complex robotic systems with wheels [29]. The natural state space is SE₂ with coordinates (x, y, θ) where x and y are Cartesian coordinates that represent the position of the sleigh within the plane and θ is the sleigh's orientation. The Lie algebra is composed of matrices of the form A as stated in (4.2). For this system, the Hamiltonian is $H = \frac{1}{2}p^T M^{-1}p$, where $p = (p_x, p_y, p_{\theta})$ and

(4.2)
$$A = \begin{pmatrix} 0 & \dot{\theta} & \dot{x} \\ -\dot{\theta} & 0 & \dot{y} \\ 0 & 0 & 0 \end{pmatrix} \qquad M = \begin{pmatrix} m & 0 & -ma\sin\theta \\ 0 & m & ma\cos\theta \\ -ma\sin\theta & ma\cos\theta & I + ma^2 \end{pmatrix}.$$

The Hamiltonian is left invariant, i.e., $H(g,p) = H(e,g^{-1}p)$ for any $g \in SE_2$, where e is the identity matrix in SE₂. It is easier to work in the Lagrangian formulation with $L = \frac{1}{2}\dot{g}M\dot{g}$. Then left translation to identity is

$$\zeta = g^{-1}\dot{g} = \begin{pmatrix} \dot{x}\cos\theta + \dot{y}\sin\theta\\ -\dot{x}\sin\theta + \dot{y}\cos\theta\\ \dot{\theta} \end{pmatrix}$$



(a) The initial density is centered around (1, 0).



(c) Around t = 5 the mass starts to concentrate in stripes around the Zeno point (0, 0).



(b) Immediately after the start of the simulation, at t = 1, the mass is attracted towards the impact surface x = 0.



(d) As time becomes large, this effect is stronger.

Figure 12: Snapshots of the evolution of an initial density under the hybrid transfer operator PDE (4.1) corresponding the the elastic bouncing ball.



Figure 13: Schematic drawing of the Chaplygin sleigh, where a is the distance from the center of mass to the knife edge.

The first component is the velocity parallel to the edge, the second is the perpendicular velocity and the third is the angular velocity of the sleigh. The dimension can further be reduced by taking the nonholonomic constraint $-\dot{x}\sin\theta + \dot{y}\cos\theta = 0$ into consideration. This is exactly the second component of ζ . Let $v = \dot{x}\cos\theta + \dot{y}\sin\theta$ and $\omega = \dot{\theta}$. Then the equations of motion can be written in the (v, ω) coordinates as

$$\begin{cases} \dot{v} = a\omega^2 \\ \dot{\omega} = -\frac{ma}{I + ma^2}v\omega \end{cases}$$

Now assume that the impacts are angle dependent, i.e. $\Sigma = \{(x, y, \theta) \in SE_2 : \theta \in \{\pm \theta_0\}\}$ which is a right coset of \mathbb{R}^2 , who is in turn a codimension 1 subgroup of SE₂. In the example of a simplified vehicle, this is equivalent to having the wheel turn suddenly after reaching a critical angle. The reset map in the reduced coordinates is:

$$\begin{cases} v^+ = v^-\\ \omega^+ = -\omega^- \end{cases}$$

Apart from v and ω we need another variable to determine where impacts happen. This is the role of q in Theorem 3.22. $\mathbb{R}\setminus SE_2 \approx \mathbb{S}^1$ so $q = \theta \in \mathbb{S}^1$. For the local section $\sigma(\theta) = (0, 0, \theta)$, the equations of motion (3.16) are equivalent to $\dot{\theta} = \omega$ and the reset map for θ is identity, so $\theta^+ = \theta^-$. Putting everything together, the reduced hybrid dynamics are given by

Continuous: Discrete:

$$\dot{v} = a\omega^2$$
 $v^+ = v^-$
 $\dot{\omega} = -\frac{ma}{I + ma^2}v\omega$ $\omega^+ = -\omega^-$
 $\dot{\theta} = \omega$ $\theta^+ = \theta^-$.

Plugging this vector field into (3.6) and using the usual measure on SE₂: $\mu = dv \wedge d\omega \wedge d\theta$, we obtain the PDE for the transfer operator

$$\frac{\partial u}{\partial t} - \frac{\partial u}{\partial v} a\omega^2 - \frac{ma}{I + ma^2} v\omega \frac{\partial v}{\partial \omega} + \omega \frac{\partial v}{\partial \theta} - \frac{ma}{I + ma^2} vu = 0 \qquad \text{if } \theta \neq \theta_0;$$
$$u(t^+, v, -\omega, \theta) = u(t^-, v, \omega, \theta) \qquad \text{if } \theta = \theta_0.$$

The dynamics of the Chaplygin sleigh can be further reduced by using the conservation of energy propriety and restricting to an energy surface. Let E be the constant energy of the sleigh, which can be written in terms of v and ω as

$$E = \frac{1}{2}mv^{2} + \frac{1}{2}(I + ma^{2})\omega^{2},$$

We can express ω in terms of v as $\omega = \pm \sqrt{C_1 - C_2 \omega^2}$ where $C_1 = 2E/(I + ma^2)$ and $C_2 = m/(I + ma^2)$. There are two branches of the solution to $\omega^2 = C_1 - C_2 v^2$. When an impact happens, the branch is changed, leading us to the 2 dimensional system

$$\begin{cases} \dot{v} = aC_1 - aC_2v^2\\ \dot{\theta} = \pm\sqrt{C_1 - C_2v^2} = X(v) \end{cases} \quad \text{if } \theta \neq \theta_0, \quad \text{with} \quad \begin{cases} v^+ = v^-\\ \theta^+ = \theta^- \end{cases} \quad \text{if } \theta = \theta_0; \end{cases}$$



Figure 14: State space for the Chaplygin sleigh restricted to an energy surface. The two halves of the cylinder are stacked on top of each other when displaying $u(t, v, \theta)$, the solution to the Hybrid transfer PDE (3.6).

This is a Filippov system where the vector field changes as we cross the lines $\theta = \pm \theta_0$. The state space looks like two copies of $\mathbb{R} \times \mathbb{S}^1$ glued along the lines $\theta = \pm \theta_0$ as in Figure 14.

Moreover, the PDE for the Frobenius-Perron operator is

$$\frac{\partial u}{\partial t} + a(C_1 - C_2 v^2) \frac{\partial u}{\partial v} + X(v) \frac{\partial u}{\partial \theta} + 2aC_2 vu = 0$$
$$u(t^+, \omega, \pm \theta_0) = u(t^-, -\omega, \pm \theta_0)$$

In the plots below (Figure 15), the two copies of $\mathbb{R} \times \mathbb{S}^1$ are identified and the value of the solution is the sum of the value on each copy.

4.3. Matrix groups. Consider the case where the configuration space is $Q = G = \operatorname{GL}_n(\mathbb{R})$, with impact surface $\Sigma = \operatorname{SL}_n(\mathbb{R})$. The Lie algebra of G is $\mathfrak{gl}_n(\mathbb{R}) \cong \mathbb{R}^{n \times n}$ with Lie bracket [A, B] = BA - AB and structure coefficients C_{ij}^k for $i, j, k \in \{1, \ldots, n^2\}$. The dual of the Lie algebra consists of matrices $p \in \mathbb{R}^{n \times n}$ with the dual pairing given by $\langle p, \xi \rangle = \sum_{ij} p_{ij} \xi_{ij}$. The pullback of left translation on dual vectors can be written in terms of matrix multiplication as $\ell_A^* p = A^T p$.

Let $H(A, P) = \frac{1}{2} \operatorname{tr}(A^T P P^T A)$. This Hamiltonian is left-invariant as $H(I_n, A^T P) = H(A, P)$. Throughout this section I_n denotes the identity in $\operatorname{GL}_n(\mathbb{R})$. Let $\zeta := A^T P \in \mathfrak{gl}_n^*$ and denote by $h: \mathfrak{gl}_n^* \to \mathbb{R}$ the Hamiltonian restricted to the dual of the Lie algebra; in coordinates $h(\zeta) = \frac{1}{2} \operatorname{tr}(\zeta \zeta^T)$. To determine the continuous component of the reduced motion, we view $\zeta \in \mathfrak{gl}_n^*$ as a row vector in \mathbb{R}^{n^2} and $X, Y \in \mathfrak{gl}_n$ as column vectors in \mathbb{R}^{n^2} .

The coadjoint representation can be written in terms of the structure coefficients as

$$\mathrm{ad}_X^*(\zeta)(Y) = \zeta([X,Y]) = C_{ij}^k X^i Y^j \zeta_k \implies (\mathrm{ad}_X^*(\zeta))_j = C_{ij}^k X^i \zeta_k$$

Therefore the continuous equations of motion are $\dot{\zeta}_i = C_{ij}^k (\zeta^T)^i \zeta_k$.

For the discrete jumps, let $\Sigma = \mathrm{SL}_n(\mathbb{R})$ be the impact surface, and note that $\mathrm{SL}_n(\mathbb{R}) = s^{-1}(0)$, where $s(A) = \det(A) - 1$. Let $\mathrm{adj}(A)$ denote the coadjoint of A, so $ds|_A = \mathrm{adj}(A)^T$.



(a) The initial density is $\rho(v,\theta)=e^{-v^2-\theta^2}$





(b) Immediately, the mass is pulled towards the right side, which corresponds to the stable fixed point $v^* = \sqrt{C_1/C_2}$.



(c) A sharp peak begins to form in the upper right corner. After an impact, θ rapidly decreases so some of the mass gathers around lower θ values.

(d) After a longer time, a delta-like peak forms in the upper right corner.

Figure 15: Evolution of an initial density under the reduced hybrid Chaplygin sleigh equations of motion.

Then, since $\Sigma = \operatorname{SL}_n(\mathbb{R}) \cdot I_n$ and $g_0 = I_n$, the formula (3.22) for this case gives

$$\Delta \zeta = -\frac{2}{n} \operatorname{tr}(\zeta) I_n.$$

Remark 4.1. Let $f(\zeta) = \operatorname{tr}(\zeta)$. Then $df = I_n d\zeta$ which commutes with any dh for any Hamiltonian h. Thus, the trace is a Casimir for the bracket on \mathfrak{gl}_n^* . Moreover, across impacts $\operatorname{tr}(\zeta^+) = -\operatorname{tr}(\zeta^-)$.

The extra variable that keeps track of whether impact occur is $q = \det(A) - 1$. Its equations of motion are given by (3.19)

$$\begin{cases} \dot{q} = (q+1)\mathrm{tr}(\zeta) \\ q^+ = q^- \end{cases}$$



Figure 16: Sample trajectories of the 2-dimensional system (4.3) which determines exactly when impacts happen for *n*-dimensional Lie-Poisson impact system with $G = \operatorname{GL}_n$, and $\Sigma = \operatorname{SL}_n$. Initially, $\operatorname{tr}(\zeta) = -2$, and $q = \det(A) - 1$ is sampled uniformly from [-3, 3].

The q variable system can also be solved exactly, using the fact that $tr(\zeta) = C$ is always constant; we obtain $q(t) = q_0 e^{Ct} - 1$.

Therefore, to determine when impacts occur we need to keep track of two scalars: the trace of the lie algebra element, and the determinant of the group element. Independently on the dimension of the configuration space, impacts are completely determined by

(4.3)
$$\begin{cases} q(t) = q_0 e^{Ct} - 1 \\ C(t) = C \end{cases} \quad \text{if } q \neq 0; \qquad \begin{cases} q^+ = q^- \\ C^+ = -C^- \end{cases} \quad \text{if } q = 0.$$

No matter the initial $tr(\zeta)$ and det(A), there is at most one impact on any trajectory (see Figure 16).

Let n = 2, and denote the components of dh by $(\alpha^1 \alpha^2 \alpha^3 \alpha^4)$. Then ad_{dh}^* is a 4×4 matrix depending on dh

$$\mathrm{ad}_{dh}^{*} = \begin{pmatrix} 0 & -\alpha^{3} & \alpha^{2} & 0 \\ -\alpha^{2} & \alpha^{1} - \alpha^{4} & 0 & \alpha^{2} \\ \alpha^{3} & 0 & \alpha^{4} - \alpha^{1} & -\alpha^{3} \\ 0 & \alpha^{3} & -\alpha^{2} & 0 \end{pmatrix}$$

For $h = \frac{1}{2} \operatorname{tr}(\zeta \zeta^T)$, $dh = \sum_i \zeta_{\operatorname{Id}} \zeta_i$ and the equations of motion are

$$(4.4) \qquad (\dot{\zeta}_1 \quad \dot{\zeta}_2 \quad \dot{\zeta}_3 \quad \dot{\zeta}_4) = (\zeta_1 \quad \zeta_2 \quad \zeta_3 \quad \zeta_4) \begin{pmatrix} 0 & -\zeta_3 & \zeta_2 & 0 \\ -\zeta_2 & \zeta_1 - \zeta_4 & 0 & \zeta_2 \\ \zeta_3 & 0 & \zeta_4 - \zeta_1 & -\zeta_3 \\ 0 & \zeta_3 & -\zeta_2 & 0 \end{pmatrix} \\ = (-\zeta_2^2 + \zeta_3^2 \quad (\zeta_1 - \zeta_4)(\zeta_2 - \zeta_3) \quad (\zeta_1 - \zeta_4)(\zeta_2 - \zeta_3) \quad \zeta_2^2 - \zeta_3^2)$$

and discrete transitions follow the rule $\zeta^+ = \begin{pmatrix} -\zeta_4^- & \zeta_2^- & \zeta_3^- & -\zeta_1 \end{pmatrix}$.

The system (4.4) can be solved exactly! Let $\zeta_1 + \zeta_4 = \operatorname{tr}(\zeta) = C$ and $\zeta_2 - \zeta_3 = D$. Then, $\dot{C} = \dot{D} = 0$ so C and D are constants. Plugging this back into (4.4), the equations of motion can be reduced to a 2-dimensional linear system

which can be solved analytically to give the following solution to (4.4):

$$\begin{aligned} \zeta_1 &= D^2 t + \zeta_1^0 \cos(2Dt) + \zeta_3^0 \sin(2Dt), \\ \zeta_2 &= D - CDt + \zeta_1^0 \sin(2Dt) + \zeta_3^0 \cos(2Dt), \\ \zeta_4 &= C - D^2 t - \zeta_1^0 \cos(2Dt) - \zeta_3^0 \sin(2Dt), \end{aligned}$$

where $D = \zeta_3^0 - \zeta_2^0$ and $C = \zeta_1^0 + \zeta_4^0$.

Now, consider the problem of finding the flow of some density under the dynamics. For this we need to find (3.6) for $\operatorname{GL}_2(\mathbb{R})$. Given the measure $\mu = d\zeta_1 \wedge d\zeta_2 \wedge d\zeta_3 \wedge d\zeta_4 \wedge dq$, we obtain that the hybrid Jacobian is $\mathcal{J}^X_{\mu} = -1$ and the divergence $\operatorname{div}_{\mu}(X) = \operatorname{tr}(\zeta) = C$. Hence, (3.6) becomes

$$\begin{cases} \partial_t u + (2D\zeta_3 + D^2)\partial_{\zeta_1} u + (2D\zeta_1 - DC)\partial_{\zeta_2} u + (2D\zeta_1 - DC)\partial_{\zeta_3} u & \text{if } q \neq 0; \\ -(2D\zeta_3 + D^2)\partial_{\zeta_4} u + C(q+1)\partial_q u = -Cu & \text{if } q \neq 0; \\ u(t^+, \zeta^+, 0) = -u(t^-, \zeta^-, 0) & \text{if } q = 0. \end{cases}$$

4.4. Modeling the spread of disease with human intervention. Consider an SIR model as in [35], governed by the following equations of motion

$$\begin{cases} \dot{S} = \mu N - \frac{\beta SI}{N} - \mu S\\ \dot{I} = \frac{\beta SI}{N} - \gamma I - \mu I - \delta I\\ \dot{R} = \gamma I - \mu R \end{cases}$$

where μ represents the natural birth and death rate, γ is the recovery rate, β is the rate of infection, and δ is the death rate due to the disease. Now, consider the following scenario: after the number of infected individuals reaches a specific threshold fraction of the population α , the government decides to intervene and treats a fraction f of the infected individuals all at once. After this, the government lets the disease evolve again with its usual dynamics. Mathematically, the intervention is represented by the impact map

$$\begin{cases} S^+ = S^- \\ I^+ = (1-f)I^- \\ R^+ = R^- + fI^- \end{cases}$$

and impact surface $\Sigma = \mathbb{R}_+ \times \{\alpha N\} \times \mathbb{R}_+ \cong \mathbb{R}^2 \subset \mathbb{R}^3$.

We are interested in the long term behaviour of this population. What percent of the individuals survive this disease? What are the best values for α and f under which $\lim_{t\to\infty} N(t)$ is maximized?

One interesting aspect about this example is that the hybrid Jacobian is no longer 1, which is not surprising since the equations do not come from a Hamiltonian.

$$\Delta^* i_X \mu = -(1-f)I\left(\frac{\beta S}{N} - \gamma - \mu - \delta\right) dS \wedge dR = (1-f)\iota_S^* i_X \mu$$

Hence, the hybrid Jacobian $\mathcal{J}^X_{\mu}(\Delta) = 1 - f$, so the hybrid Frobenius Perron-PDE (3.6) becomes

$$\begin{split} \frac{\partial u}{\partial t} + \left(\mu N - \frac{\beta SI}{N} - \mu S\right) \frac{\partial u}{\partial S} + \left(\frac{\beta SI}{N} - (\gamma + \mu + \delta)I\right) \frac{\partial u}{\partial I} \\ &+ (\gamma I - \mu R) \frac{\partial u}{\partial R} - \left(\beta \frac{I - S}{N} + 3\mu + \delta + \gamma\right) u = 0, \qquad I \neq \alpha N; \\ &u(t^+, S, \alpha N(1 - f), R) = (1 - f)u(t^-, S, \alpha N, R), \qquad I = \alpha N. \end{split}$$

Assume we are only interested in the proportion of susceptible, infected and recovered individuals, and not in the total size of the population. Under this assumption we can further reduce the dimension by normalization. Let i = I/N, s = S/N, and r = R/N. Then i + r + s = 1, the hybrid equations of motion become

$$\begin{aligned} \dot{s} &= \mu - \mu s - (\beta - \delta) s i \\ \dot{i} &= \beta s i + \delta i^2 - (\gamma + \mu + \delta) i \end{aligned} \quad \text{if } i \neq \alpha; \qquad \qquad s^+ = s^- \\ i^+ &= (1 - f) i^- \end{aligned} \quad \text{if } i = \alpha, \end{aligned}$$

and the hybrid Frobenius-Perron PDE is

$$(4.5) \qquad \frac{\partial u}{\partial t} + (\mu - \mu s - (\beta - \delta)si)\frac{\partial u}{\partial s} + (\beta si + \delta i^2 - (\gamma + \mu + \delta)i)\frac{\partial u}{\partial i} \\ = (2\mu - \beta(s - i) + \gamma + \delta - 3\delta i)u, \\ u(t^+, s, \alpha(1 - f)) = \frac{-\beta\alpha s + \delta\alpha^2 - (g + \mu + \delta)\alpha}{-\beta\alpha s(1 - f) + \delta\alpha^2(1 - f)^2 - (g + \mu + \delta)\alpha(1 - f)}u(t^-, s, \alpha)$$

Consider an initial density $\rho(s,i) = e^{-(s-0.5)^2 - 10(i-1+s)^2}$. Figure 17 shows the evolution of this density under the PDE given in (4.5). Table 2 shows the values of the parameters used during the simulation. Note that since $i + s = 1 - r \leq 1$ the configuration space for this model is the lower triangle of $[0, 1] \times [0, 1]$.

Parameter
$$\beta$$
 γ δ μ f α Value0.10.10.20.10.30.7

Table 2: Values of the parameters of the SIR model used in the simulations shown in Figure 17.

Although the simulations show that the percentage of infected population is likely to be low, this does not imply that the intervention was successful. It might be the case that the size of the population decreases so much, that the individuals are mostly recovered. In particular for initial percentage of infected individuals greater than 0.7 the population quickly becomes totally infected and the number of individuals N decreases drastically. In the future we would like to study in detail the entire 3 dimensional dynamics and the effect of the disease on the population count.

5. Conclusions. This paper contributes to the theoretical understanding of hybrid systems in two ways. First, it presents a continuity equation for the Frobenius-Perron operator (Theorem 3.12). This simplifies the problem of computing the time evolution of a given density under



along the line s + i = 1.

the population becomes infected less than 20%. and eventually dies.



ulation, which is concentrated number of infected individuals by acerbated and we obtain a pop-30% when it reaches 70\%. If I is ulation where the percentage of initially greater than 70%, then infected individuals is likely to be

Figure 17: Evolution of initial density $\rho = e^{-(s-0.5)^2 - 10(i-1+s)^2}$ under the hybrid transfer PDE (4.5) corresponding to the SIR model with human intervention.

the underlying dynamics, to the quest of finding the solution of a partial differential equation. Second, it develops a theory of reduction for hybrid systems where the state space is a Lie group (Theorem 3.22). This is particularly useful in high dimensional systems, where even solving equation (3.6) becomes difficult.

We identify several future research directions. Although the Hybrid Frobenius-Perron PDE offers great theoretical insight into the analysis of the transfer operator, the numerical methods employed for solving it should be further improved. A particular issue comes into play due to the fact that characteristics do not always stay within the predefined grid. In our current approach, when that happens, we integrate over the entire characteristic up to time 0. However, this increases the computation time and becomes problematic when a large number of the characteristics suffer from this issue. Therefore, better methods to deal with this situation are needed. Moreover, different approaches for solving (3.6) such as finite differences and finite volumes should be studied.

In order for Hybrid Lie-Poisson reduction to be valid, the impact surface needs to be the right coset of a codimension 1 normal Lie subgroup of the configuration space. These proprieties restrict the types of hybrid systems we can consider for reduction. In the future, we wish to study the instances where the conditions mentioned above are not satisfied. Even if reduction to n+1 dimensions might not be plausible without these conditions, there might still be some k < n such that the system can be reduced to n + k dimensions.

Finally, we are interested in the inverse problem, where we would like to find the underlying hybrid dynamics, using the evolution of a given density under the Frobenius-Perron operator. Suppose we can sample points from the flow of some initial distribution at any given time. These points will give an approximation to the solution of the (3.6). Can we tell whether the dynamics is hybrid just by looking at these points? Can we find an analytical expression for

the ODEs that generate this dynamics?

Appendix A. Proof of the transverse property.

Lemma A.1 (Transverse property). Let $\Delta^X_*: TM|_{\mathcal{S}} \to TM|_{\Delta(\mathcal{S})}$ be the augmented differential and let $\iota_{\mathcal{S}}: \mathcal{S} \hookrightarrow M$ and $\iota_{\Delta(\mathcal{S})}: \Delta(\mathcal{S}) \hookrightarrow M$ denote the respective inclusion maps. Then,

(A.1)
$$\det_{\mu} \Delta_*^X = \det_{\iota_{\mathcal{S}}^* i_X \mu \to \iota_{\Delta(\mathcal{S})}^* i_X \mu} \det_{\Delta(\mathcal{S})} \iota_{X\mu}.$$

Proof. Fix $x \in S$ and let $v^i \in T_x M$ for i = 1, ..., n where $n = \dim M$. Then, we can decompose v^i into $v^i = u^i + \alpha^i X_x$ where $u^i \in T_x S$ and $\alpha^i \in \mathbb{R}$. By the definition of det_{μ}, as stated in Definiton (3.4), we know that

(A.2)
$$\left(\det_{\mu} \Delta_{*}^{X}\right) \mu(v^{1}, \dots, v^{n}) = \mu_{\Delta(x)} \left(\Delta_{*}^{X} v^{1}, \dots, \Delta_{*}^{X} v^{n}\right)$$

Expanding v^i through the mentioned decomposition: $v^i = u^i + \alpha^i X_x$

$$= \mu_{\Delta(x)} \left(\Delta_*^X \left(u^1 + \alpha^1 X_x \right), \dots, \Delta_*^X \left(u^n + \alpha^n X_x \right) \right)$$

Since is Δ^X_* a linear map and $\Delta^X_*(X_x) = X_{\Delta(x)}$ by definition

$$= \mu_{\Delta(x)} \left(\Delta_* \left(u^1 \right) + \alpha^1 X_{\Delta(x)}, \dots, \Delta_* \left(u^n \right) + \alpha^n X_{\Delta(x)} \right)$$

By the linearity of k-forms, we can expand the previous expression as follows

(A.3)
$$= \mu_{\Delta(x)} \left(\Delta_* u^1, \Delta_* u^2_x, \dots, \Delta_* u^n \right)$$
$$+ \sum_{k=1}^n (-1)^{k-1} \alpha^k \mu_{\Delta(x)} \left(\Delta_* u^1_x, \dots, \widehat{\Delta_* u^k}, X_{\Delta(x)}, \dots, \Delta_* u^n \right)$$

Since $T_x \mathcal{S}$ is n-1 dimensional and each $u^i \in T_x \mathcal{S}$, the set $\{u^i\}_{i=1}^n$ is linearly dependent. Then, since k-forms with linearly dependent inputs are equal to zero, we know that (A.3) vanishes, so

$$=\sum_{k=1}^{n}(-1)^{k-1}\alpha^{k}\mu_{\Delta(x)}(\Delta_{*}u^{1},\ldots,\widehat{\Delta_{*}u^{k}},X_{\Delta(x)},\ldots,\Delta_{*}u^{n})$$

Rewriting using the fact that $\mu_{\Delta(x)}$ is an alternating product

$$=\sum_{k=1}^{n}(-1)^{k-1}\alpha^{k}\mu_{\Delta(x)}(X_{\Delta(x)},\Delta_{*}u_{x}^{1},\ldots,\widehat{\Delta_{*}u^{k}},\ldots,\Delta_{*}u^{n})$$
$$=\sum_{k=1}^{n}(-1)^{k-1}\alpha^{k}i_{X}\mu_{\Delta(x)}(\Delta_{*}u^{1},\ldots,\widehat{\Delta_{*}u^{k}},\ldots,\Delta_{*}u^{n})$$

Before we solve for $\det_{\mu} \Delta_*^X$, we will rewrite $\mu(v^1, \ldots, v^{n-1})$, the term that $\det_{\mu} \Delta_*^X$ is multiplied by in (A.2). Yet again using the decomposition $v^i = u^i + \alpha^i X_x$ and a process similar to that of rewriting the right-hand side of (A.2), yields

$$\mu(v^{1}, \dots, v^{n-1}) = \sum_{k=1}^{n} \mu_{x}(u^{1}, \dots, \widehat{u^{k}}, \alpha^{k}X_{x}, \dots, u^{n})$$
$$= \sum_{k=1}^{n} (-1)^{k-1} \alpha^{k} i_{X} \mu(u^{1}, \dots, \widehat{u^{k}}, \dots, u^{n}).$$

Now, we can isolate $\det_{\mu} \Delta_*^X$ from all other terms in (A.2),

(A.4)
$$\det_{\mu} \Delta_{*}^{X} = \frac{\sum_{k=1}^{n} (-1)^{k-1} \alpha^{k} i_{X} \mu_{\Delta(x)} (\Delta_{*} u^{1} \dots, \widehat{\Delta_{*} u^{k}}, \dots, \Delta_{*} u^{n})}{\sum_{k=1}^{n} (-1)^{k-1} \alpha^{k} i_{X} \mu(u^{1}, \dots, \widehat{u^{k}}, \dots, u^{n-1})}.$$

We have finished deriving the left-hand side of (A.1). For computing the right-hand side, we must carefully choose which vectors we plug into the form determinant definition. With a later part of the proof in mind, we modify the earlier set of u^k vectors and choose

$$\begin{cases} \{\alpha^{k}u^{1}, \dots, \widehat{u^{k}}, \dots, u^{n}\} & \text{ for } k \in \{2, 3, \dots, n\}; \\ \{\alpha^{1}u_{x}^{2}, u_{x}^{3}, \dots, u^{n}\} & \text{ for } k = 1, \end{cases}$$

both of which we will denote by the top case since the proofs are identical. The reasoning behind this choice becomes apparent in (A.5). Now computing the the right-hand side of (A.1),

$$\begin{aligned} \det(\Delta_*) & \alpha^k i_X \mu(u^1, \dots, \widehat{u^k}, \dots, u^n) \\ &= \det(\Delta_*) & i_S^*(i_X \mu) (\alpha^k u^1, \dots, \widehat{u^k}, \dots, u^n) \\ &= \det(\Delta_*) & i_S^*(i_X \mu) (\alpha^k u^1, \dots, \widehat{u^k}, \dots, u^n) \\ &= \Delta^* i_{\Delta(S)}^*(i_X \mu) (\alpha^k u^1, \dots, \widehat{\Delta_* u^k}, \dots, u^n) \\ &= i_X \mu_{\Delta(x)} (\alpha^k \Delta_* u^1, \dots, \widehat{\Delta_* u^k}, \dots, \Delta_* u^n) \end{aligned}$$
 Since $u^k \in T_x \mathcal{S} \subset T_x M$

$$\begin{aligned} &= \Delta^* i_{\Delta(S)}^*(i_X \mu) (\alpha^k u^1, \dots, \widehat{\Delta_* u^k}, \dots, u^n) \\ &= i_X \mu_{\Delta(x)} (\alpha^k \Delta_* u^1, \dots, \widehat{\Delta_* u^k}, \dots, \Delta_* u^n) \end{aligned}$$
 Definition of the form determinant

$$\begin{aligned} &= i_X \mu_{\Delta(x)} (\alpha^k \Delta_* u^1, \dots, \widehat{\Delta_* u^k}, \dots, \Delta_* u^n) \end{aligned}$$
 Since $\Delta_* u^k \in T_{\Delta(x)} (\Delta(\mathcal{S})) \subset T_{\Delta(x)} M$

Since equality still holds if we multiply both sides by $(-1)^{k-1}$, we can then sum over k to get

(A.5)
$$\det(\Delta_*) \sum_{\substack{i_{\mathcal{S}}^* \alpha^k i_X \mu \to i_{\Delta(\mathcal{S})}^* i_X \mu}} \sum_{k=1}^n (-1)^{k-1} i_X \mu_{\Delta(x)} \left(u^1, \dots, \widehat{u^k}, \dots, u^n \right)$$
$$= \sum_{k=1}^n (-1)^{k-1} \alpha^k i_X \mu_{\Delta(x)} \left(\Delta_* u^1, \dots, \widehat{\Delta_* u^k}, \dots, \Delta_* u^n \right).$$

Finally, isolating the determinant and looking back at previous equations, we get that

$$\det(\Delta_*)_{i_{\mathcal{S}}^*i_{X}\mu \to i_{\Delta(\mathcal{S})}^*i_{X}\mu} = \frac{\sum_{k=1}^{n} (-1)^{k-1} i_{X} \mu_{\Delta(x)} (\Delta_* u^1, \dots, \dot{\Delta}_* (u^{\bar{k}}), \dots, \Delta_* u^n)}{\sum_{k=1}^{n} (-1)^{k-1} i_{X} \mu (u^1, \dots, \dot{u^k}, \dots, u^n)} = (A.4) = \det_{\mu} (\Delta_*^X). \quad \blacksquare$$

Appendix B. Proof of the form determinant and inverse form determinant relation.

Lemma B.1 (Relation between form determinant and inverse form determinant). Given a diffeomorphism $f: M \to N$ and volume forms $\mu \in \Omega^n(M)$ and $\eta \in \Omega^n(N)$, the relation between det f_* and det f_*^{-1} is given by

$$\det_{\eta \to \mu}(f_*) = \frac{1}{\det_{\mu \to \eta}(f_*^{-1}) \circ f} \qquad or \qquad \det_{\mu \to \eta}(f_*^{-1}) = \frac{1}{\det_{\eta \to \mu}(f_*)} \circ f^{-1}.$$

Proof. Let $\mu \in \Omega^n(M)$ and $\eta \in \Omega^n(N)$ be volume forms. Recall that, by definition (3.4), the determinant of $f_*^{-1}: TN \to TM$ is the $C^{\infty}(N)$ function $\det_{\eta \to \mu}(\Delta_*^{-1})$ such that

(B.1)
$$\det_{\eta \to \mu} (f_*^{-1})\eta = (f^{-1})^* \mu.$$

Since μ and η are non-singular, their form determinants are nonzero, so we can divide by the determinants to get

$$\mu = \left(\det_{\mu \to \eta}(f_*)\right)^{-1} f^* \eta \quad \text{and} \quad \eta = \left(\det_{\mu \to \eta}(f_*^{-1})\right)^{-1} (f^{-1})^* \mu$$

As the above functions are inverses of each other, they have the property that $f^{-1} \circ f = \mathrm{Id}_M$. Taking the pullback of both sides of this property gives us

(B.2)
$$(f^{-1} \circ f)^* = f^* \circ (f^{-1})^* = \mathrm{Id}_M^*,$$

Now that we have the pullbacks written out, we get

$$\begin{split} \mu &= f^* \circ (f^{-1})^* \mu & \text{By (B.2)} \\ &= f^* \left(\det_{\eta \to \mu} (f_*^{-1}) \eta \right) & \text{By the definition of the form determinant of } (f^{-1})^* \\ &= \left(\det_{\eta \to \mu} (f_*^{-1}) \circ f \right) (f^* \eta) & \text{Property of pullback on top forms [22, Proposition 14.20]} \\ &= \left(\det_{\eta \to \mu} (f_*^{-1}) \circ f \right) \det_{\mu \to \eta} (f_*) \mu & \text{By the definition of } \det_{\mu \to \eta} f_* \end{split}$$

Finally, dividing gives us

$$\det_{\mu \to \eta}(f_*) = \frac{1}{\det_{\mu \to \eta}(f_*^{-1}) \circ f} \quad \text{and} \quad \det_{\eta \to \mu}(f_*^{-1}) = \frac{1}{\det_{\mu \to \eta}(f_*)} \circ f^{-1}.$$

For unconstrained or holonomically constrained systems with a stationary impact surface, $\mathcal{J}^X_\mu(\Delta) = c^4$. We show this by explicitly computing $\Delta^* i_X \mu = \mathcal{J}^X_\mu(\Delta) \iota^* i_X \mu$ which turns out to be a relatively simple computation since all but one 1-form is equal to zero.

Theorem B.2. Let $H: T^*Q \to \mathbb{R}$ be a natural Hamiltonian and $S \subset Q$ the impact surface. Let Δ be the inelastic impact map given by

$$\Delta(x,p) = \left(x, p - (1+c^2)\frac{p(\nabla h)}{dh(\nabla h)}dh\right).$$

Then, the hybrid Jacobian is simply $\mathcal{J}_{\omega^n}^{X_H}(\Delta) = c^4$.

Proof. To calculate the hybrid Jacobian for inelastic impact systems without constraints, we must isolate $\mathcal{J}^X_{\mu}(\Delta)$ from $\Delta^* i_X \mu = \mathcal{J}^X_{\mu}(\Delta) \iota^* i_X \mu$, where $\mu = \omega^n$ and $\omega = dx^i \wedge dp_i$ is our symplectic form on T^*Q . First, calculating the exterior derivative of the Hamiltonian for later use

$$dH = \frac{\partial H}{\partial x^k} dx^k + \frac{\partial H}{\partial p_k} dx^k$$

$$= \frac{\partial}{\partial x^k} \left(\frac{1}{2} g^{ij} p_i p_j + V(x) \right) dx^k + \frac{\partial}{\partial p_k} \left(\frac{1}{2} g^{ij} p_i p_j + V(x) \right) dp_k$$

$$= \left(\frac{1}{2} \frac{\partial g^{ij}}{\partial x^k} p_i p_j + \frac{\partial V(x)}{\partial x^k} \right) dx^k + \left(\frac{1}{2} g^{ij} \frac{\partial p_i}{\partial p_k} p_j + \frac{1}{2} g^{ij} p_i \frac{\partial p_j}{\partial p_k} \right) dp_k$$

$$= \left(\frac{1}{2} \frac{\partial g^{ij}}{\partial x^k} p_i p_j + \frac{\partial V(x)}{\partial x^k} \right) dx^k + \left(\frac{1}{2} g^{ij} \delta_i^k p_j + \frac{1}{2} g^{ij} p_i \delta_j^k \right) dp_k$$

$$= \left(\frac{1}{2} \frac{\partial g^{ij}}{\partial x^k} p_i p_j + \frac{\partial V(x)}{\partial x^k} \right) dx^k + g^{kj} p_j dp_k$$

Now to calculate $i_X \omega$ for the equation for the hybrid Jacobian. Since $i_X \omega = dH$, we know that

$$\begin{split} i_X \omega^n &= i_X (\omega \wedge \ldots \wedge \omega) \\ &= dH \wedge \omega^{n-1} \\ &= \left(\left(\frac{1}{2} \frac{\partial g^{ij}}{\partial x^k} p_i p_j + \frac{\partial V(x)}{\partial x^k} \right) dx^k + g^{kj} p_j dp_k \right) \\ &\quad \wedge \left(\sum_{k=1}^n dx^1 \wedge dp_1 \wedge \ldots \wedge dx^{\widehat{k} \wedge dp_k} \wedge \ldots \wedge dx^n \wedge dp_n \right) \\ &= \left(\frac{1}{2} \frac{\partial g^{ij}}{\partial x^k} p_i p_j + \frac{\partial V(x)}{\partial x^k} \right) \sum_{k=1}^n dx^1 \wedge dp_1 \wedge \ldots \wedge dx^k \wedge \widehat{dp_k} \wedge \ldots \wedge dx^n \wedge dp_n \\ &\quad + g^{kj} p_j \sum_{k=1}^n dx^1 \wedge dp_1 \wedge \ldots \wedge \widehat{dx^k} \wedge dp_k \wedge \ldots \wedge dx^n \wedge dp_n \end{split}$$

Since we have the quasi-smooth dependence property, we can choose the local coordinates about any point in the gurad so that $S = \{(x^1, \ldots, x^n) \in \mathbb{R}^n : s(x^1, \ldots, x^n) = x^n\}$, which implies $ds = dx^n$. Consequentially, all terms in $i_X \omega$ that contain dx^n are equal to zero on S, so

(B.3)
$$\iota^* i_X \omega^n = g^{nj} p_j dx^1 \wedge dp_1 \wedge \ldots \wedge \widehat{dx^n} \wedge dp_n$$

and from this, we can see that the only term that we will need to calculate is $\partial H/\partial p_n$ since it is the only term that remains in (B.3). Writing the Hamiltonian, we have that

$$H^{+} - V(x) = \frac{1}{2}g\left(p - (1 - c^{2})\frac{p(\nabla s)}{ds(\nabla s)}ds, p - (1 - c^{2})\frac{p(\nabla s)}{ds(\nabla s)}ds\right)$$
$$= \frac{1}{2}g\left(p, p\right) - (1 - c^{2})\frac{p(\nabla s)}{ds(\nabla s)}g\left(p, ds\right) + \frac{1}{2}\left((1 - c^{2})\frac{p(\nabla s)}{ds(\nabla s)}\right)^{2}g\left(ds, ds\right)$$

and taking the derivative of H as needed,

$$\begin{aligned} \frac{\partial}{\partial p_n} (H^+ - V(x)) &= \frac{1}{2} g\left(\frac{\partial p}{\partial p_n}, p\right) + \frac{1}{2} g\left(p, \frac{\partial p}{\partial p_n}\right) - \frac{1}{2} \frac{\partial p(\nabla s)}{\partial p_n} \frac{1 - c^2}{ds(\nabla s)} g\left(p, ds\right) \\ &- \frac{1}{2} (1 - c^2) \frac{p(\nabla s)}{ds(\nabla s)} g\left(\frac{\partial p}{\partial p_n}, ds\right) + \frac{1}{2} \left(1 - c^2\right)^2 \frac{\partial}{\partial p_n} \left(\frac{p(\nabla s)}{ds(\nabla s)}\right)^2 g\left(ds, ds\right) \\ &= \frac{1}{2} g\left(\frac{\partial p}{\partial p_n}, p\right) + \frac{1}{2} g\left(p, \frac{\partial p}{\partial p_n}\right) - \frac{1}{2} \frac{\partial p(\nabla s)}{\partial p_n} \frac{1 - c^2}{ds(\nabla s)} g\left(p, ds\right) \\ &- \frac{1}{2} (1 - c^2) \frac{p(\nabla s)}{ds(\nabla s)} g\left(\frac{\partial p}{\partial p_n}, ds\right) + \frac{1}{2} \left(1 - c^2\right)^2 \frac{\partial}{\partial p_n} \left(\frac{p(\nabla s)}{ds(\nabla s)}\right)^2 ds(\nabla s) \\ &= g\left(\frac{\partial p}{\partial p_n}, p\right) - (1 - c^2) \frac{p(\nabla s)}{ds(\nabla s)} g\left(\frac{\partial p}{\partial p_n}, ds\right) + \frac{(1 - c^2)^2}{ds(\nabla s)} \frac{\partial}{\partial p_n}. \end{aligned}$$

Explicitly writing out the derivatives of (B.4),

$$g\left(\frac{\partial p}{\partial p_n}, p\right) = g^{ij} \frac{\partial p_i}{\partial p_n} p_j = g^{ij} \delta_i^n p_j = g^{nj} p_j = v^n$$
$$\frac{\partial \left(p(\nabla s)\right)}{\partial p_n} = \frac{\partial \left(g^{ij} p_i \partial_j h\right)}{\partial p_n} = g^{ij} \frac{\partial p_i}{\partial p_n} \partial_j h = g^{ij} \delta_i^n \partial_j h = g^{nj} \partial_j h = (\nabla s)^n.$$

Now plugging those derivatives in,

$$\implies \frac{\partial H^+}{\partial p_n} = v^n - (\nabla s)^n \frac{1 - c^2}{ds(\nabla s)} g\left(p, ds\right) + \frac{(1 - c^2)^2}{ds(\nabla s)} (\nabla s)^n p(\nabla s)$$

$$= v^n - (\nabla s)^n \frac{1 - c^2}{ds(\nabla s)} g\left(p, ds\right) + \frac{1 - 2c^2 + c^4}{ds(\nabla s)} (\nabla s)^n p(\nabla s)$$

$$= v^n - (\nabla s)^n \frac{1 - c^2}{ds(\nabla s)} p(\nabla s) + \frac{1 - 2c^2 + c^4}{ds(\nabla s)} (\nabla s)^n p(\nabla s)$$

$$= v^n + \frac{c^4 - 1}{ds(\nabla s)} (\nabla s)^n p(\nabla s)$$

Relating this back to our original equation, we find that

As previously mentioned, $ds = dx^n$ so then $ds(\nabla s) = dx^n(\nabla s) = (\nabla s)^n$. Furthermore, $p(\nabla s) = g^{ij} p_i \partial_j s = ds(v) = dx^n(v) = v^n$. Thus, we can say that

$$\mathcal{J}^X_\mu(\Delta) = 1 + \frac{c^4 - 1}{v^n (\nabla s)^n} (\nabla s)^n v^n$$
$$= 1 + c^4 - 1$$
$$= c^4$$

This result is the simplest of inelastic collisions, but we believe that it is likely to carry over to nonholonmic systems with inelastic collisions so that $\mathcal{J}^X_\mu(\Delta) = c^4$ in general. However, this has yet to be shown and is reserved for future works.

Acknowledgments. We would like to thank Mark Walth and Dr. Richard Rand for insightful conversations and inspiration for this work. We would additionally like to thank Mallory Gaspard for assistance on the numerical aspects of this project. Finally, we would like to thank Cornell University for hosting this REU and making this research possible.

REFERENCES

- A. AMES AND S. SASTRY, Hybrid cotangent bundle reduction of simple hybrid mechanical systems with symmetry, 2006 American Control Conference, (2006), p. 6.
- [2] A. AMES, H. ZHENG, R. GREGG, AND S. SASTRY, <u>Is there life after zeno?</u> taking executions past the breaking (zeno) point, American Control Conference (ACC), 2006 (2006), p. 2652–2657, https://doi.org/10.1109/ACC.2006.1656623.
- [3] A. D. AMES AND S. SASTRY, <u>Hybrid geometric reduction of hybrid systems</u>, in Proceedings of IEEE Conference on Control Applications, IEEE, 2006, pp. 923–929.
- [4] A. BLOCH AND B. BROGLIATO, <u>Nonholonomic mechanics and control</u>, Appl. Mech. Rev., 57 (2004), pp. B3–B3.
- M. BRANICKY, Introduction to Hybrid Systems, Handbook of Networked and Embedded Control Systems. Control Engineering. Birkhäuser Boston, 2005, https://doi.org/10.1007/0-8176-4404-0 5.
- [6] M. BUDIŠIĆ, R. MOHR, AND I. MEZIĆ, <u>Applied koopmanism</u>, Chaos: An Interdisciplinary Journal of Nonlinear Science, 22 (2012), p. 047510, <u>https://doi.org/10.1063/1.4772195</u>.
- [7] S. CHAPLYGIN, On the motion of nonholonomic systems. the reducing-multiplier theorem, Regul. Chaot. Dyn., 13 (2008), pp. 369–376.
- [8] W. CLARK, Invariant Measures, Geometry, and Control of Hybrid and Nonholonomic Dynamical Systems, PhD thesis, University of Michigan, 2020.
- W. CLARK AND A. BLOCH, <u>Invariant forms in hybrid and impact systems and a taming of zeno</u>, Arch Rational Mech Anal, 247 (2023).
- [10] W. CLARK, M. OPREA, AND A. GRAVEN, <u>A geometric approach to optimal control of hybrid and impulse</u> systems, 2021.
- [11] L. J. COLOMBO, M. DE LEÓN, M. E. E. IRAZÚ, AND A. LÓPEZ-GORDÓN, <u>Generalized hybrid momentum</u> maps and reduction by symmetries of forced mechanical systems with inelastic collisions, 2021.
- [12] K. EVAN, H. JASON A., E. TIMOTHY P., AND L. ANTHONY R., Symmetry of energy divergence anomalies associated with the el niño-southern oscillation, Atmosphere, (2018).
- [13] M. E. EYREA IRAZÚ, L. COLOMBO, AND A. BLOCH, <u>Reduction by symmetries of simple hybrid mechanical systems</u>, IFAC-PapersOnLine, 54 (2021), pp. 94–99. 7th IFAC Workshop on Lagrangian and Hamiltonian Methods for Nonlinear Control LHMNC 2021.
- [14] I. GELFAND AND S. FOMIN, Calculus of Variations, Dover, 1991.
- [15] R. GOEBEL, R. G. SANFELICE, AND A. R. TEEL, <u>Hybrid dynamical systems</u>, IEEE Control Systems Magazine, 29 (2009), pp. 28–93, https://doi.org/10.1109/MCS.2008.931718.

- [16] M. GOLUBITSKY, I. STEWART, P.-L. BUONO, AND J. COLLINS, <u>Symmetry in locomotor central pattern</u> generators and animal gaits, Nature, 401 (1999), pp. 693–695.
- [17] M. JERROLD E. AND R. TUDOR, Itroduction to Mechanics and Symmetry, Springer, 1998.
- [18] S. D. JOHNSON, <u>Simple hybrid systems</u>, International Journal of Bifurcation and Chaos, 4 (1994), pp. 1655–1665.
- [19] O. JUAN-PABLO AND R. TUDOR S., <u>Cotangent bundle reduction</u>, Encyclopedia of Mathematical Physics, (2005).
- [20] A. KATOK AND B. HASSEBLATT, Introduction to the modern theory of dynamical systems, Cambridge University Press, 1995.
- [21] A. LASOTA AND M. C. MACKEY, <u>Chaos, Fractals, and Noise</u>, Applied Mathematical Sciences, Springer New York, 1985.
- [22] J. M. LEE, <u>Smooth Manifolds</u>, Springer New York, New York, NY, 2003, pp. 1–29, https://doi.org/10. 1007/978-0-387-21752-9_1, https://doi.org/10.1007/978-0-387-21752-9_1.
- [23] G. MAANI, Z. RAY, Z. MINGHAN, E. L. CHIEN, L. TZU-YUAN, T. WANGLI, L. TINGJUN, L. TIANYI, AND S. JINGWEI, Progress in symmetry preserving robot perception and control through geometry and learning, Front. Robot. AI, (2022).
- [24] J. MARSDEN AND A. WEINSTEIN, Coadjoint orbits, vortices and clebsch variables for incompressible fluids, North Holland Publishing Company, (1983).
- [25] J. E. MARSDEN, S. PEKARSKY, AND S. SHKOLLER, <u>Discrete euler-poincaré and lie-poisson equations</u>, Nonlinearity, 12 (1999), pp. 1647–1662, https://doi.org/10.1088/0951-7715/12/6/314, https://doi.org/ 10.1088%2F0951-7715%2F12%2F6%2F314.
- [26] J. E. MARSDEN AND T. S. RATIU, Introduction to mechanics and symmetry, Springer, 1998.
- [27] A. MAUROY, Y. SUSUKI, AND I. MEZIĆ, Koopman operator in systems and control, Springer, 2020.
- [28] Y. OR AND O. HALVANI, <u>Nonholonomic dynamics of the twistcar vehicle: asymptotic analysis and hybrid dynamics of frictional skidding</u>, Nonlinear Dynamics, (March, 2022).
- [29] K. P. S. AND T. DIMITRIS P., Oscillations, se(2)-snakes and motion control: A study of the roller racer, Dynamical Systems An International Journal, (December, 2001).
- [30] M. RHODES AND V. PUTKARADZE, <u>Trajectory tracing in figure skating</u>, Nonlinear Dyn, 110 (2022), pp. 3031–3044.
- [31] S. SINHA, S. P. NANDANOORI, AND E. YEUNG, Computationally efficient learning of large scale dynamical systems: A koopman theoretic approach, in 2020 IEEE International Conference on Communications, Control, and Computing Technologies for Smart Grids (SmartGridComm), 2020, pp. 1–6, https: //doi.org/10.1109/SmartGridComm47815.2020.9302960.
- [32] S. STERNBERG, Ralph abraham and jerrold e. marsden, foundations of mechanics, Bulletin (New Series) of the American Mathematical Society, 2 (1980).
- [33] L. W. TU, <u>Differential geometry: connections, curvature, and characteristic classes</u>, vol. 275, Springer, 2017.
- [34] D. ZHAO AND S. REVZEN, <u>Multi-legged steering and slipping with low dof hexapod robots</u>, Bioinspiration & Biomimetics, 15 (2020).
- [35] C. ZUZANA, K. JANA, AND R. DMITRII, RACHINSKII ABD SAMIHA C., <u>Global dynamics of sir model</u> with switched transmission rate, Journal of Mathematical Biology, (2020).