Aryabhata and the Construction of the First Trigonometric Table

Vijay A. Singh^{[1](#page-0-0)}

Visiting Professor, Physics Department, Centre for for Excellence in Basic Sciences, Kalina, Santa-Cruz East, Mumbai-400098, India

Aneesh Kumar Dhirubhai Ambani International School, BKC, Bandra East, Mumbai-400098, India

Few among us would know that the first mention of the sine and the enumeration of the first sine table are to be credited to Aryabhata. The method to generate this relies on the sine difference formula which is derived using ingenious arguments based on similar triangles. We describe how this was done. In order to make our presentation pedagogical we take a unit circle and radians instead of the (now) archaic notation in the Aryabhatiya and its commentators. We suggest a couple of exercise problems and invite the enterprising student to try their hands. We also point out that his sine and the second sine difference identities are related to the finite difference calculus we now routinely use to calculate derivatives and second derivatives. An understanding of these trigonometric identities and preparation of the sine table will enable a student to get an appreciation of the path breaking work of Aryabhata.

Keywords: Aryabhata, Sine, Sine Table, Finite Difference Calculus

I Introduction

We routinely look up a table of sines or punch on our calculator to obtain the value of a trigonometric function for a given angle. Little do we realize that a debt of gratitude is owed to the fifth century Indian savant Aryabhata who first showed the way. Aryabhata enumerated the table of sines for closely spaced angles. His methods were based on general trigonometric identities and lend themselves to extensions. The first mention of the sine function is to be found in his (one and only) seminal work the Aryabhatiya (499 CE). Aryabhata describes it in poetic terms as the half bow-string or Ardha-Jya. This is illustrated in Fig. 1. The arrow or saar is related to the cosine function. This is not the only example of poetry intruding into his mathematics. To describe the fact, heretical and revolutionary for those times and for a long time afterwards, that the earth is rotating and the sun is stationary, Aryabahta evokes the tranquil metaphor of a boat floating down the river and the stationary land mass which seems to move backwards. The *Aryabhatiya* is in verse form and there are over a 100 of them. They are written fully respecting the norms of grammar and metre. Aryabhata was a poet as well.

The Aryabhatiya consists of 121 cryptic verses, dense and laden with meaning [\[1,](#page-9-0) [2\]](#page-9-1). The work is divided into 4 parts or *padas*: the *Gitikapada* (13 verses), the mathematics or Ganitapada (33 verses), the Kalkriyapada (25 verses) and the astronomy or Golapada (50 verses). The astronomy is better known. There are two verses in the

¹ emailid: physics.sutra@gmail.com

mathematics Ganitapada describing the solution of the linear Diophantine equation. This has received due recognition. Our focus here is on the trigonometry part in the Ganitapada which in our opinion has suffered neglect and is a pioneering achievement of this savant.

Virtually every major Indian mathematician has commented on the Aryabhatiya (499 CE). Often it is in terms of a formal Bhasya (Commentary). Table I lists some of these. Notable among them is the voluminous work (Maha Bhashya) of the 15th century mathematician Nilakantha Somaiyaji (1444 CE - 1544 CE) who was part of the Kerala school which, beginning with Madhava (1350 CE - 1420 CE), founded the calculus of trigonometric functions. Our presentation relies on Somaiyaji's commentary [\[3\]](#page-9-2) and the works of Kripa Shankar Shukla and K. V. Sarma [\[2\]](#page-9-1) as well as of P. P. Divakaran [\[4\]](#page-9-3).

In this article we describe the trigonometric identities used by Aryabhata to obtain the table of sines. This entails taking the difference of the sine of two closely spaced angles and then taking the second sine difference. We next show that these identities are the same as the finite difference calculus one uses nowadays to numerically obtain the first derivative and second derivative of sine functions. We follow this up with a brief discussion.

The Indian mathematical tradition is largely word based. Results are mentioned and derivations are omitted. The Aryabhatiya (499 CE) with a little over a 100 cryptic, super-compressed verses of dense mathematics is a prime example. Our approach is pedagogical and one which will help the student and teacher to appreciate this pioneering work. Hence we shall take some liberties and describe the great master's work in terms of unit circle and radians instead of degrees and minutes. A couple of exercises in the end will help one get a more hands on understanding.

Table 1: A host of eminent mathematicians have commented on the Aryabhatiya written CE 499. Some, like Brahmagupta (600 CE) or Bhaskara II (1100 CE) have not written a specific commentary but dwelt extensively on it. The above is an abbreviated list of specific commentaries and the dates are approximate. Our main source for this is the work of K. S. Shukla and K. V. Sarma [\[2\]](#page-9-1) which cites around 20 commentaries.

II The Ardha-Jya or Sine Function

As mentioned earlier the Aryabhatiya has some 121 verses out of which 33 verses belong to the mathematics section (Ganitapada). Aryabhata works with, for the first time in the history of mathematics, the sine function. It is the half-chord AP of the unit circle in Fig. 1.

$$
sin(\theta) = \frac{AP}{OA}
$$

= AP (OA = 1)

The circle maybe large or small; correspondingly AP and OA maybe large or small, but the l.h.s. is a function of θ and is **invariant**. Further, all metrical properties related to the circle can be derived using trigonometric functions and the Pythagorean theorem (also described as the Baudhayana or Diagonal theorem [\[4\]](#page-9-3)). For example the geometric property of a triangle can be related to the arcs of the circumscribing circle using the sine and cosine functions. Or the diagonals of the inscribed quadrilateral can be related to its sides. (If a recent proof of the this Pythagorean theorem using the the law of sines holds up to scrutiny then all metrical properties of a circle can by obtained by trigonometry alone [\[5\]](#page-9-4).) By emphasizing the role of this half-chord Aryabhata endowed circle geometry with metrical properties. This alone may qualify him as the founder of trigonometry. But he did more.

Figure 1: The bow superposed on the unit circle. Half the bow string or half chord AP is $sin(\theta)$ as defined by Aryabhata. OP is $cos(\theta)$ while $PX = 1-cos(\theta)$ is called the saar. See text for comments.

It may also help to note that the length of the half-chord AP is very close to the length of the arc AX only when the angle is small (e.g. $\sin(\theta) \approx \theta$ if θ is small and in radians). This was known to Aryabhata in all likelihood by inspection. Similarly the sine of 90° is 1, since then the half-chord is the same as the radius. In the 9th verse of the Ganitapada he uses the property of an equilateral triangle and obtains the sine of 30° as $1/2$.

We pause to note that Aryabhata also states the value of π as 62832/20000 in the 10th verse. This value is 3.416 and he is careful enough to state that this is proximate (Asanna) which means we can obtain better and truer values for π presumably with more effort^2 effort^2 .

²The word *Asanna* or proximate is to be distinguished from *Sthula* which is approximate or roughly equal

III The Difference Formula for Sine and Cosine

The 12th verse of the *Ganitapada* plays a central role in the tabulation of the sine function. It is cryptic and to unravel its meaning we first need to obtain the difference formula for the sine. The presentation below relies on a number of sources: (i) The commentary of Nilakantha Somaiyaji [\[3\]](#page-9-2); (ii) the treatment of Shukla and Sarma [\[2\]](#page-9-1); (iii) and for the sake of ease of understanding we follow Divarkaran [\[4\]](#page-9-3) and take a unit circle as opposed to a radius of [3](#page-3-0)438 by earlier workers³.

Figure 2 depicts a quadrant of the unit circle where $OX = OY = 1$. The arcs XA, XB and XC trace angles θ , $\theta + \phi$ and $\theta - \phi$ respectively. The half-chords AP, BQ and CR are the corresponding sine functions. We drop a perpendicular CS from the circumference onto the half-chord BQ as shown. According to his commentator Nilakantha Somaiyaji [\[3\]](#page-9-2), Aryabhata obtained the relationship between the difference in the trigonometric functions by demonstrating that the two triangles BSC and OP A are similar and then exploiting this ingenuously. We trace his line of reasoning in the Appendix where we derive the following relations

$$
sin(\theta + \phi) - sin(\theta - \phi) = 2sin(\phi)cos(\theta)
$$
 (1)

The difference in the sines is proportional to the cosine of the mean angle.

$$
cos(\theta + \phi) - cos(\theta - \phi) = -2sin(\phi)sin(\theta)
$$
 (2)

The difference in the cosines is proportional to the (negative) of the sine of the mean angle.

IV The Sine Table

Aryabhata obtained the values of the sines at fixed angles between 0 and $\pi/2$ thus generating the sine table for $\pi/48 = 3.75$ degrees, 7.5 degrees up to 90 degrees. This table is stated in verse 12 of the first chapter, the Gitikapada. The table has been used by Indian astronomers (and astrologers) in some form or another since 499 CE up to the present. We shall see how the table was generated.

Let us take $\phi = \epsilon/2$ where ϵ is small. We take $\theta = (n-1/2)\epsilon$ where n is a positive integer from 1 to N. To fix our ideas $\epsilon = \pi/48 = 3.75^0 = 225$ and $N = 24$. We re-state the sine and cosine difference formulae (Eqs. [\(1\)](#page-3-1) and [\(2\)](#page-3-2)) from the previous section

$$
\delta s_n = s_n - s_{n-1} = 2s_{1/2}c_{(n-1/2)}
$$
\n(3)

$$
\delta c_n = c_n - c_{n-1} = -2s_{1/2}s_{(n-1/2)}
$$
\n(4)

where the symbol s_n stands for $sin(n\epsilon)$, c_n stands for $cos(n\epsilon)$ and $s_{1/2}$ for $sin(\epsilon/2)$. The above is a pair of coupled equations and it was Aryabhata's insight to take the second difference, namely

$$
\delta^2 s_n = \delta s_n - \delta s_{n-1} = 2s_{1/2}(c_{(n-1/2)} - c_{(n-3/2)})
$$

= $-4s_{1/2}^2 s_{n-1}$ on using Eq. (4) (5)

³One radian is 3438 minutes and we remind the reader that 2π radians is 180 degrees and 1 degree is 60 minutes.

Figure 2: Derivation of the sine difference relation. The figure depicts the quadrant of a unit circle of radii $OX = OY = 1$. The half-chords AP, BQ and CR are $sin(\theta)$, $sin(\theta + \phi)$ and $sin(\theta - \phi)$ respectively. It is worth noting that (later) we shall take ϕ to be a small angle.

Thus the second difference of the sines is proportional to the sine itself. The next step is to represent the r.h.s in terms of a recursion. We observe s_n on the r.h.s. of Eq.[\(5\)](#page-3-4) may be written as $s_n = s_n - s_{n-1} + s_{n-1} - s_{n-2} + s_{n-2} - \dots = \delta s_{n-1} + \delta s_{n-2} + \dots$ Thus

$$
\delta s_n - \delta s_{n-1} = -4s_{1/2}^2 \sum_{m=1}^{n-1} \delta s_m \tag{6}
$$

Thus we get a recursion relation where the second difference of the sines is expressed in terms of all previously obtained first sine differences. To initiate the recursion we need δs_1 which is $s_1 - s_0 = \sin(\epsilon) - \sin(0) \approx \epsilon$ since for small angles the half-chord and the corresponding arc are equal as stated in the previous section.

Using the recursion relation of the sine, we can generate the celebrated sine table of Aryabhata, taking $\pi = 3.1416$ and $\sin(\epsilon) = \epsilon = 0.0654$ (= 225²).

Table [2](#page-6-0) depicts some typical values of the sine function as well as the value of the sine multiplied by 3438 (the so called 'Rsine' of Aryabhata). We can see that this matches Aryabhata's celebrated sine table, up to \pm 1 minute. For example, $\theta = \pi/6$ gives 1719 minutes. For comparison, we also show the modern values of sine up to four decimal places. Note that Aryabhata takes angles up to $\pi/2$ and seemed aware of the fact that going further was unnecessary given the periodic nature of the sine function.

$\overline{\theta}$	$sin(\theta)$ Aryabhata	$\sin(\theta)$ (minutes)	$\sin(\theta)$ Modern
$\pi/48$	0.0654	225	0.0654
$2\pi/48$	0.1305	449	0.1305
$3\pi/48$	0.1951	671	0.1951
$4\pi/48$	0.2588	890	0.2588
$5\pi/48$	0.3214	1105	0.3214
$6\pi/48$	0.3827	1315	0.3827
$7\pi/48$	0.4423	1520	0.4423
$8\pi/48$	0.5000	1719	0.5000
$9\pi/48$	0.5556	1910	0.5556
$10\pi/48$	0.6088	2093	0.6088
$11\pi/48$	0.6594	2267	0.6593
$12\pi/48$	0.7072	2431	0.7071
$13\pi/48$	0.7519	2585	0.7518
$14\pi/48$	0.7935	2728	0.7934
$15\pi/48$	0.8316	2859	0.8315
$\frac{16\pi}{48}$	0.8662	2978	0.8660
$17\pi/48$	0.8971	3084	0.8969
$18\pi/48$	0.9241	3177	0.9239
$19\pi/48$	0.9472	3256	0.9469
$20\pi/48$	0.9662	3322	0.9659
$21\pi/48$	0.9812	3373	0.9808
$22\pi/48$	0.9919	3410	0.9914
$23\pi/48$	0.9983	3432	0.9979
$\overline{24}\pi/48$	1.0005	3439	1.0000

Table 2: Table of values of sine using the Aryabhatan method, taking episilon $=\pi/48$ $(= 3.75^{\circ} = 225^{\circ})$ and pi=3.1416 and comparison with the modern day values. In column 3 we also quote values in minutes as done in Verse 12, Gitika chapter of the Aryabhatiya [\[1,](#page-9-0) [2\]](#page-9-1).

V Finite Difference Calculus

Of greater relevance is the fact that the sine (or cosine) difference formulae foreshadow finite difference calculus, a popular numerical technique in this age of computation. Rewriting Eqs. [\(1\)](#page-3-1) and [\(2\)](#page-3-2) with $\phi = \epsilon$,

$$
\frac{\sin(\theta + \epsilon) - \sin(\theta - \epsilon)}{2\sin(\epsilon)} = \cos(\theta) \tag{7}
$$

$$
\frac{\cos(\theta + \epsilon) - \cos(\theta - \epsilon)}{2\sin(\epsilon)} = -\sin(\theta) \tag{8}
$$

Aryabhata took ϵ to be $\pi/48$. But he also stated that its value is *yateshtani* or as per our wish (Verse 11, $Ganitapada$). Some took it to be $\pi/96$ and others like Brahmagupta took it as $\pi/12$ or 15°. If we take ϵ to be sufficiently small we have our classic formula for finite difference calculus. Noting that $2 \sin(\epsilon/2) \approx \epsilon$ we have the finite difference version of the derivative of sine

$$
\frac{\delta sin(\theta)}{\delta \theta} = cos(\theta)
$$

and similarly for the cosine.

$$
\frac{\delta cos(\theta)}{\delta \theta} = -sin(\theta)
$$

Let us understand this with an example. We know that $\text{sine}(37^0)$ is close to 0.6 and sine(30^0) is 0.5. The difference in angle is 7^0 which in radians is 0.122. Thus the derivative of sine of the median angle 33.5° from Eq. [\(7\)](#page-6-1) is

$$
\delta \sin(\theta) / \delta \theta = (0.6 - 0.5) / 0.122 = .82
$$

Looking up the sine table or the calculator yields $cos(33.5) = 0.83$. Similarly Eqs. [\(5\)](#page-3-4) yields the second derivatives namely

$$
\delta^2 \sin(\theta) / \delta^2 \theta \approx -\sin(\theta)
$$

$$
\delta^2 \cos(\theta) / \delta^2 \theta \approx -\cos(\theta)
$$

The above are now called central difference approximations to the derivative and the second derivative. Aryabhata does not mention the term finite difference calculus (let alone calculus). But similar methods are now used to numerically solve our differential equations. A student can readily recognize the above as a standard solution of the classical simple harmonic oscillator. Note also that Newton's II Law and the famous Schrodinger equation of quantum mechanics are both second order differential equations.

VI Discussion

One can discern a continuity in Indian mathematics, howsoever tenuous, from pre-Vedic times (< 1000 BCE) up and until 1800s. A striking example is the influence of the Aryabhatiya (499 CE) on major Indian mathematicians who followed him including Madhava (1350 CE) who founded Calculus [\[4\]](#page-9-3); as also the influence on Aryabhata of the mathematics which preceded him [\[6,](#page-9-5) [7\]](#page-10-0).

Aryabhata describes the sine function "poetically" as the Ardha-Jya or half bow (see Fig. 1). To reiterate, he seemed aware that the sine of (i) zero is zero; (ii) small angle is itself, or the small arc is almost equal to the half chord; (iii) 30 degrees is $1/2$ (Ganitapada verse 9); (iv) and 90 degrees is unity. Further, that the sine function is periodic so he prudently stops the enumeration of the sine for angles greater than 90 degrees. Then, in a remarkably insightful way, he laid down the recursion relation for sine differences which enables one to generate the sine table. It is this work, more than his solution to the linear Diophantine equation (verses 31 and 32 of the *Ganitapada*) which establishes him as a genius and one of the brightest stars in the firmament of world mathematics.

The sine table can also be generated using the half angle formula. This was demonstrated in the Panchasiddhantika a text written barely 50 years after the appearance of the Aryabhatiya [\[8\]](#page-10-1). As pointed out, a feature of the Aryabhata's difference relation is how contemporaneous it is. It can be easily recognized as finite difference calculus.

It also led to the development of the calculus of trigonometric functions by the Madhava (1350 CE) and his disciples along the banks of the Nila river in Kerala. This school is variously called the Nila [\[4\]](#page-9-3) and even as the Aryabhata school [\[7\]](#page-10-0). Another aspect to note is that Bhaskara II (1100 CE) used the canonical 2π /96 division of the great circle to carry out discrete integration and obtain the (correct) expressions for the surface area and volume of the sphere. Jyesthdeva of the Nila (or Aryabhta) school in his work Yuktibhasa derived the same results using calculus (circa 1500 CE). Aryabhata can legitimately be called the founder of trigonometry.

To sum up, the Arybhatiya exercised a tremendous influence over Indian mathematicians and, for over a thousand years. For a book with just over a 100 pithy verses, its legacy remains unparalleled in the scientific world. We hope that our article will give our young audience an introduction to his work and will serve as an inspiration.

Acknowledgement: One of the authors (VAS) would place on record the many useful discussions he had with Prof. P. P. Divakaran.

Appendix: The Derivation of the sine and cosine difference formula

As stated in the text Figure 2 depicts a quadrant of the unit circle where $OX =$ $OY = 1$. The arcs XA, XB and XC trace angles θ , $\theta + \phi$ and $\theta - \phi$ respectively. The half-chords AP, BQ and CR are the corresponding sine functions. We drop a perpendicular CS from the circumference onto the half-chord BQ as shown.

We show that the two triangles BSC and OPA are similar. By construction $\angle BSC$ and $\angle OPA$ are each 90 degrees. Note $OB = OC = 1$ (unit radius) and hence $\triangle OBC$ is isosceles. This implies that $\angle OBC = 90^{\circ} - \phi$. Also $\angle OBQ = 90^{\circ} - \phi - \theta$. Hence $\angle SBC = \angle OBC - \angle OBQ = \theta$. Therefore, $\angle SBC = \angle POA = \theta$. This establishes the similarity of the two triangles by the angle-angle test.

$$
\frac{BS}{OP} = \frac{BC}{OA}
$$

Now $OA = 1$ (unit circle), $OP = cos(\theta)$ and $BC = 2 sin(\phi)$. By inspection BS $= BQ - CR = sin(\theta + \phi) - sin(\theta - \phi)$. This yields the sine difference formula (Eq. [\(1\)](#page-3-1)

$$
sin(\theta + \phi) - sin(\theta - \phi) = 2sin(\phi)cos(\theta)
$$

Thus the difference in the sines is proportional to the cosine of the mean angle. We can also obtain the cosine difference formula (Eq. [\(1\)](#page-3-1) by noting that

$$
\frac{CS}{AP} = \frac{BC}{OA}
$$

Note $AP = sin(\theta)$ and $CS = OR - OQ = cos(\theta - \phi) - cos(\theta + \phi)$. Hence

$$
cos(\theta + \phi) - cos(\theta - \phi) = -2sin(\phi)sin(\theta)
$$

The difference in the cosines is proportional to the (negative) of the sine of the mean angle. We pause to note that prima facie the two triangles we considered appear unrelated. A hallmark of Indian mathematics is strong geometric intuition and this dates back to the Sulbasutra circa 800 BCE. Another is the reliance on the "rule of three" (trirasikam) - here we employ a simple version of it namely, if $a/b = c$ then $a = b \times c$.

EXERCISES

- 1. We can generate the sine table as per Aryabhata's suggestion but not exactly using the same value for ϵ he used. We choose $\epsilon = \pi/80 \approx 0.0393$ which is the same as 2.25⁰. We take $sin(\epsilon) \approx \epsilon$. If you have a simple calculator generate all values of sine from 2.25 to 18 degrees in equal steps using Eq. [\(6\)](#page-5-0). Alternatively if you have a programmable calculator or a computer generate all values of sine from 2.25 to 90 degrees. Compare with the results your calculator will otherwise yield.
- 2. In the last section reference is made of the text Panchasiddhantika wherein the half angle forumla is mentioned, viz.

$$
cos(2\theta) = 1 - 2sin^2(\theta)
$$

How would you (i) derive this by a geometrical construction; (i) employ this to generate the sine table?

References

- [1] "Aryabhatiya", Walter E. Clark, University of Chicago Press (1930).
- [2] "Aryabhatiya", Kripashanker Shukla and K V Sarma, Vols. I and II, Indian National Science Academy Publication (1976). The Section III has been shaped by a critical reading of certain verses from the Ganitapada by these authors.
- [3] "Aryabhatiya" with the Bhasya (Commentary) of Nilakantha Somaiyaji, Parts I and II ed. Sambasiva Sastri, Government of Travancore, Trivandrum (1930, 1931); Part III, ed. Suranad Kunjan Pillai, University of Kerala, Trivandrum (1957). This work is in Sanskrit and there is no English (or Hindi) translation of this seminal text to the best of our knowledge. The other works mentioned herein are in English.
- [4] "The Mathematics of India", P. P. Divakaran, Hindustan Book Agency (2018). The book has shaped this article in ways covert and overt.
- [5] "Two New Proofs of the Pythagorean Theorem - Part I", Shailesh Shirali, At Right Angles, Issue 16, pages 7-17, (July 2023).
- [6] "The History of Hindu Mathematics's", Bibhutibhusan Datta and Avadhesh Narayan Singh, Vols. I and II, Asia Publishing House, Delhi (1935 and 1938). A pioneering book on Indian Mathematics written in Pre-Independence India.
- [7] "Geometry in Ancient and Medieval India", Sarasvati Amma, Motilal Banarsidas, Delhi (1999). It has detailed discussions worth looking at. It also uncovers the element of continuity in the Indian mathematical traditions from ancient times to the pre-British era.
- [8] "Panchasiddhantika" of Varahamihira with translation and notes by T. S. Kuppanna Sastry, P.P.S.T. Foundation, Madras (1993).

Prof. Vijay A. Singh has been faculty at IIT Kanpur (1984-2014) and HBCSE, Tata Institute for Fundamental Research (2005-2015) where he was the National Coordinator of both the Science Olympiads and the National Initiative on Undergraduate Science for a decade. He is a Fellow, National Academy of Sciences, India and was President, the Indian Association of Physics Teachers (2019-21). He is currently a Visiting Professor CEBS, Mumbai. (emailid: physics.sutra@gmail.com)

Aneesh Kumar is Standard XII student at the Dhirubhai Ambani International School, Mumbai, who is keenly interested in Mathematics and Physics. (emailid: aneesh.kumar11235@gmail.com)