

Deformations and desingularizations of conically singular associative submanifolds

Gorapada Bera

2023-11-17

Abstract

The proposals of Joyce [Joy18], and Doan and Walpuski [DW19] on counting closed associative submanifolds of G_2 -manifolds depend on various conjectural transitions. This article contributes to the transitions arising from the degenerations of associative submanifolds into conically singular (CS) associative submanifolds. First, we study the moduli space of CS associative submanifolds with singularities at a finite number of points locally modeled on associative cones in \mathbf{R}^7 and their transversality in a co-closed G_2 -manifold as well as in a path of co-closed G_2 -structures. We prove that for a generic co-closed G_2 -structure and for a generic path of co-closed G_2 -structures, there are no CS associative submanifolds having singularities modeled on cones with stability-index greater than 0 and 1, respectively. We establish that associative cones with links null-torsion holomorphic curves in S^6 have stability-index greater than 4, and all special Lagrangian cones in \mathbf{C}^3 have stability-index greater than equal to 1 with equality only holding for the Harvey-Lawson T^2 -cone and a transverse pair of planes. Second, we study the desingularizations of CS associative submanifolds in a one parameter family of co-closed G_2 -structures. Consequently, we derive desingularization results relating the above transitions for CS associative submanifolds with Harvey-Lawson T^2 -cone singularity and for associative submanifolds with transverse self-intersection.

1 Introduction: Main results

Joyce [Joy18], and Doan and Walpuski [DW19] have made proposals of constructing enumerative invariants of G_2 -manifolds by counting closed associative submanifolds. It has been observed that the counting (possibly with sign) of associative submanifolds does not lead to an invariant due to various transitions (see Figure 3) that may occur along a generic path of G_2 -structures. One type of such transitions is the degenerations of associative submanifolds into conically singular associative submanifolds. In particular, it has been conjectured (see Joyce [Joy18, Conjecture 4.4, Conjecture 5.3]) that at least the following transitions can occur along a generic path of co-closed G_2 -structures ϕ_t on a 7-dimensional manifold Y .

1. Three families of embedded closed associative submanifolds, P_t^1 with $-T < t < 0$ and P_t^2, P_t^3 with $0 < t < T$ in (Y, ϕ_t) , converge in the sense of currents to an associative submanifold P with a Harvey-Lawson (HL) T^2 -cone singularity at x in (Y, ϕ_0) , as $t \rightarrow 0$.

The P_t^i are diffeomorphic to the Dehn filling of $P^o := P \setminus B(x)$ along simple closed curves $\mu_i \subset \partial P^o \cong T^2$ which satisfy $\mu_1 \cdot \mu_2 = \mu_2 \cdot \mu_3 = \mu_3 \cdot \mu_1 = -1$.




Figure 1: Three associatives arising from a singular associative.

2. Two families of embedded closed associative submanifolds, P_t with $0 \neq t \in (-T, T)$ and $P_t^\#$ with $0 < t < T$ in (Y, ϕ_t) , converge in the sense of currents to an associative submanifold P with a self intersection in (Y, ϕ_0) , as $t \rightarrow 0$. The $P_t^\#$ are diffeomorphic to the connected sums $P_t \# (S^1 \times S^2)$ if P_t is connected, and otherwise to $P_t^+ \# P_t^-$, where $P_t = P_t^+ \amalg P_t^-$.




Figure 2: Birth of an associative out of an associative with self intersection.

One of our main goals is to confirm the above two transitions, provided there is an associative submanifold with a self intersection (see Theorem 1.29) and an associative submanifold with a Harvey-Lawson T^2 cone singularity (see Theorem 1.21), respectively. More generally, we prove a desingularization theorem (see Theorem 1.16) for conically singular associative submanifolds under a certain hypothesis in (1.17). This is done in a 1-parameter family of co-closed G_2 -structures by gluing rescaled asymptotically conical associative submanifolds of \mathbf{R}^7 . Consequently we derive the first transition (see Theorem 1.21) and partially derive the second transition (see Theorem 1.29). We would like to point out that desingularization theorems for conically singular special Lagrangian submanifolds in Calabi-Yau manifolds and coassociative submanifolds of G_2 -manifolds have been studied by Joyce [Joy04c; Joy04d] and Lotay [Lot09; Lot14] respectively.

To make further progress in the proposals of Joyce [Joy18], and Doan and Walpuski [DW19] about enumerative invariants of G_2 -manifolds one has to figure out all possible degenerations of closed associative submanifolds that can occur in a generic d -parameter family of co-closed G_2 -structures with $d = 0, 1$. It is known in geometric measure theory (see Theorem 2.28) that the associative submanifolds can only degenerate into an associative integral current with a singular set of Hausdorff dimension at most 1. The regularity of this singular set is not clear yet. If all the tangent cones are Jacobi integrable multiplicity 1 associative cones with smooth

link, then the associative integral current is conically singular with isolated singular points (see Definition 3.53). It is therefore natural to ask the following question.

Question 1.1. What are all the possible conically singular (CS) associative submanifolds that may appear in a generic d -parameter family of co-closed G_2 -structures with $d = 0, 1$?

To answer this question we study the deformation theory of conically singular associative submanifolds. We explain that (see Theorem 1.9) the index of the deformation operator can be expressed completely in terms of certain non-negative integer associated the tangent cones, we define this to be the stability index (see Definition 1.4). In particular, CS associative submanifolds having one singularity modeled on cones with stability-index 0 and 1 can only appear in a generic 0 and 1-parameter family of co-closed G_2 -structures, respectively (see Theorem 1.11). Therefore we ask the following subsequent question.

Question 1.2. What are all the associative cones in \mathbf{R}^7 with stability index equal to 0 and 1?

We establish that the Harvey-Lawson T^2 -cone and a union of two transverse special Lagrangian planes have stability index 1 (see Theorem 1.7). Also we prove that all other special Lagrangian cones in \mathbf{C}^3 have stability index strictly greater than 1 with the help of a result obtained by Haskins [Has04b]. Furthermore, we prove that all associative cones in \mathbf{R}^7 whose links are null-torsion holomorphic curves in S^6 (see Bryant [Bry82]) have stability index strictly greater than 4 with the help of a result obtained by Madnick [Mad21]. In particular, this holds for all associative cones whose link in S^6 is of genus 0 but not a totally geodesic sphere. We also see that associative cones with link of genus 1 always have stability index greater than equal to 1.

We now summarize all the main results obtained in this article.

Moduli space of conically singular associative submanifolds

Since our aim is to study the deformation theory of conically singular associative submanifolds and the associative cones play an important role there, we have chosen to study first associative cones in \mathbf{R}^7 . These are the cones whose links in (S^6, J) are pseudoholomorphic curves (see Section 3.1). For this purpose, we express the moduli space of pseudoholomorphic curves in S^6 locally as the zero set of a nonlinear map (see Definition 3.19). We prove the following theorem in Section 3.1.

Theorem 1.3. *Let Σ be a closed holomorphic curve in (S^6, J) . Then a neighbourhood of Σ in the moduli space of holomorphic curves \mathcal{M}^{hol} is homeomorphic to the zero set of a smooth map (obstruction map/Kuranishi map)*

$$\text{ob}_\Sigma : \mathcal{J}_\Sigma \rightarrow \text{coker}(\mathbf{D}_\Sigma + 2J),$$

where the operator \mathbf{D}_Σ is defined in (3.16) and \mathcal{J}_Σ is an open neighbourhood of 0 in $\ker(\mathbf{D}_\Sigma + 2J)$. Moreover the index of the operator $\mathbf{D}_\Sigma + 2J$ is zero and there is a J -anti-linear isomorphism $\gamma_\Sigma : \overline{\text{Hom}}_{\mathbf{C}}(T\Sigma, N\Sigma) \rightarrow N\Sigma$ such that the normal Cauchy-Riemann operator $\bar{\partial}_\nabla^N$ (see Definition 3.8) satisfies

$$\gamma_\Sigma \circ \bar{\partial}_\nabla^N = \mathbf{D}_\Sigma + 2J.$$

For each associative cone C in \mathbf{R}^7 we associate integer invariants $\text{s-ind}(C)$, called stability index and $\text{s-ind}_\Delta(C)$, called Δ -stability index (see Definition 1.4). It turns out that the negative of these stability indices are essentially the virtual dimensions of the moduli spaces of conically singular associative submanifolds.

Definition 1.4. Let C be an associative cone in \mathbf{R}^7 . Denote the link by Σ , which is a closed J -holomorphic curve in S^6 . Let d_λ be as in Definition 3.21. Let $\Sigma = \amalg_{j=1}^l \Sigma_j$ be the decomposition into connected components. Denote the cone of Σ_j by C_j . Let H_j be the maximal subgroup of G_2 which fixes Σ_j under the standard action of G_2 on S^6 . H_j is called the **symmetry group** of C_j . The **stability index** of C is defined by

$$\text{s-ind}(C) := \frac{d_{-1}}{2} + \sum_{-1 < \lambda \leq 1} d_\lambda - 7 - \sum_{j=1}^l (\dim G_2 - \dim H_j).$$

The associative cone C is called **rigid** if all the infinitesimal deformations of its link are induced by the G_2 -action, that is,

$$d_1 = \sum_{j=1}^l (\dim G_2 - \dim H_j).$$

We denote by $\mathcal{M}_\bullet^{\text{hol}}$ the moduli space of closed connected holomorphic curves in S^6 . The moduli spaces $\mathcal{M}_\bullet^{\text{hol}}$ and \mathcal{M}^{hol} might not be manifolds. Therefore we fix a decomposition Δ of $\mathcal{M}_\bullet^{\text{hol}}$ into a countable disjoint union of G_2 -invariant manifolds¹, that is,

$$(1.5) \quad \mathcal{M}_\bullet^{\text{hol}} \stackrel{\Delta}{=} \bigsqcup_{k \in I} \mathcal{Z}_k$$

where the indexing set I is countable, and $G_2 \cdot \mathcal{Z}_k = \mathcal{Z}_k$ and \mathcal{Z}_k are finite dimensional submanifolds of \mathcal{S} (see Definition 3.7). Let \mathcal{Z}_j , $j = 1, \dots, l$, be the manifolds in the above decomposition Δ containing Σ_j . Set $\mathcal{Z} := \prod_{j=1}^l \mathcal{Z}_j$. We define the Δ -stability index of C by

$$\text{s-ind}_\Delta(C) := \frac{d_{-1}}{2} + \sum_{-1 < \lambda \leq 1} d_\lambda - 7 - \sum_{j=1}^l \dim \mathcal{Z}_j. \quad \spadesuit$$

Remark 1.6. Note that the definition of Δ -stability index depends on the decomposition Δ in (1.5). It is also clear that $\text{s-ind}(C) \geq \text{s-ind}_\Delta(C)$. We will see in Remark 3.32 that if C is not a 3-plane then $\text{s-ind}_\Delta(C) \geq 0$. If an associative cone C is rigid then for any decomposition of $\mathcal{M}_\bullet^{\text{hol}}$ we have $\text{s-ind}_\Delta(C) = \text{s-ind}(C)$. \clubsuit

We prove the following theorem in Section 3.1.

Theorem 1.7. *Let C be an associative cone in \mathbf{R}^7 with link $\Sigma \subset S^6$.*

(i) *If genus of Σ is 1 then $\text{s-ind}(C) \geq \text{s-ind}_\Delta(C) \geq 1$.*

¹Since $\mathcal{M}_\bullet^{\text{hol}}$ is an analytic space, there exists always such a decomposition.

(ii) If Σ is a null torsion holomorphic curve in S^6 (see Example 3.5) then

$$\text{s-ind}(C) \geq \text{s-ind}_\Delta(C) > 4.$$

In particular this holds for any holomorphic curve of genus 0 in S^6 which is not a totally geodesic sphere.

(iii) If C is the Harvey-Lawson T^2 -cone (see Example 3.4) or a union of two special Lagrangian planes with transverse intersection at the origin (see Example 3.3) then

$$\text{s-ind}_\Delta(C) = \text{s-ind}(C) = 1.$$

(iv) Let C be a special Lagrangian cone in \mathbf{C}^3 which is not a plane. Then²

$$\text{s-ind}_\Delta(C) \geq \frac{b^1(\Sigma)}{2} + b^0(\Sigma) - 1 \geq 1$$

with equality if and only if C is the Harvey-Lawson T^2 -cone or a union of two special Lagrangian planes with transverse intersection at the origin.

Let (Y, ϕ) be a co-closed G_2 -manifold (see Definition 2.4). We denote by \mathcal{P} the space of all co-closed G_2 -structures on Y and by \mathcal{S} the space of all smooth paths $[0, 1] \rightarrow \mathcal{P}$. We consider conically singular (CS) associative submanifolds with isolated singularities at finite number of points in (Y, ϕ) (see Definition 3.53). These singularities are locally modeled on associative cones in \mathbf{R}^7 . The moduli space of all conically singular associative submanifolds in (Y, ϕ) is denoted by $\mathcal{M}_{\text{cs}}^\phi$ (see Definition 3.75). Given a path of co-closed G_2 -structures $\phi \in \mathcal{S}$, the 1-parameter moduli space of all closed embedded associative submanifolds is denoted by $\mathcal{M}_{\text{cs}}^{\phi_t}$ (see Definition 3.100). More explicitly,

$$\mathcal{M}_{\text{cs}}^\phi = \{(t, P) : t \in [0, 1], P \in \mathcal{M}_{\text{cs}}^{\phi_t}\}.$$

Both these moduli spaces are equipped with weighted C^∞ topology (see Definition 3.79). We will now decompose the moduli spaces $\mathcal{M}_{\text{cs}}^\phi$ and $\mathcal{M}_{\text{cs}}^{\phi_t}$ as a countable union of sub moduli spaces whose deformation theory will be studied.

Definition 1.8. Fix a decomposition $\mathcal{M}_{\bullet}^{\text{hol}} \triangleq \sqcup_{k \in I} \mathcal{Z}_k$ as in Definition 1.4. Let $\mathcal{Z} = \prod_{i=1}^m \mathcal{Z}_i$ where $\mathcal{Z}_i = \prod_{j=1}^l \mathcal{Z}_i^j$ with \mathcal{Z}_i^j is one of the manifold in the decomposition. We define by $\mathcal{M}_{\text{cs}, \mathcal{Z}}^\phi$ all CS associative submanifolds whose asymptotic cones C_i have links $\Sigma_i = \sqcup_{j=1}^l \Sigma_i^j$ with $\Sigma_i^j \in \mathcal{Z}_i^j$. We also define

$$\mathcal{M}_{\text{cs}, \mathcal{Z}}^\phi := \{(t, P) : t \in [0, 1], P \in \mathcal{M}_{\text{cs}, \mathcal{Z}}^{\phi_t}\}.$$

We write the moduli spaces $\mathcal{M}_{\text{cs}}^\phi$ and $\mathcal{M}_{\text{cs}}^{\phi_t}$ as a countable union of sub moduli spaces, namely

$$\mathcal{M}_{\text{cs}}^\phi = \bigcup_{\mathcal{Z}} \mathcal{M}_{\text{cs}, \mathcal{Z}}^\phi \quad \text{and} \quad \mathcal{M}_{\text{cs}}^{\phi_t} = \bigcup_{\mathcal{Z}} \mathcal{M}_{\text{cs}, \mathcal{Z}}^{\phi_t}.$$

Here the unions run over all possible $\mathcal{Z} = \prod_{i=1}^m \mathcal{Z}_i$ where $\mathcal{Z}_i = \prod_{j=1}^l \mathcal{Z}_i^j$ as above. \spadesuit

²The stability index in Definition 1.4 is not same as the stability index defined for special Lagrangian cones in [Joy04b, Definition 3.6].

We prove the following theorem in Section 3.5 about local Kuranishi models for the above moduli spaces.

Theorem 1.9. *Let \mathcal{Z} , $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ and $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ be as in Definition 1.8. Let $P \in \mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ be a conically singular associative submanifold in a co-closed G_2 -manifold (Y, ϕ) whose singularities are modeled on cones $C_i \in \mathcal{Z}_i$, $i = 1, \dots, m$ (see Definition 3.53). Let $\phi : [0, 1] \rightarrow \mathcal{P}$ be a path of co-closed G_2 structures such that $\phi(t_0) = \phi$. Then there exist open neighbourhoods $\tilde{\mathcal{I}}_{P,\mathcal{Z}}$ and $\bar{\mathcal{I}}_{P,\mathcal{Z}}$ of 0 in $\ker \tilde{\mathbf{D}}_{P,\mu,\mathcal{Z}}$ and $\ker \bar{\mathbf{D}}_{P,\mu,\mathcal{Z}}$ respectively (where $\tilde{\mathbf{D}}_{P,\mu,\mathcal{Z}}$ and $\bar{\mathbf{D}}_{P,\mu,\mathcal{Z}}$ are defined in Definition 3.98 and Definition 3.100 respectively and μ is chosen as in Definition 3.79) such that*

(i) *the moduli space $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ near P is homeomorphic to $\text{ob}_{P,\mathcal{Z}}^{-1}(0)$, the zero set of a smooth map*

$$\text{ob}_{P,\mathcal{Z}} : \tilde{\mathcal{I}}_{P,\mathcal{Z}} \rightarrow \text{coker } \tilde{\mathbf{D}}_{P,\mu,\mathcal{Z}}.$$

Moreover, $\text{index } \tilde{\mathbf{D}}_{P,\mu,\mathcal{Z}} = - \sum_{i=1}^m \text{s-ind}_\Delta(C_i)$.

(ii) *the 1-parameter moduli space $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ near (t_0, P) is homeomorphic to $\text{ob}_{t_0,P,\mathcal{Z}}^{-1}(0)$, the zero set of a smooth map*

$$\text{ob}_{t_0,P,\mathcal{Z}} : \bar{\mathcal{I}}_{P,\mathcal{Z}} \rightarrow \text{coker } \bar{\mathbf{D}}_{P,\mu,\mathcal{Z}}.$$

Moreover, $\text{index } \bar{\mathbf{D}}_{P,\mu,\mathcal{Z}} = \text{index } \tilde{\mathbf{D}}_{P,\mu,\mathcal{Z}} + 1 = - \sum_{i=1}^m \text{s-ind}_\Delta(C_i) + 1$.

The moduli spaces $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ and $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ are not always smooth manifolds. The following definition will make these moduli spaces smooth in a generic situation.

Definition 1.10. Let \mathcal{Z} , $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ and $\mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ be as in Definition 1.8. We define $\mathcal{P}_{\text{cs},\mathcal{Z}}^{\text{reg}}$ to be the subset consisting of all $\phi \in \mathcal{P}$ which has the property that for all CS associative $P \in \mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ the linear operator

$$\tilde{\mathbf{D}}_{P,\mu,\mathcal{Z}} \text{ is surjective.}$$

Similarly $\mathcal{P}_{\text{cs},\mathcal{Z}}^{\text{reg}}$ to be the subset consisting of all $\phi \in \mathcal{P}$ which has the property that for all $(t_0, P) \in \mathcal{M}_{\text{cs},\mathcal{Z}}^\phi$ the linear operator

$$\bar{\mathbf{D}}_{P,\mu,\mathcal{Z}} \text{ is surjective.}$$

The operators $\tilde{\mathbf{D}}_{P,\mu,\mathcal{Z}}$ and $\bar{\mathbf{D}}_{P,\mu,\mathcal{Z}}$ are defined in Definition 3.98 and Definition 3.100 respectively. In addition we define

$$\mathcal{P}_{\text{cs},\Delta}^{\text{reg}} := \bigcap_{\mathcal{Z}} \mathcal{P}_{\text{cs},\mathcal{Z}}^{\text{reg}} \text{ and } \mathcal{P}_{\text{cs},\Delta}^{\text{reg}} := \bigcap_{\mathcal{Z}} \mathcal{P}_{\text{cs},\mathcal{Z}}^{\text{reg}}.$$

Here the intersections run over all possible $\mathcal{Z} = \prod_{i=1}^m \mathcal{Z}_i$ where $\mathcal{Z}_i = \prod_{j=1}^l \mathcal{Z}_i^j$ as in Definition 1.8 and hence are countable intersections. \spadesuit

We prove the following theorem about generic transversality of the above moduli spaces in Section 3.6. It also tells us about what type of singularity model cones appear in a generic co-closed G_2 -structures and as well as in a generic path of co-closed G_2 -structures.

Theorem 1.11. *Let $\mathcal{Z} = \prod_{i=1}^m \mathcal{Z}_i$ where $\mathcal{Z}_i = \prod_{j=1}^l \mathcal{Z}_i^j$ as in Definition 1.8. Then the subsets $\mathcal{P}_{\text{cs}, \mathcal{Z}}^{\text{reg}}$, $\mathcal{P}_{\text{cs}, \Delta}^{\text{reg}}$ in \mathcal{P} , and $\mathcal{P}_{\text{cs}, \mathcal{Z}'}^{\text{reg}}$, $\mathcal{P}_{\text{cs}, \Delta}^{\text{reg}}$ in \mathcal{P}' are comeager, respectively. In particular, the following holds. Let C_i , $i = 1, \dots, m$ be associative cones in \mathbf{R}^7 having links $\Sigma_i = \sqcup_{j=1}^l \Sigma_i^j$ with $\Sigma_i^j \in \mathcal{Z}_i^j$.*

- (i) *If $\sum_{i=1}^m \text{s-ind}_\Delta(C_i) > 0$, then for any co-closed G_2 -structure $\phi \in \mathcal{P}_{\text{cs}, \Delta}^{\text{reg}}$ the moduli space $\mathcal{M}_{\text{cs}}^\phi$ contains no conically singular associative submanifolds having singularities modeled on cones with links in a neighbourhood of Σ_i in \mathcal{Z}_i .*
- (ii) *If $\sum_{i=1}^m \text{s-ind}_\Delta(C_i) > 1$, then for any path of co-closed G_2 -structures $\phi \in \mathcal{P}_{\text{cs}, \Delta}^{\text{reg}}$ the moduli space $\mathcal{M}_{\text{cs}}^\phi$ contains no conically singular associative submanifolds having singularities modeled on cones with links in a neighbourhood of Σ_i in \mathcal{Z}_i .*

Remark 1.12. We conclude from Theorem 1.11 that if $\phi \in \mathcal{P}_{\text{cs}, \Delta}^{\text{reg}}$ then the moduli space $\mathcal{M}_{\text{cs}}^\phi$ essentially can contain only conically singular associative submanifolds having singularities modeled on cones C with $\text{s-ind}_\Delta(C) = 0$. Similarly, if $\phi \in \mathcal{P}_{\text{cs}, \Delta}^{\text{reg}}$ then the moduli space $\mathcal{M}_{\text{cs}}^\phi$ essentially can contain only conically singular associative submanifolds having singularities modeled on cones C with $\text{s-ind}_\Delta(C) = 0$ or 1. ♣

Desingularizations of conically singular associative submanifolds

Let L be an asymptotically conical associative submanifold in \mathbf{R}^7 (see Definition 3.42).

Definition 1.13. The dilation action of \mathbf{R}^+ on \mathbf{R}^7 generates a **canonical Fueter section** $\hat{s}_L \in C^\infty(NL)$, that is, $\mathbf{D}_L \hat{s}_L = 0$ (see Definition 2.23, Remark 2.24). This \hat{s}_L vanishes at infinity. ♣

Let P be a conically singular associative submanifold in a co-closed G_2 -manifold (Y, ϕ) with singularity at a single point modeled on a cone C whose link is $\Sigma \subset S^6$ (see Definition 3.53). Let L be an asymptotically conical associative submanifold in \mathbf{R}^7 with the asymptotic cone C . The operators \mathbf{D}_P and \mathbf{D}_L are defined in Definition 2.23. The space $\ker \mathbf{D}_{P, \lambda}$ and $\ker \mathbf{D}_{L, \lambda}$ can be expressed informally as follows:

$$\ker \mathbf{D}_{P, \lambda} := \{u \in C^\infty(NP) : \mathbf{D}_P u = 0, u = O(r^\lambda) \text{ as } r \rightarrow 0\}$$

and

$$\ker \mathbf{D}_{L, \lambda} := \{u \in C^\infty(NL) : \mathbf{D}_L u = 0, u = O(r^\lambda) \text{ as } r \rightarrow \infty\}.$$

Definition 1.14. We define the **matching kernel** \mathcal{K}^m by

$$\mathcal{K}^m := \{(u_L, u_P) \in \ker \mathbf{D}_{L, 0} \oplus \ker \mathbf{D}_{P, 0} : i_\infty u_L = i_0 u_P\},$$

where the maps i_∞ and i_0 are the asymptotic limit maps $i_{L, 0}$ and $i_{P, 0}$ respectively, defined in Definition 3.93. ♣

Let $\phi \in \mathcal{P}$ be a path of co-closed G_2 -structures on Y and $t_0 \in (0, 1)$ such that $\phi_{t_0} := \phi(t_0) = \phi$. We set $\phi_t := \phi(t)$. The G_2 -structures ϕ_t induces the 4-forms ψ_t (see Definition 2.2).

Definition 1.15. We define the **extended matching kernel** by

$$\tilde{\mathcal{K}}^m := \{(u_L, u_P, t) \in \ker \mathbf{D}_{L,0} \oplus C_{P,0}^\infty \oplus \mathbf{R} : \mathbf{D}_P u_P + t \hat{f}_P = 0, i_\infty u_L = i_0 u_P\},$$

where \hat{f}_P is as in Definition 3.100 and $C_{P,0}^\infty := \{u \in C^\infty(NP) : u = O(1) \text{ as } r \rightarrow 0\}$. \spadesuit

We prove the following desingularization theorem in Section 4.3.

Theorem 1.16. *Let P be a conically singular associative submanifold in a co-closed G_2 -manifold (Y, ϕ) with singularity at a single point modeled on a cone C . Let L be an asymptotically conical associative submanifold in \mathbf{R}^7 with the asymptotic cone C and rate $\nu < 0$. Let $\phi \in \mathcal{P}$ be a path of co-closed G_2 -structures on Y and $t_0 \in (0, 1)$ such that $\phi_{t_0} = \phi$. Assume that, the matching kernel \mathcal{K}^m and the extended matching kernel $\tilde{\mathcal{K}}^m$ both are one dimensional, that is,*

$$(1.17) \quad \tilde{\mathcal{K}}^m = \mathcal{K}^m = \langle \hat{s}_L \rangle_{\mathbf{R}}.$$

Then there exist $\tilde{\varepsilon}_0 > 0$, a continuous function $t : [0, \tilde{\varepsilon}_0] \rightarrow [0, 1]$ with $t(0) = t_0$ and smooth closed embedded associative submanifolds $\tilde{P}_{\varepsilon, t(\varepsilon)}$ in $(Y, \phi_{t(\varepsilon)})$ for all $\varepsilon \in (0, \tilde{\varepsilon}_0)$ such that $\tilde{P}_{\varepsilon, t(\varepsilon)} \rightarrow P$ as $\varepsilon \rightarrow 0$ in the sense of integral currents.

Now we will apply Theorem 1.16 to desingularize CS associative submanifold with Harvey-Lawson T^2 -cone singularity and associative submanifold with a transverse unique self intersection.

Desingularizations for CS associatives with Harvey-Lawson T^2 -cone singularity:

We start with CS associative submanifold with Harvey-Lawson T^2 -cone singularity. Let $\phi \in \mathcal{P}_{\text{CS}}^{\text{reg}}$ (see Definition 1.10) be a path of co-closed G_2 -structures on Y and $t_0 \in (0, 1)$. Let P be a conically singular associative submanifold in (Y, ϕ_{t_0}) with Harvey-Lawson T^2 -cone singularity at a single point (see Example 3.4). We prove in Lemma 4.26 that $\ker \mathbf{D}_{P,-1} = \langle \hat{v}_P \rangle_{\mathbf{R}}$ and

$$(1.18) \quad a := \langle \hat{v}_P, \hat{f}_P \rangle_{L^2} \neq 0.$$

where \hat{f}_P is as in Definition 3.100.

Definition 1.19. $\mathcal{P}^\bullet \subset \mathcal{P}_{\text{CS}}^{\text{reg}}$ consists of all $\phi \in \mathcal{P}_{\text{CS}}^{\text{reg}}$ such that for all CS associative submanifolds P in (Y, ϕ_{t_0}) for some $t_0 \in (0, 1)$ with Harvey-Lawson T^2 -cone singularity at a single point, the asymptotic limit of \hat{v}_P has the form

$$(1.20) \quad i_{P,-1} \hat{v}_P = b_1 \xi_1 + b_2 \xi_2, \quad b_1 \neq 0, b_2 \neq 0, b_1 \neq b_2.$$

where ξ_1, ξ_2 are in (3.48) of Example 3.47. \spadesuit

Note that $\frac{b_1}{a}, \frac{b_2}{a}$ are independent of various choices of \hat{v}_P . We hope that $\mathcal{P}^\bullet \subset \mathcal{P}_{\text{CS}}^{\text{reg}}$ is comeager. We prove the following desingularization theorem for conically singular associatives with Harvey-Lawson T^2 -cone singularity at a single point in Section 4.4.

Theorem 1.21. *Let $\phi \in \mathcal{P}_{\text{cs}}^{\text{reg}}$ be a path of co-closed G_2 -structures on Y and $t_0 \in (0, 1)$. Let P be a conically singular associative submanifold of (Y, ϕ_{t_0}) in $\mathcal{M}_{\text{cs}}^\phi$ with Harvey-Lawson T^2 -cone singularity at a single point x . There exist $\tilde{\varepsilon}_0 > 0$, three continuous functions $t^i : [0, \tilde{\varepsilon}_0] \rightarrow [0, 1]$ with $t^i(0) = t_0$, $i = 1, 2, 3$ and smooth closed embedded associative submanifolds $\tilde{P}_{\varepsilon, t^i(\varepsilon)}$ in $(Y, \phi_{t^i(\varepsilon)})$ for all $\varepsilon \in (0, \tilde{\varepsilon}_0)$ such that $\tilde{P}_{\varepsilon, t^i(\varepsilon)} \rightarrow P$ as $\varepsilon \rightarrow 0$ in the sense of integral currents. The $\tilde{P}_{\varepsilon, t^i(\varepsilon)}$ are diffeomorphic to the Dehn filling of $P^\circ := P \setminus B_\varepsilon(x)$ along simple closed curves $\mu_i \subset \partial P^\circ \cong T^2$ which satisfy $\mu_1 \cdot \mu_2 = \mu_2 \cdot \mu_3 = \mu_3 \cdot \mu_1 = -1$. Furthermore, if $\phi \in \mathcal{P}^\bullet$ then there is a constant $c \neq 0$ such that*

$$(1.22) \quad \begin{aligned} t^1(\varepsilon) &= t_0 - \frac{cb_2}{a}\varepsilon^2 + o(\varepsilon^2), & t^2(\varepsilon) &= t_0 + \frac{cb_1}{a}\varepsilon^2 + o(\varepsilon^2), \\ & & \text{and } t^3(\varepsilon) &= t_0 + \frac{c(b_2 - b_1)}{a}\varepsilon^2 + o(\varepsilon^2). \end{aligned}$$

We would like to point out that our proof of (1.22) is influenced by [Joy18, Remark 5.4 a)].

Desingularizations for associative submanifolds with transverse intersection:

We will now move on to state the main results obtained in this article about desingularization of associative submanifold with a transverse unique self intersection. Let $\phi \in \mathcal{P}_{\text{cs}}^{\text{reg}}$ be a path of co-closed G_2 -structures on Y and $t_0 \in (0, 1)$. Let $P \in \mathcal{M}_{\text{cs}}^\phi$ be an associative submanifold in (Y, ϕ_{t_0}) with a transverse unique self intersection. In other words P is a conically singular associative with singularity at a unique point, which is modeled on a union of two transverse associative planes Π_\pm in \mathbf{R}^7 so that $B\Pi_0 = \Pi_+$, $B\Pi_\theta = \Pi_-$ and

$$\mathbf{R}^7 = \langle \mathbf{n} \rangle_{\mathbf{R}} \oplus B\Pi_0 \oplus B\Pi_\theta$$

is an orientation compatible splitting as in Example 3.33 and $B \in G_2$. There is an asymptotic limit map (see Lemma 3.70 and Example 3.33)

$$(1.23) \quad \{u \in C_{P,0}^\infty : \mathbf{D}Pu = O(r) \text{ as } r \rightarrow 0\} \rightarrow \Pi_+^\perp \oplus \Pi_-^\perp, \quad u \mapsto (u^+(0), u^-(0)).$$

It turns out $\hat{f}_P = O(r)$ as $r \rightarrow 0$ (see Definition 3.100).

Definition 1.24. $\mathcal{P}^\dagger \subset \mathcal{P}_{\text{cs}}^{\text{reg}}$ consists of all $\phi \in \mathcal{P}_{\text{cs}}^{\text{reg}}$ such that all associative submanifolds P in $(Y, \phi(t_0))$ for some $t_0 \in (0, 1)$ with a transverse unique self intersection are (rigid) unobstructed in $(Y, \phi(t_0))$ as immersed associative submanifolds. In other words, $\ker \mathbf{D}_{P,0} = \{0\}$ and there exists $\hat{v}_P \in C_{P,0}^\infty$ such that

$$(1.25) \quad \mathbf{D}_P \hat{v}_P = \hat{f}_P, \quad \text{and } a := (\hat{v}_P^+(0) - \hat{v}_P^-(0)) \cdot \mathbf{n} \neq 0.$$

The above is also equivalent to saying that there exists a family of immersed closed associative submanifolds

$$(1.26) \quad \{P_t : |t - t_0| \ll 1, P_{t_0} = P, P_t \text{ is embedded closed associative } \forall t \neq t_0\}. \quad \spadesuit$$

Definition 1.27. $\mathcal{P}^\ddagger \subset \mathcal{P}^\dagger$ consists of all $\phi \in \mathcal{P}^\dagger$ such that for all associative submanifolds P in $(Y, \phi(t_0))$ for some $t_0 \in (0, 1)$ with a transverse unique self intersection, there exists $\hat{u}_P \in \ker \mathbf{D}_{P, -2}$ such that

$$(1.28) \quad i_{P, -2} \hat{u}_P = (B\xi^+, B\xi^-), \quad \text{and } b := (\hat{u}_P - B\xi^+)(0) \cdot \mathbf{n} - (\hat{u}_P - B\xi^-)(0) \cdot \mathbf{n} \neq 0,$$

where ξ^\pm are defined in (3.45) of Example 3.44 and B is as in Example 3.33. \spadesuit

Note that choosing $-\mathbf{n}$ instead of \mathbf{n} will swap Π_+ and Π_- and therefore the definitions of a and b are well defined. We hope that $\mathcal{P}^\dagger \subset \mathcal{P}_{\text{cs}}^{\text{reg}}$ and $\mathcal{P}^\ddagger \subset \mathcal{P}_{\text{cs}}^{\text{reg}}$ are comeager subsets. We prove the following desingularization theorem for associative submanifolds with a transverse unique self intersection in Section 4.5.

Theorem 1.29. *Let $\phi \in \mathcal{P}^\dagger$ be a path of co-closed G_2 -structures on Y and $t_0 \in (0, 1)$. Let P be an associative submanifold of (Y, ϕ_{t_0}) in $\mathcal{M}_{\text{cs}}^\phi$ with a transverse unique self intersection at x . There exist $\tilde{\varepsilon}_0 > 0$, a continuous function $t : [0, \tilde{\varepsilon}_0] \rightarrow [0, 1]$ with $t(0) = t_0$ and smooth closed embedded associative submanifolds $\tilde{P}_{\varepsilon, t(\varepsilon)}$ in $(Y, \phi_{t(\varepsilon)})$ for all $\varepsilon \in (0, \tilde{\varepsilon}_0)$ such that $\tilde{P}_{\varepsilon, t(\varepsilon)} \rightarrow P$ as $\varepsilon \rightarrow 0$ in the sense of integral currents. The $\tilde{P}_{\varepsilon, t(\varepsilon)}$ are diffeomorphic to the connected sums $P_t \# (S^1 \times S^2)$ if P_t is connected, and otherwise to $P_t^+ \# P_t^-$, where $P_t = P_t^+ \amalg P_t^-$. Here P_t is as in (1.26).*

Remark 1.30. We discuss in Remark 4.32 following [Joy18, Remark 4.5 a)] that if $\phi \in \mathcal{P}^\ddagger$ then one would expect that there is a constant $c \neq 0$ such that

$$(1.31) \quad t(\varepsilon) = t_0 - \frac{cb}{a} \varepsilon^3 + o(\varepsilon^3).$$

We would like to point out that Nordström [Nor13] had previously proven Theorem 1.29, but did not establish (1.31). Although we have enhanced his proof by making pregluing error smaller, still it lacks required estimates to prove (1.31). We hope that a further modification by deforming Lawlor neck might result in obtaining the expression (1.31). \clubsuit

Acknowledgements. I am grateful to my PhD supervisor Thomas Walpuski for suggesting the problems solved in this article. I am also indebted to him for his constant encouragement and advice during this work. Additionally, I extend my thanks to him, Johannes Nordström, and Jason Lotay for their meticulous review of the draft and valuable feedback. I also thank to Dominik Gutwein and Viktor Majewski for their careful reading and helpful comments on the draft. This material is based upon work supported by the Simons Collaboration on Special Holonomy in Geometry, Analysis, and Physics.

2 Preliminaries: G_2 -manifolds and associative submanifolds

In this section we review definitions and basic facts about G_2 -manifolds, more generally almost G_2 -manifolds and associative submanifolds, which are important to understand this article. To delve further into these topics, we refer the reader to [SW17; Joy07; KLL20; Har90] and other relevant sources mentioned throughout the discourse.

2.1 G_2 -manifolds

The group G_2 is the automorphism group of the normed division algebra of octonions O , i.e. $G_2 := \text{Aut}(O) \subset \text{SO}(7)$. This is a simple, compact, connected, simply connected Lie group of dimension 14. Furthermore there exists a fibration $\text{SU}(3) \hookrightarrow G_2 \rightarrow S^6$. Writing $O = \text{Re } O \oplus \text{Im } O \cong \mathbf{R} \oplus \mathbf{R}^7$, we define the **cross-product** $\times : \Lambda^2 \mathbf{R}^7 \rightarrow \mathbf{R}^7$ by

$$(u, v) \mapsto u \times v := \text{Im}(uv).$$

The 3-form $\phi_e \in \Lambda^3(\mathbf{R}^7)^*$ is defined by

$$\phi_e(u, v, w) := g_e(u \times v, w),$$

where $g_e : S^2(\mathbf{R}^7) \rightarrow \mathbf{R}$ is $g_e(u, v) = -\text{Re}(uv)$, the standard Euclidean metric on \mathbf{R}^7 . These are related by an important identity

$$(2.1) \quad \iota_u \phi_e \wedge \iota_v \phi_e \wedge \phi_e = 6g_e(u, v) \text{vol}_{g_e}.$$

More explicitly, here $\{e^1, \dots, e^7\}$ is an oriented orthonormal frame on \mathbf{R}^7 such that

$$\phi_e = e^{123} - e^{145} - e^{167} - e^{246} - e^{275} - e^{347} - e^{356},$$

where $e^{ijk} := e^i \wedge e^j \wedge e^k$. The Lie group G_2 can also be expressed as

$$G_2 := \{A \in \text{GL}(\mathbf{R}^7) : A^* \phi_e = \phi_e\}.$$

There is also a 4-form $\psi_e := *_{g_e} \phi_e \in \Lambda^4(\mathbf{R}^7)^*$, which also can be defined by

$$\psi_e(u, v, w, z) := g_e([u, v, w], z),$$

where $[\cdot, \cdot, \cdot] : \Lambda^3(\mathbf{R}^7) \rightarrow \mathbf{R}^7$ is the **associator**, defined as follows:

$$[u, v, w] := (u \times v) \times w + \langle v, w \rangle u - \langle u, w \rangle v.$$

Definition 2.2. A G_2 -structure on a 7-dimensional manifold Y is a principal G_2 -bundle over Y which is a reduction of the frame bundle $\text{GL}(Y)$.

An **almost G_2 -manifold**³ is a 7-dimensional manifold Y equipped with a G_2 -structure or equivalently equipped with a **definite 3-form** $\phi \in \Omega^3(Y)$, that is, the bilinear form $G_\phi : S^2 TY \rightarrow \Lambda^7(T^*Y)$ defined by

$$G_\phi(u, v) := \iota_u \phi \wedge \iota_v \phi \wedge \phi$$

is definite. ♠

A G_2 -structure ϕ on Y defines uniquely a Riemannian metric g_ϕ and a volume form vol_{g_ϕ} on Y satisfying the identity (2.1). Moreover it defines

- a **cross product** $\times : \Lambda^2(TY) \rightarrow TY$,

³A 7-dimensional manifold is an almost G_2 -manifold if and only if it is spin, see [Gra69, Theorem 3.1-3.2]

- an **associator** $[\cdot, \cdot, \cdot] : \Lambda^3(TY) \rightarrow TY$,
- a **4-form** $\psi := *_{g_\phi} \phi \in \Omega^4(Y)$.

A G_2 -structure is also equivalent to having a definite 4-form $\psi \in \Omega^4(Y)$ with an orientation on Y ; see [Hito1, Section 8.4]. Here ψ is **definite** means the bilinear form $G_\psi : S^2 T^*Y \rightarrow \Lambda^7(T^*Y) \otimes \Lambda^7(T^*Y)$ defined by

$$G_\psi(u, v) := \iota_u \psi \wedge \iota_v \psi \wedge \psi$$

is definite. In this definition, the 4-form ψ is considered as a section of $\Lambda^3(TY) \otimes \Lambda^7(T^*Y) \cong \Lambda^4(T^*Y)$ and the contraction $\iota_u \psi$ is a section of $\Lambda^2(TY) \otimes \Lambda^7(T^*Y)$.

Definition 2.3. A G_2 -**manifold** is a 7-dimensional manifold Y equipped with a torsion-free G_2 -structure, that is, equipped with a definite 3-form $\phi \in \Omega^3(Y)$ such that $\nabla_{g_\phi} \phi = 0$, or equivalently⁴

$$d\phi = 0 \text{ and } d\psi = 0. \quad \spadesuit$$

Since the moduli space of torsion free G_2 -structures over Y is a smooth manifold of dimension $b^3(Y)$ [Joy96a, Part I, Theorem C], usually we need to consider the following larger spaces of G_2 -structures for enumerative theories [DS11, Section 3.2; Joy18, Section 2.5].

Definition 2.4. A G_2 -structure ϕ is called a **co-closed G_2 -structure** if $d\psi = 0$, where $\psi := *_{g_\phi} \phi$.

An almost G_2 -manifold (Y, ϕ) is called a **co-closed G_2 -manifold** if ϕ is a co-closed G_2 -structure.

A G_2 -structure ϕ is called **tamed**⁵ by a closed 3-form $\tau \in \Omega^3(Y)$ if for all $x \in Y$ and $u, v, w \in T_x Y$ with $[u, v, w] = 0$ and $\phi(u, v, w) > 0$ we have $\tau(u, v, w) > 0$. \spadesuit

Example 2.5. (\mathbf{R}^7, ϕ_e) is a G_2 -manifold. \bullet

Example 2.6. Let $(X, \omega_I, \omega_J, \omega_K)$ be a hyperkähler 4-manifold. The manifolds $\mathbf{R}^3 \times X$ and $T^3 \times X$ are G_2 -manifolds with G_2 -structure

$$\phi := dt^1 \wedge dt^2 \wedge dt^3 + dt^1 \wedge \omega_I + dt^2 \wedge \omega_J + dt^3 \wedge \omega_K,$$

where (t^1, t^2, t^3) are the coordinates of \mathbf{R}^3 . \bullet

Example 2.7. Let (Z, ω, Ω) be a Calabi-Yau 3-fold, where ω is a Kähler form and Ω is a holomorphic volume form on Z satisfying

$$\frac{\omega^3}{3!} = -\left(\frac{i}{2}\right)^3 \Omega \wedge \bar{\Omega}.$$

The product with the unit circle, $Y := S^1 \times Z$ is a G_2 -manifold with the G_2 -structure

$$\phi := dt \wedge \omega + \operatorname{Re} \Omega, \quad \psi = \frac{1}{2} \omega \wedge \omega + dt \wedge \operatorname{Im} \Omega,$$

where t denotes the coordinate on S^1 . In this case the holonomy group $\operatorname{Hol}(Y, g_\phi) \subset \operatorname{SU}(3)$. \bullet

⁴The equivalence was established by Fernández and Gray [FG82, Theorem 5.2]

⁵This definition can be ignored for this article; it is used only in Theorem 2.28 to bound the volume of associative sub-manifolds for compactness.

Remark 2.8. Any G_2 -manifold (Y, ϕ) admits a nowhere vanishing parallel spinor and therefore the metric g_ϕ is Ricci-flat [LM89, pg. 321; Hit74, Theorem 1.2]. A compact G_2 -manifold Y has holonomy exactly equal to G_2 if and only if $\pi_1(Y)$ is finite [Joy07, Proposition 11.2.1]. ♣

Example 2.9. Bryant [Bry87], Bryant and Salamon [BS89] first constructed local and complete manifolds with holonomy equal to G_2 respectively. Joyce [Joy96b] first constructed compact manifolds with holonomy equal to G_2 by smoothing flat T^7/Γ where Γ is a finite group of isometries of T^7 . This has been generalized later by Joyce and Karigiannis [JK17]. Kovalev [Kov03] introduced the twisted connected sum construction of G_2 -manifolds which glues a suitable matching pair of asymptotically cylindrical G_2 -manifolds. This construction was later improved by Kovalev and Lee [KL11] and Corti, Haskins, Nordström, and Pacini [CHNP15] to produce hundreds of thousands examples of compact manifolds with holonomy equal to G_2 . •

Example 2.10. Nearly parallel G_2 -manifolds (i.e. co-closed G_2 -manifolds (Y, ϕ) satisfying $d\phi = \lambda\psi$ for some constant $\lambda \in \mathbf{R}$) are examples of co-closed G_2 -manifolds. •

2.2 Associative submanifolds

In any almost G_2 -manifold we can consider a special class of 3-dimensional calibrated submanifolds, called associative submanifolds, which are the main objects of study in this article. These were first invented by Harvey and Lawson [HL82].

Definition 2.11. Let (Y, ϕ) be an almost G_2 -manifold. A 3-dimensional oriented submanifold P of Y is called an **associative submanifold**⁶ if it is semi-calibrated by the 3-form ϕ , that is, $\phi|_P$ is the volume form vol_{P, g_ϕ} on P , or equivalently ⁷ $[u, v, w] = 0$, for all $x \in P$ and $u, v, w \in T_x P$. ♣

Example 2.12. In Example 2.6, \mathbf{R}^3 and T^3 are associative submanifolds. •

Example 2.13. Let Z be a Calabi-Yau 3-fold and $S^1 \times Z$ be the G_2 -manifold as in Example 2.7. For any holomorphic curve $\Sigma \subset Z$ and special Lagrangian⁸ $L \subset Z$ the 3-dimensional submanifolds $S^1 \times \Sigma$ and $\{e^{i\theta}\} \times L$ are associative submanifolds of $S^1 \times Z$ for all $e^{i\theta} \in S^1$. •

Example 2.14. Joyce [Joy96a, Section 4.2] has produced examples of closed associative submanifolds which are the fixed point loci of G_2 -involutions in his generalized Kummer constructions. Recently new examples of associative submanifolds in generalized Kummer constructions have been constructed by Dwivedi, Platt, and Walpuski [DPW23]. •

Example 2.15. Examples of closed associative submanifolds in the twisted connected sum (TCS) G_2 -manifolds first constructed by Corti, Haskins, Nordström, and Pacini [CHNP15, Section 5, Section 7.2.2] from closed holomorphic curves and closed special Lagrangians in asymptotically cylindrical (ACyl) Calabi-Yau submanifolds. The author in [Ber22] has constructed more examples of closed associative submanifolds in the twisted connected sum G_2 -manifolds. These

⁶If ϕ is a calibration (i.e. $d\phi = 0$) and P is compact then it is a minimal submanifold and volume minimizing in its homology class [HL82, Theorem 4.2].

⁷The equivalence follows from the identity $|u \wedge v \wedge w|^2 = \phi(u, v, w)^2 + |[u, v, w]|^2$, see [HL82, section IV, Theorem 1.6]

⁸i.e. L is calibrated by $\text{Re } \Omega$.

are obtained from ACyl holomorphic curves and ACyl special Lagrangians in ACyl Calabi-Yau submanifolds using a gluing construction. It is also expected that this gluing construction can be used to produce infinitely many closed associative submanifolds in a certain TCS G_2 -manifold discussed by the physicists [BDHLMS18]. \bullet

Remark 2.16. Examples of associative submanifolds of nearly parallel G_2 -manifolds have been constructed by Lotay [Lot12] in S^7 , Kawai [Kaw15] in the squashed S^7 and Ball and Madnick [BM20] in the Berger space. \clubsuit

The following definitions are useful to set up the deformation theory for associative submanifolds in this article, which also have been discussed in [Ber22].

Definition 2.17. Let M be any submanifold of a manifold Y . The **normal bundle** $\pi : NM \rightarrow M$ is characterised by the exact sequence

$$(2.18) \quad 0 \rightarrow TM \rightarrow TY|_M \rightarrow NM \rightarrow 0.$$

The tangent bundle TNM fits into the exact sequence

$$0 \rightarrow \pi^*NM \xrightarrow{i} TNM \xrightarrow{d\pi} \pi^*TM \rightarrow 0.$$

This induces an inclusion of the space normal vector fields of M into the space vector fields of NM ,

$$\tilde{\bullet} : C^\infty(NM) \hookrightarrow \text{Vect}(NM), \quad u \mapsto \tilde{u} := i(\pi^*u).$$

Here $\pi^*u \in C^\infty(\pi^*NM)$ is the pull back section. Observe that, $[\tilde{u}, \tilde{v}] = 0$ for all $u, v \in C^\infty(NM)$.

For any section $u \in C^\infty(NM)$ we define the **graph of u** by

$$\Gamma_u := \{(x, u(x)) \in NM : x \in M\}.$$

For any section $u \in C^\infty(NM)$, the graph, Γ_u is a submanifold of NM and the bundle $\pi^*NM|_{\Gamma_u}$ fits into the split exact sequence

$$(2.19) \quad 0 \longrightarrow T\Gamma_u \xrightarrow{\quad \overset{d(u \circ \pi)}{\curvearrowright} \quad} TNM|_{\Gamma_u} \xrightarrow{1-d(u \circ \pi)} \pi^*NM|_{\Gamma_u} \longrightarrow 0.$$

In particular, this induces an isomorphism $N\Gamma_u \cong \pi^*NM|_{\Gamma_u}$. Moreover, the composition $T\Gamma_u \rightarrow TNM|_{\Gamma_u} \xrightarrow{d\pi} \pi^*TM|_{\Gamma_u}$ is an isomorphism.

The choice of a Riemannian metric g on Y induces a splitting of the exact sequence (2.18), that is,

$$TY|_M = TM \perp NM.$$

Denote by \cdot^\parallel and \cdot^\perp the projections onto the first and second summands respectively. The Levi-Civita connection ∇ on $TY|_M$ decomposes as

$$\nabla = \begin{bmatrix} \nabla^\parallel & -\Pi^* \\ \Pi & \nabla^\perp \end{bmatrix}.$$

Here $\Pi \in \text{Hom}(S^2TM, NM)$ is the **second fundamental form** of M , ∇^\parallel is the Levi-Civita connection on M and ∇^\perp is the **normal connection** on NM .

A **tubular neighbourhood map** of M is a diffeomorphism between an open neighbourhood V_M of the zero section of the normal bundle NM of M which is convex in each fiber and an open neighbourhood U_M (tubular neighbourhood) of M in Y ,

$$\Upsilon_M : V_M \rightarrow U_M$$

that takes the zero section 0 to M and the composition

$$NM \rightarrow 0^*TNM \xrightarrow{d\Upsilon_M} T\Upsilon_M|_M \rightarrow NM$$

is the identity.

Let $\Upsilon_M : V_M \rightarrow U_M$ be a tubular neighbourhood map of M . We define

$$C^\infty(V_M) := \{u \in C^\infty(NM) : \Gamma_u \subset V_M\}.$$

Let $u \in C^\infty(V_M)$. Denote by M_u the submanifold $\Upsilon_M(\Gamma_u)$ of Y . Then there is a **canonical bundle isomorphism**

$$(2.20) \quad \begin{array}{ccc} NM & \xrightarrow{\Theta_u^M} & NM_u \\ \downarrow & & \downarrow \\ M & \xrightarrow{\Upsilon_M \circ u} & M_u. \end{array}$$

induced by the following commutative diagram of bundle isomorphisms.

$$\begin{array}{ccc} NM & \xrightarrow{\Theta_u^M} & NM_u \\ \pi^* \left(\downarrow u^* \right) & & \uparrow d\Upsilon_M \\ \pi^* NM|_{\Gamma_u} & \xleftarrow{1-d(u \circ \pi)} & N\Gamma_u. \end{array} \quad \spadesuit$$

Notation 2.21. By abusing notation we will always denote the above \tilde{u} again by u . ▶

Definition 2.22. Let (Y, ϕ) be an almost G_2 -manifold. Let \mathcal{S}_3 be the set of all 3-dimensional oriented, closed smooth submanifolds of Y .

We define the C^k -**topology** on the set \mathcal{S}_3 by specifying a basis, which is a collection of all the sets of the form $\{\Upsilon_P(\Gamma_u) : u \in \mathcal{V}_P^k\}$, where $P \in \mathcal{S}_3$, Υ_P is a tubular neighbourhood map of P , and \mathcal{V}_P^k is an open set in $C^\infty(V_P)$, whose topology is induced by the C^k -norm on $C^\infty(NP)$.

The C^∞ -**topology** on the set \mathcal{S}_3 is the inverse limit topology of C^k -topologies on it, that is, a set is open with C^∞ -topology if it is open for every C^k -topology.

The **moduli space** \mathcal{M}^ϕ of closed associative submanifolds in (Y, ϕ) is the subset of all submanifolds in \mathcal{S}_3 which are associatives. The C^∞ -topology on \mathcal{M}^ϕ is the subspace topology of \mathcal{S}_3 . Let \mathcal{P} be the set of all co-closed G_2 -structures on Y . Denote the universal moduli space of closed associative submanifolds by

$$\mathcal{M} := \{(\phi, P) \in \mathcal{P} \times \mathcal{S}_3 : P \in \mathcal{M}^\phi\}.$$

Equip \mathcal{P} with the C^∞ topology. The topology of $\mathcal{M} \subset \mathcal{P} \times \mathcal{S}_3$ is then given by the subspace topology of the product C^∞ -topologies.

Let \mathcal{P} be the space of paths $\phi : [0, 1] \rightarrow \mathcal{P}$ which are smooth as sections over $[0, 1] \times Y$. Set $\phi_t := \phi(t)$. Define the **1-parameter moduli space** of closed associative submanifolds by the fiber product

$$\mathcal{M}^\phi := [0, 1] \times_{\mathcal{P}} \mathcal{M} \cong \{(t, P) \in [0, 1] \times \mathcal{S}_3 : P \in \mathcal{M}^{\phi_t}\}.$$

The topology on \mathcal{M}^ϕ is the fiber product topology, which is same as the subspace topology of the product topology of $[0, 1] \times \mathcal{S}_3$. \spadesuit

Definition 2.23. Let M be an associative submanifold (compact or noncompact) of an almost G_2 -manifold (Y, ϕ) . The operator $\mathbf{D}_M : C^\infty(NM) \rightarrow C^\infty(NM)$ is defined by

$$\langle \mathbf{D}_M v, w \rangle_{L^2} := \int_M \left\langle \sum_{i=1}^3 e_i \times \nabla_{M, e_i}^\perp v, w \right\rangle + \int_M \iota_w \nabla_v \psi,$$

for all $v, w \in C_c^\infty(NM)$. Here NM is the normal bundle of M and ∇_M^\perp is the normal connection and $\{e_1, e_2, e_3\}$ is an oriented local orthonormal frame for TM with respect to the metric g_ϕ .

The operator \mathbf{D}_M is formally self adjoint if $d\psi = 0$ (see Proposition 3.88). If (Y, ϕ) is a G_2 -manifold then $\nabla \psi = 0$ and \mathbf{D}_M is a Dirac operator, called **Fueter operator**. A closed associative submanifold M of a co-closed G_2 -manifold is called **rigid** or equivalently **unobstructed** if $\ker \mathbf{D}_M = \{0\}$. \spadesuit

Remark 2.24. The operator \mathbf{D}_M is the linearization of a nonlinear map \mathfrak{F}_M^ϕ which controls the deformation theory of associative submanifolds (see Proposition 3.86). Let $\Upsilon_M : V_M \rightarrow U_M \subset Y$ be a tubular neighbourhood map. The map $\mathfrak{F}_M^\phi : C^\infty(V_M) \rightarrow C^\infty(NM)$ is defined by

$$(2.25) \quad \langle \mathfrak{F}_M^\phi u, w \rangle_{L^2} = \int_{\Gamma_u} \iota_w \Upsilon_M^* \psi, \quad u \in C^\infty(V_M), w \in C_c^\infty(NM)$$

The notation w in the integrand is the extension vector field of $w \in C^\infty(NM)$ in the tubular neighbourhood as in Notation 2.21. The associative submanifolds can also be thought of as critical points of a functional [DT98, Section 8; DW19, Section 2.3] on the space of submanifolds. The differential of this functional is a 1-form which is locally of the form \mathfrak{F}_M^ϕ . This led Doan and Walpuski [DW19] to make a proposal of constructing Floer homology groups whose chain complex is generated by associative submanifolds. \clubsuit

The following theorem summarizes the deformation theory of closed associative submanifolds.

Theorem 2.26 ([McL98, Theorem 5.2; Joy18, Theorem 2.12; DW19, Theorem 2.20, Proposition 2.23, Section 2.7]). *Let P be an associative submanifold of a co-closed G_2 -manifold (Y, ϕ) and $\phi : [0, 1] \rightarrow \mathcal{P}$ be a path of co-closed G_2 structures such that $\phi(t_0) = \phi$. Then there exists an open neighbourhood \mathcal{J}_P of 0 in $\ker \mathbf{D}_P$ such that*

- (i) *the moduli space \mathcal{M}^ϕ near P is homeomorphic to $\text{ob}_P^{-1}(0)$, zero set of a smooth map (obstruction map/Kuranishi map)*

$$\text{ob}_P : \mathcal{J}_P \rightarrow \text{coker } \mathbf{D}_P.$$

Moreover, there is a comeager subset $\mathcal{P}^{\text{reg}} \subset \mathcal{P}$ such that for all $\phi \in \mathcal{P}^{\text{reg}}$ the moduli space \mathcal{M}^ϕ is a 0-dimensional manifold and consisting only of unobstructed (rigid) closed associative submanifolds.

- (ii) there is a comeager subset $\mathcal{P}^{\text{reg}} \subset \mathcal{P}$ such that for all $\phi \in \mathcal{P}^{\text{reg}}$ the moduli space \mathcal{M}^ϕ is a 1-dimensional manifold and there is a discrete subset $I_o \subset [0, 1]$ having the property that
- for each $t \in [0, 1] \setminus I_o$ the moduli space \mathcal{M}^{ϕ_t} is consisting only unobstructed (rigid) closed associative submanifolds.
 - for each $\hat{t} \in I_o$ the moduli space $\mathcal{M}^{\phi_{\hat{t}}}$ is consisting only closed associative submanifolds \hat{P} having $\dim \ker \mathbf{D}_{\hat{P}} \leq 1$. If $P_o \in \mathcal{M}^{\phi_{\hat{t}}}$ and $\dim \ker \mathbf{D}_{P_o} = 1$ then there exist non zero constants a, b such that $\text{ob}_{(\hat{t}, P_o)}$ can be written as

$$\text{ob}_{(\hat{t}, P_o)}(t, x) = at + bx^2 + \text{higher order terms.}$$




Figure 3: 1-parameter moduli space of associative submanifolds.

The above Theorem 2.26 leads us to the following natural question, which is still open.

Question 2.27 (Joyce [Joy18, Conjecture 2.16]). Let (Y, ϕ) be a compact almost G_2 -manifold and $\tau \in \Omega^3(Y)$ be a closed 3-form. Let \mathcal{P}_τ be the space of all co-closed G_2 -structures that are tamed by τ . Does there exist a comeager subset $\mathcal{P}_\tau^\diamond \subset \mathcal{P}_\tau$ such that for all $\phi \in \mathcal{P}_\tau^\diamond$ the moduli space \mathcal{M}^ϕ is compact?

If (Y, ϕ) is a compact almost G_2 -manifold which is tamed by $\tau \in \Omega^3(Y)$, then there is a constant $c > 0$ such that for every closed associative submanifold P in (Y, ϕ) we have (see [DS11, Section 3.2; Joy18, Section 2.5])

$$\text{vol}(P, g_\phi) \leq c[\tau] \cdot [P].$$

Therefore, we can use the following theorem of geometric measure theory to get a Federer-Fleming compactification of \mathcal{M}^ϕ . For a discussion on the proof of the following theorem in the special Lagrangian context we refer the reader to [Joy04a, Section 6; DW21, Section 4].

Theorem 2.28 (Simon [Sim83, section 6, Section 32], Spolaor [Spo15], Adams and Simon [AS88, Theorem 1], Joyce [Joy04a, Theorem 6.8]). *Let P_n be a sequence of closed associative submanifolds in a sequence of compact almost G_2 -manifolds (Y, ϕ_n) which are tamed by a fixed $\tau \in \Omega^3(Y)$. Assume ϕ_n converges to ϕ in C^∞ -topology. Then after passing to a subsequence P_n converges (in the sense of currents) to a closed integral current P_∞ which is calibrated by ϕ . Moreover*

- (i) *the Hausdorff dimension of the singular set of P_∞ is at most 1.*
- (ii) *if all the tangent cones of P_∞ are Jacobi integrable⁹ multiplicity 1 associative cone with smooth link, then P_∞ is a conically singular associative submanifold of (Y, ϕ) , in the sense of Definition 3.53.*

The above Theorem 2.28(ii) gives a hint why we chose to study in this article the conically singular associative submanifolds.

3 Moduli space of conically singular associative submanifolds

3.1 Associative cones

Definition 3.1. Let Σ be a 2-dimensional closed submanifold of $S^6 \subset \mathbf{R}^7$. Define the inclusion map $\iota : (0, \infty) \times \Sigma \rightarrow \mathbf{R}^7$ by $\iota(r, \sigma) = r\sigma$. A cone C with link Σ is the image of ι in \mathbf{R}^7 . The Euclidean metric on \mathbf{R}^7 induces a metric g_Σ on Σ and a cone metric g_C on C , that is, $g_C = dr^2 + r^2g_\Sigma$. Furthermore, it induces a metric g_{NC} and a connection ∇_C^\perp on the normal bundle NC of the cone C . Let $\pi : \mathbf{R}^7 \setminus \{0\} \rightarrow S^6$ be the projection. Then $NC = \pi^*(N\Sigma)$, pullback of the normal bundle $N\Sigma$ of Σ in S^6 , $g_{NC} = r^2\pi^*g_{N\Sigma}$ and $\nabla_C^\perp = \pi^*\nabla_\Sigma^\perp$.

The standard almost complex structure on S^6 , $J : T_x S^6 \rightarrow T_x S^6$ is defined by the standard cross product ‘ \times ’ on \mathbf{R}^7 as follows:

$$J(v) = \partial_r \times v$$

where $x \in S^6$, $v \in T_x S^6 \subset \mathbf{R}^7$.

If the cone C is an associative submanifold then we call it an **associative cone**. This is equivalent to saying that the link Σ is a pseudoholomorphic curve in the almost complex manifold (S^6, J) . ♠

Example 3.2. Any special Lagrangian cone in \mathbf{C}^3 is an associative cone in $\mathbf{R}^7 = \mathbf{R} \oplus \mathbf{C}^3$ and its link is special Legendrian in S^5 . Example 3.3 and Example 3.4 describes two examples of special Lagrangian cones in \mathbf{C}^3 that are important for the desingularization theorems (see Theorem 1.29 and Theorem 1.21) discussed in this article. For more examples of special Lagrangian cones, see [Haso4a; HK07; Joy01; Joy02]. •

Example 3.3. (Transverse pair of SL planes [Joy03, page 328]) Let C_\times be the union of a pair of special Lagrangian (SL) planes in \mathbf{C}^3 with transverse intersection at the origin. Then there exist

⁹i.e. A cone in \mathbf{R}^7 with smooth link $\Sigma \subset S^6$ is called Jacobi integrable if for every $v_\Sigma \in \ker(D_\Sigma + 2J) \subset C^\infty(N\Sigma)$, $\{\exp(tv_\Sigma) : |t| \ll 1\}$ is a 1-parameter family of holomorphic curves in (S^6, J) . Here ‘exp’ is the exponential map with respect to the round metric on S^6 , and $D_\Sigma + 2J$ is the deformation operator controlling the deformation theory of holomorphic curves in (S^6, J) (see Proposition 3.86).

a $B \in SU(3)$ and unique $\theta_1, \theta_2, \theta_3 \in (0, \pi)$ satisfying $\theta_1 \leq \theta_2 \leq \theta_3$ and $\theta_1 + \theta_2 + \theta_3 = \pi$ such that $C_\times = B\Pi_0 \cup B\Pi_\theta$, where

$$\Pi_0 = \mathbf{R}^3 \quad \text{and} \quad \Pi_\theta := \text{diag}(e^{i\theta_1}, e^{i\theta_2}, e^{i\theta_3}) \cdot \mathbf{R}^3.$$

We define $\Pi_+ := B\Pi_0$ and $\Pi_- := B\Pi_\theta$. Note that Π_\pm are uniquely determined. •

Example 3.4. (Harvey-Lawson T^2 -cone [HL82, Theorem 3.1])

$$\begin{aligned} C_{HL} &:= \{(z_1, z_2, z_3) \in \mathbf{C}^3 : |z_1| = |z_2| = |z_3|, z_1 z_2 z_3 \in (0, \infty)\} \\ &= \{r(e^{i\theta_1}, e^{i\theta_2}, e^{-i(\theta_1+\theta_2)}) \in \mathbf{C}^3 : r \in (0, \infty), \theta_1, \theta_2 \in [0, 2\pi)\} \end{aligned}$$

is a special Lagrangian cone in \mathbf{C}^3 with link $\Sigma_{HL} = C_{HL} \cap S^5$ isometric to the flat Clifford torus T^2 . •

Example 3.5. (Null torsion holomorphic curves [Bry82, Section 4]) We will be following here the exposition about null torsion holomorphic curves given in [Mad21]. Let Σ be a closed holomorphic curve in (S^6, J) . Then the characteristic $SU(3)$ connection $\tilde{\nabla}$ (see (3.9)) induces holomorphic structures on $TS|_\Sigma, T\Sigma$ and $N\Sigma$. The second fundamental form of Σ is the obstruction $\Pi \in \text{Ext}^1(N\Sigma, T\Sigma) \cong H^0(\Sigma, K_\Sigma^2 \otimes N\Sigma)$ (by Serre duality) to the holomorphic splitting of the following exact sequence:

$$0 \rightarrow T\Sigma \rightarrow TS|_\Sigma \rightarrow N\Sigma \rightarrow 0.$$

Moreover, $\Pi \neq 0$ if and only if Σ is not a totally geodesic S^2 . In this case, denote the effective divisor of the zero set of Π by Z . We define a holomorphic line bundle L_B by the following exact sequence:

$$(3.6) \quad 0 \rightarrow L_N := K_\Sigma^{-2} \otimes \mathcal{O}(Z) \hookrightarrow N\Sigma \rightarrow L_B \rightarrow 0.$$

The torsion on Σ is the obstruction $\text{III} \in \text{Ext}^1(L_B, L_N) \cong H^0(\Sigma, K_\Sigma^3 \otimes \mathcal{O}(-Z) \otimes L_B)$ (by Serre duality) to the holomorphic splitting of the exact sequence (3.6).

If Σ is **null-torsion** (i.e. $\Pi \neq 0, \text{III} = 0$), then there is a holomorphic isomorphism

$$L_B \cong K_\Sigma^3 \otimes \mathcal{O}(-Z) \quad \text{and} \quad \text{Area}(\Sigma) = 4\pi b \geq 24\pi,$$

where $b = -c_1(L_B) = 3\chi(\Sigma) + [Z]$. Moreover, no null torsion holomorphic curves in S^6 are contained in a totally geodesic S^5 . If Σ is genus zero and not a totally geodesic S^2 , then it must be a null-torsion holomorphic curve. Bryant [Bry82, Theorem 4.10] and later Rowland [Row99] proved that closed Riemann surfaces of any genus can be conformally embedded as a null-torsion J -holomorphic curve in S^6 . •

There are more examples of associative cones which are not special Lagrangians discussed in [Lot11, Section 7; Loto7b].

3.1.1 Moduli space of J -holomorphic curves in S^6 .

Definition 3.7. Let \mathcal{S} be the set of all 2-dimensional oriented, closed smooth submanifolds of S^6 . Equip \mathcal{S} with C^∞ -topology in the same way as in Definition 2.22. The **moduli space** \mathcal{M}^{hol} of embedded holomorphic curves in (S^6, J) is the subset of all submanifolds Σ in \mathcal{S} which are J -holomorphic. The topology on \mathcal{M}^{hol} is the subspace topology of the above. \spadesuit

Let Σ be a pseudoholomorphic curve in S^6 , that is, $\Sigma \in \mathcal{M}^{\text{hol}}$. We denote the complex structure on Σ by j , which is just the restriction of J . Let $\Upsilon_\Sigma : V_\Sigma \rightarrow U_\Sigma$ be a tubular neighbourhood map of Σ . For $u \in C^\infty(V_\Sigma)$ denote the submanifold $\Upsilon_\Sigma(\Gamma_u)$ by Σ_u . Note that Σ_u is J -holomorphic if and only if u satisfies the following **non-linear Cauchy-Riemann equation**

$$0 = \bar{\partial}_J u := \frac{1}{2}(du + \Upsilon_\Sigma^* J(u) \circ du \circ j) \in C^\infty(\overline{\text{Hom}}_{\mathbb{C}}(T\Sigma, u^*TV_\Sigma)).$$

The linearization of the nonlinear map $\bar{\partial}_J : C^\infty(V_\Sigma) \rightarrow C^\infty(\overline{\text{Hom}}_{\mathbb{C}}(T\Sigma, u^*TV_\Sigma))$ at the zero section, is described in [MS12, Proposition 3.1.1]. This is the linear map $\mathfrak{d}_{\Sigma, J} : C^\infty(N\Sigma) \rightarrow C^\infty(\overline{\text{Hom}}_{\mathbb{C}}(T\Sigma, TV_{\Sigma|_\Sigma}))$ defined by

$$\mathfrak{d}_{\Sigma, J} \xi := \frac{1}{2}(\nabla_\Sigma \xi + J \circ (\nabla_\Sigma \xi) \circ j + \nabla_\xi J \circ j), \quad \xi \in C^\infty(N\Sigma).$$

The tangential component of $\mathfrak{d}_{\Sigma, J}$ can be discarded for the deformation theory. The following normal component actually controls the deformation theory.

Definition 3.8. For $\Sigma \in \mathcal{M}^{\text{hol}}$ the **normal Cauchy-Riemann operator** $\bar{\partial}_\nabla^N : C^\infty(N\Sigma) \rightarrow C^\infty(\overline{\text{Hom}}_{\mathbb{C}}(T\Sigma, N\Sigma)) \cong \Omega^{0,1}(\Sigma, N\Sigma)$ is defined by

$$\bar{\partial}_\nabla^N \xi := \frac{1}{2}(\nabla_\Sigma^\perp \xi + J \circ (\nabla_\Sigma^\perp \xi) \circ j + \nabla_\xi^\perp J \circ j). \quad \spadesuit$$

The moduli space of holomorphic curves usually studied using the non-linear Cauchy-Riemann map (see [MS12, Theorem 3.1.5]) but here we study this in a different way. On \mathbb{R}^7 , the Euclidean metric is $g_e = dr^2 + r^2 g$ and the standard G_2 -structure $\phi_e, \psi_e = *_g \phi_e$ can be written as

$$\phi_e = r^2 dr \wedge \omega + r^3 \text{Re } \Omega, \quad \psi_e = \frac{r^4}{2} \omega^2 - r^3 dr \wedge \text{Im } \Omega$$

where $\omega(X, Y) := g(JX, Y)$, and Ω is a nowhere vanishing complex $(3, 0)$ -form. Together they give an $SU(3)$ -structure on S^6 . Also

$$\text{vol}_{\mathbb{R}^7} = r^6 dr \wedge \text{vol}_{S^6} \quad \text{and} \quad \text{vol}_{S^6} = \frac{\omega^3}{6} = \frac{i}{8} \Omega \wedge \bar{\Omega} = \frac{1}{4} \text{Re } \Omega \wedge \text{Im } \Omega.$$

In particular, $*_g \text{Re } \Omega = \text{Im } \Omega$, $*_g \text{Im } \Omega = -\text{Re } \Omega$. Moreover, $d\phi_e = 0$ and $d\psi_e = 0$ on \mathbb{R}^7 are equivalent to the following equations on S^6 respectively:

$$d\omega = 3 \text{Re } \Omega \quad \text{and} \quad d \text{Im } \Omega = -2\omega^2.$$

That means S^6 with this $SU(3)$ structure is a **nearly Kähler** manifold. Let ∇ be the Levi-Civita connection on S^6 with respect to the round metric g . The **characteristic $SU(3)$ connection**

$$(3.9) \quad \tilde{\nabla}_u v := \nabla_u v + \frac{1}{2}(\nabla_u J)Jv,$$

satisfies $\tilde{\nabla}g = 0$, $\tilde{\nabla}J = 0$ and $\text{Hol}(\tilde{\nabla}) \subset SU(3)$. But $\tilde{\nabla}$ is not torsion free.

Definition 3.10. Let Σ be a J -holomorphic embedded curve in S^6 . The **characteristic Cauchy-Riemann operator** on $N\Sigma$, $\bar{\partial}_{\tilde{\nabla}}^N : C^\infty(N\Sigma) \rightarrow \Omega^{0,1}(N\Sigma)$ is defined to be the induced Cauchy-Riemann operator from the characteristic $SU(3)$ connection $\tilde{\nabla}$, that is,

$$\bar{\partial}_{\tilde{\nabla}}^N \xi := \frac{1}{2}(\tilde{\nabla}^\perp \xi + J \circ (\tilde{\nabla}^\perp \xi) \circ j). \quad \spadesuit$$

Lemma 3.11 (McDuff and Salamon [MS12, Lemma C.7.1]). *For any vector fields u, v, w on S^6 ,*

- (i) $(\nabla_u J)Jv = -J(\nabla_u J)v$ and $g((\nabla_u J)v, w) = -g((\nabla_u J)w, v)$,
- (ii) $(\nabla_u J)v = -(\nabla_v J)u$ and the torsion $T_{\tilde{\nabla}}(u, v) = \frac{1}{4}N_J(u, v) = (\nabla_u J)Jv$,
- (iii) $3g((\nabla_u J)v, w) = d\omega(u, v, w) = 3 \text{Re } \Omega(u, v, w)$

Definition 3.12. The J -antilinear multiplication map, $\times_{S^6} : TS^6 \times TS^6 \rightarrow TS^6$ is defined by the orthogonal projection on TS^6 of the cross product in \mathbf{R}^7 , or equivalently for all vector fields u, v, w on S^6 ,

$$g(u \times_{S^6} v, w) = \text{Re } \Omega(u, v, w), \text{ or equivalently } u \times_{S^6} v = (\nabla_u J)v. \quad \spadesuit$$

Remark 3.13. Let Σ be an oriented smooth surface in S^6 and C be the cone in \mathbf{R}^7 with link Σ . The following are equivalent: (i) Σ is J -holomorphic, (ii) C is associative, (iii) for all $u, v \in T\Sigma$ and $w \in TS^6$, $\text{Re } \Omega(u, v, w) = 0$, (iv) for all $u, v \in T\Sigma$, $u \times_{S^6} v = 0$, (v) for all $u \in T\Sigma$ and $v \in N\Sigma$, $u \times v \in N\Sigma$. \clubsuit

Proposition 3.14. *Let Σ be an embedded J -holomorphic curve in S^6 . Then for all $\xi \in C^\infty(N\Sigma)$*

$$\bar{\partial}_{\tilde{\nabla}}^N \xi = \bar{\partial}_{\nabla}^N \xi - (\nabla^\perp J)J\xi.$$

Proof. By Lemma 3.11 we have for all $\xi \in C^\infty(N\Sigma)$,

$$J \circ \tilde{\nabla} \xi \circ j = J \circ \nabla \xi \circ j + \frac{1}{2}J(\nabla J)J\xi \circ j = J \circ \nabla \xi \circ j + \frac{1}{2}(\nabla J)J\xi.$$

The proposition now follows from (3.9). \blacksquare

Definition 3.15. Let Σ be an embedded J -holomorphic curve in Z . By Lemma 3.11 and Remark 3.13, the map $\gamma_\Sigma : T\Sigma \rightarrow \text{End}_{\mathbb{C}}(N\Sigma)$ given by

$$\gamma_\Sigma(f_\Sigma)(v_\Sigma) := f_\Sigma \times v_\Sigma, \quad \forall v_\Sigma \in N\Sigma, \quad f_\Sigma \in T\Sigma,$$

is a skew symmetric J -anti-linear Clifford multiplication. Moreover the normal bundle $N\Sigma$ together with the metric $g_{N\Sigma}$, Clifford multiplication γ_Σ and the metric connection $\tilde{\nabla}^\perp := \tilde{\nabla}_\Sigma^\perp$ is a Dirac bundle, that is, $\tilde{\nabla}^\perp \gamma_\Sigma = 0$. The associated Dirac operator is given by

$$(3.16) \quad \mathbf{D}_\Sigma := \sum_{i=1}^2 f_i \times \tilde{\nabla}_{f_i}^\perp,$$

where $\{f_i\}$ is a local orthonormal oriented frame on Σ . Note that \mathbf{D}_Σ is J -anti-linear. Moreover the map γ_Σ induces a J -anti-linear isomorphism

$$\gamma_\Sigma : \overline{\text{Hom}}_{\mathbb{C}}(T\Sigma, N\Sigma) \rightarrow N\Sigma,$$

which is defined by

$$\gamma_\Sigma(f_\Sigma^* \otimes v_\Sigma) = \gamma_\Sigma(f_\Sigma)(v_\Sigma). \quad \spadesuit$$

Remark 3.17. γ_Σ satisfies the following commutative diagram:

$$\begin{array}{ccc} C^\infty(N\Sigma) & \xrightarrow{\bar{\partial}_{\tilde{\nabla}^\perp}^N} & C^\infty(\overline{\text{Hom}}_{\mathbb{C}}(T\Sigma, N\Sigma)) \\ & \searrow \mathbf{D}_\Sigma & \downarrow \gamma_\Sigma \\ & & C^\infty(N\Sigma). \end{array} \quad \clubsuit$$

Proposition 3.18. $\mathbf{D}_\Sigma = \sum_{i=1}^2 f_i \times \nabla_{f_i}^\perp - J$ and $\gamma_\Sigma \circ \bar{\partial}_{\tilde{\nabla}^\perp}^N = \mathbf{D}_\Sigma + 2J$.

Proof. First we prove that $\gamma_\Sigma((\nabla^\perp J)J) = -2J$. For a local oriented orthonormal frame $\{f_1, f_2 = jf_1\}$ on Σ we have $\nabla_{f_i}^\perp J = \gamma_\Sigma f_i$ because of Definition 3.12 and Lemma 3.11. Therefore

$$\gamma_\Sigma((\nabla^\perp J)J) = \sum_{i=1}^2 f_i \times (\nabla_{f_i}^\perp J)J = \sum_{i=1}^2 (\gamma_\Sigma f_i)^2 J = -2J.$$

Now the first equality in the proposition follows from (3.16) and (3.9). The second equality follows from Proposition 3.14 and Remark 3.17. \blacksquare

Definition 3.19. For $\Sigma \in \mathcal{S}$, we define $\mathcal{F} : C^\infty(V_\Sigma) \rightarrow C^\infty(N\Sigma)$ by

$$\langle \mathcal{F}(u), v \rangle_{L^2} := \int_{\Gamma_u} \iota_v(Y_\Sigma^* \text{Re } \Omega)$$

where $u \in C^\infty(V_\Sigma)$ and $v \in C^\infty(N\Sigma)$. The notation v in the integrand is the extension vector field of $v \in C^\infty(N\Sigma)$ in the tubular neighbourhood as in Notation 2.21. \spadesuit

Proposition 3.20. For $u \in C^\infty(V_\Sigma)$, we have $\mathcal{F}(u) = 0$ if and only if $\Sigma_u := \Upsilon_\Sigma(\Gamma_u)$ is J -holomorphic. If $\Sigma \in \mathcal{M}^{\text{hol}}$, then the linearization of \mathcal{F} at zero, $d\mathcal{F}_0 : C^\infty(N\Sigma) \rightarrow C^\infty(N\Sigma)$ is given by

$$d\mathcal{F}_0 = J\mathbf{D}_\Sigma - 2J = J(\mathbf{D}_\Sigma + 2J).$$

This is a formally self adjoint first order elliptic operator of index 0.

Proof. The first part follows from Remark 3.13. Let $\{f_1, f_2 = Jf_1\}$ be a local oriented orthonormal frame for $T\Sigma$. For $u, v \in C^\infty(N\Sigma)$,

$$\frac{d}{dt}\Big|_{t=0} \langle \mathcal{F}(tu), v \rangle_{L^2} = \frac{d}{dt}\Big|_{t=0} \int_{\Gamma_{tu}} \iota_v(\Upsilon_\Sigma^* \text{Re } \Omega) = \int_\Sigma \mathcal{L}_u \iota_v(\Upsilon_\Sigma^* \text{Re } \Omega).$$

This is same as $\int_\Sigma \iota_v \mathcal{L}_u(\Upsilon_\Sigma^* \text{Re } \Omega) + \iota_{[u,v]}(\Upsilon_\Sigma^* \text{Re } \Omega)$. As, $[u, v] = 0$ (by Notation 2.21) and $\Sigma \in \mathcal{M}^{\text{hol}}$, this is equal to

$$\begin{aligned} & \int_\Sigma \iota_v(\Upsilon_\Sigma^* \text{Re } \Omega)(\nabla_{f_1} u, f_2) + \iota_v(\Upsilon_\Sigma^* \text{Re } \Omega)(f_1, \nabla_{f_2} u) + \iota_v \nabla_u(\Upsilon_\Sigma^* \text{Re } \Omega) \\ &= \int_\Sigma -\langle f_2 \times \nabla_{f_1}^\perp u, v \rangle + \langle f_1 \times \nabla_{f_2}^\perp u, v \rangle - \iota_v(u \wedge \Upsilon_\Sigma^* \omega) \quad (\text{as } \nabla \text{Re } \Omega = -\omega \wedge \cdot) \\ &= \int_\Sigma \langle J(f_1 \times \nabla_{f_1}^\perp u), v \rangle + \langle J(f_2 \times \nabla_{f_2}^\perp u), v \rangle - \langle u, v \rangle = \int_\Sigma \langle (JD_\Sigma - 2\text{Id})u, v \rangle. \end{aligned}$$

Therefore $d\mathcal{F}_0 = JD_\Sigma - 2\text{Id}$. Since D_Σ is a J -anti-linear (this follows from the fact that $\tilde{\nabla}J = 0$ and γ_Σ is J -anti-linear), self adjoint Dirac operator, it proves the last part of the proposition. ■

Proof of Theorem 1.3. In Proposition 3.18 we see that $\gamma_\Sigma \circ \bar{\partial}_V^N = D_\Sigma + 2J$. Extending the nonlinear map \mathcal{F} to Hölder spaces we get a smooth map

$$\mathcal{F} : C^{2,\gamma}(V_\Sigma) \rightarrow C^{1,\gamma}(N\Sigma).$$

Proposition 3.20 implies that the linearization of \mathcal{F} at zero is an elliptic operator and hence is Fredholm. By implicit function theorem applied to \mathcal{F} we obtain the map ob_Σ as stated in the theorem (see [DK90, Proposition 4.2.19]). We need to only prove that, if $u \in C^{2,\gamma}(V_\Sigma)$ with $\mathcal{F}(u) = 0$, then $u \in C^\infty(V_\Sigma)$. To prove this, we observe

$$0 = D_\Sigma(\mathcal{F}(u)) = a(u, \nabla_\Sigma^\perp u)(\nabla_\Sigma^\perp)^2 u + b(u, \nabla_\Sigma^\perp u)$$

Since $a(u, \nabla_\Sigma^\perp u) \in C^{1,\gamma}$ and $b(u, \nabla_\Sigma^\perp u) \in C^{1,\gamma}$, by Schauder elliptic regularity (see [Joy07, Theorem 1.4.2]) we obtain $u \in C^{3,\gamma}$. By repeating this argument we get higher regularity, which completes the proof of the theorem. ■

3.1.2 Fueter operator on the cone C

The operator controlling the deformation theory for AC and CS associatives will be AC and CS uniformly elliptic operator asymptotic to a conical elliptic operator (see Definition 3.64). This conical operator is the Fueter operator on the asymptotic associative cone. The Fredholm theory of AC and CS uniformly elliptic operator has been studied by Lockhart and McOwen [LM85], see also [Mar02; KL20]. It suggests that we must study the indicial roots and homogeneous kernels of this Fueter operator.

Let C be an associative cone in \mathbf{R}^7 with link Σ . The Fueter operator (see Definition 2.23) on the cone C , $D_C : C^\infty(NC) \rightarrow C^\infty(NC)$ is

$$D_C = \sum_{i=1}^3 e_i \times \nabla_{C, e_i}^\perp,$$

where $\{e_i\}$ is a local oriented orthonormal frame on C and ∇_C^\perp is the normal connection induced by the Levi-Civita connection on \mathbf{R}^7 .

Definition 3.21. For $\lambda \in \mathbf{R}$ we define the **homogeneous kernel**¹⁰ of rate λ by

$$V_\lambda := \{r^{\lambda-1}v_\Sigma \in C^\infty(NC) : v_\Sigma \in C^\infty(N_{S^6}\Sigma), \mathbf{D}_C(r^{\lambda-1}v_\Sigma) = 0\}$$

and

$$d_\lambda := \dim V_\lambda.$$

The set of **indicial roots** is defined by

$$\mathcal{D}_C := \{\lambda \in \mathbf{R} : d_\lambda \neq 0\}. \quad \spadesuit$$

Proposition 3.22. Let v_Σ be an element in $C^\infty(N_{S^6}\Sigma)$ and $\lambda \in \mathbf{R}$. Then the following hold

(i) $\mathbf{D}_C = J\partial_r + \frac{1}{r}\mathbf{D}_\Sigma + \frac{2}{r}J.$

(ii) $\mathbf{D}_C(r^{\lambda-1}v_\Sigma) = r^{\lambda-2}(\mathbf{D}_\Sigma v_\Sigma + (\lambda+1)Jv_\Sigma).$

(iii) $\mathbf{D}_C(r^{\lambda-1}(\log r)^j v_\Sigma) = r^{\lambda-2}(\log r)^j (\mathbf{D}_\Sigma v_\Sigma + (\lambda+1)Jv_\Sigma) + jr^{\lambda-2}(\log r)^{j-1}Jv_\Sigma.$

(iv) $\mathbf{D}_\Sigma(Jv_\Sigma) = -J\mathbf{D}_\Sigma v_\Sigma.$

(v) $V_\lambda = \{r^{\lambda-1}v_\Sigma : \mathbf{D}_\Sigma v_\Sigma = -(\lambda+1)Jv_\Sigma\} = \{r^{\lambda-1}v_\Sigma : (J\mathbf{D}_\Sigma)v_\Sigma = (\lambda+1)v_\Sigma\}$

(vi) $JV_{-1+\lambda} = V_{-1-\lambda}$ and $d_{-1+\lambda} = d_{-1-\lambda}$ for all $\lambda \in \mathbf{R}$.

Proof. We have $\nabla_C^\perp = dr \otimes \partial_r + \nabla_\Sigma^\perp$. For a local oriented orthonormal frame $\{f_i\}$ on $\Sigma \subset S^6$ and $v(r, \sigma) \in C^\infty(NC)$, applying Proposition 3.18 we compute

$$\begin{aligned} \mathbf{D}_C v(r, \sigma) &= \partial_r \times \nabla_{C, \partial_r}^\perp v(r, \sigma) + \frac{1}{r^2} \sum_{i=1}^2 f_i \times \nabla_{C, f_i}^\perp v(r, \sigma) \\ &= \partial_r \times_{S^6} \partial_r v(r, \sigma) + \frac{1}{r} \partial_r \times_{S^6} v(r, \sigma) + \frac{1}{r} \sum_{i=1}^2 f_i \times_{S^6} \nabla_{\Sigma, f_i}^\perp v(r, \sigma) \\ &= J\partial_r v(r, \sigma) + \frac{1}{r}\mathbf{D}_\Sigma v(r, \sigma) + \frac{2}{r}J. \end{aligned}$$

This proves (i). Now (ii) and (iii) follows from (i), indeed

$$\mathbf{D}_C(r^{\lambda-1}(\log r)^j v_\Sigma) = r^{\lambda-2}(\log r)^j ((\lambda-1)Jv_\Sigma + \mathbf{D}_\Sigma v_\Sigma + 2Jv_\Sigma) + jr^{\lambda-2}(\log r)^{j-1}Jv_\Sigma.$$

Finally, (v) follows from (ii) and (vi) follows from (iv). And (iv) follows from the fact that $\tilde{\nabla}J = 0$ and γ_Σ is J -anti-linear. \blacksquare

Proposition 3.23. Let $m \in \mathbf{N}_0$ and $v(r, \sigma) = \sum_{j=0}^m r^{\lambda-1}(\log r)^j v_{\Sigma, j} \in C^\infty(NC)$ and $v_{\Sigma, j} \in C^\infty(N_{S^6}\Sigma)$.

If $\mathbf{D}_C v(r, \sigma) = 0$ then $m = 0$ and therefore $v(r, \sigma) = r^{\lambda-1}v_{\Sigma, 0}$.

¹⁰Note that, $|r^{\lambda-1}v_\Sigma|_{g_{NC}} = r^\lambda |v_\Sigma|_{g_\Sigma}$.

Proof. If $\mathbf{D}_C v(r, \sigma) = 0$, then by Proposition 3.22(iii) and comparing the coefficients of $r^{\lambda-2}(\log r)^{j-1}$, $j \geq 1$ we see that

$$\mathbf{D}_\Sigma v_{\Sigma, m} + (\lambda + 1)Jv_{\Sigma, m} = 0 \text{ and } jJv_{\Sigma, j} + \mathbf{D}_\Sigma v_{\Sigma, j-1} + (\lambda + 1)Jv_{\Sigma, j-1} = 0.$$

Therefore,

$$\begin{aligned} m\|v_{\Sigma, m}\|_{L^2(\Sigma)}^2 &= -\langle Jv_{\Sigma, m}, \mathbf{D}_\Sigma v_{\Sigma, m-1} + (\lambda + 1)Jv_{\Sigma, m-1} \rangle_{L^2(\Sigma)} \\ &= \langle J(\mathbf{D}_\Sigma v_{\Sigma, m} + (\lambda + 1)Jv_{\Sigma, m}), v_{\Sigma, m-1} \rangle_{L^2(\Sigma)} = 0. \end{aligned}$$

Here we have used Proposition 3.22(iv). The proof is complete by backwards induction starting with $j = m$. \blacksquare

Remark 3.24. Although it follows from [LM85, Equation 1.11] that a general element in the homogeneous kernel is of the form $\sum_{j=0}^m r^{\lambda-1}(\log r)^j v_{\Sigma, j}$, Proposition 3.23 implies that it must be of the form we have defined in Definition 3.21. We also see that there is a canonical one to one correspondence between the indicial roots and eigenvalues of the self adjoint elliptic operator $J\mathbf{D}_\Sigma - 1$. In particular they are countable, discrete and will have finite intersection with any closed bounded interval of \mathbf{R} . Moreover, Proposition 3.22(vi) implies that the indicial roots and homogeneous kernels are symmetric with respect to -1 . \clubsuit

Definition 3.25. Let L be a special Lagrangian submanifold in $\mathbf{C}^3 \subset \mathbf{R} \oplus \mathbf{C}^3$. Set $e_1 = (1, 0) \in \mathbf{R} \oplus \mathbf{C}^3$. We define the isometry $\Phi_L : C^\infty(NL) \rightarrow \Omega^0(L, \mathbf{R}) \oplus \Omega^1(L, \mathbf{R})$ by

$$\Phi_L(v) := (\langle e_1, v \rangle, -(e_1 \times v)^b).$$

Also we define $\check{\mathbf{D}}_L : \Omega^0(L, \mathbf{R}) \oplus \Omega^1(L, \mathbf{R}) \rightarrow \Omega^0(L, \mathbf{R}) \oplus \Omega^1(L, \mathbf{R})$ by $\check{\mathbf{D}}_L := \Phi_L \mathbf{D}_L \Phi_L^{-1}$. \spadesuit

The operator $\check{\mathbf{D}}_L$ can be written as (for the proof see [Ber22, Proposition 3.67])

$$\check{\mathbf{D}}_L = \begin{bmatrix} 0 & d^* \\ d & *d \end{bmatrix}.$$

A direct computation yields the following lemma.

Lemma 3.26. *Let C be a cone in \mathbf{C}^3 with link Σ , then for $(f_\Sigma, h_\Sigma, \sigma_\Sigma) \in \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^1(\Sigma, \mathbf{R})$ we have*

- (i) $d(r^\lambda f_\Sigma) = \lambda r^{\lambda-1} f_\Sigma dr + r^\lambda d_\Sigma f_\Sigma$.
- (ii) $d(r^\lambda h_\Sigma dr + r^{\lambda+1} \sigma_\Sigma) = -r^\lambda dr \wedge d_\Sigma h_\Sigma + (\lambda + 1)r^\lambda dr \wedge \sigma_\Sigma + r^{\lambda+1} d_\Sigma \sigma_\Sigma$.
- (iii) $*dr = r^2 *_\Sigma 1$, $*\sigma_\Sigma = -dr \wedge *_\Sigma \sigma_\Sigma$, $*(dr \wedge \sigma_\Sigma) = *_\Sigma \sigma_\Sigma$, $*d_\Sigma \sigma_\Sigma = r^{-2} dr$.
- (iv) $d^*(r^\lambda h_\Sigma dr + r^{\lambda+1} \sigma_\Sigma) = -(\lambda + 2)r^{\lambda-1} h_\Sigma + r^{\lambda-1} d_\Sigma^* (\sigma_\Sigma)$.
- (v) $*d(r^\lambda h_\Sigma dr + r^{\lambda+1} \sigma_\Sigma) = -r^\lambda *_\Sigma d_\Sigma h_\Sigma + (\lambda + 1)r^\lambda *_\Sigma \sigma_\Sigma + r^{\lambda-1} (*_\Sigma d_\Sigma \sigma_\Sigma) dr$.

Proposition 3.27. *If C is a special Lagrangian cone in \mathbf{C}^3 whose link is Σ , then*

$$V_\lambda \cong \begin{cases} \{\sigma_\Sigma \in \Omega^1(\Sigma, \mathbf{R}) : \Delta_\Sigma \sigma_\Sigma = 0\} & \text{if } \lambda = -1, \\ \{(f_\Sigma, h_\Sigma) \in \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^0(\Sigma, \mathbf{R}) : (f_\Sigma, h_\Sigma) \text{ satisfies (3.28)}\} & \text{if } \lambda \neq -1, \end{cases}$$

$$(3.28) \quad \Delta_\Sigma f_\Sigma = \lambda(\lambda + 1)f_\Sigma, \quad \Delta_\Sigma h_\Sigma = (\lambda + 2)(\lambda + 1)h_\Sigma.$$

Proof. Since C is a special Lagrangian,

$$V_\lambda \cong \{(f_\Sigma, h_\Sigma, \sigma_\Sigma) \in \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^1(\Sigma, \mathbf{R}) : (f_\Sigma, h_\Sigma, \sigma_\Sigma) \text{ satisfies (3.29)}\},$$

$$(3.29) \quad \begin{cases} d^*(r^\lambda h_\Sigma dr + r^{\lambda+1} \sigma_\Sigma) = 0 \\ d(r^\lambda f_\Sigma) + *d(r^\lambda h_\Sigma dr + r^{\lambda+1} \sigma_\Sigma) = 0. \end{cases}$$

By Lemma 3.26 we obtain that (3.29) is equivalent to the following:

$$(3.30) \quad \begin{cases} (\lambda + 2)h_\Sigma = d_\Sigma^{*\Sigma}(\sigma_\Sigma) \\ \lambda f_\Sigma = -*_\Sigma d_\Sigma(\sigma_\Sigma) \\ (\lambda + 1)*_\Sigma \sigma_\Sigma = -d_\Sigma f_\Sigma + *_\Sigma d_\Sigma h_\Sigma. \end{cases}$$

This yields the required proposition. ■

Corollary 3.31. *If C is a special Lagrangian cone in \mathbf{C}^3 whose link is Σ , then*

$$V_\lambda \cong \begin{cases} \{h_\Sigma \in \Omega^0(\Sigma, \mathbf{R}) : \Delta_\Sigma h_\Sigma = (\lambda + 2)(\lambda + 1)h_\Sigma\} & \text{if } \lambda \in (-1, 0) \\ H^0(\Sigma, \mathbf{R}) \oplus \{h_\Sigma \in \Omega^0(\Sigma, \mathbf{R}) : \Delta_\Sigma h_\Sigma = 2h_\Sigma\} & \text{if } \lambda = 0 \\ H^1(\Sigma, \mathbf{R}) & \text{if } \lambda = -1 \\ \{f_\Sigma \in \Omega^0(\Sigma, \mathbf{R}) : \Delta_\Sigma f_\Sigma = (\lambda + 1)\lambda f_\Sigma\} & \text{if } \lambda \in (-2, -1). \end{cases} \quad \blacksquare$$

Remark 3.32. Since d_1 is the dimension of the space of infinitesimal deformations of Σ in S^6 ,

$$\frac{d_{-1}}{2} + \sum_{-1 < \lambda < 1} d_\lambda - 7 \leq \text{s-ind}_\Delta(C) \leq \text{s-ind}(C).$$

If the associative cone C is not a 3-plane then all the translations by vectors in \mathbf{R}^7 yield a 7-dimensional subspace of V_0 and hence $d_0 \geq 7$. Indeed, there are no non-trivial translations which preserves the cone C because 0 is the unique singular point of C . Thus if C is not a 3-plane,

$$\frac{d_{-1}}{2} \leq \text{s-ind}_\Delta(C) \leq \text{s-ind}(C). \quad \clubsuit$$

Example 3.33. (Pair of transverse associative planes) Let Π_{\pm} be a pair of associative planes in \mathbf{R}^7 with transverse intersection at the origin that is $\Pi_+ \cap \Pi_- = \{0\}$. Set $C_{\times} := \Pi_+ \cup \Pi_-$ and $e_1 := (1, 0, \dots, 0) \in \mathbf{R}^7 = \mathbf{R} \oplus \mathbf{C}^3$. We choose a unit vector $\mathbf{n} \in \mathbf{R}^7$ orthogonal to both Π_{\pm} so that we have an orientation compatible splitting

$$\mathbf{R}^7 = \langle \mathbf{n} \rangle_{\mathbf{R}} \oplus \Pi_+ \oplus \Pi_-.$$

Here the orientations of Π_{\pm} are given by the restrictions of the standard 3-form ϕ_e and orientation of $\langle \mathbf{n} \rangle_{\mathbf{R}}$ is given by \mathbf{n} . If we choose $-\mathbf{n}$ instead of \mathbf{n} in the above, we interchange the role of Π_+ and Π_- . Then there exists $B \in G_2$ such that

$$\mathbf{n} = Be_1, \quad B\Pi_0 = \Pi_+, \quad B\Pi_{\theta} = \Pi_-,$$

where Π_0 and Π_{θ} are special Lagrangian planes in \mathbf{C}^3 as in Example 3.3. By Corollary 3.31 we have

$$V_{\lambda} \cong \begin{cases} 0 & \text{if } \lambda \in [-1, 0) \cup (0, 1) \\ (\mathbf{R} \oplus \mathbf{R}) \oplus (\mathbf{R}^3 \oplus \mathbf{R}^3) & \text{if } \lambda = 0 \\ (\mathbf{R}^3 \oplus \mathbf{R}^5) \oplus (\mathbf{R}^3 \oplus \mathbf{R}^5) & \text{if } \lambda = 1 \end{cases}$$

If H is the symmetry group of $\Pi_0 = \mathbf{R}^3$ for the standard action of G_2 on \mathbf{R}^7 , then $H \cong SO(4)$. Hence C_{\times} is rigid and

$$\text{s-ind}_{\Delta}(C_{\times}) = \text{s-ind}(C_{\times}) = \frac{d-1}{2} + \sum_{-1 < \lambda \leq 1} d_{\lambda} - 2(\dim G_2 - \dim H) - 7 = 0 + 8 + 16 - 16 - 7 = 1. \bullet$$

Example 3.34. (Harvey-Lawson cone) Let C_{HL} be the Harvey-Lawson special Lagrangian cone with link the flat Clifford tori in \mathbf{C}^3 as given in Example 3.4. Then by Corollary 3.31 and [Mar02, Section 6.3.4, p. 132] we have

$$V_{\lambda} \cong \begin{cases} \mathbf{R}^2 & \text{if } \lambda = -1 \\ 0 & \text{if } \lambda \in (-1, 0) \cup (0, 1) \\ \mathbf{R} \oplus \mathbf{R}^6, & \text{if } \lambda = 0 \\ \mathbf{R}^6 \oplus \mathbf{R}^6 & \text{if } \lambda = 1 \end{cases}$$

If H is the symmetry group of C_{HL} for the standard action of G_2 on \mathbf{R}^7 , then $H \cong U(1)^2$. Hence C_{HL} is rigid and

$$\text{s-ind}_{\Delta}(C_{HL}) = \text{s-ind}(C_{HL}) = \frac{d-1}{2} + \sum_{-1 < \lambda \leq 1} d_{\lambda} - (\dim G_2 - \dim H) - 7 = 1 + 19 - 12 - 7 = 1 \bullet$$

Proposition 3.35. *Let C be a connected special Lagrangian cone in \mathbf{C}^3 whose link Σ is not a totally geodesic S^2 . Then*

$$\text{s-ind}_{\Delta}(C) \geq \frac{b^1(\Sigma)}{2} \geq 1$$

with equality if and only if C is C_{HL} , the Harvey-Lawson T^2 -cone (up to special unitary equivalence).

Proof. As Σ is not a totally geodesic S^2 , C is not contained in any hyperplane [Haso4b, Lemma 3.13] in \mathbb{C}^3 . Moreover the genus of Σ is at least 1 (see [Haso4a, Theorem 2.7]). The space of real linear functions on \mathbb{C}^3 induces a 6 dimensional subspace of the 2-eigenspace of Δ_Σ . Therefore, by Corollary 3.31 we have (see Remark 3.32) $d_0 \geq 7$ and

$$\text{s-ind}_\Delta(C) \geq \frac{d_{-1}}{2} = \frac{b^1(\Sigma)}{2} \geq 1.$$

In Example 3.34 we see $\text{s-ind}_\Delta(C_{HL}) = 1$. If $\text{s-ind}_\Delta(C) = 1$, then $b^1(\Sigma) = 1$, $d_0 = 7$ and $d_\lambda = 0$ for all $\lambda \in (-1, 0) \cup (0, 1)$. This implies that the first eigenvalue of Δ_Σ is 2 with multiplicity 6. Then by a theorem of Haskins [Haso4b, Theorem A] C is C_{HL} , the Harvey-Lawson T^2 -cone (up to special unitary equivalence). ■

Proposition 3.36. *Let Σ be a null-torsion holomorphic curve in S^6 and let C be the cone in \mathbb{R}^7 with link Σ . Then*

$$\text{s-ind}_\Delta(C) \geq 5.$$

In particular, this inequality holds for any genus zero holomorphic curve in S^6 except a totally geodesic S^2 .

The proof of this proposition needs the following small preparation.

Definition 3.37. Let Σ be a J -holomorphic curve in S^6 . The **Jacobi operator** $\mathcal{L}_\Sigma : C^\infty(N\Sigma) \rightarrow C^\infty(N\Sigma)$ is

$$\mathcal{L}_\Sigma = (\nabla_\Sigma^\perp)^* \nabla_\Sigma^\perp + \sum_{i=1}^2 (R(f_i, \cdot) f_i)^\perp - \sum_{i,j=1}^2 \langle \Pi(f_i, f_j), \cdot \rangle \Pi(f_i, f_j) \quad \spadesuit$$

Proposition 3.38. *Let Σ be a J -holomorphic curve in S^6 . Then the Jacobi operator \mathcal{L}_Σ satisfies*

$$\mathcal{L}_\Sigma = (\mathbf{D}_\Sigma + J)(\mathbf{D}_\Sigma + 2J) = (J\mathbf{D}_\Sigma)^2 - (J\mathbf{D}_\Sigma) - 2\text{Id}.$$

Moreover, $\text{spec}(\mathcal{L}_\Sigma) = \{\lambda^2 + \lambda - 2 : \lambda \in \mathcal{D}_C\}$, where \mathcal{D}_C is defined in Definition 3.21. Denote by $E_{\mathcal{L}_\Sigma}^\delta$ the δ eigenspace of \mathcal{L}_Σ . Then

$$E_{\mathcal{L}_\Sigma}^\delta \cong V_{\frac{-3+\sqrt{9+4\delta}}{2}} \oplus V_{\frac{-3-\sqrt{9+4\delta}}{2}} \cong V_{\frac{-3+\sqrt{9+4\delta}}{2}} \oplus V_{\frac{-1+\sqrt{9+4\delta}}{2}}$$

In particular, $E_{\mathcal{L}_\Sigma}^{-2} \cong V_{-1} \oplus V_0$ and $E_{\mathcal{L}_\Sigma}^0 \cong V_0 \oplus V_1$.

Proof. We denote the Jacobi operator for the cone C of Σ in \mathbb{R}^7 by \mathcal{L}_C . Gayet [Gay14, Theorem 2.8, Appendix 5.3] proved that

$$\mathcal{L}_C = \mathbf{D}_C^2.$$

Now for all $v_\Sigma \in C^\infty(N_{S^6}\Sigma)$ we have $\mathcal{L}_C v_\Sigma = r^{-2} \mathcal{L}_\Sigma v_\Sigma$. Therefore by Proposition 3.22(ii) we conclude that

$$\mathbf{D}_C^2 = r^{-2} (\mathbf{D}_\Sigma + J)(\mathbf{D}_\Sigma + 2J). \quad \blacksquare$$

Proof of Proposition 3.36. Let \hat{J} be an almost complex structure on $N\Sigma$ defined by the following exact sequence of complex vector bundles (see Example 3.5):

$$0 \rightarrow (L_N, J) \xrightarrow{\alpha} (N\Sigma, \hat{J}) \xrightarrow{\beta} (L_B, -J) \rightarrow 0.$$

Madnick [Mad21] has proved that Σ is null-torsion iff $\nabla^\perp \hat{J} = 0$ and in that case

$$\mathcal{L}_\Sigma = 2\bar{\partial}_{\nabla^\perp, \hat{J}}^* \bar{\partial}_{\nabla^\perp, \hat{J}} - 2$$

where $\bar{\partial}_{\nabla^\perp, \hat{J}}$ is the Cauchy-Riemann operator induced by ∇^\perp and \hat{J} on $N\Sigma$. Moreover,

$$(3.39) \quad \dim E_{\mathcal{L}_\Sigma}^{-2} \geq \text{index } \bar{\partial}_{\nabla^\perp, \hat{J}} = 2c_1(N\Sigma, \hat{J}) + 2\chi(\Sigma) = -4c_1(L_B) = 4b \geq 24.$$

Here $b = \frac{\text{Area}(\Sigma)}{4\pi} \geq 6$. By Proposition 3.38 we see that $\dim E_{\mathcal{L}_\Sigma}^{-2} = d_{-1} + d_0$ and therefore we have

$$\text{s-ind}_\Delta(C) \geq \frac{d_{-1} + d_0}{2} - 7 \geq 2b - 7 \geq 5. \quad \blacksquare$$

Proof of Theorem 1.7. (ii) is proved in Proposition 3.36, (iii) is computed in Example 3.33 and Example 3.34, (iv) is proved in Proposition 3.35. We will prove now (i). Consider the short exact sequence (3.6). If genus of Σ is 1 then $K_\Sigma \cong \mathcal{O}$ and therefore $\mathcal{O}(Z) \otimes L_B = L_N \otimes L_B \cong \mathcal{O}$. Assume Σ is not a null-torsion holomorphic curve, then the zero set of III induces an effective divisor, in particular $\deg L_B \geq [Z]$. Therefore $\deg L_B = [Z] = 0$ and hence $L_B \cong \mathcal{O}$ and $L_N \cong \mathcal{O}$. This implies that there is a non zero section ν_Σ of $N\Sigma$ such that $\bar{\partial}_{\nabla}^N \nu_\Sigma = 0$ and hence $d_{-1} \geq 2$. Moreover, considering the long exact sequence of sheaf cohomologies corresponding to (3.6) we obtain

$$0 \rightarrow \mathbb{C} \rightarrow H^0(\Sigma, N\Sigma) \rightarrow \mathbb{C} \xrightarrow{\text{III}} \mathbb{C} \rightarrow H^1(\Sigma, N\Sigma) \rightarrow \mathbb{C} \rightarrow 0.$$

Since $\text{III} \neq 0$ we obtain more precisely that $d_{-1} = 2 \dim_{\mathbb{C}} H^0(\Sigma, N\Sigma) = 2$. \blacksquare

Remark 3.40. Let Σ be a J -holomorphic curve in S^6 . The **Morse index** of the Jacobi operator \mathcal{L}_Σ is defined by $\text{Ind } \mathcal{L}_\Sigma := \sum_{\delta < 0} \dim E_{\mathcal{L}_\Sigma}^\delta$. Now Proposition 3.38 implies that

$$\text{Ind } \mathcal{L}_\Sigma = d_{-1} + 2 \sum_{-1 < \lambda < 0} d_\lambda + \sum_{0 \leq \lambda < 1} d_\lambda.$$

If genus of Σ is 1 then the proof of Theorem 1.7 implies that $\text{Ind } \mathcal{L}_\Sigma \geq 9$. If Σ is a null-torsion holomorphic curve in S^6 then \mathcal{L}_Σ does not have an eigenvalue less than -2 . Therefore by (3.39) the Morse index of \mathcal{L}_Σ

$$\text{Ind } \mathcal{L}_\Sigma \geq 4b + \sum_{0 < \lambda < 1} d_\lambda \geq 24.$$

Comparing the above observations with a generalized Willmore type conjecture made by Kusner and Wang [KW18, Remark 3.6 (1)] we make the following conjecture that if an associative cone C in \mathbb{R}^7 is not a plane then $\text{s-ind}(C) \geq \text{s-ind}_\Delta(C) \geq 1$ and equality if and only if C is the Harvey-Lawson T^2 -cone or a union of two special Lagrangian planes with transverse intersection at the origin. This has been confirmed in Theorem 1.7 provided C is a special Lagrangian cone in \mathbb{C}^3 . \clubsuit

3.2 Asymptotically conical (AC) associative submanifolds

Definition 3.41. Let C be a cone in \mathbf{R}^7 with link $\Sigma \subset S^6$, that is, $C = \iota((0, \infty) \times \Sigma)$. Let $\Upsilon_\Sigma : V_\Sigma \subset N_{S^6}\Sigma \rightarrow U_\Sigma$ be a **tubular neighbourhood map** for Σ . We define a tubular neighbourhood map

$$\Upsilon_C : V_C \rightarrow U_C$$

for C as follows. Define V_C at (r, σ) by rV_Σ and $\Upsilon_C(u(r, \sigma)) := r\Upsilon_\Sigma(r^{-1}u(r, \sigma))$ and $U_C := \text{im } \Upsilon_C$. \spadesuit

Observe that, for each $\varepsilon > 0$, the scaling map $s_\varepsilon : \mathbf{R}^7 \rightarrow \mathbf{R}^7, x \mapsto \varepsilon x$ induces an action $s_{\varepsilon*}$ on NC . Moreover, Υ_C is equivariant with respect to these actions, that is,

$$\Upsilon_C(s_{\varepsilon*}u) = s_\varepsilon \Upsilon_C u.$$

Definition 3.42. Let C be a cone in \mathbf{R}^7 with link Σ and $\Sigma = \amalg_{i=1}^m \Sigma_i$ be the decomposition into components and C_i be the cone with link Σ_i . An oriented three dimensional submanifold L of \mathbf{R}^7 is called **asymptotically conical (AC) submanifold** with cone C and rate $\nu := (\nu_1, \dots, \nu_m)$ with $\nu_i < 1$ if there exist

- a real number $R_\infty > 1$, a compact submanifold with boundary K_L of L ,
- a diffeomorphism $\Psi_L : (R_\infty, \infty) \times \Sigma \rightarrow L_\infty := L \setminus K_L \subset \mathbf{R}^7$,
- $\Psi_L - \iota$ is a section of the normal bundle NC over $\iota((R_\infty, \infty) \times \Sigma) \subset C$ lying in the tubular neighbourhood V_C of C and

$$|(\nabla_C^\perp)^k(\Psi_L - \iota)| = O(r^{\nu_i - k}),$$

for all $k \in \mathbf{N} \cup \{0\}$ as $r \rightarrow \infty$.

Here ∇_C^\perp is the normal connection on NC induced from Levi-Civita connection on \mathbf{R}^7 and $|\cdot|$ is taken with respect to the normal metric on NC and the cone metric on C . The cone C is called the **asymptotic cone** of L and L_∞ is the **end** of L .

L is called an **asymptotically conical (AC) associative submanifold** with rate ν and cone C if L is associative and an asymptotically conical submanifold with rate ν and cone C in \mathbf{R}^7 as above. \spadesuit

Remark 3.43. If L is an AC associative submanifold, then C is automatically an associative cone in \mathbf{R}^7 (see [Lot11, Proposition 2.15]). An AC submanifold of rate $\nu < 1$ is also an AC Riemannian manifold with rate $\nu - 1$ and also AC submanifold of any rate ν' with $\nu' \leq \nu < 1$. Deformation theory of AC associative submanifolds has been studied by Lotay [Lot11]. \clubsuit

There are many examples of AC special Lagrangians in \mathbf{C}^3 , see [Joy01; Joy02], but we mention only two types of AC special Lagrangians in Example 3.44 and Example 3.47 which are important for this article. For examples of AC associatives which are not AC special Lagrangians, see [Lot11, Section 7; Loto7b].

Example 3.44. (Lawlor neck, [Law89; Joy18, Section 4.1; Har90, pg. 139-143]) Let $\Pi_- := \Pi_0 = \mathbf{R}^3$ and $\Pi_+ := \Pi_\theta$ be a pair of transverse special Lagrangian planes in \mathbf{C}^3 as in Example 3.3. Define

$$\iota^+ : \mathbf{R}^+ \times S^2 \rightarrow \Pi_\theta, \quad \iota^+(r, \sigma) = r \operatorname{diag}(e^{i\theta_1}, e^{i\theta_2}, e^{i\theta_3}) \cdot \sigma$$

and $\iota^- : \mathbf{R}^+ \times S^2 \rightarrow \Pi_0, \quad \iota^-(r, \sigma) = r\sigma$. For any $A > 0$ there exists a unique triple of positive real numbers (a_1, a_2, a_3) such that $A = \frac{4\pi}{3 \cdot \sqrt{a_1 a_2 a_3}}$ and $\theta_k = \theta_k(\infty)$, where

$$\theta_k(y) = a_k \int_{-\infty}^y \frac{dx}{(1 + a_k x^2) \sqrt{P(x)}}, \quad y \in \mathbf{R}, \quad k = 1, 2, 3, \quad P(x) := \frac{\prod_{j=1}^3 (1 + a_j x^2) - 1}{x^2}.$$

The **Lawlor neck** $L_{\theta, A}$ is defined as follows:

$$L_{\theta, A} := \{(z_1(y)\sigma_1, z_2(y)\sigma_2, z_3(y)\sigma_3) \in \mathbf{C}^3 : y, \sigma_1, \sigma_2, \sigma_3 \in \mathbf{R}, \sigma_1^2 + \sigma_2^2 + \sigma_3^2 = 1\}$$

where for $k = 1, 2, 3$, $z_k(y) := e^{i\theta_k(y)} \sqrt{a_k^{-1} + y^2}$. The Lawlor neck $L_{\theta, A}$ is a smooth special Lagrangian submanifold [Har90, Theorem 7.78] diffeomorphic to $\mathbf{R} \times S^2$. The map

$$\begin{aligned} \Phi_{L_{\theta, A}} : \mathbf{R} \times S^2 &\longrightarrow L_{\theta, A} \\ (y, (\sigma_1, \sigma_2, \sigma_3)) &\mapsto (z_1(y)\sigma_1, z_2(y)\sigma_2, z_3(y)\sigma_3) \end{aligned}$$

is a diffeomorphism. Define $\Phi_{L_{\theta, A}}^\pm : \mathbf{R}^+ \times S^2 \longrightarrow L_{\theta, A}$ by $\Phi_{L_{\theta, A}}^\pm(r, \sigma) = \Phi_{L_{\theta, A}}(\pm r, \sigma)$. For every $\varepsilon > 0$, we observe that the rescaled Lawlor neck

$$\varepsilon L_{\theta, A} = L_{\theta, \varepsilon^3 A}.$$

Therefore, Lawlor neck for us is always $L_{\theta, A}$ with the normalization $A = 1$, if not mentioned otherwise.

We define $\xi^- \in C^\infty(N\Pi_0|_{\Pi_0 \setminus \{0\}})$ and $\xi^+ := \operatorname{diag}(e^{i\theta_1}, e^{i\theta_2}, e^{i\theta_3}) \cdot \xi^- \in C^\infty(N\Pi_\theta|_{\Pi_\theta \setminus \{0\}})$ as follows:

$$(3.45) \quad \xi^-(x_1, x_2, x_3) := r^{-3}(ix_1, ix_2, ix_3), \quad \text{where } r = \sqrt{x_1^2 + x_2^2 + x_3^2}.$$

Observe that, $D_{\Pi_\pm} \xi^\pm = 0$ and $\xi^\pm \in V_{-2}$. The Lawlor neck $L_{\theta, A}$ is an AC special Lagrangian submanifold asymptotic to $\Pi_0 \cup \Pi_\theta$ with rate $\nu = -2$. Moreover, the asymptotic normal sections can be written as

$$(3.46) \quad \Psi_{L_{\theta, 1}}^\pm - \iota^\pm = \xi^\pm + O(r^{-4}).$$

The $\Psi_{L_{\theta, 1}}^\pm$ are defined (up to normalizations) as follows:

$$\Psi_{L_{\theta, 1}}^-(\operatorname{Re} \Phi_{L_{\theta, 1}}^-) = \Phi_{L_{\theta, 1}}^- \quad \text{and} \quad \Psi_{L_{\theta, 1}}^+(\operatorname{Re} \operatorname{diag}(e^{-i\theta_1}, e^{-i\theta_2}, e^{-i\theta_3}) \Phi_{L_{\theta, 1}}^+) = \Phi_{L_{\theta, 1}}^+. \quad \bullet$$

Example 3.47. (Harvey-Lawson AC special Lagrangians, [HL82, Theorem 3.1; Joy18, Section 5.1]) Let $C := C_{HL}$ be the Harvey-Lawson T^2 -cone as in Example 3.4 and let ι be the inclusion

map in \mathbf{R}^7 . For any positive real number $a > 0$, the **Harvey-Lawson special Lagrangian 3-folds** are defined by

$$\begin{aligned} L_a^1 &:= \{(z_1, z_2, z_3) \in \mathbf{C}^3 : |z_1|^2 - a = |z_2|^2 = |z_3|^2, z_1 z_2 z_3 \in [0, \infty)\}, \\ L_a^2 &:= \{(z_1, z_2, z_3) \in \mathbf{C}^3 : |z_1|^2 = |z_2|^2 - a = |z_3|^2, z_1 z_2 z_3 \in [0, \infty)\}, \\ L_a^3 &:= \{(z_1, z_2, z_3) \in \mathbf{C}^3 : |z_1|^2 = |z_2|^2 = |z_3|^2 - a, z_1 z_2 z_3 \in [0, \infty)\}. \end{aligned}$$

Each $L_a^k, k = 1, 2, 3$ is a smooth special Lagrangian in \mathbf{C}^3 [HL82, Section III.3.A] diffeomorphic to $S^1 \times \mathbf{C}$. Define the diffeomorphisms

$$\begin{aligned} \Phi_{L_a^1} : \mathbf{R}^+ \times T^2 &\longrightarrow L_a^1, & (r, e^{i\theta_1}, e^{i\theta_2}) &\mapsto (e^{i\theta_1} \sqrt{r^2 + a}, r e^{i\theta_2}, r e^{-i(\theta_1 + \theta_2)}), \\ \Phi_{L_a^2} : \mathbf{R}^+ \times T^2 &\longrightarrow L_a^2, & (r, e^{i\theta_1}, e^{i\theta_2}) &\mapsto (r e^{-i(\theta_1 + \theta_2)}, e^{i\theta_1} \sqrt{r^2 + a}, r e^{i\theta_2}), \\ \Phi_{L_a^3} : \mathbf{R}^+ \times T^2 &\longrightarrow L_a^3, & (r, e^{i\theta_1}, e^{i\theta_2}) &\mapsto (r e^{i\theta_2}, r e^{-i(\theta_1 + \theta_2)}, e^{i\theta_1} \sqrt{r^2 + a}). \end{aligned}$$

For every $\varepsilon > 0$, we have¹¹

$$\varepsilon L_a^k = L_{\varepsilon^2 a}^k.$$

Each L_a^k is an AC special Lagrangian submanifold asymptotic to C_{HL} with rate $\nu = -1$. Moreover, the asymptotic normal sections can be written as

$$(3.48) \quad \Psi_{L^k} - \iota = \xi_k + O(r^{-2}),$$

where $\xi_k \in V_{-1}$ and $\xi_3 = -\xi_1 - \xi_2$. •

Definition 3.49. Let $\rho : (0, \infty) \rightarrow [0, 1]$ be a smooth function such that

$$(3.50) \quad \rho(r) = \begin{cases} 1, & r \leq 1 \\ 0, & r \geq 2. \end{cases}$$

For every $s > 0$, $\rho_s : (0, \infty) \rightarrow [0, 1]$ is defined by $\rho_s(r) := \rho(s^{-1}r)$. Let L be an AC submanifold with cone C as in Definition 3.42. We define another AC submanifold L_C which is diffeomorphic to L

$$L_C := K_L \cup \Upsilon_C(\rho_{R_\infty}(\Psi_L - \iota)).$$

We define $C_\infty := \iota((2R_\infty, \infty) \times \Sigma) \subset L_C$ and $K_{L_C} := L_C \setminus C_\infty$. ♠

Definition 3.51. Let L be an AC submanifold with cone C as in Definition 3.42 and Υ_C be as in Definition 3.41. We choose a tubular neighbourhood map

$$\Upsilon_{L_C} : V_{L_C} \rightarrow U_{L_C}$$

of L_C such that V_{L_C} and Υ_{L_C} agree with V_C and Υ_C respectively on C_∞ .

There is a section β in $V_{L_C} \subset NL_C$ which is zero on K_L and $\Upsilon_{L_C}(\Gamma\beta)$ is L , and

$$|(\nabla_{C_i}^\perp)^k \beta| = O(r^{\nu_i - k})$$

for all $k \in \mathbf{N} \cup \{0\}$ as $r \rightarrow \infty$, where ∇_C^\perp is the normal connection on NL_C induced from Levi-Civita connection on \mathbf{R}^7 . ♠

¹¹Therefore, we always assume L_a^k with the normalization $a = 1$, if not mentioned otherwise.




Figure 4: Construction of AC submanifold L_C from an AC submanifold L .

3.3 Conically singular (CS) associative submanifolds

In this section we study conically singular (CS) associative submanifolds with singularity at one point and modeled on an associative cone in \mathbf{R}^7 . Similar work for special Lagrangians and for coassociatives have been done by Joyce [Joy04b] and Lotay [Loto7a].

Definition 3.52. Let (Y, ϕ) be an almost G_2 -manifold (see Definition 2.2) and let p be a point in Y . A G_2 -framing at p is a linear isomorphism $v : \mathbf{R}^7 \rightarrow T_p Y$ such that $v^*(\phi(p)) = \phi_e$ where ϕ_e is the standard G_2 structure on \mathbf{R}^7 . A G_2 -coordinate system at p with G_2 -framing v is a diffeomorphism

$$\Upsilon : B_R(0) \subset \mathbf{R}^7 \rightarrow U$$

for some constant $0 < R < 1$ and some open set U containing p in Y satisfying $\Upsilon(0) = p$ and $d\Upsilon_0 = v$.

Two G_2 -coordinate systems Υ_1 and Υ_2 at p are called **equivalent** if they have the same G_2 -framing at p . \spadesuit

Definition 3.53. Let P be a compact subset of an almost G_2 -manifold Y and $\text{sing}(P) := \{p_1, \dots, p_m\}$ be a finite set points in P such that $\hat{P} := P \setminus \text{sing}(P)$ is a smooth three dimensional oriented submanifold of Y . We call \hat{P} the **smooth part** of P and $\text{sing}(P)$ the **singular set of points** of P . Let $\{C_1, \dots, C_m\}$ be a set of cones in \mathbf{R}^7 . Denote the link of C_i by Σ_i and inclusion map into \mathbf{R}^7 by ι_i . Let

$$\Upsilon^i : B_R(0) \rightarrow U \subset Y$$

be G_2 -coordinate systems with G_2 -framings v_i at p_i , $i = 1, \dots, m$.

(1) P is said to be a **conically singular (CS) submanifold** with singularities at p_i modeled on cones C_i and rates μ_i with $1 < \mu_i \leq 2$ if there exist

- $\varepsilon_0 > 0$ with $2\varepsilon_0 < R$ and a compact submanifold with boundary K_P of P ,
- smooth embeddings

$$\Psi_P^i : (0, 2\varepsilon_0) \times \Sigma_i \rightarrow B_R(0), \quad i = 1, 2, \dots, m$$

such that $\bigcup_{i=1}^m \Upsilon^i \circ \Psi_P^i : (0, 2\varepsilon_0) \times \Sigma_i \rightarrow Y$ is a diffeomorphism onto $\hat{P} \setminus K_P$,

- $\Psi_P^i - \iota_i$ is a smooth section of the normal bundle NC_i over $\iota((0, 2\varepsilon_0) \times \Sigma_i)$ which lies in V_{C_i} and

$$|(\nabla_{C_i}^\perp)^k(\Psi_P^i - \iota_i)| = O(r^{\mu_i - k})$$

for all $k \in \mathbf{N} \cup \{0\}$ as $r \rightarrow 0$, $i = 1, 2, \dots, m$.

Here $\nabla_{C_i}^\perp$ is the normal connection on NC_i induced from Levi-Civita connection on \mathbf{R}^7 and $|\cdot|$ is taken with respect to the normal metric on NC_i and cone metric on C_i .

- (2) If all the above sections $\Psi_P^i - \iota_i$ are zero sections then P is said to be **end conical singular (ECS) submanifold**.
- (3) The cone $\tilde{C}_i := v_i(C_i) \subset T_{p_i}Y$ is said to be the **tangent cone** of P at p_i .
- (4) P is said to be a **conically singular (CS) associative submanifold** if it is conically singular submanifold as above and \hat{P} is an associative submanifold of (Y, ϕ) . \clubsuit

Remark 3.54. We make some remarks about the above definition.

- (i) A CS submanifold with rates μ is also a CS submanifold with all rates μ' such that $1 < \mu'_i \leq \mu_i \leq 2$ for all $i = 1, 2, \dots, m$.
- (ii) A CS submanifold with rates μ is a conically singular Riemannian manifold with rates $\mu - 1$ for the metric g_ϕ induced by ϕ . Indeed

$$|(\nabla_{C_i}^\perp)^k((Y^i \circ \Psi_P^i)^* g_\phi - g_{C_i})| \lesssim \sum_{j=0}^{k+1} |(\nabla_{C_i}^\perp)^j(\Psi_P^i - \iota_i)|.$$

- (iii) $\mu_i > 1$ also implies that if P is a CS associative submanifold as above, then each cone C_i is also an associative cone in \mathbf{R}^7 . Indeed, the associator on C_i

$$|[\cdot, \cdot, \cdot]_{C_i}| = O(r^{\mu_i - 1}), \text{ as } r \rightarrow 0.$$

But $[\cdot, \cdot, \cdot]_{C_i}$ is dilation invariant and therefore $[\cdot, \cdot, \cdot]_{C_i} = 0$.

- (iv) $\mu_i \leq 2$ guarantees that the above definition of CS submanifold is independent of the choice of a G_2 -coordinate system in an equivalence class. Moreover it is independent of the choice of a G_2 -framing representing the tangent cone. \clubsuit

3.4 Linear analysis on CS and AC associative submanifolds

By considering appropriate weighted function spaces of sections of vector bundles over CS and AC submanifolds (where the weights represent decay rates of those sections) we obtain a Fredholm theory for CS and AC elliptic operators if the weights avoid the wall of critical weights or rates. The index of these operators changes when weights lie on various connected components of the complement of the wall of critical rates. This theory originally appeared in Lockhart-McOwen [LM85]. A very good exposition can be found in [Maro2], [KL20].

Notation 3.55. For a conically singular manifold P we will denote by NP the normal bundle of $\hat{P} = P \setminus \text{sing}(P)$. \blacktriangleright

3.4.1 Weighted function spaces

Let P be a CS submanifold as in Definition 3.53 and L be an AC submanifold as in Definition 3.42 asymptotic to $C = \coprod_{i=1}^m C_i$.

Notation 3.56. We denote

$$M := \begin{cases} P & \text{if } P \text{ is a CS submanifold} \\ L & \text{if } L \text{ is a AC submanifold.} \end{cases}$$

and

$$\eta := \begin{cases} \mu & \text{if } P \text{ is a CS submanifold with rate } \mu \\ \nu & \text{if } L \text{ is a AC submanifold with rate } \nu. \end{cases} \quad \blacktriangleright$$

Notation 3.57. We would like to define weighted Sobolev and Hölder spaces with rate $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_m) \in \mathbf{R}^m$. We say $\lambda <, =, > \lambda'$ if for each $i = 1, 2, \dots, m$ we have $\lambda_i <, =, > \lambda'_i$ respectively. For any $s \in \mathbf{R}$, $\lambda + s := (\lambda_1 + s, \lambda_2 + s, \dots, \lambda_m + s)$ and set $|\lambda| := \sum_{i=1}^m |\lambda_i|$. \blacktriangleright

Definition 3.58. For each $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_m) \in \mathbf{R}^m$, a **weight function** on \hat{P} , $w_{P,\lambda} : \hat{P} \rightarrow (0, \infty)$, is a smooth function on \hat{P} such that if $x = \Upsilon \circ \Psi_P(r, \sigma)$ with $(r, \sigma) \in (0, 2\varepsilon_0) \times \Sigma_i$ then

$$w_{P,\lambda}(x) = r^{-\lambda_i}.$$

The weight function on L , $w_{L,\lambda} : L \rightarrow (0, \infty)$ is defined by

$$w_{L,\lambda}(x) := (1 + |x|)^{-\lambda_i}.$$

Let $\lambda \in \mathbf{R}^m$ and $k \geq 0$, $p \geq 1$, $\gamma \in (0, 1)$. For a continuous section u of NM we define the **weighted L^∞ norm** and the **weighted Hölder semi-norm** respectively by

$$\|u\|_{L_{M,\lambda}^\infty} := \|w_{M,\lambda}u\|_{L^\infty(NM)}, \quad [u]_{C_{M,\lambda}^{0,\gamma}} := [w_{M,\lambda-\gamma}u]_{C^{0,\gamma}(NM)}.$$

For a continuous section u of NM with k continuous derivatives we define the **weighted C^k norm** and the **weighted Hölder norm** respectively by

$$\|u\|_{C_{M,\lambda}^k} := \sum_{j=0}^k \|(\nabla_M^\perp)^j u\|_{L_{M,\lambda-j}^\infty}, \quad \|u\|_{C_{M,\lambda}^{k,\gamma}} := \|u\|_{C_{M,\lambda}^k} + [(\nabla_M^\perp)^k u]_{C_{M,\lambda-k}^{0,\gamma}}$$

and, the **weighted Sobolev norm** by

$$\|u\|_{W_{M,\lambda}^{k,p}} := \left(\sum_{j=0}^k \int_M |w_{M,\lambda-j}(\nabla_M^\perp)^j u|^p w_{M,3} dV_M \right)^{\frac{1}{p}}.$$

Here, the connection ∇_M^\perp on NM is the projection of the Levi-Civita connection induced by g_ϕ for the decomposition $TY|_M = TM \oplus NM$ and the $|\cdot|$ is with respect to the metric g_ϕ . We denote always $W_{M,\lambda}^{0,p}$ by $L_{M,\lambda}^p$.

We define the **weighted Hölder space** $C_{M,\lambda}^{k,\gamma}$ to be the space of continuous sections of NM with k continuous derivatives and finite weighted Hölder norm $\|\cdot\|_{C_{M,\lambda}^{k,\gamma}}$. This is a Banach space but not separable.

We define the **weighted Sobolev space** $W_{M,\lambda}^{k,p}$, the **weighted C^k -space** $C_{M,\lambda}^k$, the **weighted L^p -space** $L_{M,\lambda}^p$, the **weighted L^∞ -space** $L_{M,\lambda}^\infty$ to be the completion of the space of compactly supported smooth sections of NM , namely $C_c^\infty(NM)$ with respect to the weighted Hölder norm $\|\cdot\|_{C_{M,\lambda}^{k,\gamma}}$, the weighted Sobolev norm $\|\cdot\|_{W_{M,\lambda}^{k,p}}$, the weighted C^k -norm $\|\cdot\|_{C_{M,\lambda}^k}$, the weighted L^p -norm $\|\cdot\|_{L_{M,\lambda}^p}$, the weighted L^∞ -norm $\|\cdot\|_{L_{M,\lambda}^\infty}$ respectively. These are all separable Banach spaces. Moreover, we define the **weighted C^∞ -space** $C_{M,\lambda}^\infty$ by

$$C_{M,\lambda}^\infty := \bigcap_{k=0}^{\infty} C_{M,\lambda}^k \quad \spadesuit$$

Remark 3.59. The spaces $W_{M,\lambda}^{k,2}$ are all Hilbert spaces with inner product coming from the polarization of the norm. We also note that

$$L^p(NM) = L_{M,-\frac{3}{p}}^p \quad \clubsuit$$

Proposition 3.60. *Let $k, l \geq 0, p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$ and $\lambda, \lambda_1, \lambda_2 \in \mathbf{R}^m$. Then the L^2 -inner product map*

$$W_{M,\lambda_1}^{k,p} \times W_{M,\lambda_2}^{l,q} \xrightarrow{\langle \cdot, \cdot \rangle_{L^2}} \mathbf{R}$$

is continuous provided $M = P$ (CS) with $\lambda_1 + \lambda_2 \geq -3$, or $M = L$ (AC) with $\lambda_1 + \lambda_2 \leq -3$. Moreover, this L^2 -inner product yields a Banach space isomorphism

$$(L_{M,\lambda}^p)^* \cong L_{M,-3-\lambda}^q$$

Proof. For $u \in L_{M,\lambda_1}^p$ and $v \in L_{M,\lambda_2}^q$ we consider

$$\begin{aligned} \langle u, v \rangle_{L^2} &= \int_M (w_{M,\lambda_1} u w_{M,\frac{3}{p}}) (w_{M,\lambda_2} v w_{M,\frac{3}{q}}) w_{M,-\lambda_1-\lambda_2-3} \text{Vol}_M \\ &\lesssim \|u\|_{W_{M,\lambda_1}^{k,p}} \|v\|_{W_{M,\lambda_2}^{l,q}}, \end{aligned}$$

provided $M = P$ (CS) with $\lambda_1 + \lambda_2 \geq -3$, or $M = L$ (AC) with $\lambda_1 + \lambda_2 \leq -3$. This proves that the L^2 -inner product map is continuous. The isomorphism stated in the proposition follows from the standard fact that $\langle \cdot, \cdot \rangle_{L^2} : L^p \times L^q \rightarrow \mathbf{R}$ is a dual pairing. \blacksquare

The following is a weighted embedding and compactness theorem [Maroz, Theorem 4.17, Theorem 4.18].

Proposition 3.61. *Let $\lambda, \lambda_1, \lambda_2 \in \mathbf{R}^m, k, l \in \mathbf{N} \cup \{0\}, p, q \geq 1, \alpha, \beta \in (0, 1)$. Then*

- i) *If $k \geq l$ and $k - \frac{n}{p} \geq l - \frac{n}{q}$, the inclusion $W_{M,\lambda_1}^{k,p} \hookrightarrow W_{M,\lambda_2}^{l,q}$ is a continuous embedding, provided for $M = L$ (AC) either $p \leq q, \lambda_1 \leq \lambda_2$ or $p > q, \lambda_1 < \lambda_2$, and for $M = P$ (CS) either $p \leq q, \lambda_1 \geq \lambda_2$ or $p > q, \lambda_1 > \lambda_2$.*

- ii) If $k + \alpha \geq l + \beta$, the inclusion $C_{M,\lambda_1}^{k,\alpha} \hookrightarrow C_{M,\lambda_2}^{l,\beta}$ is a continuous embedding, provided $\lambda_1 \leq \lambda_2$ for $M = L$ (AC) and $\lambda_1 \geq \lambda_2$ for $M = P$ (CS).
- iii) If $k - \frac{n}{p} \geq l + \alpha$, the inclusion $W_{M,\lambda}^{k,p} \hookrightarrow C_{M,\lambda}^{l,\alpha}$ is a continuous embedding.
- iv) All of the above embeddings are compact provided all the inequalities in the corresponding hypotheses are strict inequalities.

3.4.2 AC and CS elliptic operator on AC and CS associatives

Let P be a CS associative submanifold as in Definition 3.53 and L be an AC associative submanifold as in Definition 3.42 asymptotic to $C = \coprod_{i=1}^m C_i$. We continue denoting M as either P or L and η as μ for CS and ν for AC submanifolds. For any associative cone C in \mathbf{R}^7 with link Σ , the Fueter operator \mathbf{D}_C is a conical operator (see [Maro2, Section 4.3.2] for precise definition), that is after the identification of the cone \mathbf{R}^7 with the cylinder $\mathbf{R} \times S^6$ by substituting $r = e^t$ we obtain that (see Proposition 3.22)

$$(3.62) \quad r^2 \mathbf{D}_C r^{-1} = Jr \partial_r + (\mathbf{D}_\Sigma + J) = J \partial_t + (\mathbf{D}_\Sigma + J)$$

is a translation invariant operator.

Definition 3.63. Let $\{C_1, \dots, C_m\}$ be a set of associative cones. Denote $C := \coprod_{i=1}^m C_i$. Then the set of critical rates \mathcal{D}_C is defined by

$$\mathcal{D}_C := \{(\lambda_1, \lambda_2, \dots, \lambda_m) \in \mathbf{R}^m : \lambda_i \in \mathcal{D}_{C_i} \text{ for some } i\},$$

where \mathcal{D}_{C_i} is the set of all indicial roots of \mathbf{D}_{C_i} defined in Definition 3.21. We call \mathcal{D}_C the **wall of critical rates** in \mathbf{R}^m . We define

$$V_\lambda := \bigoplus_{i=1}^m V_{\lambda_i}, \quad d_{\lambda_i} := \dim V_{\lambda_i} \text{ and } d_\lambda := \sum_{i=1}^m d_{\lambda_i}.$$

where V_{λ_i} are defined in Definition 3.21. ♠

Definition 3.64. Let $\mathbf{D}_M : C_c^\infty(NM) \rightarrow C_c^\infty(NM)$ be a first order, formally self-adjoint elliptic operator. It is called **asymptotically conical (AC) uniformly elliptic** operator for $M = L$ (AC) and **conically singular (CS) uniformly elliptic** operator for $M = P$ (CS) respectively, asymptotic to the conical operators \mathbf{D}_{C_i} over the ends of M if the operator $r^2 \mathbf{D}_M r^{-1}$ is an asymptotically translation invariant uniformly elliptic operator asymptotic to $J \partial_t + (\mathbf{D}_{\Sigma_i} + J)$ (see [Maro2, Section 4.3.2]) and after the identifications by canonical bundle isomorphisms

$$\mathbf{D}_M = \mathbf{D}_{C_i} + O(r^{\eta_i - 1}).$$

The operator \mathbf{D}_M has canonical extensions to weighted function spaces and we denote them as follows:

$$\mathbf{D}_{M,\lambda}^{k,p} : W_{M,\lambda}^{k+1,p} \rightarrow W_{M,\lambda-1}^{k,p}, \quad \mathbf{D}_{M,\lambda}^{k,\gamma} : C_{M,\lambda}^{k+1,\gamma} \rightarrow C_{M,\lambda-1}^{k,\gamma} \quad \spadesuit$$

The following proposition is about elliptic regularity statements for CS or AC uniformly elliptic operator \mathbf{D}_M [Maro2, Theorem 4.6].

Proposition 3.65. *Suppose $\lambda \in \mathbf{R}^m$, $k \geq 0$, $p > 1$, $\gamma \in (0, 1)$. Let, $u, v \in L^1_{\text{loc}}$ be two locally integrable sections of NM such that u is a weak solution of the equation $\mathbf{D}_M u = v$.*

i) *If $v \in C^{k, \gamma}_{M, \lambda-1}$, then $u \in C^{k+1, \gamma}_{M, \lambda}$ is a strong solution and there exists a constant $C > 0$ such that*

$$\|u\|_{C^{k+1, \gamma}_{M, \lambda}} \leq C \left(\|\mathbf{D}_{M, \lambda}^{k, \gamma} u\|_{C^{k, \gamma}_{M, \lambda-1}} + \|u\|_{L^\infty_{M, \lambda}} \right).$$

ii) *If $v \in W^{k, p}_{M, \lambda-1}$, then $u \in W^{k+1, p}_{M, \lambda}$ is a strong solution and there exists a constant $C > 0$ such that*

$$\|u\|_{W^{k+1, p}_{M, \lambda}} \leq C \left(\|\mathbf{D}_{M, \lambda}^{k, p} u\|_{W^{k, p}_{M, \lambda-1}} + \|u\|_{L^\infty_{M, \lambda}} \right).$$

Proposition 3.66. *Suppose $k \geq 0$, $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$ and $\lambda_1, \lambda_2 \in \mathbf{R}^m$. For all $u \in W^{k+1, p}_{M, \lambda_1}$, $v \in W^{k+1, q}_{M, \lambda_2}$ we have $\langle \mathbf{D}_{M, \lambda_1}^{k, p} u, v \rangle_{L^2} = \langle u, \mathbf{D}_{M, \lambda_2}^{k, q} v \rangle_{L^2}$, provided $M = P$ (CS) with $\lambda_1 + \lambda_2 \geq -2$, or $M = L$ (AC) with $\lambda_1 + \lambda_2 \leq -2$.*

Proof. The result is true for $u, v \in C_c^\infty(NM)$ and therefore the general statement follows from Proposition 3.60. \blacksquare

Definition 3.67. Let C be a connected associative cone. We can define the Banach spaces $W^{k, p}_{C, \lambda}$, $C^{k, \gamma}_{C, \lambda}$ and all others over C similar to Section 3.4.1 replacing M by C , NM by NC and weight function w_M by $w_{C, \lambda} : C \rightarrow \mathbf{R}$ where $w_{C, \lambda}(r, \sigma) = r^{-\lambda}$, $\lambda \in \mathbf{R}$. \spadesuit

The following lemma follows from [Dono2, Lemma 3.1, Proposition 3.4, Section 3.3.1] and [MP78, Theorem 5.1] using the observation above in (3.62).

Lemma 3.68. *Let C be a connected associative cone. The conical Fueter operator $\mathbf{D}_C : C^{k+1, \gamma}_{C, \lambda} \rightarrow C^{k, \gamma}_{C, \lambda-1}$ is invertible if and only if $\lambda \in \mathbf{R} \setminus \mathcal{D}_C$. Moreover any element $u \in \ker \mathbf{D}_C$ has a L^2 -decomposition*

$$u = \sum_{\lambda \in \mathcal{D}_C} r^{\lambda-1} u_{\Sigma, \lambda}.$$

where $u_{\Sigma, \lambda}$ are λ -eigensections of $J\mathbf{D}_\Sigma - \mathbf{1}$, that is $r^{\lambda-1} u_{\Sigma, \lambda} \in V_\lambda$.

Proposition 3.69. *Let $\lambda \in \mathbf{R}^m$, $k \geq 0$, $p > 1$, $\gamma \in (0, 1)$. Then $\mathbf{D}_{M, \lambda}^{k, p}$ and $\mathbf{D}_{M, \lambda}^{k, \gamma}$ are Fredholm for all $\lambda \in \mathbf{R}^m \setminus \mathcal{D}_C$. Moreover, for all $\lambda \in \mathbf{R}^m$, $\text{Ker } \mathbf{D}_{M, \lambda}^{k, p} = \text{Ker } \mathbf{D}_{M, \lambda}^{k, \gamma}$ is finite dimensional, independent of k, p and γ .*

Proof. The operators $\mathbf{D}_{M, \lambda}^{k, p}$ and $\mathbf{D}_{M, \lambda}^{k, \gamma}$ are Fredholm with weight $\lambda \in \mathbf{R}^m \setminus \mathcal{D}_C$ follows from [Dono2, Proposition 3.6, Section 3.3.1] or [LM85, Theorem 6.2]. Independence of k, p, γ follows from Proposition 3.65 and Proposition 3.61. \blacksquare

Lemma 3.70. Let $\lambda \in \mathbf{R}^m$ and λ_1, λ_2 be two elements in $\mathbf{R}^m \setminus \mathcal{D}_C$ with $\lambda_1 < \lambda < \lambda_2$ for $M = L$ (AC) and $\lambda_2 < \lambda < \lambda_1$ for $M = P$ (CS), $|\lambda_2 - \lambda_1| \leq |\eta - 1|$ and there are no other indicial roots in between λ_1, λ_2 except possibly λ . We define the set

$$\mathcal{S}_{\lambda_2} := \{u \in C_{M, \lambda_2}^{k+1, Y} : \mathbf{D}_{M, \lambda_2}^{k, Y} u \in C_{M, \lambda_1 - 1}^{k, Y}\}.$$

Define a linear map $e_{M, \lambda} : V_\lambda \rightarrow W_{M, \lambda_2}^{k+1, P}$ (under the identifications of normal bundles NM and NC_i by the canonical bundle isomorphisms over the ends) by

$$e_{M, \lambda}(w) := \begin{cases} \bigoplus_{i=1}^m \rho_{\varepsilon_0} w_i & \text{if } M = P \text{ (CS)} \\ \bigoplus_{i=1}^m (1 - \rho_{R_\infty}) w_i & \text{if } M = L \text{ (AC)} \end{cases}$$

Then there exists a unique linear map (**asymptotic map**), $\tilde{i}_{M, \lambda} : \mathcal{S}_{\lambda_2} \rightarrow V_\lambda$ such that for any $u \in \mathcal{S}_{\lambda_2}$ we have

$$u - e_{M, \lambda} \circ \tilde{i}_{M, \lambda} u \in C_{M, \lambda_1}^{k, Y}.$$

Moreover, the following statements hold.

(i) $\mathcal{S}_{\lambda_2} \subset C_{M, \lambda}^{k+1, Y}$ and $\text{Ker } \mathbf{D}_{M, \lambda_2}^{k, Y} = \text{Ker } \mathbf{D}_{M, \lambda}^{k, Y}$, and $\text{Ker } \tilde{i}_{M, \lambda} = C_{M, \lambda_1}^{k+1, Y}$. Moreover,

$$\text{Ker}\{i_{M, \lambda} := \tilde{i}_{M, \lambda} : \text{Ker } \mathbf{D}_{M, \lambda_2}^{k, Y} \rightarrow V_\lambda\} = \text{Ker } \mathbf{D}_{M, \lambda_1}^{k, Y}$$

(ii) $\mathcal{S}_{\lambda_2} = C_{M, \lambda_1}^{k+1, Y} + \text{im } e_{M, \lambda}$ and the restriction of $\mathbf{D}_{M, \lambda_2}^{k, Y}$, denoted by

$$\widehat{\mathbf{D}}_{M, \lambda_1}^{k, Y} : C_{M, \lambda_1}^{k+1, Y} + \text{im } e_{M, \lambda} \rightarrow C_{M, \lambda_1 - 1}^{k, Y}$$

has the property that $\text{Ker } \mathbf{D}_{M, \lambda_2}^{k, Y} = \text{Ker } \widehat{\mathbf{D}}_{M, \lambda_1}^{k, Y}$ and $\text{Coker } \mathbf{D}_{M, \lambda_2}^{k, Y} \cong \text{Coker } \widehat{\mathbf{D}}_{M, \lambda_1}^{k, Y}$.

(iii) (**Wall crossing formula**) $\text{index } \mathbf{D}_{M, \lambda_2} = \text{index } \mathbf{D}_{M, \lambda_1} + d_\lambda$.

Proof. The proof of this lemma is similar to [Ber22, Lemma 3.38]. It has also been proved in [KL20, Proposition 4.21 - Corollary 4.24]. We will only show the existence of the asymptotic map $\tilde{i}_{M, \lambda}$. Given $u \in \mathcal{S}_{\lambda_2}$ we define $\tilde{u} \in \bigoplus_{i=1}^m C^\infty(NC_i)$ by

$$\tilde{u} := \begin{cases} \bigoplus_{i=1}^m \tilde{u}_i := \rho_{\varepsilon_0} u & \text{if } M = P \text{ (CS)} \\ \bigoplus_{i=1}^m (1 - \rho_{R_\infty}) u & \text{if } M = L \text{ (AC)}. \end{cases}$$

Denote any of the asymptotic cone by C . Since $\mathbf{D}_M = \mathbf{D}_C + O(r^{\eta-1})$ and $|\lambda_2 - \lambda_1| \leq |\eta - 1|$, therefore $\mathbf{D}_C \tilde{u} \in C_{C, \lambda_1 - 1}^\infty$. Lemma 3.68 implies that there exists a unique $v \in C_{C, \lambda_1}^\infty$ such that $\mathbf{D}_C(\tilde{u} - v) = 0$. We define the asymptotic map $\tilde{i}_{M, \lambda}$ as follows:

$$(3.71) \quad \tilde{i}_{M, \lambda}(u) := r^{\lambda-1}(\tilde{u} - v)_{\Sigma, \lambda}. \quad \blacksquare$$

Proposition 3.72. $\text{Ker } \mathbf{D}_{M, \lambda}^{k, Y}$ is independent of λ in each connected component of $\lambda \in \mathbf{R}^m \setminus \mathcal{D}_C$. Moreover, for all $\lambda \in \mathbf{R}^m \setminus \mathcal{D}_C$, we have

(i) $\text{Coker } \mathbf{D}_{M,\lambda} \cong \text{Ker } \mathbf{D}_{M,-2-\lambda}$. If $\lambda \geq -\frac{1}{2}$ for $M = L$ (AC) or $\lambda \leq -\frac{1}{2}$ for $M = P$ (CS), then $\text{Ker } \mathbf{D}_{M,-2-\lambda}$ is equal to $\text{Coker } \mathbf{D}_{M,\lambda}$.

(ii) If $s > 0$ for $M = L$ (AC) or $s < 0$ for $M = P$ (CS) such that -1 is the possibly only critical rates in between $-1 - s$ and $-1 + s$, then

$$\text{index } \mathbf{D}_{M,-1+s} = \dim \text{Ker } \mathbf{D}_{M,-1+s} - \dim \text{Ker } \mathbf{D}_{M,-1-s} = \frac{d_{-1}}{2}$$

(iii) $\text{index } \mathbf{D}_{L,\lambda} = -\text{index } \mathbf{D}_{P,\lambda}$ and

$$\text{index } \mathbf{D}_{L,\lambda} = \sum_{\lambda_i \geq -1} \left(\frac{d_{-1,i}}{2} + \sum_{\zeta_i \in \mathcal{D}_{C_i} \cap (-1, \lambda_i)} d_{\zeta_i} \right) - \sum_{\lambda_i < -1} \left(\frac{d_{-1,i}}{2} + \sum_{\zeta_i \in \mathcal{D}_{C_i} \cap (\lambda_i, -1)} d_{\zeta_i} \right).$$

Proof. The first statement is a direct consequence of part (iii) of previous lemma. Now first part of (i) follows from Proposition 3.60 and Proposition 3.69, and if λ as in second part of (i) we use the fact that $C_{M,-2-\lambda}^{k,Y} \subset C_{M,\lambda}^{k,Y}$. To see (ii), observe from (i) and the wall crossing formula of Lemma 3.70 that

$$\dim \text{Ker } \mathbf{D}_{M,-1+s} - \dim \text{Ker } \mathbf{D}_{M,-1-s} = \dim \text{Ker } \mathbf{D}_{M,-1-s} - \dim \text{Ker } \mathbf{D}_{M,-1+s} + d_{-1}.$$

Finally, (iii) is an immediate consequences of (ii) and the wall crossing formula stated in the Lemma 3.70. \blacksquare

Proposition 3.73 (Marshall [Mar02, section 5, Section 5.2.3, Table 5.1]). *Let L be a AC special Lagrangian submanifold in \mathbb{C}^3 . Let Σ be the link of the asymptotic cone. Then*

$$\dim \ker \mathbf{D}_{L,-1} = b^1(L) + b^0(\Sigma) - 1.$$

3.5 Moduli space of conically singular (CS) associative submanifolds

Definition 3.74. For a G_2 -structure ϕ on Y , $\mathcal{F}_\phi := \{(p, v) : v \text{ is a } G_2\text{-framing at } p \text{ in } Y\}$ is a principal G_2 -bundle over Y . Let \mathcal{P} be the set of all co-closed G_2 structures on Y . Then

$$\mathcal{F} := \bigcup_{\phi \in \mathcal{P}} \mathcal{F}_\phi \xrightarrow{\Pi} \mathcal{P} \times Y$$

is a principal G_2 -bundle over $\mathcal{P} \times Y$. \spadesuit

Definition 3.75. We define the universal space of conically singular (CS) submanifolds,

$$\mathcal{S}_{\text{cs}} := \{(\phi, P) : P \text{ is a CS submanifold in } (Y, \phi), \phi \in \mathcal{P}\}.$$

and the universal **moduli space** \mathcal{M}_{cs} of CS associative submanifolds,

$$\mathcal{M}_{\text{cs}} := \{(\phi, P) \in \mathcal{S}_{\text{cs}} : P \text{ is a CS associative submanifold}\}.$$

There is a canonical projection map $\pi : \mathcal{S}_{\text{cs}} \rightarrow \mathcal{P}$ and we define the **moduli space** of CS associative submanifolds in (Y, ϕ) by

$$\mathcal{M}_{\text{cs}}^\phi := \pi^{-1}(\phi) \cap \mathcal{M}_{\text{cs}}. \quad \spadesuit$$

We will now define a topology on \mathcal{M}_{cs} , called weighted C^∞ -topology. The topology will be defined by specifying a basis. The open sets in this basis will be constructed using end conical (EC) tubular neighbourhood maps of only end conical singular (ECS) submanifolds (see Definition 3.76). Indeed given an CS submanifold we will chose an ECS submanifold with an EC tubular neighbourhood containing the CS submanifold (see Definition 3.77). Then we vary the ECS submanifolds and the EC tubular neighbourhood maps. This will be done by choosing suitable diffeomorphisms near the singular points.

Definition 3.76. Let P be an end conical singular (ECS) submanifold of Y with end cone C as in Definition 3.53. Let $\Upsilon_C : V_C \rightarrow U_C$ be the conical tubular neighbourhood map of C as in Definition 3.41. A tubular neighbourhood map $\Upsilon_P : V_P \rightarrow U_P$ of P is called **end conical (EC)** if V_P and Υ_P agree with $\Upsilon_*(V_C)$ and $\Upsilon \circ \Upsilon_C \circ \Upsilon_*^{-1}$ on $\Upsilon(\iota((0, \varepsilon_0) \times \Sigma))$ respectively. ♣

Definition 3.77. Let P be a CS submanifold as in Definition 3.53. We define an end conical singular submanifold P_C which is diffeomorphic to \hat{P} but conical on the ends as follows

$$P_C := K_P \cup \left(\bigcup_{i=1}^m (\Upsilon^i \circ \Upsilon_{C_i})((1 - \rho_{\varepsilon_0})(\Psi_P^i - \iota_i)) \right).$$

Here ρ_{ε_0} is the cut off function defined in (3.50).

We define

$$C_{i,\varepsilon_0} := \iota((0, \varepsilon_0) \times \Sigma_i) \text{ and } K_{P_C} := P_C \setminus \bigcup_{i=1}^m \Upsilon^i(C_{i,\varepsilon_0}).$$




Figure 5: Construction of ECS submanifold P_C from a CS submanifold P (here $m = 1$).

Definition 3.78. Let P be an CS submanifold as in Definition 3.53. We choose a ECS submanifold as in Definition 3.77. Let

$$\Upsilon_{P_C} : V_{P_C} \rightarrow U_{P_C}$$

be an EC tubular neighbourhood map of P_C as in Definition 3.76. There is a section α in $V_{P_C} \subset NP_C$ which is zero on K_P and $\Upsilon_*^i \circ (\Psi_P^i - \iota_i) = \alpha \circ \Upsilon^i$ such that $\Upsilon_{P_C}(\Gamma_\alpha)$ is P . ♣

Denote

$$\begin{aligned} V_{C_{i,\varepsilon_0}} &:= V_{C_i|C_{i,\varepsilon_0}}, & U_{C_{i,\varepsilon_0}} &:= U_{C_i|C_{i,\varepsilon_0}}, \\ P_{C_{i,\varepsilon_0}} &:= P_{C_i|Y^i(C_{i,\varepsilon_0})}, & V_{P_{C_{i,\varepsilon_0}}} &:= V_{P_C|Y^i(C_{i,\varepsilon_0})}. \end{aligned}$$

The following commutative diagram helps us to keep track of the definitions above.

$$\begin{array}{ccccc} & & \Psi_{P^i-t_i} & & \\ & & \curvearrowright & & \\ C_{i,\varepsilon_0} & \xleftarrow{\quad} & V_{C_{i,\varepsilon_0}} & \xrightarrow{\Upsilon_{C_i}} & U_{C_{i,\varepsilon_0}} \\ & \downarrow \Upsilon^i & \downarrow \Upsilon_*^i & & \downarrow \Upsilon^i \\ P_{C_{i,\varepsilon_0}} & \xleftarrow{\quad} & V_{P_{C_{i,\varepsilon_0}}} = \Upsilon_*^i(V_{C_{i,\varepsilon_0}}) & \xrightarrow{\Upsilon_{P_C}} & Y \setminus K_Y. \\ & \curvearrowleft \alpha & & & \end{array}$$

We will now define the weighted C^∞ -topology on the moduli space \mathcal{M}_{cs} by specifying a basis.

Definition 3.79 (Weighted C^∞ -topology). Let $P \in \mathcal{M}_{\text{cs}}^\phi$ be a CS associative submanifold with singularities at p_i modeled on cones C_i , $i = 1, \dots, m$ as in Definition 3.53. Let Σ_i be the link of C_i . Set,

$$\tau_0 := (\phi, p_1, \dots, p_m, 0, \dots, 0).$$

We choose a ECS submanifold as in Definition 3.77. Let $\Upsilon_{P_C} : V_{P_C} \rightarrow U_{P_C}$ be an EC tubular neighbourhood map of P_C as in Definition 3.76. Since the singular points p_i and associative cones C_i are allowed to vary we need to also vary the P_C and Υ_{P_C} . This is done as follows.

By Theorem 1.3, there are tubular neighbourhood maps $\Upsilon_{\Sigma_i} : V_{\Sigma_i} \rightarrow U_{\Sigma_i}$ of Σ_i and obstruction maps $\text{ob}_{\Sigma_i} : \mathcal{J}_{\Sigma_i} \rightarrow \mathcal{O}_{\Sigma_i}$. Let U_ϕ and U_{p_i} be sufficiently small neighbourhoods of ϕ in \mathcal{P} and p_i in $Y^i(B(0, \varepsilon_0)) \subset Y$ respectively. Set

$$U_{\tau_0} := U_\phi \times \left(\prod_{i=1}^m U_{p_i} \right) \times \left(\prod_{i=1}^m \mathcal{J}_{\Sigma_i} \right).$$

The inclusion map of normal vector fields in Definition 2.17 induces smooth families of vector fields

$$(3.80) \quad \mathcal{J}_{\Sigma_i} \xrightarrow{\tilde{\cdot}} \text{Vect}(V_{\Sigma_i}) \xrightarrow{d\Upsilon_{\Sigma_i}} \text{Vect}(U_{\Sigma_i}) \hookrightarrow \text{Vect}(S^6) \xrightarrow{\rho_{\varepsilon_0}} \text{Vect}(B(0, R)).$$

Lets denote the vector fields by v_{ξ_i} whenever $\xi_i \in \mathcal{J}_{\Sigma_i}$. There are also another smooth family of vector fields

$$(3.81) \quad U_\phi \times U_{p_i} \rightarrow \text{Vect}(B(0, R))$$

defined by $(\phi', p'_i) \mapsto \rho_{\varepsilon_0} v_{\phi', p'_i}$ which is as follows. Given $\phi' \in U_\phi$ and $p'_i \in U_{p_i}$ there is a $v_{\phi', p'_i} \in M_7(\mathbf{R})$ satisfying $v_{\phi', p'_i} \cdot (\Upsilon^i)^* \phi(p_i) = (\Upsilon^i)^* \phi'(p'_i)$. The (3.80) and (3.81) together yield a smooth family of vector fields

$$(3.82) \quad U_\phi \times \left(\prod_{i=1}^m U_{p_i} \right) \times \left(\prod_{i=1}^m \mathcal{J}_{\Sigma_i} \right) \rightarrow \prod_{i=1}^m \text{Vect}(B(0, R)),$$

that is,

$$\tau := (\phi_\tau, (p_i^\tau), (\xi_i^\tau)) \mapsto (v_i^\tau := \rho_{\varepsilon_0} v_{\phi_\tau, p_i^\tau} + v_{\xi_i^\tau}).$$

The time-1 flows of the family of vector fields in (3.82) defines a smooth family of diffeomorphisms,

$$\Phi : U_\phi \times \left(\prod_{i=1}^m U_{p_i} \right) \times \left(\prod_{i=1}^m \mathcal{J}_{\Sigma_i} \right) \rightarrow \prod_{i=1}^m \text{Diff}(B(0, R)), \quad \tau \mapsto (\Phi_i^\tau)$$

In particular, the map Φ satisfies the following:

- (i) $\Upsilon_i^\tau := \Upsilon^i \circ \Phi_i^\tau$ is a G_2 -coordinate system at p_i^τ for the G_2 -structure ϕ_τ ,
- (ii) $\Phi_i^{\tau_0}$ is identity on $B(0, R)$ and Φ_i^τ is equal to identity outside $B(0, 2\varepsilon_0)$ for all $\tau \in U_{\tau_0}$.

For each τ in U_{τ_0} , we define

$$P_C^\tau := \left(\bigcup_{i=1}^m \Upsilon_i^\tau \circ (\Upsilon^i)^{-1}(P_C) \right) \cup K_P \text{ and}$$

$$\Upsilon_{P_C}^\tau := \left(\bigcup_{i=1}^m \Upsilon_i^\tau \circ (\Upsilon^i)^{-1} \circ \Upsilon_{P_C} \right) \bigcup \Upsilon_{P_C|K_P} : V_{P_C} \rightarrow Y.$$

We choose a rate $\mu > 1$ satisfying $(1, \mu_i) \cap \mathcal{D}_{C_\tau^i} = \emptyset$ for all $\tau \in U_{\tau_0}$, where C_τ^i are associative cones with links $\Upsilon_{\Sigma_i}(\xi_i^\tau)$, $\xi_i^\tau \in \mathcal{J}_{\Sigma_i}$. Here U_{τ_0} is always chosen small enough so that we can choose such a μ . Finally, the basis of the **weighted C^∞ -topology** on the moduli space \mathcal{M}_{cs} is a collection of all sets of the form

$$\{(\phi_\tau, \Upsilon_{P_C}^\tau(\Gamma_u)) : u \in \mathcal{V}_{P_C, \mu}, \tau \in \mathcal{V}_{\tau_0}\}$$

where \mathcal{V}_{τ_0} is an open set in U_{τ_0} and $\mathcal{V}_{P_C, \mu}$ is an open set in $C_\mu^\infty(V_{P_C}) := C_{P_C, \mu}^\infty \cap C^\infty(V_{P_C})$. Anyway this definition will be independent of such choices of μ (see Remark 3.101). \spadesuit

The following definition identifies two normal bundles NP_C and NP .

Definition 3.83. Let $P \in \mathcal{M}_{\text{cs}}$ be a CS associative submanifold with an asymptotic cone C . Let P_C be an ECS submanifold as in Definition 3.77 and Υ_{P_C} be an EC tubular neighbourhood map as in Definition 3.76. Then there is a **canonical** bundle isomorphism as in Definition 2.17,

$$\Theta_P^C : NP_C \rightarrow NP. \quad \spadesuit$$

Definition 3.84. Let $P \in \mathcal{M}_{\text{cs}}$ be an CS associative submanifold. Define $\mathfrak{F} : C_\mu^\infty(V_{P_C}) \times U_{\tau_0} \rightarrow C^\infty(NP_C)$ by for all $u \in C_\mu^\infty(V_{P_C})$, $\tau \in U_{\tau_0}$ and $w \in C_c^\infty(NP_C)$,

$$\langle \mathfrak{F}(u, \tau), w \rangle_{L^2} := \int_{\Gamma_u} \iota_w \Upsilon_{P_C}^* \psi_\tau.$$

The notation w in the integrand is the extension vector field of $w \in C^\infty(NP_C)$ in the tubular neighbourhood as in Notation 2.21. The L^2 inner product we choose here is coming from the canonical bundle isomorphism $\Theta_P^C : NP_C \rightarrow NP$ as in Definition 3.83 and the induced metric on NP of g_ϕ . \spadesuit




Figure 6: Construction of P_C^τ , a small deformation of P_C (here, $m = 1$).

Remark 3.85. Given a CS associative P with $\alpha \in C_\mu^\infty(V_{P_C})$ representing P as in Definition 3.76, $\mathfrak{F}(\alpha, \tau_0) = 0$ where τ_0 is as in Definition 3.79. Moreover, for $u \in C_\mu^\infty(V_{P_C})$ and $\tau \in U_{\tau_0}$ we have $\Upsilon_{P_C^\tau}(\Gamma_u) \in \mathcal{M}_{\text{cs}}$ iff $\mathfrak{F}(u, \tau) = 0$. \clubsuit

Proposition 3.86. For $\tau \in U_{\tau_0}$, the linearization of $\mathfrak{F}_\tau := \mathfrak{F}(\cdot, \tau)$ at $u \in C_{P_C, \mu}^\infty(V_{P_C})$,

$$d\mathfrak{F}_{\tau|u} : C_{P_C, \mu}^\infty \rightarrow C_{P_C, \mu-1}^\infty$$

is given for all $v \in C_{P_C, \mu}^\infty$ and $w \in C_c^\infty(NP)$ by $\langle d\mathfrak{F}_{\tau|u}(v), w \rangle = \int_{\Gamma_u} \iota_w \mathcal{L}_v(\Upsilon_{P_C^\tau}^* \psi_\tau)$. This is same as

$$\int_{\Gamma_u} \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \langle [e_2, e_3, \nabla_{e_1} v], w \rangle + \int_{\Gamma_u} \langle H_{\Gamma_u}, v \rangle \iota_w(\Upsilon_{P_C^\tau}^* \psi_\tau) + \iota_w \nabla_v(\Upsilon_{P_C^\tau}^* \psi_\tau) + \iota_{\nabla_w v}(\Upsilon_{P_C^\tau}^* \psi_\tau),$$

where $\{e_1, e_2, e_3\}$ is a local orthonormal frame for $T\Gamma_u$ and the associator $[\cdot, \cdot, \cdot]$ is defined with respect to $\Upsilon_{P_C^\tau}^* \phi_\tau$.

Proof. For a family $\{u + tv \in C_\mu^\infty(V_{P_C}) : |t| \ll 1\}$ we have

$$\frac{d}{dt} \Big|_{t=0} \mathcal{F}_{u+tv}(w) = \frac{d}{dt} \Big|_{t=0} \int_{\Gamma_{u+tv}} \iota_w(\Upsilon_{P_C^\tau}^* \psi_\tau) = \int_{\Gamma_u} \mathcal{L}_v \iota_w(\Upsilon_{P_C^\tau}^* \psi_\tau) = \int_{\Gamma_u} \iota_w \mathcal{L}_v(\Upsilon_{P_C^\tau}^* \psi_\tau) + \iota_{[v, w]}(\Upsilon_{P_C^\tau}^* \psi_\tau).$$

As $[v, w] = 0$ (see Notation 2.21), this is further equal to the following:

$$\begin{aligned} & \int_{\Gamma_u} \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \iota_w(\Upsilon_{P_C^\tau}^* \psi_\tau)(\nabla_{e_1} v, e_2, e_3) + \int_{\Gamma_u} \iota_{\nabla_w v}(\Upsilon_{P_C^\tau}^* \psi_\tau) + \iota_w \nabla_v(\Upsilon_{P_C^\tau}^* \psi_\tau) \\ &= \int_{\Gamma_u} \left\langle \sum_{\substack{\text{cyclic} \\ \text{permutations}}} [e_2, e_3, \nabla_{\Gamma_u, e_1}^\perp v], w \right\rangle \\ &+ \int_{\Gamma_u} \iota_w(\Upsilon_{P_C^\tau}^* \psi_\tau)(\nabla_{\Gamma_u, e_1}^\parallel v, e_2, e_3) + \iota_{\nabla_w v}(\Upsilon_{P_C^\tau}^* \psi_\tau) + \iota_w \nabla_v(\Upsilon_{P_C^\tau}^* \psi_\tau). \quad \blacksquare \end{aligned}$$

Corollary 3.87. *Let $u \in C_\mu^\infty(V_{P_C})$ and $\tau \in U_{\tau_0}$. If $\Upsilon_{P_C^\tau}(\Gamma_u) \in \mathcal{M}_{\text{CS}}$, a CS associative submanifold then $d\mathfrak{F}_{\tau|u}$ is given by the following: for all $v \in C_{P_C, \mu}^\infty$ and $w \in C_c^\infty(NP_C)$,*

$$\langle d\mathfrak{F}_{\tau|u}(v), w \rangle_{L^2} = \int_{\Gamma_u} \left\langle \sum_{\substack{\text{cyclic} \\ \text{permutations}}} [e_2, e_3, \nabla_{e_1} v], w \right\rangle + \int_{\Gamma_u} \iota_w \nabla_v (\Upsilon_{P_C^\tau}^* \psi_\tau). \quad \blacksquare$$

Proposition 3.88. *For $\tau \in U_{\tau_0}$, the linearization of \mathfrak{F}_τ at $u \in C_{P_C, \mu}^\infty(V_{P_C})$, $d\mathfrak{F}_{\tau|u}$ is a formally self adjoint first order differential operator.*

Proof. For all $v, w \in C_c^\infty(NP_C)$, the difference $d\mathfrak{F}_{\tau|u}(v)(w) - d\mathfrak{F}_{\tau|u}(w)(v)$ is

$$\int_{\Gamma_u} \mathcal{L}_v \iota_w (\Upsilon_{P_C^\tau}^* \psi_\tau) - \mathcal{L}_w \iota_v (\Upsilon_{P_C^\tau}^* \psi_\tau) = \int_{\Gamma_u} \iota_w \mathcal{L}_v (\Upsilon_{P_C^\tau}^* \psi_\tau) + \iota_{[v, w]} (\Upsilon_{P_C^\tau}^* \psi_\tau) - \mathcal{L}_w \iota_v (\Upsilon_{P_C^\tau}^* \psi_\tau).$$

As $[v, w] = 0$ and $d\psi_\tau = 0$, this is equal to $\int_{\Gamma_u} \iota_w \mathcal{L}_v (\Upsilon_{P_C^\tau}^* d\psi_\tau) - d(\iota_w \iota_v (\Upsilon_{P_C^\tau}^* \psi_\tau)) = 0$. \blacksquare

Definition 3.89. Given a CS associative P with $\alpha \in C_\mu^\infty(V_{P_C})$ representing P as in Definition 3.76 and τ_0 is as in Definition 3.79, we define the differential operator $\mathfrak{L}_P : C_c^\infty(NP_C) \rightarrow C_c^\infty(NP_C)$ by

$$\mathfrak{L}_P u := d\mathfrak{F}_{\tau_0|_\alpha}(u). \quad \spadesuit$$

Proposition 3.90. *Given a CS associative P with $\alpha \in C_\mu^\infty(V_{P_C})$ representing P as in Definition 3.76 we have*

$$\mathfrak{L}_P = (\Theta_P^C)^{-1} \circ \mathbf{D}_P \circ \Theta_P^C,$$

where $\mathbf{D}_P : C_c^\infty(NP) \rightarrow C_c^\infty(NP)$ is the Fueter operator defined in Definition 2.23.

Proof. For all $v, w \in C_c^\infty(NP_C)$ we have,

$$\begin{aligned} \langle \mathfrak{L}_P v, w \rangle_{L^2} &= \langle \Theta_P^C \mathfrak{L}_P v, \Theta_P^C w \rangle_{L^2(NP)} \\ &= \int_{\Gamma_\alpha} \left\langle \sum_{\substack{\text{cyclic} \\ \text{permutations}}} [e_2, e_3, \nabla_{e_1}^\perp v], w \right\rangle + \int_{\Gamma_\alpha} \iota_w \nabla_v (\Upsilon_{P_C}^* \psi) \\ &= \int_P \left\langle \sum_{\substack{\text{cyclic} \\ \text{permutations}}} [e_2, e_3, \nabla_{e_1}^\perp (\Theta_P^C v)], \Theta_P^C w \right\rangle + \int_P \iota_{\Theta_P^C w} \nabla_{\Theta_P^C v} \psi \\ &= \langle \mathbf{D}_P \Theta_P^C v, \Theta_P^C w \rangle_{L^2(NP)}. \end{aligned}$$

The equality before the last equality holds, because $w - \Theta_P^C w \in TP$ and $v - \Theta_P^C v \in TP$, and $[\cdot, \cdot, \cdot]_P = 0$. \blacksquare

Proposition 3.91. *The operator \mathfrak{L}_P is a conically singular uniformly elliptic operator asymptotic to the conical operators \mathbf{D}_{C_i} .*

Proof. Since \mathbf{D}_P is elliptic and by Proposition 3.90 $\mathfrak{L}_P = (\Theta_P^C)^{-1} \circ \mathbf{D}_P \circ \Theta_P^C$, we obtain that \mathfrak{L}_P is elliptic.

Without loss of generality we assume that $m = 1$ and $C = C_i$. It remains to prove that \mathfrak{L}_P is CS asymptotic to \mathbf{D}_C . We substitute below $r = e^t$ and denote $C_t := (t, \infty) \times \Sigma$. For all $v, w \in C_c^\infty(NC_t)$ we have

$$\begin{aligned} \langle r^2 d\mathfrak{F}_{\tau_0|_\alpha} \Upsilon_*(r^{-1}v), \Upsilon_* w \rangle_{L^2} &= \int_{C_t} r \iota_w \mathcal{L}_v(\Upsilon_C^* \Upsilon^* \psi) \\ &= \int_{C_t} r \iota_w \mathcal{L}_v(\Upsilon_C^* \psi_0) + O(e^{-t}) \|v\|_{W^{1,2}(NC)} \|w\|_{L^2(NC)}. \end{aligned}$$

By Proposition 3.95 (i) we get

$$\langle \Upsilon^* \mathfrak{L}_P v, w \rangle_{L^2(NC)} = \langle \mathbf{D}_C v, w \rangle_{L^2(NC)} + O(r^{\mu-1}) \|v\|_{W^{1,2}(NC)} \|w\|_{L^2(NC)}.$$

This completes the proof of the proposition. \blacksquare

Definition 3.92. Let P be a CS associative submanifold and $\mathbf{D}_P : C_c^\infty(NP) \rightarrow C_c^\infty(NP)$ be the Fueter operator defined in Definition 2.23. Proposition 3.91 implies that it is a conically singular uniformly elliptic operator. Therefore it has the following canonical extensions:

$$\mathbf{D}_{P,\lambda}^{k,p} : W_{P,\lambda}^{k+1,p} \rightarrow W_{P,\lambda-1}^{k,p}, \quad \mathbf{D}_{P,\lambda}^{k,Y} : C_{P,\lambda}^{k+1,Y} \rightarrow C_{P,\lambda-1}^{k,Y}.$$

Similarly we have canonical extensions of the operator \mathfrak{L}_P to weighted function spaces:

$$\mathfrak{L}_{P,\lambda}^{k,p} : W_{P_C,\lambda}^{k+1,p} \rightarrow W_{P_C,\lambda-1}^{k,p}, \quad \mathfrak{L}_{P,\lambda}^{k,Y} : C_{P_C,\lambda}^{k+1,Y} \rightarrow C_{P_C,\lambda-1}^{k,Y}. \quad \spadesuit$$

Definition 3.93. Proposition 3.91 and Proposition 3.90 imply that \mathbf{D}_P is also a CS uniformly elliptic operator and asymptotic to the conical operator \mathbf{D}_C . We define the asymptotic limit map

$$i_{P,\lambda} : \text{Ker } \mathbf{D}_{P,\lambda} \rightarrow V_\lambda$$

to be the map $i_{P,\lambda}$ defined in Lemma 3.70 for \mathbf{D}_P . \spadesuit

Definition 3.94. For all $\tau \in U_{\tau_0}$, the nonlinear map $Q_\tau : C_\mu^\infty(V_{P_C}) \rightarrow C^\infty(NP_C)$ is defined by

$$Q_\tau := \mathfrak{F}_\tau - d\mathfrak{F}_{\tau_0} - \mathfrak{F}_\tau(0). \quad \spadesuit$$

Proposition 3.95. For all $\tau \in U_{\tau_0}$ and $u, v \in C_{P_C,\mu}^\infty(V_{P_C})$, and $\eta \in C_{P_C,\mu}^\infty$ we have

- (i) $|d\mathfrak{F}_{\tau_u}(\eta) - d\mathfrak{F}_{\tau_v}(\eta)| \lesssim (w_{P_C,1}|u - v| + |\nabla^\perp(u - v)|)(w_{P_C,1}|\eta| + |\nabla^\perp \eta|).$
- (ii) $|Q_\tau(u) - Q_\tau(v)| \lesssim (w_{P_C,1}|u| + |\nabla^\perp u| + w_{P_C,1}|v| + |\nabla^\perp v|)(w_{P_C,1}|u - v| + |\nabla^\perp(u - v)|).$
- (iii) $\|Q_\tau(u)\|_{C_{P_C,\mu-1}^0} \lesssim \|Q_\tau(u)\|_{C_{P_C,2\mu-2}^0} \lesssim \|u\|_{C_{P_C,\mu}^1}^2.$

Proof. Since τ is fixed by abusing notation we denote ψ_τ by ψ . For all $w \in C_c^\infty(NP_C)$ and $\eta \in C_{P_C, \mu}^\infty$ we write

$$\langle d\mathfrak{F}_{\tau|u}(\eta) - d\mathfrak{F}_{\tau|v}(\eta), w \rangle = \int_0^1 \frac{d}{dt} \left(d\mathcal{F}_{\tau|tu+(1-t)v}(\eta)(w) \right) dt$$

and using Proposition 3.86 this becomes

$$\int_0^1 \left(\frac{d}{dt} \int_{\Gamma_{tu+(1-t)v}} L_\eta \iota_w (\Upsilon_{P_C}^* \psi) \right) dt = \int_0^1 \int_{\Gamma_{tu+(1-t)v}} L_{(u-v)} L_\eta \iota_w (\Upsilon_{P_C}^* \psi) dt.$$

As, $[u - v, w] = 0$ and $[\eta, w] = 0$ the last expression is same as

$$\int_0^1 \int_{\Gamma_{tu+(1-t)v}} \iota_w L_{(u-v)} L_\eta (\Upsilon_{P_C}^* \psi) dt.$$

The required estimate in (i) now follows from Lemma A.1. The estimate in (ii) follows from (i) after writing

$$Q_\tau(u) - Q_\tau(v) = \int_0^1 dQ_{\tau|tu+(1-t)v}(u-v) dt = \int_0^1 (d\mathfrak{F}_{\tau|tu+(1-t)v}(u-v) - d\mathfrak{F}_{\tau|0}(u-v)) dt.$$

Finally (iii) follows from (ii). Indeed, substituting $v = 0$ we have

$$w_{P_C, 2\mu-2} |Q_\xi(u)| \lesssim w_{P_C, 2\mu-2} (w_{P_C, 1} |u| + |\nabla^\perp u|)^2 \lesssim (w_{P_C, \mu} |u| + w_{P_C, \mu-1} |\nabla^\perp u|)^2.$$

Since $\mu > 1$ therefore $w_{P_C, \mu-1} |Q_\xi(u)| \lesssim w_{P_C, 2\mu-2} |Q_\xi(u)|$. This completes the proof. \blacksquare

Proposition 3.96. For all $\tau \in U_{\tau_0}$ and $u, v \in C_{P_C, \mu}^{k+1, \gamma}(V_{P_C})$ we have

$$\|Q_\tau(u) - Q_\tau(v)\|_{C_{P_C, \mu-1}^{k, \gamma}} \lesssim \|Q_\tau(u) - Q_\tau(v)\|_{C_{P_C, 2\mu-2}^{k, \gamma}} \lesssim \|u - v\|_{C_{P_C, \mu}^{k+1, \gamma}} (\|u\|_{C_{P_C, \mu}^{k+1, \gamma}} + \|v\|_{C_{P_C, \mu}^{k+1, \gamma}}).$$

Proof. Since τ is fixed by abusing notation we again denote ψ_τ by ψ . With the above notation and appropriate product operation ‘ \cdot ’, one can express $\mathcal{L}_u \mathcal{L}_v \psi$ formally as a quadratic polynomial

$$\mathcal{L}_u \mathcal{L}_v \psi = O(f_1) \cdot u \cdot v + O(f_2) \cdot (u \cdot \nabla^\perp v + v \cdot \nabla^\perp u) + \psi \cdot \nabla^\perp u \cdot \nabla^\perp v$$

where $O(f_1) := \psi \cdot \nabla B + \psi \cdot B \cdot B + B \cdot \nabla \psi + \nabla^2 \psi + R \cdot \psi$ and $O(f_2) := \nabla \psi + B \cdot \psi$. With this observation and a similar computations as in Lemma A.1 one can prove the proposition. \blacksquare

Lemma 3.97. The linearization of $\mathfrak{F}(\alpha, \phi, \cdot)$ at $(p_1, \dots, p_m, 0, \dots, 0)$ is

$$L_1 : \bigoplus_{i=1}^m (\mathbf{R}^7 \oplus V_{1,i}) \rightarrow C_{P_C, \mu-1}^\infty,$$

which is defined by

$$L_1(n_1, \dots, n_m) = \sum_{i=1}^m \mathfrak{Q}_P(\rho_{\varepsilon_0} \Upsilon_*^i n_i).$$

Proof. Without loss of generality we assume that $m = 1$ and $C = C_i$. Suppose $\xi \in V_1$ and for each t small, $\tau_t := (\phi, p, t\xi)$. For all $w \in C_c^\infty(NP_C)$ we have

$$\langle L_1(\xi), w \rangle_{L^2} = \int_{\Gamma_\alpha} \frac{d}{dt} \Big|_{t=0} \iota_w(\Upsilon_{P_C}^* \psi) = \int_{\Gamma_\alpha} \iota_w \mathcal{L}_{(\rho_{\varepsilon_0} \xi)}(\Upsilon_{P_C}^* \psi)$$

The last equality follows from Definition 3.79.

Suppose $\hat{p} \in \mathbf{R}^7$ and for each t small, $\tau_t := (\phi, \Upsilon(t\hat{p}), 0)$. For all $w \in C_c^\infty(NP_C)$ we have

$$\langle L_1(\hat{p}), w \rangle_{L^2} = \int_{\Gamma_\alpha} \frac{d}{dt} \Big|_{t=0} \iota_w(\Upsilon_{P_C}^* \psi) = \int_{\Gamma_\alpha} \iota_w \mathcal{L}_{(\rho_{\varepsilon_0} \hat{p})}(\Upsilon_{P_C}^* \psi)$$

The last equality follows from Definition 3.79. The lemma now follows from the proof of Proposition 3.86. \blacksquare

Definition 3.98. Let P be a CS associative as in Definition 3.53 asymptotic to C_i , $i = 1, \dots, m$. Denote the cone of Σ_i by C_i . Let $\Sigma_i = \sqcup_{j=1}^l \Sigma_i^j$ be the decomposition into connected components. Let $\mathcal{M}^{\text{hol}} = \sqcup_{k \in I} \mathcal{Z}_k$ be a decomposition of the moduli space of closed holomorphic curves in S^6 as in Definition 1.4. Let \mathcal{Z}_i^j be the manifold in the above decomposition containing Σ_i^j . Set $\mathcal{Z}_i = \prod_{j=1}^l \mathcal{Z}_i^j$ as in Definition 1.8. We also denote by Σ_i the product $\prod_{j=1}^l \Sigma_i^j \in \mathcal{Z}_i$. Since $T_{\Sigma_i} \mathcal{Z}_i \subset T_0 \mathcal{J}_{\Sigma_i} = V_{1,i}$, we define

$$\tilde{\mathfrak{L}}_{P,\mu,\mathcal{Z}} : C_{P_C,\mu}^\infty \oplus \left(\bigoplus_{i=1}^m (\mathbf{R}^7 \oplus T_{\Sigma_i} \mathcal{Z}_i) \right) \rightarrow C_{P_C,\mu-1}^\infty$$

to be the restriction of the linearization of $\mathfrak{F}(\cdot, \phi, \cdot)$ at $(\alpha, \phi, p_1, \dots, p_m, 0, \dots, 0)$, that is,

$$\tilde{\mathfrak{L}}_{P,\mu,\mathcal{Z}}(u, n_1, \dots, n_m) = \mathfrak{L}_P u + L_1(n_1, \dots, n_m).$$

The operators \mathfrak{L}_P and L_1 are as in Definition 3.89 and Lemma 3.97. The operator $\tilde{\mathfrak{D}}_{P,\mu,\mathcal{Z}} : C_{P,\mu}^\infty \oplus \left(\bigoplus_{i=1}^m (\mathbf{R}^7 \oplus T_{\Sigma_i} \mathcal{Z}_i) \right) \rightarrow C_{P,\mu-1}^\infty$ is the operator $\tilde{\mathfrak{L}}_{P,\mu,\mathcal{Z}}$ under the identification map Θ_P^C (see Definition 3.83), that is,

$$\tilde{\mathfrak{D}}_{P,\mu,\mathcal{Z}}(u, n_1, \dots, n_m) := \Theta_P^C \tilde{\mathfrak{L}}_{P,\mu,\mathcal{Z}}((\Theta_P^C)^{-1} u, n_1, \dots, n_m) = \mathfrak{D}_P u + \Theta_P^C L_1(n_1, \dots, n_m) \quad \spadesuit$$

Lemma 3.99. *The linearization of $\mathfrak{F}(\alpha, \cdot, p_1, \dots, p_m, 0, \dots, 0)$ at ϕ is*

$$L_2 : T_\phi \mathcal{P} \rightarrow C_{P_C,\mu-1}^\infty,$$

which is defined by

$$\langle L_2(\hat{\phi}), w \rangle_{L^2} = \int_{\Gamma_\alpha} \iota_w(\Upsilon_{P_C}^* \hat{\psi}) + \sum_{i=1}^m \langle \mathfrak{L}_P(\rho_{\varepsilon_0} \Upsilon_*^i X_\phi^i), w \rangle_{L^2},$$

where $\hat{\psi} := \frac{d}{dt} \Big|_{t=0} \psi_t$, $\psi_t = *_{\phi_t} \phi_t$ and $\phi_t := \phi + t\hat{\phi}$. The vector fields $X_\phi^i \in M_7(\mathbf{R})$ are $v_{\phi+\hat{\phi}, p_i}$ as in Definition 3.79.

Proof. Without loss of generality we assume that $m = 1$ and $C = C_i$. Suppose $\hat{\phi} \in T_\phi \mathcal{P}$ and for each t small, $\tau_t := (\phi_t, p, 0)$. For all $w \in C_c^\infty(NP_C)$ we have,

$$\langle L_2(\hat{\phi}), w \rangle_{L^2} = \int_{\Gamma_\alpha} \frac{d}{dt} \Big|_{t=0} \iota_w(\Upsilon_{P_C}^* \psi_t) = \int_{\Gamma_\alpha} \iota_w(\Upsilon_{P_C}^* \hat{\psi}) + \int_{\Gamma_\alpha} \iota_w \mathcal{L}_{(\rho_{\varepsilon_0} X_{\hat{\phi}})}(\Upsilon_{P_C}^* \psi). \quad \blacksquare$$

Definition 3.100. Let \mathcal{P} be the space of paths $\phi : [0, 1] \rightarrow \mathcal{P}$ which are smooth as section over $[0, 1] \times Y$. Equip \mathcal{P} with the C^∞ topology. Define the 1-parameter moduli space of CS associatives by the fiber product

$$\mathcal{M}_{\text{CS}}^\phi := [0, 1] \times_{\mathcal{P}} \mathcal{M}_{\text{CS}}. \quad \spadesuit$$

If $\phi \in \mathcal{P}$ and $(t_0, P) \in \mathcal{M}_{\text{CS}}^\phi$, we denote

$$\hat{\phi} := \frac{d}{dt} \Big|_{t=t_0} \phi_t, \quad f_P := L_2(\hat{\phi}) \in C^\infty(NP_C), \quad \hat{f}_P := \Theta_P^C f_P \in C^\infty(NP)$$

where L_2 and Θ_P^C are as in Lemma 3.99 and Definition 3.83 respectively. We define

$$\tilde{\mathfrak{Q}}_{P,\mu,\mathcal{Z}} : \mathbf{R} \oplus C_{P_C,\mu}^\infty \oplus \left(\bigoplus_{i=1}^m (\mathbf{R}^7 \oplus T_{\Sigma_i} \mathcal{Z}_i) \right) \rightarrow C_{P_C,\mu-1}^\infty$$

by $\tilde{\mathfrak{Q}}_{P,\mu,\mathcal{Z}}(t, \tilde{u}) = \tilde{\mathfrak{Q}}_{P,\mu,\mathcal{Z}} \tilde{u} + t f_P$, where $\tilde{\mathfrak{Q}}_{P,\mu,\mathcal{Z}}$ is defined in Definition 3.98. The operator $\bar{\mathfrak{D}}_{P,\mu,\mathcal{Z}} : \mathbf{R} \oplus C_{P_C,\mu}^\infty \oplus \left(\bigoplus_{i=1}^m (\mathbf{R}^7 \oplus T_{\Sigma_i} \mathcal{Z}_i) \right) \rightarrow C_{P_C,\mu-1}^\infty$ is the operator $\tilde{\mathfrak{Q}}_{P,\mu,\mathcal{Z}}$ under the identification map Θ_P^C , that is,

$$\bar{\mathfrak{D}}_{P,\mu,\mathcal{Z}}(t, \tilde{u}) = \tilde{\mathfrak{D}}_{P,\mu,\mathcal{Z}} \tilde{u} + t \hat{f}_P,$$

where $\tilde{\mathfrak{D}}_{P,\mu,\mathcal{Z}}$ is again defined in Definition 3.98

Proof of Theorem 1.9. By Proposition 3.96, Proposition 3.86 and Lemma 3.97 we conclude that the extension of the map $\mathcal{F}_\phi := \mathcal{F}(\cdot, \phi, \cdot)$ to weighted Hölder spaces

$$\mathfrak{F}_{\phi,\mathcal{Z}} : C_{P_C,\mu}^{2,\gamma}(V_{P_C}) \times \prod_{i=1}^m U_{p_i} \times \prod_{i=1}^m \mathcal{Z}_i \cap \mathcal{J}_{\Sigma_i} \rightarrow C_{P_C,\mu-1}^{1,\gamma}$$

is a well-defined smooth map. Here the rate μ is chosen as in Definition 3.79. By Proposition 3.69 we have that the linearization of $\mathfrak{F}_{\phi,\mathcal{Z}}$ at $(\alpha, p_1, \dots, p_m, 0, \dots, 0)$, $\tilde{\mathfrak{Q}}_{P,\mu,\mathcal{Z}}$ is a Fredholm operator. Moreover, Proposition 3.72 implies that

$$\text{index } \tilde{\mathfrak{Q}}_{P,\mu,\mathcal{Z}} = - \sum_{i=1}^m \left(\frac{d_{-1,i}}{2} + \sum_{\lambda_i \in \mathcal{D}_{C_i} \cap (-1,1]} d_{\lambda_i} \right) + \sum_{i=1}^m (7 + \dim \mathcal{Z}_i) = - \sum_{i=1}^m \text{s-ind}_\Delta(C_i).$$

Applying the implicit function theorem to $\mathfrak{F}_{\phi,\mathcal{Z}}$ and Proposition 3.65 of elliptic regularity we obtain the existence of $\text{ob}_{P,\mathcal{Z}}$ in (i) (see [DK90, Proposition 4.2.19]) of the theorem except the following. If $u \in C_{P_C,\mu}^{2,\gamma}(V_{P_C})$ with $\mathfrak{F}_{\phi_0,\mathcal{Z}}(u, \xi) = 0$ for some $\xi \in \prod_{i=1}^m U_{p_i} \times \prod_{i=1}^m \mathcal{Z}_i \cap \mathcal{J}_{\Sigma_i}$, then $u \in C_{P_C,\mu}^\infty(V_{P_C})$. To prove this, we observe

$$0 = \mathfrak{L}_P \mathfrak{F}_{\phi,\mathcal{Z}}(u, \xi) = A_\xi(u, \nabla_{P_C}^\perp u) (\nabla_{P_C}^\perp)^2 u + B_\xi(u, \nabla_{P_C}^\perp u)$$

Since $A_\xi(u, \nabla_{P_C}^\perp u), B_\xi(u, \nabla_{P_C}^\perp u) \in C_{P_C, \mu}^{1, \gamma}$, by a weighted version of Schauder elliptic regularity for second order operator (similar to Proposition 3.65) we obtain $u \in C_{P_C, \mu}^{3, \gamma}$. By repeating this argument we get higher regularity. This completes the proof of (i).

The proof of (ii) is similar to (i). Indeed, we replace $\mathfrak{F}_{\phi, \mathcal{Z}}$ by $\mathfrak{F}_{\phi, \mathcal{Z}}$ as follows. If $\phi \in \mathcal{P}$ and $t_0 \in [0, 1]$, then there exists an interval $I \subset [0, 1]$ containing t_0 such that $\phi(I) \subset U_{\phi_{t_0}}$ which was defined in Definition 3.79. We define

$$\mathfrak{F}_{\phi, \mathcal{Z}} : I \times C_{P_C, \mu}^{2, \gamma}(V_{P_C}) \times \prod_{i=1}^m U_{p_i} \times \prod_{i=1}^m \mathcal{Z}_i \cap \mathcal{J}_{\Sigma_i} \rightarrow C_{P_C, \mu-1}^{1, \gamma}$$

by $\mathfrak{F}_{\phi, \mathcal{Z}}(t, \cdot) := \mathfrak{F}_{\phi_t, \mathcal{Z}}$, $t \in I$. The linearization of $\mathfrak{F}_{\phi, \mathcal{Z}}$ is $\bar{\mathfrak{Q}}_{P, \mu, \mathcal{Z}}$ as in Definition 3.100. The remaining proof is left to the reader. \blacksquare

Remark 3.101. If μ' lying in the same connected component of μ in $(1, 2) \setminus \mathcal{D}_C$, then $u \in C_{P_C, \mu}^\infty(V_{P_C})$ satisfying $\mathfrak{F}_\tau(u) = 0$ implies $u \in C_{P_C, \mu'}^\infty(V_{P_C})$. To see this we write $d\mathfrak{F}_{\tau|_0}(u) = -Q_\tau(u) - \mathfrak{F}_\tau(0)$. By Proposition 3.95 we see that $d\mathfrak{F}_{\tau|_0}$ is also a CS uniformly elliptic operator asymptotic to \mathbf{D}_C . Also we have $Q_\tau(u) \in C_{P_C, 2\mu-2}^{k, \gamma}$ and $\mathfrak{F}_\tau(0) \in C_{P_C, 1}^{k, \gamma}$. Therefore by Lemma 3.70 we can conclude that $u \in C_{P_C, \mu'}^\infty(V_{P_C})$. \clubsuit

3.6 Proof of genericity results: Floer's C_ε space and Taubes' trick

Proof of Theorem 1.11. We prove that $\mathcal{P}_{\text{cs}, \mathcal{Z}}^{\text{reg}}$ is comeager in \mathcal{P} in Lemma 3.106. The proof of the fact that $\mathcal{P}_{\text{cs}, \mathcal{Z}}^{\text{reg}}$ is comeager in \mathcal{P} is similar. Then (i) and (ii) in Theorem 1.11 are direct consequences of Theorem 1.9. \blacksquare

To prove Lemma 3.106 we would like to use the Sard-Smale theorem applied to the map

$$\mathfrak{F}_{\mathcal{Z}} : W_{P_C, \mu}^{k+1, p}(V_{P_C}) \times U_\phi \times \prod_{i=1}^m U_{p_i} \times \prod_{i=1}^m \mathcal{Z}_i \cap \mathcal{J}_{\Sigma_i} \rightarrow W_{P_C, \mu-1}^{k, p},$$

We have chosen Sobolev spaces over Hölder spaces as the former is separable. The reader may observe that all the above analysis will also go through with Sobolev spaces. A serious problem here is that $U_\phi \subset \mathcal{P}$ is not a Banach manifold. The standard way to deal with this is to consider \mathcal{P}_k , the space of all C^k co-closed G_2 -structures on Y to enlarge U_ϕ to a Banach manifold. This comes with a drawback that the map \mathfrak{F} will have only finitely many derivatives and it needs extra effort to check exactly how many in order to state the regularity of the moduli space. To avoid all this we will instead use Floer's C_ε space $\mathcal{P}_\varepsilon \subset \mathcal{P}$ of co-closed G_2 -structures. Since the Sard-Smale theorem yields the genericity results in \mathcal{P}_ε instead of \mathcal{P} we use Taubes' trick of exhausting the moduli spaces by countable compact subsets (see Definition 3.102) to conclude the genericity results in \mathcal{P} . For more details on this idea, see [Wen21].

Definition 3.102. For every $N, m \in \mathbb{N}$ and $\mathcal{Z} = \prod_{i=1}^m \mathcal{Z}_i$, we define $\mathcal{M}_{\text{cs}, N, \mathcal{Z}}^\phi \subset \mathcal{M}_{\text{cs}, \mathcal{Z}}^\phi$ to be the set of all CS associative submanifolds P in (Y, ϕ) with singularities at p_i , cones C_i with links $\Sigma_i \in \mathcal{Z}_i$, rates $\mu_i \in [1 + \frac{1}{N}, 2 - \frac{1}{N}]$, G_2 -coordinate systems $\Upsilon^i : B(0, R) \rightarrow Y$ and embeddings $\Upsilon_p^i : (0, 2\varepsilon_0) \times \Sigma_i \rightarrow B(0, R)$, $i = 1, \dots, m$ satisfying

- $1 > R \geq 4\epsilon_0 \geq \frac{1}{N}$, $|\nabla^k \Pi_{\Sigma_i}| \leq N$, $\text{Vol}(\Sigma_i) \leq N$ for all $k = 0, \dots, 3$,
- $d_{\mathcal{M}^{\text{hol}}}(\Sigma_i, \overline{\mathcal{Z}_i} \setminus \mathcal{Z}_i) \geq \frac{1}{N}$,
- $|\nabla^k \Pi_{K_P}| \leq N$ for all $k = 0, \dots, 3$ and $\text{Vol}(P) \leq N$,
- $|\nabla^k \Upsilon^i| \leq N$ and $|(\nabla_C^\perp)^k(\Psi_P^i - \iota)| \leq Nr^{\mu_i - k}$ for all $k = 0, \dots, 3$.

We define $\mathcal{P}_{\text{cs}, N, \mathcal{Z}}^{\text{reg}} \subset \mathcal{P}$ to be the set of all $\phi \in \mathcal{P}$ for which every $P \in \mathcal{M}_{\text{cs}, N, \mathcal{Z}}^\phi$ has the property that $\overline{\mathbb{D}}_{P, \mu, \mathcal{Z}}$ is surjective. \spadesuit

Remark 3.103. Obviously,

$$\mathcal{P}_{\text{cs}, \mathcal{Z}}^{\text{reg}} = \bigcap_{N \in \mathbb{N}} \mathcal{P}_{\text{cs}, N, \mathcal{Z}}^{\text{reg}} \subset \mathcal{P}. \quad \clubsuit$$

The following lemma about convergence of CS associative submanifolds can be deduced from [She95].

Lemma 3.104. *Let $\phi_n \in \mathcal{P}$ be a sequence of G_2 -structures on Y converging to ϕ in \mathcal{P} with C^∞ topology. Fix $N \in \mathbb{N}$ and \mathcal{Z} , and let $P_n \in \mathcal{M}_{\text{cs}, N, \mathcal{Z}}^{\phi_n}$ be a sequence of CS associative submanifolds. Then there is a subsequence of P_n converging to a CS associative $P \in \mathcal{M}_{\text{cs}, N, \mathcal{Z}}^\phi$ with weighted C^∞ -topology defined in Definition 3.79.*

Lemma 3.105. *For every $N \in \mathbb{N}$ and \mathcal{Z} , $\mathcal{P}_{\text{cs}, N, \mathcal{Z}}^{\text{reg}}$ is open in \mathcal{P} .*

Proof. The complement of $\mathcal{P}_{\text{cs}, N, \mathcal{Z}}^{\text{reg}}$ is closed. Indeed, it follows from Lemma 3.104 and the fact that surjectivity is an open condition. \blacksquare

Lemma 3.106. *For every $N \in \mathbb{N}$ and \mathcal{Z} , $\mathcal{P}_{\text{cs}, N, \mathcal{Z}}^{\text{reg}}$ is dense in \mathcal{P} . Hence $\mathcal{P}_{\text{cs}, \mathcal{Z}}^{\text{reg}}$ is comeager in \mathcal{P} .*

To prove Lemma 3.106 we need Floer's C_ϵ space.

Definition 3.107. Let $E \rightarrow M$ be a vector bundle over a compact Riemannian manifold M with or without boundary. For each integer $k \geq 0$, we denote $C^k(E)$ the Banach space of C^k -sections of E . Let \mathcal{E} be the set of all sequences $(\epsilon_k)_{k=0}^\infty$ of positive real numbers with $\epsilon_k \rightarrow 0$. For each $\epsilon \in \mathcal{E}$, Floer's C_ϵ space of sections of E is defined by

$$C_\epsilon(E) := \{s \in C^\infty(E) : \|s\|_{C_\epsilon} := \sum_{k=0}^\infty \epsilon_k \|s\|_{C^k} < \infty\}.$$

A pre-order $<$ on \mathcal{E} is defined by

$$\epsilon < \epsilon' \text{ iff } \limsup_{k \rightarrow \infty} \frac{\epsilon_k}{\epsilon'_k} < \infty. \quad \spadesuit$$

Remark 3.108. As M is compact, different C^k norms for different smooth atlases of M are equivalent but the norm on Floer's C_ϵ space is an infinite sum, therefore it might not be equivalent. But this not a big concern as no statement in any theorem will mention this space, it will only be used inside the proofs and where we can fix an atlas. \clubsuit

Lemma 3.109 ([Wen19, Appendix B]). *The Floer's C_ε spaces $C_\varepsilon(E)$ have the following properties:*

- (i) $C_\varepsilon(E)$ is a separable Banach space and $C_\varepsilon(E) \hookrightarrow C^\infty(E)$ is continuous. Moreover, $C_{\varepsilon'}(E) \hookrightarrow C_\varepsilon(E)$ is a continuous embedding if $\varepsilon < \varepsilon'$.
- (ii) For every countable subset \mathbf{Q} of $C^\infty(E)$ there exists an $\varepsilon_0 \in \mathcal{E}$ such that $\mathbf{Q} \subset C_\varepsilon(E)$ for all $\varepsilon < \varepsilon_0$. In particular,

$$C^\infty(E) = \bigcup_{\varepsilon \in \mathcal{E}} C_\varepsilon(E).$$

Lemma 3.110 (Doan and Walpuski [DW19, Proposition A.2]). *Let P be a CS associative in (Y, ϕ) with singularities at p and rate $\mu \in (1, 2) \setminus \mathcal{D}_C$. Then for all $w \in \text{coker } \mathfrak{L}_{P, \mu} = \ker \mathfrak{L}_{P, -2-\mu}$ there exists a 3-form $\hat{\phi} \in T_\phi \mathcal{P}$ supported away from p such that*

$$\langle L_2(\hat{\phi}), w \rangle_{L^2} \neq 0, \quad L_2 \text{ is defined in Lemma 3.99.}$$

Proof. By unique continuation, there exists an open set U in the interior of K_P on which w does not vanish identically. Let V be a tubular neighbourhood of U in Y . Choose a real valued smooth function f supported in V such that $df(w) \geq 0$ on U and there exists a point in U where $df(w) > 0$. Let $\Theta \in \Omega^3(Y)$ be an extension of the 3-form vol_U and $\iota_w d\Theta|_V = 0$. Then there exists a 3-form $\hat{\phi} \in T_\phi \mathcal{P}$ supported in V such that $\hat{\psi} = d(f\Theta) \in \Omega^4(Y)$, where $\hat{\psi}$ is as in Lemma 3.99. By Definition 3.79 we have $X_{\hat{\phi}} = 0$ and Lemma 3.99 implies that

$$\langle L_2(\hat{\phi}), w \rangle_{L^2} = \int_{\Gamma_\alpha} \iota_w(\Upsilon_{P_C}^* \hat{\psi}) = \int_U df(w) \text{vol}_U > 0. \quad \blacksquare$$

Proof of Lemma 3.106. For each $\varepsilon \in \mathcal{E}$, let $\Omega_\varepsilon^3(Y)$ be the Floer's C_ε space of smooth 3-forms on Y . We define

$$\mathcal{P}_\varepsilon := \mathcal{P} \cap \Omega_\varepsilon^3(Y) \quad \text{and} \quad \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}} := \{(\phi, P) \in \mathcal{M}_{\text{cs}, \mathcal{Z}} : \phi \in \mathcal{P}_\varepsilon, P \in \mathcal{M}_{\text{cs}, \mathcal{Z}}^\phi\}.$$

Suppose $(\phi, P) \in \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}$. Let $p_i, i = 1, \dots, m$ be the singular points of P and $\alpha \in C_\mu^\infty(V_{P_C})$ representing P as in Definition 3.76. Set $\tau_0 := (\alpha, \phi, p_1, \dots, p_m, 0, \dots, 0)$ and

$$U_{\tau_0, \varepsilon, \mathcal{Z}} := (U_\phi \cap \mathcal{P}_\varepsilon) \times \prod_{i=1}^m U_{p_i} \times \prod_{i=1}^m \mathcal{Z}_i.$$

Let $\mathfrak{F}_{\varepsilon, \mathcal{Z}} : W_\mu^{2,p}(V_{P_C}) \times U_{\tau_0, \varepsilon, \mathcal{Z}} \rightarrow W_{P_C, \mu-1}^{1,p}$ be the restriction of \mathfrak{F} . We define

$$\mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}^{\text{reg}} := \{(\phi, P) \in \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}} : d\mathfrak{F}_{\varepsilon, \mathcal{Z}}|_{\tau_0} \text{ is surjective}\}.$$

By the Implicit function theorem [MS12, Theorem A.3.3] we obtain that $\mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}^{\text{reg}}$ is a separable Banach manifold. Moreover by [MS12, Lemma A.3.6] we see that the canonical projection map $\pi_{\varepsilon, \mathcal{Z}} : \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}^{\text{reg}} \rightarrow \mathcal{P}_\varepsilon$ is a Fredholm map. Therefore again by [MS12, Theorem A.3.3] we conclude that there exists a comeager subset $\mathcal{P}_{\varepsilon, \text{cs}, \mathcal{Z}}^{\text{reg}} \subset \mathcal{P}_\varepsilon$ having the property that for each $\phi \in \mathcal{P}_{\varepsilon, \text{cs}, \mathcal{Z}}^{\text{reg}}$ the linear operator $\tilde{\mathbf{D}}_{P, \mu, \mathcal{Z}}$ is surjective for all P with $(\phi, P) \in \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}^{\text{reg}}$.

With this preparation we are now ready to prove the lemma. Suppose $\phi_0 \in \mathcal{P}$. We can choose ε sufficiently small so that $\phi_0 \in \mathcal{P}_\varepsilon$. By Lemma 3.110 and Lemma 3.109 we can conclude that if $P \in \mathcal{M}_{\text{cs}, \mathcal{Z}}^{\phi_0}$ then $(\phi_0, P) \in \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}^{\text{reg}}$ for all sufficiently small ε . Since $\mathcal{P}_{\varepsilon, \text{cs}, \mathcal{Z}}^{\text{reg}}$ is dense in \mathcal{P}_ε , there exists a sequence $\phi_n \in \mathcal{P}_{\varepsilon, \text{cs}, \mathcal{Z}}^{\text{reg}}$ converging to ϕ_0 in C_ε -topology and hence in C^∞ -topology. Our claim is that $\phi_n \in \mathcal{P}_{\text{cs}, N, \mathcal{Z}}^{\text{reg}}$ for all n sufficiently large. If not, then there is a subsequence $P_n \in \mathcal{M}_{\text{cs}, N, \mathcal{Z}}^{\phi_n}$ such that $\widetilde{\mathcal{D}}_{P_n, \mu_n, \mathcal{Z}}$ are not surjective. By Lemma 3.104 we obtain that P_n converges to a CS associative P in (Y, ϕ_0) . Since $(\phi_0, P) \in \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}^{\text{reg}}$ therefore $(\phi_n, P_n) \in \mathcal{M}_{\text{cs}, \varepsilon, \mathcal{Z}}^{\text{reg}}$ for all n sufficiently large which contradicts to the fact that $\widetilde{\mathcal{D}}_{P_n, \mu_n, \mathcal{Z}}$ are not surjective. \blacksquare

4 Desingularizations of conically singular associative submanifolds

Let (Y, ϕ) be a co-closed G_2 -manifold. We will glue a conically singular (CS) associative submanifold P in (Y, ϕ) with singularity at one point and rescaled asymptotically conical (AC) associative submanifolds εL in \mathbf{R}^7 with $\varepsilon > 0$ to construct closed approximate associative submanifolds P_ε . We have seen in Theorem 2.26 that for generic co-closed G_2 structures the moduli space of closed associative submanifolds is 0-dimensional and therefore we can not expect to deform the P_ε to associative submanifolds \tilde{P}_ε (desingularization) in (Y, ϕ) . In this section we will see that under some hypothesis we can do this in a 1-parameter family of co-closed G_2 -structures¹².

4.1 Approximate associative desingularizations

Let P be a CS associative submanifold of a co-closed G_2 -manifold (Y, ϕ) with a singularity at p , rate $\mu \in (1, 2]$, cone C as in Definition 3.53. Let L be an AC associative submanifold in \mathbf{R}^7 with the same cone C and rate $\nu < 1$ as in Definition 3.42. The real numbers $R, R_\infty, \varepsilon_0$ are as in Definition 3.42 and Definition 3.53. Let P_C, NP_C, K_{P_C} be as in Definition 3.77 and L_C, NL_C, K_{L_C} be as in Definition 3.49.

Let $\phi \in \mathcal{P}$ be a path of co-closed G_2 -structures and $t_0 \in (0, 1)$ such that $\phi(t_0) = \phi_{t_0} = \phi$. Set $\tau_0 := (\phi, p, C)$. Let U_{τ_0} be as in Definition 3.79. Let $T > 0$ be sufficiently small such that for each $t \in (t_0 - T, t_0 + T)$ we have

$$\tau_t := (\phi_t, p, C) \in U_{\tau_0}.$$

Let $Y^{\tau_t}, P_C^{\tau_t}$ and $Y_{P_C^{\tau_t}}$ be as in Definition 3.79.

Definition 4.1. For any sufficiently small $\varepsilon > 0$, we define a closed 3-dimensional submanifold $P_{\varepsilon, t, C}$ of Y as follows. For a real number $0 < q < 1$ we define a real number, $\delta := (\varepsilon R_\infty)^q$. Then for sufficiently small ε we have

$$\varepsilon R_\infty < 2\varepsilon R_\infty < \delta < 2\delta < \delta^{\frac{1}{2}} < 2\delta^{\frac{1}{2}} < \varepsilon_0 < 2\varepsilon_0 < R.$$

We use the notation $A(a, b)$ for the annulus $\{x \in \mathbf{R}^7 : a < |x| < b\}$. For any $t \in (t_0 - T, t_0 + T)$, we define

$$P_{\varepsilon, t, C} := P_{\varepsilon, t, C}^+ \cup P_{\varepsilon, t, C}^-$$

¹²One can generalize this by using similar analysis to multiple isolated points and glue that many asymptotically conical associative submanifolds in \mathbf{R}^7 but in that case one need to desingularize in higher dimensional parameter space of co-closed G_2 structures, which we will not cover in this article.




Figure 7: AC submanifold εL_C and CS submanifold $P_C^{\tau_t}$.

where

$$P_{\varepsilon,t,C}^+ := \Upsilon^{\tau_t}(\varepsilon K_{L_C} \cup (C \cap A(\varepsilon R_\infty, 2\delta))) \quad \text{and} \quad P_{\varepsilon,t,C}^- := \Upsilon^{\tau_t}(C \cap A(\delta, 2\varepsilon_0)) \cup K_{P_C}.$$

Here $(\Upsilon^{\tau_t})^{-1}(P_{\varepsilon,t,C}^+) \subset \varepsilon L_C$ and $P_{\varepsilon,t,C}^- \subset P_C^{\tau_t}$. ♠




Figure 8: $P_{\varepsilon,t,C}$: gluing of εL_C and $P_C^{\tau_t}$.

Notation 4.2. If $t = t_0$ then we will now on omit the subscript t . ▶

Definition 4.3. The normal bundles $N(\varepsilon L_C)$ and NP_C can be glued to get a normal bundle

$$NP_{\varepsilon,C} = NP_{\varepsilon,C}^+ \cup NP_{\varepsilon,C}^-.$$

The scaling $s_\varepsilon : \mathbf{R}^7 \rightarrow \mathbf{R}^7$, $x \mapsto \varepsilon x$ induces an isomorphism $s_{\varepsilon*} : NL_C \rightarrow N(\varepsilon L_C)$ defined by

$$s_{\varepsilon*}v(x) := \varepsilon v(\varepsilon^{-1}x).$$

We have a tubular neighbourhood map

$$\Upsilon_{P_{\varepsilon,t,C}} := \Upsilon_{P_{\varepsilon,t,C}^+} \cup \Upsilon_{P_{\varepsilon,t,C}^-} : V_{P_{\varepsilon,C}} \rightarrow Y$$

where $(\Upsilon^{\tau_t})^{-1} \circ \Upsilon_{P_{\varepsilon,t,C}^+} \circ \Upsilon_*^{\tau_t}$ is the restriction of $s_\varepsilon \circ \Upsilon_{L_C} \circ s_{\varepsilon*}^{-1}$ and $\Upsilon_{P_{\varepsilon,t,C}^-}$ is the restriction of $\Upsilon_{P_C^{\tau_t}}$. ♠

Definition 4.4 (Approximate associative desingularization). Let $\alpha \in C^\infty(NP_C)$ and $\beta \in C^\infty(NL_C)$ be as in Definition 3.78 and Definition 3.51 representing P and L respectively. We define the **approximate associative desingularization** $P_{\varepsilon,t}^1$ by

$$P_{\varepsilon,t}^1 := \Upsilon_{P_{\varepsilon,t},C}(\alpha_\varepsilon^1),$$

where $\alpha_\varepsilon^1 \in C^\infty(V_{P_{\varepsilon,C}})$ is

$$(4.5) \quad \alpha_\varepsilon^1 := \rho_\delta \Upsilon_*(s_{\varepsilon*}\beta) + (1 - \rho_\delta)\alpha.$$

If there exists $\lambda_0 < \nu$ such that

- (i) $(\lambda_0, \nu] \cap \mathcal{D} = \{\lambda_1, \dots, \lambda_l\}$ with $\lambda_1 < \dots < \lambda_l$,
- (ii) there exist $s_0 > 0$ and $\alpha_i \in \ker \mathfrak{Q}_{P,\lambda_i}$, $\beta_i \in V_{\lambda_i}$ $i = 1, \dots, l$ such that

$$|(\nabla_C^\perp)^k(\beta - \sum_{i=1}^l \beta_i)| = O(r^{\max\{\lambda_0, 2\nu-1\}-k}) \text{ as } r \rightarrow \infty \text{ for all } k \in \mathbb{N} \cup \{0\}$$

and

$$|(\nabla_C^\perp)^k(\Upsilon_{P_C}^* \alpha_i - \beta_i)| = O(r^{\lambda_i + s_0 - k}) \text{ as } r \rightarrow 0 \text{ for all } k \in \mathbb{N} \cup \{0\},$$

we can define the **improved approximate associative desingularization** $P_{\varepsilon,t}^2$ by

$$P_{\varepsilon,t}^2 := \Upsilon_{P_{\varepsilon,t},C}(\alpha_\varepsilon^2),$$

where

$$(4.6) \quad \alpha_\varepsilon^2 := \rho_\delta \Upsilon_*(s_{\varepsilon*}\beta) + (1 - \rho_\delta) \left(\sum_{i=1}^l \varepsilon^{1-\lambda_i} \alpha_i + \alpha \right). \quad \spadesuit$$

Notation 4.7. We denote by $P_{\varepsilon,t}$ either $P_{\varepsilon,t}^1$ or $P_{\varepsilon,t}^2$ and by α_ε either α_ε^1 or α_ε^2 as in Definition 4.4.

We denote

$$v_0 := \max\{\lambda_0, 2\nu - 1\}$$

and

$$c_q := \min\{(1-q)(1-\nu), q(\mu-1)\}.^{13} \quad \blacktriangleright$$

Definition 4.8. The map $\tilde{\mathfrak{F}}_\varepsilon : C^\infty(V_{P_{\varepsilon,C}}) \times (t_0 - T, t_0 + T) \rightarrow C^\infty(NP_{\varepsilon,C})$ is defined by

$$\langle \tilde{\mathfrak{F}}_\varepsilon(u, t), w \rangle_{L^2} := \int_{\Gamma_u} \iota_w \Upsilon_{P_{\varepsilon,t},C}^* \psi_t$$

where $t \in (t_0 - T, t_0 + T)$, $u \in C^\infty(V_{P_{\varepsilon,C}})$ and $w \in C^\infty(NP_{\varepsilon,C})$. The notation w in the integrand is the extension vector field of $w \in C^\infty(NP_{\varepsilon,C})$ in the tubular neighbourhood as in Notation 2.21. \spadesuit

¹³It implies that $\varepsilon^{c_q} \lesssim \max\{\varepsilon^{1-\nu}\delta^{\nu-1}, \delta^{\mu-1}\} \lesssim \varepsilon^{c_q}$.




Figure 9: $P_{\epsilon, t}$: approximate associative desingularization.

Notation 4.9. We denote $\tilde{\mathfrak{F}}_\epsilon(\cdot, t)$ by $\mathfrak{F}_{\epsilon, t}$. If $t = t_0$ we omit the subscript t . ▶

Definition 4.10. The linearization of \mathfrak{F}_ϵ at $\alpha_\epsilon \in C^\infty(V_{P_{\epsilon, C}})$ is denoted by $\mathfrak{L}_{P_\epsilon}$, that is,

$$\mathfrak{L}_{P_\epsilon} := d\tilde{\mathfrak{F}}_\epsilon|_{\alpha_\epsilon} : C^\infty(NP_{\epsilon, C}) \rightarrow C^\infty(NP_{\epsilon, C}). \quad \spadesuit$$

Definition 4.11. Let $\beta \in C^\infty(NL_C)$ be as in Definition 3.51 representing L . We define the differential operator $\mathfrak{L}_L : C_c^\infty(NL_C) \rightarrow C_c^\infty(NL_C)$ by

$$\mathfrak{L}_L u := d\tilde{\mathfrak{F}}_\beta^{AC}(u),$$

where the map $\tilde{\mathfrak{F}}^{AC} : C_{L, v}^\infty(V_{L_C}) \rightarrow C^\infty(NL_C)$ is defined by

$$\langle \tilde{\mathfrak{F}}^{AC}(u), w \rangle_{L^2} := \int_{\Gamma_u} \iota_w \Upsilon_{L_C}^* \psi_\epsilon, \quad u \in C^\infty C_{L, v}^\infty(V_{L_C}), w \in C_c^\infty(NL_C). \quad \spadesuit$$

Remark 4.12. \mathfrak{L}_L is a AC uniformly elliptic operator asymptotic to \mathbf{D}_C . Moreover, there is a identification map $\Theta_L^C : NL_C \rightarrow NL$ similar to Θ_P^C (see Definition 3.83, Definition 3.92) such that

$$\mathbf{D}_L = \Theta_L^C \mathfrak{L}_L (\Theta_L^C)^{-1}. \quad \clubsuit$$

Proposition 4.13. $\mathfrak{L}_{P_\epsilon}$ is a formally self adjoint elliptic operator for all sufficiently small ϵ .

Proof. $\mathfrak{L}_{P_\epsilon}$ is a formally self adjoint operator follows from same arguments as in Proposition 3.88. It remains to prove that $\mathfrak{L}_{P_\epsilon}$ is an elliptic operator. We denote the restrictions of $\mathfrak{L}_{P_\epsilon}$ over $P_{\epsilon, C}^\pm$ by $\mathfrak{L}_{P_\epsilon^\pm}$. By Proposition 3.95 we obtain that

$$\mathfrak{L}_{P_\epsilon^-} = d\tilde{\mathfrak{F}}_{\tau_0}|_\alpha + O(\epsilon^{\epsilon q}) = \mathfrak{L}_P + O(\epsilon^{\epsilon q}).$$

Similar to Proposition 3.95 for AC associatives in \mathbf{R}^7 and the fact that $\Upsilon^* \psi = (\Upsilon^* \psi)(0) + O(r)$, we get

$$\mathfrak{L}_{P_\epsilon^+} = \Upsilon_* \mathfrak{L}_{\epsilon L} \Upsilon_*^{-1} + O(\epsilon^{\epsilon q}) = \epsilon^{-1} (\Upsilon_* S_{\epsilon^*}) \mathfrak{L}_L (\Upsilon_* S_{\epsilon^*})^{-1} + O(\epsilon^{\epsilon q}).$$

Since \mathfrak{L}_P and \mathfrak{L}_L are elliptic, therefore $\mathfrak{L}_{P_\epsilon}$ is an elliptic operator for all sufficiently small ϵ . ■

4.2 Weighted function spaces and estimates

Definition 4.14. For each $l \in \mathbf{R}$, a **weight function** $w_{\varepsilon,l} : P_{\varepsilon,C} \rightarrow (0, \infty)$ is any smooth function on $P_{\varepsilon,C}$ such that if $x = Y(r, \sigma) \in Y(B(0, R))$ then

$$w_{\varepsilon,l}(x) = (\varepsilon + r)^{-l}.$$

Let $k \geq 0$, $\gamma \in (0, 1)$. For a continuous section u of $NP_{\varepsilon,C}$ we define the **weighted L^∞ norm** and the **weighted Hölder semi-norm** respectively by

$$\|u\|_{L_{\varepsilon,l}^\infty} := \|w_{\varepsilon,l}u\|_{L^\infty(NP_{\varepsilon,C})}, \quad [u]_{C_{\varepsilon,l}^{0,\gamma}} := [w_{\varepsilon,l-\gamma}u]_{C^{0,\gamma}(NP_{\varepsilon,C})}.$$

For a continuous section u of $NP_{\varepsilon,C}$ with k continuous derivatives we define the **weighted C^k norm** and the **weighted Hölder norm** respectively by

$$\|u\|_{C_{\varepsilon,l}^k} := \sum_{j=0}^k \|(\nabla_{P_{\varepsilon,C}}^\perp)^j u\|_{L_{\varepsilon,l-j}^\infty}, \quad \|u\|_{C_{\varepsilon,l}^{k,\gamma}} := \|u\|_{C_{\varepsilon,l}^k} + [(\nabla_{P_{\varepsilon,C}}^\perp)^k u]_{C_{\varepsilon,l-k}^{0,\gamma}}.$$

We define the **weighted Hölder space** $C_{\varepsilon,l}^{k,\gamma}$, the **weighted C^k -space** $C_{\varepsilon,l}^k$ and the **weighted L^∞ -space** $L_{\varepsilon,l}^\infty$ to be the $C^{k,\gamma}$, C^k and L^∞ spaces with the weighted Hölder norm $\|\cdot\|_{C_{\varepsilon,l}^{k,\gamma}}$, weighted C^k -norm $\|\cdot\|_{C_{\varepsilon,l}^k}$ and weighted L^∞ -norm $\|\cdot\|_{L_{\varepsilon,l}^\infty}$ respectively. \spadesuit

Definition 4.15. The map $\tilde{\mathfrak{F}}_\varepsilon : C^\infty(V_{P_{\varepsilon,C}}) \times (t_0 - T, t_0 + T) \rightarrow C^\infty(NP_{\varepsilon,C})$ can be written as

$$\tilde{\mathfrak{F}}_\varepsilon(u, t) := \mathfrak{L}_{P_\varepsilon} u + (t - t_0)f_\varepsilon + \tilde{Q}_\varepsilon(u, t) + e_\varepsilon,$$

where $e_\varepsilon := \tilde{\mathfrak{F}}_\varepsilon(\alpha_\varepsilon) \in C^\infty(NP_{\varepsilon,C})$ and

$$f_\varepsilon := \left. \frac{d\tilde{\mathfrak{F}}_{\varepsilon,t}(\alpha_\varepsilon)}{dt} \right|_{t=t_0} \in C^\infty(NP_{\varepsilon,C}). \quad \spadesuit$$

Proposition 4.16 (Schauder estimate). *There exists a constant $C > 0$ such that for all $u \in C_{\varepsilon,l}^{k+1,\gamma}$ we have*

$$\|u\|_{C_{\varepsilon,l}^{k+1,\gamma}} \leq C \left(\|\mathfrak{L}_{P_\varepsilon} u\|_{C_{\varepsilon,l-1}^{k,\gamma}} + \|u\|_{L_{\varepsilon,l}^\infty} \right).$$

Proof. For $u \in C_{\varepsilon,l}^{k+1,\gamma}$, define u_\pm by restricting u over $P_{\varepsilon,C}^\pm$. Using the Schauder estimates in Proposition 3.65 and Proposition 3.95 we obtain

$$\|u_-\|_{C_{\varepsilon,l}^{k+1,\gamma}} \lesssim \|u_-\|_{C_{P,l}^{k+1,\gamma}} \lesssim \|\mathfrak{L}_{P_\varepsilon} u_-\|_{C_{P,l-1}^{k,\gamma}} + \|u_-\|_{L_{P,l}^\infty} \lesssim \|\mathfrak{L}_{P_\varepsilon} u_-\|_{C_{\varepsilon,l-1}^{k,\gamma}} + \varepsilon^{\zeta_q} \|u_-\|_{C_{\varepsilon,l}^{k+1,\gamma}} + \|u_-\|_{L_{\varepsilon,l}^\infty}$$

Similarly

$$\begin{aligned} \|u_+\|_{C_{\varepsilon,l}^{k+1,\gamma}} &\lesssim \varepsilon^{-l+1} \|s_{\varepsilon_*}^{-1} \Upsilon_*^{-1} u_+\|_{C_{L,l}^{k+1,\gamma}} \lesssim \varepsilon^{-l+1} \|\mathfrak{L}_L s_{\varepsilon_*}^{-1} \Upsilon_*^{-1} u_+\|_{C_{L,l-1}^{k,\gamma}} + \varepsilon^{-l+1} \|s_{\varepsilon_*}^{-1} \Upsilon_*^{-1} u_+\|_{L_{L,l}^\infty} \\ &\lesssim \varepsilon^{-1} \|\Upsilon_* s_{\varepsilon_*} \mathfrak{L}_L s_{\varepsilon_*}^{-1} \Upsilon_*^{-1} u_+\|_{C_{\varepsilon,l-1}^{k,\gamma}} + \|u_+\|_{L_{\varepsilon,l}^\infty} \\ &\lesssim \|\mathfrak{L}_{P_\varepsilon} u_+\|_{C_{\varepsilon,l-1}^{k,\gamma}} + \varepsilon^{\zeta_q} \|u_+\|_{C_{\varepsilon,l}^{k+1,\gamma}} + \|u_+\|_{L_{\varepsilon,l}^\infty}. \quad \blacksquare \end{aligned}$$

Proposition 4.17 (Error estimate). *The error $e_\varepsilon = \mathfrak{F}_\varepsilon(\alpha_\varepsilon)$ can be estimated as follows:*

$$\|\mathfrak{F}_\varepsilon(\alpha_\varepsilon)\|_{C_{\varepsilon,-1}^{k,Y}} \lesssim \begin{cases} \delta^\mu & \text{if } \alpha_\varepsilon = \alpha_\varepsilon^1 \text{ and } 0 < q \leq \frac{1-\nu}{\mu-\nu} \\ \delta^\mu + \varepsilon^{1-\nu_0} \delta^{\nu_0} + \varepsilon^{2(1-\nu)} & \text{if } \alpha_\varepsilon = \alpha_\varepsilon^2 \text{ and } \frac{\nu-\nu_0}{\nu+\mu-1+s_0-\nu_0} \leq q < 1 \end{cases}$$

where $\nu_0 = \max\{\lambda_0, 2\nu - 1\}$ and $\delta = \varepsilon^q$.

Proof. Let ϕ_e be the standard G_2 -structure on \mathbf{R}^7 and C be a cone. Since $|\phi - \Upsilon_*\phi_e| = O(r)$ on $B(0, R)$ therefore over $P_{\varepsilon,C} \cap \Upsilon(B(0, R))$ we have

$$|\mathfrak{F}_\varepsilon(\alpha_\varepsilon)| \lesssim |\mathfrak{F}_\varepsilon(\alpha_\varepsilon) - \mathfrak{F}_\varepsilon(\alpha_\varepsilon, \Upsilon_*\phi_e)| \lesssim |\mathfrak{F}_\varepsilon(\alpha_\varepsilon, \Upsilon_*\phi_e)| + r.$$

Case 1: Suppose $\alpha_\varepsilon = \alpha_\varepsilon^1$. Then over $P_{\varepsilon,C} \setminus \Upsilon(B(0, 2\delta))$, $\alpha_\varepsilon = \alpha$ and hence $\mathfrak{F}_\varepsilon(\alpha_\varepsilon) = 0$. Over $P_{\varepsilon,C} \cap \Upsilon(B(0, \delta))$ we have $\alpha_\varepsilon = \Upsilon_*(s_{\varepsilon*}\beta)$ and therefore $\mathfrak{F}_\varepsilon(\alpha_\varepsilon, \Upsilon_*\phi_e) = 0$. Over $P_{\varepsilon,C} \cap \Upsilon(A(\delta, 2\delta))$ we have

$$|\mathfrak{F}_\varepsilon(\alpha_\varepsilon, \Upsilon_*\phi_e)| \lesssim |\nabla\alpha_\varepsilon| \lesssim r^{\mu-1} + \varepsilon^{1-\nu} r^{\nu-1}.$$

Here the last inequality uses the assumption that $0 < q \leq \frac{1-\nu}{\mu-\nu}$ where $\delta = (\varepsilon R_\infty)^q$. From these estimates we conclude that $\|\mathfrak{F}_\varepsilon(\alpha_\varepsilon)\|_{C_{\varepsilon,-1}^0} \lesssim \delta^\mu$. A similar computation with Hölder seminorm and higher derivatives will prove that $\|\mathfrak{F}_\varepsilon(\alpha_\varepsilon)\|_{C_{\varepsilon,-1}^{k,Y}} \lesssim \delta^\mu$.

Case 2: Suppose $\alpha_\varepsilon = \alpha_\varepsilon^2$. Over $P_{\varepsilon,C} \setminus \Upsilon(B(0, 2\delta))$ we have

$$|\mathfrak{F}_\varepsilon(\alpha_\varepsilon)| \lesssim \left(r^{-1} \left| \sum_{i=1}^l \varepsilon^{1-\lambda_i} \alpha_i \right| + \left| \sum_{i=1}^l \varepsilon^{1-\lambda_i} \nabla \alpha_i \right| \right)^2 \lesssim \sum_{i=1}^l (\varepsilon^{-1} r)^{2(\lambda_i-1)} \lesssim (\varepsilon^{-1} r)^{2(\nu-1)}.$$

Over $P_{\varepsilon,C} \cap \Upsilon(A(\delta, 2\delta))$ we have $\mathfrak{F}_\varepsilon(\alpha_\varepsilon, \Upsilon_*\phi_e) = \mathbf{D}_C(\alpha_\varepsilon) + Q_C(\alpha_\varepsilon)$. The term $|\mathbf{D}_C(\alpha_\varepsilon)|$ can be estimated as follows.

$$\begin{aligned} |\mathbf{D}_C(\alpha_\varepsilon)| &\lesssim |\nabla \rho_\delta| |s_{\varepsilon*}\beta - \sum_{i=1}^l \varepsilon^{1-\lambda_i} \alpha_i - \alpha| + |\mathbf{D}_C(s_{\varepsilon*}\beta)| + |\mathbf{D}_C(\sum_{i=1}^l \varepsilon^{1-\lambda_i} \alpha_i + \alpha)| \\ &\lesssim (\varepsilon^{-1} r)^{\nu_0-1} + r^{\mu-1} + (\varepsilon^{-1} r)^{2(\nu-1)} + r^{\mu-1} \sum_{i=1}^l \varepsilon^{1-\lambda_i} r^{\lambda_i+s_0-1} + r^{2(\mu-1)}. \end{aligned}$$

If $q \geq \frac{\nu-\nu_0}{\nu+\mu-1+s_0-\nu_0}$ then we have

$$\varepsilon^{1-\lambda_i} r^{\lambda_i+\mu+s_0-2} \lesssim \varepsilon^{1-\nu} r^{\nu+\mu+s_0-2} \lesssim (\varepsilon^{-1} r)^{\nu_0-1}.$$

The term $|Q_C(\alpha_\varepsilon)|$ can be estimated as follows.

$$|Q_C(\alpha_\varepsilon)| \lesssim (r^{-1} |\alpha_\varepsilon| + |\nabla \alpha_\varepsilon|)^2 \lesssim (\varepsilon^{-1} r)^{2(\nu-1)} + r^{2(\mu-1)}.$$

Hence

$$\|\mathfrak{F}_\varepsilon(\alpha_\varepsilon)\|_{C_{\varepsilon,-1}^0} \lesssim \delta^\mu + \varepsilon^{1-\nu_0} \delta^{\nu_0} + \varepsilon^{2(1-\nu)}.$$

A similar computation with Hölder seminorm and higher derivatives will also prove that $\|\mathfrak{F}_\varepsilon(\alpha_\varepsilon)\|_{C_{\varepsilon,-1}^{k,Y}} \lesssim \delta^\mu + \varepsilon^{1-\nu_0} \delta^{\nu_0} + \varepsilon^{2(1-\nu)}$. \blacksquare

Proposition 4.18 (Quadratic estimate). For any $\varepsilon > 0$ there exists $0 < T_\varepsilon < T$ such that for all $u, v \in C^\infty(V_{P_\varepsilon, C})$, $\eta \in C^\infty(NP_{\varepsilon, C})$ and $t_1, t_2 \in (t_0 - T_\varepsilon, t_0 + T_\varepsilon)$ the following estimates hold.

$$(i) \quad |d\mathfrak{F}_{\varepsilon|u}(\eta) - d\mathfrak{F}_{\varepsilon|v}(\eta)| \lesssim (w_{\varepsilon,1}|u - v| + w_{\varepsilon,0}|\nabla^\perp(u - v)|)(w_{\varepsilon,1}|\eta| + |w_{\varepsilon,0}\nabla^\perp\eta|),$$

$$(ii) \quad \|Q_\varepsilon(u) - Q_\varepsilon(v)\|_{C_{\varepsilon,-1}^{k,Y}} \lesssim \varepsilon^{-1}\|u - v\|_{C_{\varepsilon,0}^{k+1,Y}} \left(\|u - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,Y}} + \|v - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,Y}} \right).$$

$$(iii) \quad \|\tilde{Q}_\varepsilon(u, t_1) - \tilde{Q}_\varepsilon(v, t_2)\|_{C_{\varepsilon,-1}^{k,Y}} \\ \lesssim \varepsilon^{-1} \left(\|u - v\|_{C_{\varepsilon,0}^{k+1,Y}} + |t_1 - t_2| \right) \left(\|u - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,Y}} + \|v - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,Y}} + |t_1 - t_0| + |t_2 - t_0| \right).$$

Proof. The proofs are exactly like Proposition 3.95, Proposition 3.96. An important observation here is that one can express $\mathcal{L}_u\mathcal{L}_v\psi$ over $V_{P_\varepsilon, C}$ formally as a quadratic polynomial

$$\mathcal{L}_u\mathcal{L}_v\psi = O(f_{1,\varepsilon}) \cdot u \cdot v + O(f_{2,\varepsilon}) \cdot (u \cdot \nabla^\perp v + v \cdot \nabla^\perp u) + \psi \cdot \nabla^\perp u \cdot \nabla^\perp v$$

where $O(f_{1,\varepsilon}) = O(w_{\varepsilon,1})$ and $O(f_{2,\varepsilon}) = O(w_{\varepsilon,0})$. Finally to see (ii) we write $Q_\varepsilon(u) - Q_\varepsilon(v)$ as

$$\int_0^1 dQ_{\varepsilon|tu+(1-t)v}(u - v)dt = \int_0^1 (d\mathfrak{F}_{\varepsilon|tu+(1-t)v}(u - v) - d\mathfrak{F}_{\varepsilon|\alpha_\varepsilon}(u - v))dt$$

and (ii) follows from (i). To prove (iii) we only observe that $\tilde{Q}_\varepsilon(u, t_1) - \tilde{Q}_\varepsilon(v, t_2)$ is equal to

$$\int_0^1 (d\tilde{\mathfrak{F}}_{\varepsilon|(tu+(1-t)v, tt_1+(1-t)t_2)}(u - v, 0) - d\tilde{\mathfrak{F}}_{\varepsilon|(\alpha_\varepsilon, tt_1+(1-t)t_2)}(u - v, 0))dt \\ + \int_0^1 (d\tilde{\mathfrak{F}}_{\varepsilon|(\alpha_\varepsilon, tt_1+(1-t)t_2)}(u - v, 0) - d\tilde{\mathfrak{F}}_{\varepsilon|(\alpha_\varepsilon, t_0)}(u - v, 0))dt \\ + \int_0^1 (d\tilde{\mathfrak{F}}_{\varepsilon|(tu+(1-t)v, tt_1+(1-t)t_2)}(0, t_1 - t_2) - d\tilde{\mathfrak{F}}_{\varepsilon|(tu+(1-t)v, t_0)}(0, t_1 - t_2))dt \\ + \int_0^1 (d\tilde{\mathfrak{F}}_{\varepsilon|(tu+(1-t)v, t_0)}(0, t_1 - t_2) - d\tilde{\mathfrak{F}}_{\varepsilon|(\alpha_\varepsilon, t_0)}(0, t_1 - t_2))dt. \quad \blacksquare$$

Definition 4.19. Denote the asymptotic limit maps (see Lemma 3.70) $i_{P,0}$ and $i_{L,0}$ by i_P and i_L respectively. We define

(i) the matching kernel \mathcal{K}^m by

$$\mathcal{K}^m := \{(u_L, u_P) \in \ker \mathfrak{Q}_{L,0} \times \ker \mathfrak{Q}_{P,0} : i_L(u_L) = i_P(u_P)\},$$

(ii) the approximate kernel of $\mathfrak{Q}_{P_\varepsilon}$ by

$$\mathcal{K}_\varepsilon^m := \{\rho_{\frac{\delta}{2}}\Upsilon_*(s_{\varepsilon*}u_L) + (1 - \rho_{2\delta})u_P : (u_L, u_P) \in \mathcal{K}^m\},$$

(iii) $\mathcal{X}_\varepsilon^{k+1,Y} := \{u \in C_{\varepsilon,0}^{k+1,Y} : \langle (\Upsilon_*s_{\varepsilon*})^{-1}u, u_L \rangle_{L_{K_L}^2} = \langle u, u_P \rangle_{L_{K_P}^2} = 0 \ \forall (u_L, u_P) \in \mathcal{K}^m\}. \quad \spadesuit$

Proposition 4.20 (Linear estimate I). Fix a real number $\omega > 0$. For any sufficiently small $\varepsilon > 0$, there exists a constant $C_\omega > 0$ independent of ε such that for all $u \in \mathcal{X}_\varepsilon^{k+1,Y}$, we have

$$\|u\|_{C_{\varepsilon,0}^{k+1,Y}} \leq C_\omega \varepsilon^{-\omega} \|\mathfrak{L}_{P_\varepsilon} u\|_{C_{\varepsilon,-1}^{k,Y}}.$$

Proof. By Schauder estimate in Proposition 4.16 it is enough to prove that

$$\|u\|_{L_{\varepsilon,0}^\infty} \leq C_\omega \varepsilon^{-\omega} \|\mathfrak{L}_{P_\varepsilon} u\|_{C_{\varepsilon,-1}^{k,Y}}.$$

We prove this by contradiction. If this is not true then there exist a decreasing sequence $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and u_n in $\mathcal{X}_{\varepsilon_n}^{k+1,Y}$ such that

$$\|u_n\|_{L_{\varepsilon_n,0}^\infty} = 1, \quad \varepsilon_n^{-\omega} \|\mathfrak{L}_{P_{\varepsilon_n}} u_n\|_{C_{\varepsilon_n,-1}^{k,Y}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Denote the restrictions of u_n over $P_{\varepsilon_n,C}^\pm$ by u_n^\pm . We define $u_{n,P} := u_n^-$ and $u_{n,L} := \varepsilon_n (\Upsilon_* s_{\varepsilon_n})^{-1} u_n^+$. By Schauder estimate in Proposition 4.16 we have $\|u_n\|_{C_{\varepsilon_n,0}^{k+1,Y}}$ is bounded and hence $\|u_{n,P}\|_{C_{P,0}^{k+1,Y}}$, $\|u_{n,L}\|_{C_{L,0}^{k+1,Y}}$ are also bounded. The Arzelà-Ascoli theorem implies that there exist subsequences which we call again $u_{n,P}$ and $u_{n,L}$, and there exist u_P in $C_{P,0}^{k+1,\frac{Y}{2}}$, u_L in $C_{L,0}^{k+1,\frac{Y}{2}}$ such that $\mathfrak{L}_P u_P = 0$, $\mathfrak{L}_L u_L = 0$, and

$$\|u_{n,P} - u_P\|_{C_{P,\text{loc}}^{k+1,\frac{Y}{2}}} \rightarrow 0 \quad \text{and} \quad \|u_{n,L} - u_L\|_{C_{L,\text{loc}}^{k+1,\frac{Y}{2}}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Moreover, by elliptic regularity in Proposition 3.65 we get, $u_P \in C_{P,0}^{k+1,Y}$ and $u_L \in C_{L,0}^{k+1,Y}$. By taking further subsequences if necessary we prove the following which leads to the contradiction:

- (i) $\|u_{n,P} - u_P\|_{C_{P_{\varepsilon_n},C^0}^{k+1,Y}} \rightarrow 0$ as $n \rightarrow \infty$,
- (ii) $\|u_{n,L} - u_L\|_{C_{L_{\varepsilon_n^{-1}}\delta_n,C^0}^{k+1,Y}} \rightarrow 0$ as $n \rightarrow \infty$ where $L_{\varepsilon_n^{-1}\delta_n,C} := s_{\varepsilon_n}^{-1} \Upsilon^{-1} P_{\varepsilon_n,C}^+ \subset L_C$,
- (iii) $u_P = 0$ and $u_L = 0$. Hence, $\|u_n\|_{L_{\varepsilon_n,0}^\infty} \leq \|u_n^+\|_{L_{P_{\varepsilon_n},C}^\infty} + \|u_n^-\|_{L_{P_{\varepsilon_n},C}^\infty} \rightarrow 0$ as $n \rightarrow \infty$.

To prove (i), we fix a real number p with $1 < p < \frac{1}{q}$. For sufficiently small $s > 0$ we have

$$\begin{aligned} \|\mathfrak{L}_P((1 - \rho_{\delta_n^p})u_n)\|_{C_{P,-s-1}^{k,Y}} &\lesssim (\varepsilon_n^{\zeta_{pq}} + \varepsilon_n^{\zeta_q}) \|u_n\|_{C_{0,\varepsilon_n}^{k+1,Y}} + \|\mathfrak{L}_{P_{\varepsilon_n}}(1 - \rho_{\delta_n^p})u_n\|_{C_{\varepsilon_n,-s-1}^{k,Y}} \\ &\lesssim (\varepsilon_n^{\zeta_{pq}} + \varepsilon_n^{\zeta_q} + \delta_n^{ps}) \|u_n\|_{C_{0,\varepsilon_n}^{k+1,Y}} + \|\mathfrak{L}_{P_{\varepsilon_n}} u_n\|_{C_{\varepsilon_n,-s-1}^{k,Y}} \\ &\lesssim \varepsilon_n^{\zeta_{pq}} + \varepsilon_n^{\zeta_q} + \delta_n^{ps} + \|\mathfrak{L}_{P_{\varepsilon_n}} u_n\|_{C_{\varepsilon_n,-1}^{k,Y}} \lesssim \varepsilon_n^{\zeta_{pq}} + \varepsilon_n^{\zeta_q} + \delta_n^{ps} + \varepsilon_n^\omega. \end{aligned}$$

Therefore there exists $v_{n,P} \in \ker \mathfrak{L}_{P,-s} = \ker \mathfrak{L}_{P,0}$ satisfying

$$\|(1 - \rho_{\delta_n^p})u_n - v_{n,P}\|_{C_{P,-s}^{k+1,Y}} \lesssim \varepsilon_n^{\zeta_{pq}} + \varepsilon_n^{\zeta_q} + \delta_n^{ps} + \varepsilon_n^\omega$$

and hence $\|u_{n,P} - v_{n,P}\|_{C_{P_{\varepsilon_n}, C^0}^{k+1, \gamma}} \lesssim (\varepsilon_n^{\zeta_{pq}} + \varepsilon_n^{\zeta_q} + \delta_n^{ps} + \varepsilon_n^\omega) \delta_n^{-s} \rightarrow 0$ as $n \rightarrow \infty$. As, $\ker \mathfrak{Q}_{P,0}$ is finite dimensional, the norms $\|\cdot\|_{C_{K_P}^{k+1, \frac{\gamma}{2}}}$ and $\|\cdot\|_{C_{P,0}^{k+1, \gamma}}$ are equivalent. Taking further subsequence yields $\|v_{n,P} - u_P\|_{C_{K_P}^{k+1, \frac{\gamma}{2}}} \rightarrow 0$ and hence $\|v_{n,P} - u_P\|_{C_{P,0}^{k+1, \gamma}} \rightarrow 0$ as $n \rightarrow \infty$.

To prove (ii), we define $\tilde{u}_{n,L} := \varepsilon_n (\Upsilon_{*s_{\varepsilon_n}})^{-1} (\rho_{\frac{1}{\delta_n}} u_n)$. For sufficiently small $s > 0$ we have

$$\begin{aligned} \|\mathfrak{Q}_L \tilde{u}_{n,L}\|_{C_{L,s-1}^{k,\gamma}} &\lesssim \varepsilon_n^{s-1} \|\Upsilon_{*s_{\varepsilon_n}} \mathfrak{Q}_L s_{\varepsilon_n}^{-1} \Upsilon_*^{-1} (\rho_{\frac{1}{\delta_n}} u_n)\|_{C_{\varepsilon_n, s-1}^{k,\gamma}} \\ &\lesssim \varepsilon_n^s (\varepsilon_n^{\zeta_q} + \delta_n^{\frac{\mu-1}{2}}) \|\rho_{\frac{1}{\delta_n}} u_n\|_{C_{\varepsilon_n, s}^{k+1, \gamma}} + \varepsilon_n^s \|\mathfrak{Q}_{P_{\varepsilon_n}} (\rho_{\frac{1}{\delta_n}} u_n)\|_{C_{\varepsilon_n, s-1}^{k,\gamma}} \\ &\lesssim \varepsilon_n^s \delta_n^{-\frac{s}{2}} \|u_n\|_{C_{\varepsilon_n, 0}^{k+1, \gamma}} + \varepsilon_n^s \delta_n^{-\frac{s}{2}} \|\mathfrak{Q}_{P_{\varepsilon_n}} u_n\|_{C_{\varepsilon_n, -1}^{k,\gamma}} \lesssim \varepsilon_n^s \delta_n^{-\frac{s}{2}} + \varepsilon_n^s \delta_n^{-\frac{s}{2}} \varepsilon_n^\omega. \end{aligned}$$

Then as in (i) there exists $v_{n,L} \in \ker \mathfrak{Q}_{L,s} = \ker \mathfrak{Q}_{L,0}$ satisfying

$$\begin{aligned} \|u_{n,L} - v_{n,L}\|_{C_{L,0}^{k+1, \gamma}} &\lesssim (\varepsilon_n^{-1} \delta_n)^s (\varepsilon_n^s \delta_n^{-\frac{s}{2}} + \varepsilon_n^s \delta_n^{-\frac{s}{2}} \varepsilon_n^\omega) \\ &\lesssim \delta_n^{\frac{s}{2}} + \delta_n^{\frac{s}{2}} \varepsilon_n^\omega \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Again as in (i) by taking further subsequence if necessary we get, $\|v_{n,L} - u_L\|_{C_{K_L}^{k+1, \frac{\gamma}{2}}} \rightarrow 0$ and

$\|v_{n,L} - u_L\|_{C_{L,0}^{k+1, \gamma}} \rightarrow 0$ as $n \rightarrow \infty$.

It remains to prove (iii). For each $\sigma \in \Sigma$ we have,

$$i_P u_P(\sigma) = \Upsilon_*^{-1} \lim_{n \rightarrow \infty} u_{n,P}(\Upsilon\left(\frac{3\delta_n}{2}, \sigma\right)) = \lim_{n \rightarrow \infty} u_{n,L}\left(\frac{3\varepsilon_n^{-1}\delta_n}{2}, \sigma\right) = i_L u_L(\sigma).$$

Since $u_n \in \mathcal{X}_{\varepsilon_n}^{k+1, \gamma}$ therefore as $n \rightarrow \infty$ we get

$$\|u_P\|_{L_{K_P}^2} = \langle u_P - u_{n,P}, u_P \rangle_{L_{K_P}^2} \lesssim \|u_{n,P} - u_P\|_{L_{P,0}^\infty} \|u_P\|_{L_{P,0}^\infty} \rightarrow 0$$

and hence $u_P = 0$. Similarly we get $u_L = 0$. ■

Definition 4.21. We define the extended matching kernel

$$\tilde{\mathcal{K}}^m := \{(u_L, u_P, t) \in \ker \mathfrak{Q}_{L,0} \oplus C_{P,0}^{k+1, \gamma} \oplus \mathbf{R} : \mathfrak{Q}_P u_P + t f_P = 0, i_L u_L = i_P u_P\}$$

where f_P is defined in Definition 3.100. ♠

Proposition 4.22 (Linear estimate II). *If the extended matching kernel $\tilde{\mathcal{K}}^m$ is equal to the matching kernel \mathcal{K}^m and $\omega > 0$ be any real number, then for any sufficiently small $\varepsilon > 0$ there exists a constant $\tilde{C}_\omega > 0$ independent of ε such that for all $u \in \mathcal{X}_\varepsilon^{k+1, \gamma}$ and $t \in \mathbf{R}$, we have*

$$\|u\|_{C_{\varepsilon,0}^{k+1, \gamma}} + |t| \leq \tilde{C}_\omega \varepsilon^{-\omega} \|\mathfrak{Q}_{P_\varepsilon} u + t f_\varepsilon\|_{C_{\varepsilon,-1}^{k,\gamma}}.$$

Proof. Let f_ε^\pm be the restrictions of f_ε over $P_{\varepsilon,C}^\pm$. Then similar to the proof of Proposition 4.17 we obtain that

$$\|f_\varepsilon^- - f_P\|_{C_{P_{\varepsilon,C}^-}^{k,Y}} \lesssim \varepsilon^{c_q} \quad \text{and} \quad \|f_\varepsilon^+\|_{C_{P_{\varepsilon,C}^+}^{k,Y}} \lesssim \varepsilon^{c_q}.$$

Claim 1: For all $u \in \mathcal{X}_\varepsilon^{k+1,Y}$ with $\|u\|_{C_{\varepsilon,0}^{k+1,Y}} \leq 1$ we have

$$1 \leq \tilde{C}_\omega \varepsilon^{-\omega} \|\mathfrak{Q}_{P_\varepsilon} u - f_\varepsilon\|_{C_{\varepsilon,-1}^{k,Y}}.$$

We will prove this by contradiction. Suppose it is not true. Then there exist sequences $\varepsilon_n \rightarrow 0$ and $u_n \in \mathcal{X}_{\varepsilon_n}^{k+1,Y}$ such that $\|u_n\|_{C_{\varepsilon_n,0}^{k+1,Y}} \leq 1$ and $\varepsilon_n^{-\omega} \|\mathfrak{Q}_{P_{\varepsilon_n}} u_n - f_{\varepsilon_n}\|_{C_{\varepsilon_n,-1}^{k,Y}}$ converges to 0 as $n \rightarrow \infty$. After defining $u_{n,P}$ and $u_{n,L}$ as in Proposition 4.20 a similar argument as in there yields a smooth section $u_P \in C_{P,0}^{k+1,Y}$ and $u_L \in C_{L,0}^{k+1,Y}$ such that

$$\mathfrak{Q}_P u_P = f_P, \quad \|u_{n,P} - u_P\|_{C_{P,\text{loc}}^{k+1,\frac{Y}{2}}} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

and $\mathfrak{Q}_L u_L = 0$, $\|u_{n,L} - u_L\|_{C_{L,\text{loc}}^{k+1,\frac{Y}{2}}} \rightarrow 0$ as $n \rightarrow \infty$. Moreover, we can prove as in Proposition 4.20 the following.

- (i) $\|u_{n,P} - u_P\|_{C_{P_{\varepsilon_n,C}^0}^{k+1,Y}} \rightarrow 0$ as $n \rightarrow \infty$,
- (ii) $\|u_{n,L} - u_L\|_{C_{L_{\varepsilon_n^{-1}\delta_n,C}^0}^{k+1,Y}} \rightarrow 0$ as $n \rightarrow \infty$ where $L_{\varepsilon_n^{-1}\delta_n,C} := s_{\varepsilon_n}^{-1} \Upsilon^{-1} P_{\varepsilon_n,C}^+ \subset L_C$,
- (iii) $i_P u_P = i_L u_L$. This contradicts to the assumption.

To prove (i), we fix a real number p with $1 < p < \frac{1}{q}$. For sufficiently small $s > 0$ we have

$$\begin{aligned} & \|\mathfrak{Q}_P(1 - \rho_{\delta_n^p})(u_n - u_P)\|_{C_{P,-s-1}^{k,Y}} \\ & \leq \varepsilon_n^{c_{pq}} + \varepsilon_n^{c_q} + \|\mathfrak{Q}_{P_{\varepsilon_n}}(1 - \rho_{\delta_n^p})u_n - \mathfrak{Q}_P(1 - \rho_{\delta_n^p})u_P\|_{C_{\varepsilon_n,-s-1}^{k,Y}} \\ & \leq \varepsilon_n^{c_{pq}} + \varepsilon_n^{c_q} + \delta_n^{ps} + \|(1 - \rho_{\delta_n^p})(\mathfrak{Q}_{P_{\varepsilon_n}} u_n - f_P)\|_{C_{\varepsilon_n,-s-1}^{k,Y}} \\ & \leq \varepsilon_n^{c_{pq}} + \varepsilon_n^{c_q} + \delta_n^{ps} + \|\mathfrak{Q}_{P_{\varepsilon_n}} u_n - f_{\varepsilon_n}\|_{C_{\varepsilon_n,-1}^{k,Y}} + \|(1 - \rho_{\delta_n^p})(f_{\varepsilon_n} - f_P)\|_{C_{\varepsilon_n,-1}^{k,Y}} \\ & \lesssim \varepsilon_n^{c_{pq}} + \varepsilon_n^{c_q} + \delta_n^{ps} + \varepsilon_n^\omega. \end{aligned}$$

Now (i) follows as in (i) of Proposition 4.20. Since $\delta_n^{\frac{s}{2}} \|\rho_{\frac{1}{\delta_n}} f_{\varepsilon_n}\|_{C_{\varepsilon_n,-1}^{k,Y}} \rightarrow 0$ as $n \rightarrow \infty$, (ii) follows from (ii) of Proposition 4.20. Again the proof of (iii) is also similar to (iii) of Proposition 4.20.

Claim 2: For all $u \in \mathcal{X}_\varepsilon^{k+1,Y}$ and $t \in \mathbf{R}$ with $\|u\|_{C_{\varepsilon,0}^{k+1,Y}} = 1$ and $|t| \leq 1$ we have

$$1 \leq \tilde{C}_\omega \varepsilon^{-\omega} \|\mathfrak{Q}_{P_\varepsilon} u - t f_\varepsilon\|_{C_{\varepsilon,-1}^{k,Y}}.$$

Suppose it is not true. Then there exist sequences $\varepsilon_n \rightarrow 0$ and $u_n \in \mathcal{X}_{\varepsilon_n}^{k+1,Y}$, $t_n \in \mathbf{R}$ such that $\|u_n\|_{C_{\varepsilon_n,0}^{k+1,Y}} = 1$, $t_n \rightarrow t_\infty$ and $\varepsilon_n^{-\omega} \|\mathfrak{Q}_{P_{\varepsilon_n}} u_n - t_n f_{\varepsilon_n}\|_{C_{\varepsilon_n,-1}^{k,Y}}$ converges to 0 as $n \rightarrow \infty$. If $t_\infty \neq 0$ then

similar arguments to Claim 1 replacing f_{ε_n} by $t_n f_{\varepsilon_n}$ lead to a contradiction. If $t_\infty = 0$ then similar arguments in Proposition 4.20 leads to again a contradiction.

Observe that Claim 1 and 2 are enough to prove the proposition. ■

4.3 Proof of the desingularization theorem

To prove the desingularization theorem we solve the nonlinear PDE, $\tilde{\mathfrak{F}}_\varepsilon(u, t) = 0$ with (u, t) is very close to $(\alpha_\varepsilon, 0)$. Indeed, we use the following Lemma 4.23 which can be proved similarly to [Ber22, Lemma 4.16].

Lemma 4.23. *Let \mathcal{X}, \mathcal{Y} be two Banach spaces and $\mathcal{X}_1 \subset \mathcal{X}$ be a Banach subspace. Let $V \subset \mathcal{X}$ be a neighbourhood of $0 \in \mathcal{X}$. Let $x_0 \in V$. Let $F : V \rightarrow \mathcal{Y}$ be a smooth map of the form¹⁴*

$$F(x) = L(x) + Q(x) + F(x_0)$$

such that:

- $L : \mathcal{X}_1 \rightarrow \mathcal{Y}$ is an invertible operator and there exists a constant $c_L > 0$ such that for all $x \in \mathcal{X}_1$, $\|x\|_{\mathcal{X}} \leq c_L \|Lx\|_{\mathcal{Y}}$.
- $Q : V \rightarrow \mathcal{Y}$ is a smooth map and there exists a constant $c_Q > 0$ such that for all $x_1, x_2 \in V$,

$$\|Q(x_1) - Q(x_2)\|_{\mathcal{Y}} \leq c_Q \|x_1 - x_2\|_{\mathcal{X}} (\|x_1 - x_0\|_{\mathcal{X}} + \|x_2 - x_0\|_{\mathcal{X}}).$$

If $\|F(x_0)\|_{\mathcal{Y}} \leq \frac{1}{10c_L^2 c_Q}$ and $B(x_0, \frac{1}{5c_L c_Q}) \subset V$, then there exists an unique $x \in x_0 + \mathcal{X}_1$ with

$$\|x - x_0\|_{\mathcal{X}} \leq c_L \|F(x_0)\|_{\mathcal{Y}}, \quad F(x) = 0.$$

Proof of Theorem 1.16. Let α_ε be α_ε^1 as in Definition 4.4. Let T_ε be as in Proposition 4.18. The map

$$\tilde{\mathfrak{F}}_\varepsilon : C_{\varepsilon,0}^{k+1,\mathcal{Y}}(V_{P_\varepsilon,C}) \times (t_0 - T_\varepsilon, t_0 + T_\varepsilon) \rightarrow C_{\varepsilon,-1}^{k,\mathcal{Y}}$$

can be written as $\tilde{\mathfrak{F}}_\varepsilon(u, t_0 + t) = \mathfrak{L}_{P_\varepsilon} u + t f_\varepsilon + \tilde{Q}_\varepsilon(u, t_0 + t) + \mathfrak{F}_\varepsilon(\alpha_\varepsilon)$. Since the matching kernel \mathcal{K}^m is one dimensional and the index of $\mathfrak{L}_{P_\varepsilon}$ is zero, by Proposition 4.20 and Proposition 4.22 we obtain that

$$\tilde{L}_\varepsilon : \mathcal{X}_\varepsilon^{k+1,\mathcal{Y}} \oplus \mathbb{R} \rightarrow C_{\varepsilon,-1}^{k,\mathcal{Y}}$$

defined by $\tilde{L}_\varepsilon(u, t) := \mathfrak{L}_{P_\varepsilon} u + t f_\varepsilon$, is invertible and

$$\|u\|_{C_{\varepsilon,0}^{k+1,\mathcal{Y}}} + |t| \leq C_{\tilde{L}_\varepsilon} \|\tilde{L}_\varepsilon(u, t)\|_{C_{\varepsilon,-1}^{k,\mathcal{Y}}} \quad \text{where } C_{\tilde{L}_\varepsilon} = O(\varepsilon^{-\omega}).$$

Again by Proposition 4.18 if $t_1, t_2 \in (-T_\varepsilon, T_\varepsilon)$, then we have

$$\begin{aligned} & \|\tilde{Q}_\varepsilon(u, t_0 + t_1) - \tilde{Q}_\varepsilon(v, t_0 + t_2)\|_{C_{\varepsilon,-1}^{k,\mathcal{Y}}} \\ & \leq C_{Q_\varepsilon} \left(\|u - v\|_{C_{\varepsilon,0}^{k+1,\mathcal{Y}}} + |t_1 - t_2| \right) \left(\|u - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,\mathcal{Y}}} + \|v - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,\mathcal{Y}}} + |t_1| + |t_2| \right), \end{aligned}$$

¹⁴Note that $Q(x_0) = -L(x_0)$.

where $C_{Q_\varepsilon} := O(\varepsilon^{-1})$. Since $\nu < 0$, by Proposition 4.17 we have,

$$\|\tilde{\mathfrak{F}}_\varepsilon(\alpha_\varepsilon)\|_{C_{\varepsilon,-1}^{k,Y}} \leq \frac{1}{10C_{L_\varepsilon}^2 C_{Q_\varepsilon}} = O(\varepsilon^{1+2\omega})$$

for sufficiently small ω . Hence by Lemma 4.23, we have for all sufficiently small $\varepsilon > 0$, there exist $t(\varepsilon) \in (t_0 - T, t_0 + T)$ and smooth closed associative submanifold $\tilde{P}_{\varepsilon,t(\varepsilon)} := Y_{P_{\varepsilon,t(\varepsilon)},C}(\tilde{\alpha}_\varepsilon)$ in $(Y, \phi_{t(\varepsilon)})$, that is $\tilde{\mathfrak{F}}_\varepsilon(\tilde{\alpha}_\varepsilon, t(\varepsilon)) = 0$. Moreover, if $\frac{1}{\mu} < q \leq \frac{1-\nu}{\mu-\nu}$ then

$$\|\tilde{\alpha}_\varepsilon - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,Y}} + |t(\varepsilon) - t_0| \lesssim \varepsilon^{q\mu-\omega}.$$

Finally we see that $\tilde{P}_{\varepsilon,t(\varepsilon)} \rightarrow P$ in the sense of currents as $\varepsilon \rightarrow 0$. Indeed, for any 3-form $\xi \in \Omega^3(Y)$ we have

$$\left| \int_{\tilde{P}_{\varepsilon,t(\varepsilon)}} \xi - \int_P \xi \right| \leq \left| \int_{\tilde{P}_{\varepsilon,t(\varepsilon)}} \xi - \int_{P_{\varepsilon,t(\varepsilon)}} \xi \right| + \left| \int_{P_{\varepsilon,t(\varepsilon)}} \xi - \int_P \xi \right| \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad \blacksquare$$

Remark 4.24. Observe that if there exists $\lambda_0 < \nu$ satisfying (i) and (ii) of Definition 4.4 then we can consider $\alpha_\varepsilon = \alpha_\varepsilon^2$ in the proof of Theorem 1.16. Note also that for both cases in the proof of Theorem 1.16 $(\tilde{\alpha}_\varepsilon, t(\varepsilon))$ satisfies the following estimates.

$$(4.25) \quad \begin{aligned} & \|\tilde{\alpha}_\varepsilon - \alpha_\varepsilon\|_{C_{\varepsilon,0}^{k+1,Y}} + |t(\varepsilon) - t_0| \\ & \lesssim \begin{cases} \varepsilon^{q\mu-\omega}, & \text{if } \alpha_\varepsilon = \alpha_\varepsilon^1 \text{ and } \frac{1}{\mu} < q \leq \frac{1-\nu}{\mu-\nu} \\ \varepsilon^{q\mu-\omega} + \varepsilon^{1+(q-1)\nu_0-\omega}, & \text{if } \alpha_\varepsilon = \alpha_\varepsilon^2 \text{ and } \frac{\nu-\nu_0}{\nu+\mu-1+s_0-\nu_0} \leq q < 1. \end{cases} \end{aligned}$$

Indeed these estimates follows from Lemma 4.23, Proposition 4.22 and Proposition 4.17. \clubsuit

4.4 Desingularizations for CS associatives with Harvey-Lawson T^2 -cone singularity

Lemma 4.26. *Let $\phi \in \mathcal{P}_{\text{cs}}^{\text{reg}}$ be a path of co-closed G_2 -structures and $P \in \mathcal{M}_{\text{cs}}^\phi$ be a CS associative with singularity at only one point. Let $L \subset \mathbf{R}^7$ be an AC associative with the same asymptotic cone C of P . If $d_{-1} = 2$, then*

$$\tilde{\mathcal{K}}^m = \mathcal{K}^m \cong \ker \mathfrak{L}_{L,-1}.$$

Moreover, $\text{s-ind}(C) = 1$, $\dim \ker \mathfrak{L}_{P,-1} = 1$ and $\ker \mathfrak{L}_{P,-1} = \langle v_P \rangle_{\mathbf{R}}$ such that $\langle v_P, f_P \rangle_{L^2} \neq 0$.

Proof. Since $\phi \in \mathcal{P}_{\text{cs}}^{\text{reg}}$, for any CS associative P and any deformation family \mathcal{Z} of C the operator

$$\bar{\mathfrak{Q}}_{P,\mu,\mathcal{Z}}^{k,Y} : \mathbf{R} \oplus C_{P,\mu}^{k+1,Y} \oplus \mathbf{R}^7 \oplus T_{\Sigma}\mathcal{Z} \rightarrow C_{P,\mu-1}^{k,Y}$$

defined in Definition 3.100 is surjective. Therefore by Lemma 3.70 we obtain that

$$\bar{\mathfrak{Q}}_{P,-1+s}^{k,Y} : \mathbf{R} \oplus C_{P,-1+s}^{k+1,Y} \rightarrow C_{P,-2+s}^{k,Y}$$

defined by $\bar{\mathfrak{Q}}_{P,-1+s}^{k,Y}(t, u) = \mathfrak{L}_P u + t f_P$, is surjective for all sufficiently small $s > 0$. Theorem 1.11 implies that the stability index of the asymptotic cone C satisfies $\text{s-ind}(C) \leq 1$. Since $d_{-1} = 2$,

therefore $V_\lambda = 0$ for all non-zero $\lambda \in (-1, 1)$ and $\text{s-ind}(C) = 1$. By Proposition 3.72 we have index $\mathfrak{L}_{P, -1+s} = -1$. Hence, $\ker \mathfrak{L}_{P, -1} = \text{coker } \mathfrak{L}_{P, -1+s} = \langle v_P \rangle$ is one dimensional satisfying $\langle v_P, f_P \rangle_{L^2} \neq 0$ and $\ker \mathfrak{L}_{P, -1+s} = 0$. This implies there does not exist any $u_P \in C_{P,0}^{k+1,Y}$ such that $\mathfrak{L}_P u_P = f_P$. Indeed, Proposition 3.66 would imply

$$0 = \langle \mathfrak{L}_P v_P, u_P \rangle_{L^2} = \langle v_P, \mathfrak{L}_P u_P \rangle_{L^2} = \langle v_P, f_P \rangle_{L^2} \neq 0.$$

Hence, $\tilde{\mathcal{K}}^m = \mathcal{K}^m$. Also we have $\ker \mathfrak{L}_{P,0} \subset \ker \mathfrak{L}_{P, -1+s} = \{0\}$ and therefore again by Lemma 3.70 we obtain that the matching kernel $\mathcal{K}^m \cong \ker \mathfrak{L}_{L, -1}$. \blacksquare

Proof of Theorem 1.21. We have seen in Example 3.34 that the Harvey-Lawson cone has $d_{-1} = 2$. Therefore by Lemma 4.26 and Proposition 3.73 we obtain that for each Harvey-Lawson AC special Lagrangian $L^i := L_1^i$, $i = 1, 2, 3$ (see Example 3.47) the extended matching kernel and the matching kernel are one dimensional. Indeed,

$$\dim \tilde{\mathcal{K}}^m = \dim \mathcal{K}^m = \dim \ker \mathfrak{L}_{L, -1}^i = b^1(S^1 \times \mathbb{C}) + b^0(T^2) - 1 = 1.$$

By Theorem 1.16 we obtain that for sufficiently small $\varepsilon > 0$, there exists $t^i(\varepsilon)$ and smooth closed associative $\tilde{P}_{\varepsilon, t^i(\varepsilon)}$ in $(Y, \phi_{t^i(\varepsilon)})$ such that $\tilde{P}_{\varepsilon, t^i(\varepsilon)} \rightarrow P$ in the sense of currents as $\varepsilon \rightarrow 0$. $\tilde{P}_{\varepsilon, t^i(\varepsilon)}$ is diffeomorphic to $P_{\varepsilon, t^i(\varepsilon)}$. Since L^i are obtained by Dehn filling of $C_{HL}^o := C_{HL} \setminus B(0, 1)$ along simple closed curves μ_i as mentioned in the theorem, therefore it is diffeomorphic to the Dehn filling of P^o as required in the theorem (this was observed in [DW19, Remark 3.6]).

It remains to prove that the leading order term of $t^i(\varepsilon) - t_0$ is of the required form if $\phi \in \mathcal{P}^\bullet$. Let $v_P \in \ker \mathfrak{L}_{P, -1-s}$ be as in Lemma 4.26. Then

$$i_{P, -1} \hat{v}_P = b_1 \xi_1 + b_2 \xi_2, \quad b_1 \neq 0, b_2 \neq 0, b_1 \neq b_2 \text{ where } \xi_1, \xi_2 \text{ are in (3.48) of Example 3.47.}$$

We will prove for $t(\varepsilon) = t^1(\varepsilon)$ only as the others follow from similar arguments. We use the notation $\alpha_\varepsilon = \alpha_\varepsilon^1 = \rho_\delta \Upsilon_*(s_{\varepsilon*} \beta) + (1 - \rho_\delta) \alpha$ and $(\tilde{\alpha}_\varepsilon, t(\varepsilon))$ from Theorem 1.16, which satisfies

$$\tilde{\mathfrak{F}}_\varepsilon(\tilde{\alpha}_\varepsilon, t(\varepsilon)) = \mathfrak{L}_{P_\varepsilon}(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon) + (t(\varepsilon) - t_0) f_\varepsilon + Q_\varepsilon(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon, t(\varepsilon) - t_0) + \mathfrak{F}_\varepsilon(\alpha_\varepsilon) = 0.$$

Define $\mathring{P}_\varepsilon := P_C \setminus Y(B(0, 2\varepsilon R_\infty))$ and $\mathring{P}_\delta := P_C \setminus Y(B(0, \delta))$. Over \mathring{P}_δ ,

$$\mathfrak{F}_\varepsilon(\alpha_\varepsilon) = \mathfrak{L}_P(\alpha_\varepsilon - \alpha) + Q_P(\alpha_\varepsilon - \alpha).$$

Therefore,

$$(4.27) \quad 0 = \langle \tilde{\mathfrak{F}}_\varepsilon(\tilde{\alpha}_\varepsilon, t(\varepsilon)), v_P \rangle_{L^2_{\mathring{P}_\varepsilon}} = (t(\varepsilon) - t_0) \langle f_\varepsilon, v_P \rangle_{L^2_{\mathring{P}_\varepsilon}} + \langle \mathfrak{L}_P \alpha_\varepsilon, v_P \rangle_{L^2_{\mathring{P}_\delta}} + e_\varepsilon^1 + e_\varepsilon^2 + e_\varepsilon^3 + e_\varepsilon^4 + e_\varepsilon^5$$

where $e_\varepsilon^1 := \langle \mathfrak{L}_{P_\varepsilon}(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon), v_P \rangle_{L^2_{\mathring{P}_\varepsilon}}$, $e_\varepsilon^2 := \langle Q_\varepsilon(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon, t(\varepsilon) - t_0), v_P \rangle_{L^2_{\mathring{P}_\varepsilon}}$, and

$$e_\varepsilon^3 := \langle Q_P(\alpha_\varepsilon - \alpha), v_P \rangle_{L^2_{\mathring{P}_\delta}}, \quad e_\varepsilon^4 = -\langle \mathfrak{L}_P(\alpha), v_P \rangle_{L^2_{\mathring{P}_\varepsilon}} + \langle \mathfrak{L}_P(\alpha), v_P \rangle_{L^2_{\mathring{P}_\delta \setminus \mathring{P}_\varepsilon}}, \quad e_\varepsilon^5 = \langle \mathfrak{F}_\varepsilon(\alpha_\varepsilon), v_P \rangle_{L^2_{\mathring{P}_\varepsilon \setminus \mathring{P}_\delta}}.$$

By Proposition 3.88 we obtain that

$$(4.28) \quad \langle \mathfrak{L}_P \alpha_\varepsilon, v_P \rangle_{L^2_{\mathring{P}_\delta}} = \int_{\partial \mathring{P}_\delta} \iota_{\alpha_\varepsilon} \iota_{v_P} \Upsilon_{P_C}^* \psi = \int_{\partial \mathring{P}_\delta} \iota_{s_{\varepsilon*} \beta} \iota_{b_1 \xi_1 + b_2 \xi_2} \Upsilon^* \psi(0) + O(\varepsilon^{2+q}).$$

Now,

$$(4.29) \quad \int_{\partial \mathring{P}_\delta} \iota_{s_\varepsilon * \beta} \iota_{b_1 \xi_1 + b_2 \xi_2} \Upsilon^* \psi(0) = \int_{\partial B(0, \delta) \cap C} \iota_{s_\varepsilon * \xi_1} \iota_{b_1 \xi_1 + b_2 \xi_2} \Upsilon^* \psi(0) + O(\varepsilon^{3-q}),$$

and

$$(4.30) \quad \int_{\partial B(0, \delta) \cap C} \iota_{s_\varepsilon * \xi_1} \iota_{b_1 \xi_1 + b_2 \xi_2} \Upsilon^* \psi(0) = \varepsilon^2 \int_{\Sigma} \langle \partial_r \times \xi_1, b_1 \xi_1 + b_2 \xi_2 \rangle = b_2 \varepsilon^2 \int_{\Sigma} \langle \partial_r \times \xi_1, \xi_2 \rangle.$$

We will now estimate the remaining terms in (4.27). Using (4.25) we compute

$$\begin{aligned} |e_\varepsilon^1| &= |\langle \mathfrak{Q}_{P_\varepsilon}(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon) - \mathfrak{Q}_P(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon), v_P \rangle_{L^2_{P_\varepsilon}}| + |\langle \mathfrak{Q}_P(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon), v_P \rangle_{L^2_{P_\varepsilon}}| \\ &\lesssim \varepsilon^{2+q\mu-\omega} + \int_{\partial \mathring{P}_\varepsilon} |\iota_{\tilde{\alpha}_\varepsilon - \alpha_\varepsilon} \iota_{v_P} \Upsilon_{P_C}^* \psi| \lesssim \varepsilon^{1+q\mu-\omega} \end{aligned}$$

and

$$|e_\varepsilon^2| \lesssim \varepsilon^{2q\mu-2\omega} |\log \varepsilon| + \varepsilon^{q\mu-\omega} |t(\varepsilon) - t_0| + |t(\varepsilon) - t_0|^2 |\log \varepsilon| \lesssim \varepsilon^{2q\mu-2\omega} |\log \varepsilon|.$$

We can further compute

$$|e_\varepsilon^3| \lesssim \varepsilon^{2q\mu}, \quad |e_\varepsilon^5| \lesssim \varepsilon^{3q}$$

and

$$|e_\varepsilon^4| \lesssim \int_{\partial \mathring{P}_\varepsilon} |\iota_\alpha \iota_{v_P} \Upsilon_{P_C}^* \psi| + \varepsilon^{2q\mu} \lesssim \varepsilon^{1+\mu} + \varepsilon^{2q\mu}.$$

Hence by choosing ω sufficiently small and $q > \frac{2}{3}$, we obtain

$$(4.31) \quad |e_\varepsilon^1| + |e_\varepsilon^2| + |e_\varepsilon^3| + |e_\varepsilon^4| + |e_\varepsilon^5| = o(\varepsilon^2).$$

Since $\langle f_\varepsilon, v_P \rangle_{L^2_{P_\varepsilon}} = \langle f_P, v_P \rangle_{L^2} + o(1)$, therefore combining (4.27), (4.28), (4.29), (4.30) and (4.31) we obtain

$$t^1(\varepsilon) = t(\varepsilon) = t_0 - \frac{cb_2}{\langle f_P, v_P \rangle_{L^2}} \varepsilon^2 + o(\varepsilon^2),$$

where $c = \int_{\Sigma} \langle J \xi_1, \xi_2 \rangle \neq 0$. Similarly we can prove for $t^2(\varepsilon)$ and $t^3(\varepsilon)$. Only thing to notice here is that $\xi_3 = -\xi_1 - \xi_2$ as in Example 3.47. \blacksquare

4.5 Desingularizations for associative submanifolds with transverse intersection

Let $P \in \mathcal{M}_{cs}^\phi$ be an associative submanifold in (Y, ϕ_{t_0}) with a transverse unique self intersection. In other words, P is a conically singular associative with singularity at a unique point, which is modeled on a union of two transverse associative planes.

Proof of Theorem 1.29. In Example 3.33 we see that the union of two transverse associative planes has $d_{-1} = 0$ and $V_\lambda = \{0\}$ for all $\lambda \in (-1, 0)$. If $\phi \in \mathcal{P}^\dagger$ (see Definition 1.24), Proposition 3.73 implies that for the special Lagrangian Lawlor neck L (see Example 3.44) the extended matching kernel and the matching kernel are one dimensional. Indeed,

$$\dim \mathcal{K}^m = \dim \ker \mathfrak{Q}_{L, -1} = b^1(S^2 \times \mathbf{R}) + b^0(S^2 \amalg S^2) - 1 = 1.$$

By Theorem 1.16 we obtain that for sufficiently small $\varepsilon > 0$, there exists $t(\varepsilon) \in (t_0 - T_\varepsilon, t_0 + T_\varepsilon)$ and smooth closed associative $\tilde{P}_{\varepsilon, t(\varepsilon)}$ in $(Y, \phi_{t(\varepsilon)})$ such that $\tilde{P}_{\varepsilon, t(\varepsilon)} \rightarrow P$ in the sense of currents as $\varepsilon \rightarrow 0$. $\tilde{P}_{\varepsilon, t(\varepsilon)}$ are diffeomorphic to $P_{\varepsilon, t(\varepsilon)}$ and hence are diffeomorphic to the connected sums as mentioned in the theorem. \blacksquare

Remark 4.32 (Discussion on leading order term). It remains to prove that the leading order term of $t(\varepsilon) - t_0$ is of the required form if $\phi \in \mathcal{P}^\sharp$ (see Definition 1.27). We can consider

$$\alpha_\varepsilon = \alpha_\varepsilon^2 := \rho_\delta \Upsilon_*(s_{\varepsilon*} \beta) + (1 - \rho_\delta)(\varepsilon^3 u_P + \alpha).$$

Here α and β represents P and L respectively and $u_P := (\Theta_P^C)^{-1} \hat{u}_P \in \ker \mathfrak{L}_{P, -2}$ with

$$\beta_1^\pm = B\xi^\pm, l = 1, \lambda_1 = \nu = -2, \lambda_0 = -4, \mu = 2, s_0 = 2, q = \frac{2}{5}.$$

By abusing notation we denote $B\xi^\pm$ by simply ξ^\pm . We use the notation $(\tilde{\alpha}_\varepsilon, t(\varepsilon))$ from Theorem 1.16 following Remark 4.24, which satisfies

$$(4.33) \quad \tilde{\mathfrak{F}}_\varepsilon(\tilde{\alpha}_\varepsilon, t(\varepsilon) - t_0) = \mathfrak{L}_{P_\varepsilon}(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon) + (t(\varepsilon) - t_0)f_\varepsilon + Q_\varepsilon(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon, t(\varepsilon)) + \mathfrak{F}_\varepsilon(\alpha_\varepsilon) = 0.$$

Define $\mathring{P}_\varepsilon := P_C \setminus \Upsilon(B(0, 2\varepsilon R_\infty))$ and $\mathring{P}_\delta := P_C \setminus \Upsilon(B(0, \delta))$. Over \mathring{P}_δ ,

$$\mathfrak{F}_\varepsilon(\alpha_\varepsilon) = \mathfrak{L}_P(\alpha_\varepsilon - \alpha) + Q_P(\alpha_\varepsilon - \alpha).$$

Therefore,

$$(4.34) \quad 0 = \langle \tilde{\mathfrak{F}}_\varepsilon(\tilde{\alpha}_\varepsilon, t(\varepsilon)), u_P \rangle_{L^2_{\mathring{P}_\varepsilon}} = (t(\varepsilon) - t_0) \langle f_\varepsilon, u_P \rangle_{L^2_{\mathring{P}_\varepsilon}} + \langle \mathfrak{L}_P \alpha_\varepsilon, u_P \rangle_{L^2_{\mathring{P}_\delta}} + e_\varepsilon^1 + e_\varepsilon^2 + e_\varepsilon^3 + e_\varepsilon^4 + e_\varepsilon^5$$

where $e_\varepsilon^1 := \langle \mathfrak{L}_{P_\varepsilon}(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon), u_P \rangle_{L^2_{\mathring{P}_\varepsilon}}$, $e_\varepsilon^2 := \langle Q_\varepsilon(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon, t(\varepsilon) - t_0), u_P \rangle_{L^2_{\mathring{P}_\varepsilon}}$, and

$$e_\varepsilon^3 := \langle Q_P(\alpha_\varepsilon - \alpha), u_P \rangle_{L^2_{\mathring{P}_\delta}}, \quad e_\varepsilon^4 = -\langle \mathfrak{L}_P(\alpha), u_P \rangle_{L^2_{\mathring{P}_\delta}} + \langle \mathfrak{L}_P(\alpha), u_P \rangle_{L^2_{\mathring{P}_\varepsilon \setminus \mathring{P}_\delta}}, \quad e_\varepsilon^5 = \langle \mathfrak{F}_\varepsilon(\alpha_\varepsilon), u_P \rangle_{L^2_{\mathring{P}_\varepsilon \setminus \mathring{P}_\delta}}.$$

By Proposition 3.88 we obtain that $\langle \mathfrak{L}_P \alpha_\varepsilon, u_P \rangle_{L^2_{\mathring{P}_\delta}}$ is

$$(4.35) \quad \int_{\partial \mathring{P}_\delta} \iota_{\alpha_\varepsilon} \iota_{u_P} \Upsilon_{P_C}^* \psi = \varepsilon^3 \int_{\partial B_{\Pi^+}(0, \delta)} \iota_{\xi^+} \iota_{u_P - \xi^+} \Upsilon^* \psi - \varepsilon^3 \int_{\partial B_{\Pi^-}(0, \delta)} \iota_{\xi^-} \iota_{u_P - \xi^-} \Upsilon^* \psi + o(\varepsilon^3).$$

Now,

$$(4.36) \quad \int_{\partial B_{\Pi^\pm}(0, \delta)} \iota_{\xi^\pm} \iota_{u_P - \xi^\pm} \Upsilon^* \psi = \int_{\partial B_{\Pi^\pm}(0, \delta)} \iota_{\xi^\pm} \iota_{u_P - \xi^\pm} \psi_e + o(\varepsilon)$$

and

$$(4.37) \quad \int_{\partial B_{\Pi^\pm}(0, \delta)} \iota_{\xi^\pm} \iota_{u_P - \xi^\pm} \psi_e = (u_P - \xi^\pm)(0) \cdot \mathbf{n} + o(\varepsilon).$$

Define $v_P = (\Theta_P^C)^{-1} \hat{v}_P$. Then $\mathfrak{L}_P v_P = f_P$. We observe $\langle f_\varepsilon, u_P \rangle_{L^2_{\hat{P}_\varepsilon}} = \langle f_P, u_P \rangle_{L^2_{\hat{P}_\delta}} + o(1)$. Therefore, similar to (4.35), (4.36), (4.37) we obtain

$$(4.38) \quad \begin{aligned} \langle f_P, u_P \rangle_{L^2_{\hat{P}_\delta}} &= \int_{\partial B_{\Pi^+}(0, \delta)} \iota_{v_P} \iota_{\xi^+} \psi_e - \int_{\partial B_{\Pi^-}(0, \delta)} \iota_{v_P} \iota_{\xi^-} \psi_e + o(1) \\ &= (v_P^+(0) - v_P^-(0)) \cdot \mathbf{n} + o(1). \end{aligned}$$

We can further compute

$$(4.39) \quad |e_\varepsilon^3| + |e_\varepsilon^4| + |e_\varepsilon^5| = o(\varepsilon^3).$$

If $|e_\varepsilon^1| + |e_\varepsilon^2| = o(\varepsilon^3)$ then we get the leading order expression as in (1.31). However it is **not** clear from our desingularization theorem why it should hold. With the help of (4.25), we can only express that:

$$\begin{aligned} |e_\varepsilon^1| &= |\langle \mathfrak{L}_{P_\varepsilon}(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon) - \mathfrak{L}_P(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon), u_P \rangle_{L^2_{\hat{P}_\varepsilon}}| + |\langle \mathfrak{L}_P(\tilde{\alpha}_\varepsilon - \alpha_\varepsilon), u_P \rangle_{L^2_{\hat{P}_\varepsilon}}| \\ &\lesssim o(\varepsilon^3) + \left| \int_{\partial B_{\Pi^+}(0, 2\varepsilon R_\infty)} \iota_{\tilde{\alpha}_\varepsilon - \alpha_\varepsilon} \iota_{\xi^+} \Upsilon^* \psi - \int_{\partial B_{\Pi^-}(0, 2\varepsilon R_\infty)} \iota_{\tilde{\alpha}_\varepsilon - \alpha_\varepsilon} \iota_{\xi^-} \Upsilon^* \psi \right|. \quad \clubsuit \end{aligned}$$

A Appendix

Lemma A.1. *Let M be a 3-dimensional oriented submanifold of an almost G_2 -manifold (Y, ϕ) . Let $\Upsilon_M : V_M \subset NM \rightarrow Y$ be a tubular neighbourhood map. There exists a constant $C > 0$ such that if $w \in C^\infty(NM)$ and $u, v, s \in C^\infty(V_M)$ then over $\Gamma_s := \text{graph } s \subset V_M$ we have¹⁵*

$$|\iota_w \mathcal{L}_u \mathcal{L}_v \psi| \lesssim |w| \left(f_1 |u| |v| + f_2 (|u| |\nabla v| + |v| |\nabla u|) + |\nabla u| |\nabla v| |\psi| \right),$$

with

$$\begin{aligned} f_1 &:= |\psi| |\nabla B| + |\psi| |B|^2 + |B| |\nabla \psi| + |\nabla^2 \psi| + |R| |\psi| \\ f_2 &:= |\nabla \psi| + |B| |\psi|. \end{aligned}$$

Here R is the Riemann curvature tensor and $B : C^\infty(NM) \times C^\infty(NM) \rightarrow C^\infty(TV_M)$ is defined by $B(u, v) := \nabla_u v$

Proof. Let $\{e_1, e_2, e_3\}$ be a local orthonormal frame for $T\Gamma_s$. For a torsion free connection we have

$$\mathcal{L}_v \psi(w, e_1, e_2, e_3) = \nabla_v \psi(w, e_1, e_2, e_3) + \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \psi(w, \nabla_{e_1} v, e_2, e_3) + \psi(\nabla_w v, e_1, e_2, e_3).$$

¹⁵By abusing notation we denote again by u the extension of the vector field u over NM by translating fiberwise. We also denote $\Upsilon_M^* \psi$ by ψ . The notation \mathfrak{L} stands for the Lie derivative.

And also by definition we have

$$\mathcal{L}_u \mathcal{L}_v \psi(w, e_1, e_2, e_3) = u(\mathcal{L}_v \psi(w, e_1, e_2, e_3)) - \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \mathcal{L}_v \psi(w, [u, e_1], e_2, e_3).$$

Now,

$$\begin{aligned} u(\nabla_v \psi(w, e_1, e_2, e_3)) &= \nabla_u \nabla_v \psi(w, e_1, e_2, e_3) \\ &+ \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \nabla_v \psi(w, \nabla_{e_1} u + [u, e_1], e_2, e_3) + \nabla_v \psi(\nabla_w u, e_1, e_2, e_3). \end{aligned}$$

And,

$$\begin{aligned} u(\psi(w, \nabla_{e_1} v, e_2, e_3)) &= \nabla_u \psi(w, \nabla_{e_1} v, e_2, e_3) + \psi(w, \nabla_u \nabla_{e_1} v, e_2, e_3) + \psi(w, \nabla_{e_1} v, \nabla_u e_2, e_3) \\ &+ \psi(w, \nabla_{e_1} v, e_2, \nabla_u e_3) + \psi(\nabla_w u, \nabla_{e_1} v, e_2, e_3). \end{aligned}$$

Furthermore,

$$\begin{aligned} u(\psi(\nabla_w v, e_1, e_2, e_3)) &= \nabla_u \psi(\nabla_w v, e_1, e_2, e_3) + \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \psi(\nabla_w v, \nabla_u e_1, e_2, e_3) \\ &+ \psi(\nabla_w \nabla_u v + R(u, w)v, e_1, e_2, e_3). \end{aligned}$$

And

$$\begin{aligned} &\mathcal{L}_v \psi(w, [u, e_1], e_2, e_3) \\ &= \psi(w, \nabla_{[u, e_1]} v, e_2, e_3) + \psi(\nabla_w v, [u, e_1], e_2, e_3) \\ &+ \psi(w, [u, e_1], \nabla_{e_2} v, e_3) + \psi(w, [u, e_1], e_2, \nabla_{e_3} v) + \nabla_v \psi(w, [u, e_1], e_2, e_3). \end{aligned}$$

Finally, putting together after some cancellations we obtain

$$\begin{aligned} |\mathcal{L}_u \mathcal{L}_v \psi(w, e_1, e_2, e_3)| &\lesssim |w|(|\nabla_u \nabla_v \psi| + |\nabla \psi| |\nabla u| |v|) + |\nabla \psi| |v| |\nabla_w u| \\ &+ |w| |\nabla \psi| |u| |\nabla v| + |\psi| |\nabla v| |\nabla_w u| \\ &+ |\nabla_w v| (|\nabla \psi| |u| + |\psi| |\nabla u|) + |\psi| (|\nabla_w \nabla_u v| + |R| |w| |u| |v|) \\ &+ |w| |\psi| \left(\sum_{i=1}^3 |\nabla_{e_i}(\nabla_u v)| + |R| |u| |v| + |\nabla u| |\nabla v| \right). \end{aligned}$$






Observe that,






$$\begin{aligned} |\nabla_u \nabla_v \psi| &\lesssim |u| |v| (|B| |\nabla \psi| + |\nabla^2 \psi|), \\ |\nabla_{e_i}(\nabla_u v)| &\lesssim |\nabla B| |u| |v| + |B| |u| |\nabla v| + |B| |v| |\nabla u|, \\ |\nabla_w(\nabla_u v)| &\lesssim |w| |u| |v| (|\nabla B| + |B|^2), \end{aligned}$$





Therefore,

$$|\iota_w \mathcal{L}_u \mathcal{L}_v \psi| \lesssim f_1 |w| |u| |v| + f_2 |w| (|u| |\nabla^\perp v| + |v| |\nabla^\perp u|) + |w| |\nabla^\perp u| |\nabla^\perp v| |\psi|. \quad \blacksquare$$









References


- [AS88] David Adams and Leon Simon. *Rates of asymptotic convergence near isolated singularities of geometric extrema*. *Indiana Univ. Math. J.* 37.2 (1988), pp. 225–254. DOI: 10.1512/iumj.1988.37.37012.  (cit. on p. 18)
- [BM20] G. Ball and J. Madnick. *Associative Submanifolds of the Berger Space*. 2020. arXiv: 2003.13169. to appear in *Communications in Analysis and Geometry* (cit. on p. 14)
- [Ber22] Gorapada Bera. *Associative submanifolds in twisted connected sum G_2 -manifolds* (Sept. 2022). arXiv: 2209.00156.  (cit. on pp. 13, 14, 25, 39, 62)
- [BDHLMs18] A. P. Braun, M. Del Zotto, J. Halverson, M. Larfors, D. R. Morrison, and S. Schäfer-Nameki. *Infinitely many M_2 -instanton corrections to M -theory on G_2 -manifolds*. *Journal of High Energy Physics* 2018.9 (2018). DOI: 10.1007/jhep09(2018)077 (cit. on p. 14)
- [BS89] R. L. Bryant and S. M. Salamon. *On the construction of some complete metrics with exceptional holonomy*. *Duke Mathematical Journal* 58.3 (1989), pp. 829–850. DOI: 10.1215/S0012-7094-89-05839-0. MR: 1016448 (cit. on p. 13)
- [Bry87] R. L. Bryant. *Metrics with exceptional holonomy*. *Annals of Mathematics* 126.3 (1987), pp. 525–576. DOI: 10.2307/1971360. MR: 916718. Zbl: 0637.53042 (cit. on p. 13)
- [Bry82] Robert L. Bryant. *Submanifolds and special structures on the octonians*. *Journal of Differential Geometry* 17.2 (Jan. 1982). DOI: 10.4310/jdg/1214436919.  (cit. on pp. 3, 19)
- [CHNP15] A. Corti, M. Haskins, J. Nordström, and T. Pacini. *G_2 -manifolds and associative submanifolds via semi-Fano 3-folds*. *Duke Mathematical Journal* 164.10 (2015), pp. 1971–2092. DOI: 10.1215/00127094-3120743. MR: 3369307. Zbl: 06486366 (cit. on p. 13)
- [DW19] A. Doan and T. Walpuski. *On counting associative submanifolds and Seiberg–Witten monopoles*. *Pure and Applied Mathematics Quarterly* 15.4 (2019), pp. 1047–1133. DOI: 10.4310/PAMQ.2019.v15.n4.a4. arXiv: 1712.08383.  (cit. on pp. 1, 2, 16, 51, 64)
- [DW21] A. Doan and T. Walpuski. *Castelnuovo’s bound and rigidity in almost complex geometry*. *Advances in Mathematics* 379 (2021). DOI: 10.1016/j.aim.2020.107550. arXiv: 1809.04731. MR: 4199270. Zbl: 07300460.  (cit. on p. 17)
- [Dono2] S. K. Donaldson. *Floer homology groups in Yang–Mills theory*. Cambridge Tracts in Mathematics 147. With the assistance of M. Furuta and D. Kotschick. Cambridge: Cambridge University Press, 2002, pp. viii+236. DOI: 10.1017/CBO9780511543098. MR: 1883043. Zbl: 0998.53057 (cit. on p. 38)
- [DK90] S. K. Donaldson and P. B. Kronheimer. *The geometry of four-manifolds*. Oxford Mathematical Monographs. New York, 1990. MR: MR1079726. Zbl: 0904.57001 (cit. on pp. 23, 49)

- [DS11] S. K. Donaldson and E. P. Segal. *Gauge theory in higher dimensions, II. Surveys in differential geometry*. Vol. 16. Geometry of special holonomy and related topics. International Press, 2011, pp. 1–41. arXiv: 0902.3239. MR: 2893675. Zbl: 1256.53038 (cit. on pp. 12, 17)
- [DT98] S. K. Donaldson and R. P. Thomas. *Gauge theory in higher dimensions. The geometric universe*. Oxford University Press, 1998, pp. 31–47. MR: 1634503. Zbl: 0926.58003.  (cit. on p. 16)
- [DPW23] Shubham Dwivedi, Daniel Platt, and Thomas Walpuski. *Associative Submanifolds in Joyce’s Generalised Kummer Constructions. Communications in Mathematical Physics* 401.3 (Aug. 2023), pp. 2327–2353. DOI: 10.1007/s00220-023-04716-7.  (cit. on p. 13)
- [FG82] M. Fernández and A. Gray. *Riemannian manifolds with structure group G_2* . *Ann. Mat. Pura Appl. (4)* 132 (1982), 19–45 (1983). DOI: 10.1007/BF01760975. MR: 696037. Zbl: 0524.53023 (cit. on p. 12)
- [Gay14] D. Gayet. *Smooth moduli spaces of associative submanifolds. The Quarterly Journal of Mathematics* 65.4 (2014), pp. 1213–1240. DOI: 10.1093/qmath/hat042. MR: 3285769. Zbl: 1310.53049 (cit. on p. 28)
- [Gra69] A. Gray. *Vector cross products on manifolds. Transactions of the American Mathematical Society* 141 (1969), pp. 465–504. MR: 0243469. Zbl: 0182.24603 (cit. on p. 11)
- [Har90] F. R. Harvey. *Spinors and calibrations*. Vol. 9. Perspectives in Mathematics. Boston, MA: Academic Press Inc., 1990, pp. xiv+323 (cit. on pp. 10, 31)
- [HL82] R. Harvey and H. B. Lawson Jr. *Calibrated geometries. Acta Math.* 148 (1982), pp. 47–157. DOI: 10.1007/BF02392726. MR: MR666108. Zbl: 0584.53021 (cit. on pp. 13, 19, 31, 32)
- [Haso4a] Mark Haskins. *Special Lagrangian cones. Amer. J. Math.* 126.4 (2004), pp. 845–871.  (cit. on pp. 18, 27)
- [Haso4b] Mark Haskins. *The geometric complexity of special Lagrangian T^2 -cones. Invent. Math.* 157.1 (2004), pp. 11–70. DOI: 10.1007/s00222-003-0348-x.  (cit. on pp. 3, 27, 28)
- [HK07] Mark Haskins and Nikolaos Kapouleas. *Special Lagrangian cones with higher genus links. Invent. Math.* 167.2 (2007), pp. 223–294. DOI: 10.1007/s00222-006-0010-5.  (cit. on p. 18)
- [Hit74] N. Hitchin. *Harmonic spinors. Advances in Math.* 14 (1974), pp. 1–55. MR: 0358873 (cit. on p. 13)
- [Hito1] N. J. Hitchin. *Stable forms and special metrics. Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*. Vol. 288. Contemp. Math. Providence, RI: Amer. Math. Soc., 2001, pp. 70–89. arXiv: math/0107101. MR: 1871001. Zbl: 1004.53034 (cit. on p. 12)

- [Joy96a] D. D. Joyce. *Compact 8-manifolds with holonomy Spin(7)*. *Invent. Math.* 123.3 (1996), pp. 507–552. DOI: 10.1007/s002220050039. MR: MR1383960. Zbl: 0858.53037 (cit. on pp. 12, 13)
- [Joy96b] D. D. Joyce. *Compact Riemannian 7-manifolds with holonomy G_2 . II*. *Journal of Differential Geometry* 43.2 (1996), pp. 329–375. DOI: 10.4310/jdg/1214458110. MR: MR1424428. Zbl: 0861.53023 (cit. on p. 13)
- [Joy03] D. D. Joyce. *Special Lagrangian submanifolds with isolated conical singularities. V. Survey and applications*. *Journal of Differential Geometry* 63.2 (2003), pp. 279–347. MR: 2015549.  (cit. on p. 18)
- [Joy04a] D. D. Joyce. *Special Lagrangian submanifolds with isolated conical singularities. I. Regularity*. *Annals of Global Analysis and Geometry* 25.3 (2004), pp. 201–251. DOI: 10.1023/B:AGAG.0000023229.72953.57. MR: 2053761 (cit. on pp. 17, 18)
- [Joy04b] D. D. Joyce. *Special Lagrangian submanifolds with isolated conical singularities. II. Moduli spaces*. *Annals of Global Analysis and Geometry* 25.4 (2004), pp. 301–352. DOI: 10.1023/B:AGAG.0000023230.21785.8d. MR: 2054572 (cit. on pp. 5, 33)
- [Joy04c] D. D. Joyce. *Special Lagrangian submanifolds with isolated conical singularities. III. Desingularization, the unobstructed case*. *Annals of Global Analysis and Geometry* 26.1 (2004), pp. 1–58. DOI: 10.1023/B:AGAG.0000023231.31950.cc. MR: 2054578 (cit. on p. 2)
- [Joy04d] D. D. Joyce. *Special Lagrangian submanifolds with isolated conical singularities. IV. Desingularization, obstructions and families*. *Annals of Global Analysis and Geometry* 26.2 (2004), pp. 117–174. DOI: 10.1023/B:AGAG.0000031067.19776.15. MR: 2070685 (cit. on p. 2)
- [Joy18] D. D. Joyce. *Conjectures on counting associative 3-folds in G_2 -manifolds*. *Modern geometry: a celebration of the work of Simon Donaldson*. Proceedings of Symposia in Pure Mathematics 99. American Mathematical Society, 2018, pp. 97–160. DOI: 10.1090/pspum/099/01739. arXiv: 1610.09836. MR: 3838881. Zbl: 1448.53061 (cit. on pp. 1, 2, 9, 10, 12, 16, 17, 31)
- [JK17] D.D. Joyce and S. Karigiannis. *A new construction of compact torsion-free G_2 -manifolds by gluing families of Eguchi–Hanson spaces*. *Journal of Differential Geometry* 117.2 (2017), pp. 255–343. DOI: 10.4310/jdg/1612975017. arXiv: 1707.09325. Zbl: 1464.53067.  (cit. on p. 13)
- [Joy02] Dominic Joyce. *Special Lagrangian m -folds in C^m with symmetries*. *Duke Math. J.* 115.1 (2002), pp. 1–51. DOI: 10.1215/S0012-7094-02-11511-7.  (cit. on pp. 18, 30)
- [Joy01] Dominic D. Joyce. *Evolution equations for special Lagrangian 3-folds in C^3* . *Ann. Global Anal. Geom.* 20.4 (2001), pp. 345–403. DOI: 10.1023/A:1013034620426.  (cit. on pp. 18, 30)

- [Joy07] Dominic D. Joyce. *Riemannian holonomy groups and calibrated geometry*. Vol. 12. Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, 2007, pp. x+303. [↗](#) (cit. on pp. 10, 13, 23)
- [KLL20] Spiro Karigiannis, Naichung Conan Leung, and Jason D. Lotay, eds. *Lectures and Surveys on G_2 -Manifolds and Related Topics*. Springer US, 2020. DOI: 10.1007/978-1-0716-0577-6. [↗](#) (cit. on p. 10)
- [KL20] Spiro Karigiannis and Jason D. Lotay. *Deformation theory of G_2 conifolds*. *Communications in Analysis and Geometry* 28.5 (2020), pp. 1057–1210. DOI: 10.4310/cag.2020.v28.n5.a1. [↗](#) (cit. on pp. 23, 34, 39)
- [Kaw15] K. Kawai. *Some associative submanifolds of the squashed 7–sphere*. *The Quarterly Journal of Mathematics* 66.3 (2015), pp. 861–893. DOI: 10.1093/qmath/hav021. MR: 3396095. Zbl: 1326.53070 (cit. on p. 14)
- [Kov03] A. Kovalev. *Twisted connected sums and special Riemannian holonomy*. *Journal für die Reine und Angewandte Mathematik* 565 (2003), pp. 125–160. DOI: 10.1515/crll.2003.097. MR: MR2024648. Zbl: 1043.53041 (cit. on p. 13)
- [KL11] A. Kovalev and N.-H. Lee. *$K3$ surfaces with non-symplectic involution and compact irreducible G_2 –manifolds*. *Mathematical Proceedings of the Cambridge Philosophical Society* 151.2 (2011), pp. 193–218. DOI: 10.1017/S030500411100003X. MR: 2823130. Zbl: 1228.53064 (cit. on p. 13)
- [KW18] Rob Kusner and Peng Wang. *On the index of minimal 2-tori in the 4-sphere* (Mar. 2018). eprint: 1803.01615. [↗](#) (cit. on p. 29)
- [Law89] Gary Lawlor. *The angle criterion*. *Invent. Math.* 95.2 (1989), pp. 437–446. DOI: 10.1007/BF01393905. [↗](#) (cit. on p. 31)
- [LM89] H. B. Lawson Jr. and M.-L. Michelsohn. *Spin geometry*. Vol. 38. Princeton Mathematical Series. Princeton, NJ: Princeton University Press, 1989, pp. xii+427. MR: 1031992. Zbl: 0688.57001 (cit. on p. 13)
- [LM85] R. B. Lockhart and R. C. McOwen. *Elliptic differential operators on noncompact manifolds*. *Annali della Scuola Normale Superiore di Pisa* 12.3 (1985), pp. 409–447. MR: 837256. Zbl: 0615.58048. [↗](#) (cit. on pp. 23, 25, 34, 38)
- [Loto7a] J. D. Lotay. *Coassociative 4–folds with conical singularities*. *Comm. Anal. Geom.* 15.5 (2007), pp. 891–946. MR: 2403190. [↗](#) (cit. on p. 33)
- [Loto9] J. D. Lotay. *Desingularization of coassociative 4–folds with conical singularities*. *Geometric and Functional Analysis* 18.6 (2009), pp. 2055–2100. DOI: 10.1007/s00039-009-0711-1. MR: 2491698 (cit. on p. 2)
- [Lot14] J. D. Lotay. *Desingularization of coassociative 4–folds with conical singularities: obstructions and applications*. *Transactions of the American Mathematical Society* 366.11 (2014), pp. 6051–6092. DOI: 10.1090/S0002-9947-2014-06193-X. MR: 3256193 (cit. on p. 2)

- [Lot12] J.D. Lotay. *Associative submanifolds of the 7–sphere*. *Proceedings of the London Mathematical Society* 105.6 (2012), pp. 1183–1214. DOI: 10.1112/plms/pdso29. MR: 3004102. Zbl: 1268.53019 (cit. on p. 14)
- [Lot11] Jason D. Lotay. *Asymptotically conical associative 3-folds*. *Q. J. Math.* 62.1 (2011), pp. 131–156. DOI: 10.1093/qmath/hap036.  (cit. on pp. 19, 30)
- [Loto7b] Jason Dean Lotay. *Calibrated submanifolds of \mathbf{R}^7 and \mathbf{R}^8 with symmetries*. *Q. J. Math.* 58.1 (2007), pp. 53–70. DOI: 10.1093/qmath/halo15.  (cit. on pp. 19, 30)
- [Mad21] Jesse Madnick. *The Second Variation for Null-Torsion Holomorphic Curves in the 6-Sphere* (Jan. 2021). arXiv: 2101.09580.  (cit. on pp. 3, 19, 29)
- [Maroz] S. Marshall. *Deformations of special Lagrangian submanifolds*. DPhil. Thesis. University of Oxford, 2002.  (cit. on pp. 23, 27, 34, 36, 37, 40)
- [MP78] V. G. Maz'ya and B. A. Plamenevskii. *Estimates in L_p and in Hölder classes, and the Miranda–Agmon maximum principle for the solutions of elliptic boundary value problems in domains with singular points on the boundary*. *Mathematische Nachrichten* 81 (1978), pp. 25–82. MR: 0492821 (cit. on p. 38)
- [MS12] D. McDuff and D. Salamon. *J–holomorphic curves and symplectic topology*. Second. Vol. 52. American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2012, pp. xiv+726. MR: 2954391. Zbl: 1272.53002 (cit. on pp. 20, 21, 52)
- [McL98] R. C. McLean. *Deformations of calibrated submanifolds*. *Communications in Analysis and Geometry* 6.4 (1998), pp. 705–747. DOI: 10.4310/CAG.1998.v6.n4.a4. MR: 1664890. Zbl: 0929.53027 (cit. on p. 16)
- [Nor13] J. Nordström. *Desingularising intersecting associatives*. unpublished. 2013 (cit. on p. 10)
- [Row99] Todd Westley Rowland. *Smooth holomorphic curves in S^6* . Thesis (Ph.D.)–The University of Chicago. ProQuest LLC, Ann Arbor, MI, 1999, p. 55.  (cit. on p. 19)
- [SW17] D. A. Salamon and T. Walpuski. *Notes on the octonions*. *Proceedings of the 23rd Gökova Geometry–Topology Conference*. 2017, pp. 1–85. arXiv: 1005.2820. MR: 3676083. Zbl: 06810387.  (cit. on p. 10)
- [She95] Zhong Min Shen. *A convergence theorem for Riemannian submanifolds*. *Trans. Amer. Math. Soc.* 347.4 (1995), pp. 1343–1350. DOI: 10.2307/2154814.  (cit. on p. 50)
- [Sim83] L. Simon. *Lectures on geometric measure theory*. Vol. 3. Proceedings of the Centre for Mathematical Analysis, Australian National University. Canberra: Australian National University Centre for Mathematical Analysis, 1983. MR: 756417. Zbl: 0546.49019 (cit. on p. 18)
- [Spo15] Luca Spolaor. *Almgren’s type regularity for Semicalibrated Currents* (Nov. 2015). arXiv: 1511.07705.  (cit. on p. 18)

- [Wen19] C. Wendl. *Lectures on Symplectic Field Theory*. 2019. arXiv: 1612.01009 (cit. on p. 51)
- [Wen21] C. Wendl. *How I learned to stop worrying and love the Floer C_ε space*. 2021.  (cit. on p. 50)