

# Scalar relativistic effects with Multiwavelets: Implementation and benchmark

Anders Brakestad,<sup>†,‡</sup> Stig Rune Jensen,<sup>†,‡,¶</sup> Christian Tantardini,<sup>†,‡,§</sup> Quentin  
Pitteloud,<sup>†,‡</sup> Peter Wind,<sup>†,‡</sup> Jānis Užulis,<sup>||</sup> Andris Gulans,<sup>||</sup> Kathrin Helen  
Hopmann,<sup>‡</sup> and Luca Frediani<sup>\*,†,‡</sup>

<sup>†</sup>*Hylleraas Centre for Quantum Molecular Sciences, UiT The Arctic University of Norway,  
9037 Tromsø, Norway*

<sup>‡</sup>*Department of Chemistry, UiT The Arctic University of Norway, 9037 Tromsø, Norway*

<sup>¶</sup>*First authorship is shared between AB and SRJ*

<sup>§</sup>*Department of Materials Science and NanoEngineering, Rice University, Houston, Texas  
77005, United States of America.*

<sup>||</sup>*Department of Physics, University of Latvia, Jelgavas iela 3, Riga, LV-1004, Latvia*

E-mail: luca.frediani@uit.no

## Abstract

The importance of relativistic effects in quantum chemistry is widely recognized, not only for heavier elements but throughout the periodic table. At the same time, relativistic effects are strongest in the nuclear region, where the description of electrons through linear combination of atomic orbitals becomes more challenging. Furthermore, the choice of basis sets for heavier elements is limited compared to lighter elements where precise basis sets are available. Thanks to the framework of multiresolution analysis, multiwavelets provide an appealing alternative to overcome this challenge: they lead to robust error control and adaptive algorithms that automatically refine the

basis set description until the desired precision is reached. This allows to achieve a proper description of the nuclear region.

In this work we extended the Multiwavelet-based code **MRChem** to the scalar zero-order regular approximation (ZORA) framework. We validated our implementation comparing the total energies for a small set of elements and molecules. To confirm the validity of our implementation, we compared both against a radial numerical code for atoms and the plane-wave based code **exciting**.

# Introduction

In his famous Nobel Price Lecture, P.A.M. Dirac stated that with the advent of quantum mechanics all fundamental problems of chemistry were in principle solved<sup>1</sup>. It is however interesting that he did not seem to realize that relativity would also play a role, despite his own fundamental contribution to combining relativity and quantum mechanics<sup>2,3</sup>. It is now widely accepted that the correct description of the electronic structure of atoms and molecules requires the inclusion of relativistic effects. This is particularly relevant for the core- and the innermost valence shell-electrons of heavier elements, which move at sizeable fractions of the speed of light. A wide number of chemical properties and phenomena, such as the yellow colour of gold<sup>4,5</sup>, the liquid state of mercury<sup>5,6</sup>, the functioning of lead-acid batteries<sup>5,7</sup>, the catalytic behavior of cobalt<sup>5,8-10</sup>, the difficulties to describe fluorinated compounds with DFT<sup>11-13</sup> all depend on relativistic effects. There is a wide spectrum of methods to include relativistic effects in electronic structure calculations. They range from the use of effective core potential (ECP)<sup>14-17</sup>, also known as pseudopotentials in the solid state physics community<sup>18</sup>, to the solution of the 4-component Dirac equation. In between these two extremes are a number of methods with different degrees of accuracy and complexity<sup>19</sup>.

The ECP method collapses a given number of core-electrons and the nucleus to provide a core potential for the valence electron. This has the dual advantage of reducing the number of explicit electrons and at the same time including relativistic effects implicitly. On the other hand, if core electronic properties such as high pressure chemistry, core-electron spectroscopy<sup>20</sup> and nuclear magnetic resonance (NMR) shielding constants<sup>9</sup> are considered, ECP methods fall short, and all-electron calculations are necessary. The four-component Dirac equation<sup>2</sup> constitutes the starting point for the treatment of relativistic effects, and the full Breit Hamiltonian<sup>21-25</sup> is the the most complete treatment of relativistic effects for a many-electron system. On the other hand, it is also the most computationally expensive: spinorbitals are 4-component complex functions. Spin is no longer a good quantum number in a relativistic framework and the simplified picture of two electrons with opposite spin shar-

ing the same orbital is no longer valid<sup>19,26,27</sup>. Most operators couple the different components of a spinorbital, leading roughly to a factor 100 in computational cost, because coupling four complex functions requires  $8 \times 8$  matrices instead of a real scalar<sup>28–31</sup>. Although progress has been made to make 4-component calculations faster and easily available<sup>28–31</sup>, it is still convenient to attempt approximations that promise great reduction in the computational cost at the price of reduced accuracy. The first step in such a hierarchy of approximations is the elimination of the small components through a Foldy–Wouthuysens (FWs) transformation<sup>32,33</sup>. The choice of transformation leads to different kinds of methods, such as the Pauli Hamiltonian<sup>34–36</sup>, the Regular Approximations (RAs)<sup>37–39</sup>, the Douglas-Kroll-Hess (DKH) Hamiltonian<sup>40–44</sup>, or the exact two-component (X2C) Hamiltonian method<sup>19,45–53</sup>. Further reduction to a scalar method is also possible for the Pauli Hamiltonian and the RAs<sup>37–39</sup>. In particular, the RAs has two interesting features: (1) the decoupling part of the FWs transformation can be expanded in a convergent series to recover the exact elimination, and (2) the renormalization part can be exactly incorporated in the wave function. The ZORA<sup>34,37,54–56</sup> is the simplest form of Hamiltonian keeping only the zeroth order in both parts. The reduction to a scalar method is carried out by applying the Dirac identity and discarding the spin-orbit term. The advantage of ZORA is to keep most of the standard algorithms of quantum chemistry in their original form just re-scaling the kinetic energy by a function that includes the potential energy (see below Theory section)<sup>34,37,54–56</sup>. The numerical treatment of this rescaling function can be challenging, because it displays a cusp at each nucleus<sup>34,37,54–56</sup>. Standard approaches, based on atomic orbital expansions, can struggle to get an accurate description in this region. The issue is further aggravated for heavier nuclei, where such a correction is important, and at the same time the number of basis sets available is more limited and less is known about their true precision<sup>57,58</sup>.

Some efforts to assess the precision of ZORA for all-electron calculations of heavier elements have been undertaken using Gaussian type orbitals (GTOs), but it is anyway challenging to assess the precision without an external reference<sup>57,58</sup>. One such option for hydrogen-

like ions ( $\text{He}^+$ ,  $\text{Ne}^{9+}$ ,  $\text{Ar}^{17+}$ , and so on) is constituted by the scaling properties of the ZORA Hamiltonian in a two-component framework, which yields the (scaled) exact Dirac energies for such a system<sup>59</sup>. However, for many-electron systems, and/or for a scalar relativistic approach, assessing the true precision of a GTO basis is challenging and several basis sets are developed to include relativistic effects. Examples of all-electron relativistic basis sets include the universal Gaussian basis set (UGBS)<sup>60</sup>, the atomic natural orbitals basis sets<sup>61–65</sup>, the X2C basis sets<sup>66</sup>, and the segmented all-electron relativistic contracted (SARC) basis sets<sup>67–71</sup>. Importantly, such basis sets have to be fitted to the chosen Hamiltonian (ZORA, DKH). This comes on top of the required fitting of the basis set to a given electronic structure method, and if relevant, to a particular property<sup>71</sup>. Although some of the known all-electron relativistic basis sets may provide good results for a particular method and property, their transferability is limited. The large number of available GTO basis sets is an indication that no single basis set is good enough to describe all properties of interest to sufficient precision<sup>58</sup>.

In recent years, multiwavelets (MWs)<sup>72</sup> have emerged as a powerful alternative to traditional local basis sets<sup>73</sup>. Their foundation based on multiresolution analysis<sup>74</sup> leads to a basis set that is not empirically parameterized. Robust error control<sup>75–77</sup> means that the user can set a finite but arbitrary target precision, and adaptive algorithms<sup>78–80</sup> ensure that the representation of molecular orbitals is automatically refined until the required precision is reached. Instead of a plethora of bases to choose from, the user only needs to set the requested precision and the polynomial order of the basis, providing in practice a robust black-box, where straightforward numerical considerations guide the user’s choice. MWs have proven reliable and robust in providing Hartree-Fock (HF) and density functional theory (DFT) benchmark results for energies<sup>73</sup> and properties,<sup>81</sup> and have ventured both towards post-HF methods<sup>82</sup> and relativistic treatments.<sup>83</sup>

In this work we present a MW implementation of the ZORA method in the **MRChem** code. To verify the correctness of the implementation, we consider two alternative methods: a

radial atomic solver implemented as a separate package<sup>84</sup> and linearized augmented plane wave (LAPW) implemented in the electronic-structure code `exciting`<sup>85</sup>. Both of these approaches provide means for systematic improvement of the precision and are capable of yielding total energies approaching the complete basis set (CBS) limit as demonstrated in Refs. 84 and 86.

## Theory and implementation

### From the Dirac equation to the ZORA Hamiltonian

We will briefly expose how the ZORA<sup>34,37,54–56</sup> is derived starting from the 4-component Dirac equation of an electron, which describes a 4-component spinor  $\Psi$ , subject to a potential  $V$ :

$$(\beta c^2 + V + c\boldsymbol{\alpha} \cdot \mathbf{p}) \Psi = E\Psi, \quad (1)$$

In the above equation  $\mathbf{p}$  is the momentum operator,  $c$  is the speed of light, atomic units ( $m_e = 1$ ,  $\hbar = 1$ ,  $e = -1$ ) are assumed and  $\boldsymbol{\alpha}$  and  $\beta$  are defined as follows:

$$\beta = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}, \quad \boldsymbol{\alpha} = \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{pmatrix},$$

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (2)$$

The first step towards ZORA consists in applying the FW transformation to the Dirac Hamiltonian of Eq. (1):

$$H^{FW} = U^\dagger H^D U, \quad (3)$$

where the transformation matrix  $U = W_1 W_2$  is a product of a decoupling matrix  $W_1$  and a renormalization matrix  $W_2$ :

$$W_1 = \begin{pmatrix} 1 & -R^\dagger \\ R & 1 \end{pmatrix}, \quad W_2 = \begin{pmatrix} 1/\sqrt{1+R^\dagger R} & 0 \\ 0 & 1/\sqrt{1+RR^\dagger} \end{pmatrix}, \quad (4)$$

and  $R$  is the exact coupling between the large and the small components of a 4-spinorbits:

$$R = \frac{1}{2c^2 - V + E} c \boldsymbol{\sigma} \cdot \mathbf{p}. \quad (5)$$

The inverse potential term depends on the eigenvalue  $E$ , and this dependence can be expanded in a Taylor series as follows:

$$R = \frac{1}{2c^2 - V} \left( 1 + \frac{E}{2c^2 - V} \right)^{-1} \boldsymbol{\sigma} \cdot \mathbf{p} = \frac{1}{2c^2 - V} \sum_{k=0}^{\infty} \frac{(-E)^k}{(2c^2 - V)^k} \boldsymbol{\sigma} \cdot \mathbf{p}. \quad (6)$$

.

Restricting the expansion to the zero-th order leads to RA. Once the renormalization  $W_2$  is also considered, the following 2-component Hamiltonian is obtained:

$$H^{RA} = \frac{1}{\sqrt{1+R^\dagger R}} \left[ V + \boldsymbol{\sigma} \cdot \mathbf{p} \frac{c^2}{2c^2 - V} \boldsymbol{\sigma} \cdot \mathbf{p} \right] \frac{1}{\sqrt{1+R^\dagger R}}. \quad (7)$$

The ZORA Hamiltonian is finally obtained by Taylor-expanding the renormalization operator and retaining only the zero-order term:

$$H^{ZORA} = V + \boldsymbol{\sigma} \cdot \mathbf{p} \frac{c^2}{2c^2 - V} \boldsymbol{\sigma} \cdot \mathbf{p}. \quad (8)$$

Using the Dirac identity

$$(\boldsymbol{\sigma} \cdot \mathbf{A})(\boldsymbol{\sigma} \cdot \mathbf{B}) = \mathbf{A} \cdot \mathbf{B} + i \boldsymbol{\sigma} \cdot (\mathbf{A} \times \mathbf{B}), \quad (9)$$

we can separate the scalar-relativistic and spin-orbit contributions to the kinetic energy covered by the first and the second terms, respectively. Here, we keep only the scalar-relativistic part and obtain the following Hamiltonian:

$$H_{SR}^{ZORA} = V + \mathbf{p} \frac{c^2}{2c^2 - V} \cdot \mathbf{p} = V + \frac{1}{2} \mathbf{p} \kappa \cdot \mathbf{p} \quad (10)$$

where in the last expression we have implicitly defined  $\kappa = (1 - \frac{V}{2c^2})^{-1}$ . Given the ZORA Hamiltonian, a Kohn-Sham (KS) DFT implementation is then obtained by replacing the non-relativistic kinetic energy operator with its ZORA counterpart:

$$\left( \frac{1}{2} \mathbf{p} \cdot \kappa \mathbf{p} + V \right) \varphi_i = \epsilon_i \varphi_i \quad (11)$$

It should be noted that in practical implementations the potential defining  $\kappa$  is usually *not* the full KS potential. Given the form of  $\kappa$ , the most important contribution is the nuclear attraction. Introducing Coulomb ( $\hat{J}$ ) and exchange and correlation ( $\hat{V}_{xc}$ ) is possible, but the corresponding operator has to be recomputed numerically at each iteration. Additionally, operations such as function multiplications are difficult to perform in traditional linear combination atomic orbital basis representations. A common choice for GTO calculations is to use a fixed atomic potential, which is called the atomic-ZORA approximation<sup>87</sup>. The ZORA kinetic operator can then be pre-computed and used throughout the calculation. Atomic-ZORA has the advantage of being gauge-invariant<sup>87</sup>.

## ZORA equations in a Multiwavelet framework

To obtain a MW implementation of the ZORA eigenvalue problem, it is necessary to transform the differential equation into an integral equation, in analogy with the non-relativistic case<sup>75,77</sup>. The standard KS equations can be concisely written as follows:

$$\hat{F} \varphi_i = \sum_j F_{ij} \varphi_j, \quad (12)$$



where  $\hat{F}$  is the Fock operator,  $\varphi_i$  refer to an occupied molecular orbital, and  $F_{ij}$  are the matrix elements of the Fock operator between two occupied orbitals, assuming a general non-canonical (non-diagonal) form.

Within the framework of KS-DFT, the Fock operator consists of the kinetic energy  $\hat{T}$ , the nuclear attraction  $\hat{V}_{nuc}$ , the Coulomb repulsion  $\hat{J}$ , the Hartree-Fock exchange  $\hat{K}$  scaled by some numerical factor  $\lambda \in [0, 1]$ , and the exchange and correlation potential  $\hat{V}_{xc}$ :

$$\hat{F} = \hat{T} + \hat{V}_{nuc} + \hat{J} - \lambda \hat{K} + \hat{V}_{xc}. \quad (13)$$

In the non-relativistic domain, the coupled KS differential equations (12) can be rewritten in integral form,<sup>88,89</sup> by making use of the bound-state Helmholtz kernel

$$\varphi_i = -2\hat{G}_{\mu_i} \left( \hat{V} \varphi_i - \sum_{j \neq i} F_{ij} \varphi_j \right), \quad (14)$$

where  $\hat{G}_{\mu_i} = [-\nabla^2 - \mu_i^2]^{-1}$  is the integral convolution operator associated with the bound-state Helmholtz kernel  $G(r) = e^{-\mu_i r}/r$ , using  $\mu_i = \sqrt{-2F_{ii}}$  (i.e. the diagonal elements of the Fock matrix).

In the ZORA Hamiltonian, the kinetic energy operator becomes:

$$\hat{T} = \mathbf{p} \cdot \frac{c^2}{2c^2 - V_Z} \mathbf{p} = -\frac{c^2}{2c^2 - V_Z} \nabla^2 - \nabla \frac{c^2}{2c^2 - V_Z} \cdot \nabla = -\frac{1}{2} \kappa \nabla^2 - \frac{1}{2} \nabla \kappa \cdot \nabla. \quad (15)$$

where  $V_Z = \hat{V}_{nuc} + \hat{J} + \hat{V}_{xc}$ . Including  $\hat{J}$  and  $\hat{V}_{xc}$  does not pose any issue in the MW framework, other than the computational overhead of having to update the potential at every iteration, because all potentials are anyway treated on an equal footing using a numerical grid. The non-local Hartree-Fock exchange, in turn, does not seem to contribute significantly to  $V_Z$  as shown in Ref. 90.

Inserting the ZORA kinetic operator into Eq. (12) we obtain:

$$\left(-\frac{1}{2}\kappa\nabla^2 - \frac{1}{2}\nabla\kappa \cdot \nabla + V\right)\varphi_i = \sum_j F_{ij}\varphi_j. \quad (16)$$

In order to make use of the same framework as the non-relativistic implementation of Eq. (14), it is necessary to isolate the Laplacian and the diagonal element of the sum on the right-hand side, which together make up the bound-state Helmholtz operator  $\hat{G} = (\nabla^2 + 2F_{ii})^{-1}$ . This is achieved first by division by  $\kappa$ , recalling that  $\kappa^{-1} = 1 - V_Z/2c$ . The following integral equation is obtained:

$$\varphi_i = -2\hat{G}_{\mu_i} \left[ -\frac{1}{2} \frac{\nabla\kappa}{\kappa} \cdot \nabla \varphi_i + \left( \frac{V}{\kappa} + \frac{V_Z}{2c^2} F_{ii} \right) \varphi_i - \frac{1}{\kappa} \sum_{j \neq i} F_{ij} \varphi_j \right]. \quad (17)$$

When  $c \rightarrow \infty$ ,  $\kappa \rightarrow 1$  and  $\nabla\kappa \rightarrow 0$ , and the non-relativistic form as in Eq. (14) is recovered. Eq.17 can therefore be seen as a *level-shifted* version of its non-relativistic counterpart. Although it cannot be expected that the iterative solution of Eq. (17) will work for arbitrary shifts, the approach is justified by recalling that  $\kappa \simeq 1$  almost everywhere, except close to the nuclei. Our tests indicate that it becomes more difficult to converge the above equation when the ZORA contribution becomes larger, either *physically* by going down the periodic table, or *artificially* by letting  $c \rightarrow 0$ . To overcome the convergence issues for heavier nuclei (5<sup>th</sup> row of the periodic table) we have therefore introduced a finite nucleus model, as described in Section . For elements beyond the 5<sup>th</sup> row, one has to keep in mind that ZORA becomes questionable<sup>91</sup>.

An alternative approach to obtain the desired  $(\nabla^2 + 2F_{ii})$  term from Eq. (16) is to add a Laplacian term  $(\frac{1}{2}\nabla^2)$  directly, thus avoiding division by  $\kappa$ . Our tests indicate that the strategy presented above works better, despite the additional singularity introduced. The main reason seems to be that the former strategy removes the Laplacian altogether, which is ill-conditioned in the discontinuous MW basis, whereas the latter keeps part of it on the right-hand side.

## Implementation with Multiwavelets

An implementation of the ZORA method as outlined above, is currently in a development version of the **MRChem** package<sup>92</sup>, and is expected to appear in the next official release v1.2. **MRChem** is a numerical quantum chemistry code based on a MW framework, in which all functions and operators are represented on their own fully adaptive multi-resolution numerical real-space grid. This allows for efficient all-electron treatment of medium to large molecules (hundreds of atoms) at Self Consistent Field (SCF) level of theory (both HF and DFT).

The  $\kappa$  function is computed as a point-wise map of the chosen ZORA potential  $V_Z$  through

$$\kappa(r) = \frac{1}{1 - V_Z(r)/2c^2}, \quad (18)$$

and similarly for its inverse

$$\kappa^{-1}(r) = 1 - V_Z(r)/2c^2. \quad (19)$$

Both functions are represented on their own adaptive numerical grid, and subsequently treated as standard multiplicative potential operators in the SCF procedure of solving Eq. (17).

## Point nucleus models

In a MW framework, the singularity of the nuclear potential can lead to numerical problems when  $V_{nuc}$  is computed and used. In the non-relativistic case, the issue is circumvented by replacing the analytic  $1/r$  potential with a smoothed approximation<sup>75</sup>:

$$u(r) = \frac{\text{erf}(r)}{r} + \frac{1}{3\sqrt{\pi}} \left( e^{-r^2} + 16e^{-4r^2} \right), \quad (20)$$

and then parameterized as  $u(r/s)/s$ , where  $s$  is a scalar smoothing parameter. The smoothing parameter depends on the nuclear charge  $Z$  and the desired precision  $\epsilon$  as explained in Ref

$$s = \left( \frac{0.00435\epsilon}{Z^5} \right)^{1/3}. \quad (21)$$

The above prescription constitutes a *numerical* smoothing of the nucleus to avoid accidental infinities in the representations. It should not be confused with *physical* finite nucleus models<sup>93</sup>, which are common in relativistic methods. This numerical smoothing is much sharper than the common finite nucleus models, and it is meant to yield results which are – within the requested precision – equivalent to using a point charge.

## Finite nucleus models

In order to overcome the numerical issues faced by the pointwise nuclei, we have introduced the Gaussian nuclear model, as described by Visscher *et al.*<sup>93</sup>. Not only does the model overcome the numerical problems of pointwise charges, but it is also a sounder physical description of larger nuclei. The nuclear charge is modelled as a Gaussian distribution

$$\rho^G(R) = \rho_0^G e^{-\xi R^2} \quad (22)$$

with the normalisation prefactor  $\rho_0^G = eZ(\xi/\pi)^{3/2}$  and the parameter  $\xi$  related to the the root-mean-square radius of the nucleus via the expression  $\xi = 3/\langle R^2 \rangle$ . To determine  $\langle R^2 \rangle$ , we apply the empirical formula for Ref. 94:

$$\sqrt{\langle R^2 \rangle} = (0.836A^{1/3} + 0.570)fm \quad (23)$$

where  $A$  is the nuclear mass number and  $fm$  is a femtometer length unit ( $10^{-15}$  m). We note that a choice of the nuclear mass  $A$  is depends on whether the abundance of isotopes is taken in to account. To avoid the ambiguity, we use the tabulated values of  $\sqrt{\langle R^2 \rangle}$  provided in Ref. 93.

The potential for a Gaussian charge distribution can be represented in an analytic form

by Visscher *et al.*<sup>93</sup>:

$$V^G(R) = -\frac{eZ}{R}\text{erf}(\sqrt{\xi}R), \quad (24)$$

which is the solution to the Poisson equation for the charge density defined by Eq. (22).

Among the different possibilities presented in Ref. 93, we have chosen the Gaussian model for the present work because it is simpler than the more realistic Fermi model and because it is implemented in all GTO codes. The goal of the present work is the validation of the implementation. It is therefore less relevant which model is used, provided that it is consistent throughout the software packages employed.

## Methods for verification

To verify correctness of the ZORA implementation in **MRChem**, we use two other codes: a numerical atomic solver<sup>84</sup> and an all-electron full-potential LAPW code **exciting**. In this section, we briefly introduce them and provide details on the implementation of new features needed for making a direct comparison with **MRChem**.

The atomic solver assumes atoms with spherically symmetric densities, and the mono-electronic wave functions are represented as  $\varphi_{nlm}(\mathbf{r}) = u_{nl}(r)Y_{lm}(\hat{r})$ , where  $u_{nl}(r)$  is a radial function defined on a one-dimensional radial grid. The SCF solver uses a similar approach as the MW framework described above. It reduces the non-relativistic Kohn-Sham equation to the integral form as follows:

$$\varphi_{nlm}(\mathbf{r}) = -2\hat{G}_{\epsilon_{nl}}\left(\hat{V}\varphi_{nlm}(\mathbf{r})\right). \quad (25)$$

This equation matches Eq. (14), with vanishing off-diagonal terms of the Fock matrix, because the canonical representation is used. The radial solver originally supported only non-relativistic calculations: to implement ZORA, we used Eq. (17) in the canonical form, i.e., replacing  $F_{ii}$  with  $\epsilon_{nl}$  and setting  $F_{ij} = 0$  for  $i \neq j$ . This approach avoids the evaluation of second derivatives and the corresponding numerical noise which accumulates during the

self-consistency iterations.

To consider systems beyond atoms, we use **exciting**. In a nutshell, the code relies on partitioning the unit cell into non-overlapping atomic spheres and the interstitial region. In the atomic spheres, the wavefunctions are expressed in terms of atomic-like orbitals that are updated during the self-consistence cycle. In the interstitial region, one represents the wavefunctions with plane waves using the smoothness of the Kohn-Sham potential. Based on such an approach, we use two types of basis functions: (i) augmented plane-waves (APWs)<sup>95,96</sup> and (ii) local-orbitals (LOs)<sup>97</sup>. Each APW combines a plane wave in the interstitial region with atomic-like orbitals in the spheres, whereas each LO is a linear combination of two atomic-like orbitals in one particular sphere and strictly zero everywhere else. As shown in Ref. 86, it is possible to obtain a systematic convergence of the total energies simply by increasing the number of APWs and LOs.

The ZORA Hamiltonian is already available in the released version of the code, whereas the smeared nucleus feature described in Sec. is implemented in the present work. Aside from adopting the electrostatic potential due to a Gaussian charge density, we also revise the evaluation of the following integral:

$$E_M = \int \rho_n(\mathbf{r}) V_C(\mathbf{r}) d\mathbf{r}. \quad (26)$$

If a nucleus at the site  $\mathbf{R}_0$  is defined as a point charge, its density distribution is  $\rho_n(\mathbf{R}) = Z\delta(\mathbf{R} - \mathbf{R}_0)$  leading to  $E_M = ZV_C(\mathbf{R}_0)$ . For smeared nuclei with  $\rho_n(\mathbf{R}) = \rho^G(\mathbf{R})$  (see Eq. 24), Eq. 26 is evaluated as an integral on a radial grid.

## Computational details

All calculations were performed with the PBE functional<sup>98</sup> using the XCFun<sup>99</sup> library, the LibXC<sup>100</sup> library and the native implementation in **MRChem**, the atomic solver and **exciting**, respectively. Being aware of slight inconsistencies in the PBE parameters employed in these

libraries, we set  $\beta = 0.06672455060314922$  and  $\mu = 0.066725\frac{\pi^2}{3}$  as defined in XCFun and use these values in calculations with all three codes.

For all MW calculations, a development version of **MRChem** has been employed. Interpolating polynomials of 9<sup>th</sup> order were used with a simulation box size of 128  $a_0$ . The numerical precision thresholds used are summarized in Table 1.

Table 1: **MRChem** precision parameters.

Parameter	Value	Explanation
<code>world_prec</code>	1.0e-6	Overall numerical precision
<code>energy_thrs</code>	1.0e-6	Convergence threshold total energy
<code>orbital_thrs</code>	1.0e-4	Convergence threshold maximum orbital residual

All **exciting** calculations were performed using large cubic unit cell with a side length of 25  $a_0$  for atoms and 30  $a_0$  for molecules. The electrostatic interaction of the periodic images was eliminated by introducing the truncation of the Coulomb interaction following the approach explained in Ref. 81. Aside from acquiring the isolated limit, this adjustment allows us to obtain ZORA energies consistent with both **MRChem** and the atomic solver. The unmodified electrostatic potential for periodic densities is defined uniquely except for an additive constant which introduces an ambiguity in the ZORA energies<sup>34,37,54–56</sup>. The truncation of the Coulomb interaction makes the potential unique and thus removes this ambiguity completely<sup>34,37,54–56</sup>. The canonical orbitals were expressed in terms of local orbitals and LAPWs with the cutoff  $R_{\text{MT}}K_{\text{max}}$  sufficient to ensure a few  $\mu\text{Ha}$  precision. The specific settings in the case of each atom and molecule are stored in Table 2 and a set of input and output data files are available in the repository.

Table 2: LAPW Parameters and structural data used in the calculations of the considered molecules and atoms.  $R_{\text{MT}}^{\text{min}}G_{\text{max}}$  is the product of the smallest muffin-tin radius and the largest reciprocal lattice vector.

Material	$R_{\text{MT}}^{\text{min}}G_{\text{max}}$	$R_{\text{MT}}[a_0]$	Bond length [ $\text{\AA}$ ]
CaO	11	1.63 / 1.40	1.8221 <sup>101</sup>
CuH	10	1.40 / 1.00	1.4626 <sup>101</sup>
SrO	13	1.63 / 1.40	1.9050 <sup>102</sup>
Cu <sub>2</sub>	14	1.8	2.2197 <sup>101</sup>
AgH	11	1.68 / 1.20	1.6180 <sup>101</sup>
I <sub>2</sub>	14	1.6	2.6630 <sup>102</sup>
He	11	2	
Ne	12	2	
Ar	13	2	
Kr	13	2	
Xe	15	2	

The calculations with the atomic solver employed a 5<sup>th</sup> order polynomial on a radial grid with the innermost and outermost points  $r_{\text{min}} = 10^{-8} a_0$  and  $r_{\text{max}} = 35 a_0$ . The number of radial points was set to 5000, which is fully sufficient to guarantee a sub- $\mu\text{Ha}$  precision.

The total energies of diatomic molecules were calculated in **MRChem** and **exciting** without geometry relaxation using the internuclear distances given in Table 2. The bond lengths in Table 2, are taken from the NIST<sup>101</sup> database with the exception of SrO and I<sub>2</sub>, which are private communication from not published work.

## Results and discussion

### Relative contributions of the ZORA terms

The current **MRChem** implementation is able to include all local contributions of the potential into  $\hat{V}_Z$ . It is therefore possible to measure their relative weights for a given atom and how their contribution changes with the nuclear charge. We have performed a series of calculations for the noble gases from helium to xenon, with all seven possibilities: one contribution only, two contributions and all three. The results are summarized in Fig. 1. The nuclear



potential  $\hat{V}_{nuc}$  is the largest contribution, as expected. It is followed by the Coulomb term  $\hat{J}$ , and the exchange and correlation potential  $\hat{V}_{xc}$  is the smallest one. Moreover the relativistic correction increases roughly with the fourth power of the nuclear charge as expected,<sup>5</sup> and the nuclear term becomes progressively more dominant for heavier atoms.




Figure 1: Relative contributions to the total energy (in Hartree) for all seven possible ZORA operators, compared to the non-relativistic total energy, for the noble gases He – Xe. Note that the scales on the y-axes are different for each sub-plot.

## Validation

To validate the ZORA implementation in **MRChem**, we perform total energy calculations of noble gas atoms and a small set of molecules broadly covering the first five rows of the periodic table (H-Xe). In the case of the atoms, we apply three different types of calculations: the atomic solver, LAPWs and MWs. For the diatomic molecules we compare the results obtained using MW and LAPW by means of **MRChem** and **exciting**, respectively.

To assess the potential agreement which can be achieved between the various codes, we

have performed non-relativistic calculations with all three codes, using point-charge nuclei (numerically smoothed as described in Section in the case of MWs). The results are summarized in Table 3. We have then obtained a root mean square deviation (RMSD) of the relative error between MWs and LAPWs equal to  $5.21 \cdot 10^{-8}$ , and between MWs and atomic solver equal to  $8.07 \cdot 10^{-10}$ . We concluded that the three methods are in very good agreement, setting the mark for what can be expected in the ZORA domain, using a Gaussian nuclear charge distribution as in Equation (24). The ZORA results are summarized in Table 4 showing a RMSD for the relative errors between MWs and LAPWs equal to  $3.95 \cdot 10^{-8}$ , and between MWs and atomic solver equal to  $6.93 \cdot 10^{-9}$ , in line with what has been observed for the non-relativistic regime, thus confirming the validity of the implementation. In terms of absolute errors, we find that most discrepancies in the total energies are within 10 microHartrees, and only in two cases (AgH and I<sub>2</sub>) the differences are larger, yet they do not exceed 21 microHartrees. This level of agreement is well below the so called *chemical accuracy threshold* (1 kcal/mol).

Finally, we find that introducing the relativistic corrections in the multiwavelet formalism as expressed in Eq. 17 leads to a minor increase in **MRChem** runtimes. Therefore we anticipate that the analysis of the performance (runtimes and parallel scaling) given in Ref.<sup>103</sup> remains valid in the ZORA case.

Table 3: Non-relativistic total energies (given in Hartrees) obtained using three different codes and their relative differences. In all cases, the point-like nucleus model is used.

Species	AtomSolver	exciting	MRChem	Relative difference	
				MRChem vs AtomSolver	MRChem vs exciting
He	-2.892935499	-2.892935485	-2.892935497	-4.1e-10	4.4e-09
Ne	-128.866433504	-128.866433222	-128.866433465	-3.1e-10	1.9e-09
Ar	-527.346141146	-527.346138312	-527.346140237	-1.7e-09	3.7e-09
Kr	-2753.416138121	-2753.416137130	-2753.416137698	-0.1e-09	0.2e-09
Xe	-7234.233259457	-7234.233258510	-7234.233259210	-0.03e-09	0.09e-09
CaO	-	-752.562050270	-752.562058080	-	1.0e-08
CuH	-	-1640.901716750	-1640.901726830	-	6.1e-09
SrO	-	-3208.161711950	-3208.161721780	-	3.1e-09
Cu <sub>2</sub>	-	-3280.675839820	-3280.675845520	-	1.7e-09
AgH	-	-5200.162245050	-5200.162259390	-	2.8e-09
I <sub>2</sub>	-	-13840.158307100	-13840.158328190	-	1.5e-09

Table 4: Scalar-relativistic ZORA total energies (given in Hartrees) obtained using three different codes and their relative differences. In all cases, the smeared nucleus model is used.

Species	AtomSolver	exciting	MRChem	Relative difference	
				MRChem vs AtomSolver	MRChem vs exciting
He	-2.893118631	-2.893118615	-2.89311862	-2.2e-09	3.2e-09
Ne	-129.089335224	-129.089335102	-129.08933496	-2.1e-09	-1.1e-09
Ar	-530.224039154	-530.224036217	-530.22403777	-2.6e-09	2.9e-09
Kr	-2810.049763806	-2810.049761830	-2810.04975504	-3.1e-09	-2.4e-09
Xe	-7564.596665119	-7564.596661910	-7564.59655428	-1.5e-08	-1.4e-08
CaO	-	-757.195492387	-757.195491088	-	-1.7e-09
CuH	-	-1663.238492560	-1663.238503660	-	6.7e-09
SrO	-	-3279.826587400	-3279.826596402	-	2.7e-09
Cu <sub>2</sub>	-	-3325.342851280	-3325.342843306	-	-2.4e-09
AgH	-	-5380.042000350	-5380.042003292	-	0.5e-09
I <sub>2</sub>	-	-14448.747786900	-14448.747761310	-	-1.8e-09

## Conclusions

We have formulated the ZORA method in a form compatible with multiwavelets and implemented it in the MRChem program. The validity and precision of the implementation has been tested against a radial, numerical atomic code and a plane wave code, showing excellent agreement. The current study was done with the specific idea to validate method and theory with a small benchmark: atoms and diatomics, covering broadly the periodic table up to and including 5<sup>th</sup>-row elements. The model is also capable of dealing with all parts of the electronic potential self-consistently, with the exception of the HF exchange, which is the only one which cannot be expressed in closed form.

## Acknowledgement

This work was supported by the Norwegian Research Council through a Centre of Excellence grant (Hylleraas Centre 262695), a FRIPRO grant (ReMRChem 324590), by the Tromsø Research Foundation (TFS2016KHH), and by UNINETT Sigma2 through grants of computer time (nn9330k and nn4654k). Jānis Użulis acknowledges funding provided by the project "Strengthening of the capacity of doctoral studies at the University of Latvia within the framework of the new doctoral model", identification No. 8.2.2.0/20/I/006. Andris Gulans acknowledges funding provided by European Regional Development Fund via the Central Finance and Contracting Agency of Republic of Latvia under the grant agreement 1.1.1.5/21/A/004. We would like to thank Dr. Susi Lehtola from Department of Chemistry at the University of Helsinki in Finland for the useful feedback to the original draft of the manuscript.

## References

- (1) Dirac, P. A. Theory of electrons and positrons. *Nobel Lecture* **1933**, 12, 320–325.

- (2) Dirac, P. A. M. The Quantum Theory of the Electron. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* **1928**, *117*, 610–624.
- (3) Simões, A. Dirac’s Claim and the Chemists. *Physics in Perspective* **2002**, *4*, 253–266.
- (4) Bartlett, N. Relativistic effects and the chemistry of gold. *Gold Bulletin* **1998**, *31*, 22–25.
- (5) Pyykkö, P. Relativistic Effects in Chemistry: More Common Than You Thought. *Annual Review of Physical Chemistry* **2012**, *63*, 45–64.
- (6) Keeton, S.; Loucks, T. Electronic structure of mercury. *Physical Review* **1966**, *152*, 548.
- (7) Ahuja, R.; Blomqvist, A.; Larsson, P.; Pyykkö, P.; Zaleski-Ejgierd, P. Relativity and the lead-acid battery. *Physical Review Letters* **2011**, *106*, 018301.
- (8) Demissie, T. B.; Garabato, B. D.; Ruud, K.; Kozłowski, P. M. Mercury Methylation by Cobalt Corrinoids: Relativistic Effects Dictate the Reaction Mechanism. *Angewandte Chemie International Edition* **55**, 11503–11506.
- (9) Vícha, J.; Marek, R.; Straka, M. High-Frequency  $^{13}\text{C}$  and  $^{29}\text{Si}$  NMR Chemical Shifts in Diamagnetic Low-Valence Compounds of TlI and PbII: Decisive Role of Relativistic Effects. *Inorganic Chemistry* **2016**, *55*, 1770–1781.
- (10) Lein, M.; Rudolph, M.; Hashmi, S. K.; Schwerdtfeger, P. Homogeneous Gold Catalysis: Mechanism and Relativistic Effects of the Addition of Water to Propyne. *Organometallics* **2010**, *29*, 2206–2210.
- (11) Boualili, F. Z.; Nemouchi, M.; Godefroid, M.; Jönsson, P. Weak correlation and strong relativistic effects on the hyperfine interaction in fluorine. *Physical Review A* **2021**, *104*, 062813.

- (12) Mattsson, S.; Paulus, B.; Redeker, F. A.; Beckers, H.; Riedel, S.; Müller, C. The Crystal Structure of  $\alpha$ -F2: Solving a 50 Year Old Puzzle Computationally. *Chemistry – A European Journal* **2019**, *25*, 3318–3324.
- (13) Tantardini, C.; Jalolov, F. N.; Kvashnin, A. G. Crystal Structure Evolution of Fluorine under High Pressure. *The Journal of Physical Chemistry C* **2022**, *126*, 11358–11364.
- (14) Hellmann, H. A new approximation method in the problem of many electrons. *The Journal of Chemical Physics* **1935**, *3*, 61–61.
- (15) Gombás, P. Über die metallische Bindung. *Zeitschrift für Physik* **1935**, *94*, 473–488.
- (16) Xu, X.; Truhlar, D. G. Accuracy of Effective Core Potentials and Basis Sets for Density Functional Calculations, Including Relativistic Effects, As Illustrated by Calculations on Arsenic Compounds. *Journal of Chemical Theory and Computation* **2011**, *7*, 2766–2779.
- (17) Stoll, H.; Metz, B.; Dolg, M. Relativistic energy-consistent pseudopotentials—Recent developments. *Journal of Computational Chemistry* **2002**, *23*, 767–778.
- (18) Heine, V. In *The Pseudopotential Concept*; Ehrenreich, H., Seitz, F., Turnbull, D., Eds.; Solid State Physics; Academic Press, 1970; Vol. 24; pp 1–36.
- (19) Saue, T. Relativistic Hamiltonians for chemistry: A primer. *ChemPhysChem* **2011**, *12*, 3077–3094.
- (20) Rudek, B. et al. Relativistic and resonant effects in the ionization of heavy atoms by ultra-intense hard X-rays. *Nature Communications* **2018**, *9*, 4200.
- (21) Breit, G. An Interpretation of Dirac’s Theory of the Electron. *Proceedings of the National Academy of Sciences* **1928**, *14*, 553–559.
- (22) Breit, G. Dirac’s Equation and the Spin-Spin Interactions of Two Electrons. *Physical Review* **1932**, *39*, 616–624.

- (23) Moss, R. *Advanced molecular quantum mechanics: an introduction to relativistic quantum mechanics and the quantum theory of radiation*; Springer Science & Business Media, 2012.
- (24) Dyall, K. G.; Fægri Jr, K. *Introduction to relativistic quantum chemistry*; Oxford University Press, 2007.
- (25) Helgaker, T.; Coriani, S.; Jørgensen, P.; Kristensen, K.; Olsen, J.; Ruud, K. Recent Advances in Wave Function-Based Methods of Molecular-Property Calculations. *Chemical Reviews* **2012**, *112*, 543–631.
- (26) Jacob, C. R.; Reiher, M. Spin in density-functional theory. *International Journal of Quantum Chemistry* **2012**, *112*, 3661–3684.
- (27) Marian, C. M. Spin–orbit coupling and intersystem crossing in molecules. *Wiley Interdisciplinary Reviews: Computational Molecular Science* **2012**, *2*, 187–203.
- (28) Repisky, M.; Komorovsky, S.; Kadek, M.; Konecny, L.; Ekström, U.; Malkin, E.; Kaupp, M.; Ruud, K.; Malkina, O. L.; Malkin, V. G. ReSpect: Relativistic spectroscopy DFT program package. *The Journal of Chemical Physics* **2020**, *152*, 184101.
- (29) Saue, T. et al. The DIRAC code for relativistic molecular calculations. *The Journal of Chemical Physics* **2020**, *152*, 204104.
- (30) Zhang, Y. et al. BDF: A relativistic electronic structure program package. *The Journal of Chemical Physics* **2020**, *152*, 064113.
- (31) Belpassi, L.; De Santis, M.; Quiney, H. M.; Tarantelli, F.; Storchi, L. BERTHA: Implementation of a four-component Dirac–Kohn–Sham relativistic framework. *The Journal of Chemical Physics* **2020**, *152*, 164118.
- (32) Foldy, L. L.; Wouthuysen, S. A. On the Dirac Theory of Spin 1/2 Particles and Its Non-Relativistic Limit. *Physical Review* **1950**, *78*, 29–36.

- (33) Foldy, L. L. The Electromagnetic Properties of Dirac Particles. *Physical Review* **1952**, *87*, 688–693.
- (34) Chang, C.; Pelissier, M.; Durand, P. Regular two-component Pauli-like effective Hamiltonians in Dirac theory. *Physica Scripta* **1986**, *34*, 394.
- (35) Kang, H.; Uhlmann, G. Inverse problems for the Pauli Hamiltonian in two dimensions. *Journal of Fourier Analysis and Applications* **2004**, *10*, 201–215.
- (36) Goldman, T. Gauge invariance, time-dependent Foldy-Wouthuysen transformations, and the Pauli Hamiltonian. *Physical Review D* **1977**, *15*, 1063.
- (37) van Lenthe, E.; Baerends, E.-J.; Snijders, J. G. Relativistic total energy using regular approximations. *The Journal of Chemical Physics* **1994**, *101*, 9783–9792.
- (38) Dyal, K. G.; van Lenthe, E. Relativistic regular approximations revisited: An infinite-order relativistic approximation. *The Journal of Chemical Physics* **1999**, *111*, 1366–1372.
- (39) Mohri, M.; Nederhof, M.-J. *Robustness in language and speech technology*; Springer, 2001; pp 153–163.
- (40) Douglas, M.; Kroll, N. M. Quantum electrodynamical corrections to the fine structure of helium. *Annals of Physics* **1974**, *82*, 89–155.
- (41) Hess, B. A. Applicability of the no-pair equation with free-particle projection operators to atomic and molecular structure calculations. *Physical Review A* **1985**, *32*, 756.
- (42) Hess, B. A. Relativistic electronic-structure calculations employing a two-component no-pair formalism with external-field projection operators. *Physical Review A* **1986**, *33*, 3742.
- (43) Jansen, G.; Heß, B. A. Revision of the Douglas-Kroll transformation. *Physical Review A* **1989**, *39*, 6016.



- (44) Reiher, M. Douglas–Kroll–Hess Theory: a relativistic electrons-only theory for chemistry. *Theoretical Chemistry Accounts* **2006**, *116*, 241–252.
- (45) Kutzelnigg, W.; Liu, W. Quasirelativistic theory equivalent to fully relativistic theory. *The Journal of Chemical Physics* **2005**, *123*, 241102.
- (46) Liu, W.; Peng, D. Infinite-order quasirelativistic density functional method based on the exact matrix quasirelativistic theory. *The Journal of Chemical Physics* **2006**, *125*, 044102.
- (47) Liu, W.; Peng, D. Exact two-component Hamiltonians revisited. *The Journal of Chemical Physics* **2009**, *131*, 031104.
- (48) Iliáš, M.; Saue, T. An infinite-order two-component relativistic Hamiltonian by a simple one-step transformation. *The Journal of Chemical Physics* **2007**, *126*, 064102.
- (49) Peng, D.; Liu, W.; Xiao, Y.; Cheng, L. Making four- and two-component relativistic density functional methods fully equivalent based on the idea of “from atoms to molecule”. *The Journal of Chemical Physics* **2007**, *127*, 104106.
- (50) Peng, D.; Mikkelsen, N.; Weigend, F.; Reiher, M. An efficient implementation of two-component relativistic exact-decoupling methods for large molecules. *The Journal of chemical physics* **2013**, *138*, 184105.
- (51) Liu, W. Ideas of relativistic quantum chemistry. *Molecular Physics* **2010**, *108*, 1679–1706.
- (52) Liu, W. Advances in relativistic molecular quantum mechanics. *Physics Reports* **2014**, *537*, 59–89.
- (53) Liu, W. Big picture of relativistic molecular quantum mechanics. *National Science Review* **2016**, *3*, 204–221.

- (54) Filatov, M.; Cremer, D. On the physical meaning of the ZORA Hamiltonian. *Molecular Physics* **2003**, *101*, 2295–2302.
- (55) Heully, J.-L.; Lindgren, I.; Lindroth, E.; Lundqvist, S.; Martensson-Pendrill, A.-M. Diagonalisation of the Dirac Hamiltonian as a basis for a relativistic many-body procedure. *Journal of Physics B: Atomic and Molecular Physics* **1986**, *19*, 2799.
- (56) van Lenthe, E.; Baerends, E.-J.; Snijders, J. G. Relativistic regular two-component Hamiltonians. *The Journal of Chemical Physics* **1993**, *99*, 4597–4610.
- (57) Pantazis, D. A.; Neese, F. All-electron basis sets for heavy elements. *WIREs Computational Molecular Science* **2014**, *4*, 363–374.
- (58) Güell, M.; Luis, J. M.; Sola, M.; Swart, M. Importance of the basis set for the spin-state energetics of iron complexes. *The Journal of Physical Chemistry A* **2008**, *112*, 6384–6391.
- (59) van Leeuwen, R.; van Lenthe, E.; Baerends, E. J.; Snijders, J. G. Exact solutions of regular approximate relativistic wave equations for hydrogen-like atoms. *The Journal of Chemical Physics* **1994**, *101*, 1272–1281.
- (60) de Castro, E. V. R.; Jorge, F. E. Accurate universal Gaussian basis set for all atoms of the Periodic Table. *The Journal of Chemical Physics* **1998**, *108*, 5225–5229.
- (61) Zobel, J. P.; Widmark, P.-O.; Veryazov, V. The ANO-R Basis Set. *Journal of Chemical Theory and Computation* **2020**, *16*, 278–294.
- (62) Zobel, J. P.; Widmark, P.-O.; Veryazov, V. Correction to “The ANO-R Basis Set”. *Journal of Chemical Theory and Computation* **2021**, *17*, 3233–3234.
- (63) Roos, B.; Veryazov, V.; Widmark, P.-O. Relativistic atomic natural orbital type basis sets for the alkaline and alkaline-earth atoms applied to the ground-state potentials for the corresponding dimers. *Theoretical Chemistry Accounts* **2004**, *111*, 345–351.

- (64) Roos, B. O.; Lindh, R.; Malmqvist, P.-Å.; Veryazov, V.; Widmark, P.-O. Main Group Atoms and Dimers Studied with a New Relativistic ANO Basis Set. *The Journal of Physical Chemistry A* **2004**, *108*, 2851–2858.
- (65) Roos, B. O.; Lindh, R.; Malmqvist, P.-Å.; Veryazov, V.; Widmark, P.-O. New Relativistic ANO Basis Sets for Transition Metal Atoms. *The Journal of Physical Chemistry A* **2005**, *109*, 6575–6579.
- (66) Pollak, P.; Weigend, F. Segmented Contracted Error-Consistent Basis Sets of Double- and Triple- $\zeta$  Valence Quality for One- and Two-Component Relativistic All-Electron Calculations. *Journal of Chemical Theory and Computation* **2017**, *13*, 3696–3705.
- (67) Pantazis, D. A.; Neese, F. All-electron scalar relativistic basis sets for the lanthanides. *Journal of Chemical Theory and Computation* **2009**, *5*, 2229–2238.
- (68) Pantazis, D. A.; Neese, F. All-electron scalar relativistic basis sets for the actinides. *Journal of Chemical Theory and Computation* **2011**, *7*, 677–684.
- (69) Pantazis, D. A.; Neese, F. All-electron scalar relativistic basis sets for the 6 p elements. *Theoretical Chemistry Accounts* **2012**, *131*, 1–7.
- (70) Aravena, D.; Neese, F.; Pantazis, D. A. Improved segmented all-electron relativistically contracted basis sets for the lanthanides. *Journal of Chemical Theory and Computation* **2016**, *12*, 1148–1156.
- (71) Rolfes, J. D.; Neese, F.; Pantazis, D. A. All-electron scalar relativistic basis sets for the elements Rb–Xe. *Journal of Computational Chemistry* **2020**, *41*, 1842–1849.
- (72) Alpert, B. K. A Class of Bases in  $L^2$  for the Sparse Representation of Integral Operators. *SIAM Journal on Mathematical Analysis* **1993**, *24*, 246–262.
- (73) Jensen, S. R.; Saha, S.; Flores-Livas, J. A.; Huhn, W.; Blum, V.; Goedecker, S.;

- Frediani, L. The elephant in the room of density functional theory calculations. *The Journal of Physical Chemistry Letters* **2017**, *8*, 1449–1457.
- (74) Keinert, F. In *Encyclopedia of Complexity and Systems Science*; Meyers, R. A., Ed.; Springer New York, 2009; pp 5841–5858.
- (75) Harrison, R. J.; Fann, G. I.; Yanai, T.; Gan, Z.; Beylkin, G. Multiresolution quantum chemistry: Basic theory and initial applications. *The Journal of Chemical Physics* **2004**, *121*, 11587–11598.
- (76) Kato, T.; Yokoi, Y.; Sekino, H. Basis set limit computation of dynamic polarizability at near-resonance region. *International Journal of Quantum Chemistry* **2012**, *113*, 286–289.
- (77) Jensen, S. R.; Saha, S.; Flores-Livas, J. A.; Huhn, W.; Blum, V.; Goedecker, S.; Frediani, L. The Elephant in the Room of Density Functional Theory Calculations. *The Journal of Physical Chemistry Letters* **2017**, *8*, 1449–1457.
- (78) Alpert, B.; Beylkin, G.; Gines, D.; Vozovoi, L. Adaptive Solution of Partial Differential Equations in Multiwavelet Bases. *Journal of Computational Physics* **2002**, *182*, 149–190.
- (79) Beylkin, G.; Cheruvu, V.; Pérez, F. Fast adaptive algorithms in the non-standard form for multidimensional problems. *Applied and Computational Harmonic Analysis* **2008**, *24*, 354–377.
- (80) Frediani, L.; Fossgaard, E.; Flå, T.; Ruud, K. Fully adaptive algorithms for multivariate integral equations using the non-standard form and multiwavelets with applications to the Poisson and bound-state Helmholtz kernels in three dimensions. *Molecular Physics* **2013**, *111*, 1143–1160.

- (81) Jensen, S. R.; Flå, T.; Jonsson, D.; Monstad, R. S.; Ruud, K.; Frediani, L. Magnetic properties with multiwavelets and DFT: the complete basis set limit achieved. *Physical Chemistry Chemical Physics* **2016**, *18*, 21145–21161.
- (82) Bischoff, F. A.; Harrison, R. J.; Valeev, E. F. Computing many-body wave functions with guaranteed precision: The first-order Møller-Plesset wave function for the ground state of helium atom. *The Journal of Chemical Physics* **2012**, *137*, 104103.
- (83) Anderson, J.; Sundahl, B.; Harrison, R.; Beylkin, G. Dirac-Fock calculations on molecules in an adaptive multiwavelet basis. *The Journal of Chemical Physics* **2019**, *151*, 234112.
- (84) Užulis, J.; Gulans, A. Radial Kohn–Sham problem via integral-equation approach. *Journal of Physics Communications* **2022**, *6*, 085002.
- (85) Gulans, A.; Kontur, S.; Meisenbichler, C.; Nabok, D.; Pavone, P.; Rigamonti, S.; Sagmeister, S.; Werner, U.; Draxl, C. exciting: a full-potential all-electron package implementing density-functional theory and many-body perturbation theory. *Journal of Physics: Condensed Matter* **2014**, *26*, 363202.
- (86) Gulans, A.; Kozhevnikov, A.; Draxl, C. Microhartree precision in density functional theory calculations. *Physical Review B* **2018**, *97*, 161105.
- (87) Knuth, F.; Carbogno, C.; Atalla, V.; Blum, V.; Scheffler, M. All-electron formalism for total energy strain derivatives and stress tensor components for numeric atom-centered orbitals. *Computer Physics Communications* **2015**, *190*, 33–50.
- (88) Kalos, M. H. Monte Carlo Calculations of the Ground State of Three- and Four-Body Nuclei. *Physical Review* **1962**, *128*, 1791–1795.
- (89) Beylkin, G.; Mohlenkamp, M. J. Numerical operator calculus in higher dimensions. *Proceedings of the National Academy of Sciences* **2002**, *99*, 10246–10251.

- (90) Faas, S.; van Lenthe, J. H.; Hennum, A. C.; Snijders, J. G. An ab initio two-component relativistic method including spin–orbit coupling using the regular approximation. *The Journal of Chemical Physics* **2000**, *113*, 4052–4059.
- (91) Visscher, L.; Saue, T. Approximate relativistic electronic structure methods based on the quaternion modified Dirac equation. *The Journal of Chemical Physics* **2000**, *113*, 3996–4002.
- (92) MRCHEM program package **2020**. DOI: 10.5281/zenodo.3786996.
- (93) Visscher, L.; Dylla, K. Dirac-Fock Atomic Electronic Structure Calculations Using Different Nuclear Charge Distributions. *Atomic Data and Nuclear Data Tables* **1997**, *67*, 207–224.
- (94) Johnson, W.; Soff, G. The lamb shift in hydrogen-like atoms,  $1 \leq Z \leq 110$ . *Atomic Data and Nuclear Data Tables* **1985**, *33*, 405–446.
- (95) Slater, J. C. Wave functions in a periodic potential. *Physical Review* **1937**, *51*, 846.
- (96) Andersen, O. K. Linear methods in band theory. *Physical Review B* **1975**, *12*, 3060.
- (97) Sjöstedt, E.; Nordström, L.; Singh, D. J. An alternative way of linearizing the augmented plane-wave method. *Solid State Communications* **2000**, *114*, 15–20.
- (98) Perdew, J. P.; Burke, K.; Ernzerhof, M. Generalized Gradient Approximation Made Simple. *Physical Review Letters* **1996**, *77*, 3865–3868.
- (99) Ekström, U.; Visscher, L.; Bast, R.; Thorvaldsen, A. J.; Ruud, K. Arbitrary-Order Density Functional Response Theory from Automatic Differentiation. *Journal of Chemical Theory and Computation* **2010**, *6*, 1971–1980.
- (100) Lehtola, S.; Steigemann, C.; Oliveira, M. J.; Marques, M. A. Recent developments in libxc — A comprehensive library of functionals for density functional theory. *SoftwareX* **2018**, *7*, 1–5.

- (101) Huber, K. P.; Herzberg, G. H. *Constants of Diatomic Molecules (data prepared by Jean W. Gallagher and Russell D. Johnson, III) in NIST Chemistry Web-Book, NIST Standard Reference Database Number 69, Eds. P.J. Linstrom and W.G. Mallard, National Institute of Standards and Technology, Gaithersburg MD, 20899, <https://doi.org/10.18434/T4D303>, (retrieved November 24, 2023); 2023.*
- (102) Brakestad, A. private communication.
- (103) Wind, P.; Bjørgve, M.; Brakestad, A.; Gerez S., G. A.; Jensen, S. R.; Eikås, R. D. R.; Frediani, L. MRChem Multiresolution Analysis Code for Molecular Electronic Structure Calculations: Performance and Scaling Properties. *Journal of Chemical Theory and Computation* **2023**, *19*, 137–146, PMID: 36410396.

## TOC Graphic

