Flat hypercomplex nilmanifolds are H-solvable

Yulia Gorginyan

October 5, 2023

Abstract

We say that a hypercomplex nilpotent Lie algebra is \mathbb{H} -solvable if there exists a sequence of \mathbb{H} -invariant subalgebras

$$\mathfrak{g}_1^{\mathbb{H}} \supset \mathfrak{g}_2^{\mathbb{H}} \supset \cdots \supset \mathfrak{g}_{k-1}^{\mathbb{H}} \supset \mathfrak{g}_k^{\mathbb{H}} = 0,$$

such that $[\mathfrak{g}_i^{\mathbb{H}}, \mathfrak{g}_i^{\mathbb{H}}] \subset \mathfrak{g}_{i+1}^{\mathbb{H}}$. Let $N = \Gamma \backslash G$ be a hypercomplex nilmanifold with flat Obata connection and $\mathfrak{g} = \mathrm{Lie}(G)$. We prove that the Lie algebra \mathfrak{g} is \mathbb{H} -solvable.

Contents

1	Introduction		2
	1.1	Affine manifolds	2
	1.2	Hypercomplex affine nilmanifolds	3
	1.3	\mathbb{H} -solvable Lie algebras and algebraic holonomy	5
2	Pre	liminaries: Nilpotent Lie groups and algebras	7
3	Algebraic holonomy group		8
	3.1	Left equivariant vector bundles	8
	3.2	Algebraic holonomy group	11
	3.3	Maltsev completion	13
	3.4	Unipotent holonomy group and H-solvability	16

1 Introduction

1.1 Affine manifolds

A manifold X together with a torsion-free flat connection ∇ is called **an affine manifold**. Equivalently, an affine manifold X is a manifold with an atlas such that all translation maps between charts are in $Aff(\mathbb{R}^n)$ ([FGH], [Sh]).

Recall that a group of affine transformations $\mathrm{Aff}(\mathbb{R}^n)$ is a semidirect product $\mathrm{GL}_n(\mathbb{R}) \ltimes \mathbb{R}^n$. If $f \in \mathrm{Aff}(\mathbb{R}^n)$ then for any $x \in \mathbb{R}^n$ there exist a linear part $A \in \mathrm{GL}(\mathbb{R}^n)$ and a translation part $t \in \mathbb{R}^n$ such that f(x) = Ax + t. The linearization map is the natural homomorphism $l : \mathrm{Aff}(\mathbb{R}^n) \longrightarrow \mathrm{GL}(\mathbb{R}^n)$ given by the formula l(f) = A.

An affine manifold X is called **complete** if its universal cover is affine equivalent to \mathbb{R}^n , i.e. $X = \mathbb{R}^n/\Gamma$, where $\Gamma \subset \text{Aff}(\mathbb{R}^n)$ is a discrete subgroup of the affine transformation group.

There are three famous open questions related to affine geometry. In the historical context, the first was the conjecture of Chern:

Conjecture 1.1: (Chern) The Euler characteristic of a compact affine manifold vanishes [Gol].

Kostant and Sullivan proved Chern's conjecture for the compact complete affine manifolds [KS]. Also, it was proven by Bruno Klingler [Klin] in the case when a compact affine manifold admits a parallel volume form. As far as we know, the general case of Chern's conjecture remains open.

The conjecture of Markus links the existence of a parallel volume form with the completeness of a compact affine manifold.

Conjecture 1.2: (Markus, 1962)[Mar] A compact affine manifold X admits parallel volume form if and only if the manifold X is complete.

Recall that a subgroup Γ of an affine group $Aff(\mathbb{R}^n)$ is called **crystallographic** [A] if its action on \mathbb{R}^n is properly discontinuous, free, and cocompact.

Recall that a group G is said to have a certain property P virtually if G contains a subgroup H of finite index which has the property P.

Theorem 1.3: (Bieberbach, 1911) [Bieb] Every discrete subgroup Γ of an isometry group Isom(\mathbb{R}^n) is virtually abelian. Every crystallographic subgroup of Isom(\mathbb{R}^n) is virtually a translation group. For a given n there exists only a finite number of such groups $\Gamma \subset \text{Isom}(\mathbb{R}^n)$ up to a conjugation.

One of the ways to generalize the theorem of Bieberbach is to consider an affine transformation group $\mathrm{Aff}(\mathbb{R}^n)$ instead of $\mathrm{Isom}(\mathbb{R}^n)$.

Conjecture 1.4: (Auslander, 1964)[Aus] Every crystallographic subgroup of an affine group is virtually solvable, i.e. contains a solvable subgroup of finite index.

Auslander's conjecture was proven by D. Fried and W. Goldman [FG] in the case n = 3 and the result was refined up to dimension n = 6 by H. Abels, G.A. Margulis, and G.A. Soifer [AMS]. Also, Auslander's conjecture is true in a case when the monodromy group preserves a metric of signature (1, n) [GK] and (2, n) [AMS2].

1.2 Hypercomplex affine nilmanifolds

Recall that a nilmanifold N is a compact quotient of a connected simply connected nilpotent Lie group G by a lattice subgroup Γ , which acts on the group G from the left. We denote the quotient as $N = \Gamma \backslash G$. It is called an affine nilmanifold if G has a left-invariant affine structure, i.e. the flat torsion-free connection ∇ such that $L_g^*(\nabla) = \nabla$, where $L_g : G \longrightarrow G$ is the left-translation.

Let $\pi: \widetilde{X} \longrightarrow X$ be the universal cover of a manifold X and $\pi_1(X)$ the fundamental group. An affine immersion $D: \widetilde{X} \longrightarrow \mathbb{R}^n$ is called **the developing map**. By a well-known theorem ([OV], [Gol1]) D exists and is uniquely defined up to an affine automorphism.

Definition 1.5: The affine holonomy representation is a unique homomorphism $h: \pi_1(X) \longrightarrow \operatorname{Aff}(\mathbb{R}^m)$ which satisfies $D \circ \gamma = h(\gamma) \circ D$ for every $\gamma \in \pi_1(X)$. The affine holonomy group $\mathcal{H} := \operatorname{Im}(h)$ is the image of the homomorphism h in $\operatorname{Aff}(\mathbb{R}^n)$. The linear holonomy group is the image of the affine holonomy group under the linearization map, $\mathcal{L} := l(\mathcal{H}) \subset \operatorname{GL}(\mathbb{R}^n)$.

Definition 1.6: A closed Lie subgroup $G \subset GL(V)$ is called a linear algebraic group if G is given by a system of polynomial equations.

Definition 1.7: A representation of a linear algebraic group is called **unipotent** if, in a certain basis, its image is the group of strictly upper-triangular matrices.

Let X be a compact affine manifold whose linear holonomy representation is unipotent. Then X admits a parallel volume form. The converse was proven by Goldman, Fried, and Hirsch, [FGH, Theorem A]:

Theorem 1.8: Let X be a compact affine manifold with a parallel volume form. Assume that its affine holonomy group is nilpotent. Then the linear holonomy representation is unipotent.

We will use the following reformulation of Theorem 1.8:

Theorem 1.9: Let X be a compact affine manifold with a parallel volume form. Assume that its fundamental group is nilpotent. Then the monodromy representation is unipotent.

We are interested in affine nilmanifolds which also possess a hypercomplex structure. First, recall a definition of a complex nilmanifold.

Definition 1.10: Let G be a Lie group equipped with a left-invariant complex structure. **A complex nilmanifold** is a pair $(N = \Gamma \setminus G, I)$, where N is a nilmanifold and the complex structure I is obtained from the corresponding left-invariant complex structure on G.

An almost hypercomplex manifold X is a smooth manifold equipped with three endomorphisms I, J and K of the tangent bundle satisfying the quaternionic relations $I^2 = J^2 = K^2 = -\operatorname{Id}$ and IJ = K. When almost complex structures are integrable, the quadruple (X, I, J, K) is called **a hypercomplex manifold.**

Definition 1.11: Let G be a nilpotent Lie group with a left-invariant hypercomplex structure I, J, K and $\Gamma \subset G$ a cocompact lattice. Then $(N = \Gamma \setminus G, I, J, K)$ is called **a hypercomplex nilmanifold**.

M. Obata showed [Ob] that on a hypercomplex manifold M, there exists a unique torsion-free connection ∇ preserving the complex structures: $\nabla I = \nabla J = \nabla K = 0$. It is called **the Obata connection**.

It could be written in the following form

$$\nabla_X Y = \frac{1}{2}([X, Y] + I[IX, Y] - J[X, JY] + K[IX, JY]), \tag{1.1}$$

for any $X, Y \in TM$ [Sol].

The existence of a parallel volume form on a hypercomplex nilmanifold is guaranteed by the following theorem:

Theorem 1.12: [BDV, Theorem 3.2] Let $N = \Gamma \backslash G$ be a hypercomplex nilmanifold, $n = \dim_{\mathbb{C}} G$. Then G admits a left-invariant, non-zero, holomorphic section Ω of the canonical bundle $\Lambda^{n,0}G$. Moreover, $\nabla \Omega = 0$, where ∇ is the Obata connection.

1.3 ℍ-solvable Lie algebras and algebraic holonomy

An operator I on a real Lie algebra \mathfrak{g} is called a **complex structure operator** if $I^2 = -\operatorname{Id}$ and the $\sqrt{-1}$ -eigenspace $\mathfrak{g}^{1,0}$ is a Lie subalgebra in the complexification $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \otimes \mathbb{C}$.

Remark 1.13: Take an endomorphism $I \in \text{End } \mathfrak{g}$, $I^2 = -\text{Id}$, and extend it to the left-invariant almost complex structure on the Lie group G. Then I is complex if and only if $[\mathfrak{g}^{1,0},\mathfrak{g}^{1,0}] \subset \mathfrak{g}^{1,0}$. In other words, this definition is compatible with Definition 1.10.

A hypercomplex structure on a Lie algebra \mathfrak{g} is a triple of complex structure operators I, J and K on \mathfrak{g} satisfying the quaternionic relations.

Definition 1.14: Let \mathfrak{g} be a nilpotent hypercomplex Lie algebra. Define inductively \mathbb{H} -invariant Lie subalgebras:

$$\mathfrak{g}_i^{\mathbb{H}} := \mathbb{H}[\mathfrak{g}_{i-1}^{\mathbb{H}}, \mathfrak{g}_{i-1}^{\mathbb{H}}],$$

where $\mathfrak{g}_0^{\mathbb{H}} = \mathfrak{g}$ and

$$\mathfrak{g}_1^{\mathbb{H}}:=\mathbb{H}[\mathfrak{g},\mathfrak{g}]=[\mathfrak{g},\mathfrak{g}]+I[\mathfrak{g},\mathfrak{g}]+J[\mathfrak{g},\mathfrak{g}]+K[\mathfrak{g},\mathfrak{g}].$$

It is natural to study if $\mathfrak{g}_1^{\mathbb{H}}$ is a proper subalgebra in \mathfrak{g} .

A hypercomplex nilpotent Lie algebra \mathfrak{g} is called \mathbb{H} -solvable if the following sequence terminates for some $k \in \mathbb{Z}_{>0}$:

$$\mathfrak{g}_1^{\mathbb{H}} \supset \mathfrak{g}_2^{\mathbb{H}} \supset \dots \supset \mathfrak{g}_{k-1}^{\mathbb{H}} \supset \mathfrak{g}_k^{\mathbb{H}} = 0.$$
 (1.2)

Conjecture 1.15: Let (\mathfrak{g}, I, J, K) be a nilpotent hypercomplex Lie algebra. Then it is \mathbb{H} -solvable.

Let $(N = \Gamma \backslash G, I, J, K)$ be a hypercomplex nilmanifold. Consider a complex nilmanifold (N, L) with a general complex structure L = aI + bJ + cK, where $(a, b, c) \in S^2$, obtained from the hypercomplex structure. It is natural to try to describe complex submanifolds in (N, L). We solved this problem in the case of complex curves under the additional assumption on the Lie algebra $\mathfrak{g} = \mathrm{Lie}(G)$ [Gor]. Precisely, if the corresponding Lie algebra \mathfrak{g} is \mathbb{H} -solvable, then there are no complex curves in a complex nilmanifold (N, L) for the general complex structure L.

In this work, we show that in the case when a hypercomplex nilmanifold $\Gamma \backslash G$ admits the flat Obata connection, then $\mathfrak{g} = \operatorname{Lie}(G)$ is \mathbb{H} -solvable. The main argument we use relies on the notion of an algebraic holonomy group, which we define below (Definition 3.19).

Example 1.16: It was shown in [DF] that all 8-dimensional hypercomplex nilpotent Lie algebras are Obata-flat. However, in the same paper, I. Dotti and A. Fino presented an example of the 3-step nilpotent hypercomplex Lie algebra of dimension 12 which has non-zero curvature of the Obata connection.

Let \mathfrak{g} be a nilpotent Lie algebra over the field \mathbb{R} . A rational structure in \mathfrak{g} is a subalgebra $\mathfrak{g}_{\mathbb{Q}} \subset \mathfrak{g}$ defined over the rational numbers such that $\mathfrak{g}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{R} = \mathfrak{g}$. Assume that a Lie algebra \mathfrak{g} admits a rational structure $\mathfrak{g}_{\mathbb{Q}}$. By [CG] it happens if and only if there exists a nilmanifold $\Gamma \backslash G$ such that $\mathfrak{g}_{\mathbb{Q}} := \operatorname{span}_{\mathbb{Q}} \langle \log \Gamma \rangle$.

In the case when a Lie group G admits a left-invariant hypercomplex structure with the flat Obata connection, we prove that $\mathfrak{g}_i^{\mathbb{H}}$ is a proper subalgebra of \mathfrak{g}_{i-1} using the following approach.

A connection on a Lie algebra \mathfrak{g} is an \mathbb{R} -linear map

$$\nabla: \mathfrak{g} \longrightarrow \mathfrak{g}^* \otimes \mathfrak{g}$$

such that $X \mapsto \nabla_X \in \text{End}(\mathfrak{g})$. Note that a connection on a Lie algebra \mathfrak{g} is the same as a left-invariant connection on a Lie group G (Claim 3.6). This notion can be generalized to arbitrary G-equivariant vector bundle on a Lie group G, giving the algebraic version of an equivariant connection (see (3.1)).

The curvature tensor R of the connection ∇ defined in the following way:

$$R(X,Y) = [\nabla_X, \nabla_Y] - \nabla_{[X,Y]},$$

where $X, Y \in \mathfrak{g}$. The connection ∇ is called **flat** if R = 0. Notice that left invariant flat connections on G are equivalent to the representations of a Lie algebra on itself considered as a vector space (Claim 3.9).

In the end, we prove the following theorem:

Theorem 1.17: Let (N, I, J, K) be a hypercomplex nilmanifold with the flat Obata connection. Then the corresponding Lie algebra is \mathbb{H} -solvable.

2 Preliminaries: Nilpotent Lie groups and algebras

Let G be a Lie group. Define inductively the descending chain of normal subgroups of G

$$G = G_0 \supset G_1 \supset G_2 \supset \dots \supset G_k \supset \dots , \tag{2.1}$$

where $G_j := [G_{j-1}, G]$ is the subgroup generated by the elements of the form $xyx^{-1}y^{-1}$, $x \in G_{j-1}$, $y \in G$.

Definition 2.1: A Lie group G is called a **nilpotent Lie group** if (2.1) terminates for some $k \in \mathbb{Z}_{>0}$, i.e. $G_k = \{e\}$.

The descending series of a real Lie algebra $\mathfrak g$ is the chain of ideals defined as follows:

$$\mathfrak{g}_0 \supset \mathfrak{g}_1 \supset \cdots \supset [\mathfrak{g}_k, \mathfrak{g}] \supset \cdots,$$

 $\mathfrak{g}_0 = \mathfrak{g} \text{ and } \mathfrak{g}_k = [\mathfrak{g}_{k-1}, \mathfrak{g}].$

Definition 2.2: A Lie algebra \mathfrak{g} is called **nilpotent** if $\mathfrak{g}_s = 0$ for some $s \in \mathbb{Z}_{>0}$.

Let \mathfrak{g} be a real Lie algebra and \mathfrak{g}^* its dual space. For any $\alpha \in \mathfrak{g}^*$ and $\xi, \theta \in \mathfrak{g}$ the Chevalley–Eilenberg differential $d: \mathfrak{g}^* \longrightarrow \Lambda^2 \mathfrak{g}^*$ is defined as follows

$$d\alpha(\xi,\theta) = -\alpha([\xi,\theta]). \tag{2.2}$$

It extends to a finite-dimensional complex

$$0 \longrightarrow \mathfrak{g}^* \longrightarrow \Lambda^2 \mathfrak{g}^* \longrightarrow \cdots \longrightarrow \Lambda^{2n} \mathfrak{g}^* \longrightarrow 0$$
 (2.3)

by the Leibniz rule: $d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^{\tilde{\alpha}} \alpha \wedge d\beta$, where $\alpha, \beta \in \mathfrak{g}^*$. The identity $d^2 = 0$ follows from the Jacobi identity [CE].

Notice that the kernel of a closed 1-form is an ideal in the Lie algebra \mathfrak{g} . According to the definition of the Chevalley–Eilenberg differential, any closed 1-form $\alpha \in \mathfrak{g}^*$ vanishes on the commutator ideal $[\mathfrak{g},\mathfrak{g}] \subseteq \ker \alpha$.

Remark 2.3: The intersection of all kernels of closed 1-forms $\alpha \in \mathfrak{g}^*$

$$\Sigma := \bigcap_{\alpha \in \mathfrak{g}^*, d\alpha = 0} \ker \alpha$$

also forms an ideal in the Lie algebra \mathfrak{g} which coincides with the commutator ideal $[\mathfrak{g},\mathfrak{g}]$.

Recall that **a distribution** on a smooth manifold N is a sub-bundle $\Sigma \subset TN$. The distribution is called **involutive** if it is closed under the Lie bracket. **A leaf** of the distribution Σ is a maximal connected, immersed submanifold $L \subset N$ such that L tangent to Σ at each point. If Σ is involutive, then the set of all its leaves is called **a foliation**.

3 Algebraic holonomy group

3.1 Left equivariant vector bundles

Recall that a map $\varphi: X \longrightarrow Y$ of two manifolds X and Y with an action of a group G is called G-equivariant if $\varphi(g \cdot x) = g \cdot \varphi(x)$ for any $g \in G$, $x \in X$. A vector bundle $\pi: \mathbb{B} \longrightarrow X$ on a manifold X is called **a** G-equivariant vector bundle if its total space is equipped with a G-action such that the

projection π is a G-equivariant map and the action of G is linear on the fibers, i.e. $\mathbb{B}_x \longrightarrow \mathbb{B}_{qx}$ is a linear map for all $g \in G$.

Let G be a Lie group, $L_g: G \longrightarrow G$ the left translation $(h \mapsto g \cdot h)$, and B a finite-dimensional vector space with a basis $\{b_1, \ldots, b_n\}$.

Definition 3.1: Let $\mathfrak{g} = \text{Lie}(G)$ be a Lie algebra and $\mathfrak{g}^* = \text{Hom}(\mathfrak{g}, \mathbb{R})$ its dual. An algebraic connection is an \mathbb{R} -linear map

$$\nabla: B \longrightarrow \mathfrak{g}^* \otimes B$$
.

We use the following notation

$$\nabla(b) = \sum_{i} \theta_{i} \otimes A_{i}(b) \tag{3.1}$$

where $\theta_i \in \mathfrak{g}^*$ and $A_i \in \text{End}(B)$, $b \in B$.

Let $d: \mathfrak{g}^* \longrightarrow \Lambda^2 \mathfrak{g}^*$ be the Chevalley-Eilenberg differential. We define **the curvature** of the connection ∇ by the following formula

$$\Theta_{\nabla} := d\omega + \omega \wedge \omega, \tag{3.2}$$

where
$$\omega := \sum_i \theta_i \otimes A_i$$
, $d\omega = \sum_i d\theta_i \otimes A_i$ and $\omega \wedge \omega = \sum_{i < j} \theta_i \wedge \theta_j \otimes [A_i, A_j]$.

Consider a vector bundle $\pi: \mathbb{B} \longrightarrow G$ on a Lie group G. From now we denote the fiber at the identity $e \in G$ by $\mathbb{B}_e = B$.

Definition 3.2: A left equivariant vector bundle $\pi : \mathbb{B} \longrightarrow G$ over a Lie group G is a G-equivariant vector bundle π on a Lie group G with an action given by the left translations. A left invariant section s is determined by its value s_e at $e \in G$. This gives a map

$$\mathbb{L}: \mathbb{B}_e \longrightarrow H^0(G, \mathbb{B}), \quad s = \mathbb{L}(s_e),$$

$$\mathbb{L}(s_e)(g) = s(g) := (L_g)_*(s_e).$$
(3.3)

identified the fiber \mathbb{B}_e with the space $H^0(G,\mathbb{B})$ of all G-invariant sections of \mathbb{B} .

Remark 3.3: All tensor powers of a *G*-equivariant vector bundle are *G*-equivariant.

Claim 3.4: The category of left equivariant vector bundles on a Lie group G is equivalent to the category of vector spaces.

Proof: The group G acts on itself by the left multiplication freely and transitively. Any left equivariant bundle \mathbb{B} is trivial as a vector bundle and there is a basis of left invariant sections $\{s_1, \ldots, s_n\}$, such that $s_i = \mathbb{L}(b_i)$. For each left equivariant vector bundle the vector space of its left invariant sections gives a functor to the category of vector spaces. Conversely, given a vector space, consider it as a space of sections evaluated at the identity $e \in G$, and then extend them via all left translations (3.3) to obtain a left equivariant vector bundle.

Now we are able to give a definition of an invariant connection on the bundle \mathbb{B} . We use an algebraic connection (3.1) on the fiber $B = \mathbb{B}_e$.

Definition 3.5: Let $\pi : \mathbb{B} \longrightarrow X$ be a G-equivariant vector bundle over a manifold X. A connection ∇ on \mathbb{B} is called **equivariant** if it defines a left invariant differential operator

$$\nabla (\mathbb{L}(s_e)) = \mathbb{L}((\nabla s)_e), \tag{3.4}$$

where ∇s is understood as a section of the left-equivariant bundle $\Lambda^1(G) \otimes \mathbb{B}$, \mathbb{L} is the left-translation map defined in (3.3), and $(\nabla s)_e$ means the value of the section in $e \in G$.

Definition 3.7: Let $\theta \in \mathfrak{g}^*$ and $b \in B$. Define a map

$$d_{\nabla}: \mathfrak{g}^* \otimes_{\mathbb{R}} B \longrightarrow \Lambda^2 \mathfrak{g}^* \otimes_{\mathbb{R}} B$$

by the following formula:

$$d_{\nabla}(\theta \otimes b) = d\theta \otimes b - \theta \wedge \nabla b. \tag{3.5}$$

This defines the curvature of an algebraic connection

$$\Theta = d_{\nabla}^2 : B \longrightarrow \Lambda^2 \mathfrak{g}^* \otimes B.$$

As in (3.2) one can also write $\Theta = d\omega + \omega \wedge \omega$.

Claim 3.8: The curvature of a left equivariant connection \mathbb{V} on a G-equivariant bundle $\pi: \mathbb{B} \longrightarrow X$ is an \mathbb{R} -linear map $\Theta_{\mathbb{V}}: \mathbb{B} \longrightarrow \Lambda^2(X) \otimes \mathbb{B}$ given by $\Theta_{\mathbb{V}}(s) := \mathbb{L}(d^2_{\nabla}(s_e))$.

We say that bundle \mathbb{B} is **flat** if $\Theta_{\mathbb{W}} = 0$.

Claim 3.9: Let $\pi: \mathbb{B} \longrightarrow G$ be a left equivariant vector bundle over a Lie group. Then the bundle is flat if and only if $\nabla_X: \mathfrak{g} \longrightarrow \operatorname{End}(B)$ is a Lie algebra representation.

Proof: Assume that $\nabla : \mathfrak{g} \longrightarrow B^* \otimes B \cong \operatorname{End}(B)$ is a representation and the map is given as follows $X \mapsto \nabla_X$. Then for any $X, Y \in \mathfrak{g}$ we have $[\nabla_X, \nabla_Y] - \nabla_{[X,Y]} = 0$, hence by Claim 3.8 the curvature vanishes. Conversely, if $\Theta_{\overline{\mathbb{W}}} = 0$, then $\Theta_{\nabla} = [\nabla_X, \nabla_Y] - \nabla_{[X,Y]} = 0$, hence ∇_X is a representation.

Consider a left-equivariant vector bundle $\pi: \mathbb{B} \longrightarrow G$ over a Lie group G. Let Γ be a discrete subgroup of a Lie group G which acts on G from the left and $q: G \longrightarrow \Gamma \backslash G$ the quotient map. Denote by \mathbb{B}_{Γ} the induced vector bundle $\pi_{\Gamma}: \mathbb{B}_{\Gamma} \longrightarrow \Gamma \backslash G$ on a manifold $\Gamma \backslash G$ obtained by taking the fiberwise quotients by Γ such that the following diagram is commutative

$$\mathbb{B} \xrightarrow{q_{\Gamma}} \mathbb{B}_{\Gamma} \\
\downarrow^{\pi} \qquad \downarrow^{\pi_{\Gamma}} \\
G \xrightarrow{q} \Gamma \backslash G.$$

Given a vector bundle (\mathbb{B}, \mathbb{W}) with an equivariant connection \mathbb{W} , we denote by \mathbb{W}^{Γ} the induced connection on a bundle \mathbb{B}_{Γ} .

3.2 Algebraic holonomy group

Recall that there is a bijection between the set of isomorphism classes of flat vector bundles over X and the set of conjugacy classes of homomorphisms $\varphi : \pi_1(X) \longrightarrow \operatorname{GL}(\mathbb{R}^n)$ (It is called "the Riemann-Hilbert correspondence", see e.g. [OV]).

Let $p: E \longrightarrow M$ be a vector bundle over a manifold M and $\delta: [0,1] \longrightarrow M$ a smooth path in M. An equation $\nabla_{\dot{\delta}(t)} v = 0$ defines a parallel transport

of a section v of $E_{|_{\delta(t)}}$ along the curve δ . The parallel transport along a loop $\delta: [0,1] \longrightarrow M$ such that $x = \delta(0) = \delta(1)$ gives an endomorphism $p_{\delta} \in \operatorname{End}(E_x)$ of the fiber E_x .

Definition 3.10: Let $\nabla : E \longrightarrow \Lambda^1(M) \otimes E$ be a connection on a vector bundle $\pi : E \longrightarrow G$. **The holonomy group** of the connection ∇ is the group of linear transformations of a fiber E_x given by all parallel translations along all smooth loops based at x:

$$\mathcal{Hol}_{\nabla} = \{ p_{\delta} \mid p_{\delta} \in \operatorname{End}(E_x) \}.$$

Definition 3.11: The holonomy group of a flat connection ∇ is called **the** monodromy group of ∇ .

When G is a Lie subgroup $\mathrm{GL}(n,K), K=\mathbb{R}$ or \mathbb{C} , its Lie algebra is canonically identified with the corresponding Lie subalgebra in $\mathrm{Mat}(n,K)$. In this case, recall that the exponential map $\exp:\mathfrak{g}\longrightarrow G$ is defined as follows

$$X \mapsto e^{tX} = \operatorname{Id} + \frac{t}{1!}X + \frac{t^2}{2!}X^2 + \dots + \frac{t^k}{k!}X^k + \dots, \ t \in \mathbb{R}, X \in \mathfrak{g},$$
 (3.6)

and its local inverse is given by a logarithm

$$1 + \gamma \mapsto \log(1 + \gamma) = \gamma - \frac{\gamma^2}{2} + \frac{\gamma^3}{3} - \dots + (-1)^{k+1} \frac{\gamma^k}{k} + \dots, \tag{3.7}$$

for all $\gamma \in U \subset G$ in a sufficiently small open neighborhood U of the identity $e \in G$.

Let G be a simply connected nilpotent Lie group (it is diffeomorphic to a Euclidean space \mathbb{R}^n [VGO, Corollary 2, p. 53]) and Γ a cocompact lattice in G, i.e. the left quotient $\Gamma \backslash G$ is a nilmanifold.

Theorem 3.12: Let (\mathbb{B}, \mathbb{W}) be a flat left equivariant vector bundle over a nilpotent Lie group G, Γ a cocompact lattice and $(\mathbb{B}_{\Gamma}, \mathbb{W}^{\Gamma})$ the corresponding flat bundle on the nilmanifold $\Gamma \backslash G$. Then the monodromy group $\mathcal{Hol}_{\mathbb{W}^{\Gamma}}$ of the connection \mathbb{W}^{Γ} is generated by $e^{\log \gamma}$, where $\gamma \in \Gamma$.

Proof: Let $\delta:[0,1] \longrightarrow G$ be a 1-parametric subgroup in a Lie group G, such that $\delta(0) = e$, $\delta(1) \in \Gamma$ and tangent to a vector $X = \log \gamma$. Then the solution of a linear differential equation $\nabla_X b = 0$ along the curve γ is given by the exponents $e^{tX} = e^{t \log \gamma}$.

3.3 Maltsev completion

In this section, we assume that a group G is a finitely generated torsion-free nilpotent group.

Definition 3.13: [Mal] A nilpotent group G is called **Maltsev complete** if for each $g \in G$ and for all $n \in \mathbb{Z}_{>0}$ the equation $x^n = g$ has solutions in G.

Definition 3.14: [Mal] Let Γ be a subgroup of a Maltsev complete nilpotent group G. Then the set $\hat{\Gamma} = \{g \in G \mid g^n \in \Gamma, n \in \mathbb{Z}\} \subset G$ is called **the Maltsev completion** of a group Γ . In [Mal], the definition of the Maltsev completion was given over the field \mathbb{Q} but it could be done over any field k of characteristic zero.

Definition 3.15: [GH] Let k be a field of characteristic zero. **The Maltsev completion functor** \mathcal{M}_k is a functor from the category of finitely generated torsion-free nilpotent groups to the category of unipotent algebraic k-groups. If G is a finitely generated nilpotent group, then $\mathcal{M}_k(G) := \overline{\Phi(G)}$, where

$$\Phi: G \longrightarrow \operatorname{End}(B)$$

is a faithful unipotent representation and $\overline{\Phi(G)}$ is the Zariski closure of the image, which lies in the subgroup of upper-triangular matrices.

Remark 3.16: We will also denote the rational Maltsev completion of a group G by \hat{G} , as it was introduced in Definition 3.13.

Below we are listing some properties of the Maltsev completion [GH]:

Properties 3.17:

- 0. A connected nilpotent Lie group is Maltsev complete;
- 2. Let Γ be a subgroup of a nilpotent torsion-free Lie group G. By definition, $\mathcal{M}_k(\Gamma) \subset G$ is a Maltsev complete group. Its isomorphism class does not depend on G and the embedding of Γ to G;
- 3. The Maltsev completion $\mathcal{M}_k(\Gamma)$ is a minimal complete subgroup which contains Γ ;
- 4. The Maltsev completion $\mathcal{M}_k(\Gamma)$ is equipped with a natural structure of an algebraic group over k.

Claim 3.18: The functor $\mathcal{M}_{\mathbb{Q}}$ provides a bijection between the finite-dimensional unipotent representations of Γ over \mathbb{Q} and the finite-dimensional \mathbb{Q} -representations of $\hat{\Gamma}$, considered as an algebraic group:

$$\Gamma \xrightarrow{\Phi} \hat{\Gamma}$$

$$End(B)$$

Moreover, the image $\Phi(\Gamma)$ is Zariski dense in $\hat{\Phi}(\hat{\Gamma})$.

Proof: [GH]. ■

$$\mathcal{Hol}_{\nabla\!\!\!\!\!\!\nabla^{\Gamma}} \cong \Phi(\Gamma). \tag{3.8}$$

The notion of a holonomy group has a geometric nature. We define an algebraic version of it as follows.

Definition 3.19: Let B be a vector space, \mathfrak{g} a Lie algebra, and $\nabla : B \longrightarrow \mathfrak{g}^* \otimes B$ an algebraic connection (3.1). **An algebraic holonomy group** \mathcal{Hol}^a_{∇} is a subgroup of GL(B) generated by the matrix exponents:

$$\mathcal{H}\!o\ell_{\nabla}^{a} := \langle e^{t\nabla_{X}} \mid t \in \mathbb{R}, \text{ for all } X \in \mathfrak{g} \rangle.$$

Remark 3.20: We assume that an equivariant connection ∇ on a G-equivariant vector bundle B is flat. In this case, by Claim 3.9 the associated algebraic connection defines a representation of a Lie algebra $X \mapsto \nabla_X \in \text{End}(B)$.

Let G be a nilpotent group and \hat{G} its rational Maltsev completion, which is a rational algebraic group, i.e. it can be written as $\hat{G} = \operatorname{Spec}(A)$, where A is its ring of regular functions.

Remark 3.21: In Definition 3.15 we defined the functor of Maltsev completion \mathcal{M}_k with coefficients in a field k. The real Maltsev completion $\mathcal{M}_{\mathbb{R}}$ can be equivalently defined as follows:

$$\hat{G} \otimes_{\mathbb{Q}} \mathbb{R} := \operatorname{Spec}(A \otimes_{\mathbb{Q}} \mathbb{R}) = \mathscr{M}_{\mathbb{R}}(G).$$

Theorem 3.22: Let (\mathbb{B}, \mathbb{W}) be a flat left equivariant vector bundle over a nilpotent Lie group $G, \Gamma \subset G$ a cocompact lattice, and $(\mathbb{B}_{\Gamma}, \mathbb{W}^{\Gamma})$ the induced bundle over the nilmanifold $\Gamma \backslash G$. Suppose that $\nabla : B \longrightarrow \mathfrak{g}^* \otimes B$ is the algebraic connection associated with \mathbb{W} . Assume that the monodromy representation $\Phi : \Gamma \longrightarrow \operatorname{End}(B)$ is unipotent.

Then

$$\overline{\mathcal{H}ol_{\nabla^{\Gamma}}} = \mathcal{H}ol_{\nabla}^{a}, \tag{3.9}$$

where $\overline{\mathcal{H}ol_{\nabla^{\Gamma}}} \subseteq GL(B)$ denotes the Zariski closure of the monodromy group of $(\mathbb{B}_{\Gamma}, \nabla^{\Gamma})$.

Proof: Consider the chain of embeddings

where $\widehat{\mathcal{H}ol}_{\overline{\mathbb{W}}^{\Gamma}}$ is the rational Maltsev completion of the group $\mathcal{H}ol_{\overline{\mathbb{W}}^{\Gamma}}$.

Notice that $\mathcal{H}ol_{\mathbb{W}^{\Gamma}} \cong \Phi(\Gamma)$ is Zariski dense in $\widehat{\mathcal{H}ol_{\mathbb{W}^{\Gamma}}}$ by [BrVer, Lemma 3.11] and $\widehat{\mathcal{H}ol_{\mathbb{W}^{\Gamma}}}$ is Zariski dense in $\widehat{\mathcal{H}ol_{\mathbb{W}^{\Gamma}}} \otimes_{\mathbb{Q}} \mathbb{R}$. Hence,

$$\overline{\mathcal{H}ol_{\mathbf{\nabla}^{\Gamma}}} = \widehat{\mathcal{H}ol_{\mathbf{\nabla}^{\Gamma}}} \otimes_{\mathbb{Q}} \mathbb{R}. \tag{3.11}$$

By Definition 3.19, the algebraic holonomy group $\mathcal{H}o\ell_{\nabla}^{a} \subseteq \widehat{\mathcal{H}o\ell_{\nabla}}^{\Gamma}$ is a real group, which also contains $\mathcal{H}o\ell_{\nabla}^{\Gamma}$. Therefore, it has to be isomorphic to $\widehat{\mathcal{H}o\ell_{\nabla}}^{\Gamma} \otimes_{\mathbb{Q}} \mathbb{R}$, because of the minimality of the real Maltsev completion.

Corollary 3.23: Let $\Gamma \setminus G$ be a hypercomplex nilmanifold with the flat Obata conection ∇^{Ob} (1.1) on TG. Then the action of the algebraic holonomy on $\mathfrak{g} = \operatorname{Lie}(G)$ is unipotent.

Proof: Let ∇ be a flat algebraic connection on the Lie algebra $\mathfrak{g} = T_e G$, associated with ∇^{Ob} . Define the action $\mathcal{H}\!o\ell^a_{\nabla} \times \mathfrak{g} \longrightarrow \mathfrak{g}$ on \mathfrak{g} as follows:

$$(e^{\nabla_X}, Y) \mapsto e^{\nabla_X} \cdot Y, \tag{3.12}$$

where $X, Y \in \mathfrak{g}$. By Theorem 1.9, the monodromy representation $\mathcal{H}ol_{\nabla^{\Gamma}}$ is unipotent. From Theorem 3.22 it follows that $\mathcal{H}ol_{\nabla^{\Gamma}}$ is Zariski dense in $\mathcal{H}ol_{\nabla}^a$. Hence $\mathcal{H}ol_{\nabla}^a$ also unipotent. \blacksquare

3.4 Unipotent holonomy group and ℍ-solvability

Let (G, I, J, K) be a nilpotent Lie group with a left-invariant hypercomplex structure. In what follows we denote by ∇ the Obata connection (1.1), which we assumed to be flat, by \mathcal{Hol}_{∇} and \mathcal{Hol}_{∇}^a its holonomy (monodromy) and algebraic holonomy groups respectively.

Let \mathfrak{g} be a nilpotent hypercomplex Lie algebra. Consider the smallest \mathbb{H} -invariant subspace of \mathfrak{g} containing the commutator ideal $[\mathfrak{g}, \mathfrak{g}]$:

$$\mathfrak{g}_1^{\mathbb{H}} := \mathbb{H}[\mathfrak{g}, \mathfrak{g}] = [\mathfrak{g}, \mathfrak{g}] + I[\mathfrak{g}, \mathfrak{g}] + J[\mathfrak{g}, \mathfrak{g}] + K[\mathfrak{g}, \mathfrak{g}].$$

Since $\mathbb{H}[\mathfrak{g},\mathfrak{g}]$ contains the commutator ideal, it is an ideal, hence a subalgebra of \mathfrak{g} .

Define inductively the H-invariant Lie subalgebras:

$$\mathfrak{g}_i^{\mathbb{H}} := \mathbb{H}[\mathfrak{g}_{i-1}^{\mathbb{H}}, \mathfrak{g}_{i-1}^{\mathbb{H}}], i \in \mathbb{Z}_{\geqslant 2}.$$

Claim 3.24: Let (\mathfrak{g}, I, J, K) be a hypercomplex nilpotent Lie algebra with the flat Obata connection. Then $\nabla_X Y \in \mathfrak{g}_{i+1}^{\mathbb{H}}$ for any $X, Y \in \mathfrak{g}_i^{\mathbb{H}}$.

Proof: Suppose that $X, Y \in \mathfrak{g}_i^{\mathbb{H}}$. Then by (1.1) $\nabla_X Y \in \mathfrak{g}_{i+1}^{\mathbb{H}}$.

For each $i \in \mathbb{Z}_{\geq 0}$, $X \in \mathfrak{g}_i^{\mathbb{H}}$, the elements $e^{\nabla_X} \in \operatorname{End}(\mathfrak{g}_i^{\mathbb{H}})$ generate the algebraic holonomy group, associated to the Lie subalgebra $\mathfrak{g}_i^{\mathbb{H}}$:

$$\mathcal{H}ol_{\nabla^i}^a = \langle e^{t\nabla_X} \mid t \in \mathbb{R}, X \in \mathfrak{g}_i^{\mathbb{H}} \rangle.$$

Remark 3.25: Notice that $\mathcal{Hol}_{\nabla^i}^a$ acts on $\mathfrak{g}_i^{\mathbb{H}}$ as in (3.12) and it could happen that it is not a subgroup or even a subset of the algebraic holonomy group \mathcal{Hol}_{∇}^a . Indeed, the action of ∇_X on $\mathfrak{g}_i^{\mathbb{H}}$ could be trivial, but act non-trivially on \mathfrak{g} for some $X \in \mathfrak{g}_i^{\mathbb{H}}$. However, the action of $\mathcal{Hol}_{\nabla^i}^a$ on $\mathfrak{g}_i^{\mathbb{H}}$ coincides with the action of the corresponding subgroup of \mathcal{Hol}_{∇}^a . Therefore, it is unipotent on $\mathfrak{g}_i^{\mathbb{H}}$ if \mathcal{Hol}_{∇}^a is unipotent.

The following theorem is the main result of this paper.

Theorem 3.26: Let (\mathfrak{g}, I, J, K) be a hypercomplex nilpotent Lie algebra with the flat Obata connecton. Assume that the algebraic monodromy representation is unipotent. Then \mathfrak{g} is \mathbb{H} -solvable.

Proof: The action of $\mathcal{Hol}_{\nabla^i}^a$ on the Lie subalgebra $\mathfrak{g}_i^{\mathbb{H}}$ defined as in (3.12). By the assumption, the representation of the algebraic holonomy on \mathfrak{g} is unipotent, hence, by Remark 3.25, the action of $\mathcal{Hol}_{\nabla^i}^a$ is unipotent on the $\mathfrak{g}_i^{\mathbb{H}}$.

Let $\varphi : \mathcal{Hol}_{\nabla^i}^a \longrightarrow \operatorname{End}((\mathfrak{g}_i^{\mathbb{H}})^*)$ be the action of the algebraic holonomy group on the dual Lie algebra $(\mathfrak{g}_i^{\mathbb{H}})^*$. Every unipotent representation of a Lie algebra has an invariant vector. This implies the existence of a non-zero Obata-parallel 1-form for $\alpha \in \Lambda^1(\mathfrak{g}_i^{\mathbb{H}})^*$. Again, we consider the holonomy action on $(\mathfrak{g}_i^{\mathbb{H}})^*$ and for each i we obtain a non-zero parallel (hence, closed¹) 1-form $\alpha_i \in \Lambda^1(\mathfrak{g}_i^{\mathbb{H}})^*$.

The intersection of kernels $\Sigma_{\alpha_i} = \ker \alpha_i \cap \ker I\alpha_i \cap \ker J\alpha_i \cap \ker K\alpha_i$ gives an \mathbb{H} -invariant foliation, which contains $\mathfrak{g}_i^{\mathbb{H}}$ as a proper subspace (Remark 2.3). This implies $\mathfrak{g}_{i+1}^{\mathbb{H}} \subsetneq \mathfrak{g}_i^{\mathbb{H}}$. The sequence $\mathfrak{g} \subsetneq \mathfrak{g}_1^{\mathbb{H}} \subsetneq \ldots$ terminates in finitely many steps since \mathfrak{g} is finite-dimensional. \blacksquare

Corollary 3.27: Let (N, I, J, K) be a hypercomplex nilmanifold with flat Obata connection ∇ . Then the Lie algebra $\mathfrak{g} = \text{Lie}(G)$ is \mathbb{H} -solvable.

Proof: By Corollary 3.23, the representation of the algebraic holonomy on \mathfrak{g} is unipotent. \blacksquare

References

- [A] H. Abels, Properly discontinuous groups of affine transformations: a survey, Geom. Ded. 87 (2001), no. 1-3, 309–333. (Cited on page 2.)
- [AMS] H. Abels, G. A. Margulis, G. A. Soifer, *The Auslander conjecture for dimension less then* 7, arXiv:2011.12788 (Cited on page 3.)
- [AMS2] H. Abels, G. A. Margulis, G. A. Soifer, The linear part of an affine group acting properly discontinuously and leaving a quadratic form invariant, (English summary) Geom. Dedicata 153 (2011), 1–46. (Cited on page 3.)
- [Aus] L. Auslander, The structure of complete locally affine manifolds, Topology 3 1964 suppl. 1 131–139 (Cited on page 3.)

¹For any torsion-free connection ∇ and any 1-form β we have the equality $d\beta = \text{Alt}(\nabla\beta)$, where Alt is the skew-symmetrization map.

- [BDV] M. L. Barberis, I. G. Dotti, M. Verbitsky, Canonical bundles of complex nilmanifolds, Math. Res. Lett. 16 (2009), no. 2, 331–347. (Cited on page 5.)
- [Bieb] L. Bieberbach, Über die Bewegungsgruppen der Euklidischen Raume, Mathematische Annalen, 70 (3) (1911): 297–336 (Cited on page 3.)
- [BrVer] M. Brandenbursky, M. Verbitsky, Non-commutative Barge-Ghys quasimorphisms, https://doi.org/10.48550/arXiv.2212.12958 (Cited on page 15.)
- [CE] C. Chevalley, S. Eilenberg, "Cohomology theory of Lie groups and Lie algebras", Transactions of the American Mathematical Society, 63 (1): 85–124, 1948 (Cited on page 8.)
- [CG] L. Corwin, F. P. Greenleaf, Representations of Nilpotent Lie Groups and Their Applications. Part I, Basic Theory and Examples, Cambridge Univ. Press, Cambridge, UK, 1990 (Cited on page 6.)
- [DGMS] P. Delign, P. Griffiths, J. Morgan, D. Sullivan, *Real homotopy theory of Kähler manifolds*, In: Invent. Math. **29** (1975), no. 3, 245–274. (Not cited.)
- [DF] I. G. Dotti, A. Fino, Hypercomplex nilpotent Lie groups, Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000), 310–314, Contemp. Math., 288, Amer. Math. Soc., Providence, RI, 2001. differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000), 310–314, Contemp. Math., 288, Amer. Math. Soc., Providence, RI, 2001. (Cited on page 6.)
- [FG] D. Fried, W. Goldman, Three-dimensional affine crystallographic groups, Adv. Math. 47 (1983), 1–49. (Cited on page 3.)
- [FGH] D. Fried, W. Goldman, M. W. Hirsch Affine manifolds with nilpotent holonomy, Commentarii Mathematici Helvetici, volume 56, pages 487-523, 1981 (Cited on pages 2 and 4.)
- [Gol] W. Goldman, Two papers which changed my life: Milnor's seminal work on flat manifolds and flat bundles, "Frontiers in Complex Dynamics: In Celebration of John Milnor's 80th Birthday" A. Bonifant, M. Lyubich,

- S. Sutherland (eds.), Princeton Univ. Press) DG:1108.0216 (Cited on page 2.)
- [Gol1] W. Goldman, Geometric Structures on Manifolds, Graduate Studies in Mathematics, Volume: 227; 2022; 409 pp, AMS (Cited on page 3.)
- [GK] W. Goldman, Y. Kamishima, The fundamental group of a compact flat Lorentz space form is virtually polycyclic, Journal of Differential Geometry. 19 (1): 233–240, 1984 (Cited on page 3.)
- [Gor] Yu. Gorginian, Complex curves in hypercomplex nilmanifolds with H-solvable Lie algebras, arXiv:2207.12561 (Cited on page 6.)
- [GH] Grunewald, Fritz (D-BONN); O'Halloran, Joyce (1-WIM) Nilpotent groups and unipotent algebraic groups, J. Pure Appl. Algebra 37 (1985), no. 3, 299–313. (Cited on pages 13 and 14.)
- [Ob] M. Obata, Affine connections on manifolds with almost complex, quaternionic or Hermitian structure, Jap. J. Math., 26 (1955), 43-79. (Cited on page 4.)
- [OV] L. Ornea, M. Verbitsky, *Principles of Locally Conformally Kahler Geometry*, arXiv:2208.07188 (Cited on pages 3 and 11.)
- [Klin] B. Klingler, Chern's conjecture for special affine manifolds, Annals of Mathematics 186 (2017), 1-27. (Cited on page 2.)
- [KS] B. Kostant, D. Sullivan, The Euler characteristic of a compact affine space form is zero, Bull. Amer. Math. Soc. 81 (1975) (Cited on page 2.)
- [Mal] A. I. Maltsev, On a class of homogeneous spaces, Izv. Akad. Nauk. Armyan. SSSR Ser. Mat. 13 (1949), 201-212. (Cited on page 13.)
- [Mar] L. Markus, Cosmological models in differential geometry, Mimeographed Notes, Univ. of Minnesota (1962), p. 58. (Cited on page 2.)
- [Sh] H. Shima, *The geometry of Hessian structures*, World Scientific Publishing Co. Pte. Ltd., Singapore, (2007) (Cited on page 2.)
- [Sol] A. Soldatenkov, *Holonomy of the Obata connection on SU(3)*, nt. Math. Res. Notices (2012), Vol. 2012 (15), 3483-3497 (Cited on page 5.)

[VGO] È. B. Vinberg, V. V. Gorbatsevich, A. L. Onishchik, Structure of Lie groups and Lie algebras, Lie groups and Lie algebras – 3, Itogi Nauki i Tekhniki. Ser. Sovrem. Probl. Mat. Fund. Napr., 41, VINITI, Moscow, 1990, 5–253 (Cited on page 12.)

Yulia Gorginyan

INSTITUTO NACIONAL DE MATEMÁTICA PURA E APLICADA (IMPA)

ESTRADA DONA CASTORINA, 110

Jardim Botânico, CEP 22460-320

RIO DE JANEIRO, RJ - BRASIL

ALSO:

LABORATORY OF ALGEBRAIC GEOMETRY,

NATIONAL RESEARCH UNIVERSITY (HSE),

DEPARTMENT OF MATHEMATICS, 6 USACHEVA STR.

Moscow, Russia

iuliia.gorginian@impa.br