

A Fixed-Parameter Tractable Algorithm for Counting Markov Equivalence Classes with the same Skeleton ^{*}

Vidya Sagar Sharma[†]

Causal DAGs (also known as Bayesian networks) are a popular tool for encoding conditional dependencies between random variables. In a causal DAG, the random variables are modeled as vertices in the DAG, and it is stipulated that every random variable is independent of its ancestors conditioned on its parents. It is possible, however, for two different causal DAGs on the same set of random variables to encode exactly the same set of conditional dependencies. Such causal DAGs are said to be *Markov equivalent*, and equivalence classes of Markov equivalent DAGs are known as *Markov Equivalent Classes* (MECs). Beautiful combinatorial characterizations of MECs have been developed in the past few decades, and it is known, in particular that all DAGs in the same MEC must have the same “skeleton” (underlying undirected graph) and v-structures (induced subgraph of the form $a \rightarrow b \leftarrow c$).

These combinatorial characterizations also suggest several natural algorithmic questions. One of these is: given an undirected graph G as input, how many distinct Markov equivalence classes have the skeleton G ? Much work has been devoted in the last few years to this and other closely related problems. However, to the best of our knowledge, a polynomial time algorithm for the problem remains unknown.

In this paper, we make progress towards this goal by giving a fixed parameter tractable algorithm for the above problem, with the parameters being the treewidth and the maximum degree of the input graph G . The main technical ingredient in our work is a construction we refer to as *shadow*, which lets us create a “local description” of long-range constraints imposed by the combinatorial characterizations of MECs.

1 Introduction

A graphical model is used to graphically represent a set of conditional independence relations between random variables. In the literature, both directed and undirected variations have been utilized to model various types of dependency structures. In this paper, our focus is on studying graphical models represented by directed acyclic graphs (DAGs), also called the Bayesian graphical model, which effectively conveys conditional independence relations and causal influences between random variables through the use of directed acyclic graphs (Pearl, 2009). This model is well-studied and has been found extensive applications in many fields, such as material science (Ren et al., 2020), game theory (Kearns, Littman, and Singh, 2001), and biology (Friedman, 2004; Finegold and Drton, 2011).

In the Bayesian graphical model, a probability distribution over a set of random variables V is said to satisfy a DAG G with vertices V if and only if for every $w \in V$, w is independent of the set of all its non-descendants conditional on the set of its parents (the definitions of “parent” and “non-descendant” are provided in Section 2). The set of probability distributions that satisfy G is denoted by $\text{Markov}(G)$. Further, for disjoint $A, B, S \subseteq V$, G is said to *entail* that A is independent of B given S (written $G \models A \perp B | S$) if and only if A is independent of B given S in every probability distribution in $\text{Markov}(G)$. Two DAGs are said to be *Markov equivalent* if both entail the same set of such conditional independence relations. Verma and Pearl (1990) gave an elegant graphical characterization of this equivalence: two DAGs are Markov equivalent if and only if they have the same *skeleton* (underlying undirected graph) and the same set of v-structures (induced

^{*}Accepted to the Proceedings of the 38th Annual AAAI Conference on Artificial Intelligence (AAAI 2024)

[†]Tata Institute of Fundamental Research, Mumbai. Email: vidyasagartifr@gmail.com.

subgraphs of the form $a \rightarrow b \leftarrow c$). DAGs that are Markov equivalent to each other are said to belong to the same *Markov equivalence class* (MEC) (see Figure 1).




Figure 1: Markov equivalent DAGs and MECs with the same skeleton. D_1, D_2 , and D_3 are Markov equivalent, while D_4 is not equivalent to D_1, D_2 , and D_3 . The MEC $M_1 = \{D_1, D_2, D_3\}$ contains D_1, D_2 and D_3 , and its graphical representation is the union of D_1, D_2 , and D_3 . The MEC M_2 contains only D_4 , and its graphical representation matches D_4 . The MECs M_1 and M_2 share G as their skeleton, and in fact are the only MECs with skeleton G . Both M_1 and M_2 entail a conditional independence relation of the form $B \perp C \mid S$, where in M_1 , $S = \{C\}$, and in M_2 , $S = \emptyset$.

As an MEC consists of Markov equivalent DAGs, it uniquely represents the set of conditional independence relations represented by the DAGs it contains and is graphically represented by a partially directed graph which is the graphical union of the DAGs it contains. We treat an MEC and its graphical representation as the same. Since all the DAGs that belong to an MEC have the same skeleton and the set of v-structures, an MEC is uniquely determined by its skeleton and v-structures. [Andersson, Madigan, and Perlman \(1997\)](#) gave a necessary and sufficient condition for a partially directed graph to be an MEC (see [Theorem 2.18](#) below). For convenience, the graph representing an MEC is often considered synonymous with the MEC itself, and both are referred to as an ‘‘MEC’’.

[Meek \(1995\)](#) formulated rules to construct an MEC based on knowledge about the conditional independence relations among random variables. For a set of conditional independence relations involving a set of random variables V and represented by an MEC M , [Meek \(1995, p. 3\)](#) also showed that two random variables A and B are non-adjacent in the skeleton of M if and only if there exists $S \subseteq V \setminus \{A, B\}$ such that M entails that $A \perp B \mid S$. This implies that two MECs M_1 and M_2 sharing the same skeleton entail sets of conditional independence relations \mathcal{M}_1 and \mathcal{M}_2 , respectively, which have the following important relation: For two random variables A and B , \mathcal{M}_1 contains a conditional independent relation indicating that A is independent of B given some $S_1 \subseteq V \setminus \{A, B\}$ if and only if \mathcal{M}_2 also contains a conditional independence relation signifying that A is independent of B given some $S_2 \subseteq V \setminus \{A, B\}$ (S_1 and S_2 may not be same). In other words, MECs with the same skeleton can be related by an equivalence relation that has not just a graphical representation (i.e., that they have the same underlying undirected graph), but also a natural statistical one (the one given above).

This connection between MECs with the same skeleton motivates the problem of understanding this class. In particular, a natural question to ask is: how many MECs are in this class?

Our Contributions We now formalize the problem outlined above. Our input consists of a connected undirected graph G with n nodes. Our objective is to determine the count of MECs that have G as their skeleton. The primary contribution of this paper is the introduction of a fixed-parameter tractable (FPT) algorithm. This algorithm, when given an undirected graph G , computes the number of MECs whose skeletons match G . The algorithm’s parameters are the degree and the treewidth of the input undirected graph. (An algorithm is said to be FPT with respect to a parameter if there is a constant c and a computable function f such that the algorithm’s runtime on instances of size n for which the value of the parameter is k is bounded above by $f(k) \cdot n^c$: the crucial point here is that the degree of the polynomial in n does not depend upon the

parameter k (Cygan et al., 2015).) Our main result presents an algorithm capable of counting the MECs associated with an input undirected graph G with n nodes and having a degree of δ and a treewidth of k . This counting can be achieved in $O(n(2^{O(k^4\delta^4)} + n^2))$ time. Importantly, the runtime of our algorithm remains polynomially bounded when the parameters δ and k are both bounded above by constants. As an illustrative example, our algorithm demonstrates polynomial runtime for tree graphs with bounded degrees.

As of now, we do not know the precise computational complexity of this problem, and we consider our result as a first step towards a complete resolution of this question. This mirrors the situation of the problem of counting DAGs of an MEC, where initially an algorithm that was exponential in the degree of the graph was given by Ghassami et al. (2019) and then improved by Talvitie and Koivisto (2019) who gave a fixed parameter tractable algorithm for the problem. Finally, a polynomial algorithm was provided by Wienöbst, Bannach, and Liśkiewicz (2021). We hope that the techniques introduced in our paper will be useful in the further study for the problem.

Related Work The problem of counting MECs with a given number of nodes (instead of a skeleton) has received extensive attention in the literature. Gillispie and Perlman (2001) developed a computer program for computing the number of MECs with n nodes. Gillispie and Perlman (2002) created a computer program to enumerate Markov equivalence classes, studying class size distributions and the number of edges for graphs up to 10 vertices. They also observed that the ratio of DAGs to the number of MECs seems to asymptotically converge to around 3.7. Steinsky (2003) presented a recursive formula for counting Markov equivalence classes of size 1. Gillispie (2006) provided a recursive algorithm for counting MECs of any size. He, Jia, and Yu (2013) analyze the set of MECs with n nodes by constructing a Markov chain on the space of MECs and show that most edges of an MECs are directed. More recently, Schmid and Sly (2022) show that the expected ratio of the number of DAGs and the number of MECs approaches a positive constant when the number of nodes goes to infinity.

All of the previous results focused on the class of MECs with a given number of nodes, and not with a given skeleton. Radhakrishnan, Solus, and Uhler (2016) focused on counting MECs with the same skeleton. They classified MECs based on the number of v -structures present and derived a generating function for counting MECs. They experimentally demonstrated that the generating function varies for graphs with the same number of vertices. In subsequent work, Radhakrishnan, Solus, and Uhler (2018) delved further into the problem of counting MECs with the same skeleton. They explored generating functions for specific graph structures (e.g., path graphs, cycle graphs, star graphs, and bi-star graphs) and provided tight lower and upper bounds for the number of MECs in any tree.

We are not aware of any progress on the problem of counting MECs for general graphs, and to the best of our knowledge, this paper is the first to introduce a fixed-parameter tractable algorithm for this problem.

Technical Overview: For an undirected graph G , we address the problem of counting Markov Equivalence Classes (MECs) of G , i.e., counting MECs that have the skeleton G . A trivial brute force approach is to iterate through each possible directed acyclic graph (DAG) having the skeleton G . Since each DAG with the skeleton G is a member of a unique MEC of G , and each MEC of G is a nonempty set of DAGs with the skeleton G , we count the different MECs to which these DAGs belong. This approach yields the count of MECs with the skeleton G . However, due to the fact that the number of DAGs with the skeleton G is $2^{|E_G|}$ (E_G is the set of edges of G), the runtime of this approach becomes exponential in the size of G .

In this paper, we provide a fixed-parameter tractable (FPT) algorithm to count MECs of G such that the run time of our algorithm is exponential in terms of the parameters, degree, and treewidth of the input graph, but is polynomial in the size of the input graph, where the degree of the polynomial does not depend on the parameters.

We coin a new term “shadow” (Definition 2.36). A shadow of an MEC M on $X \subseteq V_M$ is a triple (O, P_1, P_2) , where O is the induced subgraph of M on X , $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ is a function that for any $(u, v), (x, y) \in E_O$, answers whether there exists a triangle-free path (Definition 2.23) of length three or more in M from (u, v) to (x, y) , and $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ (V_O is the vertex set of O) is a function that for any $((u, v), w) \in E_O \times V_O$, answers whether there exists a triangle-free path of length three or more in M from

(u, v) to w . A triangle-free path of a graph is a path of the graph such that for any three consecutive nodes a, b and c of the path, there is no edge in the graph with endpoints a and c . A triangle-free path starts from an edge (u, v) if the first two nodes of the path are u and v . A triangle-free path ends on an edge (x, y) if the last two nodes of the path are x and y . A triangle-free path ends on a vertex w if the last node of the path is w .

With a slight abuse of notation, for an undirected graph G , we define a triple (O, P_1, P_2) as a shadow of G if O is a partial MEC of G (a graph with skeleton G and obeys items 1 to 3 of Theorem 2.18, Definition 2.21), and $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ and $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ are two functions. We show that for any undirected graph G , for any $X \subseteq V_G$, summation of $|\text{MEC}(G, O, P_1, P_2)|$, the size of the set of MECs of G having shadow (O, P_1, P_2) , over all shadows of $G[X]$ equals to the number of MECs of G (Lemma 2.40). This reduces the problem of counting MECs of an undirected graph G into counting MECs of G with shadow (O, P_1, P_2) for each possible shadow (O, P_1, P_2) of $G[X]$ for some $X \subseteq V_G$. It is the idea of “shadow”, a nontrivial concise representation of an MEC, which is the key for the construction of the fixed-parameter tractable algorithm (alg. 5).

We provide a recursive algorithm (alg. 7) to solve the reduced problem. For this, we find a relation between the MECs of an undirected graph G and the MECs of an induced subgraph G' of G . We note that for an MEC M of G , $M[V_{G'}]$ may not be an MEC of G' . But, we found that there exists a unique MEC M' of G that has the same set of v-structure as $M[V_{G'}]$ (Lemma 2.42). We call the MEC M' as the projection of M on $V_{G'}$ (Definition 2.41). We find a graphical resemblance between an MEC M of G and its projection (Lemma 2.45) that if an edge is directed in the projection then it is also directed in the MEC.

For the recursion, we divide the graph using its tree decomposition. Suppose T is a tree decomposition of G and S_1 is the root node of T . We pick a neighbor S_2 of T . We cut the edge $S_1 - S_2$ of T such that we get two induced subtrees T_1 and T_2 with roots S_1 and S_2 , respectively, and representing the induced subgraphs G_1 and G_2 of G , respectively. From the tree decomposition properties, we have (a) G is the union of G_1 and G_2 , and (b) $I = V_{G_1} \cap V_{G_2}$ is a vertex separator of G . We start with finding a graphical relation between an MEC M of G and its projections M_1 and M_2 on V_{G_1} and V_{G_2} respectively. Let (O, P_1, P_2) be the shadow of M on $S_1 \cup S_2 \cup N(S_1 \cup S_2, G)$ ($N(X, G)$ is the set of neighbors of the vertices in X in G), (O_1, P_{11}, P_{12}) be the shadow of M_1 on $S_1 \cup N(S_1, G_1)$, and (O_2, P_{21}, P_{22}) be the shadow of M_2 on $S_2 \cup N(S_2, G_2)$. We find a graphical resemblance between the shadow of M , M_1 and M_2 . We find necessary conditions that are obeyed by the shadows (Lemma 3.2). We later show that the conditions are also sufficient (Lemma 3.34). That means, if we have shadows (O, P_1, P_2) of $G[S_1 \cup S_2 \cup N(S_1 \cup S_2, G)]$, (O_1, P_{11}, P_{12}) of $G_1[S_1 \cup N(S_1, G_1)]$, and (O_2, P_{21}, P_{22}) of $G_2[S_2 \cup N(S_2, G_2)]$ that obeys the necessary and sufficient conditions then for each pair of MECs (M_1, M_2) such that M_1 is an MEC of G_1 with shadow (O_1, P_{11}, P_{12}) and M_2 is an MEC of G_2 with shadow (O_2, P_{21}, P_{22}) there exists a unique MEC M of G with shadow (O, P_1, P_2) . This relation shows that if we have the knowledge of $|\text{MEC}(G_1, O_1, P_{11}, P_{12})|$ for each shadow (O_1, P_{11}, P_{12}) of $G_1[S_1 \cup N(S_1, G_1)]$, and $|\text{MEC}(G_2, O_2, P_{21}, P_{22})|$ for each shadow (O_2, P_{21}, P_{22}) of $G_2[S_2 \cup N(S_2, G_2)]$ then we can compute $|\text{MEC}(G, O, P_1, P_2)|$ for each shadow (O, P_1, P_2) of $G[S_1 \cup S_2 \cup N(S_1 \cup S_2, G)]$ (Lemma 3.63).

Lemma 3.63 provides us an approach to compute $|\text{MEC}(G, O, P_1, P_2)|$ for each shadow (O, P_1, P_2) of $G[S_1 \cup S_2 \cup N(S_1 \cup S_2, G)]$. But, for the recursion, we need to compute $|\text{MEC}(G, O, P_1, P_2)|$ for each shadow (O, P_1, P_2) of $G[S_1 \cup N(S_1, G)]$. Since a shadow of $G[S_1 \cup S_2 \cup N(S_1 \cup S_2, G)]$ contains more information than a shadow of $G[S_1 \cup N(S_1, G)]$, we can trim them to fulfill our purpose. We define the projection of shadow (Definition 3.64) for this. A shadow (O', P'_1, P'_2) of $G[S_1 \cup N(S_1, G)]$ is said to be a projection of a shadow (O, P_1, P_2) of $G[S_1 \cup S_2 \cup N(S_1 \cup S_2, G)]$ if O' is an induced subgraph of O , for all $((x, y), (u, v)) \in E_{O'} \times E_{O'}$, $P'_1((x, y), (u, v)) = P_1((x, y), (u, v))$, and for all $((u, v), w) \in E_{O'} \times V_{O'}$, $P'_2((u, v), w) = P_2((u, v), w)$. If a shadow (O', P'_1, P'_2) of $G[S_1 \cup N(S_1, G)]$ is a projection of a shadow (O, P_1, P_2) of $G[S_1 \cup S_2 \cup N(S_1 \cup S_2, G)]$ then if M is an MEC of G with shadow (O, P_1, P_2) then M has also (O', P'_1, P'_2) as its shadow. For each shadow (O', P'_1, P'_2) of $G[S_1 \cup N(S_1, G)]$, this leads us to count MECs of G with shadow (O', P'_1, P'_2) when for each shadow (O, P_1, P_2) of $G[S_1 \cup S_2 \cup N(S_1 \cup S_2, G)]$, we have the knowledge of $|\text{MEC}(G, O, P_1, P_2)|$ (Lemma 3.65).

For the recursion, we cut an edge of the tree decomposition of G to construct induced subgraphs G_1 and G_2 . Since the number of edges of the tree decomposition of G is linear in the size of G , the number

of times we recursively call our algorithm is linear in the size of the input graph. The number of possible shadows is exponential in the treewidth and the degree of the input graph. The computation of the number of MECs having a specific shadow also takes time exponential in terms of the treewidth and the degree of the input graph. The overall time complexity (Theorem 5.3) of the algorithm is $O(n(2^{O(k^4\delta^4)} + n^2))$, where the parameters k and δ are the treewidth and the degree of the input graph, respectively, and n is the number of nodes of the input graph.

Paper Structure: In Section 2, we define the terminologies used in this paper. In Section 2.1, we go through basic graph theory terminologies (eg: path, cycle, chordal graphs, chain graphs, the union of graphs, tree decompositions, LBFS ordering, skeleton, v-structure), Markov equivalence class, and related results. In Section 2.2, we define the terminologies that are newly introduced in this paper, eg: Markov union of graphs, partial MECs, triangle-free path, a shadow of an MEC, projection of an MEC, and related results. In Section 3, we formally define the problem, and provide the necessary and sufficient conditions for the shadows (Sections 3.1 and 3.4). In Section 3.3, we provide a modified version of LBFS algorithm. In Section 3.2, we define derived path functions and extension of shadows. In Section 3.5, we provide a formula to compute $|\text{MEC}(G, O, P_1, P_2)|$. In Section 4, we provide a fixed parameter tractable algorithm to count MECs of an undirected graph. Section 5 analyzes the time complexity of the algorithms discussed in this paper. In Section 6, we discuss the open problems.

2 Preliminary

2.1 Old existing definitions and related results

The following definitions and observations are well-studied terms and results in graph theory and graphical models.

Basic graph terminologies: A graph G is a pair (V, E) , where V is said to be the set of vertices of G , and $E \subseteq V \times V$ is said to be the set of edges of G . For $u, v \in V$, if $(u, v), (v, u) \in E$, then we say there is an undirected edge between u and v , denoted as $u - v$. For $u, v \in V$, if $(u, v) \in E$ and $(v, u) \notin E$, then we say there is a directed edge from u to v , denoted as $u \rightarrow v$. For $u, v \in V$, when we say that (u, v) is an edge in G , it means that either there exists an undirected edge $u - v$ in G or there exists a directed edge $u \rightarrow v \in G$.

For a graph G , we denote V_G as the set of vertices of G , and E_G as the set of edges of G . A partially directed graph is a graph in which some edges are directed and the remaining edges are undirected. If all the edges of a graph are undirected, then the graph is undirected. If all the edges of a graph are directed, then the graph is directed. u is said to be a neighbor of v in a graph G if either $(u, v) \in E_G$ or $(v, u) \in E_G$. $N(X, G)$ is the set of neighbors of X in the graph G , i.e., $N(X, G) = \{v : \exists u \in X \text{ and either } (u, v) \in E_G \text{ or } (v, u) \in E_G\}$. The degree of a node is the number of neighbors of the node. The degree of a graph is the maximum degree of its nodes.

Definition 2.1 (Path, chord of a path, descendant, parent). For a graph G , a sequence $P = (u_1, u_2, \dots, u_{l-1}, u_l)$ of distinct vertices is said to be a **path** from u_1 to u_l if for all $1 \leq i < l$, $(u_i, u_{i+1}) \in E_G$ (i.e., either $u_i - u_{i+1} \in E_G$ or $u_i \rightarrow u_{i+1} \in E_G$). The **length of the path** is denoted by l . P is said to be an **undirected path** of G if for all $1 \leq i < l$, $u_i - u_{i+1} \in E_G$. Otherwise, if the path contains directed edges, it is called a **directed path** of G . For an undirected path $P = (u_1, u_2, \dots, u_{l-1}, u_l)$ of G , we denote the **reverse of path** P as \bar{P} , i.e., $\bar{P} = (u_l, u_{l-1}, \dots, u_2, u_1)$. For a path P of G , an edge in G between two non-adjacent nodes of P is said to be a **chord** of P . A path P is said to be a **chordless path** of a graph G if G does not have any chord of P , i.e., between any two non-adjacent nodes of P there is no edge in G . In other words, if $P = (u_1, u_2, \dots, u_{l-1}, u_l)$ is a chordless path of G then for $1 \leq i < j \leq l$, if $j \neq i + 1$, then neither $u_i - u_j \in E_G$, nor $u_i \rightarrow u_j \in E_G$, nor $u_j \rightarrow u_i \in E_G$. In a directed graph G , v is said to be a **descendant** of u if there exists a path from u to v . For a directed edge $y \rightarrow x$, y is said to be a **parent** of x .

For a path $P = (u_1, u_2, \dots, u_{l-1}, u_l)$, if the first two nodes of P are u and v (i.e., $u_1 = u$ and $u_2 = v$), and the last two nodes of P are x and y (i.e., $u_{l-1} = x$ and $u_l = y$), then we say P is a **path from** (u, v) **to**

(x, y) (note that the boundary cases $(u, v) = (x, y)$ and $u \neq v = x \neq y$ are both allowed by this definition). Similarly, if the first two nodes of P are u and v , and the last node of P is w then we say P is a **path from (u, v) to w** (again note that the degenerate case $v = w$ is allowed by this definition).

Definition 2.2 (Cycle, chord of a cycle, DAG). For a graph G , a *cycle* C is a sequence $(u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ of vertices in G such that $l \geq 3$, and both (u_1, u_2, \dots, u_l) and (u_l, u_1) are paths in G . The edge between two consecutive nodes of the sequence is said to be an edge of the cycle C . An edge between two non-adjacent nodes of a cycle is said to be a *chord of the cycle*. $C = (u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ is said to be a **chordless cycle** if it does not have any chord, i.e., $l \geq 4$ and both $P_1 = (u_1, u_2, \dots, u_{l-1})$ and $P_2 = (u_2, \dots, u_{l-1}, u_l)$ are chordless paths in G . C is said to be an **undirected cycle** of G if all the edges of the cycle are undirected, i.e., for all $1 \leq i \leq l$, $u_i - u_{i+1} \in E_G$. C is said to be a **directed cycle** if at least one edge of C is directed, i.e., there exists an i such that $1 \leq i \leq l$ and $u_i \rightarrow u_{i+1} \in E_G$. A directed graph is said to be a **directed acyclic graph (DAG)** if the graph does not have a cycle.

Definition 2.3 (Chordal Graph). An undirected graph G is said to be a *chordal graph* if there does not exist a chordless cycle in G .

Definition 2.4 (Chain Graph). A graph G is said to be a *chain graph* if there does not exist a directed cycle in G .

Proposition 2.5. *For every chain graph G , there always exists a topological ordering τ of the vertices of G such that for $u \rightarrow v \in E_G$, $\tau(u) < \tau(v)$.*

Proof. We can construct a topological ordering τ by first creating a topological ordering τ_1 of the vertices of the directed subgraph of G and then placing the nodes in τ_1 into τ in the same order as they appear in τ_1 . Next, we add the remaining nodes of G in any arbitrary order to complete τ . \square

Proposition 2.6. *Let G be a chain graph, C be an undirected connected component of G , and $u, v \in C$. Then $u \rightarrow v \notin G$.*

Proof. As both u and v belong to the same undirected connected component of G , there must exist an undirected path connecting them. If $u \rightarrow v \in G$, then combining this directed edge with the undirected path forms a directed cycle in G , which contradicts the definition of a chain graph. Thus, we conclude that $u \rightarrow v \notin G$. This completes the proof. \square

Proposition 2.7. *Let G be a graph with a chordal skeleton. If $C = (u_1, u_2, \dots, u_l, u_1)$ is a directed cycle in G , then there exists a directed cycle $C' = (a, b, c, a)$ in G of length three such that a, b, c are in the cycle C .*

Proof. Consider the smallest directed cycle C' in G that contains vertices of C . Such a cycle exists because C exists. We aim to show that C' must be of length three.

Suppose, for the sake of contradiction, that C' has more than three vertices, so it can be expressed as $C' = (v_1, v_2, \dots, v_m, v_1)$. Since the skeleton of G is chordal, there must exist a chord in C' . Let's assume this chord is between vertices v_i and v_j , where $i < j$, and v_i and v_j are not adjacent in C .

There are three possible cases: either $v_i - v_j$, $v_i \rightarrow v_j$, or $v_i \leftarrow v_j$. In each of these cases, either $(v_1, v_2, \dots, v_i, v_j, v_{j+1}, v_m, v_1)$ or $(v_i, v_{i+1}, \dots, v_j, v_i)$ forms a directed cycle in G of length less than m , which contradicts the minimality of C' . Therefore, we conclude that C' must be of length three.

This completes the proof of Proposition 2.7. \square

Definition 2.8 (Union of graphs). Let G_1 and G_2 be two graphs. G is said to be the *union* of G_1 and G_2 if $V_G = V_{G_1} \cup V_{G_2}$ and $E_G = E_{G_1} \cup E_{G_2}$.

Definition 2.9 (Tree decomposition). A *tree decomposition* of an undirected graph G is a pair (\mathcal{X}, T) , where $\mathcal{X} = \{X_1, X_2, \dots, X_l\}$ is a family of subsets of V_G , and T is a tree whose nodes are subsets X_i , satisfying the following properties:

1. $V_G = \bigcup_{i=1}^l X_i$.

2. For every edge $(u, v) \in E_G$, there is a subset X_i such that $u, v \in X_i$.
3. For any node $v \in V_G$, the set of nodes $\{X_i \mid v \in X_i\}$ induces a connected subgraph in T . This is equivalent to if X_i and X_j both contain a node v , then all nodes in the path between X_i and X_j contain v .

Property 2.10 is an intrinsic property of tree decomposition.

Property 2.10 (Intersection property of tree decomposition, Diestel (2005)). *Let G be an undirected graph, and $(\mathcal{X} = \{X_1, X_2, \dots, X_l\}, T)$ is a tree decomposition of G . For any edge $X_i - X_j \in E_T$, $X_i \cap X_j$ is a vertex separator of G .*

Definition 2.11 (Perfect elimination ordering (PEO)). Let G be an undirected graph. Let τ be an ordering of its vertices. τ is said to be a *perfect elimination ordering* of G if for all nodes v of G , the subgraph induced in G by the neighbors of v that come before v in τ forms a clique, i.e., for $u - v, w - v \in E_G$, if $\tau(u) < \tau(v)$ and $\tau(w) < \tau(v)$ then $u - w \in E_G$.

The following result is a classical result on the chordal graphs:

Proposition 2.12 (Dirac (1961)). *For a graph G , there exists a PEO if, and only if, G is a chordal graph.*

Definition 2.13 (LBFS Ordering). For a chordal graph G , Rose, Tarjan, and Lueker (1976) construct a lexicographical breadth-first search (LBFS) algorithm that returns a PEO ordering of a chordal graph. An ordering of a graph G that can be returned by the LBFS algorithm is said to be an *LBFS ordering* of G . A modified LBFS algorithm is provided in Alg. 3.

The following observations on chordal graphs come from Rose, Tarjan, and Lueker (1976).

Observation 2.14. *Let τ be an LBFS ordering of a chordal graph G . For $x, y, z \in V_G$, if both x and y are adjacent to z (i.e., $x - z, y - z \in E_G$) and both are of lesser rank than z in τ (i.e., $\tau(x) < \tau(z)$ and $\tau(y) < \tau(z)$) then x and y are also adjacent in G (i.e., $x - y \in E_G$).*

Proof. An LBFS ordering of a chordal graph G is known to be a perfect elimination ordering of G (Rose, Tarjan, and Lueker, 1976). Given that τ represents an LBFS ordering of a chordal graph G , we can deduce that the subgraph induced by the neighbors of z that appear before z in τ forms a clique. Consequently, if x and y are neighbors of z and satisfy $\tau(x) < \tau(z)$ and $\tau(y) < \tau(z)$ (i.e., both appear before z in τ), then x and y must belong to the same clique, implying that $(x, y) \in E_G$. \square

Observation 2.15. *Let G be an undirected chordal graph, and τ be an LBFS ordering of the vertices of G . For vertices $u, v \in V_G$ if $P = (p_1 = u, p_2, \dots, p_l = v)$ is a chordless path from u to v in G then there exists $1 \leq i \leq l$ such that that $\tau(p_1) > \tau(p_2) > \dots > \tau(p_i)$, and $\tau(p_i) < \tau(p_{i+1}) \dots < \tau(p_l)$ (note that when $i = 1$ or $i = l$ then one or the other of the above series of inequalities is empty). In other words, there are no “internal local maxima” of τ in P : τ is either strictly increasing, or strictly decreasing, or has a unique global minimum in the interior of P .*

Proof. There cannot exist vertices p_{j-1}, p_j, p_{j+1} such that $\tau(p_{j-1}) < \tau(p_j)$, and $\tau(p_{j+1}) < \tau(p_j)$. Otherwise, from Observation 2.14, $p_{j-1} - p_{j+1} \in E_G$, which further implies that P is not a chordless path. \square

Definition 2.16 (Skeleton (Verma and Pearl, 1990)). For a graph G , the *skeleton* of G , denoted as $skel(G)$ or G_U , is the underlying undirected graph of G . More formally, the skeleton of a graph G is an undirected graph obtained by replacing each directed edge $u \rightarrow v$ of G with an undirected edge $u - v$.

Definition 2.17 (v-structure (Verma and Pearl, 1990)). For a graph G , an induced subgraph of G of the form $a \rightarrow b \leftarrow c$ is said to be a v-structure of G . We denote the set of v-structures of G by $\mathcal{V}(G)$.

MEC: A directed acyclic graph (DAG) represents a set of conditional independence relations between random variables. Two DAGs are said to be Markov equivalent if both represent the same set of conditional independence relations between the random variables. A Markov equivalent class (MEC) is a set of DAGs that are Markov equivalent. An MEC can be represented by a complete partial directed acyclic graph (CPDAG) which is a union of the DAGs contained by that MEC (Andersson, Madigan, and Perlman, 1997). With a slight abuse of terminology, we equate the graph which represents an MEC with the MEC itself, and refer to both as an “MEC”. An MEC contains a unique set of v-structures. For an MEC, if we have given the skeleton of the MEC and its set of v-structures, we can construct the MEC using Meek’s rules (Meek, 1995). A DAG and its corresponding MEC both have the same skeleton and the same set of v-structures. If we have a DAG then using Meek’s rules, we can construct the MEC to which the DAG belongs.

Andersson, Madigan, and Perlman (1997) give the following necessary and sufficient conditions for a graph to be an MEC.




Figure 2: Strongly protected $u \rightarrow v$.

Theorem 2.18 (Andersson, Madigan, and Perlman (1997)). *A graph G is an MEC if, and only if,*

1. G is a chain graph.
2. For every chain component τ of G , G_τ is chordal, i.e., every undirected connected component of G is chordal.
3. The configuration $a \rightarrow b - c$ does not occur as an induced subgraph of G .
4. Every directed edge $u \rightarrow v \in G$ is strongly protected in G , i.e., $u \rightarrow v$ is a part of at least one of the subgraphs of G as shown in Figure 2.

2.2 New Terminologies and Related Results

In this paper, we introduce the following definitions and observations, which play a crucial role in simplifying our main result. Firstly, we define synchronous graphs, the Markov union of synchronous graphs, and partial MECs. Additionally, we introduce new notations used in this paper to represent these concepts. In the latter part of this subsection, we conduct a detailed study of three important terms: (a) triangle-free path, (b) the shadow of an MEC and a graph, and (c) the projection of an MEC. We thoroughly examine these three terms and also explore the relevant results and implications of these newly introduced terminologies in the context of our paper.

Definition 2.19 (Synchronous Graphs). Two graphs G and H are said to be *synchronous graphs* if there do not exist vertices $x, y \in V_G \cap V_H$ such that $x \rightarrow y \in E_G$ and $y \rightarrow x \in E_H$.

Definition 2.20 (Markov Union of Synchronous Graphs). Let G_1, G_2, \dots, G_l be pairwise synchronous graphs. The *Markov union* of G_1, G_2, \dots, G_l is a graph G , denoted by $G = U_M(G_1, G_2, \dots, G_l)$, such that

$\text{skeleton}(G)$ is the skeleton of the union of G_1, G_2, \dots, G_l (see Definition 2.8 for the definition of the graph union), and for any edge $u - v \in \text{skeleton}(G)$, $u \rightarrow v \in G$ if $u \rightarrow v \in G_i$ for any G_i . More formally, $V_G = \bigcup_{i=1}^l V_{G_i}$, the set of directed edges of G is $D_G = \{u \rightarrow v : u \rightarrow v \in E_{G_i} \text{ for some } 1 \leq i \leq l\}$, and the set of undirected edges of G is $U_G = \{u - v : u - v \in E_{G_i} \text{ for some } 1 \leq i \leq l, \text{ and there does not exist a } j \text{ such that } u \rightarrow v \in E_{G_j} \text{ or } v \rightarrow u \in E_{G_j}\}$. In other words, if $u \rightarrow v$ is a directed edge in any graph G_i , then $u \rightarrow v$ is also a directed edge in the Markov union graph, and an undirected edge $u - v$ of G_i becomes an undirected edge of G only if neither $u \rightarrow v$ nor $v \rightarrow u$ is part of any graph in the union. Since the graphs are pairwise synchronous, if one graph G_i contains a directed edge $u \rightarrow v$, then no other graph G_j can contain the directed edge $v \rightarrow u$.

Definition 2.21 (Partial MEC¹). A graph M is said to be a *partial MEC* if it satisfies the following conditions:

1. M is a chain graph.
2. Undirected connected components of M are chordal.
3. There does not exist a subgraph of the form $u \rightarrow v - w \in M$.

i.e., a partial MEC obeys items 1 to 3 of Theorem 2.18. In other words, a partial MEC is a chain graph with chordal undirected connected components and no subgraph of the form $u \rightarrow v - w$.

Definition 2.21 implies that every MEC is a partial MEC, and a partial MEC is an MEC if it also obeys item 4 of Theorem 2.18.

Notation 2.22. We say an MEC M is an MEC of an undirected graph G if $\text{skeleton}(M) = G$. Similarly, we say a partial MEC M is a partial MEC of an undirected graph G if $\text{skeleton}(M) = G$. For an undirected graph G , we denote $\text{MEC}(G)$ as the set of MECs of G , and $\text{PMEC}(G)$ as the set of partial MECs of G .

2.2.1 Triangle-Free Path

Definition 2.23 (Triangle-free path). For a graph G , a path $P = (u_1, u_2, \dots, u_{l-1}, u_l)$ is said to be a *triangle-free path* of G if for any node u_i in P , there is no edge between the two adjacent nodes of u_i in P . More formally, for any $1 \leq i \leq l - 2$, neither $u_i - u_{i+2} \in E_G$, nor $u_i \rightarrow u_{i+2} \in E_G$, nor $u_{i+2} \rightarrow u_i \in E_G$, i.e., $u_i - u_{i+2} \notin \text{skeleton}(G)$.

Note that if $v \rightarrow u \in E_G$ then there cannot be a triangle-free path with the first two nodes u and v (if $v \rightarrow u \in E_G$ then from Definition 2.1, there cannot be a path with the first two nodes u and v , as neither $u - v \in E_G$ nor $u \rightarrow v \in E_G$).

Observation 2.24. For a graph G , every chordless path $P = (u_1, u_2, \dots, u_l)$ of G is a triangle-free path of G .

Proof. Since P is a chordless path, there cannot be an edge between two non-adjacent nodes of P . This implies there cannot be an edge between u_i and u_{i+2} for any $1 \leq i \leq l - 2$. This further implies P is a triangle-free path of G . \square

For our purposes, the utility of the notion of a triangle-free path comes from the following partial converse.

Proposition 2.25. Let G be an undirected chordal graph, and let P be a triangle-free path in G . Then, P is also a chordless path.

¹The same notion appeared under the name ‘Restricted Chain Graph’ in the earlier work of [Van Der Zander and Liskiewicz \(2016\)](#); [Bang and Didelez \(2023\)](#). We want to thank an anonymous reviewer for pointing out the references.

Proof. Suppose, if possible, that $P = (u_0, u_1, \dots, u_\ell)$ is a triangle-free path in G that is not a chordless path. Choose a chord $u_i - u_j$ (assuming $i < j$ without loss of generality) of P with the smallest possible value of $j - i$. Since P is a triangle-free path, we must have $j - i \geq 3$. However, in that case $C = (u_i, u_{i+1}, \dots, u_j, u_i)$ is a simple cycle in G of length at least four. Since G is chordal, this cycle must have a chord. By the construction of C , this chord must be of the form $u_a - u_b$ with $i \leq a < b \leq j$ and $1 < b - a < j - i$. But then, $u_a - u_b$ is a chord of P which contradicts the minimal choice of $j - i$. We, therefore, conclude that P must be a chordless path. \square

The following corollary of this partial converse is also useful.

Corollary 2.26. *Let G be an undirected chordal graph, and suppose that u_1 and u_2 are adjacent in G . Suppose that $P = (v_1 = u_1, v_2, v_3, \dots, v_\ell)$ is a triangle-free path in G which contains u_2 . Then, it must be the case that u_2 is the second vertex in P (i.e., $v_2 = u_2$).*

Proof. By Proposition 2.25, P is a chordless path in G , and hence no v_i for $i > 2$ can be adjacent to $v_1 = u_1$ (otherwise, $v_1 - v_i$ would be a chord of P). Since u_2 is adjacent to u_1 and is contained in P , this only leaves open the possibility $v_2 = u_2$. \square

The following observation extends the above to MECs.

Observation 2.27. *Let G be a chain graph with chordal undirected components, and let P be an undirected triangle-free path in G . Then, P is an undirected chordless path in G .*

Proof. Since P is undirected, it must be contained wholly in some undirected chordal component H of G . Thus, Proposition 2.25 implies that P must in fact be a chordless path in H (and therefore also in G). \square

The following observation shows the transitive nature of triangle-free paths in a chain graph.

Observation 2.28 (Concatenation of Triangle-Free Paths). *Consider a chain graph G with chordal undirected components. Let $u, v, x, y, w \in V_G$ be (possibly non-distinct) vertices of G . Suppose $P_1 = (a_1 = u, a_2 = v, \dots, a_{l-1} = x, a_l = y)$ and $P_2 = (b_1 = x, b_2 = y, \dots, b_{m-1}, b_m = w)$ are triangle-free paths in G from (u, v) to (x, y) and from (x, y) to w , respectively. Then, their concatenation $P := (a_1 = u, a_2 = v, \dots, a_{l-1} = x, a_l = y, b_3, b_4, \dots, b_{m-1}, b_m = w)$ is a triangle-free path in G from (u, v) to w .*

Proof. From the definition of P , we observe that (i) for any two consecutive vertices p and q in P , either the edge $p \rightarrow q$ or the edge $p - q$ exists in G because p and q must be consecutive in either P_1 or P_2 , both of which are paths in G , and (ii) for any three consecutive vertices p, q , and r in P , the vertices p, q , and r are distinct, and p and r are not adjacent in G because p, q , and r must be consecutive in one of P_1 or P_2 , both of which are triangle-free paths.

Thus, to show that P is a triangle-free path in G it only remains to show that all vertices in P are distinct. Suppose, if possible, that this is not the case. Then, there must exist a segment of P of the form $C := (c_1 = p, c_2, \dots, c_{k-1}, c_k = p)$ in which the vertices c_1, c_2, \dots, c_{k-1} are distinct. From the observations above about P , we see that C must be a cycle in G . Since G is a chain graph, C cannot have any directed edges. Therefore, C must be entirely contained within an undirected component H of G . Furthermore, from our second observation, C must have at least four nodes. If C does not have a chord in H then C is a chordless cycle in G , which contradicts the assumption that the undirected connected components of G are chordal. Suppose C has a chord (c_i, c_j) . We choose such (c_i, c_j) with $i < j$ and $j - i$ being minimal. In this case, we have another cycle $C' := (c_i, c_{i+1}, \dots, c_j, c_{j+1} = c_i)$. We have selected c_i and c_j such that there is no edge between any two non-adjacent nodes of C' (due to the minimality of $j - i$). According to our second observation, C' must have at least four nodes. This makes C' a chordless cycle, contradicting the assumption that the undirected components of G are chordal.

We conclude that all vertices in P are distinct. Combined with the above two observations about P , this shows that P is a triangle-free path in G . \square

The following observation is an easy consequence of the fact that the reverse of an undirected path is also a path.

Observation 2.29. For a graph G , if $P = (u_1 = u, u_2 = v, \dots, u_{l-1} = x, u_l = y)$ is an undirected triangle-free path from (u, v) to (x, y) in G then $\bar{P} = (u_l = y, u_{l-1} = x, \dots, u_2 = v, u_1 = u)$, the reverse of the path P , is also a triangle-free path from (y, x) to (v, u) .

Proof. Suppose that \bar{P} is not a triangle-free path, then there must exist nodes u_i and u_{i+2} such that there is an edge between u_i and u_{i+2} . But, then, P is also not a triangle-free path, which is a contradiction. \square

The following observation comes from the item 3 of Theorem 2.18.

Observation 2.30. Let M be an MEC, and $P = (u_1 = x, u_2 = y, \dots, u_{l-1} = u, u_l = v)$ be a triangle-free path from (x, y) to (u, v) in M . If $u - v \in M$ then P is an undirected chordless path in M , more specifically $x - y \in M$.

Proof. Since P is a path, for each $1 \leq i \leq l - 1$, either $u_i - u_{i+1} \in M$ or $u_i \rightarrow u_{i+1} \in M$. Suppose Observation 2.30 is not true. Then, pick the highest i such that $u_i \rightarrow u_{i+1} \in M$. i must be less than $l - 1$, as $u_{l-1} - u_l \in M$. But then M has an induced subgraph $u_i \rightarrow u_{i+1} - u_{i+2}$, which is a contradiction from Item 3 of Theorem 2.18. This implies all the edges of P are undirected. This further implies P is an undirected triangle-free path in M . From Observation 2.27, P is a chordless path in M (from Theorem 2.18, M is a chain graph with chordal undirected components). \square

Observations 2.24 and 2.28 and Proposition 2.25 imply the following:

Lemma 2.31. Let G be an undirected chordal graph. Suppose $P_1 = (a_1 = u, a_2 = v, \dots, a_{l-1} = x, a_l = y)$ and $P_2 = (b_1 = x, b_2 = y, \dots, b_{m-1} = u, b_m = w)$ are chordless paths in G from (u, v) to (x, y) and from (x, y) to w , respectively. Then, their concatenation $P := (a_1 = u, a_2 = v, \dots, a_{l-1} = x, a_l = y, b_3, b_4, \dots, b_{m-1}, b_m = w)$ is a chordless path in G from (u, v) to w .

Proof. From Observation 2.24, P_1 and P_2 are triangle-free paths in G . From Observation 2.28, the path P , concatenation of P_1 and P_2 , is a triangle-free path in G . And, from Proposition 2.25, P is a chordless path in G . \square

We now introduce a triangle-free path variant of a source node.

Definition 2.32 (Canonical Source Node). Let G be a graph. A node $s \in V_G$ is said to be a *canonical source node* of G if there does not exist a directed edge $u \rightarrow v \in G$ such that there exists a triangle-free path from (u, v) to s in G .

In particular, note that the existence of an edge of the form $u \rightarrow s$ also prohibits s from being a canonical source node, due to the presence of the (degenerate) triangle-free path (u, s) from (u, s) to s .

Definition 2.33 (Certificate for a non-canonical source node). Let G be a graph. For any non-canonical source node $z \in V_G$, we say x is a *certificate* of z if there exists a $y \in V_G$ such that $x \rightarrow y \in G$ and there exists a triangle-free path from (x, y) to z . In other words, x certifies that z is not a canonical source node in G .

The following observation shows that self-certification is not possible.

Observation 2.34. Let G be a graph, and $u \in V_G$ such that u is not a canonical source node. Then u cannot be a certificate of itself.

Proof. Suppose that, if possible, u is a certificate of itself. Then from Definition 2.33, there exists a $v \in V_G$ such that $u \rightarrow v \in G$ and there exists a triangle-free path $Q = (u_1 = u, u_2 = v, \dots, u_l = u)$ from (u, v) to u . But, then from the definition of a path (Definition 2.1), Q is not a path. Hence, Q is not a triangle-free path. Therefore, u cannot be a certificate of itself. \square

We now provide an algorithm that for an input chain graph G with chordal undirected connected components, outputs a pair of functions (P_1, P_2) such that for two distinct edges (u, v) and (x, y) in G , $P_1((u, v), (x, y)) = 1$ if there exists a triangle-free path from (u, v) to (x, y) in G , else $P_1((u, v), (x, y)) = 0$. And, for an edge (u, v) and a vertex w of G , $P_2((u, v), w) = 1$ if $v \neq w$, and there exists a triangle-free path from (u, v) to w in G , else $P_2((u, v), w) = 0$.

Algorithm 1: TFP(G)

Input : A chain graph G with chordal undirected connected components
Output : (P_1, P_2) such that
 $P_1 : E_G \times E_G \rightarrow \{0, 1\}$ and $P_2 : E_G \times V_G \rightarrow \{0, 1\}$ such that
(a) for $(u, v), (x, y) \in E_G$, $P_1((u, v), (x, y)) = 1$ if $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) in G else $P_1((u, v), (x, y)) = 0$, and
(b) for $((u, v), w) \in E_G \times V_G \rightarrow \{0, 1\}$, $P_2((u, v), w) = 1$ if $v \neq w$ and there exists a triangle-free path from (u, v) to w in G else $P_2((u, v), w) = 0$.

- 1 Construct a function $P_1 : E_G \times E_G \rightarrow \{0, 1\}$.
- 2 Construct a function $P_2 : E_G \times V_G \rightarrow \{0, 1\}$.
- 3 **foreach** $((u, v), (x, y)) \in E_G \times E_G$ **do**
- 4 | $P_1((u, v), (x, y)) \leftarrow 0$
- 5 **end**
- 6 **foreach** $((u, v), w) \in E_G \times V_G$ **do**
- 7 | $P_2((u, v), w) \leftarrow 0$
- 8 **end**
- 9 **foreach** *triple* (u, v, w) **do**
- 10 | **if** (u, v, w) *is a triangle-free path in* G **then**
- 11 | | $P_1((u, v), (v, w)) = 1$
- 12 | | $P_2((u, v), w) = 1$
- 13 | **end**
- 14 **end**
- 15 **while** $\exists (u, v), (x, y), (z_1, z_2) \in E_G$ *such that* $P_1((u, v), (x, y)) = 0$, $(u, v) \neq (x, y)$, *and*
 $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$ **do**
- 16 | $P_1((u, v), (x, y)) = 1$
- 17 **end**
- 18 **while** $\exists (u, v), (z_1, z_2) \in E_G$ *and* $w \in V_G$ *such that* $P_2((u, v), w) = 0$, $v \neq w$, *and*
 $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$ **do**
- 19 | $P_2((u, v), w) = 1$
- 20 **end**
- 21 **return** (P_1, P_2)

Lemma 2.35. *Let G be a chain graph with chordal undirected connected components. For input G , Alg. 1 returns (P_1, P_2) , a tuple of functions, such that (a) $P_1 : E_G \times E_G \rightarrow \{0, 1\}$ and for $(u, v), (x, y) \in E_G$, $P_1((u, v), (x, y)) = 1$ if and only if $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) in G , and (b) $P_2 : E_G \times V_G \rightarrow \{0, 1\}$, and for $((u, v), w) \in E_G \times V_G$, $P_2((u, v), w) = 1$ if and only if $v \neq w$ and there exists a triangle-free path from (u, v) to w in G .*

Proof. We will prove the correctness of P_1 and P_2 separately.

Correctness of P_1 : We first prove that for all $(u, v), (x, y) \in E_G$ such that $(u, v) \neq (x, y)$ and there exists a triangle-free path $Q = (u_1 = u, u_2 = v, \dots, u_{l-1} = x, u_l = y)$ from (u, v) to (x, y) in G , we have $P_1((u, v), (x, y)) = 1$. We will use induction on the length l of the triangle-free path. Since $(u, v) \neq (x, y)$, l must be greater than one.

If $l = 2$, then $v = x$, and $Q = (u, v = x, y)$ is a triangle-free path from (u, v) to (x, y) in G . Therefore, while running lines 9-14 of Alg. 1, we have $P_1((u, v), (x, y)) = 1$. This implies that for all pairs of distinct edges $(u, v), (x, y) \in E_G$, if there exists a triangle-free path from (u, v) to (x, y) of length two then $P_1((u, v), (x, y)) = 1$.

Let us assume that for some $k \geq 2$, for all pairs of distinct edges $(u, v), (x, y) \in E_G$, if there exists a triangle-free path from (u, v) to (x, y) in G of length at most k then $P_1((u, v), (x, y)) = 1$. We now show that for $(u, v), (x, y) \in E_G$ such that $(u, v) \neq (x, y)$, if there exists a triangle-free path $Q = (u_1 = u, u_2 =$

$v, \dots, u_{l-1} = x, u_l = y$ from (u, v) to (x, y) in G of length $l = k + 1$, then $P_1((u, v), (x, y)) = 1$.

Take two subpaths of Q : $Q_1 = (u_1 = u, u_2 = v, u_3)$ and $Q_2 = (u_2 = v, \dots, u_{l-1} = x, u_l = y)$. Both paths must be triangle-free paths of length at most k . Therefore, from our assumption, $P_1((u_1 = u, u_2 = v), (u_2, u_3)) = P_1((u_2, u_3), (u_{l-1} = x, u_l = y)) = 1$. But then, while running lines 15-17, we get $P_1((u, v), (x, y)) = 1$. This completes the induction. Thus, we have proven that for all $(u, v), (x, y) \in E_G$ such that $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) in G , we have $P_1((u, v), (x, y)) = 1$.

We now prove that for all $(u, v), (x, y) \in E_G$, if $P_1((u, v), (x, y)) = 1$ then $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) in G . Suppose that this is not true. Then there must exist a pair of edges $(u, v), (x, y) \in E_G$ such that $P_1((u, v), (x, y)) = 1$ and either $(u, v) = (x, y)$ or there does not exist a triangle-free path from (u, v) to (x, y) in G .

For all pairs of edges $(u, v), (x, y) \in E_G$, Alg. 1 initializes $P_1((u, v), (x, y)) = 0$ (lines 3-5). Pick the first such pair $(u, v), (x, y) \in E_G$ for which $P_1((u, v), (x, y)) = 1$ and either $(u, v) = (x, y)$ or there does not exist a triangle-free path from (u, v) to (x, y) , i.e., for all the pairs $(u', v'), (x', y') \in E_G$, for which the value of $P_1((u', v'), (x', y'))$ has been updated to 1 before the update of $P_1((u, v), (x, y))$, $(u', v') \neq (x', y')$ and there exists a triangle-free path from (u', v') to (x', y') . In the algorithm, for any pair of edges (u, v) and (x, y) , Alg. 1 updates P_1 either at lines 9-14, or at lines 15-17. If the value of $P_1((u, v), (x, y))$ has been updated while running lines 9-14 of Alg. 1 then we have $v = x$ and $(u, v = x, y)$ is a triangle-free path. This further implies $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) .

And, if $P_1((u, v), (x, y))$ has been updated while running lines 15-17 of Alg. 1 then we have $(u, v) \neq (x, y)$ and there exists a $(z_1, z_2) \in E_G$ such that $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$. Since $P_1((u, v), (z_1, z_2))$ and $P_1((z_1, z_2), (x, y))$ have been updated before the update of $P_1((u, v), (x, y))$, therefore, from our assumption, (a) $(u, v) \neq (z_1, z_2)$ and there exists a triangle-free path $Q_1 = (u_1 = u, u_2 = v, u_{l-1} = z_1, u_l = z_2)$ from (u, v) to (z_1, z_2) , and (b) $(z_1, z_2) \neq (x, y)$ and $Q_2 = (v_1 = z_1, v_2 = z_2, \dots, v_{m-1} = x, v_m = y)$ is a triangle-free path from (z_1, z_2) to (x, y) , respectively. But, then from Observation 2.28, concatenation of Q_1 and Q_2 gives us a triangle-free path from (u, v) to (x, y) . Thus, $P_1((u, v), (x, y)) = 1$ correctly depicts that $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) . This completes the proof of correctness of P_1 .

Correctness of P_2 : We first prove that for all $((u, v), w) \in E_G \times V_G$ such that $v \neq w$ and there exists a triangle-free path $Q = (u_1 = u, u_2 = v, \dots, u_l = w)$ from (u, v) to w in G we have $P_2((u, v), w) = 1$. We prove this using induction on the length l of the triangle-free path. Since $v \neq w$, l must be greater than one.

If $l = 2$, then $Q = (u, v, w)$ is a triangle-free path from (u, v) to w in G . Therefore, while running lines 9-14 of Alg. 1, we get $P_2((u, v), w) = 1$. This implies that for all $((u, v), w) \in E_G \times V_G$, if there exists a triangle-free path from (u, v) to w in G of length two then $P_2((u, v), w) = 1$.

Let us assume that for some $k \geq 2$, for all $((u, v), w) \in E_G \times V_G$, if $v \neq w$ and there exists a triangle-free path from (u, v) to w in G of length at most k then $P_2((u, v), w) = 1$. We now show that for all $((u, v), w) \in E_G \times V_G$ such that $v \neq w$, if there exists a triangle-free path $Q = (u_1 = u, u_2 = v, \dots, u_l = w)$ from (u, v) to w in G of length $l = k + 1$, then $P_2((u, v), w) = 1$.

Take two subpaths of Q : $Q_1 = (u_1 = u, u_2 = v, u_3)$ and $Q_2 = (u_2 = v, \dots, u_{l-1}, u_l = w)$. Since Q is a triangle-free path, therefore, the subpath Q_1 is a triangle-free path from $(u_1 = u, u_2 = v)$ to (u_2, u_3) . Similarly, the subpath Q_2 is a triangle-free path from (u_2, u_3) to $u_l = w$ of length k . Since the length of Q_1 is two, therefore $(u_1, u_2) \neq (u_2, u_3)$. Therefore, from the correctness of P_1 , $P_1((u_1 = u, u_2 = v), (u_2, u_3)) = 1$. Since the length of Q_2 is k , therefore, $u_3 \neq u_l$ (otherwise Q_2 is of length $1 < k$, a contradiction). Therefore, from our assumption, $P_2((u_2, u_3), u_l = w) = 1$. But, then while running lines 18-20, we get $P_2((u, v), w) = 1$ (since $P_1((u_1 = u, u_2 = v), (u_2, u_3)) = P_2((u_2, u_3), u_l = w) = 1$). This completes the induction. Thus, we prove that for all $((u, v), w) \in E_G \times V_G$ such that $v \neq w$ and there exists a triangle-free path from (u, v) to w in G , we have $P_2((u, v), w) = 1$.

We now prove that for all $((u, v), w) \in E_G \times V_G$, if $P_2((u, v), w) = 1$ then $v \neq w$ and there exists a triangle-free path from (u, v) to w in G . Suppose this is not true. Then there must exist $((u, v), w) \in E_G \times V_G$ such that $P_2((u, v), w) = 1$ and either $v = w$ or there does not exist a triangle-free path from (u, v) to w in G .

For all the pair $((u, v), w) \in E_G \times V_G$, Alg. 1 initializes $P_2((u, v), w) = 0$ (lines 3-5). If possible, pick the first such $((u, v), w) \in E_G \times V_G$ for which $P_2((u, v), w) = 1$ and either $v = w$ or there does not exist a triangle-free path from (u, v) to w in G , i.e., for all $((u', v'), w') \in E_G \times V_G$ for which the value of $P_2((u', v'), w')$

has been updated to 1 before the update of $P_2((u, v), w)$, we have $v' \neq w'$ and there exists a triangle-free path from (u', v') to w' . In the algorithm, for any $((u, v), w) \in E_G \times V_G$, Alg. 1 updates P_2 either at lines 9-14, or at lines 18-20. Suppose the value of $P_2((u, v), w)$ has been updated while running lines 9-14 of Alg. 1 then we must have (u, v, w) as a triangle-free path from (u, v) to w . Also, $v \neq w$ (otherwise, (u, v, w) is even not a path).

Now we move to the remaining possibility. Suppose $P_2((u, v), w)$ has been updated while running lines 18-20 of Alg. 1 then we must have $v \neq w$ and there exists a $(z_1, z_2) \in E_G$ such that $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$. We have shown the correctness of P_1 earlier. Therefore, $P_1((u, v), (z_1, z_2)) = 1$ implies that $(u, v) \neq (z_1, z_2)$ and there exist triangle-free paths $Q_1 = (u_1 = u, u_2 = v, u_{l-1} = z_1, u_l = z_2)$ from (u, v) to (z_1, z_2) . Also, $P_2((z_1, z_2), w)$ has been already updated. Therefore, from our assumption, there exists a triangle-free path $Q_2 = (v_1 = z_1, v_2 = z_2, \dots, v_{m-1}, v_m = w)$ from (z_1, z_2) to w . But, then from Observation 2.28, concatenation of Q_1 and Q_2 gives us a triangle-free path from (u, v) to w . Thus, $P_2((u, v), w) = 1$ correctly depicts that $v \neq w$ and there exists a triangle-free path from (u, v) to w in G . This completes the proof of Lemma 2.35. \square

2.2.2 Shadow of an MEC

Definition 2.36 (Shadow of an MEC). Let M be an MEC of G , $Y \subseteq V_G$, $O \in \text{PMEC}(G[Y])$ (i.e., O be a partial MEC with skeleton $G[Y]$), and $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ and $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ be two functions. We define (O, P_1, P_2) as a shadow of the MEC M on Y , denoted by $(O, P_1, P_2) = \text{shadow}(M, Y)$, if the following occur:

1. $M[Y] = O$, i.e., O is an induced subgraph of M on Y .
2. For $(u, v), (x, y) \in E_O$, $P_1((u, v), (x, y)) = 1$ if $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) in M , else $P_1((u, v), (x, y)) = 0$.
3. For $(u, v) \in E_O$ and $w \in V_O$, $P_2((u, v), w) = 1$, if $v \neq w$ and there exists a triangle-free path from (u, v) to w in M , else $P_2((u, v), w) = 0$.

With a slight abuse of notation, we say (O, P_1, P_2) as a shadow of an undirected graph G , denoted by $(O, P_1, P_2) \in \text{shadow}(G)$, if $O \in \text{PMEC}(G)$, and $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ and $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ are two functions.

Remark 2.37 (P_1 does not determine P_2). Note that P_2 is not necessarily determined by P_1 . This is because there can exist $(u, v) \in E_O$ and $w \in Y$ such that there exists a triangle-free path from (u, v) to w in M , i.e., $P_2((u, v), w) = 1$, but for which there does not exist any edge $(x, w) \in E_O$ such that $P_1((u, v), (x, w)) = 1$. This can happen when every triangle-free path $P = (u_1 = u, u_2 = v, \dots, u_{l-1}, u_l = w)$ in M from (u, v) to w has $u_{l-1} \notin Y$.

Definition 2.38 ($\text{MEC}(G, O, P_1, P_2)$). Let G be an undirected graph, $Y \subseteq V_G$, and $(O, P_1, P_2) \in \text{shadow}(G[Y])$. We define $\text{MEC}(G, O, P_1, P_2)$ to be the set of MECs M of G for which (O, P_1, P_2) is a shadow of M on Y . More formally, $\text{MEC}(G, O, P_1, P_2) := \{M : M \in \text{MEC}(G) \text{ and } (O, P_1, P_2) = \text{shadow}(M, V_O)\}$.

The following observation is intuitive from Definition 2.36.

Observation 2.39. *Let G be an undirected graph, and let M be an MEC of G . For $Y \subseteq V_G$, if (O, P_1, P_2) is a shadow of the MEC M on Y , then (O, P_1, P_2) is a shadow of the undirected graph $G[Y]$.*

Proof. Suppose (O, P_1, P_2) is the shadow of M on Y . According to Definition 2.36, $O = M[Y]$, and $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ and $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ are two functions. Thus, to establish that (O, P_1, P_2) is a shadow of $G[Y]$, we only need to demonstrate that O is a partial MEC.

Since M is an MEC, as stated in Theorem 2.18, M is a chain graph with chordal undirected connected component and does not possess any induced subgraph of the form $u \rightarrow v - w$. It's worth noting that an induced subgraph of a chain graph with chordal undirected connected component is also a chain graph with

chordal undirected connected component. Furthermore, if M has no induced subgraph of the form $u \rightarrow v - w$, then its induced subgraphs must also lack such a structure.

This implies that O also adheres to the conditions specified in Theorem 2.18. Thus, by Definition 2.21, O qualifies as a partial MEC. This concludes the proof of Observation 2.39. \square

Definition 2.38 gives us a way to partition the set of MECs of any undirected graph G . Lemma 2.40 partitions the set of MECs of G based on its shadow.

Lemma 2.40. *Let G be an undirected graph. Then, for any $Y \subseteq V_G$,*

$$|\text{MEC}(G)| = \sum_{(O, P_1, P_2) \in \text{shadow}(G[Y])} |\text{MEC}(G, O, P_1, P_2)| \quad (1)$$

Proof. From Definition 2.38, for each $M \in \text{MEC}(G)$, there exists a unique shadow (O, P_1, P_2) of M on Y . From Observation 2.39, $(O, P_1, P_2) \in \text{shadow}(G[Y])$. This proves Lemma 2.40. \square

2.2.3 Projection of an MEC

Definition 2.41 (Projection of an MEC). Let G be an undirected graph, $S \subseteq V_G$, $G' = G[S]$ be an induced subgraph of G , M be an MEC of G , and M' be an MEC of G' . We say M' is a *projection* of M on S , denoted as $\mathcal{P}(M, S) = M'$, if $\mathcal{V}(M') = \mathcal{V}(M[S])$.

With slight overloading of the function \mathcal{P} , for two subsets $S_1, S_2 \subseteq V_G$, we define (M_1, M_2) as a projection of M on (S_1, S_2) , denoted as: $\mathcal{P}(M, S_1, S_2) = (M_1, M_2)$, if $M_1 = \mathcal{P}(M, S_1)$, and $M_2 = \mathcal{P}(M, S_2)$.

Lemma 2.42. *Let G be an undirected graph, and M be an MEC of G . For any $S \subseteq V_G$, there exists a unique MEC M' of the graph $G[S]$ such that $\mathcal{P}(M, S) = M'$.*

Proof. As a reminder from the introduction, it is well-known that every DAG corresponds to a unique MEC, and the set of v-structures in a DAG is identical to the set of v-structures in its corresponding MEC. We first show the existence of such an MEC M' . Let D be a DAG such that $D \in M$. The set of v-structures contained by $D[S]$ equals $\mathcal{V}(M[S])$ (since both D and M have the same set of v-structures). Let M' be an MEC of $G[S]$ such that $D[S] \in M'$. The set of v-structures of M' must be the same as the set of v-structures of $D[S]$, i.e. $\mathcal{V}(M') = \mathcal{V}(M[S])$. The uniqueness of M' comes from the fact that if two MECs of a graph have the same set of v-structure then both are the same. \square

Lemma 2.42 further implies that the following corollaries:

Corollary 2.43. *Let G be an undirected graph, and M be an MEC of G . For $S_1, S_2 \subseteq V_G$, there exists a unique tuple $(M_1, M_2) \in \text{MEC}(G[S_1]) \times \text{MEC}(G[S_2])$ such that $\mathcal{P}(M, S_1, S_2) = (M_1, M_2)$.*

Proof. From Definition 2.41, $M_1 = \mathcal{P}(M, S_1)$ and $M_2 = \mathcal{P}(M, S_2)$. From Lemma 2.42, $\mathcal{P}(M, S_1)$ and $\mathcal{P}(M, S_2)$ both are unique. This proves the uniqueness of (M_1, M_2) . \square

Corollary 2.44. *Let G be an undirected graph, $S \subseteq V_G$, $M \in \text{MEC}(G)$, and $M' \in \text{MEC}(G[S])$. If $\mathcal{P}(M, S) = M'$ then for any $S' \subseteq S$, $\mathcal{V}(M[S']) = \mathcal{V}(M'[S'])$, and $\mathcal{P}(M, S') = \mathcal{P}(M', S')$.*

Proof. Since $S' \subseteq S$, for $u, v, w \in S'$, if a v-structure $u \rightarrow v \leftarrow w \in \mathcal{P}(M', S')$ then from Definition 2.41, the v-structures also belongs to M' and $\mathcal{P}(M, S')$. Similarly, for $u, v, w \in S'$, if a v-structure $u \rightarrow v \leftarrow w \in \mathcal{P}(M, S')$ then from Definition 2.41, the v-structures also belongs to M' , and $\mathcal{P}(M', S)$. This implies $\mathcal{V}(M, S') = \mathcal{V}(M', S')$. From Definition 2.41, this further implies that $\mathcal{P}(M, S') = \mathcal{P}(M', S')$. \square

The following lemma shows some structural similarity between an MEC M and its projection M' .

Lemma 2.45. *Let G be a graph, $S \subseteq V_G$, $M \in \text{MEC}(G)$, and $\mathcal{P}(M, S) = M'$. If $u \rightarrow v \in M'$, then $u \rightarrow v \in M$.*

Proof. Meek (1995) gives a set of rules using which, if we have knowledge about the v-structure of an MEC, then we can construct the MEC. According to Definition 2.41, $\mathcal{V}(M') \subseteq \mathcal{V}(M)$. Therefore, if an edge can be directed using Meek's rule in M' , then it must be directed in M (using the same Meek's rule). This proves Lemma 2.45. However, we have another proof. This proof is more rigorous and has better clarity. We are presenting this proof because this approach of proving will be used multiple times in this paper.

Let us assume that Lemma 2.45 is not true, i.e., there exists an edge $u \rightarrow v \in M'$ such that $u \rightarrow v \notin M$. Since M' is a chain graph (from item 1 of Theorem 2.18), from Proposition 2.5, there exists a topological ordering τ of the vertices of M' such that if $u \rightarrow v \in M'$ then $\tau(u) < \tau(v)$. Pick 2 vertices $u, v \in V_{M'}$ such that $u \rightarrow v \in M'$, $u \rightarrow v \notin M$, and for any edge $u' \rightarrow v' \in M'$, if $u' \rightarrow v' \notin M$ then either $\tau(v') > \tau(v)$, or $v = v'$ and $\tau(u') \leq \tau(u)$. The chosen vertices u and v imply that the following claim:

Claim 2.46. *For any edge $u' \rightarrow v' \in M'$, if $\tau(v') < \tau(v)$, or $v = v'$ and $\tau(u) < \tau(u')$ then $u' \rightarrow v' \in M$.*

Since $u \rightarrow v \in M'$, it must be strongly protected in M' (item 4 of Theorem 2.18). Then, $u \rightarrow v$ is part of one of the subgraphs shown in fig. 2. We go through each possibility and show that in each case, $u \rightarrow v \in M$, contradicting our assumption.

Case-1: $u \rightarrow v$ is a part of item-(a) of fig. 2, i.e., $w \rightarrow u \rightarrow v \in M'$. Then, $\tau(w) < \tau(u) < \tau(v)$. From Claim 2.46, $w \rightarrow u \in M$. $u - v \notin M$, as it violates item 3 of Theorem 2.18. Neither $v \rightarrow u \in M$, otherwise, $w \rightarrow u \leftarrow v$ is a v-structure in M . Since M' is a projection of M , and $u, v, w \in V_{M'}$, from Definition 2.41, if the v-structure $w \rightarrow u \leftarrow v \in M$ then $w \rightarrow u \leftarrow v \in M'$. But, this is a contradiction as $u \rightarrow v \in M'$. Thus the only option that remains is $u \rightarrow v \in M$.

Case-2: $u \rightarrow v$ is a part of item-(b) of fig. 2, i.e., $u \rightarrow v \leftarrow w \in M'$. Since $u \rightarrow v \leftarrow w$ is a v-structure, from Definition 2.41, $u \rightarrow v \leftarrow w \in M$, as M' is a projection of M . This implies that $u \rightarrow v \in M$.

Case-3: $u \rightarrow v$ is a part of item-(c) of fig. 2, i.e., $u \rightarrow w \rightarrow v \leftarrow u \in M'$. Then, $\tau(u) < \tau(w) < \tau(v)$. From Claim 2.46, $u \rightarrow w \rightarrow v \in M$. If $u - v \in M$, or $v \rightarrow u \in M$ then (u, v, w, u) is a directed cycle in M , contradicting item 1 of Theorem 2.18. This implies $u \rightarrow v \in M$.

Case-4: $u \rightarrow v$ is a part of item-(d) of fig. 2. Since $w \rightarrow v \leftarrow w'$ is a v-structure, from Definition 2.41, $w \rightarrow v \leftarrow w' \in M$, as M' is a projection of M . If $v - u \in M$, or $v \rightarrow u \in M$ then $w \rightarrow u \leftarrow w' \in M$, otherwise M has a directed cycle (w, v, u, w) (if $w \rightarrow u \notin M$), or (w', v, u, w') (if $w' \rightarrow u \notin M$), which violates item 1 of Theorem 2.18. Since, $w \rightarrow u \leftarrow w'$ is a v-structure of M , and $w, u, w' \in V_{M'}$, from Definition 2.41, $w \rightarrow u \leftarrow w' \in M'$, which is a contradiction, as $w - u - w' \in M'$. Thus the only option that remains is $u \rightarrow v \in M$.

We have shown that $u \rightarrow v \in M$ in each case. This proves Lemma 2.45. □

Lemma 2.45 further implies the following corollaries:

Corollary 2.47. *Let G be a graph, $S \subseteq V_G$, $M \in \text{MEC}(G)$, and $\mathcal{P}(M, S) = M'$. If $u \rightarrow v \in M$, and $u, v \in S$ then either $u \rightarrow v \in M'$, or $u - v \in M'$.*

Proof. Suppose $u, v \in S$ and $u \rightarrow v \in M$. From the construction of projection, $\text{skeleton}(M') = \text{skeleton}(M[S])$. This implies if $u \rightarrow v \in M$ then either $u \rightarrow v \in M'$ or $v \rightarrow u \in M'$ or $u - v \in M'$. But, if $v \rightarrow u \in M'$ then from Lemma 2.45, $v \rightarrow u \in M$, a contradiction. This implies either $u \rightarrow v \in M'$, or $u - v \in M'$. □

Corollary 2.48. *Let G be a graph, $S \subseteq V_G$, $M \in \text{MEC}(G)$, and $\mathcal{P}(M, S) = M'$. If $u - v \in M$, and $u, v \in S$ then $u - v \in M'$.*

Proof. Suppose $u, v \in S$ and $u - v \in M$. From the construction of projection, $\text{skeleton}(M') = \text{skeleton}(M[S])$. This implies if $u - v \in M$ then either $u \rightarrow v \in M'$ or $v \rightarrow u \in M'$ or $u - v \in M'$. From Lemma 2.45, if there is a directed edge between u and v in M' then M also has a directed edge between u and v , a contradiction. This implies $u - v \in M'$. □

Corollary 2.49. *Let M be an MEC and M' be a projection of M . Let $P = (u_1, u_2, \dots, u_l)$ be a triangle-free path in M such that all the nodes of P are in M' , i.e., for all $1 \leq i \leq l$, $u_i \in V_{M'}$ then P is a triangle-free path in M' .*

Proof. Suppose $P = (u_1, u_2, \dots, u_l)$ is a triangle-free path in M such that all the nodes of P are in M' . From Lemma 2.45 and Corollary 2.48, P is a path in M' . P must be a triangle-free path in M' , otherwise, if there exist u_{i-1}, u_i, u_{i+1} such that there is an edge between u_{i-1} and u_{i+1} in M' (of any form undirected or directed i.e., either $u_{i-1} - u_{i+1} \in M'$, or $u_{i-1} \rightarrow u_{i+1} \in M'$, or $u_{i+1} \rightarrow u_{i-1} \in M'$) then even in M there is an edge between u_{i-1} and u_{i+1} (since M' is a projection of M , if $u - v \in \text{skeleton}(M')$ then $u - v \in \text{skeleton}(M[V_M])$). This implies P is not a triangle-free path in M , a contradiction. This implies P is a triangle-free path in M' . \square

The following corollary is a partial converse of Corollary 2.49.

Corollary 2.50. *Let M be an MEC and M' be a projection of M . Let $P = (u_1, u_2, \dots, u_l)$ be a triangle-free path in M' . If $u_2 \rightarrow u_1 \notin M$ then P is also a triangle-free path in M .*

Proof. Suppose that $u_2 \rightarrow u_1 \notin M$. Since P is a triangle-free path in M' , if P is a path in M then it is a triangle-free path in M . We show that P is a path in M . Suppose that P is not a path in M . Then there exists an edge $u_i \leftarrow u_{i+1} \in M$. Pick the least i such that $u_i \leftarrow u_{i+1} \in M$. Clearly, $i \geq 2$ (from our assumption $u_2 \rightarrow u_1 \notin M$). Since P is a triangle-free path in M' , there does not exist an edge in between u_{i-1} and u_{i+1} . This implies that either $u_{i-1} - u_i \leftarrow u_{i+1} \in M$ or $u_{i-1} \rightarrow u_i \leftarrow u_{i+1} \in M$. From item 2 of Theorem 2.18, $u_{i-1} - u_i \leftarrow u_{i+1} \notin M$. From Definition 2.41, $\mathcal{V}(M') = \mathcal{V}(M[V_{M'}])$. This implies if the v-structure $u_{i-1} \rightarrow u_i \leftarrow u_{i+1} \in M$ then we have $u_{i-1} \rightarrow u_i \leftarrow u_{i+1} \in M'$, as $u_{i-1}, u_i, u_{i+1} \in V_{M'}$. But, then P is not a path in M' (because $u_i \leftarrow u_{i+1} \in M'$), which is a contradiction. This implies that P is a path in M . \square

The following corollary is an extension of Corollary 2.50.

Corollary 2.51. *Let M be an MEC and M' be a projection of M . If $P = (u_1 = x, u_2 = y, u_3, \dots, u_{l-1} = u, u_l = v)$ is a triangle-free path from (x, y) to (u, v) in M' , and $x \rightarrow y \in M$ then for all $1 \leq i < l$, $u_i \rightarrow u_{i+1} \in M$.*

Proof. For $i = 1$, Corollary 2.51 is true as $u_1 \rightarrow u_2 \in M$. Pick the least $i > 1$ such that $u_i \rightarrow u_{i+1} \notin M$. If $u_i - u_{i+1} \in M$ then $u_{i-1} \rightarrow u_i - u_{i+1} \in M$ which disobeys item 2 of Theorem 2.18. And, if $u_i \leftarrow u_{i+1} \in M$ then $u_{i-1} \rightarrow u_i \leftarrow u_{i+1}$ is a v-structure in M . This implies that $u_{i-1} \rightarrow u_i \leftarrow u_{i+1} \in M'$, as from from Definition 2.41, $\mathcal{V}(M[V_{M'}]) = \mathcal{V}(M')$. But, then, P is not a path in M' , a contradiction. This implies that no such i exists, i.e., for all i , $u_i \rightarrow u_{i+1} \in M$. \square

The following proposition is due to the non-existence of a directed cycle in an MEC.

Proposition 2.52. *Let M be an MEC, and \mathcal{C} be an undirected connected component of M . There cannot be a directed edge between two vertices of \mathcal{C} , i.e., if $u, v \in \mathcal{C}$ and there is an edge between u and v in M then $u - v \in M$.*

Proof. Suppose that $u, v \in \mathcal{C}$, and $u \rightarrow v \in M$. Since $u, v \in \mathcal{C}$, there exists an undirected path from v to u . Combining the undirected path with $u \rightarrow v$ gives a directed cycle in M , which contradicts item 1 of Theorem 2.18. This implies if there is an edge between u and v then it must be undirected. This proves Proposition 2.52. \square

The following lemma shows the unidirectional nature of triangle-free paths in an MEC. We use this later for the verification of an MEC.

Lemma 2.53. *Let M be an MEC of an undirected graph G . For $(u, v), (x, y) \in E_G$ such that $(u, v) \neq (x, y)$, if there exists a triangle-free path from (u, v) to (x, y) in M then there cannot exist a triangle-free path from (x, y) to (u, v) .*

Proof. Suppose, if possible, that there exist triangle-free paths P_1 from (u, v) to (x, y) and P_2 from (x, y) to u (P_2 is a prefix of the hypothesised triangle-free path from (x, y) to (u, v)). Then by Observation 2.28, the concatenation Q of P_1 and P_2 (in the sense of Observation 2.28) is a triangle-free path from (u, v) to

u . But this is a contradiction, since by the definition of path, Definition 2.1, no vertex can appear twice in a path. This implies there cannot exist a triangle-free path from (x, y) to u . Thus, there cannot exist a triangle-free path from (x, y) to (u, v) . \square

We now describe our approach to counting MECs of an undirected graph.

3 Counting MECs of a Graph

Let G be an undirected graph. We want to count MECs that have the same skeleton as G , i.e., $|\text{MEC}(G)|$. We provide a formal description of the problem.

Problem 1 (Counting MECs of an Undirected Graph). *Input:* An undirected graph G .

Output: $|\text{MEC}(G)|$.

We present a recursive algorithm to solve Problem 1. For this recursion, we divide the input graph using its tree decomposition. We construct a tree decomposition $(X = \{X_1, X_2, \dots, X_s\}, T)$ of G . We select node X_1 as the root node of T . For each node X_i in T , we denote the subtree of T rooted at X_i as T_i . The induced subgraph of G represented by T_i is denoted as G_i . This implies that $T = T_1$, $G = G_1$, and our goal is to count the MECs of G_1 .

In the subtree T_i , let the children of X_i be denoted as $(X_{i1}, X_{i2}, \dots, X_{il})$ (i.e., for T_i , we consider X_{i1} as the first child of X_i , X_{i2} as the second child of X_i , and so on). For any $0 \leq j \leq l$, we denote T_i^j as the induced subtree of T_i that contains node X_i and the nodes of $T_{i1}, T_{i2}, \dots, T_{ij}$. In other words, T_i^0 contains only one node, X_i , and $T_i^l = T_i$. The induced subgraph of G represented by T_i^j is denoted as G_i^j . If the number of children of X_1 is p , then $T_1^p = T_1 = T$, and $G_1^p = G_1 = G$, and **we want to count MECs of G_1^p** . Refer to Figure 3 for further clarity.

Let $X_{i1}, X_{i2}, \dots, X_{il}$ be the children of X_i in the tree T_i . For $1 \leq j \leq l$, Observation 3.1 establishes a relationship between G_i^j , G_i^{j-1} , and G_{ij} .

Observation 3.1. *Let G be an undirected graph, and $(X = \{X_1, X_2, \dots, X_s\}, T)$ be a tree decomposition of G . For $1 \leq i \leq s$, let $X_{i1}, X_{i2}, \dots, X_{il}$ be the children of X_i in the tree T_i . Then, for all $j > 0$:*

1. G_i^{j-1} and G_{ij} are induced subgraphs of G_i^j .
2. $G_i^j = G_i^{j-1} \cup G_{ij}$.
3. $X_i \subseteq V_{G_i^{j-1}}$.
4. $X_{ij} \subseteq V_{G_{ij}}$.
5. $I_j = X_i \cap X_{ij}$ is a vertex separator of G_i^j that separates $V_{G_i^{j-1}} \setminus I_j$ and $V_{G_{ij}} \setminus I_j$.
6. $V_{G_i^{j-1}} \cap V_{G_{ij}} = X_i \cap X_{ij} = I_j$

Proof. From the construction of G_i^j, G_i^{j-1} and G_{ij} , items 1 to 4 are true. Since T_i^j is a tree decomposition of G_i^j , $I_j = X_i \cap X_{ij}$ serves as a vertex separator of G_i^j that separates $V_{G_i^{j-1}} \setminus I_j$ and $V_{G_{ij}} \setminus I_j$, validating item 5. The validation of item 6 comes from the fact that $X_i \subseteq V_{G_i^{j-1}}$, $X_{ij} \subseteq V_{G_{ij}}$, and $X_i \cap X_{ij}$ is a vertex separator of G_i^j that separates $V_{G_i^{j-1}} \setminus I_j$ and $V_{G_{ij}} \setminus I_j$. \square

Remember from the above discussion that we want to count MECs of G_1^p , where p is the number of children of the root node X_1 in the tree decomposition of G . Observation 3.1 shows that if we cut the edge $X_i - X_{ij}$ in the tree T_i^j then we get two subtrees T_i^{j-1} and T_{ij} of T_i . This further gives us two induced subgraphs G_i^{j-1} and G_{ij} of G_i^j that have the properties items 1 to 6 of Observation 3.1. We establish a relation between the MECs of G_i^j, G_i^{j-1} , and G_{ij} that if we have the knowledge about (a) $|\text{MEC}(G_i^{j-1}, O_1, P_{11}, P_{12})|$




Figure 3: Example: G is an undirected graph, and T is a tree decomposition of G where: $X_1 = \{1, 2, 3\}$, $X_2 = \{2, 4, 5\}$, $X_3 = \{2, 3, 6\}$, $X_4 = \{3, 7, 8\}$, $X_5 = \{5, 9, 10\}$, $X_6 = \{7, 11, 12\}$, $X_7 = \{8, 13, 14\}$, $X_8 = \{12, 15, 16\}$, and $X_9 = \{14, 17\}$. T_4 , T_6 , and T_7 are induced subtrees of T rooted at X_4 , X_6 , and X_7 , respectively. We assume X_6 and X_7 are the first and second children of X_4 . T_4^0 is an induced subgraph of T containing only the node X_4 . T_4^1 is an induced subtree of T containing node X_4 and the nodes of T_6 , and T_4^2 is an induced subtree of T_4 containing node X_4 and the nodes of T_6 and T_7 . G_4 is the induced subgraph of G represented by T_4 . G_4^0 is the induced subgraph of G represented by T_4^0 . G_4^1 is the induced subgraph of G represented by T_4^1 . G_4^2 is the induced subgraph of G represented by T_4^2 . Since $T_4 = T_4^2$, it follows that $G_4 = G_4^2$.

for each shadow (O_1, P_{11}, P_{12}) of $G_i^{j-1}[X_i \cup N(X_i, G_i^{j-1})]$, and (b) $|\text{MEC}(G_{i_j}, O_2, P_{21}, P_{22})|$ for each shadow (O_2, P_{21}, P_{22}) of $G_{i_j}[X_{i_j} \cup N(X_{i_j}, G_{i_j})]$ then we can compute $|\text{MEC}(G_i^j, O, P_1, P_2)|$ for each shadow (O, P_1, P_2) of $G_i^j[X_i \cup N(X_i, G_i^j)]$. This gives us a recursive method to compute $|\text{MEC}(G_1^p, O, P_1, P_2)|$ for each shadow (O, P_1, P_2) of G_1^p on $X_1 \cup N(X_1, G_1^p)$. Then, with the assistance of Lemma 2.40, we can effectively count the MECs of $G = G_1^p$.

We first show the relation between $\text{MEC}(G_i^j)$, $\text{MEC}(G_i^{j-1})$ and $\text{MEC}(G_{i_j})$, using which we construct a recursive algorithm to count the MECs of G . To understand this relation, we first go through a similar simple scenario. Let us consider an undirected graph H . Let H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1, S_2 be the subsets of V_{H_1} and V_{H_2} , respectively, such that both S_1 and S_2 contains nodes of I , i.e., $S_1 \cap S_2 = I$. Here, H resembles G_i^j , H_1 resembles G_i^{j-1} , H_2 resembles G_{i_j} , S_1 resembles X_i , S_2 resembles X_{i_j} , and I resembles I_j . We show some relation between MECs of H, H_1 , and H_2 .

For given shadows (O_1, P_{11}, P_{12}) of $H_1[S_1 \cup N(S_1, H_1)]$, (O_2, P_{21}, P_{22}) of $H_2[S_2 \cup N(S_2, H_2)]$ and (O, P_1, P_2) of $H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)]$, and MECs $M \in \text{MEC}(H, O, P_1, P_2)$, $M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12})$, and $M_2 \in \text{MEC}(H_2, O_2, P_{21}, P_{22})$, we first show a necessary condition for the shadow (O, P_1, P_2) such that $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$.

3.1 Necessary condition for a shadow of an MEC

Lemma 3.2. *Let H be an undirected graph, and let H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1 and S_2 be subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let $Y = S_1 \cup S_2 \cup N(S_1 \cup S_2, H)$, $Y_1 = S_1 \cup N(S_1, H_1)$, and $Y_2 = S_2 \cup N(S_2, H_2)$. Let M , M_1 , and M_2 be the MECs of H , H_1 , and H_2 , respectively. Let $(O, P_1, P_2) = \text{shadow}(M, Y)$, $(O_1, P_{11}, P_{12}) = \text{shadow}(M_1, Y_1)$, and $(O_2, P_{21}, P_{22}) = \text{shadow}(M_2, Y_2)$. If $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$, then:*

1. For $a \in \{1, 2\}$, if $u \rightarrow v \in O_a$, then $u \rightarrow v \in O$.
2. For $a \in \{1, 2\}$, $\mathcal{V}(O_a) = \mathcal{V}(O[V_{O_a}])$ (see Definition 2.17 for $\mathcal{V}(G)$).
3. For $a \in \{1, 2\}$, if $u - v \in O_a$, then $u \rightarrow v \in O$ if, and only if, either of the following occurs:
 - (a) $u \rightarrow v$ is strongly protected in O .
 - (b) There exists $x - y \in O_a$ such that $x \rightarrow y \in O$, and $P_{a1}((x, y), (u, v)) = 1$ (i.e., there exists a triangle-free path from (x, y) to (u, v) in M_a with $(x, y) \neq (u, v)$).
 - (c) There exists $x - y \in O_a$ such that $x \rightarrow y \in O$, and $P_{a2}((x, y), v) = P_{a2}((v, u), x) = 1$ (i.e., $y \neq v$, $u \neq x$, and there exist triangle-free paths in M_a from (x, y) to v , and from (v, u) to x).
4. For $((u, v), (x, y)) \in E_O \times E_O$, $P_1((u, v), (x, y)) = 1$ if, and only if, $(u, v) \neq (x, y)$ and at least one of the following occurs:
 - (a) There exists a triangle-free path from (u, v) to (x, y) in O .
 - (b) For $a \in \{1, 2\}$, $P_{a1}((u, v), (x, y)) = 1$.
 - (c) There exists $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$.
5. For $((u, v), w) \in E_O \times V_O$, $P_2((u, v), w) = 1$ if, and only if, $v \neq w$ and at least one of the following occurs:
 - (a) There exists a triangle-free path from (u, v) to w in O .
 - (b) For $a \in \{1, 2\}$, $P_{a2}((u, v), w) = 1$.
 - (c) There exists $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$.

Proof of Lemma 3.2. Let $M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12})$, $M_2 \in \text{MEC}(H_2, O_2, P_{21}, P_{22})$, and $M \in \text{MEC}(H, O, P_1, P_2)$ be such that $\mathcal{P}(M, V_{M_1}, V_{M_2}) = (M_1, M_2)$.

Proof of item 1: Suppose for $a \in \{1, 2\}$, $u \rightarrow v \in O_a$. From the construction, $u, v \in V_{O_a} = S_a \cup N(S_a, H_a) \subseteq S_1 \cup S_2 \cup N(S_1 \cup S_2, H)$. From Definition 2.38, O_a is an induced subgraph of M_a on $S_a \cup N(S_a, H_a)$. This implies that $u \rightarrow v \in M_a$. Since M_a is a projection of M , from Lemma 2.45, $u \rightarrow v \in M$. From Definition 2.38, $O = M[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)]$. This implies that $u \rightarrow v \in O$. \square

Proof of item 2. From our assumption, for $a \in \{1, 2\}$, $M_a \in \text{MEC}(H_a, O_a, P_{a1}, P_{a2})$. This implies (O_a, P_{a1}, P_{a2}) is the shadow of M_a on $V_{O_a} = S_a \cup N(S_a, H_a)$. Then, from item 1 of Definition 2.36, $O_a = M_a[S_a \cup N(S_a, H_a)]$. This implies that $\mathcal{V}(O_a) = \mathcal{V}(M_a[S_a \cup N(S_a, H_a)])$. Since M_a is a projection of M onto V_{H_a} , and $S_a \cup N(S_a, H_a) \subseteq V_{H_a}$, Corollary 2.44 implies that $\mathcal{V}(M_a[S_a \cup N(S_a, H_a)]) = \mathcal{V}(M[S_a \cup N(S_a, H_a)])$. Furthermore, considering that O is an induced subgraph of M over $S_1 \cup S_2 \cup N(S_1 \cup S_2, H)$, and given that $S_a \cup N(S_a, H_a) \subseteq S_1 \cup S_2 \cup N(S_1 \cup S_2, H)$, we can conclude that $\mathcal{V}(M[S_a \cup N(S_a, H_a)]) = \mathcal{V}(O[S_a \cup N(S_a, H_a)])$. Thus, we have successfully established the relationship $\mathcal{V}(O_a) = \mathcal{V}(O[S_a \cup N(S_a, H_a)])$. This further implies that $\mathcal{V}(O_a) = \mathcal{V}(O[V_{O_a}])$, as from the construction, $V_{O_a} = S_a \cup N(S_a, H_a)$. \square

Proof of item 3. We present a more robust result than item 3 from Lemma 3.2.

Claim 3.3. For $a \in \{1, 2\}$, for an edge $u - v \in M_a$, $u \rightarrow v \in M$ if and only if either of the following occurs:

1. $u \rightarrow v$ is strongly protected in O .
2. There exists an undirected edge $x - y \in O_a$ such that $x \rightarrow y \in O$, and there exists a triangle-free path $Q = (u_1 = x, u_2 = y, \dots, u_{l-1} = u, u_l = v)$ from (x, y) to (u, v) in M_a where $l \geq 3$.
3. There exists an undirected edge $x - y \in O_a$ such that $x \rightarrow y \in O$, and there exist triangle-free paths $Q_1 = (u_1 = x, u_2 = y, \dots, u_l = v)$ from (x, y) to v in M_a , and $Q_2 = (v_1 = v, v_2 = u, \dots, v_m = x)$ from (v, u) to x in M_a where $l, m \geq 3$.

Proof. Observations 3.4 to 3.6 prove the \rightarrow direction of Claim 3.3.

Observation 3.4. If $u \rightarrow v$ is strongly protected in O , then $u \rightarrow v \in M$.

Proof. The fact that $u \rightarrow v$ is strongly protected in O implies that $u \rightarrow v \in O$. From our assumption, $M \in \text{MEC}(H, O, P_1, P_2)$. From Definition 2.38, this implies (O, P_1, P_2) is the shadow of M on Y . Therefore, from item 1 of Definition 2.36, O is an induced subgraph of M on Y . Thus, $u \rightarrow v \in O$ implies that $u \rightarrow v \in M$. \square

Observation 3.5. For $u - v \in M_a$, if there exists $x - y \in O_a$ such that there is a triangle-free path $Q = (u_1 = x, u_2 = y, \dots, u_{l-1} = u, u_l = v)$ from (x, y) to (u, v) in M_a , and $x \rightarrow y \in O$, then for all $1 \leq i < l$, $u_i \rightarrow u_{i+1} \in M$, more specifically, $u \rightarrow v \in M$.

Proof. Suppose $u - v, x - y \in M_a$, $x \rightarrow y \in O$, and $Q = (u_1 = x, u_2 = y, \dots, u_{l-1} = u, u_l = v)$ is a triangle-free path from (x, y) to (u, v) in M_a . Since O is an induced subgraph of M , and $x \rightarrow y \in O$, it follows that $x \rightarrow y \in M$. Then, according to Corollary 2.51, for all $1 \leq i < l$, $u_i \rightarrow u_{i+1} \in M$, and more specifically, $u \rightarrow v \in M$. \square

Observation 3.6. For $u - v \in M_a$, if there exists an undirected edge $x - y \in O_a$ such that $x \rightarrow y \in O$, and in M_a , there exist triangle-free paths $Q_1 = (u_1 = x, u_2 = y, \dots, u_l = v)$ from (x, y) to v in M_a , and $Q_2 = (v_1 = v, v_2 = u, \dots, v_m = x)$ from (v, u) to x in M_a , where $l, m \geq 3$, then $u \rightarrow v \in M$.

Proof. Suppose $u - v, x - y \in O_a$, $x \rightarrow y \in O$, $Q_1 = (u_1 = x, u_2 = y, \dots, u_l = v)$ is a triangle-free path from (x, y) to v in M_a , and $Q_2 = (v_1 = v, v_2 = u, \dots, v_m = x)$ is a triangle-free path from (v, u) to x in M_a , with $l, m \geq 3$.

Since O is an induced subgraph of M and $x \rightarrow y \in O$, it follows that $x \rightarrow y \in M$. Then, according to Corollary 2.50, Q_1 is a triangle-free path from (x, y) to v in M . Since $x \rightarrow y \in M$, Q_1 is a directed path in M .

Q_2 cannot be a path in M , otherwise, we would obtain a directed cycle in M by concatenating Q_1 and Q_2 , which contradicts item 1 of Theorem 2.18. But then, from Corollary 2.50, if $u \rightarrow v \notin M$ then Q_2 is a triangle-free path in M , a contradiction. This implies $u \rightarrow v \in M$. This completes the proof of Observation 3.6. \square

We now prove the \leftarrow direction of Claim 3.3. W.l.o.g., let $a = 1$. Since M is a chain graph, $M[V_{M_1}]$ is also a chain graph. Then from Proposition 2.5, there must exist an ordering τ of the vertices of M_1 such that for $u, v \in V_{M_1}$, if $u \rightarrow v \in M$ then $\tau(u) < \tau(v)$. Pick a least rank node v in τ such that for any edge $x - y \in O_1$, if $x \rightarrow y \in M$, and $\tau(y) < \tau(v)$ then $x \rightarrow y$ obeys Claim 3.3 (i.e., $x \rightarrow y$ either follows item 1 of Claim 3.3, or it follows item 2 of Claim 3.3, or it follows item 3 of Claim 3.3). After picking such a node v , pick highest rank node u such that $u - v \in M_1$, $u \rightarrow v \in M$, and for any node $z \in V_{M_1}$ such that $\tau(u) < \tau(z) < \tau(v)$, if $z - v \in M_1$, and $z \rightarrow v \in M$ then $z \rightarrow v$ obeys Claim 3.3. We show that $u \rightarrow v$ also obeys Claim 3.3. Our selection of $u - v \in M_1$ implies the following claim:

Claim 3.7. For any edge $u' - v' \in M_1$ such that $u' \rightarrow v' \in M$, if $\tau(v') < \tau(v)$, or $v' = v$ and $\tau(u) < \tau(u')$ then $u' \rightarrow v'$ obeys Claim 3.3, i.e., either

1. $u' \rightarrow v'$ is strongly protected in O , or
2. there exists $x - y \in O_1$ such that $x \rightarrow y \in O$ and there exists a triangle-free path from (x, y) to (u', v') in M_1 of length greater than one (i.e., the path containing more than two nodes), or
3. there exists $x - y \in O_1$ such that $x \rightarrow y \in O$ and there exist triangle-free paths in M_1 from (x, y) to v' , and from (v', u') to x such that each triangle-free path is of length greater than one.

Since M is an MEC, and $u \rightarrow v \in M$, from item 4 of Theorem 2.18, $u \rightarrow v$ must be strongly protected in M . This implies that $u \rightarrow v$ is a part of one of the induced subgraphs in fig. 2. We show that $u \rightarrow v$ obeys Claim 3.3 in each such possible induced subgraph of M . We now go through all the possibilities.

Case 1: Suppose $u \rightarrow v$ is strongly protected in M because it is part of an induced subgraph $w \rightarrow u \rightarrow v \in M$ (similar to Figure 2.a). There are two possibilities: either $w \in V_{H_1}$ or $w \notin V_{H_1}$.

If $w \notin V_{H_1}$, then $w \in V_{H_2} \setminus I$. This implies that $u \in I$, as $u \in V_{H_1}$ is a neighbor of $w \in V_{H_2} \setminus I$ in H , and I is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. This, in turn, implies that $u, v, w \in I \cup N(I, H) \subseteq V_O$, and the induced subgraph is also a part of O . Thus, $u \rightarrow v$ is strongly protected in O , as shown in fig. 2.a. Let's now consider the other possibility.

Suppose $w \in V_{H_1}$. Since M_1 is a projection of M onto V_{H_1} , we can deduce, from Corollary 2.47 and item 3 of Theorem 2.18, that $w - u \in M_1$. Also, as $w \rightarrow u \rightarrow v \in M$ and by the construction of τ , we have $\tau(w) < \tau(u) < \tau(v)$. According to Claim 3.7, $w \rightarrow u$ satisfies Claim 3.3, specifically, either (1) item 1, (2) item 2, or (3) item 3. We will explore each possibility and demonstrate that in each case, $u \rightarrow v$ obeys Claim 3.3.

1. Suppose $w \rightarrow u$ satisfies item 1. This implies that $w \rightarrow u$ is strongly protected in O , and thus, $w \rightarrow u \in O$. In this scenario, $u \rightarrow v$ obeys item 2, since there exists a triangle-free path $P = (w, u, v)$ from $w - u$ to $u - v$ in M_1 (with length greater than one), and $w \rightarrow u \in O$.

2. Suppose $w \rightarrow u$ satisfies item 2. In this case, there exists an undirected edge $x - y \in O_1$ such that $x \rightarrow y \in O$, and a triangle-free path $Q = (u_1 = x, u_2 = y, \dots, u_{l-1} = w, u_l = u)$ from (x, y) to (w, u) in M_1 , with $l \geq 3$. We first demonstrate that Q does not contain v . According to Observation 2.30, Q forms a chordless path in M_1 . If Q contains v , then there must be some $i < l - 1$ where $u_i = v$ (since $u_l = u$ and $u_{l-1} = w$). This would lead to a contradiction as Q would no longer be a chordless path. Thus, Q does not contain v .

As $Q' = (w, u, v)$ is a triangle-free path from (w, u) to (u, v) in M_1 , we can use Observation 2.28 to conclude that $Q'' = (u_1 = x, u_2 = y, \dots, u_{l-1} = w, u_l = u, v)$ is a triangle-free path from (x, y) to (u, v) in M_1 (by concatenating Q and Q'). Consequently, $u \rightarrow v$ satisfies item 2 as Q'' is a triangle-free path in M_1 (of length greater than one, given that $l \geq 3$), and $x \rightarrow y \in O$.

3. Suppose $w \rightarrow u$ satisfies item 3. This implies that there exists an undirected edge $x - y \in M_1$, where $x \rightarrow y \in O$, and there are triangle-free paths $Q_1 = (u_1 = x, u_2 = y, \dots, u_l = u)$ from (x, y) to u in M_1 , and $Q_2 = (v_1 = u, v_2 = w, \dots, v_m = x)$ from (u, w) to x in M_1 , with both l and m are greater than two. No edge in Q_1 and Q_2 is directed in M_1 . A directed edge in Q_1 or Q_2 implies a directed cycle in M_1 (which we get by concatenating Q_1 and Q_2), contradicting item 1 of Theorem 2.18.

Therefore, both v and the nodes in Q_1 and Q_2 belong to the same undirected connected component \mathcal{C} of M_1 , and both Q_1 and Q_2 are undirected triangle-free paths in \mathcal{C} . By Proposition 2.25, both Q_1 and Q_2 are chordless paths in \mathcal{C} .

Now, let's consider $Q_3 = (u_1 = x, u_2 = y, \dots, u_l = u, u_{l+1} = v)$. If Q_3 is a chordless path in \mathcal{C} , it is also a chordless path in M_1 , as \mathcal{C} is an induced subgraph of M_1 . In this case, $u \rightarrow v$ satisfies item 2, as Q_3 is a triangle-free path in M_1 .

However, if Q_3 is not a chordless path in \mathcal{C} , this implies that there must exist an index $i \leq l - 1$ where an edge exists between u_i and v in \mathcal{C} . According to Proposition 2.52, $u_i - v$ is an undirected edge in \mathcal{C} . Since Q_1 is a triangle-free path in M_1 and $x \rightarrow y \in M$, from Corollary 2.51, Q_1 forms a fully

directed path in M . Moreover, as Q_1 is fully directed and $u \rightarrow v \in M$, Q_3 is also fully directed in M . Since Q_3 passes through u_i , there exists a directed path from u_i to v in M . This implies that $u_i \rightarrow v \in M$. Otherwise, M would have a directed cycle $(u_i, u_{i+1}, \dots, u_l = u, v, u_i)$, which contradicts item 1 of Theorem 2.18, as M is an MEC.

If $i < l - 1$, then $u_i \rightarrow v \leftarrow u_l$ forms a v-structure in M (since there is no edge between u_i and u_l in M , as P_1 is a chordless path). However, this leads to a contradiction because it would imply that $\mathcal{V}(M[V_{M_1}]) \neq \mathcal{V}(M_1)$ (as $u_i \rightarrow v \leftarrow u_l \in M$ while $u_i - v - u_l \in M_1$). But, from Definition 2.41, since M_1 is a projection of M , both M_1 and $M[V_{M_1}]$ must have the same set of v-structures. Thus, the only possible value of i is $l - 1$. In this case, $Q_4 = (u_1 = x, u_2 = y, \dots, u_{l-1}, v)$ is a chordless path in M_1 . Additionally, since $Q_5 = (v, u, w)$ is a triangle-free path in M_1 from (v, u) to (u, w) , and Q_2 is a triangle-free path in M_1 from (u, w) to x , we can use Observation 2.28 to show that $Q_6 = (v, u, w, v_3, v_4, \dots, v_m = x)$ is a triangle-free path from (v, u) to (w, x) in M_1 . Both Q_4 and Q_6 have lengths greater than one. Hence, $u \rightarrow v$ satisfies item 3 of Claim 3.3 as Q_4 is a triangle-free path from (x, y) to v in M_1 , Q_6 is a triangle-free path from (v, u) to x in M_1 , and $x \rightarrow y \in O$.

This implies that in all the cases of Case 1, $u \rightarrow v$ obeys item 3.

Case 2: Consider the scenario where $u \rightarrow v$ is strongly protected in M because it is part of an induced subgraph $u \rightarrow v \leftarrow w \in M$ (similar to Figure 2.b). There are two possibilities: either $w \notin V_{H_1}$ or $w \in V_{H_1}$.

If $w \notin V_{H_1}$, as demonstrated in Case 1, the induced subgraph is also an induced subgraph of O . Therefore, $u \rightarrow v$ is strongly protected in O , and it obeys item 1 of Claim 3.3. Let's now consider the other possibility.

Suppose $w \in V_{H_1}$. Then, $v \rightarrow u \leftarrow w$ is a v-structure in M such that $u, v, w \in V_{M_1}$. Since M_1 is a projection of M onto V_{H_1} , from Definition 2.41, the set of v-structures of M_1 and the set of v-structures of $M[V_{M_1}]$ must be the same. This implies $u \rightarrow v \leftarrow w \in M_1$. This leads to a contradiction since our assumption is that $u - v \in M_1$. Hence, it implies that $w \notin V_{H_1}$.

Case 3: Suppose $u \rightarrow v$ is strongly protected in M because it is part of an induced subgraph $u \rightarrow w \rightarrow v \leftarrow u \in M$ (same as Figure 2.c). There are two possibilities: either $w \notin V_{H_1}$ or $w \in V_{H_1}$.

If $w \notin V_{H_1}$, as demonstrated in Case 1, the induced subgraph is also an induced subgraph of O . Therefore, $u \rightarrow v$ is strongly protected in O , and it adheres to item 1 of Claim 3.3. Let's now consider the other possibility.

Suppose $w \in V_{H_1}$. Since $u \rightarrow w$ and $w \rightarrow v \in M$, and since M_1 is a projection of M onto V_{H_1} , we can deduce from Corollary 2.47 that either $u \rightarrow w \in M_1$ or $u - w \in M_1$, and either $w \rightarrow v \in M_1$ or $w - v \in M_1$.

Given that $u - v \in M_1$, if $u \rightarrow w \in M_1$ or $w \rightarrow v \in M_1$, it would lead to a directed cycle (u, w, v, u) in M_1 , which contradicts item 1 of Theorem 2.18. Thus, we conclude that $u - w$ and $w - v \in M_1$. As $u \rightarrow w$ and $w \rightarrow v \in M$, based on the ordering in τ , it holds that $\tau(u) < \tau(w) < \tau(v)$. Referring to Claim 3.7, we see that both $u \rightarrow w$ and $w \rightarrow v$ adhere to Claim 3.3. This implies that either $u \rightarrow w$ obeys item 1, or it follows item 2, or it satisfies item 3. Let's examine each of these possibilities.

1. Suppose $u \rightarrow w$ obeys item 1 of Claim 3.3, i.e., $u \rightarrow w$ is strongly protected in O . This implies $u, w \in V_O$ and $u \rightarrow w \in O$. As shown earlier, $w \rightarrow v$ also obeys Claim 3.3.

This implies either $w \rightarrow v$ obeys item 1 of Claim 3.3, or it obeys item 2, or it obeys item 3 of Claim 3.3. We will consider each possibility.

- (a) Suppose $w \rightarrow v$ obeys item 1 of Claim 3.3, i.e., $w \rightarrow v$ is strongly protected in O . This implies $w, v \in V_O$ and $w \rightarrow v \in O$. Since $u, v, w \in V_O$, $u \rightarrow v \in M$, and O is an induced subgraph of M , we have $u \rightarrow v \in O$. This shows that $u \rightarrow v$ is also strongly protected in O , as it is part of the induced subgraph $u \rightarrow w \rightarrow v \leftarrow u \in O$. This implies $u \rightarrow v$ obeys item 1 of Claim 3.3.
- (b) Suppose $w \rightarrow v$ obeys item 2 of Claim 3.3, i.e., there exists an edge $x - y \in O_1$ such that there is a triangle-free path $Q = (u_1 = x, u_2 = y, \dots, u_{l-1} = w, u_l = v)$ from (x, y) to (w, v) in M_1 , with $l \geq 3$ and $x \rightarrow y \in O$.

According to Observation 2.30, Q is an undirected chordless path in M_1 . We first demonstrate that Q does not contain u . Suppose Q contains u , then for some $i \leq l - 2$, $u_i = u$ (since $u_l = v$

and $u_{l-1} = w$). However, this contradicts the fact that Q is a chordless path, as $u - v \in M_1$. Hence, Q does not contain u . Since $u - v \in M_1$, nodes of Q and u are in the same undirected connected chordal component of M_1 .

We now show that $u_{l-2} - u \in M_1$. From Observation 3.5, $u_{l-2} \rightarrow w \in M$, since Q is a triangle-free path in M_1 , and $x \rightarrow y \in M$ (because O is an induced subgraph of M and $x \rightarrow y \in O$). This implies there must be an edge between u_{l-2} and u , otherwise, $u_{l-2} \rightarrow w \leftarrow u$ would form a v-structure in M . Since M_1 is a projection of M on V_{H_1} and $u_{l-2}, w, u \in V_{H_1}$, if $u_{l-2} \rightarrow w \leftarrow u$ is a v-structure in M , then $u_{l-2} \rightarrow w \leftarrow u \in M_1$. But this contradicts the construction where $w - u \in M_1$. Therefore, there must be an edge between u_{l-2} and u . Since u_{l-2} and u are in the same undirected connected component of M_1 , from Proposition 2.6, $u_{l-2} - u \in M_1$.

We now consider the least i such that $u_i - u \in M_1$. If $i > 1$, then $Q' = (u_1 = x, u_2 = y, \dots, u_i, u, v)$ forms a chordless path from (x, y) to (u, v) , as v cannot be a neighbor of any u_i due to Q being a chordless path and $i \leq l - 2$. This implies that $u \rightarrow v$ obeys item 2 of Claim 3.3. If $i = 1$, then $Q_1 = (v, u, x)$ forms a triangle-free path from (v, u) to x in M_1 . Therefore, (u, v) obeys item 3 of Claim 3.3, as Q is a triangle-free path from (x, y) to v in M_1 (of length greater than one), Q_1 is a triangle-free path from (v, u) to x in M_1 (of length greater than one), and $x \rightarrow y \in O$. This concludes the analysis of this case, showing that in each possibility $u \rightarrow v$ obeys Claim 3.3.

- (c) Suppose $w \rightarrow v$ obeys item 3 of Claim 3.3, i.e., there exists an edge $x - y \in O_1$ such that $x \rightarrow y \in O$, and there exist triangle-free paths $Q_1 = (u_1 = x, u_2 = y, \dots, u_l = v)$ from (x, y) to v in M_1 , and $Q_2 = (v_1 = v, v_2 = w, \dots, v_m = x)$ from (v, w) to x in M_1 such that $l, m \geq 3$.

Q_1 and Q_2 must both be undirected paths in M_1 , otherwise combining Q_1 and Q_2 would create a directed cycle in M_1 . This implies that all nodes in Q_1 , all nodes in Q_2 , and u belong to the same undirected connected chordal component \mathcal{C} of M_1 . This further implies that Q_1 and Q_2 are the triangle-free paths of a chordal graph \mathcal{C} . Then, from Proposition 2.25, Q_1 and Q_2 are chordless paths of M_1 .

There are two possibilities: either u is in Q_1 , or u is not in Q_1 . If u is in Q_1 , then $u_{l-1} = u$, as Q_1 is a chordless path in \mathcal{C} . Then, Q_1 is a triangle-free path of M_1 from (x, y) to (u, v) . That means (u, v) obeys item 2 of Claim 3.3, as Q_1 is a triangle-free path from (x, y) to (u, v) in M_1 and $x \rightarrow y \in O$.

Now we consider the other possibility. Suppose u is not in Q_1 . There are two possibilities: either there is an edge between u_{l-1} and u in M , or there is no edge between them in M . If there is no edge between them in M , then $u_{l-1} \rightarrow v \leftarrow u$ forms a v-structure in M . Since M_1 is a projection of M and $u_{l-1}, v, u \in M_1$, the v-structure $u_{l-1} \rightarrow v \leftarrow u \in M$ implies $u_{l-1} \rightarrow v \leftarrow u \in M_1$. This implies $u \rightarrow v \in M_1$. However, this is a contradiction, as $u - v \in M_1$. This implies there is an edge between u_{l-1} and u in M . From Corollaries 2.47 and 2.48, there is an edge between u_{l-1} and u in M_1 as well. Since u_{l-1} and u are in the same undirected connected component of M_1 , therefore, from Proposition 2.6, $u_{l-1} - u \in M_1$.

We now consider a path $Q_3 := (u_1 = x, u_2 = y, \dots, u_{l-1}, u)$, obtained by replacing u_l with u in the chordless path Q_1 . There are two possibilities: either Q_3 is not a chordless path in \mathcal{C} , or Q_3 is a chordless path in \mathcal{C} . We show that in both the possibility $u \rightarrow v$ obeys Claim 3.3.

- i. Suppose Q_3 is not a chordless path in \mathcal{C} . Then there must exist a k such that $k < l - 1$ and $u_k - u \in \mathcal{C}$, because Q_1 is a chordless path and Q_3 is not a chordless path in \mathcal{C} . Pick the least k such that $u_k - u \in \mathcal{C}$. If $k > 1$, then this implies that $Q'_3 := (u_1 = x, u_2 = y, \dots, u_k, u, v)$ is a chordless path from (x, y) to (u, v) in \mathcal{C} (since Q_1 is a chordless path and there cannot be an edge between v and any u_i s in Q_3). This further implies that $u \rightarrow v$ obeys item 2 of Claim 3.3, as $x \rightarrow y \in O$ and Q'_3 is a triangle-free path from (x, y) to (u, v) in M_1 (since \mathcal{C} is an undirected connected component of M_1 and every chordless path is a triangle-free path). If $k = 1$, then $Q''_3 := (v, u, x)$ is a triangle-free path from (v, u) to x in M_1 (since Q_1 is a chordless path and there cannot be an edge between x and v). This implies that $u \rightarrow v$ obeys item 3 of Claim 3.3, due to the existence of triangle-free paths Q_1, Q''_3 , and $x \rightarrow y \in O$. This shows that in each possibility of this subcase, $u \rightarrow v$ obeys Claim 3.3.

- ii. Suppose Q_3 is a chordless path in \mathcal{C} . We claim that $Q_4 := (u, v_2 = w, v_3, \dots, v_m = x)$ (obtained by replacing the first vertex v of Q_2 by u) is not a chordless path in \mathcal{C} .

Proof of the claim. Suppose that Q_4 is a chordless path in \mathcal{C} . Since Q_3 is a chordless path in \mathcal{C} and \mathcal{C} is an undirected connected component of M_1 , from Observation 2.24, Q_3 is a triangle-free path in M_1 . Since $x \rightarrow y \in O$, and Q_3 is a triangle-free path in M_1 , from Observation 3.5, Q_3 is a directed path in M . Also, since $u \rightarrow w \in M$ and Q_4 is a chordless path in M_1 , from Corollary 2.51, Q_4 is a directed path in M . This creates a directed cycle in M by concatenating Q_3 and Q_4 . However, this is a contradiction, as stated in item 2 of Theorem 2.18, M cannot have a directed cycle. This proves that Q_4 is not a chordless path in M_1 . \square

Since Q_2 is a chordless path and Q_4 is not a chordless path, there must exist a $k > 2$ such that $u - v_k \in M_1$. Pick the highest k , i.e., for all $j > k$, $u - v_j \notin M_1$. Then $Q_5 := (u, v_k, v_{k+1}, \dots, v_m = x)$ is a chordless path in \mathcal{C} . Since Q_2 is a chordless path in M_1 , there cannot be an edge between v and v_i for $i \geq k > 2$. This further implies that $Q'_5 := (v, u, v_k, v_{k+1}, \dots, v_m = x)$ (obtained by adding v to Q_5) is a chordless path from (v, u) to x in M_1 . This implies that $u \rightarrow v$ obeys item 3 of Claim 3.3, since $x \rightarrow y \in O$, Q_1 is a triangle-free path from (x, y) to v in M_1 , and Q'_5 is a triangle-free path from (v, u) to x in M_1 (every chordless path is a triangle-free path).

This concludes the analysis of all possibilities in this case, showing that in each case $u \rightarrow v$ obeys Claim 3.3.

2. Suppose $u \rightarrow w$ obeys item 2 of Claim 3.3, i.e., there exists an edge $x - y \in O_1$ such that there exists a triangle-free path $Q := (u_1 = x, u_2 = y, \dots, u_{l-1} = u, u_l = w)$ from (x, y) to (u, w) in M_1 , and $x \rightarrow y \in O$. From Observation 2.30, Q is a chordless path in M_1 . We first show that Q does not contain v .

Suppose Q contains v . Then $u_{l-2} = v$, since Q is a chordless path. Then from Observation 3.5, $v \rightarrow u \in M$ (since Q is a chordless path from (x, y) to (u, w) , $x \rightarrow y \in O$, $u_{l-2} = v$, and $u_{l-1} = u$). However, this is a contradiction, as $u \rightarrow v \in M$. This implies v is not in Q .

Q not containing v implies that $Q_1 := (u_1 = x, u_2 = y, \dots, u_{l-1} = u, v)$, obtained by replacing w with v in Q , is a path in M_1 . We show that Q_1 is a chordless path in M_1 . This implies that $u \rightarrow v$ obeys item 2 of Claim 3.3, as $x \rightarrow y \in O$, and Q_1 is a triangle-free path from (x, y) to (u, v) in M_1 (every chordless path is a triangle-free path) of length greater than one (since $l \geq 3$). Thus, the only thing we need to complete this subcase is to show Q_1 is a chordless path in M_1 .

Suppose Q_1 is not a chordless path in M_1 . $Q' := (u_1 = x, u_2 = y, \dots, u_{l-1} = u)$, a subpath of the chordless path Q , is a chordless path from $x - y$ to u . Since Q' is a chordless path, and Q_1 is not a chordless path, there must exist an $i < l - 1$ such that $u_i - v \in M_1$. As $x \rightarrow y \in O$ and Q is a triangle-free path from (x, y) to (u, w) , from Observation 3.5, each edge of Q is directed in M . Since $u \rightarrow v \in M$, $Q'_1 := (u_i, u_{i+1}, \dots, u_{l-1} = u, v)$ is a directed path in M . This implies that $u_i \rightarrow v \in M$, otherwise, we get a directed cycle $C = (u_i, u_{i+1}, \dots, u_{l-1}, v, u_i)$ in M , which contradicts item 1 of Theorem 2.18. Since Q is a chordless path, there is no edge between w and u_i (since $i < l - 1$). However, then $w \rightarrow v \leftarrow u_i$ forms a v-structure in M . Since M_1 is a projection of M on V_{H_1} , and $w, v, u_i \in V_{H_1}$, from Definition 2.41, $w \rightarrow v \leftarrow u_i \in M_1$. But, this is a contradiction, as $w - v \in M_1$. This implies that Q_1 is a chordless path in M_1 .

As argued above this concludes the proof for the case when $u \rightarrow w$ obeys item 2 of Claim 3.3. We now move to the final possibility when $u \rightarrow w$ obeys item 3 of Claim 3.3.

3. Suppose $u \rightarrow w$ obeys item 3 of Claim 3.3, i.e., there exists an edge $x - y \in O_1$ such that in M_1 , there exist triangle-free paths $Q_1 := (u_1 = x, u_2 = y, \dots, u_l = w)$ from (x, y) to w , and $Q_2 := (v_1 = w, v_2 = u, \dots, v_m = x)$ from (w, u) to x . Q_1 and Q_2 both must be undirected paths in M_1 , otherwise, combining

Q_1 and Q_2 gives a directed cycle in M_1 . This implies that the nodes in Q_1 , nodes in Q_2 , and v , all belong to the same undirected connected chordal component \mathcal{C} of M_1 . This further implies that Q_1 and Q_2 are the triangle-free paths of a chordal graph \mathcal{C} . Then, from Proposition 2.25, Q_1 and Q_2 are chordless paths of \mathcal{C} . We now show that v is neither in Q_1 nor in Q_2 .

Suppose Q_1 contains v , i.e., for some $1 \leq i < l$, $u_i = v$. From Observation 3.5, each edge of Q_1 is directed in M , as Q_1 is a triangle-free path from (x, y) to (u_{l-1}, w) in M_1 , and $x \rightarrow y \in O$. This implies that $(u_i = v, u_{i+1}, \dots, u_l = w, v)$ forms a directed cycle in M , as from the construction, $w \rightarrow v \in M$. However, this contradicts item 1 of Theorem 2.18. Therefore, Q_1 does not contain v .

Suppose Q_2 contains v . Since Q_2 is a chordless path in M_1 , and $w - v \in M_1$, if Q_2 contains v , then $v_2 = v$. But, $v_2 = u$, which is a contradiction. Therefore, Q_2 does not contain v .

We now show that either $Q_3 := (u_1 = x, u_2 = y, \dots, u_l = w, v)$ (obtained by adding v to Q_1) or $Q_4 := (u_1 = x, u_2 = y, \dots, u_{l-1}, v)$ (obtained by replacing w with v in Q_1) is a chordless path in \mathcal{C} , and $Q_5 = (v, v_2 = u, v_3, \dots, v_m = x)$ (obtained by replacing the node w with v in Q_2) is a chordless path in \mathcal{C} . This will be our main ingredient to prove that $u \rightarrow v$ obeys Claim 3.3.

Suppose neither Q_3 nor Q_4 is a chordless path in \mathcal{C} . Since Q_3 is not a chordless path in \mathcal{C} and Q_1 is a chordless path in \mathcal{C} , for some $i \leq l-1$, $u_i - v \in \mathcal{C}$. Pick the least i . If $i \neq l-1$ then we get an undirected chordless cycle $(u_i, u_{i+1}, \dots, u_{l-1}, u_l, v, u_i)$ in the chordal graph \mathcal{C} , a contradiction. This implies $u_{l-1} - v \in \mathcal{C}$. Similarly, since Q_4 is not a chordless path in \mathcal{C} and Q_1 is a chordless path in \mathcal{C} , $u_{l-2} - v \in \mathcal{C}$. Since Q_1 is a triangle-free path in M_1 and $x \rightarrow y \in O$, from Observation 3.5, $u_{l-2} \rightarrow u_{l-1}, u_{l-1} \rightarrow w \in M$. Then, $u_{l-2} \rightarrow v \in M$, otherwise, we get a directed cycle $C = (u_{l-2}, u_{l-1}, w, v, u_{l-2})$ in M (remember $w \rightarrow v \in M$), which is a contradiction from item 1 of Theorem 2.18. But, then, $u_{l-2} \rightarrow v \leftarrow w$ is a v-structure in M (since Q_1 is a chordless path, there is no edge between u_{l-2} and w). Since M_1 is a projection of M on V_{H_1} , and $u_{l-2}, v, w \in V_{H_1}$, from Definition 2.41, $u_{l-2} \rightarrow v \leftarrow w \in M_1$. But $w - v$ is an undirected edge in M_1 , a contradiction. Thus, our assumption that neither Q_3 nor Q_4 is a chordless path in \mathcal{C} is wrong. This implies either Q_3 or Q_4 is a chordless path in \mathcal{C} . We now show that Q_5 is a chordless path in \mathcal{C} .

Since Q_2 is a triangle-free path in \mathcal{C} , from Observation 2.30, Q_2 is a chordless path in \mathcal{C} . This implies that the subpath $Q'_5 := (v_2 = u, v_3, \dots, v_m = x)$ of Q_2 is a chordless path in \mathcal{C} . Suppose Q_5 is not a chordless path in \mathcal{C} . Then, there exists a v_i such that $i > 2$, and $v - v_i \in \mathcal{C}$. Pick the highest i such that $v_i - v \in \mathcal{C}$. Then, $Q_6 := (v, v_i, v_{i+1}, \dots, v_m = x)$ is a chordless path from (v, v_i) to x in \mathcal{C} . Then, $v_i \rightarrow v \in M$. Otherwise, from Corollary 2.50, Q_6 is a triangle-free path in M , and the concatenation of Q_1 , $w \rightarrow v$, and Q_6 yield a directed cycle in M , contradicting item 1 of Theorem 2.18. Since Q_2 is a chordless path in M_1 , there cannot be an edge between w and v_i (as $i > 2$). This implies that $w \rightarrow v \leftarrow v_i$ is a v-structure in M . Since M_1 is a projection of M on V_{H_1} , and $w, v, v_i \in V_{H_1}$, from Definition 2.41, the v-structure $w \rightarrow v \leftarrow v_i \in M_1$. This gives a contradiction, as $w - v \in M_1$. This implies that Q_5 is a chordless path in M_1 .

Thus, in this case, there exists a triangle-free path from (x, y) to v , in the form of either Q_3 or Q_4 (from Observation 2.24, every chordless path is a triangle-free path) such that $x \rightarrow y \in O$, and there exists a triangle-free path from (v, u) to x (in the form of Q_5). Since $l, m \geq 3$, the length of all the paths Q_3, Q_4 , and Q_5 is greater than one. This implies that $u \rightarrow v$ obeys item 3 of Claim 3.3.

Case 4: Suppose $u \rightarrow v$ is strongly protected in M because it is part of an induced subgraph $u \rightarrow v \leftarrow w - u - w' \rightarrow v$ (same as Figure 2.d). Both w and w' cannot be in V_{H_1} , otherwise the v-structure $w \rightarrow v \leftarrow w' \in M_1$, as M_1 is a projection of M on V_{H_1} , and $w, v, w' \in V_{H_1}$. But, then, $u - v \notin M_1$, otherwise, it creates a directed cycle (w, v, u, w) , as from Corollary 2.48, $u - w \in M_1$. This implies that both w and w' cannot be in V_{H_1} . And, if w or w' is not in V_{H_1} then as we have seen in Case 1, the induced subgraph is an induced subgraph of O , i.e., $u \rightarrow v$ is strongly protected in O (as it is part of an induced subgraph of O as shown in fig. 2.d), and $u \rightarrow v$ obeys item 1 of Claim 3.3.

We show that in all the possibilities, $u \rightarrow v$ obeys Claim 3.3. This completes the proof of \leftarrow direction of Claim 3.3. This further completes the proof of Claim 3.3. \square

If there exists a triangle-free path P from (u, v) to (x, y) of length more than one then $(u, v) \neq (x, y)$. Similarly, if Q is a triangle-free path from (u, v) to w of length more than one then $v \neq w$. This shows that Claim 3.3 implies item 3 of Lemma 3.2. \square

Proof of item 4. Observations 3.8 to 3.10 prove \rightarrow of item 4.

Observation 3.8. *If $(u, v) \neq (x, y)$ and there exists a triangle-free path in O from (u, v) to (x, y) , then $P_1((u, v), (x, y)) = 1$.*

Proof. Since (O, P_1, P_2) is the shadow of M , as defined in item 1 of Definition 2.36, O is an induced subgraph of M . Moreover, according to item 2 of Definition 2.36, $P_1((u, v), (x, y))$ answers whether there exists a triangle-free path in M from (u, v) to (x, y) . If there exists a triangle-free path P in O from (u, v) to (x, y) , then P is also a triangle-free path in M , and therefore, $P_1((u, v), (x, y)) = 1$. \square

Observation 3.9. *For $(u, v), (x, y) \in E_O$, if $(u, v) \neq (x, y)$ and for any $a \in \{1, 2\}$, there exists a triangle-free path P in M_a from (u, v) to (x, y) , i.e., $P_{a1}((u, v), (x, y)) = 1$, then P is also a triangle-free path in M , i.e., $P_1((u, v), (x, y)) = 1$.*

Proof. Since (O, P_1, P_2) is the shadow of M , as defined in item 1 of Definition 2.36, O is an induced subgraph of M . Moreover, according to item 2 of Definition 2.36, $P_1((u, v), (x, y))$ answers whether there exists a triangle-free path in M from (u, v) to (x, y) . For $a \in \{1, 2\}$, suppose $(u, v) \neq (x, y)$ and P is a triangle-free path from (u, v) to (x, y) in M_a . Since $(u, v) \in E_O$, it follows that $v \rightarrow u \notin O$. This implies $v \rightarrow u \notin M$. Then, from Corollary 2.50, P is a triangle-free path in M , i.e., $P_1((u, v), (x, y)) = 1$. \square

Observation 3.10. *For $u, v, x, y, z_1, z_2 \in V_O$, if $(u, v) \neq (x, y)$, $P_1((u, v), (z_1, z_2)) = 1$, and $P_1((z_1, z_2), (x, y)) = 1$, then $P_1((u, v), (x, y)) = 1$.*

Proof. Since (O, P_1, P_2) is the shadow of M , as mentioned in item 2 of Definition 2.36, $P_1((u, v), (x, y))$ answers whether there exists a triangle-free path in M from (u, v) to (x, y) . Suppose $u, v, x, y, z_1, z_2 \in V_O$ such that $(u, v) \neq (x, y)$, and $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$. This implies there exist triangle-free paths $Q_1 = (u_1 = u, u_2 = v, \dots, u_{l-1} = z_1, u_l = z_2)$ from (u, v) to (z_1, z_2) in M and $Q_2 = (v_1 = z_1, v_2 = z_2, \dots, v_{m-1} = x, v_m = y)$ in M , from (z_1, z_2) to (x, y) in M . Then, according to Observation 2.28, $P = (u_1 = u, u_2 = v, \dots, u_{l-1} = z_1, u_l = z_2, v_3, v_4, \dots, v_{m-1} = x, v_m = y)$ is a triangle-free path from (u, v) to (x, y) in M . Thus, $P_1((u, v), (x, y)) = 1$. \square

We now prove \leftarrow of item 4.

Observation 3.11. *For $((u, v), (x, y)) \in E_O \times E_O$, if $P_1((u, v), (x, y)) = 1$ then $(u, v) \neq (x, y)$ and either of the following occurs:*

1. *There exists a triangle-free path from (u, v) to (x, y) in O .*
2. *For $a \in \{1, 2\}$, $P_{a1}((u, v), (x, y)) = 1$ (i.e., there exists a triangle-free path from (u, v) to (x, y) in M_a).*
3. *There exists $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$ (i.e., there exist triangle-free paths in M from (u, v) to (z_1, z_2) , and from (z_1, z_2) to (x, y)).*

Proof. Suppose for some $((u, v), (x, y)) \in E_O \times E_O$, $P_1((u, v), (x, y)) = 1$, then since (O, P_1, P_2) is the shadow of M on V_O , from Definition 2.36, $(u, v) \neq (x, y)$ and there exists a triangle-free path $P = (u_1 = u, u_2 = v, \dots, u_{l-1} = x, u_l = y)$ from (u, v) to (x, y) in M . If P is completely in O (i.e., all the nodes of P belong to V_O) then P is a triangle-free path in $O = M[V_O]$ (from Definition 2.36, O is an induced subgraph of M). This obeys item 1 of Observation 3.11. If all the nodes of P belongs to V_{H_a} for some $a \in \{1, 2\}$ then from Corollary 2.49, P is a triangle-free path in M_a , i.e., $P_{a1}((u, v), (x, y)) = 1$. This shows if all the nodes of P belongs to V_{H_a} for some $a \in \{1, 2\}$ then item 2 of Observation 3.11 is obeyed.

Suppose P is neither completely in O , nor completely in M_1 , nor completely in M_2 . There are two possibilities either $u_3 \in V_O$ or $u_3 \notin V_O$. We deal with each possibility. If $u_3 \in V_O$ then we have $(v, u_3) \in E_O$

such that (a) $Q_1 = (u_1 = u, u_2 = v, u_3)$, the subpath of P from u_1 to u_3 , is a triangle-free path from (u, v) to (v, u_3) in M , i.e., $P_1((u, v), (v, u_3)) = 1$ (from Definition 2.36, as (O, P_1, P_2) is the shadow of M), and (b) $Q_2 = (u_2 = v, u_3, \dots, u_{l-1} = x, u_l = y)$, the subpath of P from u_2 to u_l , is a triangle-free path from (v, u_3) to (x, y) in M , i.e., $P_1((v, u_3), (x, y)) = 1$ (note that $(v, u_3) \neq (x, y)$, otherwise P is completely in O , contradicting our assumption). This obeys item 3 of Observation 3.11.

Suppose $u_3 \notin V_O$. Since P is a triangle-free path in M , either $u - v \in M$ or $u \rightarrow v \in M$. Since $(u, v) \in E_O$, either $u - v \in O$ or $u \rightarrow v \in O$. Then, from Claim 3.12, either $u, v \in V_{O_1}$ or $u, v \in V_{O_2}$, or both.

Claim 3.12. *If $u - v \in O$ or $u \rightarrow v \in O$ then either $u, v \in V_{O_1}$ or $u, v \in V_{O_2}$.*

Proof. From the construction, $V_O = V_{O_1} \cup V_{O_2}$, and $I = V_{O_1} \cap V_{O_2}$ is a vertex separator of O . W.l.o.g., suppose $u \in V_{O_1}$. If $v \in V_{O_1}$ then we are done. Suppose, $v \notin V_{O_1}$, i.e., $v \in V_O \setminus V_{O_1} = V_{O_2} \setminus I$. Then, $u \in I$, as I is a vertex separator that separates $V_O \setminus V_{O_1}$ and $V_O \setminus V_{O_2}$. But, then, $u, v \in V_{O_2}$, as $I \subseteq V_{O_2}$. \square

W.l.o.g., suppose $u, v \in V_{O_1}$. Then, there must exist a node z in P such that $z \notin V_{M_1}$, otherwise, from Corollary 2.49, P is a triangle-free path in M_1 , contradicting our assumption. Pick the first such z in P , i.e., all the nodes before z in P are in V_{M_1} . Let z_2 be the predecessor of z in P , and z_1 be the predecessor of z_2 in P . Since z is the first node in P which is not in V_{M_1} , $z \in V_{M_2} \setminus V_{M_1}$, and $z_2 \in V_{M_1}$. Since $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} = V_{M_1}$ and $V_{H_2} = V_{M_2}$, therefore, $z_2 \in I$, and $z \in V_{M_2} \setminus I$. This implies that $z, z_1, z_2 \in V_O$, as $V_O \supseteq I \cup N(I, O)$. Since u_3 is not in O , we can say that $z_1 = u_k$ for some $k > 3$. Then, $(z_1, z_2) \in E_O$, and (a) $Q_1 = (u_1 = u, u_2 = v, \dots, u_k = z_1, u_{k+1} = z_2)$, the subpath of P from u to z_2 , is a triangle-free path from (u, v) to (z_1, z_2) in M such that $((u_1, u_2) \neq (z_1, z_2))$ (since $k > 3$), i.e., $P_1((u_1, u_2), (z_1, z_2)) = 1$, and (b) $Q_2 = (u_k = z_1, u_{k+1} = z_2, \dots, u_{l-1} = x, u_l = y)$, the subpath of P from z_1 to y , is a triangle-free path from (z_1, z_2) to (x, y) in M such that $(z_1, z_2) \neq (x, y)$ (if $(z_1, z_2) = (x, y)$ then P is completely in M_1 , contradicting our assumption), i.e., $P_1((z_1, z_2), (x, y)) = 1$. This obeys item 3 of Observation 3.11. This implies that in all the possibilities, Observation 3.11 is obeyed. This completes the proof. \square

This completes the proof of item 4. \square

Proof of item 5. Proof of item 5 is similar to the proof of item 4. For completeness, we are giving the proof. Observations 3.13 to 3.15 prove \leftarrow of item 5.

Observation 3.13. *If $v \neq w$, and there exists a triangle-free path in O from (u, v) to w , then $P_2((u, v), w) = 1$.*

Proof. Suppose $v \neq w$. If there exists a triangle-free path P in O from (u, v) to w , then P is also a triangle-free path in M . This follows from item 1 of Definition 2.36, which states that O is an induced subgraph of M . Consequently, according to item 3 of Definition 2.36, which specifies that $P_2((u, v), w) = 1$ if $v \neq w$ and there exists a triangle-free path from (u, v) to w , we conclude that $P_2((u, v), w) = 1$. \square

Observation 3.14. *For $((u, v), w) \in E_O \times V_O$, if $v \neq w$, and for some $a \in \{1, 2\}$, there exists a triangle-free path P in M_a , from (u, v) to w , i.e., $P_{a2}((u, v), w) = 1$, then P is also a triangle-free path in M , i.e., $P_2((u, v), w) = 1$.*

Proof. Without loss of generality, let us assume $a = 1$. Suppose $((u, v), w) \in E_O \times V_O$, $v \neq w$, and P is a triangle-free path from (u, v) to w in M_1 (i.e., $P_{12}((u, v), w) = 1$). Since $(u, v) \in E_O$, therefore, $(u, v) \in E_M$. This follows from item 1 of Definition 2.36, which states that O is an induced subgraph of M . This further implies that $v \rightarrow u \notin M$. Consequently, from Corollary 2.50, we deduce that P is also a triangle-free path in M . Thus, according to item 3 of Definition 2.36, which states that $P_2((u, v), w) = 1$ if $v \neq w$ and there exists a triangle-free path from (u, v) to w , we conclude that $P_2((u, v), w) = 1$. \square

Observation 3.15. *For $u, v, w, z_1, z_2 \in V_O$, if $v \neq w$, $P_1((u, v), (z_1, z_2)) = 1$, and $P_2((z_1, z_2), w) = 1$, then $P_2((u, v), w) = 1$.*

Proof. Suppose $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$. From items 1 and 2 of Definition 2.36, there exists triangle-free paths $Q_1 = (u_1 = u, u_2 = v, \dots, u_{l-1} = z_1, u_l = z_2)$ from (u, v) to (z_1, z_2) in M (as $P_1((u, v), (z_1, z_2)) = 1$), and $Q_2 = (v_1 = z_1, v_2 = z_2, \dots, v_{m-1}, v_m = w)$ from (z_1, z_2) to w in M (as $P_2((z_1, z_2), w) = 1$). Then, from Observation 2.28, $P = (u_1 = u, u_2 = v, \dots, u_{l-1} = z_1, u_l = z_2, v_3, v_4, \dots, v_{m-1}, v_m = w)$ is a triangle-free path from (u, v) to w in M . Thus, from item 2 of Definition 2.36, $P_2((u, v), w) = 1$. \square

We now prove \leftarrow of item 5.

Observation 3.16. For $((u, v), w) \in E_O \times V_O$, if $P_2((u, v), w) = 1$ then $v \neq w$, and either of the following occurs:

1. There exists a triangle-free path from (u, v) to w in O .
2. For some $a \in \{1, 2\}$, $P_{a2}((u, v), w) = 1$ (i.e., there exists a triangle-free path from (u, v) to w).
3. There exists $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$ (i.e., there exist triangle-free paths in M from (u, v) to (z_1, z_2) , and from (z_1, z_2) to w).

Proof. From item 3 of Definition 2.36, if $P_2((u, v), w) = 1$, then $v \neq w$, and there exists a triangle-free path $P = (u_1 = u, u_2 = v, \dots, u_l = w)$ from (u, v) to w in M .

Similar to the proof of Observation 3.11, either

1. P is a triangle-free path in O , or
2. P is a triangle-free path in M_1 or M_2 , i.e., for some $a \in \{1, 2\}$, $P_{a2}((u, v), w) = 1$, or
3. there exists $(z_1, z_2) \in E_O$ such that $(u, v) \neq (z_1, z_2)$, $w \neq z_2$, and there exist triangle-free paths in M from (u, v) to (z_1, z_2) and from (z_1, z_2) to w , i.e., $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$.

This completes the proof. \square

This completes the proof of \leftarrow of item 5. This further completes the proof of item 5. \square

This completes the proof of Lemma 3.2. \square

3.2 Derived Path Function and Extension of Shadows

We note that P_1 and P_2 in Lemma 3.2 are not independent. Knowing $O, P_{11}, P_{12}, P_{21}$, and P_{22} , we can compute P_1 and P_2 using items 4 and 5 of Lemma 3.2 (this is shown in Lemma 3.18). This motivates us to define (P_1, P_2) as a derived path function of $(O, P_{11}, P_{12}, P_{21}, P_{22})$. Definition 3.17 formally define this.

Definition 3.17 (Derived Path Function). Let H, H_1 and H_2 be undirected graphs such that H_1 and H_2 are induced subgraphs of H , and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let $S_1 \subseteq V_{H_1}$ and $S_2 \subseteq V_{H_2}$ such that $S_1 \cap S_2 = I$. For $a \in \{1, 2\}$, let (O_a, P_{a1}, P_{a2}) be a shadow of H_a on $S_a \cup N(S_a, H_a)$ and $O \in \text{PMEC}(H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)])$ such that O obeys items 1 to 3 of Lemma 3.2. Let $P_1 : E_O \times E_O \rightarrow \{0, 1\}$, and $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ be two functions. We call (P_1, P_2) the derived path function of $(O, P_{11}, P_{12}, P_{21}, P_{22})$, denoted as $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$, if it is computed by following the below steps:

1. Step 1 (Initialization):

- (a) $\forall ((u, v), (x, y)) \in E_O \times E_O$, if $(u, v) \neq (x, y)$ and there exists a triangle-free path from (u, v) to (x, y) in O then $P_1((u, v), (x, y)) = 1$ else $P_1((u, v), (x, y)) = 0$.
- (b) $\forall ((u, v), w) \in E_O \times V_O$, if $v \neq w$ and there exists a triangle-free path from (u, v) to w in O then $P_2((u, v), w) = 1$ else $P_2((u, v), w) = 0$.

2. Step 2 (Update 1):

- (a) $\forall((u, v), (x, y)) \in E_O \times E_O$ such that $(u, v) \neq (x, y)$ and $P_1((u, v), (x, y)) = 0$, if for some $a \in \{1, 2\}$, $P_{a1}((u, v), (x, y)) = 1$ then update P_1 with $P_1((u, v), (x, y)) = 1$.
- (b) $\forall((u, v), w) \in E_O \times V_O$ such that $v \neq w$ and $P_2((u, v), w) = 0$, if for any $a \in \{1, 2\}$, $P_{a2}((u, v), w) = 1$ then update P_2 with $P_2((u, v), w) = 1$.

3. Step 3 (Update 2: add transitivity):

Repeat the following until no further change in (P_1, P_2) :

- (a) $\forall((u, v), (x, y)) \in E_O \times E_O$ such that $(u, v) \neq (x, y)$, and $P_1((u, v), (x, y)) = 0$, if there exists $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$ then update P_1 with $P_1((u, v), (x, y)) = 1$.
- (b) $\forall((u, v), w) \in E_O \times V_O$, such that $v \neq w$ and $P_2((u, v), w) = 0$, if there exists $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$ then update P_2 with $P_2((u, v), w) = 1$.

We emphasize again that P_2 is not in general determined by P_1 ; see Remark 2.37 following Definition 2.36.

Algorithm 2: DPF($O, P_{11}, P_{12}, P_{21}, P_{22}$)

Input : A graph O , and four functions P_{11}, P_{12}, P_{21} and P_{22} such that H, H_1 and H_2 are undirected graphs such that H_1 and H_2 are induced subgraphs of H , and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$, $S_1 \subseteq V_{H_1}$ and $S_2 \subseteq V_{H_2}$ such that $S_1 \cap S_2 = I$, for $a \in \{1, 2\}$, (O_a, P_{a1}, P_{a2}) is a shadow of H_a on $S_a \cup N(S_a, H_a)$ and $O \in \text{PMEC}(H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)])$ such that O obeys items 1 to 3 of Lemma 3.2.

Output : $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$.

```

1  $(P_1, P_2) \leftarrow \text{TFP}(O)$ 
2 foreach  $a \in \{1, 2\}$  do
3   foreach  $((u, v), (x, y)) \in E_O \times E_O$  such that  $(u, v) \neq (x, y)$  do
4     if  $P_{a1}((u, v), (x, y)) = 1$  then
5        $P_1((u, v), (x, y)) = 1$ 
6     end
7   end
8   foreach  $((u, v), w) \in E_O \times V_O$  such that  $v \neq w$  do
9     if  $P_{a2}((u, v), w) = 1$  then
10       $P_2((u, v), w) = 1$ 
11    end
12  end
13 end
14 while  $\exists((u, v), (x, y)) \in E_O \times E_O$  such that  $(u, v) \neq (x, y)$ ,  $P_1((u, v), (x, y)) = 0$ , and
15 there exists  $(z_1, z_2) \in E_O$  such that  $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$  do
16    $P_1((u, v), (x, y)) = 1$ 
17 end
18 while  $\exists((u, v), w) \in E_O \times V_O$  such that  $v \neq w$ ,  $P_2((u, v), w) = 0$ , and
19 there exists  $(z_1, z_2) \in E_O$  such that  $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$  do
20    $P_2((u, v), w) = 1$ 
21 end
22 return  $(P_1, P_2)$ 

```

We construct Alg. 2 to compute $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$. Line 1 performs the initialization part (Step 1 of Definition 3.17). Lines 2-13 do the first update of P_1 and P_2 (Step 2 of Definition 3.17). Lines 14-21 do the final update of P_1 and P_2 (Step 3 of Definition 3.17).

The following lemma shows that if (O, P_1, P_2) is the shadow of an MEC, and (O_1, P_{11}, P_{12}) and (O_2, P_{21}, P_{22}) are the shadows of two projections of the MEC then knowing the value of $O, P_{11}, P_{12}, P_{21}$ and P_{22} , we can compute (P_1, P_2) using alg. 2, as $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$.

Lemma 3.18. *Let H be an undirected graph, and H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1, S_2 be the subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let $Y = S_1 \cup S_2 \cup N(S_1 \cup S_2, H)$, $Y_1 = S_1 \cup N(S_1, H_1)$, and $Y_2 = S_2 \cup N(S_2, H_2)$. Let M, M_1, M_2 be the MECs of H, H_1, H_2 , respectively, (O, P_1, P_2) be the shadow of M on Y , (O_1, P_{11}, P_{12}) be the shadow of M_1 on Y_1 , and (O_2, P_{21}, P_{22}) be the shadow of M_2 on Y_2 . If $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$ then $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$, i.e., (P_1, P_2) is the derived path function of $(O, P_{11}, P_{12}, P_{21}, P_{22})$.*

Proof. Let $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22}) = (P'_1, P'_2)$. We show that $(P_1, P_2) = (P'_1, P'_2)$. We first show that $P_1 = P'_1$. Since (O, P_1, P_2) is a shadow of M , therefore, from Definition 2.41, the domain of P_1 is $E_O \times E_O$ and the co-domain of P_1 is $\{0, 1\}$. Also, from line 1 of alg. 2, since $(P'_1, P'_2) = \text{TFP}(O)$, therefore, from alg. 1, the domain of P'_1 is $E_O \times E_O$ and the co-domain of P'_1 is $\{0, 1\}$. This implies both P_1 and P'_1 have the same domain and co-domain. Thus, to prove $P_1 = P'_1$, it is sufficient to show that (a) $\forall((u, v), (x, y)) \in E_O \times E_O$, if $P'_1((u, v), (x, y)) = 1$ then $P_1((u, v), (x, y)) = 1$, and (b) $\forall((u, v), (x, y)) \in E_O \times E_O$, if $P_1((u, v), (x, y)) = 1$ then $P'_1((u, v), (x, y)) = 1$.

Claim 3.19. *For all $((u, v), (x, y)) \in E_O \times E_O$, if $P'_1((u, v), (x, y)) = 1$, then $P_1((u, v), (x, y)) = 1$.*

Proof. From Definition 3.17, P'_1 reached its final version after following steps 1, 2, and 3 of Definition 3.17.

We first show that at the end of step 1, for any $((u, v), (x, y)) \in E_O$, if $P'_1((u, v), (x, y)) = 1$, then $P_1((u, v), (x, y)) = 1$. From line 1 of alg. 2, $(P'_1, P'_2) = \text{TFP}(O)$. Therefore, from Lemma 2.35, if $P'_1((u, v), (x, y)) = 1$, then there exists a triangle-free path from (u, v) to (x, y) in O . From item 1 of Definition 2.36, O is an induced subgraph of M . This implies there exists a triangle-free path from (u, v) to (x, y) in M . Then, from item 2 of Definition 2.36, $P_1((u, v), (x, y)) = 1$. This implies that at the end of the initialization step (i.e., step 1) of Definition 3.17, if $P'_1((u, v), (x, y)) = 1$, then $P_1((u, v), (x, y)) = 1$.

We now show that at step 2 (update 1) of Definition 3.17, for any $((u, v), (x, y)) \in E_O \times E_O$, if P'_1 is updated with $P'_1((u, v), (x, y)) = 1$, then $P_1((u, v), (x, y)) = 1$. Suppose for some $((u, v), (x, y)) \in E_O \times E_O$, P'_1 is updated with $P'_1((u, v), (x, y)) = 1$ at step 2 of Definition 3.17. From Definition 3.17, this implies that $(u, v) \neq (x, y)$ and for some $a \in \{1, 2\}$, $P_{a1}((u, v), (x, y)) = 1$. Since P_1 obeys item 4 of Lemma 3.2, from item 4b of item 4 of Lemma 3.2, $P_1((u, v), (x, y)) = 1$. This implies that if for some $((u, v), (x, y)) \in E_O \times E_O$, P'_1 is updated with $P'_1((u, v), (x, y)) = 1$ at step 2 of Definition 3.17, then $P_1((u, v), (x, y)) = 1$. This further implies that at the start of step 3, for any $((u, v), (x, y)) \in E_O \times E_O$, if $P'_1((u, v), (x, y)) = 1$, then $P_1((u, v), (x, y)) = 1$.

Suppose while running step 3, if at some iteration, for some $((u, v), (x, y)) \in E_O \times E_O$, we update P'_1 with $P'_1((u, v), (x, y)) = 1$. Then $P_1((u, v), (x, y)) = 1$ provided that before that iteration, for all $((u', v'), (x', y')) \in E_O \times E_O$, if $P'_1((u', v'), (x', y')) = 1$, then $P_1((u', v'), (x', y')) = 1$. From Definition 3.17, if at step 3 of Definition 3.17, for any $((u, v), (x, y)) \in E_O \times E_O$, P'_1 is updated with $P'_1((u, v), (x, y)) = 1$, then $(u, v) \neq (x, y)$ and there exists a $(z_1, z_2) \in E_O$ such that $P'_1((u, v), (z_1, z_2)) = P'_1((z_1, z_2), (x, y)) = 1$. Since at the start of the iteration, $P'_1((u, v), (z_1, z_2)) = P'_1((z_1, z_2), (x, y)) = 1$, therefore, from our assumption, $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$. Since P_1 follows item 4 of Lemma 3.2, from item 4c of item 4 of Lemma 3.2, $P_1((u, v), (x, y)) = 1$. This proves that even at the end of step 3, for all $((u, v), (x, y)) \in E_O \times E_O$, if $P'_1((u, v), (x, y)) = 1$, then $P_1((u, v), (x, y)) = 1$. This proves Claim 3.19. \square

Claim 3.20. *For all $((u, v), (x, y)) \in E_O \times E_O$, if $P_1((u, v), (x, y)) = 1$, then $P'_1((u, v), (x, y)) = 1$.*

Proof. Let us construct a directed acyclic graph (DAG) D such that $V_D = E_O$, and for $(u, v), (x, y) \in E_O$, $(u, v) \rightarrow (x, y) \in E_D$ if, and only if, $P_1((u, v), (x, y)) = 1$. We first show that D is a DAG.

Since P_1 obeys item 4 of Lemma 3.2, if $P_1((u, v), (x, y)) = 1$, then $(u, v) \neq (x, y)$. This implies that D does not contain any self-loop. Suppose D is not a DAG. Then, pick the smallest length cycle $((u_1, v_1), (u_2, v_2), \dots, (u_n, v_n), (u_1, v_1))$ in D for some $n \geq 2$. This implies $P_1((u_1, v_1), (u_2, v_2)) =$

$P_1((u_2, v_2), (u_3, v_3)) = \dots = P_1((u_{n-1}, v_{n-1}), (u_n, v_n)) = P_1((u_n, v_n), (u_1, v_1)) = 1$. Since P_1 obeys item 4 of Lemma 3.2, therefore, from item 4c of item 4 of Lemma 3.2, $P_1((u_1, v_1), (u_2, v_2)) = P_1((u_2, v_2), (u_3, v_3)) = \dots = P_1((u_{n-1}, v_{n-1}), (u_n, v_n)) = 1$ implies $P_1((u_1, v_1), (u_n, v_n)) = 1$. This further implies that in M , there exist triangle-free paths from (u_1, v_1) to (u_n, v_n) , and from (u_n, v_n) to (u_1, v_1) . But, this contradicts Lemma 2.53. This implies that D does not have any cycles.

We now show that if $(u, v) \rightarrow (x, y) \in E_D$, then $P'_1((u, v), (x, y)) = 1$, which completes the proof of Claim 3.20 (remember, from the construction of D , $(u, v) \rightarrow (x, y) \in E_D$ if $P_1((u, v), (x, y)) = 1$). Let τ be an ordering on the vertices of D such that if $(u, v) \rightarrow (x, y) \in E_D$, then $\tau((u, v)) < \tau((x, y))$ (the existence of τ is because D is a DAG).

Suppose there exists $(u, v) \rightarrow (x, y) \in E_D$ such that $P'_1((u, v), (x, y)) = 0$. Pick such a $(u, v) \rightarrow (x, y) \in E_D$ such that for all $(u', v') \rightarrow (x', y') \in E_D$,

$$\text{if } \tau(x', y') < \tau(x, y), \text{ or } (x', y') = (x, y) \text{ and } \tau(u, v) < \tau(u', v'), \text{ then } P'_1((u', v'), (x', y')) = 1. \quad (2)$$

From the construction of D , since $(u, v) \rightarrow (x, y) \in E_D$, therefore, $P_1((u, v), (x, y)) = 1$. Since P_1 obeys item 4 of Lemma 3.2, therefore, either (a) there exists a triangle-free path from (u, v) to (x, y) in O , or (b) for some $a \in \{1, 2\}$, $P_{a1}((u, v), (x, y)) = 1$, or (c) there exists a $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$. If $P_1((u, v), (x, y)) = 1$ due to (a) or (b), then from steps 1 and 2 of Definition 3.17, $P'_1((u, v), (x, y)) = 1$. If $P_1((u, v), (x, y)) = 1$ due to (c), then there exists a $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$. This implies $\tau((u, v)) < \tau((z_1, z_2)) < \tau((x, y))$. Then from eq. (2), $P'_1((u, v), (z_1, z_2)) = 1$ (because $\tau((z_1, z_2)) < \tau((x, y))$), and $P'_1((z_1, z_2), (x, y)) = 1$ (because $\tau((u, v)) < \tau((z_1, z_2))$). But, then $P'_1((u, v), (x, y)) = 1$, updated at step 3 of Definition 3.17. This implies that for all $(u, v), (x, y) \in E_O$, if $P_1((u, v), (x, y)) = 1$, then $P'_1((u, v), (x, y)) = 1$. This completes the proof of Claim 3.20. \square

Claims 3.19 and 3.20 proves $P_1 = P'_1$.

We now prove $P_2 = P'_2$. Since (O, P_1, P_2) is a shadow of M , therefore, according to Definition 2.41, the domain of P_2 is $E_O \times V_O$, and the co-domain of P_1 is $\{0, 1\}$. Also, from line 1 of alg. 2, since $(P'_1, P'_2) = \text{TFP}(O)$, therefore, based on alg. 1, the domain of P'_2 is $E_O \times V_O$, and the co-domain of P'_1 is $\{0, 1\}$. This implies that both P_2 and P'_2 have the same domain and co-domain. Thus, to prove $P_1 = P'_1$, it is sufficient to show that (a) for all $((u, v), w) \in E_O \times V_O$, if $P_2((u, v), w) = 1$, then $P'_2((u, v), w) = 1$, and (b) for all $((u, v), w) \in E_O \times V_O$, if $P'_2((u, v), w) = 1$, then $P_2((u, v), w) = 1$. The proof of (a) is similar to Claim 3.19, and the proof of (b) is similar to Claim 3.20. For completeness, we provide the proof.

Claim 3.21. $\forall ((u, v), w) \in E_O \times V_O$, if $P'_2((u, v), w) = 1$ then $P_2((u, v), w) = 1$.

Proof. From Definition 3.17, P'_2 reached its final version after following steps 1, 2, and 3 of Definition 3.17.

Firstly, we demonstrate that at the end of step 1, for any $((u, v), w) \in E_O \times V_O$, if $P'_2((u, v), w) = 1$, then $P_2((u, v), w) = 1$. From line 1 of alg. 2, we have $(P'_1, P'_2) = \text{TFP}(O)$. Therefore, from Lemma 2.35, if $P'_2((u, v), w) = 1$, then $v \neq w$, and there exists a triangle-free path from (u, v) to w in O . Since P_2 satisfies item 5 in Lemma 3.2, from item 5a, we have $P_2((u, v), w) = 1$. This implies that at the end of the initialization step (i.e., step 1) of Definition 3.17, if $P'_2((u, v), w) = 1$, then $P_2((u, v), w) = 1$.

Next, we demonstrate that if at step item 2 (update 1) of Definition 3.17, for any $((u, v), w) \in E_O \times V_O$, P'_2 is updated with $P'_2((u, v), w) = 1$, then $P_2((u, v), w) = 1$.

Suppose for some $((u, v), w) \in E_O \times V_O$, P'_2 is updated with $P'_2((u, v), w) = 1$ at step 2 of Definition 3.17. From Definition 3.17, this implies that $v \neq w$ and for some $a \in \{1, 2\}$, $P_{a2}((u, v), w) = 1$. Since P_2 satisfies Lemma 3.2, from item 5b of item 5 of Lemma 3.2, we have $P_2((u, v), w) = 1$. This implies that if for some $((u, v), w) \in E_O \times V_O$, P'_2 is updated with $P'_2((u, v), w) = 1$ at step 2, then $P_2((u, v), w) = 1$. This further implies that at the start of step 3, for any $((u, v), w) \in E_O \times V_O$, if $P'_2((u, v), w) = 1$, then $P_2((u, v), w) = 1$.

Suppose while running step 3, if at some iteration, for some $((u, v), w) \in E_O \times V_O$, we update P'_2 with $P'_2((u, v), w) = 1$. Then $P_2((u, v), w) = 1$ provided that before that iteration, for all $((u', v'), w') \in E_O \times V_O$, if $P'_2((u', v'), w') = 1$, then $P_2((u', v'), w') = 1$. From Definition 3.17, if at step 3 of Definition 3.17, for any $((u, v), w) \in E_O \times V_O$, P'_2 is updated with $P'_2((u, v), w) = 1$, then $v \neq w$, and there exists a $(z_1, z_2) \in E_O$

such that $P'_1((u, v), (z_1, z_2)) = P'_2((z_1, z_2), w) = 1$. Since we have already shown that $P_1 = P'_1$, therefore, $P_1((u, v), (z_1, z_2)) = 1$. Also, since at the start of the iteration, we have $P'_2((z_1, z_2), w) = 1$, therefore, from our assumption, $P_2((z_1, z_2), w) = 1$. Since P_2 follows Lemma 3.2, from item 5c of item 5 of Lemma 3.2, we have $P_2((u, v), w) = 1$. This proves that even at the end of step 3, for all $((u, v), w) \in E_O \times V_O$, if $P'_2((u, v), w) = 1$ then $P_2((u, v), w) = 1$. This completes the proof of Claim 3.21. \square

Claim 3.22. $\forall((u, v), w) \in E_O \times V_O$, if $P_2((u, v), w) = 1$ then $P'_2((u, v), w) = 1$.

Proof. We show that for all $((u, v), w) \in E_O \times V_O$, if $P_2((u, v), w) = 1$ then $P'_2((u, v), w) = 1$. Let us construct a directed acyclic graph (DAG) D such that $V_D = E_O$, and for $(u, v), (x, y) \in E_O$, $(u, v) \rightarrow (x, y) \in E_D$ if, and only if, $P_1((u, v), (x, y)) = 1$. We have shown in the proof of Claim 3.20 that D is a DAG (read the first two paragraphs of Claim 3.20 for the proof that D is a DAG). Let τ be an ordering on the vertices of D such that if $(u, v) \rightarrow (x, y) \in E_D$ then $\tau((u, v)) < \tau((x, y))$ (existence of τ is because D is a DAG). If possible, suppose there exists a $((u, v), w) \in E_O \times V_O$ such that $P_2((u, v), w) = 1$ and $P'_2((u, v), w) = 0$. For w , pick such a (u, v) which is highest in τ , i.e.,

$$\forall(u', v') \in E_O \text{ such that } \tau((u, v)) < \tau((u', v')), \text{ if } P_2((u', v'), w) = 1 \text{ then } P'_2((u', v'), w) = 1. \quad (3)$$

Since P_2 obeys item 5 of Lemma 3.2, therefore, either (a) there exists a triangle-free path from (u, v) to w in O , or (b) for any $a \in \{1, 2\}$, $P_{a2}((u, v), w) = 1$, and $v \rightarrow u \notin O$, or (c) there exists a $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$. If $P_2((u, v), w) = 1$ due to (a) or (b) then from items 1 and 2 of Definition 3.17, $P'_2((u, v), w) = 1$. And, if $P_2((u, v), w) = 1$ due to (c) then there exists a $(z_1, z_2) \in E_O$ such that $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$. Since, we have shown that $P_1 = P'_1$, therefore, $P'_1((u, v), (z_1, z_2)) = P_1((u, v), (z_1, z_2)) = 1$. Also, since $P_1((u, v), (z_1, z_2)) = 1$, therefore, $\tau((u, v)) < \tau((z_1, z_2))$. Then from eq. (3), $P'_2((z_1, z_2), w) = 1$. But, then $P'_2((u, v), w) = 1$, updated at step 3 of Definition 3.17. This implies that for all $((u, v), w) \in E_O \times V_O$, if $P_2((u, v), w) = 1$ then $P'_2((u, v), w) = 1$. This completes the proof of Claim 3.22. \square

Claims 3.21 and 3.22 proves $P_2 = P'_2$. This implies $(P_1, P_2) = (P'_1, P'_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$. This completes the proof of Lemma 3.18. \square

We define the following, to summarize the necessary conditions provided by Lemma 3.2.

Definition 3.23 (Valid derived path function). (P_1, P_2) is said to be a “valid derived path function” of $(O, P_{11}, P_{12}, P_{21}, P_{22})$ if:

1. $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$, and
2. For all $(u, v), (x, y) \in E_O$, if $P_1((u, v), (x, y)) = 1$ then $P_1((x, y), (u, v)) = 0$.

Definition 3.24 (Extension). Let H be an undirected graph, and H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1 and S_2 be the subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let $A = H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)]$, $B_1 = H_1[S_1 \cup N(S_1, H_1)]$, and $B_2 = H_2[S_2 \cup N(S_2, H_2)]$. Let $O_1 \in \text{PMEC}(B_1)$ and $O_2 \in \text{PMEC}(B_2)$ be partial MECs. For each $a \in \{1, 2\}$, let $P_{a1} : E_{O_a} \times E_{O_a} \rightarrow \{0, 1\}$, and $P_{a2} : E_{O_a} \times V_{O_a} \rightarrow \{0, 1\}$ be two functions. We define $O \in \text{PMEC}(A)$ to be an *extension* of (O_1, P_{11}, P_{12}) and (O_2, P_{21}, P_{22}) , denoted as $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, if

1. For $a \in \{1, 2\}$, for $u, v \in S_a \cup N(S_a, H_a)$, if $u \rightarrow v \in O_a$ then $u \rightarrow v \in O$.
2. For $a \in \{1, 2\}$, $\mathcal{V}(O_a) = \mathcal{V}(O, S_a \cup N(S_a, H_a))$.
3. For $a \in \{1, 2\}$, for $u - v \in O_a$, $u \rightarrow v \in O$ if, and only if, at least one of the following occurs:
 - (a) $u \rightarrow v$ is strongly protected in O .
 - (b) $\exists x - y \in O_a$ such that $x \rightarrow y \in O$, and $P_{a1}((x, y), (u, v)) = 1$.
 - (c) $\exists x - y \in O_a$ such that $x \rightarrow y \in O$, $P_{a2}(x, y, v) = 1$, and $P_{a2}(v, u, x) = 1$.

4. $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$ is valid (see Definition 3.23 for valid derived path function).

With a slight abuse of notation, we say (O, P_1, P_2) as an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, if

1. O obeys items 1 to 4 of Definition 3.24, and
2. $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$.

Since P_1 in Lemma 3.2 obeys item 2 of Definition 3.23 (from Lemma 2.53), therefore, Lemma 3.18 and Definitions 3.17, 3.23 and 3.24 summarises Lemma 3.2 as follows:

Lemma 3.25. *Let H be an undirected graph, and H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1, S_2 be the subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let $Y = S_1 \cup S_2 \cup N(S_1 \cup S_2, H)$, $Y_1 = S_1 \cup N(S_1, H_1)$, and $Y_2 = S_2 \cup N(S_2, H_2)$. Let M, M_1, M_2 be MECs of H, H_1, H_2 , respectively. Let (O, P_1, P_2) be the shadow of M on Y , (O_1, P_{11}, P_{12}) be the shadow of M_1 on Y_1 , and (O_2, P_{21}, P_{22}) be the shadow of M_2 on Y_2 . If $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$ then*

1. $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, and
2. $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, O_2, P_{21}, P_{22})$.

We show in Section 3.4 that the necessary conditions in Lemma 3.25 are also sufficient. We then use this result along with Lemma 3.25 to construct an algorithm that counts MECs of H . To prove this, we need a modified version of the LBFS algorithm of [Rose, Tarjan, and Lueker \(1976\)](#). In the next subsection, we introduce a modified version of the LBFS algorithm. We will also see some properties of the new LBFS algorithm. All these results will be used in Section 3.4 to show that the necessary conditions provided in Lemma 3.25 are sufficient.

3.3 Lexicographical breadth-first search (LBFS) algorithm

In this section, we present a modified version of the LBFS algorithm. Let H be an undirected graph, and let H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H . For each $a \in \{1, 2\}$, let S_a be a subset of V_{H_a} such that $I \subseteq S_a$, and let M_a be an MEC of H_a . The shadow of M_a on $S_a \cup N(S_a, H_a)$ is denoted as (O_a, P_{a1}, P_{a2}) . Consider $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$ (i.e., O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$; refer to Definition 3.24 for the definition of extension).

We introduce a modified LBFS algorithm, Alg. 3. This algorithm takes as input (a) O and (b) \mathcal{C} , an undirected connected component of M_a for $a \in \{1, 2\}$. Alg. 3 returns an LBFS ordering of \mathcal{C} that obeys O , i.e., if τ is the returned LBFS ordering of \mathcal{C} , then for any $u, v \in V_{\mathcal{C}}$ such that if $u \rightarrow v \in O$ then $\tau(u) < \tau(v)$.

Alg. 3 is a variation of the LBFS algorithm introduced by [Rose, Tarjan, and Lueker \(1976\)](#). Instead of selecting a vertex arbitrarily, Alg. 3 chooses a canonical source node (Definition 2.32) at line 5 of Alg. 3. The crucial point to verify is the existence of a canonical source node in $U_M(\mathcal{C}, O)[X]$, where $U_M(\mathcal{C}, O)$ represents the Markov union of \mathcal{C} and O (see Definition 2.20 for the definition of the Markov union), and $X \subseteq V_{\mathcal{C}}$.

Claim 3.26. *There always exists a canonical source vertex in $U_M(\mathcal{C}, O)[X]$, where $X \subseteq V_{\mathcal{C}}$.*

Proof. Suppose $U_M(\mathcal{C}, O)[X]$ does not contain any canonical source vertices. Definition 2.33 implies that if a node u is not a canonical source vertex in $U_M(\mathcal{C}, O)[X]$ then there exists a node v that certifies that u is not a canonical source vertex. And the certification is an edge $v \rightarrow v' \in U_M(\mathcal{C}, O)[X]$ and a triangle-free path from (v, v') to u in $U_M(\mathcal{C}, O)[X]$. If $U_M(\mathcal{C}, O)[X]$ does not have contain any canonical source nodes, then, since X has a finite number of nodes, there exists a sequence $(u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ such that u_1, u_2, \dots, u_l are distinct, and for each $1 \leq i \leq l$, u_i is a certificate of u_{i+1} . Now, we will prove the following claim, which is an important component in demonstrating that we will not obtain any such sequence. This, in turn, implies that there always exists a canonical source node in $U_M(\mathcal{C}, O)[X]$.

Claim 3.27. *In $U_M(\mathcal{C}, O)[X]$, suppose u, v , and w are three distinct vertices such that u is a certificate of v , and v is a certificate of w . Then, either $u \rightarrow v \in O$, or u is a certificate of w .*

Algorithm 3: LBFS(\mathcal{C}, O): A modified version of the Lexicographic BFS (Rose, Tarjan, and Lueker (1976)). If the algorithm is executed with O as a null graph the algorithm performs a normal LBFS.

Input : (a) An undirected connected chordal graph \mathcal{C} , and (b) a chain graph O such that there exist three undirected graphs H, H_1 and H_2 such that

- (i) $V_{H_1} \cap V_{H_2} = I$ is a vertex separator of H ,
- (ii) there exist $S_1 \in V_{H_1}$ and $S_2 \in V_{H_2}$ such that $I = V_{S_1} \cap V_{S_2}$,
- (iii) M_1 and M_2 are MECs of H_1 and H_2 , respectively,
- (iv) for each $a \in \{1, 2\}$, (O_a, P_{a1}, P_{a2}) is the shadow of M_a on $S_a \cup N(S_a, H_a)$,
- (v) $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, and
- (vi) \mathcal{C} is an undirected connected component of M_1 or M_2 .

Output : an LBFS ordering τ of \mathcal{C} that is consistent with O

- 1 $\mathcal{S} \leftarrow$ sequence of sets initialized with $(V_{\mathcal{C}})$
- 2 $\tau \leftarrow$ empty list
- 3 **while** \mathcal{S} is non-empty **do**
- 4 $X \leftarrow$ first non-empty set of \mathcal{S}
- 5 $v \leftarrow$ arbitrary canonical source node of $U_M(\mathcal{C}, O)[X]$
- 6 Add vertex v to the end of τ .
- 7 Replace the set X in the sequence \mathcal{S} by the set $X \setminus \{v\}$.
- 8 $N(v) \leftarrow \{x \mid x \notin \tau \text{ and } v - x \in E_{\mathcal{C}}\}$
- 9 Denote the current \mathcal{S} by (S_1, \dots, S_k)
- 10 Replace each S_i by $S_i \cap N(v, \mathcal{C}), S_i \setminus N(v, \mathcal{C})$.
- 11 Remove all empty sets from \mathcal{S} .
- 12 **end**
- 13 **return** τ

Proof. Suppose in $U_M(\mathcal{C}, O)[X]$, u is a certificate of v , and v is a certificate of w . Then, from Definition 2.33, since u is a certificate of v , there must exist a node $u' \in X$ such that $u \rightarrow u' \in U_M(\mathcal{C}, O)[X]$, and there exists a triangle-free path $Q_1 = (x_1 = u, x_2 = u', \dots, x_l = v)$ from (u, u') to v in $U_M(\mathcal{C}, O)[X]$. Similarly, since v is a certificate of w , there exists a node $v' \in X$ such that $v \rightarrow v' \in U_M(\mathcal{C}, O)[X]$, and there exists a triangle-free path $Q_2 = (y_1 = v, y_2 = v', \dots, y_m = w)$ from (v, v') to w in $U_M(\mathcal{C}, O)[X]$.

We first show that for any triangle-free path P in $U_M(\mathcal{C}, O)[X]$, P is a triangle-free path in \mathcal{C} . From Definition 2.20, if $u \rightarrow v \in U_M(\mathcal{C}, O)$, then either $u \rightarrow v \in \mathcal{C}$ or $u \rightarrow v \in O$. Since \mathcal{C} is an undirected graph (recall that \mathcal{C} is an undirected connected component of M_a), therefore, $u \rightarrow v \in O$. From the construction, for $Y = V_{\mathcal{C}} \cap V_O$, $\text{skeleton}(\mathcal{C}[Y]) = \text{skeleton}(O[Y])$. Since $X \subseteq V_{\mathcal{C}}$ ($X \in \mathcal{S}$, and in Alg. 3, the elements of \mathcal{S} are the subsets of $V_{\mathcal{C}}$), therefore, if $u \rightarrow v \in U_M(\mathcal{C}, O)[X]$, then $u \rightarrow v \in O$ and $u - v \in \mathcal{C}$. This further implies that each triangle-free path in $U_M(\mathcal{C}, O)[X]$ is a triangle-free path in \mathcal{C} .

The above discussion implies that Q_1 and Q_2 are triangle-free paths in \mathcal{C} and $u \rightarrow u', v \rightarrow v' \in O$.

If $l = 2$, i.e., the length of Q_1 is one, then $u' = v$, and $u \rightarrow v \in O$, and we are done.

Suppose $l > 2$. We show that there exists a triangle-free path from (u, u') to w in $U_M(\mathcal{C}, O)[X]$. As shown above, Q_1 and Q_2 are the triangle-free paths in \mathcal{C} . Since \mathcal{C} is an undirected connected component of an MEC M_a , therefore, from item 2 of Theorem 2.18, \mathcal{C} is chordal. Then from Observation 2.27, Q_1 and Q_2 are chordless paths of \mathcal{C} . Pick the least $i < l$ such that for some $1 \leq j \leq m$, $x_i - y_j \in \mathcal{C}$. After this, pick the highest j such that $x_i - y_j \in \mathcal{C}$. The existence of such an edge is due to $x_{l-1} - y_1 \in \mathcal{C}$ (in Q_1 , $x_{l-1} - y_1$ is an edge as $x_l = y_1$).

If $i \neq 1$, then $Q = (x_1 = u, x_2 = u', \dots, x_i, y_j, \dots, y_m = w)$ is a triangle-free path from (u, u') to w in \mathcal{C} . We now show that Q is also a triangle-free path in $U_M(\mathcal{C}, O)[X]$.

Since Q_1 and Q_2 are triangle-free paths in $U_M(\mathcal{C}, O)[X]$, to show that Q is a triangle-free path, the only thing we have to show is $y_j \rightarrow x_i \notin O$. If both x_i and y_j are not nodes of V_O , then there cannot be $y_j \rightarrow x_i \in O$. Suppose $y_j, x_i \in V_O$. Since Q is a triangle-free path in \mathcal{C} , the subpath $Q' = (x_1 = u, x_2 =$

u', \dots, x_i, y_j) of Q is also a triangle-free path in \mathcal{C} . Since \mathcal{C} is an undirected connected component of M_a , a triangle-free path in \mathcal{C} is a triangle-free path in M_a . This implies $P_{a1}((u, u'), (x_i, y_j)) = 1$. Since O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, from item 3b of item 3 of Definition 3.24, $x_i \rightarrow y_j \in O$ (because $P_{a1}((u, u'), (x_i, y_j)) = 1$ and $u \rightarrow u' \in O$). This shows we get the required triangle-free path in $U_M(\mathcal{C}, O)[X]$, Q , from (u, u') to w .

Thus, the only possibility that remains is $i = 1$. We show that this case cannot occur. Suppose, if possible, that $i = 1$. Recall that $l > 2$. Pick the least j such that $x_1 - y_j \in \mathcal{C}$. Since Q_1 is a chordless path in \mathcal{C} , and $l > 2$, j must be greater than 1, since $y_1 = x_l$. Then $Q = (y_1 = v, y_2 = v', \dots, y_j, x_1 = u)$ is a chordless path in \mathcal{C} , from (v, v') to u . As \mathcal{C} is an undirected connected component of M_a , and Q_1 and Q are triangle-free paths in \mathcal{C} from (u, u') to v and from (v, v') to u , respectively, therefore, $P_{a2}((u, u'), v) = P_{a2}((v, v'), u) = 1$. Since O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, therefore, from item 3c of item 3, $v' \rightarrow v \in O$ (as $u \rightarrow u' \in O$ and $P_{a2}((u, u'), v) = P_{a2}((v, v'), u) = 1$). But, this is a contradiction, as from our assumption, $v \rightarrow v' \in O$. This shows that $i = 1$ does not occur. This completes the proof of Claim 3.27. \square

We now prove the following claim to complete the proof of Claim 3.26.

Claim 3.28. *There cannot exist a sequence $(u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ such that for each $1 \leq i \leq l$, u_i is a certificate of u_{i+1} .*

We will prove Claim 3.28 using induction on l . From Observation 2.34, we know that self-certification is not possible, implying that $l \geq 2$.

Base Case: $l = 2$. Assume, for the sake of contradiction, that Claim 3.28 is false for $l = 2$. In other words, there exist nodes u_1 and u_2 such that u_1 is a certificate of u_2 , and u_2 is a certificate of u_1 . This implies the existence of edges $u_1 - v_1$ and $u_2 - v_2$ in \mathcal{C} such that $u_1 \rightarrow v_1$ and $u_2 \rightarrow v_2$ in O . Additionally, there exist triangle-free paths Q_1 from (u_1, v_1) to u_2 and Q_2 from (u_2, v_2) to u_1 in \mathcal{C} . Since \mathcal{C} is an undirected connected component of the MEC M_a (for some $a \in \{1, 2\}$), \mathcal{C} is an undirected chordal graph (cf. item 2 of Theorem 2.18). Thus, Q_1 and Q_2 are triangle-free paths M_a .

This implies that $v_1 \neq u_2$ and $v_2 \neq u_1$. If, for instance, $v_1 = u_2$, then by applying Corollary 2.26 to the undirected chordal graph \mathcal{C} , the adjacency of $v_1 = u_2$ and u_1 and the triangle-free path Q_2 from (u_2, v_2) to u_1 would imply $v_2 = u_1$ as well. Similarly, if $v_2 = u_1$, it would imply $v_1 = u_2$. Thus, we must have both $v_1 = u_2$ and $v_2 = u_1$, which leads to the existence of both $u_1 \rightarrow u_2$ and $u_2 \rightarrow u_1$ in O , contradicting our initial assumption. Therefore, it follows that $v_1 \neq u_2$ and $v_2 \neq u_1$.

Given that O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, we can apply item 3c of item 3 of Definition 3.24. This reveals that $u_1 \rightarrow v_1 \in O$ together with the triangle-free paths Q_1 from (u_1, v_1) to u_2 (with $v_1 \neq u_2$) and Q_2 from (u_2, v_2) to u_1 (with $v_2 \neq u_1$) imply $v_2 \rightarrow u_2 \in O$. This, however, contradicts the assumption that $u_2 \rightarrow v_2 \in O$. Thus, we have shown that Claim 3.28 holds for $l = 2$.

Induction: $l \geq 3$. Assume that Claim 3.28 is true for $l = k - 1$, where $k \geq 3$. We will now use Claim 3.27 to prove Claim 3.28 for $l = k$.

Assume, for the sake of contradiction, that Claim 3.28 is not true for a sequence $(u_1, u_2, \dots, u_k, u_{k+1} = u_1)$. According to Claim 3.27, for each $1 \leq i \leq l - 1$, either u_i is a certificate for u_{i+2} or $u_i \rightarrow u_{i+1} \in O$, and for $i = l$, either u_l is a certificate for u_2 or $u_l \rightarrow u_1 \in O$. If, for any $1 \leq i \leq k - 1$, u_i is a certificate for u_{i+2} , it would create a sequence $(u_1, u_2, \dots, u_i, u_{i+2}, \dots, u_k, u_{k+1} = u_1)$ of $k - 1$ distinct nodes that violates our induction hypothesis. Similarly, if u_k is a certificate for u_2 , it would create another sequence $(u_2, u_3, \dots, u_k, u_2)$ of $k - 1$ distinct nodes that violates our induction hypothesis. Thus, the only possibility is that for each $1 \leq i \leq k$, $u_i \rightarrow u_{i+1} \in O$. However, this leads to the presence of a directed cycle in O , which is a contradiction because O is a chain graph (as per item 1 of Definition 2.21). Therefore, we have proven Claim 3.28 for all l by induction.

This completes the proof of Claim 3.26. \square

The following observation and its corollaries show that the output ordering is consistent with O .

Observation 3.29. Let \mathcal{C} be an undirected connected component of M_a for $a \in \{1, 2\}$. Suppose Alg. 3 is called for \mathcal{C} and O . And, at the beginning of an iteration of the while loop (lines 3-12 of Alg. 3), let $\tau = (a_1, a_2, \dots, a_r)$ and $\mathcal{S} = (S_1, S_2, \dots, S_l)$. Then, for any triangle-free path $P = (u_1, u_2, \dots, u_m)$ of \mathcal{C} such that $u_1 \rightarrow u_2 \in O$, the following statements hold for all $1 \leq i < j \leq m$,

1. If $u_j \in \tau$, then $u_i \in \tau$ and u_i arrived in τ before u_j .
2. If $u_j \in S_x$ for some $1 \leq x \leq l$, then either $u_i \in \tau$, or $u_i \in S_y$ for $y \leq x$.
3. If $u_1 \in \tau$, then at most one vertex of P is in S_1 .

Before proving Observation 3.29, we go through the following claim, which is a necessary ingredient for the proof of Observation 3.29.

Claim 3.30. If $P = (u_1, u_2, \dots, u_l)$ is a triangle-free path in \mathcal{C} and $u_1 \rightarrow u_2 \in O$ then P is a triangle-free path in $U_M(\mathcal{C}, O)$.

Proof. Suppose P is not a triangle-free path in $U_M(\mathcal{C}, O)$ then there must exist an i such that $u_i \leftarrow u_{i+1} \in U_M(\mathcal{C}, O)$. Since \mathcal{C} is an undirected graph, if there exists a directed edge in $U_M(\mathcal{C}, O)$ then the directed edge must belong to O . This implies $u_i \leftarrow u_{i+1} \in O$. From our assumption, $u_1 \rightarrow u_2 \in O$. This implies $i > 1$. From our assumption, P is a triangle-free path in \mathcal{C} , an undirected connected component of M_a for some $a \in \{1, 2\}$. Also, from the construction, $V_{M_a} \cap V_O = V_{O_a}$. This implies $u_1, u_2, u_i, u_{i+1} \in V_{O_a}$. Since O_a is an induced subgraph of M_a , $(u_1, u_2), (u_i, u_{i+1}) \in E_{O_a}$. This implies $P_{a1}((u_1, u_2), (u_i, u_{i+1})) = 1$, as a subpath of P from (u_1, u_2) to (u_i, u_{i+1}) is also a triangle-free path in M_a , and $(u_1, u_2) \neq (u_i, u_{i+1})$ because $i > 1$. Since $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, from item 3b, $u_i \rightarrow u_{i+1} \in O$, a contradiction. This implies P is a triangle-free path in $U_M(\mathcal{C}, O)$. \square

Proof of Observation 3.29. We will establish the proof through induction. Initially, at the start of the first iteration of the while loop in Alg. 3, we have $\mathcal{S} = (V_{\mathcal{C}})$, and τ is an empty list. Consequently, Observation 3.29 is satisfied at this stage.

Now, let's assume that at the beginning of some iteration of the while loop, $\tau = (a_1, a_2, \dots, a_r)$ and $\mathcal{S} = (S_1, S_2, \dots, S_l)$, and that Observation 3.29 is upheld. We aim to demonstrate that, even at the end of this iteration of the while loop, Observation 3.29 continues to hold.

During the run of the iteration, we do the following: At line 5, we select a canonical source node v from $U_M(\mathcal{C}, O)[S_1]$. We remove vertex v from S_1 and add it to τ (lines 6-7). Subsequently, we update each set S_i as follows: S_i is replaced by $S_i \cap N(v, \mathcal{C})$ and $S_i \setminus N(v, \mathcal{C})$ (line 10). This results in the modified $\mathcal{S} = (S_{11}, S_{12}, \dots, S_{l1}, S_{l2})$, where $S_{11} = (S_1 \setminus v) \cap N(v, \mathcal{C})$, $S_{12} = (S_1 \setminus v) \setminus N(v, \mathcal{C})$, and for $i > 1$, $S_{i1} = S_i \cap N(v, \mathcal{C})$, and $S_{i2} = S_i \setminus N(v, \mathcal{C})$. And, in the end, we remove the empty sets in \mathcal{S} (line 11).

We now demonstrate that if a triangle-free path $P = (u_1, u_2, \dots, u_m)$ satisfies $u_1 \rightarrow u_2 \in O$ and adheres to Observation 3.29 at the start of the iteration, it will continue to satisfy items 1 to 3 of Observation 3.29 even at the end of this iteration. There are two possibilities: either P contains the picked node v (the canonical source node picked at step-5), or it does not. We go through both possibilities one by one.

1. Suppose that P contains the picked node v . Let $u_a = v$, for some $1 \leq a \leq m$. There are two cases: either $a = 1$ (i.e., v is the first node of P), or $a > 1$.

- (a) Suppose $a = 1$. Since $u_a = v$ has been picked from S_1 , this implies that at the start of the iteration, u_1 is in S_1 . Since, at the start of the iteration, P obeys Observation 3.29, item 1 of Observation 3.29 implies that at the start of the iteration, none of the nodes of P are in τ .

At the end of the iteration, u_1 is in τ , and all other nodes of P remain in some set of \mathcal{S} . We will now demonstrate that for all possible pairs of nodes u_i and u_j , where $1 \leq i < j \leq m$, at the end of the iteration, u_i and u_j satisfy items 1 and 2 of Observation 3.29. For $1 = i < j \leq m$, u_i and u_j satisfy items 1 and 2 of Observation 3.29, as at the end of the iteration, u_i is in τ and u_j is in some set of \mathcal{S} .

Suppose $1 < i < j \leq m$. Then, at the start of the iteration, u_i and u_j are in some set S_x and S_y of \mathcal{S} , respectively. Since at the start of the iteration, u_i and u_j satisfy item 2 of Observation 3.29, we have $x \leq y$.

If $x < y$, then at the end of the iteration, the set in which u_i lies (either S_{x1} or S_{x2}) comes before the set in which u_j lies (either S_{y1} or S_{y2}). Thus, if $x < y$, then at the end of the iteration, u_i and u_j satisfy items 1 and 2 of Observation 3.29.

Suppose $x = y$, i.e., at the start of the iteration, both u_i and u_j lie in the same set S_x of \mathcal{S} . Since P is a triangle-free path of a chordal graph \mathcal{C} , from Proposition 2.25, P is a chordal graph. This implies that the only neighbor of $u_1 = v$ in P is u_2 . Therefore, if $i = 2$, then u_i is a neighbor of v and at the end of the iteration, it will move to set S_{x1} ; otherwise, u_i is not a neighbor of v and at the end of the iteration, it will move to S_{x2} . Since $1 < i < j$, j cannot be equal to 2. This implies u_j cannot be a neighbor of v and at the end of the iteration, it will move to S_{x2} . This shows that at the end of the iteration, the set in which u_i lies does not come after the set in which u_j lies in \mathcal{S} . This shows that at the end of the iteration, u_i and u_j satisfy items 1 and 2 of Observation 3.29.

We now show that at the end of the iteration, item 3 of Observation 3.29 is also satisfied.

As shown above, at the end of the iteration, u_1 is in τ , and all the remaining nodes of P are in some set of \mathcal{S} . We have shown that at the end of the iteration, P obeys items 1 and 2 of Observation 3.29. If at the end of the iteration, there does not exist any node of P that is in the first set of \mathcal{S} , then item 3 of Observation 3.29 is satisfied. Suppose at the end of the iteration, there exists a node of P that lies in the first set of \mathcal{S} . Pick the highest i such that at the end of the iteration, u_i is in the first set of \mathcal{S} . If $i = 2$, then at the end of the iteration, only one node of P is in the first set of \mathcal{S} . This implies that at the end of the iteration, item 3 of Observation 3.29 is satisfied. Suppose $i > 2$, then, from item 2 of Observation 3.29, at the end of the iteration, u_2, u_3, \dots, u_i all are in the first set of \mathcal{S} . This further implies even at the start of the iteration, u_2, u_3, \dots, u_i are in the same set of \mathcal{S} , say S_x . But, we know that among these nodes only u_2 is a neighbor of $u_1 = v$. This implies at the end of the iteration, u_2 must be in S_{x1} , and all the remaining nodes u_3, \dots, u_i must be in S_{x2} . This contradicts our assumption that at the end of the iteration, u_2, u_3, \dots, u_i are in the same set of \mathcal{S} . This implies $i > 2$ does not occur. This implies even at the end of the iteration, item 3 of Observation 3.29 is obeyed. This implies that in all the possibilities of this case, items 1 to 3 of Observation 3.29 are satisfied. We will now move on to the second case.

- (b) Suppose $a > 1$. Since $u_a = v$ has been picked from S_1 , this implies that at the start of the iteration, u_a is in S_1 . From item 2 of Observation 3.29, it follows that at the start of the iteration, u_1 must be in either S_1 or in τ . If u_1 is in S_1 at the start of the iteration, then since u_a is also in S_1 , item 2 of Observation 3.29 implies that at the start of the iteration, the nodes between u_1 and u_a in P (i.e., $u_1, u_2, u_3, \dots, u_{a-1}, u_a$) are all in S_1 . This implies $Q = (u_1, u_2, \dots, u_a)$ is a triangle-free path in $\mathcal{C}[S_1]$. But, then, from Claim 3.30, u_a is not a canonical source vertex in $U_M(\mathcal{C}, O)[S_1]$, due to the existence of Q and $u_1 \rightarrow u_2 \in O$. But, this is a contradiction, as we have picked u_a at line 5 of alg. 3 because it is a canonical source vertex in $U_M(\mathcal{C}, O)[S_1]$. Therefore, u_1 cannot be in S_1 , but must be in τ . Then, item 3 of Observation 3.29 implies that at the start of the iteration, the only node of P in S_1 is $u_a = v$.

From item 2 of Observation 3.29, it follows that at the start of the iteration, u_1, u_2, \dots, u_{a-1} are in τ . And, at the end of the iteration, u_a also joins τ .

We will now show that, for all possible pairs of nodes u_i and u_j , where $1 \leq i < j \leq m$, at the end of the iteration, u_i and u_j obey items 1 and 2 of Observation 3.29.

If at the start of the iteration, both u_i and u_j are in τ , then even at the end of the iteration, they remain in τ and obey items 1 and 2. If at the start of the iteration, u_i is in τ and u_j is in some set of \mathcal{S} , then at the end of the iteration, either (a) both u_i and u_j are in τ (this case occurs when $u_j = u_a = v$) and u_i joins τ before u_j , or (b) u_i is in τ , and u_j is in some set of \mathcal{S} (this case occurs when $j \neq a$). Both possibilities obey items 1 and 2.

Suppose at the start of the iteration, both u_i and u_j are in some sets S_x and S_y of \mathcal{S} , respectively. Since at the start of the iteration, u_i and u_j obey Observation 3.29, $x \leq y$.

If $i = a$, i.e., $u_i = v$, then at the end of the iteration, the pair u_i and u_j obey items 1 and 2, as at the end of the iteration, u_i joins τ and u_j remains in some set of \mathcal{S} .

Suppose $u_i \neq v$. If $x < y$, then even at the end of the iteration, the set in which u_i lies (either S_{x1} or S_{x2}) comes before the set in which u_j lies (either S_{y1} or S_{y2}). Suppose $x = y$. Then, at the start of the iteration, both u_i and u_j are in the same set S_x .

Similar to the previous case, P is a chordless path. Therefore, there can be at most two neighbors of $v = u_a$ in P : one is u_{a-1} , and another is u_{a+1} (if $a < m$). As discussed earlier, at the start of the iteration, u_{a-1} is in τ . This, in turn, implies that either none of u_i and u_j are neighbors of $u_a = v$, or only u_i is a neighbor of v (this scenario arises when $i = a + 1$).

If neither is a neighbor of v , then at the end of the iteration, both are in S_{x2} . If u_i is a neighbor of v , then at the end of the iteration, $u_i \in S_{x1}$, and $u_j \in S_{x2}$. This implies the set in which u_i lies does not come after the set in which u_j lies in \mathcal{S} . This shows that, in all the possibilities of this case, items 1 and 2 of Observation 3.29 are obeyed.

Similar to the previous case, we can show that at the end of the iteration, the only possible node of P which can be in the first set of P is u_{a+1} . This validates item 3 of Observation 3.29. Thus, we show that in this case, items 1 to 3 of Observation 3.29 are obeyed. We now move to the next possibility.

2. Suppose P does not contain v . This implies that no node of P moves to τ in this iteration. We will now demonstrate that, at the end of the iteration, for all pairs of nodes u_i and u_j in P such that $1 \leq i < j \leq m$, Observation 3.29 is satisfied. To prove item 1 of Observation 3.29, we assume that at the end of the iteration, u_j is in τ . To prove item 2 of Observation 3.29, we assume that at the end of the iteration, u_j is in some set of \mathcal{S} . Finally, we prove item 3 of Observation 3.29.

Suppose at the end of the iteration, u_j is in τ . This implies that even at the start of the iteration, $u_j \in \tau$, as no node of P moved to τ in this iteration. Since at the start of the iteration, P adheres to Observation 3.29, according to item 1 of Observation 3.29, u_i has joined τ before u_j . Hence, at the end of the iteration, item 1 of Observation 3.29 is upheld.

Suppose at the end of the iteration, u_j is in some set of \mathcal{S} . Then, even at the start of the iteration, u_j must be in some set, say S_y , of \mathcal{S} . Since at the start of the iteration, P obeys Observation 3.29, from item 2 of Observation 3.29, at the start of the iteration, either u_i is in τ or u_i is in some set, say S_x , of \mathcal{S} such that $x \leq y$.

If at the start of the iteration, u_i is in τ then even at the end of the iteration, u_i is in τ , and u_i and u_j obey item 2 of Observation 3.29. Now, suppose at the start of the iteration, u_i is in some set S_x of \mathcal{S} .

If $x < y$, then at the end of the iteration, u_i is in a set of \mathcal{S} (either S_{x1} or S_{x2}) that comes before the set in which u_j belongs (either S_{y1} or S_{y2}), as both S_{x1} and S_{x2} come before S_{y1} and S_{y2} . This implies that Observation 3.29 is obeyed in this case.

If $x = y$, then either both u_i and u_j belong to the same set S_{x1} (if both are neighbors of v) or both u_i and u_j belong to the same set S_{x2} (if neither of them is a neighbor of v) or $u_i \in S_{x1}$ and $u_j \in S_{x2}$ (if u_i is a neighbor of v , and u_j is not a neighbor of v) or $u_j \in S_{x1}$ and $u_i \in S_{x2}$ (if u_j is a neighbor of v , and u_i is not a neighbor of v). The first three possibilities obey Observation 3.29, as in the first and second cases, u_i and u_j both are in the same set of \mathcal{S} , and in the third case, u_i is in S_{x1} , which comes before S_{x2} in which u_j belongs. We now show that the last possibility cannot occur.

Suppose u_j is a neighbor of v , and u_i is not a neighbor of v . Pick the least k such that u_k comes after u_i in P , and u_k is a neighbor of v . Such a u_k exists due to the presence of u_j .

Then, $Q = (u_1, u_2, \dots, u_k, v)$ is a triangle-free path in \mathcal{C} such that $u_1 \rightarrow u_2 \in O$.

Now, from our induction hypothesis, at the start of the iteration, all the triangle-free paths of \mathcal{C} obey Observation 3.29. This implies Q also obeys Observation 3.29 at the start of the iteration.

v has been picked in this iteration implying that at the start of the iteration, $v \in S_1$. From item 2 of Observation 3.29, at the start of the iteration, all other nodes of Q are either in τ or in S_1 . Since u_i is

not in τ at the start of the iteration, this implies that at the start of the iteration, u_i is in S_1 along with v . Then, item 3 of Observation 3.29 implies that at the start of the iteration, all the nodes of Q are in S_1 .

But then, v cannot be a canonical source node in $U_M(\mathcal{C}, O)[S_1]$ due to the existence of a triangle-free path Q in $U_M(\mathcal{C}, O)[S_1]$ and $u_1 \rightarrow u_2 \in O$. The fact that Q is a triangle-free path follows from Claim 3.30.

This implies this possibility cannot occur. This shows that in this case, items 1 and 2 of Observation 3.29 are obeyed.

We now show that at the end of the iteration, P also obeys item 3 of Observation 3.29. Since in this iteration, the picked node v is not a node of P , this implies that the set of nodes of P in τ at the start of the iteration is the same as the nodes of P in τ at the end of the iteration. If none of the nodes of P are in τ , then item 2 is satisfied.

Suppose a node of P is in τ . Since at the end of the iteration, P obeys items 1 and 2 of Observation 3.29, there exists some $i \geq 1$ such that at the end of the iteration, u_1, u_2, \dots, u_i are in τ and each of the remaining nodes of P is in some set of \mathcal{S} . If at the end of the iteration, no node of P is in the first set of \mathcal{S} , then item 3 is obeyed.

Suppose at the end of the iteration, there exists a node of P in the first set of \mathcal{S} . Pick the highest j such that at the end of the iteration, u_j is in the first set of \mathcal{S} . Since at the end of the iteration, P obeys items 1 and 2 of Observation 3.29, at the end of the iteration, there must be $u_{i+1}, u_{i+2}, \dots, u_j$ all are in the first set of \mathcal{S} . If $j = i + 1$, then item 3 is obeyed, as in that case there exists only a single node of P in the first set of \mathcal{S} . We now show that $j = i + 1$.

Suppose $j > i + 1$. Then, at the start of the iteration when u_i has been picked and has been moved to τ , $u_{i+1}, u_{i+2}, \dots, u_j$ must be in the same set, say S_x , of \mathcal{S} (because from alg. 3, once two nodes of \mathcal{C} gets separated and move to two different set of \mathcal{S} , they never unite). Since P is a triangle-free path of a chordal graph \mathcal{C} , from Proposition 2.25, P is a chordless path. This implies that only u_{i+1} is a neighbor of u_i and rest of the nodes $u_{i+2}, u_{i+3} \dots, u_j$ are not the neighbors of u_i in P . This further implies that at the end of the iteration in which u_i has been picked at step 5 of alg. 3, u_{i+1} moves to the set S_{x1} of \mathcal{S} and u_j moves to S_{x2} of \mathcal{S} . This further implies that even at the end of the current iteration, u_{i+1} and u_j are in different sets of \mathcal{S} , a contradiction. This implies $j > i + 1$ cannot occur. This shows that at the end of the iteration, item 3 of Observation 3.29 is obeyed.

This demonstrates that in all scenarios, at the end of the iteration, items 1 to 3 of Observation 3.29 are true. Thus, Observation 3.29 is validated. \square

Observation 3.29 implies the following corollaries:

Corollary 3.31. *For input \mathcal{C} and O , let τ be an LBFS ordering returned by Alg. 3. If (u_1, u_2, \dots, u_l) is a chordless path in \mathcal{C} such that $u_1 \rightarrow u_2 \in O$, then $\tau(u_1) < \tau(u_2) < \dots < \tau(u_l)$.*

Corollary 3.32. *For input \mathcal{C} and O , let τ be an LBFS ordering returned by Alg. 3. For $a - b \in \mathcal{C}$, if $a \rightarrow b \in O$, then $\tau(a) < \tau(b)$.*

As discussed earlier, Alg. 3 is a modified LBFS ordering of the LBFS ordering given by Rose et al. (Rose, Tarjan, and Lueker (1976)). Instead of picking a node arbitrarily at line 5 of Alg. 3, we choose an arbitrary canonical source node. Since there always exists a canonical source node (as shown in Claim 3.26), Alg. 3 outputs an LBFS ordering. Corollary 3.32 demonstrates that the ordering returned by Alg. 3 is consistent with O . Lemma 3.33 summarizes this.

Lemma 3.33. *Let H be an undirected graph, H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H . For each $i \in \{1, 2\}$, let S_i be a subset of V_{H_i} such that $I \subseteq S_i$, M_i be an MEC of H_i , and the shadow of M_i on $S_i \cup N(S_i, H_i)$ be (O_i, P_{i1}, P_{i1}) . Let $O \in \mathcal{E}(O_1, P_{11}, P_{12}, P_{12}, O_2, P_{21}, P_{22})$. For $i \in \{1, 2\}$, let \mathcal{C} be a universal critical component of M_i . For inputs \mathcal{C} and O , Alg. 3 returns an LBFS ordering τ that satisfies O , i.e., for $u \rightarrow v \in O$ such that $u, v \in V_{\mathcal{C}}$, $\tau(u) < \tau(v)$.*

3.4 Sufficient condition for a shadow of an MEC of H

In this subsection, we show that items 1 and 2 of Lemma 3.25 constitutes the required sufficient condition.

Lemma 3.34. *Let H be an undirected graph, and H_1 and H_2 be the induced subgraph of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1, S_2 be the subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let M_1 be an MEC of H_1 , and M_2 be an MEC of H_2 . Let (O_1, P_{11}, P_{12}) be the shadow of M_1 on $S_1 \cup N(S_1, H_1)$, and (O_2, P_{21}, P_{22}) be the shadow of M_2 on $S_2 \cup N(S_2, H_2)$. Let $A = H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)]$ and $O \in \text{PMEC}(A)$. If O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$ then there exists a unique MEC M such that $M[V_O] = O$, and $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$. If $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22}) = (P_1, P_2)$ then the shadow of M on V_O is (O, P_1, P_2) .*

Algorithm 4: Construct MEC

Input : Three graphs M_1, M_2 , and O such that
 H is an undirected graph and
 H_1 and H_2 are two induced subgraphs of H such that
 $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H .
 S_1 and S_2 are subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$.
 M_1 and M_2 are MECs of H_1 and H_2 , respectively.
 $\forall a \in \{1, 2\}$, the shadow of M_a on $S_a \cup N(S_a, H_a)$ is (O_a, P_{a1}, P_{a2}) , and
 $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$.

Output: M : an MEC of $H_1 \cup H_2$ such that
 $M \in \text{MEC}(H, O, P_1, P_2)$, $M[V_O] = O$, and $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$.

- 1 $M \leftarrow U_M(M_1, M_2, O)$
- 2 $I \leftarrow V_{M_1} \cap V_{M_2}$
- 3 **foreach** $i \in \{1, 2\}$ **do**
- 4 **foreach** undirected connected component \mathcal{C} of M_i **do**
- 5 $\tau \leftarrow \text{LBFS}(\mathcal{C}, O)$.
- 6 Denote a vertex v of \mathcal{C} as u_i if $\tau(v) = i$.
- 7 $a \leftarrow 2$
- 8 **while** $a \leq |V_{\mathcal{C}}|$ **do**
- 9 $b \leftarrow a - 1$
- 10 **while** $b \geq 1$ **do**
- 11 **if** $\exists c$ such that $c < b$, and $u_c - u_b - u_a$ is an induced subgraph of \mathcal{C} , and
 $u_c \rightarrow u_b - u_a \in M$, or $\exists c$ such that $b < c < a$, and $u_b - u_c - u_a - u_b \in \mathcal{C}$, and
 $u_b \rightarrow u_c \rightarrow u_a - u_b \in M$ **then**
- 12 Replace the edge $u_b - u_a$ in M with $u_b \rightarrow u_a$.
- 13 **end**
- 14 $b = b - 1$
- 15 **end**
- 16 $a = a + 1$
- 17 **end**
- 18 **end**
- 19 **end**
- 20 **return** M

Proof. Suppose $M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12})$, $M_2 \in \text{MEC}(H_2, O_2, P_{21}, P_{22})$, $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, and $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$. We show that there exists a unique MEC M such that $M[V_O] = O$, $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$, and the shadow of M on V_O is (O, P_1, P_2) . We first construct such an MEC M using alg. 4. Alg. 4 follows the following steps to construct the MEC:

1. Initialize $M = U_M(M_1, M_2, O)$ (line-1 of alg. 4), a Markov union (Definition 2.20) of M_1, M_2 and O .
2. For each $i \in \{1, 2\}$ (lines 3-19 of alg. 4), for each undirected connected component \mathcal{C} of M_i (lines 4-18 of alg. 4), we do the following:
 - (a) Construct an LBFS ordering τ of \mathcal{C} using alg. 3 for input \mathcal{C} and O (line 5 of alg. 4).
 - (b) For $u - v \in M$ such that $\tau(u) < \tau(v)$, replace $u - v$ with $u \rightarrow v$ (line 12 of alg. 4) if either
 - i. there exists a vertex $x \in \mathcal{C}$ such that $\tau(x) < \tau(u)$, and there exists $x - u - v$, an induced subgraph of \mathcal{C} , such that $x \rightarrow u \in M$ (line 11), or
 - ii. there exists a vertex $x \in \mathcal{C}$ such that $\tau(u) < \tau(x) < \tau(v)$, and $u \rightarrow x, x \rightarrow v \in M$ (line 11).

We begin by demonstrating that alg. 4 correctly executes steps 1 and 2. The implementation of step 1 can be found at line 1 of alg. 4. Next, we verify that for each $i \in \{1, 2\}$ and for every undirected connected component \mathcal{C} of M_i , steps 2a and 2b are executed correctly.

Step 2a is implemented at line 5 of alg. 4, where we construct an LBFS ordering τ for \mathcal{C} . To demonstrate that step 2b is also executed correctly, we utilize Corollary 3.42 to provide assurance in the accuracy of alg. 4.

Let's suppose that step 2b is not accurately executed by alg. 4. This would imply the existence of an x such that either (a) $x - u - v$ forms an induced subgraph of \mathcal{C} , with $\tau(x) < \tau(u) < \tau(v)$, and $x \rightarrow u - v \in M$, or (b) $u - x - v - u$ forms an induced subgraph of \mathcal{C} , with $\tau(u) < \tau(x) < \tau(v)$, and $u \rightarrow x, x \rightarrow v - u \in M$.

Let's address both cases individually:

Case (a): Assuming there exists an x such that $x - u - v$ forms an induced subgraph of \mathcal{C} , $\tau(x) < \tau(u) < \tau(v)$, and $x \rightarrow u - v \in M$. According to Corollary 3.42, $x \rightarrow u$ was added to M either during initialization or when $a = \tau(u)$ and $b = \tau(x)$. This implies that when we have $a = \tau(v)$ and $b = \tau(u)$, $x \rightarrow u$ is in M . However, during the execution of lines 11–13 (when we have $a = \tau(v)$ and $b = \tau(u)$), $u \rightarrow v$ is added to M , leading to a contradiction. Therefore, this case cannot occur.

Case (b): Assuming there exists an x such that $u - x - v - u$ forms an induced subgraph of \mathcal{C} , with $\tau(u) < \tau(x) < \tau(v)$, and $u \rightarrow x, x \rightarrow v - u \in M$. According to Corollary 3.42, $u \rightarrow x$ was added to M either during initialization or when $a = \tau(x)$ and $b = \tau(u)$, and $x \rightarrow v$ was added to M either during initialization or when $a = \tau(v)$ and $b = \tau(x)$. This implies that when we have $a = \tau(v)$ and $b = \tau(u)$, both $u \rightarrow x$ and $x \rightarrow u$ are in M . However, during the execution of lines 11–13 (when we have $a = \tau(v)$ and $b = \tau(u)$), $u \rightarrow v$ is added to M , again leading to a contradiction. Therefore, this case also cannot occur.

In conclusion, we have demonstrated that 2b is executed correctly by alg. 4. This further affirms that alg. 4 accurately implements steps 1 and 2.

Lemmas 3.35 to 3.39 validate M .

Lemma 3.35. $M[V_O] = O$.

Lemma 3.36. M is an MEC of H .

Lemma 3.37. $\mathcal{P}(M, V_{M_1}, V_{M_2}) = (M_1, M_2)$.

Lemma 3.38. If there exists an MEC M' of H such that $M'[V_O] = O$, and $\mathcal{P}(M', V_{M_1}, V_{M_2}) = (M_1, M_2)$ then $M' = M$.

Lemma 3.39. If $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22}) = (P_1, P_2)$ then $M \in \text{MEC}(H, O, P_1, P_2)$.

Lemma 3.36 shows that the graph returned by alg. 4 is an MEC. Lemma 3.35 shows that O is an induced subgraph of M , i.e., $M[V_O] = O$, and Lemma 3.37 shows that M_1 and M_2 are projections of M . Lemma 3.38 shows that there exists a unique MEC with O as its induced subgraph, and M_1 and M_2 as its projections. Additionally, Lemma 3.39 shows that if $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$, then the shadow of M on V_O is (O, P_1, P_2) . This also implies that given $O, P_{11}, P_{12}, P_{21}$, and P_{22} , we can compute the shadow of M on V_O .

Proofs of Lemmas 3.35 to 3.39 are provided below. Given these lemmas, Lemma 3.34 follows immediately. \square

We now prove Lemma 3.35. In preparation for proving Lemma 3.35, we first prove the following observation. The proof of Lemma 3.35 appears on page 49.

Observation 3.40. *For $i \in \{1, 2\}$, for any undirected connected component \mathcal{C} of M_i , at line-5 of alg. 4, let τ be the LBFS ordering returned by alg. 3. Then for $u - v \in \mathcal{C}$, at any iteration of the while loop (lines 8-17) of alg. 4, if $u \rightarrow v \in M$ then either of the following occurs:*

1. $u \rightarrow v \in O$
2. $\exists a \rightarrow b \in O$ such that there exists a chordless path from (a, b) to (u, v) in \mathcal{C} such that $(a, b) \neq (u, v)$.
3. $\exists a \rightarrow b \in O$ such that there exist chordless paths P and Q such that P is a chordless path from (a, b) to v in \mathcal{C} , and Q is a chordless path from (v, u) to a in \mathcal{C} , such that $b \neq v$ and $u \neq a$.

Proof. We start with the following observation:

Observation 3.41. *For $i \in \{1, 2\}$, if $u - v \in M_i$ and $u \rightarrow v \in U_M(M_1, M_2, O)$ then $u \rightarrow v \in O$.*

Proof. From Definition 2.20, if $u - v \in M_1$ and $u \rightarrow v \in U_M(M_1, M_2, O)$ then either $u \rightarrow v \in M_2$ or $u \rightarrow v \in O$. $u \rightarrow v \in M_2$ implies that $u, v \in V_{M_1} \cap V_{M_2} = I$. Since (O_2, P_{21}, P_{22}) is the shadow of M_2 on V_{O_2} , therefore, from item 1 of Definition 2.36, $O_2 = M_2[V_{O_2}]$. This implies $u \rightarrow v \in O_2$ as from the construction, $I \subseteq V_{O_2}$. Since $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, therefore, from item 1 of Definition 3.24, we have $u \rightarrow v \in O$. Thus, we can say that if $u - v \in M_1$ and $u \rightarrow v \in U_M(M_1, M_2, O)$ then $u \rightarrow v \in O$. Similarly, if $u - v \in M_2$ and $u \rightarrow v \in U_M(M_1, M_2, O)$ then $u \rightarrow v \in O$. This completes the proof of Observation 3.41. \square

Let us number the undirected connected components of M_1 and M_2 in the sequence in which they are considered on line 4 as $\mathcal{C}_1, \mathcal{C}_2, \dots$. Suppose that Observation 3.40 is true for all \mathcal{C}_β with $\beta < \alpha$. We now prove by induction on the iteration of the while loop (lines 8-17) of alg. 4 corresponding to the connected component \mathcal{C}_α that the Observation 3.40 is true for $\mathcal{C} = \mathcal{C}_\alpha$ as well.

Base case: We first show that at the start of the first iteration of the while loop corresponding to $\mathcal{C} = \mathcal{C}_\alpha$, it must be the case that for any edge $u - v \in \mathcal{C}$, if $u \rightarrow v \in M$ then $u \rightarrow v \in O$. This will establish that Observation 3.40 holds at the start of the first iteration of the while loop corresponding to the undirected connected chordal component $\mathcal{C} = \mathcal{C}_\alpha$.

Suppose, if possible, at the start of the first iteration of the while loop, there exists $u - v \in \mathcal{C}$ such that $u \rightarrow v \in M$. There are two possibilities: either $u \rightarrow v$ is added at the initialization step, i.e., at line 1 of alg. 4, or $u \rightarrow v$ is added to M during the processing of some \mathcal{C}_γ with $\gamma < \alpha$ at lines 3-19 of alg. 4. At the initialization step, $M = U_M(M_1, M_2, O)$. \mathcal{C} is an undirected connected component of M_i for some $i \in \{1, 2\}$. This implies $u - v \in M_i$. Then from Observation 3.41, if $u \rightarrow v \in U_M(M_1, M_2, O)$ then $u \rightarrow v \in O$. This validates Observation 3.40. We now go through the other possibility that $u \rightarrow v$ has been added to M during the processing of some \mathcal{C}_γ with $\gamma < \alpha$.

W.l.o.g., let us assume that \mathcal{C}_α is an undirected connected component of M_1 . Then \mathcal{C}_γ cannot be an undirected connected component of M_1 (two distinct undirected connected components of a graph do not share any vertex). This implies \mathcal{C}_γ is an undirected connected component of M_2 . This further implies $u - v$ is in both M_1 and M_2 . That means $u, v \in V_{M_1} \cap V_{M_2} = I \subseteq V_O$. Since $u - v \in \mathcal{C}_\gamma$, an undirected connected component of M_2 , therefore, $u - v \in M_2$. Also, since $u, v \in I \subseteq V_{O_2}$, and O_2 is an induced subgraph of M_2 (from item 1 of Definition 2.36 as (O_2, P_{21}, P_{22}) is a shadow of M_2), therefore, $u - v \in O_2$. Since we assumed that Observation 3.40 is true for such a \mathcal{C}_γ , one of the three conditions (items 1 to 3) of Observation 3.40 must be true for $u - v$, as $u - v$ is considered as an edge in \mathcal{C}_γ .

If item 1 of Observation 3.40 holds then we get $u \rightarrow v \in O$. If one of item 2 or item 3 of Observation 3.40 holds, then in the undirected connected chordal component \mathcal{C}_γ of M_2 , there exists an edge $x - y \in \mathcal{C}_\gamma$ such that $x \rightarrow y \in O$, and either (a) $(x, y) \neq (u, v)$ and there exists a chordless path from (x, y) to (u, v) in \mathcal{C}_γ (due to item 2 of Observation 3.40), or (b) $y \neq v$, $u \neq x$, and there exist chordless paths Q_1 and Q_2 in \mathcal{C}_γ such that Q_1 is a chordless path from (x, y) to v , and Q_2 is a chordless path from (v, u) to x (due to item 3

of Observation 3.40). Since C_γ is an undirected connected component of M_2 , therefore, a chordless path in C_γ is a chordless path of M_2 . And, from Observation 2.24, every chordless path is a triangle-free path. This implies chordless paths in C_γ are triangle-free paths of M_2 . Also, since $(x, y) \in C_\gamma$, an undirected connected component of M_2 , and $x \rightarrow y \in O$, this implies $x, y \in V_{M_2} \cap V_O = V_{O_2}$. Since (O_2, P_{21}, P_{22}) is the shadow of M_2 on V_{O_2} , from Definition 2.36, O_2 is an induced subgraph of M_2 . This implies $(x, y) \in E_{O_2}$, and if there exists a chordless path in C_γ , from (x, y) to (u, v) , then $P_{21}((x, y), (u, v)) = 1$, and if there exist chordless paths in C_γ , from (x, y) to v and from (v, u) to x then $P_{22}((x, y), v) = P_{22}((v, u), x) = 1$. Since $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, therefore, in both the possibilities, from item 3 of Definition 3.24, we again have $u \rightarrow v \in O$. This shows that for any $u - v \in C_\alpha$, when C_α is processed at lines 4-18 of alg. 4, at the start of the first iteration of the while loop at lines 8-17, if $u \rightarrow v \in M$ then $u \rightarrow v \in O$. This establishes the base case of the induction.

Induction step: Suppose now that at the start of an iteration of the while loop (at lines 8-17 of alg. 4), Observation 3.40 is valid, that means, at the start of the iteration, for any edge $u' - v' \in \mathcal{C}$, if $u' \rightarrow v' \in M$ then $u' \rightarrow v'$ follows Observation 3.40. We show that even at the end of the iteration, Observation 3.40 is valid, i.e., for any $u - v \in \mathcal{C}$, during the iteration, if $u \rightarrow v$ is added to M then the new directed edge $u \rightarrow v$ obeys Observation 3.40. During the iteration, let there exists an edge $u - v \in \mathcal{C}$ such that $u \rightarrow v$ is added to M in the while loop while processing $\mathcal{C} = \mathcal{C}_\alpha$. We show that $u \rightarrow v$ obeys Observation 3.40. Let τ be the LBFS ordering of \mathcal{C} returned by alg. 3 at line 5 of alg. 4 for the input \mathcal{C} and O . From lines 11-13 of alg. 4, if $u \rightarrow v$ is added to M then at the start of the iteration, either of the following has occurred: (a) there exists an induced subgraph $x - u - v \in \mathcal{C}$ such that $\tau(x) < \tau(u) < \tau(v)$, and $x \rightarrow u - v \in M$, or (b) there exists an induced subgraph $u - x - v - u \in \mathcal{C}$ such that $\tau(u) < \tau(x) < \tau(v)$, $u \rightarrow x, x \rightarrow v \in M$ and $u - v \in M$. We go through each case and show that in each case $u \rightarrow v$ obeys Observation 3.40.

1. Suppose that during the iteration, $u - v \in M$ is replaced with $u \rightarrow v$ because there exists an induced subgraph $x - u - v \in \mathcal{C}$ such that $\tau(x) < \tau(u) < \tau(v)$, and at the start of the iteration, $x \rightarrow u - v \in M$. By the induction hypothesis, we have that $x \rightarrow u \in M$ because one of the following holds: (i) $x \rightarrow u \in O$, or (ii) $\exists a \rightarrow b \in O$ such that there exists a chordless path $(p_1 = a, p_2 = b, \dots, p_{l-1} = x, p_l = u)$ from $a - b$ to $x - u$ in \mathcal{C} (with $(a, b) \neq (x, u)$), or (iii) $\exists a \rightarrow b \in O$ such that there exists a chordless path from $a - b$ to u , and a chordless path from $u - x$ to a in \mathcal{C} (with $b \neq u$ and $x \neq a$). We start with the first possibility.

(a) Suppose that $x \rightarrow u \in M$ because $x \rightarrow u \in O$. Then the new directed edge $u \rightarrow v$ obeys item 2 of Observation 3.40, due to the existence of the chordless path (x, u, v) in \mathcal{C} and $x \rightarrow u \in O$. We now go through the second possibility.

(b) Suppose that $x \rightarrow u \in M$ because $\exists a \rightarrow b \in O$ such that there exists a chordless path $Q = (p_1 = a, p_2 = b, \dots, p_{l-1} = x, p_l = u)$ from $a - b$ to $x - u$ in \mathcal{C} (with $(a, b) \neq (x, u)$). By the hypothesis of this case, $Q' = (x, u, v)$ is a chordless path in \mathcal{C} (remember in this case, $x - u - v$ is an induced subgraph of \mathcal{C}). Then, from Lemma 2.31, $P = (p_1 = a, p_2 = b, \dots, p_{l-1} = x, p_l = u, v)$, concatenation of Q and Q' , is a chordless path in \mathcal{C} .

Thus, we have a chordless path P in \mathcal{C} from (a, b) to (u, v) with $(a, b) \neq (u, v)$ (Q being a path precludes $a = u$) and $a \rightarrow b \in O$. This further implies that the new directed edge $u \rightarrow v$ obeys item 2 of Observation 3.40. We now move to the third possibility.

(c) Suppose that $x \rightarrow u \in M$ because $\exists a \rightarrow b \in O$ such that there exists a chordless path $Q_1 = (p_1 = a, p_2 = b, \dots, p_l = u)$ from (a, b) to u in \mathcal{C} , and a chordless path $Q_2 = (q_1 = u, q_2 = x, \dots, q_m = a)$ from (u, x) to a in \mathcal{C} (with $b \neq u$ and $x \neq a$).

Since Q_1 is a chordless path and $a \rightarrow b \in O$, according to Corollary 3.31, we have $\tau(a) = \tau(p_1) < \tau(p_2) < \tau(p_3) < \dots < \tau(p_l) = \tau(u)$.

Now, we claim that

$$\text{For } 1 \leq i \leq l - 2, v - p_i \notin \mathcal{C}. \quad (4)$$

To see this, suppose, if possible, that for some $i \leq l-2$, $v - p_i \in \mathcal{C}$. As argued above, $\tau(p_i) < \tau(u)$, and by the hypothesis for this case, we have $\tau(u) < \tau(v)$. Then, from Observation 2.14, we must have $p_i - u \in \mathcal{C}$. Since $i \leq l-2$, this constitutes a chord of Q_1 and contradicts the fact that Q_1 is a chordless path. This establishes eq. (4).

Due to eq. (4), only the following two options are now allowed: either (i) for each $1 \leq i \leq l-1$, $v - p_i \notin \mathcal{C}$, or (ii) $v - p_{l-1} \in \mathcal{C}$ and for each $1 \leq i \leq l-2$, $v - p_i \notin \mathcal{C}$.

If the first option holds, then $(p_1 = a, p_2 = b, \dots, p_{l-1}, p_l = u, v)$ is a chordless path (because Q_1 is a chordless path) from (a, b) to (u, v) with $(a, b) \neq (u, v)$ ($a = u$ is precluded by Q_1 being a path), and this implies that $u \rightarrow v$ obeys item 2 of Observation 3.40.

In case the second option holds, $Q_3 = (p_1 = a, p_2 = b, \dots, p_{l-1}, v)$ is a chordless path from (a, b) to v . Since $\tau(p_2) = \tau(b) < \tau(u)$ (as argued above) and $\tau(u) < \tau(v)$ (by the hypothesis for Case 1), we also have $b \neq v$.

Now, we claim that

$$Q_4 = (v, q_1 = u, q_2 = x, \dots, q_m = a) \text{ is a chordless path in } \mathcal{C} \text{ from } (v, u) \text{ to } a. \quad (5)$$

By the hypothesis of this case, $Q'_2 = (v, u, x)$ is a chordless path in \mathcal{C} (remember $x - u - v$ is an induced subgraph of \mathcal{C} in this case). From Lemma 2.31, $Q_4 = (v, q_1 = u, q_2 = x, q_3, \dots, q_m = a)$, concatenation of Q'_2 and Q_2 , is a chordless path in \mathcal{C} . Also, Q_2 being a path precludes $u = a$.

We thus obtain a chordless path Q_4 from (v, u) to a (with $u \neq a$). Along with the chordless path Q_3 from (a, b) to v (with $b \neq v$ and $a \rightarrow b \in O$), this implies that $u \rightarrow v$ obeys item 3 of Observation 3.40.

We now move to the second case.

2. Suppose that during the iteration, $u - v \in M$ is replaced by $u \rightarrow v$ because there exists an induced subgraph $u - x - v - u \in \mathcal{C}$ such that $\tau(u) < \tau(x) < \tau(v)$, $u \rightarrow x, x \rightarrow v \in M$ and $u - v \in M$. Again, by the induction hypothesis, we have that $u \rightarrow x \in M$ because one of the following holds: (i) $u \rightarrow x \in O$, or (ii) $\exists a \rightarrow b \in O$ such that there exists a chordless path $(p_1 = a, p_2 = b, \dots, p_{l-1} = u, p_l = x)$ from (a, b) to (u, x) in \mathcal{C} (with $(a, b) \neq (u, x)$), or (iii) $\exists a \rightarrow b \in O$ such that there exists a chordless path from (a, b) to v , and a chordless path from (v, x) to a in \mathcal{C} (with $b \neq v$ and $x \neq a$). We again go through each possibility.

- (a) Suppose $u \rightarrow x \in M$ because $u \rightarrow x \in O$. Again, by the induction hypothesis, we have that $x \rightarrow v \in M$ because one of the following holds: (i) $x \rightarrow v \in O$, or (ii) $\exists c \rightarrow d \in O$ such that there exists a chordless path $(q_1 = c, q_2 = d, \dots, q_{m-1} = x, q_m = v)$ from (c, d) to (x, v) in \mathcal{C} (with $(c, d) \neq (x, v)$), or (iii) $\exists c \rightarrow d \in O$ such that there exists a chordless path from (c, d) to v , and a chordless path from (v, x) to c in \mathcal{C} (with $d \neq v$ and $x \neq c$). We go through each sub-case.

- i. Let's assume that $x \rightarrow v \in M$ due to the fact that $x \rightarrow v \in O$. The presence of both $u \rightarrow x$ and $x \rightarrow v$ in O implies that u, v , and x are all elements of V_O .

At the start of the iteration, we have $u - v \in M$. This implies even after the initialization step (line-1), we have $u - v \in M$. From the construction of M , $\text{skeleton}(M[V_O]) = \text{skeleton}(O)$. That means $u - v \in \text{skeleton}(O)$. Since O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, by Definition 3.24, O is a partial MEC. Then, from item 1 of Definition 2.21, O is a chain graph. This implies $u \rightarrow v \in O$ (or else O would contain a directed cycle (u, w, v, u)).

But, then, from the initialization of M (remember M is initialized with $U_M(M_1, M_2, O)$), $u \rightarrow v \in M$, a contradiction. This contradiction implies that this particular sub-case cannot occur. We will now proceed to examine the remaining sub-cases.

- ii. Let's assume that $x \rightarrow v \in M$ because there exists $c \rightarrow d \in O$ such that there is a chordless path $Q = (p_1 = c, p_2 = d, \dots, p_{l-2}, p_{l-1} = x, p_l = v)$ from (c, d) to (x, v) in \mathcal{C} (with $(c, d) \neq (x, v)$). It's worth noting that, due to the hypothesis of this case, u and v are adjacent, while c and v are not (since Q is a chordless path). This implies $c \neq u$.

According to Corollary 3.31, we have $\tau(c) = \tau(p_1) < \tau(p_2) < \dots < \tau(p_{l-1}) = \tau(x) < \tau(p_l) = \tau(v)$. Since $\tau(p_{l-2}) < \tau(x)$, and $\tau(u) < \tau(x)$ (as per the hypothesis of this case), Observation 2.14 implies that $u - p_{l-2} \in \mathcal{C}$ (since x is adjacent to both u and p_{l-2} in \mathcal{C}).

Let $1 \leq i \leq l-2$ be the smallest number such that $u - p_i \in \mathcal{C}$. If $i > 1$, then, as Q is a chordless path, $Q' = (p_1 = c, p_2 = d, \dots, p_i, u, v)$ is a chordless path in \mathcal{C} from (c, d) to (u, v) with $(c, d) \neq (u, v)$ (since $c \neq u$). Since $c \rightarrow d \in O$, this implies that $u \rightarrow v$ satisfies item 2 of Observation 3.40.

If $i = 1$, then $Q_1 = (v, u, c)$ is a chordless path from (v, u) to c (c is not adjacent to v in \mathcal{C} because Q is a chordless path). Since $Q = (p_1 = c, p_2 = d, \dots, p_{l-1} = x, p_l = v)$ is a chordless path from (c, d) to v (with $d \neq v$) and Q_1 is a chordless path from (v, u) to c (with $c \neq u$), $u \rightarrow v$ satisfies item 3 of Observation 3.40.

Therefore, in this subcase, $u \rightarrow v$ complies with Observation 3.40. We will now consider the remaining sub-case.

- iii. Suppose $x \rightarrow v \in M$ because $\exists c \rightarrow d \in O$ such that there exists a chordless path $Q_1 = (q_1 = c, q_2 = d, \dots, q_m = v)$ from (c, d) to v in \mathcal{C} (in which $d \neq v$), and a chordless path $Q_2 = (p_1 = v, p_2 = x, \dots, p_l = c)$ from (v, x) to c in \mathcal{C} (in which $c \neq x$). Note that c is not adjacent to v (since Q_1 is a chordless path), and u is adjacent to v (by the hypothesis of this case). We thus also have $u \neq c$.

We now claim that there exists a chordless path Q from (v, u) to c in \mathcal{C} . This further implies that $u \rightarrow v$ obeys item 3 of Observation 3.40 due to the existence of chordless paths Q_1 from (c, d) to v (with $d \neq v$) and Q from (v, u) to c (with $u \neq c$) such that $c \rightarrow d \in O$.

Proof of the claim. There are two possibilities: either $\tau(p_3) < \tau(p_2)$, or $\tau(p_3) > \tau(p_2)$. We go through both possibilities and show that in each possibility there exists a chordless path from (v, u) to c .

Suppose $\tau(p_3) < \tau(p_2 = x)$. Since we also have $\tau(u) < \tau(x)$ (by the hypothesis of this case), and since x is adjacent to both u and p_3 , Observation 2.14 implies that $u - p_3 \in \mathcal{C}$. Pick the highest i such that $u - p_i \in \mathcal{C}$. Since $u - p_3 \in \mathcal{C}$, i must be greater than 2. Then, $Q = (v, u, p_i, \dots, p_l = c)$ is a chordless path from (v, u) to c (since v is adjacent to u and Q_1 is a chordless path). We now move to the other possibility.

Suppose $\tau(p_3) > \tau(p_2 = x)$. Since Q_2 is a chordless path in \mathcal{C} , Observation 2.15 implies that $\tau(p_2 = x) < \tau(p_3) < \dots < \tau(p_l = c)$, i.e., $\tau(x) < \tau(c)$. From the hypothesis of this case, $\tau(u) < \tau(x)$. This implies that $\tau(u) < \tau(c)$. We now claim that u must be adjacent to every vertex in the chordless path $Q_1 = (q_1 = c, q_2 = d, \dots, q_m = v)$, and in particular to c . Since Q_1 is a chordless path and $c \rightarrow d \in O$, therefore, from Corollary 3.31, $\tau(q_1) = \tau(c) < \tau(q_2) = \tau(d) < \dots < \tau(q_{m-1}) < \tau(q_m) = \tau(v)$. Since $\tau(u) < \tau(c)$, we have $\tau(u) < \tau(q_i) < \tau(q_{i+1})$ for every $1 \leq i \leq m-1$. Suppose if possible that there exists an i such that $q_i - u \notin \mathcal{C}$, then pick the highest such i . Since u is adjacent to $v = q_m$ in \mathcal{C} (by the hypothesis of this case), we must have $i \leq m-1$. Further, by the choice of i , we must have that u is adjacent to q_{i+1} . But, then, since $\tau(q_i) < \tau(q_{i+1})$ and $\tau(u) < \tau(q_{i+1})$, Observation 2.14 implies that $q_i - u \in \mathcal{C}$ (since q_i and u both are adjacent to q_{i+1}), which is a contradiction to the choice of i . Thus, u is adjacent to every vertex in Q_1 , and in particular to c . This implies that $Q = (c, u, v)$ is a chordless path in \mathcal{C} (as argued above v and c are not adjacent and $u \neq c$).

Thus, we show that in both possibilities there exists a chordless path Q from (v, u) to c . \square

We now move to the next possibility.

- (b) Let's assume that $u \rightarrow x \in M$ because there exists an edge $a - b \in \mathcal{C}$ such that $a \rightarrow b \in O$, and there exists a chordless path $Q = (p_1 = a, p_2 = b, \dots, p_{l-1} = u, p_l = x)$ from (a, b) to (u, x) in \mathcal{C} (with $(a, b) \neq (u, x)$). Note that we trivially also have $a \neq u$, since Q is a path.

Now, by Corollary 3.31 applied to Q , we also have $\tau(p_1) = \tau(a) < \tau(p_2) = \tau(b) < \dots < \tau(p_{l-1}) < \tau(p_l)$.

We now claim that for each $1 \leq i \leq l-2$, $v - p_i \notin \mathcal{C}$. To see this, suppose, if possible, that v is adjacent to p_i for some $1 \leq i \leq l-2$. Then, since $\tau(p_i) < \tau(x)$, $\tau(x) < \tau(v)$ (by the hypothesis for Case 2), and since v is adjacent to both x and p_i , Observation 2.14 would imply that $x - p_i \in \mathcal{C}$. But this would be a chord in the chordless path Q , thereby leading to a contradiction. We conclude that for each $1 \leq i \leq l-2$, $v - p_i \notin \mathcal{C}$. This implies that $(p_1 = a, p_2 = b, \dots, p_{l-1} = u, v)$ is a chordless path from (a, b) to (u, v) with $(a, b) \neq (u, v)$ (since $a \neq u$).

This further implies that in this case, $u \rightarrow v$ adheres to item 2 of Observation 3.40.

- (c) Suppose that $u \rightarrow x \in M$ because there exists an edge $a - b \in \mathcal{C}$ such that $a \rightarrow b \in O$, and there exists a chordless path $Q_1 = (p_1 = a, p_2 = b, \dots, p_l = x)$ from (a, b) to x in \mathcal{C} (with $b \neq x$), and a chordless path $Q_2 = (q_1 = x, q_2 = u, \dots, q_m = a)$ from (x, u) to a in \mathcal{C} (with $u \neq a$). By Corollary 3.31 applied to Q_1 , we have $\tau(p_1) = \tau(a) < \tau(p_2) < \dots < \tau(p_{l-1}) < \tau(p_l) = \tau(x)$. Since $\tau(q_m) = \tau(a) < \tau(x) = \tau(q_1)$, Observation 2.15 applied to the LBFS ordering τ and the chordless path Q_2 of the undirected chordal graph \mathcal{C} then also implies that we must have $\tau(q_i) < \tau(x)$ for every $2 \leq i \leq m$. By the hypothesis of Case 2, $\tau(v) < \tau(x)$. Therefore, for all i , $\tau(q_i) < \tau(v)$. We now claim that

$$\text{For } 3 \leq i \leq m, v - q_i \notin \mathcal{C}. \quad (6)$$

To see this, suppose, if possible, that for some $3 \leq i \leq m$, $v - q_i \in \mathcal{C}$. As argued above, we have $\tau(q_i) < \tau(v)$ and $\tau(q_1) < \tau(v)$. Therefore, from Observation 2.14, $q_1 - q_i \in \mathcal{C}$ (as both q_1 and q_i are adjacent to v). Since $3 \leq i \leq m$, this constitutes a chord of Q_2 and thus contradicts that Q_2 is a chordless path. This establishes eq. (6). Due to eq. (6), we have that $Q_3 = (v, q_2 = u, q_3, \dots, q_m = a)$ (which we get by replacing q_1 with v in Q_2) is a chordless path in \mathcal{C} .

We now show the existence of a chordless path from (a, b) to v in \mathcal{C} . Since Q_1 is a chordless path, and $a \rightarrow b \in O$, from Corollary 3.31, $\tau(p_1 = a) < \tau(p_2 = b) < \dots < \tau(p_l = x)$. Suppose for some $i \leq l-2$, $v - p_i \in \mathcal{C}$. Then, from Observation 2.14, $p_i - p_l \in \mathcal{C}$, as $\tau(p_i) < \tau(p_l = x) < \tau(v)$ (from our hypothesis of this case, $\tau(x) < \tau(v)$), which further implies that Q_1 is not a chordless path, which is a contradiction. This implies that for all $i \leq l-2$, $v - p_i \notin \mathcal{C}$. Thus, we get that either $Q_4 := (p_1 = a, p_2 = b, \dots, p_{l-1}, v)$ is a chordless path from (a, b) to v (if $p_{l-1} - v \in \mathcal{C}$), or else $Q_4 = (p_1 = a, p_2 = b, \dots, p_l, v)$ is a chordless path from (a, b) to v (if $p_{l-1} - v \notin \mathcal{C}$). Note that in either case, we also have $b \neq v$ since $\tau(b) = \tau(p_2) < \tau(x)$ and also $\tau(x) < \tau(v)$ (by the hypothesis for Case 2).

Existence of the chordless paths Q_3 and Q_4 implies that $u \rightarrow v$ obeys item 3 of Observation 3.40.

This concludes the second case.

We have thus shown that in all cases $u \rightarrow v$ obeys Observation 3.40. This completes the induction, and hence also the proof of Observation 3.40. \square

Observation 3.40 implies the following corollary.

Corollary 3.42. *For $i \in \{1, 2\}$, let \mathcal{C} be an undirected connected component of M_i . Let τ be the LBFS ordering returned at line 5 of alg. 4 in the iteration of the second foreach loop (lines 4-18 of alg. 4) when \mathcal{C} is considered. For an undirected edge $u - v$ in \mathcal{C} , if $u \rightarrow v \in M$ then either $u \rightarrow v$ is added to M in the initialization step (line 1 of alg. 4) or in the second foreach loop (lines 4-18 of alg. 4) when \mathcal{C} is considered, in the iteration of the first while loop (lines 8-17 of alg. 4) when $a = \tau(v)$ and in the iteration of the second while loop (lines 10-15 of alg. 4) when $b = \tau(u)$.*

Proof. Without loss of generality, let's assume $i = 1$, i.e., \mathcal{C} is an undirected connected component of M_1 . From the construction of M , it's evident that if $u \rightarrow v \in M$, then it's either added at the initialization step (line 1 of alg. 4), or $u - v$ is in some undirected connected component \mathcal{C}' of M_1 or M_2 , and it gets directed at line 12 when \mathcal{C}' is considered in the second foreach loop (lines 4-18 of alg. 4).

If $u \rightarrow v$ is added at the initialization step, then we are done. Suppose $u \rightarrow v$ is not added at the initialization step; we'll now show that $\mathcal{C} = \mathcal{C}'$.

Suppose $\mathcal{C} \neq \mathcal{C}'$. Then, \mathcal{C}' must be an undirected connected component of M_2 , as two different undirected connected component of a graph cannot share a node, and both \mathcal{C} and \mathcal{C}' contain $u - v$. This implies $u, v \in V_{M_1} \cap V_{M_2} = I \subseteq V_O$.

Since $u - v \in \mathcal{C}'$ and $u \rightarrow v \in M$, $u \rightarrow v$ must obey Observation 3.40. We now show that this implies $u \rightarrow v \in O$.

If item 1 of Observation 3.40 holds, then we have $u \rightarrow v \in O$. If one of item 2 or item 3 of Observation 3.40 holds, then in the undirected connected chordal component \mathcal{C}' of M_2 , there exists an edge $x - y \in \mathcal{C}'$ such that $x \rightarrow y \in O$, and either (a) $(x, y) \neq (u, v)$ and there exists a chordless path from (x, y) to (u, v) in \mathcal{C}' (due to item 2 of Observation 3.40), or (b) $y \neq v$, $u \neq x$, and there exist chordless paths Q_1 and Q_2 in \mathcal{C}' such that Q_1 is a chordless path from (x, y) to v , and Q_2 is a chordless path from (v, u) to x (due to item 3 of Observation 3.40).

Since \mathcal{C}' is an undirected connected chordal component of M_2 (from item 2 of Theorem 2.18), a chordless path in \mathcal{C}' is a chordless path of M_2 . And, from Observation 2.24, every chordless path in \mathcal{C}' is a triangle-free path. This implies chordless paths in \mathcal{C}' are triangle-free paths of M_2 . Also, since $(x, y) \in \mathcal{C}'$, an undirected connected component of M_2 , and $x \rightarrow y \in O$, this implies $x, y \in V_{M_2} \cap V_O = V_{O_2}$.

Since (O_2, P_{21}, P_{22}) is the shadow of M_2 on V_{O_2} , from Definition 2.36, O_2 is an induced subgraph of M_2 . This implies $(x, y) \in E_{O_2}$, and if there exists a chordless path in \mathcal{C}' , from (x, y) to (u, v) , then $P_{21}((x, y), (u, v)) = 1$, and if there exist chordless paths in \mathcal{C}' , from (x, y) to v and from (v, u) to x then $P_{22}((x, y), v) = P_{22}((v, u), x) = 1$.

Since $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, therefore, in both possibilities, from item 3 of Definition 3.24, we again have $u \rightarrow v \in O$.

But, if $u \rightarrow v \in O$, then $u \rightarrow v$ has been added to M at the initialization step, contradicting our assumption. This implies $\mathcal{C} = \mathcal{C}'$. This further implies $u \rightarrow v$ is added to M at the iteration of the second foreach loop (lines 3–19) when \mathcal{C} is considered. It is obvious from the algorithm that in that iteration, $u \rightarrow v$ is added to M when $a = \tau(v)$ and $b = \tau(u)$. This completes the proof of Corollary 3.42. \square

The following corollary is implied by Corollaries 3.31 and 3.32 when applied to Observation 3.40.

Corollary 3.43. *For $i \in \{1, 2\}$, and any undirected connected component \mathcal{C} of M_i at line-5 of alg. 4, let τ be the LBFS ordering returned by alg. 3. Then, for $u - v \in \mathcal{C}$, at any iteration of the while loop (lines 8-17) of alg. 4, if $u \rightarrow v \in M$, then either of the following occurs:*

1. $u \rightarrow v$ is strongly protected in O , and $\tau(u) < \tau(v)$, or
2. $\exists a \rightarrow b \in O$ such that there exists a chordless path $Q = (p_1 = a, p_2 = b, \dots, p_{l-1} = u, p_l = v)$ from (a, b) to (u, v) in \mathcal{C} such that $(a, b) \neq (u, v)$ and $\tau(p_1) < \tau(p_2) < \dots < \tau(p_{l-1}) < \tau(p_l)$. In particular, $\tau(u) < \tau(v)$.
3. $\exists a \rightarrow b \in O$ such that there exists a chordless paths Q_1 and Q_2 such that $Q_1 = (p_1 = a, p_2 = b, \dots, p_{l-1}, p_l = v)$ is a chordless path from (a, b) to v in \mathcal{C} and Q_2 is a chordless path from (v, u) to a in \mathcal{C} such that $b \neq v$ and $u \neq a$, and $\tau(p_1 = a) < \tau(p_2 = b) < \dots < \tau(p_{l-1}) < \tau(p_l = v)$, and $\tau(z) < \tau(v)$ for every node $z \neq v$ in Q_2 (in particular, $\tau(u) < \tau(v)$).

Proof. We prove the relative ordering of nodes (mentioned in items 1 to 3 of Corollary 3.43) at the end of the proof. We first prove the remaining parts of the corollary. Let for $i \in \{1, 2\}$, \mathcal{C} be an undirected connected component of M_i , and $u - v \in \mathcal{C}$. From Observation 3.40, if $u \rightarrow v \in M$ then either (a) $u \rightarrow v \in O$ (item 1 of Observation 3.40), or (b) there exists an edge $a - b \in \mathcal{C}$ such that there exists a chordless path from (a, b) to (u, v) in \mathcal{C} (item 2 of Observation 3.40), or (c) there exists an edge $a - b \in \mathcal{C}$ such that there exist chordless paths in \mathcal{C} from (a, b) to v and from (v, u) to a (item 3 of Observation 3.40).

Suppose $u - v \in \mathcal{C}$, and $u \rightarrow v \in M$ because it obeys item 1 of Observation 3.40, i.e., $u \rightarrow v \in O$. This implies $u, v \in V_{M_i} \cap V_O = I \subseteq V_{O_i}$. As (O_i, P_{i1}, P_{i2}) is the shadow of M_i on V_{O_i} , therefore, from

item 1 of Definition 2.36, O_i is an induced subgraph of M_i . Since $u, v \in V_{O_i}$ and $u - v \in \mathcal{C}$, an undirected connected component of M_i , therefore, $u - v \in O_i$. Since $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, therefore, O obeys Definition 3.24. Item 3 of Definition 3.24 implies that if $u \rightarrow v \in O$ then either (i) $u \rightarrow v$ is strongly protected in O , or (ii) there exists an edge $a - b \in O_i$ such that $P_{i1}((a, b), (u, v)) = 1$, or (iii) there exists an edge $a - b \in O_i$ such that $P_{i2}((a, b), v) = P_{i2}((v, u), a) = 1$.

If $u \rightarrow v$ is strongly protected in O then it obeys item 1 of Corollary 3.43.

Suppose $u \rightarrow v \in O$ and there exists an edge $a - b \in O_i$ such that $P_{i1}((a, b), (u, v)) = 1$. We first show that $a - b$ and $u - v$ are in the same undirected connected component \mathcal{C} of M_i . Since $P_{i1}((a, b), (u, v)) = 1$ this implies $(a, b) \neq (u, v)$ and there exists a triangle-free path $Q = (q_1 = a, q_2 = b, \dots, q_{l-1} = u, q_l = v)$ from (a, b) to (u, v) in M_i . We claim that all the edges of Q are undirected. If this is not true then pick a directed edge $u_j \rightarrow u_{j+1}$ of Q such that j is maximum. Since $u - v$ is an undirected edge in M_i , therefore, $j < l - 1$. But, then $u_j \rightarrow u_{j+1} - u_{j+2}$ is an induced subgraph of M_i , contradicting item 3 of Theorem 2.18 (remember M_i is an MEC). This validates our claim that all the edges of Q are undirected. This implies Q is a triangle-free path in the undirected connected component \mathcal{C} of M_i . From item 2 of Theorem 2.18, \mathcal{C} is a chordal graph. Therefore, from Proposition 2.25, Q is a chordless path in \mathcal{C} . This implies $u \rightarrow v$ obeys item 2 of Corollary 3.43.

Suppose $u \rightarrow v \in O$ because there exists an edge $a - b \in O_i$ such that $P_{i2}((a, b), v) = P_{i2}((v, u), a) = 1$. This implies there exist triangle-free paths Q_1 and Q_2 in M_i such that Q_1 is a triangle-free path from (a, b) to v , and Q_2 is a triangle-free path from (v, u) to a . All the edges in Q_1 and Q_2 must be undirected, otherwise, by concatenating Q_1 and Q_2 we get a directed cycle in M_i , contradicting item 1 of Theorem 2.18, as M_i is an MEC. This implies Q_1 and Q_2 are triangle-free paths in \mathcal{C} , an undirected connected component of M_i . From item 2 of Theorem 2.18, \mathcal{C} is a chordal graph. Therefore, from Proposition 2.25, Q_1 and Q_2 are chordless paths in \mathcal{C} . This implies $u \rightarrow v$ obeys item 3 of Corollary 3.43.

Suppose $u \rightarrow v \in M$ because it obeys item 2 of Observation 3.40. Then, $u \rightarrow v$ obeys item 2 of Corollary 3.43. Similarly, if $u \rightarrow v \in M$ because it obeys item 3 of Observation 3.40. Then, $u \rightarrow v$ obeys item 3 of Corollary 3.43.

The ranking of the nodes comes from Observation 2.15 and Corollary 3.32. If $u \rightarrow v \in O$ then from Corollary 3.32, $\tau(u) < \tau(v)$. In the chordless path of item 2 of Corollary 3.43, $a \rightarrow b \in O$. From Corollary 3.32, $\tau(a) < \tau(b)$. Then, from Observation 2.15, the nodes of the chordless path follows the order: $\tau(p_1 = a) < \tau(p_2 = b) < \dots < \tau(p_{l-1}) < \tau(p_l)$. Similarly, the nodes in the chordless path P_1 of item 3 of Corollary 3.43 follow the order: $\tau(p_1 = a) < \tau(p_2 = b) < \dots < \tau(p_{l-1}) < \tau(p_l = v)$. In particular, this also implies $\tau(a) < \tau(v)$. Thus, Observation 2.15 implies that for all the nodes $z \neq v$ of the chordless path P_2 of item 3 of Corollary 3.43, we have $\tau(z) < \tau(v)$. This completes the proof of Corollary 3.43. \square

We are now ready to prove Lemma 3.35.

Proof of Lemma 3.35. We initialize M as $U_M(M_1, M_2, O)$ (line 1 of alg. 4). This implies that after the initialization step, if $u \rightarrow v \in O$ then $u \rightarrow v \in M$. Throughout the algorithm, we only convert an undirected edge into a directed one (lines 11-13). This implies that if $u \rightarrow v \in O$ then $u \rightarrow v$ belongs to the graph M returned by the algorithm. To complete the proof, we have to show that even for any undirected edge $u - v \in O$, M returned by alg. 4 has $u - v$.

Suppose, if possible, that there is an edge $u - v$ of O that is directed as $u \rightarrow v$ in M . If $u - v \in O$ then $u - v \in \text{skeleton}(O)$. From the construction, $\text{skeleton}(O) = \text{skeleton}(O_1) \cup \text{skeleton}(O_2)$. This implies either $u - v \in \text{skeleton}(O_1)$ or $u - v \in \text{skeleton}(O_2)$ or both.

Without loss of generality, let us assume that $u - v \in \text{skeleton}(O_1)$. Since $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$ and $u - v \in O$, from item 1 of Definition 3.24, $u - v \in O_1$. This implies $u - v \in M_1$ as O_1 is an induced subgraph of M_1 (from item 1 of Definition 2.36, as (O_1, P_{11}, P_{12}) is the shadow of M_1 on V_{O_1}).

This further implies there exists an undirected connected component \mathcal{C} of M_1 that contains $u - v$. Since from our assumption, $u \rightarrow v \in M$ and $u \rightarrow v \notin O$, therefore, from Observation 3.40, either (a) there exists $a - b \in \mathcal{C}$ such that $a \rightarrow b \in O$ and there exists a chordless path from (a, b) to (u, v) in \mathcal{C} , or (b) there exists $a - b \in \mathcal{C}$ such that there exists chordless paths in \mathcal{C} from (a, b) to v and from (v, u) to a .

In both of the possibilities, $a \rightarrow b \in O$ and $a - b \in M_1$ (remember \mathcal{C} is an undirected connected component of M_1). This implies $a, b \in V_{M_1} \cap V_O = I \subseteq V_{O_1}$. This further implies $a - b \in O_1$ as O_1 is an induced subgraph of M_1 . In the first possibility, since there is a chordless path from (a, b) to (u, v) in \mathcal{C} , therefore, there is a chordless path from (a, b) to (u, v) in M_1 .

From Observation 2.24, this further implies that there is a triangle-free path from (a, b) to (u, v) in M_1 . That means $P_{11}((a, b), (u, v)) = 1$. But, then from item 3 of Definition 3.24, $u \rightarrow v \in O$, a contradiction. Similarly, in the second possibility, we have $a - b \in O_1$ and there exist triangle-free paths from (a, b) to v and from (v, u) to a in M_1 .

This implies $P_{22}((a, b), v) = P_{22}((v, u), a) = 1$. But, then from item 3 of Definition 3.24, $u \rightarrow v \in O$, a contradiction. This implies that for any undirected edge $u - v \in O$, M returned by alg. 4 has $u - v$. The above discussion implies that $M[V_O] = O$. \square

The following observations can be easily derived from Corollary 3.43. These observations will be used later, and if preferred, the reader can skip ahead to Lemma 3.47 directly.

Observation 3.44. *For $a \in \{1, 2\}$, and for any undirected connected component \mathcal{C} of M_a , let τ be the LBFS ordering obtained from alg. 3 when it is called at line 5 of alg. 4. Then, for $u - v \in \mathcal{C}$, if $u \rightarrow v \in O$ then $u \rightarrow v \in M$, and $\tau(u) < \tau(v)$.*

Proof. At the initialization step, line 1, of alg. 4, if $u \rightarrow v \in O$, then $u \rightarrow v \in M$. From item 1 of Corollary 3.43, it follows that $\tau(u) < \tau(v)$. \square

Observation 3.45. *For $a \in \{1, 2\}$, and for any undirected connected component \mathcal{C} of M_a , let τ be the LBFS ordering returned by alg. 3 when it is called at line 5 of alg. 4. Then, for $u - v \in \mathcal{C}$, if $u \rightarrow v \in M$, then $\tau(u) < \tau(v)$.*

Proof. Without loss of generality, let us assume $a = 1$. For $u - v \in \mathcal{C}$, if $u \rightarrow v \in M$, then either it got directed at the initialization step (line 1 of alg. 4), or it gets directed at lines 11-13 of alg. 4.

At the initialization step, $M = U_M(M_1, M_2, O)$. Therefore, if $u \rightarrow v \in M$ and $u \rightarrow v \notin M_1$, then either $u \rightarrow v \in M_2$ or $u \rightarrow v \in O$. If $u \rightarrow v \in M_2$, then $u, v \in V_{M_1} \cap V_{M_2} = I \cup V_{O_2}$. Since O_2 is an induced subgraph of M_2 , therefore $u \rightarrow v \in O_2$.

Since O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, from Definition 3.24, $u \rightarrow v \in O$. This further implies that if $u \rightarrow v \in M$, then either $u \rightarrow v \in O$ or it gets directed during the run of lines 11-13 of alg. 4.

If $u \rightarrow v \in O$, then from Observation 3.44, $\tau(u) < \tau(v)$. And, if $u \rightarrow v$ is directed during the run of lines 11-13 of alg. 4, then from items 2 and 3 of Corollary 3.43, $\tau(u) < \tau(v)$. \square

Observation 3.46. *For $a \in \{1, 2\}$, and for any undirected connected component \mathcal{C} of M_a , let τ be the LBFS ordering obtained from alg. 3. Then, for $u - v \in \mathcal{C}$, if there exists $a - b \in \mathcal{C}$ such that either $a \rightarrow b \in M$ or $a \rightarrow b \in O$, and there exists a chordless path $Q = (u_1 = a, u_2 = b, \dots, u_{l-1} = u, u_l = v)$ from (a, b) to (u, v) in \mathcal{C} then $\tau(u_1) < \tau(u_2) < \dots < \tau(u_l)$, and for all $1 \leq i \leq l - 1$, $u_i \rightarrow u_{i+1} \in M$. In particular, $u \rightarrow v \in M$.*

Proof. Note that when $a \rightarrow b \in O$, from the initialization step (line 1) of alg. 3, we have $a \rightarrow b \in M$. Thus, we can assume that $a \rightarrow b \in M$.

Now, since $Q = (u_1 = a, u_2 = b, \dots, u_{l-1} = u, u_l = v)$ is a chordless path from (a, b) to (u, v) in \mathcal{C} , and $a \rightarrow b \in M$, Observation 3.45 implies that $\tau(u_1 = a) < \tau(u_2 = b)$. Since Q is a chordless path in the undirected chordal graph \mathcal{C} and τ is an LBFS ordering of \mathcal{C} , Observation 2.15 implies that we must then have $\tau(u_1) < \tau(u_2) < \dots < \tau(u_{l-1}) < \tau(u_l)$.

Since $u_1 = a$ and $u_2 = b$, we have $u_1 \rightarrow u_2 \in M$. Lines 8-17 of alg. 4 show that for $1 < i < l$, in the iteration of the while loop at line 8 when $a = i + 1$, before the end of the iteration of the while loop at line 10 when $b = i$, we get $u_i \rightarrow u_{i+1} \in M$. Because, if at the initialization step, we do not have $u_i \rightarrow u_{i+1}$ in M then due to the existence of the induced subgraph $u_{i-1} \rightarrow u_i - u_{i+1}$ in M , the if condition at line 11 is satisfied and we get $u_i \rightarrow u_{i+1} \in M$ at line 12. Thus, we conclude that for all $1 \leq i \leq l - 1$, $u_i \rightarrow u_{i+1} \in M$. \square

We now prove Lemma 3.36.

Proof of Lemma 3.36. We first show that M is an MEC. Then we show that $\text{skeleton}(M) = \text{skeleton}(H)$. Together these two imply that M is an MEC of H . Theorem 2.18 provides necessary and sufficient conditions for an MEC. Lemmas 3.47 to 3.50 validate that M obeys the required conditions mentioned in Theorem 2.18 to be an MEC. This further implies that M is an MEC.

Lemma 3.47. M is a chain graph.

Lemma 3.48. Every undirected connected component of M is chordal.

Lemma 3.49. M does not have an induced subgraph of the form $a \rightarrow b - c$.

Lemma 3.50. Every directed edge $u \rightarrow v \in M$ is strongly protected in M .

Proofs of Lemmas 3.47 to 3.50 are given below.

We now prove that $\text{skeleton}(M) = H$. From the initialization step (line 1 of alg. 4), $M = U_M(M_1, M_2, O)$. The remaining lines of alg. 4 orient the undirected edges of M and do not change the skeleton of M . Therefore, $\text{skeleton}(M) = \text{skeleton}(M_1) \cup \text{skeleton}(M_2) \cup \text{skeleton}(O)$.

From the construction, $\text{skeleton}(M_1) = \text{skeleton}(H_1)$, $\text{skeleton}(M_2) = \text{skeleton}(H_2)$, and $\text{skeleton}(H) = \text{skeleton}(H_1) \cup \text{skeleton}(H_2) \cup \text{skeleton}(O)$. This implies $\text{skeleton}(M) = \text{skeleton}(H)$. Thus, M is an MEC of H , as it is an MEC with H as its skeleton. \square

We will now prove Lemmas 3.47 to 3.50. If preferred, readers may choose to bypass the proofs and proceed directly to Lemma 3.37, where we demonstrate that M_1 and M_2 are projections of M .

First, we present the proof of Lemma 3.47. The proof of Lemma 3.48 can be found on page 60, the proof of Lemma 3.49 is on page 61, and the proof of Lemma 3.50 is on page 61.

Proof of Lemma 3.47. The following observations, to be proved later, imply that M is a chain graph.

Observation 3.51. For $a \in \{1, 2\}$, $M[V_{H_a}]$ is a chain graph.

Observation 3.52. M does not have any chordless cycle of length greater than 3.

Suppose M is not a chain graph. Then M has a directed cycle. Choose a cycle $C = (u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ of the smallest length. l must be greater than two.

Suppose $l = 3$. We claim that either all the nodes of the cycle are in M_1 (i.e., $u_1, u_2, u_3 \in V_{M_1}$) or all the nodes of the cycle are in M_2 (i.e., $u_1, u_2, u_3 \in V_{M_2}$).

Suppose this is not true. Then there exists a node in the cycle that is in M_1 but not in M_2 , and there exists a node in the cycle that is in M_2 but not in M_1 . Without loss of generality, let us assume that u_1 is in M_1 but not in M_2 (i.e., $u_1 \in V_{M_1} \setminus V_{M_2} = V_{M_1} \setminus I$, where $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H separating $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$). Then u_2 and u_3 , the neighbors of u_1 in C , must be in M_1 (neither u_2 nor u_3 can be in $V_{M_2} \setminus I$ since there is no edge between $V_{M_1} \setminus I$ and $V_{M_2} \setminus I$, as they are separated by I). This validates our claim.

From Observation 3.51, there cannot be a directed cycle in $M[V_{M_1}]$ or $M[V_{M_2}]$. Therefore, C must have a length greater than three.

Suppose $l > 3$. Since we have chosen a cycle of the smallest length, C must be a chordless cycle. If C had a chord (an edge between two non-adjacent nodes of the cycle), we would obtain a directed cycle of smaller length in M , regardless of whether the chord is directed or undirected. But from Observation 3.52, M cannot have any chordless cycle. This implies this case cannot occur.

Thus, we have shown that M does not have any directed cycle, i.e., M is a chain graph. This completes the proof of Lemma 3.47. \square

Proof of Observation 3.51. If $M[V_{H_a}]$ is not a chain graph, then there exists a directed cycle $C = (u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ in M such that each node of the cycle is in V_{H_a} .

From the construction of M , for $p, q \in V_{H_a}$, if there is an edge $p \rightarrow q$ in M , then either $p \rightarrow q \in M_a$ or $p - q \in M_a$, and if $p - q \in M$, then $p - q \in M_a$. This implies C is also a cycle in M_a . This further implies

that none of the edges in C are directed in M_a , otherwise M_a contains a directed cycle, contradicting item 1 of Theorem 2.18.

This implies all the edges of the cycle are undirected edges in M_a , and all the nodes of the cycle are part of the same undirected connected component \mathcal{C} of M_a . This further implies that for any undirected connected component \mathcal{C} of M_a , $M[V_{\mathcal{C}}]$ contains a directed cycle. From Observation 3.53, this is not possible.

Observation 3.53. *For $a \in \{1, 2\}$, for any undirected connected component \mathcal{C} of M_a , there does not exist a directed cycle in $M[V_{\mathcal{C}}]$.*

Proof of Observation 3.53 is given below. This completes the proof of Observation 3.51. \square

Proof of Observation 3.53. Suppose for some $a \in \{1, 2\}$, for some undirected connected component \mathcal{C} of M_a , there exist a directed cycle in $M[V_{\mathcal{C}}]$. Pick a directed cycle $C = (u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ in $M[V_{\mathcal{C}}]$ of the smallest length. We first show that l cannot be greater than three.

Suppose $l > 3$. Then, C must be a chordless cycle, because if C has a chord then whatever the direction of the chord, we get a directed cycle of smaller length. From the construction of M , $\text{skeleton}(M[V_{\mathcal{C}}]) = \text{skeleton}(M_a[V_{\mathcal{C}}]) = \mathcal{C}$. This further implies C is a chordless cycle in \mathcal{C} . But, since \mathcal{C} is a chordal graph (from item 2 of Theorem 2.18, as \mathcal{C} is an undirected connected component of MEC M_a), it cannot contain any chordless cycle, a contradiction. Therefore, if there exists any directed cycle in $M[V_{\mathcal{C}}]$ then the shortest length directed cycle must contain only three vertices.

There are three possibilities: either all the edges of the cycle are directed, or at most two edges of the cycle are directed, or only one edge of the cycle is directed.

Suppose there exist vertices $x, y, z \in \mathcal{C}$ such that $x - y - z - x \in \mathcal{C}$ and $x \rightarrow y \rightarrow z \rightarrow x \in M$, i.e., all edges of the cycle are directed. Let τ be the LBFS ordering of \mathcal{C} we get at line 5 of alg. 4. From Observation 3.45, for $u - v \in V_{\mathcal{C}}$, if $u \rightarrow v \in M$ then $\tau(u) < \tau(v)$. This implies that $x \rightarrow y \rightarrow z \rightarrow x$ cannot occur in M , otherwise, we have $\tau(x) < \tau(y) < \tau(z) < \tau(x)$, a contradiction.

Suppose there exist vertices $x, y, z \in \mathcal{C}$ such that $x - y - z - x \in \mathcal{C}$ and $x \rightarrow y \rightarrow z - x \in M$ (two edges of the cycle are directed). Then from Observation 3.45, $\tau(x) < \tau(y) < \tau(z)$. And, while processing the edge $x - z$ on lines 11-13 of alg. 4, we replace $x - z$ with $x \rightarrow z$. This implies that $x \rightarrow y \rightarrow z - x$ cannot occur in M . We thus have the following: if vertices x, y, z in \mathcal{C} form a directed cycle C in M , then C has exactly one directed edge. The only possibility that remains therefore is of a cycle of the form $x \rightarrow y - z - x$. Observation 3.54 shows that even this possibility cannot occur.

Observation 3.54. *For $a \in \{1, 2\}$, let \mathcal{C} be an undirected connected component of M_a , and M be the graph returned by alg. 4 for input M_1, M_2 and O . There does not exist vertices $x, y, z \in \mathcal{C}$ such that $x - y - z - x \in \mathcal{C}$ and $x \rightarrow y - z - x \in M$.*

This completed the proof of Observation 3.53. \square

Proof of Observation 3.54. Let τ be the LBFS ordering returned by alg. 3 at line 5 of alg. 4. From Observation 3.45, for $p - q \in V_{\mathcal{C}}$, if $p \rightarrow q \in M$, then $\tau(p) < \tau(q)$. We proceed by structuring the analysis based on the relative ordering of x, y , and z in τ . Claim 3.55 lists all such potential orderings and shows that none of them are allowed, thus completing the proof of Observation 3.54.

Claim 3.55. *There do not exist $u, v, w \in \mathcal{C}$ such that $\tau(u) < \tau(v) < \tau(w)$, $u - v - w - u \in \mathcal{C}$ and either $u \rightarrow v - w - u \in M$ or $v \rightarrow w - u - v \in M$ or $u \rightarrow w - v - u \in M$.*

We establish the validity of Claim 3.55 using induction on $\tau(w)$.

Base Case ($\tau(w) = 2$): This case is vacuously true, as any node in τ must have a rank greater than or equal to 1. Therefore, there are no nodes u and v satisfying $1 \leq \tau(u) < \tau(v) < \tau(w) = 2$.

Inductive Hypothesis: Suppose Claim 3.55 is true for $\tau(w) = k$, where $k \geq 2$.

Inductive Step: We now demonstrate that Claim 3.55 holds for $\tau(w) = k + 1$.

Suppose, for the sake of argument, that there exist $u, v, w \in \mathcal{C}$ such that $\tau(u) < \tau(v) < \tau(w) = k + 1$, and either $u \rightarrow v - w - u \in M$, or $u - v \rightarrow w - u \in M$, or $u - v - w \leftarrow u \in M$.

We select u to be the node with the maximum possible rank, i.e., for any u' with $\tau(u) < \tau(u')$, there does not exist any v satisfying $\tau(u') < \tau(v) < \tau(w) = k + 1$ and either $u' \rightarrow v - w - u \in M$, or $u' - v \rightarrow w - u \in M$, or $u' - v - w \leftarrow u \in M$ hold. We then select v to be the node with the minimum possible rank, i.e., for any v' with $\tau(u) < \tau(v') < \tau(v) < \tau(w) = k + 1$, neither $u \rightarrow v' - w - u \in M$, nor $u - v' \rightarrow w - u \in M$, nor $u - v' - w \leftarrow u \in M$ hold.

The following claims show that neither $u \rightarrow v - w - u \in M$, nor $u - v \rightarrow w - u \in M$, nor $u - v - w \leftarrow u \in M$ is satisfied:

Claim 3.56. $u \rightarrow v - w - u \notin M$.

Claim 3.57. $u - v \rightarrow w - u \notin M$.

Claim 3.58. $u - v - w \leftarrow u \notin M$.

Proofs of Claims 3.56 to 3.58 are given below. These claims validate Claim 3.55. \square

Proof of Claim 3.56. Suppose $u \rightarrow v - w - u \in M$. From Corollary 3.43, either of the following has occurred:

1. $u \rightarrow v$ is strongly protected in O .
2. There exists $a \rightarrow b \in O$ such that $(a, b) \neq (u, v)$ and there exists a chordless path from (a, b) to (u, v) in \mathcal{C} .
3. There exists $a \rightarrow b \in O$ such that $b \neq v$, $u \neq a$ and there exist chordless paths in \mathcal{C} from (a, b) to v and from (v, u) to a .

We go through each possibility.

1. Assume that $u \rightarrow v$ is strongly protected in O . Consequently, $u \rightarrow v$ is part of one of the induced subgraphs of O , as depicted in fig. 2.
 - (a) Consider the scenario where $u \rightarrow v$ is strongly protected due to its presence in an induced subgraph $p \rightarrow u \rightarrow v \in O$ (similar to Figure 2.a). Keep in mind that u, v , and w are members of \mathcal{C} , an undirected connected component of M_a for some $a \in \{1, 2\}$. We need to examine whether p belongs to M_a or not.

Suppose $p \notin M_a$. Without loss of generality, let's assume $a = 1$. From the construction, if $p \notin V_{M_1}$, then u , a neighbor of p in M , belongs to $I = V_{M_1} \cap V_{M_2}$, as $u \in V_{M_1}$ (recall that u belongs to \mathcal{C} , an undirected connected component of M_a), and I is a vertex separator of H that separates $V_{M_1} \setminus I$ and $V_{M_2} \setminus I$. However, this implies $u, v, w \in I \cup N(I, H) \subseteq V_O$. Consequently, $u \rightarrow v - w - u \in O$, as from Lemma 3.35, $M[V_O] = O$. Nevertheless, this contradicts item 1 of Definition 2.21, as O is a partial MEC. Thus, the only remaining possibility is $p \in M_a$.

Suppose $p \in M_a$. Since $u, p \in V_{M_a}$ and $p \rightarrow u \in M$, from the construction of M , either $p \rightarrow u \in M_a$ or $p - u \in M_a$. If $p \rightarrow u \in M_a$, then $p \rightarrow u - v$ is an induced subgraph in M_a , a contradiction from item 2 of Theorem 2.18. This implies $p - u \in M$, which means $p \in \mathcal{C}$.

From Observation 3.44, $\tau(p) < \tau(u)$. This further implies $\tau(p) < \tau(u) < \tau(v) < \tau(w)$ (from our hypothesis $\tau(u) < \tau(v) < \tau(w)$). Since $p, v \in \mathcal{C}$, and there is no edge between p and v in M , from the construction of M , there must be no edge between p and v in \mathcal{C} . This also implies $p - w \notin \mathcal{C}$; otherwise, since $\tau(v) < \tau(w)$, $\tau(p) < \tau(w)$, and $v - w \in \mathcal{C}$, from Observation 2.14, $p - v \in \mathcal{C}$, leading to a contradiction. This implies that $p \rightarrow u - w \in M$. However, this is not possible as while running lines 11-13 of alg. 4, we get $u \rightarrow w \in M$. Hence, this case does not occur.

Considering the above analysis, it is evident that neither $p \in M_a$ nor $p \notin M_a$ holds true. Consequently, this case cannot occur. We will now proceed to explore other possibilities.

(b) Suppose $u \rightarrow v$ is strongly protected because it is part of an induced subgraph $u \rightarrow v \leftarrow p \in O$ (same as Figure 2.b). Similar to the above case, the only possibility we need to consider is when $p \in \mathcal{C}$. Since $u, v, p \in V_O$ as well as in \mathcal{C} , an undirected connected component of M_a , therefore, $u, v, w \in O_a = V_O \cap V_{M_a}$. Since $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, therefore, from item 2 of Definition 3.24, the v-structure $u \rightarrow v \leftarrow p$ also belongs to O_a . Since (O_a, P_{a1}, P_{a2}) is a shadow of M_a on V_{O_a} , therefore, from item 1 of Definition 2.36, O_a is an induced subgraph of M_a . This further implies $u \rightarrow v \leftarrow p \in M_a$. But, this is a contradiction as $u - v \in M_a$ (since \mathcal{C} is an undirected connected component of M_a , if $u - v \in \mathcal{C}$ then $u - v \in M_a$). This contradiction implies that this case also does not occur.

(c) Suppose $u \rightarrow v$ is strongly protected in O because it is part of an induced subgraph $u \rightarrow p \rightarrow v \leftarrow u \in O$ (same as Figure 2.c). Similar to the above Case 1.a, the only possibility we need to consider is when $p \in \mathcal{C}$. Then from Observation 3.45, $\tau(u) < \tau(p) < \tau(v)$. Combining this with our induction hypothesis, we get $\tau(u) < \tau(p) < \tau(v) < \tau(w)$. There are two possibilities: either $p - w \notin \mathcal{C}$, or $p - w \in \mathcal{C}$.

If $p - w \notin \mathcal{C}$ then while running lines 11-13 of alg. 4, we get $v \rightarrow w \in M$ (because $\tau(p) < \tau(v) < \tau(w)$ and $p \rightarrow v - w$ is an induced subgraph in M). This means that this case does not occur as $v - w \in M$.

Suppose then that $p - w \in \mathcal{C}$. Then there are three possibilities: either $p - w \in M$, or $p \leftarrow w \in M$, or $p \rightarrow w \in M$. We will show that none of the three can occur.

- i. If $p - w \in M$, then we would have a triplet (p, v, w) that satisfies the conditions $\tau(u) < \tau(p) < \tau(v) < \tau(w)$, and $p \rightarrow v - w - p \in M$. However, this directly contradicts our choice of u as the vertex with the maximal $\tau(u)$ (recall that we have selected u as a node with maximum rank in τ such that there exists a v satisfying $\tau(u) < \tau(v) < \tau(w) = k + 1$ and $u \rightarrow v - w - u \in M$). Hence, it is concluded that $p - w \notin M$.
- ii. Since $\tau(p) < \tau(w)$, therefore, according to Observation 3.45, $w \rightarrow p \notin M$.
- iii. If $p \rightarrow w \in M$ then from lines 11-13 of alg. 4, $u \rightarrow w \in M$ (due to $\tau(u) < \tau(p) < \tau(w)$, and $u \rightarrow p \rightarrow w - u \in M$). However, this leads to a contradiction as $u - w \in M$.

This analysis demonstrates that none of the three possibilities can occur, thereby ruling out the possibility of case $p - w \in \mathcal{C}$. With this, we conclude that this case also does not occur.

(d) Suppose $u \rightarrow v$ is strongly protected in O because it is part of an induced subgraph $u \rightarrow v \leftarrow p - u - p' \rightarrow v$ (same as Figure 2.d). Similar to above case 1.a, the only possibility we need to consider is when $p, p' \in \mathcal{C}$. Similar to Case 1.b, the v-structure $p \rightarrow v \leftarrow p'$ in O implies that the v-structure also belongs to M_a , i.e., $p \rightarrow v \leftarrow p' \in M_a$. But, since \mathcal{C} is an undirected connected component of M_a , and $p, v \in \mathcal{C}$, therefore, from Proposition 2.6, $p \rightarrow v$ cannot be in M_a . This leads to a contradiction. Hence, it can be deduced that this particular case also does not come to pass.

We show that $u \rightarrow v$ is a part of none of the induced subgraphs shown in fig. 2. This implies that this possibility does not occur, i.e., $u \rightarrow v$ is not strongly protected in O . We now move to the other possibilities.

2. Suppose $u \rightarrow v \in O$ because of the existence of $x - y \in \mathcal{C}$ such that $x \rightarrow y \in O$, and there exists a chordless path $Q = (p_1 = x, p_2 = y, \dots, p_{l-1} = u, p_l = v)$ from (x, y) to (u, v) in \mathcal{C} . From Observation 3.46, $p_{l-2} \rightarrow p_{l-1} = p_{l-2} \rightarrow u \in M$. We thus have the induced sub-graph $p \rightarrow u \rightarrow v$ in M with $p = p_{l-2}$. This case is now exactly the same as the first case of the first possibility, and hence as proved there, cannot occur.
3. Suppose there exists $x - y \in O$ such that $x \rightarrow y \in O$, and there exist chordless paths in \mathcal{C} : $Q_1 = (p_1 = x, p_2 = y, \dots, p_l = v)$ from (x, y) to v and a chordless path $Q_2 = (q_1 = v, q_2 = u, \dots, q_m = x)$ from (v, u) to x in \mathcal{C} , such that $y \neq v$ and $u \neq x$.

Note that by Observation 3.46, we then have

$$\tau(p_i) < \tau(v) \text{ and } p_i \rightarrow p_{i+1} \text{ for every } 1 \leq i \leq l-1. \quad (7)$$

There are two possibilities: either $Q'_1 = (p_1 = x, p_2 = y, \dots, p_l = v, w)$ is a chordless path in \mathcal{C} , or Q'_1 is not a chordless path in \mathcal{C} . If Q'_1 were a chordless path, then from Observation 3.46, $v \rightarrow w \in M$. This is a contradiction, since by hypothesis, we have $v - w \in M$. Thus, Q'_1 cannot be a chordless path.

We now show that $Q := (p_1 = x, p_2 = y, \dots, p_{l-1}, w)$ (obtained by replacing the node p_l with w in Q_1) is a chordless path.

Since Q_1 is a chordless path in \mathcal{C} , and Q'_1 is not a chordless path in \mathcal{C} , there must exist an edge $p_i - w \in \mathcal{C}$ for some $i < l$. As observed in eq. (7), for every $i \leq l-1$, $\tau(u_i) < \tau(v) < \tau(w)$ (the second inequality comes from the hypothesis of Claim 3.55). Thus, for any $i \leq l-2$, if $p_i - w \in \mathcal{C}$, then from Observation 2.14, $p_i - v \in \mathcal{C}$. But, then Q_1 is not a chordless path, a contradiction. This implies that the neighbor of w in Q_1 , responsible for ensuring that Q'_1 is not a chordless path, must be p_{l-1} . This further implies that Q is a chordless path.

Note now that since $\tau(p_{l-1}) < \tau(v) < \tau(w)$ and $\tau(u) < \tau(w)$, Observation 2.14 implies that $p_{l-1} - u \in \mathcal{C}$ (since w is adjacent to both p_{l-1} and u). We will now show that

$$p_{l-1} \rightarrow w \in M \text{ and } u \rightarrow p_{l-1} \notin M. \quad (8)$$

The first follows from Observation 3.46 since Q as constructed above is a chordless path in \mathcal{C} from (x, y) to (p_{l-1}, w) where $x \rightarrow y \in M$. Suppose, if possible, that $u \rightarrow p_{l-1} \in M$. Then, by Observation 3.45, and the observation preceding eq. (8), $\tau(u) < \tau(p_{l-1}) < \tau(w)$. Thus, while running lines 11-13 of alg. 4, we get $u \rightarrow w \in M$ (as $\tau(u) < \tau(p_{l-1}) < \tau(w)$ and $u \rightarrow p_{l-1}, p_{l-1} \rightarrow w \in M$). This is a contradiction, as $u - w \in M$. This implies that $u \rightarrow p_{l-1} \notin M$.

We now show that there exists some $i \geq 2$ for which $q_{i+1} \rightarrow q_i \in M$. Recall that $Q_2 = (q_1 = v, q_2 = u, \dots, q_m = x)$ is a chordless path from (v, u) to x in \mathcal{C} , an undirected connected component of M_a , as guaranteed by the hypothesis for this case.

From the construction, $\text{skeleton}(M[V_{\mathcal{C}}]) = \mathcal{C}$. Since $u - p_{l-1} \in \mathcal{C}$, therefore, $u - p_{l-1} \in \text{skeleton}(M)$. This implies from eq. (8), either $p_{l-1} \rightarrow u \in M$ or $p_{l-1} - u \in M$. Thus, from eq. (7), if we do not have $q_{i+1} \rightarrow q_i \in M$ for any $i \geq 2$, then M has a directed cycle contained in the sequence $C_1 := (p_1 = x, p_2, \dots, p_{l-1}, q_2 = u, q_3, \dots, q_m = x)$. From Proposition 2.7, there must thus exist a directed cycle of the form $C = (a, b, c)$ such that a, b, c are nodes of the cycle C_1 .

From eq. (7), $\tau(p_1 = q_m) < \tau(v)$, therefore, from Observation 2.15, for path Q_2 , for $i > 1$, $\tau(q_i) < \tau(v)$. Also, from eq. (7), for $1 \leq i < l$, $\tau(p_i) < \tau(v)$ since $\tau(v) < \tau(w)$. Therefore, for $1 \leq i < l$, $\tau(p_i) < \tau(w)$, and for $i > 1$, $\tau(q_i) < \tau(w)$. This implies each node of the cycle C_1 has a rank lower than the rank of w in τ . Therefore, the rank of the highest rank node of the cycle C in τ is less than the rank of w .

However, this is a contradiction, as per our induction hypothesis, there cannot be any such directed cycle with the rank of the highest rank node being less than $k+1$. This validates that for some $i \geq 2$, $q_{i+1} \rightarrow q_i \in M$.

Since Q_2 is an undirected chordless path in \mathcal{C} , therefore, $\bar{Q}_2 = (q_m, q_{m-1}, \dots, q_2, q_1)$, the reverse of the path Q_2 , is also a chordless path. Consequently, $Q_3 = (q_{i+1}, q_i, \dots, q_2, q_1)$, a subpath of \bar{Q}_2 , is also a chordless path in \mathcal{C} . Since $q_{i+1} \rightarrow q_i \in M$, according to Observation 3.46, $\tau(q_{i+1}) < \tau(q_i) < \dots, \tau(q_2) < \tau(q_1)$, and for each $i+1 \leq j \leq 2$, $q_j \rightarrow q_{j-1} \in M$.

We thus have the induced sub-graph $p \rightarrow u \rightarrow v$ in M with $p = q_3$ (remember $q_2 = u$ and $q_1 = v$). This case is now exactly the same as the first case of the first possibility, and hence as proved there, this case cannot occur.

The above discussion shows that there is no possibility for $u \rightarrow v - w - u \in M$. □

Proof of Claim 3.57. Suppose that $u - v \rightarrow w - u \in M$. From Corollary 3.43, either of the following has occurred:

1. $v \rightarrow w$ is strongly protected in O .
2. There exists $a \rightarrow b \in O$ such that $(a, b) \neq (v, w)$, and there exists a chordless path from (a, b) to (v, w) in \mathcal{C} .
3. There exists $a \rightarrow b \in O$ such that there exist chordless paths in \mathcal{C} from (a, b) to w and from (w, v) to a such that $b \neq w$ and $v \neq a$.

Let's examine each possibility.

1. Suppose $v \rightarrow w$ is strongly protected in O . This implies that $v \rightarrow w$ is part of one of the induced subgraphs of O , as illustrated in fig. 2.

- (a) Assume that $v \rightarrow w$ is strongly protected due to its presence in an induced subgraph $p \rightarrow v \rightarrow w \in O$ (similar to Figure 2.a). Similar to the first case of the first possibility of the proof of Claim 3.56, the only relevant scenario is when p belongs to \mathcal{C} .

From the construction, $\text{skeleton}(M[V_{\mathcal{C}}]) = \mathcal{C}$. This implies $p - v \in \mathcal{C}$ and $p - w \notin \mathcal{C}$. Given that $p - v \in \mathcal{C}$ and $p \rightarrow v \in M$, we can infer from Observation 3.45 that $\tau(p) < \tau(v)$. Given that $u - v \in \mathcal{C}$ and $\tau(u) < \tau(v)$, we can deduce from Observation 2.14 that $p - u \in \mathcal{C}$. Since $\text{skeleton}(M[V_{\mathcal{C}}]) = \mathcal{C}$, therefore, either (1) $p - u \in M$ or (2) $p \rightarrow u \in M$ or (3) $u \rightarrow p \in M$. We will show that none of the three can occur.

- i. If $p - u \in M$, then M would possess a directed cycle (p, v, u, p) with the highest rank node v having a rank less than $\tau(w)$, thereby contradicting our induction hypothesis.
- ii. If $p \rightarrow u \in M$ then using Observation 3.45, we have $\tau(p) < \tau(u)$. However, since $p - w \notin \mathcal{C}$ and $\tau(p) < \tau(u) < \tau(w)$, applying lines 11-13 of alg. 4 would lead to $u \rightarrow w \in M$, which contradicts $u - w \in M$.
- iii. And, if $u \rightarrow p \in M$ then using Observation 3.45, we have $\tau(u) < \tau(p)$. Therefore, since $\tau(u) < \tau(p) < \tau(v)$ and $u \rightarrow p \rightarrow v - u \in M$, applying lines 11-13 of alg. 4 would lead to $u \rightarrow v \in M$, which contradicts $u - v \in M$.

Hence, based on the analysis above, it becomes evident that none of these scenarios are feasible. Therefore, we conclude that this case cannot occur.

- (b) Assume that $v \rightarrow w$ is strongly protected due to the presence in an induced subgraph $v \rightarrow w \leftarrow p \in O$ (as shown in Figure 2.b). Similar to the second case of the first possibility of the proof of Claim 3.56, this case does not occur.

- (c) Suppose $v \rightarrow w$ is strongly protected in O because it is part of an induced subgraph $v \rightarrow p \rightarrow w \leftarrow v \in O$ (same as Figure 2.c). Similar to the third case of the first possibility of the proof of Claim 3.56, $p \in \mathcal{C}$. From Lemma 3.35, $v \rightarrow p \rightarrow w \leftarrow v \in M$. From the construction, $\text{skeleton}(M[V_{\mathcal{C}}]) = \mathcal{C}$. This implies $p - w \in \mathcal{C}$ (as $p \rightarrow w \in M$ and $p, w \in \mathcal{C}$). From Observation 3.45, $\tau(v) < \tau(p) < \tau(w)$. From Observation 2.14, this implies $u - p \in \mathcal{C}$ (since $p - w, u - w \in \mathcal{C}, \tau(p) < \tau(w)$ and $\tau(u) < \tau(w)$). Since $u - p \in \mathcal{C}$ and $\text{skeleton}(M[V_{\mathcal{C}}]) = \mathcal{C}$, therefore, either (1) $u - p \in M$, or (2) $u \rightarrow p \in M$, or (3) $p \rightarrow u \in M$. We show that none of these cases occur.

- i. If $u - p \in M$ then M have a directed cycle $u - v \rightarrow p - u$ such that the rank of the highest rank node of the cycle is less than $\tau(w)$, which contradicts our induction hypothesis. This implies $u - p \notin M$.
- ii. Suppose $u \rightarrow p \in M$. Then, since $u \rightarrow p, p \rightarrow w \in M$, while running lines 11-13 of alg. 4, we get $u \rightarrow w \in M$, a contradiction, as $u - w \in M$. This implies $u \rightarrow p \notin M$ does not occur.
- iii. Since $\tau(u) < \tau(p)$, from Observation 3.45, $p \rightarrow u \notin M$.

The above analysis shows that neither $u - p \in M$ nor $u \rightarrow p \in M$ nor $p \rightarrow u \in M$. This implies that this case does not occur.

- (d) Suppose $v \rightarrow w$ is strongly protected in O because it is part of an induced subgraph $v \rightarrow w \leftarrow p - v - p' \rightarrow w$ (same as Figure 2.d). Similar to the fourth case of the first possibility of the proof of Claim 3.56, this case does not occur.
2. Suppose $v \rightarrow w \in M$ because of the existence of some $x - y \in \mathcal{C}$ such that $x \rightarrow y \in O$, and there exists a chordless path $P = (p_1 = x, p_2 = y, \dots, p_{l-1} = v, p_l = w)$ from (x, y) to (v, w) in \mathcal{C} . From Observation 3.46, $p_{l-2} \rightarrow p_{l-1} \in M$. We thus have the induced sub-graph $p \rightarrow v \rightarrow w$ in M with $p = p_{l-2}$. This case is now exactly the same as the first case of the first possibility, and hence as proved there, cannot occur.
3. Suppose there exists $x - y \in O$ such that $x \rightarrow y \in O$, and there exists a chordless path in \mathcal{C} : $Q_1 = (p_1 = x, p_2 = y, \dots, p_l = w)$ from (x, y) to w in \mathcal{C} , and $Q_2 = (q_1 = w, q_2 = v, \dots, q_m = x)$ from (w, v) to x in \mathcal{C} such that $y \neq w$ and $v \neq x$. According to Observation 3.46, we have $\tau(p_{l-1}) < \tau(w)$, and $p_{l-1} \rightarrow w \in M$. Considering that $\tau(p_{l-1}) < \tau(w)$, $\tau(v) < \tau(w)$, and $\tau(u) < \tau(w)$, it follows from Observation 2.14 that $p_{l-1} - u$ and $p_{l-1} - v$ are in \mathcal{C} .

There are two possibilities: either $v \rightarrow p_{l-1} \in M$, or $v \rightarrow p_{l-1} \notin M$. Let's examine both possibilities:

- (a) Suppose $v \rightarrow p_{l-1} \in M$. Since $u - p_{l-1} \in \mathcal{C}$, we have either $u - p_{l-1} \in M$, or $u \rightarrow p_{l-1} \in M$, or $p_{l-1} \rightarrow u \in M$.
- i. Suppose $u - p_{l-1} \in M$. This would result in a directed cycle (u, v, p_{l-1}, u) in M where the node with the highest rank, u , has a rank lower than $\tau(w)$, contradicting our induction hypothesis.
 - ii. Suppose $u \rightarrow p_{l-1} \in M$. Since $p_{l-1} \rightarrow w \in M$, running lines 11-13 of alg. 4 would yield $u \rightarrow w \in M$. This contradicts the assumption $u - w \in M$. Thus, $u \rightarrow p_{l-1} \notin M$.
 - iii. From the hypothesis of Claim 3.55, $\tau(u) < \tau(v)$. Since from our assumption, $v \rightarrow p_{l-1} \in M$, therefore, from Observation 3.45, $\tau(v) < \tau(p_{l-1})$. This implies $\tau(u) < \tau(p_{l-1})$. Then, from Observation 3.45, $p_{l-1} \rightarrow u \notin M$.

The above analysis shows that neither $u - p_{l-1} \in M$ nor $u \rightarrow p_{l-1} \in M$ nor $p_{l-1} \rightarrow u \in M$.

- (b) Suppose $v \rightarrow p_{l-1} \notin M$. Then for some $i \geq 2$, $q_{i+1} \rightarrow q_i$; otherwise, a directed cycle $(p_1 = x, p_2, \dots, p_{l-1}, q_2 = u, q_3, \dots, q_m = x)$ with a maximum rank lower than $\tau(w)$ would exist in M . From Proposition 2.7, this would lead to a directed cycle in M of length 3 with a maximum rank lower than $\tau(w)$, contradicting our induction hypothesis. Hence, for some $i \geq 2$, $q_{i+1} \rightarrow q_i \in M$. Since Q_2 is an undirected chordless path in \mathcal{C} , therefore, $\bar{Q}_2 = (q_m, q_{m-1}, \dots, q_2, q_1)$, the reverse of the path Q_2 , is also a chordless path. Consequently, $Q_3 = (q_{i+1}, q_i, \dots, q_2, q_1)$, a subpath of \bar{Q}_2 , is also a chordless path in \mathcal{C} . Since $q_{i+1} \rightarrow q_i \in M$, according to Observation 3.46, $\tau(q_{i+1}) < \tau(q_i) < \dots, \tau(q_2) < \tau(q_1)$, and for each $i + 1 \leq j \leq 2$, $q_j \rightarrow q_{j-1} \in M$. We thus have the induced sub-graph $p \rightarrow v \rightarrow w$ in M with $p = q_3$ (remember Q_2 is a chordless path, $q_1 = w$ and $q_2 = v$). This case is now exactly the same as the first case of the first possibility, and hence as proved there, cannot occur.

The above discussion shows a contradiction in each possibility, thereby demonstrating that $u - v \rightarrow w - u \notin M$. This establishes the truth of Claim 3.57. □

Proof of Claim 3.58. Suppose $u - v - w \leftarrow u \in M$. From Corollary 3.43, either of the following occurred:

1. $u \rightarrow w$ is strongly protected in O .
2. There exists $a \rightarrow b \in O$ such that $(a, b) \neq (u, w)$ and there exists a chordless path from (a, b) to (u, w) in \mathcal{C} .

3. There exists $a \rightarrow b \in O$ such that there exist chordless paths in \mathcal{C} from (a, b) to w and from (w, u) to a such that $b \neq w$ and $u \neq a$.

We will consider each possibility.

1. $u \rightarrow w$ is strongly protected in O . Then, $u \rightarrow w$ is part of one of the induced subgraphs of O in fig. 2.

- (a) Suppose $u \rightarrow w$ is strongly protected because it is part of an induced subgraph $p \rightarrow u \rightarrow w \in O$ (same as Figure 2.a). Similar to the first case of the first possibility of the proof of Claim 3.56, the only possibility we need to consider is when $p \in \mathcal{C}$. From Observation 3.45, $\tau(p) < \tau(u)$. If $p - v \notin \mathcal{C}$, then while running lines 11-13 of alg. 4, we get $u \rightarrow v \in M$, a contradiction, as $p \rightarrow u - v$ is an induced subgraph in M and $\tau(p) < \tau(u) < \tau(v)$ (second inequality comes from the hypothesis of Claim 3.55). However, this is a contradiction as $u - v \in M$. This implies $p - v \in \mathcal{C}$. There are three possibilities: either $v - p \in M$, or $v \rightarrow p \in M$, or $p \rightarrow v \in M$. Let's consider each one.

- i. If $v - p \in M$ then we have a directed cycle (p, u, v, p) in M such that the rank of the highest rank node of the cycle (which is v) is less than the rank of w , this contradicts the induction hypothesis of Claim 3.55. Hence implies $v - p \notin M$.
- ii. From Observation 3.45, $v \rightarrow p \notin M$ as $\tau(p) < \tau(v)$.
- iii. Suppose $p \rightarrow v \in M$. But, then while running lines 11-13 of alg. 4, we get $v \rightarrow w \in M$, a contradiction, as $p \rightarrow v - w$ is an induced subgraph in M and $\tau(p) < \tau(v) < \tau(w)$.

We have shown that in all the possibilities of this subcase, we get a contradiction. Therefore, this subcase cannot occur.

- (b) Suppose that $u \rightarrow w$ is strongly protected because it is part of an induced subgraph $u \rightarrow w \leftarrow p \in O$ (same as Figure 2.b). Similar to the second case of the first possibility of the proof of Claim 3.56, this case does not occur.

- (c) Suppose $u \rightarrow w$ is strongly protected in O because it is part of an induced subgraph $u \rightarrow p \rightarrow w \leftarrow u \in O$ (same as Figure 2.c). Similar to the third case of the first possibility of the proof of Claim 3.56, the only possibility we need to consider is when $p \in \mathcal{C}$. From Observation 3.45, $\tau(u) < \tau(p) < \tau(w)$. From Observation 2.14, $v - p \in \mathcal{C}$ (since $p - w, v - w \in \mathcal{C}, \tau(p) < \tau(w)$ and $\tau(v) < \tau(w)$). From the construction, $\text{skeleton}(M[V_{\mathcal{C}}]) = \mathcal{C}$. Therefore, $v - p \in \text{skeleton}(M)$. This implies either $v - p \in M$ or $v \rightarrow p \in M$ or $p \rightarrow v \in M$. We go through each possibility.

- i. Suppose $v - p \in M$. Then we get a directed cycle (u, p, v, u) in M such that the highest rank node (which could be either v or p) in the cycle is less than the rank of w in τ , this contradicts the induction hypothesis of Claim 3.55. This implies $v - p \notin M$.
- ii. Suppose $v \rightarrow p \in M$. Then from Observation 3.45, $\tau(v) < \tau(p)$. Then while running lines 11-13 of alg. 4, we get $v \rightarrow w \in M$ (as $\tau(v) < \tau(p) < \tau(w)$ and $v \rightarrow p, p \rightarrow w \in M$). But, this is a contradiction as $v - w \in M$. This implies $v \rightarrow p \notin M$.
- iii. Suppose $p \rightarrow v \in M$. Then from Observation 3.45, $\tau(p) < \tau(v)$. Then while running lines 11-13 of alg. 4, we get $u \rightarrow v \in M$ (as $\tau(u) < \tau(p) < \tau(v)$ and $u \rightarrow p, p \rightarrow v \in M$). But, this is a contradiction as $u - v \in M$. This implies $p \rightarrow v \notin M$.

The above analysis shows that none of the possibilities can occur. This implies that this case does not occur.

- (d) Suppose $u \rightarrow w$ is strongly protected in O because it is part of an induced subgraph $u \rightarrow w \leftarrow p - u - p' \rightarrow w$ (same as Figure 2.d). Similar to the fourth case of the first possibility of the proof of Claim 3.56, this case does not occur.

2. Suppose $u \rightarrow w \in M$ because of the existence of some $x - y \in \mathcal{C}$ such that $x \rightarrow y \in O$, and there exists a chordless path $P = (p_1 = x, p_2 = y, \dots, p_{l-1} = u, p_l = w)$ from (x, y) to (u, w) in \mathcal{C} . From Observation 3.46, $p_{l-2} \rightarrow p_{l-1} \in M$. We thus have the induced sub-graph $p \rightarrow u \rightarrow w$ in M with $p = p_{l-2}$. This case is now exactly the same as the first case of the first possibility, and hence as proved there, cannot occur.

3. Suppose there exists $x - y \in O$ such that $x \rightarrow y \in O$, and there exist chordless paths in \mathcal{C} : $Q_1 = (p_1 = x, p_2 = y, \dots, p_l = w)$ from (x, y) to w in \mathcal{C} , and $Q_2 = (q_1 = w, q_2 = u, \dots, q_m = x)$ from (w, u) to x such that $y \neq w$ and $u \neq x$. From Observation 3.46, $\tau(p_{l-1}) < \tau(w)$, and $p_{l-1} \rightarrow w \in M$. Since $\tau(p_{l-1}) < \tau(w)$, $\tau(v) < \tau(w)$, and $\tau(u) < \tau(w)$ (the last two inequality comes from the hypothesis of Claim 3.55), therefore, from Observation 2.14, $u - p_{l-1}, v - p_{l-1} \in \mathcal{C}$. From the construction, $\text{skeleton}(M[V_{\mathcal{C}}]) = \mathcal{C}$. This implies $u - p_{l-1}, v - p_{l-1} \in \text{skeleton}(M)$. There are two possibilities: either $u \rightarrow p_{l-1} \in M$, or $u \rightarrow p_{l-1} \notin M$ (i.e., either $u - p_{l-1} \in M$ or $p_{l-1} \rightarrow u \in M$). We go through each possibility.

- (a) Suppose $u \rightarrow p_{l-1} \in M$. Since $v - p_{l-1} \in \text{skeleton}(M)$, either we have $v - p_{l-1} \in M$ or $v \rightarrow p_{l-1} \in M$ or $v \leftarrow p_{l-1} \in M$.
- i. Suppose $v - p_{l-1} \in M$. Then we have a directed cycle (u, p_{l-1}, v, u) in M such that the rank of the highest rank node in the cycle is bigger than $\tau(w)$ in τ . This contradicts the induction hypothesis of Claim 3.55. This implies $v - p_{l-1} \notin M$.
 - ii. Suppose $v \rightarrow p_{l-1} \in M$. Then, since $p_{l-1} \rightarrow w \in M$, therefore, while running lines 11-13 of alg. 4, we get $v \rightarrow w \in M$ (as $v \rightarrow p_{l-1}, p_{l-1} \rightarrow w \in M$). But, from our assumption, $v - w \in M$. This implies that $v \rightarrow p_{l-1} \notin M$.
 - iii. Suppose $v \leftarrow p_{l-1} \in M$. Then, while running lines 11-13 of alg. 4, we get $u \rightarrow v \in M$ (as $u \rightarrow p_{l-1} \in M, p_{l-1} \rightarrow v \in M$). But, from our assumption $u - v \in M$. This implies that $v \leftarrow p_{l-1} \notin M$.

The above analysis shows that when $u \rightarrow p_{l-1} \in M$, neither $v - p_{l-1} \in M$ nor $v \rightarrow p_{l-1} \in M$ nor $v \leftarrow p_{l-1} \in M$. This implies $u \rightarrow p_{l-1} \in M$ does not occur.

- (b) Suppose $u \rightarrow p_{l-1} \notin M$. Then for some $i \geq 2$, $q_{i+1} \rightarrow q_i$; otherwise, a directed cycle $(p_1 = x, p_2, \dots, p_{l-1}, q_2 = u, q_3, \dots, q_m = x)$ with a maximum rank lower than $\tau(w)$ would exist in M . From Proposition 2.7, this would lead to a directed cycle in M of length 3 with a maximum rank lower than $\tau(w)$, contradicting the induction hypothesis of Claim 3.55. Hence, for some $i \geq 2$, $q_{i+1} \rightarrow q_i \in M$. Since Q_2 is an undirected chordless path, $\bar{Q}_2 = (q_m, q_{m-1}, \dots, q_2, q_1)$, the reverse of the path Q_2 , is also a chordless path. Consequently, $Q_3 = (q_{i+1}, q_i, \dots, q_2, q_1)$, a subpath of \bar{Q}_2 , is also a chordless path in \mathcal{C} . Since $q_{i+1} \rightarrow q_i \in M$, according to Observation 3.46, $\tau(q_{i+1}) < \tau(q_i) < \dots, \tau(q_2) < \tau(q_1)$, and for each $i + 1 \leq j \leq 2$, $q_j \rightarrow q_{j-1} \in M$. We thus have the induced sub-graph $p \rightarrow u \rightarrow w$ in M with $p = q_3$ (remember Q_2 is a chordless path, $q_1 = w$ and $q_2 = u$). This case is now exactly the same as the first case of the first possibility, and hence as proved there, cannot occur.

This indicates a contradiction in all possibilities, thereby demonstrating that $u - v - w \leftarrow u \notin M$. This establishes the truth of Claim 3.58. □

Proof of Observation 3.52. Let $C = (u_1, u_2, \dots, u_l, u_{l+1} = u_1)$ be a chordless cycle in M . If C lies entirely within M_1 (i.e., each node of C belongs to V_{M_1}), then by Observation 3.51, C is undirected. However, from the construction of M , (a) for any $u, v \in V_{M_1}$, if $u - v \in M$, then $u - v \in M_1$ (due to the initialization step, line 1 of alg. 4, a directed edge in M_1 is a directed edge in M), and (b) if there is no chord in C in M then there is no chord in C in M_1 as well (because all the nodes of C are in M_1 and $\text{skeleton}(M[V_{M_1}]) = \text{skeleton}(M_1)$). This implies that C is an undirected chordless cycle in M_1 , which contradicts item 2 of Theorem 2.18. Hence, the nodes in C cannot be exclusively within M_1 or M_2 . Thus, there must exist a node u_i in C that is in $V_{M_1} \setminus I$, a node u_j in C that is in $V_{M_2} \setminus I$, and a node u_k in C that is in I , as $I = V_{M_1} \cap V_{M_2}$ is a vertex separator of $H = \text{skeleton}(M)$ that separates $V_{M_1} \setminus I$ and $V_{M_2} \setminus I$.

Without loss of generality, let's assume that $u_1 \in I$. Therefore, $u_1, u_2, u_l \in V_O$ since u_2 and u_l are neighbors of u_1 , and $I \cup N(I, H) \subseteq V_O$. Since O is an induced subgraph of M (from Lemma 3.35), and C is a cycle in M , for any edge (u_i, u_{i+1}) in C , if $u_i, u_{i+1} \in V_O$, then $(u_i, u_{i+1}) \in O$. More specifically, $(u_1, u_2), (u_l, u_1) \in E_O$.

We enumerate the edges in C present in O in the order they appear starting from (u_1, u_2) , ending with (u_l, u_1) . This enumeration is represented as $L = ((x_1, y_1), (x_2, y_2), \dots, (x_m, y_m))$, where $(x_1, y_1) = (u_1, u_2)$ and $(x_m, y_m) = (u_l, u_1)$. Notably, $m \geq 2$ since the list contains the edges (u_1, u_2) and (u_l, u_1) . Additionally, let $(x_{m+1}, y_{m+1}) = (x_1, x_2)$, and define $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$. We present the following claims for the edges in this enumeration:

1. For $1 \leq i \leq m$, if $y_i = x_{i+1}$, then (x_i, y_i, y_{i+1}) is a triangle-free path in O , and $P_1((x_i, y_i), (x_{i+1}, y_{i+1})) = 1$.
2. For $1 \leq i \leq m$, if $y_i \neq x_{i+1}$, then either there exists a triangle-free path from (x_i, y_i) to (x_{i+1}, y_{i+1}) in M_1 , or there exists a triangle-free path from (x_i, y_i) to (x_{i+1}, y_{i+1}) in M_2 , and $P_1((x_i, y_i), (x_{i+1}, y_{i+1})) = 1$.

Proof of the claim. For any $1 \leq i \leq m$ where $y_i = x_{i+1}$, due to the chordless nature of C and the presence of (x_i, y_i) and (x_{i+1}, y_{i+1}) as edges in O , it follows that (x_i, y_i, y_{i+1}) is a triangle-free path in O . Consequently, considering that $(P_1, P_2) = \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$, by item 1 of Definition 3.17, $P_1((x_i, y_i), (x_{i+1}, y_{i+1}))$ is initialized to 1 due to the presence of a triangle-free path from (x_i, y_i) to (x_{i+1}, y_{i+1}) in O . This validates the first part of the claim.

Suppose for some $1 \leq i \leq m$, $y_i \neq x_{i+1}$. Then, y_i cannot be part of I . Otherwise, the neighbors of y_i would be within O ($I \cup N(I, H) \subseteq V_O$). That means, for $1 \leq j \leq l$, if $y_i = u_j$ then the next node u_{j+1} in C is also in V_O . This implies (u_j, u_{j+1}) is in O , as C is a cycle in M , $u_j, u_{j+1} \in V_O$, and O is an induced subgraph of M . This implies the next edge after (x_i, y_i) in L is $(u_j, u_{j+1}) = (x_{i+1}, y_{i+1})$. But, then $y_i = x_{i+1}$, a contradiction. This implies that y_i must either be in $V_{M_1} \setminus I$ or in $V_{M_2} \setminus I$. Without loss of generality, assume that $y_i \in V_{M_1} \setminus I$. Consequently, x_{i+1} must not be in $V_{M_2} \setminus I$, or else the path between y_i and x_{i+1} in C would contain an I node, as I serves as a vertex separator between $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. This implies that within the path from y_i to x_{i+1} in C , there exists an edge from C in O due to the containment of $I \cup N(I, H) \subseteq V_O$ and the fact that C is a cycle in M , and O is an induced subgraph of M . Similarly, x_{i+1} must not be in I ; otherwise, the edge right before (x_{i+1}, y_{i+1}) in C would be in O as well (since $I \cup N(I, H) \subseteq V_O$), but the edge prior to (x_{i+1}, y_{i+1}) in O is (x_i, y_i) . This implies that (x_i, y_i) is right before (x_{i+1}, y_{i+1}) in C , leading to $y_i = x_{i+1}$, contradicting the assumption. Hence, $x_{i+1} \in V_{M_1} \setminus I$, which further implies $y_{i+1} \in V_{M_1}$ since a neighbor of a node in $V_{M_1} \setminus I$ within V_O must belong to V_{M_1} , and y_{i+1} is the neighbor of x_{i+1} and lies in V_O (due to $(x_{i+1}, y_{i+1}) \in O$). Thus, x_i, y_i, x_{i+1} , and y_{i+1} all belong to M_1 . Furthermore, the path from (x_i, y_i) to (x_{i+1}, y_{i+1}) must exclusively lie within M_1 , otherwise there would be an edge in O between (x_i, y_i) and (x_{i+1}, y_{i+1}) in C , a contradiction. With our construction of M , for any $u, v \in V_{M_1}$, if $(u, v) \in M$, then $(u, v) \in M_1$. This implies that the chordless path from (x_i, y_i) to (x_{i+1}, y_{i+1}) in C is also a chordless path in M_1 , meaning $P_{a1}((x_i, y_i), (x_{i+1}, y_{i+1})) = 1$ (since every chordless path is a triangle-free path). Also, $y_1 \rightarrow x_i \notin M$, otherwise C wouldn't possess the edge (x_i, y_i) . Therefore, by item 2 of Definition 3.17, $P_1((x_i, y_i), (x_{i+1}, y_{i+1})) = 1$. \square

It follows that for all $1 \leq i \leq m$, $P_1((x_i, y_i), (x_{i+1}, y_{i+1})) = 1$. Now, we claim that $P_1((x_1, y_1), (x_m, y_m)) = 1$. Suppose this isn't true and select the smallest $j \leq m$ for which $P_1((x_1, y_1), (x_j, y_j)) = 0$. Given the previous discussion, since $P_1((x_i, y_i), (x_{i+1}, y_{i+1})) = 1$ for all $1 \leq i \leq m$, it implies that $j > 2$. This further leads to $P_1((x_1, y_1), (x_{j-1}, y_{j-1})) = P_1((x_{j-1}, y_{j-1}), (x_j, y_j)) = 1$. But, from item 3 of Definition 3.17, we get $P_1((x_1, y_1), (x_i, y_i)) = 1$, a contradiction. Hence, $P_1((x_1, y_1), (x_m, y_m)) = 1$. Additionally, from our earlier claim, $P_1((x_m, y_m), (x_{m+1}, y_{m+1})) = (x_1, y_1) = 1$.

According to the assumption of Lemma 3.34, O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$. Thus, by Definition 3.24, $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$ is valid (as per Definition 3.23). However, the fact that $P_1((x_1, y_1), (x_m, y_m)) = P_1((x_m, y_m), (x_1, y_1)) = 1$ demonstrates that $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$ is invalid, leading to a contradiction. Consequently, the existence of a chordless cycle in M is not possible. \square

Proof of Lemma 3.48. Assume, for the sake of contradiction, that Lemma 3.48 is false, implying the existence of an undirected connected component denoted as C in M that is not chordal. Since C is not chordal, there must exist an undirected chordless cycle $C = (u_1, u_2, \dots, u_l, u_{l+1} = u_1)$, where l is greater than or

equal to 4. This implies that no edges exist between non-adjacent nodes of C within \mathcal{C} . We consider two possibilities: (a) there is an edge between non-adjacent nodes of C in M , but it is directed; or (b) no edge, either directed or undirected, exists between non-adjacent nodes of C in M .

Let's analyze these cases. Firstly, suppose there is a directed edge between two non-adjacent nodes of C in M . This would create a directed cycle in M , contradicting Lemma 3.47 which states that M is a chain graph.

Next, consider the second possibility, where there are no edges, neither directed nor undirected, between non-adjacent nodes in M . As a result, C would be a chordless cycle in M . However, this contradicts Observation 3.52, which establishes that M does not contain any chordless cycles.

Consequently, neither of these cases is feasible. Hence, we conclude that Lemma 3.48 holds true, and there are no undirected connected components in M that are not chordal. \square

Proof of Lemma 3.49. We will first demonstrate that there are no induced subgraphs of the form $a \rightarrow b - c$ within $M[V_{M_1}]$ and $M[V_{M_2}]$.

Let's consider the scenario where there is an induced subgraph of the form $a \rightarrow b - c$ within $M[V_{M_i}]$ for some $i \in \{1, 2\}$. Based on the construction of M , either $a \rightarrow b - c$ or $a - b - c$ is an induced subgraph of M_i . According to item 3 of Theorem 2.18, $a \rightarrow b - c$ cannot be an induced subgraph of M_i .

If $a - b - c \in M_i$, then a , b , and c all belong to an undirected connected component \mathcal{C} of M_i , and (a, b, c) forms a chordless path from (a, b) to (b, c) within \mathcal{C} . Now, considering Observation 3.46, since (a, b, c) is a chordless path and $a \rightarrow b \in M$, it follows that $b \rightarrow c \in M$. This leads to a contradiction since we assumed that $b - c \in M$. Thus, we can conclude that there cannot exist an induced subgraph of the form $a \rightarrow b - c$ within $M[V_{M_i}]$.

The above analysis implies that if there exists an induced subgraph $a \rightarrow b - c$ in M , then all the nodes of the induced subgraph cannot belong exclusively to V_{M_1} or V_{M_2} . This implies that at least one node must be in $V_{M_1} \setminus I$ (where $I = V_{H_1} \cap V_{H_2}$), and one node must be in $V_{M_2} \setminus I$. Since I acts as a separator between $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$, the induced subgraph must contain at least one node from I . Therefore, we have two possibilities: (i) $a \in V_{M_1} \setminus I$, $b \in I$, and $c \in V_{M_2} \setminus I$, or (ii) $a \in V_{M_2} \setminus I$, $b \in I$, and $c \in V_{M_1} \setminus I$. In both cases, b belongs to I .

Consequently, we have $a, b, c \in I \cup N(I, H) \subseteq V_O$. This implies that the induced subgraph is in O as O is an induced subgraph of M . However, this contradicts the fact that O is a partial MEC, as stated in item 3 of Definition 2.21. Nonetheless, O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, and Definition 3.24 implies that O is a partial MEC. This creates a contradiction. Hence, there cannot be any such induced subgraph in M . Thus, Lemma 3.49 is validated. \square

Proof of Lemma 3.50. Suppose $u \rightarrow v \in M$. Since, from the construction, $\text{skeleton}(M) = \text{skeleton}(M_1) \cup \text{skeleton}(M_2) = H_1 \cup H_2$, there are two possibilities: either $u - v \in E_{H_1}$ or $u - v \in E_{H_2}$. Without loss of generality, we assume $u - v \in E_{H_1}$. From the construction of M , either $u \rightarrow v \in M_1$ or $u - v \in M_1$. Because if $v \rightarrow u \in M_1$ then due to the initialization step, line 1 of alg. 4, we have $v \rightarrow u \in M$, which contradicts our assumption that $u \rightarrow v \in M$. We will show that in both cases, $u \rightarrow v$ is strongly protected in M . We will consider each possibility in turn:

1. Suppose $u \rightarrow v \in M_1$. Then $u \rightarrow v$ is strongly protected in M_1 according to item 4 of Theorem 2.18. As per the construction of M , all the directed edges in M_1 are also directed in M (due to line 1 of alg. 4). Now, let's consider the induced subgraphs shown in (a), (b), (c), and (d) of fig. 2.

If $u \rightarrow v$ is strongly protected in M_1 due to its inclusion in an induced subgraph shown in (a), (b), or (c) of fig. 2, then the fact that all the edges in those induced subgraphs are directed implies that the induced subgraph is also an induced subgraph of M (due to line 1 of alg. 4). This further implies $u \rightarrow v$ is also strongly protected in M .

Now, let's focus on the case where $u \rightarrow v$ is strongly protected in M_1 because it is part of the induced subgraph shown in fig. 2-(d). In this induced subgraph, the edges $w \rightarrow v$, $w' \rightarrow v$, and $u \rightarrow v$ are directed, indicating that $w \rightarrow v$, $w' \rightarrow v$, and $u \rightarrow v$ are also present in M due to the construction of M . To establish the strong protection of $u \rightarrow v$ in M , we consider different cases. If $w - u$ and $w' - u$

are present in M , then $u \rightarrow v$ is part of the induced subgraph shown in fig. 2-(d), making it strongly protected in M . If $u \rightarrow w$ or $u \rightarrow w'$ exists in M , then again $u \rightarrow v$ is part of an induced subgraph, either as shown in fig. 2-(c) with $u \rightarrow w \rightarrow v \leftarrow u$, or as shown in fig. 2-(c) with $u \rightarrow w' \rightarrow v \leftarrow u$. In both cases, $u \rightarrow v$ is strongly protected in M .

The remaining possibility is that $w \rightarrow u$ or $w' \rightarrow u$ are present in M . Since $w - u - w'$ is present in M_1 , it implies that w , u , and w' belong to the same undirected connected component \mathcal{C} of M_1 . Now, considering the fact that $w - u - w'$ forms a chordless path in \mathcal{C} , according to Observation 3.46, if $w \rightarrow u$ exists in M , then $u \rightarrow w'$ must also be present in M . Similarly, if $w' \rightarrow u \in M$, then $u \rightarrow w \in M$. And, as shown above, if $u \rightarrow w$ or $u \rightarrow w'$ exists in M , then $u \rightarrow v$ is strongly protected.

In conclusion, we have shown that in all possible cases, if $u \rightarrow v$ is strongly protected in M_1 , then it is also strongly protected in M . Hence, the claim is verified.

2. Suppose $u - v \in M_1$ and $u \rightarrow v \in M$. From Observation 3.40, $u \rightarrow v \in M$ either because (a) $u \rightarrow v$ is strongly protected in O , (b) there exists $x \rightarrow y \in O$ with $(u, v) \neq (x, y)$ and a chordless path from (x, y) to (u, v) in M_1 , or (c) there exists $x \rightarrow y \in O$ and chordless paths in M_1 from (x, y) to v and from (v, u) to x , with $y \neq v$ and $u \neq x$. We will consider each possibility in detail.

- (a) If $u \rightarrow v$ is strongly protected in O , then since O is an induced subgraph of M (from Lemma 3.35), if $u \rightarrow v$ is strongly protected in O , then it is strongly protected in M .
- (b) Suppose there exists an edge $x \rightarrow y \in O$ with a chordless path $Q = (u_1 = x, u_2 = y, \dots, u_{l-1} = u, u_l = v)$ from (x, y) to (u, v) in \mathcal{C} where $(x, y) \neq (u, v)$. According to Observation 3.46, $u_{l-2} \rightarrow u_{l-1} \rightarrow u_l \in M$. This implies that $u \rightarrow v$ is strongly protected in M as it is part of an induced subgraph shown in fig. 2-(a).
- (c) Suppose there are chordless paths $Q_1 = (u_1 = x, u_2 = y, \dots, u_l = u)$ in M_1 from (x, y) to u , and $Q_2 = (v_1 = v, v_2 = u, \dots, v_m = x)$ from (v, u) to x , where $y \neq u$ and $u \neq x$. All the edges of Q_1 and Q_2 must be undirected in M_1 , otherwise, concatenating Q_1 and Q_2 would create a directed cycle in M_1 , contradicting item 1 of Theorem 2.18. This implies that all nodes in Q_1 and Q_2 belong to an undirected connected component \mathcal{C} of M_1 .

Let τ be the LBFS ordering returned by alg. 3 when called at line 5 of alg. 4 for the undirected connected component \mathcal{C} of M_1 . According to Observation 3.46, $\tau(u_1) < \tau(u_2) < \dots < \tau(u_l)$, and for all $1 \leq i < l - 1$, $u_i \rightarrow u_{i+1} \in M$. According to Observation 3.45, $\tau(u) < \tau(v)$. Based on Observation 2.14, $u_{l-1} - u \in \mathcal{C}$.

If $u \rightarrow u_{l-1} \in M$, then $u \rightarrow v$ is strongly protected in M due to the induced subgraph $u \rightarrow u_{l-1} \rightarrow v \leftarrow u$ (as shown in fig. 2-(c)).

Suppose $u \rightarrow u_{l-1} \notin M$. Then there must exist $j \geq 2$ such that $v_{j+1} \rightarrow v_j \in M$, otherwise, $C = (u_1 = x, u_2, \dots, u_{l-1}, v_2, \dots, v_m = x)$ is a directed cycle in M , contradicting Lemma 3.47. In this case, $u_{j+1} \rightarrow u_j \in M$ for some $j \geq 2$. Then $Q'_2 = (u_{j+1}, u_j, \dots, u_2, u_1)$ is a chordless path and $u_{j+1} \rightarrow u_j \in M$. Then, according to Observation 3.46, $v_3 \rightarrow v_2 \rightarrow v_1 \in M$. This implies that $u \rightarrow v$ is strongly protected in M as it is part of an induced subgraph $v_3 \rightarrow v_2 \rightarrow v_1 = v$ (as shown in fig. 2-(a)).

Thus, we have shown that in all possible cases, $u \rightarrow v$ is strongly protected in M . This verifies the claim.

This completes the proof of Lemma 3.50. □

Proof of Lemma 3.37. Without loss of generality, we prove that $\mathcal{P}(M, V_{H_1}) = M_1$. Similarly, we can claim that $\mathcal{P}(M, V_{H_2}) = M_2$, which completes the proof of Lemma 3.37.

Since M_1 is an MEC of H_1 , it suffices to show that $\mathcal{V}(M_1) = \mathcal{V}(M[V_{H_1}])$ (from Definition 2.41). From the initialization step, all directed edges of M_1 are also directed edges of M . This implies $\mathcal{V}(M_1) \subseteq \mathcal{V}(M[V_{H_1}])$

as from the construction of M , $\text{skeleton}(M[V_{M_1}]) = \text{skeleton}(M_1)$. Thus, to establish $\mathcal{V}(M_1) = \mathcal{V}(M[V_{H_1}])$, we only need to demonstrate $\mathcal{V}(M[V_{H_1}]) \subseteq \mathcal{V}(M_1)$.

Suppose $\mathcal{V}(M[V_{H_1}]) \not\subseteq \mathcal{V}(M_1)$. This implies the existence of $u, v, w \in V_{H_1}$ such that $u \rightarrow v \leftarrow w \in M$ and $u \rightarrow v \leftarrow w \notin M_1$. From the construction of M , if $u \rightarrow v \in M$, then either $u \rightarrow v \in M_1$ or $u - v \in M_1$ (because if $v \rightarrow u \in M_1$ from line 1 of alg. 4, $v \rightarrow u \in M$). Similarly, if $w \rightarrow v \in M$, then either $w \rightarrow v \in M_1$ or $w - v \in M_1$. Since $u \rightarrow v \leftarrow w \notin M_1$, either $u \rightarrow v - w \in M_1$ or $u - v \leftarrow w \in M_1$ or $u - v - w \in M_1$. From item 3 of Theorem 2.18, the first two possibilities are not possible. Thus, the only remaining possibility is that $u - v - w$ is an induced subgraph of M . However, this contradicts Observation 3.46, as if $u \rightarrow v \in M$, then $v \rightarrow w \in M$, which contradicts the presence of $w \rightarrow v \in M$. This confirms that there are no such $u, v, w \in V_{H_1}$ for which $\mathcal{V}(M[V_{H_1}]) \not\subseteq \mathcal{V}(M_1)$.

Thus, we have established $\mathcal{V}(M[V_{H_1}]) \subseteq \mathcal{V}(M_1)$. Combined with the earlier inclusion that $\mathcal{V}(M_1) \subseteq \mathcal{V}(M[V_{H_1}])$, we conclude that $\mathcal{V}(M_1) = \mathcal{V}(M[V_{H_1}])$.

Hence, we have successfully shown that $\mathcal{P}(M, V_{H_1}) = M_1$, completing the proof of Lemma 3.37. \square

Proof of Lemma 3.38. Suppose there exists an MEC M' such that $M'[V_O] = O$, and $\mathcal{P}(M', V_{M_1}, V_{M_2}) = (M_1, M_2)$. We will show that $M' = M$.

We know that each MEC of a graph has a unique set of v-structures. Our aim is to demonstrate that both M and M' contain the same set of v-structures. This implies that $M = M'$.

From Lemma 3.35, we have $M[V_O] = O$. Also, from Lemma 3.37, we know that $\mathcal{P}(M, V_{M_1}, V_{M_2}) = (M_1, M_2)$. Since both M and M' are MECs of H , and $\text{skeleton}(H) = \text{skeleton}(H_1) \cup \text{skeleton}(H_2)$, it follows that for any v-structure $u \rightarrow v \leftarrow w$ that belongs to either M or M' , one of the following cases holds: (a) $u, v, w \in V_{M_1}$, (b) $u, v, w \in V_{M_2}$, (c) $v \in I = V_{M_1} \cap V_{M_2}$, $u \in V_{M_1} \setminus I$, and $w \in V_{M_2} \setminus I$, (d) $u \in V_{M_2} \setminus I$, $v \in I$, and $w \in V_{M_1} \setminus I$.

We now address each of these cases.

For case (a), if $u \rightarrow v \leftarrow w$ is a v-structure in M with $u, v, w \in V_{M_1}$, since M_1 is a projection of M on V_{M_1} , the v-structure $u \rightarrow v \leftarrow w$ also belongs to M_1 . Similarly, as M_1 is a projection of M' on V_{M_1} , the same v-structure belongs to M' . A similar argument can be made for v-structures in M' . The same argument holds for v-structures when $u, v, w \in V_{M_2}$ (i.e., for case (b)).

For case (c), if $u \rightarrow v \leftarrow w$ is a v-structure in M with $u \in V_{M_1} \setminus I$, $v \in I$, and $w \in V_{M_2} \setminus I$, this v-structure is in O as $I \cup N(I, H) \subseteq V_O$, and O is an induced subgraph of M . Since O is also an induced subgraph of M' , the v-structure also belongs to M' . A similar argument can be made for v-structures in M' with $u \in V_{M_1} \setminus I$, $v \in I$, and $w \in V_{M_2} \setminus I$. The same argument holds for v-structures when $u \in V_{M_2} \setminus I$, $v \in I$, and $w \in V_{M_1} \setminus I$ (i.e., for case (d)).

The above analysis demonstrates that a v-structure $u \rightarrow v \leftarrow w$ is present in M if, and only if, it is also present in M' . This implies that both M and M' have the same set of v-structures, and therefore, $M = M'$. This completes the proof of Lemma 3.38. \square

Proof of Lemma 3.39. Let $\text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22}) = (P_1, P_2)$, and (O', P'_1, P'_2) be the shadow of M on V_O . We aim to show $(O, P_1, P_2) = (O', P'_1, P'_2)$. By definition (Definition 2.36), $O' = M[V_O]$. Furthermore, from Lemma 3.35, $O = M[V_O]$. Thus, we conclude that $O = O'$.

We are given that the shadow of M_1 on $S_1 \cup N(S_1, H_1)$ is (O_1, P_{11}, P_{12}) , and the shadow of M_2 on $S_2 \cup N(S_2, H_2)$ is (O_2, P_{21}, P_{22}) . Also, we are given $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$. Furthermore, from Lemma 3.37, we have $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$. Therefore, based on Lemma 3.25, we conclude that $(P'_1, P'_2) = (P_1, P_2)$. This shows that (O, P_1, P_2) is the shadow of M on V_O , i.e., $M \in \text{MEC}(H, O, P_1, P_2)$. \square

3.5 Finding Size of $\text{MEC}(H, O, P_1, P_2)$

We use Definition 3.24 to define a one to one function f_{O, P_1, P_2} . We use the function for the computation of $|\text{MEC}(H, O, P_1, P_2)|$.

Definition 3.59. Let H be an undirected graph, and let H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Define

subsets S_1 and S_2 of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let $A = H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)]$, $B_1 = H_1[S_1 \cup N(S_1, H_1)]$, and $B_2 = H_2[S_2 \cup N(S_2, H_2)]$. Consider $(O, P_1, P_2) \in \text{shadow}(A)$. Define $T_1 = \text{MEC}(H, O, P_1, P_2)$ and $T_2 = \{(M_1, M_2) : \exists (O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22}) \text{ such that } (O_1, P_{11}, P_{12}) \in \text{shadow}(B_1), (O_2, P_{21}, P_{22}) \in \text{shadow}(B_2), (O, P_1, P_2) \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22}), M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12}), \text{ and } M_2 \in \text{MEC}(H_2, O_2, P_{21}, P_{22})\}$. The function $f_{O, P_1, P_2} : T_1 \rightarrow T_2$ is defined as $f_{O, P_1, P_2}(M) = (M_1, M_2)$ if $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$.

Lemma 3.60. f_{O, P_1, P_2} is bijective.

Proof. We first prove \rightarrow of Lemma 3.60.

Claim 3.61. For each $M \in \text{MEC}(H, O, P_1, P_2)$, there exists a unique (M_1, M_2) such that $f_{O, P_1, P_2}(M) = (M_1, M_2)$.

Proof. Let $M \in \text{MEC}(H, O, P_1, P_2)$. From Definition 2.38, M is an MEC of H and its shadow on V_O is (O, P_1, P_2) . Let $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$. From Corollary 2.43, there exists a unique (M_1, M_2) such that $\mathcal{P}(M, V_{H_1}, V_{H_2}) = (M_1, M_2)$. Thus, we only have to show that $(M_1, M_2) \in T_2$. Let $(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1)$ and $(O_2, P_{21}, P_{22}) \in \text{shadow}(B_2)$. Therefore, from Definition 2.38, $M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12})$ and $M_2 \in \text{MEC}(H_2, O_2, P_{21}, P_{22})$. From Lemma 3.25, $(O, P_1, P_2) \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$. From the definition of T_2 (given at Definition 3.59), $(M_1, M_2) \in T_2$. Thus, from Definition 3.59, $f_{O, P_1, P_2}(M) = (M_1, M_2)$. □

We now prove \leftarrow of Lemma 3.60.

Claim 3.62. Let (O_1, P_{11}, P_{12}) be a shadow of B_1 and (O_2, P_{21}, P_{22}) be a shadow of B_2 such that (O, P_1, P_2) is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$. For each $M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12})$ and $M_2 \in \text{MEC}(H_2, O_2, P_{21}, P_{22})$, there exists a unique $M \in \text{MEC}(H, O, P_1, P_2)$ such that $f_{O, P_1, P_2}(M) = (M_1, M_2)$.

Proof. Lemma 3.34 proves this. □

Claims 3.61 and 3.62 prove Lemma 3.60. □

Lemma 3.60 implies that the following:

Lemma 3.63. Let H be an undirected graph, and H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1 and S_2 be the subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let $A = H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)]$, $B_1 = H_1[S_1 \cup N(S_1, H_1)]$, and $B_2 = H_2[S_2 \cup N(S_2, H_2)]$. For any $(O, P_1, P_2) \in \text{shadow}(A)$,

$$|\text{MEC}(H, O, P_1, P_2)| = \sum_{\substack{(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1) \\ (O_2, P_{21}, P_{22}) \in \text{shadow}(B_2) \\ (O, P_1, P_2) \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})}} |\text{MEC}(H_1, O_1, P_{11}, P_{12})| \times |\text{MEC}(H_2, O_2, P_{21}, P_{22})| \quad (9)$$

Proof. From Lemma 3.60, $|T_1| = |T_2|$. Since $T_1 = \text{MEC}(H, O, P_1, P_2)$, therefore, $|T_1| = |\text{MEC}(H, O, P_1, P_2)|$. Since T_2 contains tuples (M_1, M_2) such that for some tuple $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$ such that $(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1)$, $(O_2, P_{21}, P_{22}) \in \text{shadow}(B_2)$, and $(O, P_1, P_2) \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, $M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12})$ and $M_2 \in \text{MEC}(H_2, O_2, P_{21}, P_{22})$. This implies

$$T_2 = \bigcup_{\substack{(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1) \\ (O_2, P_{21}, P_{22}) \in \text{shadow}(B_2) \\ (O, P_1, P_2) \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})}} \text{MEC}(H_1, O_1, P_{11}, P_{12}) \times \text{MEC}(H_2, O_2, P_{21}, P_{22}). \quad (10)$$

We show that sets $\text{MEC}(H_1, O_1, P_{11}, P_{12}) \times \text{MEC}(H_2, O_2, P_{21}, P_{22})$ are disjoint. Suppose sets $\text{MEC}(H_1, O_1, P_{11}, P_{12}) \times \text{MEC}(H_2, O_2, P_{21}, P_{22})$ are not disjoint. Then there exists (M_1, M_2) such that $(M_1, M_2) \in \text{MEC}(H_1, O_1, P_{11}, P_{12}) \times$

$\text{MEC}(H_2, O_2, P_{21}, P_{22})$ as well as $(M_1, M_2) \in \text{MEC}(H_1, O'_1, P'_{11}, P'_{12}) \times \text{MEC}(H_2, O'_2, P'_{21}, P'_{22})$. This implies $M_1 \in \text{MEC}(H_1, O_1, P_{11}, P_{12})$ as well as $M_1 \in \text{MEC}(H_1, O'_1, P'_{11}, P'_{12})$. But, from Definition 2.36, shadow of M on V_{B_1} is unique, i.e., $(O_1, P_{11}, P_{12}) = (O'_1, P'_{11}, P'_{12})$. Similarly, we can say that $(O_2, P_{21}, P_{22}) = (O'_2, P'_{21}, P'_{22})$. This shows $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22}) = (O'_1, P'_{11}, P'_{12}, O'_2, P'_{21}, P'_{22})$. This implies that the sets $\text{MEC}(H_1, O_1, P_{11}, P_{12}) \times \text{MEC}(H_2, O_2, P_{21}, P_{22})$ are disjoint. This further implies

$$|T_2| = \sum_{\substack{(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1) \\ (O_2, P_{21}, P_{22}) \in \text{shadow}(B_2) \\ (O, P_1, P_2) \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})}} |\text{MEC}(H_1, O_1, P_{11}, P_{12})| \times |\text{MEC}(H_2, O_2, P_{21}, P_{22})|. \quad (11)$$

This proves Lemma 3.63. \square

As discussed earlier, H resembles G_i^j , H_1 resembles G_i^{j-1} , H_2 resembles G_{i_j} , and S_1 and S_2 resemble X_i and X_{i_j} , respectively. Let $A = G_i^j[X_i \cup X_{i_j} \cup N(X_i \cup X_{i_j}, G_i^j)]$, $A' = G_i^j[X_i \cup N(X_i, G_i^j)]$, $B_1 = G_i^{j-1}[X_i \cup N(X_i, G_i^{j-1})]$, and $B_2 = G_{i_j}[X_{i_j} \cup N(X_{i_j}, G_{i_j})]$. Lemma 3.63 implies that if for all $(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1)$ and $(O_2, P_{21}, P_{22}) \in \text{shadow}(B_2)$, we have the knowledge about $|\text{MEC}(G_i^{j-1}, O_1, P_{11}, P_{12})|$ and $|\text{MEC}(G_{i_j}, O_2, P_{21}, P_{22})|$ then for each $(O, P_1, P_2) \in \text{shadow}(A)$, we can compute $|\text{MEC}(G_i^j, O, P_1, P_2)|$. But, for the recursion, we need $|\text{MEC}(H, O', P'_1, P'_2)|$ for each $(O', P'_1, P'_2) \in \text{shadow}(A')$. As (O, P_1, P_2) contains more information than (O', P'_1, P'_2) , we define projection of (O, P_1, P_2) to get $|\text{MEC}(H, O', P'_1, P'_2)|$ for each required (O', P'_1, P'_2) .

Definition 3.64 (Projection of a shadow). Let H be an undirected graph. S and S' are two vertex subsets of H such that $S' \subseteq S \subseteq V_H$. Let $(O, P_1, P_2) \in \text{shadow}(G[S])$ and $(O', P'_1, P'_2) \in \text{shadow}(G[S'])$. We say (O', P'_1, P'_2) is a projection of (O, P_1, P_2) on S' denoted as $\mathcal{P}(O, P_1, P_2, S') = (O', P'_1, P'_2)$ if

1. $O[S'] = O'$,
2. for $((u, v), (x, y)) \in E_{G[S']} \times E_{G[S']}$, $P'_1((u, v), (x, y)) = P_1((u, v), (x, y))$, and
3. for $((u, v), w) \in E_{G[S']} \times V_{G[S']}$, $P'_2((u, v), w) = P_2((u, v), w)$.

We now compute $|\text{MEC}(H, O', P'_1, P'_2)|$.

Lemma 3.65. Let H be an undirected graph, S and S' are two vertex subsets of H such that $S' \subseteq S \subseteq V_H$, $A = H[S]$, and $A' = H[S']$. Let $(O', P'_1, P'_2) \in \text{shadow}(A')$. Then,

$$|\text{MEC}(H, O', P'_1, P'_2)| = \sum_{\substack{(O, P_1, P_2) \in \text{shadow}(A) \\ (O', P'_1, P'_2) = \mathcal{P}(O, P_1, P_2, S')}} |\text{MEC}(H, O, P_1, P_2)| \quad (12)$$

Proof. We first show the following:

$$\text{MEC}(H, O', P'_1, P'_2) = \bigcup_{\substack{(O, P_1, P_2) \in \text{shadow}(A) \\ (O', P'_1, P'_2) = \mathcal{P}(O, P_1, P_2, S')}} \text{MEC}(H, O, P_1, P_2) \quad (13)$$

Let g be a function with L.H.S. of eq. (13) (i.e., $\text{MEC}(H, O', P'_1, P'_2)$) as its domain, and R.H.S. of eq. (13) as its co-domain. g is defined as $g(M) = M$.

Claim 3.66. g is bijective.

Proof. We first prove the \rightarrow of Claim 3.66.

Claim 3.67. For each $M \in \text{MEC}(H, O', P'_1, P'_2)$, there exists a unique $(O, P_1, P_2) \in \text{shadow}(A)$ such that $\mathcal{P}(O, P_1, P_2, S') = (O', P'_1, P'_2)$, and $M \in \text{MEC}(H, O, P_1, P_2)$.

Proof. $M \in \text{MEC}(H, O', P'_1, P'_2)$. From Definition 2.36, there exists a unique triple (O, P_1, P_2) such that $(O, P_1, P_2) \in \text{shadow}(A)$ and $M \in \text{MEC}(H, O, P_1, P_2)$. Only thing that we have to check is whether $\mathcal{P}(O, P_1, P_2) = (O', P'_1, P'_2)$ or not. From Definition 2.38, $O = M[S]$, and $O' = M[S']$. Since $S' \subseteq S$, $O[S'] = O'$. Similarly, since $S' \subseteq S$, $E_{A'} \times E_{A'} \subseteq E_A \times E_A$. This implies for all $((u, v), (x, y)) \in E_{A'} \times E_{A'}$, $P'_1((u, v), (x, y)) = P_1((u, v), (x, y))$ (as both answers whether there exists a triangle-free path from (u, v) to (x, y) in M). Similarly, for all $((u, v), w) \in E_{A'} \times V_{A'}$, $P'_2((u, v), w) = P_2((u, v), w)$. This further implies that $\mathcal{P}(O, P_1, P_2) = (O', P'_1, P'_2)$ (from Definition 3.64). This proves the claim. \square

We now prove the \leftarrow of Claim 3.66.

Claim 3.68. *Suppose there exists a triplet (O, P_1, P_2) such that $(O, P_1, P_2) \in \text{shadow}(A)$ and $\mathcal{P}(O, P_1, P_2, S') = (O', P'_1, P'_2)$. Then, for each $M \in \text{MEC}(H, O, P_1, P_2)$, $M \in \text{MEC}(H, O', P'_1, P'_2)$.*

Proof. Let $M \in \text{MEC}(H, O, P_1, P_2)$. From Definition 2.38, $M[S] = O$. Since $\mathcal{P}(O, P_1, P_2, S') = (O', P'_1, P'_2)$, $M[S'] = O[S'] = O'$. Since $S' \subseteq S$, $E_{A'} \times E_{A'} \subseteq E_A \times E_A$. This implies that for all $((u, v), (x, y)) \in E_{A'} \times E_{A'}$, $P'_1((u, v), (x, y)) = P_1((u, v), (x, y))$, as $\mathcal{P}(O, P_1, P_2, S') = (O', P'_1, P'_2)$. This implies for each $((u, v), (x, y)) \in E_{A'} \times E_{A'}$, P'_1 correctly depicts that whether there is a triangle-free path from (u, v) to (x, y) in M . Similarly, for all $((u, v), w) \in E_{A'} \times V_{A'}$, $P'_2((u, v), w)$ correctly depicts that whether there is a triangle-free path from (u, v) to w in M . The above discussion implies that (O', P'_1, P'_2) is the shadow of M on S' (from Definition 3.64). This implies that $M \in \text{MEC}(H, O', P'_1, P'_2)$. \square

Claims 3.67 and 3.68 prove Claim 3.66. \square

Claim 3.66 proves eq. (13). From Definition 2.36, an MEC M of H belongs to a unique $\text{MEC}(H, O, P_1, P_2)$. This implies that sets in the union in the R.H.S. of eq. (13) are disjoint. This further implies eq. (12). This completes the proof of Lemma 3.65. \square

Combining Lemmas 3.63 and 3.65, we get the following lemma:

Lemma 3.69. *Let H be an undirected graph, and H_1 and H_2 be two induced subgraphs of H such that $H = H_1 \cup H_2$, and $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. Let S_1 and S_2 be the subsets of V_{H_1} and V_{H_2} , respectively, such that $S_1 \cap S_2 = I$. Let $S = S_1 \cup S_2 \cup N(S_1 \cup S_2, H)$, and $S' = S_1 \cup N(S_1, H)$, $A = H[S]$, $A' = H[S']$, $B_1 = H_1[S_1 \cup N(S_1, H_1)]$, and $B_2 = H_2[S_2 \cup N(S_2, H_2)]$. Then, for any partial MEC $(O', P'_1, P'_2) \in \text{shadow}(A')$,*

$$|\text{MEC}(H, O', P'_1, P'_2)| = \sum_{\substack{(O, P_1, P_2) \in \text{shadow}(A) \\ (O_1, P_{11}, P_{12}) \in \text{shadow}(B_1) \\ (O_2, P_{21}, P_{22}) \in \text{shadow}(B_2) \\ (O, P_1, P_2) \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22}) \\ (O', P'_1, P'_2) = \mathcal{P}(O, P_1, P_2, S')}} |\text{MEC}(H_1, O_1, P_{11}, P_{12})| \times |\text{MEC}(H_2, O_2, P_{21}, P_{22})|. \quad (14)$$

4 Algorithm for counting MECs of a graph

Using Lemma 3.69, we can compute $|\text{MEC}(H, O', P'_1, P'_2)|$ for any shadow $(O', P'_1, P'_2) \in \text{shadow}(H[S_1 \cup N(S_1, H)])$. Since H resembles G_i^j , the above discussion implies that we can count the MECs of G_i^j . This gives us a recursive method to compute the MECs of G .

Based on the above discussion, we construct alg. 5 to count MECs of any undirected connected graph G . Alg. 5 takes an undirected graph G as its input, and returns $|\text{MEC}(G)|$ as the output.

Lemma 4.1. *Let G be an undirected connected graph. Given input G , alg. 5 returns $|\text{MEC}(G)|$.*

Algorithm 5: Count-MEC(G)

Input : An undirected connected graph G
Output : $|MEC(G)|$

- 1 $(\mathcal{X} = \{X_1, X_2, \dots, X_l\}, T) \leftarrow$ a tree decomposition of G .
- 2 $F = \text{Count}(G, (\mathcal{X}, T), X_1)$;
- 3 $sum \leftarrow 0$
- 4 $A \leftarrow G[X_1 \cup N(X_1, G)]$
- 5 **foreach** $(O, P_1, P_2) \in \text{shadow}(A)$ **do**
- 6 | $sum = sum + F(O, P_1, P_2)$
- 7 **end**
- 8 **return** sum

Algorithm 6: BRUTEFORCECOUNT(G)

Input : An undirected connected graph G
Output : A function $F : \text{shadow}(G) \rightarrow \mathbb{Z}$ such that for each $(O, P_1, P_2) \in \text{shadow}(G)$,
 $F(O, P_1, P_2) = |MEC(G, O, P_1, P_2)|$

- 1 $F : \text{shadow}(G) \rightarrow \mathbb{Z}$
- 2 **foreach** $(O, P_1, P_2) \in \text{shadow}(G)$ **do**
- 3 | $F(O, P_1, P_2) \leftarrow 0$
- 4 **end**
- 5 **foreach** $O \in MEC(G)$ **do**
- 6 | $(P_1, P_2) \leftarrow \text{TFP}(O)$
- 7 | $F(O, P_1, P_2) = 1$
- 8 **end**
- 9 **return** F

Algorithm 7: Count($G, (\mathcal{X}, T), R_1$)

Input : An undirected graph G , a tree decomposition (\mathcal{X}, T) of G , and a root node R_1 of T
Output : A function $F : \text{shadow}(G) \rightarrow \mathbb{Z}$ such that for each $(O, P_1, P_2) \in \text{shadow}(G[R_1 \cup N(R_1, G)])$,
 $F(O, P_1, P_2) = |\text{MEC}(G, O, P_1, P_2)|$

```
1 if  $\mathcal{X} = \{R_1\}$  then
2   | return BRUTEFORCECOUNT( $G$ )
3 else
4   |  $R_2 \leftarrow$  last child of  $R_1$  in  $T$ .
5   | Cut the edge  $R_1 - R_2$  of  $T$ .
6   |  $T_1 \leftarrow$  induced subtree of  $T$  containing  $R_1$ 
7   |  $T_2 \leftarrow$  induced subtree of  $T$  containing  $R_2$ 
8   |  $G_1 \leftarrow$  induced subtree of  $G$  represented by  $T_1$ 
9   |  $G_2 \leftarrow$  induced subtree of  $G$  represented by  $T_2$ 
10  |  $\mathcal{X}_1 \leftarrow$  subset of  $X$  containing nodes of  $T_1$ 
11  |  $\mathcal{X}_2 \leftarrow$  subset of  $X$  containing nodes of  $T_2$ 
12  |  $F_1 \leftarrow$  Count( $G_1, (\mathcal{X}_1, T_1), R_1$ )
13  |  $F_2 \leftarrow$  Count( $G_2, (\mathcal{X}_2, T_2), R_2$ )
14  |  $A' \leftarrow G[R_1 \cup N(R_1, G)]$ 
15  |  $A \leftarrow G[R_1 \cup R_2 \cup N(R_1 \cup R_2, G)]$ 
16  |  $B_1 \leftarrow G_1[R_1 \cup N(R_1, G_1)]$ 
17  |  $B_2 \leftarrow G_2[R_2 \cup N(R_2, G_2)]$ 
18  |  $F : \text{shadow}(A') \rightarrow \mathbb{Z}$ 
19  | foreach  $(O', P'_1, P'_2) \in \text{shadow}(A')$  do
20  |   |  $F(O', P'_1, P'_2) \leftarrow 0$ 
21  |   end
22  | foreach  $O \in \text{PMEC}(A)$  do
23  |   | foreach  $(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1)$  do
24  |     | foreach  $(O_2, P_{21}, P_{22}) \in \text{shadow}(B_2)$  do
25  |       | if  $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$  then
26  |         |   |  $(P_1, P_2) \leftarrow \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$ 
27  |         |   |  $(O', P'_1, P'_2) \leftarrow \mathcal{P}(O, P_1, P_2, R_1 \cup N(R_1, G))$ .
28  |         |   |  $F(O', P'_1, P'_2) = F(O', P'_1, P'_2) + F_1(O_1, P_{11}, P_{12}) \times F_2(O_2, P_{21}, P_{22})$ 
29  |         |   end
30  |         | end
31  |         | end
32  |         end
33  |       end
34 end return  $F$ 
```

Algorithm 8: Is.Extension

Input : An MECs O , and two triples (O_1, P_{11}, P_{12}) and (O_2, P_{21}, P_{22}) such that H is an undirected graph, H_1 and H_2 are induced subgraphs of H such that $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H that separates $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$, for $a \in \{1, 2\}$, $S_a \subseteq V_{H_a}$ such that $S_1 \cap S_2 = I$, for $a \in \{1, 2\}$, $(O_a, P_{a1}, P_{a2}) \in \text{shadow}(H_a[S_a \cup N(S_a, H_a)])$, and $O \in \text{PMEC}(H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)])$

Output : 1 : if $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$,
0 : otherwise.

```
1 foreach  $a \in \{1, 2\}$  do
2   | if  $\exists u \rightarrow v \in O_a$  such that  $u \rightarrow v \notin O$  then
3   |   | return 0.
4   | end
5 end
6 foreach  $a \in \{1, 2\}$  do
7   | foreach  $u, v, w \in V_{O_a}$  do
8   |   | if  $u \rightarrow v \leftarrow w$  is a  $v$ -structure in  $O$  but not a  $v$ -structure in  $O_a$  then
9   |   |   | return 0.
10  |   | end
11  | end
12 end
13 foreach  $a \in \{1, 2\}$  do
14  | foreach  $u - v \in O_a$  do
15  |   | if  $u \rightarrow v \in O$  then
16  |   |   | if  $u \rightarrow v$  is not strongly protected in  $O$  then
17  |   |   |   | if  $\nexists x - y \in O_a$  such that  $x \rightarrow y \in O$  and  $P_{a1}((x, y), (u, v)) = 1$  then
18  |   |   |   |   | if  $\nexists x - y \in O_a$  such that  $x \rightarrow y \in O$  and  $P_{a2}((x, y), v) = P_{a2}((v, u), x) = 1$  then
19  |   |   |   |   |   | return 0.
20  |   |   |   |   | end
21  |   |   |   | end
22  |   |   | end
23  |   | end
24  |   | if  $u - v \in O$  then
25  |   |   | if  $\exists x - y \in O_a$  such that  $x \rightarrow y \in O$  and  $P_{a1}((x, y), (u, v)) = 1$  then
26  |   |   |   | return 0.
27  |   |   |   | end
28  |   |   | if  $\exists x - y \in O_a$  such that  $x \rightarrow y \in O$  and  $P_{a2}((x, y), v) = P_{a2}((v, u), x) = 1$  then
29  |   |   |   | return 0.
30  |   |   |   | end
31  |   |   | end
32  |   | end
33 end
34  $(P_1, P_2) \leftarrow \text{DPF}(O, P_{11}, P_{12}, P_{21}, P_{22})$ 
35 foreach  $((u, v), (x, y)) \in E_O \times E_O$  do
36  | if  $(u, v) \neq (x, y)$  and  $P_1((u, v), (x, y)) = P_1((x, y), (u, v)) = 1$  then
37  |   | return 0
38  | end
39 end
40 return 1.
```

Algorithm 9: Projection(O, P_1, P_2, X)

Input : A partial MEC O , two functions P_1 and P_2 , and $X \subseteq V_O$ such that
 $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ and $P_2 : E_O \times V_O \rightarrow \{0, 1\}$

Output: $\mathcal{P}(O, P_1, P_2, X)$

```
1  $O' \leftarrow O[X]$ 
2  $P'_1 : E_{O'} \times E_{O'} \rightarrow \{0, 1\}$ 
3  $P'_2 : E_{O'} \times X \rightarrow \{0, 1\}$ 
4 foreach  $((u, v), (x, y)) \in E_{O'} \times E_{O'}$  do
5   |  $P'_1((u, v), (x, y)) \leftarrow P_1((u, v), (x, y))$ 
6 end
7 foreach  $((u, v), w) \in E_{O'} \times V_{O'}$  do
8   |  $P'_2((u, v), w) \leftarrow P_2((u, v), w)$ 
9 end
10 return  $(O', P'_1, P'_2)$ 
```

Proof. Alg. 5 constructs a tree decomposition $(\mathcal{X} = \{X_1, X_2, \dots, X_l\}, T)$ of G (at line 1 of alg. 5). It then invokes alg. 7 with input G , (\mathcal{X}, T) , and X_1 . Let $A = G[X_1 \cup N(X_1, G)]$. Lemma 4.2 demonstrates that alg. 7 returns a function F with domain $\text{shadow}(A)$ and co-domain \mathbb{Z} such that for each triplet $(O, P_1, P_2) \in \text{shadow}(A)$, $F(O, P_1, P_2) = |\text{MEC}(G, O, P_1, P_2)|$. Lines 5–7 of alg. 5 implement Lemma 2.40 to obtain $|\text{MEC}(G)|$. This confirms the validity of Lemma 4.1. \square

Lemma 4.2. *Let G be an undirected graph, (\mathcal{X}, T) be a tree decomposition of G , $R_1 \in \mathcal{X}$ be the root node of T , and $A' = G[R_1 \cup N(R_1, G)]$. For input G , (\mathcal{X}, T) , and R_1 , alg. 7 returns a function F with domain $\text{shadow}(A')$ and co-domain \mathbb{Z} such that for any triplet $(O, P_1, P_2) \in \text{shadow}(A')$, $F(O, P_1, P_2) = |\text{MEC}(G, O, P_1, P_2)|$.*

Proof. If the tree T has only one node, the root node R_1 , i.e., $\mathcal{X} = \{R_1\}$, then alg. 7 calls alg. 6. Lemma 4.3 shows that alg. 6 returns the required function F . If T has more than one node, we pick the last child R_2 of R_1 in T (line-4 of alg. 7), and cut the edge $R_1 - R_2$ of T (line-5 of alg. 7). This gives us two induced subgraphs T_1 and T_2 of T such that T_1 contains R_1 , and T_2 contains R_2 (lines 6-7), and two induced subgraphs G_1 and G_2 of G such that T_1 represents G_1 , and T_2 represents G_2 . The two trees T_1 and T_2 have sets of nodes \mathcal{X}_1 and \mathcal{X}_2 , respectively (lines 10-11 of alg. 7). We define $A = G[R_1 \cup R_2 \cup N(R_1 \cup R_2, G)]$, $A' = G[R_1 \cup N(R_1, G)]$, $B_1 = G_1[R_1 \cup N(R_1, G)]$, and $B_2 = G_2[R_2 \cup N(R_2, G)]$ for simplification of our analysis (lines 14-17 of alg. 7). We observe that $G = G_1 \cup G_2$, and $R_1 \cap R_2$ is a vertex separator of G (from Property 2.10). We find that this scenario resembles what we discussed after Observation 3.1. Here, G resembles H , G_1 and G_2 resemble H_1 and H_2 respectively, R_1 and R_2 resemble S_1 and S_2 . Thus, to implement Lemma 3.69, we need the functions F_1 and F_2 such that for any triplet $(O_1, P_{11}, P_{12}) \in \text{shadow}(B_1)$, $F_1(O_1, P_{11}, P_{12}) = |\text{MEC}(G_1, O_1, P_{11}, P_{12})|$, and for any triplet $(O_2, P_{21}, P_{22}) \in \text{shadow}(B_2)$, $F_2(O_2, P_{21}, P_{22}) = |\text{MEC}(G_2, O_2, P_{21}, P_{22})|$. Recursively, we compute these two functions F_1 and F_2 (lines 12-13 of alg. 7). After obtaining F_1 and F_2 , we implement Lemma 3.69 (lines 22-32) to compute the required function F . \square

Lemma 4.3. *Let G be an undirected graph. For input G , alg. 6 returns a function F such that for any triplet (O, P_1, P_2) where $(O, P_1, P_2) \in \text{shadow}(G)$, and $F(O, P_1, P_2) = |\text{MEC}(G, O, P_1, P_2)|$.*

Proof. We initialize the output function F with $F(O, P_1, P_2) = 0$ for all triplets $(O, P_1, P_2) \in \text{shadow}(G)$ (lines 2–4). For each MEC O of G , we compute $\text{TFP}(O)$ and update F with $F(O, P_1, P_2) = 1$ (lines 6 and 7). According to Lemma 2.35, $\text{TFP}(O)$ provides a pair of functions (P_1, P_2) such that

1. $P_1 : E_O \times E_O \rightarrow \{0, 1\}$, and $P_1((u, v), (x, y)) = 1$ if and only if $(u, v) \neq (x, y)$ and there exists a triangle-free path exists from (u, v) to (x, y) in O , and
2. $P_2 : E_O \times V_O \rightarrow \{0, 1\}$, and $P_2((u, v), w) = 1$ if and only if $v \neq w$ and a triangle-free path exists from (u, v) to w in O .

This implies that for any triple $(O, P_1, P_2) \in \text{shadow}(G)$, $F(O, P_1, P_2) = 1$ if O is an MEC of G and (O, P_1, P_2) is the shadow of O on V_G ; otherwise, $F(O, P_1, P_2) = 0$. We now demonstrate that for each $(O, P_1, P_2) \in \text{shadow}(G)$, $F(O, P_1, P_2) = |\text{MEC}(G, O, P_1, P_2)|$. According to Definition 2.38, $|\text{MEC}(G, O, P_1, P_2)|$ is the count of MECs M such that (O, P_1, P_2) is the shadow of M on V_O . Since for any partial MEC $O \in \text{PMEC}(G)$, $V_O = V_G$, if $M \in \text{MEC}(G, O, P_1, P_2)$, then $M = O$ and $(P_1, P_2) = \text{TFP}(O)$. This implies that if O is not an MEC of G , then $|\text{MEC}(G, O, P_1, P_2)| = 0$ regardless of P_1 and P_2 , and if O is an MEC of G , then if $(P_1, P_2) = \text{TFP}(O)$, then $|\text{MEC}(G, O, P_1, P_2)| = 1$, else $|\text{MEC}(G, O, P_1, P_2)| = 0$. This demonstrates that for each $(O, P_1, P_2) \in L$, $F(O, P_1, P_2) = |\text{MEC}(G, O, P_1, P_2)|$. This completes the proof of Lemma 4.3. \square

Lemma 4.4. *Let O be a partial MEC, $G = \text{skeleton}(O)$, $X \subseteq V_G$, and $P_1 : E_G \times E_G \rightarrow \{0, 1\}$ and $P_2 : E_G \times V_G \rightarrow \{0, 1\}$ be two functions. For the input O, P_1, P_2 , and X , alg. 9 outputs $\mathcal{P}(O, P_1, P_2, X)$.*

Proof. Let alg. 9 return (O', P'_1, P'_2) . As per line 1 in Definition 2.41, $O' = O[X]$. This demonstrates that O satisfies item 1 in Definition 3.64. Lines 4–6 of alg. 9 demonstrate that P'_1 satisfies item 2 in Definition 3.64. Similarly, lines 7–9 of alg. 9 show that P'_2 satisfies item 3 in Definition 3.64. Thus, according to Definition 3.64, $(O', P'_1, P'_2) = \mathcal{P}(O, P_1, P_2, X)$. \square

Lemma 4.5. *Let H be an undirected graph, H_1 and H_2 be two induced subgraphs of H such that $I = V_{H_1} \cap V_{H_2}$ is a vertex separator of H , separating $V_{H_1} \setminus I$ and $V_{H_2} \setminus I$. For $a \in \{1, 2\}$, let $S_a \subseteq V_{H_a}$ such that $S_1 \cap S_2 = I$. For each $a \in \{1, 2\}$, let $(O_a, P_{a1}, P_{a2}) \in \text{shadow}(H_a[S_a \cup N(S_a, H_a)])$. Additionally, let $O \in \text{PMEC}(H[S_1 \cup S_2 \cup N(S_1 \cup S_2, H)])$.*

For the input $O, O_1, O_2, P_{11}, P_{12}, P_{21}$, and P_{22} , alg. 8 returns 1 if $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, otherwise it returns 0.

Proof. By the definition of extension, Definition 3.24, if $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$, then O satisfies items 1 to 4 of Definition 3.24.

Lines 1–5 handle cases where O does not satisfy item 1 of Definition 3.24, causing alg. 8 to return 0.

If O satisfies lines 1–5 then each directed edge in O_1 and O_2 is also a directed edge in O . This implies for $a \in \{1, 2\}$, $\mathcal{V}(O_a) \subseteq \mathcal{V}(O[V_{O_a}])$, since from the construction, $\text{skeleton}(O_a) = \text{skeleton}(O[V_{O_a}])$. Lines 6–12 deal with the verification of $\mathcal{V}(O[V_{O_a}]) \subseteq \mathcal{V}(O_a)$. If O does not satisfy $\mathcal{V}(O[V_{O_a}]) \subseteq \mathcal{V}(O_a)$, then alg. 8 returns 0. Therefore, if O passes lines 1–12 and moves ahead then O satisfies item 2 of Definition 3.24.

Lines 13–33 cover cases where O does not satisfy item 3 of Definition 3.24, causing alg. 8 to return 0.

Lines 15–23 and lines 24–31 check the conditions for item 3 of Definition 3.24.

Lines 35–39 address cases where O does not satisfy item 4 of Definition 3.24, leading alg. 8 to return 0.

This implies that if alg. 8 returns 1 for the inputs $O, O_1, O_2, P_{11}, P_{12}, P_{21}, P_{22}$, then O satisfies items 1 to 4 of Definition 3.24. By Definition 3.24, this further implies that $O \in \mathcal{E}(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$. This concludes the proof of Lemma 4.5. \square

5 Time Complexity

Lemma 5.1. *For an input chain graph G with chordal undirected connected components, the time complexity of alg. 1 is $O(|E_G|^5)$.*

Proof. The running time of lines 3–5 in alg. 1 is $O(|E_G|^2)$, as the first foreach loop iterates $|E_G|^2$ times, and each iteration takes $O(1)$ time. Similarly, the running time of lines 6–8 is $O(|V_G| \cdot |E_G|)$, since the second foreach loop iterates $|V_G| \cdot |E_G|$ times, and each iteration takes $O(1)$ time. Furthermore, the running time of lines 9–14 is $O(|V_G|^3)$, as the third foreach loop iterates $|V_G|^3$ times, and each iteration takes $O(1)$ time.

In each iteration of the while loop at lines 15–17, the algorithm updates P_1 by setting $P_1((u, v), (x, y)) = 1$ for a pair of edges (u, v) and (x, y) where $P_1((u, v), (x, y)) = 0$ at the iteration's beginning. As the number of such pairs is at most $|E_G|^2$, the while loop iterates at most $|E_G|^2$ times. Each iteration takes $O(|E_G|^3)$ time to find three edges $(u, v), (x, y), (z_1, z_2)$ satisfying conditions such as $P_1((u, v), (x, y)) = 0$, $(u, v) \neq (x, y)$, and $P_1((u, v), (z_1, z_2)) = P_1((z_1, z_2), (x, y)) = 1$. This implies that the running time of lines 15–17 is $O(|E_G|^5)$.

Similarly, in each iteration of the while loop at lines 18–20, the algorithm updates P_2 by setting $P_2((u, v), w) = 1$ for an edge (u, v) and a node w , provided that $P_2((u, v), w) = 0$ at the start of the iteration. The number of such edge-node pairs can be at most $|V_G| \cdot |E_G|$, causing the while loop to iterate at most $|V_G| \cdot |E_G|$ times. Each iteration takes $O(|V_G| \cdot |E_G|^2)$ time to find two edges (u, v) and (z_1, z_2) and a node w satisfying conditions such as $P_2((u, v), w) = 0$, $v \neq w$, and $P_1((u, v), (z_1, z_2)) = P_2((z_1, z_2), w) = 1$. Thus, the running time complexity of lines 18–20 is $O(|V_G|^2 \cdot |E_G|^3)$.

Combining the above, the total running time complexity of the algorithm is $O(|E_G|^5)$. \square

Lemma 5.2. *For the input elements $O, P_{11}, P_{12}, P_{21}$, and P_{22} , the time complexity of alg. 2 is $O(|E_O|^5)$.*

Proof. Line 1 of alg. 2 takes $O(|E_O|^5)$ time as it calls alg. 1. Lines 3–7 run for $O(|E_O|^2)$ time, where the foreach loop at line 3 iterates $O(|E_O|^2)$ times, with each iteration taking $O(1)$ time. Similarly, lines 8–11 run for $O(|V_O| \cdot |E_O|)$ time. The foreach loop at line 8 iterates for $O(|V_O| \cdot |E_O|)$ time, and each iteration takes $O(1)$ time. Thus, the overall running time of lines alg. 2–13 is $O(|E_O|^2)$.

Similar to alg. 1, the running time complexity of lines 14–17 and lines 18–21 is $O(|E_O|^5)$ and $O(|V_O|^2 \cdot |E_O|^3)$, respectively. Therefore, the time complexity of alg. 2 is $O(|E_O|^5)$. \square

Theorem 5.3. *For an input undirected graph G , the time complexity of alg. 5 is $O(n(2^{O(k^4 \cdot \delta^4)} + n^2))$, where δ represents the degree of G , and k denotes the treewidth of G .*

Proof. For a graph of treewidth k , we can utilize Korhonen’s algorithm (Korhonen (2022)) to construct a tree decomposition (\mathcal{X}, T) of width $k \leq d \leq 2k + 1$ at line 1 of alg. 5. Korhonen provides an algorithm (Korhonen (2022)) that, for a given n -vertex graph G of treewidth k , outputs a tree decomposition of G of width d such that $k \leq d \leq 2k + 1$ in time $2^{O(k)} \cdot n$.

At line 2, we compute the required function F using alg. 7. From Lemma 5.5, it takes $O(n(2^{O(d^4 \cdot \delta^4)} + n^2))$ time. Since the width of the tree decomposition is d , the size of X_1 is $d + 1$. Also, as the degree of the graph is δ , the number of nodes in $X_1 \cup N(X_1, G)$ is at most $(1 + \delta)(1 + d)$. Therefore, the size of A (at line alg. 5) is $O(d^2 \delta^2)$, and the number of partial MECs of A is $O(3^{d^2 \delta^2})$, as for each edge $u - v \in A$, in a partial $O \in \text{PMEC}(A)$, there are three possibilities: either $u \rightarrow v \in O$ or $v \rightarrow u \in O$ or $u - v \in O$.

For a partial MEC O of A , (a) the number of distinct functions $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ is at most $2^{|E_O|^2} = 2^{O(d^4 \delta^4)}$, and (b) the number of distinct functions $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ is at most $2^{|E_O| \cdot |V_O|} = 2^{O(d^3 \delta^3)}$.

This implies the number of shadows $(O, P_1, P_2) \in \text{shadow}(A)$ is $O(2^{O(d^4 \delta^4)})$. This implies that the foreach loop at line 5 runs for $O(2^{O(d^4 \delta^4)})$ iterations. Each iteration of the foreach loop (line 6) takes $O(1)$ time. Thus, the total running time of lines 5–7 is $O(2^{O(d^4 \delta^4)})$.

Since $k \leq d \leq 2k + 1$, the total running time of the algorithm is $O(n(2^{O(k^4 \cdot \delta^4)} + n^2))$. \square

Lemma 5.4. *For an input undirected graph G , the time complexity of alg. 6 is $O(2^{O(|E_G|^2)})$.*

Proof. For any edge $u - v \in E_G$, in a partial MEC O of G , there are three possibilities: either $u - v \in O$, or $u \rightarrow v \in O$, or $v \rightarrow u \in O$. Therefore, the number of partial MECs of G is at most $3^{|E_G|}$. For any partial MEC O of G , the number of different functions $P_1 : E_O \times E_O \rightarrow \{0, 1\}$ is $2^{|E_O|^2} = O(2^{|E_G|^2})$. Similarly, for any partial MEC O of G , the number of different functions $P_2 : E_O \times V_O \rightarrow \{0, 1\}$ is $2^{|E_O| \cdot |V_O|} = O(2^{|E_G| \cdot |V_G|})$. This implies the number of shadows of G is $O(2^{|E_G|^2})$. Therefore, the construction of F and its initialization (lines 1–4 of alg. 6) takes $O(2^{O(|E_G|^2)})$ time.

As discussed above, the number of partial MECs with skeleton G is $3^{|E_G|}$. Since each MEC is a partial MEC, therefore the number of MECs with skeleton G is also $3^{|E_G|}$. Therefore, the number of iterations in the second foreach loop (lines 5–8) is at most $3^{|E_G|}$. Verifying whether O is an MEC or not, at line 5, will take $O(|V_G| + |E_G|)$ time, as verifying each condition of items 1 to 4 will take $O(|V_G| + |E_G|)$ time. From Lemma 5.1, the runtime of line 6 is $O(|E_G|^5)$. Therefore, the runtime of the second foreach loop (lines 5–8) is $O(3^{|E_G|} \times |E_G|^5)$. Thus, the time complexity of alg. 6 is $O(2^{O(|E_G|^2)})$. \square

Lemma 5.5. *For the input elements $G, (\mathcal{X}, T)$, and R_1 , the time complexity of alg. 7 is $O(n(2^{O(d^4 \cdot \delta^4)} + n^2))$, where d is the width of the tree decomposition (\mathcal{X}, T) of G and δ is the degree of G .*

Proof. If $\mathcal{X} = \{R_1\}$, then the tree T has only one node R_1 , and $V_G = R_1$. If the width of T is d , then $|V_G| = d + 1$, and $|E_G| = O(d^2)$. From Lemma 5.4, the time complexity to compute `BRUTEFORCECOUNT`(G) is $O(2^{O(d^4)})$. Therefore, the runtime of lines 1–3 is $O(2^{O(d^4)})$.

The time complexity to run lines 4–11 and lines 14–17 is $O(|V_G| + |E_G|)$. If the width of T is d , then the size of each R_1 and R_2 is $d + 1$. And, if the degree of G is δ , then the size of each $R_1 \cup N(R_1, G_1)$, $R_2 \cup N(R_2, G_2)$, $R_1 \cup N(R_1, G)$, and $R_1 \cup R_2 \cup N(R_1 \cup R_2, G)$ is $O(d\delta)$. This implies that the number of edges in A, A', B_1 , and B_2 is $O(d^2\delta^2)$. Therefore, as explained in Theorem 5.3, the size of each $\text{shadow}(A)$, $\text{shadow}(A')$, $\text{shadow}(B_1)$, and $\text{shadow}(B_2)$ is $O(2^{O(d^4\delta^4)})$.

The initialization of F at lines 19–21 will take $O(2^{O(d^4\delta^4)})$ as the size of L' is $O(2^{O(d^4\delta^4)})$.

Since the number of different PMECs of A is $O(3^{O(d^2\delta^2)})$, the for loop at line 22 runs for $O(3^{O(d^2\delta^2)})$ iterations. Since the sizes of $\text{shadow}(B_1)$ and $\text{shadow}(B_2)$ are $O(2^{O(d^4\delta^4)})$ (as computed in Lemma 5.4), the for loops at lines 23 and 24 run for $O(2^{O(d^4\delta^4)})$ iterations. From Lemma 5.6, the runtime of verifying whether O is an extension of $(O_1, P_{11}, P_{12}, O_2, P_{21}, P_{22})$ or not (at line 25) is $O(|E_O|^5) = O(d^{10}\delta^{10})$ (as discussed above the number of edges in O is $O(d^2\delta^2)$). From Lemma 5.2, the runtime of line 26 is $O(|E_O|^5) = O(d^{10}\delta^{10})$ time. Therefore, the overall runtime of lines 22–32 takes $O(2^{O(d^4\delta^4)} + |V_G| + |E_G|)$ time.

At lines 12–13, `alg. 7` calls itself. Each time `alg. 7` calls itself, it cuts one of the edges of T and breaks it into two disjoint parts T_1 and T_2 . Since the number of edges in the tree decomposition T is $O(|V_G|)$ ([Bodlaender and Kloks \(1996\)](#)), the number of times `alg. 7` calls itself is $O(|V_G|)$. This implies that the time complexity to run `alg. 7` is $O(|V_G| \cdot (2^{O(d^4\delta^4)} + |V_G| + |E_G|)) = O(n(2^{O(d^4\delta^4)} + n^2))$. This completes the proof. \square

Lemma 5.6. *For the input elements $O, (O_1, P_{11}, P_{12})$, and (O_2, P_{21}, P_{22}) , the time complexity of `alg. 8` is $O(|E_O|^5)$.*

Proof. Considering the input constraint provided at `alg. 8`, the size of O_1 and O_2 cannot exceed the size of O (since $\text{skeleton}(O_1)$ and $\text{skeleton}(O_2)$ are induced subgraphs of $\text{skeleton}(O)$). Therefore, the time complexity to run lines 1–5 is $O(|E_O|)$. And, the time complexity to run lines 6–12 is $O(|V_O|^3)$. The verification of an edge of O to check if it is strongly protected (line 16) will take cumulatively $O(|V_O| + |E_O|)$ time. The verification of the existence of an edge $x - y \in O_a$ such that $x \rightarrow y \in O$ and $P_{a1}((x, y), (u, v)) = 1$ (lines 17 and 25) takes $O(|E_O|)$ time. The verification of the existence of an edge $x - y \in O_a$ such that $x \rightarrow y \in O$ and $P_{a2}((x, y), v) = P_{a2}((v, u), x) = 1$ (lines 18 and 28) takes $O(|E_O|)$ time. The foreach loop at lines 7–11 runs for $O(|E_O|)$ iterations. Therefore, the time complexity of running lines 7–11 is $O(|E_O|^2)$ time. From Lemma 5.2, the runtime to construct a derived path function (P_1, P_2) at line 34 is $O(|E_O|^5)$ time. The foreach loop present at lines 35–39 runs for $O(|E_O|^2)$ iterations. Each iteration of the foreach loop takes $O(1)$ time. Therefore, the running of lines 35–39 takes $O(|E_O|^2)$ time. The above discussion implies that the time complexity of `alg. 8` is $O(|E_O|^5)$. \square

Lemma 5.7. *For the input elements O, P_1, P_2 , and X , the time complexity of `alg. 9` is $O(|E_O|^2)$.*

Proof. The construction of the induced subgraph O' at line 1 takes $O(|V_O| + |E_O|)$ time. The construction of the functions P'_1 and P'_2 at lines 2 and 3, respectively, takes $O(|E_O|^2)$ and $O(|V_O| \cdot |E_O|)$ time. The number of iterations the for loop at line 4–6 runs is $O(|E_{O'}|^2)$, and the running time of each iteration is $O(1)$. Similarly, the number of iterations the for loop at line 7–9 runs is $O(|V_{O'}| \cdot |E_{O'}|)$, and the running time of each iteration is $O(1)$. Therefore, the running time of lines 4–6 and lines 7–9 is $O(|E_{O'}|^2)$ and $O(|V_{O'}| \cdot |E_{O'}|)$ time, respectively. Since O' is an induced subgraph of O , it follows that $|E_O| \geq |E_{O'}|$ and $|V_O| \geq |V_{O'}|$. This further implies that the time complexity of `alg. 9` is $O(|E_O|^2)$. \square

Theorem 5.3 implies the following:

Theorem 5.8. *For an undirected connected graph G , there exists an algorithm that counts the number of MECs with skeleton G in $O(n(2^{O(k^4\delta^4)} + n^2))$ time, where k and δ are the treewidth and the degree of G , respectively.*

6 Conclusion and Open Problems

We provide a fixed parameter tractable algorithm with runtime $O(n(2^{O(k^4\delta^4)} + n^2))$ for the problem of counting the number of MECs with a given skeleton, where n is the number of nodes in the input graph and k and δ are the treewidth and the degree of the input graph.

The main problem left open by this work is to either provide a fully polynomial time algorithm for this problem, or else to prove that it is computationally hard in general. An intermediate open problem towards the goal of understanding the complexity fully is to improve the runtime of the fixed-parameter tractable algorithm itself: perhaps using different parameters.

As stated in the introduction, we consider our results to be a first step towards understanding the computational complexity of this problem. However, we hope that the definitions and techniques introduced in this paper will be helpful in further study of the problem.

References

- Andersson, S. A.; Madigan, D.; and Perlman, M. D. 1997. A Characterization of Markov Equivalence Classes for Acyclic Digraphs. *Annals of Statistics*, 25(2): 505–541.
- Bodlaender, H. L.; and Kloks, T. 1996. Efficient and constructive algorithms for the pathwidth and treewidth of graphs. *Journal of Algorithms*, 21(2): 358–402.
- Cygan, M.; Fomin, F. V.; Kowalik, L.; Lokshantov, D.; Marx, D.; Pilipczuk, M.; Pilipczuk, M.; and Saurabh, S. 2015. *Parameterized Algorithms*. Springer International Publishing.
- Diestel, R. 2005. Graph theory 3rd ed. *Graduate texts in mathematics*, 173(33): 12.
- Dirac, G. A. 1961. On Rigid Circuit Graphs. In *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, volume 25, 71–76. Springer.
- Finegold, M.; and Drton, M. 2011. Robust Graphical Modeling of Gene Networks using Classical and Alternative t -Distributions. *Annals of Applied Statistics*, 1057–1080.
- Friedman, N. 2004. Inferring Cellular Networks using Probabilistic Graphical Models. *Science*, 303(5659): 799–805.
- Ghassami, A.; Salehkaleybar, S.; Kiyavash, N.; and Zhang, K. 2019. Counting and Sampling from Markov Equivalent DAGs using Clique Trees. In *Proceedings of the 33rd AAAI Conference on Artificial Intelligence (AAAI 2019)*, volume 33, 3664–3671.
- Gillispie, S. B. 2006. Formulas for counting acyclic digraph Markov equivalence classes. *Journal of Statistical Planning and Inference*, 136(4): 1410–1432.
- Gillispie, S. B.; and Perlman, M. D. 2001. Enumerating Markov equivalence classes of acyclic digraph models. *arXiv preprint [arXiv:1301.2272](https://arxiv.org/abs/1301.2272)*.
- Gillispie, S. B.; and Perlman, M. D. 2002. The Size Distribution for Markov Equivalence Classes of Acyclic Digraph Models. *Artificial Intelligence*, 141(1-2): 137–155.
- He, Y.; Jia, J.; and Yu, B. 2013. Reversible MCMC on Markov equivalence classes of sparse directed acyclic graphs. *The Annals of Statistics*, 41(4): 1742–1779. Publisher: Institute of Mathematical Statistics.
- Kearns, M.; Littman, M. L.; and Singh, S. 2001. Graphical Models for Game Theory. In *Proceedings of the 17th Conference on Uncertainty in Artificial Intelligence conference (UAI 2001)*, 253–260.
- Korhonen, T. 2022. A single-exponential time 2-approximation algorithm for treewidth. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, 184–192. IEEE.

- Meek, C. 1995. Causal Inference and Causal Explanation with Background Knowledge. In *Proceedings of the 11th Conference on Uncertainty in Artificial Intelligence (UAI 1995)*, 403–410.
- Pearl, J. 2009. *Causality: Models, Reasoning and Inference*. Cambridge University Press.
- Radhakrishnan, A.; Solus, L.; and Uhler, C. 2016. Counting Markov Equivalence Classes by Number of Immoralities. *arXiv:1611.07493*.
- Radhakrishnan, A.; Solus, L.; and Uhler, C. 2018. Counting Markov equivalence classes for DAG models on trees. *Discrete Applied Mathematics*, 244: 170–185.
- Ren, Z.; Oviedo, F.; Thway, M.; Tian, S. I. P.; Wang, Y.; Xue, H.; Dario Perea, J.; Layurova, M.; Heumueller, T.; Birgersson, E.; Aberle, A. G.; Brabec, C. J.; Stangl, R.; Li, Q.; Sun, S.; Lin, F.; Peters, I. M.; and Buonassisi, T. 2020. Embedding Physics Domain Knowledge into a Bayesian Network Enables Layer-by-Layer Process Innovation for Photovoltaics. *NPJ Computational Materials*, 6(1): 9.
- Rose, D. J.; Tarjan, R. E.; and Lueker, G. S. 1976. Algorithmic Aspects of Vertex Elimination on Graphs. *SIAM Journal on Computing*, 5(2): 266–283.
- Schmid, D.; and Sly, A. 2022. On the number and size of Markov equivalence classes of random directed acyclic graphs. *arXiv preprint arXiv:2209.04395*.
- Steinsky, B. 2003. Enumeration of labelled chain graphs and labelled essential directed acyclic graphs. *Discrete mathematics*, 270(1-3): 267–278.
- Talvitie, T.; and Koivisto, M. 2019. Counting and Sampling Markov Equivalent Directed Acyclic Graphs. In *Proceedings of the 33rd AAAI Conference on Artificial Intelligence (AAAI 2019)*, volume 33, 7984–7991.
- Van Der Zander, B.; and Liskiewicz, M. 2016. Separators and Adjustment Sets in Markov Equivalent DAGs Graphs. In *Proceedings of the 30th AAAI Conference on Artificial Intelligence (AAAI 2016)*, volume 30, 3315–3321.
- Bang, Christine W.; and Didelez, V. 2023. Do We Become Wiser with Time? On Causal Equivalence with Tiered Background Knowledge. In *Proceedings of the 39th Conference on Uncertainty in Artificial Intelligence (UAI 2023)*, 119–129.
- Verma, T.; and Pearl, J. 1990. Equivalence and synthesis of causal models. In *Proceedings of the Sixth Annual Conference on Uncertainty in Artificial Intelligence*, 255–270.
- Wienöbst, M.; Bannach, M.; and Liśkiewicz, M. 2021. Polynomial-Time Algorithms for Counting and Sampling Markov Equivalent DAGs. In *Proceedings of the 35th AAAI Conference on Artificial Intelligence (AAAI 2021)*, 12198–12206.