


Expressive Quantale-Valued Logics for Coalgebras: an Adjunction-Based Approach

Harsh Beohar  

University of Sheffield, United Kingdom

Sebastian Gurke  

Universität Duisburg-Essen, Germany

Barbara König  

Universität Duisburg-Essen, Germany

Karla Messing  

Universität Duisburg-Essen, Germany

Jonas Forster  

Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany

Lutz Schröder  

Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany

Paul Wild  

Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany

Abstract

We address the task of deriving fixpoint equations from modal logics characterizing behavioural equivalences and metrics (summarized under the term *conformances*). We rely on an earlier work that obtains Hennessy-Milner theorems as corollaries to a fixpoint preservation property along Galois connections between suitable lattices. We instantiate this to the setting of coalgebras, in which we spell out the compatibility property ensuring that we can derive a behaviour function whose greatest fixpoint coincides with the logical conformance. We then concentrate on the linear-time case, for which we study coalgebras based on the machine functor living in Eilenberg-Moore categories, a scenario for which we obtain a particularly simple logic and fixpoint equation. The theory is instantiated to concrete examples, both in the branching-time case (bisimilarity and behavioural metrics) and in the linear-time case (trace equivalences and trace distances).

2012 ACM Subject Classification Theory of computation → Modal and temporal logics

Keywords and phrases modal logics, coalgebras, behavioural equivalences, behavioural metrics, linear-time semantics, Eilenberg-Moore categories

Digital Object Identifier 10.4230/LIPIcs...XX

Funding The first author was supported by the EPSRC NIA Grant EP/X019373/1, while the remaining authors were supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 434050016 (SpeQt).

1 Introduction

Behavioural equivalences (such as bisimilarity and trace equivalence) are an important technique to identify states with the same behaviour in a transition system [38]. They have been complemented by notions of behavioural metrics [13, 37, 9] measuring the distance between states, in particular in a quantitative setting. We work in a coalgebraic setting [31] that allows us to answer generic questions about behavioural equivalences and metrics, parametrized over various branching types (non-deterministic, probabilistic, weighted, etc.).



© H. Beohar, S. Gurke, B. König, K. Messing, J. Forster, L. Schröder, P. Wild;
licensed under Creative Commons License CC-BY 4.0

Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

There are various ways to characterize behavioural equivalences or metrics, which we illustrate using trace equivalence as an example: (i) *direct specification*: Two states x, y are trace equivalent if they admit the same traces; (ii) *logic*: x, y are trace equivalent if they cannot be distinguished in a modal logic based on diamond modalities and the constant *true*; (iii) *fixpoint equation*: x, y are trace equivalent if the pair $(\{x\}, \{y\})$ is contained in the greatest fixpoint of the bisimulation function on the determinized transition system; (iv) *games*: there is an attacker-defender game characterizing the equivalence.

Our focus is on (ii) and (iii), and our take is quite different from the usual approach: Instead of first defining the behavioural equivalence/metric and then setting up an expressive logic for it, we start by defining the logic and derive a fixpoint equation from the logic. Fixpoint equations are of great interest since efficient algorithms for computing behavioural conformances are almost always based on fixpoint characterizations; in future work, we aim to exploit such characterizations for algorithmic purposes. However, for a given logic, corresponding fixpoint equations do not always exist, and we give conditions for ensuring that they do. We use the Galois connection approach from [4] as a starting point, and instead of instantiating it for each case study, we propose a generic coalgebraic framework. By employing fibrations (resp. indexed categories) [16, 17], we parameterize over the notion of *conformance* (e.g. equivalence, metric) that is our focus of attention. Moreover, we parametrize over a quantale in which both conformances and formulae take their values.

One interest, particularly, is in linear-time notions of conformance (such as trace/language equivalences and their quantitative cousins), for which we work in an Eilenberg-Moore category where the coalgebras live. We exploit the generalized powerset construction [19, 33] and characterize those (trace) logics that can be turned into suitable fixpoint equations on the determinized coalgebra, using a notion of compatibility [4] that has its roots in up-to techniques [29]. We also study the relation of compatibility to the notion of depth-1 separation used in (quantitative) graded logics [10, 12].

More concretely, we work with coalgebras of the form $c: X \rightarrow FTX$, living in some category \mathbf{C} , where a monad T intuitively specifies the implicit branching (or side effects) and a functor F describes the explicit branching type. For instance, for a non-deterministic automaton we choose $T = \mathcal{P}$ and $F = _ \Sigma \times \mathbf{2}$. We fix an *EM-law* $\zeta: TF \Rightarrow FT$ [19] allowing us to obtain a determinized coalgebra, i.e., a coalgebra of the form $c^\# : TX \rightarrow FTX$ that can be viewed as a coalgebra in the Eilenberg-Moore category of T . We can then define a logic function log that is defined on sets of \mathcal{V} -valued predicates on X and whose least fixpoint induces a behavioural conformance on TX . Alternatively, given the determinization $c^\#$, we can – in a fibrational style – define a conformance on TX as the greatest fixpoint of a Kantorovich lifting followed by a reindexing via $c^\#$. The aim is to show that both conformances coincide.

We allow arbitrary constants in the logic, which – in particular in the linear-time case – are able to add extra distinguishing power to the logic. Along the way we give an answer to the question of why, unlike branching-time logics, linear-time logics often do not need any additional (boolean) operators, only modalities and constants.

As examples we consider bisimilarity and branching-time pseudometrics for probabilistic transition systems, as well as linear-time conformances such as trace equivalence and trace distance.

Roadmap. After reviewing preliminaries in Section 2, we summarize the approach based on Galois connections (adjunctions) in Section 3. The instantiation to generic coalgebras is presented in Section 4 and a concrete quantale-valued branching-time logic spelled out in Section 5. Section 6 specializes to coalgebras in Eilenberg-Moore categories, leading to

results strengthening those for the general case. Finally, Section 7 details the linear-time case studies mentioned above, and we conclude in Section 8.

2 Preliminaries

We recall some basic definitions and facts on lattices, quantales, generalized distances, coalgebra, monads and their algebras and on indexed categories. We do assume basic familiarity with category theory (e.g. [1]).

2.1 Lattices, Fixpoints and Galois Connections

A *complete lattice* $(\mathbb{L}, \sqsubseteq)$ consists of a set \mathbb{L} with a partial order \sqsubseteq such that each $Y \subseteq \mathbb{L}$ has a least upper bound $\bigsqcup Y$ (also called supremum, join) and a greatest lower bound $\bigsqcap Y$ (also called infimum, meet). The Knaster-Tarski theorem [35] guarantees that any monotone function $f: \mathbb{L} \rightarrow \mathbb{L}$ on a complete lattice \mathbb{L} has a *least fixpoint* μf and a *greatest fixpoint* νf .

Let \mathbb{L}, \mathbb{B} be two lattices. A *Galois connection* from \mathbb{L} to \mathbb{B} is a pair $\alpha \dashv \gamma$ of monotone functions $\alpha: \mathbb{L} \rightarrow \mathbb{B}, \gamma: \mathbb{B} \rightarrow \mathbb{L}$ such that for all $\ell \in \mathbb{L}, b \in \mathbb{B}$: $\alpha(\ell) \sqsubseteq b \iff \ell \sqsubseteq \gamma(b)$.

A *closure* $\text{cl}: \mathbb{L} \rightarrow \mathbb{L}$ is a monotone, idempotent and extensive (i.e. $\forall x \in \mathbb{L} \ x \sqsubseteq \text{cl}(x)$) function on a lattice. A *co-closure* is monotone, idempotent and extensive wrt. \supseteq . Given a Galois connection $\alpha \dashv \gamma$, $\gamma \circ \alpha$ is always a closure and $\alpha \circ \gamma$ a co-closure.

2.2 Quantales and Generalized Distances

► **Definition 1.** A (*unital, commutative*) *quantale* $(\mathcal{V}, \otimes, 1)$, or just \mathcal{V} , is a complete lattice with an associative, commutative operation $\otimes: \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$ with unit 1 that distributes over arbitrary (possibly infinite) joins \bigvee . If 1 is the top element of \mathcal{V} , then \mathcal{V} is *integral*.

In a quantale \mathcal{V} , the functor $- \otimes y$ has a right adjoint $[y, -]$ for every $y \in \mathcal{V}$; that is, $x \otimes y \leq z \iff x \leq [y, z]$ for all $x, y, z \in \mathcal{V}$.

- **Example 2.** 1. The Boolean algebra $\mathbf{2}$ with $\otimes = \wedge$ and unit 1 is an integral quantale; for $y, z \in \mathbf{2}$, we have $[y, z] = y \rightarrow z$.
2. The complete lattice $[0, 1]$ ordered by the reversed order of the reals, i.e. $\leq = \geq_{\mathbb{R}}$, and equipped with truncated addition $r \otimes s = \min(r + s, 1)$, is an integral quantale; for $r, s \in [0, 1]$, we have $[r, s] = s \dot{-} r = \max(s - r, 0)$ (truncated subtraction).

Given a quantale \mathcal{V} , a *directed (\mathcal{V} -valued) pseudometric (on X)* is a function $d: X \times X \rightarrow \mathcal{V}$ such that (i) $\forall x \in X \ d(x, x) \geq 1$ (reflexivity); (ii) $\forall x, y, z \in X \ d(x, z) \geq d(x, y) \otimes d(y, z)$ (transitivity/triangle inequality). Moreover, d is a *pseudometric* if additionally, (iii) $\forall x, y \in X \ d(x, y) = d(y, x)$ (symmetry). We write $\mathbf{DPMet}_{\mathcal{V}}(X)$ to denote the lattice of all directed pseudometrics on X , while for pseudometrics we use $\mathbf{PMet}_{\mathcal{V}}(X)$. Given $d_X \in \mathbf{DPMet}_{\mathcal{V}}(X), d_Y \in \mathbf{DPMet}_{\mathcal{V}}(Y)$, a function $f: X \rightarrow Y$ is *non-expansive* (wrt. d_X, d_Y) if $d_X(x, x') \leq d_Y(f(x), f(x'))$ for all $x, x' \in X$. Note that due to the choice of order in the quantale, the inequality in the definitions above is reversed wrt. the standard definitions that are typically given in the order on the reals. As originally observed by Lawvere [25], one may see directed \mathcal{V} -valued pseudometrics as \mathcal{V} -enriched categories, or just \mathcal{V} -categories, and non-expansive functions as \mathcal{V} -functors.

2.3 Coalgebras and Eilenberg-Moore Categories

Given a functor $F: \mathbf{C} \rightarrow \mathbf{C}$, an F -coalgebra (X, c) (or simply c) consists of an object $X \in \mathbf{C}$ and a \mathbf{C} -arrow $c: X \rightarrow FX$. In the paradigm of *universal coalgebra* [31], we understand X as the state space of a transition system, F as specifying the branching type of the system, and c as a transition map that assigns to each state a collection of successors structured according to F . For instance, when $\mathbf{C} = \mathbf{Set}$ is the category of sets and functions, then the powerset functor \mathcal{P} assigns to each set its powerset, and \mathcal{P} -coalgebras are just sets equipped with a transition relation. On the other hand, the (finitely supported) distribution functor \mathcal{D} assigns to each set X the set of finitely supported probability distributions on X , given in terms of maps $p: X \rightarrow [0, 1]$ with finite support such that $\sum_{x \in X} p(x) = 1$. A \mathcal{D} -coalgebra thus is precisely a Markov chain.

Recall that a *monad* $(T, \eta: \text{Id} \Rightarrow T, \mu: TT \Rightarrow T)$ on \mathbf{C} , usually denoted by just T , consists of a functor $T: \mathbf{C} \rightarrow \mathbf{C}$ and natural transformations $\eta: \text{Id} \Rightarrow T$ (the *unit*) and $\mu: TT \Rightarrow T$ (the *multiplication*), subject to certain coherence laws. Monads abstractly capture notions of algebraic theory, with TX being thought of as terms modulo provable equality over variables in X . Correspondingly, a T -algebra (X, a) consists of an object X of \mathbf{C} and a \mathbf{C} -arrow $a: TX \rightarrow X$ such that $a \circ \eta_X = \text{id}_X$ and $a \circ Ta = a \circ \mu_X$; we may think of T -algebras as algebras for the algebraic theory encapsulated by T . A homomorphism between T -algebras $(X, a), (Y, b)$ is a \mathbf{C} -arrow $f: X \rightarrow Y$ such that $b \circ Tf = f \circ a$. The *Eilenberg-Moore category* of T , denoted $\mathbf{EM}(T)$, is the category of T -algebras and their homomorphisms. There is a free-forgetful adjunction $L \dashv R: \mathbf{C} \rightarrow \mathbf{EM}(T)$, where the forgetful functor R maps an algebra to its underlying \mathbf{C} -object and L maps an object $X \in \mathbf{C}$ to the free algebra (TX, μ_X) .

A (*monad-over-functor*) *distributive law* (or *EM-law*) of a monad T over a functor F is a natural transformation $\zeta: TF \Rightarrow FT$ satisfying $\zeta_X \circ \eta_{FX} = F\eta_X$ and $\zeta_X \circ \mu_{FX} = F\mu_X \circ \zeta_{TX} \circ T\zeta_X$. This is equivalent to saying that the assignment $\tilde{F}(X, a) = (FX, Fa \circ \zeta_X)$ defines a *lifting* $\tilde{F}: \mathbf{EM}(T) \rightarrow \mathbf{EM}(T)$ of F (where *lifting* means that $R\tilde{F} = FR$). Then, the *determinization* [33] of a coalgebra $c: X \rightarrow FTX$ in \mathbf{C} is the transpose $c^\# : LX \rightarrow \tilde{F}LX$ of c under $L \dashv R$. More concretely the determinization can be obtained as $c^\# = F\mu_X \circ \zeta_{TX} \circ Tc$. For instance, when $FX = X^\Sigma \times \mathbf{2}$ and $T = \mathcal{P}$, then this yields exactly the standard powerset construction for the determinization of non-deterministic automata.

2.4 Indexed Categories and Fibrations

Our aim is to equip objects of a category with additional information, e.g., consider sets with (equivalence) relations or metrics. Formally, this is done by working with fibrations, in particular we will consider fibrations arising from the Grothendieck construction for indexed categories [16, 17]. For us it is sufficient to consider as indexed categories functors $\Phi: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Pos}$, where \mathbf{Pos} is the category of posets (ordered by \preceq) with monotone maps. Such functors induce a fibration $U: \int \Phi \rightarrow \mathbf{C}$ where U is the forgetful functor and $\int \Phi$ is the category whose objects and arrows are characterized as follows:

$$\frac{X \in \mathbf{C} \wedge d \in \Phi X}{(X, d) \in \int \Phi} \quad \frac{X \xrightarrow{f} Y \in \mathbf{C} \wedge d \preceq (\Phi f)d'}{(X, d) \xrightarrow{f} (Y, d')} \in \int \Phi$$

Here, $f^* = \Phi f$ is also called the *reindexing operation*, and d is called a *conformance*.

Typical examples are functors $\Phi: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Pos}$ mapping a set X to the lattice of equivalence relations or pseudometrics on X .

3 Adjoint Logic: the General Framework

We summarize previous results on relating logical and behavioural conformances using Galois connections [4]. These results are based on the following well-known property that shows how fixpoints are preserved by Galois connections (e.g. [3, 7, 8]). The formulation of these properties involves a notion of compatibility studied for coinductive up-to techniques [29].

► **Definition 3.** Let $\log, \text{cl}: \mathbb{L} \rightarrow \mathbb{L}$ be monotone endofunctions on a partial order $(\mathbb{L}, \sqsubseteq)$. Then \log is *cl-compatible* if $\log \circ \text{cl} \sqsubseteq \text{cl} \circ \log$.

► **Theorem 4** ([4]). Let $\alpha: \mathbb{L} \rightarrow \mathbb{B}$, $\gamma: \mathbb{B} \rightarrow \mathbb{L}$ be a Galois connection between complete lattices \mathbb{L}, \mathbb{B} , and let $\log: \mathbb{L} \rightarrow \mathbb{L}$, $\text{beh}: \mathbb{B} \rightarrow \mathbb{B}$ be monotone. Then the following holds.

1. If $\text{beh} = \alpha \circ \log \circ \gamma$ then $\alpha(\mu \log) \sqsubseteq \mu \text{beh}$.
2. If $\alpha \circ \log = \text{beh} \circ \alpha$, then $\alpha(\mu \log) = \mu \text{beh}$. If \log reaches its fixpoint in ω steps, i.e., $\mu \log = \log^\omega(\perp)$, then so does beh .
3. Let $\text{cl} = \gamma \circ \alpha$ be the closure operator of the Galois connection, and suppose that $\text{beh} = \alpha \circ \log \circ \gamma$. If \log is *cl-compatible*, then $\alpha(\mu \log) = \mu \text{beh}$.

► **Remark 5.** In fact, there is a weaker notion than compatibility that ensures the same result, i.e., $\alpha(\mu \log) = \mu \text{beh}$. In particular, it is sufficient to show the following condition:

- $\log(\text{cl}(\ell)) \sqsubseteq \text{cl}(\log(\ell))$ for all $\ell \in \mathbb{S} \subseteq \mathbb{L}$ where \mathbb{S} is an invariant of \log , i.e. $\log[\mathbb{S}] \subseteq \mathbb{S}$, $\perp \in \mathbb{S}$, and \mathbb{S} is closed under directed joins. (If the fixpoint is reached in ω steps, closure under directed joins is not required.)

We apply this in a scenario where \mathbb{L} consists of logical formulas (or more precisely their semantics, in the shape of sets of definable predicates) and \mathbb{B} consists of conformances. These Galois connections will be contra-variant when we consider the quantalic ordering in \mathbb{B} . Then, \log is the “logic function” that adds a layer of modalities and propositional operators to a given set of predicates, so that $\mu \log$ is the semantics of the set of formulas of the logic. On the other hand, beh is the “behaviour function”, whose greatest fixpoint νbeh (remember the contra-variance) is behavioural conformance.

► **Example 6.** The simplest Galois connection used in [4] for characterizing behavioural equivalence is between $\mathbb{L} = \mathcal{P}(\mathbf{2}^X)$ (sets of predicates on X) and $\mathbb{B} = \mathbf{PMet}_2(X)$ (equivalences on X), where α maps every set of predicates to the equivalence relation induced by it and γ maps an equivalence to the set of predicates closed under it.

Moving to pseudometrics we obtain a Galois connection between $\mathbb{L} = \mathcal{P}([0, 1]^X)$ (sets of real-valued predicates on X) and $\mathbb{B} = \mathbf{PMet}_{[0,1]}(X)$ (pseudometrics on X) where α maps every set of functions $X \rightarrow [0, 1]$ to the least pseudometric making all these functions non-expansive and γ takes a pseudometric and produces all its non-expansive functions.

So first, define a logical universe \mathbb{L} and a logic function $\log: \mathbb{L} \rightarrow \mathbb{L}$. Second, choose a suitable Galois connection $\alpha \dashv \gamma$ to a behaviour universe \mathbb{B} and show that \log is *cl-compatible*. Third, derive the behaviour function $\text{beh} = \alpha \circ \log \circ \gamma: \mathbb{B} \rightarrow \mathbb{B}$. From the results above, we automatically obtain the equality $\alpha(\mu \log) = \mu \text{beh}$, which tells us that logical and behavioural equivalence respectively distance coincide (Hennessy-Milner theorem).

4 Adjoint Logic for Coalgebras

In this section we will describe a general framework where the adjoint logic is instantiated to the setting of coalgebraic modal logic.

4.1 Setting up the Adjunction

One can generalize from Example 6 and instead of a set X take an object in a (locally small) category \mathbf{C} . Furthermore we fix an object $\Omega \in \mathbf{C}$ (the *truth value* object), which in all our applications will be a quantale. Predicates are represented by the indexed category $\mathbf{C}(_, \Omega)$; thus, sets of predicates (lattice \mathbb{L}) are given by the indexed category $\mathcal{P} \circ \mathbf{C}(_, \Omega)$ (where the order is inclusion). In addition, we use an indexed category Φ specifying the notion of conformance on X (lattice \mathbb{B}) and work with the following assumptions:

- A1** Each fibre ΦX is a poset having arbitrary meets (thus, a complete lattice) and the reindexing map preserves these meets (i.e. $\int \Phi$ has fibred limits).
A2 Let $d_\Omega \in \Phi \Omega$ be a fixed conformance on the truth value object Ω .

For an arrow $f \in \mathbf{C}(X, Y)$, we write f^\bullet for the reindexing in $\mathbf{C}(_, \Omega)$ ($f^\bullet g = g \circ f$, where $g \in \mathbf{C}(Y, \Omega)$) and f^* for the reindexing in Φ ($f^* = \Phi f$).

► **Theorem 7.** *Let X be an object of \mathbf{C} . Under Assumptions **A1** and **A2**, there is a dual adjoint situation (contravariant Galois connection) $\alpha_X \dashv \gamma_X$ between the underlying fibres:*

$$\begin{aligned} \alpha_X: \mathcal{P}(\mathbf{C}(X, \Omega)) &\rightarrow \Phi(X)^{op} & S \subseteq \mathbf{C}(X, \Omega) &\mapsto \bigwedge_{k \in S} k^*(d_\Omega) \\ \gamma_X: \Phi(X)^{op} &\rightarrow \mathcal{P}(\mathbf{C}(X, \Omega)) & d \in \Phi X &\mapsto \{k \in \mathbf{C}(X, \Omega) \mid d \preceq k^*(d_\Omega)\}. \end{aligned}$$

More concretely: α_X, γ_X are both antitone ($S \subseteq S' \implies \alpha_X(S) \succeq \alpha_X(S')$, $d \preceq d' \implies \gamma_X(d) \supseteq \gamma_X(d')$) and we have $d \preceq \alpha_X(S) \iff S \subseteq \gamma_X(d)$ for $d \in \Phi X$ and $S \in \mathcal{P}\mathbf{C}(X, \Omega)$.

Thus, for $X \in \mathbf{C}$, the fibres $\mathcal{P}\mathbf{C}(X, \Omega)$ and $(\Phi X)^{op}$ will take the role of \mathbb{B} and \mathbb{L} (respectively) as in [4, Theorem 3.2]. Moreover, Theorem 7 will be instantiated to obtain the desired Galois connections between predicates and conformances for our case studies.

► **Example 8.** Let $\mathbf{C} = \mathbf{Set}$ and $\Omega = \mathcal{V}$ be a quantale. We consider $\Phi X = \mathbf{DPMet}_{\mathcal{V}}(X)$ (resp., $\Phi X = \mathbf{PMet}_{\mathcal{V}}(X)$) with the order \preceq on ΦX induced by the pointwise lifting of the order \leq on \mathcal{V} . The reindexing functor f^* for a function $f: X \rightarrow Y$ is given by $f^*d = d \circ (f \times f)$; thus, satisfying **A1**. As conformance d_Ω on \mathcal{V} we take the internal hom $[_, _]$ (resp., its symmetrization: $d_\Omega(x, y) = [x, y] \wedge [y, x]$); thus, satisfying **A2**. Then we have $\alpha_X \dashv \gamma_X: \mathcal{P}(\mathbf{Set}(X, \mathcal{V})) \rightleftarrows \mathbf{DPMet}_{\mathcal{V}}(X)$, where:

$$\begin{aligned} \alpha_X(S)(x, x') &= \bigwedge_{h \in S} d_\Omega(h(x), h(x')) \\ \gamma_X(d) &= \{h: X \rightarrow \mathcal{V} \mid \forall_{x, x' \in X} d(x, x') \leq d_\Omega(h(x), h(x'))\} \end{aligned}$$

In both cases, α_X assigns to a set of maps the greatest (directed) pseudometric making all these maps non-expansive, while γ_X maps a pseudometric all its non-expansive maps.

4.2 Characterizing Closure

Given that the key condition imposed on the logic function in Theorem 4 is compatibility with the closure of the Galois connection, it is important to understand how this closure operates. In the setting of Theorem 7 we can characterize the closure in terms of non-expansive propositional operators, provided that γ is natural. We note first that α is always natural:

► **Proposition 9.** *In Theorem 7, the transformation α is natural in $X \in \mathbf{C}$, that is, for $f \in \mathbf{C}(X, Y)$, we have $\alpha_X \circ \mathcal{P}(f^\bullet) = f^* \circ \alpha_Y$.*

For the right adjoint γ , naturality need not hold in general. It does hold for **Set** and generalized (directed) metrics over the quantales in Example 2. A counterexample can however be constructed for $\mathbf{C} = \mathbf{EM}(\mathcal{P})$ and $\mathcal{V} = [0, 1]$ (see [4]).

If γ is natural, then we can characterize the closure $\gamma \circ \alpha$ using the internal language of indexed categories. To this end, suppose that \mathbf{C} has (small) products. Then for every $S \subseteq \mathbf{C}(X, \Omega)$, we have a unique tupling $\langle S \rangle: X \rightarrow \Omega^S$ such that $\pi_k \circ \langle S \rangle = k$ for all $k \in S$, where $\pi_k: \Omega^S \rightarrow \Omega$ is the product projection for k .

► **Lemma 10.** *The right adjoint γ in Theorem 7 is laxly natural, i.e. $\mathcal{P}(f^\bullet) \circ \gamma_Y \subseteq \gamma_X \circ f^*$ for $f: X \rightarrow Y \in \mathbf{C}$. If γ is natural (i.e. the inclusion is an equality) and \mathbf{C} has products, then $\gamma_X(\alpha_X(S)) = \mathcal{P}(\langle S \rangle^\bullet)(\gamma_{\Omega^S}(d_{\Omega^S}))$, where $d_{\Omega^S} = \bigwedge_{k \in S} \pi_k^* d_\Omega$.*

This result can be interpreted as follows: $\gamma_{\Omega^S}(d_{\Omega^S})$ is the set of all non-expansive functions $\Omega^S \rightarrow \Omega$, hence all non-expansive operators of arbitrary arity on Ω . Reindexing via $\langle S \rangle$ means to combine all predicates in S via those operators, hence we describe the closure under all non-expansive operations on Ω .

4.3 Towards a Generic Logic Function

Since our slogan is to generate the behaviour function from the logic function, we start by setting up our logical framework first. Following [4, 11], we adopt a semantic approach to defining a (modal) logic, i.e., we specify the operators (including modalities) as a transformation of predicates; formally, as a (natural) transformation \mathbf{log} of type $\mathcal{PC}(_, \Omega) \Rightarrow \mathcal{PC}(_, \Omega)$. The idea is that the logic function \mathbf{log}_X adds one “layer” of modal depth; in particular, the least fixpoint $\mu \mathbf{log}_X$ of \mathbf{log}_X can be seen as the set of (interpretations of) all modal formulas.

In particular, we require

A3 Fix a family $(ev_\lambda \in \mathbf{C})_{\lambda \in \Lambda}$ of evaluation maps $ev_\lambda: F\Omega \rightarrow \Omega$.

As noted in [32], such evaluation maps – commonly used in coalgebraic modal logic – correspond to natural transformations of type $\mathbf{C}(_, \Omega) \rightarrow \mathbf{C}(F_, \Omega)$ by the Yoneda lemma.

► **Proposition 11.** *A family of evaluation maps $(ev_\lambda)_{\lambda \in \Lambda}$ induces a natural transformation $\Lambda: \mathcal{PC}(_, \Omega) \Rightarrow \mathcal{PC}(F_, \Omega)$ given by $S \mapsto \{\lambda_X(h) \mid \lambda \in \Lambda, h \in S\}$, where $\lambda_X(h) = ev_\lambda \circ Fh$.*

Apart from modalities, a logic typically needs operators and constants. We do not consider constants as 0-ary operators, which allows us to distinguish between operators that arise as in Lemma 10 from the closure of the Galois connection (that is, non-expansive operators) and the remaining (constant) operators that bring additional distinguishing power. This is, for instance, needed in the case of trace equivalence on a determinized transition system to distinguish the empty set of states from sets of states having no transitions. We need an additional (constant) predicate for this task that can neither be provided by the closure nor by a constant modality (see Section 7.2).

A4 We assume a set $\Theta_X \subseteq \mathbf{C}(X, \Omega)$ of *constants* (which is later restricted to consist of free extensions of constant maps).

To model the propositional operators, we introduce a closure cl'_X :

A5 For each $X \in \mathbf{C}$ we assume that there is a closure $cl'_X: \mathcal{PC}(X, \Omega) \rightarrow \mathcal{PC}(X, \Omega)$ (not necessarily natural), specifying the propositional operators.

We say that cl'_X is a *subclosure* of cl_X whenever $\text{cl}'_X \subseteq \text{cl}_X$, which means that the propositional operators implemented by cl'_X are already contained in the closure induced by the Galois connection (cf. Lemma 10).

Now we can define the logic function for a coalgebra c as

$$\log_X = \mathcal{P}(c^\bullet) \circ \Lambda_X \circ \text{cl}'_X \cup \Theta_X : \mathcal{PC}(X, \Omega) \rightarrow \mathcal{PC}(X, \Omega).$$

Its least fixpoint contains all predicates that can be described by modal formulas.¹

► **Example 12.** Let $\mathbf{C} = \mathbf{Set}$, $c: X \rightarrow FX$ and $\Phi X = \mathbf{PMet}_{\mathcal{V}}(X)$. Recall the Galois connection from Example 8 and consider the following two examples, where in both cases no constants are needed, i.e., Θ_X is empty; thus, **A4** vacuously holds.

1. *Bisimilarity on (unlabelled) transition systems:* we let $F = \mathcal{P}_{fin}$ (finite powerset functor), $\mathcal{V} = 2$, and consider the evaluation map $ev_{\diamond} : \mathcal{P}_{fin}2 \rightarrow 2$ encoding the usual diamond modality: $ev_{\diamond}(U) = 1 \iff 1 \in U$. This can be extended to a logic by choosing as cl' (Assumption **A5**) the closure under all (finitary) Boolean operators.

2. *Behavioural metrics for probabilistic transition systems with termination:* we let $FX = \mathcal{D}X + 1$ (where $1 = \{\checkmark\}$) and $\mathcal{V} = ([0, 1], \geq_{\mathbb{R}})$. Define two evaluation maps: $ev_E : \mathcal{D}[0, 1] + 1 \rightarrow [0, 1]$ corresponds to expectation, i.e. $ev_E(p) = \sum_{r \in [0, 1]} r \cdot p(r)$ if $p \in \mathcal{D}X$ (0 otherwise). Furthermore, $ev_* : \mathcal{D}[0, 1] + 1 \rightarrow [0, 1]$ with $ev_*(p) = 1$ if $p = \checkmark$ (0 otherwise). We extend this to a logic by defining cl' (Assumption **A5**): we add as operators the constant 1, $\min(\varphi, \varphi')$, $1 - \varphi$ and $\varphi \dot{-} q$ for a rational q (where φ is a formula), as in similar logics for probabilistic transition systems [37].

To ensure that the requirements of Section 3 are met, we have to show compatibility of the logic function. To this end, we introduce the notion of compatibility of cl' .

► **Definition 13.** Given a closure $\text{cl}'_X : \mathcal{PC}(X, \Omega) \rightarrow \mathcal{PC}(X, \Omega)$, we say that cl'_X is *compatible* if the map $\Lambda_X \circ \text{cl}'_X$ (for each $X \in \mathbf{C}$) is compatible with the closure $\text{cl}_X = \gamma_X \circ \alpha_X$ induced by the adjoint situation in Theorem 7, i.e., $\Lambda_X \circ \text{cl}'_X \circ \text{cl}_X \subseteq \text{cl}_{FX} \circ \Lambda_X \circ \text{cl}'_X$.

The following results hold under Assumptions **A1-A5** and thus, we avoid stating them in various lemma/theorem statements.

► **Proposition 14.** *For a given compatible closure cl'_X , the above logic function \log_X is cl_X -compatible, i.e., $\log_X \circ \text{cl}_X \subseteq \text{cl}_X \circ \log_X$.*

We will now study equivalent conditions and special cases in which compatibility holds. First, it is easy to see that $\text{cl}' = \text{cl}$ is always compatible, but typically introduces infinitary operators. Moreover, if cl' is the identity (that is, there are no propositional operators), then compatibility of cl' reduces to cl -compatibility of Λ .

► **Lemma 15.** *Let cl'_X be a subclosure of cl_X . It holds that cl'_X is compatible if and only if $\alpha_{FX} \circ \Lambda_X \circ \text{cl}'_X \preceq \alpha_{FX} \circ \Lambda_X \circ \text{cl}_X$.*

We next adapt the separation property establishing expressiveness of *graded logics* w.r.t. *graded semantics* [10, 12] to the present setting, an additional twist being that the conformance w.r.t. which modalities must be separating is the one induced by the modalities themselves.

¹ In our setup a formula is either a constant or starts with a modality, which still results in an expressive logic. One could slightly modify \log_X and obtain all formulas by adding another closure cl' .

► **Definition 16** (Depth-1 self-separation). A set $S \subseteq \mathbf{C}(X, \Omega)$ of predicates is *initial* for $d \in \Phi X$ if $\alpha_X(S) = d$. Let cl'_X be a subclosure of cl_X . The *depth-1 self-separation* property holds if for every S that is closed under cl'_X (i.e., $S = \text{cl}'_X(S)$) and initial for d , it follows that $\Lambda_X(S)$ is initial for $\kappa_X(d)$ where $\kappa_X = \alpha_{FX} \circ \Lambda_X \circ \gamma_X$.

► **Lemma 17.** *Let cl'_X be a subclosure of cl_X . Then cl'_X is compatible if and only if the depth-1 self-separation property holds.*

Finally, we study a sufficient condition on evaluation maps ensuring cl -compatibility of Λ .

► **Lemma 18.** *If each evaluation map ev_λ arises as a natural transformation $\eta: F \Rightarrow \text{Id}$ or $\eta: F \Rightarrow \Omega$ (Ω is the constant functor mapping every object to Ω), that is $ev_\lambda = \eta_\lambda$, then Λ is compatible with cl .*

► **Example 19.** We establish compatibility for the logics considered in Example 12. In branching-time logics in general, depth-1 self-separation usually boils down to establishing a Stone-Weierstraß type property saying that if $S \subseteq \mathbf{C}(X, \Omega)$ is initial and closed under cl'_X , then S is dense in $\mathbf{C}(X, \Omega)$, for suitably restricted X [11]. For finitary set functors such as \mathcal{P}_{fin} or \mathcal{D} , it suffices to prove self-separation on finite X . Additional details are as follows.

1. In the case of unlabelled transition systems, we are given an equivalence relation R on a finite set X , a set $S \subseteq \mathbf{Set}(X, 2)$ that is initial for R and closed under Boolean combinations, and $A, B \in \mathcal{P}_{fin}(X)$ that are distinguished by some predicate $\diamond_X f: \mathcal{P}_{fin}(X) \rightarrow 2$ where f is invariant under R . We then have to show that A, B are distinguished by $\diamond g$ for some $g \in S$. But by functional completeness of Boolean logic and because X is finite, S is in fact the set of *all* R -invariant functions $X \rightarrow 2$, so we can just take $g = f$.


2. The argument is similar for probabilistic transition systems, with some additional considerations necessitated by the quantitative setting. We are now given a set $S \subseteq \mathbf{Set}(X, [0, 1])$ that is initial for d and closed under propositional operators as per Example 12.2. By a variant of the Stone-Weierstraß theorem, this implies that S is dense in the space of non-expansive maps $(X, d) \rightarrow [0, 1]$ (see [42]), which means that in an argument as in the previous item, we can take g to range over functions in S that approximate the given non-expansive function $f: (X, d) \rightarrow [0, 1]$ arbitrarily closely, using additionally that the predicate lifting induced by ev_E as in Example 12(2) is non-expansive [41].

4.4 Towards a Generic Behaviour Function

Building on the previous section, we define the behaviour function $\text{beh}_X: \Phi X \rightarrow \Phi X$ as $\text{beh}_X = \alpha_X \circ \log_X \circ \gamma_X$ and – under the assumption of compatibility – we have² $\alpha_X(\mu \log_X) = \nu \text{beh}_X$. In other words, the notions of logical and behavioural conformances coincide.

This motivates a closer investigation of νbeh_X : in what sense does it coincide with known behavioural equivalences or metrics? Defining a behavioural conformance (that is, an element of ΦX) in a fibrational setting is typically done by taking the greatest fixpoint of a function defined in two steps: the lifting of a conformance ΦX to $\Phi(FX)$, followed by a reindexing via c . Here, we consider Kantorovich-style [2] or codensity [21] liftings based on the evaluation maps. Kantorovich liftings have originally been used to lift metrics on a set X to metrics of probability distributions over X . In the probabilistic case, an alternative

² Note that the adjunction defined in Theorem 7 is contravariant. Hence the least fixpoint of beh from Theorem 4 becomes the greatest fixpoint νbeh_X wrt. the lattice order \preceq .



■ **Figure 1** The adjoint setup with $\log_X = \mathcal{P}(c^*) \circ \Lambda_X \circ \text{cl}'_X \cup \Theta_X$ and $\text{beh}_X = c^* \circ \kappa_X \cup \alpha_X(\Theta_X)$.

characterization is given via optimal transport plans in transportation theory (earth mover's distance) [39].

We use the natural transformation Λ introduced earlier and consider the composite $\kappa_X = \alpha_{FX} \circ \Lambda_X \circ \gamma_X$, the mentioned Kantorovich lifting. If $F = \mathcal{D}$ and the evaluation map is expectation, then we obtain exactly the classical Kantorovich lifting.

Given a coalgebra $c: X \rightarrow FX$, we use the *behaviour function* $\text{beh}_X = c^* \circ \kappa_X \wedge \alpha_X(\Theta_X)$; here, Θ_X is a set of constants as in Section 4.3 (Asumption **A4**).

► **Example 20.** We derive the behaviour functions for Examples 12 and 19.

1. In the case $F = \mathcal{P}$ and $\mathcal{V} = 2$, the lifting $\kappa_X(R) \subseteq \mathcal{P}X \times \mathcal{P}X$ of an equivalence relation $R \subseteq X \times X$ is the Egli-Milner lifting, i.e. $U \kappa_X(R) V \iff \forall x \in U \exists y \in V x R y \wedge \forall y \in V \exists x \in U x R y$. It is well-known that the greatest fixpoint of $\text{beh}_X = c^* \circ \kappa_X$ is precisely Park-Milner bisimilarity.

2. In the case $FX = \mathcal{D}X + 1$ and $\mathcal{V} = [0, 1]$, we obtain the lifting $\kappa_X(d) \in \mathbf{PMet}(\mathcal{D}X + 1)$ of a pseudometric $d \in \mathbf{PMet}(X)$. It is easy to see that $\kappa_X(d)(p_1, p_2)$ is the distance given by the classical Kantorovich lifting of d if $p_1, p_2 \in \mathcal{D}X$. If $p_1 = \surd = p_2$, then the distance is 0, otherwise 1. The least fixpoint (under the usual order on $[0, 1]$) of the behaviour function $\text{beh}_X = c^* \circ \kappa_X$ agrees with standard notions of bisimulation distance (e.g. [37]).

We conclude the section by showing that behaviour functions defined in this way are actually the ones obtained from the logic function. For the diagram underlying the proof see Figure 1.

► **Theorem 21.** *Assume that cl'_X is a subclosure of cl_X and compatible. For a set Θ_X of constants and a coalgebra $c: X \rightarrow FX$, the logic function $\log_X = \mathcal{P}(c^*) \circ \Lambda_X \circ \text{cl}'_X \cup \Theta_X$ induces $\text{beh}_X = c^* \circ \kappa_X \wedge \alpha_X(\Theta_X)$, i.e., $\alpha_X \circ \log_X \circ \gamma_X = \text{beh}_X$.*

Putting everything together via Theorem 4, if cl'_X is a compatible closure, then we have $\alpha_X(\mu \log_X) = \nu \text{beh}_X$, that is, logical conformance coincides with behavioural conformance.

5 Logics for Quantale-valued Simulation Distances

We next consider a quantitative modal logic \mathcal{L}_Λ that we show to be expressive for similarity distance (a behavioural directed metric) under certain conditions. Throughout this section, our working category \mathbf{C} is \mathbf{Set} , we have a fixed functor F on \mathbf{Set} , and a fixed quantale $\Omega = \mathcal{V}$ with distance $d_\mathcal{V} = [_, _]$ being the internal hom. Furthermore $\Phi X = \mathbf{DPMet}_\mathcal{V}(X)$. In this section we assume naturality of γ , which holds for the quantales given in Example 2.

$$\varphi \in \mathcal{L}_\Lambda ::= \bigwedge_{i \in I} \varphi_i \mid \varphi \otimes v \mid d_\mathcal{V}(v, \varphi) \mid [\lambda] \varphi \quad (\text{for } v \in \mathcal{V}, \lambda \in \Lambda, I \in \mathbf{Set})$$

This logic is the positive fragment of quantale-valued coalgebraic modal logic [40, 11], and generalizes logics for real-valued simulation distance [41] to the quantal setting. The first three operators are regarded as propositional operators of cl' , while the $[\lambda]$ are the modalities. We do not use explicit constants ($\Theta_X = \emptyset$), but note that constant truth \top is included as the empty meet. On a coalgebra $c: X \rightarrow FX \in \mathbf{Set}$, we interpret a formula φ as a function $\llbracket \varphi \rrbracket: X \rightarrow \mathcal{V}$ as usual (by structural induction on terms). Note that we do not allow negation, as we aim to characterize similarity distance. Disjunction could be included but, as in the two-valued case [38], is not needed to characterize simulation. Meet is infinitary, so the logic function log does not reach its least fixpoint in ω steps.

The next results show that the three operators (as well as join) are all non-expansive and hence cl' is a subclosure of cl (cf. Lemma 10), which moreover is compatible.

► **Proposition 22.** *Infinitary meets, infinitary joins, negative scaling $d_{\mathcal{V}}(v, _)$, and positive scaling $_ \otimes v$ (for $v \in \mathcal{V}$) are non-expansive.*

► **Proposition 23.** *If each $\lambda \in \Lambda$ is sup-preserving (i.e. $\lambda_X(\bigvee P) = \bigvee \mathcal{P}(\lambda_X)(P)$, for every subset $P \subseteq \mathbf{Set}(X, \mathcal{V})$), then the sub-closure cl' of cl as above is compatible.*

Diamond-like modalities (for powerset or fuzzy powerset) are typically sup-preserving. In such cases, Theorem 21 yields expressiveness of the above logic for similarity distance, defined as the greatest fixpoint of $\text{beh}_X(d) = c^* \circ \kappa_X(d)$ where κ_X is the directed Kantorovich lifting.

6 The Adjoint Setup in an Eilenberg-Moore Category

While we have seen in the examples of the previous sections that the framework can be instantiated to coalgebras living in \mathbf{Set} , thus providing Hennessy-Milner theorems for bisimilarity, we are now interested in tackling trace equivalences and trace metrics. To this end, we work in Eilenberg-Moore categories [33], which also allows us to determinize a coalgebra using the generalized powerset construction (cf. 2.3).


In particular, we instantiate the adjoint setup to the category $\mathbf{EM}(T)$ of T -algebras (for some monad T on \mathbf{C}), provide conditions guaranteeing compatibility, and characterize the behaviour function. Furthermore, taking inspiration from [32], we also introduce a general syntax for modal formulas that can be interpreted over coalgebras living in $\mathbf{EM}(T)$. As introduced in Section 2.3, we fix a coalgebra $c: X \rightarrow FTX$ living in \mathbf{C} and its determinization $c^\#: LX \rightarrow \tilde{F}LX$ in $\mathbf{EM}(T)$ via a distributive law $\zeta: TF \Rightarrow FT$.

We assume an indexed category $\Psi: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Pos}$ (that has fibred limits) and lift it to the category $\mathbf{EM}(T)$ of T -algebras by postcomposition, that is, $\Phi = \Psi \circ R$ (thus ensuring **A1**):

$$\mathbf{EM}(T)^{\text{op}} \xrightarrow{R} \mathbf{C}^{\text{op}} \xrightarrow{\Psi} \mathbf{Pos}.$$

Here, R is the forgetful functor in the free-forgetful adjunction $L \dashv R: \mathbf{C} \rightarrow \mathbf{EM}(T)$ from Section 2.3. To handle **A2**, we fix a truth value object $\Omega \in \mathbf{C}$ equipped with a T -algebra structure $o: T\Omega \rightarrow \Omega$ and $d_\Omega \in \Phi\Omega$. These assumptions ensure that Theorem 7 becomes applicable. We will denote the reindexing for Φ by $_*$, while we overload the notation and specify the reindexing in both \mathbf{C} and $\mathbf{EM}(T)$ by $_*$.

We focus on free algebras $LX = (TX, \mu_X)$ (over $X \in \mathbf{C}$) and apply Theorem 7 to the above-mentioned indexed category $\Psi \circ R$, which gives the adjoint situations depicted by shaded rectangles in Figure 2. We note that the middle hom-set $\mathbf{EM}(T)(LX, (\Omega, o))$ is isomorphic to $\mathbf{C}(X, \Omega)$ at the top left – due to the free-forgetful adjunction – with the bijection between the respective powersets witnessed by α', γ' . This allows us to define the logic function on the lattice $\mathcal{P}(\mathbf{C}(X, \Omega))$, which is a simpler structure than $\mathcal{P}(\mathbf{EM}(T)(LX, (\Omega, o)))$. In particular, formulas can then be evaluated directly on the state space X .



■ **Figure 2** The adjoint setup for algebras, where $\text{beh}_{LX} = (c^\#)^* \circ \kappa_{LX} \wedge \alpha_{LX} (\Theta_{LX})$ and $\text{log}'_X = \mathcal{P}(c^\bullet) \circ \Lambda'_X \cup \Theta_X$.

6.1 Logic and Behaviour Function for Coalgebras in Eilenberg-Moore

Recall from Section 4.3 that we need evaluation maps in the working category to define a logic function. So, to ensure **A3**, we assume a set Λ of evaluation maps for \tilde{F} , i.e., a family $(\tilde{F}(\Omega, o) \xrightarrow{ev_\lambda} (\Omega, o) \in \mathbf{EM}(T))_{\lambda \in \Lambda}$ of algebra homomorphisms. More concretely, a \mathbf{C} -arrow $ev_\lambda: F\Omega \rightarrow \Omega$ is such an algebra homomorphism if it satisfies $o \circ T ev_\lambda = ev_\lambda \circ Fo \circ \zeta_\Omega$.

As in Section 4, this induces a natural transformation Λ . Since every homomorphism is also a map in \mathbf{C} , we can define Λ'_X , the predicate lifting on $\mathcal{PC}(X, \Omega)$:

$$\mathcal{PC}(X, \Omega) \xrightarrow{\alpha'_X} \mathcal{PEM}(T)(LX, (\Omega, o)) \xrightarrow{\Lambda_{LX}} \mathcal{PEM}(T)(\tilde{F}LX, (\Omega, o)) \xrightarrow{\mathcal{P}(R)} \mathcal{PC}(FTX, \Omega).$$

Note that Λ' is a natural transformation (since α, Λ are natural transformations and R is a functor); on components it can be easily characterized as follows:

► **Lemma 24.** *We have that $\Lambda'_X(S) = \{ev_\lambda \circ Fo \circ FTh \mid h \in S\}$ where $S \subseteq \mathbf{C}(X, \Omega)$.*

That is, $\Lambda'_X(S)$ is obtained by first lifting the predicates from $\mathbf{C}(X, \Omega)$ to $\mathbf{C}(TX, \Omega)$ via the evaluation map $o: T\Omega \rightarrow \Omega$ and then to $\mathbf{C}(FTX, \Omega)$ via $ev_\lambda: F\Omega \rightarrow \Omega$. This process can be seen as applying a “double modality” for T and F .

We can now invoke the results of the previous chapter and assume that Λ is compatible with the closure induced by the adjunction, that is, we work without propositional operators (hence cl' , as mentioned in Assumption **A5**, is the identity), only constants, at first sight a strong property. We will however see in the next section that this always holds when F is a machine functor and we choose suitable evaluation maps.

The next theorem focusses on free algebras and is partly a corollary of Proposition 14 and Theorem 21. However there is a new component: instead of defining the logic function on (free) Eilenberg-Moore categories, reindexing via the determinized coalgebra $c^\#$, it is possible – as indicated above – to define it directly on arrows of type $X \rightarrow \Omega$ living in \mathbf{C} , reindexing with c . This coincides with the view that formulas should be evaluated on states in X rather than elements of TX . The diagram in Figure 2 outlines how to show this result.

► **Theorem 25.** *We fix a coalgebra $(c: X \rightarrow FTX) \in \mathbf{C}$. Assume that Λ_{LX} is compatible with the closure cl_{LX} and fix $\Theta_{LX} \subseteq \mathbf{EM}(T)(LX, \Omega)$ to ensure that **A4** holds.*

1. *Then the logic function $\text{log}_{LX} = \mathcal{P}((c^\#)^*) \circ \Lambda_{LX} \cup \Theta_{LX}$ is cl_{LX} -compatible.*

2. For the behaviour function $\text{beh}_{LX} = (c^\#)^* \circ \kappa_{LX} \wedge \alpha_{LX}(\Theta_{LX})$ (where $\kappa_{LX} = \alpha_{\bar{F}LX} \circ \Lambda_{LX} \circ \gamma_{LX}$), we have $\alpha_{LX}(\mu \log_{LX}) = \nu \text{beh}_{LX}$.
3. Now define another logic function $\log'_X = \mathcal{P}(c^\bullet) \circ \Lambda'_X \cup \Theta'_X$ with $\Theta_{LX} = \alpha'_X(\Theta'_X)$. It holds that $\alpha'_X \circ \log'_X \circ \gamma'_X = \log_{LX}$ and we obtain $\alpha_{LX}(\alpha'_X(\mu \log'_X)) = \nu \text{beh}_X$.

We hence consider a simple logic \mathcal{L}_{EM} for $\mathbf{EM}(T)$, where T is a monad on \mathbf{Set} :

$$\varphi \in \mathcal{L}_{EM} ::= \theta \mid [\lambda]\varphi \quad (\text{where } \theta \in \Theta, \lambda \in \Lambda)$$

Given a coalgebra $c: X \rightarrow FTX \in \mathbf{Set}$, each formula $\varphi \in \mathcal{L}_{EM}$ is interpreted as a function $\llbracket \varphi \rrbracket: X \rightarrow \Omega$, which is defined by structural induction as follows:

- Let $\varphi = \theta$. Then $\llbracket \varphi \rrbracket$ is given by a predefined constant $X \rightarrow \Omega$.
- Let $\varphi = [\lambda]\varphi'$. Then $\llbracket \varphi \rrbracket = ev_\lambda \circ Fo \circ FT\llbracket \varphi' \rrbracket \circ c$ (see definition of Λ'_X in Lemma 24).

► **Corollary 26.** *Under the requirements of Theorem 25, the logic \mathcal{L}_{EM} is expressive for the behavioural conformance beh_{LX} , i.e., $\alpha_{LX}(\alpha'_X(\{\llbracket \varphi \rrbracket \mid \varphi \in \mathcal{L}_{EM}\})) = \nu \text{beh}_{LX}$.*

6.2 The Machine Functor

Our next aim is to show that the machine functor has certain natural evaluation maps ensuring that the predicate lifting is cl-compatible (one of the conditions of Theorem 25). Throughout this section, we restrict ourselves to a monad T on \mathbf{Set} and fix the *machine functor* $M = _{}^\Sigma \times B$ with $\Sigma \in \mathbf{Set}$ and $(B, b) \in \mathbf{EM}(T)$. Since all monads in \mathbf{Set} are strong and B is endowed with a T -algebra structure $b: TB \rightarrow B$, there is a canonical distributive law ζ [18, Exercise 5.4.4]:

$$\zeta_X: T(X^\Sigma \times B) \xrightarrow{\langle a \rightarrow T(\pi_a \circ \pi_1), b \circ T\pi_2 \rangle} (TX)^\Sigma \times B, \quad (1)$$

where $(\pi_i)_{i \in \{1,2\}}$ are the usual projections and $\pi_a: X^\Sigma \rightarrow X$ is the evaluation map ($\pi_a(g) = g(a)$ where $g: \Sigma \rightarrow X$). Now let \tilde{M} be the lifting of M to $\mathbf{EM}(T)$, induced by the ζ . We observe that the evaluation maps suggested by it arise from natural transformations in the sense of Lemma 18.

- **Proposition 27. 1.** *Let $a \in \Sigma$. Then $\eta_a: \tilde{M} \Rightarrow Id$ given by the composite $MX \xrightarrow{\pi_1} X^\Sigma \xrightarrow{\pi_a} X$ is a natural transformation.*
2. *Let $f: (B, b) \rightarrow (\Omega, o)$ be a homomorphism. Then $\eta'_f: \tilde{M} \Rightarrow \Omega$ given by the composite $MX \xrightarrow{\pi_2} B \xrightarrow{f} \Omega$ is a natural transformation.*

Thus, $ev_a = \eta_a$, $ev_f = \eta'_f$ satisfy the properties of Lemma 18, and if each evaluation map is of this form, then Λ is cl-compatible.

6.3 Alternative Formulation of Kantorovich Lifting

The behaviour function in Section 6.1 is based on the generalized Kantorovich lifting κ [2], which works as follows: given a pseudometric d on Y (here $Y = TX$), generate *all* non-expansive functions $Y \rightarrow \Omega$ wrt. d , lift these functions to $FY \rightarrow \Omega$ and from there generate a pseudometric on FY . However, κ_{LX} – since it is defined in an Eilenberg-Moore category – works subtly differently: it takes *all non-expansive functions that are algebra homomorphisms*. This looks natural in the categorical setting, but may pose problems if we implement the procedures. The standard (probabilistic) Kantorovich lifting can for instance be computed based on the Kantorovich-Rubinstein duality, by determining optimal transport plans [39].

Here, both types of liftings coincide at least on relevant metrics. To show this result, we first define an alternative way of lifting, as opposed to defining the lifting on T -algebra maps. Applying Theorem 7 on Ψ (rather than on $\Psi \circ R$) gives the adjunction $\alpha_X^{\mathbf{C}} \dashv \gamma_X^{\mathbf{C}}: \mathcal{PC}(X, \Omega) \rightleftarrows \Psi X^{\text{op}}$. Now consider the lifting $\kappa_{TX}^{\mathbf{C}} = \alpha_{FTX}^{\mathbf{C}} \circ \Lambda_{TX} \circ \gamma_{TX}^{\mathbf{C}}$, where $\Lambda_{TX}: \mathcal{PC}(TX, \Omega) \rightarrow \mathcal{PC}(FTX, \Omega)$ is defined identically to Λ_{LX} .

► **Theorem 28.** *Assume that d is preserved by the co-closure, i.e. $d = \alpha_{LX}(\gamma_{LX}(d))$, the co-closure $\alpha_X^{\mathbf{C}} \circ \gamma_X^{\mathbf{C}}$ is the identity, and each evaluation map ev_λ arises from some natural transformation either of type $F \Rightarrow Id$ or $F \Rightarrow \Omega$. Then the two liftings $\kappa_{LX}, \kappa_{TX}^{\mathbf{C}}$ coincide on d , i.e., $\kappa_{LX}(d) = \kappa_{TX}^{\mathbf{C}}(d)$.*

The condition $d = \alpha_{LX}(\gamma_{LX}(d))$ is a necessary, but not a serious restriction: this property is typically satisfied by the \top metric and is preserved by the behaviour function. Hence during fixpoint iteration this invariant is preserved and the greatest fixpoints of beh_X based on either version of the Kantorovich lifting coincide (if the fixpoint is reached in ω steps).

In addition, if $\mathbf{C} = \mathbf{Set}$ and \mathcal{V} is an integral quantale, it can easily be shown that the co-closure $\alpha_X^{\mathbf{C}} \circ \gamma_X^{\mathbf{C}}$ is the identity (see Lemma 35 in the appendix). This enables us to concretely spell out the behaviour function for the case of the machine functor $M = _ \Sigma \times B$, provided that the conformances ΨX are (directed) pseudometrics.

► **Theorem 29.** *Assume that $\mathbf{C} = \mathbf{Set}$, $\Psi X = \mathbf{DPMet}_{\mathcal{V}}(X)$ (resp. $\Psi X = \mathbf{PMet}_{\mathcal{V}}(X)$) for an integral quantale \mathcal{V} and let $d_{\mathcal{V}} = [_, _]$ (resp. the symmetrized variant of $[_, _]$). Let $d: LX \times LX \rightarrow \mathcal{V}$ be a pseudometric that is preserved by the co-closure $\alpha_{LX} \circ \gamma_{LX}$. Assume that M is the machine functor and the family of evaluation maps is*

$$\{ev_a \mid a \in \Sigma\} \cup \{ev_f \mid f \in \mathcal{F} \subseteq \mathbf{EM}(T)((B, b), (\mathcal{V}, o))\}.$$

Then the corresponding behaviour function $\text{beh}_{LX}: \Psi(LX) \rightarrow \Psi(LX)$ is defined as follows: let $t_1, t_2 \in LX$ with $c^\#(t_i) = (b_i, g_i) \in B \times LX^\Sigma$:

$$\text{beh}_{LX}(d)(t_1, t_2) = \bigwedge_{a \in \Sigma} d(g_1(a), g_2(a)) \wedge \bigwedge_{f \in \mathcal{F}} d_{\mathcal{V}}(f(b_1), f(b_2)) \wedge \bigwedge_{\theta \in \Theta_{LX}} d_{\mathcal{V}}(\theta(t_1), \theta(t_2)),$$

The above function beh is co-continuous and fixpoint iteration terminates after ω steps.

7 Case Studies for the Linear-time Case

7.1 Workflow

We recall the parameters of our framework and set out a workflow that we follow in our case studies. Let F be a machine functor and T a monad on a category \mathbf{C} .

- Model systems as coalgebras of type $c: X \rightarrow FTX$ with a distributive law $\zeta: TF \Rightarrow FT$.
- Fix a truth value object $(\Omega, o) \in \mathbf{EM}(T)$ and $d_\Omega \in \Psi\Omega$.
- Define a fibration (indexed category) $\Phi = \Psi \circ R: \mathbf{EM}(T)^{\text{op}} \rightarrow \mathbf{Pos}$ by fixing an indexed category $\Psi: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Pos}$ to define the conformances.
- Fix a set Λ of evaluation maps (predicate liftings) as homomorphisms $\tilde{F}(\Omega, o) \rightarrow (\Omega, o)$.
- Fix a set of constants $\Theta_X \subseteq \mathbf{C}(X, \Omega)$.

Note that the last four conditions correspond to Assumptions **A1-A5**, which are necessary to set up a logic and a behaviour function beh as defined in Section 6.1 and guarantee expressiveness of the resulting logic. Whenever we choose $\Psi X = \mathbf{DPMet}_{\mathcal{V}}(X)$ or $\Psi X = \mathbf{PMet}_{\mathcal{V}}(X)$ for an integral quantale \mathcal{V} , we can rely on the characterization of the fixpoint equation in Theorem 29.

7.2 Trace Equivalence for Labelled Transition Systems

We spell out a simple case study: trace equivalence [38] for labelled transition systems. The main ingredients are summarized in the table below.

$\mathbf{C} = \mathbf{Set}, F = _^\Sigma, T = \mathcal{P}$ $(B = 1)$ $c: X \rightarrow (\mathcal{P}X)^\Sigma$ $c^\#: \mathcal{P}X \rightarrow (\mathcal{P}X)^\Sigma$	Logic: evaluation maps: ev_a (Prop. 27(1)) constants $\Theta_X = \{1\}$, constant 1-function formulas: $\varphi = [a_1] \cdots [a_n]1$
$\Omega = \mathbf{2}$ (Ex. 2(1)) $o: \mathcal{P}\mathbf{2} \rightarrow \mathbf{2}$ supremum $\Psi(X)$: equivalences on X d_Ω : equality on Ω	Behaviour function: $\mathbf{beh}_{\mathcal{P}X}: \Psi(\mathcal{P}X) \rightarrow \Psi(\mathcal{P}X)$ $\mathbf{beh}_{\mathcal{P}X}(R)(U, V) = (U = \emptyset \Leftrightarrow V = \emptyset) \wedge$ $\quad \forall a \in \Sigma (c^\#(U)(a), c^\#(V)(a)) \in R$

The modality $[a]$ boils down to the standard diamond modality (due to the definition of o); a state $x \in X$ satisfies $\varphi = [a_1] \cdots [a_n]1$ iff there exists a trace $a_1 \cdots a_n$ from x . The constant 1 is needed to start building formulas and to distinguish the empty set from a non-empty set on $LX = \mathcal{P}X$. (Note that $\Theta_{LX} = \alpha'_X(\Theta_X) = \{\tilde{1}\}$ with $\tilde{1}(Y) = 0$ iff $Y = \emptyset$.) Its role cannot be taken by a constant modality or an operator, since those have to be homomorphisms in $\mathbf{EM}(\mathcal{P})$, hence sup-preserving.

Expressiveness of trace logic $\mathcal{L}_{\mathbf{EM}}$ now directly follows from Corollary 26.

7.3 Trace Distance for Probabilistic Automata

A *probabilistic automaton* [30] is a quadruple (X, Σ, μ, p) where for each state $x \in X$ and each possible action $a \in \Sigma$ there is a probability distribution $\mu_{x,a}$ on the possible successors in X , and where each state $x \in X$ has a payoff value $p(x) \in [0, 1]$. Following [33, 34], we model them as coalgebras in the Eilenberg-Moore setting as detailed in the table below.

$\mathbf{C} = \mathbf{Set}, F = _^\Sigma \times [0, 1], T = \mathcal{D}$ $(B = [0, 1], b \text{ expectation})$ $c: X \rightarrow (\mathcal{D}X)^\Sigma \times [0, 1]$ $c^\#: \mathcal{D}X \rightarrow (\mathcal{D}X)^\Sigma \times [0, 1]$	Logic: evaluation maps: ev_a (Prop. 27(1)) $ev_*(f, r) = r$ (Prop. 27(2)) constants $\Theta_X = \emptyset$ formulas: $\varphi = [a_1] \cdots [a_n]*$
$\Omega = [0, 1]$ (Ex. 2(2)) $o: \mathcal{D}[0, 1] \rightarrow [0, 1]$ expectation $\Psi(X) = \mathbf{PMet}_{[0,1]}(X)$ $d_\Omega(r, s) = r - s $	Behaviour function: $\mathbf{beh}_{\mathcal{D}X}: \Psi(\mathcal{D}X) \rightarrow \Psi(\mathcal{D}X)$ $\mathbf{beh}_{\mathcal{D}X}(d)(p_1, p_2) = \max\{\sup_{a \in \Sigma} d(g_1(a), g_2(a)), d_\Omega(r_1, r_2)\}$

Thus, given a formula $\varphi = [a_1] \cdots [a_n]*$ and a state $x \in X$, $\llbracket \varphi \rrbracket(x)$ gives us the expected payoff after choosing actions according to the word $a_1 \cdots a_n$. The distance of two states x_1, x_2 is hence the supremum of the difference of payoffs, over all words.

Expressiveness again follows from Corollary 26.

7.4 Directed Fuzzy Trace Distance

We now consider directed trace distances for weighted transition systems over a generic quantale.

We work with the “fuzzy” monad $T = \mathcal{P}_\mathcal{V}$ (aka \mathcal{V} -valued powerset monad [15, Remark 1.2.3]) on \mathbf{Set} that is defined as $\mathcal{P}_\mathcal{V} = \mathcal{V}^X$ on objects and as $Tf(g)(y) = \bigvee_{f(x)=y} g(x)$ (for $f: X \rightarrow Y$) on arrows. Its unit $\eta_X: X \rightarrow \mathcal{P}_\mathcal{V}X$ is given by $\eta_X(x)(x') = 1$ if $x = x'$ and 0 (the empty join) otherwise. Multiplication $\mu_X: \mathcal{P}_\mathcal{V}\mathcal{P}_\mathcal{V}X \rightarrow \mathcal{P}_\mathcal{V}X$ is defined as

XX:16 Expressive Quantale-Valued Logics for Coalgebras: an Adjunction-Based Approach

$\mu_X(G)(x) = \bigvee_{g \in \mathcal{P}_{\mathcal{V}}X} G(g) \otimes g(x)$. Note that for $\mathcal{V} = 2$ (cf. Example 2(1)) T corresponds to the powerset monad \mathcal{P} .

$\mathbf{C} = \mathbf{Set}, F = _ \Sigma, T = \mathcal{P}_{\mathcal{V}}$ $(B = 1)$ $c: X \rightarrow (\mathcal{P}_{\mathcal{V}}X)^{\Sigma}$ $c^{\#}: \mathcal{P}_{\mathcal{V}}X \rightarrow (\mathcal{P}_{\mathcal{V}}X)^{\Sigma}$	Logic: evaluation maps: ev_a (Prop. 27(1)) constants $\Theta_X = \{1\}$, constant 1-function formulas: $\varphi = [a_1] \cdots [a_n]1$
$\Omega = \mathcal{V}, o: \mathcal{P}_{\mathcal{V}}\mathcal{V} \rightarrow \mathcal{V}$ $g \mapsto \bigvee_{v \in \mathcal{V}} g(v) \otimes v$ $\Psi(X) = \mathbf{DPMet}_{\mathcal{V}}(X)$ $d_{\Omega}(v, v') = [v, v']$	Behaviour function: $\text{beh}_{\mathcal{P}_{\mathcal{V}}X}: \Psi(\mathcal{P}_{\mathcal{V}}X) \rightarrow \Psi(\mathcal{P}_{\mathcal{V}}X)$ $\text{beh}_{\mathcal{P}_{\mathcal{V}}X}(d)(g_1, g_2) = \bigwedge_{a \in \Sigma} d(c^{\#}(g_1)(a), c^{\#}(g_2)(a)) \wedge$ $\left[\bigvee_{x \in X} g_1(x), \bigvee_{x \in X} g_2(x) \right]$

Evaluating a formula $\varphi = [a_1] \cdots [a_n]1$ on a state $x_0 \in X$ results in

$$\llbracket \varphi \rrbracket(x_0) = \bigvee \left\{ \bigotimes_{0 \leq i < n} c(x_i)(a_{i+1})(x_{i+1}) \mid x_1 \dots x_n \in X^n \right\}.$$

This follows directly from structural induction on φ , from distributivity and from the evaluation of the modality $[a]\varphi'$:

$$\llbracket [a]\varphi' \rrbracket(x) = \bigvee_{y \in X} c(x)(a)(y) \otimes \llbracket \varphi' \rrbracket(y).$$

Intuitively we check how well x can match the trace $a_1 \dots a_n$, where $c(x)(a)(y)$ measures the degree to which x can make an a -transition to y .

The second part of the minimum in the definition of beh stems from the constants $\Theta_X = \{1\}$, since $\Theta_{LX} = \alpha'_X(\Theta_X) = \{\bar{1}\}$ with $\bar{1}(h) = \bigvee_{x \in X} h(x)$ for $h: X \rightarrow \mathcal{V}$. Without it, the fixpoint iteration would stabilize at the constant 1-pseudometric.

Expressiveness again follows from Corollary 26. Expressiveness of a logic for *symmetric* fuzzy trace distance has already been shown in previous work [12].

8 Conclusion

Related work. By now there is a large number of papers considering coalgebraic semantics beyond branching-time, for instance [14, 33, 26, 6]. Furthermore in the same period quite a wealth of results on the treatment of behavioural metrics in coalgebraic generality has been published [37, 2, 22, 11]. However, there is little work combining both linear-time semantics and behavioural metrics in the setting of coalgebra. In that respect we want to mention [12] that is based on the graded monad framework [26] and which investigates exactly this combination. However, different from the present paper, the focus is on the expressiveness of the logics with respect a graded semantics (that intuitively specifies the traces of a state). Hence, using the classification of the introduction, it studies the relationship of (i) and (ii).

Unlike other approaches, our main focus is on exploiting an adjunction (Galois connection) and fixpoint preservation results to obtain Hennessy-Milner theorems “for free”. We start by setting up a logic, characterizing the behavioural equivalence, and investigate under which circumstances we can derive a corresponding fixpoint characterization. The fixpoint equation might be defined on an infinite state space, but often there are finitary techniques that can be employed, such as reducing the state space to a finite subset, linear programming, up-to techniques, etc. In particular, for systems as in Section 7.4 we are working on promising results (based on [3]) for deriving bounds for behavioural distances via finite witnesses using up-to techniques, even for infinite state spaces. The algorithmic angle of our approach is not

yet fully worked out in the present paper but establishing fixpoint equations as we do here is a necessary first step in this direction.

Note that our concept deviates from the dual adjunction approach [20, 23, 24, 27] to coalgebraic modal logic. There the functor on the “logic universe” characterizes the *syntax* of the logics, while the semantics is instead given by a natural transformation. Nevertheless, it complements (at least when restricted to the classical case of Boolean predicates) the recent approach [36] that combines fibrations in the dual adjunction setup since having contravariant Galois connections between fibres (at a ‘local’ level) is equivalent to having dual adjunctions between certain fibred categories (at a ‘global’ level). It is unclear how to establish this correspondence in the setting of quantitative \mathcal{V} -valued predicates.

Future work. Currently the operators of the logic, given by cl' , are rather generic, although we instantiated them in special cases to ensure expressiveness (see in particular Sections 5 and 6.2). We envision a general theory to ensure expressiveness of the logics, similar to Post’s functional completeness theorem [28], which characterizes complete sets of operators for the boolean case. This question is strongly related to the notion of an approximating family in [22] that has again close connections to compatibility as discussed in [4].

We will also study the condition requiring that a conformance (pseudometric) is preserved by the co-closure ($d = \alpha_{LX}(\gamma_{LX}(d))$). Previous results [4] suggest that this is related to the notion of (metric) congruence, as e.g. defined in [5], but the connection seems to be non-trivial.

Another avenue of research is to further investigate the quantale-valued logic for the branching case introduced in Section 5, to extend it to the undirected case and restrict to finitary operators.

References

- 1 Jiří Adámek, Horst Herrlich, and George E. Strecker. *Abstract and concrete categories: The joy of cats*. Wiley, 1990. Republished in: Reprints in Theory and Applications of Categories, No. 17 (2006) pp. 1–507.
- 2 Paolo Baldan, Filippo Bonchi, Henning Kerstan, and Barbara König. Coalgebraic behavioral metrics. *Logical Methods in Computer Science*, 14(3), 2018. Selected Papers of the 6th Conference on Algebra and Coalgebra in Computer Science (CALCO 2015).
- 3 Paolo Baldan, Barbara König, and Tommaso Padoan. Abstraction, up-to techniques and games for systems of fixpoint equations. In *Proc. of CONCUR ’20*, volume 171 of *LIPICs*, pages 25:1–25:20. Schloss Dagstuhl – Leibniz Center for Informatics, 2020.
- 4 Harsh Beohar, Sebastian Gurke, Barbara König, and Karla Messing. Hennessy-Milner theorems via Galois connections. In *Proc. of CSL ’23*, volume 252 of *LIPICs*, pages 12:1–12:18. Schloss Dagstuhl – Leibniz Center for Informatics, 2023.
- 5 Filippo Bonchi, Alexandra Silva, and Ana Sokolova. The Power of Convex Algebras. In *Proc. of CONCUR ’17*, volume 85 of *LIPICs*, pages 23:1–23:18. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2017.
- 6 Corina Cirstea. From branching to linear time, coalgebraically. *Fundamenta Informaticae*, 150(3-4):379–406, 2017.
- 7 Patrick Cousot and Radhia Cousot. Systematic design of program analysis frameworks. In *Proc. of POPL ’79*, pages 269–282. ACM Press, 1979.
- 8 Patrick Cousot and Radhia Cousot. Temporal abstract interpretation. In Mark N. Wegman and Thomas W. Reps, editors, *Proc. of POPL ’00*, pages 12–25. ACM, 2000.
- 9 Josée Desharnais, Vineet Gupta, Radha Jagadeesan, and Prakash Panangaden. Metrics for labelled Markov processes. *Theoretical Computer Science*, 318:323–354, 2004.

- 10 Ulrich Dorsch, Stefan Milius, and Lutz Schröder. Graded monads and graded logics for the linear time - branching time spectrum. In Wan J. Fokkink and Rob van Glabbeek, editors, *Concurrency Theory, CONCUR 2019*, volume 140 of *LIPICs*, pages 36:1–36:16. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2019.
- 11 Jonas Forster, Sergey Goncharov, Dirk Hofmann, Pedro Nora, Lutz Schröder, and Paul Wild. Quantitative Hennessy-Milner Theorems via Notions of Density. In *Proc. of CSL '23*, volume 252 of *LIPICs*, pages 22:1–22:20, 2023.
- 12 Jonas Forster, Lutz Schröder, Paul Wild, Harsh Beohar, Sebastian Gurke, Barbara König, and Karla Messing. Graded semantics and graded logics for Eilenberg-Moore coalgebras, 2023. arXiv:2307.14826.
- 13 Alessandro Giacalone, Chi-Chang Jou, and Scott A. Smolka. Algebraic reasoning for probabilistic concurrent systems. In Manfred Broy and Cliff B. Jones, editors, *Programming concepts and methods*, pages 443–458. North-Holland, 1990.
- 14 Ichiro Hasuo, Bart Jacobs, and Ana Sokolova. Generic trace semantics via coinduction. *Logical Methods in Computer Science*, 3(4:11):1–36, 2007.
- 15 Dirk Hofmann, Gavin J. Seal, and Walter Tholen, editors. *Lax algebras*, page 145–283. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2014.
- 16 Bart Jacobs. *Categorical Logic and Type Theory*, volume 141 of *Studies in Logic and the Foundations of Mathematics*. Elsevier, 1st edition, Jan 1999.
- 17 Bart Jacobs. Predicate logic for functors and monads. Available from author’s website, 2010.
- 18 Bart Jacobs. *Introduction to Coalgebra: Towards Mathematics of States and Observation*. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 2016.
- 19 Bart Jacobs, Alexandra Silva, and Ana Sokolova. Trace semantics via determinization. In *Proc. of CMCS '12*, pages 109–129. Springer, 2012. LNCS 7399.
- 20 Bartek Klin. Coalgebraic modal logic beyond sets. In *Proc. of MFPS '07*, volume 173 of *ENTCS*, pages 177–201, 2007.
- 21 Yuichi Komorida, Shin-ya Katsumata, Nick Hu, Bartek Klin, and Ichiro Hasuo. Codensity games for bisimilarity. In *Proc. of LICS '19*, pages 1–13, 2019.
- 22 Yuichi Komorida, Shin-ya Katsumata, Clemens Kupke, Jurriaan Rot, and Ichiro Hasuo. Expressivity of quantitative modal logics: Categorical foundations via codensity and approximation. In *Proc. LICS '21*, pages 1–14. IEEE, 2021.
- 23 Clemens Kupke and Dirk Pattinson. Coalgebraic semantics of modal logics: An overview. *Theoretical Computer Science*, 412:5070–5094, 2011.
- 24 Clemens Kupke and Jurriaan Rot. Expressive logics for coinductive predicates. In *Proc. of CSL '20*, volume 152 of *LIPICs*, pages 26:1–26:18. Schloss Dagstuhl – Leibniz Center for Informatics, 2020.
- 25 F. William Lawvere. Metric spaces, generalized logic, and closed categories. *Rendiconti del Seminario Matematico e Fisico di Milano*, 43(1):135–166, 1973. Republished in Reprints in Theory and Applications of Categories 1 (2002), 1–37.
- 26 Stefan Milius, Dirk Pattinson, and Lutz Schröder. Generic trace semantics and graded monads. In *Proc. of CALCO '15*, volume 35 of *LIPICs*, pages 253–269. Schloss Dagstuhl – Leibniz-Zentrum fuer Informatik, 2015.
- 27 Dusko Pavlovic, Michael Mislove, and James Worrell. Testing semantics: Connecting processes and process logics. In *Proc. of AMAST '06*, pages 308–322. Springer, 2006. LNCS 4019.
- 28 Francis Jeffrey Pelletier and Norman M. Martin. Post’s functional completeness theorem. *Notre Dame Journal of Formal Logic*, 31(2), 1990.
- 29 Damien Pous. Complete lattices and up-to techniques. In *Proc. of APLAS '07*, pages 351–366. Springer, 2007. LNCS 4807.
- 30 Michael O. Rabin. Probabilistic automata. *Information and Control*, 6(3):230–245, 1963.
- 31 Jan Rutten. Universal coalgebra: A theory of systems. *Theoretical Computer Science*, 249:3–80, 2000.

- 32 Lutz Schröder. Expressivity of coalgebraic modal logic: The limits and beyond. *Theoretical Computer Science*, 390(2):230–247, 2008. Foundations of Software Science and Computational Structures.
- 33 Alexandra Silva, Filippo Bonchi, Marcello M. Bonsangue, and Jan J. M. M. Rutten. Generalizing the powerset construction, coalgebraically. In *Proc. of FSTTCS '10*, volume 8, pages 272–283. Schloss Dagstuhl – Leibniz Center for Informatics, 2010. LIPIcs.
- 34 Alexandra Silva, Filippo Bonchi, Marcello M. Bonsangue, and Jan J. M. M. Rutten. Generalizing determinization from automata to coalgebras. *Logical Methods in Computer Science*, 9(1:09), 2013.
- 35 Alfred Tarski. A lattice-theoretical fixpoint theorem and its applications. *Pacific Journal of Mathematics*, 5:285–309, 1955.
- 36 Ruben Turkenburg, Harsh Beohar, Clemens Kupke, and Jurriaan Rot. Forward and Backward Steps in a Fibration. In Paolo Baldan and Valeria de Paiva, editors, *10th Conference on Algebra and Coalgebra in Computer Science (CALCO 2023)*, volume 270 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 6:1–6:18, Dagstuhl, Germany, 2023. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- 37 Franck van Breugel and James Worrell. A behavioural pseudometric for probabilistic transition systems. *Theoretical Computer Science*, 331:115–142, 2005.
- 38 Rob van Glabbeek. The linear time – branching time spectrum I. In J.A. Bergstra, A. Ponse, and S.A. Smolka, editors, *Handbook of Process Algebra*, chapter 1, pages 3–99. Elsevier, 2001.
- 39 Cédric Villani. *Optimal Transport – Old and New*, volume 338 of *A Series of Comprehensive Studies in Mathematics*. Springer, 2009.
- 40 Paul Wild and Lutz Schröder. A quantified coalgebraic van Benthem theorem. In *Proc. of FOSSACS '21*, volume 12650 of *LNCS*, pages 551–571. Springer, 2021.
- 41 Paul Wild and Lutz Schröder. Characteristic logics for behavioural hemimetrics via fuzzy lax extensions. *Logical Methods in Computer Science*, 18(2), 2022.
- 42 Paul Wild, Lutz Schröder, Dirk Pattinson, and Barbara König. A van Benthem theorem for fuzzy modal logic. In Anuj Dawar and Erich Grädel, editors, *Logic in Computer Science, LICS 2018*, pages 909–918. ACM, 2018.

A Proofs for Section 2 (Preliminaries)

► **Lemma 30.** *Let \mathcal{V} be a quantale. Then for $z \in \mathcal{V}$ it holds that $[1, z] = z$. If in addition the quantale is integral, we have $[z, 1] = 1$*

Proof. As right adjoint of the associative operation the internal hom of a quantale is concretely defined as follows:

$$[y, z] = \bigvee \{x' \in \mathcal{V} \mid x' \otimes y \leq z\}.$$

Hence we can simply use this characterization and obtain:

$$[1, z] = \bigvee \{x' \in \mathcal{V} \mid x' \otimes 1 \leq z\} = \bigvee \{x' \in \mathcal{V} \mid x' \leq z\} = z.$$

If in addition 1 is the largest element, we have:

$$[z, 1] = \bigvee \{x' \in \mathcal{V} \mid x' \otimes z \leq 1\} = \bigvee \mathcal{V} = 1. \quad \blacktriangleleft$$

► **Lemma 31.** *Let $d \in \mathbf{DPMet}_{\mathcal{V}}(X)$ and define $d_{\mathcal{V}} = [_, _]: \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$. Fix $y \in X$ and define a function $h: X \rightarrow \mathcal{V}$ as $h(x) = d(y, x)$. Then h is non-expansive wrt. $d, d_{\mathcal{V}}$.*

The same holds if $d \in \mathbf{PMet}_{\mathcal{V}}(X)$ and $d_{\mathcal{V}}$ is defined as $d_{\mathcal{V}}(x, y) = [x, y] \wedge [y, x]$.

Whenever \mathcal{V} is integral, in both cases it holds for all $x \in X$ that $d_{\mathcal{V}}(h(y), h(x)) = d(y, x)$

Proof. We start with a directed metric $d \in \mathbf{DPMet}_{\mathcal{V}}(X)$ and $d_{\mathcal{V}} = [_, _]$ and our aim is to show that $d \preceq d_{\mathcal{V}} \circ (h \times h)$:

$$\begin{aligned} d \preceq d_{\mathcal{V}} \circ (h \times h) &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \leq d_{\mathcal{V}}(h(x_1), h(x_2)) \\ &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \leq [h(x_1), h(x_2)] \\ &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \otimes h(x_1) \leq h(x_2) \\ &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \otimes d(y, x_1) \leq d(y, x_2) \\ &\iff \forall_{x_1, x_2 \in X} d(y, x_1) \otimes d(x_1, x_2) \leq d(y, x_2) \end{aligned}$$

Here we use the fact that $x \otimes y \leq z \iff x \leq [y, z]$ (adjunction) and the commutativity of \otimes . The last statement is true due to the triangle inequality and hence non-expansiveness of h follows.

Whenever $d_{\Omega} = [_, _]$, $d_{\mathcal{V}}(h(y), h(x))$ simplifies to $d_{\mathcal{V}}(d(y, y), d(y, x)) = [1, d(y, x)] = d(y, x)$, using the fact that \mathcal{V} is integral (and hence $d(y, y) = 1$) and Lemma 30.

We now take an undirected metric $d \in \mathbf{PMet}_{\mathcal{V}}(X)$ and define $d_{\mathcal{V}}(x, y) = [x, y] \wedge [y, x]$. Again our aim is to show that $d \preceq d_{\mathcal{V}} \circ (h \times h)$:

$$\begin{aligned} d \preceq d_{\mathcal{V}} \circ (h \times h) &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \leq d_{\mathcal{V}}(h(x_1), h(x_2)) \\ &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \leq [h(x_1), h(x_2)] \wedge [h(x_2), h(x_1)] \\ &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \leq [h(x_1), h(x_2)] \text{ and } d(x_1, x_2) \leq [h(x_2), h(x_1)] \\ &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \otimes h(x_1) \leq h(x_2) \text{ and } d(x_1, x_2) \otimes h(x_2) \leq h(x_1) \\ &\iff \forall_{x_1, x_2 \in X} d(x_1, x_2) \otimes d(y, x_1) \leq d(y, x_2) \text{ and } d(x_1, x_2) \otimes d(y, x_2) \leq d(y, x_1) \\ &\iff \forall_{x_1, x_2 \in X} d(y, x_1) \otimes d(x_1, x_2) \leq d(y, x_2) \text{ and } d(y, x_2) \otimes d(x_2, x_1) \leq d(y, x_1) \end{aligned}$$

Here we use again the adjunction and the commutativity of \otimes . In addition we need symmetry of d . The last statements are again true due to the triangle inequality and hence non-expansiveness of h follows.

In this case $d_{\mathcal{V}}(h(y), h(x))$ simplifies to the undirected case we have $d_{\mathcal{V}}(d(y, y), d(y, x)) = [1, d(y, x)] \wedge [d(y, x), 1] = d(y, x) \wedge 1 = d(y, x)$, based on Lemma 30, where we again use that the quantale is integral. ◀

B

 Proofs for Section 3 (Adjoint Logic: the General Framework)

Combining logic functions results in the combination of the corresponding behaviour functions, which is essential in establishing Hennessy-Milner theorems compositionally.

► **Proposition 32** ([4]). *Let $\log_1, \log_2, \text{cl}: \mathbb{L} \rightarrow \mathbb{L}$ be monotone functions on a complete lattice \mathbb{L} such that \log_1, \log_2 are cl-compatible. Then $\log_1 \sqcup \log_2$ and $\log_1 \circ \log_2$ are also cl-compatible.*

Further, let $\text{beh}_i = \alpha \circ \log_i \circ \gamma$ be the induced behaviour functions ($i = 1, 2$). Then the behaviour functions of $\log_1 \sqcup \log_2$ and $\log_1 \circ \log_2$ are $\text{beh}_1 \sqcup \text{beh}_2$ and $\text{beh}_1 \circ \text{beh}_2$, respectively. Every constant function k and the identity are c-compatible. Their behaviour functions are the constant function $b \mapsto \alpha(\ell)$ (where ℓ is the constant value of k) respectively the co-closure $\alpha \circ \gamma$.

C

 Proofs for Section 4 (Adjoint Logic for Coalgebras)

► **Theorem 7.** *Let X be an object of \mathbf{C} . Under Assumptions **A1** and **A2**, there is a dual adjoint situation (contravariant Galois connection) $\alpha_X \dashv \gamma_X$ between the underlying fibres:*

$$\begin{aligned} \alpha_X: \mathcal{P}(\mathbf{C}(X, \Omega)) &\rightarrow \Phi(X)^{op} & S \subseteq \mathbf{C}(X, \Omega) &\mapsto \bigwedge_{k \in S} k^*(d_\Omega) \\ \gamma_X: \Phi(X)^{op} &\rightarrow \mathcal{P}(\mathbf{C}(X, \Omega)) & d \in \Phi X &\mapsto \{k \in \mathbf{C}(X, \Omega) \mid d \preceq k^*(d_\Omega)\}. \end{aligned}$$

More concretely: α_X, γ_X are both antitone ($S \subseteq S' \implies \alpha_X(S) \succeq \alpha_X(S')$, $d \preceq d' \implies \gamma_X(d) \supseteq \gamma_X(d')$) and we have $d \preceq \alpha_X(S) \iff S \subseteq \gamma_X(d)$ for $d \in \Phi X$ and $S \in \mathcal{P}\mathbf{C}(X, \Omega)$.

Proof. First we argue that α_X, γ_X are order preserving (in this case antitone). For α_X , let $S \subseteq S'$. Clearly,

$$\alpha_X(S) = \bigwedge_{h \in S} h^*d_\Omega \succeq \bigwedge_{h \in S'} h^*d_\Omega = \alpha_X(S').$$

For γ_X , let $d \preceq d'$ and let $h \in \gamma_X(d')$. I.e., $d' \preceq h^*d_\Omega$; thus, $d \preceq h^*d_\Omega$. Hence, $h \in \gamma_X(d)$.

Now it remains to show that $d \preceq \alpha_X(S) \iff S \subseteq \gamma_X(d)$, for every $d \in \Phi X$ and $S \in \mathcal{P}\mathbf{C}(X, \Omega)$.

$$\begin{aligned} d \preceq \alpha_X(S) &\iff d \preceq \bigwedge_{h \in S} h^*d_\Omega \iff \forall_{h \in S} d \preceq h^*d_\Omega \\ &\iff \forall_{h \in S} h \in \gamma_X(d) \iff S \subseteq \gamma_X(d). \end{aligned} \quad \blacktriangleleft$$

► **Lemma 10.** *The right adjoint γ in Theorem 7 is laxly natural, i.e. $\mathcal{P}(f^\bullet) \circ \gamma_Y \subseteq \gamma_X \circ f^*$ for $f: X \rightarrow Y \in \mathbf{C}$. If γ is natural (i.e. the inclusion is an equality) and \mathbf{C} has products, then $\gamma_X(\alpha_X(S)) = \mathcal{P}(\langle S^\bullet \rangle)(\gamma_{\Omega^S}(d_{\Omega^S}))$, where $d_{\Omega^S} = \bigwedge_{k \in S} \pi_k^*d_\Omega$.*

Proof. In order to show $\mathcal{P}(f^\bullet) \circ \gamma_Y \subseteq \gamma_X \circ f^*$ we fix $d \in \Psi Y$ and $h \in \mathcal{P}(f^\bullet)(\gamma_Y(d))$. This implies that $h = k \circ f$ where $k \in \gamma_Y(d)$, hence $d \preceq k^*d_\Omega$. From this we can infer $h^*d_\Omega = (k \circ f)^*d_\Omega = f^*(k^*d_\Omega) \succeq f^*d$. Hence $h \in \gamma_X(f^*d)$ follows.

XX:22 Expressive Quantale-Valued Logics for Coalgebras: an Adjunction-Based Approach

Now suppose that γ is natural. We observe that $\langle S \rangle^* d_\Omega^S = \alpha_X S$ for every $S \subseteq \mathbf{C}(X, \Omega)$.

$$\begin{aligned}
 \langle S \rangle^* d_\Omega^S &= \langle S \rangle^* \bigwedge_{k \in S} \pi_k^* d_\Omega && \because \Phi \text{ has indexed limits} \\
 &= \bigwedge_{k \in S} \langle S \rangle^* \pi_k^* d_\Omega && \because \text{reindexing } (g \circ f)^* = f^* \circ g^* \\
 &= \bigwedge_{k \in S} (\pi_k \circ \langle S \rangle)^* d_\Omega && \because \text{universal property of products} \\
 &= \bigwedge_{k \in S} k^* d_\Omega = \alpha_X S.
 \end{aligned}$$

Now the result follows directly from the naturality of γ , i.e.,

$$\mathcal{P}(\langle S \rangle^\bullet) \gamma_\Omega^S (d_\Omega^S) = \gamma_X \langle S \rangle^* (d_\Omega^S) = \gamma_X \alpha_X S. \quad \blacktriangleleft$$

► **Proposition 11.** *A family of evaluation maps $(ev_\lambda)_{\lambda \in \Lambda}$ induces a natural transformation $\Lambda: \mathcal{PC}(_, \Omega) \Rightarrow \mathcal{PC}(F_, \Omega)$ given by $S \mapsto \{\lambda_X(h) \mid \lambda \in \Lambda, h \in S\}$, where $\lambda_X(h) = ev_\lambda \circ Fh$.*

Proof. Naturality of Λ amounts to showing that, given $f: X \rightarrow Y$ in \mathbf{C} and $S' \subseteq \mathbf{C}(X, \Omega)$:

$$\begin{aligned}
 \mathcal{P}(Ff)^\bullet(\Lambda_Y S') &= \mathcal{P}(Ff)^\bullet \{ev_\lambda \circ Fh \mid h \in S'\} = \{ev_\lambda \circ Fh \circ Ff \mid h \in S'\} \\
 &= \{ev_\lambda \circ F(h \circ f) \mid h \in S'\} = \{ev_\lambda \circ Fk \mid k \in \mathcal{P}f^\bullet(S')\} = \Lambda_X(\mathcal{P}f^\bullet(S')) \quad \blacktriangleleft
 \end{aligned}$$

► **Lemma 33.** *Let $cl_X = \gamma_X \circ \alpha_X$ be the closure obtained from the Galois connection in Theorem 7 and let $cl'_X: \mathcal{PC}(X, \Omega) \rightarrow \mathcal{PC}(X, \Omega)$ be some closure. Then $cl'_X \subseteq cl_X$ is equivalent to $cl_X = cl'_X \circ cl_X$ and equivalent to $cl_X = cl_X \circ cl'_X$.*

Proof. Assume that $cl'_X \subseteq cl_X$. We obtain $cl_X \subseteq cl'_X \circ cl_X$, $cl_X \subseteq cl_X \circ cl'_X$ trivially, since cl'_X is a closure and satisfies $id_{\mathbf{C}(X, \Omega)} \subseteq cl'_X$. Furthermore

$$cl'_X \circ cl_X \subseteq cl_X \circ cl_X = cl_X,$$

similarly for the other equality ($cl_X = cl_X \circ cl'_X$).

Now assume that $cl'_X \circ cl_X = cl_X$. Then, again by using the fact that cl_X is a closure:

$$cl'_X \subseteq cl'_X \circ cl_X = cl_X.$$

If on the other hand $cl_X = cl_X \circ cl'_X$ we can argue similarly by observing

$$cl'_X \subseteq cl_X \circ cl'_X = cl_X. \quad \blacktriangleleft$$

► **Proposition 14.** *For a given compatible closure cl'_X , the above logic function \log_X is cl_X -compatible, i.e., $\log_X \circ cl_X \subseteq cl_X \circ \log_X$.*

Proof. We use Proposition 32 to prove compatibility of \log in a compositional manner. Let $\log_1 = \mathcal{P}(c^\bullet) \circ \Lambda_X \circ cl'_X$ and $\log_2 = \Theta_X$, i.e., $\log_X = \log_1 \cup \log_2$.

First, we derive

$$\begin{aligned}
 \log_1 \circ cl_X &= \mathcal{P}(c^\bullet) \circ \Lambda_X \circ cl'_X \circ cl_X && (\because cl'_X \text{ is compatible}) \\
 &\subseteq \mathcal{P}(c^\bullet) \circ cl_{FX} \circ \Lambda_X \circ cl'_X \\
 &= \mathcal{P}(c^\bullet) \circ \gamma_{FX} \circ \alpha_{FX} \circ \Lambda_X \circ cl'_X && (\because \text{lax naturality of } \gamma) \\
 &\subseteq \gamma_X \circ c^* \circ \alpha_{FX} \circ \Lambda_X \circ cl'_X && (\because \text{naturality of } \alpha) \\
 &= \gamma_X \circ \alpha_X \circ \mathcal{P}(c^\bullet) \circ \Lambda_X \circ cl'_X \\
 &= cl_X \circ \log_1.
 \end{aligned}$$

Note that \log_2 is a constant function (thus cl -compatible) and thus Proposition 32 is applicable. ◀

► **Lemma 15.** *Let cl'_X be a subclosure of cl_X . It holds that cl'_X is compatible if and only if $\alpha_{FX} \circ \Lambda_X \circ \text{cl}'_X \preceq \alpha_{FX} \circ \Lambda_X \circ \text{cl}_X$.*

Proof. Since we assume that cl'_X is a subclosure of cl_X , compatibility of cl' simplifies to

$$\Lambda \circ \text{cl}_X \subseteq \text{cl}_{FX} \circ \Lambda_X \circ \text{cl}'_X = \gamma_{FX} \circ \alpha_{FX} \circ \Lambda_X \circ \text{cl}'_X.$$

Due to the properties of a Galois connection this is equivalent to

$$\alpha_{FX} \circ \Lambda_X \circ \text{cl}'_X \preceq \alpha_{FX} \circ \Lambda_X \circ \text{cl}_X. \quad \blacktriangleleft$$

► **Lemma 17.** *Let cl'_X be a subclosure of cl_X . Then cl'_X is compatible if and only if the depth-1 self-separation property holds.*

Proof. We first show that the depth-1 self-separation property implies compatibility of cl'_X , by proving the inequality of Lemma 15. So let $T \subseteq \mathbf{C}(X, \Omega)$. Define $S = \text{cl}'_X(T)$ and observe that S is closed under cl' . Put $d = \alpha_X(S)$; by definition, S is initial for d . By the depth-1 self-separation property, it follows that $\Lambda_X(S)$ is initial for $\kappa_X(d) = \alpha_{FX}(\Lambda_X(\gamma_X(d)))$. Hence

$$\begin{aligned} \alpha_{FX}(\Lambda_X(\text{cl}'_X(T))) &= \alpha_{FX}(\Lambda_X(S)) = \alpha_{FX}(\Lambda_X(\gamma_X(d))) = \alpha_{FX}(\Lambda_X(\gamma_X(\alpha_X(S)))) \\ &= \alpha_{FX}(\Lambda_X(\text{cl}_X(\text{cl}'_X(T)))) = \alpha_{FX}(\Lambda_X(\text{cl}_X(T))), \end{aligned}$$

using the fact that cl' is a subclosure of cl . From Lemma 15 we infer compatibility of cl'_X .

We now show the other direction, i.e., compatibility of cl' implies depth-1 self-separation. Let $S \subseteq \mathbf{C}(X, \Omega)$ be a set that is closed under cl'_X and S is initial for d , i.e., $\alpha_X(d) = S$. We have to show that $\Lambda_X(S)$ is initial for $\kappa_X(d)$, i.e., that $\alpha_{FX}(\Lambda_X(S)) = \alpha_{FX}(\Lambda_X(\gamma_X(d)))$. The inequality $\alpha_{FX}(\Lambda_X(S)) \succeq \alpha_{FX}(\Lambda_X(\gamma_X(d)))$ follows trivially from the properties of the (contra-variant) Galois connection, since $S \subseteq \gamma_X(\alpha_X(S)) = \gamma_X(d)$. For the other direction we use Lemma 15, which gives us:

$$\begin{aligned} \alpha_{FX}(\Lambda_X(S)) &= \alpha_{FX}(\Lambda_X(\text{cl}'_X(S))) \preceq \alpha_{FX}(\Lambda_X(\text{cl}_X(S))) \\ &= \alpha_{FX}(\Lambda_X(\gamma_X(\alpha_X(S)))) = \alpha_{FX}(\Lambda_X(\gamma_X(d))). \quad \blacktriangleleft \end{aligned}$$

► **Lemma 18.** *If each evaluation map ev_λ arises as a natural transformation $\eta: F \Rightarrow \text{Id}$ or $\eta: F \Rightarrow \Omega$ (Ω is the constant functor mapping every object to Ω), that is $ev_\lambda = \eta_\lambda$, then Λ is compatible with cl .*

Proof. We denote the components of the natural transformation by $ev_{\lambda, X}$, hence the evaluation map itself corresponds to $ev_{\lambda, \Omega}$.

Let $X \in \mathbf{C}$ and $S \subseteq \mathbf{C}(X, \Omega)$. We need to show that $\Lambda_X \text{cl}_X(S) \subseteq \text{cl}_{FX} \circ \Lambda_X(S)$.

So assume $ev_\lambda \circ Fk \in \Lambda_X \circ \text{cl}_X(S)$ for some $\lambda \in \Lambda$ and $k \in \text{cl}_X(S)$ (i.e., $\alpha_X(S) \preceq k^*(d_\Omega)$). Now we do the case distinction based on the type of ev_λ .

■ Let $ev_\lambda: F \Rightarrow \Omega$. Then we find

$$\begin{aligned} \alpha_{FX}(\Lambda_X(S)) &= \bigwedge_{\lambda' \in \Lambda, h \in S} (ev_{\lambda', \Omega} \circ Fh)^*(d_\Omega) && (\because ev_{\lambda'} \text{ is natural}) \\ &= \bigwedge_{\lambda' \in \Lambda, h \in S} ev_{\lambda', X}^*(d_\Omega) \\ &= \bigwedge_{\lambda' \in \Lambda} ev_{\lambda', X}^*(d_\Omega) \\ &\preceq ev_{\lambda, X}^*(d_\Omega) \end{aligned}$$

Thus, $ev_{\lambda, X} = ev_{\lambda, \Omega} \circ Fk \in \text{cl}_{FX}(\Lambda_X(S))$.

XX:24 Expressive Quantale-Valued Logics for Coalgebras: an Adjunction-Based Approach

- Let $ev_\lambda: F \Rightarrow \text{Id}$. Using $k \in \text{cl}_X(S)$ we have $\alpha_X(S) \preceq k^*(d_\Omega)$ and since reindexing functor are order preserving, so we find that

$$\begin{aligned}
 ev_{\lambda,X}^*(\alpha_X(S)) &\preceq ev_{\lambda,X}^*k^*(d_\Omega) \\
 \iff ev_{\lambda,X}^*\left(\bigwedge_{h \in S} h^*(d_\Omega)\right) &\preceq ev_{\lambda,X}^*k^*(d_\Omega) \\
 \iff \bigwedge_{h \in S} (ev_{\lambda,X}^*h^*(d_\Omega)) &\preceq ev_{\lambda,X}^*k^*(d_\Omega) \\
 \iff \bigwedge_{h \in S} (h \circ ev_{\lambda,X})^*(d_\Omega) &\preceq (k \circ ev_{\lambda,X})^*(d_\Omega) \\
 \iff \bigwedge_{h \in S} (ev_{\lambda,\Omega} \circ Fh)^*(d_\Omega) &\preceq (ev_{\lambda,\Omega} \circ Fk)^*(d_\Omega)
 \end{aligned}$$

Moreover $\alpha_{FX}\Lambda_X(S) = \bigwedge_{\lambda' \in \Lambda, h \in S} (ev_{\lambda',\Omega} \circ Fh)^*(d_\Omega) \preceq \bigwedge_{h \in S} (ev_{\lambda,\Omega} \circ Fh)^*(d_\Omega)$ and thus using the above (derived) inequality we find

$$\alpha_{FX}\Lambda_X(S) \preceq \bigwedge_{h \in S} (ev_{\lambda,\Omega} \circ Fh)^*(d_\Omega) \preceq (ev_{\lambda,\Omega} \circ Fk)^*(d_\Omega)$$

Thus, $ev_{\lambda,\Omega} \circ Fk \in \text{cl}_{FX}(\Lambda_X(S))$. ◀

► **Theorem 21.** *Assume that cl'_X is a subclosure of cl_X and compatible. For a set Θ_X of constants and a coalgebra $c: X \rightarrow FX$, the logic function $\log_X = \mathcal{P}(c^\bullet) \circ \Lambda_X \circ \text{cl}'_X \cup \Theta_X$ induces $\text{beh}_X = c^* \circ \kappa_X \wedge \alpha_X(\Theta_X)$, i.e., $\alpha_X \circ \log_X \circ \gamma_X = \text{beh}_X$.*

Proof. Define \log_1, \log_2 as in the proof of Proposition 14. Then we have:

$$\begin{aligned}
 \alpha_X \circ \log_X \circ \gamma_X &= \alpha_X \circ \log_1 \circ \gamma_X \wedge \alpha_X \circ \log_2 \circ \gamma_X \\
 &= \alpha_X \circ \mathcal{P}(c^\bullet) \circ \Lambda_X \circ \text{cl}'_X \circ \gamma_X \wedge \alpha_X \circ \Theta_X \circ \gamma_X \\
 &\quad (\because \text{cl}' \circ \text{cl} = \text{cl}, \text{Lemma 33}) \\
 &= \alpha_X \circ \mathcal{P}(c^\bullet) \circ \Lambda_X \circ \gamma_X \wedge \alpha_X(\Theta_X) \\
 &\quad (\because \alpha \text{ is natural}) \\
 &= c^* \circ \alpha_{FX} \circ \Lambda_X \circ \gamma_X \wedge \alpha_X(\Theta_X) \\
 &= \text{beh}_X.
 \end{aligned}$$
◀

D Proofs for Section 5 (Logics for Quantale-valued Simulation Distances)

► **Proposition 22.** *Infinitary meets, infinitary joins, negative scaling $d_{\mathcal{V}}(v, _)$, and positive scaling $_ \otimes v$ (for $v \in \mathcal{V}$) are non-expansive.*

Proof. For the non-expansiveness of arbitrary meets, let $v_i, w_i \in \mathcal{V}$ ($i \in I$). We need to show that $d_{\mathcal{V}}(v_i, w_i) \leq d_{\mathcal{V}}(\bigwedge_{i \in I} v_i, \bigwedge_{i \in I} w_i)$. By reflexivity we have $d_{\mathcal{V}}(v_i, w_i) \leq d_{\mathcal{V}}(v_i, w_i)$ for each $i \in I$. Then, by $v \otimes _ \dashv d_{\mathcal{V}}(v, _)$, we have $d_{\mathcal{V}}(v_i, w_i) \otimes v_i \leq w_i$ for each i . Thus, taking infima both sides and since infimum is order preserving we get

$$\bigwedge_{i \in I} (d_{\mathcal{V}}(v_i, w_i) \otimes v_i) \leq \bigwedge_{i \in I} w_i.$$

Moreover since $v \otimes _$ is order preserving we have

$$d_{\mathcal{V}}(v_i, w_i) \otimes \bigwedge_{i \in I} v_i \leq \bigwedge_{i \in I} (d_{\mathcal{V}}(v_i, w_i) \otimes v_i).$$

So by transitivity of \leq we get $d_{\mathcal{V}}(v_i, w_i) \otimes \bigwedge_{i \in I} v_i \leq \bigwedge_{i \in I} w_i$, which by $v \otimes _ \dashv d_{\mathcal{V}}(v, _)$ is equivalent to $d_{\mathcal{V}}(v_i, w_i) \leq d_{\mathcal{V}}(\bigwedge_{i \in I} v_i, \bigwedge_{i \in I} w_i)$.

For the non-expansiveness of arbitrary joins, we prove, using distributivity of \otimes over join:

$$\begin{aligned} & \forall_{i \in I} d_{\mathcal{V}}(v_i, w_i) \leq d_{\mathcal{V}}(v_i, w_i) \\ \iff & \forall_{i \in I} d_{\mathcal{V}}(v_i, w_i) \otimes v_i \leq w_i \\ \implies & \bigvee_{i \in I} (d_{\mathcal{V}}(v_i, w_i) \otimes v_i) \leq \bigvee_{i \in I} w_i \\ \iff & d_{\mathcal{V}}(v_i, w_i) \otimes \bigvee_{i \in I} v_i \leq \bigvee_{i \in I} w_i \\ \iff & d_{\mathcal{V}}(v_i, w_i) \leq d_{\mathcal{V}}(\bigvee_{i \in I} v_i, \bigvee_{i \in I} w_i). \end{aligned}$$

For non-expansiveness of negative scaling, observe that $d_{\mathcal{V}}(v_1, v_2) \leq d_{\mathcal{V}}(d_{\mathcal{V}}(v, v_1), d_{\mathcal{V}}(v, v_2))$ by the triangle inequality. From commutativity of \otimes it follows that $d_{\mathcal{V}}(v_1, v_2) \otimes d_{\mathcal{V}}(v, v_1) \leq d_{\mathcal{V}}(v, v_2)$. In the last step we use the adjunction property to obtain: $d_{\mathcal{V}}(v, v_1) \otimes d_{\mathcal{V}}(v_1, v_2) \leq d_{\mathcal{V}}(v, v_2)$.

For non-expansiveness of positive scaling, we again start from $d_{\mathcal{V}}(v_1, v_2) \leq d_{\mathcal{V}}(v_1, v_2)$ (reflexivity). Then, using the adjunction property, we obtain $d_{\mathcal{V}}(v_1, v_2) \otimes v_1 \leq v_2$. This implies that $d_{\mathcal{V}}(v_1, v_2) \otimes v_1 \otimes v \leq v_2 \otimes v$. And again using the adjunction property we can conclude that $d_{\mathcal{V}}(v_1, v_2) \leq d_{\mathcal{V}}(v_1 \otimes v, v_2 \otimes v)$. ◀

In order to show compatibility of cl'_X , we need the following result that rewrites a predicate $p \in \text{cl}_X$ in a certain normal form.

▶ **Proposition 34.** *Let $S \subseteq \text{Set}(X, \mathcal{V})$. Then every non-expansive predicate $p \in \text{cl}_X(S)$ can be rewritten as $\bigvee_{x \in X} p_x$ where $p_x(y) = (\bigwedge_{h \in S} d_{\mathcal{V}}(h(x), h(y))) \otimes p(x)$.*

Proof. Remember that we assume γ_X being natural. I.e., for any $p \in \text{cl}_X(S)$ there is a non-expansive function $op: (\mathcal{V}^S, d_{\mathcal{V}}^S) \rightarrow (\mathcal{V}, d_{\mathcal{V}})$ such that $p = op \circ \langle S \rangle$ (cf. Lemma 10). In particular, the function op is given by the (enriched) left Kan extension $\text{Lan}_{\langle S \rangle} p$ of p along $\langle S \rangle$. Since \mathcal{V} is cocomplete and has tensors (given by \otimes), thus using the coend description of left Kan extension:

$$\text{Lan}_{\langle S \rangle} p(u) = \bigvee_{x \in X} d_{\mathcal{V}}^S(\langle S \rangle x, u) \otimes p(x) = \bigvee_{x \in X} \left(\bigwedge_{h \in S} d_{\mathcal{V}}(h(x), u(h)) \right) \otimes p(x),$$

where $u: S \rightarrow \mathcal{V}$.

In particular, for $y \in X$ we have

$$p(y) = \text{Lan}_{\langle S \rangle} p(\langle S \rangle y) = \bigvee_{x \in X} \left(\bigwedge_{h \in S} d_{\mathcal{V}}(h(x), h(y)) \right) \otimes p(x) = \bigvee_{x \in X} p_x(y). \quad \blacktriangleleft$$

▶ **Proposition 23.** *If each $\lambda \in \Lambda$ is sup-preserving (i.e. $\lambda_X(\bigvee P) = \bigvee \mathcal{P}(\lambda_X)(P)$, for every subset $P \subseteq \text{Set}(X, \mathcal{V})$), then the sub-closure cl' of cl as above is compatible.*

Proof. Since cl' is a subclosure of cl , it suffices to show that $\Lambda_X(\text{cl}_X(S)) \subseteq \text{cl}_{FX}(\Lambda_X(\text{cl}'_X(S)))$ (for $S \subseteq \mathbf{Set}(X, \mathcal{V})$). Assume $\lambda_X(p) \in \Lambda_X(\text{cl}_X(S))$, for some $\lambda \in \Lambda, p \in \text{cl}_X(S)$. Using the previous proposition we find

$$\lambda_X(p) = \lambda_X\left(\bigvee_{x \in X} p_x\right) = \bigvee_{x \in X} \lambda_X(p_x)$$

Clearly, $p_x \in \text{cl}'_X(S)$ because $d_{\mathcal{V}}(h(x), h(_)) \in \text{cl}'_X(S)$ (for $h \in S$) and cl'_X is closed under arbitrary meets and positive scaling. Thus $\lambda_X(p_x) \in \Lambda_X \text{cl}'_X(S)$ (for each $x \in X$); whence $\lambda_X(p_x) \in \text{cl}_{FX} \Lambda_X \text{cl}'_X(S)$. And since full closure (in particular cl_{FX}) is closed under arbitrary joins, we conclude that $\lambda_X(p) \in \text{cl}_{FX} \Lambda_X \text{cl}'_X(S)$. ◀

E

 Proofs for Section 6 (The Adjoint Setup in an Eilenberg-Moore Category)

► **Lemma 24.** *We have that $\Lambda'_X(S) = \{ev_\lambda \circ Fo \circ FTh \mid h \in S\}$ where $S \subseteq \mathbf{C}(X, \Omega)$.*

Proof. Given S , α'_X maps every $(h: X \rightarrow \Omega) \in S$ to $o \circ Lh = o \circ Th: LX \rightarrow \Omega$. (Note that o equals $\epsilon_{(\Omega, o)}$, the component for (Ω, o) of the co-unit.) Λ_{LX} maps this arrow to $ev_\lambda \circ \tilde{F}(o \circ Lh) = ev_\lambda \circ Fo \circ FTh$. ◀

► **Theorem 25.** *We fix a coalgebra $(c: X \rightarrow FTX) \in \mathbf{C}$. Assume that Λ_{LX} is compatible with the closure cl_{LX} and fix $\Theta_{LX} \subseteq \mathbf{EM}(T)(LX, \Omega)$ to ensure that **A4** holds.*

1. *Then the logic function $\log_{LX} = \mathcal{P}((c^\#)^\bullet) \circ \Lambda_{LX} \cup \Theta_{LX}$ is cl_{LX} -compatible.*
2. *For the behaviour function $\text{beh}_{LX} = (c^\#)^* \circ \kappa_{LX} \wedge \alpha_{LX}(\Theta_{LX})$ (where $\kappa_{LX} = \alpha_{\tilde{F}LX} \circ \Lambda_{LX} \circ \gamma_{LX}$), we have $\alpha_{LX}(\mu \log_{LX}) = \nu \text{beh}_{LX}$.*
3. *Now define another logic function $\log'_X = \mathcal{P}(c^\bullet) \circ \Lambda'_X \cup \Theta'_X$ with $\Theta_{LX} = \alpha'_X(\Theta'_X)$. It holds that $\alpha'_X \circ \log'_X \circ \gamma'_X = \log_{LX}$ and we obtain $\alpha_{LX}(\alpha'_X(\mu \log'_X)) = \nu \text{beh}_X$.*

Proof. The functions $\log_{LX}, \text{beh}_{LX}$ are instantiations of the templates in Section 4, where cl'_{LX} is the identity. Hence cl_{LX} -compatibility of \log_{LX} follows directly from Proposition 14 and preservation of fixpoints from Theorem 21.

In order to prove that $\alpha'_X \circ \log'_X \circ \gamma'_X = \log_{LX}$, we first show that $\alpha'_X \circ \mathcal{P}(c^\bullet) \circ \Lambda'_X \circ \gamma'_X = \mathcal{P}((c^\#)^\bullet) \circ \Lambda_{LX}$. Since α'_X, γ'_X are inverse to each other (the adjunction is an equivalence), this is equivalent to showing $\mathcal{P}(c^\bullet) \circ \Lambda'_X = \gamma'_X \circ \mathcal{P}((c^\#)^\bullet) \circ \Lambda_{LX} \circ \alpha'_X$. Hence it is sufficient to show that $\gamma'_X \circ \mathcal{P}((c^\#)^\bullet) = \mathcal{P}(c^\bullet) \circ \mathcal{P}(R)$, since this implies

$$\gamma'_X \circ \mathcal{P}((c^\#)^\bullet) \circ \Lambda_{LX} \circ \alpha'_X = \mathcal{P}(c^\bullet) \circ \mathcal{P}(R) \circ \Lambda_{LX} \circ \alpha'_X = \mathcal{P}(c^\bullet) \circ \Lambda'_X.$$

Hence let $S \subseteq \mathbf{EM}(T)(\tilde{F}LX, (\Omega, o))$. The map $\mathcal{P}(c^\bullet) \circ \mathcal{P}(R)$ transforms all $g \in S$ (which are of type $g: FTX \rightarrow \Omega$ as arrows in \mathbf{C}) to $g \circ c: X \rightarrow \Omega$ in \mathbf{C} . Instead $\gamma'_X \circ \mathcal{P}((c^\#)^\bullet)$ transforms such g first to EM maps $g \circ c^\#: TX \rightarrow \Omega$ and then to $g \circ c^\# \circ \eta_X: X \rightarrow \Omega$ living in \mathbf{C} (note that η_X is the unit of free-forgetful adjunction $L \dashv R: \mathbf{C} \rightarrow \mathbf{EM}(T)$). Then $g \circ c = g \circ c^\# \circ \eta_X$ follows, since $c^\#$ is the transpose of c .

From this $\alpha'_X \circ \log'_X \circ \gamma'_X = \log_{LX}$ follows, since α'_X preserves unions and $\Theta_{LX} = \alpha'_X(\Theta'_X)$.

This implies $\alpha'_X \circ \log'_X = \log_{LX} \circ \alpha'_X$, since α'_X, γ'_X are inverse to each other. Hence $\alpha'_X(\mu \log'_X) = \mu \log_{LX}$ according to Theorem 4.

We can conclude by observing that $\alpha_{LX}(\alpha'_X(\mu \log'_X)) = \alpha_{LX}(\mu \log_{LX}) = \nu \text{beh}_X$. ◀

► **Proposition 27. 1.** *Let $a \in \Sigma$. Then $\eta_a: \tilde{M} \Rightarrow Id$ given by the composite $MX \xrightarrow{\pi_1} X^\Sigma \xrightarrow{\pi_a} X$ is a natural transformation.*

2. Let $f: (B, b) \rightarrow (\Omega, o)$ be a homomorphism. Then $\eta'_f: \tilde{M} \Rightarrow \mathbf{\Omega}$ given by the composite $MX \xrightarrow{\pi_2} B \xrightarrow{f} \Omega$ is a natural transformation.

Thus, $ev_a = \eta'_\Omega$, $ev_f = \eta'_\Omega$ satisfy the properties of Lemma 18, and if each evaluation map is of this form, then Λ is cl-compatible.

Proof. First remember that every monad on **Set** is strong.

Note that in both the cases it is straightforward to see that the evaluation maps are natural transformations of type $M \Rightarrow \text{Id}$ or $M \Rightarrow \mathbf{\Omega}$; thus, it suffices to show that ev_a and ev_f are algebra maps. This is because the lifted functor \tilde{M} and M are the same on arrows; thus, naturality of ev_a, ev_f follows directly from the naturality squares in **Set**.

For the first item, consider the following diagram with $(A, a) \in \mathbf{EM}(T)$:

$$\begin{array}{ccccc}
T(A^\Sigma \times B) & \xrightarrow{T\pi_1} & T(A^\Sigma) & \xrightarrow{T\pi_a} & TA \\
\langle a \mapsto T(\pi_a \circ \pi_1), b \circ T\pi_2 \rangle \downarrow & & a \mapsto T\pi_a \downarrow & \nearrow \pi'_a & \downarrow a \\
(TA)^\Sigma \times B & & (TA)^\Sigma & & \\
a^\Sigma \times B \downarrow & & a^\Sigma \downarrow & & \\
A^\Sigma \times B & \xrightarrow{\pi_1} & A^\Sigma & \xrightarrow{\pi_a} & A
\end{array}$$

where π_a, π'_a are the obvious evaluation maps. Note that the map $a \mapsto T\pi_a$ is the unique map that makes the triangle commute. Moreover a^Σ is the unique map that makes the ‘lower’ square commute. Lastly, the rectangle on the left commutes since $(a \mapsto T\pi_a) \circ T\pi_1 = a \mapsto T(\pi_a \circ \pi_1)$.

For the second item, consider the following diagram with $(A, a) \in \mathbf{EM}(T)$ and $(B, b) \xrightarrow{f} (\Omega, o) \in \mathbf{EM}(T)$. The square on the right commutes since f is an homomorphism, while the rectangle on the left obviously commutes.

$$\begin{array}{ccccc}
T(A^\Sigma \times B) & \xrightarrow{T\pi_2} & TB & \xrightarrow{Tf} & T\Omega \\
\langle a \mapsto T(\pi_a \circ \pi_1), b \circ T\pi_2 \rangle \downarrow & & \downarrow b & & \downarrow o \\
(TA)^\Sigma \times B & & & & \\
a^\Sigma \times \text{id}_B \downarrow & & \downarrow & & \\
A^\Sigma \times B & \xrightarrow{\pi_2} & B & \xrightarrow{f} & \Omega
\end{array}$$

► **Theorem 28.** Assume that d is preserved by the co-closure, i.e. $d = \alpha_{LX}(\gamma_{LX}(d))$, the co-closure $\alpha_X^{\mathbf{C}} \circ \gamma_X^{\mathbf{C}}$ is the identity, and each evaluation map ev_λ arises from some natural transformation either of type $F \Rightarrow \text{Id}$ or $F \Rightarrow \mathbf{\Omega}$. Then the two liftings $\kappa_{LX}, \kappa_{TX}^{\mathbf{C}}$ coincide on d , i.e., $\kappa_{LX}(d) = \kappa_{TX}^{\mathbf{C}}(d)$.

Proof. Note that each evaluation map ev_λ , $\lambda \in \Lambda$ arises from a natural transformation either of type $F \Rightarrow \text{Id}$ or $F \Rightarrow \mathbf{\Omega}$. Let $K \subseteq \Lambda$ denote those predicate liftings that are of the second

type. Now consider the following derivation:

$$\begin{aligned}
 \kappa_{LX}(d) &= \alpha_{\tilde{F}LX}(\Lambda_{LX}(\gamma_{LX}(d))) \\
 &= \alpha_{\tilde{F}LX}\left(\bigcup_{\lambda \in \Lambda} \mathcal{P}(\lambda_{LX})(\gamma_{LX}(d))\right) \\
 &= \alpha_{\tilde{F}LX}(\{ev_{\lambda, \Omega} \circ \tilde{F}h \mid h \in \gamma_{LX}(d) \wedge \lambda \in \Lambda\}) && \because \text{naturality of } ev_{\lambda} \\
 &= \alpha_{\tilde{F}LX}\left(\{h \circ ev_{\lambda, TX} \mid h \in \gamma_{LX}(d) \wedge \lambda \notin K\} \right. \\
 &\quad \left. \cup \{ev_{\lambda, TX} \mid \lambda \in K\}\right) && \text{definition of } _ \bullet \\
 &= \alpha_{\tilde{F}LX}\left(\{ev_{\lambda, TX}^\bullet(h) \mid h \in \gamma_{LX}(d) \wedge \lambda \notin K\} \right. \\
 &\quad \left. \cup \{ev_{\lambda, TX} \mid \lambda \in K\}\right) \\
 &= \alpha_{\tilde{F}LX}\left(\bigcup_{\lambda \notin K} \mathcal{P}(ev_{\lambda, TX}^\bullet)(\gamma_{LX}(d)) \cup \bigcup_{\lambda \in K} \{ev_{\lambda, TX}\}\right) && \because \alpha \text{ is left adjoint} \\
 &= \bigwedge_{\lambda \notin K} \alpha_{\tilde{F}LX}(\mathcal{P}(ev_{\lambda, TX}^\bullet)(\gamma_{LX}(d))) \wedge \bigwedge_{\lambda \in K} ev_{\lambda, TX}^*(d_\Omega) && \because \text{naturality of } \alpha \\
 &= \bigwedge_{\lambda \notin K} ev_{\lambda, TX}^*(\alpha_{LX}(\gamma_{LX}(d))) \wedge \bigwedge_{\lambda \in K} ev_{\lambda, TX}^*(d_\Omega) && \because \alpha_{LX}(\gamma_{LX}(d)) = d \\
 &= \bigwedge_{\lambda \notin K} ev_{\lambda, TX}^*(d) \wedge \bigwedge_{\lambda \in K} ev_{\lambda, TX}^*(d_\Omega) && \because \alpha_{TX}^C \circ \gamma_{TX}^C = \text{id} \\
 &= \bigwedge_{\lambda \notin K} ev_{\lambda, TX}^*(\alpha_{TX}^C(\gamma_{TX}^C(d))) \wedge \alpha_{FTX}^C(\{ev_{\lambda, TX} \mid \lambda \in K\}) && \because \text{naturality of } \alpha^C \\
 &= \bigwedge_{\lambda \notin K} \alpha_{FTX}^C(\mathcal{P}(ev_{\lambda, TX}^\bullet)(\gamma_X^C(d))) \wedge \alpha_{FTX}^C(\{ev_{\lambda, TX} \mid \lambda \in K\}) \\
 &= \alpha_{FTX}^C\left(\bigcup_{\lambda \notin K} \mathcal{P}(ev_{\lambda, TX}^\bullet)(\gamma_X^C(d)) \cup \{ev_{\lambda, TX} \mid \lambda \in K\}\right) \\
 &= \alpha_{FTX}^C \circ \Lambda_{TX} \circ \gamma_X^C(d) \\
 &= \kappa_{TX}^C(d). \quad \blacktriangleleft
 \end{aligned}$$

► **Lemma 35.** Assume that $\mathbf{C} = \mathbf{Set}$ and $\Psi X = \mathbf{DPMet}_\mathcal{V}(X)$ (resp. $\Psi = \mathbf{PMet}_\mathcal{V}(X)$) for an integral quantale Ω . Let let $d_\Omega = [_, _]$ (resp. the symmetrized variant of $[_, _]$). Then $\alpha_X^C \circ \gamma_X^C$ is the identity.

Proof. In order to show that $\alpha_X^C \circ \gamma_X^C = \text{id}_{\Psi X}$, we first observe that $\alpha_X^C \circ \gamma_X^C \preceq \text{id}_{\Psi(TX)}$ follows from the properties of the adjunction (Galois connection). In order to show equality, i.e., $\alpha_X^C(\gamma_X^C(d)) = d$ for $d \in \Psi(TX)$, we prove that – given $t_1, t_2 \in TX$ – there exists $h \in \gamma_X^C(d)$ such that $d(t_1, t_2) = h^*d_\Omega(t_1, t_2)$. According to Lemma 31 $h: TX \rightarrow \Omega$, defined as $h(t) = d(t_1, t)$, is non-expansive wrt. d and hence contained in $\gamma_X^C(d)$. Furthermore $h^*d_\Omega(t_1, t_2) = d_\Omega(h(t_1), h(t_2)) = d(t_1, t_2)$, using Lemma 31. ◀

► **Theorem 29.** Assume that $\mathbf{C} = \mathbf{Set}$, $\Psi X = \mathbf{DPMet}_\mathcal{V}(X)$ (resp. $\Psi X = \mathbf{PMet}_\mathcal{V}(X)$) for an integral quantale \mathcal{V} and let $d_\mathcal{V} = [_, _]$ (resp. the symmetrized variant of $[_, _]$). Let $d: LX \times LX \rightarrow \mathcal{V}$ be a pseudometric that is preserved by the co-closure $\alpha_{LX} \circ \gamma_{LX}$. Assume that M is the machine functor and the family of evaluation maps is

$$\{ev_a \mid a \in \Sigma\} \cup \{ev_f \mid f \in \mathcal{F} \subseteq \mathbf{EM}(T)((B, b), (\mathcal{V}, o))\}.$$

Then the corresponding behaviour function $\text{beh}_{LX} : \Psi(LX) \rightarrow \Psi(LX)$ is defined as follows: let $t_1, t_2 \in LX$ with $c^\#(t_i) = (b_i, g_i) \in B \times LX^\Sigma$:

$$\text{beh}_{LX}(d)(t_1, t_2) = \bigwedge_{a \in \Sigma} d(g_1(a), g_2(a)) \wedge \bigwedge_{f \in \mathcal{F}} d_{\mathcal{V}}(f(b_1), f(b_2)) \wedge \bigwedge_{\theta \in \Theta_{LX}} d_{\mathcal{V}}(\theta(t_1), \theta(t_2)),$$

Proof. According to our previous results (Theorem 21) the behaviour function has the form $\text{beh}_{LX} = (c^\#)^* \circ \kappa_{LX} \wedge \alpha_{LX}(\Theta_{LX})$.

With the definition of α in Theorem 7 is straightforward to see that

$$\alpha_{LX}(\Theta_{LX})(t_1, t_2) = \bigwedge_{\theta \in \Theta_{LX}} \theta^*(d_{\mathcal{V}})(t_1, t_2) = \bigwedge_{\theta \in \Theta_{LX}} d_{\mathcal{V}}(\theta(t_1), \theta(t_2)),$$

matching the third part of the definition of beh_{LX} given in the statement of the proposition.

For the first and second part we have to unravel κ_{LX} : according to Theorem 28 we have

$$\kappa_{LX} = \kappa_{TX}^{\mathbf{C}} = \alpha_{FTX}^{\mathbf{C}} \circ \Lambda_{TX} \circ \gamma_{TX}^{\mathbf{C}}.$$

Hence, given $s_1, s_2 \in TX$ where $s_i = (g_i, b_i)$, $g : \Sigma \rightarrow TX$, $b_i \in B$:

$$\begin{aligned} \kappa_{LX}(d)(s_1, s_2) &= \alpha_{FTX}^{\mathbf{C}}(\Lambda_{TX}(\gamma_{TX}^{\mathbf{C}}(d)))(s_1, s_2) \\ &= \bigwedge \{k^*(d_{\mathcal{V}})(s_1, s_2) \mid k \in \Lambda_{TX}(\gamma_{TX}^{\mathbf{C}}(d))\} \\ &= \bigwedge \{d_{\mathcal{V}}(k(s_1), k(s_2)) \mid k = \text{ev}_\lambda \circ Fh, h \in \gamma_{TX}^{\mathbf{C}}(d)\} \\ &= \bigwedge \{d_{\mathcal{V}}(\text{ev}_\lambda \circ Fh(s_1), \text{ev}_\lambda \circ Fh(s_2)) \mid h : TX \rightarrow \mathcal{V}, d \preceq h^*d_{\mathcal{V}}\} \\ &= \bigwedge_{a \in \Sigma} \{d_{\mathcal{V}}(\text{ev}_a \circ Fh(s_1), \text{ev}_a \circ Fh(s_2)) \mid h : TX \rightarrow \mathcal{V}, d \preceq h^*d_{\mathcal{V}}\} \wedge \\ &\quad \bigwedge_{f \in \mathcal{F}} \{d_{\mathcal{V}}(\text{ev}_f \circ Fh(s_1), \text{ev}_f \circ Fh(s_2)) \mid h : TX \rightarrow \mathcal{V}, d \preceq h^*d_{\mathcal{V}}\} \end{aligned}$$

The property $d \preceq h^*d_{\mathcal{V}}$ states that h is non-expansive (wrt. $d, d_{\mathcal{V}}$). We consider the two parts of the meet separately:

- First part of meet: assume that $\lambda = a \in \Sigma$. Then

$$\text{ev}_a \circ Fh(s_i) = \text{ev}_a \circ Fh(g_i, b_i) = h(g_i(a))$$

By non-expansiveness of h it is clear that always $d_{\mathcal{V}}(h(g_1(a)), h(g_2(a))) \geq d(g_1(a), g_2(a))$. In fact we can choose h such that equality holds: we set $h(x) = d(g_1(a), _)$ and infer from Lemma 31 that it is non-expansive, i.e., that $d \preceq h^*d_{\mathcal{V}}$. Observe that $h(g_1(a)) = d(g_1(a), g_1(a)) = 1$ (remember that d is reflexive and the quantale is integral). In addition $h(g_2(a)) = d(g_1(a), g_2(a))$. Hence we obtain

$$d_{\mathcal{V}}(h(g_1(a)), h(g_2(a))) = d(g_1(a), g_2(a)),$$

again with Lemma 31.

Hence the infimum is reached in $d(g_1(a), g_2(a))$ and the first part of the meet simplifies to

$$\bigwedge_{a \in \Sigma} d(g_1(a), g_2(a)).$$

XX:30 Expressive Quantale-Valued Logics for Coalgebras: an Adjunction-Based Approach

- Second part of meet: assume that $\lambda = f \in \mathcal{F}$. Then

$$ev_f \circ Fh(s_i) = ev_f \circ Fh(g_i, b_i) = f(b_i).$$

This means that the infimum is independent of the choice of h and simplifies to

$$\bigwedge_{f \in \mathcal{F}} d_{\mathcal{V}}(f(b_1), f(b_2)).$$

Now set $s_i = c^\#(t_i)$ to include the reindexing via $c^\#$ and one obtains the characterization of beh_{LX} . ◀