

On the generalized Friedrichs-Lee model with multiple discrete and continuous states

Zhiguang Xiao*

College of Physics, Sichuan University, Chengdu 610065, P. R. China

Zhi-Yong Zhou†

School of Physics, Southeast University, Nanjing 211189, P. R. China

Abstract

In this study, we present several improvements of the non-relativistic Friedrichs-Lee model with multiple discrete and continuous states and still retain its solvability. Our findings establish a solid theoretical basis for the exploration of resonance phenomena in scenarios involving the presence of multiple interfering states across various channels. The scattering amplitudes associated with the continuous states naturally adhere to coupled-channel unitarity, rendering this framework particularly valuable for investigating hadronic resonant states appearing in multiple coupled channels. Moreover, this generalized framework exhibits a wide-range applicability, enabling investigations into resonance phenomena across diverse physical domains, including hadron physics, nuclear physics, optics, and cold atom physics, among others.

arXiv:2310.14937v1 [hep-ph] 23 Oct 2023

* xiaozg@scu.edu.cn

† zhouzhy@seu.edu.cn, corresponding author

I. INTRODUCTION

Unstable states constitute a ubiquitous phenomenon in contemporary physics, manifesting across various disciplines such as molecular physics, nuclear physics and particle physics. In the realm of hadronic physics, the prevalence of unstable resonances is particularly notable within the context of strong interactions, where new resonant states are frequently encountered and documented. These resonances assume significant significance in unraveling the fundamental characteristics of hadrons and their interactions, perpetuating their investigation as a vibrant research area within the field of particle physics.

To explore the characteristics of unstable states across diverse branches of physics, several models sharing a similar conceptual framework have independently emerged. Among these models, the Friedrichs model stands as a simple non-relativistic Hamiltonian that couples a bare discrete state to a bare continuous state [1]. Within this model, the solutions for unstable generalized eigenstates can be rigorously obtained and expressed in terms of the bare states. In the realm of quantum field theory, the Lee model was developed to investigate the properties of field renormalization [2]. This model considers two nucleon states, denoted as N and V , which can be converted to each other by absorbing or emitting a bosonic θ particle through the processes $V \rightleftharpoons N + \theta$. Analogous models can be found in various domains, such as the Jaynes-Cummings model in quantum optics [3] and the Anderson model in condensed matter physics [4]. In this article, we collectively refer to these models as the Friedrichs-Lee (FL) model, highlighting their shared conceptual foundation. The generalized eigenstates of the full interacting Hamiltonian within the FL model can be explicitly determined in terms of the original discrete state and the continuum states.

The original Friedrichs-Lee model, which involves only one discrete and one continuous state, is often considered as a toy model due to its simplicity. It is usually employed to comprehend the properties of bound states, virtual states and resonant states that appear in the scattering processes. When the bare discrete state is above the continuum threshold, its pole position moves to the second sheet and become a pair of resonance poles. If the bare discrete state is below the threshold, there would be an accompanied virtual state pole on the second sheet when the interaction is turned on. Besides these states generated from the bare discrete states, there could also be dynamically generated states from the singularities of the interaction vertices [5]. The mathematical background of describing the unstable states is the Rigged Hilbert Space (RHS) quantum mechanics [6–8], rather than the conventional Hilbert space. In the RHS quantum mechanics, the Hamiltonian H , as an Hermitian operator, could have generalized complex eigenvalues and the related eigenstates corresponding to the pole of the S -matrix that lies on the unphysical sheet of the analytically continued energy plane, commonly referred to as the Gamow states. The Friedrichs model was also extended to include more continuous or discrete states and with a more realistic interaction vertex function. As a result, it finds extensive application in a wide range of realistic scenarios, particularly in the study of hadronic scattering processes [9–14]. Furthermore, coupled-channel models sharing similar spirits have demonstrated success in describing a variety of resonance phenomena in different physical systems [15–25]. The widespread applicability and efficacy of these models in describing resonance phenomena render them as powerful tools in studying the properties of unstable states in different physical contexts.

In the hadron physics, the usual effective field theory calculation of the scattering amplitude encounter challenges pertaining to unitarity and analyticity. The perturbative S matrix generally fails to generate bound states or resonance poles on the analytically continued Riemann surface of the energy plane. To address this, various unitarization methods are used, such as the K -matrix method. The typical K -matrix parameterization of the S -matrix like $S = \frac{1-iK}{1+iK}$ lacks a dynamical origin and enforces unitarity by manual intervention. However, this parametrization does not guarantee the absence of unphysical spurious poles, including those situated in the the complex energy plane of the first Riemann sheet, which violates causality. In contrast, the Friedrichs-Lee model achieves unitarity as a consequence of its dynamics, and the Hermitian property of the Hamiltonian ensures the absence of spurious poles in the first Riemann sheet. These are the immediate advantages of these kinds of models over the K -matrix parameterization.

While there have been notable achievements in the application of such models, there remain areas that warrant further improvement. A prominent challenge pertains to the incorporation of contact interactions, such as the four-point vertex, which is not generally solvable. In refs. [10, 26], a particular form of separable interaction involving the continuum states is introduced, where the interaction vertex function between the discrete states and the continuum

also appears as the factors of the separable interaction between two continuum states. The Hamiltonian is

$$\begin{aligned}
H = & \sum_{i=1}^D M_i |i\rangle \langle i| + \sum_{i=1}^N \int_{a_i}^{\infty} d\omega \omega |\omega; i\rangle \langle \omega; i| \\
& + \sum_{i,j=1}^C v_{ij} \left(\int_{a_i}^{\infty} d\omega f_i(\omega) |\omega; i\rangle \right) \left(\int_{a_j}^{\infty} d\omega f_j^*(\omega) \langle \omega; j| \right) \\
& + \sum_{j=1}^D \sum_{i=1}^C \left[u_{ji}^* |j\rangle \left(\int_{a_i}^{\infty} d\omega f_i^*(\omega) \langle \omega; i| \right) + u_{ji} \left(\int_{a_i}^{\infty} d\omega f_i(\omega) |\omega; i\rangle \right) \langle j| \right], \tag{1}
\end{aligned}$$

where the form factor $f_i(\omega)$ is associated with the continuum state $|\omega; i\rangle$. There are two aspects which could be improved for this model. First, the interaction between the discrete states $|j\rangle$ and the continuum could be extended to a general function $f_{ij}(\omega)$, for a more realistic description of the strong interaction in the real world. Using the quark pair creation (QPC) model as an example, the interaction between a meson and their decay products is expressed as a complicated integration between the wave function for the three states and the pair production vertex [12, 27]. Secondly, the interaction between the continuum states does not need to be factorized using the same factor as the interaction between the discrete state and the continuum. In this paper, we will demonstrate that after factorizing the continuum-continuum interaction independent of the interaction vertex between the discrete-continuum interaction, the model remains exactly solvable. In principle, the extra continuum-continuum interaction should be the residue interaction after subtracting the s -channel intermediate discrete state contribution, which could have no relation with the discrete-continuum interaction. Whether this interaction can be expressed as a separable potential remains an open question. Nevertheless, as an exactly solvable model, the solution is always valuable. Moreover, in general, a generic interaction potential between continuum states could be expanded using a series of general separable basis. In fact, one can also expand both the discrete-continuum interaction vertex and continuum-continuum interaction vertices using the same function basis. Thus, the study of such separable potentials may have broader physical application. In this paper, our focus lies on these two kinds of improvements: the inclusion of multiple bare discrete and continuum states in the FL model and incorporation of a more general discrete-continuum interaction, along with various separable continuum-continuum interactions. By rigorously solving the eigenstates for the Hamiltonian, we obtain the “in” and “out” states, the scattering S-matrix, discrete state solution, and other mathematical physics properties. Our aim is to establish a solid foundation for the further phenomenological applications of the FL model by including these additional physical features.

We organize the paper as follows: In Section II, the solution of the FL model with more general interactions between discrete states and continuum states is derived. Section III discusses the case with extra separable continuum-continuum interactions. Section IV is devoted to studying the case when the interaction potential between continuum states could be approximated by a sum of separable potentials and consider the cases when both the continuum-continuum potential and continuum-discrete potentials are approximated by a truncated series. In section V, as an application, we consider some simple examples and discuss the behavior of the discrete states after turning on various interaction. Section VI is the conclusion.

II. THE EXTENDED FRIEDRICHS-LEE MODEL WITH MULTIPLE DISCRETE STATES AND CONTINUUM STATES

First, we are going to consider a system with D kinds of discrete states and C kinds of continuum states, where C and D denotes the numbers of the continuum states and the discrete states respectively. If there is no interaction, the mass of the j -th discrete state $|j\rangle$ is M_j , while the energy spectrum of the n -th continuum state ranges in $[a_n, \infty)$ with the threshold energy a_n . The interaction between the j -th discrete state and the n -th continuum state can be generally represented by a coupling function $f_{jn}(\omega)$. The full Hamiltonian can be expressed as

$$H = H_0 + H_I, \tag{2}$$

where the free Hamiltonian H_0 could be written down explicitly as

$$H_0 = \sum_{i=1}^D M_i |i\rangle \langle i| + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \omega |\omega; n\rangle \langle \omega; n|, \tag{3}$$

and the interaction part H_I reads

$$H_I = \sum_{j=1}^D \sum_{n=1}^C \left[|j\rangle \left(\int_{a_n}^{\infty} d\omega f_{jn}^*(\omega) \langle \omega; n| \right) + \left(\int_{a_n}^{\infty} d\omega f_{jn}(\omega) |\omega; n\rangle \right) \langle j| \right]. \quad (4)$$

The free eigenstates are supposed to be orthogonal to each other and the normalization conditions satisfy $\langle i|j\rangle = \delta_{ij}$, $\langle i|\omega; n\rangle = 0$ and $\langle \omega; n|\omega'; n'\rangle = \delta(\omega - \omega')\delta_{nn'}$. For simplicity, we first suppose that there is no degenerate threshold and no degenerate discrete states. In fact, if there are degenerate states with the same threshold and the same interactions with the other states, the corresponding solutions will also be degenerate with the same expression after the interactions are turned on, and we will take them as one state with degenerate degrees of freedom just like different magnetic quantum numbers when there is no magnetic field. If the states with degenerate threshold take part in different interactions, the following discussion will not be modified too much. We will come back to this case later.

The general solution for the energy eigenvalue problem $H|\Psi(E)\rangle = E|\Psi(E)\rangle$ can be represented as a linear combination of the discrete states and the continuum states,

$$|\Psi(E)\rangle = \sum_{i=1}^D \alpha_i(E) |i\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \psi_n(E, \omega) |\omega; n\rangle, \quad (5)$$

where the $\alpha_i(E)$ and $\psi_n(E, \omega)$ functions are defined as the coefficient functions of the discrete states and the continuum states respectively. By substituting this ansatz into the eigenvalue equation, and carefully examining the coefficients preceding the discrete states and the continuum states, we can derive two distinct sets of equations,

$$(M_j - E)\alpha_j(E) + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega f_{jn}^*(\omega) \psi_n(E, \omega) = 0, \quad \text{for } j = 1, \dots, D \quad (6)$$

$$\sum_{j=1}^D \alpha_j(E) f_{jn}(\omega) + (\omega - E)\psi_n(E, \omega) = 0, \quad \text{for } n = 1, \dots, C, \text{ and } \omega > a_n. \quad (7)$$

An important observation to make is that the formula exhibits a nontrivial complexity, which does not appear in the single-channel scenario. Specifically, for a given energy range $a_l < \omega < a_{l+1}$, there are only l equations present in Eqs. (7).

Consequently, the eigenvalue problem yields both continuum solutions and discrete solutions. These solutions correspond to different regimes of the spectrum, which will be addressed carefully in the following.

1. The continuum state solutions

When the energy E is above the highest threshold, that means, $E > a_C$, there will be C continuum states when the interactions are turned on, so the m -th continuum solution will be

$$|\Psi_m(E)\rangle = \sum_{i=1}^D \alpha_{mi}(E) |i\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \psi_{mn}(E, \omega) |\omega; n\rangle, \quad m = 1, 2, \dots, C \quad (8)$$

However, when the energy E is lower than the highest threshold, e.g., $E \in [a_l, a_{l+1})$, $l < C$, there will be l degenerate continuum eigenstates, $m = 1, 2, \dots, l$, and the other states are not well-defined below their thresholds and are set to $\mathbf{0}$. To remove the ambiguity of the degenerate states, it is required that when the interaction is turned off, i.e. $f_{jm}(\omega) \rightarrow 0$, $|\Psi_m\rangle$ tends to the free continuum state $|E; m\rangle$. We expect that, when the eigenvalue $E \in [a_1, a_2]$, we can solve $\alpha_{1,i}$ and $\psi_{1,i}$ in $|\Psi_1(E)\rangle$, and then analytically extend these parameters to $E \in [a_2, a_3]$ to solve $|\Psi_2(E)\rangle$, and so on. In this way the eigenfunctions can be uniquely determined. From Eq. (6,7) in terms of the coefficients in Eq. (8), the coefficient function $\psi_{mn}(E, \omega)$ before the continuum state in different energy regions could be expressed as

$$\text{(for } n \leq l) \quad \psi_{mn}(E, \omega) = \gamma_n \delta_{mn} \delta(\omega - E) + \frac{1}{E - \omega \pm i0} \sum_{j=1}^D \alpha_{mj}(E) f_{jn}(\omega),$$

$$\text{(for } n > l) \quad \psi_{mn}(E, \omega) = \frac{1}{E - \omega \pm i0} \sum_{j=1}^D \alpha_{mj}(E) f_{jn}(\omega).$$

This equation could be concisely written down in one equation by using the Heaviside step function $\Theta(x)$,

$$\psi_{mn}^{\pm}(E, \omega) = \gamma_n \delta_{mn} \delta(\omega - E) \Theta(E - a_n) + \frac{1}{E - \omega \pm i0} \sum_{j=1}^D f_{jn}(\omega) \alpha_{mj}(E). \quad (9)$$

Notice that ψ_{mn}^{\pm} is a generalized function, and in order to distinguish between different integral contours, we have included $\pm i0$ in the denominator of the integral in Eq.(8). The ψ^+ state corresponds to the coefficient for the in-state while ψ^- corresponds to those of the out-state. For the convenience of the future discussions, we will omit the superscripts \pm in the notations. It should be understood that the appropriate superscript can be easily inferred based on the context. In the cases where there is a need to explicitly indicate the in-state or out-state, we will make use of the superscript accordingly.

Inserting this equation back into Eq.(6), we can obtain the equations for the coefficient functions $\alpha_{mk}(E)$ for $m = 1, 2, \dots, l$

$$-\sum_{k=1}^D \alpha_{mk}(E) \left[\delta_{kj}(E - M_j) - \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \frac{f_{kn}(\omega) f_{jn}^*(\omega)}{E - \omega \pm i0} \right] + \sum_{n=1}^C \gamma_m(E) \delta_{mn} f_{jn}^*(E) = 0. \quad (10)$$

With many different discrete states and continuum ones involved in, the representation becomes much more complex than the simplest version. In fact, the formula and the derivation procedure could be simplified by introducing the matrix form. In the following, the matrices are represented in bold faces and the dot symbol “ \cdot ” represents the matrix product, and the matrix element is expressed in the form like $(\boldsymbol{\eta})_{ij}$. For example, Eq. (10) could be written down in the matrix form as

$$-\boldsymbol{\alpha}^{\pm}(E) \cdot \boldsymbol{\eta}_{\pm}(E) + \boldsymbol{\gamma}(E) \cdot \mathbf{f}^{\dagger}(E) = 0,$$

where $\boldsymbol{\alpha}$ and \mathbf{f} are the $C \times D$ and $D \times C$ matrices for the coefficients α_{mk} and f_{jm} , respectively. The matrix $\boldsymbol{\gamma}$ is defined as a diagonal matrix of dimension $C \times C$

$$(\boldsymbol{\gamma})_{mn}(E) = \gamma_n \delta_{mn} \Theta(E - a_n),$$

whose diagonal elements γ_n could be different in principle for different n and the values could be determined by the normalization conditions. The $\boldsymbol{\eta}_{\pm}$ matrix, the inverse of resolvent function matrix, is of dimension $D \times D$ and every matrix element reads

$$(\boldsymbol{\eta}_{\pm})_{kj}(E) = (E - M_j) \delta_{kj} - \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \frac{f_{kn}(\omega) f_{jn}^*(\omega)}{E - \omega \pm i0}. \quad (11)$$

In general, the determinant of $\boldsymbol{\eta}$ matrix does not vanish for $a_l < E < a_{l+1}$, the matrix $\boldsymbol{\alpha}^{\pm}$ can be represented as

$$\boldsymbol{\alpha}^{\pm}(E) = \boldsymbol{\gamma}(E) \cdot \mathbf{f}^{\dagger}(E) \cdot \boldsymbol{\eta}_{\pm}^{-1}(E).$$

Inserting this result into eq. (9), the coefficient functions ψ_{mn} before the continuum states can be obtained in matrix representation

$$\psi^{\pm}(E, \omega) = \gamma \delta(\omega - E) + \frac{1}{E - \omega \pm i0} \boldsymbol{\gamma}(E) \cdot \mathbf{f}^{\dagger}(E) \cdot \boldsymbol{\eta}_{\pm}^{-1}(E) \cdot \mathbf{f}(\omega).$$

The solution of the continuum eigenstate can then be expressed as

$$\begin{aligned} |\Psi_m^{\pm}(E)\rangle &= \sum_{i=1}^D \alpha_{mi}^{\pm}(E) |i\rangle + \sum_{n=1}^C \int d\omega \psi_{mn}^{\pm}(E, \omega) |\omega; n\rangle \\ &= \gamma_m \Theta(E - a_m) |E, m\rangle + \sum_{k=1}^D (\boldsymbol{\gamma}(E) \cdot \mathbf{f}^{\dagger}(E) \cdot \boldsymbol{\eta}_{\pm}^{-1}(E))_{mk} \left(|k\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \frac{f_{kn}(\omega)}{E - \omega \pm i0} |\omega; n\rangle \right) \end{aligned} \quad (12)$$

Notice that in the energy region $a_l < E < a_{l+1}$, the wave function $|\Psi_m^{\pm}\rangle$ for $m > l$ should vanish. Another required condition is that, when the coupling function f_{jn} vanishes, $|\Psi_m^{\pm}(E)\rangle$ tends to $|E, m\rangle$. Therefore, the

coefficient γ_m is determined to be 1. It can be checked that the normalization satisfies $\langle \Psi_m^\pm(E) | \Psi_n^\pm(E') \rangle = \delta(E - E') \delta_{mn}$. Actually, in the point view of the scattering theory, $|\Psi^+\rangle$ is the “in” state and $|\Psi^-\rangle$ is the “out” state, so the S matrix can be obtained by inner product of the “in” state and the “out” state as

$$\begin{aligned} \langle \Psi_m^-(E) | \Psi_n^+(E') \rangle &= \gamma_m^* \gamma_n \delta(E - E') - 2\pi i \delta(E - E') (\boldsymbol{\gamma}(E') \cdot \mathbf{f}^\dagger(E') \cdot \boldsymbol{\eta}_+^{-1\dagger}(E) \cdot \mathbf{f}(E) \cdot \boldsymbol{\gamma}^\dagger(E))_{nm} \\ &= \delta(E - E') [\boldsymbol{\gamma} \cdot (1 - 2\pi i \mathbf{f}^T(E) \cdot \boldsymbol{\eta}_+^{-1}(E) \cdot \mathbf{f}^*(E)) \cdot \boldsymbol{\gamma}]_{nm}. \end{aligned} \quad (13)$$

The $\boldsymbol{\eta}_\pm(E)$ function can be analytically extended to the complex E plane with $\eta_+(E)$ and $\eta_-(E)$ coinciding with $\boldsymbol{\eta}(E)$ on the upper edge and lower edge of the real axis above the thresholds, respectively. We can also define the analytically continued S matrix

$$\mathbf{S} = 1 - 2\pi i \mathbf{f}^T(E) \cdot \boldsymbol{\eta}^{-1}(E) \cdot \mathbf{f}^*(E), \quad (14)$$

where E is analytically continued to the complex energy plane and only when E is real and on the upper edge of the cut above the lowest threshold a_1 is the S matrix the physical one. Given the presence of C continuous states with distinct thresholds, it is a general result that there exist 2^C different Riemann sheets for the analytically continued S matrix. Since only the Riemann sheets nearest to the physical region affect the physical S -matrix the most, we label the m -th sheet as the Riemann sheet continued from the physical region (a_m, a_{m+1}) , where the first sheet where the physical S matrix resides is called physical sheet by convention.

It is worth pointing out that the formula of scattering matrix Eq. (14) has important phenomenological applications. For example, in studying the particle-particle scattering processes, the two particles that collides or those final states (usually called as channels in the scattering experiments) form continuum states, while the intermediate resonance states are regarded as the discrete states. The (n, m) -th element of scattering matrix in Eq. (14) could describe the scattering amplitudes from the channel of n -th continuum state to the m -th channel. The coupled-channel unitarity is naturally satisfied among all the related scattering amplitudes due to the obvious relation $\mathbf{S}\mathbf{S}^\dagger = I$. Furthermore, once the coupling function between the discrete and continuum state is reliably described by some dynamical models, the physical observables, such as the cross sections, could be predicted or calculated [28].

2. The discrete state solutions:

In Eqs. (6, 7), if the eigenvalue $E \notin [a_n, \infty)$ for $n = 1, \dots, C$, there is no need to introduce the $\pm i0$ in the denominator of the integrand and we have

$$\psi_n(E, \omega) = \frac{1}{E - \omega} \sum_j f_{jn}(\omega) \alpha_j(E), \quad (\text{for } n = 1, \dots, C) \quad (15)$$

$$(\boldsymbol{\alpha}(E) \cdot \boldsymbol{\eta}(E))_j = \sum_{k=1}^D \alpha_k(E) \left[\delta_{kj} (M_j - E) - \sum_{m=1}^C \int_{a_m}^{\infty} d\omega \frac{f_{km}(\omega) f_{jm}^*(\omega)}{\omega - E} \right] = 0. \quad (\text{for } j = 1, \dots, D) \quad (16)$$

In order to obtain nonzero solutions of $\alpha_k(E)$, it is necessary to satisfy the condition $\det \boldsymbol{\eta}(E) = 0$. This condition implies that there may exist discrete energy solutions for this equation, which in general correspond to the poles of the S -matrix elements. If there exist solutions on the first sheet, they must reside on the real axis below the lowest threshold. Additionally, solutions can also be found on the unphysical sheets, which may corresponds to complex conjugate resonance poles on the complex energy plane or to virtual state poles located on the real axis below the lowest threshold. There would be at least D discrete solutions which tend to the bare discrete states, i.e. $\alpha_k^{(l)} \rightarrow \delta_{kl}$ and $E \rightarrow M_l$ for $l = 1, 2, \dots, D$, as all the coupling function $f_{ln} \rightarrow 0$. Furthermore, it is possible for other dynamically generated states that do not go to the bare states when the interactions are switched off. In general, the solutions does not exhibit degeneracy, indicating that the poles for S matrix are just simple poles. If the degenerate solutions occur for $\det \boldsymbol{\eta}(E) = 0$, it implies that two or more poles may coincide and form a higher order pole. This situation is considered to be accidental and only occurs for some special coupling functions. For the purposes of our discussion, we will not consider this special case and assume that the solutions are non-degenerate. Then for each energy solution E_i , we can also find the eigenvector $\alpha_k^{(i)}(E_i)$ and $\psi_n^{(i)}(E_i, \omega)$, and the wave function of discrete state is expressed as

$$|\Psi^{(i)}(E_i)\rangle = \sum_{j=1}^D \alpha_j^{(i)}(E_i) (|j\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \frac{f_{jn}(\omega)}{E_i - \omega} |\omega; n\rangle). \quad (17)$$




FIG. 1. The integral contour for the resonance solution.

When E_i lies on the real axis of the first Riemann Sheet below the lowest threshold, this wave function corresponds to a bound state. In this case, the integrals in $\eta(E_i)$ and $\alpha_k^{(i)}(E_i)$ are real. The normalization for this state is well-defined and the $\alpha_k^{(i)}(E_i)$ can be chosen such that

$$\begin{aligned}
 1 &= \sum_{jk}^D \alpha_k^{(i)}(E_i) \left(\delta_{jk} + \sum_{m=1}^C \int_{a_m} d\omega \frac{f_{km}(\omega) f_{jm}^*(\omega)}{(E_i - \omega)^2} \right) \alpha_j^{(i)*}(E_i) \\
 &= \sum_{jk}^D \alpha_k^{(i)}(E_i) \eta'_{kj}(E_i) \alpha_j^{(i)*}(E_i),
 \end{aligned} \tag{18}$$

where $\eta'_{kj}(E)$ being the derivative of $\eta_{kj}(E)$ w.r.t E . Actually, this equation has a probabilistic explanation. The first term on the right-hand side of the equal sign represents the probability of finding the bare discrete states in the bound state, while the second one represents those of finding the bare continuum states in it. If we define

$$Z_k^{(i)} = |\alpha_k^{(i)}(E_i)|^2, \quad X_m^{(i)} = \sum_{k,j=1}^D \alpha_k^{(i)}(E_i) \left(\int_{a_m} d\omega \frac{f_{km}(\omega) f_{jm}^*(\omega)}{(E_i - \omega)^2} \right) \alpha_j^{(i)*}(E_i), \tag{19}$$

then, $Z^{(i)} \equiv \sum_k Z_k^{(i)}$ is called the elementariness and $X^{(i)} \equiv \sum_{m=1}^C X_m^{(i)}$ is called the compositeness for the bound state. When the solution E_i resides on the unphysical sheet, it is necessary to deform the integral contour to bypass the pole position in different integrals, as illustrated in Fig. 1. For resonance poles on the m -th sheet, the integral contour for the first m -th integral should be deformed accordingly, following the contour shown in Fig. 1. In such cases, the usual definition of the normalization may not be well-defined, as the integral contour for the pole and its conjugate pole are not consistent. Therefore, it becomes necessary to define the normalization through the inner product of the state and its conjugate state which corresponds to the conjugate pole. The resulting normalization is similar to Eq. (18) with E_i replaced by the pole position on the unphysical sheet and the integral contour suitably deformed. However, it is important to note that the probabilistic interpretation of each term in the sum will no longer hold, as the terms may not be real or positive for poles on the unphysical plane.

Now we come to the case with degenerate threshold. If there are different continuum states with the same threshold a_n , with degeneracy h_n , we need to add another label κ to the continuum to denote the different continuum states sharing the same threshold, $|\omega, n\kappa\rangle$. Thus all the indices in the equations labelling the continuum states would include the additional indices κ to label the degenerate states, for example, $f_{in}(\omega)$, α_{ni} , γ_m , ψ_{mn} , Ψ_m become $f_{i,n\kappa}$, $\alpha_{n\kappa,i}$, $\gamma_{m\kappa}$, $\psi_{m\kappa,n\kappa'}$. There will be $\mathfrak{h} = \sum_{n=1}^C h_n$ continuum states. The sum over the continuum states also needs to sum over the κ . The matrix γ is defined as $\gamma_{m\kappa,n\kappa'}(E) = \delta_{m\kappa,n\kappa'} \Theta(E - a_n)$. \mathbf{f} matrix becomes a $D \times \mathfrak{h}$ matrix and $\boldsymbol{\eta}$ is still a $D \times D$ matrix. With all these changes, the previous discussion and equations can be smoothly used in this case.

III. A MORE GENERAL MODEL WITH A SEPARABLE CONTINUUM-CONTINUUM INTERACTION

In the previous case, we considered a scenario where a bare continuum state is only coupled to the bare discrete states but not to the other continuum states. However, when the direct interactions between continuum states become

significant, it is more appropriate to include the corresponding term in the interaction Hamiltonian H_I . Analytically solving the Hamiltonian with a general continuum-continuum interaction is generally not feasible. Therefore, in this section, we will focus on the case with a separable interaction, which still allows for an exactly solvable solution.

The Hamiltonian, including D discrete states and C continua with factorizable self-interacting contact terms, can be expressed as

$$\begin{aligned}
H = & \sum_{i=1}^D M_i |i\rangle\langle i| + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \omega |\omega; n\rangle\langle \omega; n| \\
& + \sum_{m,n=1}^C v_{mn} \left(\int_{a_m}^{\infty} d\omega g_m(\omega) |\omega; m\rangle \right) \left(\int_{a_n}^{\infty} d\omega g_n^*(\omega) \langle \omega; n| \right) \\
& + \sum_{j=1}^D \sum_{n=1}^C \left[|j\rangle \left(\int_{a_n}^{\infty} d\omega f_{jn}^*(\omega) \langle \omega; n| \right) + \left(\int_{a_n}^{\infty} d\omega f_{jn}(\omega) |\omega; n\rangle \right) \langle j| \right]. \tag{20}
\end{aligned}$$

In this case, the new coupling constants v_{mn} between two continuum states have been incorporated in the interaction terms, and $v_{mn} = v_{nm}^*$ is satisfied to meet the Hermiticity requirement. The form-factor functions $g_n(\omega)$ are involved in the interaction between two continuum states, and $f_{jn}(\omega)$ represents the interaction vertex between the j -th discrete state and the n -th continuum state. In the case where the coupling constant matrix v_{mn} is degenerate, certain continuum states may decouple from the contact interaction. This allow us to choose a suitable set of continuum basis states in which the decoupled states do not appear in the contact interaction terms. Consequently, the number of g_m functions required is reduced, and the upper limit of the sum involving these functions becomes less than C . However, it is important to note that although these states decouple in the contact terms, they may remain to be coupled with the discrete states. Nevertheless, this consideration does not significantly alter the subsequent discussion. For the sake of simplicity, we assume that the coupling constant matrix v_{mn} is non-degenerate.

Similar to the previous case, we are also going to solve the Hamiltonian eigenfunction $H|\Psi(E)\rangle = E|\Psi(E)\rangle$. The eigenstate of the Hamiltonian with eigenvalue E can be expanded in terms of the discrete states and the continuum states as

$$|\Psi(E)\rangle = \sum_{i=1}^D \alpha_i(E) |i\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \psi_n(E, \omega) |\omega; n\rangle. \tag{21}$$

Inserting this ansatz into the eigenvalue equation and projecting to the discrete eigenstates or the continuum ones, one can find two sets of equations

$$(M_j - E)\alpha_j(E) + A_j(E) = 0, \quad j = 1, \dots, D \tag{22}$$

$$\sum_{j=1}^D \alpha_j(E) f_{jn}(\omega) + (\omega - E)\psi_n(E, \omega) + \sum_{m=1}^C v_{nm} B_m(E) g_n(\omega) = 0, \quad n = 1, \dots, C, \quad \text{and } \omega > a_n \tag{23}$$

where, two new integration functions $A_j(E)$ and $B_n(E)$ have been defined as

$$A_j(E) \equiv \sum_{n=1}^C \int_{a_n}^{\infty} d\omega f_{jn}^*(\omega) \psi_n(E, \omega), \quad B_n(E) \equiv \int_{a_n}^{\infty} d\omega g_n^*(\omega) \psi_n(E, \omega).$$

Since we have assumed that the continuum-continuum coupling constants v_{mn} s are not degenerate, there are C independent $B_n(E)$ functions. On the contrary, if v_{mn} matrix is degenerate, the only change is that there will be fewer g_n and B_n functions.

Similarly as discussed previously, if the eigenvalue $E \in [a_l, a_{l+1})$ for $l < C$, there should be l continuum solutions $|\Psi_m(E)\rangle$, $m = 1, 2, \dots, l$ with the same eigenvalue E . As the interactions are gradually deactivated, it is required that these continuum solutions tends to well-defined states $|E, m\rangle$. This ensures that in the absence of interactions, the continuum solutions can be uniquely determined as the continuum states $|E, m\rangle$, thus eliminating any ambiguity in their characterization. This requirement guarantees a smooth transition from the interacting system to the non-interacting system.

Under these specific conditions, the l continuum state solutions for $a_l < E < a_{l+1}$ coincide with the first l states for $E > a_C$. Consequently, it is sufficient to solve for the solutions when $E > a_C$, and then the first l solutions can be

obtained for the range $E < a_l$. For each continuum state $|\Psi_m(E)\rangle$ with $E > a_C$, the corresponding coefficients are denoted as α_{jm} , $\psi_{nm}(E, \omega)$, A_{jm} , and B_{nm} as in eqs. (21), (22), and (23). Then, one could obtain

$$\alpha_{jm}^{\pm}(E) = \frac{1}{E - M_j} A_{jm}^{\pm}(E), \quad (24)$$

$$\psi_{nm}^{\pm}(E, \omega) = \delta_{nm} \delta(\omega - E) + \frac{1}{E - \omega \pm i0} \left(\sum_{j=1}^D \frac{A_{jm}^{\pm}(E) f_{jn}(\omega)}{E - M_j} + \sum_{n'=1}^C v_{nn'} B_{n'm}^{\pm}(E) g_n(\omega) \right). \quad (25)$$

The procedure of solving the equation Eq. (25) is straightforward but intricate. The strategy is to apply the operations $\sum_n \int_{a_n} d\omega f_{jn}^*(E) \times$ on the left-hand side of eq. (24) and $\sum_n v_{nm}^* \int d\omega g_n^*(\omega) \times$ on the left-hand side of eq. (25). After this, we can derive the following expressions:

$$0 = f_{jm}^*(E) - \sum_{j'=1}^D \frac{A_{j'm}(E)}{E - M_{j'}} \left(\delta_{j'j} (E - M_{j'}) - \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{j'n}(\omega) f_{jn}^*(\omega)}{E - \omega \pm i0} \right) + \sum_{n'=1}^C \left(B_{n'm}(E) \sum_{n=1}^C v_{nn'} \int_{a_n} \frac{f_{jn}^*(\omega) g_n(\omega)}{E - \omega \pm i0} \right) \quad (26)$$

$$0 = v_{mm'}^* g_m^*(E) + \sum_{j=1}^D \alpha_{jm} \sum_{n=1}^C v_{nm'}^* \int_{a_n} \frac{f_{jn}(\omega) g_n^*(\omega)}{E - \omega \pm i0} - \sum_{n'=1}^C B_{n'm}(E) \left(v_{n'm'}^* - \sum_{n=1}^C v_{nm'}^* v_{nn'} \int_{a_n} \frac{g_n(\omega) g_n^*(\omega)}{E - \omega \pm i0} \right). \quad (27)$$

The matrix representation proves to be a valuable tool in simplifying the derivation process and achieving concise results. In this context, we introduce matrix \mathbf{Y} and \mathbf{F} with $(C + D) \times C$ dimension, matrices \mathbf{V}^A and \mathbf{V}^B , with dimensions $D \times D$ and $C \times C$ respectively, matrices \mathbf{V}^{AB} and \mathbf{V}^{BA} , having dimensions $D \times C$ and $C \times D$ respectively, and finally, a $(C + D) \times (C + D)$ matrix \mathbf{M} encompassing \mathbf{V}^A , \mathbf{V}^B , \mathbf{V}^{AB} and \mathbf{V}^{BA} as follows

$$\begin{aligned} (\mathbf{Y})_{jm} &= \alpha_{jm} = \frac{A_{jm}(E)}{E - M_j}, \quad (\mathbf{F})_{jm} = f_{jm}^*, \quad \text{for } m = 1, \dots, C; j = 1, \dots, D, \\ (\mathbf{Y})_{mn} &= B_{m-D, n}, \quad (\mathbf{F})_{mn} = v_{n, m-D}^* g_n^*, \quad \text{for } n = 1, \dots, C; m = D + 1, \dots, D + C, \\ (\mathbf{V}^A)_{ij} &= \delta_{ij} (E - M_i) - \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{jn}(\omega) f_{in}^*(\omega)}{E - \omega \pm i0}, \quad \text{for } i = 1, \dots, D; j = 1, \dots, D, \\ (\mathbf{V}^B)_{mn} &= \left(v_{mn} - \sum_{l=1}^C v_{ml} v_{ln} \int_{a_m} \frac{g_l(\omega) g_l^*(\omega)}{E - \omega \pm i0} \right), \quad \text{for } m = 1, \dots, C; n = 1, \dots, C, \\ (\mathbf{V}^{AB})_{im} &= - \sum_{n=1}^C v_{n, m} \int_{a_n} \frac{f_{in}^*(\omega) g_n(\omega)}{E - \omega \pm i0}, \quad \text{for } i = 1, \dots, D; m = 1, \dots, C, \\ (\mathbf{V}^{BA})_{mj} &= - \sum_{n=1}^C v_{nm}^* \int_{a_n} \frac{f_{j, n}(\omega) g_n^*(\omega)}{E - \omega \pm i0}, \quad \text{for } j = 1, \dots, D; m = 1, \dots, C, \\ \mathbf{M}_{IJ} &= \begin{pmatrix} \mathbf{V}^A(E) & \mathbf{V}^{AB}(E) \\ \mathbf{V}^{BA}(E) & \mathbf{V}^B(E) \end{pmatrix}_{IJ}, \quad \text{for } I, J = 1, \dots, C + D \end{aligned} \quad (29)$$

Similar to the coefficients α and ψ , we have omitted the superscript “ \pm ” in the notations of the matrices \mathbf{Y} , \mathbf{V} , and \mathbf{M} , which can be inferred from the surrounding contexts. With these matrices, the two equations (26) and (27) above can be expressed in matrix form

$$\mathbf{M} \cdot \mathbf{Y} = \mathbf{F}. \quad (30)$$

or in component form

$$\sum_{j=1}^D V_{ij}^A(E) \alpha_{jm}(E) + \sum_{n=1}^C V_{in}^{AB}(E) B_{nm}(E) = f_{im}^*(E), \quad (31)$$

$$\sum_{j=1}^D V_{nj}^{BA}(E) \alpha_{jm}(E) + \sum_{n'=1}^C V_{nn'}^B(E) B_{n'm}(E) = g_m^*(E) \delta_{m, n}. \quad (32)$$

Before further proceeding, let us look at some properties of these matrices. From the relation $v_{mn}^* = v_{nm}$, we can observe the following symmetric properties:

$$\begin{aligned} (\mathbf{V}^{A+})_{ij}^* &= (\mathbf{V}^{A-})_{ji}, & (\mathbf{V}^{B+})_{mn}^* &= (\mathbf{V}^{B-})_{nm}, & (\mathbf{V}^{AB+})_{jn}^* &= (\mathbf{V}^{BA-})_{nj}, \\ \mathbf{M}^{+\dagger} &= \mathbf{M}^-. \end{aligned} \quad (33)$$

In the case where $v_{mn}(\omega)$, $f_{jm}(\omega)$ and $g_n(\omega)$ are real functions, the corresponding function matrices possess the real analyticity property. As a result, they can be analytically continued to the entire complex E plane and satisfy the Schwartz reflection property. Moreover, the analytically continued function matrices can relate the $+i0$ and $-i0$ counterparts, representing the limits on the upper and lower edges along the real axis above the threshold. In the cases that $v_{mn}(\omega)$, $f_{jm}(\omega)$ and $g_n(\omega)$ are complex, the function matrices will no longer be real analytic, but the determinant of the matrix \mathbf{M} , denoted as $\det \mathbf{M}$, remains real analytic. Thus the analytically continued determinant exhibits the Schwartz reflection symmetry, $\det \mathbf{M}(z) = \det \mathbf{M}^*(z^*)$.

In general, $\det \mathbf{M}$ is nonzero for real E values above the lowest threshold. As a result, \mathbf{M} possesses an inverse, and \mathbf{Y} can be obtained by $\mathbf{Y} = \mathbf{M}^{-1} \cdot \mathbf{F}$. Then, by inserting $A_{jm}(E)$ or $\alpha_{jm}(E)$ and B_{nm} into Eq. (25), one obtains the coefficients ψ_{nm}^\pm , and the continuum eigenstates are solved to be

$$|\Psi_m^\pm(E)\rangle = |E, m\rangle + \sum_{j=1}^D \alpha_{jm}^\pm(E) \left(|j\rangle + \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{jn}(\omega)}{E - \omega \pm i0} |\omega; n\rangle \right) + \sum_{n, n'=1}^C v_{nn'} B_{n'm}^\pm(E) \int_{a_n} d\omega \frac{g_n(\omega)}{E - \omega \pm i0} |\omega; n\rangle \quad (34)$$

for $E > a_m$. Upon comparison with Eq. (12), this solution is different only in the last term, stemming from the presence of separable potential. Importantly, it can be confirmed that the solution retains the previous normalization condition, $\langle \Psi_m^\pm(E) | \Psi_n^\pm(E') \rangle = \delta(E - E') \delta_{mn}$. This normalization condition guarantees the orthogonality of the wave functions, ensuring their compatibility and consistency within the framework of the problem.

The S -matrix can be obtained as

$$S_{mn}(E, E') = \delta_{mn} \delta(E - E') - 2\pi i \delta(E - E') \left(\sum_{I=1, J=1}^{D+C} (\mathbf{F}^\dagger)_{mI} (\mathbf{M}^+)_{IJ}^{-1} (\mathbf{F}_{Jn}) \right), \quad (35)$$

or

$$\mathbf{S}(E, E') = \mathbf{I} \delta(E - E') - 2\pi i \delta(E - E') \mathbf{F}^\dagger \cdot (\mathbf{M}^+)^{-1} \cdot \mathbf{F} \quad (36)$$

in a simplified matrix form. For a more thorough derivation of the normalization and meticulous calculation of the S -matrix, please refer to the appendix A, where we provide a detailed presentation of the calculations, offering a comprehensive and in-depth derivation of the normalization condition and the S -matrix.

Subsequently, our attention turns towards the derivation of discrete eigenstates. The eigenvalues for the discrete states does not coincide with the spectrum of the continuum states. Thus, using the condition $E \notin [a_i, \infty)$ for $i = 1, \dots, C$, one can solve Eq.(23) and obtain

$$\alpha_j(E) = \frac{1}{E - M_j} A_j(E), \quad (37)$$

$$\psi_n(E, \omega) = \frac{1}{E - \omega} \left(\sum_{j=1}^D \alpha_j(E) f_{jn}(\omega) + \sum_{n'=1}^C v_{nn'} B_{n'}(E) g_n(\omega) \right). \quad (38)$$

By multiplying Eq. (38) with $f_{jn}^*(\omega)$ and $v_{nm}^* g_n^*(\omega)$ separately, and subsequently summing over n and integrating w.r.t. the variable ω , we arrive at the following expressions:

$$0 = - \sum_{j'=1}^D \frac{A_{j'}(E)}{E - M_{j'}} \left(\delta_{j'j} (E - M_{j'}) - \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{j'n}(\omega) f_{jn}^*(\omega)}{E - \omega} \right) + \sum_{n'=1}^C \left(B_{n'}(E) \sum_{n=1}^C v_{nn'} \int_{a_n} \frac{f_{jn}^*(\omega) g_n(\omega)}{E - \omega} \right), \quad (39)$$

$$0 = \sum_{j=1}^D \sum_{n=1}^C v_{nm}^* \frac{A_j(E)}{E - M_j} \int_{a_n} \frac{f_{jn}(\omega) g_n^*(\omega)}{E - \omega} - \sum_{n'=1}^C B_{n'}(E) \left(v_{n'm}^* - \sum_{n=1}^C v_{nm}^* v_{nn'} \int_{a_n} \frac{g_n(\omega) g_n^*(\omega)}{E - \omega} \right). \quad (40)$$

The expressions obtained in these two equations deviate from Eqs. (26) and (27) by the absence of the first terms on the right hand side. Analogous to the definition in Eq. (29), we can introduce the matrices \mathbf{V}^A , \mathbf{V}^{AB} , \mathbf{V}^{BA} , \mathbf{V}^B , and

\mathbf{M} as the analytic continuation of the matrices in Eq. (29) and

$$\mathbf{X}^T = \left(\frac{A_1}{E - M_1}, \dots, \frac{A_D}{E - M_D}, B_1, \dots, B_C \right) = (\alpha_1, \dots, \alpha_D, \dots, B_1, \dots, B_C).$$

Then Eqs.(39) and (40) can be expressed as

$$\mathbf{M} \cdot \mathbf{X} = 0.$$

To obtain nonzero solutions for the vector \mathbf{X} , it is essential to satisfy the condition that the determinant of \mathbf{M} is equal to zero, i.e., $\det \mathbf{M}(E) = 0$. By analytically continuing this equations to different Riemann sheets and solving it on each sheet, we can determine the generalized discrete eigenvalues.

Once the generalized eigenvalues are determined, the vector \mathbf{X} can be solved for each eigenvalue. Substituting the solutions for α_j and B_n into Eq. (38), we obtain the discrete solution from Eq. (21) for each generalized energy eigenvalue,

$$|\Psi^{(i)}(E_i)\rangle = \sum_{j=1}^D \alpha_j^{(i)}(E_i) (|j\rangle) + \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{jn}(\omega)}{E_i - \omega} |\omega; n\rangle + \sum_{n, n'=1}^C v_{nn'} B_{n'}^{(i)}(E_i) \int_{a_n} d\omega \frac{g_n(\omega)}{E_i - \omega} |\omega; n\rangle, \quad (41)$$

where the superscript (i) denotes the i -th discrete solution. The solution for \mathbf{X} is only determined up to a normalization. On the first Riemann sheet, the zeros of the $\det \mathbf{M}$ can only be located on the real axis below a_1 due to the hermicity of the Hamiltonian. These zeros correspond to the discrete eigenvalues E_b . It is possible for the associated states to have a finite norm, and we can impose the normalization condition on the coefficients to ensure

$$\begin{aligned} 1 &= \sum_{i=1}^D |\alpha_i(E_b)|^2 + \sum_{n=1}^C \int_{a_j}^{\infty} d\omega \frac{1}{(E_b - \omega)^2} \left| \sum_{j=1}^D \alpha_j(E_b) f_{jn}(\omega) + \sum_{n'=1}^C v_{nn'} B_{n'}(E_b) g_n(\omega) \right|^2 \\ &= \mathbf{X}(E_b) \cdot \mathbf{M}'(E_b) \cdot \mathbf{X}^*(E_b), \end{aligned} \quad (42)$$

where $\mathbf{M}'(E)$ is the derivative of the matrix w.r.t. E . Within the framework described earlier, each term in the summation can be interpreted as the probability of finding the corresponding bare state within the bound state. However, there could also be complex energy solutions present on different unphysical sheets. As the determinant of \mathbf{M} is a real analytic function, these complex eigenvalue solutions appear as complex conjugate pairs.

As already mentioned, there exist 2^C distinct Riemann sheets. However, for our specific purposes, we focus solely on solutions E_R that reside on the lower half Riemann sheet closest to the physical sheet. These solutions have a significant impact on the physical S -matrix elements. Since E_R lies on a nearby unphysical sheet, the evaluation of the matrix value of \mathbf{M} at this point requires deforming the integral contours to the corresponding sheet around E_R defined in the matrix \mathbf{V} and \mathbf{M} in Eq. (29) as illustrated in Fig. 1 [5]. Also in the state solution Eq. (41), the integral contours are also deformed similarly. The normalization requirement of these states may resemble Eq. (42), but with E_b replaced by E_R , and the integral contour adjusted accordingly following the deformation depicted in Fig. 1. However, it is important to note that there is no probabilistic explanations for each terms in the sum, as they may not be real. Additionally, there can also be real solutions below the lowest threshold a_1 on unphysical sheets, which correspond to virtual states. Similar to the resonant states, the corresponding integral contours should be deformed in Eq. (41) and in Eq. (29) for these states.

IV. APPROXIMATING A GENERAL POTENTIAL USING SEPARABLE POTENTIALS

In scenarios where the interaction potential can be reasonably approximated as separable potentials, wherein the potential can be expressed as the product of two components associated with the ingoing and outgoing states, respectively, the problem can be effectively addressed and hold practical significance. Now the problem is how to approximate a potential using the separable potentials. Before addressing this problem, let us review how a general contact interaction can arise in the elastic and nonelastic scattering.

As discussed in the previous section, the Hamiltonian of a most general model with multiple discrete states and

continuum states and their interactions can be expressed as

$$\begin{aligned}
H &= \sum_{i=1}^D M_i |i\rangle \langle i| + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \omega |\omega; n\rangle \langle \omega; n| \\
&+ \sum_{m,n=1}^C \int_{a_m}^{\infty} d\omega' \int_{a_n}^{\infty} d\omega V_{mn}(\omega', \omega) |\omega'; m\rangle \langle \omega; n| \\
&+ \sum_{j=1}^D \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \left(f_{jn}^*(\omega) |j\rangle \langle \omega; n| + f_{jn}(\omega) |\omega; n\rangle \langle j| \right).
\end{aligned}$$

In the context of nonrelativistic two-particle scattering, i.e. when the in-state and out-state are composed of the same two-particle content, in the angular momentum representation, these continuum states can be expressed as $|\omega, n\rangle = \sqrt{\mu p} |p, JM; lS\rangle$, where J, M, l, S are the quantum numbers for total angular momentum, magnetic angular momentum, relative orbital angular momentum, and total spin, respectively, and are collectively denoted using n and a_n are the threshold. Here, p represents the radial momentum, μ is the reduced mass and $\omega = \frac{p^2}{2\mu} + a_n$ represents the total energy. The normalization of the continuum states is chosen such that the inner product between two continuum states is given by $\langle \omega', n' | \omega, n \rangle = \delta_{n'n} \delta(\omega - \omega')$. The momentum space potential $V_{mn}(\omega', \omega)$ can be derived from the coordinate space potential $V(\mathbf{r})$. For simplicity, we consider only the rotational invariant potential. The potential function $V_{mn}(\omega', \omega)$ arises from the matrix elements $V_{n'n}(\omega', \omega) \equiv \langle \omega', n' | V | \omega, n \rangle = \sqrt{\mu' p'} \sqrt{\mu p} \langle p' J M l' S' | V | p J M l S \rangle$. For simplicity, we look at the case when the in-states and out-states correspond to the same spinless particles, the momentum-space potential $V(k', k)$ and be expressed in terms of the coordinate-space potential $V(r)$ as follows:

$$\langle k', l', m' | V | k, l, m \rangle = \frac{\delta_{l'l} \delta_{m'm}}{k' k} \int r^2 dr \frac{2}{\pi} \hat{j}_l(k'r) V(r) \hat{j}_l(kr), \quad (43)$$

where the $\hat{j}(z)$ represents the Riccati-Bessel function.

In the cases where the in-states and out-states can have different particle compositions, we can generalize the potential $V(\omega', \omega)$ accordingly. In addition to the angular momentum quantum numbers, the labels n and n' can also denote the different particle compositions $|\omega, n\rangle$. If the potential in coordinate space, $V(\mathbf{r}', \mathbf{r})$, in the center-of-mass system, is invariant under rotation, it can be expressed as a function of $\mathbf{r}^2, \mathbf{r}'^2$ and $\mathbf{r} \cdot \mathbf{r}'$. Here, \mathbf{r} and \mathbf{r}' represent the position of the in-state and out-state relative coordinates, respectively. In the case of spinless particle system, the matrix elements for in-states and out-states can be expressed as follows:

$$\begin{aligned}
V_{n'n}(\omega', \omega) &\equiv \langle n' \omega' l m | V | n \omega l m \rangle = \int d\mathbf{r} d\mathbf{r}' \langle n', \omega' l m | \mathbf{r}' \rangle V(\mathbf{r}', \mathbf{r}) \langle \mathbf{r} | n, \omega l m \rangle \\
&= \frac{2}{\pi} \left(\frac{\mu' \mu}{p' p} \right)^{1/2} \int dr dr' \hat{j}_l(pr) \hat{j}_l(p'r') \tilde{V}_l(r', r), \\
\tilde{V}_l(r', r) \delta_{ll'} \delta_{mm'} &= r r' \int d\Omega d\Omega' Y_{l'm'}^*(\hat{\mathbf{r}}') Y_{lm}(\hat{\mathbf{r}}) \tilde{V}(\mathbf{r}', \mathbf{r}),
\end{aligned}$$

where $\hat{\mathbf{r}} = \mathbf{r}/r$. The Wigner-Ekart theorem has been employed to account for the spherical symmetry of the potential $V(\mathbf{r}', \mathbf{r})$. When the in-state and out state can have spins, the total angular momentum are conserved but the orbital angular momentum could be different. We can include the different orbital angular momentum and total spin quantum numbers lS and $l'S'$ into n and n' to label different in-states and out-states,

$$\begin{aligned}
V_{n'n}(\omega', \omega) &\equiv \langle n' \omega' J M | V | n \omega J M \rangle \\
&= \frac{2}{\pi} \left(\frac{\mu' \mu}{p' p} \right)^{1/2} \int dr dr' \sum_{l' S S'} \hat{j}_l(pr) \hat{j}_{l'}(p'r') \tilde{V}_{l' S S'}^{J M}(r', r), \\
\tilde{V}_{l' S S'}^{J M}(r', r) &= r r' \sum_{m m' m_s m'_s} \int d\Omega d\Omega' Y_{l'm'}^*(\hat{\mathbf{r}}') Y_{lm}(\hat{\mathbf{r}}) \tilde{V}_{S S'}(\mathbf{r}', \mathbf{r}) C_{lm S m_s}^{J M} C_{l' m' S' m'_s}^{J M},
\end{aligned}$$

where $C_{lm S m_s}^{J M}$ is the Clebsch-Gordon coefficients. To make further progress, we also suppose that the potential is square integrable for both ω' and ω , and the same for the interaction vertex between the discrete states and the continuum states $f_{jn}(\omega)$, that is

$$\int_{a_m} d\omega' \int_{a_n} d\omega |V_{mn}(\omega', \omega)|^2 = \text{finite}, \quad \int_{a_n} d\omega |f_{jn}(\omega)|^2 = \text{finite}.$$

There are no exact solutions for Hamiltonian for general potentials $V_{mn}(\omega', \omega)$. However, it is well-known that such a potential can be expanded using a sum of separable potentials [29]. For continuum states $|\omega, m\rangle$, we can choose a set of complete base functions $\tilde{g}_{m\rho}(\omega)$, with $\int_{a_m} d\omega \tilde{g}_{m\rho}^*(\omega) \tilde{g}_{m\delta}(\omega) = \delta_{\rho\delta}$ and $\sum_{\delta} \tilde{g}_{m\delta}(\omega') \tilde{g}_{m\delta}(\omega) = \delta(\omega' - \omega)$. The basis sets for different continuum states, i.e. for different m and n , do not need to be the same. The potential $V_{mn}(\omega', \omega)$ can then be expanded as

$$V_{mn}(\omega', \omega) = \sum_{\rho\delta} v_{mn,\rho\delta} \tilde{g}_{m\rho}(\omega) \tilde{g}_{n\delta}^*(\omega).$$

In the following we will use the Greek letter ρ, δ to label the basis, and repeated Greek letters are summed over without explicit sum symbol and the sum symbol for the Latin letters would still be left explicit. The coefficient matrix composed of $v_{mn,\rho\delta}$ is hermitian $v_{mn,\rho\delta} = v_{nm,\delta\rho}^*$ and is supposed to be non-degenerate. In general, there are infinite number of bases, the sum of δ and ρ is up to infinity. Since the expansion coefficients $v_{mn,\rho\delta}$ are small at large enough order, one can make an approximation and truncate the series to a finite order N , i.e. $v_{mn,\rho\delta} = 0$ for $\rho, \delta > N$. Then, the general Hamiltonian for multiple continuum states and discrete states can be recast as

$$\begin{aligned} H = & \sum_{i=1}^D M_i |i\rangle \langle i| + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \omega |\omega; n\rangle \langle \omega; n| \\ & + \sum_{m,n=1}^C v_{mn,\rho\delta} \left(\int_{a_m}^{\infty} d\omega' \tilde{g}_{m\rho}(\omega') |\omega'; m\rangle \right) \left(\int_{a_n}^{\infty} d\omega \tilde{g}_{n\delta}^*(\omega) \langle \omega; n| \right) \\ & + \sum_{j=1}^D \sum_{n=1}^C \left[|j\rangle \left(\int_{a_n}^{\infty} d\omega f_{jn}^*(\omega) \langle \omega; n| \right) + \left(\int_{a_n}^{\infty} d\omega f_{jn}(\omega) |\omega; n\rangle \right) \langle j| \right]. \end{aligned} \quad (44)$$

The general eigenstate for this eigenvalue problem can be expanded using the bare discrete states and the bare continuum states

$$|\Psi(E)\rangle = \sum_{i=1}^D \alpha_i(E) |i\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \psi_n(E, \omega) |\omega; n\rangle. \quad (45)$$

Similar to previous sections, for $|\Psi_m\rangle$, the corresponding α_i and ψ_n will have another index m , i.e. α_{im} and ψ_{nm} .

With the same procedures as the previous section, the approximate properly normalized continuum states can be solved as

$$\begin{aligned} |\Psi_m^{\pm}(E)\rangle = & |E; m\rangle + \sum_{j=1}^D \alpha_{jm}^{\pm}(E) (|j\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \frac{f_{jn}(\omega)}{E - \omega \pm i0} |\omega; n\rangle) \\ & + \left(\sum_{n,n'=1}^C v_{nn',\delta'\rho} \psi_{n'm\rho}^{\pm}(E) \int_{a_n}^{\infty} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0} |\omega; n\rangle \right), \end{aligned} \quad (46)$$

where $\alpha_{im}^{\pm}(E)$ and $\psi_{n'm\rho}^{\pm}(E) \equiv \int_{a_n}^{\infty} \psi_{n'm}(\omega) \tilde{g}_{n\rho}(\omega)$ can be obtained in Eq.(B11). The S -matrix can then be obtained,

$$\langle \Psi_m^-(E) | \Psi_n^+(E') \rangle = \delta_{mn} \delta(E - E') - 2\pi i \delta(E - E') \left(\tilde{\mathbf{F}}_m^{\dagger} \cdot (\tilde{\mathbf{M}}^+)^{-1} \cdot \tilde{\mathbf{F}}_m \right), \quad (47)$$

where the matrix $\tilde{\mathbf{M}}^+$ with dimension $(D + NC) \times (D + NC)$ and vector $\tilde{\mathbf{F}}_m$ with dimension $(D + NC)$ are defined in Eq.(B7). The detailed calculation is left to appendix B. The discrete eigenvalues can be obtained by solving the equation $\det \tilde{\mathbf{M}}(E) = 0$. The discrete state corresponding to eigenvalue E_i can be expressed as

$$|\Psi^{(i)}(E_i)\rangle = \sum_{j=1}^D \alpha_j^{(i)}(E_i) (|j\rangle + \int_{a_n}^{\infty} d\omega \frac{f_{jn}(\omega)}{E_i - \omega} |\omega; n\rangle) + \sum_{n'=1}^C v_{nn',\delta'\rho} \psi_{n'm\rho}^{(i)}(E_i) \int_{a_n}^{\infty} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E_i - \omega} |\omega; n\rangle, \quad (48)$$

where the integral contour needs to be deformed for E_i on unphysical sheets as before.

In fact, we could go further and also expand the interaction function $f_{jm}(\omega)$ using the same set of basis $\tilde{g}_{m\delta}(\omega)$ as in the corresponding contact interaction involving the continuum state $|\omega, m\rangle$,

$$f_{jm}(\omega) = \sum_{\delta} f_{jm\delta} \tilde{g}_{m\delta}(\omega), \quad f_{jm\delta} = \int d\omega f_{jm}(\omega) \tilde{g}_{m\delta}^*(\omega),$$

and also make an approximation by truncating the series to the N -th order the same as in the contact terms, that is, $f_{jm\delta} = 0$ for $\delta > N$. This may reduce the dimension of the matrix $\tilde{\mathbf{M}}$ and may also simplify the numerical calculation. Then, the general Hamiltonian for multiple continuum states and discrete states can be recast as

$$\begin{aligned}
H = & \sum_{i=1}^D M_i |i\rangle\langle i| + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \omega |\omega; n\rangle\langle \omega; n| \\
& + \sum_{m,n=1}^C v_{mn,\rho\delta} \left(\int_{a_m}^{\infty} d\omega' \tilde{g}_{m\rho}(\omega') |\omega'; m\rangle \right) \left(\int_{a_n}^{\infty} d\omega \tilde{g}_{n\delta}^*(\omega) \langle \omega; n| \right) \\
& + \sum_{j=1}^D \sum_{n=1}^C \left[f_{jn\delta}^* |j\rangle \left(\int_{a_n}^{\infty} d\omega \tilde{g}_{n\delta}^*(\omega) \langle \omega; n| \right) + f_{jn\delta} \left(\int_{a_n}^{\infty} d\omega \tilde{g}_{n\delta}(\omega) |\omega; n\rangle \right) \langle j| \right]. \tag{49}
\end{aligned}$$

This case is more like the cases discussed in [10, 26], where the same form factor comes with the continuum both in the discrete-continuum and continuum-continuum interaction. Using the eigenstate ansatz

$$\begin{aligned}
|\Psi(E)\rangle = & \sum_{i=1}^D \alpha_i(E) |i\rangle + \int_{a_n} d\omega \psi_n(E, \omega) |\omega; n\rangle \\
= & \sum_{i=1}^D \alpha_i(E) |i\rangle + \sum_{n=1}^C \psi_{n\delta}(E) \int_{a_n} d\omega \tilde{g}_{n\delta}(\omega) |\omega; n\rangle, \tag{50}
\end{aligned}$$

one can proceed solving the eigenvalue problem similarly to the previous section, of which the details will be left to appendix C. The properly normalized continuum state can be solved and expressed as

$$|\Psi_m^\pm(E)\rangle = \sum_{i=1}^D \alpha_{im}^\pm(E) |i\rangle + |E, m\rangle + \sum_{n=1}^C \sum_{n'=1}^C \psi_{n'm\rho}^\pm(E) V_{n\delta', n'\rho}(E) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0} |\omega; n\rangle, \tag{51}$$

where the α_{im}^\pm and $\psi_{n'm\rho}^\pm$ can be solved from Eq.(C9). The S -matrix can be obtained

$$\begin{aligned}
\langle \Psi_m^-(E) | \Psi_n^+(E') \rangle = & \delta_{mn} \delta(E - E') - 2\pi i \delta(E - E') \sum_{n=1}^C \sum_{n'=1}^C \psi_{n'm\rho}^{-*}(E') V_{n\delta', n'\rho}^*(E') \tilde{g}_{n\delta'}^*(E) \\
= & \delta_{mn} \delta(E - E') - 2\pi i \delta(E - E') (\tilde{\mathbf{F}}_m^\dagger \cdot (\tilde{\mathbf{W}}^+)^{-1} \cdot \tilde{\mathbf{F}}_n), \tag{52}
\end{aligned}$$

where the $NC \times NC$ matrix $\tilde{\mathbf{W}}^+$ and NC dimensional vector $\tilde{\mathbf{F}}_m$ are defined in Eq. (C8) and Eq. (C7). Similar to previous section, the generalized energy eigenvalues for the discrete state can be obtained from the $\det \tilde{\mathbf{M}}(E) = 0$, and for each eigenvalue E_i , $\psi_{n\rho}^{(i)}$ can be solved from $\tilde{\mathbf{M}} \cdot \tilde{\mathbf{Y}} = 0$, where the matrix $\tilde{\mathbf{M}}$ and $\tilde{\mathbf{Y}}$ are defined in Eq. (C6) and Eq. (C7). Then we have the discrete eigenstates,

$$|\Psi^{(i)}(E_i)\rangle = \sum_{n'=1}^C \psi_{n'\rho}^{(i)}(E_i) \left[\sum_{j=1}^D \frac{f_{jn'\rho}^*}{E_i - M_j} |j\rangle + V_{n\delta', n'\rho}(E_i) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E_i - \omega} |\omega; n\rangle \right], \tag{53}$$

with

$$\tilde{\mathbf{Y}}^{(i)\dagger}(E_i) \cdot \mathbf{V}(E_i) \cdot \tilde{\mathbf{W}}'(E_i) \cdot \mathbf{V}(E_i) \cdot \tilde{\mathbf{Y}}^{(i)}(E_i) = 1,$$

where $\tilde{\mathbf{W}}'(E)$ being the derivative of the matrix w.r.t. E and the integral contours being deformed for resonances and virtual states as before.

V. DISCUSSION

With the exact solution when there are contact interactions, we can try to discuss the effect of turning on some small coupling to the mass of the discrete states. The basic consideration is like this: At the leading order where there is no interaction, the discrete state is determined by $E - \mu = 0$, and the bare mass is the solution. When there




FIG. 2. T

he real part of the dispersive integral $\text{Re}G_1(E)$ for $|f(\omega)|^2 = \sqrt{\omega - a_1}e^{-E/\Lambda}$, with $\Lambda = 5$, $a_1 = 0$.

is some interaction with coupling constant λ turned on, an equation like $E - \mu + \lambda\chi(E) = 0$ needs to be considered where $\chi(E)$ is real and small near $E = \mu$. The next-to-leading-order solution would be $E = \mu - \lambda\chi(\mu) + O(\lambda^2)$, and from the sign of $\chi(\mu)$ we can determine the tendency of behavior of the solution. If we know that the $\chi(E)$ is a positive or negative monotonic function we can also determine whether the solution is below or above μ for large λ . For example, it is a positive increasing function, it is easy to see that the solution will be smaller than μ for any positive λ . When the leading order equation $E - \mu = 0$ is replaced by another equation, $\zeta(E) = 0$ where ζ is real and monotonic in a sufficiently large region near the zero point $E = \mu$, similar analysis can also be done.

We need also to look at the behavior of the dispersion integral like $F(E) = \int_a^\infty d\omega \frac{|f(\omega)|^2}{E - \omega + i0}$. When $E < a$, it is a pure negative decreasing function. When $E > a$, the imaginary part is $-\pi|f(E)|^2$ which is purely negative for $E > a$ and the real part is the principal value integral in $F(E)$. The real part is negative and increasing with E near the threshold and goes through zero point E_0 and reaches a positive maximum and then goes down towards zero when E tends to infinity. See Fig. 2 for an example. Usually, the integrand includes a phase space factor $\rho(E) \propto \sqrt{E - a_1}$ which suppresses the integrand near the threshold a , which causes E_0 to be high. See Fig. 2 for the case with $|f(\omega)|^2 = \sqrt{\omega - a_1}e^{-E/\Lambda}$. We can see that only when E is going up near the Λ which characterizes the inverse of the interaction range, $\text{Re}F(E)$ becomes positive.

First, let us look at the case when there are only two continuum, to see how turning on small coupling between the two continua changes the discrete spectrum of the coupled channel system. In this case, the \mathbf{M} matrix becomes

$$\mathbf{M} = \begin{pmatrix} v_{11}(1 - v_{11}G_1(E)) - |v_{12}|^2G_2(E) & v_{12}(1 - v_{11}G_1(E)) - v_{12}v_{22}G_2(E) \\ v_{21}(1 - v_{11}G_1(E)) - v_{22}v_{21}G_2(E) & v_{22}(1 - v_{22}G_2(E)) - |v_{12}|^2G_1(E) \end{pmatrix}, \quad (54)$$

$$G_i = \int_{a_i}^\infty d\omega \frac{|g_i(\omega)|^2}{E - \omega + i\epsilon},$$

where a_1 and a_2 are the thresholds for the two channels with $a_1 < a_2$. It is well known that when there is an attractive interaction in the single channel case there could be a bound state. This can be understood by looking at the \mathbf{M} matrix when there is only one continuum,

$$M_{11} = v_{11}(1 - v_{11}G_1(E)).$$

Since $G_1(E) < 0$ for $E < a_1$, the zero point of this M_{11} below the threshold a_1 can appear only when v_{11} is negative, namely at E_0 . When there is the second channel coupled with this channel, a simple result can be reached is that when $v_{22} = 0$ and only the interaction between the two channels is left $v_{12} \neq 0$, the bound state will go deeper below the threshold. This can be proved by directly calculating the $\det \mathbf{M}$ matrix and obtaining

$$\det \mathbf{M} = -|v_{12}|^2 \left(1 - v_{11}G_1(E) - |v_{12}|^2G_1(E)G_2(E) \right).$$

It is easy to see that since the last term in the bracket is negative and the term $-v_{11}G_1(E)$ is also negative and become less negative with smaller E for $E < a_1$, E should be more smaller to have a zero point of $\det \mathbf{M}$, denoted as E_{0b} . When v_{22} is also turned on, the result depends on the sign of v_{22} . Its contribution to the $\det \mathbf{M}$ at the zero point E_{0b} can be expressed as

$$-|v_{12}|^2v_{22}G_2(E) \left(1 - v_{22}G_2(E) \right) - |v_{12}|^2v_{22}^2(G_2(E))^2 = -|v_{12}|^2v_{22}G_2(E).$$

It is easy to see that when the v_{22} is negative, that is, more attractive, this term contributes negatively to $\det \mathbf{M}$ for $E < a_1$, thus also making the bound state more deeper, whereas when v_{22} is positive, the bound state moves oppositely from E_{0b} . Thus with both v_{12} and positive v_{22} turned on, whether the bound state is moving up or down

from E_0 depends on the competing of the effect from v_{22} and v_{12} . Since v_{12} only appears in the form of $|v_{12}|^2$, the phase or sign of v_{12} is irrelevant to this result in this case.

Then let us look at the effect of turning on the first channel to the discrete spectrum of the second channel. Similar to previous case, when there is only an attractive interaction in the second channel, i.e. $v_{22} < 0$ with $v_{11} = v_{12} = 0$, there could be a dynamically generated bound state (E_b) below a_2 and we suppose that it is also above a_1 . Then, we turn on the v_{12} , that is, when $v_{11} = 0$ and $v_{12} \neq 0$, we have

$$\det \mathbf{M} = -|v_{12}|^2 \left(1 - v_{22}G_2(E) - |v_{12}|^2 G_1(E)G_2(E) \right).$$

Since $E_b > a_1$, the $G_1(E)$ factor in the last term contributes a imaginary $-\pi i |g_1(\omega)|^2$ factor, and combining with the other factors produces a negative imaginary part of $|v_{12}|^2$ order in the bracket. Then $\det M = 0$ requires the state corresponding to E_b moves onto the complex plane so that the term $-v_{22}G_2(E)$ generates a negative imaginary part of order v_{12}^2 to cancel the imaginary part of the last term in the bracket. Thus, the pole moves continuously to the second Riemann sheet. Whether the mass of the pole is moving up or down depends on the real part of the $G_1(E)$ of the last term, $\text{Re}G_1(E_b) = P.V. \int_{a_1}^{\infty} d\omega \frac{|g_1(\omega)|^2}{E_b - \omega}$: if it is negative(positive) the mass will go up(down) since $G_2(E)$ is negative for $E < a_1$ and the last term in the bracket is positive(negative). This happens to the example form factor in Fig. 2 when the Λ is much larger than a_2 , and $\text{Re}G_1(E)$ is negative and monotonically increasing near E_b , and then turning on the v_{12} will cause the mass of the dynamically generated state in second channel going down. When v_{11} is also turned on, similar to previous case, at E_b , the term of order v_{11} in $\det \mathbf{M}$ is

$$-|v_{12}|^2 v_{11} G_1(E_b).$$

Thus, if $\text{Re}G_1(E_b)$ is negative, a negative v_{11} will cause the mass of the dynamical state becoming smaller because of the negative real part of above term. This effect seems to propagate the attractive interaction of the first channel to the second channel. In contrast, a positive v_{11} tends to make the mass of the state larger, which will compete with the effect of the v_{12} to determine the final effect. Notice that if $\text{Re}G_1(E_b)$ is positive the effect will be the opposite. In the example of exponential form factor case, this would happen if the typical interaction scale Λ is small compared to the second threshold a_2 and $E_0 < a_2$, and then the bound state can be located between E_0 and a_2 .

Besides the negative v_{11} and v_{22} , pure v_{12} may also generate bound states. If $v_{11} = v_{22} = 0$, $\det \mathbf{M} = -|v_{12}|^2 (1 - |v_{12}|^2 G_1(E)G_2(E))$. Since $G_1(E)$ and $G_2(E)$ both are negative below the first threshold, there could also be a solution to $\det \mathbf{M} = 0$ when $|v_{12}|^2$ is large enough. When $|v_{12}|^2$ becomes smaller, the bound state would go up through the threshold to the second Riemann sheet and becomes a virtual state or a resonance. Thus only turning on v_{12} is equivalent to have more attractive interaction when there are only two continuum states.

Next, we add a discrete state with bare mass μ with coupling vertex functions to the two continua, $f_1(\omega)$ and $f_2(\omega)$. Now the \mathbf{M} matrix is

$$\mathbf{M} = \begin{pmatrix} (E - \mu) - \mathcal{F}_1(E) - \mathcal{F}_2(E) & -v_{11}\mathcal{F}_1^g(E) - v_{21}\mathcal{F}_2^g(E) & -v_{12}\mathcal{F}_1^g(E) - v_{22}\mathcal{F}_2^g(E) \\ -v_{11}\mathcal{F}_1^{g\dagger}(E) - v_{21}\mathcal{F}_2^{g\dagger}(E) & v_{11}(1 - v_{11}G_1(E)) - |v_{12}|^2 G_2(E) & v_{12}(1 - v_{11}G_1(E)) - v_{12}v_{22}G_2(E) \\ -v_{12}\mathcal{F}_1^{g\dagger}(E) - v_{22}\mathcal{F}_2^{g\dagger}(E) & v_{21}(1 - v_{11}G_1(E)) - v_{22}v_{21}G_2(E) & v_{22}(1 - v_{22}G_2(E)) - |v_{12}|^2 G_1(E) \end{pmatrix}, \quad (55)$$

$$\text{where } \mathcal{F}_n = \int_{a_n} d\omega \frac{f_n(\omega)f_n^*(\omega)}{E - \omega + i0}, \quad \mathcal{F}_n^g = \int_{a_n} d\omega \frac{f_n^*(\omega)g_n(\omega)}{E - \omega + i0}, \quad \mathcal{F}_n^{g\dagger} = \int_{a_n} d\omega \frac{f_n(\omega)g_n^*(\omega)}{E - \omega + i0}.$$

When there is only the interaction between discrete state and continua, i.e. $v_{11} = 0, v_{12} = 0, v_{22} = 0$, we need only to look at the $M_{11} \equiv (E - \mu) - \mathcal{F}_1(E) - \mathcal{F}_2(E) = 0$. When $\mu < a_1$, since $\mathcal{F}_1(\mu)$ and $\mathcal{F}_2(\mu)$ are negative, the solution, denoted as E_μ , would be smaller than μ , which means that when the discrete state is below both continuum, turning on the interaction between discrete states and the continuum makes the discrete states go deeper below the thresholds. When $a_1 < \mu < a_2$ and suppose $\text{Re}\mathcal{F}_1(\mu) < 0$ as in the exponential form factor with large Λ , a weak interaction between the discrete state and the two continua will also make the mass of the discrete state smaller. Since $\mathcal{F}_1(\mu)$ has an negative imaginary part, the discrete state solution would develop a negative imaginary part and move to the complex plane of the second Riemann sheet. When $\mu > a_2$, similar discussion can be done. When $\text{Re}(\mathcal{F}_{1,2}(\mu)) < 0$, they will play the role of attractive interactions to push mass of the discrete state down. The negative imaginary parts of $\mathcal{F}_{1,2}(\mu)$ cause the solution goes down the the third sheet of the Riemann sheet.

Now, we gradually turn on $v_{11} \neq 0$ and leave $v_{12} = v_{22} = 0$. Then we need to look at the zero point of the determinant of the first 2×2 submatrix, denoted as \mathbf{M}_{12}

$$\det \mathbf{M}_{12} = ((E - \mu) - \mathcal{F}_1 - \mathcal{F}_2)(v_{11}(1 - v_{11}G_1(E))) - (v_{11}^2 \mathcal{F}_1^g \mathcal{F}_1^{g\dagger}).$$

We consider $E_\mu < a_1$ first, where E_μ is the solution for $M_{11} = 0$, i.e. the discrete state μ is renormalized to E_μ by turning on f_1 and f_2 . Then, the simplest case is $v_{11} > 0$, when no bound state is developed by pure continuum-continuum interaction. Since at $E = E_\mu$ the last term $(v_{11}\mathcal{F}_1^g(E_\mu))^2$ is positive and $(v_{11}(1 - v_{11}G_1(E_\mu))) > 0$, $\mathcal{F}_{1,2}(E_\mu) < 0$, turning on v_{11} has the opposite effect as $\mathcal{F}_{1,2}$ to the state corresponding to E_μ . So, the state is moving upward from the previous solution E_μ towards the threshold. If $v_{11} < 0$, when $|v_{11}|$ is small enough, $(v_{11}(1 - v_{11}G_1(E_\mu))) < 0$, and the negativity of the last term will cause the discrete states corresponding to E_μ to move downward. For larger $|v_{11}|$ there could also be a bound state at E_b come down from the threshold a_1 , generated from the first channel continuum interaction when $f_1(\omega) = 0$, and it satisfies $(1 - v_{11}G_1(E_b)) = 0$, and $E_b < a_1$. Then we can also look at how switching on a small interaction $f_1(\omega)$ between the discrete state and the first continuum affects the bound state E_b . Since $E_\mu < E_b$ the factor $(E_b - \mu) - \mathcal{F}_1(E_b) - \mathcal{F}_2(E_b) > 0$, the negativity of the last term causes the bound state corresponding to E_b to move up towards the threshold. Thus in this case the last term seems to play a role of a repulsive interaction.

Next we look at the case $a_1 < \mu < a_2$, $v_{11} \neq 0$, $v_{12} = v_{22} = 0$ and $\mathcal{F}_2(\mu) < 0$, and $\text{Re}\mathcal{F}_1(\mu) < 0$ is supposed. Then, at $E = \mu$ we have

$$\text{Re det } \mathbf{M}_{12}(\mu) = (-\text{Re}\mathcal{F}_1(\mu) - \mathcal{F}_2(\mu))(v_{11}(1 - v_{11}\text{Re}G_1(\mu))) - v_{11}^2(\text{Re}\mathcal{F}_1^g(\mu))^2. \quad (56)$$

When $v_{11} > 0$ and $\text{Re}G_1(\mu) < 0$, the factor $\text{Re}(v_{11}(1 - v_{11}G_1(\mu))) > 0$, the negativity of the last term $-(\text{Re}\mathcal{F}_1^g(\mu))^2$ plays an opposite role than the first term which is positive. Thus, while turning on the discrete-continuum interaction $f_{1,2}$, which generating the first term, pushes the mass of the discrete state down, turning on a positive v_{11} tends to pull it up towards the threshold a_2 . This can be understood since it introduces a repulsive interaction in the continuum-continuum interaction. However, if $\text{Re}G_1(\mu) > 0$ such that $(1 - v_{11}\text{Re}G_1(\mu)) < 0$, which happens when Λ is much smaller than a_2 and $\mu > E_0$ where $G_1(E_0) = 0$, it could happen that a repulsive interaction $v_{11} > 0$ tends to push the discrete state mass down. When $v_{11} < 0$ and $|v_{11}|$ is small enough, such that $(v_{11}(1 - v_{11}\text{Re}G_1(\mu))) < 0$, the last term has the same sign as the first term, and thus similar to the first term, it pushes the mass of the discrete state down. If $\mu > a_2$, then both $\mathcal{F}_{1,2}$ have imaginary parts and there will be an additional term $-4\pi^2 v_{11}^2 |f_2(\mu)|^2 |g_1(\mu)|^2$ in Eq.(56) which is always negative, and thus it will have the similar effect as the last term $-v_{11}^2(\text{Re}\mathcal{F}_1^g(\mu))^2$.

Let us then discuss the effect of nonzero v_{12} and set $v_{11} = v_{22} = 0$. Now the \mathbf{M} matrix becomes

$$\mathbf{M} = \begin{pmatrix} (E - \mu) - \mathcal{F}_1(E) - \mathcal{F}_2(E) & -v_{21}\mathcal{F}_2^g(E) & -v_{12}\mathcal{F}_1^g(E) \\ -v_{21}^*\mathcal{F}_2^{g\dagger}(E) & -|v_{12}|^2 G_2(E) & v_{12} \\ -v_{12}^*\mathcal{F}_1^{g\dagger}(E) & v_{21} & -|v_{12}|^2 G_1(E) \end{pmatrix},$$

and

$$\begin{aligned} \det \mathbf{M} = & |v_{12}|^2 \left(-((E - \mu) - \mathcal{F}_1(E) - \mathcal{F}_2(E))(1 - |v_{12}|^2 G_1(E)G_2(E)) \right. \\ & + |v_{12}|^2 (\mathcal{F}_2^g(E)\mathcal{F}_2^{g\dagger}(E)G_1(E) + \mathcal{F}_1^g(E)\mathcal{F}_1^{g\dagger}(E)G_2(E)) \\ & \left. + v_{12}^*\mathcal{F}_1^{g\dagger}(E)\mathcal{F}_2^g(E) + v_{12}\mathcal{F}_1^g(E)\mathcal{F}_2^{g\dagger}(E) \right). \end{aligned} \quad (57)$$

This time v_{12} not only appears in $|v_{12}|^2$, but also has linear terms. There could be two kinds of bound states. The first kind of bound state could be originate from the bare discrete state at $E = \mu$, $\mu < a_1$. When $v_{12} = 0$, the bare state at $E = \mu$ is renormalized to $E_\mu < a_1$, which satisfies $(E_\mu - \mu) - \mathcal{F}_1(E_\mu) - \mathcal{F}_2(E_\mu) = 0$. There could also be another kind of bound state from the effective attractive interaction of v_{12} . When there is no discrete-continuum interaction, a bound state located at $E_b < a_1$ can be generated from the solution of $1 - |v_{12}|^2 G_1(E)G_2(E) = 0$, as already discussed previously. For $E < a_1$, the $|v_{12}|^2$ term on the second line of Eq. (57) is always negative and decreasing with E , and the $v_{12}^{(*)}$ linear terms becomes

$$2\text{Re}[v_{12}\mathcal{F}_1^{g\dagger}(E)\mathcal{F}_2^g(E)].$$

We first consider a special case that the last line are too small compared with the second line and can be ignored, for example, $|\mathcal{F}_2^g| \ll |\mathcal{F}_1^g|$ and $|\mathcal{F}_2^g| \ll |G_2|$. Then when $|v_{12}|^2$ is small enough, the factor $1 - |v_{12}|^2 G_1(E_\mu)G_2(E_\mu) > 0$ for $E < E_\mu$, and then the effect of the pure negative second line is to push the discrete state from E_μ down away from the threshold. When $|v_{12}|^2$ is large enough, such that the bound state E_b is generated and coming down from the threshold a_1 , since $E_\mu < E_b$, we still have $1 - |v_{12}|^2 G_1(E_\mu)G_2(E_\mu) > 0$. So, the discrete state generated from the bare state always goes away from the threshold. We also have $(E_b - \mu) - \mathcal{F}_1(E_b) - \mathcal{F}_2(E_b) > 0$, and the effect of the second line is to pull the bound state from E_b towards the threshold a_1 . So, the second line plays the role of an attractive interaction for the bound state generated from the bare discrete state, but as a repulsive interaction for

the bound state generated in the continuum-continuum interaction. If the last two terms on the last line can not be ignored, they will bring complexity to the discussion. If it is negative, it plays the same role as the second line, and if it is positive it will play the opposite role and competes with the second line. In fact, since it is of order v_{12} , it may have larger contribution than the second term for small v_{12} . If both the interaction vertices f_i and g_i are the real positive exponential ones like in the previous example, the sign of the last line is determined by $\text{Re}v_{12}$.

In more complicated cases, the results may be intricate, and may not have a simple picture. The previous cases serve as examples for the analysis of the effects of the different interaction in various situations and qualitative understanding the behavior of the pole positions.

VI. CONCLUSION

This paper presents several improvements to the Friedrichs model, aiming to provide a more comprehensive description of coupled channel scattering in real-world scenarios. Firstly, we investigate situations involving multiple discrete states and continuum states, focusing on the general interaction between these discrete states and the continuum states. Secondly, we consider the inclusion of contact interactions among the continuum states, employing a more general separable interaction that is independent of the interaction between the discrete states and the continuum states. Notably, this extended model remains exactly solvable. Thirdly, we address scenarios where the square integrable interaction between the continuum states takes a non-separable form, rendering it non-solvable. However, we propose an approach to approximate this potential by expanding it in terms of a chosen basis set, effectively expressing it as a truncated series of separable potentials. Consequently, at a finite order, this potential becomes solvable. To simplify the analysis, we also suggest utilizing the same basis set for expanding both the discrete-continuum interaction and the continuum-continuum potential. A few simple examples are discussed to analyse the behaviors of the masses of the discrete states when different interaction are turned on, which may be helpful in qualitatively understanding the spectrum in the coupled channel system.

This discussion establishes a theoretical foundation for the application of the Friedrichs model in various contexts, including hadron physics and other areas involving coupled channel scattering and intermediate resonances. To utilize the model effectively, one must first model the interaction between the discrete-continuum and continuum-continuum components. Subsequently, the continuum-continuum potential can be approximated using a series of separable potentials, enabling resonance searches or S -matrix calculations. An advantageous aspect of this model is the automatic preservation of unitarity in the S -matrix, while avoiding the presence of spurious poles on the first Riemann sheet. In contrast, the conventional K -matrix parameterization lacks control over spurious poles on the physical sheet.

However, a remaining challenge lies in determining the continuum-continuum interaction in a reasonable manner. Further research is required to develop suitable approaches for obtaining this interaction term in a manner that meets the physical expectations and provides reliable results.

ACKNOWLEDGMENTS

This work is supported by China National Natural Science Foundation under contract No. 11975075, No. 12375132, No. 12375078, No. 11575177, No.11947301, and No.12335002. This work is also supported by “the Fundamental Research Funds for the Central Universities”.

Appendix A: The detailed derivation of the normalization and the S matrix in section III

The normalization of the continuum state using the coefficients in (24) and (25) can be calculated as follows,

$$\begin{aligned}
\langle \Psi_m^\pm(E) | \Psi_n^\pm(E') \rangle &= \sum_{i=1}^D \alpha_{im}^*(E) \alpha_{in}(E') + \delta_{mn} \delta(E - E') \\
&+ \frac{1}{E - E' \mp i0} \left(\sum_{j=1}^D \frac{A_{jm}^*(E) f_{jn}^*(E')}{E - M_j} + \sum_{n'=1}^C v_{nn'}^* B_{n'm}^*(E) g_n^*(E') \right) \\
&+ \frac{1}{E' - E \pm i0} \left(\sum_{j=1}^D \frac{A_{jm}(E') f_{jn}(E)}{E' - M_j} + \sum_{n'=1}^C v_{nn'} B_{n'm}(E') g_n(E) \right) \\
&+ \sum_{m'=1}^C \int_{a_{m'}} d\omega \frac{1}{E - \omega \mp i0} \frac{1}{E' - \omega \pm i0} \left(\sum_{j=1}^D \frac{A_{jm}^*(E) f_{jm'}^*(\omega)}{E - M_j} + \sum_{n'=1}^C v_{m'n'}^* B_{n'm}^*(E) g_{m'}^*(\omega) \right) \\
&\quad \frac{1}{E' - E \pm i0} \left(\frac{1}{E - \omega \mp i0} - \frac{1}{E' - \omega \pm i0} \right) \\
&\times \left(\sum_{j'=1}^D \frac{A_{j'n}(E') f_{j'm'}(\omega)}{E' - M_{j'}} + \sum_{n''=1}^C v_{m'n''} B_{n''n}(E') g_{m'}(\omega) \right). \tag{A1}
\end{aligned}$$

Notice that we have omitted the \pm superscripts in the coefficients α_{im} , A_{jm} and B_{mn} since they all have the same superscript of \pm . Using the definitions in Eq. (29), Eqs. (31) and (32), the last two lines can be reduced as

$$\begin{aligned}
&\frac{1}{E' - E \pm i0} \left[\sum_{j,j'=1}^D \alpha_{jm}^{\pm*}(E) (\delta_{jj'}(E - E') - V_{jj'}^{\pm*}(E) + V_{jj'}^\pm(E')) \alpha_{j'n}^\pm(E') \right. \\
&\quad + \sum_{j=1}^D \sum_{n''=1}^C \alpha_{jm}^{\pm*}(E) \left(- \sum_{m'} V_{m'j}^{\pm BA^*}(E) v_{m'n''} + V_{jn''}^{\pm AB}(E') \right) B_{n''n}^\pm(E') \\
&\quad + \sum_{j'=1}^D \sum_{n'=1}^C B_{n'm}^{\pm*} \left(-V_{j'n'}^{\pm AB^*}(E) + \sum_{m'=1}^C v_{m'n'}^* V_{j'm'}^{\pm BA} \right) \alpha_{j'n}^\pm(E') \\
&\quad \left. + \sum_{n',n'',m'=1}^C B_{n'm}^{\pm*}(E) \left(-V_{m'n'}^{\pm B^*}(E) v_{m'n''} + v_{m'n'}^* V_{m'n''}^{\pm B}(E') \right) B_{n''n}^\pm(E') \right] \\
&= - \sum_{j=1}^D \alpha_{jm}^{\pm*}(E) \alpha_{jn}^\pm(E) + \frac{1}{E' - E \pm i0} \left[\sum_{j'=1}^D \left(-f_{j'm}(E) \right) \alpha_{j'n}^\pm(E') + \alpha_{j'm}^{\pm*}(E) f_{j'n}^*(E') \right. \\
&\quad \left. + \sum_{n',m'=1}^C \left(-\delta_{mm'} g_m(E) v_{m'n'} B_{n'n}^\pm(E') + B_{n'm}^{\pm*}(E) v_{m'n'}^* g_{m'}^*(E') \delta_{m'n} \right) \right].
\end{aligned}$$

Thus, they cancel with the terms in Eq. (A1) except for the $\delta(E - E')$ term. The S -matrix can also be obtained by

$$\begin{aligned}
\langle \Psi_m^-(E) | \Psi_n^+(E') \rangle &= \sum_{i=1}^D \sum_{i=1}^D \alpha_{im}^{-*}(E) \alpha_{in}^+(E') + \delta_{mn} \delta(E - E') \\
&+ \left(\frac{1}{E - E' - i0} - 2\pi i \delta(E - E') \right) \left(\sum_{j=1}^D \frac{A_{jm}^{-*}(E) f_{jn}^*(E')}{E - M_j} + \sum_{n'=1}^C v_{nn'}^* B_{n'm}^{-*}(E) g_n^*(E') \right) \\
&+ \frac{1}{E' - E + i0} \left(\sum_{j=1}^D \frac{A_{jm}^+(E') f_{jn}(E)}{E' - M_j} + \sum_{n'=1}^C v_{nn'} B_{n'm}^+(E') g_n(E) \right) \\
&+ \sum_{m'=1}^C \int_{a_{m'}} d\omega \frac{1}{E - \omega + i0} \frac{1}{E' - \omega + i0} \left(\sum_{j=1}^D \frac{A_{jm}^{-*}(E) f_{jm'}^*(\omega)}{E - M_j} + \sum_{n'=1}^C v_{m'n'}^* B_{n'm}^{-*}(E) g_{m'}^*(\omega) \right) \\
&\quad \frac{1}{E' - E + i0} \left(\frac{1}{E - \omega + i0} - \frac{1}{E' - \omega + i0} \right) \\
&\times \left(\sum_{j'=1}^D \frac{A_{j'n}^+(E') f_{j'm'}(\omega)}{E' - M_{j'}} + \sum_{n''=1}^C v_{m'n''} B_{n''n}^+(E') g_{m'}(\omega) \right). \tag{A2}
\end{aligned}$$

We have used $\frac{1}{E - \omega + i0} \frac{1}{E' - \omega + i0} = \frac{1}{E' - E + i0} \left(\frac{1}{E - \omega - i0} - \frac{1}{E' - \omega + i0} \right) - 2\pi i \delta(E - \omega) \frac{1}{E' - \omega + i0} = \frac{1}{E' - E + i0} \left(\frac{1}{E - \omega + i0} - \frac{1}{E' - \omega + i0} \right)$. Since $\delta(E' - E) \left(\frac{1}{E - \omega + i0} - \frac{1}{E' - \omega + i0} \right) = 0$, the $i0$ in the first factor does not have any effect. The last two lines can be reduced to

$$\begin{aligned}
&\frac{1}{E' - E + i0} \left[\sum_{j,j'=1}^D \alpha_{jm}^{-*}(E) (\delta_{jj'}(E - E') - V_{j'j}^{-*}(E) + V_{jj'}^+(E')) \alpha_{jn}^+(E') \right. \\
&\quad + \sum_{j=1}^D \sum_{n'=1}^C \alpha_{jm}^{-*}(E) \left(- \sum_{m'} V_{m'j}^{-BA*}(E) v_{m'n''} + V_{jn''}^{+AB}(E') \right) B_{n'n}^+(E') \\
&\quad + \sum_{j'=1}^D \sum_{n'=1}^C B_{n'm}^{-*} \left(-V_{j'n'}^{-AB*}(E) + \sum_{m'=1}^C v_{m'n'}^* V_{j'm'}^{+BA} \right) \alpha_{j'n}^+(E') \\
&\quad \left. + \sum_{n',n'',m'=1}^C B_{n'm}^{-*}(E) \left(-V_{m'n'}^{-B*}(E) v_{m'n''} + v_{m'n'}^* V_{m'n''}^{+B}(E') \right) B_{n''n}^+(E') \right] \\
&= - \sum_{j=1}^D \alpha_{jm}^{-*}(E) \alpha_{jn}^+(E) + \frac{1}{E' - E + i0} \left[\sum_{j'=1}^D \left(-f_{j'm}(E) \right) \alpha_{j'n}^+(E') + \alpha_{j'm}^{-*}(E) f_{j'n}^*(E') \right. \\
&\quad \left. + \sum_{n',m'=1}^C \left(-\delta_{mm'} g_m(E) v_{m'n'} B_{n'n}^+(E') + B_{n'm}^{-*}(E) v_{m'n'}^* g_{m'}^*(E') \delta_{m'n} \right) \right].
\end{aligned}$$

These terms cancel with the other terms except the terms with $\delta(E - E')$. Notice that for $E = E'$ and is real, the final sum inside the square bracket will be zero,

$$\begin{aligned}
&\left[\sum_{j'=1}^D \left(-f_{j'm}(E) \alpha_{j'n}^+(E) + \alpha_{j'm}^{-*}(E) f_{j'n}^*(E) \right) \right. \\
&\quad \left. + \sum_{n',m'=1}^C \left(-\delta_{mm'} g_m(E) v_{m'n'} B_{n'n}^+(E) + B_{n'm}^{-*}(E) v_{m'n'}^* g_{m'}^*(E) \delta_{m'n} \right) \right] = 0,
\end{aligned}$$

which can be derived directly from Eqs. (31) and (32). Then the S matrix can be derived:

$$\begin{aligned}
S_{mn}(E, E') &= \delta(E - E') - 2\pi i \delta(E - E') \left(\sum_{j=1}^D \frac{A_{jm}^{-*}(E) f_{jn}^*(E')}{E - M_j} + \sum_{n'=1}^C v_{nn'}^* B_{n'm}^{-*}(E) g_n^*(E') \right) \\
&= \delta(E - E') - \pi i \delta(E - E') \left(\sum_{j=1}^D (\alpha_{jm}^{-*}(E) f_{jn}^*(E') + f_{j'm}(E) \alpha_{j'n}^{\pm}(E')) \right. \\
&\quad \left. + \sum_{n'=1}^C (v_{nn'}^* B_{n'm}^{-*}(E) g_n^*(E') + v_{mn'} B_{n'n}^+(E) g_m(E')) \right).
\end{aligned}$$

From the definition of \mathbf{Y} and \mathbf{F} in Eq. (28) and solving \mathbf{Y} from Eq.(30), we can reformulate the previous equation using the matrices and obtain Eq. (35).

Appendix B: Solving the approximate contact potential

This section we provide the details for solving the eigenstate problem for the Hamiltonian in section IV. After approximating the general contact potential as the sum of the separable potentials, the approximated Hamiltonian for multiple continuum states and discrete states is shown in Eq.(44) which is copied here for completeness

$$\begin{aligned}
H &= \sum_{i=1}^D M_i |i\rangle \langle i| + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \omega |\omega; n\rangle \langle \omega; n| \\
&\quad + \sum_{m,n=1}^C v_{mn,\rho\delta} \left(\int_{a_m}^{\infty} d\omega' \tilde{g}_{m\rho}(\omega') |\omega'; m\rangle \right) \left(\int_{a_n}^{\infty} d\omega \tilde{g}_{n\delta}^*(\omega) \langle \omega; n| \right) \\
&\quad + \sum_{j=1}^D \sum_{n=1}^C \left[|j\rangle \left(\int_{a_n}^{\infty} d\omega f_{jn}^*(\omega) \langle \omega; n| \right) + \left(\int_{a_n}^{\infty} d\omega f_{jn}(\omega) |\omega; n\rangle \right) \langle j| \right]. \tag{B1}
\end{aligned}$$

The general eigenstate for this eigenvalue problem can be expanded using the bare discrete states and the bare continuum states

$$|\Psi(E)\rangle = \sum_{i=1}^D \alpha_i(E) |i\rangle + \sum_{n=1}^C \int_{a_n}^{\infty} d\omega \psi_n(E, \omega) |\omega; n\rangle.$$

The proceeding derivation goes in parallel with the process in section III. With this ansatz, the eigenvalue problem can be reduced to the following equations

$$\begin{aligned}
(M_j - E) \alpha_j(E) + A_j(E) &= 0, \quad j = 1, \dots, D \\
\sum_{j=1}^D \alpha_j(E) f_{jn}(\omega) + (\omega - E) \psi_n(E, \omega) + \sum_{m=1}^C v_{nm,\rho\delta} \psi_{m\delta}(E) \tilde{g}_{n\rho}(\omega) &= 0, \quad n = 1, \dots, C, \quad \text{and } \omega > a_n
\end{aligned}$$

where we have defined

$$A_j(E) = \sum_{n=1}^C \int_{a_n}^{\infty} d\omega f_{jn}^*(\omega) \psi_n(E, \omega), \quad \psi_{n\delta}(E) = \int d\omega \psi_n(E, \omega) \tilde{g}_{n\delta}^*(\omega). \tag{B2}$$

There are C continuum eigenstate solutions and $|\Psi(E)\rangle$, α_i , $\psi_{n\delta}$ and $A_j(E)$ need another index m to denote different continuum solutions, i.e. $|\Psi_m(E)\rangle$, $\alpha_{im}(E)$, $\psi_{nm\delta}(E)$, and $A_{jm}(E)$. Similar to section III, we require that $|\Psi_m(E)\rangle$ tends to $|E, m\rangle$ as the interactions are turned off and consider the C continuum solutions for $E > a_C$. Then the above equations can be reduced to

$$\alpha_{jm}^{\pm}(E) = \frac{1}{E - M_j} A_{jm}^{\pm}(E) \tag{B3}$$

$$\psi_{nm}^{\pm}(E, \omega) = \delta_{nm} \delta(E - \omega) + \sum_{j=1}^D \frac{\alpha_{jm}^{\pm}(E) f_{jn}(\omega)}{E - \omega \pm i0} + \sum_{n'=1}^C v_{nn',\delta'\rho} \frac{\psi_{n'm\rho}^{\pm}(E) \tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0}. \tag{B4}$$

By applying the operation $\sum_n \int_{a_n} d\omega f_{jn}^*(\omega) \times$ and the operations $\sum_{n\delta} v_{nm',\rho'\delta}^* \int_{a_n} d\omega \tilde{g}_{n\delta}^*(\omega) \times$ on (B4) respectively, we obtain

$$0 = -f_{jm}^*(E) + \sum_{j'=1}^D \left((E - M_{j'}) \delta_{jj'} - \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{j'n}(\omega) f_{jn}^*(\omega)}{E - \omega \pm i0} \right) \frac{A_{j'm}^\pm(E)}{E - M_{j'}} - \sum_{n',n=1}^C v_{nn',\delta'\rho} \psi_{n'm\rho}^\pm(E) \int_{a_n} d\omega \frac{f_{jn}^*(\omega) \tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0} \quad (\text{B5})$$

$$0 = -v_{mm',\delta\rho}^* \tilde{g}_{m\delta}^*(E) - \sum_{j=1}^D \alpha_{jm}^\pm(E) \sum_{n=1}^C v_{nm',\delta\rho}^* \int_{a_n} d\omega \frac{f_{jn}(\omega) \tilde{g}_{n\delta}^*(\omega)}{E - \omega \pm i0} \\ + \sum_{n'}^C \psi_{n'm\rho}^\pm(E) \left[v_{n'm',\rho\rho'}^* - \sum_{n=1}^C v_{nm',\delta\rho}^* v_{nn',\delta'\rho} \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega) \tilde{g}_{n\delta}^*(\omega)}{E - \omega \pm i0} \right]. \quad (\text{B6})$$

These two equations corresponds to previous Eq. (26) and (27). Notice that the differences are Greek letters and the sums here. Similar to the vectors and matrices defined in Eqs.(28)-(29), we define

$$(\tilde{\mathbf{F}}_m)_{m'\rho'} = v_{mm',\delta\rho'}^* \tilde{g}_{m\delta}^*(E), \quad (\tilde{\mathbf{F}}_m)_j = f_{jm}^*(E), \quad m = 1, \dots, C; j = 1, \dots, D; \delta = 1, 2, \dots, N \\ (\tilde{\mathbf{Y}}_m(E))_j = \alpha_{jm}(E), \quad (\tilde{\mathbf{Y}}_m)_{n'\rho} = \psi_{n'm\rho}^\pm(E), \quad j = 1, \dots, D; m, n' = 1, \dots, C; \rho = 1, 2, \dots, N \quad (\text{B7})$$

$$\tilde{\mathbf{V}}_{m'\rho',n'\rho}^{BB} = v_{n'm',\rho\rho'}^* - \sum_{n=1}^C v_{nm',\delta\rho}^* v_{nn',\delta'\rho} \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega) \tilde{g}_{n\delta}^*(\omega)}{E - \omega \pm i0}, \quad m', n' = 1, 2, \dots, C; \rho', \rho = 1, 2, \dots, N \\ \tilde{\mathbf{V}}_{jj'}^{AA} = (E - M_{j'}) \delta_{jj'} - \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{j'n}(\omega) f_{jn}^*(\omega)}{E - \omega \pm i0}, \quad j, j' = 1, 2, \dots, D \\ \tilde{\mathbf{V}}_{j,n'\rho}^{AB} = - \sum_{n=1}^C v_{nn',\delta'\rho} \int_{a_n} d\omega \frac{f_{jn}^*(\omega) \tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0}, \quad j = 1, \dots, D; n' = 1, \dots, C; \rho = 1, 2, \dots, N \\ \tilde{\mathbf{V}}_{m'\rho',j}^{BA} = - \sum_{n=1}^C v_{nm',\delta\rho}^* \int_{a_n} d\omega \frac{f_{jn}(\omega) \tilde{g}_{n\delta}^*(\omega)}{E - \omega \pm i0}, \quad j = 1, \dots, D; m' = 1, \dots, C; \rho' = 1, 2, \dots, N \\ \tilde{\mathbf{M}}_{IJ} = \begin{pmatrix} \tilde{\mathbf{V}}_{ij}^{AA}(E) & \tilde{\mathbf{V}}_{i,n'\rho}^{AB}(E) \\ \tilde{\mathbf{V}}_{m'\rho',j}^{BA}(E) & \tilde{\mathbf{V}}_{m'\rho',n'\rho}^{BB}(E) \end{pmatrix}_{IJ}. \quad IJ = 1, \dots, D, D+1, \dots, D+NC \quad (\text{B8})$$

We still have $\tilde{\mathbf{M}}^{+\dagger} = \tilde{\mathbf{M}}^-$. With these matrices, Eqs. (B5) and (B6) can be expressed as

$$\tilde{\mathbf{M}} \cdot \tilde{\mathbf{Y}}_m = \tilde{\mathbf{F}}_m$$

or

$$\sum_{j=1}^D \tilde{\mathbf{V}}_{ij}^{AA}(E) \alpha_{jm}(E) + \sum_{n'=1}^C \tilde{\mathbf{V}}_{i,n'\rho}^{AB}(E) \psi_{n'm\rho}(E) = f_{im}^*(E), \quad (\text{B9})$$

$$\sum_{j=1}^D \tilde{\mathbf{V}}_{n\delta,j}^{BA}(E) \alpha_{jm}(E) + \sum_{n'=1}^C \tilde{\mathbf{V}}_{n\delta,n'\rho}^{BB}(E) \psi_{n'm\rho}(E) = v_{nn',\delta\rho}^*(E) \tilde{g}_{m\rho}^*(E). \quad (\text{B10})$$

As before $\tilde{\mathbf{M}}$ is still independent of m , but $\tilde{\mathbf{F}}_m$ depends on m . If there are infinite number of bases, the matrix $\tilde{\mathbf{M}}$ and vector $\tilde{\mathbf{F}}_m$ and $\tilde{\mathbf{Y}}$ are infinite dimensional. Now we have supposed that the bases chosen are well enough, and have made a truncation to a finite order N of the expansion of the potential $V_{nn'}$ i.e. $v_{nn',\delta\rho} = 0$ for $\delta, \rho > N$. Then $\tilde{\mathbf{M}}$ is a $(D + NC) \times (D + NC)$ matrix. In general, the matrix $\tilde{\mathbf{M}}$ is non-degenerate for $E > a_m$, and $\tilde{\mathbf{Y}}_m$ can be solved,

$$\tilde{\mathbf{Y}}_m(E) = \tilde{\mathbf{M}}^{-1} \cdot \tilde{\mathbf{F}}_m. \quad (\text{B11})$$

With all the $\psi_{nm\rho}^\pm(E)$ and $\alpha_{jm}^\pm(E)$ at hand, the approximate continuum solutions are solved as

$$\begin{aligned} |\Psi_m^\pm(E)\rangle &= \sum_{i=1}^D \alpha_{im}^\pm(E) |i\rangle + |E; m\rangle + \sum_{n=1}^C \int_{a_n} d\omega \frac{1}{E - \omega \pm i0} \left(\sum_{j=1}^D \alpha_{jm}^\pm(E) f_{jn}(\omega) + \sum_{n'=1}^C v_{nn',\delta'\rho} \psi_{n'm\rho}^\pm(E) \tilde{g}_{n\delta'}(\omega) \right) |\omega; n\rangle \\ &= |E; m\rangle + \sum_{j=1}^D \alpha_{jm}^\pm(E) \left(|j\rangle + \sum_{n=1}^C \int_{a_n} d\omega \frac{f_{jn}(\omega)}{E - \omega \pm i0} |\omega; n\rangle \right) + \left(\sum_{n,n'=1}^C v_{nn',\delta'\rho} \psi_{n'm\rho}^\pm(E) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0} |\omega; n\rangle \right). \end{aligned} \quad (\text{B12})$$

It can be checked that the normalization is $\langle \Psi_m^\pm(E) | \Psi_n^\pm(E') \rangle = \delta_{mn} \delta(E - E')$. The S -matrix can be obtained,

$$\langle \Psi_m^-(E) | \Psi_n^+(E') \rangle = \delta_{mn} \delta(E - E') - 2\pi i \delta(E - E') \left(\tilde{\mathbf{F}}_m^\dagger \cdot (\tilde{\mathbf{M}}^+)^{-1} \cdot \tilde{\mathbf{F}}_m \right). \quad (\text{B13})$$

For discrete eigenvalues and discrete eigenstates, there will not be the delta function in Eq. (B4), and we have equations,

$$\tilde{\mathbf{M}} \cdot \tilde{\mathbf{Y}} = 0,$$

where $\tilde{\mathbf{Y}}$ is defined similar to Eq. (B7) without subindex m . The generalized energy eigenvalues for the discrete state can be obtained from the $\det \tilde{\mathbf{M}}(E) = 0$ and the eigenvector $\tilde{\mathbf{Y}}$ can be obtained with proper normalization chosen as in previous section. Then the i -th discrete state can be expressed as

$$\begin{aligned} |\Psi^{(i)}(E_i)\rangle &= \sum_{i=1}^D \alpha_j^{(i)}(E_i) \left(|j\rangle + \int_{a_n} d\omega \frac{f_{jn}(\omega)}{E_i - \omega} |\omega; n\rangle \right) + \sum_{n'=1}^C v_{nn',\delta'\rho} \psi_{n'm\rho}^{(i)}(E_i) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E_i - \omega} |\omega; n\rangle, \\ &\text{with } \tilde{\mathbf{Y}}^{(i)}(E_i) \cdot \tilde{\mathbf{M}}'(E_i) \cdot \tilde{\mathbf{Y}}^{(i)*}(E_i) = 1 \end{aligned} \quad (\text{B14})$$

where $\tilde{\mathbf{M}}'(E)$ is the derivative of the matrix w.r.t. E and the integral contour needs to be deformed for E_i on unphysical sheets as before.

Appendix C: Solving the eigenvalue problem with both approximate vertex functions and contact interactions

This section serves to solve the eigenstates for the Hamiltonian in Eq.(49) where both the contact potential and the vertex are expanded using the function bases $\tilde{g}_{n\delta}$, which we reproduce here for completeness

$$\begin{aligned} H &= \sum_{i=1}^D M_i |i\rangle \langle i| + \sum_{n=1}^C \int_{a_n}^\infty d\omega \omega |\omega; n\rangle \langle \omega; n| \\ &+ \sum_{m,n=1}^C v_{mn,\rho\delta} \left(\int_{a_m}^\infty d\omega' \tilde{g}_{m\rho}(\omega') |\omega'; m\rangle \right) \left(\int_{a_n}^\infty d\omega \tilde{g}_{n\delta}^*(\omega) \langle \omega; n| \right) \\ &+ \sum_{j=1}^D \sum_{n=1}^C \left[f_{jn\delta}^* |j\rangle \left(\int_{a_n}^\infty d\omega \tilde{g}_{n\delta}^*(\omega) \langle \omega; n| \right) + f_{jn\delta} \left(\int_{a_n}^\infty d\omega \tilde{g}_{n\delta}(\omega) |\omega; n\rangle \right) \langle j| \right]. \end{aligned} \quad (\text{C1})$$

This case is more like the cases discussed in [10, 26], where the same form factor comes with the continuum both in the discrete-continuum and continuum-continuum interaction. Using the eigenstate ansatz

$$\begin{aligned} |\Psi(E)\rangle &= \sum_{i=1}^D \alpha_i(E) |i\rangle + \int_{a_n} d\omega \psi_n(E, \omega) |\omega; n\rangle \\ &= \sum_{i=1}^D \alpha_i(E) |i\rangle + \sum_{n=1}^C \psi_{n\delta}(E) \int_{a_n} d\omega \tilde{g}_{n\delta}(\omega) |\omega; n\rangle, \end{aligned} \quad (\text{C2})$$

and proceeding in solving the eigenvalue problem similarly to the previous section, one finds that A_j in Eq. (B2) becomes

$$A_j(E) = \sum_{n=1}^C f_{jn\delta}^* \psi_{n\delta}(E).$$

For the m -th continuum solution $|\Psi_m(E)\rangle$, the coefficients α_i , ψ_n and $\psi_{n\delta}$ take another subindex m and Eq. (B3) and (B4) becomes

$$\alpha_{jm}^\pm(E) = \frac{1}{E - M_j} A_{jm}^\pm(E) = \frac{1}{E - M_j} \sum_{n'=1}^C f_{jn'\rho}^* \psi_{n'm\rho}^\pm(E), \quad (\text{C3})$$

$$\psi_{nm}^\pm(E, \omega) = \delta_{nm} \delta(E - \omega) + \sum_{n'=1}^C \psi_{n'm\rho}^\pm(E) V_{n\delta', n'\rho}(E) \frac{\tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0}, \quad (\text{C4})$$

where $V_{n\delta', n'\rho}(E)$ is defined as the matrix elements of a $NC \times NC$ matrix \mathbf{V}

$$(\mathbf{V}(E))_{n\delta', n'\rho} = \sum_{j=1}^D \frac{f_{jn\delta'} f_{jn'\rho}^*}{E - M_j} + v_{nn', \delta'\rho},$$

which is supposed to be non-degenerate for general E . Multiplying $\tilde{g}_{n\delta}^*(\omega)$ to above equation (C4) and integrating w.r.t ω , we get

$$\delta_{nm} \tilde{g}_{n\delta}^*(E) = \sum_{n'=1}^C \psi_{n'm\rho}^\pm(E) \left[\delta_{nn'} \delta_{\rho\delta} - V_{n\delta', n'\rho}(E) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega) \tilde{g}_{n\delta}^*(\omega)}{E - \omega \pm i0} \right]. \quad (\text{C5})$$

Define $NC \times NC$ matrix $\tilde{\mathbf{M}}$ and NC dimensional vector $\tilde{\mathbf{F}}_m, \tilde{\mathbf{Y}}_m$ as

$$\tilde{\mathbf{M}}_{n\delta, n'\rho} = \delta_{nn'} \delta_{\rho\delta} - V_{n\delta', n'\rho}(E) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega) \tilde{g}_{n\delta}^*(\omega)}{E - \omega \pm i0}, \quad (\text{C6})$$

$$(\tilde{\mathbf{F}}_m)_{n\delta} = \delta_{nm} \tilde{g}_{n\delta}^*(E), \quad (\tilde{\mathbf{Y}}_m)_{n'\rho} = \psi_{n'm\rho}^\pm(E). \quad (\text{C7})$$

Then Eq.(C5) can be expressed as $\tilde{\mathbf{F}}_m = \tilde{\mathbf{M}} \cdot \tilde{\mathbf{Y}}_m$. Notice that $\tilde{\mathbf{M}}$ is independent of the m -th solution, but $\tilde{\mathbf{F}}_m$ depends on m . To see the real analyticity of $\det \tilde{\mathbf{M}}$, we define

$$(\tilde{\mathbf{W}}^\pm)_{n\delta, n'\rho} = (\tilde{\mathbf{M}}^\pm \cdot \mathbf{V}^{-1})_{n\delta, n'\rho} = \mathbf{V}_{n\delta, n'\rho}^{-1} - \delta_{nn'} \int_{a_n} d\omega \frac{\tilde{g}_{n\rho}(\omega) \tilde{g}_{n\delta}^*(\omega)}{E - \omega \pm i0}, \quad (\text{C8})$$

and then we have $\tilde{\mathbf{M}}^\pm = \tilde{\mathbf{W}}^\pm \cdot \mathbf{V}$. Since $(\tilde{\mathbf{W}}^\pm(E))^\dagger = \tilde{\mathbf{W}}^\mp(E)$ and $\mathbf{V}(E)$ is hermitian for real E , $\det \tilde{\mathbf{M}}^{\pm\dagger}(E) = \det \tilde{\mathbf{M}}^\mp(E)$. So the analytically continued $\det \tilde{\mathbf{M}}(E)$ with $\det \tilde{\mathbf{M}}^+(E)$ and $\det \tilde{\mathbf{M}}^-(E)$ on the upper and lower edge of the cut above the threshold satisfies the Schwartz reflection relation, $\det \tilde{\mathbf{M}}^+(E) = \det \tilde{\mathbf{M}}^*(E^*)$.

Then, in general, the matrix $\tilde{\mathbf{M}}$ is non-degenerate for $E > a_m$, and $\tilde{\mathbf{Y}}_m$ can be solved,

$$\psi_{nm\rho}^\pm(E) = (\tilde{\mathbf{Y}}_m(E))_{n\rho} = (\tilde{\mathbf{M}}^{-1} \cdot \tilde{\mathbf{F}}_m)_{n\rho} = (\tilde{\mathbf{M}}^{-1})_{n\rho, m\delta} \tilde{g}_{m\delta}(E). \quad (\text{C9})$$

With all the $\psi_{nm\rho}^\pm(E)$ for $\rho \leq N$ at hand, all the $\alpha_{jm}^\pm(E)$ and $\psi_{nm}^\pm(E, \omega)$ can also be obtained. Then, the continuum state can be approximated as

$$|\Psi_m^\pm(E)\rangle = \sum_{i=1}^D \alpha_{im}^\pm(E) |i\rangle + |E, m\rangle + \sum_{n=1}^C \sum_{n'=1}^C \psi_{n'm\rho}^\pm(E) V_{n\delta', n'\rho}(E) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E - \omega \pm i0} |\omega; n\rangle. \quad (\text{C10})$$

It can be checked that the normalization is $\langle \Psi_m^\pm(E) | \Psi_n^\pm(E') \rangle = \delta_{mn} \delta(E - E')$. The S -matrix can be obtained

$$\begin{aligned} \langle \Psi_m^-(E) | \Psi_n^+(E') \rangle &= \delta_{mn} \delta(E - E') - 2\pi i \delta(E - E') \sum_{n=1}^C \sum_{n'=1}^C \psi_{n'm\rho}^{-*}(E') V_{n\delta', n'\rho}^*(E') \tilde{g}_{n\delta'}^*(E) \\ &= \delta_{mn} \delta(E - E') - 2\pi i \delta(E - E') (\tilde{\mathbf{F}}_m^\dagger \cdot (\tilde{\mathbf{W}}^+)^{-1} \cdot \tilde{\mathbf{F}}_n). \end{aligned} \quad (\text{C11})$$

Similar to previous section, the generalized energy eigenvalues for the discrete state can be obtained from the $\det \tilde{\mathbf{M}}(E) = 0$, and for each eigenvalue E_i , $\psi_{n\rho}^{(i)}$ can be solved from $\tilde{\mathbf{M}} \cdot \tilde{\mathbf{Y}} = 0$. Then we have the discrete eigenstates,

$$|\Psi^{(i)}(E_i)\rangle = \sum_{n'=1}^C \psi_{n'\rho}^{(i)}(E_i) \left[\sum_{j=1}^D \frac{f_{jn'\rho}^*}{E_i - M_j} |j\rangle + \sum_{n=1}^C V_{n\delta', n'\rho}(E_i) \int_{a_n} d\omega \frac{\tilde{g}_{n\delta'}(\omega)}{E_i - \omega} |\omega; n\rangle \right], \quad (\text{C12})$$

$$\text{with } \tilde{\mathbf{Y}}^{(i)\dagger}(E_i) \cdot \mathbf{V}(E_i) \cdot \tilde{\mathbf{W}}'(E_i) \cdot \mathbf{V}(E_i) \cdot \tilde{\mathbf{Y}}^{(i)}(E_i) = 1$$

with integral contour deformed for resonances and virtual states as before.

-
- [1] K. O. Friedrichs, *Commun. Pure Appl. Math.* **1**, 361 (1948).
 - [2] T. D. Lee, *Phys. Rev.* **95**, 1329 (1954).
 - [3] E. T. Jaynes and F. W. Cummings, *Proc. IEEE* **51**, 89 (1963).
 - [4] P. Facchi, H. Nakazato, and S. Pascazio, *Phys. Rev. Lett.* **86**, 2699 (2001).
 - [5] Z. Xiao and Z.-Y. Zhou, *Phys. Rev. D* **94**, 076006 (2016), arXiv:1608.00468 [hep-ph].
 - [6] A. Bohm and M. Gadella, *Dirac Kets, Gamow Vectors and Gel'fand Triplets*, edited by A. Bohm and J. D. Dollard, *Lecture Notes in Physics*, Vol. 348 (Springer Berlin Heidelberg, 1989).
 - [7] O. Civitarese and M. Gadella, *Phys. Rep.* **396**, 41 (2004).
 - [8] T. Petrosky, I. Prigogine, and S. Tasaki, *Physica* **173A**, 175 (1991).
 - [9] Z. Xiao and Z.-Y. Zhou, *J. Math. Phys.* **58**, 062110 (2017), arXiv:1608.06833 [hep-ph].
 - [10] Z. Xiao and Z.-Y. Zhou, *J. Math. Phys.* **58**, 072102 (2017), arXiv:1610.07460 [hep-ph].
 - [11] Z.-Y. Zhou and Z. Xiao, *Phys. Rev. D* **96**, 054031 (2017), [Erratum: *Phys. Rev. D* **96**, 099905 (2017)], arXiv:1704.04438 [hep-ph].
 - [12] Z.-Y. Zhou and Z. Xiao, *Phys. Rev. D* **97**, 034011 (2018), arXiv:1711.01930 [hep-ph].
 - [13] Z.-Y. Zhou and Z. Xiao, *Eur. Phys. J. C* **81**, 551 (2021), arXiv:2008.08002 [hep-ph].
 - [14] Z.-Y. Zhou and Z. Xiao, *Eur. Phys. J. C* **80**, 1191 (2020), arXiv:2008.02684 [hep-ph].
 - [15] E. Eichten, K. Gottfried, T. Kinoshita, K. Lane, and T.-M. Yan, *Phys. Rev. D* **21**, 203 (1980).
 - [16] N. A. Tornqvist, *Z. Phys. C* **68**, 647 (1995), arXiv:hep-ph/9504372.
 - [17] Yu. S. Kalashnikova, *Phys. Rev. D* **72**, 034010 (2005), arXiv:hep-ph/0506270 [hep-ph].
 - [18] B.-Q. Li, C. Meng, and K.-T. Chao, *Phys. Rev. D* **80**, 014012 (2009), arXiv:0904.4068 [hep-ph].
 - [19] P. G. Ortega, J. Segovia, D. R. Entem, and F. Fernandez, *Phys. Rev. D* **81**, 054023 (2010), arXiv:0907.3997 [hep-ph].
 - [20] X. Cao and H. Lenske, (2014), arXiv:1410.1375 [nucl-th].
 - [21] F. Giacosa, *Acta Phys. Polon. Supp.* **13**, 83 (2020), arXiv:1904.10368 [hep-ph].
 - [22] T. Wolkanowski, M. Sołtysiak, and F. Giacosa, *Nucl. Phys. B* **909**, 418 (2016), arXiv:1512.01071 [hep-ph].
 - [23] T. Wolkanowski, F. Giacosa, and D. H. Rischke, *Phys. Rev. D* **93**, 014002 (2016), arXiv:1508.00372 [hep-ph].
 - [24] B. Wang, Z.-W. Liu, and X. Liu, *Phys. Rev. D* **99**, 036007 (2019), arXiv:1812.04457 [hep-ph].
 - [25] Z. Yang, G.-J. Wang, J.-J. Wu, M. Oka, and S.-L. Zhu, *Phys. Rev. Lett.* **128**, 112001 (2022), arXiv:2107.04860 [hep-ph].
 - [26] T. Sekihara, T. Hyodo, and D. Jido, *PTEP* **2015**, 063D04 (2015), arXiv:1411.2308 [hep-ph].
 - [27] L. Micu, *Nucl. Phys. B* **10**, 521 (1969).
 - [28] Z.-Y. Zhou, C.-Y. Li, and Z. Xiao, (2023), arXiv:2304.07052 [hep-ph].
 - [29] A. Sitenko, *Lecture in Scattering Theory* (Pergamon Press, 1971).