

Coarse correlated equilibria in linear quadratic mean field games and application to an emission abatement game

Luciano Campi^a

Federico Cannerozzi^b

Fanny Cartellier^c

November 8, 2023

Abstract

Coarse correlated equilibria (CCE) are a good alternative to Nash equilibria (NE), as they arise more naturally as outcomes of learning algorithms and they may exhibit higher payoffs than NE. CCEs include a device which allows players' strategies to be correlated without any cooperation, only through information sent by a mediator. We develop a methodology to concretely compute mean field CCEs in a linear-quadratic mean field game framework. We compare their performance to mean field control solutions and mean field NE (usually named MFG solutions). Our approach is implemented in the mean field version of an emission abatement game between greenhouse gas emitters. In particular, we exhibit a simple and tractable class of mean field CCEs which allows to outperform very significantly the mean field NE payoff and abatement levels, bridging the gap between the mean field NE and the social optimum obtained by mean field control.

Keywords: Mean field games, coarse correlated equilibrium, mean field Nash equilibrium, mean field control, emissions' abatement game.

2020 AMS subject classification: 91A16, 49N80, 49N10, 91B76.

1 Introduction

Mean field games (MFGs) have been introduced in mid 2000s in [23] and independently in [20]. They arise as limit systems of large dynamic symmetric games with interactions of mean field type. In the limit, the concept of Nash equilibrium translates into a fixed point problem in the space of flows of measures. This equilibrium concept is commonly defined as an MFG solution, for two main reasons. On the one hand, approximate Nash equilibria with vanishing approximation error can be constructed starting from such an MFG solution (see, e.g., [8, 10, 22]). On the other hand, Nash equilibria (NEs) for the N -player game can be shown to converge to such MFG solutions (see, e.g., [21, 22]). In this sense, MFG solutions can be considered as the infinitely many players analogue of Nash equilibria, so that one can (and we will) refer to commonly called MFG solutions also as mean field Nash equilibria (mean field NE, for short).

Despite their popularity, Nash equilibria present some flaws. First, they raise numerical complexity issues, see for instance [15]. Second, it is well-known in game theory that agents are proved

^aDepartment of Mathematics "Federigo Enriques", University of Milan, Via Saldini 50, 20133, Milan, Italy. E-mail address: luciano.campi@unimi.it

^bDepartment of Mathematics "Federigo Enriques", University of Milan, Via Saldini 50, 20133, Milan, Italy. E-mail address: federico.cannerozzi@unimi.it

^cCentre of Research in Economics and Statistics (CREST), CNRS, École polytechnique, GENES, ENSAE Paris, Institut Polytechnique de Paris, 91120 Palaiseau, France. E-mail address: fanny.cartellier@ensae.fr.

to actually behave according to a Nash equilibrium only under strong rationality assumptions. Finally, they can be highly inefficient compared to social optimum. As an alternative to Nash equilibria, correlated equilibria (CEs) and coarse correlated equilibria (CCEs) have been introduced in game theory literature. They can be understood as a generalization of the notion of Nash equilibrium by the introduction of a correlation device, which allows agents to adopt correlated strategies without any cooperation. While CEs were introduced by Aumann in 1974 [1], CCEs were introduced in [18] and explicitly by [25] as a generalization of CEs. CCEs have been shown in game theory and computational literature to arise naturally from no-regret adaptive learning procedures ([19],[29, Section 17.4]). Moreover, they are computationally “easier” as shown by [15]. Finally, they are shown to be able to outperform NE payoffs in standard game theory [14, 24] even in situations where correlated equilibria cannot, for instance in potential games [28]. For these reasons in this paper we focus on CCEs.

CCEs can be interpreted as follows in an N -player setting. A moderator, or correlation device, picks a strategy profile for the N players randomly according to some publicly known distribution; then, she recommends it privately to the players. Before the lottery is run, each player has to decide whether to commit to the moderator recommendation (whatever it will be), assuming that all other players commit, only knowing the lottery distribution. If a player commits, then she is communicated in private her (and only her) selected strategy, and must follow it. Instead, if a player deviates, she will do so without any information on the outcome of the lottery, assuming that all other players follow the private suggestion they receive. A lottery is a CCE if every player prefers to commit rather than unilaterally deviate, assuming that all others do commit. CCEs are a generalization of Aumann’s notion of correlated equilibria (see [1, 2]), since in the latter each player is asked to commit to moderator’s lottery after having seen her suggested strategy.

Lately, correlated and coarse correlated equilibria have made their appearance in MFG literature. Bonnesini et al. [4, 5, 9] establish existence and convergence results for correlated equilibria in mean field games with discrete time and finite state and action spaces. A second group of papers by Muller et al. [26, 27] considers both CEs and CCEs in a similar setting. In addition, they provide an extensive discussion of learning algorithms for both types of equilibria in MFG. Lastly, in [7], CCEs have been introduced in both continuous time stochastic differential games and mean field games. The notion of coarse correlated solution to the MFG is justified by proving an approximation result. An existence result is also proved, by means of a minimax theorem. Although its generality, this result is not constructive, and the question of how to construct coarse correlated solutions to MFG is left open.

This paper’s goal is to develop a methodology for computing mean field CCEs, and to effectively compare them to mean field NEs and mean field control (MFC) solutions (see [12, 13] for an insightful discussion on the differences between such two notions and a quantitative comparison). For this reason, we do not consider the N -player game, but we limit our analysis to the mean field game. Since we search for explicit solutions, we restrict our analysis to linear-quadratic stochastic MFGs, working in a setting closely related to [16]. Applying our methodology to a toy model, we show that mean field CCEs indeed allow to significantly outperform the mean field NE in terms of payoffs under identified conditions.

We propose a notion of mean field CCE which is strongly inspired by the notion of coarse correlated solution to the MFG of [7]. As for a mean field NE, our notion of mean field CCEs is any suitable pair made of a strategy and a flow of moments, with the following important differences. The flow of moments can be stochastic, and the strategy can be correlated to the flow of moments even without the presence of a common noise, as it is the case in this paper. The way they are correlated is chosen by the moderator at the beginning of the game as part of the equilibrium. We call such pairs *correlated flows*. In few words, any of such correlated flows is a mean field CCE if the representative player has no incentive to deviate before knowing the

flow realisation, and if the flow is consistent, i.e., at any time t the flow of moments equals the conditional expectation of the representative player's state given the whole flow of moments up to terminal time.

Our main contributions can be summarised as follows:

- After focusing on a suitable class of suggested strategies and flows of moments verifying the consistency condition, we reduce the search of a mean field CCE to an inequality involving only the law of stochastic flow of moments at the equilibrium.
- We compare the payoffs of mean field CCEs with those of mean field NEs and MFC solutions. We show that the MFC optimal payoff is the unattainable upper bound for all mean field CCEs and provide a condition on the law of the stochastic flow of moments so that mean field CCEs in a specific class yield a higher payoff than mean field NE.
- Finally, we apply our results to an emission abatement game between countries, inspired by environmental economics literature on international environmental agreements [3, 14]. We show that it is possible to build simple mean field CCEs that both yield much higher payoffs than the mean field NE and guarantee higher average abatement levels.

The application also shows an additional interest of CCEs, which is to help a regulator not only to lead the population to a more optimal payoff than the free-riding NE, but also or otherwise to lead it to match other and potentially payoff-conflicting targets, such as the abatement level of players in this application. To the best of our knowledge, no attempt has been made so far to identify CCEs analytically in a mean field game, nor to explore and illustrate their potential in outperforming the payoffs of mean field Nash equilibria.

The rest of the paper is organised as follows: in Section 2 we state the assumptions, which will be in force throughout the whole paper, and give the definition of mean field CCE. In Section 3, we develop the methodology for computing mean field CCEs, while in Section 4 the comparison between mean field CCEs, MFC solutions and mean field NEs is carried out. In Section 5, we apply the results of the previous sections to the abatement game, and we analyse and explore the resulting characterization of the set of mean field CCEs which outperform the payoff of the unique mean field NE. Finally, we collect in the Appendix the most standard proofs, which we choose to include for the sake of completeness.

2 Setting

Let $T > 0$ be a fixed time horizon. Let $d, k \in \mathbb{N}$. For $n \in \mathbb{N}$, denote by \mathcal{S}^n the set of $n \times n$ symmetric matrices and by I_n the identity matrix in \mathcal{S}^n . We are going to work under the following set of assumptions.

Assumptions A. *Consider the following vector valued or matrix valued functions:*

- (1) $A, \sigma \in L^\infty([0, T]; \mathbb{R}^{d \times d});$
- (2) $B \in L^\infty([0, T]; \mathbb{R}^{d \times k});$
- (3) $Q, \bar{Q}, \tilde{Q} \in L^\infty([0, T]; \mathcal{S}^d), R \in \mathcal{C}([0, T]; \mathcal{S}^k), H, \bar{H}, \tilde{H} \in \mathcal{S}^d;$
- (4) $H, \bar{H}, \tilde{H} \geq 0, Q_t \geq d_1 I_d$ for every $t \in [0, T], d_1 \geq 0, R_t \geq d_2 I_k$ for every $t \in [0, T], d_2 > 0;$
- (5) $S \in L^\infty([0, T]; \mathbb{R}^{k \times d}), \sup_{t \in [0, T]} |S_t|^2 < d_1 d_2$ if $d_1 > 0, S_t = 0$ for every $t \in [0, T]$ otherwise;

(6) $L, q \in L^\infty([0, T]; \mathbb{R}^d)$, $r \in L^\infty([0, T]; \mathbb{R}^k)$.

Let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a complete filtered probability space satisfying usual assumptions, let W be a d dimensional \mathbb{F} -Brownian motion and let ξ be an \mathbb{R}^d -valued \mathcal{F}_0 -measurable random variable with law ν . Denote by ν_1 and ν_2 the first and second moments of ν respectively. Suppose that ξ and W are independent. Throughout the paper, we assume the following assumption:

Assumption U. *The σ -algebra \mathcal{F}_0 is large enough to support a \mathcal{F}_0 -measurable uniform random variable independent of ξ and W .*

In the following, we denote by $\mathbb{F}^1 = (\mathcal{F}_t^1)_{t \in [0, T]}$ the filtration generated by ξ and W , which we assume without loss of generality to satisfy the usual conditions.

Given an arbitrary filtration \mathbb{G} , we will use the standard notation $\mathbb{H}^2(\mathbb{G})$ for the set of all \mathbb{G} -progressively measurable \mathbb{R}^k -valued processes $\alpha = (\alpha_t)_{t \in [0, T]}$ such $\mathbb{E}[\int_0^T |\alpha_t|^2 dt] < \infty$.

We introduce the notion of correlated flow and mean field coarse correlated equilibrium.

Definition 1 (Correlated flow). A correlated flow is a pair (λ, μ) satisfying the following properties:

- i) $\lambda = (\lambda_t)_{t \in [0, T]}$ is a process in $\mathbb{H}^2(\mathbb{F})$.
- ii) $\mu = (\mu_t)_{t \in [0, T]}$ is an \mathcal{F}_0 -measurable $\mathcal{C}([0, T]; \mathbb{R}^d)$ -random variable.
- iii) μ is independent of both ξ and W .

We refer to λ as the *recommended strategy* and to μ as the *random flow of moments*.

We can interpret a correlated flow (λ, μ) as follows: moderator's lottery is run before the game starts and independently of the idiosyncratic shocks that determine the random evolution of representative player's state. This is made possible by Assumption U, which allows for some independent extra randomness. We stress that, while the recommended strategy λ is correlated both to ξ and W and to μ , μ is independent of the initial datum and the noise. We will sometimes use the equivalent expressions "correlated strategy" or "suggested strategy" to refer to λ .

Let us consider a correlated flow (λ, μ) . We now assign dynamics and payoff functional. We consider a state variable with linear dynamics given by

$$dX_t = (A_t X_t + B_t \lambda_t) dt + \sigma_t dW_t, \quad X_0 = \xi, \quad (2.1)$$

and a linear-quadratic payoff functional

$$\begin{aligned} \mathfrak{J}(\lambda, \mu) = \mathbb{E} \left[\int_0^T \left(\langle L_t, \mu_t \rangle - \frac{1}{2} \langle \bar{Q}_t \mu_t, \mu_t \rangle \right) - \left(\frac{1}{2} \langle Q_t X_t, X_t \rangle + \langle \tilde{Q}_t X_t, \mu_t \rangle + \langle q_t, X_t \rangle + \frac{1}{2} \langle R_t \lambda_t, \lambda_t \rangle \right. \right. \\ \left. \left. + \langle S_t X_t, \lambda_t \rangle + \langle r_t, \lambda_t \rangle \right) dt - \frac{1}{2} \langle \bar{H} \mu_T, \mu_T \rangle - \left(\frac{1}{2} \langle H X_T, X_T \rangle + \langle \tilde{H} X_T, \mu_T \rangle \right) \right]. \end{aligned} \quad (2.2)$$

When needed, we will stress the dependence of the process X on the control λ by using the notation X^λ .

Now, in order to move to the definition of mean field CCE, two cases must be distinguished. If the representative player decides to trust the mediator and therefore accepts to follow her recommendation λ before knowing it, the dynamics is given by equation (2.1), and the player gets the reward $\mathfrak{J}(\lambda, \mu)$. If instead she decides to deviate, she uses a strategy $\beta \in \mathbb{H}^2(\mathbb{F}^1)$, her state dynamics is given by equation (2.1) with β instead of λ , and her reward is $\mathfrak{J}(\beta, \mu)$. Observe

that when she deviates, her strategy β is measurable only with respect to the initial datum and the idiosyncratic noise, since she has no information on the outcome of the moderator's lottery. The deviating player can only use her knowledge of the law of the correlated flow (λ, μ) , which is assumed to be publicly known. As a consequence, when deviating, the state process X of the representative player is independent of the random flow of moments μ , which, however, still appears in her payoff.

Definition 2 (Mean field coarse correlated equilibrium). A correlated flow (λ, μ) is a mean field CCE if the following holds:

- (i) Optimality: for every deviation $\beta \in \mathbb{H}^2(\mathbb{F}^1)$, it holds

$$\mathfrak{J}(\lambda, \mu) \geq \mathfrak{J}(\beta, \mu). \quad (2.3)$$

- (ii) Consistency: let $X = (X_t)_{t \in [0, T]}$ be the solution to equation (2.1) with the control process λ . For every time $t \in [0, T]$, μ_t is a version of the conditional expectation of X_t given μ , that is,

$$\mu_t = \mathbb{E}[X_t | \mu] \quad \mathbb{P}\text{-a.s.} \quad \forall t \in [0, T]. \quad (2.4)$$

The definition of mean field CCE has two fundamental differences with the usual definition of mean field NE. First of all, as already mentioned, the optimality condition features an asymmetry between the suggested strategy, which belongs to $\mathbb{H}^2(\mathbb{F})$, and deviating player's strategies, which belong to the smaller class $\mathbb{H}^2(\mathbb{F}^1)$, since the former depends also on the information used by the moderator to run her lottery while the latter does not. As for the consistency condition, we notice that, coherently with μ being stochastic, it is formulated in terms of conditional expectations, although no common noise is present. It should be interpreted in the following way: if all players commit to the mediator's lottery outcomes before knowing them, then the flow of measures should arise from aggregation of the individual behaviors. In the mean field limit, the influence of the idiosyncratic noise on the flow of moments vanishes, while the influence of moderator's lottery does not. Therefore, μ stays stochastic and its stochasticity should derive from moderator's lottery only. We refer to [7] for more considerations and a deeper analysis of the connection with the N -player game.

Remark 1. The reader might have noticed that μ does not appear in the state dynamics (2.1). While computing mean field NEs and MFC solutions in the linear-quadratic case with the flow of moments in the dynamics is standard, computing mean field CCEs can be more delicate when μ appears in the state dynamics. We refer to Section 3.1 and to Remark 4 therein for more explanations.

Remark 2. In [7], moderator's lottery was modeled in the following way: an auxiliary probability space was chosen by the moderator to support the extra randomness for her lottery. As a consequence, the recommended strategy, dynamics and payoff were naturally defined on a suitable product space supporting ξ, W and such extra randomness. Here, thanks to Assumption **U**, the given filtration \mathbb{F} is already big enough to allow for any extra randomization the moderator might want to use. In both formulations moderator's lottery is run independently of ξ and W and deviations are measurable with respect to ξ and W only. Moreover, while [7] considers a stochastic flow of measures, and the consistency condition is given in terms of conditional probabilities, here it is enough to consider a flow of moments and conditional expectations, due to the linear-quadratic structure of the MFG.

3 Computing mean field coarse correlated equilibria

The set of coarse correlated equilibria is typically very wide and it is difficult to characterize in a continuous time setting. We therefore focus on a tractable class of correlated flows for which we are able to characterize a sufficient condition for being a mean field CCE. To do so, we adopt the following procedure:

- We fix a correlated flow (λ, μ) . We suppose that the representative player does not commit to the moderator's lottery and we compute her best deviating strategy $\hat{\beta}$, i.e.

$$\hat{\beta} = \arg \max_{\beta \in \mathbb{H}^2(\mathbb{R}^1)} \mathfrak{J}(\beta, \mu).$$

This is the content of Proposition 3.1. Observe that $\hat{\beta}$ will depend upon the law of (λ, μ) itself, but not on its actual realization.

- We define a parameterised class of correlated flows (λ, μ) of similar shape as the best deviating strategy $\hat{\beta}$ so that the consistency condition (2.4) is fulfilled. The correlation is due to a suitable random parameter δ . This is accomplished in subsection 3.2.
- Finally, for (λ, μ) in such a class, with corresponding parameter δ , we express the optimality condition

$$\mathfrak{J}(\lambda, \mu) \geq \mathfrak{J}(\hat{\beta}, \mu)$$

as an inequality involving the law of μ and δ only. Such an inequality is established in Theorem 3.3.

As a result, we reduce the search for a mean field CCE to finding a law for μ and δ that verifies an optimality inequality. The choice of focusing on a class of correlated flows with shape similar to the best deviation allows for explicit analytical comparison between the two payoffs in the optimality condition (2.3).

Remark 3. Interestingly, the outlined procedure does not involve the usual two steps procedure used to compute mean field NEs: first, optimize with a fixed flow of moments and, second, perform a fixed point argument to determine the flow. Indeed, we first impose the consistency condition and then we verify the optimality condition, more in line with an MFC fashion. This sheds light on one important feature of mean field CCEs: they can be regarded as a middle ground between mean field NE and MFC solutions. The comparison will be carried out in Section 4, and in Section 5 through the study of a simple yet important example.

3.1 Deviating player's optimization problem

Suppose that the representative player does not commit to the lottery. Therefore, as anticipated in Section 2, she chooses a strategy on her own before the moderator sends his recommendation, hence in particular without any information on the realisation of the correlated flow. The only information she has about (λ, μ) is the joint law of the pair itself, which is assumed to be publicly known. Due to the linear-quadratic structure of the MFG and the fact that any admissible deviation β is independent of μ , it turns out that knowing the expectation of μ_t for all $t \in [0, T]$ is enough.

Since the term $\int_0^T (\langle L_t, \mu_t \rangle - \frac{1}{2} \langle \bar{Q}_t \mu_t, \mu_t \rangle) dt - \langle \bar{H} \mu_T, \mu_T \rangle$ in (2.2) can be viewed as an uncontrolled constant for the deviating player's optimization problem, we can focus on the equivalent optimization problem

$$\min_{\beta \in \mathbb{H}^2(\mathbb{R}^1)} \mathfrak{J}'(\beta, \mu),$$

where

$$\begin{aligned} \mathfrak{J}'(\beta, \mu) = & \mathbb{E} \left[\int_0^T \left(\frac{1}{2} \langle Q_t X_t, X_t \rangle + \langle \tilde{Q}_t X_t, \mu_t \rangle + \langle q_t, X_t \rangle + \frac{1}{2} \langle R_t \beta_t, \beta_t \rangle + \langle S_t X_t, \beta_t \rangle + \langle r_t, \beta_t \rangle \right) dt \right. \\ & \left. + \frac{1}{2} \langle H X_T, X_T \rangle + \langle \tilde{H} X_T, \mu_T \rangle \right] \end{aligned} \quad (3.1)$$

under the constraint

$$dX_t = (A_t X_t + B_t \beta_t) dt + \sigma_t dW_t, \quad X_0 = \xi. \quad (3.2)$$

Since $\beta \in \mathbb{H}^2(\mathbb{F}^1)$, it follows that X is \mathbb{F}^1 -adapted, and therefore is independent of the flow of moments μ , which implies that deviating player's payoff can be written as:

$$\begin{aligned} \mathfrak{J}'(\beta, \mu) = & \int_0^T \mathbb{E} \left[\mathbb{E} \left[\frac{1}{2} \langle Q_t X_t, X_t \rangle + \langle \tilde{Q}_t X_t, \mu_t \rangle + \langle q_t, X_t \rangle + \frac{1}{2} \langle R_t \beta_t, \beta_t \rangle + \langle S_t X_t, \beta_t \rangle \right. \right. \\ & \left. \left. + \langle r_t, \beta_t \rangle \mid \mathcal{F}_t^1 \right] \right] dt + \mathbb{E} \left[\mathbb{E} \left[\frac{1}{2} \langle H X_T, X_T \rangle + \langle \tilde{H} X_T, \mu_T \rangle \mid \mathcal{F}_T^1 \right] \right] \\ = & \mathbb{E} \left[\int_0^T \left(\frac{1}{2} \langle Q_t X_t, X_t \rangle + \langle \tilde{Q}_t \mathbb{E}[\mu_t], X_t \rangle + \langle q_t, X_t \rangle + \frac{1}{2} \langle R_t \beta_t, \beta_t \rangle + \langle S_t X_t, \beta_t \rangle + \langle r_t, \beta_t \rangle \right) dt \right. \\ & \left. + \frac{1}{2} \langle H X_T, X_T \rangle + \langle \tilde{H} \mathbb{E}[\mu_T], X_T \rangle \right]. \end{aligned} \quad (3.3)$$

This is now a standard linear quadratic control problem, which can be solved by the stochastic maximum principle.

Proposition 3.1 (Optimal strategy for the deviating player). *Let ϕ , ψ and θ be the solutions of the following ODEs:*

$$\begin{cases} \dot{\phi}_t + \phi_t A_t + A_t^\top \phi_t + Q_t - (\phi_t B_t + S_t^\top) R_t^{-1} (B_t^\top \phi_t + S_t) = 0, & \phi_T = H, \\ \dot{\psi}_t + A_t^\top \psi_t + \tilde{Q}_t - (\phi_t B_t + S_t^\top) R_t^{-1} B_t^\top \psi_t = 0, & \psi_T = \tilde{H}, \\ \dot{\theta}_t + \psi_t \frac{d\mathbb{E}[\mu_t]}{dt} + A_t^\top \theta_t + q_t - (\phi_t B_t + S_t^\top) R_t^{-1} (B_t^\top \theta_t + r_t) = 0, & \theta_T = 0. \end{cases} \quad (3.4)$$

There exists a unique optimal strategy for the deviating player, which is given by

$$\hat{\beta}_t = -R_t^{-1} ((B_t^\top \phi_t + S_t) X_t + B_t^\top \psi_t \mathbb{E}[\mu_t] + B_t^\top \theta_t + r_t). \quad (3.5)$$

We postpone the proof to the Appendix. We observe only that the optimal control is actually feedback in the state X_t and in the expectation $\mathbb{E}[\mu_t]$. Moreover, while the functions ϕ and ψ do not depend upon μ or its expectation, the flow of expectations $\mathbb{E}[\mu_t]$ appears in the equation for θ , through its time derivative $\frac{d\mathbb{E}[\mu_t]}{dt}$.

Remark 4. This first step towards calculating mean field CCEs requires a filtering procedure, since the deviating player does not observe the actual realisation of μ . If the dynamics of the deviating player were dependent on μ , this step would require a much more involved analysis. Indeed, the state process X and μ would not be independent, even if $\beta \in \mathbb{H}^2(\mathbb{F}^1)$, which would lead to considering the projections on \mathbb{F}^1 of the processes X^i , $X^i X^j$ and $X^j \mu^i$, $1 \leq i, j \leq d$. This is why we have opted for a flow-free state dynamics, and postponed the analysis of the more general case to future research.

3.2 Correlated flow

We now consider a class of correlated flows (λ, μ) with a similar structure as the deviating player's best strategy $\hat{\beta}$ in (3.5). Our goal is to easily compare the payoff functionals $\mathfrak{J}'(\lambda, \mu)$ and $\mathfrak{J}'(\hat{\beta}, \mu)$. Hence we use the same functions ϕ and ψ , whereas we replace $\mathbb{E}[\mu_t]$ with μ_t itself and the term $R_t^{-1}(B_t^\top \theta_t + r_t)$ with a free parameter $\delta = (\delta_t)_{t \in [0, T]}$. Given any such δ , we define μ so that the consistency condition (2.4) is satisfied, so that we will be left with taking care of the optimality condition only.

More precisely, let \mathcal{G} be the set of all correlated flows (λ, μ) defined as

$$\begin{aligned} \lambda_t &= -R_t^{-1}((B_t^\top \phi_t + S_t)X_t + B_t^\top \psi_t \mu_t + \delta_t), \\ \dot{\mu}_t &= (A_t - B_t R_t^{-1}(B_t^\top \phi_t + S_t + B_t^\top \psi_t))\mu_t - B_t R_t^{-1} \delta_t, \quad \mu_0 = \nu_1, \end{aligned} \quad (3.6)$$

where $\delta = (\delta_t)_{t \in [0, T]}$ is any process in $\mathbb{H}^2(\mathcal{F}_0)$ independent of ξ and W , and ϕ and ψ are as in (3.4). The parameter δ represents the extra source of randomness in the correlated flow with respect to ξ and W .

Lemma 3.2. *Any correlated flow $(\lambda, \mu) \in \mathcal{G}$ satisfies the consistency condition (2.4).*

Proof. Let $(\lambda, \mu) \in \mathcal{G}$ corresponding to some $\delta \in \mathbb{H}^2(\mathcal{F}_0)$ independent of ξ and W . To ease the notation, set

$$\Phi_t = R_t^{-1}(B_t^\top \phi_t + S_t), \quad \Psi_t = R_t^{-1} B_t^\top \psi_t, \quad \Theta_t = R_t^{-1}(B_t^\top \theta_t + r_t). \quad (3.7)$$

Notice that μ satisfies the measurability requests of Definition 1. The dynamics of the representative player state is given by

$$\begin{aligned} dX_t &= ((A_t - B_t \Phi_t)X_t - B_t(\Psi_t \mu_t + R_t^{-1} \delta_t)) dt + \sigma_t dW_t, \\ X_0 &= \xi, \end{aligned} \quad (3.8)$$

which implies that the process $(\mu_t - X_t)_{t \in [0, T]}$ satisfies the stochastic differential equation

$$d(\mu_t - X_t) = (A_t - B_t \Phi_t)(\mu_t - X_t) dt - \sigma_t dW_t, \quad (3.9)$$

Since $\delta \in \mathbb{H}^2(\mathcal{F}_0)$, equation (3.8) admits a unique continuous adapted solution X satisfying $\mathbb{E}[\sup_{t \in [0, T]} |X_t|^2] < \infty$. Since ξ , W and δ are independent by assumption, by taking the conditional expectation with respect to μ in (3.9), we get

$$d\mathbb{E}[\mu_t - X_t | \mu] = (A_t - B_t \Phi_t) \mathbb{E}[\mu_t - X_t | \mu] dt, \quad \mathbb{E}[\mu_0 - X_0 | \mu] = \mu_0 - \mathbb{E}[\xi] = 0, \quad \mathbb{P}\text{-a.s.},$$

which implies $\mathbb{E}[\mu_t - X_t | \mu] = 0$ \mathbb{P} -a.s. for every t , i.e. (2.4). \square

Remark 5. Although the structure of the class \mathcal{G} is simple and quite specific, we will see later in the application section (Section 5) that it is rich enough to contain a large set of mean field CCEs with some desirable properties, such as significantly outperforming the mean field NE.

3.3 Optimality condition

Let $(\lambda, \mu) \in \mathcal{G}$. Since consistency has already been verified in Lemma 3.2, the goal is now to restate the optimality condition (2.3) in terms of quantities dependent upon the law of μ and δ only.

Theorem 3.3. Let $(\lambda, \mu) \in \mathcal{G}$ corresponding to some $\delta \in \mathbb{H}^2(\mathcal{F}_0)$ independent of ξ and W . Let Φ, Ψ and Θ be given by in (3.7). Set

$$M_t = Q_t + \Phi_t^\top R_t \Phi_t - 2\Phi_t^\top S_t, \quad N_t = \tilde{Q}_t + \Psi_t^\top R_t \Phi_t - \Psi_t^\top S_t, \quad G_t = \Psi_t^\top R_t \Psi_t. \quad (3.10)$$

Let $f(\mu) = (f_t(\mu))_{t \in [0, T]}$ be given by

$$\begin{cases} \dot{f}_t(\mu) = (A_t - B_t \Phi_t) f_t(\mu) + B_t (\Psi_t (\mu_t - \mathbb{E}[\mu_t]) + R_t^{-1} \delta_t - \Theta_t), & 0 \leq t \leq T, \\ f_0(\mu) = 0. \end{cases} \quad (3.11)$$

Then, (λ, μ) is a mean field CCE if and only if the following condition is satisfied:

$$\begin{aligned} & \int_0^T \left(\mathbb{E}[\langle N_t(\mu_t - \mathbb{E}[\mu_t]), \mu_t - \mathbb{E}[\mu_t] \rangle] + \frac{1}{2} (\mathbb{E}[\langle G_t \mu_t, \mu_t \rangle] - \langle G_t \mathbb{E}[\mu_t], \mathbb{E}[\mu_t] \rangle) \right. \\ & \quad \left. + \frac{1}{2} \mathbb{E}[\langle R_t^{-1} \delta_t, \delta_t \rangle] - \frac{1}{2} \langle R_t \Theta_t, \Theta_t \rangle \right) dt + \mathbb{E}[\langle \tilde{H}(\mu_T - \mathbb{E}[\mu_T]), \mu_T - \mathbb{E}[\mu_T] \rangle] \\ & \leq \int_0^T \left(\frac{1}{2} (\mathbb{E}[\langle M_t(\mu_t + f_t(\mu)), \mu_t + f_t(\mu) \rangle] - \mathbb{E}[\langle M_t \mu_t, \mu_t \rangle]) + \mathbb{E}[\langle N_t f_t(\mu), \mathbb{E}[\mu_t] \rangle] \right. \\ & \quad \left. + \langle q_t - \Phi_t^\top r_t, \mathbb{E}[f_t(\mu)] \rangle + \langle B_t^\top (\phi_t + \psi_t) \mathbb{E}[\mu_t], \Theta_t \rangle - \mathbb{E}[\langle B_t^\top (\phi_t + \psi_t) \mu_t, R_t^{-1} \delta_t \rangle] \right. \\ & \quad \left. + \mathbb{E}[\langle B_t^\top \phi_t f_t(\mu), \Theta_t \rangle] - \mathbb{E}[\langle r_t, \Theta_t - R_t^{-1} \delta_t \rangle] \right) dt \\ & \quad + \frac{1}{2} (\mathbb{E}[\langle H(\mu_T + f_T(\mu)), \mu_T + f_T(\mu) \rangle] - \mathbb{E}[\langle H \mu_T, \mu_T \rangle]) + \langle \tilde{H} \mathbb{E}[f_T(\mu)], \mathbb{E}[\mu_T] \rangle. \end{aligned} \quad (3.12)$$

Proof. Since $(\lambda, \mu) \in \mathcal{G}$ satisfies the consistency condition (2.4) by Lemma 3.2, we focus on the optimality condition (2.3). This is equivalent to verifying that

$$\mathfrak{J}'(\lambda, \mu) \leq \mathfrak{J}'(\hat{\beta}, \mu), \quad (3.13)$$

with $\hat{\beta}$ given by (3.5) and $\mathfrak{J}'(\lambda, \mu)$ and $\mathfrak{J}'(\hat{\beta}, \mu)$ are defined by (3.3). Denote by $\hat{X} = (\hat{X}_t)_{t \in [0, T]}$ the state of the deviating player when she uses the strategy $\hat{\beta}$ defined in (3.5), i.e.

$$d\hat{X}_t = ((A_t - B_t \Phi_t) \hat{X}_t - B_t (\Psi_t \mathbb{E}[\mu_t] + \Theta_t)) dt + \sigma_t dW_t, \quad \hat{X}_0 = \xi,$$

and by $X = (X_t)_{t \in [0, T]}$ the state of the representative player corresponding to the correlated flow (3.6), i.e.

$$dX_t = ((A_t - B_t \Phi_t) X_t - B_t (\Psi_t \mu_t + R_t^{-1} \delta_t)) dt + \sigma_t dW_t, \quad X_0 = \xi.$$

We rewrite the cost functionals $\mathfrak{J}'(\lambda, \mu)$ and $\mathfrak{J}'(\hat{\beta}, \mu)$ by taking advantage of the explicit form of λ and $\hat{\beta}$ and functions (3.10):

$$\begin{aligned} \mathfrak{J}'(\lambda, \mu) = & \mathbb{E} \left[\int_0^T \left(\frac{1}{2} \langle M_t X_t, X_t \rangle + \langle N_t X_t, \mu_t \rangle + \frac{1}{2} \langle G_t \mu_t, \mu_t \rangle + \langle q_t - \Phi_t^\top r_t, X_t \rangle \right. \right. \\ & \left. \left. + \langle (R_t \Phi_t - S_t) X_t, R_t^{-1} \delta_t \rangle + \langle R_t \Psi_t \mu_t, R_t^{-1} \delta_t \rangle - \langle \Psi_t^\top r_t, \mu_t \rangle \right. \right. \\ & \left. \left. + \frac{1}{2} \langle R_t^{-1} \delta_t, \delta_t \rangle - \langle r_t, R_t^{-1} \delta_t \rangle \right) dt + \frac{1}{2} \langle H X_T, X_T \rangle + \langle \tilde{H} X_T, \mu_T \rangle \right], \end{aligned}$$

and

$$\begin{aligned} \mathfrak{J}'(\hat{\beta}, \mu) = & \mathbb{E} \left[\int_0^T \left(\frac{1}{2} \langle M_t \hat{X}_t, \hat{X}_t \rangle + \langle N_t \hat{X}_t, \mathbb{E}[\mu_t] \rangle + \frac{1}{2} \langle G_t \mathbb{E}[\mu_t], \mathbb{E}[\mu_t] \rangle + \langle q_t - \Phi_t^\top r_t, \hat{X}_t \rangle \right. \right. \\ & \left. \left. + \langle (R_t \Phi_t - S_t) \hat{X}_t, \Theta_t \rangle + \langle R_t \Psi_t \mathbb{E}[\mu_t], \Theta_t \rangle - \langle \Psi_t^\top r_t, \mathbb{E}[\mu_t] \rangle \right. \right. \\ & \left. \left. + \frac{1}{2} \langle R_t \Theta_t, \Theta_t \rangle - \langle r_t, \Theta_t \rangle \right) dt + \frac{1}{2} \langle H \hat{X}_T, \hat{X}_T \rangle + \langle \tilde{H} \hat{X}_T, \mathbb{E}[\mu_T] \rangle \right] \end{aligned}$$

By Itô's formula, we get

$$d(\hat{X}_t - X_t) = (A_t - B_t \Phi_t)(\hat{X}_t - X_t)dt + B_t (\Psi_t(\mu_t - \mathbb{E}[\mu_t]) + R_t^{-1} \delta_t - \Theta_t) dt, \quad \hat{X}_0 - X_0 = 0,$$

so that it holds

$$\hat{X}_t = X_t + f_t(\mu), \quad 0 \leq t \leq T, \quad \mathbb{P}\text{-a.s.} \quad (3.14)$$

In particular, we note that $f(\mu)$ is $\sigma(\mu)$ -measurable. Then, we have

$$\begin{aligned} \mathbb{E}[\langle M_t \hat{X}_t, \hat{X}_t \rangle] &= \mathbb{E}[\langle M_t(X_t + f_t(\mu)), X_t + f_t(\mu) \rangle] \\ &= \mathbb{E}[\langle M_t X_t, X_t \rangle] + \mathbb{E}[\langle M_t(\mu_t + f_t(\mu)), \mu_t + f_t(\mu) \rangle] - \mathbb{E}[\langle M_t \mu_t, \mu_t \rangle], \end{aligned}$$

where we have used the fact that X satisfies the consistency condition (2.4). Therefore, we have

$$\begin{aligned} \mathfrak{J}'(\hat{\beta}, \mu) &= \mathbb{E} \left[\int_0^T \left(\frac{1}{2} \langle M_t X_t, X_t \rangle + \frac{1}{2} \langle M_t(\mu_t + f_t(\mu)), \mu_t + f_t(\mu) \rangle - \frac{1}{2} \langle M_t \mu_t, \mu_t \rangle + \langle N_t X_t, \mathbb{E}[\mu_t] \rangle \right. \right. \\ &\quad + \langle N_t f_t(\mu), \mathbb{E}[\mu_t] \rangle + \frac{1}{2} \langle G_t \mathbb{E}[\mu_t], \mathbb{E}[\mu_t] \rangle + \langle q_t - \Phi_t^\top r_t, X_t \rangle + \langle q_t - \Phi_t^\top r_t, f_t(\mu) \rangle \\ &\quad + \langle (R_t \Phi_t - S_t) X_t, \Theta_t \rangle + \langle (R_t \Phi_t - S_t) f_t(\mu), \Theta_t \rangle + \langle R_t \Psi_t \mathbb{E}[\mu_t], \Theta_t \rangle - \langle \Psi_t^\top r_t, \mathbb{E}[\mu_t] \rangle \\ &\quad + \frac{1}{2} \langle R_t \Theta_t, \Theta_t \rangle - \langle r_t, \Theta_t \rangle \left. \right) dt + \frac{1}{2} \langle H X_T, X_T \rangle + \frac{1}{2} \langle H(\mu_T + f_T(\mu)), \mu_T + f_T(\mu) \rangle - \frac{1}{2} \langle H \mu_T, \mu_T \rangle \\ &\quad + \langle \tilde{H} X_T, \mathbb{E}[\mu_T] \rangle + \langle \tilde{H} f_T(\mu), \mathbb{E}[\mu_T] \rangle \left. \right]. \end{aligned}$$

Since the correlated flow (λ, μ) satisfies the consistency condition (2.4), and noticing that

$$\begin{aligned} \mathbb{E}[\langle N_t \mu_t, \mathbb{E}[\mu_t] - \mu_t \rangle] &= -\mathbb{E}[\langle N_t(\mathbb{E}[\mu_t] - \mu_t), \mathbb{E}[\mu_t] - \mu_t \rangle], \\ \mathbb{E}[\langle (R_t \Phi_t - S_t) \mu_t, \Theta_t \rangle] &= \langle (R_t \Phi_t - S_t) \mathbb{E}[\mu_t], \Theta_t \rangle, \end{aligned}$$

we obtain

$$\begin{aligned} \mathfrak{J}'(\hat{\beta}, \mu) - \mathfrak{J}'(\lambda, \mu) &= \mathbb{E} \left[\int_0^T \left(\frac{1}{2} \langle M_t(\mu_t + f_t(\mu)), \mu_t + f_t(\mu) \rangle - \frac{1}{2} \langle M_t \mu_t, \mu_t \rangle \right. \right. \\ &\quad - \langle N_t(\mathbb{E}[\mu_t] - \mu_t), \mathbb{E}[\mu_t] - \mu_t \rangle + \langle N_t f_t(\mu), \mathbb{E}[\mu_t] \rangle + \frac{1}{2} \langle G_t \mathbb{E}[\mu_t], \mathbb{E}[\mu_t] \rangle - \frac{1}{2} \langle G_t \mu_t, \mu_t \rangle \\ &\quad + \langle q_t - \Phi_t^\top r_t, f_t(\mu) \rangle + \langle (R_t(\Phi_t + \Psi_t) - S_t) \mathbb{E}[\mu_t], \Theta_t \rangle - \langle (R_t(\Phi_t + \Psi_t) - S_t) \mu_t, R_t^{-1} \delta_t \rangle \\ &\quad + \langle (R_t \Phi_t - S_t) f_t(\mu), \Theta_t \rangle + \frac{1}{2} \langle R_t \Theta_t, \Theta_t \rangle - \frac{1}{2} \langle R_t^{-1} \delta_t, \delta_t \rangle - \langle r_t, \Theta_t - R_t^{-1} \delta_t \rangle \left. \right) dt \\ &\quad + \frac{1}{2} \langle H(\mu_T + f_T(\mu)), \mu_T + f_T(\mu) \rangle - \frac{1}{2} \langle H \mu_T, \mu_T \rangle \\ &\quad - \langle \tilde{H}(\mathbb{E}[\mu_T] - \mu_T), \mathbb{E}[\mu_T] - \mu_T \rangle + \langle \tilde{H} f_T(\mu), \mathbb{E}[\mu_T] \rangle \left. \right]. \end{aligned}$$

Therefore, the correlated flow (λ, μ) defined by (3.6) is a correlated flow if and only if the RHS above is non-negative. By rearranging the terms and using the equalities

$$R_t \Phi_t - S_t = B_t^\top \phi_t, \quad R_t \Psi_t = B_t^\top \psi_t, \quad (3.15)$$

we get condition (3.12). \square

The condition for a correlated flow of class \mathcal{G} to be a mean field CCE is now reduced to an optimality condition which only depends on the population average state and the correlating device of the mediator, i.e. on the joint law of (δ, μ) . Even though the inequality looks quite long, it can become very tractable and easy to interpret when one specifies some class of dynamics for μ as done in Section 5.

4 Comparison with MFC solution and mean field NE

In this section, we analyze the relationship between mean field CCEs, mean field NEs and MFC solutions. In more detail, we prove the following results:

- We compute the MFC solution $\hat{\alpha}^{MFC}$ and we show that it is unique and optimal in the broader class of controls $\mathbb{H}^2(\mathbb{F})$. This is accomplished in Proposition 4.1 and Lemma 4.2. Then, we show that no mean field CCE can outperform the payoff of the MFC solution. Moreover, if the MFC solution is not a mean field NE, we establish that the MFC payoff is unattainable by a mean field CCE. This is accomplished in Theorem 4.3.
- As for mean field NE, we show in Proposition 4.4 that there exists a unique mean field NE in our setting. Then, we show that for a correlated flow (λ, μ) to be a mean field CCE different from the mean field NE, it is necessary that the flow of moments μ is stochastic, which is the content of Theorem 4.5.
- Finally, Theorem 4.6 gives a condition so that a mean field CCE $(\lambda, \mu) \in \mathcal{G}$ yields a higher payoff than the mean field NE.

We remark that the results in the first two points above are fully general, in the sense that they do not restrict to correlated flows in the class \mathcal{G} defined by (3.6), while the condition on a mean field CCE (λ, μ) to outperform the payoff of the mean field NE is provided only for correlated flows in \mathcal{G} .

We recall here for reader's convenience the definitions of both mean field NE and MFC solution.

Definition 3. We say that a pair $(\hat{\alpha}, \hat{m}) \in \mathbb{H}^2(\mathbb{F}^1) \times \mathcal{C}([0, T]; \mathbb{R}^d)$ is a mean field Nash equilibrium if the following properties hold:

- (i) Optimality: $\hat{\alpha}$ maximizes $\mathfrak{J}(\cdot, \hat{m})$ over $\mathbb{H}^2(\mathbb{F}^1)$, i.e.,

$$\mathfrak{J}(\hat{\alpha}, \hat{m}) = \max_{\beta \in \mathbb{H}^2(\mathbb{F}^1)} \mathfrak{J}(\beta, \hat{m}). \quad (4.1)$$

- (ii) Consistency: let $X^{NE} = (X_t^{NE})_{t \in [0, T]}$ be the solution to equation (2.1) with the control process $\hat{\alpha}$. For every time $t \in [0, T]$, \hat{m}_t equals the expectation of X_t^{NE} , i.e.,

$$\hat{m}_t = \mathbb{E}[X_t^{NE}], \quad \forall t \in [0, T]. \quad (4.2)$$

Definition 4. For any $\beta \in \mathbb{H}^2(\mathbb{F}^1)$, let X^β be the solution of equation (2.1) with β instead of λ . Denote by $\mathbb{E}[X^\beta] = (\mathbb{E}[X_t^\beta])_{t \in [0, T]}$ the corresponding flow of first order moments. We say that a strategy $\hat{\alpha}^{MFC}$ is a MFC solution, if

$$\mathfrak{J}(\hat{\alpha}^{MFC}, \mathbb{E}[X^{\hat{\alpha}^{MFC}}]) = \max_{\beta \in \mathbb{H}^2(\mathbb{F}^1)} \mathfrak{J}(\beta, \mathbb{E}[X^\beta]). \quad (4.3)$$

4.1 Comparison with MFC solution

In this subsection we compare the expected payoffs of mean field CCEs and the MFC solution. In our setting, there exists a unique MFC solution. Since computations are very standard, we postpone them to the Appendix.

Proposition 4.1. Let ϕ^{MFC} and θ^{MFC} be the solutions of the following equations:

$$\begin{cases} \dot{\phi}_t^{MFC} + \phi_t^{MFC} A_t + A_t^\top \phi_t^{MFC} + (Q_t + 2\tilde{Q}_t + \bar{Q}_t) - (\phi_t^{MFC} B_t + S_t^\top) R_t^{-1} (B_t^\top \phi_t^{MFC} + S_t) = 0, \\ \phi_T^{MFC} = H + 2\tilde{H} + \bar{H}, \\ \dot{\theta}_t^{MFC} + A_t^\top \theta_t^{MFC} + q_t - L_t - (\phi_t^{MFC} B_t + S_t^\top) R_t^{-1} (B_t^\top \theta_t^{MFC} + r_t) = 0, \\ \theta_T^{MFC} = 0. \end{cases} \quad (4.4)$$

Define \bar{A} and \bar{B} as

$$\bar{A}_t = A_t - B_t R_t^{-1} B_t \phi_t^{MFC} - B_t R_t^{-1} S_t, \quad \bar{B}_t = B_t R_t^{-1} (B_t^\top \theta_t^{MFC} + r_t). \quad (4.5)$$

Let $\bar{\psi}$ and $\bar{\theta}$ be the solutions of the following equations:

$$\begin{cases} \dot{\bar{\psi}}_t + \bar{A}_t^\top \bar{\psi}_t + A_t^\top \bar{\psi}_t + (\bar{Q}_t + 2\tilde{Q}_t) - (\phi_t B_t + S_t^\top) R_t^{-1} B_t^\top \bar{\psi}_t = 0, & \bar{\psi}_T = \bar{H} + 2\tilde{H}, \\ \dot{\bar{\theta}}_t - \bar{\psi}_t \bar{B}_t + A_t^\top \bar{\theta}_t + q_t - (\phi_t B_t + S_t^\top) R_t^{-1} (B_t^\top \bar{\theta}_t + r_t) = 0, & \bar{\theta}_T = 0. \end{cases} \quad (4.6)$$

Let ϕ be the solution of the matrix Riccati equation in (3.4). There exists a unique MFC solution $\hat{\alpha}^{MFC}$, which is given by

$$\dot{\hat{\alpha}}_t^{MFC} = -R_t^{-1} ((B_t^\top \phi_t + S_t) X_t^{MFC} + B_t^\top \bar{\psi}_t \bar{x}_t^{MFC} + (B_t^\top \bar{\theta}_t + r_t)), \quad (4.7a)$$

$$\dot{\bar{x}}_t^{MFC} = (A_t - B_t R_t^{-1} B_t \phi_t^{MFC} - B_t R_t^{-1} S_t) \bar{x}_t^{MFC} - B_t R_t^{-1} (B_t^\top \theta_t^{MFC} + r_t), \quad \bar{x}_0^{MFC} = \nu_1, \quad (4.7b)$$

where X^{MFC} is the solution of

$$\begin{cases} dX_t^{MFC} = ((A_t - R_t^{-1} (B_t^\top \phi_t + S_t)) X_t^{MFC} - R_t^{-1} B_t^\top \bar{\psi}_t \bar{x}_t^{MFC} - R_t^{-1} (B_t^\top \bar{\theta}_t + r_t)) dt + \sigma_t dW_t, \\ X_0 = \xi. \end{cases} \quad (4.8)$$

In particular, it holds $\bar{x}_t^{MFC} = \mathbb{E}[X_t^{MFC}]$ for every $t \in [0, T]$.

Showing that no mean field CCE can outperform the payoff of the MFC solution requires first to show that the MFC solution is actually optimal over the larger control set $\mathbb{H}^2(\mathbb{F})$, as it is done in the following preliminary lemma:

Lemma 4.2. Let $\hat{\alpha}^{MFC}$ be the solution of the MFC problem. Then, for any β in $\mathbb{H}^2(\mathbb{F})$, $\beta \neq \hat{\alpha}$, it holds

$$\mathfrak{J}(\hat{\alpha}^{MFC}, \bar{x}^{MFC}) > \mathfrak{J}(\beta, \mathbb{E}[X^\beta]). \quad (4.9)$$

Proof. To ease the notation, we set

$$\mathfrak{J}^{MFC}(\alpha) := \mathfrak{J}(\alpha, \mathbb{E}[X^\alpha]), \quad (4.10)$$

where the process X has dynamics given by (2.1), for any $\alpha \in \mathbb{H}^2(\mathbb{F})$. We observe that

$$\begin{aligned} & \mathbb{E} \left[\frac{1}{2} \langle \bar{Q}_t \mathbb{E}[X_t], \mathbb{E}[X_t] \rangle + \langle \tilde{Q}_t X_t, \mathbb{E}[X_t] \rangle + \frac{1}{2} \langle Q_t X_t, X_t \rangle + \frac{1}{2} \langle R_t \alpha_t, \alpha_t \rangle + \langle S_t X_t, \alpha_t \rangle \right] \\ &= \mathbb{E} \left[\frac{1}{2} \langle (\bar{Q}_t + 2\tilde{Q}_t) \mathbb{E}[X_t], \mathbb{E}[X_t] \rangle + \frac{1}{2} \langle Q_t X_t, X_t \rangle + \frac{1}{2} \langle R_t \alpha_t, \alpha_t \rangle + \langle S_t X_t, \alpha_t \rangle \right] \\ &= \mathbb{E} \left[\frac{1}{2} \langle (\bar{Q}_t + 2\tilde{Q}_t + Q_t) \mathbb{E}[X_t], \mathbb{E}[X_t] \rangle + \frac{1}{2} \langle Q_t (X_t - \mathbb{E}[X_t]), (X_t - \mathbb{E}[X_t]) \rangle \right. \\ & \quad \left. + \frac{1}{2} \langle R_t \alpha_t, \alpha_t \rangle + \langle S_t (X_t - \mathbb{E}[X_t]), \alpha_t \rangle + \langle S_t \mathbb{E}[X_t], \alpha_t \rangle \right]. \end{aligned}$$

By Assumptions **A**, this equality implies that the running payoff in the functional \mathfrak{J}^{MFC} is strictly concave jointly in $\mathbb{E}[X_t]$, $X_t - \mathbb{E}[X_t]$ and α_t , for every $t \in [0, T]$. Since \mathfrak{J}^{MFC} is also upper semi-continuous, this implies that the maximum exists and that it is unique over the broader class $\mathbb{H}^2(\mathbb{F})$.

We are left to show that the maximum point is indeed $\hat{\alpha}^{MFC}$. For the sake of clarity, we set

$$\begin{aligned} f(t, x, m, a) &= \langle L_t, m \rangle - \frac{1}{2} \langle \bar{Q}_t m, m \rangle - \frac{1}{2} \langle Q_t x, x \rangle - \langle \tilde{Q}_t x, m \rangle - \langle q_t, x \rangle \\ &\quad - \frac{1}{2} \langle R_t a, a \rangle - \langle S_t x, a \rangle - \langle r_t, a \rangle, \\ g(x, m) &= - \left(\frac{1}{2} \langle \bar{H} m, m \rangle + \frac{1}{2} \langle H x, x \rangle + \langle \tilde{H} x, m \rangle \right). \end{aligned} \quad (4.11)$$

Let β in $\mathbb{H}^2(\mathbb{F})$. We define the following process:

$$\tilde{\beta}_t = \mathbb{E}[\beta_t | \mathcal{F}_t^1], \quad t \in [0, T]. \quad (4.12)$$

Since \mathbb{F}^1 satisfies the usual assumptions, $\tilde{\beta}$ can be taken \mathbb{F}^1 -progressively measurable (see, e.g., [6, Section 2]). Let $\tilde{X} = (\tilde{X}_t)_{t \in [0, T]}$ be the solution of

$$d\tilde{X}_t = (A_t \tilde{X}_t + B_t \tilde{\beta}_t) dt + \sigma_t dW_t, \quad \tilde{X}_0 = \xi.$$

Then, using the explicit expression for the solution \tilde{X} of the SDE above, it can be shown by direct computation that

$$\tilde{X}_t = \mathbb{E}[X_t^\beta | \mathcal{F}_t^1] \quad \mathbb{P}\text{-a.s.}, \quad t \in [0, T].$$

Due to the concave linear quadratic structure of f , we have the following:

$$\begin{aligned} \mathfrak{J}^{MFC}(\beta) &= \mathbb{E} \left[\int_0^T f(t, X_t^\beta, \mathbb{E}[X_t^\beta], \beta_t) dt + g(X_T^\beta, \mathbb{E}[X_T^\beta]) \right] \\ &= \int_0^T \mathbb{E} \left[\mathbb{E} \left[f(t, X_t^\beta, \mathbb{E}[X_t^\beta], \beta_t) \mid \mathcal{F}_t^1 \right] \right] dt + \mathbb{E} \left[\mathbb{E} \left[g(X_T^\beta, \mathbb{E}[X_T^\beta]) \mid \mathcal{F}_T^1 \right] \right] \\ &\leq \mathbb{E} \left[\int_0^T f(t, \mathbb{E}[X_t^\beta | \mathcal{F}_t^1], \mathbb{E}[X_t^\beta], \mathbb{E}[\beta_t | \mathcal{F}_t^1]) dt + g(\mathbb{E}[X_T^\beta | \mathcal{F}_T^1], \mathbb{E}[X_T^\beta]) \right] \\ &= \mathbb{E} \left[\int_0^T f(t, \tilde{X}_t, \mathbb{E}[\tilde{X}_t], \tilde{\beta}_t) dt + g(\tilde{X}_T, \mathbb{E}[\tilde{X}_T]) \right] = \mathfrak{J}^{MFC}(\tilde{\beta}), \end{aligned}$$

where we have used the fact that $\mathbb{E}[X_t^\beta] = \mathbb{E}[\tilde{X}_t]$ for every time t . Since $\tilde{\beta}$ belongs to $\mathbb{H}^2(\mathbb{F}^1)$, Proposition 4.1 implies

$$\mathfrak{J}^{MFC}(\beta) \leq \mathfrak{J}^{MFC}(\tilde{\beta}) \leq \mathfrak{J}^{MFC}(\hat{\alpha}^{MFC}).$$

By strict concavity, we deduce that the inequality is strict for any $\beta \neq \hat{\alpha}$. \square

In the next theorem we prove that the MFC solution provides an upper bound to the payoffs of any mean field CCEs. Moreover, this upper bound can not be attained unless the MFC solution is a mean field NE.

Theorem 4.3 (No outperformance over the MFC solution). *Let (λ, μ) a mean field CCE. Then, the following holds:*

- (i) *If $\mathfrak{J}(\lambda, \mu) \geq \mathfrak{J}(\hat{\alpha}^{MFC}, \bar{x}^{MFC})$, then $(\lambda, \mu) = (\hat{\alpha}^{MFC}, \bar{x}^{MFC})$, so $\mathfrak{J}(\lambda, \mu) = \mathfrak{J}(\hat{\alpha}^{MFC}, \bar{x}^{MFC})$;*

(ii) If the MFC solution is not a mean field NE, then $\mathfrak{J}(\lambda, \mu) < \mathfrak{J}(\hat{\alpha}^{MFC}, \bar{x}^{MFC})$. In particular, the MFC solution is not a mean field CCE either.

Proof of (i). By using the payoff functional \mathfrak{J}^{MFC} defined by (4.10), the payoffs' inequality reads as

$$\mathfrak{J}(\lambda, \mu) \geq \mathfrak{J}(\hat{\alpha}^{MFC}, \bar{x}^{MFC}) = \mathfrak{J}^{MFC}(\hat{\alpha}^{MFC}). \quad (4.13)$$

We reformulate the MFC problem weakly, by taking advantages of the results of [11, Paragraph 6.6]. We define the set $\mathcal{A} \subseteq \mathcal{P}(\mathbb{R}^d \times \mathcal{C}([0, T]; \mathbb{R}^d) \times L^2([0, T]; \mathbb{R}^k))$ of admissible probability measures in the following way: take any filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ satisfying the usual assumptions, equipped with a d -dimensional \mathbb{F} -Brownian motion W and an \mathcal{F}_0 -measurable random variable ξ independent of W . Let $\alpha \in \mathbb{H}^2(\mathbb{F})$, which we regard as random variable taking values in $L^2([0, T]; \mathbb{R}^k)$. Let $X = X^\alpha$ be the solution of

$$dX_t^\alpha = (A_t X_t^\alpha + B_t \alpha_t) dt + dW_t, \quad X_0^\alpha = \xi. \quad (4.14)$$

Then, a probability measure P belongs to \mathcal{A} if $P = \mathbb{P} \circ (\xi, X^\alpha, \alpha)^{-1}$. For any $P \in \mathcal{A}$, set $\bar{x}_t = \int_{\mathbb{R}^d} y (P \circ x_t^{-1})(dy)$. By recalling the definitions of f and g in (4.11), define the payoff functional

$$\begin{aligned} \mathcal{J}(P) &= \int_{\mathbb{R}^d \times \mathcal{C}([0, T], \mathbb{R}^d) \times L^2([0, T], \mathbb{R}^k)} \left(\int_0^T f(t, x_t, \bar{x}_t, a) dt + g(x_T, \bar{x}_T) \right) P(dz, dx, da) \\ &= \mathbb{E}^{\mathbb{P}} \left[\int_0^T \left(f(t, X_t, \mathbb{E}^{\mathbb{P}}[X_t], \alpha_t) \right) dt + g(X_T, \mathbb{E}^{\mathbb{P}}[X_T]) \right]. \end{aligned} \quad (4.15)$$

By [11, Theorem 6.37], there exists a probability measure P^* in \mathcal{A} so that

$$\mathcal{J}(P^*) \geq \mathcal{J}(P) \quad \forall P \in \mathcal{A}. \quad (4.16)$$

Let $(\xi, X^{MFC}, \hat{\alpha}^{MFC})$ be the MFC solution given by Proposition 4.1 and let \hat{P} be its law. Let (λ, μ) be a mean field CCE and $(\xi, X^\lambda, \lambda)$ be the corresponding initial state, state process and correlated strategy. We show the following properties:

1. The maximum point P^* is unique and it is equal to \hat{P} .
2. For every m in the support of μ , there exists a version of the regular conditional probability of $(\xi, \lambda, X^\lambda)$ given $\mu = m$; if we set $P^m = \mathbb{P}((\xi, \lambda, X^\lambda) \in \cdot \mid \mu = m)$, then P^m belongs to \mathcal{A} , and it holds

$$\mathfrak{J}(\lambda, \mu) = \int_{\mathcal{C}([0, T]; \mathbb{R}^d)} \mathcal{J}(P^m) \rho(dm),$$

where ρ denotes the law of μ .

3. We use the above equality to show that $\mu = \bar{x}^{MFC}$ \mathbb{P} -a.s. and deduce $\lambda = \hat{\alpha}^{MFC}$ $d\mathbb{P} \otimes dt$ -a.e..

As for point 1, let P^* be the admissible probability measure that maximizes \mathcal{J} . Let $(\Omega^*, \mathcal{F}^*, \mathbb{F}^*, \mathbb{P}^*)$, W^* , ξ^* , α^* and X^* be so that $P^* = \mathbb{P}^* \circ (\xi^*, \alpha^*, X^*)^{-1}$. By applying Proposition 4.1 in this probability space, there exists an optimal control $\hat{\beta}$ which maximizes \mathfrak{J} over $\mathbb{H}^2(\mathbb{F}^*)$. Since the flow of moments of $X^{\hat{\beta}}$ is still given by (4.7b) and (4.8) admits a strong solution, we have $\mathbb{P}^* \circ (\xi^*, X^{\hat{\beta}}, \hat{\beta})^{-1} = \mathbb{P} \circ (\xi, X^{MFC}, \hat{\alpha}^{MFC})^{-1} = \hat{P}$. Therefore, we can conclude that

$$\begin{aligned} \mathcal{J}(P^*) &= \mathbb{E}^{\mathbb{P}^*} \left[\int_0^T \left(f(t, X_t^*, \mathbb{E}^{\mathbb{P}^*}[X_t^*], \alpha_t^*) \right) dt + g(X_T^*, \mathbb{E}^{\mathbb{P}^*}[X_T^*]) \right] \\ &\geq \mathbb{E}^{\mathbb{P}^*} \left[\int_0^T \left(f(t, X_t^{\hat{\beta}}, \mathbb{E}^{\mathbb{P}^*}[X_t^{\hat{\beta}}], \hat{\beta}_t) \right) dt + g(X_T^{\hat{\beta}}, \mathbb{E}^{\mathbb{P}^*}[X_T^{\hat{\beta}}]) \right] = \mathcal{J}(\hat{P}), \end{aligned}$$

with the inequality being strict if $\beta^* \neq \hat{\beta}$. This shows point 1.

As for point 2, we can suppose without loss of generality that $(\Omega, \mathcal{F}, \mathbb{P})$ is a Polish probability space. We note that the state process X^λ is adapted to the filtration generated by ξ , W and λ , which is countably generated. This implies that there exists a version of the regular conditional probability of \mathbb{P} given $\mu = m$, that we denote by \mathbb{P}^m . Since ξ and W are independent of μ , it is straightforward to see that W is a Brownian motion under \mathbb{P}^m as well, the law of ξ under \mathbb{P}^m is ν and that X^λ still satisfies equation (2.1). Let $P^m = \mathbb{P}^m \circ (\xi, X^\lambda, \lambda)^{-1}$ and observe that P^m belongs to \mathcal{A} for ρ -a.e m in $\mathcal{C}([0, T], \mathbb{R}^d)$. The consistency condition implies that $\mathbb{E}^{\mathbb{P}^m} [X_t^\lambda] = m_t$ for ρ -a.e. m , which in turn implies that

$$\begin{aligned} \mathfrak{J}(\lambda, \mu) &= \mathbb{E}^{\mathbb{P}} \left[\int_0^T \left(f(t, X_t^\lambda, \mu_t, \lambda_t) \right) dt + g(X_T, \mu_T) \right] \\ &= \mathbb{E}^{\mathbb{P}} \left[\mathbb{E}^{\mathbb{P}} \left[\int_0^T \left(f(t, X_t^\lambda, \mu_t, \lambda_t) \right) dt + g(X_T, \mu_T) \middle| \mu \right] \right] \\ &= \int_{\mathcal{C}([0, T], \mathbb{R}^d)} \mathbb{E}^{\mathbb{P}^m} \left[\int_0^T \left(f(t, X_t^\lambda, \mathbb{E}^{\mathbb{P}^m} [X_t^\lambda], \lambda_t) \right) dt + g(X_T, \mathbb{E}^{\mathbb{P}^m} [X_T^\lambda]) \right] \rho(dm) \\ &= \int_{\mathcal{C}([0, T], \mathbb{R}^d)} \mathcal{J}(P^m) \rho(dm). \end{aligned}$$

By (4.13) and (4.16), we have

$$\int_{\mathcal{C}([0, T], \mathbb{R}^d)} (\mathcal{J}(\hat{P}) - \mathcal{J}(P^m)) \rho(dm) \leq 0, \quad \mathcal{J}(\hat{P}) \geq \mathcal{J}(P^m),$$

which implies $\mathcal{J}(\hat{P}) = \mathcal{J}(P^m)$ for ρ -a.e. m . Since \hat{P} is the unique maximizer of \mathcal{J} by point 1, we get $P^m = \hat{P}$ ρ -a.e.. In particular, this implies

$$m_t = \mathbb{E}^{\mathbb{P}^m} [X_t^\lambda] = \int_{\mathbb{R}^d} y(P^m \circ x_t)^{-1}(dy) = \int_{\mathbb{R}^d} y(\hat{P} \circ x_t)^{-1}(dy) = \mathbb{E}[X_t^{MFC}] = \bar{x}_t^{MFC} \text{ for } \rho\text{-a.e. } m.$$

Thus, μ is a.s. equal to \bar{x}^{MFC} , so that the consistency condition (2.4) for the mean field CCE (λ, μ) rewrites as $\bar{x}_t^{MFC} = \mathbb{E}[X_t^\lambda]$. Therefore, we have

$$\mathfrak{J}(\lambda, \mu) = \mathbb{E}^{\mathbb{P}} \left[\int_0^T \left(f(t, X_t^\lambda, \mathbb{E}[X_t^\lambda], \lambda_t) \right) dt + g(X_T, \mathbb{E}[X_T^\lambda]) \right] = \mathfrak{J}^{MFC}(\lambda).$$

Since, by Lemma 4.2, $\hat{\alpha}^{MFC}$ is unique, the previous equality implies that λ is equal to $\hat{\alpha}^{MFC} d\mathbb{P} \otimes dt$ -a.e., which concludes the proof. \square

Proof of (ii). Let us assume that the MFC solution $(\hat{\alpha}^{MFC}, \bar{x}^{MFC})$ is not a mean field NE (see upcoming Definition 3). By item (i) of Theorem 4.3, every mean field CCE yields a lower payoff than the MFC solution; moreover, if there was a mean field CCE yielding the same payoff as the MFC solution, it would be the MFC solution itself. Therefore, we just need to prove that the MFC solution is not a mean field CCE.

The pair $(\hat{\alpha}^{MFC}, \bar{x}^{MFC})$ is a correlated flow which satisfies the consistency condition in the definition of mean field CCE. Moreover, since $\hat{\alpha}^{MFC} \in \mathbb{H}^2(\mathbb{F}^1)$ and \bar{x}^{MFC} is deterministic, it satisfies the consistency condition of the definition of the mean field NE as well. Since by assumption the MFC solution is not a mean field NE, it is the optimality condition (4.1) in the upcoming definition of mean field NE which is not satisfied. Therefore, there exists $\beta \in \mathbb{H}^2(\mathbb{F}^1)$ so that $\mathfrak{J}(\beta, \bar{x}^{MFC}) > \mathfrak{J}(\hat{\alpha}^{MFC}, \bar{x}^{MFC})$. Since such β is an admissible deviation to the correlated flow $(\hat{\alpha}^{MFC}, \bar{x}^{MFC})$, the optimality condition (2.3) in definition of mean field CCE is not satisfied either. This means that the MFC solution is not a mean field CCE. \square

4.2 Comparison with mean field Nash equilibria

As for mean field NE, we first show that the only mean field CCE with deterministic flow of moments is the mean field NE itself. In particular, this implies that randomization of the flow of moments is needed for mean field CCEs to reach higher payoffs than the mean field NE. Differently from the MFC case, no general outperformance result can be established for mean field CCEs. Instead, one can derive an outperformance condition for correlated flows in the class \mathcal{G} , in a similar approach as for the optimality condition in subsection 3.3.

As shown by next proposition, there exists a unique mean field NE. The proof is a standard application of the Pontryagin maximum principle approach together with the fixed point argument of [11, Chapter 4]. We include it in the Appendix for the sake of completeness.

Proposition 4.4. *Let ϕ^{NE} and θ^{NE} the solutions of the following system:*

$$\begin{cases} \dot{\phi}_t^{NE} + \phi_t^{NE} A + A^\top \phi_t^{NE} + (Q + \tilde{Q}) - (\phi_t^{NE} B + S^\top) R^{-1} (B^\top \phi_t^{NE} + S) = 0, \\ \phi_T^{NE} = H + \tilde{H}, \\ \dot{\theta}_t^{NE} + A^\top \theta_t^{NE} + q - (\phi_t^{NE} B + S^\top) R^{-1} (B^\top \theta_t^{NE} + r_t) = 0, \\ \theta_T^{NE} = 0. \end{cases} \quad (4.17)$$

Define \hat{A} and \hat{B} as

$$\hat{A}_t = A_t - B_t R_t^{-1} B_t \phi_t^{NE} - B_t R_t^{-1} S, \quad \hat{B}_t = B_t R_t^{-1} (B_t^\top \theta_t^{NE} + r_t). \quad (4.18)$$

Let $\theta^{\hat{m}}$ be the solution of the following equation:

$$\dot{\theta}_t^{\hat{m}} - \psi_t \frac{d\hat{m}_t}{dt} + A_t^\top \theta_t^{\hat{m}} + q_t - (\phi_t B_t + S^\top) R_t^{-1} (B_t^\top \theta_t^{\hat{m}} + r_t) = 0, \quad \theta_T^{\hat{m}} = 0. \quad (4.19)$$

Let ϕ be the solution of the matrix Riccati equation in (3.4). There exists a unique mean field NE $(\hat{\alpha}, \hat{m})$, which is given by

$$\dot{\hat{m}}_t = (A_t - B_t R_t^{-1} B_t \phi_t^{NE} - B_t R_t^{-1} S) \hat{m}_t - B_t R_t^{-1} (B_t^\top \theta_t^{NE} + r_t), \quad \hat{m}_0 = \nu_1, \quad (4.20a)$$

$$\hat{\alpha}_t = -R_t^{-1} ((B_t^\top \phi_t + S) X_t^{NE} + B_t^\top \psi_t \hat{m}_t + B_t^\top \theta_t^{\hat{m}} + r_t), \quad (4.20b)$$

where X^{NE} is the solution of equation (2.1) with the control $\hat{\alpha}$.

We observe that, by definition, a mean field NE is a mean field CCE with deterministic flow of measures \hat{m} . The converse is true as well, as shown by the following Theorem:

Theorem 4.5. *Let (λ, μ) be a mean field CCE with deterministic μ . Then, (λ, μ) is the mean field NE.*

Proof. We start by observing that, by using the same concavity and projections arguments as in the proof of Lemma 4.2, we have

$$\mathfrak{J}(\hat{\alpha}, \hat{m}) > \mathfrak{J}(\beta, \hat{m}) \quad \forall \beta \in \mathbb{H}^2(\mathbb{F}), \quad \beta \neq \hat{\alpha}.$$

Let (λ, μ) be a coarse correlated equilibrium with deterministic flow of moments μ . Then, the consistency condition (2.4) becomes $\mu_t = \mathbb{E}[X_t^\lambda]$ for every time t . By optimality, it holds $\mathfrak{J}(\lambda, \mu) \geq \mathfrak{J}(\beta, \mu)$ for every $\beta \in \mathbb{H}^2(\mathbb{F}^1)$. By reasoning as in the proof of Lemma 4.2, there exists a strategy $\tilde{\lambda} \in \mathbb{H}^2(\mathbb{F}^1)$ so that

$$\tilde{\lambda}_t = \mathbb{E}[\lambda_t | \mathcal{F}_t^1], \quad X_t^{\tilde{\lambda}} = \mathbb{E}[X_t^\lambda | \mathcal{F}_t^1], \quad \mathbb{P}\text{-a.s.}, \quad \forall t \in [0, T],$$

where $X^{\tilde{\lambda}}$ is the solution of equation (2.1) corresponding to the strategy $\tilde{\lambda}$. Since μ is deterministic, by exploiting the convex linear quadratic structure of the payoff functional \mathfrak{J}' , we have

$$\mathfrak{J}(\beta, \mu) \leq \mathfrak{J}(\lambda, \mu) \leq \mathfrak{J}(\tilde{\lambda}, \mu), \quad \forall \beta \in \mathbb{H}^2(\mathbb{F}^1). \quad (4.21)$$

Since μ is deterministic by assumption, the consistency condition holds true for the correlated flow $(\tilde{\lambda}, \mu)$ as well, so that (4.21) implies that $(\tilde{\lambda}, \mu)$ is itself a mean field NE. By uniqueness of the mean field NE, we deduce $(\tilde{\lambda}, \mu) = (\hat{\alpha}, \hat{m})$, so that in particular $\mu = \hat{m}$ \mathbb{P} -a.s.. Since $\hat{\alpha}$ is the unique maximizer of $\mathfrak{J}(\cdot, \hat{m})$ over $\mathbb{H}^2(\mathbb{F})$, we deduce $\mathfrak{J}(\hat{\alpha}, \hat{m}) \geq \mathfrak{J}(\lambda, \hat{m})$. Since (λ, μ) is a mean field CCE by assumption, it holds $\mathfrak{J}(\hat{\alpha}, \hat{m}) \leq \mathfrak{J}(\lambda, \hat{m})$, which, by uniqueness, implies that $\lambda = \hat{\alpha}$ $dt \otimes \mathbb{P}$ -a.e. as well. \square

Finally, consider again correlated flows (λ, μ) in the class \mathcal{G} . By using their specific structure as described in (3.6), we are able to provide a condition under which they yield a higher payoff than the mean field NE.

Theorem 4.6. *Let $\theta^{\hat{m}}$ be the solution of (4.19). Set*

$$\Theta_t^{\hat{m}} = R_t^{-1}(B_t^\top \theta_t^{\hat{m}} + r_t). \quad (4.22)$$

Let $(\lambda, \mu) \in \mathcal{G}$ corresponding to some $\delta \in \mathbb{H}^2(\mathcal{F}_0)$. Then, $\mathfrak{J}(\lambda, \mu)$ is higher than the payoff $\mathfrak{J}(\hat{\alpha}, \hat{m})$ given by the mean field NE if and only if the following inequality is satisfied:

$$\begin{aligned} & \int_0^T \left(\frac{1}{2} (\langle \bar{Q} \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle \bar{Q} \mu_t, \mu_t \rangle]) + \frac{1}{2} (\langle M_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle M_t \mu_t, \mu_t \rangle]) \right. \\ & + \frac{1}{2} (\langle G_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle G_t \mu_t, \mu_t \rangle]) + \langle N_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle N_t \mu_t, \mu_t \rangle] \\ & + \langle L_t - q_t + R_t^{-1}(B_t^\top (\phi_t + \psi_t) + S_t)^\top r_t, \mathbb{E}[\mu_t] - \hat{m}_t \rangle + \langle B_t^\top \phi_t \hat{m}_t, \Theta_t^{\hat{m}} \rangle \\ & - \mathbb{E}[\langle B_t^\top \phi_t \mu_t, R_t^{-1} \delta_t \rangle] - \mathbb{E}[\langle B_t^\top \psi_t \mu_t, R_t^{-1} \delta_t \rangle] \\ & + \langle B_t^\top \psi_t \hat{m}_t, \Theta_t^{\hat{m}} \rangle + \frac{1}{2} (\langle R_t \Theta_t^{\hat{m}}, \Theta_t^{\hat{m}} \rangle - \mathbb{E}[\langle R_t^{-1} \delta_t, \delta_t \rangle]) - \langle r_t, \Theta_t^{\hat{m}} - R_t^{-1} \mathbb{E}[\delta_t] \rangle \Big) dt \\ & + \frac{1}{2} (\langle \bar{H} \hat{m}_T, \hat{m}_T \rangle - \mathbb{E}[\langle \bar{H} \mu_T, \mu_T \rangle]) + \frac{1}{2} (\langle H \hat{m}_T, \hat{m}_T \rangle - \mathbb{E}[\langle H \mu_T, \mu_T \rangle]) \\ & + \langle \tilde{H} \hat{m}_T, \hat{m}_T \rangle - \mathbb{E}[\langle \tilde{H} \mu_T, \mu_T \rangle] \geq 0. \end{aligned} \quad (4.23)$$

The proof is similar to the one of Theorem 3.3. For the sake of completeness, we include it in the Appendix. We observe that, although the inequality (4.23) is not easy to interpret, it involves only the law of μ and its associated δ . Moreover, it can be verified separately from the optimality condition (3.12), giving some room for mean field CCEs to outperform the mean field NE payoff. This will be accomplished for the abatement game in Section 5.

5 Application to an emission abatement game

In this section we consider an emission abatement game inspired by environmental economics literature on international environmental agreements, in line with the very popular model of [3]. Previous section's findings allow us to exhibit a simple class of coarse correlated equilibria which (highly) outperforms the mean field NE in this game.

The emission abatement game has the following payoff and dynamics of the representative player state:

$$\mathfrak{J}(\alpha, \mu) = \mathbb{E} \left[\int_0^T \left(a \mu_t - \frac{b}{2} \mu_t^2 - \frac{1}{2} \alpha_t^2 - \frac{\varepsilon}{2} (\mu_t - X_t)^2 \right) dt \right], \quad (5.1a)$$

$$dX_t = \alpha_t dt + dW_t, \quad X_0 = \xi, \quad (5.1b)$$

with a, b non-negative constants and $\varepsilon > 0$. The strategy α_t represents the abatement rate of the player at time t , while X_t models the cumulated abatement over the interval $[0, t]$.

We translate a slightly modified version of the abatement game [3] into a dynamic stochastic mean field game. We follow the N -player formulation of [14] by considering symmetric players, and a normalization of the number of players is implicitly added by replacing the sum of abatement efforts by the flow of moments μ . We also add the last term in ε , inspired by further developments of this model in the literature (see [17]), which can be interpreted as a reputational cost. It appears to be necessary when one wants mean field CCEs outperforming the mean field NE at the mean field limit. Indeed, when $\varepsilon = 0$, there exists only a unique mean field CCE, corresponding to the mean field NE. This is straightforward by direct computations and can be also deduced from Proposition 5.1 (see upcoming Remark 7).

Following [3], the other terms of the payoff can be interpreted as follows. The term $a\mu_t - \frac{b}{2}\mu_t^2$, which depends solely on the mean field component μ , is the ‘‘abatement benefit’’. It represents the individual benefit of global climate change mitigation allowed by aggregate abatement efforts, with a decreasing marginal benefit. The quadratic term in the control, i.e. $-\frac{1}{2}\alpha_t^2$, is an ‘‘abatement cost’’ that the representative country privately pays for its abatement effort.

We do not claim that a mean field approximation of the abatement game of [3] is a right way to approach the problem of international environmental agreements economically. We rather use this payoff functional as a toy example that allows us to illustrate very efficiently the interest of mean field CCEs in a context of common good, and to contribute to the findings of [14].

Remark 6. Going from static to dynamic games also induces some additional assumptions that were not included in reference models [3, 14]. We chose to represent the ‘‘abatement benefit’’ as a running payoff rather than a terminal one, considering that environmental objectives are not only to reach a given level of emissions at a terminal time, but also to abate as much as possible, as early as possible.

5.1 Translation and interpretation of findings in the abatement game

In this subsection we apply the theory developed in the previous section to compute mean field CCEs in the abatement game. In the next subsection, we will make a step further and exhibit a simple but interesting subclass of correlated flows (λ, μ) which verify both the optimality inequality (3.12) and the NE outperformance inequality (4.23).

We use the setting of Section 2 with $d = k = 1$. The parameters are given by

$$A_t = 0, B_t = 1, \sigma_t = 1, L_t = a, \bar{Q}_t = b + \varepsilon, Q_t = \varepsilon, \tilde{Q}_t = -\varepsilon, R_t = 1, \quad \forall t \in [0, T], \quad (5.2)$$

and remaining parameters equal 0. According to Proposition 3.1 with the abatement game parameters as in (5.2), for a given correlated flow (λ, μ) , the best deviating strategy and the corresponding state process are given by

$$\begin{aligned} \hat{\beta}_t &= \phi_t(\mathbb{E}[\mu_t] - \hat{X}_t) - \theta_t, \\ d\hat{X}_t &= \hat{\beta}_t dt + dW_t, \quad \hat{X}_0 = \xi, \end{aligned} \quad (5.3)$$

with ϕ and θ satisfying equations

$$\begin{cases} \dot{\phi}_t + \varepsilon - \phi_t^2 = 0, & \phi_T = 0, \\ \dot{\theta}_t - \phi_t \left(\theta_t + \frac{d\mathbb{E}[\mu_t]}{dt} \right) = 0, & \theta_T = 0. \end{cases} \quad (5.4)$$

Note that ψ does not appear as in this case $\psi = -\phi$. We stress that, as only unilateral deviation is allowed, the deviating player can not act on the abatement benefit, and therefore does not consider a and b in her optimal strategy.

The family \mathcal{G} of correlated flows defined by (3.6) is composed of any correlated flow (λ, μ) so that:

$$\begin{aligned}\lambda_t &= \phi_t(\mu_t - X_t) - \delta_t, \\ \dot{\mu}_t &= -\delta_t, \quad \mu_0 = \nu_1,\end{aligned}$$

for some $\delta \in \mathbb{H}^2(\mathcal{F}_0)$ and where X solves

$$dX_t = \lambda_t dt + dW_t, \quad X_0 = \xi.$$

In particular, we note that it holds $\delta_t = -\frac{d\mu}{dt}$. Therefore, in this model, the class \mathcal{G} is composed of correlated flows (λ, μ) verifying

$$\lambda_t = \phi_t(\mu_t - X_t) + \frac{d\mu}{dt}, \quad \mathbb{E} \left[\int_0^T \left(\frac{d\mu}{dt} \right)^2 dt \right] < \infty. \quad (5.5)$$

As the correlated strategy depends on μ itself, we remark that the state variable becomes actually mean-reverting. The extra term $\frac{d\mu}{dt}$ allows the state of the representative player following the suggested strategy to satisfy the consistency condition by following the suggested variations of μ .

As shown by the next proposition, in the abatement game, the optimality condition only depends on the law of μ , the reputational cost parameter ε , and the final time horizon T .

Proposition 5.1 (Optimality condition for the abatement game). *Let (λ, μ) be a correlated flow in \mathcal{G} . Let $f(\mu) = (f_t(\mu))_{t \in [0, T]}$ be given by*

$$\begin{cases} \dot{f}_t(\mu) = - \left(\phi_t (f_t(\mu) + \mu_t - \mathbb{E}[\mu_t]) + \frac{d\mu}{dt} + \theta_t \right), & 0 \leq t \leq T, \\ f_0(\mu) = 0. \end{cases} \quad (5.6)$$

Then, (λ, μ) is a mean field CCE if and only if the following condition is satisfied:

$$\int_0^T \mathbb{E} \left[\left(\frac{d\mu}{dt} \right)^2 \right] dt \leq \int_0^T \mathbb{E} \left[(\phi_t f_t(\mu) + \theta_t)^2 + \phi_t^2 (\mu_t - \mathbb{E}[\mu_t] + f_t(\mu))^2 + (\varepsilon - \phi_t^2) f_t(\mu)^2 \right] dt. \quad (5.7)$$

Proof. Referring to (3.7) and (3.10), the auxiliary functions for the abatement game are as follows:

$$\begin{aligned}\Phi_t &= \phi_t, & \Psi_t &= -\phi_t, & \Theta_t &= \theta_t, \\ M_t &= \varepsilon + \phi_t^2, & N_t &= -\varepsilon - \phi_t^2, & G_t &= \phi_t^2.\end{aligned} \quad (5.8)$$

This implies that $f(\mu)$ given in (3.11) takes the form of equation (5.6), recalling that $\delta_t = -d\mu/dt$ by (5.5). After a few computations, we get that the optimality condition (3.12) rewrites as

$$\begin{aligned}\int_0^T \mathbb{E} [\delta_t^2] dt &\leq \int_0^T \mathbb{E} \left[(\phi_t f_t(\mu) + \theta_t)^2 + \phi_t^2 (\mu_t - \mathbb{E}[\mu_t] + f_t(\mu))^2 + (\varepsilon - \phi_t^2) f_t^2(\mu) \right. \\ &\quad \left. + 2\varepsilon(\mu_t - \mathbb{E}[\mu_t])(\mu_t - \mathbb{E}[\mu_t] + f_t(\mu)) \right] dt,\end{aligned}$$

using that

$$\mathbb{E}[\mu_t^2 - \mathbb{E}[\mu_t]^2] = \mathbb{E}[(\mu_t - \mathbb{E}[\mu_t])^2].$$

Since $f_t(\mu) = \hat{X}_t - X_t$ \mathbb{P} -a.s., for every time t by (3.14) and $f(\mu)$ is $\sigma(\mu)$ -measurable by definition, we have

$$f_t(\mu) = \mathbb{E}[f_t(\mu)|\mu] = \mathbb{E}[\hat{X}_t - X_t|\mu] = \mathbb{E}[\hat{X}_t] - \mu_t$$

where we used the consistency condition (2.4) and the fact that \hat{X} and μ are independent. This implies that

$$\mathbb{E}[(\mu_t - \mathbb{E}[\mu_t])(\mu_t - \mathbb{E}[\mu_t] + f_t(\mu))] = \mathbb{E}[\mu_t - \mathbb{E}[\mu_t]](-\mathbb{E}[\mu_t] + \mathbb{E}[\hat{X}_t]) = 0.$$

Therefore, (λ, μ) is a mean field CCE if and only if condition (5.7) is satisfied. \square

By Proposition 4.4, there exists a unique mean field NE $(\hat{\alpha}, \hat{m})$, which is given by

$$\hat{\alpha}_t = \phi_t(\hat{m}_t - X_t^{NE}), \quad (5.9a)$$

$$\hat{m}_t = \nu_1, \quad \forall t \in [0, T], \quad (5.9b)$$

since we have $\phi^{NE} = \theta^{NE} = 0$ in (4.17), which implies $\theta^{\hat{m}} = 0$ in (4.19) as well.

The mean field NE consists, on average, to null abatement, as \hat{m}_t stays constant equal to its initial value. This corresponds to a free-riding equilibrium, where everybody does as little as possible, and prefers to take advantage of the others' efforts. As a result, nobody does anything.

Remark 7. One can easily see from the optimality condition in equation (5.7) that, if $\varepsilon = 0$, the only mean field CCE is the mean field NE. Indeed, in this case $\phi_t = \theta_t = 0$ and $f_t(\mu) = d\mu/dt$, for all $t \in [0, T]$. Hence the right-hand side term in (5.7) is null, forcing $d\mu/dt = 0$, $t \in [0, T]$. As $\lambda_t = d\mu/dt$, we get $\lambda_t = \hat{\alpha}_t = 0$, $\hat{m}_t = \nu_1$, which is the mean field NE when $\varepsilon = 0$. This seems consistent with the findings of [14]. Indeed, in an equivalent N -player static deterministic game without the reputational cost ($\varepsilon = 0$), the authors find that, the more players, the less the payoff-maximising CCE outperforms the payoff of the NE. This probably comes from the fact that, at the mean field limit, there is only one mean field CCE, which is the mean field Nash equilibrium itself.

By Proposition 4.1, there exists a unique MFC solution $(\hat{\alpha}^{MFC}, \bar{x}_t^{MFC})$ which reads:

$$\hat{\alpha}_t^{MFC} = \phi_t(\bar{x}_t^{MFC} - X_t^{MFC}) - \bar{\eta}_t \bar{x}_t^{MFC} - \bar{\chi}_t, \quad (5.10a)$$

$$\dot{\bar{x}}_t^{MFC} = -\bar{\eta}_t \bar{x}_t^{MFC} - \bar{\chi}_t, \quad \bar{x}_0^{MFC} = \nu_1, \quad (5.10b)$$

with

$$\begin{cases} \dot{\bar{\eta}}_t = \bar{\eta}_t^2 - b, & \bar{\eta}_T = 0, \\ \dot{\bar{\chi}}_t = \bar{\eta}_t \bar{\chi}_t + a, & \bar{\chi}_T = 0. \end{cases} \quad (5.11)$$

The MFC solution adds to the mean-reversion two terms which depend on a and b , i.e. on the coefficients of the abatement benefit. One can note that the MFC solution and the mean field NE coincide if and only if $a = b = 0$. To the contrary, when the ‘‘common good’’ aspect of climate is accounted for in the payoff through the abatement benefit, the central planner can reach higher payoffs by preventing any inefficient free-riding behaviour. This gives some room for mean field CCEs to bridge the gap between the free-riding mean field NE and the central planner optimum.

To find mean field CCEs outperforming the mean field NE, the following condition should be fulfilled.

Proposition 5.2 (Outperformance condition over mean field NE). *Let (λ, μ) be a correlated flow in \mathcal{G} . Then,*

$$\mathfrak{J}(\lambda, \mu) \geq \mathfrak{J}(\hat{\alpha}, \hat{m}) \iff \mathbb{E} \left[\int_0^T \left(a(\mu_t - \hat{m}_t) - \frac{b}{2}(\mu_t^2 - \hat{m}_t^2) \right) dt \right] \geq \frac{1}{2} \mathbb{E} \left[\int_0^T \left(\frac{d\mu}{dt} \right)^2 dt \right].$$

Proof. By recalling the identities in (5.8), inequality (4.23) takes the following form:

$$\begin{aligned} \mathfrak{J}(\lambda, \mu) - \mathfrak{J}(\hat{\alpha}, \hat{m}) &= \int_0^T \left(\frac{1}{2}(b + \varepsilon)(\hat{m}_t^2 - \mathbb{E}[\mu_t^2]) + \frac{1}{2}(\varepsilon + \phi_t^2)(\hat{m}_t^2 - \mathbb{E}[\mu_t^2]) + \frac{1}{2}\phi_t^2(\hat{m}_t^2 - \mathbb{E}[\mu_t^2]) \right. \\ &\quad \left. - (\varepsilon + \phi_t^2)(\hat{m}_t^2 - \mathbb{E}[\mu_t^2]) + a(\mathbb{E}[\mu_t] - \hat{m}) - \phi_t \mathbb{E}[\mu_t \delta_t] + \phi \mathbb{E}[\delta_t \mu_t] - \frac{1}{2}\mathbb{E}[\delta_t^2] \right) \geq 0 \end{aligned}$$

By rearranging the terms and recalling that $\delta_t = -d\mu/dt$, we get the desired inequality. \square

The equivalence in Proposition 5.2 clearly illustrates that, when $a = b = 0$, the best payoff mean field CCE is actually the mean field NE. This was also implied by the fact that, when $a = b = 0$, the MFC solution is a mean field NE as we already mention above.

5.2 A tractable class of mean field CCEs

In this subsection we show that, when $a \neq 0$ or $b \neq 0$, the optimality and outperformance conditions are not empty, and neither is their intersection. In this case, the MFC solution is distinct from the mean field NE, which implies, according to Theorem 4.3, that the MFC solution is not a mean field CCE, as the required control does not resist any unilateral deviation which tends to a less costly free-riding option. However, by introducing correlation through correlated flows, one can manage to drive the population at quite high abatement levels, leading to more desirable social outcomes than the one of the mean field NE.

The optimality condition (5.7) is very convenient to use when one focuses on a specific class of dynamics for μ . In this subsection, we consider a subclass $\mathcal{G}_l \subseteq \mathcal{G}$ where the flows of moments are linear in time.

More precisely, let \mathcal{G}_l be the set of all correlated flows $(\lambda, \mu) \in \mathcal{G}$ such that

$$\mu_t = \nu_1 + tZ, \quad t \in [0, T], \quad (5.12)$$

for some $Z \in L^2(\mathcal{F}_0)$ independent of ξ and W . Then, for all correlated flows $(\lambda, \mu) \in \mathcal{G}_l$ we have

$$\lambda_t = \phi_t(\mu_t - X_t) + Z, \quad t \in [0, T].$$

In the rest of the paper, we will use the notations $z_1 := \mathbb{E}[Z]$, $z_2 := \mathbb{E}[Z^2]$, $\sigma_z^2 := \mathbb{V}[Z]$.

Proposition 5.3 (Optimality condition for \mathcal{G}_l). *Let $(\lambda, \mu) \in \mathcal{G}_l$. Then (λ, μ) is a mean field CCE if and only if*

$$z_1^2 c_M + \sigma_z^2 c_V \geq 0 \quad (5.13)$$

with

$$c_M = \int_0^T \left((\phi_t r_t + g_t)^2 + \varepsilon r_t^2 \right) dt - T, \quad c_V = \int_0^T \left(\phi_t^2 (v_t - t)^2 + \varepsilon v_t^2 \right) dt - T, \quad (5.14)$$

and

$$g_t = \int_t^T \phi_s e^{-\int_t^s \phi_u du} ds, \quad r_t = \int_0^t (1 - g_s) e^{-\int_s^t \phi_u du} ds, \quad v_t = \int_0^t (s\phi_s + 1) e^{-\int_s^t \phi_u du} ds. \quad (5.15)$$

Proof. For any given $(\lambda, \mu) \in \mathcal{G}_l$ we have $\frac{d\mu}{dt} = Z$ a.s. and $\mathbb{E}[\mu_t] = \nu_1 + tz_1$ so that

$$\begin{aligned} \int_0^T \mathbb{E} \left[\left(\frac{d\mu}{dt} \right)^2 \right] dt &= Tz_2, & \dot{\theta}_t &= \phi_t(\theta_t + z_1), \quad \theta_T = 0, \\ \dot{f}_t(\mu) &= -(\phi_t(f_t(\mu) + t(Z - z_1)) + Z + \theta_t), & f_0(\mu) &= 0. \end{aligned}$$

Let us set

$$p_t := \int_0^t e^{-\int_s^t \phi_u du} ds.$$

By using p_t and the auxiliary functions defined in (5.15), $f_t(\mu)$ and θ_t can be rewritten as follows:

$$\theta_t = -z_1 g_t, \quad f_t(\mu) = -Z p_t - (Z - z_1)(v_t - p_t) + z_1(p_t - r_t).$$

We compute the different terms appearing in the integral of the right-hand side of the optimality condition:

$$\begin{aligned} \mathbb{E}[(f_t(\mu))^2] &= \sigma_z^2 v_t^2 + z_1^2 r_t^2, & \mathbb{E}[(\mu_t - \mathbb{E}[\mu_t] + f_t(\mu))^2] &= \sigma_z^2 (v_t - t)^2 + z_1^2 r_t^2, \\ \mathbb{E}[(\phi_t f_t(\mu) + \theta_t)^2] &= \sigma_z^2 \phi_t^2 v_t^2 + z_1^2 (\phi_t r_t + g_t)^2. \end{aligned}$$


After summing, simplifying and factorising, the optimality condition becomes an inequality on the moments of Z as follows:

$$T z_2 \leq z_1^2 \int_0^T ((\phi_t r_t + g_t)^2 + \varepsilon r_t^2) dt + \sigma_z^2 \int_0^T (\phi_t^2 (v_t - t)^2 + \varepsilon v_t^2) dt.$$

As $z_2 = \sigma_z^2 + z_1^2$, we get (5.13) and (5.14). \square

Thanks to this simple optimality condition, the set \mathcal{G}_l of mean field CCEs can be easily explored numerically and analytically. In Figure 1a we represent the running expected payoffs (time derivative of the payoff) of a mean field CCE, the MFC solution and the mean field NE as curves, and their total payoffs as dots. Figure 1b represents the average of the state variables at each time for the same equilibria. As one can see, the mean field CCE in the figure outperforms the mean field NE in terms of both payoff and abatement levels. Moreover, Figure 1b shows that this mean field CCE also outperforms the MFC solution in terms of average level of cumulated abatement at the end of the period, i.e. $\mathbb{E}[\mu_T] > \bar{x}_T^{MFC}$.

Implications of the optimality condition for \mathcal{G}_l can be further analysed by stating some of its analytical properties.



(a) Running expected utility and payoff of a mean field CCE, the MFC solution and the mean field NE.

(b) Average level of cumulated abatement as a function of time for this same mean field CCE, the MFC solution and the mean field NE.

Figure 1: A mean field CCE in \mathcal{G}_l bridging the gap between the mean field NE and the MFC solution. Parameter values: $T = 5, a = 2, b = 1, \varepsilon = 1, \nu_1 = 0.1, z_1 = 0.6, \sigma_z^2 = 0.06$.

Proposition 5.4. *The coefficients c_M, c_V defined in Proposition 5.3 verify the following:*

(i) $c_M < 0$,

(ii) $c_V > 0$ if and only if $\varepsilon T^2 \geq 3$.

Proof. (i) We argue by contradiction. Suppose $c_M \geq 0$. By (5.13), this is equivalent to the existence of a mean field CCE in \mathcal{G}_l so that the associated random variable Z satisfies $\sigma_Z^2 = 0$ and $z_1 > 0$. Since $\sigma_Z^2 = 0$, (λ, μ) is a mean field CCE with deterministic flow of moments $\mu_t = \nu_1 + tz_1$, for any t in $[0, T]$. By Theorem 4.5, this implies that $(\lambda, \mu) = (\hat{\alpha}, \hat{m})$. Since $\hat{m}_t = \nu_1$ for every time t , this implies that $z_1 = 0$, leading to a contradiction.

(ii) We now show that $c_V > 0$ if and only if $\varepsilon T^2 \geq 3$. Since $c_M < 0$ by point (i), condition (5.13) implies that $c_V \geq 0$ if and only if there exists a mean field CCE in \mathcal{G}_l so that the associated random variable Z satisfies $z_1 = 0$ and $\sigma_z^2 > 0$. We now identify the conditions so that such a correlated flow is a mean field CCE. In particular, it verifies $\mathbb{E}[\mu_t] = \nu_1$, for all $t \in [0, T]$. By equation (5.3), the optimal strategy of the deviating player is given by $\hat{\beta}_t = \phi_t(\nu_1 - \hat{X}_t)$, where \hat{X} is deviating player's state process. Such a correlated flow is a mean field CCE if and only if $\mathfrak{J}(\lambda, \mu) \geq \mathfrak{J}(\hat{\beta}, \mu)$, which is in turn equivalent to $\mathfrak{J}'(\lambda, \mu) \leq \mathfrak{J}'(\hat{\beta}, \mu)$, where

$$\begin{aligned}\mathfrak{J}'(\hat{\beta}, \mu) &= \frac{1}{2} \mathbb{E} \left[\int_0^T \left((\varepsilon + \phi_t^2) \mathbb{E}[(\nu_1 - \hat{X}_t)^2] + \varepsilon t^2 \sigma_z^2 \right) dt \right], \\ \mathfrak{J}'(\lambda, \mu) &= \frac{1}{2} \mathbb{E} \left[\int_0^T \left((\varepsilon + \phi_t^2) \mathbb{E}[(\mu_t - X_t)^2] + \sigma_z^2 \right) dt \right].\end{aligned}$$

By computing and comparing their dynamics, it can be shown that

$$\mathbb{E}[(\mu_t - X_t)^2] = \mathbb{E}[(\nu_1 - \hat{X}_t)^2], \quad \forall t \in [0, T].$$

Therefore, (λ, μ) is a mean field CCE if and only if $\varepsilon \frac{T^3}{3} - T \geq 0$. This allows to conclude that $c_V \geq 0$ is equivalent to $\varepsilon T^2 \geq 3$. Since c_V is null if and only if $\varepsilon = 0$ and $\varepsilon \neq 0$ by assumption, we deduce that $c_V > 0$ is equivalent to $\varepsilon T^2 \geq 3$. \square

Proposition 5.4 implies that, if the reputational cost coefficient ε and time horizon T are small enough, the only mean field CCE in \mathcal{G}_l is the mean field NE. On the contrary, when T, ε are big enough, for any expectation of Z there exists a variance level so that any correlated flow with same expectation and higher variance is a mean field CCE.

5.3 Comparison with mean field NE

We have seen above that increasing the variance of Z is a way to build mean field CCEs easily. However, increasing the variance of Z comes at the cost of lowering the odds to outperform the mean field NE, as shown in the next Proposition.

Proposition 5.5 (Outperformance over the mean field NE in \mathcal{G}_l). *A correlated flow $(\lambda, \mu) \in \mathcal{G}_l$ outperforms the mean field NE in terms of payoff if and only if*

$$Tz_1(a - b\nu_1) - (z_1^2 + \sigma_z^2) \left(b \frac{T^2}{3} + 1 \right) \geq 0 \quad (5.16)$$

Proof. This result follows directly from Proposition 5.2. The inequality is assessed in the specific case of correlated flows in \mathcal{G}_l , using the following equalities:

$$\hat{m}_t = \nu_1, \quad \mathbb{E}[\mu_t] = \nu_1 + tz_1, \quad \mathbb{E}[\mu_t^2] = \nu_1^2 + 2\nu_1tz_1 + t^2(z_1^2 + \sigma_z^2), \quad \frac{d\mu}{dt} = Z, \quad t \in [0, T]. \quad (5.17)$$

\square

The optimality and outperformance conditions for correlated flows in \mathcal{G}_l in, respectively, Proposition 5.3 and Proposition 5.5, are both expressed in terms of the first and second moments of associated variable Z . This allows us to characterize analytically a region of mean field CCEs outperforming the mean field NE in the plane (z_1, σ_z^2) .

Proposition 5.6. *Assume $\varepsilon T^2 \geq 3$. Then, a correlated flow in \mathcal{G}_l is a mean field CCE outperforming the mean field NE in terms of payoff if and only if the associated random variable Z verifies*

$$-\frac{c_M}{c_V} z_1^2 \leq \sigma_z^2 \leq z_1 \frac{3T(a - b\nu_1)}{bT^2 + 3} - z_1^2. \quad (5.18)$$

Moreover, the set of mean field CCEs outperforming the mean field NE is not reduced to the mean field NE if and only if $a - b\nu_1 > 0$.

Proof. By combining Propositions 5.1, 5.4 and 5.5, we can see that a correlated flow in \mathcal{G}_l with moments z_1, σ_z^2 for Z is a mean field CCE outperforming the mean field NE in terms of expected payoff if and only if equation (5.18) is verified. Let us denote by $f, g : \mathbb{R} \rightarrow \mathbb{R}$ respectively the left hand-side and the right-hand side of that equation as function of z_1 . They are both parabola intersecting at the point $(0, 0)$. The second derivative of f, f'' , is strictly increasing as c_V is positive according to Proposition 5.4, while g'' is strictly decreasing. Simple arguments therefore imply that the region between the two curves characterized in equation (5.18), i.e.,

$$\{(x, y) \in \mathbb{R}^2 : f(x) \leq y \leq g(x)\},$$

is not equal to the point $(0, 0)$ if and only if $g'(0) > f'(0)$. This is the case if and only if $a - b\nu_1 > 0$. If the region between the two curves was reduced to the point $(0, 0)$, the only mean field CCEs outperforming the mean field NE would verify $z_1 = \sigma_z^2 = 0$, which corresponds to the mean field NE. \square

Figure 2 represents the region of mean field CCEs outperforming the payoff of the mean field NE in the plane (z_1, σ_z^2) for the same parameters as in Figure 1. The outperformance condition parabola (“upper parabola”) is represented in red, while the optimality condition parabola (“lower parabola”) is in blue.

Proposition 5.6 shows that each of the parameters a, b and ε of the payoff plays specific roles in identifying mean field CCEs that outperform the payoff of the mean field NE. The upper parabola comes from the outperformance condition and only depends on a, b while the lower parabola comes from the optimality condition and only depends on ε . The existence of the abatement benefit leaves space for more correlated flows to outperform the free-riding equilibrium payoff, as the upper parabola increases in a and decreases in b . Moreover, Figure 3 shows that the ratio $-c_M/c_V$ and hence the lower parabola is decreasing in ε . Therefore, the reputational cost helps correlated flows to be CCEs. Indeed, with a higher reputational cost, countries have more interest in staying close to one another, and therefore the correlation device is more enforcing.

Figure 4 represents the payoffs of the mean field CCEs belonging to \mathcal{G}_l and which outperform the mean field NE in terms of payoff, i.e., verifying equation (5.18). According to this graph, using the simple and tractable class of correlated flows \mathcal{G}_l , one is able to explore a large part of the payoffs attainable by mean field CCEs in this game. Indeed, mean field CCEs payoffs get pretty close to the unattainable bound provided by the MFC solution payoff, relatively to the payoff of the mean field NE.

The mean field CCEs in \mathcal{G}_l which are optimal in terms of expected payoffs can be identified analytically.




Figure 2: Region of mean field CCEs in \mathcal{G}_l which outperform the Nash equilibrium in the plane (z_1, σ_z^2) .




Figure 3: Representation of $-c_M/c_V$ in function of ε , where ε verifies $\varepsilon T^2 \geq 3$. Parameterization: $T = 5$.

Proposition 5.7. *Assume $\varepsilon T^2 \geq 3$. Then, the expected payoff of mean field CCEs in \mathcal{G}_l is maximised by a correlated flow (λ, μ) so that the associated random variable Z satisfies*

$$z_1 = \frac{T(a - b\nu_1)}{2(1 - \frac{c_M}{c_V})(b\frac{T^2}{3} + 1)}, \quad \sigma_z^2 = -z_1^2 \frac{c_M}{c_V}. \quad (5.19)$$

Proof. We note that

$$\mathfrak{J}(\lambda, \mu) = \mathfrak{J}(\hat{\alpha}, m) + \frac{T^2}{2} z_1 (a - b\nu_1) - \frac{(\sigma_z^2 + z_1^2)}{2} \left(b\frac{T^3}{3} + T \right).$$

Therefore, $\mathfrak{J}(\lambda, \mu)$ is strictly decreasing in σ_z^2 . Moreover, since $c_V > 0$ according to Proposition 5.4, the optimality condition for correlated flows in \mathcal{G}_l of Proposition 5.3 implies that (λ, μ) is a mean field CCE if and only if

$$\sigma_z^2 \geq -z_1^2 \frac{c_M}{c_V}.$$




Figure 4: Expected payoff of mean field CCEs in \mathcal{G}_l which outperform the mean field NE in terms of payoff (i.e. verifying equation (5.18)) in the 3D space with z_1 as the x-axis, σ_z^2 as the y-axis and expected payoff as the z-axis. Parameters: same as Figure 1.

As $-c_M/c_V > 0$, for any given z_1 the mean field CCE with the highest expected payoff verifies $\sigma_z^2 = -z_1^2 c_M/c_V$. From now on, let us set σ_z^2 to this value. We get

$$\mathfrak{J}(\lambda, \mu) = \mathfrak{J}(\hat{\alpha}, m) + \frac{T^2}{2} z_1 (a - b\nu_1) - \frac{z_1^2 (1 - \frac{c_M}{c_V})}{2} \left(b \frac{T^3}{3} + T \right),$$

which is a polynomial in z_1 whose maximum point is given by z_1 as in (5.19). \square

Figure 5a shows the payoffs of mean field CCEs in \mathcal{G}_l with payoff-maximizing variance for Z , i.e. verifying $\sigma_z^2 = -z_1^2 c_M/c_V$. These payoffs are expressed as a function of z_1 , on the x -axis, and they are compared to the payoffs of the MFC solution and the mean field NE, with same parameter settings as in the other figures. Figure 5b represents the average cumulated abatement over the whole time interval for the same equilibria, in the same fashion.

One can see out of Figure 5 that for “little ambitious” mean field CCEs in \mathcal{G}_l , i.e., with relatively small z_1 , there is actually a significant increase in both abatement levels and payoffs with regards to the mean field NE. However, there is a critical value of z_1 , given by the payoff-maximising value of Proposition 5.7 and represented by the grey vertical line, after which increasing abatement comes at the cost of decreasing the payoff. This is in partial contrast with the results of [14] where a much stronger trade-off was observed. The difference is due to the presence of the reputational cost.

Characterizing a surface of mean field CCEs allows any moderator to choose the mean field CCE which corresponds to its goal, which might be to maximise expected payoff, or to consider positive externalities of abatement which are not “priced” in \mathfrak{J} , and therefore to favor high abatement over maximising payoffs.

Acknowledgements

The third author has been supported by French National Research Agency (ANR) under the program “Investissements d’Avenir” (“Investments for future”), via the FCD Labex of the Institut Louis Bachelier.




Figure 5: (a) Payoff of mean field CCEs of \mathcal{G}_l with optimal variance as a function of z_1 , and (b) their average cumulated abatement $\mathbb{E}[\mu_T]$, compared to the same quantities for the MFC solution and mean field NE. Parameter values are the same as in Figure 1.

Appendix

Proof of Proposition 3.1. We follow the approach of [30, Chapter 6]. We start by noticing that the equation for ϕ is a matrix Riccati equation, which admits a unique solution $\phi \in \mathcal{C}^1([0, T], \mathcal{S}^d)$ by Chapter 6, Theorem 7.2 therein. This implies the existence and uniqueness for ψ and θ as well as they satisfy linear ODEs.

First, thanks to Assumptions **A**(4), the cost functional \mathfrak{J}' is strictly convex and therefore has a unique minimizer. Indeed, by looking at (3.3), we have

$$\begin{aligned} & \frac{1}{2} \langle Q_t X_t, X_t \rangle + \langle \tilde{Q}_t \mathbb{E}[\mu_t] + q_t, X_t \rangle + \frac{1}{2} \langle R_t \beta_t, \beta_t \rangle + \langle S_t X_t, \beta_t \rangle + \langle \tilde{Q}_t \mathbb{E}[\mu_t] + q_t, X_t \rangle + \langle r_t, \beta_t \rangle \\ & \geq \frac{1}{2} d_1 |X_t|^2 + \frac{1}{2} d_2 |\beta_t|^2 - \sup_t |S_t| |X_t| |\beta_t| + \langle \tilde{Q}_t \mathbb{E}[\mu_t] + q_t, X_t \rangle + \langle r_t, \beta_t \rangle \\ & > \frac{1}{2} d_1 |X_t|^2 + \frac{1}{2} d_2 |\beta_t|^2 - \sqrt{d_1} \sqrt{d_2} |S_t| |X_t| |\beta_t| + \langle \tilde{Q}_t \mathbb{E}[\mu_t] + q_t, X_t \rangle + \langle r_t, \beta_t \rangle. \end{aligned}$$

This inequality and the assumption $H \geq 0$ imply that the cost functional is strictly convex and lower semicontinuous, which yields that the minimizer exists and it is unique. Observe that this holds for any $(\mathbb{E}[\mu_t])_{t \in [0, T]}$, since it appears only in the linear terms $\langle \tilde{Q}_t \mathbb{E}[\mu_t] + q_t, X_t \rangle$ and $\langle r_t, \beta_t \rangle$.

We apply the stochastic maximum principle, as in [30, Chapter 6, Proposition 5.5]. In the following, for the sake of clarity, we omit the dependence on time in all the matrices appearing in the coefficients and in the cost functions. Let $\mathcal{H}(t, x, y, \beta)$ be the reduced Hamiltonian of the system, defined as

$$\mathcal{H}(t, x, y, \beta) = \langle Ax + B\beta, y \rangle + \frac{1}{2} \langle Qx, x \rangle + \langle \tilde{Q} \mathbb{E}[\mu_t] + q, x \rangle + \langle Sx, \beta \rangle + \frac{1}{2} \langle R\beta, \beta \rangle + \langle r, \beta \rangle.$$

Then, the control $\hat{\beta} = (\hat{\beta}_t)_{t \in [0, T]}$ is optimal if and only if there exists a 4-tuple $(X, \hat{\beta}, Y, Z)$ which satisfies

$$\begin{cases} dX_t = (A_t X_t + B_t \hat{\beta}_t) dt + \sigma_t dW_t, & X_0 = \xi, \\ dY_t = - \left(A_t^\top Y_t + Q_t X_t + \tilde{Q}_t \mathbb{E}[\mu_t] + q_t + S_t^\top \hat{\beta}_t \right) dt + Z_t dW_t, & Y_T = H X_T + \tilde{H} \mathbb{E}[\mu_T], \\ B_t^\top Y_t + S_t X_t + R_t \hat{\beta}_t + r_t = 0. \end{cases} \quad (5.20)$$

We make the following ansatz on Y :

$$Y_t = \phi_t X_t + \psi_t \mathbb{E}[\mu_t] + \theta_t,$$

with ϕ , ψ and θ deterministic functions taking values in $\mathbb{R}^{d \times d}$, $\mathbb{R}^{d \times d}$ and \mathbb{R}^d respectively. Since R is invertible for every time t by assumption **A**(4), by comparing the stochastic differential of the ansatz with (5.20), we find that $Z_t = \hat{\phi}_t \sigma$ and that ϕ , ψ and θ must satisfy equations (3.4). \square

Proof of Proposition 4.1. We follow the Pontryagin maximum principle approach for MFC problems of [11, Chapter 6]. Let \mathcal{H} be the Hamiltonian of the system:

$$\begin{aligned} \mathcal{H}(t, x, y, m, \alpha) = & \langle Ax + B\alpha, y \rangle - \frac{1}{2} \langle \bar{Q}m, m \rangle + \langle L, m \rangle - \left(\frac{1}{2} \langle Qx, x \rangle + \langle \tilde{Q}m + q, x \rangle \right. \\ & \left. + \langle Sx, \alpha \rangle + \frac{1}{2} \langle R\alpha, \alpha \rangle + \langle r, \alpha \rangle \right), \quad (t, x, y, m, \alpha) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^k. \end{aligned}$$

Then, a control $\hat{\alpha} = (\hat{\alpha}_t)_{t \in [0, T]}$ is optimal if and only if there exists a 4-tuple $(X, \hat{\alpha}, Y, Z)$ which satisfies

$$\begin{cases} dX_t = (A_t X_t + B_t \hat{\alpha}_t) dt + \sigma dW_t, \\ X_0 = \xi, \\ dY_t = - \left(A_t^\top Y_t + Q_t X_t + q_t + S_t^\top \hat{\alpha}_t + (\bar{Q}_t + 2\tilde{Q}_t) \mathbb{E}[X_t] - L_t \right) dt + Z_t dW_t, \\ Y_T = H X_T + (\bar{H} + 2\tilde{H}) \mathbb{E}[X_T], \\ B_t^\top Y_t + S_t X_t + R_t \hat{\alpha}_t + r_t = 0. \end{cases} \quad (5.21)$$

Set $x_t = \mathbb{E}[X_t]$, $y_t = \mathbb{E}[Y_t]$ and $a_t = \mathbb{E}[\hat{\alpha}_t]$. Then, by taking expectation, we get the following system

$$\begin{cases} \dot{x}_t = A_t x_t + B_t a_t, & x_0 = \mathbb{E}[\xi], \\ \dot{y}_t = - \left(A_t^\top y_t + (Q_t + 2\tilde{Q}_t + \bar{Q}_t) x_t + q_t - L_t + S_t^\top a_t \right), & y_T = (H + 2\tilde{H} + \bar{H}) x_T, \\ B_t^\top y_t + S_t x_t + R_t a_t + r_t = 0. \end{cases} \quad (5.22)$$

To find a solution, we make the following ansatz on y :

$$y_t = \phi_t^{MFC} x_t + \theta_t^{MFC},$$

with ϕ^{MFC} and θ^{MFC} suitable deterministic functions taking values in $\mathbb{R}^{d \times d}$ and \mathbb{R}^d respectively. Since R is invertible for every time t by assumption **A**(4), by comparing the differential of the ansatz with (5.22), we get to equations (4.4). By [30, Chapter 6, Theorem 7.2] there exists a unique solution for the matrix Riccati equation for $\phi^{MFC} \in \mathcal{C}^1([0, T], \mathcal{S}^d)$. We note that the flow of expectations $\bar{x}^{MFC} = (\bar{x}_t^{MFC})_{t \in [0, T]}$ satisfies the differential equation (4.7b).

To prove the existence of a solution to the forward backward system (5.21), we can make the ansatz

$$Y_t = \hat{\phi}_t X_t + \bar{\psi}_t \bar{x}_t^{MFC} + \bar{\theta}_t.$$

with $\hat{\phi}$, $\bar{\psi}$ and $\bar{\theta}$ deterministic functions taking values in $\mathbb{R}^{d \times d}$, $\mathbb{R}^{d \times d}$ and \mathbb{R}^d respectively. By differentiating the ansatz, comparing it with (5.21) and using the invertibility of R for any time t , we find that $Z_t = \hat{\phi}_t \sigma$ and that $\hat{\phi}$ satisfies the same equation as ϕ , so that $\hat{\phi} = \phi$, and equations (4.6) must be satisfied by $\bar{\psi}$ and $\bar{\theta}$. \square

Proof of Proposition 4.4. We follow the Pontryagin maximum principle approach together with the fixed point argument of [11, Chapter 4]. Let \mathcal{H} be the Hamiltonian of the system:

$$\begin{aligned} \mathcal{H}(t, x, y, m, \alpha) = & \langle Ax + B\alpha, y \rangle - \frac{1}{2} \langle \bar{Q}m, m \rangle + \langle L, m \rangle - \left(\frac{1}{2} \langle Qx, x \rangle + \langle \tilde{Q}m + q, x \rangle \right. \\ & \left. + \langle Sx, \alpha \rangle + \frac{1}{2} \langle R\alpha, \alpha \rangle + \langle r, \alpha \rangle \right), \quad (t, x, y, m, \alpha) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^k. \end{aligned}$$

Then, a control $\hat{\alpha} = (\hat{\alpha}_t)_{t \in [0, T]}$ is optimal if and only if there exists a 4-tuple $(X, \hat{\alpha}, Y, Z)$ which satisfies

$$\begin{cases} dX_t = (AX_t + B\hat{\alpha}_t)dt + \sigma dW_t, & X_0 = \xi, \\ dY_t = - \left(A^\top Y_t + QX_t + \tilde{Q}\hat{m}_t + q + S^\top \hat{\alpha}_t \right) dt + Z_t dW_t, & Y_T = HX_T + \tilde{H}\hat{m}_T, \\ B^\top Y_t + SX_t + R\hat{\alpha}_t + r_t = 0. \end{cases} \quad (5.23)$$

Set $x_t = \mathbb{E}[X_t]$, $y_t = \mathbb{E}[Y_t]$ and $a_t = \mathbb{E}[\hat{\alpha}_t]$. Then, the consistency condition $\mathbb{E}[X_t] = \hat{m}_t$ for every $0 \leq t \leq T$ holds if and only if the following system

$$\begin{cases} \dot{x}_t = Ax_t + Ba_t, & x_0 = \mathbb{E}[\xi], \\ \dot{y}_t = - \left(A^\top y_t + (Q + \tilde{Q})x_t + q + S^\top a_t \right), & y_T = (H + \tilde{H})x_T, \\ B^\top y_t + Sx_t + Ra_t + r_t = 0, \end{cases} \quad (5.24)$$

admits a unique solution. We make the following ansatz on y :

$$y_t = \phi_t^{NE} x_t + \theta_t^{NE},$$

with ϕ^{NE} and θ^{NE} suitable deterministic functions taking values in $\mathbb{R}^{d \times d}$ and \mathbb{R}^d respectively. Since R is invertible for every time t by assumption **A**(4), by comparing the differential of the ansatz with (5.24), we get to equations (4.17). We note that the flow of moments $\hat{m} = (\hat{m}_t)_{t \in [0, T]}$ satisfies the differential equation (4.20a).

The last step it to prove the existence of a solution to the forward backward system (5.23). We make the ansatz

$$Y_t = \hat{\phi}_t X_t + \hat{\psi}_t \hat{m}_t + \theta_t^{\hat{m}}.$$

with $\hat{\phi}$, $\hat{\psi}$ and $\theta^{\hat{m}}$ deterministic functions taking values in $\mathbb{R}^{d \times d}$, $\mathbb{R}^{d \times d}$ and \mathbb{R}^d respectively. By the same reasoning of Proposition 3.1, we find that $Z_t = \hat{\phi}_t \sigma$, that $\hat{\phi}$ and $\hat{\psi}$ satisfy the same equations as ϕ and ψ , so that $\hat{\phi} = \phi$ and $\hat{\psi} = \psi$, and that equation (4.19) must be satisfied by $\theta^{\hat{m}}$. \square

Proof of Theorem 4.6. By using $(\phi, \psi, \theta^{\hat{m}})$, we write the dynamics of the state process X^{NE} as

$$\begin{aligned} dX_t^{NE} = & \left((A_t - B_t R_t^{-1} (B_t^\top \phi_t + S)) X_t^{NE} - B_t R_t^{-1} (B_t^\top \psi_t \hat{m}_t + B_t^\top \theta_t^{\hat{m}} + r_t) \right) dt + \sigma_t dW_t \\ = & \left((A_t - B_t \Phi_t) X_t^{NE} - B_t (\Psi_t \hat{m}_t + \Theta_t^{\hat{m}}) \right) dt + \sigma_t dW_t, \end{aligned}$$

with Φ and Ψ defined by (3.10) and $\Theta^{\hat{m}}$ by (4.22). We remark that $\theta^{\hat{m}}$ and thus $\Theta^{\hat{m}}$ depend on \hat{m} through its time derivative $d\hat{m}/dt$. Let $f^{\hat{m}, \mu} = (f_t^{\hat{m}, \mu})_{t \in [0, T]}$ be the solution of

$$\dot{f}_t^{\hat{m}, \mu} = (A_t - B_t \Phi_t) f_t^{\hat{m}, \mu} + B_t (\Psi_t (\hat{m}_t - \mu_t) + \Theta_t^{\hat{m}} - \delta_t), \quad f_0^{\hat{m}, \mu} = 0.$$

By Itô's formula, we have that

$$f_t^{\hat{m}, \mu} = X_t - X_t^{NE}.$$

Since $f(\mu)$ is $\sigma(\mu)$ -measurable and X^{NE} is \mathbb{F}^1 -progressively measurable, we have both that X^{NE} and $f^{\hat{m}, \mu}$ are independent and that

$$f_t^{\hat{m}, \mu} = \mathbb{E}[f_t^{\hat{m}, \mu} | \mu] = \mathbb{E}[X_t - X_t^{NE} | \mu] = \mu_t - \hat{m}_t, \quad (5.25)$$

by consistency condition. Since it holds

$$\begin{aligned} \mathfrak{J}(\lambda, \mu) - \mathfrak{J}(\hat{\alpha}, \hat{m}) &= \int_0^T \left(\frac{1}{2} \langle \bar{Q} \hat{m}_t, \hat{m}_t \rangle - \frac{1}{2} \mathbb{E}[\langle \bar{Q} \mu_t, \mu_t \rangle] + \langle L_t, \mathbb{E}[\mu_t] - \hat{m}_t \rangle \right) dt \\ &+ \frac{1}{2} \langle \bar{H} \hat{m}_T, \hat{m}_T \rangle - \frac{1}{2} \mathbb{E}[\langle \bar{H} \mu_T, \mu_T \rangle] + \mathfrak{J}'(\hat{\alpha}, \hat{m}) - \mathfrak{J}'(\lambda, \mu), \end{aligned}$$

we focus on the difference $\mathfrak{J}'(\hat{\alpha}, \hat{m}) - \mathfrak{J}'(\lambda, \mu)$. In a very similar way as in the proof of Theorem 3.3 we obtain

$$\begin{aligned} \mathfrak{J}'(\hat{\alpha}, \hat{m}) - \mathfrak{J}'(\lambda, \mu) &= \int_0^T \left(\frac{1}{2} (\langle M_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle M_t(\hat{m}_t + f_t^{\hat{m}, \mu}), \hat{m}_t + f_t^{\hat{m}, \mu} \rangle]) \right. \\ &+ \frac{1}{2} (\langle G_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle G_t \mu_t, \mu_t \rangle]) + \langle N_t \hat{m}_t, \hat{m}_t - \mathbb{E}[\mu_t] \rangle - \mathbb{E}[\langle N_t f_t^{\hat{m}, \mu}, \mu_t \rangle] \\ &- \langle q_t - \Phi_t^\top r_t, \mathbb{E}[f_t^{\hat{m}, \mu}] \rangle - \langle \Psi_t^\top r_t, \hat{m}_t - \mathbb{E}[\mu_t] \rangle + \langle (R_t \Phi_t - S_t) \hat{m}_t, (\Theta_t^{\hat{m}} - R_t^{-1} \mathbb{E}[\delta_t]) \rangle \\ &- \mathbb{E}[\langle (R_t \Phi_t - S_t) f_t^{\hat{m}, \mu}, R_t^{-1} \delta_t \rangle] - \mathbb{E}[\langle R_t \Psi_t \mu_t, R_t^{-1} \delta_t \rangle] \\ &+ \langle R_t \Psi_t \hat{m}_t, \Theta_t^{\hat{m}} \rangle + \frac{1}{2} (\langle R_t \Theta_t^{\hat{m}}, \Theta_t^{\hat{m}} \rangle - \mathbb{E}[\langle R_t^{-1} \delta_t, \delta_t \rangle]) - \langle r_t, \Theta_t^{\hat{m}} - R_t^{-1} \mathbb{E}[\delta_t] \rangle \Big) dt \\ &+ \frac{1}{2} \langle H \hat{m}_T, \hat{m}_T \rangle - \frac{1}{2} \langle H(\hat{m}_T + \hat{f}_T(\mu)), \hat{m}_T + \hat{f}_T(\mu) \rangle + \langle \tilde{H} \hat{m}_T, \hat{m}_T - \mathbb{E}[\mu_T] \rangle - \mathbb{E}[\langle \tilde{H} \hat{f}_T(\mu), \mu_T \rangle]. \end{aligned}$$

Finally, we observe that, by using (5.25), we have

$$\begin{aligned} \langle M_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle M_t(\hat{m}_t + f_t^{\hat{m}, \mu}), \hat{m}_t + f_t^{\hat{m}, \mu} \rangle] &= \langle M_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle M_t \mu_t, \mu_t \rangle], \\ \langle N_t \hat{m}_t, \hat{m}_t - \mathbb{E}[\mu_t] \rangle - \mathbb{E}[\langle N_t f_t^{\hat{m}, \mu}, \mu_t \rangle] &= \langle N_t \hat{m}_t, \hat{m}_t \rangle - \mathbb{E}[\langle N_t \mu_t, \mu_t \rangle], \\ \langle (R_t \Phi_t - S_t) \hat{m}_t, (\Theta_t^{\hat{m}} - R_t^{-1} \mathbb{E}[\delta_t]) \rangle - \mathbb{E}[\langle (R_t \Phi_t - S_t) f_t^{\hat{m}, \mu}, R_t^{-1} \delta_t \rangle] \\ &= \langle (R_t \Phi_t - S_t) \hat{m}_t, \Theta_t^{\hat{m}} \rangle - \mathbb{E}[\langle (R_t \Phi_t - S_t) \mu_t, R_t^{-1} \delta_t \rangle], \\ \langle q_t - \Phi_t^\top r_t, \mathbb{E}[f_t^{\hat{m}, \mu}] \rangle + \langle \Psi_t^\top r_t, \hat{m}_t - \mathbb{E}[\mu_t] \rangle &= \langle q_t - (\Phi_t + \Psi_t)^\top r_t, \mathbb{E}[\mu_t] - \hat{m}_t \rangle. \end{aligned}$$

By using these identities together with (3.15), we get to (4.23). \square

References

- [1] R. J. Aumann. Subjectivity and correlation in randomized strategies. *J. Math. Econom.*, 1(1):67–96, 1974.
- [2] R. J. Aumann. Correlated equilibrium as an expression of Bayesian rationality. *Econometrica*, 55(1):1–18, 1987.
- [3] S. Barrett. Self-enforcing international environmental agreements. *Oxford economic papers*, 46(Supplement_1):878–894, 1994.

- [4] O. Bonesini. *Four essays in between Probability Theory and Financial Mathematics*. PhD thesis, Università degli Studi di Padova, Padua, Italy, 2023.
- [5] O. Bonesini, L. Campi, and M. Fischer. Correlated equilibria for mean field games with progressive strategies, 2022.
- [6] P. Brémaud and M. Yor. Changes of filtrations and of probability measures. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 45(4):269–295, 1978.
- [7] L. Campi, F. Cannerozzi, and M. Fischer. Coarse correlated equilibria for continuous time mean field games in open loop strategies, 2023.
- [8] L. Campi and M. Fischer. N -player games and mean-field games with absorption. *Ann. Appl. Probab.*, 28(4):2188–2242, 2018.
- [9] L. Campi and M. Fischer. Correlated equilibria and mean field games: a simple model. *Math. Oper. Res.*, 47(3):2240–2259, 2022.
- [10] R. Carmona and F. Delarue. Probabilistic analysis of mean-field games. *SIAM J. Control Optim.*, 51(4):2705–2734, 2013.
- [11] R. Carmona and F. Delarue. *Probabilistic Theory of Mean Field Games with Applications I*. Springer, 2018.
- [12] R. Carmona, F. Delarue, and A. Lachapelle. Control of McKean-Vlasov dynamics versus mean field games. *Math. Financ. Econ.*, 7(2):131–166, 2013.
- [13] R. Carmona, C. V. Graves, and Z. Tan. Price of anarchy for mean field games. In *CEMRACS 2017—numerical methods for stochastic models: control, uncertainty quantification, mean-field*, volume 65 of *ESAIM Proc. Surveys*, pages 349–383. EDP Sci., Les Ulis, 2019.
- [14] T. Dokka, H. Moulin, I. Ray, and S. SenGupta. Equilibrium design in an n -player quadratic game. *Review of economic design*, 2022.
- [15] I. Gilboa and E. Zemel. Nash and correlated equilibria: Some complexity considerations. *Games and Economic Behavior*, 1(1):80–93, 1989.
- [16] P. J. Graber. Linear quadratic mean field type control and mean field games with common noise, with application to production of an exhaustible resource. *Appl. Math. Optim.*, 74(3):459–486, 2016.
- [17] C. Grüning and W. Peters. Can justice and fairness enlarge international environmental agreements? *Games*, 1(2):137–158, 2010.
- [18] J. Hannan. Approximation to Bayes risk in repeated play. In *Contributions to the theory of games, vol. 3*, Annals of Mathematics Studies, no. 39, pages 97–139. Princeton University Press, Princeton, N.J., 1957.
- [19] S. Hart and A. Mas-Colell. Regret-based continuous-time dynamics. *Games Econom. Behav.*, 45(2):375–394, 2003. Special issue in honor of Robert W. Rosenthal.
- [20] M. Huang, R. P. Malhamé, and P. E. Caines. Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Commun. Inf. Syst.*, 6(3):221–251, 2006.

- [21] D. Lacker. On the convergence of closed-loop nash equilibria to the mean field game limit. *The Annals of Applied Probability*, 30(4):1693–1761, 2020.
- [22] D. Lacker and L. Le Flem. Closed-loop convergence for mean field games with common noise. *Ann. Appl. Probab.*, 33(4):2681–2733, 2023.
- [23] J.-M. Lasry and P.-L. Lions. Mean field games. *Jpn. J. Math.*, 2(1):229–260, 2007.
- [24] H. Moulin, I. Ray, and S. S. Gupta. Improving Nash by coarse correlation. *Journal of Economic Theory*, 150:852–865, 2014.
- [25] H. Moulin and J.-P. Vial. Strategically zero-sum games: the class of games whose completely mixed equilibria cannot be improved upon. *Internat. J. Game Theory*, 7(3-4):201–221, 1978.
- [26] P. Muller, R. Elie, M. Rowland, M. Lauriere, J. Perolat, S. Perrin, M. Geist, G. Piliouras, O. Pietquin, and K. Tuyls. Learning Correlated Equilibria in Mean-Field Games, 2022.
- [27] P. Muller, M. Rowland, R. Elie, G. Piliouras, J. Perolat, M. Lauriere, R. Marinier, O. Pietquin, and K. Tuyls. Learning equilibria in mean-field games: Introducing mean-field psro, 2021.
- [28] A. Neyman. Correlated equilibrium and potential games. *International Journal of Game Theory*, 26:223–227, 1997.
- [29] T. Roughgarden. *Twenty Lectures on Algorithmic Game Theory*. Cambridge University Press, 2016.
- [30] J. Yong and X. Y. Zhou. *Stochastic controls*, volume 43 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 1999. Hamiltonian systems and HJB equations.