Quantumness and quantum to classical transition in the generalized Rabi model

Wei-Feng Zhuang,^{1,2,*} Yun-Tong Yang,^{3,4} Hong-Gang Luo,^{3,4,5,†} Ming Gong,^{1,6,7,‡} and Guang-Can Guo^{1,6,7}

¹CAS Key Laboratory of Quantum Information, University of Science and Technology of China, Hefei, 230026, China

²Beijing Academy of Quantum Information Sciences, Beijing 100193, China

³School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China

⁴Lanzhou Center for Theoretical Physics & Key Laboratory of Theoretical

Physics of Gansu Province, Lanzhou University, Lanzhou 730000, China

⁶Synergetic Innovation Center of Quantum Information and Quantum Physics,

University of Science and Technology of China, Hefei, Anhui 230026, China

⁷CAS Center For Excellence in Quantum Information and Quantum Physics, University of Science and Technology of China, Hefei, Anhui 230026, China

 $(D_{2}+1, N_{2}) = 14, 2022)$

(Dated: November 14, 2023)

The quantum to classical transition (QCT) is one of the central mysteries in quantum physics. This process is generally interpreted as state collapse from measurement or decoherence from interacting with the environment. Here we define the quantumness of a Hamiltonian by the free energy difference between its quantum and classical descriptions, which vanishes during QCT. We apply this criterion to the many-body Rabi model and study its scaling law across the phase transition, finding that not only the temperature and Planck constant, but also all the model parameters are important for this transition. We show that the Jaynes-Cummings and anti Jaynes-Cummings models exhibit greater quantumness than the Rabi model. Moreover, we show that the rotating wave and anti-rotating wave terms in this model have opposite quantumness in QCT. We demonstrate that the quantumness may be enhanced or suppressed at the critical point. Finally, we estimate the quantumness of the Rabi model in current trapped ion experiments. The quantumness provides an important tool to characterize the QCT in a vast number of many-body models.

Quantum and classical physics differ fundamentally. In quantum mechanics, particles are described as waves, exhibiting phenomena such as entanglement and superposition. In contrast, classical physics precludes these characteristics, with well-defined mass points satisfying the Newton's laws. The quantum to classical transition (QCT) is explained through the orthodox Copenhagen interpretation of wave function collapse from measurement [1-4], which is then understood by the quantum decoherence theory [5-14]. The QCT is a rather subtle issue. While in principle no such boundary exists, because the bulk metals and superconductors in solids and the neutron star in astrophysics should be described by quantum mechanics; however, there are also abundant evidences that when the system temperature is high enough, or the Planck constant becomes negligible, the quantum nature may disappear. Furthermore, the importance of system size can be evidenced from macroscopic Schrödinger cat [6, 15–18]. Therefore, these factors should be among the most crucial factors for this QCT. Unfortunately, a quantitative characterization of this transition is still lacking.

Definition of quantumness: Imagine a Hamiltonian $\mathcal{H}(x,p)$, where x and p are coordinate and momentum operators, respectively. In classical physics, xp = px are the same; yet in quantum physics, they are different, with $[x,p] = i\hbar$, where \hbar is the Planck constant divided by 2π . Obviously, in quantum mechanics, x^2 and $p^{-1}xp^2xp^{-1}$ are different. Thus, when we say that a system exhibits quantum behaviors, we mean that it possesses some fea-

tures quantitatively different from the classical counterpart. Since in thermodynamics all observations can be determined by the partition function, we expect their difference can be reflected from their free energies. When these two descriptions yield the same free energy, they can not be distinguished from measurements. Based on this intuitive picture, we define the quantumness as

$$\Delta_{\rm QC} = F_{\rm Q} - F_{\rm C},\tag{1}$$

where $F_{\rm Q}$ and $F_{\rm C}$ are the free energies using quantum and classical mechanics. The QCT is exact when $\Delta_{\rm QC} = 0$, and the nature of quantum vanishes. This work examines this quality in the many-body Rabi model, which hosts quantum phase transition. Strikingly, we show that the critical point with significant quantum (or classical) fluctuation is not necessarily associated with maximal quantumness. Finally, we will estimate the role of quantumness in the current trapped ion experiments.

We can gain some insight into this problem by investigating some simple models. (A) A free particle model with $\mathcal{H} = p^2/2m$, where *m* is the particle mass. The partition functions are $Z_{\rm Q} = Z_{\rm C} = \int dx dp/(2\pi\hbar)e^{-\beta\mathcal{H}} = V\sqrt{2m\pi/\beta}$, where *V* is the real-space volume, and $\beta = 1/k_BT$, with k_B is the Boltzmann constant and *T* is the temperature. Hence $\Delta_{\rm QC} = 0$. (B) A quantum oscillator model with $\mathcal{H} = p^2/2m + m\omega^2 x^2/2$, with eigenvalues $E_n = \hbar\omega(n+1/2)$. We have $Z_{\rm Q} = \sum_{n=0}^{\infty} e^{-\beta\hbar\omega(n+1/2)}$, and $Z_{\rm C} = \int dx dp/(2\pi\hbar)e^{-\beta\mathcal{H}}$, yielding

$$\Delta_{\rm QC} = \frac{1}{24} (\hbar\omega)^2 \beta - \frac{1}{2880} (\hbar\omega)^4 \beta^3 + \cdots, \qquad (2)$$

⁵Beijing Computational Science Research Center, Beijing 100084, China



FIG. 1. (a) Energy levels of the generalized Rabi model, in which the solid and dashed lines correspond to coupling induced by rotating wave term (g_1) and anti-rotating wave term (g_2) , respectively. When $g_2 = 0$ (JC model) or $g_1 = 0$ (aJC model), this model can be solved exactly; see discussions in (I) and (II). (b) Phase transition in this model characterized by energy gaps $\delta E_n = E_n - E_0$ by tuning of g_i . (c) Dispersion of energy levels in the normal phase $(g < g_c)$, critical point (at g_c) and superradiant phase $(g > g_c)$ when $\omega/\Omega \to 0$.

when $\beta \hbar \omega \ll 1$, which is always a positive value. When $\omega = 0$, it reduces to model (A). Thus when $k_B T \gg \hbar \omega$, the quantization effect is not important, and QCT happens. In the presence of many-body interaction, the level spacing is increased and the quantumness $\Delta_{\rm QC}$ will also be increased in accordingly. (C) QCT from their statistics. One can find that the crossover between Bose-Einstein and Fermi-Dirac distributions and the Boltzmann-Maxwell distribution can be reached when $\beta \mu \ll -1$ (μ is the chemical potential). The last two models show that the system parameters are also important for QCT.

Quantumness in the generalized Rabi model: With these results in mind, we examine the physics in the generalized Rabi model [19–27]

$$\mathcal{H} = \hbar \omega a^{\dagger} a + \frac{\hbar \Omega}{2} \sigma^{z} + \hbar g_{1} a \sigma^{\dagger} + \hbar g_{2} a \sigma^{-} + \text{h.c.}, \quad (3)$$

where ω is the harmonic oscillator frequency, Ω is the spin level spacing, and g_1 , g_2 are the couplings of rotating and anti-rotating wave terms, respectively. This model has been intensively studied using trapped ions [28–35], superconducting qubits [36–39], quantum dots [40–42], and ultracold atoms [43–52], *etc.* When $g_2 = 0$ and $g_1 = 0$, it reduces to the Jaynes-Cummings (JC) model and anti-Jaynes-Cummings (aJC) model, respec-



FIG. 2. Verification of quantumness in Eq. 4. (a) JC model with $g_2 = 0$, $g_1 = 0.3$, 0.7, 1.5, with $\beta = 2$; (b) aJC model with $g_1 = 0$, $g_2 = 0.3$, 0.7, 1.5, with $\beta = 2$; (c) Rabi model with g = 0.6 (normal phase) and (d) g = 1.2 (superradiant phase), with $\beta = 5$ and $\eta = g_1/g_2$. In the Rabi model, A = 0, hence $x\Delta_{\rm QC}$ is independent of x. In all figures, we set $g_c = 1$.

tively. Recent investigations show that when $\omega/\Omega \rightarrow 0$, this model exhibits a second-order phase transition from a normal phase to a superradiant phase [29, 38, 45, 53–55]. In contrast to the models (A - C), Eq. 3 enables us to study the quantumness across the critical point. Since the rotating and anti-rotating wave terms are ubiquitous, this model enables us to explore their individual role on quantumness.

(I). JC model with $g_1 \neq 0$ and $g_2 = 0$. Let $q = g_1/g_c$, $\omega = g_{\rm c}/\sqrt{x}$, and $\Omega = g_{\rm c}\sqrt{x}$, where $x = \Omega/\omega$, such that $\omega \Omega = g_c^2$ is fixed during the variation of the parameters. This model can be solved analytically because the whole Hilbert space can be decoupled into small subspaces (Fig. 1 (a)), with eigenvalues $E_n^{\pm} = \hbar(n+1)\omega \pm \hbar \sqrt{g_1^2(n+1) + \frac{(\omega-\Omega)^2}{4}}$ for $n \ge 0$. The configuration in Fig. 1 (a) is of particular importance in the limit $\Omega \ll k_B T$, where only the lower branch E_n^- is significant. The phase transition of this model can be manifested from the vanished energy gap, as shown in Fig. 1 (b); and the corresponding spectra are shown in Fig. 1 (c). In the normal phase, we expect $E_n^- \propto n$; and in the superradiant phase, we expect $E_n^- \sim (n - n_c)^2$. In the phase boundary, we have $n_c = 0$ and $E_n^- \propto n^2$. With this feature, the summation of n can be generalized from zero to infinity (see the filled regime in Fig. 1 (c)). Using the Euler-Maclaurin formula, we can obtain the free energy in these two phases. Meanwhile, the classical free energy is obtained by defining

TABLE I. Summarized free energies in the normal phase (n) and superradiant (sr) phases. In the first column, JC, aJC refer to Jaynes-Cummings model and anti-Jaynes-Cummings model, respectively. The free energies are obtained from the partition functions $Z_{\rm C} = e^{-\beta F_{\rm C}}$ and $Z_{\rm Q} = e^{-\beta F_{\rm Q}}$ using classical and quantum treatments, and in the classical model, JC and aJC have the same free energy, because in the classical level these models are the same (upon a transformation $a \to a^*$).

Model (n/sr)	Quantum free energy $F_{\rm Q}$	Classical free energy $F_{\rm C}$
JC (n)	$\left -\frac{\hbar g_c \sqrt{x}}{2} - \frac{1}{\beta} \ln \left(\frac{\sqrt{x}}{\hbar g_c \beta (1-q^2)} \right) + \frac{\hbar g_c q^2}{2\sqrt{x}} + \frac{\hbar^2 \beta^2 g_c^2 (q^2-1)^3 + 24q^2}{24\beta (q^2-1)x} \right $	$\left -\frac{\hbar g_c \sqrt{x}}{1-\frac{1}{2}} \ln \left(\frac{\sqrt{x}}{1-\frac{1}{2}}\right)\right $
aJC (n)	$-\frac{\hbar g_c \sqrt{x}}{2} - \frac{1}{\beta} \ln \left(\frac{\sqrt{x}}{\hbar \beta g_c (1-q^2)} \right) - \frac{\hbar g_c q^2}{2\sqrt{x}} + \frac{-\hbar^2 \beta^2 g_c^2 (q^2-1)^3 + 24q^2 (2q^2+1)}{24x\beta(1-q^2)} $	$2 \beta \left(ng_{c}\beta(1-q^{2}) \right)$
JC (sr)	$\left -\frac{\hbar g_c(x(q^4x + x - 2) + 1)}{4q^2 x^{3/2}} - \frac{1}{\beta} \ln \left(q \sqrt{\frac{\pi}{\hbar \beta g_c}} x^{3/4} \right) \right $	$\left -\frac{(1+q^4)\hbar g_c}{4a^2} \sqrt{x} - \frac{1}{\beta} \ln \left(q_{\lambda} \sqrt{\frac{\pi}{\beta\hbar a}} x^{3/4} \right) \right $
aJC (sr)	$\left -\frac{\hbar g_{\rm c}(x(q^4x+x+2)+1)}{4q^2x^{3/2}} - \frac{1}{\beta}\ln\!\left(q\sqrt{\frac{\pi}{\hbar\beta g_{\rm c}}}x^{3/4}\right) \right.$	$4q$ · β ($\sqrt{\rho ng_c}$)



FIG. 3. Coefficients of A and B in these three models. The red lines in (a) - (d) are calcualted by numerial fitting of $\Delta_{\rm QC}$ at $\beta = 20, 80$ for various g_1 and g_2 , and the dashed lines are these by the analytical expressions in Eq. 5 and Eq. 6. The other parameters are: (a - b) $g_2 = 0$ for JC model; (c - d) $g_1 = 0$ for the aJC model; and (e -f) for the generalized Rabi model for $g < g_c$ and $g > g_c$. $\eta = 0$ and $\eta = 1$ correspond to the analytic results of JC and aJC model. In (e) the two curves are identical to the empirical formula of Eq. 7.

 $a = \sqrt{\omega/2\hbar}(\hat{x} + i\hat{p}/\omega)$ and $a^{\dagger} = a^*$, which is commonly used to analyze its phase transition in the limit $\omega/\Omega \to 0$ (see Eq. 2) [55–57]. Then the classical partition function is given by $Z_{\rm C} = \sum_{s=\pm} \int dx dp/(2\pi\hbar) e^{-\beta E_s(a,a^*)}$, in which $E_{\pm} = \hbar \omega |a|^2 \pm \hbar \sqrt{(a^*g_1 + ag_2)(ag_1 + a^*g_2) + \Omega^2}$. We can solve this model using the saddle point method, which is almost exact in the low temperature regime. The



FIG. 4. Quantumness $\Delta_{\rm QC}$ near the critical point in the generalized Rabi model. (a) $\eta = 0$ (for the JC model); (b) $\eta = 0.5$ (for the ideal Rabi model); (c) $\eta = 0.75$ (for the generalized Rabi model); (d) $\eta = 1$ (for the aJC model). In all figures, we have used $g_c = 1$ and $\beta = 5$.

results of $F_{\rm Q}$ and $F_{\rm C}$ for these phases are summarized in Table I, yielding

$$\Delta_{\rm QC} = \frac{A}{\sqrt{x}} + \frac{B}{x} + \cdots, \quad x = \frac{\Omega}{\omega}.$$
 (4)

If we write $\Delta_{\rm QC}(\omega) = \Delta'_{\rm QC}(0)\omega + \Delta''_{\rm QC}(0)\omega^2/2$, then $A \propto \Delta'_{\rm QC}(0)$ and $B \propto \Delta''_{\rm QC}(0)$. Thus QCT happens when $x \to \infty$. We find analytically that

(II). aJC model with $g_2 \neq 0$ and $g_1 = 0$. We define $q = g_2/g_c$, while the other parameters are the same as (I). This model can also be calculated exactly using $E_n^{\pm} = (n + \frac{1}{2})\hbar\omega \pm \frac{1}{2}(\hbar\omega - \sqrt{4\hbar^2g^2(n+1) + \hbar^2(\omega+\Omega)^2})$ for $n \geq -1$, which is slightly different from that in the JC model. The results are summarized in Table I, with quantumness the same as Eq. 4. We have

$$A = \begin{cases} -\frac{\hbar g_c q^2}{2} \\ -\frac{\hbar g_c}{2q^2} \\ 0 \end{cases}, B = \begin{cases} \frac{\hbar^2 \beta^2 g_c^2 (q^2 - 1)^3 - 24(2q^2 + q^4)}{24\beta(q^2 - 1)} & q < 1 \\ 0 & q > 1 \end{cases}.$$
(6)

The classical energies are the same (see Table I). Physically, quantumness is fully determined the discrete nature of quantum mechanics. For example, when $\omega \ll \Omega$, we have $E_n^-(\omega) = E_n^-(0) + \kappa \omega$, where $\kappa = (dE_n^-(\omega)/d\omega)$. When $\kappa \neq 0$, we naturally have the A term. Therefore, when all these terms disappear, quantumness naturally is vanishes, indicating QCT.

The results in Eqs. 5 - 6 already reveal some important and sutble difference between the JC and aJC models. The scaling law of Eq. 4 for these two cases are shown in Fig. 2 (a) and (b); and the corresponding coefficients of A and B are shown in Fig. 3 (a) to (d). In both models, the term A is always important in the whole phase regime. Especially, A will takes on a maximal value at the phase boundary. We find that A in the JC and aJC models take opposite values, showing that the signs of the quantumness in these two models are different. The negative quantumness is in stark contrast to that in the harmonic oscillator (Eq. 2). Besides, we find that the coefficient B is only important in the normal phase, while in the supperradiant phase, it becomes vanished.

(III). Generalized Rabi model with $g_1g_2 \neq 0$. Let $\eta = g_1/(g_1+g_2)$, the phase transition happens at $(g_1+g_2)^2 = \omega \Omega$. We only need to consider the regime with $g_i \geq 0$ [58]. When $\eta = 1/2$, it yields the Rabi model. This model can not be solved analytically, and we solve this model using numerical method. We find that the quantumness can still be written in the general form as Eq. 4. Our numerical results of quantumness are presented in Fig. 2 (c) - (d) and the extracted coefficients A and B are shown in Fig. 3 (e) and (f). Strikingly, we find that when $\eta = 1/2$, the quantumness is only given by the B term, with A = 0. Thus due to the counteract effect of the JC and aJC interactions, the Rabi model has the smallest quantumness. From Fig. 3 (e) we have

$$A(\eta) = A(0)(1 - \eta) + A(1)\eta,$$
(7)

where A(0) = -A(1) are the coefficients in the aJC and JC models, respectively. This relation naturally explains the vanished A in the Rabi model.

(IV). Quantumness and phase transition: It is natural to explore the quantumness across the critical point, which has the strongest fluctuations in both classical and quantum mechanics. However, their difference is not necessary to be increased. From the sign of A, we see that the contribution of JC and aJC are opposite, which can yield totally different behaviors. The numerical results for these models are presented in Fig. 4. We find that with the increase of g, the quantumness is enhanced in the JC and aJC models; however, in the Rabi model, it is suppressed near the critical point. These results suggest that the quantumness is an intrinsic property of the Hamiltonian, which can be increased or decreased by the critical fluctuations. As a result, it is only in this limit $\Delta_{\rm QC} = 0$ that the classical phase transition can happen in the general Rabi model [55].



FIG. 5. QCT in experimental Rabi model with trapped ions [59]. Here we have used $\omega = 2\pi \times 4.0$ kHz, $\Omega = 2\pi \times 100.0$ kHz, which corresponds to a ratio x = 25, and the critical point is at $g_{\rm c} = 2\pi \times 20.0$ kHz, showing that $\Delta_{\rm QC}/F_{\rm C} \sim 1\%$.

Quantumness in experiment with trapped ions: Finally, we discuss the relevance of our results in the current experiments with trapped ions implemented in the experiments in Duan group [59]. In this experiment with Rabi model, $\omega = 2\pi \times 4.0$ kHz, $\Omega = 2\pi \times 100.0$ kHz, x = 25, and the critical point is at $g_c = 2\pi \times 20.0$ kHz (at T = 0). By varying of the coupling g from 0 to 14 kHz (1.4 g_c), we estimate $F_{\rm Q}, F_{\rm C}$ and $\Delta_{\rm QC}$ in these experiments in Fig. 5 , showing that in the current setup, the quantumness is about 1% of the total classical energy. This value is small, yet nonzero, thus in experiment only a smooth crossover between the two phases can be demonstrated. We find that to observe a relative sharp transition, x must be increased to 100, yielding $\Delta_{\rm QC}/F_{\rm C}$ to be further reduced by one order of magnitude since $\Delta_{\rm QC}/F_{\rm C} \sim x^{-3/2}$, for A = 0 and $F_{\rm C} \propto \sqrt{x}$; see Table I.

To conclude, we define the quantumness Δ_{QC} to quantitatively characterize the QCT, and examine this quality in the generalized Rabi model. This interpretation does not rely on the state collapse, thus is not a feature of some particular states as discussed in quantum information science [9, 14, 60], instead, it is a feature of the Hamiltonian in thermal equilibrium. We show explicitly that not only the temperature and Planck constant, but also the system parameters are important for the QCT. This quantity can be applied to study the QCT in a broad range of many-body physical models, and clarify the roles of quantumness on observations and phase transitions, which could help to resolve the long-sought mysterious boundary between quantum and classical worlds. From this perspective, we expect this quantity to be important for these models which can not be described approximately by classical mechanics. It also has potential to be unified with the decoherence theory for QCT [5-14].

Acknowledgements: This work is supported by the National Natural Science Foundation of China (NSFC) with No. 11774328, No. 11834005, No. 12247101, the Innovation Program for Quantum Science and Technology (No. 2021ZD0301200 and No. 2021ZD0301500), and Na-

tional Key Research and Development Program of China (Grant No. 2022YFA1402704).

- * zhuangwf@baqis.ac.cn
- [†] luohg@lzu.edu.cn
- [‡] gongm@ustc.edu.cn
- Henry Pierce Stapp, "The Copenhagen interpretation," American Journal of Physics 40, 1098–1116 (1972).
- [2] Angelo Bassi, Kinjalk Lochan, Seema Satin, Tejinder P Singh, and Hendrik Ulbricht, "Models of wave-function collapse, underlying theories, and experimental tests," Rev. Mod. Phys. 85, 471 (2013).
- [3] John G Cramer, "The transactional interpretation of quantum mechanics," Rev. Mod. Phys. 58, 647 (1986).
- [4] Kristian Camilleri and Maximilian Schlosshauer, "Niels Bohr as philosopher of experiment: Does decoherence theory challenge Bohrś doctrine of classical concepts?" Stud. Hist. Philos. Sci. Part B: Stud. Hist. Philos. Mod. Phys. 49, 73–83 (2015).
- [5] Wojciech H Zurek, "Decoherence and the transition from quantum to classical," Phys. today 44, 36–44 (1991).
- [6] Serge Haroche, "Entanglement, decoherence and the quantum/classical boundary," Phys. today 51, 36–42 (1998).
- [7] D Zeh, "On the interpretation of measurement in quantum theory," Found. Phys. 1, 69–76 (1970).
- [8] Juan Pablo Paz and Wojciech Hubert Zurek, "Environment-induced decoherence and the transition from quantum to classical," in *Fundamentals of quantum information: quantum computation, communication, decoherence and all that* (Springer, 2002) pp. 77–148.
- Maximilian Schlosshauer, "Quantum decoherence," Phys. Rep. 831, 1–57 (2019).
- [10] H Dieter Zeh, "On the interpretation of measurement in quantum theory," Found. Phys. 1, 69–76 (1970).
- Wojciech H Zurek, "Pointer basis of quantum apparatus: Into what mixture does the wave packet collapse?" Phys. Rev. D 24, 1516 (1981).
- [12] Maximilian Schlosshauer, "Decoherence, the measurement problem, and interpretations of quantum mechanics," Rev. Mod. Phys. **76**, 1267 (2005).
- [13] Wojciech H Zurek, "Environment-induced superselection rules," Phys. Rev. D 26, 1862 (1982).
- [14] Wojciech Hubert Zurek, "Decoherence, einselection, and the quantum origins of the classical," Rev. Mod. Phys. 75, 715 (2003).
- [15] Wui Seng Leong, Mingjie Xin, Zilong Chen, Shijie Chai, Yu Wang, and Shau-Yu Lan, "Large array of schrödinger cat states facilitated by an optical waveguide," Nat. Commun. 11, 5295 (2020).
- [16] Maciej Lewenstein, Marcelo F Ciappina, Emilio Pisanty, Javier Rivera-Dean, Philipp Stammer, Th Lamprou, and Paraskevas Tzallas, "Generation of optical schrödinger cat states in intense laser-matter interactions," Nat. Phys. 17, 1104–1108 (2021).
- [17] Zhiling Wang, Zenghui Bao, Yukai Wu, Yan Li, Weizhou Cai, Weiting Wang, Yuwei Ma, Tianqi Cai, Xiyue Han, Jiahui Wang, et al., "A flying schrödinger's cat in multipartite entangled states," Sci. Adv. 8, eabn1778 (2022).

- [18] Marius Bild, Matteo Fadel, Yu Yang, Uwe von Lüpke, Phillip Martin, Alessandro Bruno, and Yiwen Chu, "Schrödinger cat states of a 16-microgram mechanical oscillator," Science 380, 274–278 (2023).
- [19] Yong Li, Z. D. Wang, and C. P. Sun, "Quantum criticality in a generalized Dicke model," Phys. Rev. A 74, 023815 (2006).
- [20] M. Aparicio Alcalde and B. M. Pimentel, "Path integral approach to the full Dicke model," Physica A 390, 3385– 3396 (2011).
- [21] Qing-Hu Chen, Chen Wang, Shu He, Tao Liu, and Ke-Lin Wang, "Exact solvability of the quantum Rabi model using Bogoliubov operators," Phys. Rev. A 86, 023822 (2012).
- [22] Alexander Moroz, "On solvability and integrability of the Rabi model," Ann. Phys. 338, 319–340 (2013).
- [23] Murray T. Batchelor and Huan-Qiang Zhou, "Integrability versus exact solvability in the quantum Rabi and Dicke models," Phys. Rev. A 91, 053808 (2015).
- [24] L. Bakemeier, A. Alvermann, and H. Fehske, "Quantum phase transition in the Dicke model with critical and noncritical entanglement," Phys. Rev. A 85, 043821 (2012).
- [25] Daniel Braak, Qing-Hu Chen, Murray T Batchelor, and Enrique Solano, "Semi-classical and quantum Rabi models: in celebration of 80 years," J. Phys. A: Math. Theor. 49, 300301 (2016).
- [26] F. A. Wolf, F. Vallone, G. Romero, M. Kollar, E. Solano, and D. Braak, "Dynamical correlation functions and the quantum Rabi model," Phys. Rev. A 87, 023835 (2013).
- [27] Daniel Braak, "Integrability of the Rabi model," Phys. Rev. Lett. 107, 100401 (2011).
- [28] Dingshun Lv, Shuoming An, Zhenyu Liu, Jing-Ning Zhang, Julen S. Pedernales, Lucas Lamata, Enrique Solano, and Kihwan Kim, "Quantum simulation of the quantum rabi model in a trapped ion," Phys. Rev. X 8, 021027 (2018).
- [29] Ricardo Puebla, Myung-Joong Hwang, Jorge Casanova, and Martin B Plenio, "Probing the dynamics of a superradiant quantum phase transition with a single trapped ion," Phys. Rev. Lett. 118, 073001 (2017).
- [30] Sam Genway, Weibin Li, Cenap Ates, Benjamin P Lanyon, and Igor Lesanovsky, "Generalized dicke nonequilibrium dynamics in trapped ions," Phys. Rev. Lett. **112**, 023603 (2014).
- [31] Arghavan Safavi-Naini, RJ Lewis-Swan, Justin G Bohnet, M Gärttner, KA Gilmore, JE Jordan, J Cohn, James K Freericks, Ana Maria Rey, and JJ Bollinger, "Verification of a many-ion simulator of the dicke model through slow quenches across a phase transition," Phys. Rev. Lett. **121**, 040503 (2018).
- [32] JS Pedernales, I Lizuain, S Felicetti, G Romero, L Lamata, and E Solano, "Quantum rabi model with trapped ions," Sci. Rep. 5, 15472 (2015).
- [33] Johannes Koch, Geram R Hunanyan, Till Ockenfels, Enrique Rico, Enrique Solano, and Martin Weitz, "Quantum rabi dynamics of trapped atoms far in the deep strong coupling regime," Nat. Commun. 14, 954 (2023).
- [34] Q-X Mei, B-W Li, Y-K Wu, M-L Cai, Ye Wang, Lin Yao, Z-C Zhou, and L-M Duan, "Experimental realization of the rabi-hubbard model with trapped ions," Phys. Rev. Lett. 128, 160504 (2022).
- [35] B-W Li, Q-X Mei, Y-K Wu, M-L Cai, Y Wang, L Yao, Z-C Zhou, and L-M Duan, "Observation of non-markovian spin dynamics in a jaynes-cummings-hubbard model us-

ing a trapped-ion quantum simulator," Phys. Rev. Lett. **129**, 140501 (2022).

- [36] Fumiki Yoshihara, Tomoko Fuse, Sahel Ashhab, Kosuke Kakuyanagi, Shiro Saito, and Kouichi Semba, "Superconducting qubit–oscillator circuit beyond the ultrastrong-coupling regime," Nat. Phys. 13, 44–47 (2017).
- [37] Pierre Nataf and Cristiano Ciuti, "No-go theorem for superradiant quantum phase transitions in cavity QED and counter-example in circuit QED," Nat. Commun. 1, 72 (2010).
- [38] Ri-Hua Zheng, Wen Ning, Ye-Hong Chen, Jia-Hao Lü, Li-Tuo Shen, Kai Xu, Yu-Ran Zhang, Da Xu, Hekang Li, Yan Xia, et al., "Observation of a superradiant phase transition with emergent cat states," Phys. Rev. Lett. 131, 113601 (2023).
- [39] Jinchen Zhao and Myung-Joong Hwang, "Frustrated superradiant phase transition," Phys. Rev. Lett 128, 163601 (2022).
- [40] Michael Scheibner, Thomas Schmidt, Lukas Worschech, Alfred Forchel, Gerd Bacher, Thorsten Passow, and Detlef Hommel, "Superradiance of quantum dots," Nat. Phys. 3, 106–110 (2007).
- [41] VI Yukalov and EP Yukalova, "Dynamics of quantum dot superradiance," Phys. Rev. B 81, 075308 (2010).
- [42] Valerii K Kozin, Dmitry Miserev, Daniel Loss, and Jelena Klinovaja, "Quantum phase transitions and cat states in cavity-coupled quantum dots," arXiv preprint arXiv:2310.15167 (2023).
- [43] Chris Hamner, Chunlei Qu, Yongping Zhang, JiaJia Chang, Ming Gong, Chuanwei Zhang, and Peter Engels, "Dicke-type phase transition in a spin-orbitcoupled bose–einstein condensate," Nat. Commun. 5, 4023 (2014).
- [44] Kristian Baumann, Christine Guerlin, Ferdinand Brennecke, and Tilman Esslinger, "Dicke quantum phase transition with a superfluid gas in an optical cavity," Nature 464, 1301–1306 (2010).
- [45] Kristian Baumann, Rafael Mottl, Ferdinand Brennecke, and Tilman Esslinger, "Exploring symmetry breaking at the Dicke quantum phase transition," Phys. Rev. Lett. 107, 140402 (2011).
- [46] D Schmidt, H Tomczyk, S Slama, and C Zimmermann, "Dynamical instability of a Bose-Einstein condensate in an optical ring resonator," Phys. Rev. Lett. 112, 115302 (2014).
- [47] Manuele Landini, Nishant Dogra, Katrin Kröger, Lorenz Hruby, Tobias Donner, and Tilman Esslinger, "Formation of a spin texture in a quantum gas coupled to a cavity," Phys. Rev. Lett. **120**, 223602 (2018).
- [48] Ronen M Kroeze, Yudan Guo, Varun D Vaidya, Jonathan Keeling, and Benjamin L Lev, "Spinor self-ordering of a

quantum gas in a cavity," Phys. Rev. Lett. **121**, 163601 (2018).

- [49] Markus P. Baden, Kyle J Arnold, Arne L Grimsmo, Scott Parkins, and Murray D. Barrett, "Realization of the Dicke model using cavity-assisted Raman transitions," Phys. Rev. Lett. **113**, 020408 (2014).
- [50] Nick Sauerwein, Francesca Orsi, Philipp Uhrich, Soumik Bandyopadhyay, Francesco Mattiotti, Tigrane Cantat-Moltrecht, Guido Pupillo, Philipp Hauke, and Jean-Philippe Brantut, "Engineering random spin models with atoms in a high-finesse cavity," Nat. Phys., 1–7 (2023).
- [51] Xiaotian Zhang, Yu Chen, Zemao Wu, Juan Wang, Jijie Fan, Shujin Deng, and Haibin Wu, "Observation of a superradiant quantum phase transition in an intracavity degenerate fermi gas," Science **373**, 1359–1362 (2021).
- [52] Kenji Hayashida, Takuma Makihara, Nicolas Marquez Peraca, Diego Fallas Padilla, Han Pu, Junichiro Kono, and Motoaki Bamba, "Perfect intrinsic squeezing at the superradiant phase transition critical point," Sci. Rep. 13, 2526 (2023).
- [53] Myung-Joong Hwang, Ricardo Puebla, and Martin B. Plenio, "Quantum Phase Transition and Universal Dynamics in the Rabi Model," Phys. Rev. Lett. 115, 180404 (2015).
- [54] Maoxin Liu, Stefano Chesi, Zu-Jian Ying, Xiaosong Chen, Hong-Gang Luo, and Hai-Qing Lin, "Universal Scaling and Critical Exponents of the Anisotropic Quantum Rabi Model," Phys. Rev. Lett. **119**, 220601 (2017).
- [55] Wei-Feng Zhuang, Bin Geng, Hong-Gang Luo, Guang-Can Guo, and Ming Gong, "Universality class and exact phase boundary in the superradiant phase transition," Phys. Rev. A 104, 053308 (2021).
- [56] Clive Emary and Tobias Brandes, "Quantum Chaos Triggered by Precursors of a Quantum Phase Transition: The Dicke Model," Phys. Rev. Lett. 90, 044101 (2003).
- [57] Clive Emary and Tobias Brandes, "Chaos and the quantum phase transition in the Dicke model," Phys. Rev. E 67, 066203 (2003).
- [58] We focus on $g_1 > 0$ and $g_2 > 0$. When these two coefficients have opposite sign, we can make a transformation $a \to ia$, and $\sigma^- \to i\sigma^-$ to change them to have the same sign. When $g_1 < 0$ and $g_2 < 0$, we can change $a \to -a$ to change them to be positive valued.
- [59] M-L Cai, Z-D Liu, W-D Zhao, Y-K Wu, Q-X Mei, Y Jiang, L He, X Zhang, Z-C Zhou, and L-M Duan, "Observation of a quantum phase transition in the quantum rabi model with a single trapped ion," Nat. Commun. 12, 1–8 (2021).
- [60] Kavan Modi, Aharon Brodutch, Hugo Cable, Tomasz Paterek, and Vlatko Vedral, "The classical-quantum boundary for correlations: Discord and related measures," Rev. Mod. Phys. 84, 1655 (2012).