This is the Accepted Manuscript of this published paper: Poch, O et al., 2023. Identification of Ammonium Salts on Comet 67P/C-G Surface from Infrared VIRTIS/Rosetta Data Based on Laboratory Experiments. Implications and Perspectives, in: Mennella, V., Joblin, C. (Eds.), European Conference on Laboratory Astrophysics ECLA2020, Astrophysics and Space Science Proceedings. Springer International Publishing, Cham, pp. 271–279. https://doi.org/10.1007/978-3-031-29003-9_31

Identification of ammonium salts on comet 67P/C-G surface from infrared VIRTIS/Rosetta data based on laboratory experiments. Implications and Perspectives.

Olivier Poch^{1[0000-0001-6777-8296]}, Istiqomah Istiqomah¹, Eric Quirico¹, Pierre Beck^{1,2}, Bernard Schmitt¹, Patrice Theulé³, Alexandre Faure¹, Pierre Hily-Blant¹, Lydie Bonal¹, Andrea Raponi⁴, Mauro Ciarniello⁴, Batiste Rousseau¹, Sandra Potin¹, Olivier Brissaud¹, Laurène Flandinet¹, Gianrico Filacchione⁴, Antoine Pommerol⁵, Nicolas Thomas⁵, David Kappel^{6,7}, Vito Mennella⁸, Lyuba Moroz⁷, Vassilissa Vinogradoff⁹, Gabriele Arnold⁷, Stéphane Erard¹⁰, Dominique Bockelée-Morvan¹⁰, Cédric Leyrat¹⁰, Fabrizio Capaccioni⁴, Maria Cristina De Sanctis⁴, Andrea Longobardo^{4,11}, Francesca Mancarella¹², Ernesto Palomba⁴, Federico Tosi⁴

¹ Univ. Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France
² Institut Universitaire de France, Paris, France
³ Aix-Marseille Université, CNRS, CNES, LAM, Marseille, France
⁴ IAPS, INAF, 00133 Rome, Italy
⁵ Physikalisches Institut, University of Bern, CH-3012 Bern, Switzerland
⁶ Institute of Physics and Astronomy, University of Potsdam, 14476 Potsdam, Germany
⁷ Institute for Planetary Research, DLR, 12489 Berlin, Germany
⁸ INAF–Osservatorio Astronomico di Capodimonte, Napoli, Italy
⁹ CNRS, Aix-Marseille Université, PIIM, UMR CNRS 7345, 13397 Marseille, France
¹⁰ LESIA, Obs. de Paris, PSL, CNRS, Sorbonne Univ., Univ. de Paris, 92195 Meudon, France
¹¹ DIST, Università Parthenope, 80143 Napoli, Italy
¹² Dipartimento di Matematica e Fisica "E. De Giorgi," Università del Salento, Lecce, Italy

olivier.poch@univ-grenoble-alpes.fr

Abstract. The nucleus of comet 67P/Churyumov-Gerasimenko exhibits a broad spectral reflectance feature around 3.2 μ m, which is omnipresent in all spectra of the surface, and whose attribution has remained elusive since its discovery. Based on laboratory experiments, we have shown that most of this absorption feature is due to ammonium (NH4⁺) salts mixed with the dark surface material. The depth of the band is compatible with semi-volatile ammonium salts being a major reservoir of nitrogen in the comet, which could dominate over refractory organic matter and volatile species. These salts may thus represent the long-sought reservoir of nitrogen in comets, possibly bringing their nitrogen-to-carbon ratio in agreement with the solar value. Moreover, the reflectance spectra of several asteroids are compatible with the presence of NH4⁺ salts at their surfaces. The presence of such salts, and other NH4⁺-bearing compounds on as-

teroids, comets, and possibly in proto-stellar environments, suggests that NH₄⁺ may be a tracer of the incorporation and transformation of nitrogen in ices, minerals and organics, at different phases of the formation of the Solar System.

Keywords: comets, laboratory experiments, nitrogen.

1 Introduction

The nucleus of comet 67P/Churyumov-Gerasimenko was mapped by the Visible, InfraRed and Thermal Imaging Spectrometer, Mapping Channel (VIRTIS-M) onboard the Rosetta spacecraft. The nucleus appeared almost spectrally uniform from 0.4 to 4 μ m, characterised by a very low reflectance of few percent, a positive (red) spectral slope, and a broad absorption feature from 2.9 to 3.6 μ m, centred at 3.2 μ m [1]. This absorption band is the most prominent feature and is widespread on the surface of comet 67P nucleus. Earlier analyses of these reflectance spectra suggested that the darkness and slope could be due to refractory poly-aromatic organics and opaque minerals (anhydrous sulfides and Fe-Ni alloys), but the attribution of the 3.2- μ m absorption feature has remained elusive [1, 2, 3]. It has been proposed to be mainly due to semi-volatile components of low molecular weight, most plausibly carboxylic (-COOH) bearing molecules or ammonium (NH₄⁺) ions, with contribution from C–H stretches [2, 4]. To test these hypotheses, we performed a series of laboratory measurements of the reflectance spectra of carboxylic acids or ammonium salts under simulated cometary-like conditions.

2 Laboratory measurements

Cometary dust appears to be made of aggregates of sub-micrometre-sized (sub- μ m) grains [5]. Moreover, cometary nuclei are porous (70-85 %), and their surfaces partially covered by this dust are likely to have a high porosity [6]. We have developed an experimental protocol aiming at simulating these structural properties of cometary dust, following the same methodology as described in [7].

Sub-µm grains of opaque iron sulfide (pyrrhotite, Fe_{1-x}S, 0 < x < 0.2) were dispersed in a liquid water solution containing carboxylic acids or ammonium salts. The liquid mixture was then nebulised to produce droplets that were frozen in liquid nitrogen to form icy dust particles. Finally, these particles were placed inside a thermal-vacuum chamber under cometary-like temperature (170-200 K) and high vacuum (< 10⁻⁵ mbar). Under these conditions, the water ice sublimated, and after several hours, a sublimate residue composed of a porous surface of opaque sub-µm grains mixed with a carboxylic acid or ammonium salts was obtained.

The reflectance spectra of these sublimate residues were measured from 0.4 to $4 \mu m$ using the spectro-radio-goniometer SHINE of the Cold Surfaces Spectroscopy facility of the Institut de Planétologie et d'Astrophysique de Grenoble (IPAG) [8].

3 Results

Figure 1 shows that the sublimate residue containing ≤ 17 wt% ammonium formate (NH₄⁺ HCOO⁻) exhibits an absorption band matching very closely the spectral position, asymmetric shape and maxima of absorption at 3.1 and 3.3 µm of the absorption feature observed on comet 67P.



Fig. 1. Average reflectance spectrum of comet 67P nucleus observed by VIRTIS-M (black line, S2) [4], compared to a sublimate residue composed of ≤ 17 wt% NH₄⁺ HCOO⁻ and ≥ 83 wt% opaque pyrrhotite grains measured in the laboratory at 170-200 K (red line, S1). From [9]. Reprinted with permission from AAAS.

We found that other ammonium salts $NH_4^+ X^-$ could also provide a good match to the VIRTIS spectrum (ammonium sulfate, citrate etc.), so the exact nature of X⁻ is unknown yet, and is certainly a mixture of various counter-ions. However, carboxylic acids, such as lactic acid, have very different absorption features [9]. Consequently, the 3.1- and 3.3-µm absorption bands observed on comet 67P appear to be due to the N–H stretching vibrations of NH_4^+ in ammonium salt, and several counter-ions may be expected. Additional absorption features present between 3.35 and 3.50 µm are attributed to C–H stretching vibrations of aliphatic carbonaceous compounds [4]. Other molecules have been proposed to contribute to the broad absorption, such as hydroxylated silicates [10].

The comparison of the depth of the absorption feature on the spectrum of comet 67P with spectra of mixtures of opaque grains and ammonium salts prepared and measured in the laboratory, suggests that the dust may contain several percent of ammonium salts. The abundance of ammonium salts is unknown, but could be from few percent up to 40 wt% in the cometary dust. The determination of a more precise concentration of the salts remains difficult because the band depth depends on many physical properties of the surface (grain sizes, mixing mode etc.) which may differ in

our experiments compared to the comet. Moreover, one cannot exclude that the concentration of NH_{4^+} at the surface may be different from at depth [11].

4 Conclusions and Perspectives

4.1 Independent identifications of NH₄⁺ salts by Rosetta instruments

The surface spectrum of comet 67P shown in Figure 1 is an average of all the dataset acquired at the beginning of the mission, from August to September 2014 [4, 9]. When the comet was closer than 2.25 AU from the Sun (around perihelion, so between April 2015 and January 2016), some dust particles from the coma entered the ROSINA-DFMS mass spectrometer, where they were heated at 273 K [12,13]. All fragments (NH₃, H₂S, HCl, etc.) typical of the thermal decomposition of several ammonium salts (NH₄⁺ SH⁻, NH₄⁺ Cl⁻, NH₄⁺ F⁻, NH₄⁺ CN⁻, NH₄⁺ HCOO⁻, NH₄⁺ HCOO⁻) were detected by the mass spectrometer in the hours following the entry of the dust particles ([12,13], see Rubin *et al.* in this volume for a presentation of ROSINA results). VIRTIS-M and ROSINA-DFMS instruments have thus independently identified the presence of ammonium salts in the cometary dust [9, 12]. The dust analyzer instrument onboard Rosetta, COSIMA, has not reported the detection of ammonium salts, most probably because these relatively volatile compounds have completely sublimated during the long pre-analysis storage of the dust particles at 283 K inside this instrument [14].

4.2 A new cometary reservoir of nitrogen

The 3.2-µm absorption feature observed on the entire surface of the comet 67P nucleus can be mainly attributed to ammonium salts and aliphatic organic matter. The exact concentration of the salts remains unknown. If they comprise more than about 5 wt% of the dust, these salts could be the major reservoir of nitrogen in the comet, dominating over the refractory organic matter and volatile species. Consequently, the abundance of nitrogen in comet 67P may be closer to the solar elemental composition than previously thought [9].

Moreover, NH_4^+ salts are possibly present not only on comet 67P but on a multitude of comets [12]. Indeed, the observations of higher NH_3/H_2O ratios in comae with decreasing perihelion distances (5% for D/2012 S1 (ISON) at 0.46 AU compared to 0.2% for 67P at 2 AU) [15, 12] may be explained by the relatively high sublimation temperature of NH_4^+ salts, decomposing more efficiently in NH_3 when closer to the Sun [9, 12, 16]. A significant fraction of nitrogen may thus be locked in ammonium salts in a multitude of comets.

4.3 NH4⁺ on other comets, asteroids, and in proto-stellar environments?

As mentioned earlier [17], the absorption feature of comet 67P shares similarities with other features observed around 3.1 μ m on several asteroids [9]. Could some of these absorption features be partly due to ammonium salts? Ammoniated phyllosilicates

and ammonium salts have been detected on the dwarf planet Ceres ([18], see also Raponi *et al.* in this volume). Absorption bands similar of NH_4^+ -bearing phyllosilicates [19] have also been detected on dust grains returned from the asteroid Ryugu by the Hayabusa2 mission ([20], see also Brucato *et al.* in this volume for a presentation of this mission). Could Ceres and Ryugu have inherited ammonium ions from outer Solar System objects similar to comet 67P? In addition, ammonium salts have been tentatively detected in the icy grains around proto-stars [21]. NH_4^+ -bearing compounds may thus be tracers of the incorporation and transformations of nitrogen, in ices, minerals and organics, from the proto-solar nebula to comets and asteroids. Future spectroscopic observations of such environments and bodies over an extended wavelength range up to the mid-infrared, as permitted with the James Webb Space Telescope, may provide new insights on the presence and abundance of NH_4^+ , and the nature of its counter-ions, in these various environments and objects.

4.4 Ammonium salts formation and origins

Laboratory experiments have shown that ammonium salts can be produced in/on solid ices via purely thermal reactions [22, 23, 24],

- by acid-base reactions from 10-30 K to higher temperatures: NH₃ + HCOOH → NH₄⁺ + HCOO⁻
- or nucleophilic addition of NH₃ with CO₂: $2NH_3 + CO_2 \rightarrow NH_4^+ + NH_2COO^-$.

The presence of surfaces outside and inside the ice (e.g. porous dust surfaces, and/or surfaces formed by thermal processing of ices: segregation, crystallization or sublimation) enables diffusion of the reactants and enhance the formation of these salts [22, 24]. So NH_4^+ salts can start to form early in pre- or proto-stellar ices (at 10-30 K), but their formation is probably more efficient as the temperature increases at later stages, in protoplanetary disks and in comets approaching the Sun.

4.5 Potential influences of NH4⁺ salts on planetary formation processes

Understanding the origins and evolutions of these salts may help to know how nitrogen was incorporated and distributed from interstellar environments to planetary systems. Ammonium salts may have played important roles at different stages of this evolution:

- Once being formed, they may influence the chemical evolution of icy grains mantles (reactivity, phase changes, etc.).
- If abundant enough, semi-volatile ammonium salts could enhance the stickiness of icy/dusty grains, contributing to accretion [25, 26]. Moreover, some of these nitrogen-bearing salts are solid up to more than 200 K, so they may provide nitrogen closer to the Sun than the N₂ and NH₃ snow lines [27].

- Salts lower the melting point of water in the subsurface of icy bodies (icy moons of the giant planets, Ceres), with implications for cryo-volcanism and habitability (pH, water activity) [28].
- Finally, NH₄⁺ salts are known to facilitate several prebiotic reactions occurring in liquid water, such as the phosphorylation of RNA [29].

5 Acknowledgements

O.P. acknowledges a post-doctoral fellowship from Centre National d'Etudes Spatiales (CNES). This work was supported by the CNES, the Agence Nationale de la Recherche (Classy grant, ANR-17-CE31-0004), the European Research Council (SOLARYS grant, ERC-CoG2017-771691), the Italian Space Agency and the DLR. This work takes advantage of the collaboration of the ISSI international team "Comet 67P/Churyumov-Gerasimenko Surface Composition as a Playground for Radiative Transfer Modeling and Laboratory Measurements", number 397. D.K. acknowledges DFG-grant KA 3757/2-1.

References

- Capaccioni, F., Coradini, A., Filacchione, G., et al.: The Organic-Rich Surface of Comet 67P/Churyumov-Gerasimenko as Seen by VIRTIS/Rosetta. Science 347, aaa0628 (2015). https://doi.org/10.1126/science.aaa0628
- Quirico, E., Moroz, L.V., Schmitt, B., et al.: Refractory and Semi-Volatile Organics at the Surface of Comet 67P/Churyumov-Gerasimenko: Insights from the VIRTIS/Rosetta Imaging Spectrometer. Icarus 272, 32 (2016). https://doi.org/10.1016/j.icarus.2016.02.028
- Rousseau, B., Érard, S., Beck, P., et al.: Laboratory Simulations of the Vis-NIR Spectra of Comet 67P Using Sub-Mm Sized Cosmochemical Analogues. Icarus 306, 306 (2018). https://doi.org/10.1016/j.icarus.2017.10.015
- Raponi, A., Ciarniello, M., Capaccioni, F., et al.: Infrared Detection of Aliphatic Organics on a Cometary Nucleus. Nature Astronomy 4, 5 (2020). https://doi.org/10.1038/s41550-019-0992-8
- 5. Levasseur-Regourd, A.-C., Agarwal, J., Cottin, H., et al.: Cometary Dust. Space Science Reviews 214, (2018).
- Kofman, W., Herique, A., Barbin, Y., et al.: Properties of the 67P/Churyumov-Gerasimenko Interior Revealed by CONSERT Radar. Science 349, (2015). https://doi.org/10.1007/s11214-018-0496-3
- Poch, O., Pommerol, A., Jost, B., et al.: Sublimation of Water Ice Mixed with Silicates and Tholins: Evolution of Surface Texture and Reflectance Spectra, with Implications for Comets. Icarus 267, 154 (2016). https://doi.org/10.1016/j.icarus.2015.12.017
- Brissaud, O., Schmitt, B., Bonnefoy, N., et al.: Spectrogonio Radiometer for the Study of the Bidirectional Reflectance and Polarization Functions of Planetary Surfaces 1 Design and Tests. Applied Optics 43, 1926 (2004). https://doi.org/10.1364/AO.43.001926
- Poch, O., Istiqomah, I., Quirico, E., et al.: Ammonium Salts Are a Reservoir of Nitrogen on a Cometary Nucleus and Possibly on Some Asteroids. Science 367, (2020). https://doi.org/10.1126/science.aaw7462

6

- Mennella, V., Ciarniello, M., Raponi, A., et al.: Hydroxylated Mg-Rich Amorphous Silicates: A New Component of the 3.2 µm Absorption Band of Comet 67P/Churyumov– Gerasimenko. The Astrophysical Journal Letters 897, L37 (2020). https://doi.org/10.3847%2F2041-8213%2Fab919e
- Suzuki, T., Shinnaka, Y., Majumdar, L., et al.: Possibility of Concentration of Nonvolatile Species near the Surface of Comet 67P/Churyumov-Gerasimenko. Astronomy & Astrophysics 645, A134 (2021). https://doi.org/10.1051/0004-6361/202038491
- Altwegg, K., Balsiger, H., Hänni, N., et al.: Evidence of Ammonium Salts in Comet 67P as Explanation for the Nitrogen Depletion in Cometary Comae. Nature Astronomy 4, 5 (2020). https://doi.org/10.1038/s41550-019-0991-9
- Altwegg, K., Combi, M., Fuselier, S.A., et al.: Abundant ammonium hydrosulphide embedded in cometary dust grains. Monthly Notices of the Royal Astronomical Society 516(3), 3900–3910 (2022). https://doi.org/10.1093/mnras/stac2440
- Fray, N., Bardyn, A., Cottin, H., et al.: Nitrogen-to-Carbon Atomic Ratio Measured by COSIMA in the Particles of Comet 67P/Churyumov–Gerasimenko. Monthly Notices of the Royal Astronomical Society 469, S506 (2017). https://doi.org/10.1093/mnras/stx2002
- Dello Russo, N., Vervack, R.J., Kawakita, H., et al.: The Compositional Evolution of C/2012 S1 (ISON) from Ground-Based High-Resolution Infrared Spectroscopy as Part of a Worldwide Observing Campaign. Icarus 266, 152 (2016). https://doi.org/10.1016/j.icarus.2015.11.030
- Mumma, M., Charnley, S., Cordiner, M., et al.: The Relationship of HCN, NH3, C2H6, H2O, and Ammoniated Salts in Comets: A Key Clue to Origins. EPSC-DPS Joint Meeting 2019 (2019). https://ui.adsabs.harvard.edu/abs/2019EPSC...13.1916M
- Rivkin, A.S., Howell, E.S., Emery, J.P.: Infrared Spectroscopy of Large, Low-Albedo Asteroids: Are Ceres and Themis Archetypes or Outliers?. Journal of Geophysical Research: Planets 124, 1393 (2019). https://doi.org/10.1029/2018JE005833
- Sanctis, M.C.D., Ammannito, E., Carrozzo, F.G., et al.: Ceres's Global and Localized Mineralogical Composition Determined by Dawn's Visible and Infrared Spectrometer (VIR). Meteoritics & Planetary Science 53, 1844 (2018). https://doi.org/10.1111/maps.13104
- Petit, S., Righi, D., Madejová, J.: Infrared Spectroscopy of NH4+-Bearing and Saturated Clay Minerals: A Review of the Study of Layer Charge. Applied Clay Science 34, 22 (2006). https://doi.org/10.1016/j.clay.2006.02.007
- Pilorget, C., Okada, T., Hamm, V., et al.: First Compositional Analysis of Ryugu Samples by the MicrOmega Hyperspectral Microscope. Nature Astronomy 6, 2 (2022). https://doi.org/10.1038/s41550-021-01549-z
- Boogert, A.C.A., Gerakines, P.A., Whittet, D.C.B.: Observations of the Icy Universe. Annual Review of Astronomy and Astrophysics 53, 541 (2015). https://doi.org/10.1146/annurev-astro-082214-122348
- Theulé, P., Duvernay, F., Danger, G., Thermal Reactions in Interstellar Ice: A Step towards Molecular Complexity in the Interstellar Medium, Advances in Space Research 52, 1567 (2013). https://doi.org/10.1016/j.asr.2013.06.034
- Bergner, J.B., Öberg, K.I., Rajappan, M., Fayolle, E.C.: Kinetics and Mechanisms of the Acid-Base Reaction Between NH3 and HCOOH in Interstellar Ice Analogs. The Astrophysical Journal 829, 85 (2016). https://doi.org/10.3847/0004-637X/829/2/85
- Potapov, A., Theulé, P., Jäger, C., Henning, T.: Evidence of Surface Catalytic Effect on Cosmic Dust Grain Analogs: The Ammonia and Carbon Di-oxide Surface Reaction. The Astrophysical Journal Letters 878, L20 (2019). https://doi.org/10.3847/2041-8213/ab2538

- Blum, J., Wurm, G.: The Growth Mechanisms of Macroscopic Bodies in Protoplanetary Disks. Annual Review of Astronomy and Astrophysics 46, 21 (2008). https://doi.org/10.1146/annurev.astro.46.060407.145152
- Lodders, K.: Jupiter Formed with More Tar than Ice. The Astrophysical Journal 611, 587 (2004). https://doi.org/10.1086/421970
- Kruczkiewicz, F., Vitorino, J., Congiu, E., Theulé, P., Dulieu, F.: Ammonia Snow Lines and Ammonium Salts Desorption. Astronomy & Astrophysics 652, A29 (2021). https://doi.org/10.1051/0004-6361/202140579
- Marion, G.M., Kargel, J.S., Catling, D.C., Lunine, J.I.: Modeling Ammonia–Ammonium Aqueous Chemistries in the Solar System's Icy Bodies. Icarus 220, 932 (2012). https://doi.org/10.1016/j.icarus.2012.06.016
- Burcar, B., Pasek, M., Gull, M., et al., Darwin's Warm Little Pond: A One-Pot Reaction for Prebiotic Phosphorylation and the Mobilization of Phosphate from Minerals in a Urea-Based Solvent. Angewandte Chemie International Edition 55, 13249 (2016). https://doi.org/10.1002/anie.201606239