

# A frame approach for equations involving the fractional Laplacian

Ioannis P. A. Papadopoulos\*    Timon S. Gutleb †    José A. Carrillo ‡  
Sheehan Olver §

March 1, 2024

## Abstract

Exceptionally elegant formulae exist for the fractional Laplacian operator applied to weighted classical orthogonal polynomials. We utilize these results to construct a solver, based on frame properties, for equations involving the fractional Laplacian of any power,  $s \in (0, 1)$ , on an unbounded domain in one or two dimensions. The numerical method represents solutions in an expansion of weighted classical orthogonal polynomials as well as their unweighted counterparts with a specific extension to  $\mathbb{R}^d$ ,  $d \in \{1, 2\}$ . We examine the frame properties of this family of functions for the solution expansion and, under standard frame conditions, derive an a priori estimate for the stationary equation. Moreover, we prove one achieves the expected order of convergence when considering an implicit Euler discretization in time for the fractional heat equation. We apply our solver to numerous examples including the fractional heat equation (utilizing up to a 6<sup>th</sup>-order Runge–Kutta time discretization), a fractional heat equation with a time-dependent exponent  $s(t)$ , and a two-dimensional problem, observing spectral convergence in the spatial dimension for sufficiently smooth data.

## 1 Introduction

In this work, we develop a spectral method to solve equations of the form

$$\mathcal{L}[u] := \sum_{k=1}^K \lambda_k (-\Delta)^{s_k} [u] = f, \quad (1.1)$$

where  $\lambda_k \in \mathbb{R}$  are known constants and  $s_k \in [0, 1]$ .  $(-\Delta)^s$  denotes the fractional Laplacian, as defined in the next section, and  $(-\Delta)^0 := \mathcal{I}$  is defined to be the identity operator. The domain is the whole real space  $\mathbb{R}^d$ ,  $d \in \{1, 2\}$ . For uniqueness, we seek a solution that decays as  $|x| \rightarrow \infty$ .

Many equations may reduce to (1.1) after nondimensionalization, linearization, or partial (time) discretization [6, 9, 15, 25, 34]. Solvers for (1.1) are an intermediate step for more

\*Weierstrass Institute for Applied Analysis and Stochastics, DE, [papadopoulos@wias-berlin.de](mailto:papadopoulos@wias-berlin.de);

†Department of Mathematics, University of British Columbia, CA, [gutleb@maths.ubc.ca](mailto:gutleb@maths.ubc.ca);

‡Mathematical Institute, University of Oxford, UK, [carrillo@maths.ox.ac.uk](mailto:carrillo@maths.ox.ac.uk);

§Department of Mathematics, Imperial College London, UK, [s.olver@imperial.ac.uk](mailto:s.olver@imperial.ac.uk).

complicated problems such as Newton linearizations of nonlocal Burgers-type equations [8, 21], power law absorption [57], quasi-geostrophic equations [17], fractional Keller-Segel type models [13, 26, 39, 40], or fractional porous medium flow [14]. Finite element methods are popular for approximating the solutions of these equations [11, 42]. However, since the operators are nonlocal and the solutions tend to be heavy-tailed (for instance decay at the rate of a Cauchy distribution for  $s = 1/2$ ) [58, Sec. 2], then any truncation of  $\mathbb{R}^d$  may result in artificial numerical artefacts [23, Sec. 9]. Spectral methods that construct bases with global support can be found in the literature [16, 18, 41, 43, 50, 52, 54, 55]. These alleviate the need to truncate  $\mathbb{R}^d$ ,  $d \in \{1, 2, 3\}$ , and some observe spectral convergence for sufficiently smooth data [23].

Our goal is to construct a flexible spectral method for (1.1) posed on  $\mathbb{R}^d$ ,  $d \in \{1, 2\}$ , via a frame approach [3, 4]. Our algorithm is summarized in Algorithm 1. This approach is different from other commonly used numerical methods for fractional diffusion based on finite-element, finite-difference and finite volume methods, see e.g. [1, 5, 7, 12, 22, 35, 42, 46, 56]. The main distinctive advantage compared to the previous literature is the flexibility in the choice of the family of functions for the solution expansion, the region of provable convergence, and the numerical confirmation that the algebraic tails are well approximated.

---

**Algorithm 1** Frame solver for (1.1).

---

1: **Input:**

- |   |  |
|---|--|
| $\mathcal{L}, f.$                       | ▷ Operator and right-hand side in (1.1). |
| $\Phi_N = \{\phi_1, \dots, \phi_N\}.$   | ▷ Basis for approximating $u.$           |
| $\epsilon, x_j, j \in \{1, \dots, M\}.$ | ▷ SVD tolerance and collocation points.  |

2: **Output:**

- |                                |   |
|--------------------------------|---|
| $\mathbf{u} \in \mathbb{R}^N.$ | ▷ $u \approx \sum_{i=1}^N \mathbf{u}_i \phi_i.$ |
|--------------------------------|---|
- 3: Assemble the matrix  $X_{ij} = (\mathcal{L}\phi_i)(x_j)$ ,  $i = 1, \dots, N$ ,  $j = 1, \dots, M$ .
  - 4: Assemble the vector  $\mathbf{y}_j = f(x_j)$ ,  $j = 1, \dots, M$ .
  - 5: Via an  $\epsilon$ -truncated SVD projection, compute  $\mathbf{f} \approx \operatorname{argmin}_{\mathbf{v} \in \mathbb{C}^N} \|X\mathbf{v} - \mathbf{y}\|_{\ell^2}$ .
  - 6:  $\mathbf{u} \leftarrow \mathbf{f}.$
- 

The performance of Algorithm 1 for solving (1.1) is heavily dependent on the choice of the family of functions  $\Phi_N$  that approximate  $u$ . A crucial ingredient for the construction of  $\Phi_N$  are recent results concerning the application of  $(-\Delta)^s$  to weighted and extended Jacobi,  $P_n^{(s,s)}(x)$ , and Zernike polynomials,  $Z_{n,m,j}^{(b)}(x, y)$  [32]. By fixing  $\Phi_N$ , we induce the family of functions for the right-hand side expansion  $\mathcal{L}\Phi_N = (\mathcal{L}\phi_1, \dots, \mathcal{L}\phi_N)$ . Thus the problem reduces from solving a fractional differential equation to expanding the right-hand side  $f$  in the induced family of functions. By noticing this space has the qualities of a frame, this motivates obtaining the expansion by solving a least-squares problem via an  $\epsilon$ -truncated SVD projection as done in line 5 of Algorithm 1. More precisely, in Theorem 5.2 we show that the family of functions consisting of extended and weighted Jacobi polynomials is a frame on the Hilbert space  $H_w^s(\mathbb{R})$ ,  $s \in (1/2, 1)$ , defined in Definition 5.3. Then, by utilizing Theorem 5.1, in Corollary 5.1 we deduce that the operator  $(\mathcal{I} + (-\Delta)^s)$ ,  $s \in (1/2, 1)$ , induces a family of functions for the expansion of the right-hand side that is a frame on the dual space of  $H_w^s(\mathbb{R})$ . This leads to the a priori estimate in Theorem 5.3. Although we do not show that the families of functions are frames on  $L^2(\mathbb{R})$ , we observe that if the correct family of functions is chosen, and a sufficient number of collocation

points are sampled, then the coefficients of the right-hand side expansion are well-behaved. Moreover, the coefficients of the expansion for the solution  $u$  are immediately deduced and the convergence is spectral for smooth right-hand sides.

If the number of collocation points and the number of functions in the frame are the same then, mathematically, the solver is very similar to a collocation method e.g. [54, 55]. The main differences are the flexible choices for the solution family of functions, the smaller overhead for the setup of the method, and the construction as a frame approach. A critique is that the truncated SVD of an  $m \times n$  matrix,  $m > n$ , scales at  $\mathcal{O}(mn^2)$ . For the purposes of this work, the factorization time is negligible and does not influence our results. However, the focus of future work will be to achieve close to  $\mathcal{O}(n \log^2 n)$  operations via the *AZ*-algorithm [20] and randomized linear algebra solvers for least-squares [33, 44, 45].

In Section 2 we rigorously state our problem and provide the relevant mathematical background. Then:

1. In Section 3 we explain why an expansion of the right-hand side  $f$ , as conducted in line 5 of Algorithm 1, provides an approximation for the solution.
2. We introduce the weighted Jacobi and Zernike polynomials in Section 4 and define the functions that are found in the induced family of functions for the right-hand side expansion.
3. In Section 5 we prove a number of results that contextualizes our method as a frame approach. These results motivate the use of an  $\epsilon$ -truncated SVD projection to expand  $f$ . We end with an a priori estimate in Theorem 5.3.
4. With these tools in Section 6 we provide the algorithm for the time discretization via Runge–Kutta methods, which is subtly different to a standard approach. We also prove that, in the case of a backward Euler discretization, we obtain the expected convergence rates in time and space.
5. We motivate our spectral method in several examples in Section 7 including equations with multiple fractional Laplacians with different exponents, a variable exponent, and a two-dimensional example.

## 2 Mathematical setup

Let  $W^{s,p}(\mathbb{R}^d)$ ,  $d \in \{1, 2\}$ , denote the (possibly fractional) Sobolev space [2, 24] and  $H^s(\mathbb{R}^d) := W^{s,2}(\mathbb{R}^d)$ . We denote the Lebesgue space by  $L^s(\mathbb{R}^d)$ ,  $s > 0$ . We seek a solution  $u \in H^s(\mathbb{R}^d)$  for (1.1). Let  $H^{-s}(\mathbb{R}^d) := H^s(\mathbb{R}^d)^*$  denote the dual space of  $H^s(\mathbb{R}^d)$ . Then we require  $f \in H^{-s}(\mathbb{R}^d)$ . Given a Banach space  $V$  equipped with the norm  $\|\cdot\|_V$ , we define the dual norm of the dual space  $V^*$  as follows, for any  $u^* \in V^*$ ,

$$\|u^*\|_{V^*} := \sup_{u \in V, u \neq 0} \frac{|\langle u^*, u \rangle_{V^*, V}|}{\|u\|_V}. \quad (2.1)$$

It follows that for any  $u^* \in V^*$  and  $v \in V$ ,  $\langle u^*, v \rangle_{V^*, V} \leq \|u^*\|_{V^*} \|v\|_V$ . We denote the space of bounded linear operators between the Banach spaces  $V$  and  $W$  by  $\mathcal{B}(V, W)$  and equip this space with the operator norm, for any  $\mathcal{A} \in \mathcal{B}(V, W)$ ,

$$\|\mathcal{A}\|_{\mathcal{B}(V, W)} := \sup_{u \in V, u \neq 0} \frac{\|\mathcal{A}u\|_W}{\|u\|_V}. \quad (2.2)$$

Similarly, we have that for any  $\mathcal{A} \in \mathcal{B}(V, W)$  and  $v \in V$ ,  $\|\mathcal{A}u\|_W \leq \|\mathcal{A}\|_{\mathcal{B}(V, W)}\|v\|_V$ .

In this work,  $\mathcal{I}$  denotes the identity operator and  $(-\Delta)^s$  denotes the fractional Laplacian. The fractional Laplacian has multiple definitions, many of them equivalent on unbounded domains when considering sufficiently regular functions [38] (and not equivalent in other contexts). Let  $\mathcal{S}(\mathbb{R}^d)$  denote the space of Schwartz functions on  $\mathbb{R}^d$ ,  $d \in \{1, 2\}$ . For any  $s \in (0, 1)$ , we define  $(-\Delta)^s : \mathcal{S}(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$  as [24, Sec. 3]:

$$(-\Delta)^s u(\mathbf{x}) := c_{d,s} \int_{\mathbb{R}^d} \frac{u(\mathbf{x}) - u(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|^{d+2s}} d\mathbf{y}, \text{ for a.e. } \mathbf{x} \in \mathbb{R}^d, \quad c_{d,s} := \frac{4^s \Gamma(d/2 + s)}{\pi^{d/2} |\Gamma(-s)|}. \quad (2.3)$$

Here  $\int_{\mathbb{R}^d} \cdot$  denotes the Cauchy principal value integral [37, Ch. 2.4] and  $\Gamma(\cdot)$  denotes the Gamma function. For any  $u \in H^s(\mathbb{R}^d)$ , it may be shown that  $(-\Delta)^{s/2} u \in L^2(\mathbb{R}^d)$  [24, Prop. 3.6]. The action of the fractional Laplacian may be recast in weak form,  $(-\Delta)^s : H^s(\mathbb{R}^d) \rightarrow H^{-s}(\mathbb{R}^d)$ :

$$\langle (-\Delta)^s u, v \rangle_{H^{-s}(\mathbb{R}^d), H^s(\mathbb{R}^d)} = \langle (-\Delta)^{s/2} u, (-\Delta)^{s/2} v \rangle_{L^2(\mathbb{R}^d)} \text{ for all } v \in H^s(\mathbb{R}^d). \quad (2.4)$$

Let  $\lambda > 0$ . Consider the weak form of (1.1), for a given  $f \in H^{-s}(\mathbb{R}^d)$  and for all  $v \in H^s(\mathbb{R}^d)$ ,

$$\langle (-\Delta)^{s/2} u, (-\Delta)^{s/2} v \rangle_{L^2(\mathbb{R}^d)} + \lambda \langle u, v \rangle_{L^2(\mathbb{R}^d)} = \langle f, v \rangle_{H^{-s}(\mathbb{R}^d), H^s(\mathbb{R}^d)}. \quad (2.5)$$

The existence of a unique solution  $u_* \in H^s(\mathbb{R}^d)$  of (2.5) may be shown via the Lax–Milgram theorem, e.g. [43, Sec. 2.2] and [50, Thm. 2.2]. By definition of  $H^s(\mathbb{R}^d)$ , we have that  $u_* \in L^2(\mathbb{R}^d)$ . Moreover, if  $f \in L^2(\mathbb{R}^d)$  (and thus  $(-\Delta)^s u_* \in L^2(\mathbb{R}^d)$ ) then  $u_*$  also solves [38, Th. 1.1]

$$(\lambda \mathcal{I} + (-\Delta)^s) u_* = f \text{ a.e. in } \mathbb{R}^d. \quad (2.6)$$

### 3 The spectral method

In this section we motivate Algorithm 1 as an approach for solving (1.1). In essence we are diagonalizing the operator  $\mathcal{L}$ .

**Definition 3.1** (Quasimatrix). *Throughout this work, we use boldface capital letters to denote a quasimatrix associated with a set of functions. A quasimatrix is a matrix whose “columns” are functions defined on  $\mathbb{R}^d$  [53, Lec. 5]. For instance if  $\Phi = \{\phi_i\}_{i=1}^\infty$  then*

$$\Phi(x) := (\phi_1(x) \quad \phi_2(x) \quad \dots). \quad (3.1)$$

**Proposition 3.1.** *Consider the equation (1.1), let  $H$  be a Hilbert space, and fix a right-hand side  $f \in H^*$ . Suppose that  $\mathcal{L}^{-1} \in \mathcal{B}(H^*, H)$ . Consider a finite-dimensional basis  $\Phi_N = \{\phi_1, \dots, \phi_N\}$  and define  $\Psi_N = \mathcal{L}\Phi_N = \{\mathcal{L}\phi_1, \dots, \mathcal{L}\phi_N\}$ . Consider the expansion coefficient  $\mathbf{f}$  for  $f$  in the basis  $\Psi_N$ . Then*

$$\|u - \Phi_N \mathbf{f}\|_H \leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(H^*, H)} \|f - \Psi_N \mathbf{f}\|_{H^*}. \quad (3.2)$$

*Proof.*

$$\begin{aligned} \|u - \Phi_N \mathbf{f}\|_H &= \|\mathcal{L}^{-1} \mathcal{L}u - \mathcal{L}^{-1} \mathcal{L}\Phi_N \mathbf{f}\|_H \\ &= \|\mathcal{L}^{-1} f - \mathcal{L}^{-1} \Psi_N \mathbf{f}\|_H \leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(H^*, H)} \|f - \Psi_N \mathbf{f}\|_{H^*}. \end{aligned} \quad (3.3)$$

□

Algorithm 1 is approximately finding an expansion for  $f$  via an  $\epsilon$ -truncated SVD projection and thus Proposition 3.1 implies that by setting  $\mathbf{u} = \mathbf{f}$ , then  $\|u - \Phi_N \mathbf{u}\|_H \leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(H^*, H)} \|f - \Psi_N \mathbf{f}\|_{H^*}$ .

In order to expand  $f$ , we must know the exact expressions for the functions in  $\Psi_N = \mathcal{L}\Phi_N$ . In the next section, we define functions such that, in the context of (1.1), these expressions are known.

## 4 Extended and weighted orthogonal polynomials

### 4.1 Jacobi polynomials

The Jacobi polynomials  $\{P_n^{(a,b)}(x)\}_{n \in \mathbb{N}}$  are a family of complete univariate bases of classical orthogonal polynomials on the interval  $(-1, 1)$  with weight parameters  $a, b \in \mathbb{R}$  such that  $a, b > -1$ . They are orthogonal with respect to the weight  $(1-x)^a(1+x)^b$ . Let  ${}_2F_1(\cdot, \cdot; \cdot; \cdot)$  and  $(\cdot)_n$  denote the hypergeometric function [47, Sec. 15.2] and Pochhammer symbol [47, Sec. 5.2(iii)], respectively. We pick the standard normalization such that [47, Eq. 18.5.7]

$$P_n^{(a,b)}(x) := \frac{(a+1)_n}{n!} {}_2F_1\left(-n, 1+a+b+n; a+1; \frac{1}{2}(1-x)\right). \quad (4.1)$$

Let  $(\cdot)_+$  denote the max function  $(z)_+ := \max(0, z)$  [47, p. 1.16.19]. Then we define the weighted Jacobi polynomial  $Q_n^{(a,b)}(x)$  as

$$Q_n^{(a,b)}(x) := (1-x)_+^a (1+x)_+^b P_n^{(a,b)}(x). \quad (4.2)$$

For any  $x \in \mathbb{R} \setminus [-1, 1]$ , we fix  $P_n^{(a,b)}(x) = Q_n^{(a,b)}(x) = 0$  for all  $n \in \mathbb{N}_0$ .

The extended Jacobi functions  $\tilde{P}_n^{(a,s)}(x)$ ,  $s \in (0, 1)$ ,  $a > -1$ , are defined as

$$\tilde{P}_n^{(a,s)}(x) := (-\Delta)^s Q_n^{(a,a)}(x). \quad (4.3)$$

Let  $\Gamma(\cdot)$  denote the Gamma function. The following theorem provides explicit expressions for  $\tilde{P}_n^{(a,s)}(x)$ .

**Theorem 4.1** (Row 1, Tab. 3 in [32]). *Consider  $s \in (0, 1)$  or  $s \in (-1/2, 0)$  and  $a > -1$ . Then, for any  $n \geq 0$  and  $x \in \mathbb{R}$ ,  $|x| \neq 1$ ,*

$$\tilde{P}_n^{(a,s)}(x) = 4^s \frac{\Gamma(a+n+1)}{n!} x^{n-2\lfloor \frac{n}{2} \rfloor} \times \begin{cases} \frac{\pi {}_2F_1(-a+s-\lfloor \frac{n}{2} \rfloor, n+s-\lfloor \frac{n}{2} \rfloor+\frac{1}{2}; n-2\lfloor \frac{n}{2} \rfloor+\frac{1}{2}; x^2)}{\sin(\pi(2\lfloor \frac{n}{2} \rfloor-n-s+\frac{1}{2}))\Gamma(n-2\lfloor \frac{n}{2} \rfloor+\frac{1}{2})\Gamma(-n-s+\lfloor \frac{n}{2} \rfloor+\frac{1}{2})\Gamma(a-s+\lfloor \frac{n}{2} \rfloor+1)} & |x| < 1, \\ -\frac{2^{-n-2s} \sin(\pi s)\Gamma(n+2s+1)|x|^{-2\lfloor \frac{n-1}{2} \rfloor-2s-3} {}_2F_1\left(s+\lfloor \frac{n}{2} \rfloor+1, \frac{2\lfloor \frac{n-1}{2} \rfloor+3}{2}+s; \frac{2n+3}{2}+a; \frac{1}{x^2}\right)}{\sqrt{\pi}\Gamma(\frac{2n+3}{2}+a)} & |x| > 1. \end{cases} \quad (4.4)$$

When  $a = s$ , the expression (4.4) simplifies. We find that, for  $x \in (-1, 1)$ ,  $\tilde{P}_n^{(s,s)}(x)$  is equal, up to a constant, to the corresponding Jacobi polynomial<sup>1</sup>  $P_n^{(s,s)}(x)$ .

<sup>1</sup>One could instead re-normalize  $\tilde{P}_n^{(s,s)}$  such that  $\tilde{P}_n^{(s,s)}(x) = P_n^{(s,s)}(x)$  for  $x \in (-1, 1)$ . However, this has the unfortunate consequence of making the notation more obfuscated in the later results.

**Theorem 4.2** (Row 1\*, Tab. 5 in [32]). *Consider  $s \in (0, 1)$  or  $s \in (-1/2, 0)$ . Then, for any  $n \geq 0$  and  $x \in \mathbb{R}$ ,  $|x| \neq 1$ ,*

$$\tilde{P}_n^{(s,s)}(x) = \begin{cases} \frac{4^s \Gamma(s + \lfloor \frac{n}{2} \rfloor + 1) \Gamma(n + s - \lfloor \frac{n}{2} \rfloor + \frac{1}{2})}{\lfloor \frac{n}{2} \rfloor! \Gamma(n - \lfloor \frac{n}{2} \rfloor + \frac{1}{2})} P_n^{(s,s)}(x) & |x| < 1, \\ -\frac{\sin(\pi s) x^{n-2} \lfloor \frac{n}{2} \rfloor! \Gamma(n+s+1) \Gamma(n+2s+1) |x|^{-2} \lfloor \frac{n-1}{2} \rfloor^{-2s-3} {}_2F_1(s + \lfloor \frac{n-1}{2} \rfloor + \frac{3}{2}, s + \lfloor \frac{n}{2} \rfloor + 1; n + s + \frac{3}{2}; \frac{1}{x^2})}{2^n \sqrt{\pi} n! \Gamma(n + s + \frac{3}{2})} & |x| > 1. \end{cases} \quad (4.5)$$

**Remark 4.1.** *We see that for  $x \in (-1, 1)$ , Theorem 4.2 implies that  $\tilde{P}_n^{(s,s)}$  are in fact scaled Jacobi polynomials for  $x \in (-1, 1)$ . This is a key observation in later results.*

**Remark 4.2.** *Thanks to their definition, we expect a solution family of functions consisting of extended Jacobi functions to accurately capture the algebraic decay of the solution tails. It can be shown that  $\tilde{P}_0^{(s,s)}(x) \sim |x|^{-1-2s}$  as  $|x| \rightarrow \infty$ , matching the expected decay rate for  $u \in H^s(\mathbb{R})$ , and  $\tilde{P}_n^{(s,s)}(x)$  decay at a faster rate for  $n \geq 1$ . Although this conjecture concerning the accurate representation of the algebraic tails is not proven in this work, we numerically verify it in Section 7.4.*

## 4.2 Affine transformations

In general, expanding  $u(x)$  solely in the weighted Jacobi polynomial space achieves a poor approximation. Hence, we consider expansions as sums of many families of functions centred at various points on  $\mathbb{R}$ .

**Definition 4.1** (Affine transformation). *Consider an interval  $I = [a, b] \subset \mathbb{R}$ ,  $a < b$  and the affine transformation  $y = 2/(b-a)(x - (a+b)/2)$ . We define the affine transformed function, centred at  $I$ , as  $f^I(x) = f(y)$ .*

Consider the intervals  $I_k$ ,  $k \in \{1, \dots, K\}$ , and the expansion

$$u(x) = \sum_{k=1}^K \sum_{j=0}^{\infty} \left[ \tilde{u}_j^k \tilde{P}_j^{I_k, (-s, -s)}(x) + u_j^k Q_j^{I_k, (s, s)}(x) \right]. \quad (4.6)$$

We order these affine transformation in the quasimatrix  $\Phi(x)$  as follows:

$$\Phi(x) = \left( \tilde{P}_0^{I_1, (-s, -s)}(x) \dots \tilde{P}_0^{I_K, (-s, -s)}(x) \mid Q_0^{I_1, (s, s)}(x) \dots Q_0^{I_K, (s, s)}(x) \mid \dots \right), \quad (4.7)$$

Thus (4.6) may be rewritten as  $u(x) = \Phi(x)\mathbf{u}$ .

**Remark 4.3.** *These results extend to the case where the Jacobi weight and fractional exponent do not match. The key observation is that, for  $-1 < s + t < 1$ ,*

$$(-\Delta)^t \tilde{P}_n^{(a,s)}(x) = (-\Delta)^{t+s} Q_n^{(a,a)}(x) = \tilde{P}_n^{(a,s+t)}(x). \quad (4.8)$$

*This allows one to apply the spectral method to solve general equations of the form (1.1). A numerical example is provided in Section 7.2.*

### 4.3 Generalized Zernike polynomials

The generalized Zernike polynomials,  $Z_{n,m,j}^{(b)}(x, y)$ ,  $a > -1$ , are multivariate orthogonal polynomials in Cartesian coordinates defined on the unit disk,  $\{(x, y) : x^2 + y^2 \leq 1\}$ . The subscript  $n$  denotes the polynomial degree and  $(m, j)$  denotes the Fourier mode and sign. If  $n$  is even, then  $m \in \{0, 2, \dots, n\}$  and, if  $n$  is odd, then  $m \in \{1, 3, \dots, n\}$ . If  $m = 0$ , then  $j = 1$ , otherwise  $j \in \{0, 1\}$ . The generalized Zernike polynomials are orthogonal with respect to the weight  $(1 - r^2)^b$ , where  $r^2 = x^2 + y^2$ . They may be defined via the Jacobi polynomials as:

$$V_{m,j}(x, y) := r^m \sin(m\theta + j\pi/2) \quad (4.9)$$

$$Z_{n,m,j}^{(b)}(x, y) := V_{m,j}(x, y) P_{(n-m)/2}^{(b,m)}(2r^2 - 1). \quad (4.10)$$

We define the extended Zernike functions for  $(x, y) \in \mathbb{R}^2$ ,  $r \neq 1$ , (when they exist) as

$$\tilde{Z}_{n,m,j}^{(b,s)}(x, y) := (-\Delta)^s [(1 - r^2)^b Z_{n,m,j}^{(b)}(x, y)]. \quad (4.11)$$

The following theorem provides explicit expressions for  $\tilde{Z}_{n,m,j}^{(s,s)}(x, y)$ .

**Theorem 4.3** (Row A\*\*, Tab. 7 in [32]). *Suppose that  $b = s$ ,  $s \in (-1, 1)$  and  $r^2 = x^2 + y^2$ . Then, for any  $n \geq 0$  and  $(x, y) \in \mathbb{R}^2$ ,  $r \neq 1$ ,*

$$\begin{aligned} \tilde{Z}_{n,m,j}^{(s,s)}(x, y) &= V_{m,j}(x, y) \frac{4^s \Gamma(1 + s + n)}{\Gamma(n + 1)} \\ &\times \begin{cases} \frac{\Gamma(1 + m + n + s) P_n^{(s,m)}(2r^2 - 1)}{\Gamma(1 + n + m)} & r < 1, \\ \frac{(-1)^n \Gamma(1 + m + n + s) {}_2F_1\left(n + s + 1, 1 + m + n + s; s + m + 2n + 2; \frac{1}{r^2}\right)}{\Gamma(-n - s) \Gamma(s + 1 + m + 2n + 1) r^{2(1 + m + n + s)}} & r > 1. \end{cases} \end{aligned} \quad (4.12)$$

## 5 Frames

As the family of approximating functions is not orthogonal, the expansion of a known function,  $f(x)$ , is nontrivial. In particular, the expansion need not be unique. For our purposes, we desire an expansion that approximates our known function to the required tolerance and the coefficients of the expansion are (relatively) small in magnitude.

We turn to the techniques used in frame theory [3, 4]. Essentially, we find the coefficients of the expansion that optimally interpolate the values of  $f(x)$  (or a linear operator applied to  $f$ ) at a set of collocation points in a least squares sense. Consider the collocation points  $\mathbf{x} = (x_1, \dots, x_M)$ . Let  $\Phi(x)$  denote the quasimatrix for the expansion of the solution  $u(x)$ . Pick the number of functions in the truncation  $N$ . As depicted in line 3 of Algorithm 1, the least-squares matrix is given by

$$X_{ij} = [l_i[\mathcal{L}\Phi]]_j, \quad i = 1 \dots, M, \quad j = 1, \dots, N, \quad (5.1)$$

where  $\{l_i\}$  are a set of linear operators on  $\{x_i\}$ . In this work, we solely consider the identity, i.e.  $l_i[\mathcal{L}\Phi] = (\mathcal{L}\Phi)(x_i)$ . Similarly we compute  $\mathbf{y}_i = l_i[f]$  and we solve the following least-squares problem for the expansion coefficients  $\mathbf{u}$ :

$$\min_{\mathbf{u} \in \mathbb{C}^N} \|\mathbf{X}\mathbf{u} - \mathbf{y}\|_{\ell^2}, \quad (5.2)$$

so that  $f(x) \approx \mathcal{L}\Phi(x)\mathbf{u}$  and, therefore,  $u(x) \approx \Phi(x)\mathbf{u}$  (as discussed in Section 3). In a frame setting, the least-squares matrix is often ill-conditioned for increasing  $M$  and  $N$ . However, in practice we recover suitable least squares solutions if we sufficiently oversample the collocation points and use a truncated SVD solver. The remainder of this section will focus on explaining this phenomenon.

Consider the SVD factorization  $X = U\Sigma V^\top$  with the convention where  $\Sigma$  is a square diagonal matrix with the singular values arranged in decreasing magnitude along the diagonal. Given a precision tolerance  $\epsilon$ , a truncated SVD solver finds the first singular value such that  $\Sigma_{i+1,i+1} < \epsilon < \Sigma_{i,i}$ . Then, the submatrix  $\Sigma_\epsilon \in \mathbb{R}^{i \times i}$ ,  $[\Sigma_\epsilon]_{j,j} = \Sigma_{j,j}$ ,  $j \in \{1, \dots, i\}$  is extracted. The  $\epsilon$ -truncated SVD projection of (5.2) is given by

$$\mathbf{u}_\epsilon = V_\epsilon \Sigma_\epsilon^{-1} U_\epsilon^\top \mathbf{y}, \quad (5.3)$$

where  $V_\epsilon$  and  $U_\epsilon$  are the first  $i$  columns of  $V$  and  $U$ , respectively.

**Remark 5.1** (Truncated SVD cost). *With  $M > N$  the cost of computing the SVD of  $X$  scales as  $\mathcal{O}(MN^2)$ . In the one-dimensional examples found in Section 7, we found that this cost is negligible and the bottleneck actually occurs in the assembly of  $X$  rather than the SVD factorization. However, our analytical results generalize to two and three dimensions. Here the cost of the factorization may prove prohibitive. However, for suitable frames, the solve may be implemented in  $\mathcal{O}(N \log^2 N)$  operations via the AZ-algorithm [20] and randomized linear algebra solvers for least-squares [33, 44, 45]. Achieving this speedup will be the focus of future work.*

**Remark 5.2** (Evaluation of the least-squares matrix). *A potential computational bottleneck is the assembly of the least-squares matrix  $X$  as defined in (5.1), particularly if we are required to evaluate many hypergeometric functions at many collocation points. Although a direct evaluation of many  ${}_2F_1$  functions may be slow, we note that the extended Jacobi functions satisfy the same three-term recurrence as their Jacobi polynomial counterparts. Thus one may use the forward recurrence for efficient evaluation of the extended Jacobi functions on the interval  $[-1, 1]$  and (F. W. J) Olver's and Miller's algorithm for evaluation off the interval  $[-1, 1]$ , cf. [31, Sec. 2.3], [47, Sec. 3.6] and [48, App. B]. Alternatively, off the interval one can use a continued fraction approach advocated by Gautschi [30].*

We now introduce the notion of a frame.

**Definition 5.1** (Frame). *Consider a Hilbert space  $(H, (\cdot, \cdot)_H)$ . An indexed family of functions  $\{\phi_n\} \in H$  is called a frame for  $H$  if there exist constants  $0 < c \leq C < \infty$  such that*

$$c\|f\|_H^2 \leq \sum_n |(f, \phi_n)_H|^2 \leq C\|f\|_H^2 \text{ for all } f \in H. \quad (5.4)$$

As the family of functions for the expansion of the right-hand side will vary according to the choice of (1.1), it is helpful to construct a framework whereby if one shows that the family of functions for the solution expansion is a frame (which remains fixed), then the induced family of functions for the right-hand is automatically a frame provided that the operator  $\mathcal{L}$  in (1.1) operator is well-behaved.

**Theorem 5.1** (Frames on the dual space). *Suppose that the family of functions  $\Phi = \{\phi_n\}_{n \in \mathbb{N}_0}$  is a frame on the Hilbert space  $H$ . Consider the bounded linear operator  $\mathcal{L} : H \rightarrow H^*$  where  $H^*$*



is the dual space of  $H$ . Moreover, assume that the adjoint operator  $\mathcal{L}^* : H^* \rightarrow H$  is bounded and positive-definite with constants  $M_b$  and  $M_p$ , respectively. Then  $\mathcal{L}\Phi = \{\mathcal{L}\phi_n\}_{n \in \mathbb{N}_0}$  is a frame on the Hilbert space  $H^*$ .

*Proof.* Let  $\psi_n = \mathcal{L}\phi_n$ ,  $n \in \mathbb{N}_0$ . Now consider any  $f^* \in H^*$ . The lower bound is derived as follows:

$$cM_p^2 \|f^*\|_{H^*}^2 \leq c \|\mathcal{L}^* f^*\|_H^2 \leq \sum_{n=0}^{\infty} (\mathcal{L}^* f^*, \phi_n)_H^2 = \sum_{n=0}^{\infty} (f^*, \psi_n)_{H^*}^2, \quad (5.5)$$

where  $c$  is the lower bound for the frame condition of  $\Phi$ . The first inequality follows from the positive-definiteness of the adjoint operator  $\mathcal{L}^*$  and the second inequality follows from the frame condition of  $\Phi$ . The final equality follows by the definition of the adjoint operator and  $\psi_n$ .

Similarly, let  $C$  denote the upper bound for the frame condition of  $\Phi$ , then

$$\begin{aligned} \sum_{n=0}^{\infty} (f^*, \psi_n)_{H^*}^2 &= \sum_{n=0}^{\infty} (\mathcal{L}^* f^*, \phi_n)_H^2 \\ &\leq C \|\mathcal{L}^* f^*\|_H^2 \leq C \|\mathcal{L}^*\|_{\mathcal{B}(H^*, H)}^2 \|f^*\|_{H^*}^2 = CM_b^2 \|f^*\|_{H^*}^2. \end{aligned} \quad (5.6)$$

□

## 5.1 Construction of frames for $\mathcal{L} = (\mathcal{I} + (-\Delta)^s)$

In this subsection, we show that a family of functions consisting of weighted Jacobi polynomials and their fractional Laplacian counterparts  $\Phi(x)$  are a frame on a weighted Hilbert space. Then, by utilizing [Theorem 5.1](#), we conclude that  $(\mathcal{I} + (-\Delta)^s)\Phi(x)$  is a frame on the dual space.

**Definition 5.2** (Weighted Lebesgue space). *Let  $w : [a, b] \rightarrow [c, \infty)$ ,  $c > 0$ , be a measurable function. Then, the weighted Lebesgue space  $L_w^p(a, b)$  is equipped with the norm  $\|f\|_{L_w^p(a, b)} := (\int_a^b |f|^p w \, dx)^{1/p}$ . If  $p = 2$ , then  $L_w^2(a, b)$  is a Hilbert space equipped with the inner product  $(f, g)_{L_w^2(a, b)} := \int_a^b f g w \, dx$ .*

**Definition 5.3** (Hilbert space  $H_w^s(\mathbb{R})$ ). *Let  $H_w^s(\mathbb{R})$  denote the Hilbert space*

$$H_w^s(\mathbb{R}) := \{u \in L_w^2(\mathbb{R}) : \text{supp}(u) \subseteq \text{supp}(w), (-\Delta)^{s/2} u \in L^2(\mathbb{R})\}, \quad (5.7)$$

*equipped with the inner-product*

$$(u, v)_{H_w^s(\mathbb{R})} := (u, v)_{L_w^2(\mathbb{R})} + ((-\Delta)^{s/2} u, (-\Delta)^{s/2} v)_{L^2(\mathbb{R})}.$$

*Here  $\text{supp}$  denotes the support of a function.*

**Remark 5.3.** *When considering nonlocal operators defined on  $\mathbb{R}$ , such as  $(-\Delta)^s$ , we note that even if the support of a function  $u$  is compactly contained in  $\mathbb{R}$ , this does not mean  $(-\Delta)^s u$  is compactly supported in  $\mathbb{R}$ . This is the motivation behind considering the inner-product defined on all of  $\mathbb{R}$  in [Definition 5.3](#).*

The following lemma proves orthogonality of weighted and extended Jacobi polynomials. This is heavily utilized for the frame result in [Theorem 5.2](#). A similar result may be found in [[50](#), Prop. 3.6].

**Lemma 5.1** (Orthogonality). *Let  $w(x) = (1-x^2)_+^{-s}$ ,  $s < 1$ . Then,  $\{Q_n^{(s,s)}\}_{n \in \mathbb{N}_0}$  and  $\{\tilde{P}_n^{(-s,-s)}\}_{n \in \mathbb{N}_0}$  satisfy*

$$(\tilde{P}_n^{(-s,-s)}, \tilde{P}_m^{(-s,-s)})_{H_w^s(\mathbb{R})} = \tilde{C}_{n,m} \delta_{n,m} \quad \text{and} \quad (Q_n^{(s,s)}, Q_m^{(s,s)})_{H_w^s(\mathbb{R})} = C_{n,m} \delta_{n,m}, \quad (5.8)$$

where  $\delta_{nm}$  denotes the Kronecker delta and  $C_{n,m}, \tilde{C}_{n,m} \in \mathbb{R}$ .

*Proof.* Let  $c_{s,n} \in \mathbb{R}$ ,  $n \in \mathbb{N}_0$ , denote the constants such that, for all  $x \in (-1, 1)$ ,  $\tilde{P}_n^{(s,s)}(x) = c_{s,n} P_n^{(s,s)}(x)$ . Then,

$$\begin{aligned} & (Q_n^{(s,s)}, Q_m^{(s,s)})_{H_w^s(\mathbb{R})} \\ &= (Q_n^{(s,s)}, Q_m^{(s,s)})_{L_w^2(\mathbb{R})} + ((-\Delta)^{s/2} Q_n^{(s,s)}, (-\Delta)^{s/2} Q_m^{(s,s)})_{L^2(\mathbb{R})} \\ &= (Q_n^{(s,s)}, P_m^{(s,s)})_{L^2(-1,1)} + (Q_n^{(s,s)}, (-\Delta)^s Q_m^{(s,s)})_{L^2(\mathbb{R})} \\ &= (Q_n^{(s,s)}, P_m^{(s,s)})_{L^2(-1,1)} + (Q_n^{(s,s)}, c_{s,m} P_m^{(s,s)})_{L^2(-1,1)} = C_{n,m} \delta_{n,m}. \end{aligned} \quad (5.9)$$

The result for  $\{\tilde{P}_n^{(-s,-s)}\}_{n \in \mathbb{N}_0}$  follows similarly.  $\square$

**Lemma 5.2** (Bessel's inequality, cf. Ch. 8, Sec. 4-5 in [59]). *Let  $p_n$ ,  $n \in \mathbb{N}_0$ , denote an orthonormal sequence in the inner-product space  $H$ . Then, for every  $u \in H$ , the following Bessel's inequality holds:*

$$\sum_{n=0}^{\infty} (u, p_n)_H^2 \leq \|u\|_H^2. \quad (5.10)$$

**Theorem 5.2** (Frame on  $H_w^s(\mathbb{R})$ ,  $s \in (1/2, 1)$ ). *Let  $w(x) = (1-x^2)_+^{-s}$ ,  $s \in (1/2, 1)$ . Let  $\{\hat{Q}_n^{(s,s)}\}$  and  $\{\hat{P}_n^{(s,s)}\}$  denote the orthonormalized families of functions,  $\{Q_n^{(s,s)}\}$  and  $\{\tilde{P}_n^{(s,s)}\}$ , respectively, with respect to the  $H_w^s(\mathbb{R})$  inner-product. Then, the space spanned by  $\{\hat{Q}_n^{(s,s)}\}_{n=0}^{\infty} \cup \{\hat{P}_n^{(-s,-s)}\}_{n=0}^{\infty}$  is a frame on  $H_w^s(\mathbb{R})$ .*

*Proof.* Consider any  $u \in H_w^s(\mathbb{R})$ . We note that the weighted polynomials  $\hat{Q}_n$ ,  $n \in \mathbb{N}_0$ , are dense in  $L^2(-1, 1)$  for any functions that vanish on  $-1$  and  $1$ . By a Hölder regularity result [23, Th. 8.2] since  $s > 1/2$  then  $u$  must be continuous. By definition  $u = 0$  a.e. in  $\mathbb{R} \setminus [-1, 1]$  and, therefore,  $u(\pm 1) = 0$ . Thus we may find an expansion

$$u(x) = \sum_{n=0}^{\infty} u_n \hat{Q}_n(x) \quad \text{for a.e. } x \in \mathbb{R}. \quad (5.11)$$

Moreover, since  $(\hat{Q}_n, \hat{Q}_m)_{L_w^2(-1,1)} = c_{n,m} \delta_{n,m}$  for some  $c_{n,m} \in \mathbb{R}$ , we have that

$$u_n = (c_{n,n})^{-1} (u, \hat{Q}_n)_{L_w^2(-1,1)}.$$

Now,

$$\begin{aligned} \|u\|_{H_w^s(\mathbb{R})}^2 &= \left\| \sum_{n=0}^{\infty} u_n \hat{Q}_n^{(s,s)} \right\|_{H_w^s(\mathbb{R})}^2 = \sum_{n,m=0}^{\infty} (u_n \hat{Q}_n^{(s,s)}, u_m \hat{Q}_m^{(s,s)})_{H_w^s(\mathbb{R})} \\ &= \sum_{n=0}^{\infty} u_n^2 = \sum_{n=0}^{\infty} (u, \hat{Q}_n^{(s,s)})_{H_w^s(\mathbb{R})}^2, \end{aligned} \quad (5.12)$$

where the penultimate equality follows from [Lemma 5.1](#) and the final equality follows from the orthonormality of  $\hat{Q}_n^{(s,s)}$  in  $H_w^s(\mathbb{R})$ . Hence by [\(5.12\)](#) we achieve the lower bound of the frame condition:

$$\|u\|_{H_w^s(\mathbb{R})}^2 \leq \sum_{n=0}^{\infty} \left[ (u, \hat{P}_n^{(-s,-s)})_{H_w^s(\mathbb{R})}^2 + (u, \hat{Q}_n^{(s,s)})_{H_w^s(\mathbb{R})}^2 \right]. \quad (5.13)$$

We now consider the upper bound. Note that since  $\{\hat{P}_n\}$  and  $\{\hat{Q}_n\}$  are orthonormal sequences in  $H_w^s(\mathbb{R})$  we have

$$\sum_{n=0}^{\infty} \left[ (u, \hat{P}_n^{(-s,-s)})_{H_w^s(\mathbb{R})}^2 + (\tilde{u}, \hat{Q}_n^{(s,s)})_{H_w^s(\mathbb{R})}^2 \right] \leq 2\|u\|_{H_w^s(\mathbb{R})}^2, \quad (5.14)$$

where the inequality follows from [Lemma 5.2](#).  $\square$

A basic building block for a frame suitable for expanding the solution of  $(\mathcal{I} + (-\Delta)^s)u = f$  is  $\Phi = \{\hat{Q}_n^{(s,s)}\}_{n=0}^{\infty} \cup \{\hat{P}_n^{(-s,-s)}\}_{n=0}^{\infty}$ . The operator  $\mathcal{L} = (-\Delta)^s + \mathcal{I}$  implies we expand the right-hand side  $f(x)$  in  $\Psi = \{\hat{P}_n^{(s,s)} + \hat{Q}_n^{(s,s)}\}_{n \in \mathbb{N}_0} \cup \{\hat{Q}_n^{(-s,-s)} + \hat{P}_n^{(-s,-s)}\}_{n \in \mathbb{N}_0}$ . In the next corollary we show that by utilizing [Theorem 5.1](#) and [Theorem 5.2](#), we may deduce that  $\Psi$  is also a frame.

**Corollary 5.1** (Frame on  $H_w^s(\mathbb{R})^*$ ,  $s \in (1/2, 1)$ ).  $\Psi = \{\hat{P}_n^{(s,s)} + \hat{Q}_n^{(s,s)}\}_{n \in \mathbb{N}_0} \cup \{\hat{Q}_n^{(-s,-s)} + \hat{P}_n^{(-s,-s)}\}_{n \in \mathbb{N}_0}$  is a frame on  $(H_w^s(\mathbb{R}))^*$  where  $w(x) = (1 - x^2)_+^{-s}$ ,  $s \in (1/2, 1)$ .

*Proof.* Let  $\mathcal{L} = (-\Delta)^s + \mathcal{I}$ . In [Theorem 5.2](#) it was shown that  $\Phi = \mathcal{L}^{-1}\Psi$  is a frame on  $H = H_w^s(\mathbb{R})$ . By [Theorem 5.1](#), it is sufficient to show that the adjoint  $\mathcal{L}^*$  is positive-definite and bounded on  $H^* = (H_w^s(\mathbb{R}))^*$ .

**(Bounded).** Consider any  $u^*, v^* \in H^*$ . Since  $H$  is a Hilbert space, it is reflexive. Thus, by the Riesz Representation Theorem [[27](#), Sec. D.2, Th. 2], for any  $u^* \in H^*$  there exists a unique  $u \in H$  such that  $\|u^*\|_{H^*} = \|u\|_H$  and  $\langle u^*, v \rangle_{H^*, H} = (u, v)_H$  for all  $v \in H$ . Let  $u, v \in H$  denote the unique Riesz representations of  $u^*$  and  $v^*$ , respectively. Then

$$\begin{aligned} |\langle \mathcal{L}^* v^*, u^* \rangle| &= |\langle \mathcal{L}^* v^*, u \rangle_H| = |\langle \mathcal{L} u, v^* \rangle_{H^*}| = |\langle \mathcal{L} u, v \rangle| = |(u, v)_{H^s(\mathbb{R})}| \\ &\leq \|u\|_{H^s(\mathbb{R})} \|v\|_{H^s(\mathbb{R})} \leq \|u\|_H \|v\|_H = \|u^*\|_{H^*} \|v^*\|_{H^*}, \end{aligned} \quad (5.15)$$

where the first, third, and final equalities follow by the Riesz Representation Theorem, the second equality follows by definition of the adjoint operator, the fourth equality follows from an integration by parts, the first inequality follows from the Cauchy-Schwarz inequality, and the second inequality follows since  $\text{supp}(u), \text{supp}(v) \subseteq \text{supp}(w)$  and  $w^{1/2}(x) \geq 1$  for  $x \in [-1, 1]$ . Thus

$$\|\mathcal{L}^* u^*\|_H = \sup_{v^* \neq 0} \frac{|\langle \mathcal{L}^* u^*, v^* \rangle|}{\|v^*\|_{H^*}} \leq \|u^*\|_{H^*}. \quad (5.16)$$

**(Positive-definite).** Recall that  $H = H_w^s(\mathbb{R})$ . Thus, by definition,

$$\|u\|_H^2 = (u, u)_H = ((-\Delta)^{s/2} u, (-\Delta)^{s/2} u)_{L^2(\mathbb{R})} + (w^{1/2} u, w^{1/2} u)_{L^2(\mathbb{R})}. \quad (5.17)$$

Now

$$(w^{1/2} u, w^{1/2} u)_{L^2(\mathbb{R})} = \|w^{1/2} u\|_{L^2(\mathbb{R})}^2 \leq \|w^{1/2}\|_{L^p(\mathbb{R})}^2 \|u\|_{L^q(\mathbb{R})}^2, \quad (5.18)$$

where  $1/p + 1/q = 2$ . For  $\|w^{1/2}\|_{L^p(\mathbb{R})}^2 < \infty$ , we require  $p < 2/s$ . Hence, we require  $q > 2/(1-s)$ . Now by [24, Th. 8.2] if  $q \leq \infty$ , then there exists a  $C \in \mathbb{R}$  such that  $\|u\|_{L^q(\mathbb{R})} \leq C\|u\|_{H^s(\mathbb{R})}$ . Hence, by fixing any  $q > 2/(1-s)$ , we have that

$$(w^{1/2}u, w^{1/2}u)_{L^2(\mathbb{R})} \leq C^2 \|w^{1/2}\|_{L^p(\mathbb{R})}^2 \|u\|_{H^s(\mathbb{R})}^2. \quad (5.19)$$

Therefore,

$$\begin{aligned} \|u^*\|_{H^*}^2 &= \|u\|_H^2 \leq (C^2 \|w^{1/2}\|_{L^p(\mathbb{R})}^2 + 1)(u, u)_{H^s(\mathbb{R})} = \tilde{C} \langle \mathcal{L}u, u \rangle \\ &= \tilde{C} (\mathcal{L}u, u^*)_{H^*} = \tilde{C} (u, \mathcal{L}^* u^*)_H \leq \tilde{C} \|\mathcal{L}^* u^*\|_H \|u\|_H = \tilde{C} \|\mathcal{L}^* u^*\|_H \|u^*\|_{H^*}. \end{aligned} \quad (5.20)$$

By dividing through by  $\|u^*\|_{H^*}$  in (5.20), we conclude the positive-definiteness and the result.  $\square$

We conclude this section with the following a priori estimate which constitutes the main theorem of this work.

**Theorem 5.3** (A priori estimate). *Consider (1.1) and suppose that the conditions of Theorem 5.1 hold. Let  $\Phi$  denote the quasimatrix of a frame for the Hilbert space  $H = H_w^s(\mathbb{R})$  with  $H_w^s(\mathbb{R})$  as defined in Definition 5.3. Consider the truncated quasimatrices  $\Phi_N = (\phi_1 \ \phi_2 \ \cdots \ \phi_N)$  and  $\Psi_N = \mathcal{L}\Phi_N$ . Suppose that  $\mathbf{u}_N \in \mathbb{R}^N$  is computed via Algorithm 1. Then, there exist constants  $\lambda, \kappa > 0$  such that*

$$\begin{aligned} \|u - \Phi_N \mathbf{u}_N\|_H &\leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(H^*, H)} \inf_{\mathbf{v} \in \mathbb{C}^N} \{ \|f - \Psi_N \mathbf{v}\|_{H^*} + \kappa \|f - \Psi_N \mathbf{v}\|_M + \epsilon \lambda \|\mathbf{v}\|_{\ell^2} \}, \end{aligned} \quad (5.21)$$

where  $x_j, j = 1, \dots, M$  are the collocation points and  $\|v\|_M := \sum_{j=1}^M v(x_j)^2$ .

**Remark 5.4.** *The constants  $\lambda$  and  $\kappa$  are dependent on truncated SVD tolerance  $\epsilon$ , the collocation points, and  $N$  as well as the problem and the frame. Ideally their magnitude is  $\mathcal{O}(1)$  as  $N \rightarrow \infty$  for careful choices of the collocation points, in which case we are guaranteed to reach an accuracy of  $\epsilon$  if  $N$  is taken sufficiently large. For a thorough discussion of the constants and their behaviour, we refer the reader to [4].*

*Proof of Theorem 5.3.* First note that

$$\|u - \Phi_N \mathbf{u}_N\|_H = \|\mathcal{L}^{-1}f - \mathcal{L}^{-1}\Psi_N \mathbf{u}_N\|_H \leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(H^*, H)} \|f - \Psi_N \mathbf{u}_N\|_{H^*}. \quad (5.22)$$

By Theorem 5.1,  $\Psi = \mathcal{L}\Phi$  is a frame on  $H^*$ . Moreover, by definition,  $\mathbf{u}_N$  is the vector obtained by conducting an  $\epsilon$ -truncated SVD projection for frame expansion of  $f$ , as described in (5.2). Thus, by leveraging Theorem 3.7 in [4], we conclude the result.  $\square$

**Remark 5.5** (Convergence rates). *Suppose that the right-hand side  $f$  in (1.1) is smooth and has an asymptotic algebraic decay (or faster) when  $|x| \rightarrow \infty$ . As a rule of thumb, if one constructs a solution frame  $\Phi$  containing extended and weighted Jacobi or Zernike functions, centred on various affine-transformed intervals or balls, that cover the support of  $f$  prior to where the asymptotic decay dominates, then Theorem 5.3 indicates we expect spectral convergence to the solution of (1.1). The rough argument is that the induced frame  $\Psi$  for  $f$  includes functions that are piecewise polynomials prior to the asymptotic decay, with increasing degree as  $N \rightarrow \infty$ . Moreover,  $\Psi$  also includes functions that can match an asymptotic algebraic (or faster) decay. Proving such a conjecture is beyond the scope of this work, however, we numerically verify this observation in Section 7.*

## 6 Time-dependent problems

Implementing solvers for fractional-in-space time dependent problems is notoriously difficult due to the slow decay of the solutions and the accumulation of errors in the coefficients of the expansion. If the domain is truncated then the artificial boundary layers will eventually dominate the error in the solution. Similarly, although many time-stepping schemes are initially stable, the coefficients of the spatial expansion often become increasingly larger in magnitude leading to floating point cancellation error and a breakdown in the solution for large time  $t$ .

Our spectral method may be coupled with any Runge–Kutta (RK) method allowing for an arbitrary order method in time and spectral in space. Consider the fractional-in-space time-dependent equation:

$$\partial_t u + F(t, u) = 0, \quad u(x, 0) = u_0(x), \quad (6.1)$$

where  $u(\cdot, t) \in H^s(\mathbb{R})$  for a.e.  $t \geq 0$ . For instance, in the fractional heat equation, we have  $F(t, u) = (-\Delta)^s u(x, t)$  such that (6.1) becomes

$$\partial_t u(x, t) + (-\Delta)^s u(x, t) = 0, \quad u(x, 0) = u_0(x). \quad (6.2)$$

Now given  $u(x, t_j) = v_j(x)$  and some  $t_{j+1} = t_j + \delta t$ , an  $m$ -stage RK method approximates  $u(x, t_{j+1})$  with

$$u(x, t_{j+1}) \approx v_{j+1}(x) = v_j(x) + \delta t \sum_{i=1}^m b_i k_i(x), \quad (6.3)$$

where for all  $1 \leq i \leq m$  the stages  $k_i$  satisfy

$$k_i + F\left(t + c_i \delta t, u + \delta t \sum_{l=1}^m A_{il} k_l\right) = 0. \quad (6.4)$$

The coefficients  $b_i$ ,  $c_i$  and  $A_{il}$  are chosen so that the resulting method has the required degree of accuracy as well as other favourable properties e.g. stability. (6.4) is of the form (1.1). Thus we may use the frame expansion to discretize in space as described in Algorithm 2 for (6.2).

By considering (6.2), setting  $m = A_{11} = c_1 = b_1 = 1$  in (6.3)–(6.4), and multiplying the result by  $\delta t$ , one recovers the implicit Euler discretization of (6.2):

$$v_{j+1}(x) + \delta t (-\Delta)^s v_{j+1}(x) = v_j(x). \quad (6.5)$$

Consider the Hilbert space  $H$  such that  $(-\Delta)^s : H \rightarrow H^*$  and a truncated quasimatrix frame  $\Phi_N = (\phi_1, \dots, \phi_N)$  for  $H$ . Let  $\Phi_N^* = (\mathcal{I} + \delta t (-\Delta)^s) \Phi_N$ . Then, the implicit Euler discretization of (6.2) amounts to the following: given the coefficient vector,  $\mathbf{u}_j$ , at time-step  $j$ , the next approximate time-step,  $\mathbf{u}_{j+1}$  is

$$\mathbf{u}_{j+1} = \mathbf{u}_j^* \quad \text{where} \quad \|\Phi_N^*(x) \mathbf{u}_j^* - \Phi_N(x) \mathbf{u}_j\|_{H^*} \leq h_j \|\Phi_N(x) \mathbf{u}_j\|_{H^*}, \quad (6.6)$$

for some  $h_j > 0$ . We compute  $\mathbf{u}_j^*$  by an  $\epsilon$ -truncated SVD projection.

In the remainder of this section we show that an implicit Euler discretization in time coupled with a frame expansion in space, as described in Section 3, of the fractional heat equation, results in the expected convergence, i.e.  $\mathcal{O}(\delta t) + \mathcal{O}(h)$  where  $h = \max_j h_j$ . This result is confirmed in the numerical results in Section 7 and typically, for sufficiently smooth data and large  $M$ , we have  $h \leq \max\{c_1 \epsilon, \exp(-c_2 N)\}$  for some  $c_1, c_2 > 0$ .

---

**Algorithm 2**  $m$ -stage Runge–Kutta method for the fractional heat equation (6.2).

---

1: **Input:**

- |   |  |
|---|--|
| $A \in \mathbb{R}^{m \times m}$ , $b_i$ , $i \in \{1, \dots, m\}$ . | ▷ $m$ -stage Runge–Kutta coefficients.                                 |
| $\mathbf{u}_0$  | ▷ Coefficients for the initial state $u_0(\cdot)$ .                    |
| $X, X_*$  | ▷ Least-squares matrices for $\Phi_N(x)$ and $(-\Delta)^s \Phi_N(x)$ . |
| $\delta t, J$   | ▷ Time step and maximum time step iterations.                          |

2: Assemble  $X_A = I_m \otimes X + \delta t(A \otimes X_*)$ , where  $\otimes$  denotes the Kronecker product and  $I_m$  is the  $m \times m$  identity matrix.

3: **for**  $j = 1, 2, \dots, J$  **do**

4:    $\mathbf{y} \leftarrow X_* \mathbf{u}_{j-1}$ .

5:   Via an  $\epsilon$ -truncated SVD projection, compute

$$\mathbf{k}_A \approx \operatorname{argmin}_{\mathbf{v} \in \mathbb{C}^{Nm}} \|X_A \mathbf{v} - \underbrace{(\mathbf{y}^\top \cdots \mathbf{y}^\top)^\top}_{m \text{ repeats}}\|_{\ell^2}.$$

6:   Extract the  $m$  vectors  $\{\mathbf{k}_i\}_{i=1}^m$ ,  $\mathbf{k}_A = (\mathbf{k}_1^\top \cdots \mathbf{k}_m^\top)^\top$ .

7:    $\mathbf{u}_j \leftarrow \mathbf{u}_{j-1} + \delta t \sum_{i=1}^m b_i \mathbf{k}_i$ .

8: **end for**

---

**Lemma 6.1.** Consider the Hilbert space  $H_\omega^s(\mathbb{R})$ , where  $\omega(x) = 1$  in  $\Omega \subseteq \mathbb{R}$ , for an arbitrary open interval  $\Omega$ , and  $\omega(x) = 0$  in  $\mathbb{R} \setminus \Omega$ . Let  $\mathcal{L} = \mathcal{I} + (-\Delta)^s$ . Fix any  $u^* \in H_\omega^s(\mathbb{R})$  and also denote by  $u^* \in H_\omega^s(\mathbb{R})^*$  the embedding of  $u^*$  into  $H_\omega^s(\mathbb{R})^*$ . Then, we have that

$$\|u^*\|_{H_\omega^s(\mathbb{R})^*} \leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(H_\omega^s(\mathbb{R}), H_\omega^s(\mathbb{R}))} \|u^*\|_{H_\omega^s(\mathbb{R})}. \quad (6.7)$$

*Proof.* Consider any  $u^* \in H_\omega^s(\mathbb{R})^*$ . There exists a unique  $u \in H_\omega^s(\mathbb{R})$  such that, for all  $v \in H_\omega^s(\mathbb{R})$  [27, Sec. D.2, Th. 2],

$$(u, v)_{H_\omega^s(\mathbb{R})} = \langle \mathcal{L}u, v \rangle_{H_\omega^s(\mathbb{R})^*, H_\omega^s(\mathbb{R})} = \langle u^*, v \rangle_{H_\omega^s(\mathbb{R})^*, H_\omega^s(\mathbb{R})}, \quad (6.8)$$

and  $\|u^*\|_{H_\omega^s(\mathbb{R})^*} = \|u\|_{H_\omega^s(\mathbb{R})}$ . Hence, the Riesz map is  $u = \mathcal{L}^{-1}u^*$ . Thus,

$$\|u^*\|_{H_\omega^s(\mathbb{R})^*} = \|u\|_{H_\omega^s(\mathbb{R})} = \|\mathcal{L}^{-1}u^*\|_{H_\omega^s(\mathbb{R})} \leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(H_\omega^s(\mathbb{R}), H_\omega^s(\mathbb{R}))} \|u^*\|_{H_\omega^s(\mathbb{R})}. \quad (6.9)$$

□

**Theorem 6.1** (Implicit Euler convergence). Consider the fractional heat equation (6.2) and choose the uniform time discretization points  $t_0, \dots, t_J$ . Define  $\delta t = t_1 - t_0$  and let  $\Phi_N(x)$  denote the truncated quasimatrix of a frame on the Hilbert space  $H = H_\omega^s(\mathbb{R})$  with  $H_\omega^s(\mathbb{R})$  as defined in Lemma 6.1. Let  $\mathcal{L}_{\delta t} = (\mathcal{I} + \delta t(-\Delta)^s)$  which corresponds to an implicit Euler discretization in time. Fix  $\Phi_N^*(x) = \mathcal{L}_{\delta t} \Phi_N(x)$ . Suppose that  $\|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)}$  and  $\|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)}$  are bounded and there exists a Lipschitz constant  $L > 0$  such that for any  $w_1, w_2 \in H$ ,  $\|(-\Delta)^s(w_1 - w_2)\|_{H^*} \leq L\|w_1 - w_2\|_H$ . Fix the number of collocation points  $M$  and the required SVD tolerance  $\epsilon$ . Define  $h = \max_{j \in \{0, \dots, J\}} h_j$  with  $h_j$  being defined as in (6.6). We denote the exact solution of (6.2) by  $u(x, t)$ . Suppose that we have a  $\mathbf{u}_0$  and  $0 < h_0 \leq h < 1$  such that

$$\|u_0 - \Phi_N \mathbf{u}_0\|_H \leq h_0 \|u_0\|_H. \quad (6.10)$$

Then, for some  $C_0 > 0$ , the following error bound holds:

$$\|u(\cdot, t_j) - \Phi_N \mathbf{u}_j\|_H \leq C_0 \delta t + C_1(u_0, j) C_2(h, j) h, \quad (6.11)$$

where  $C_2(h, j) = \sum_{i=0}^j (1+h)^i$  and  $C_1(u_0, j) = \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)}^j \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)}^j \|u_0\|_H$ .

*Proof.* Let  $u_j(x)$ ,  $j \in \{0, \dots, J\}$ , denote the solutions of

$$\frac{u_{j+1}(x) - u_j(x)}{\delta t} + (-\Delta)^s u_{j+1}(x) = 0, \quad u_0(x) = u(x, t_0). \quad (6.12)$$

By assumption  $(-\Delta)^s : H \rightarrow H^*$  is Lipschitz continuous and thus there exists a constant  $C_0 > 0$  independent of  $\delta t$  such that

$$\|u(\cdot, t_j) - u_j\|_H \leq C_0 \delta t. \quad (6.13)$$

(6.13) follows from classical results for implicit Euler discretizations, e.g. [28, Sec. 5] where the Euclidian norms have been replaced by  $\|\cdot\|_H$  and  $\|\cdot\|_{H^*}$  as appropriate. Hence,

$$\|u(\cdot, t_j) - \Phi_N \mathbf{u}_j\|_H \leq C_0 \delta t + \|u_j - \Phi_N \mathbf{u}_j\|_H. \quad (6.14)$$

Now, by recalling the implicit Euler time step (6.6) and utilizing Lemma 6.1,

$$\begin{aligned} \|u_j - \Phi_N \mathbf{u}_j\|_H &= \|\mathcal{L}_{\delta t}^{-1}(\mathcal{L}_{\delta t} u_j - \mathcal{L}_{\delta t} \Phi_N \mathbf{u}_j)\|_H \\ &\leq \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)} \|u_{j-1} - \Phi_N^* \mathbf{u}_j\|_{H^*} \\ &\leq \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)} (\|\Phi_N \mathbf{u}_{j-1} - \Phi_N^* \mathbf{u}_{j-1}^*\|_{H^*} + \|u_{j-1} - \Phi_N \mathbf{u}_{j-1}\|_{H^*}) \\ &\leq \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)} (h \|\Phi_N \mathbf{u}_{j-1}\|_{H^*} + \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)} \|u_{j-1} - \Phi_N \mathbf{u}_{j-1}\|_H) \\ &\leq h \sum_{i=0}^{j-1} \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)}^{j-i} \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)}^{j-i-1} \|\Phi_N \mathbf{u}_i\|_{H^*} \\ &\quad + \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)}^j \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)}^j \|u_0 - \Phi_N \mathbf{u}_0\|_H. \end{aligned} \quad (6.15)$$

We note that, for any  $i \in \{0, 1, \dots, j-1\}$ ,

$$\begin{aligned} \|\Phi_N \mathbf{u}_i\|_{H^*} &\leq \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)} \|\Phi_N \mathbf{u}_i\|_H = \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)} \|\Phi_N \mathbf{u}_{i-1}^*\|_{H^*} \\ &\leq \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)} \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)} \|\Phi_N^* \mathbf{u}_{i-1}^*\|_{H^*} \\ &\leq (1+h) \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)} \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H^*, H)} \|\Phi_N \mathbf{u}_{i-1}\|_{H^*} \\ &\leq (1+h)^i \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)}^i \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H, H^*)}^i \|\Phi_N \mathbf{u}_0\|_{H^*} \\ &\leq (1+h)^{i+1} \|\mathcal{L}_1^{-1}\|_{\mathcal{B}(H, H)}^{i+1} \|\mathcal{L}_{\delta t}^{-1}\|_{\mathcal{B}(H, H^*)}^i \|u_0\|_H, \end{aligned} \quad (6.16)$$

By assumption  $\|u_0 - \Phi_N \mathbf{u}_0\|_H \leq h \|u_0\|_H$ . Hence, by utilizing the bound in (6.16) for (6.15) and the subsequent bound of  $\|u_j - \Phi_N \mathbf{u}_j\|_H$  into (6.14), we deduce the result.  $\square$

## 7 Examples

In this section we provide several numerical examples.

**Code availability:** For reproducibility, an implementation of the solver as well as scripts to generate the plots and solutions can be found in FractionalFrames.jl [29] and archived on Zenodo [49]. The implementation is written in Julia [10] and heavily relies on the ClassicalOrthogonalPolynomials.jl [19] and HypergeometricFunctions.jl [36] packages.

## 7.1 The Gaussian

In our first example we consider the exact solution  $u(x) = e^{-x^2}$ . We note that [52, Prop. 4.2],

$$(\mathcal{I} + (-\Delta)^s)u(x) = e^{-x^2} + 4^s \frac{\Gamma(s + 1/2)}{\Gamma(1/2)} {}_1F_1(s + 1/2; 1/2; -x^2), \quad (7.1)$$

where  ${}_1F_1$  denotes the Kummer confluent hypergeometric function [47, Sec. 16.2]. We investigate (7.1) when  $s = 1/3$ .

We pick  $\bigcup_{k=1}^5 \{\tilde{P}_n^{I_k, (-1/3, -1/3)}\}_{n=0}^\infty \cup \{Q_n^{I_k, (1/3, 1/3)}\}_{n=0}^\infty$ , as the solution family of functions, where  $I_1, \dots, I_5$  are  $[-5, -3]$ ,  $[-3, -1]$ ,  $[-1, 1]$ ,  $[1, 3]$ , and  $[3, 5]$ , respectively.  $\tilde{P}_n^{I_k}(x)$  and  $Q_n^{I_k}(x)$  denote the affine-transformed extended Jacobi functions and weighted Jacobi polynomials, respectively, as defined in Section 4.1 and Definition 4.1. As the induced family of functions, for the right-hand side expansion, contains functions that are undefined for  $x \in \{-5, -3, -1, 1, 3, 5\}$ , we choose 5001 equally spaced points in  $[a + \epsilon, b - \epsilon]$ ,  $\epsilon = 10^{-2}$ , where  $a, b$  represent the endpoints of each interval as well as 5001 equally spaced points in  $[-10 + \epsilon, -5 - \epsilon]$  and  $[5 + \epsilon, 10 - \epsilon]$ . This results in 35,007 collocation points.

A semi-log plot of the convergence for the right-hand side and the solution is given in Fig. 1a. The error is measured in the  $\ell^\infty$ -norm as measured at the collocation points. We observe approximately spectral convergence as we include more functions in the expansion, with the right-hand side error stagnating at around  $\mathcal{O}(10^{-14})$  and the solution error at  $\mathcal{O}(10^{-11})$ .

Since we are not guaranteed that the family of functions is a frame on all  $\mathbb{R}$ , in Fig. 1b we investigate the behaviour of the expansion coefficients of the right-hand side. In particular we plot the  $\ell^\infty$ -norm of the coefficient vector as we increase the number of functions in the expansion. Despite the lack of a strict frame condition, we achieve bounded coefficients of magnitude  $\mathcal{O}(1)$  for  $N > 50$ .




Figure 1: (Left)  $\ell^\infty$ -norm error in the right-hand side and solution of the Jacobi approximation for (7.1) with  $s = 1/3$ . We observe approximate spectral convergence with an error stagnation at around  $\mathcal{O}(10^{-14})$  for the right-hand side and  $\mathcal{O}(10^{-11})$  for the solution. (Right)  $\ell^\infty$ -norm of the coefficient vector of the expansion. We observe that coefficient norm is small.



## 7.2 Multiple exponents

In this section we consider an equation with fractional Laplacians with different exponents. Consider the equation:

$$(\mathcal{I} + (-\Delta)^{1/3} + (-\Delta)^{1/5})u(x) = e^{-x^2} + (-\Delta)^{1/3}e^{-x^2} + (-\Delta)^{1/5}e^{-x^2}. \quad (7.2)$$

As in the previous example, the exact solution to this equation is  $u(x) = e^{-x^2}$ . The sum space we consider is  $\bigcup_{k=1}^5 \{\tilde{P}_j^{I_k, (-1/4, -1/4)}\}_{j=0}^\infty \cup \{Q_j^{I_k, (1/4, 1/4)}\}_{j=0}^\infty$  where the intervals  $I_1, \dots, I_5$  and collocation points are the same as in Section 7.1.

A semi-log plot of the convergence for the right-hand side and the solution is given in Fig. 2a. The error is measured in the  $\ell^\infty$ -norm as measured at the collocation points. We observe approximately spectral convergence until  $N = 150$  where the convergence rate stagnates. The best right-hand side error is  $\mathcal{O}(10^{-13})$  and the best solution error is  $\mathcal{O}(10^{-10})$ . We observe that as  $N$  increases the error becomes worse in the solution. As in the previous example we also plot the  $\ell^\infty$ -norm of the coefficient vector in Fig. 2b and observe that the magnitude is small, reaching  $\mathcal{O}(1)$  when  $N > 50$ . We expect that fine-tuning the weight parameters in the Jacobi polynomials would allow one to achieve faster convergence.




Figure 2: (Left) Pointwise error in the right-hand side and solution of the Jacobi approximation for (7.2). We observe approximate spectral convergence with an error stagnation at around  $\mathcal{O}(10^{-13})$  for the right-hand side and  $\mathcal{O}(10^{-10})$  for the solution. (Right)  $\ell^\infty$ -norm of the coefficient vector of the expansion. We observe that coefficient norm is small.

## 7.3 Two-dimensional Gaussian

In this example we exemplify that the spectral method extends to two-dimensional problems. Consider the equation

$$(-\Delta)^{1/2}u(x, y) = 2\Gamma(3/2)_1F_1(3/2; 1; -x^2 - y^2), \quad (7.3)$$

which has the exact solution  $u(x, y) = \exp(-x^2 - y^2)$  [52, Prop. 4.2]. We utilize a solution family of functions consisting of weighted and extended Zernike polynomials and functions (as introduced in Section 4.3), respectively, that are scaled radially on a sequence of nested disks. Let

$W_{n,m,j}^{(b)}(x, y) := (1 - r^2)^b Z_{n,m,j}^{(b)}(x, y)$ . Since the right-hand side and the solution are rotationally invariant, both functions are best approximated by the family of functions restricted to the  $(m, j) = (0, 1)$  Fourier mode and sign. Thus we use the sum space

$$\bigcup_{k=1}^K \{W_{n,0,1}^{(1/2)}(a_k x, a_k y)\} \cup \{\tilde{Z}_{n,0,1}^{(-1/2,-1/2)}(a_k x, a_k y)\}, \quad a_k \in \{1, 3/2, 2, 3, 4\}. \quad (7.4)$$

As the sum space contains functions that are undefined for  $r \in \{0, 1, 3/2, 2, 3, 4\}$ , we choose 1001 equally spaced points in the radial direction  $[a + \epsilon, b - \epsilon]$ ,  $\epsilon = 10^{-3}$ , where  $a, b$  represent the endpoints of each interval as well as 1001 equally spaced points in  $[4 + \epsilon, 10 - \epsilon]$ . We then cross product these radial collocation points with 30 equally spaced angular collocation points between 0 to  $2\pi$ . This results in 180,180 collocation points. A semi-log plot of the convergence for the solution is given in Fig. 3a. The error is measured in the  $\ell^\infty$ -norm as measured at the collocation points. The best solution error is  $\mathcal{O}(10^{-9})$ . We plot the  $\ell^\infty$ -norm of the coefficient vector in Fig. 3b and observe that the magnitude is  $\mathcal{O}(1)$ .




Figure 3: (Left) Pointwise error in the solution of the Zernike approximation for (7.3). We observe approximate spectral convergence with an error stagnation at around  $\mathcal{O}(10^{-9})$ . (Right)  $\ell^\infty$ -norm of the coefficient vector of the expansion. We observe that coefficient norm is small.

## 7.4 Fractional heat equation

We consider the time-dependent fractional heat equation (6.2) with the initial state  $u(x, 0) = (1 + x^2)^{-1}$  and  $s = 1/2$ . The explicit solution is  $u(x, t) = (1 + t)/(x^2 + (1 + t)^2)$ . We consider the time discretizations backward Euler (1<sup>st</sup> order), implicit midpoint rule (2<sup>nd</sup> order), Gauss–Legendre (4<sup>th</sup> order), and Gauss–Legendre (6<sup>th</sup> order) via Algorithm 2.

Our goal is to examine the convergence of the temporal discretization. We utilize the solution family of functions  $\bigcup_{k=1}^5 \{\tilde{P}_j^{I_k, (-1/2, -1/2)}\}_{j=1}^\infty \cup \{Q_j^{I_k, (1/2, 1/2)}\}_{j=0}^\infty$  where the intervals  $I_1, \dots, I_5$  are  $[-5, -3]$ ,  $[-3, -1]$ ,  $[-1, 1]$ ,  $[1, 3]$ , and  $[3, 5]$ , respectively. We choose 5001 equally spaced points in  $[a + \epsilon, b - \epsilon]$ ,  $\epsilon = 10^{-4}$ , where  $a, b$  represent the endpoints of each interval as well as 5001 equally spaced points in  $[-20 + \epsilon, -5 - \epsilon]$  and  $[5 + \epsilon, 20 - \epsilon]$ . This results in 35,007 collocation points. We include 250 functions in the family of functions. Expanding the initial state results in an  $\ell^\infty$ -norm error of  $3.16 \times 10^{-12}$  as measured at the collocation points. Next we measure the

relative maximum  $\ell^\infty$ -norm as measured at all the collocation points and time steps from  $t = 0$  to  $t = 1$  for time step choices  $\delta t \in \{10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}\}$ , i.e.

$$\max_{k \in \{0, 1, 2, \dots, \delta t^{-1}\}} \max_{x \in X_c} \frac{|u(x, k\delta t) - \Phi(x)\mathbf{v}_k|}{|u(x, k\delta t)|}, \quad (7.5)$$

where  $X_c$  is the set of the collocation points. The convergence results are given in the left panel in Fig. 4. We observe the expected order of convergence for the methods, with deviations occurring close to  $\mathcal{O}(10^{-10})$  where the error in the spatial discretization likely causes the degradation in the convergence. This example is numerical evidence for the result provided in Theorem 6.1. Moreover, it appears that the result extends to all implicit Runge–Kutta methods and their classical associated convergence rates. In the right panel of Fig. 4 we plot the pointwise error of the solution, computed via the 6<sup>th</sup> order Gauss–Legendre method with  $\delta t = 10^{-2}$ , at  $t = 1$  in the domain  $x \in [-10^3, 10^3]$ . We observe that the algebraic tails of the solution are accurately captured by this spectral method.




Figure 4: (Left) Convergence of the four Runge–Kutta methods for decreasing  $\delta t$  using (7.5) to measure the error. The red, blue, orange, and magenta dashed lines indicate  $\mathcal{O}(\delta t)$ ,  $\mathcal{O}(\delta t^2)$ ,  $\mathcal{O}(\delta t^4)$ , and  $\mathcal{O}(\delta t^6)$  convergence rates, respectively. We observe the expected convergence rates of all four methods. We note that the stagnation of the convergence rate at  $\mathcal{O}(10^{-10})$  is likely caused by the error in the spatial discretization. (Right) Pointwise error of the approximated solution, computed via the 6<sup>th</sup>-order Gauss–Legendre method with  $\delta t = 10^{-2}$ , at  $t = 1$  in the domain  $x \in [-10^3, 10^3]$ . We see that the algebraic tails are suitably captured.

## 7.5 Variable exponent $s$

Consider the heat equation (6.2), except now the exponent of the fractional Laplacian also evolves in time [51], i.e.

$$\partial_t u(x, t) + (-\Delta)^{s(t)} u(x, t) = 0, \quad u(x, 0) = (1 + x^2)^{-1}. \quad (7.6)$$

In this example we pick  $s(t) = s^*(t) := 1/2 - t/3$ . We choose the same intervals, collocation points, and truncation degree as in Section 7.4. Utilizing an implicit midpoint rule for the temporal discretization with  $\delta t = 10^{-2}$ , we consider the family of functions  $\bigcup_{k=1}^5 \{\tilde{P}_n^{I_k, (-s(t_n), -s(t_n))}\}_{n=1}^\infty \cup \{Q_n^{I_k, (s(t_n), s(t_n))}\}_{n=0}^\infty$ ,  $t_j = j\delta t$ ,  $j = 0, 1, \dots, 100$ , at each time step. In Fig. 5 we plot the approximation of the solution of (7.6) at  $t = 1$  together with the equivalent solutions if  $s(t) = 1/2$ ,

$1/3$ , and  $1/6$  for all  $t$ . We see that the decay of the variable exponent solution is similar to the decay where  $s = 1/3$  which is equal to  $s^*(1/2)$ .




Figure 5: The approximation solutions of (7.6) where  $s(t) = 1/2, 1/2 - t/3, 1/3$ , and  $1/6$  at  $t = 1$ .

## 8 Conclusions

In this work, we introduced a frame approach, summarized in Algorithm 1, for solving equations involving nonlocal and fractional terms such as the fractional Laplacian  $(-\Delta)^s$ ,  $s \in (0, 1)$ . One fixes a family of functions for the expansion of the solution and expands the right-hand side of the equation in the space induced by applying the equation operator to this family of functions. This diagonalizes the equation and reduces the problem from a solve of a nonlocal problem to an interpolation problem.

Under suitable conditions, we showed that if the family of functions for the solution expansion is a frame in a suitable Hilbert space, then the induced family of functions for the right-hand side expansion is a frame in the dual of the Hilbert space. By considering solution expansions in weighted Jacobi and Zernike polynomials, we utilized recent results to deduce the family of functions for the right-hand side expansion. With this knowledge, one automatically recovers favourable properties of frames, in particular, the utilization of a truncated SVD projection to find a well-posed expansion. This culminated in deriving an a priori estimate in Theorem 5.3.

This spatial discretization was coupled with Runge–Kutta methods in time in order to develop a method for time-dependent problems including the fractional heat equation as well as a problem where the exponent of the fractional Laplacian evolved in time. We proved that one recovers the optimal convergence rates under an implicit Euler discretization and observed the expected convergence rates for our chosen Runge–Kutta methods in practice, up to a 6<sup>th</sup> order method. Moreover, the method was also competitive when applied to a two-dimensional problem.

This approach is general and is not limited to the operators considered in this work. Provided one has explicit expressions of the equation operator applied to a family of functions, one may use this frame approach for a considerable number of different problems. Future work will consider generalizing this technique to be competitive in higher dimensions. The bottleneck inevitably becomes the SVD factorization to interpolate the right-hand side. However, by using randomized SVD solvers [33, 45] and the AZ-algorithm [20], we hope to considerably reduce the computational complexity.

## Acknowledgments

IPAP, TSG and SO were supported by the EPSRC grant EP/T022132/1 “Spectral element methods for fractional differential equations, with applications in applied analysis and medical imaging”. IPAP and SO were also supported by the Leverhulme Trust Research Project Grant RPG-2019-144 “Constructive approximation theory on and inside algebraic curves and surfaces”. IPAP was also supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – The Berlin Mathematics Research Center MATH+ (EXC-2046/1, project ID: 390685689). TSG was also supported by a PIMS-Simons postdoctoral fellowship, jointly funded by the Pacific Institute for the Mathematical Sciences and the Simons Foundation. JAC was supported by the Advanced Grant Nonlocal-CPD (Nonlocal PDEs for Complex Particle Dynamics: Phase Transitions, Patterns and Synchronization) of the European Research Council Executive Agency (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 883363).

## References

- [1] G. Acosta and J. P. Borthagaray, “A fractional Laplace equation: Regularity of solutions and finite element approximations,” *SIAM Journal on Numerical Analysis*, vol. 55, no. 2, pp. 472–495, 2017. DOI: [10.1137/15m1033952](https://doi.org/10.1137/15m1033952).
- [2] R. A. Adams and J. J. Fournier, *Sobolev spaces*, Second. Elsevier, 2003, ISBN: 978-0-12-044143-3.
- [3] B. Adcock and D. Huybrechs, “Frames and numerical approximation,” *SIAM Review*, vol. 61, no. 3, pp. 443–473, 2019. DOI: [10.1137/17M1114697](https://doi.org/10.1137/17M1114697).
- [4] B. Adcock and D. Huybrechs, “Frames and numerical approximation II: Generalized sampling,” *Journal of Fourier Analysis and Applications*, vol. 26, no. 6, pp. 1–34, 2020. DOI: [10.1007/s00041-020-09796-w](https://doi.org/10.1007/s00041-020-09796-w).
- [5] M. Ainsworth and C. Glusa, “Towards an efficient finite element method for the integral fractional Laplacian on polygonal domains,” in *Contemporary Computational Mathematics - A Celebration of the 80th Birthday of Ian Sloan*, Springer International Publishing, 2018, pp. 17–57. DOI: [10.1007/978-3-319-72456-0\\_2](https://doi.org/10.1007/978-3-319-72456-0_2).
- [6] H. Antil and S. Bartels, “Spectral approximation of fractional PDEs in image processing and phase field modeling,” *Computational Methods in Applied Mathematics*, vol. 17, no. 4, pp. 661–678, 2017. DOI: [10.1515/cmam-2017-0039](https://doi.org/10.1515/cmam-2017-0039).
- [7] R. Bailo, J. A. Carrillo, S. Fronzoni, and D. Gómez-Castro, *A finite-volume scheme for fractional diffusion on bounded domains*, 2023. arXiv preprint: [2309.08283](https://arxiv.org/abs/2309.08283) [[math.NA](https://arxiv.org/html/math)].
- [8] G. R. Baker, X. Li, and A. C. Morlet, “Analytic structure of two 1D-transport equations with nonlocal fluxes,” *Physica D: Nonlinear Phenomena*, vol. 91, no. 4, pp. 349–375, 1996. DOI: [10.1016/0167-2789\(95\)00271-5](https://doi.org/10.1016/0167-2789(95)00271-5).
- [9] D. A. Benson, S. W. Wheatcraft, and M. M. Meerschaert, “Application of a fractional advection-dispersion equation,” *Water Resources Research*, vol. 36, no. 6, pp. 1403–1412, 2000. DOI: [10.1029/2000WR900031](https://doi.org/10.1029/2000WR900031).

- [10] J. Bezanson, A. Edelman, S. Karpinski, and V. B. Shah, “Julia: A fresh approach to numerical computing,” *SIAM Review*, vol. 59, no. 1, pp. 65–98, 2017. DOI: [10.1137/141000671](https://doi.org/10.1137/141000671).
- [11] A. Bonito, J. P. Borthagaray, R. H. Nochetto, E. Otárola, and A. J. Salgado, “Numerical methods for fractional diffusion,” *Computing and Visualization in Science*, vol. 19, no. 5, pp. 19–46, 2018. DOI: [10.1007/s00791-018-0289-y](https://doi.org/10.1007/s00791-018-0289-y).
- [12] A. Bonito, J. P. Borthagaray, R. H. Nochetto, E. Otárola, and A. J. Salgado, “Numerical methods for fractional diffusion,” *Computing and Visualization in Science*, vol. 19, no. 5-6, pp. 19–46, 2018. DOI: [10.1007/s00791-018-0289-y](https://doi.org/10.1007/s00791-018-0289-y).
- [13] N. Bournaveas and V. Calvez, “The one-dimensional Keller-Segel model with fractional diffusion of cells,” *Nonlinearity*, vol. 23, no. 4, pp. 923–935, 2010, ISSN: 0951-7715. DOI: [10.1088/0951-7715/23/4/009](https://doi.org/10.1088/0951-7715/23/4/009).
- [14] L. Caffarelli and J. Vázquez, “Regularity of solutions of the fractional porous medium flow with exponent  $1/2$ ,” *St. Petersburg Mathematical Journal*, vol. 27, no. 3, pp. 437–460, 2016. DOI: [10.1090/spmj/1397](https://doi.org/10.1090/spmj/1397).
- [15] B. Carmichael, H. Babahosseini, S. N. Mahmoodi, and M. Agah, “The fractional viscoelastic response of human breast tissue cells,” *Physical Biology*, vol. 12, no. 4, p. 046001, 2015. DOI: [10.1088/1478-3975/12/4/046001](https://doi.org/10.1088/1478-3975/12/4/046001).
- [16] J. Cayama, C. M. Cuesta, and F. de la Hoz, “Numerical approximation of the fractional Laplacian on  $\mathbb{R}$  using orthogonal families,” *Applied Numerical Mathematics*, vol. 158, pp. 164–193, 2020. DOI: [10.1016/j.apnum.2020.07.024](https://doi.org/10.1016/j.apnum.2020.07.024).
- [17] D. Chae, A. Córdoba, D. Córdoba, and M. A. Fontelos, “Finite time singularities in a 1D model of the quasi-geostrophic equation,” *Advances in Mathematics*, vol. 194, no. 1, pp. 203–223, 2005. DOI: [10.1016/j.aim.2004.06.004](https://doi.org/10.1016/j.aim.2004.06.004).
- [18] S. Chen, J. Shen, and L.-L. Wang, “Laguerre functions and their applications to tempered fractional differential equations on infinite intervals,” *Journal of Scientific Computing*, vol. 74, no. 3, pp. 1286–1313, 2018. DOI: [10.1007/s10915-017-0495-7](https://doi.org/10.1007/s10915-017-0495-7).
- [19] *ClassicalOrthogonalPolynomials.jl*, version 0.7.2, 2023. [Online]. Available: <https://github.com/JuliaApproximation/ClassicalOrthogonalPolynomials.jl>.
- [20] V. Coppé, D. Huybrechs, R. Matthysen, and M. Webb, “The AZ algorithm for least squares systems with a known incomplete generalized inverse,” *SIAM Journal on Matrix Analysis and Applications*, vol. 41, no. 3, pp. 1237–1259, 2020. DOI: [10.1137/19M1306385](https://doi.org/10.1137/19M1306385).
- [21] A. Córdoba, D. Córdoba, and M. A. Fontelos, “Integral inequalities for the Hilbert transform applied to a nonlocal transport equation,” *Journal de Mathématiques Pures et Appliquées*, vol. 86, no. 6, pp. 529–540, 2006. DOI: [10.1016/j.matpur.2006.08.002](https://doi.org/10.1016/j.matpur.2006.08.002).
- [22] N. Cusimano, F. del Teso, L. Gerardo-Giorda, and G. Pagnini, “Discretizations of the spectral fractional Laplacian on general domains with Dirichlet, Neumann, and Robin boundary conditions,” *SIAM Journal on Numerical Analysis*, vol. 56, no. 3, pp. 1243–1272, 2018, ISSN: 0036-1429. DOI: [10.1137/17M1128010](https://doi.org/10.1137/17M1128010).
- [23] M. D’Elia, Q. Du, C. Glusa, M. Gunzburger, X. Tian, and Z. Zhou, “Numerical methods for nonlocal and fractional models,” *Acta Numerica*, vol. 29, pp. 1–124, 2020. DOI: [10.1017/S096249292000001X](https://doi.org/10.1017/S096249292000001X).

- [24] E. Di Nezza, G. Palatucci, and E. Valdinoci, “Hitchhiker’s guide to the fractional Sobolev spaces,” *Bulletin des Sciences Mathématiques*, vol. 136, no. 5, pp. 521–573, 2012. DOI: [10.1016/j.bulsci.2011.12.004](https://doi.org/10.1016/j.bulsci.2011.12.004).
- [25] Q. Du, *Nonlocal Modeling, Analysis, and Computation*. SIAM, 2019, ISBN: 978-1-61197-561-1. DOI: [10.1137/1.9781611975628](https://doi.org/10.1137/1.9781611975628).
- [26] C. Escudero, “The fractional Keller-Segel model,” *Nonlinearity*, vol. 19, no. 12, pp. 2909–2918, 2006, ISSN: 0951-7715. DOI: [10.1088/0951-7715/19/12/010](https://doi.org/10.1088/0951-7715/19/12/010).
- [27] L. C. Evans, *Partial Differential Equations*, 2nd ed. American Mathematical Society, 2010, ISBN: 978-0821849743.
- [28] I. Faragó, “Note on the convergence of the implicit Euler method,” in *Numerical Analysis and its Applications: 5th International Conference, NAA 2012, Lozenetz, Bulgaria, June 15-20, 2012, Revised Selected Papers 5*, Springer, 2013, pp. 1–11. DOI: [10.1007/978-3-642-41515-9\\_1](https://doi.org/10.1007/978-3-642-41515-9_1).
- [29] *FractionalFrames.jl*, version 0.0.3, 2024. [Online]. Available: <https://github.com/ioannisPpapadopoulos/FractionalFrames.jl>.
- [30] W. Gautschi, “Minimal solutions of three-term recurrence relations and orthogonal polynomials,” *Mathematics of Computation*, vol. 36, no. 154, pp. 547–554, 1981. DOI: [10.1090/S0025-5718-1981-0606512-6](https://doi.org/10.1090/S0025-5718-1981-0606512-6).
- [31] W. Gautschi, *Orthogonal polynomials: computation and approximation*. Oxford University Press, Apr. 2004, ISBN: 9780198506720. DOI: [10.1093/oso/9780198506720.001.0001](https://doi.org/10.1093/oso/9780198506720.001.0001).
- [32] T. S. Gutleb and I. P. A. Papadopoulos, *Explicit fractional Laplacians and Riesz potentials of classical functions*, 2023. arXiv preprint: [2311.10896 \[math.NA\]](https://arxiv.org/abs/2311.10896).
- [33] N. Halko, P.-G. Martinsson, and J. A. Tropp, “Finding structure with randomness: Probabilistic algorithms for constructing approximate matrix decompositions,” *SIAM Review*, vol. 53, no. 2, pp. 217–288, 2011. DOI: [10.1137/090771806](https://doi.org/10.1137/090771806).
- [34] Y. Hatano and N. Hatano, “Dispersive transport of ions in column experiments: An explanation of long-tailed profiles,” *Water Resources Research*, vol. 34, no. 5, pp. 1027–1033, 1998. DOI: [10.1029/98WR00214](https://doi.org/10.1029/98WR00214).
- [35] Y. Huang and A. Oberman, “Numerical methods for the fractional Laplacian: A finite difference-quadrature approach,” *SIAM Journal on Numerical Analysis*, vol. 52, no. 6, pp. 3056–3084, 2014. DOI: [10.1137/140954040](https://doi.org/10.1137/140954040).
- [36] *HypergeometricFunctions.jl*, version 0.3.10, 2023. [Online]. Available: <https://github.com/JuliaMath/HypergeometricFunctions.jl>.
- [37] F. W. King, *Hilbert Transforms: Volume 1* (Encyclopedia of Mathematics and its Applications). Cambridge University Press, 2009, vol. 1. DOI: [10.1017/CB09780511721458](https://doi.org/10.1017/CB09780511721458).
- [38] M. Kwaśnicki, “Ten equivalent definitions of the fractional Laplace operator,” *Fractional Calculus and Applied Analysis*, vol. 20, no. 1, pp. 7–51, 2017. DOI: [10.1515/fca-2017-0002](https://doi.org/10.1515/fca-2017-0002).
- [39] L. Lafleche and S. Salem, “Fractional Keller-Segel equation: Global well-posedness and finite time blow-up,” *Communications in Mathematical Sciences*, vol. 17, no. 8, pp. 2055–2087, 2019, ISSN: 1539-6746. DOI: [10.4310/CMS.2019.v17.n8.a1](https://doi.org/10.4310/CMS.2019.v17.n8.a1).



- [40] D. Li and J. Rodrigo, “Finite-time singularities of an aggregation equation in  $\mathbb{R}^n$  with fractional dissipation,” *Communications in Mathematical Physics*, vol. 287, no. 2, pp. 687–703, 2009, ISSN: 0010-3616. DOI: [10.1007/s00220-008-0669-0](https://doi.org/10.1007/s00220-008-0669-0).
- [41] H. Li, R. Liu, and L.-L. Wang, “Efficient Hermite spectral-Galerkin methods for nonlocal diffusion equations in unbounded domains,” 2021. [Online]. Available: <https://personal.ntu.edu.sg/lilian/HermiteNonlocal-Submitted.pdf>.
- [42] A. Lischke *et al.*, “What is the fractional Laplacian? A comparative review with new results,” *Journal of Computational Physics*, vol. 404, p. 109 009, 2020. DOI: [10.1016/j.jcp.2019.109009](https://doi.org/10.1016/j.jcp.2019.109009).
- [43] Z. Mao and J. Shen, “Hermite spectral methods for fractional PDEs in unbounded domains,” *SIAM Journal on Scientific Computing*, vol. 39, no. 5, A1928–A1950, 2017. DOI: [10.1137/16M1097109](https://doi.org/10.1137/16M1097109).
- [44] M. Meier and Y. Nakatsukasa, *Randomized algorithms for Tikhonov regularization in linear least squares*, 2022. arXiv preprint: [2203.07329](https://arxiv.org/abs/2203.07329) [math.NA].
- [45] M. Meier, Y. Nakatsukasa, A. Townsend, and M. Webb, *Are sketch-and-precondition least squares solvers numerically stable?* 2023. arXiv preprint: [2302.07202](https://arxiv.org/abs/2302.07202) [math.NA].
- [46] R. H. Nochetto, E. Otárola, and A. J. Salgado, “A PDE approach to fractional diffusion in general domains: A priori error analysis,” *Foundations of Computational Mathematics*, vol. 15, no. 3, pp. 733–791, 2015, ISSN: 1615-3375. DOI: [10.1007/s10208-014-9208-x](https://doi.org/10.1007/s10208-014-9208-x).
- [47] F. W. J. Olver *et al.*, *NIST Digital Library of Mathematical Functions*, <http://dlmf.nist.gov/>, Release 1.1.4 of 2022-01-15. [Online]. Available: <http://dlmf.nist.gov/>.
- [48] S. Olver and Y. Xu, “Orthogonal structure on a wedge and on the boundary of a square,” *Foundations of Computational Mathematics*, vol. 19, pp. 561–589, 2019. DOI: [10.1007/s10208-018-9393-0](https://doi.org/10.1007/s10208-018-9393-0).
- [49] I. P. A. Papadopoulos, *ioannisPApapadopoulos/FractionalFrames.jl*, version 0.0.3, Feb. 2024. DOI: [10.5281/zenodo.10710911](https://doi.org/10.5281/zenodo.10710911).
- [50] I. P. A. Papadopoulos and S. Olver, *A sparse spectral method for fractional differential equations in one-spacial dimension*. arXiv preprint: [2210.08247](https://arxiv.org/abs/2210.08247) [math.NA].
- [51] S. Patnaik, J. P. Hollkamp, and F. Semperlotti, “Applications of variable-order fractional operators: A review,” *Proceedings of the Royal Society A*, vol. 476, no. 2234, p. 20 190 498, 2020. DOI: [10.1098/rspa.2019.0498](https://doi.org/10.1098/rspa.2019.0498).
- [52] C. Sheng, J. Shen, T. Tang, L.-L. Wang, and H. Yuan, “Fast Fourier-like mapped Chebyshev spectral-Galerkin methods for PDEs with integral fractional Laplacian in unbounded domains,” *SIAM Journal on Numerical Analysis*, vol. 58, no. 5, pp. 2435–2464, 2020. DOI: [10.1137/19M128377X](https://doi.org/10.1137/19M128377X).
- [53] G. W. Stewart, *Afternotes Goes to Graduates School*. SIAM, 1998, ISBN: 978-0-89871-404-3. DOI: [10.1137/1.9781611971422](https://doi.org/10.1137/1.9781611971422).
- [54] T. Tang, L.-L. Wang, H. Yuan, and T. Zhou, “Rational spectral methods for PDEs involving fractional Laplacian in unbounded domains,” *SIAM Journal on Scientific Computing*, vol. 42, no. 2, A585–A611, 2020. DOI: [10.1137/19M1244299](https://doi.org/10.1137/19M1244299).



- [55] T. Tang, H. Yuan, and T. Zhou, “Hermite spectral collocation methods for fractional PDEs in unbounded domains,” *Communications in Computational Physics*, vol. 24, no. 4, pp. 1143–1168, 2018. DOI: [10.4208/cicp.2018.hh80.12](https://doi.org/10.4208/cicp.2018.hh80.12).
- [56] F. del Teso, “Finite difference method for a fractional porous medium equation,” *Calcolo*, vol. 51, no. 4, pp. 615–638, 2014.
- [57] B. E. Treeby and B. T. Cox, “Modeling power law absorption and dispersion for acoustic propagation using the fractional Laplacian,” *The Journal of the Acoustical Society of America*, vol. 127, no. 5, pp. 2741–2748, 2010. DOI: [10.1121/1.3377056](https://doi.org/10.1121/1.3377056).
- [58] J. L. Vázquez, “Asymptotic behaviour for the fractional heat equation in the Euclidean space,” *Complex Variables and Elliptic Equations*, vol. 63, no. 7-8, pp. 1216–1231, 2018. DOI: [10.1080/17476933.2017.1393807](https://doi.org/10.1080/17476933.2017.1393807).
- [59] K Yosida, *Functional Analysis*, 6th edition. Springer, 1980, ISBN: 978-3-540-58654-8. DOI: [10.1007/978-3-642-61859-8](https://doi.org/10.1007/978-3-642-61859-8).