

Non-Gaussian correlations in the steady-state of driven-dissipative clouds of two-level atoms

Giovanni Ferioli, Sara Pancaldi, Antoine Glicenstein, David Clément, Antoine Browaeys,^{*} and Igor Ferrier-Barbut[†]
*Université Paris-Saclay, Institut d'Optique Graduate School,
 CNRS, Laboratoire Charles Fabry, 91127, Palaiseau, France*

We report experimental measurements of the second-order coherence function $g^{(2)}(\tau)$ of the light emitted by a laser-driven dense ensemble of ^{87}Rb atoms. We observe a clear departure from the Siegert relation valid for Gaussian chaotic light. Measuring intensity and first-order coherence, we conclude that the violation is not due to the emergence of a coherent field. This indicates that the light obeys non-Gaussian statistics, stemming from non-Gaussian correlations in the atomic medium. More specifically, the steady-state of this driven-dissipative many-body system sustains high-order correlations in the absence of first-order coherence. These findings call for new theoretical and experimental explorations to uncover their origin and they open new perspectives for the realization of non-Gaussian states of light.

The properties of the light emitted by an ensemble of atoms become collective when they are placed inside a volume with a size smaller than their transition wavelength, or when they share a common electromagnetic mode, e.g. inside an optical cavity or along a waveguide. For example, superradiance is a consequence of a collective coupling to a common mode [1, 2]. This collective coupling of the emitters may induce a modification of the statistics of the emitted light and quantum correlations of the emitters' internal degrees of freedom. Relating the statistical properties of the light to the correlations inside the atomic ensemble remains, in the general case, challenging [3, 4]. In this context an outstanding goal is to stabilize non-trivial correlations in the steady-state of a driven-dissipative many-body system [5–8].

In a recent experiment [9], we observed a modification of the photon emission rate in a mode propagating along the long axis of a cigar-shaped cloud of two-level atoms strongly driven by a resonant laser. This enhancement of intensity observed *during the early dynamics* was due to the spontaneous establishment of interatomic correlations, not imparted by the driving laser but rather resulting from superradiance. The question then arises as to whether the *steady state* also features atomic correlations. Information on them may be provided by measuring intensity correlations [10–13]. In particular, a test for the statistical independence of a large number of emitters is the so-called Siegert relation [14–16]. It relates the second order coherence (intensity correlations) of $N \gg 1$ emitters $g_N^{(2)}(\tau) = \langle \hat{E}^-(t)\hat{E}^-(t+\tau)\hat{E}^+(t+\tau)\hat{E}^+(t) \rangle / \langle \hat{I}(t) \rangle^2$ to the first order coherence (field correlations) $g_N^{(1)}(\tau) = \langle \hat{E}^-(t)\hat{E}^+(t+\tau) \rangle / \langle \hat{I}(t) \rangle$, where $\hat{I}(t) = \hat{E}^-(t)\hat{E}^+(t)$ is the intensity and \hat{E}^- is the field radiated by the ensemble. This relation reads: $g_N^{(2)}(\tau) = 1 + |g_N^{(1)}(\tau)|^2$. Its validity has been tested on different platforms with statistically independent atoms generating chaotic light [15], including few atoms in cavity [17], ions [18] or dilute atomic clouds [16, 19–21]. Its violation is a marker of a correlated medium [22–24]. It could be used in experiments to test the predictions of recent theoretical works [25–28] that predict the emergence of high-order correlations in driven atomic ensembles.

Here, we measure the second order coherence $g_N^{(2)}(\tau)$ of the light emitted by cigar-shaped atomic clouds laser driven *per-*

pendicularly to their long axis. We observe a violation of the Siegert relation in steady state, revealing the presence of correlations between atoms. In particular, the violation always appears as a reduction of $g_N^{(2)}(\tau)$ for photons emitted in the mode in which the system features superradiance. Ab-initio numerical calculations for our regime of thousands of emitters are out of reach. However, the Siegert relation can be discussed without knowledge of the microscopic dynamics: It assumes that the connected correlations (or cumulants as defined in [29]) of order higher than 2 cancel, *i.e.* that the field obeys *Gaussian statistics*, and that the radiated field has *zero mean* ($\langle \hat{E}^- \rangle = 0$) [22]. Its experimental violation indicates a failure of one of these two hypotheses. We provide experimental evidence that, in our system, the average field cancels implying that the field is non-Gaussian, and emerges from a non-Gaussian steady-state of the driven atomic medium. Our observations demonstrate that non-Gaussian correlations can emerge from driven-dissipative dynamics, as was recently identified in theoretical studies of related systems [27, 28].

Our experimental platform, detailed in [30], is sketched in figure 1(a). It relies on 4 high-numerical-aperture aspheric lenses. We load up to $\simeq 5000$ ^{87}Rb atoms in a $3.4\mu\text{m}$ -waist optical dipole trap making use of gray molasses. The atomic cloud has a typical temperature of $200\mu\text{K}$, with a (calculated) radial size $\ell_{\text{rad}} \simeq 0.6\lambda$ ($1/e^2$ radius), and a measured axial size $\ell_{\text{ax}} \simeq 20 - 25\lambda$, where $\lambda = 2\pi/k = 780.2\text{nm}$ is the wavelength of the D2 transition. To isolate two internal states and produce a cloud of two-levels atoms, we perform the experiment in the presence of a magnetic field $B = 96\text{G}$ oriented perpendicularly to the atomic cloud. We prepare the atoms in the state $|g\rangle = |5S_{1/2}, F=2, m_F=2\rangle$ by hyperfine and Zeeman optical pumping. The cloud is then excited to $|e\rangle = |5P_{3/2}, F=3, m_F=3\rangle$ (D_2 transition, $\Gamma_0/2\pi = 6\text{MHz}$ and $I_{\text{sat}} \simeq 1.67\text{mW/cm}^2$) using a σ^+ -polarized laser on resonance. Contrarily to our recent work [24] but identically to [9], this beam propagates along B , *i.e.* perpendicularly to the cloud axis. The excitation beam is much larger than the cloud size, so that all atoms experience the same Rabi frequency Ω . In all experiments presented here $\Omega > 2\Gamma$ and we observe that the excited state population has reached saturation. We collect the light emitted by the cloud in two different directions:




FIG. 1. **Experimental setup and $g_N^{(2)}(\tau)$ measurements** (a) Scheme of the experiment. A cigar-shaped cloud of ^{87}Rb atoms is excited by a resonant laser beam propagating perpendicularly to its long axis. The light emitted by the cloud is collected either along its axis (\hat{u}_z , shown) or in a perpendicular direction (\hat{u}_\perp , not shown) by two avalanche photodiodes (APD1,2) arranged in a Hanbury-Brown and Twiss configuration. A time tagger (T.T.) records the photon arrival times. Inset: example of intensity $\langle \hat{I}(t) \rangle$ collected along \hat{u}_z . Red: steady-state where $g_N^{(2)}(\tau)$ is calculated. (b) $g_N^{(2)}(\tau)$ along \hat{u}_z for a dilute cloud, compared to the Siegert relation (dashed line). (c) $g_N^{(2)}(\tau)$ in the dense regime measured along \hat{u}_z (red), violating the Siegert relation. Light blue: collection along \hat{u}_\perp .

the first one is aligned along the main axis of the cloud (\hat{u}_z), the second one is perpendicular to it (\hat{u}_\perp , not aligned with the driving laser) [9, 31].

To measure $g_N^{(2)}(\tau)$, we implement a Hanbury-Brown and Twiss setup: the collected fluorescence is coupled into an optical fiber and then split by a fiber-based 50/50 beam-splitter, whose two outputs are connected to fiber-coupled avalanche photodiodes (APDs) operating in single-photon counting mode. We record the photon arrival times in each arm of the beam-splitter using a time-to-digital converter. From these, we compute $g_N^{(2)}(t_1, t_2) = n_c(t_1, t_2) / n_1(t_1)n_2(t_2)$ where $n_i(t_i)$ is the photon number detected in arm i at time t_i and $n_c(t_1, t_2)$ the number of coincidences on both arms at times t_1 and t_2 (see details in [32]). The time bin is 1 ns.

To calibrate our experiment, we first study a case where the Siegert relation is expected to hold, that is a cloud of independent atoms. To reach this regime, we release the cloud from the trap and let it expand in free flight. The radial size increases by a factor > 10 , up to $\sim 5 \mu\text{m}$. The atoms are excited by a $10 \mu\text{s}$ -long pulse of resonant light. This long duration is necessary to detect a large number of correlations. In these conditions, the intensity correlation $g_N^{(2)}(\tau)$ measured along the main axis of the cloud (see figure 1b) is in excellent agreement with the prediction of the Siegert relation without any free parameter: it features oscillations at the Rabi frequency of the laser, as expected [15]. This confirms that the correlations between atoms are negligible. This good agreement also serves as a quantitative calibration of our detection scheme. Indeed, several effects could reduce the value of $g_N^{(2)}(0)$ below 2: First, a too low time-resolution would lead to

a reduction of $g_N^{(2)}(0)$ [33]. The resolution time of the detectors is 350 ps, much shorter than atomic dynamics timescales (≥ 5 ns). Second, collecting multiple spatial modes over a solid angle larger than a coherence area would also reduce $g_N^{(2)}(0)$ [19, 34]. Here we collect the fluorescence light with an aspheric lens and project it on a single-mode optical fiber. We thus do not expect these two systematics to play a role. The fact that we measure a nearly perfect contrast in this dilute case ($g_N^{(2)}(0) = 1.98 \pm 0.03$) demonstrates that this is the case and that no systematic effects could reduce the value of $g_N^{(2)}(\tau)$.

To study the dense regime, we prepare the clouds as presented above, then switch off the trap and immediately shine a 400 ns-long pulse of resonant laser light. We then recapture the atoms in the optical tweezer and repeat this sequence up to 20 times to accumulate statistics, checking that the atom number is reduced by less than 10%. During the laser pulse, the thermal expansion is negligible, and we thus assume that the atomic distribution remains identical to the trapped one. To obtain the steady-state correlation function $g_N^{(2)}(\tau)$, we restrict the times t and $t' = t + \tau$ to a time window of 250 ns when the atomic system has reached steady state, as highlighted in the inset of figure 1(a) (more details in [32]).

Strikingly, as shown in figure 1(c), we observe in this dense regime a violation of the Siegert relation (dotted line) along the cloud axis. This is the direction along which the emission is collective and leads to superradiance during the early dynamics [9]. In this axial direction, we measure a reduction of $g_N^{(2)}(\tau)$ with respect to the dilute case for all delays τ . We even obtain $g_N^{(2)}(\tau) < 1$ at around half a period of the os-

cillation (colored area in figure 1c). Furthermore, we observe that the photon statistics depends on the direction of detection: The statistics of photons emitted perpendicularly to the cloud axis is well described by the Siegert relation. In this direction, emission is not collective and we do not observe superradiance because interferences are too weak [9]. For the same reason we do not expect the Siegert relation to be violated, as discussed in more details in [32]. We discuss this in more details in [32], and note that it was predicted to occur also in the configuration of [24] where the cloud is driven along its axis [27]. To plot this relation, we assume $g_N^{(1)}(\tau) = g_1^{(1)}(\tau)$. We have experimentally verified this assumption by measuring the first-order correlation function $g_N^{(1)}(\tau)$ in the dense regime, following the method used in [35, 36] (see details in [32]). As shown in figure 2, we find it to be in agreement with the single atom expectation.

Let us discuss the implication of the observed violation of the Siegert relation along the cloud axis and how it can reveal non-Gaussian statistics of the emitted light. If one first assumes Gaussian light statistics, all connected correlations of more than two operators cancel and the correlation of four operators then reads [29]: $\langle \hat{A}\hat{B}\hat{C}\hat{D} \rangle = \langle \hat{A}\hat{B} \rangle \langle \hat{C}\hat{D} \rangle + \langle \hat{A}\hat{C} \rangle \langle \hat{B}\hat{D} \rangle + \langle \hat{A}\hat{D} \rangle \langle \hat{B}\hat{C} \rangle - 2\langle \hat{A} \rangle \langle \hat{B} \rangle \langle \hat{C} \rangle \langle \hat{D} \rangle$. Applying this to $\hat{A} = \hat{E}^-(t) = \hat{D}^\dagger$, $\hat{B} = \hat{E}^-(t+\tau) = \hat{C}^\dagger$ [37] yields

$$g_N^{(2)}(\tau) = 1 + |g_N^{(1)}(\tau)|^2 - \frac{2|\langle \hat{E}^- \rangle|^4}{\langle \hat{I} \rangle^2} + \frac{|\langle \hat{E}^-(t)\hat{E}^-(t+\tau) \rangle|^2}{\langle \hat{I} \rangle^2}, \quad (1)$$

with t taken in steady state and $\langle \hat{E}^- \rangle$ the average electric field radiated by the cloud in steady state. The last term oscillates fast and is in general neglected [22, 23, 38]. Thus the observed violation of the Siegert relation, $g_N^{(2)}(\tau) \leq 1 + |g_N^{(1)}(\tau)|^2$ for all delays τ can only be explained in two ways. Either the field does not obey Gaussian statistics so that Eq. (1) does not apply, or the average field $\langle \hat{E}^- \rangle$ is non-zero in steady state.

The existence of such an average field in steady state would be non-trivial as it is not externally imposed by the driving laser. This laser imprints a phase factor $e^{-ik_{\text{las}} \cdot r_n}$ on atom n . The field emitted by the cloud in the direction \hat{u}_z is $\hat{E}^- = \sum_{n=1}^N \hat{\sigma}_n^+ e^{ik_{\text{las}} \cdot r_n}$. Since $k_{\text{las}} \perp \hat{u}_z$, the laser does not directly excite atomic dipoles whose radiations constructively interfere along \hat{u}_z . A non-zero average field would then result from a many-body dynamics creating a coherence along \hat{u}_z . Coherence has been observed in dilute clouds, during the late decay following the extinction of the laser excitation [39]. In order to assess if a coherent field is emitted in steady-state for our strongly driven clouds, we perform two experimental tests.

We obtain the first compelling evidence that $\langle \hat{E}^- \rangle \approx 0$ by measuring the intensity emitted by the cloud along \hat{u}_z . Figure 2(a) shows the steady-state intensity $\langle \hat{I} \rangle$ measured along \hat{u}_z as a function of the atom number N . Since the field is the sum of the radiation of the individual dipoles, a non-zero average field $\langle \hat{E}^- \rangle$ should be proportional to N . In the presence of a non-zero average field, the intensity should read $\langle \hat{I} \rangle = \sum_{i,j} \langle \hat{\sigma}_i^+ \hat{\sigma}_j^- \rangle e^{ik_{\text{las}} \cdot (r_i - r_j)} \simeq \sum_{i=1}^N \langle \hat{\sigma}_i^+ \hat{\sigma}_i^- \rangle + |\langle \hat{E}^- \rangle|^2$:




FIG. 2. **Evidence for a negligible average field.** (a) Intensity $\langle \hat{I} \rangle$ measured along \hat{u}_z in steady state versus atom number N . Error bars are standard error on the mean (s.e.m.). Dashed: linear and a quadratic scalings. (b) $|g_N^{(1)}(\tau)|$ with $\Omega \simeq 4.5\Gamma$ (red) and expectation for a single atom (black).

The average field leads to the appearance of a quadratic term ($\propto N^2$) on top of the linear scaling due to incoherent scattering. We however observe in figure 2(a) a clear linear scaling, indicating that the field radiated by the cloud has a negligible average value. From the residuals of a linear fit to the data, we obtain $|\langle \hat{E}^- \rangle|^2 / \langle \hat{I} \rangle \leq 0.17$ so that the third term in (1) could cause a reduction of $g_N^{(2)}(\tau)$ of at most 0.06. This is much smaller than the reduction of $\simeq 0.3$ that we observe. The second piece of evidence comes from the observation of the decay of the first order coherence to zero at long times, as shown in figure 2(b): $g_N^{(1)}(\tau) \rightarrow 0$ for $\tau \gg 1/\Gamma$. In the long-time limit, we expect $\langle \hat{E}^-(t)\hat{E}^+(t+\tau) \rangle = \langle \hat{E}^-(t) \rangle \langle \hat{E}^+(t+\tau) \rangle$ and hence $g_N^{(1)}(\tau) \rightarrow |\langle \hat{E}^- \rangle|^2 / \langle \hat{I} \rangle$. The data in figure 2(b) again sets a bound of about $|\langle \hat{E}^- \rangle|^2 / \langle \hat{I} \rangle \leq 0.2$. As a consequence our measurement of first-order coherence further demonstrates the fact that no average coherent field emerges. We thus have strong experimental evidence that our observation of $g_N^{(2)}(\tau) < 1 + |g_N^{(1)}(\tau)|^2$ reveals a non-Gaussian statistics of the light emitted by the cloud along its main axis.

To quantify the departure from Gaussian statistics we measure high-order connected correlations [40–43], which would cancel in the Gaussian case. The measured $g_N^{(2)}(\tau)$ can be related to the normalized two-times connected correlation $C(\tau) = \langle \hat{E}^-(t)\hat{E}^-(t+\tau)\hat{E}^+(t+\tau)\hat{E}^-(t) \rangle_c / \langle \hat{I} \rangle^2$, using the equation (derived in [32]):

$$g_N^{(2)}(\tau) = g_{\text{Gauss}}^{(2)}(\tau) + C(\tau), \quad (2)$$

where $g_{\text{Gauss}}^{(2)}(\tau) = 1 + |g_N^{(1)}(\tau)|^2 + |\langle \hat{E}^-(t)\hat{E}^-(t+\tau) \rangle|^2 / \langle \hat{I} \rangle^2$. This expression assumes $\langle \hat{E}^- \rangle = 0$ as justified above. These connected correlations quantify the lack or excess of photon pairs separated by τ with respect to the case of a Gaussian light. From Eq. (2), one can indeed show that $C(\tau) = g_{\text{Gauss}}^{(2)}(\tau)(f(\tau) - 1)$ where $f(\tau) = n_c(\tau)/n_{\text{cGauss}}(\tau)$ is the fraction of detected photon pairs separated by τ with respect to what would have been detected for Gaussian light (with the same average intensity $\langle \hat{I} \rangle$). Since the third term of $g_{\text{Gauss}}^{(2)}(\tau)$ is always positive, we get the following lower

bound: $|\mathcal{C}(\tau)| \geq 1 + |g_N^{(1)}(\tau)|^2 - g_N^{(2)}(\tau)$. This quantity can be directly extracted from the data. We show in figure 3(a) the intensity correlation $g_N^{(2)}(\tau)$ as a function of the atom number N , for $I_{\text{las}} \simeq 50I_{\text{sat}}$ ($\Omega/\Gamma_0 \simeq 5$). Figure 3(b,c) reports the values of $g_N^{(2)}(0)$, and the corresponding connected correlation $\mathcal{C}(0)$, as a function of N . We do find that at low N the data converge towards the prediction of the Siegert relation. For increasing N the disagreement grows. We also find that $g_N^{(2)}(0)$ (and $\mathcal{C}(0)$) does not vary when changing the Rabi frequency between $\Omega = 2\Gamma$ ($I/I_{\text{sat}} = 4$) and $\Omega = 10\Gamma$ ($I/I_{\text{sat}} = 200$), as shown in [32]: despite a very strong drive the data does not converge towards single atom behaviour in this range of driving strength. The inset of figure 3(c) shows how the connected correlations decay in time. We observe a maximum of correlation and a non-monotonic decay towards zero. The correlations observed in figure 3 with $\mathcal{C}(\tau) \neq 0$ indicate that second-order coherence emerges. The fact that second-order coherence is built in the absence of first-order coherence is a signature of non-Gaussian statistics. A theoretical understanding of the measured $\mathcal{C}(\tau)$ is beyond the scope of the present work. It requires a description of the atomic correlations emerging in the cloud.

In this perspective, we relate the statistics of the light field to the one of the atomic state. In ref. [9], we already observed the appearance of beyond-mean-field correlations between the atomic dipoles, *i.e.* $\langle \hat{\sigma}_i^+ \hat{\sigma}_j^- \rangle_c \neq 0$, during the early dynamics following the application of the excitation laser. To do so, we measured the intensity emitted along \hat{u}_z , whose expression in terms of atomic dipoles is $\langle \hat{I} \rangle = \sum_{ij} \langle \hat{\sigma}_i^+ \hat{\sigma}_j^- \rangle e^{-ik\hat{u}_z \cdot (\mathbf{r}_i - \mathbf{r}_j)}$. These correlations resulted from the superradiant emission along the axial direction of the cloud. Contrarily, the measurements of second order coherence presented here probe higher order correlations *in steady state*. In terms of atomic dipoles, the connected correlations of the field measured above read

$$\langle \hat{E}^- \hat{E}^- \hat{E}^+ \hat{E}^+ \rangle_c = \sum_{ijkl} \langle \hat{\sigma}_i^+ \hat{\sigma}_j^+ \hat{\sigma}_k^- \hat{\sigma}_l^- \rangle_c e^{-ik\hat{u}_z \cdot (\mathbf{r}_i - \mathbf{r}_j + \mathbf{r}_k - \mathbf{r}_l)}. \quad (3)$$

Here, an operator $\hat{\mathcal{O}}$ is taken at time t in steady state and $\hat{\mathcal{O}}'$ at time $t' = t + \tau$. Hence, the observation of non-zero connected correlations in the field implies that $\langle \hat{\sigma}_i^+ \hat{\sigma}_j^+ \hat{\sigma}_k^- \hat{\sigma}_l^- \rangle_c \neq 0$, *i.e.* the atomic medium features high-order correlations that obey non-Gaussian statistics. These high-order correlations are not externally imposed and emerge in steady-state as a result of the competition between driving and collective dissipation. This shows that in our free-space system, despite the absence of spatial order, collective dissipation can stabilize non-trivial correlations.

In conclusion, we have investigated the photon statistics of the light emitted in steady state by a dense superradiant cloud of atoms under strong driving, observing a violation of the Siegert relation. Our data support the fact that this violation is not due to the appearance of a coherent field. They rather indicate that a non-Gaussian field emerges in the steady state of this driven-dissipative system, which originates from non-Gaussian correlations between atoms in the cloud.




FIG. 3. **Atom number N dependence of correlations** (a) $g_N^{(2)}(\tau)$ versus τ and N . (b) $g_N^{(2)}(0)$ obtained by averaging $g_N^{(2)}(\tau)$ in the interval $-2 \text{ ns} \leq \tau \leq 2 \text{ ns}$. (c) Connected correlation $\mathcal{C}(0)$ as defined in the main text. Inset: example of $\mathcal{C}(\tau)$. The error bars are s.e.m.

The appearance of stable non-Gaussian correlations in steady state under strong driving is an unexpected observation. Our findings thus call for theoretical investigations to identify the mechanisms at play in this dissipative quantum many-body system, and to elucidate their relationship with superradiance. More generally, this should motivate investigations to determine whether the correlations we observed could be used as a resource to prepare non-trivial states of the field [44–46]. Experimentally, we plan to measure the quadratures of the radiated field to extract its Wigner function and determine if the non-Gaussian character we observed is accompanied by Wigner negativity [47]. Another outlook would be to extend our investigation beyond the steady state, studying for instance the photon statistics during a superradiant burst [48–50].

We acknowledge discussions with Darrick Chang, Francis Robicheaux, Tommaso Roscilde, Hans Peter Büchler, Bruno Laburthe-Tolra and Martin Robert de saint Vincent. This project has received funding from the European Research Council (Advanced grant No. 101018511, ATARAXIA), by

the Agence National de la Recherche (project DEAR and ANR-22-PETQ-0004 France 2030, project QuBitAF) and by the Region Ile-de-France in the framework of DIM SIRTEQ (projects DSHAPE and FSTOL). S.P. is funded by the Paris Saclay Quantum Center.

* antoine.browaeys@institutoptique.fr

† igor.ferrier-barbut@institutoptique.fr

- [1] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
- [2] Superradiance: An essay on the theory of collective spontaneous emission, *Physics Reports* **93**, 301 (1982).
- [3] O. Somech and E. Shahmoon, Quantum entangled states of a classically radiating macroscopic spin (2022), [arXiv:2204.05455 \[quant-ph\]](https://arxiv.org/abs/2204.05455).
- [4] O. Tziperman, G. Baranes, A. Gorlach, R. Ruimy, M. Faran, N. Gutman, A. Pizzi, and I. Kaminer, Spontaneous emission from correlated emitters (2023), [arXiv:2306.11348 \[quant-ph\]](https://arxiv.org/abs/2306.11348).
- [5] T. E. Lee, S. Gopalakrishnan, and M. D. Lukin, Unconventional magnetism via optical pumping of interacting spin systems, *Phys. Rev. Lett.* **110**, 257204 (2013).
- [6] B. Olmos, D. Yu, and I. Lesanovsky, Steady-state properties of a driven atomic ensemble with nonlocal dissipation, *Phys. Rev. A* **89**, 023616 (2014).
- [7] C. D. Parmee and N. R. Cooper, Phases of driven two-level systems with nonlocal dissipation, *Phys. Rev. A* **97**, 053616 (2018).
- [8] C. D. Parmee and J. Ruostekoski, Signatures of optical phase transitions in superradiant and subradiant atomic arrays, *Communications Physics* **3**, 205 (2020).
- [9] G. Ferioli, A. Glicenstein, F. Robicheaux, R. T. Sutherland, A. Browaeys, and I. Ferrier-Barbut, Laser-driven superradiant ensembles of two-level atoms near dicke regime, *Phys. Rev. Lett.* **127**, 243602 (2021).
- [10] H. J. Kimble, M. Dagenais, and L. Mandel, Photon antibunching in resonance fluorescence, *Phys. Rev. Lett.* **39**, 691 (1977).
- [11] F. Diedrich and H. Walther, Nonclassical radiation of a single stored ion, *Phys. Rev. Lett.* **58**, 203 (1987).
- [12] S. Mahmoodian, M. Čepulkovskis, S. Das, P. Lodahl, K. Hammerer, and A. S. Sørensen, Strongly correlated photon transport in waveguide quantum electrodynamics with weakly coupled emitters, *Phys. Rev. Lett.* **121**, 143601 (2018).
- [13] A. S. Prasad, J. Hinney, S. Mahmoodian, K. Hammerer, S. Rind, P. Schneeweiss, A. S. Sørensen, J. Volz, and A. Rauschenbeutel, Correlating photons using the collective nonlinear response of atoms weakly coupled to an optical mode, *Nature Photonics* **14**, 719 (2020).
- [14] A. Siegert, *On the fluctuations in signals returned by many independently moving scatterers* (Radiation Laboratory, Massachusetts Institute of Technology, 1943).
- [15] R. Loudon, *The quantum theory of light* (OUP Oxford, 2000).
- [16] D. Ferreira, R. Bachelard, W. Guerin, R. Kaiser, and M. Fouché, Connecting field and intensity correlations: The siegert relation and how to test it, *American Journal of Physics* **88**, 831 (2020).
- [17] M. Henrich, A. Kuhn, and G. Rempe, Transition from antibunching to bunching in cavity qed, *Phys. Rev. Lett.* **94**, 053604 (2005).
- [18] A. Kovalenko, D. Babjak, A. Lešundák, L. Podhora, L. Lachman, P. Obšil, T. Pham, O. Číp, R. Filip, and L. Slodička, Emergence of super-poissonian light from indistinguishable single-photon emitters, *Optica* **10**, 456 (2023).
- [19] S. Bali, D. Hoffmann, J. Simán, and T. Walker, Measurements of intensity correlations of scattered light from laser-cooled atoms, *Phys. Rev. A* **53**, 3469 (1996).
- [20] J. A. Grover, P. Solano, L. A. Orozco, and S. L. Rolston, Photon-correlation measurements of atomic-cloud temperature using an optical nanofiber, *Phys. Rev. A* **92**, 013850 (2015).
- [21] P. Lassègues, M. A. F. Biscassi, M. Morisse, A. Cidrim, P. G. S. Dias, H. Eneriz, R. C. Teixeira, R. Kaiser, R. Bachelard, and M. Hugbart, Transition from classical to quantum loss of light coherence, *Phys. Rev. A* **108**, 042214 (2023).
- [22] P.-A. Lemieux and D. J. Durian, Investigating non-gaussian scattering processes by using nth-order intensity correlation functions, *J. Opt. Soc. Am. A* **16**, 1651 (1999).
- [23] H. J. Carmichael, P. Drummond, P. Meystre, and D. F. Walls, Intensity correlations in resonance fluorescence with atomic number fluctuations, *Journal of Physics A: Mathematical and General* **11**, L121 (1978).
- [24] G. Ferioli, A. Glicenstein, I. Ferrier-Barbut, and A. Browaeys, A non-equilibrium superradiant phase transition in free space, *Nature Physics* **19**, 1345 (2023).
- [25] F. Robicheaux and D. A. Suresh, Beyond lowest order mean-field theory for light interacting with atom arrays, *Phys. Rev. A* **104**, 023702 (2021).
- [26] O. Rubies-Bigorda, S. Ostermann, and S. F. Yelin, Characterizing superradiant dynamics in atomic arrays via a cumulant expansion approach, *Phys. Rev. Res.* **5**, 013091 (2023).
- [27] C. D. Mink and M. Fleischhauer, Collective radiative interactions in the discrete truncated Wigner approximation, *SciPost Phys.* **15**, 233 (2023).
- [28] K. C. Stitely, F. Finger, R. Rosa-Medina, F. Ferri, T. Donner, T. Esslinger, S. Parkins, and B. Krauskopf, Quantum fluctuation dynamics of dispersive superradiant pulses in a hybrid light-matter system, *Phys. Rev. Lett.* **131**, 143604 (2023).
- [29] R. Kubo, Generalized cumulant expansion method, *Journal of the Physical Society of Japan* **17**, 1100 (1962).
- [30] A. Glicenstein, G. Ferioli, L. Brossard, Y. R. P. Sortais, D. Barredo, F. Nogrette, I. Ferrier-Barbut, and A. Browaeys, Preparation of one-dimensional chains and dense cold atomic clouds with a high numerical aperture four-lens system, *Phys. Rev. A* **103**, 043301 (2021).
- [31] A. Glicenstein, G. Ferioli, A. Browaeys, and I. Ferrier-Barbut, From superradiance to subradiance: exploring the many-body dicke ladder, *Opt. Lett.* **47**, 1541 (2022).
- [32] Supplementary information.
- [33] L. MANDEL and E. WOLF, Coherence properties of optical fields, *Rev. Mod. Phys.* **37**, 231 (1965).
- [34] E. Jakeman, C. J. Oliver, and E. R. Pike, The effects of spatial coherence on intensity fluctuation distributions of gaussian light, *Journal of Physics A: General Physics* **3**, L45 (1970).
- [35] H.-G. Hong, W. Seo, M. Lee, W. Choi, J.-H. Lee, and K. An, Spectral line-shape measurement of an extremely weak amplitude-fluctuating light source by photon-counting-based second-order correlation spectroscopy, *Opt. Lett.* **31**, 3182 (2006).
- [36] L. Ortiz-Gutiérrez, R. C. Teixeira, A. Eloy, D. F. da Silva, R. Kaiser, R. Bachelard, and M. Fouché, Mollow triplet in cold atoms, *New Journal of Physics* **21**, 093019 (2019).
- [37] This assumes that for $\tau \neq 0$ the correlators can be separated for operators taken at different times. See a discussion in Refs. [25, 51].
- [38] D. A. Steck, Quantum and atom optics, (2007).
- [39] D. C. Gold, P. Huft, C. Young, A. Safari, T. G. Walker, M. Saffman, and D. D. Yavuz, Spatial coherence of light in col-

- lective spontaneous emission, *PRX Quantum* **3**, 010338 (2022).
- [40] T. Schweigler, V. Kasper, S. Erne, I. Mazets, B. Rauer, F. Cataldini, T. Langen, T. Gasenzer, J. Berges, and J. Schmiedmayer, Experimental characterization of a quantum many-body system via higher-order correlations, *Nature* **545**, 323 (2017).
- [41] M. Rispoli, A. Lukin, R. Schittko, S. Kim, M. E. Tai, J. Léonard, and M. Greiner, Quantum critical behaviour at the many-body localization transition, *Nature* **573**, 385 (2019).
- [42] N. Stiesdal, J. Kumlin, K. Kleinbeck, P. Lunt, C. Braun, A. Paris-Mandoki, C. Tresp, H. P. Büchler, and S. Hofferberth, Observation of three-body correlations for photons coupled to a rydberg superatom, *Phys. Rev. Lett.* **121**, 103601 (2018).
- [43] P. M. Preiss, J. H. Becher, R. Klemt, V. Klinkhamer, A. Bergschneider, N. Defenu, and S. Jochim, High-contrast interference of ultracold fermions, *Phys. Rev. Lett.* **122**, 143602 (2019).
- [44] A. I. Lvovsky, P. Grangier, A. Ourjoumtsev, V. Parigi, M. Sasaki, and R. Tualle-Brouri, Production and applications of non-gaussian quantum states of light (2020), [arXiv:2006.16985 \[quant-ph\]](https://arxiv.org/abs/2006.16985).
- [45] M. Walschaers, Non-gaussian quantum states and where to find them, *PRX Quantum* **2**, 030204 (2021).
- [46] L. Lachman and R. Filip, Quantum non-gaussianity of light and atoms, *Progress in Quantum Electronics* **83**, 100395 (2022).
- [47] B. Olmos, G. Buonaiuto, P. Schneeweiss, and I. Lesanovsky, Interaction signatures and non-gaussian photon states from a strongly driven atomic ensemble coupled to a nanophotonic waveguide, *Phys. Rev. A* **102**, 043711 (2020).
- [48] S. Cardenas-Lopez, S. J. Masson, Z. Zager, and A. Asenjo-Garcia, Many-body superradiance and dynamical mirror symmetry breaking in waveguide qed, *Phys. Rev. Lett.* **131**, 033605 (2023).
- [49] S. J. Masson, I. Ferrier-Barbut, L. A. Orozco, A. Browaeys, and A. Asenjo-Garcia, Many-body signatures of collective decay in atomic chains, *Phys. Rev. Lett.* **125**, 263601 (2020).
- [50] D. Bhatti, J. von Zanthier, and G. S. Agarwal, Superbunching and nonclassicality as new hallmarks of superradiance, *Scientific Reports* **5**, 17335 (2015).
- [51] W. Verstraelen, D. Huybrechts, T. Roscilde, and M. Wouters, Quantum and classical correlations in open quantum spin lattices via truncated-cumulant trajectories, *PRX Quantum* **4**, 030304 (2023).
- [52] G. S. Agarwal, L. M. Narducci, D. H. Feng, and R. Gilmore, Intensity correlations of a cooperative system, *Phys. Rev. Lett.* **42**, 1260 (1979).
- [53] L. Pucci, A. Roy, T. S. do Espirito Santo, R. Kaiser, M. Kastner, and R. Bachelard, Quantum effects in the cooperative scattering of light by atomic clouds, *Phys. Rev. A* **95**, 053625 (2017).
- [54] R. Jones, R. Saint, and B. Olmos, Far-field resonance fluorescence from a dipole-interacting laser-driven cold atomic gas, *Journal of Physics B: Atomic, Molecular and Optical Physics* **50**, 014004 (2016).
- [55] L. Allen and J. H. Eberly, *Optical resonance and two-level atoms* (Dover, 1987).
- [56] R. T. Sutherland and F. Robicheaux, Superradiance in inverted multilevel atomic clouds, *Phys. Rev. A* **95**, 033839 (2017).
- [57] If one assumes that correlations are along the driving laser direction $\hat{u}_c \parallel \mathbf{k}_{\text{las}}$, one obtains that $\mu(\hat{u}_c) < 10^{-4}$, and thus correlations in this direction are too weak to induce collective effects, the double sum always cancels when averaging over the collection solid angle.

Supplemental Material

Data analysis and reconstruction of the correlation function $g_N^{(2)}(\tau)$

In this section, we give more details about the measurement of the second-order coherence $g_N^{(2)}(\tau)$. This correlation function is related to measurable quantities by the following expression:

$$g_N^{(2)}(t_1, t_2) = \frac{n_c(t_1, t_2)}{n_1(t_1)n_2(t_2)} \quad (\text{S1})$$

where $n_i(t_i)$ is the photon arrival rate on detector i at time t_i and $n_c(t_1, t_2)$ is the rate of two-photon coincidence at times $t = t_1$ and $t = t_2$ in detectors 1 and 2. As explained in the main text, for every excitation pulse, we record all the photons arrival times t_1 (t_2) on detector 1 (detector 2) with respect to a common trigger. We thus measure the total number of photons $N_i(t_i)$ detected by detector i at time t_i . In the same way, we measure the total number of coincidences $N_c(t_1, t_2)$. The rates n_i are related to N_i :

$$\begin{aligned} N_1(t_1) &= \varepsilon_1 N_S n_1(t_1) \\ N_2(t_2) &= \varepsilon_2 N_S n_2(t_2) \\ N_c(t_1, t_2) &= \varepsilon_1 \varepsilon_2 N_S n_c(t_1, t_2) \end{aligned} \quad (\text{S2})$$

where ε_i is the detection efficiency of detector i and N_S is the number of times the experiment is repeated. The two-times correlation function is then obtained as:

$$g_N^{(2)}(t_1, t_2) = N_S \frac{N_c(t_1, t_2)}{N_1(t_1)N_2(t_2)}. \quad (\text{S3})$$




FIG. S1. **Example of the two-time correlation function** $g_N^{(2)}(t_1, t_2)$ detected in the experiment, together with the two corresponding time-dependent photon counts detected by each APD. The black rectangle represents the region where the steady state $g_N^{(2)}(\tau)$ is extracted with $|\tau| \leq 50$ ns.

An example of matrix $g_N^{(2)}(t_1, t_2)$ is shown in figure S1, together with the respective time-dependent photon counts recorded by the two APDs. The measurements reported in the main text focus on the steady state, where $g_N^{(2)}(t_1, t_2) = g_N^{(2)}(|t_1 - t_2|) = g_N^{(2)}(\tau)$. This function is evaluated selecting the data where the atomic system has reached the steady state: we thus average over the last $\simeq 250$ ns of the excitation pulse and select $|\tau| \leq 50$ ns, a range where the number of correlations is sufficiently large. An illustrative example is reported in Fig S1.

Heterodyne measurement of the fluorescence spectrum

To further characterize the light field emitted by the cloud in steady-state, we measure the field correlation function $g_N^{(1)}(\tau)$, and show that it is compatible with $g_1^{(1)}(\tau)$ also in the dense regime. To do so, we use a heterodyne detection scheme [35, 36]. In short, a local oscillator (LO) is derived from the laser light used to excite the atoms. It is then shifted by $\omega_{LO}/(2\pi) = 110\text{MHz}$, and coupled into a fiber. We measure the intensity correlation $g_{N,HD}^{(2)}(\tau)$ obtained by combining the LO field and the one scattered by the atoms E^- into a fiber-based beamsplitter. This quantity is related to the first and second order correlations of the light field by [35]:

$$g_{HD}^{(2)}(\tau) = 1 + \alpha \left(g_N^{(2)}(\tau) - 1 \right) - \beta \cos(\omega_{LO}\tau) g_N^{(1)}(\tau) \quad (\text{S4})$$

Here, we have assumed the probe to be resonant with the atomic transition and thus $\text{Im}[g_N^{(1)}(\tau)] = 0$. The parameters α and β in Eq. S4 depend on the intensity of the emitted field (I_{SC}) and of the local oscillator (I_{LO}):

$$\alpha = \frac{\langle I_{SC} \rangle^2}{(\langle I_{SC} \rangle + \langle I_{LO} \rangle)^2}, \quad \beta = 2 \frac{\langle I_{SC} \rangle \langle I_{LO} \rangle}{(\langle I_{SC} \rangle + \langle I_{LO} \rangle)^2}. \quad (\text{S5})$$

A typical signal of $g_{HD}^{(2)}(\tau)$ is shown in figure S2(a), by demodulation of the frequency component at ω_{LO} we can obtain $g_N^{(1)}(\tau)$ as shown in the main text. Furthermore the spectrum of the light emitted by the cloud is the Fourier transform of $g_N^{(1)}(\tau)$. We thus extracted it by Fourier transforming $g_{HD}^{(2)}(\tau)$ and considering the frequency components centered around ω_{LO} .




FIG. S2. (a) Intensity correlation function measured by the heterodyning technique used to extract $g_N^{(1)}(\tau)$ shown in the main text. Here $N \simeq 3000$ atoms and $\Omega \simeq 4.5\Gamma_0$. (b) Spectrum obtained by Fourier transforming the data in (a) and selecting the component around ω_{LO} . Black line: theoretical prediction from the calculation of the resonant fluorescence spectrum of a single atom for the same Ω . Both the experimental data and the theoretical curve are normalized by setting $S(0) = 1$. The dashed lines indicate Ω .

Figure S2 reports a typical example of spectrum $S(\omega)$. The experimental findings exhibit the Mollow triplet, and are always well described by the expression of the resonant fluorescence spectrum of a single atom [15]. Varying the atom number up to $N = 5000$ and scanning the driving strength up to tens of I_{sat} we have always observed agreement with the single-atom expectations. This shows that in our experiment $g_N^{(1)}(\tau)$ is *not* modified with respect to single atom case, contrarily to $g_N^{(2)}(\tau)$, as predicted for superradiant ensembles [52]. The fact that $g^{(1)}(\tau)$ agrees with the one for a single atoms was expected for ensembles of atomic emitters (but not precisely for our system) [53, 54]. These works showed that the influence of collective behavior are very weak in the field correlation $g^{(1)}(\tau)$. Theoretical expectations are challenging in our regime but one might also expect very weak modifications, below our experimental signal to noise ratio. In any case, as $|g_N^{(1)}(\tau)|^2 > 0$ and $g_N^{(1)}(0) = 1$ by definition, if the Siegert relation applies a modification of $g_N^{(1)}(\tau)$ cannot account for the reduction of $g_N^{(2)}(\tau)$ reported in the main text.

Connected correlations

Here, following the main text, we assume $\langle \hat{E}^- \rangle = 0$. The connected correlation of order 4 is defined by [29]:

$$\begin{aligned} \langle \hat{A}\hat{B}\hat{C}\hat{D} \rangle_c &= \langle \hat{A}\hat{B}\hat{C}\hat{D} \rangle - \langle \hat{A}\hat{B} \rangle \langle \hat{C}\hat{D} \rangle - \langle \hat{A}\hat{C} \rangle \langle \hat{B}\hat{D} \rangle - \langle \hat{A}\hat{D} \rangle \langle \hat{B}\hat{C} \rangle + 2\langle \hat{A} \rangle \langle \hat{B} \rangle \langle \hat{C} \rangle \langle \hat{D} \rangle \\ &- \langle \hat{A}\hat{B}\hat{C} \rangle_c \langle \hat{D} \rangle - \langle \hat{B}\hat{C}\hat{D} \rangle_c \langle \hat{A} \rangle - \langle \hat{C}\hat{D}\hat{A} \rangle_c \langle \hat{B} \rangle - \langle \hat{D}\hat{A}\hat{B} \rangle_c \langle \hat{C} \rangle. \end{aligned} \quad (\text{S6})$$




FIG. S3. **Measurement of the connected correlations.** (a,b) Black lines: examples of $g_N^{(2)}(\tau)$ measured for $N \simeq 500$ and $\Omega \simeq 8\Gamma_0$ (a) and $N \simeq 5000$ and $\Omega \simeq 5\Gamma_0$ (b). Gray lines: Siegert relation for the same parameters. (c,d) $C(\tau)$ for the experimental parameters of (a, b) respectively.

For $\langle \hat{E}^- \rangle = 0$, the contribution of the third order connected correlations cancels, leading to Eq. (2) of the main text involving the normalized connected correlation $C(\tau)$: $C(\tau) = \langle \hat{E}^-(t)\hat{E}^-(t+\tau)\hat{E}^+(t+\tau)\hat{E}^+(t) \rangle_c / \langle \hat{I} \rangle^2$. Here $\langle \hat{E}^-(t)\hat{E}^-(t+\tau)\hat{E}^+(t+\tau)\hat{E}^+(t) \rangle_c$ is the connected correlation of the field.

Figure S3(c,d) show examples of normalized connected correlation $C(\tau)$. They have been evaluated from the intensity correlation $g_N^{(2)}(\tau)$ reported respectively in (a,b), measured for different N and Ω . In the low atom number regime, $C(\tau)$ averages to zero, while it clearly departs from 0 for large N .

Comparing on-axis to perpendicular emission

The measurements presented in the main text indicate that light collected perpendicularly to the cloud axis verifies the Siegert relation. We show here that this can be expected.

Let us first discuss the light intensity $\langle \hat{I} \rangle = \langle \hat{E}^- \hat{E}^+ \rangle$ emitted by the cloud. As in our previous work [9], using $\hat{E}^-(\hat{u}) = \sum_{n=1}^N \hat{\sigma}_n^+ e^{ik\hat{u}\cdot\mathbf{r}_n}$ the intensity in direction \hat{u} is:

$$\langle \hat{I} \rangle = I_1(\hat{u}) \left(\sum_{n=1}^N \langle \hat{n}_n^e \rangle + \sum_{n \neq m}^N \langle \hat{\sigma}_n^+ \hat{\sigma}_m^- \rangle e^{ik\hat{u}\cdot(\mathbf{r}_n - \mathbf{r}_m)} \right), \quad (\text{S7})$$

with $\hat{n}_i^e = \hat{\sigma}_i^+ \hat{\sigma}_i^-$ and $I_1(\hat{u})$ the dipole radiation pattern for a single atom. Collective effects, coming from two-atom correlations accounted for in the second term can be compared to the incoherent sum of intensities (first term). To perform this comparison one needs to assume an expression for the correlations. In a direction \hat{u}_c , the strongest possible correlations are given by phased dipoles along \hat{u}_c : $\langle \hat{\sigma}_n^+ \hat{\sigma}_m^- \rangle = \langle \hat{\sigma}_1^+ \hat{\sigma}_2^- \rangle e^{-ik\hat{u}_c\cdot(\mathbf{r}_n - \mathbf{r}_m)}$ with $\langle \hat{\sigma}_1^+ \hat{\sigma}_2^- \rangle$ identical for all pairs (n, m) , its maximum value being 1/4. In this case one can calculate the ratio of the total power emitted in 4π by the second and first term, see for instance refs. [55, 56].

This ratio $P_{\text{single}}/P_{\text{coop}} = \mu(\hat{u}_c) \times N$ defines the parameter $\mu(\hat{u}_c)$ that can be calculated using the Gaussian sizes of our cloud $\ell_{\text{rad}} \simeq 0.6\lambda$, $\ell_{\text{ax}} \simeq 20 - 25\lambda$. For $\hat{u}_c = \hat{u}_z$ one gets $\mu(\hat{u}_z) \approx 2 \times 10^{-3}$. In this case, for $N \gtrsim 1000$ correlations play a role ($P_{\text{single}}/P_{\text{coop}} > 1$) and the intensity emitted along axis is dominated by them (second term). Thus for our clouds with N up to 5000, correlations *along axis* can play a role. As we showed in [9], the correlations that spontaneously emerge in the cloud are indeed along \hat{u}_z . On the contrary, for the direction *perpendicular* to the cloud axis, the double sum in the second term cancels when integrating over a solid angle because there are no constructive interferences, and the first term (incoherent intensity) always dominates. This is why no enhancement of intensity is observed in the perpendicular direction [57].

Now for what concerns $g_N^{(2)}(\hat{u}_c) \propto \sum_{i,j,k,l} \langle \hat{\sigma}_i^+ \hat{\sigma}_j^+ \hat{\sigma}_k^- \hat{\sigma}_l^- \rangle e^{-ik\hat{u}_c\cdot(\mathbf{r}_i + \mathbf{r}_j - \mathbf{r}_k - \mathbf{r}_l)}$, the same arguments hold: perpendicularly to the cloud axis there can be no constructive interferences. In this case, the Siegert relation is easily derived by cancelling double

sums, as in [15]. This is why the fact that the Siegert relation holds radially is not surprising.

Importantly, the fact that the Siegert relation holds *does not* mean that there are no *atomic* connected correlations. Indeed for the Siegert relation to be violated, one needs atomic connected correlations *with the right phase relation* so that they constructively interfere: in equation (3) of the main text, if the phase factors average-out to zero, the sum can cancel despite the individual connected correlations not being zero. This differentiates our two directions of observation: along the cloud axis the phase factors are right so that the Siegert relation is violated, in the perpendicular direction the phases average to zero, the Siegert relation is verified.

Dependence on drive intensity.

In figure S4, we present measurements of $g_N^{(2)}(0)$ and connected correlations $C(0)$ on Rabi frequency of the drive. We observe no modification up to $\Omega = 10\Gamma$ ($I/I_{\text{sat}} = 200$): Despite a very strong drive the data does not converge towards single atom behaviour. At weak drive the emission rate is low and we could not collect enough data to measure $g_N^{(2)}$ with enough precision. This regime would be interesting to study in future experiments.




FIG. S4. **Drive dependence.** Dependence of $g_N^{(2)}(0)$ (a) and connected correlations $C(0)$ (b) on Rabi frequency. The measurements span between $I/I_{\text{sat}} = 2\Omega^2/\Gamma^2 = 4$ and $I/I_{\text{sat}} = 200$.