

Light New Physics in $B \rightarrow K^{(*)}\nu\bar{\nu}$?

Wolfgang Altmannshofer*

*Santa Cruz Institute for Particle Physics and Department of Physics,
University of California, Santa Cruz, CA 95064, USA*

Andreas Crivellin†

*Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland and
Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland*

Huw Haigh‡ and Gianluca Inguglia§

Institute of High Energy Physics, 1050, Vienna, Austria

Jorge Martin Camalich¶

*Instituto de Astrofísica de Canarias, C/ Vía Láctea, s/n E38205 - La Laguna, Tenerife, Spain and
Universidad de La Laguna, Departamento de Astrofísica, La Laguna, Tenerife, Spain*

The study of the rare decays $B \rightarrow K^{(*)}\nu\bar{\nu}$ offers a window into the dynamics operating at the electroweak scale, allowing studies of the Standard Model and searches for heavy new physics. However, the analysis of these decays is also potentially sensitive to the on-shell production of new light bosons X through the process $B \rightarrow K^{(*)}X$. In particular, Belle II has recently measured $B^+ \rightarrow K^+\nu\bar{\nu}$, finding a 2.8σ excess under the assumption of heavy new physics. Since this excess is rather localized in the kaon energy, a fit that includes the decay mode $B^+ \rightarrow K^+X$ to the kinematic distributions prefers $m_X \approx 2\text{ GeV}$ with branching fraction $\text{Br}[B \rightarrow KX] = (8.8 \pm 2.5) \times 10^{-6}$ and a significance of $\approx 3.6\sigma$. However, no excess was found in the BaBar measurements of $B \rightarrow K^{(*)}\nu\bar{\nu}$, and a global analysis of the Belle II and BaBar data leads to $\text{Br}[B \rightarrow KX] = (5.1 \pm 2.1) \times 10^{-6}$ with a reduced significance of $\approx 2.4\sigma$. We then study various simplified dark-flavoured models and present a possible UV completion based on a gauged $B_3 - L_3$ symmetry, highlighting the discovery potential of dedicated searches for $B \rightarrow K^{(*)}X$ at Belle II.

I. INTRODUCTION

The Cabibbo-Kobayashi-Maskawa (CKM) mechanism [1] of the Standard Model (SM) was established by the B factories Belle [2] and BaBar [3] to be the leading source of quark flavour violation. Furthermore, the discovery of the Higgs boson [4, 5] at the Large Hadron Collider (LHC) at CERN [6, 7] completed the SM. However, this does not exclude the existence of beyond-the-SM physics but rather only limits its possible size and strongly motivates the experimental search for it, both at the high-energy frontier and with precision observables.

Historically, indirect evidence for new particles often preceded direct discoveries. In particular, the existence of the charm quark, the W boson, the top quark, and also the Higgs were expected due to indirect measurements of the Fermi interactions, kaon mixing, electroweak precision observables, etc. In this context, semi-leptonic B meson decays are a particularly useful tool for indirect new physics (NP) searches: they have distinct and clean experimental signatures and, in general, controllable the-

oretical uncertainties as well as suppressed rates, making them sensitive probes of beyond-the-SM physics. In fact, an interesting number of anomalies, i.e. deviations from the SM predictions arose [8]. In particular, global fits to semileptonic B decays involving $b \rightarrow c\tau\nu$ and $b \rightarrow s\ell^+\ell^-$ transitions show interesting hints for NP (see Refs. [9–11] for recent reviews).

Recently, the Belle II collaboration released an analysis of the closely related flavour-changing-neutral-current (FCNC) process $B^+ \rightarrow K^+\nu\bar{\nu}$ [12, 13] finding an excess of 2.8σ over the SM hypothesis. This significance was obtained under the assumption of heavy NP [14], allowing a connection to the anomalies in $b \rightarrow c\tau\nu$ and $b \rightarrow s\ell^+\ell^-$ [15–19]. However, we will pursue a different path here: The $B \rightarrow K^+\nu\bar{\nu}$ measurement can be reinterpreted as a search for the two-body $B \rightarrow KX$ decay if the undetected particle X is stable (approximately) or decays invisibly [20–25]. [74] For this, X must be quite light, with $m_X \leq m_B - m_K$, such that it would result in a resonant feature in the spectrum of the squared invariant mass of the di-neutrino system (denoted by q^2) of $B \rightarrow K\nu\bar{\nu}$. This is different from the case of the SM, or any heavy NP contribution, where all q^2 dependence arises from the form factors, phase space, and the experimental efficiency. [75]

Actually, the study of these types of experimental signatures has recently intensified in the context of dark-flavoured sectors, which are new light particles weakly coupled to the SM fermions with a rich flavour struc-

*Electronic address: waltmann@ucsc.edu

†Electronic address: andreas.crivellin@cern.ch

‡Electronic address: huw.haigh@oeaw.ac.at

§Electronic address: gianluca.inguglia@oeaw.ac.at

¶Electronic address: jcamalich@iac.es

ture that can induce FCNC (see [26, 27] for reviews). This includes a QCD axion emerging from the breaking of horizontal flavour symmetries [20, 28–35], axion-like particles (ALPs) [36–43] and new neutral gauge bosons such as light Z' bosons in the closely related process $b \rightarrow s\ell^+\ell^-$ [44–53].

Fortunately, Belle II provides information on the q^2 spectrum in their analysis, and, in fact, it shows a peak, localized around $q^2 = 4 \text{ GeV}^2$, suggesting that the possible excess might be better described by a new light mediator than with heavy NP. Therefore, in the next section, we will consider the experimental status of the $b \rightarrow s\nu\bar{\nu}$ transitions, including previous data from BaBar [54], and perform a global analysis and recast of the data under the hypothesis of light NP. In Sec. III, we will then study a series of simplified models that could be searched for by dedicated analyses of $b \rightarrow sX$ and propose an example of a possible UV completion before we conclude in Sec. IV.

II. EXPERIMENTAL STATUS AND STATISTICAL ANALYSIS

Due to the challenges in reconstructing the events in which the visible final state only involves a kaon (for charged B meson decays) or its decay products (for neutral B meson decays), searches for $B \rightarrow K^{(*)}\nu\bar{\nu}$ have only been performed at the B -factories Belle and BaBar.[76] Here, B mesons are produced in pairs from the decay of a $\Upsilon(4S)$ resonance and various analysis techniques to search for $B \rightarrow K^{(*)}\nu\bar{\nu}$ are available. In semileptonic tagged analyses (STA), one of the B mesons is reconstructed via its semileptonic decay while the other one is used to search for $B \rightarrow K^{(*)}\nu\bar{\nu}$. Similarly, hadronic-tagged analyses (HTA) use the hadronic decay of one of the B mesons and study the decay of interest of the other one. BaBar and Belle have searched for $B \rightarrow K^{(*)}\nu\bar{\nu}$ using both techniques [54–57]. The BaBar experiment found $\text{Br}[B^+ \rightarrow K^+\nu\bar{\nu}] = 1.5^{+1.7+0.4}_{-0.8-0.2} \times 10^{-5}$ and $\text{Br}[B \rightarrow K^*\nu\bar{\nu}] = 3.8^{+2.9}_{-2.6} \times 10^{-5}$ [54], while Belle provided 90% CL upper limits to the same process at the level of 2.7×10^{-5} .

An additional analysis technique, referred to as inclusive tag analysis (ITA), already adopted by Belle and Belle II in previous studies [58, 59], allows one to reconstruct *inclusively* the decay $B^+ \rightarrow K^+\nu\bar{\nu}$ from the charged kaon. This alternative methodology negates the requirement of a coincidental fully reconstructed hadronic (or semileptonic) B -decay to tag the event, thereby providing a higher signal efficiency at the cost of reduced signal purity due to increased background levels. In the recently released results obtained by Belle II, both ITA and HTA techniques are used [14], and the results are combined. Driven by the ITA technique with its higher statistics, Belle II obtained the first evidence for the decay $B^+ \rightarrow K^+\nu\bar{\nu}$ with 3.6σ significance, measuring $\text{Br}[B^+ \rightarrow K^+\nu\bar{\nu}] = (2.4 \pm 0.7) \times 10^{-5}$. This is in 2.8σ tension with the SM prediction of $\text{Br}[B^+ \rightarrow K^+\nu\bar{\nu}]_{\text{SM}} =$

$(0.497 \pm 0.037) \times 10^{-5}$ (excluding the contribution from $B^+ \rightarrow \tau^+\nu$ with $\tau^+ \rightarrow K^+\bar{\nu}$) [60] (see also [61, 62] for other recent SM predictions that agree well with the result that we are using).[77]

This result is interesting because it might indicate not only the presence of NP in the $b \rightarrow s\nu\bar{\nu}$ transitions but even the presence of new light states. This can be seen by looking at the supplemental material that accompanies the Belle II result [14]. The post-fit distributions of events as a function of q^2 indicate that the observed excess clusters in the region around $q^2 = 4 \text{ GeV}^2$, as can be seen in the right plot of Fig. 3 in the appendix, showing the data and SM yields from the Belle II search for $B^+ \rightarrow K^+\nu\bar{\nu}$ [14]. However, to evaluate the significance of such an excess, a fit taking into account the experimental resolution and all available data, including BaBar’s where no excess has been observed, has to be performed.

Therefore, we use the differential distributions of the $B^{0,+} \rightarrow K^{0,+}\nu\bar{\nu}$ measurements of Belle II [14] and BaBar [54] under the assumption that a light resonance escaping detection is present (i.e. X is either stable, sufficiently long-lived or decays invisibly) to evaluate the combined significance for NP. Furthermore, we will use the $B \rightarrow K^*\nu\bar{\nu}$ measurement of BaBar [54] to set an upper limit on $\text{Br}[B \rightarrow K^*X]$.

We fit the NP signal to the reconstructed data by modelling the resonance X with a Gaussian distribution. This is done via a binned maximum likelihood fit, using the `pyhf` software package [63]. In the combined fit to $B^+ \rightarrow K^+\nu\bar{\nu}$ (Belle and Babar) and $B^0 \rightarrow K^0\nu\bar{\nu}$ (BaBar) data, each measurement constitutes a channel in the statistical `pyhf` model with a fully correlated signal. Similarly, in the fit to the BaBar $B^0 \rightarrow K^{0,*}\nu\bar{\nu}$ and $B^+ \rightarrow K^{+,*}\nu\bar{\nu}$ distributions, the relative signal is fixed by isospin invariance to be (approximately) equal.

While we assume that the particle X has a negligible intrinsic (physical) width, we nonetheless assign a Gaussian standard deviation of 1.5 GeV^2 to its q^2 distribution to capture the detector resolution.[78]

The fit to the Belle-II data alone results in $\text{Br}[B^+ \rightarrow K^+X] = (8.8 \pm 2.5) \times 10^{-6}$ for $m_X \approx 2 \text{ GeV}$, with a significance of 3.6σ . The inclusion of BaBar data in the fit reduces the significance to 2.4σ and

$$\text{Br}[B \rightarrow KX] = (5.1 \pm 2.1) \times 10^{-6}, \quad (1)$$

is preferred (see Fig. 5 in the appendix for details). For $B \rightarrow K^*X$, only BaBar data is available, and since there is no excess seen, an upper limit of a few times 10^{-5} , depending on m_X , can be obtained. The results of the $B \rightarrow KX$ and $B \rightarrow K^*X$ fits are depicted in the left and right panels of Fig. 1, respectively.

III. MODELS OF LIGHT NEW PHYSICS

Here, we consider two options of light particles that can lead to $B \rightarrow KX$, a light neutral vector (i.e. a Z') and flavoured axions and ALPs [20, 28–43]. In both cases, X




FIG. 1: Left: Combined fit to $\text{Br}[B \rightarrow KX]$ from Belle II and BaBar as a function of the mass of X . Right: Same for $\text{Br}[B \rightarrow K^*X]$ (only BaBar data available).

should not decay to charged SM fermions as those decays would give prominent resonant signals in e.g. $b \rightarrow sl^+\ell^-$ decays. Couplings to electrons, muons, and light quarks should be absent or sufficiently small such that the new boson is long-lived enough to decay outside the detector or has a dominant invisible decay width. [79]

Since a flavour-changing bottom-strange coupling (g_{sb}) is needed to obtain the desired decay mode, in principle constraints from $B_s - \bar{B}_s$ mixing have to be considered. In fact, for a light Z' or ALP, one can set up an operator product expansion in m_X/m_b to calculate this new physics contribution and obtain bounds on the flavour changing couplings g_{sb} [20]. However, these limits are typically weaker than the ones obtained from decays such as $B \rightarrow K^{(*)}X$ because (in contrast to the case of heavy NP) the decay rate is proportional to the quadratic (not quartic) power of the couplings which is the same scaling as for the neutral-meson mixing amplitude.

A. Light vectors (Z')

Including couplings up to dimension 6, the interaction Lagrangian is [47]

$$\mathcal{L}_{Z'} \supset \left\{ g_L^{(4)} Z'_\mu (\bar{s}\gamma^\mu P_L b) + \frac{g_L^{(5)}}{\Lambda} Z'_{\mu\nu} (\bar{s}\sigma^{\mu\nu} P_R b) + \frac{g_L^{(6)}}{\Lambda^2} \partial^\nu Z'_{\mu\nu} (\bar{s}\gamma^\mu P_L b) + \text{h.c.} \right\} + \{L \leftrightarrow R\}, \quad (2)$$

where $Z'_{\mu\nu} = \partial_\mu Z'_\nu - \partial_\nu Z'_\mu$ is the Z' field strength tensor. For later convenience, we also introduce vector and axial-vector couplings $g_V^{(d)} = g_R^{(d)} + g_L^{(d)}$ and $g_A^{(d)} = g_R^{(d)} - g_L^{(d)}$.

In this setup, we find the following $B \rightarrow KZ'$ decay rates if only one of the couplings is switched on at a time

$$\Gamma_{B \rightarrow KZ'}^{(4)} = \frac{|g_V^{(4)}|^2 m_B^3}{64\pi m_{Z'}^2} \lambda^{\frac{3}{2}} f_+, \quad (3)$$

$$\Gamma_{B \rightarrow KZ'}^{(5)} = \frac{|g_V^{(5)}|^2 m_B m_{Z'}^2}{16\pi \Lambda^2} \left(1 + \frac{m_K}{m_B}\right)^{-2} \lambda^{\frac{3}{2}} f_T, \quad (4)$$

$$\Gamma_{B \rightarrow KZ'}^{(6)} = \frac{|g_V^{(6)}|^2 m_B^3 m_{Z'}^2}{64\pi \Lambda^4} \lambda^{\frac{3}{2}} f_+, \quad (5)$$

with the phase space function

$$\lambda = 1 + \frac{m_K^4}{m_B^4} + \frac{m_{Z'}^4}{m_B^4} - 2 \left(\frac{m_K^2}{m_B^2} + \frac{m_{Z'}^2}{m_B^2} + \frac{m_K^2 m_{Z'}^2}{m_B^4} \right). \quad (6)$$

The $B \rightarrow K$ form factors f_+ and f_T can be found in Refs. [61, 64, 65] and have to be evaluated at $q^2 = m_{Z'}^2$.

Similarly, we find for the $B \rightarrow K^*Z'$ decay rates

$$\Gamma_{B \rightarrow K^*Z'}^{(4)} = \frac{m_B}{32\pi} \lambda_*^{\frac{1}{2}} \left[|g_V^{(4)}|^2 \mathcal{F}_V + |g_A^{(4)}|^2 \mathcal{F}_A \right], \quad (7)$$

$$\Gamma_{B \rightarrow K^*Z'}^{(5)} = \frac{m_B}{8\pi} \frac{m_B^2}{\Lambda^2} \lambda_*^{\frac{1}{2}} \left[|g_V^{(5)}|^2 \mathcal{F}_T + |g_A^{(5)}|^2 \mathcal{F}_{T5} \right], \quad (8)$$

$$\Gamma_{B \rightarrow K^*Z'}^{(6)} = \frac{m_B}{32\pi} \frac{m_{Z'}^4}{\Lambda^4} \lambda_*^{\frac{1}{2}} \left[|g_V^{(6)}|^2 \mathcal{F}_V + |g_A^{(6)}|^2 \mathcal{F}_A \right], \quad (9)$$




FIG. 2: *Left:* Correlations between $B \rightarrow KZ'$ and $B \rightarrow K^*Z'$ (colored lines) for the different $\bar{s}bZ'$ operators considered in this work. These are compared to the experimental data stemming from the combination of Belle-II, Babar and Belle measurements, which is represented by the red regions corresponding to $\Delta\chi^2 = 2.3$ and $\Delta\chi^2 = 6.18$. Belle's upper limit (hatched region at 2σ) and the new Belle II measurement (blue vertical band at 1σ and 2σ). *Right:* preferred regions in the $g_L - g_R$ plane. One can see that (approximately) vectorial couplings of the order of 10^{-8} are suggested by current data.

with

$$\mathcal{F}_V = \lambda_* \left(1 + \frac{m_{K^*}}{m_B}\right)^{-2} V^2, \quad (10)$$

$$\mathcal{F}_A = \lambda_* \left(1 + \frac{m_{K^*}}{m_B}\right)^2 A_1^2 + \frac{32m_{K^*}^2}{m_{Z'}^2} A_{12}^2, \quad (11)$$

$$\mathcal{F}_T = \lambda_* T_1^2, \quad (12)$$

$$\mathcal{F}_{T5} = \left(1 - \frac{m_{K^*}^2}{m_B^2}\right)^2 T_2^2 + \frac{8m_{K^*}^2 m_{Z'}^2}{m_B^2 (m_B + m_{K^*})^2} T_{23}^2, \quad (13)$$

The phase space function λ_* is defined equivalently to the $B \rightarrow KZ'$ decay, with the replacement $m_K \rightarrow m_{K^*}$. The $B \rightarrow K^*$ form factors V , A_1 , A_{12} , T_1 , T_2 , T_{23} can be found in Refs. [65, 66].

The results can be seen in the plots of Fig. 2 that show the correlation between the $B \rightarrow KZ'$ and $B \rightarrow K^*Z'$ branching ratios (left plot) and the best-fit regions in the plane of Z' couplings to left-handed and right-handed quark currents (right plot). In both plots, the mass of the Z' is fixed to the best-fit value ~ 2 GeV, c.f. the discussion in section II. One can see that couplings only to left-handed or right-handed quarks, $g_{L/R}^{(4/6)}$, generate $B \rightarrow K^*Z'$ branching ratios that exceed the experimental bounds by a factor of few. The dimension-5 dipole couplings, $g_{L/R}^{(5)}$, lead to even larger $B \rightarrow K^*Z'$. Couplings that are dominantly vectorial are needed for explaining the enhanced $B \rightarrow K\nu\bar{\nu}$ branching ratio without violating the constraints from $B \rightarrow K^*\nu\bar{\nu}$. The size of the Z' couplings is very small, at the order of 10^{-8} .

B. Axion like particle

We consider now a massive pseudoscalar or axion-like particle (ALP) a coupled with a derivative coupling to the bs current,

$$\mathcal{L}_{\text{ALP}} \supset \frac{\partial_\mu a}{2f} (\bar{s}\gamma^\mu (g_V + g_A\gamma_5)b) + \text{h.c.} \quad (14)$$

where f is the ALP decay constant and where we have started from the vectorial basis for the couplings. Note that this also covers the case of neutral (pseudo)scalars with (effective) $g_S\bar{s}b + g_P\bar{s}\gamma_5b$ couplings (without a derivative), by identifying $g_{V(A)}m_b/2f = g_{S(P)}$ as per the equations of motion.

In this setup, the $B \rightarrow Ka$ and $B \rightarrow K^*a$ decay rates are given by,

$$\Gamma_{B \rightarrow Ka} = \frac{|g_V|^2 m_B^3}{64\pi f^2} \left(1 - \frac{m_K^2}{m_B^2}\right)^2 \lambda^{\frac{1}{2}} f_0^2, \quad (15)$$

$$\Gamma_{B \rightarrow K^*a} = \frac{|g_A|^2 m_B^3}{64\pi f^2} \lambda_*^{\frac{3}{2}} A_0^2, \quad (16)$$

where λ and λ_* are now the same as for $B \rightarrow KZ'$ and $B \rightarrow K^*Z'$, respectively, but replacing $m_{Z'}$ by m_a . The form factors f_0 and A_0 did not enter the expressions for the $B \rightarrow K^{(*)}Z'$ rates but can also be found in Refs. [61, 64–66]. They are evaluated at $q^2 = m_a^2$.

As is evident from equations (15) and (16) above, only vectorial couplings are capable of explaining $B \rightarrow K\nu\bar{\nu}$




FIG. 3: Left: Histogram showing Belle II data and MC [14] used in the NP fit, with four signal regions of the discriminator output, each containing 2% signal efficiency across three q^2 bins. Right: The highest sensitivity bin (i.e. with signal discriminator 0.98 – 1) split into finer q^2 bins, showing the resonant characteristics of the observed excess for which the red line shows the best fit. Note that these data were not used for the fit but are only shown as illustrations.

while axial vector couplings are constrained by $B \rightarrow K^* \nu \bar{\nu}$. The corresponding constraints on the couplings are qualitatively similar to the Z' . In terms of the ALP decay constant, Eq. (1) implies $F_V \equiv 2f/|g_V| = 3.1_{-0.5}^{+1.0} \times 10^8$ for $m_a = 2$ GeV, while the upper limit from $B \rightarrow K^* a$ leads to $F_A \equiv 2f/|g_A| \gtrsim 1.7 \times 10^8$ GeV at 2σ .

C. $B_3 - L_3$ Symmetry

Let us outline one possible UV complete model that could give rise to $B \rightarrow KX$, which is based on a gauged $B_3 - L_3$ symmetry. This means we assume that third-generation baryon and lepton numbers are oppositely charged under a new gauged $U(1)_X$ symmetry. This charge assignment is anomaly-free and can provide an explanation of the smallness of the CKM elements $V_{cb,ts}$ and $V_{ub,td}$. The reason for this is that in unbroken $U(1)_X$, no 1–3 and 2–3 elements in the quark Yukawa couplings are allowed. This means that while all quark masses and the Cabibbo angle can be obtained from the SM Higgs after it acquires its vacuum expectation value (VEV) from renormalizable dim-4 couplings, $V_{cb,ts}$ and $V_{ub,td}$ are zero at this level. One option to obtain non-zero $V_{cb,ts}$ and $V_{ub,td}$ is to introduce additional Higgs doublets charged under $U(1)_X$ to generate $V_{cb,ts}$ and $V_{ub,td}$ via their VEVs (possibly in conjunction with vector-like quarks [67]).

Since the $U(1)_X$ gauge boson needs to have both left-handed and right-handed bs coupling to explain $B \rightarrow KX$ without violating the bounds from $B \rightarrow K^* X$, we have to add two additional Higgs doublets charged under $B_3 - L_3$ with opposite $U(1)_X$ charges [68].[80] Furthermore, a singlet under $SU(2)_L$ charged under $U(1)_X$ is needed to obtain the preferred Z' mass $m_{Z'} \simeq 2$ GeV without overshooting $\text{Br}[B \rightarrow KX]$. Finally, note that

since 2 GeV is below the bottom and tau thresholds, such a Z' boson naturally decays invisibly to tau neutrinos and thus escapes detection.

IV. CONCLUSIONS AND OUTLOOK

Motivated by the recently observed excess of 2.8σ (w.r.t. the SM prediction) in $B^+ \rightarrow K^+ \nu \bar{\nu}$ by Belle II, we studied the option of light new physics, i.e. $B \rightarrow KX$ where the new boson X escapes detection. In fact, since the excess is localized around 4 GeV^2 , the investigation of light NP hypothesis is motivated. Assuming that the “effective” width of the particle is given by the detector resolution (i.e. the physical width is small compared to the detector resolution), we found a significance of $\approx 2.4\sigma$ and a preferred branching ratio of $\text{Br}[B \rightarrow KX] = (5.1 \pm 2.1) \times 10^{-6}$ for $m_X \approx 2$ GeV once the Belle II data is combined with the corresponding BaBar result.[81] Similarly we performed a fit to $B \rightarrow K^* \nu \bar{\nu}$ where no excess is observed.

We studied the two simplified NP physics models with a light vector and a light pseudoscalar with derivative couplings (i.e. an ALP). It is found that the flavour-changing coupling to bottom and strange quarks should be dominantly vectorial to explain the excesses. Finally, we proposed an example of a UV complete model, a gauged $B_3 - L_3$ symmetry, broken by three additional scalars (two $SU(2)_L$ doublets and one singlet) that naturally leads to the desired signature.

One should note that while our fitting approach provides insightful information on possible underlying processes in $B^+ \rightarrow K^+ \nu \bar{\nu}$, it leaves room for improvements. The reason is that while the $B^+ \rightarrow K^+ X$ carries kinematic information typical of a two-body decay, the orig-




FIG. 4: BaBar data and simulated MC [54], showing $B^+ \rightarrow K^+ \nu \bar{\nu}$ (left) and $B^0 \rightarrow K^0 \nu \bar{\nu}$ (right), provided in bins of $S_B = q^2/m_B^2$. The distribution of the fitted resonance is shown in green.




FIG. 5: Best fit and associated 1σ errors for $\text{Br}[B \rightarrow K^{(*)} X]$ as a function of m_X , for the fit to the BaBar distributions (green), the Belle II distribution (blue) and the combined fit to all data (red).

inal search is not optimized for such a case, and a dedicated experimental analysis will provide a better sensitivity. However, we hope that our work motivates dedicated analyses by the B factories.

Appendix A: Details on the fit

We perform maximum likelihood fits to the BaBar [54] and Belle II [14] data using the `pyhf` software package [63]. For this, we include the essential estimated

experimental systematic uncertainties, based on those quoted in the Belle II analysis, as nuisance parameters. The signal is fit to data using binned templates of the q^2 distributions, derived from post-fit Monte Carlo (MC) distributions, including individual contributions from $B\bar{B}$ and continuum SM background and the predictions of the SM contribution to $B \rightarrow K^{(*)} \nu \nu$. The corresponding templates are shown in Fig. 3 (left) for the Belle II analysis. They contain 12 bins in total: Three q^2 bins which are repeated in four bins of the signal discriminator output. These are constructed such that the expected signal efficiency is a constant 2% in the four regions.

For each of the four fit templates shown on the left in Fig. 3, we include an overall normalization uncertainty of 10% and the associated statistical uncertainty obtained from the measured number of events. The continuum background template has an additional uncertainty of 10% from shape systematics, which we allow for each bin to fluctuate individually.[82] To validate these choices, a fit including only the SM contribution, and no injected NP signal, is first conducted. From this, $\text{Br}[B^+ \rightarrow K^+ \nu \bar{\nu}] = (2.8 \pm 0.7) \times 10^{-5}$ is found, which is in good agreement with the result of the Belle II analysis.

The BaBar data are provided in bins of $S_B = q^2/m_B^2$ and the distributions are shown in Fig. 4 with contributions from the background, the SM to $B \rightarrow K \nu \nu$ and the NP signal. The associated signal efficiency in each bin is provided and considered in the fit via scaling of the resonance template. Only statistical uncertainties are accounted for in templates of the BaBar fit. In the case of the simultaneous fit to the Belle and BaBar data, the normalisation of the $B \rightarrow K \nu \nu$ templates are fixed by the SM expectations to the channels.

The NP signal is modelled with a Gaussian distribution, with the initial pre-fit yield of the template defined as the number of excess events observed in the Belle II

data. We assume a standard deviation of 1.5 GeV^2 for this distribution, the result of this can be seen in Fig. 3, where the NP contribution is scaled to the best-fit branching fraction. This assumption is also true of the fits to BaBar distributions, where the Gaussian mean and standard deviation are scaled as the other distributions ($S_B = q^2/m_B^2$).

Note that the look-elsewhere effect here is considered negligible since the fit only takes into account three (independent) q^2 bins (see left panel of Fig. 3).

Acknowledgments

A.C. gratefully acknowledges support by the Swiss

National Science Foundation (No. PP00P21-76884). G.I. and H.H. acknowledge support by the European Research Council under the grant agreement No. 947006 - *InterLeptons*. The research of W.A. is supported by the U.S. Department of Energy grant number DE-SC0010107. J.M.C. thanks MICINN for funding through the grant “DarkMaps” PID2022-142142NB-I00.

-
- [1] M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
- [2] A. Abashian et al. (Belle), *Nucl. Instrum. Meth. A* **479**, 117 (2002).
- [3] B. Aubert et al. (BaBar), *Nucl. Instrum. Meth. A* **479**, 1 (2002), hep-ex/0105044.
- [4] P. W. Higgs, *Phys. Lett.* **12**, 132 (1964).
- [5] F. Englert and R. Brout, *Phys. Rev. Lett.* **13**, 321 (1964).
- [6] G. Aad et al. (ATLAS), *Phys. Lett. B* **716**, 1 (2012), 1207.7214.
- [7] S. Chatrchyan et al. (CMS), *Phys. Lett. B* **716**, 30 (2012), 1207.7235.
- [8] A. Crivellin and B. Mellado (2023), 2309.03870.
- [9] J. Albrecht, D. van Dyk, and C. Langenbruch, *Prog. Part. Nucl. Phys.* **120**, 103885 (2021), 2107.04822.
- [10] D. London and J. Matias, *Ann. Rev. Nucl. Part. Sci.* **72**, 37 (2022), 2110.13270.
- [11] B. Capdevila, A. Crivellin, and J. Matias (2023), 2309.01311.
- [12] W. Altmannshofer, A. J. Buras, D. M. Straub, and M. Wick, *JHEP* **04**, 022 (2009), 0902.0160.
- [13] A. J. Buras, J. Girrbach-Noe, C. Niehoff, and D. M. Straub, *JHEP* **02**, 184 (2015), 1409.4557.
- [14] E. Ganiev, (On behalf of the Belle II Collaboration). Talk given at EPS-HEP 2023 “On Radiative and Electroweak Penguin Decays”, Hamburg, DESY (<https://indico.desy.de/event/34916/contributions/-146877/attachments/84380/111798/-EWP@Belle2.EPS.pdf>) (2023).
- [15] L. Allwicher, D. Becirevic, G. Piazza, S. Rosaura-Alcaraz, and O. Sumensari (2023), 2309.02246.
- [16] P. Athron, R. Martinez, and C. Sierra (2023), 2308.13426.
- [17] T. Felkl, A. Giri, R. Mohanta, and M. A. Schmidt (2023), 2309.02940.
- [18] C.-H. Chen and C.-W. Chiang (2023), 2309.12904.
- [19] X.-G. He, X.-D. Ma, and G. Valencia (2023), 2309.12741.
- [20] J. Martin Camalich, M. Pospelov, P. N. H. Vuong, R. Ziegler, and J. Zupan, *Phys. Rev. D* **102**, 015023 (2020), 2002.04623.
- [21] A. W. M. Guerrero and S. Rigolin, *Fortsch. Phys.* **71**, 2200192 (2023), 2211.08343.
- [22] A. Datta, A. Hammad, D. Marfatia, L. Mukherjee, and A. Rashed, *JHEP* **03**, 108 (2023), 2210.15662.
- [23] S. Bruggisser, L. Grabitz, and S. Westhoff (2023), 2308.11703.
- [24] M. Abdughani and Y. Reyimuaji (2023), 2309.03706.
- [25] A. Berezhnoy and D. Melikhov (2023), 2309.17191.
- [26] J. F. Kamenik and C. Smith, *JHEP* **03**, 090 (2012), 1111.6402.
- [27] E. Goudzovski et al., *Rept. Prog. Phys.* **86**, 016201 (2023), 2201.07805.
- [28] F. Wilczek, *Phys. Rev. Lett.* **49**, 1549 (1982).
- [29] J. L. Feng, T. Moroi, H. Murayama, and E. Schnapka, *Phys. Rev. D* **57**, 5875 (1998), hep-ph/9709411.
- [30] L. Calibbi, F. Goertz, D. Redigolo, R. Ziegler, and J. Zupan, *Phys. Rev. D* **95**, 095009 (2017), 1612.08040.
- [31] Y. Ema, K. Hamaguchi, T. Moroi, and K. Nakayama, *JHEP* **01**, 096 (2017), 1612.05492.
- [32] F. Arias-Aragon and L. Merlo, *JHEP* **10**, 168 (2017), [Erratum: *JHEP* **11**, 152 (2019)], 1709.07039.
- [33] K. Choi, S. H. Im, C. B. Park, and S. Yun, *JHEP* **11**, 070 (2017), 1708.00021.
- [34] L. Calibbi, D. Redigolo, R. Ziegler, and J. Zupan, *JHEP* **09**, 173 (2021), 2006.04795.
- [35] L. Di Luzio, A. W. M. Guerrero, X. P. Díaz, and S. Rigolin, *JHEP* **06**, 046 (2023), 2304.04643.
- [36] B. Gripaios, A. Pomarol, F. Riva, and J. Serra, *JHEP* **04**, 070 (2009), 0902.1483.
- [37] E. Izaguirre, T. Lin, and B. Shuve, *Phys. Rev. Lett.* **118**, 111802 (2017), 1611.09355.
- [38] M. J. Dolan, T. Ferber, C. Hearty, F. Kahlhoefer, and K. Schmidt-Hoberg, *JHEP* **12**, 094 (2017), [Erratum: *JHEP* **03**, 190 (2021)], 1709.00009.
- [39] F. Björkeröth, E. J. Chun, and S. F. King, *JHEP* **08**, 117 (2018), 1806.00660.
- [40] M. B. Gavela, R. Houtz, P. Quilez, R. Del Rey, and O. Sumensari, *Eur. Phys. J. C* **79**, 369 (2019), 1901.02031.
- [41] A. Carmona, C. Scherb, and P. Schwaller, *JHEP* **08**, 121 (2021), 2101.07803.
- [42] M. Bauer, M. Neubert, S. Renner, M. Schnubel, and A. Thamm, *JHEP* **09**, 056 (2022), 2110.10698.
- [43] T. Ferber, A. Filimonova, R. Schäfer, and S. Westhoff, *JHEP* **04**, 131 (2023), 2201.06580.
- [44] F. Sala and D. M. Straub, *Phys. Lett. B* **774**, 205 (2017), 1704.06188.

- [45] M. K. Mohapatra and A. Giri, Phys. Rev. D **104**, 095012 (2021), 2109.12382.
- [46] A. Datta, J. Kumar, J. Liao, and D. Marfatia, Phys. Rev. D **97**, 115038 (2018), 1705.08423.
- [47] W. Altmannshofer, M. J. Baker, S. Gori, R. Harnik, M. Pospelov, E. Stamou, and A. Thamm, JHEP **03**, 188 (2018), 1711.07494.
- [48] F. Sala, Nucl. Part. Phys. Proc. **303-305**, 14 (2018), 1809.11061.
- [49] F. Bishara, U. Haisch, and P. F. Monni, Phys. Rev. D **96**, 055002 (2017), 1705.03465.
- [50] D. Borah, L. Mukherjee, and S. Nandi, JHEP **12**, 052 (2020), 2007.13778.
- [51] L. Darmé, M. Fedele, K. Kowalska, and E. M. Sessolo, JHEP **03**, 085 (2022), 2106.12582.
- [52] A. Greljo, Y. Soreq, P. Stangl, A. E. Thomsen, and J. Zupan, JHEP **04**, 151 (2022), 2107.07518.
- [53] A. Crivellin, C. A. Manzari, W. Altmannshofer, G. Inguoglia, P. Feichtinger, and J. Martin Camalich, Phys. Rev. D **106**, L031703 (2022), 2202.12900.
- [54] J. P. Lees et al. (BaBar), Phys. Rev. D **87**, 112005 (2013), 1303.7465.
- [55] P. del Amo Sanchez et al. (BaBar), Phys. Rev. D **82**, 112002 (2010), 1009.1529.
- [56] O. Lutz et al. (Belle), Phys. Rev. D **87**, 111103 (2013), 1303.3719.
- [57] J. Grygier et al. (Belle), Phys. Rev. D **96**, 091101 (2017), [Addendum: Phys.Rev.D 97, 099902 (2018)], 1702.03224.
- [58] M. T. Prim et al. (Belle), Phys. Rev. D **101**, 032007 (2020), 1911.03186.
- [59] F. Abudinén et al. (Belle-II), Phys. Rev. Lett. **127**, 181802 (2021), 2104.12624.
- [60] W. G. Parrott, C. Bouchard, and C. T. H. Davies (HPQCD), Phys. Rev. D **107**, 014511 (2023), [Erratum: Phys.Rev.D 107, 119903 (2023)], 2207.13371.
- [61] D. Bećirević, G. Piazza, and O. Sumensari, Eur. Phys. J. C **83**, 252 (2023), 2301.06990.
- [62] Y. Amhis, M. Kenzie, M. Reboud, and A. R. Wiederhold (2023), 2309.11353.
- [63] L. Heinrich, M. Feickert, G. Stark, and K. Cranmer, J. Open Source Softw. **6**, 2823 (2021).
- [64] W. G. Parrott, C. Bouchard, and C. T. H. Davies ((HPQCD collaboration)§, HPQCD), Phys. Rev. D **107**, 014510 (2023), 2207.12468.
- [65] N. Gubernari, M. Reboud, D. van Dyk, and J. Virto (2023), 2305.06301.
- [66] A. Bharucha, D. M. Straub, and R. Zwicky, JHEP **08**, 098 (2016), 1503.05534.
- [67] W. Altmannshofer, S. Gori, M. Pospelov, and I. Yavin, Phys. Rev. D **89**, 095033 (2014), 1403.1269.
- [68] A. Crivellin, G. D'Ambrosio, and J. Heeck, Phys. Rev. D **91**, 075006 (2015), 1503.03477.
- [69] H. K. Dreiner, J. Y. Günther, and Z. S. Wang (2023), 2309.03727.
- [70] I. Adachi et al. (Belle-II), Phys. Rev. Lett. **124**, 141801 (2020), 1912.11276.
- [71] T. E. Browder et al. (CLEO), Phys. Rev. Lett. **86**, 2950 (2001), hep-ex/0007057.
- [72] X.-G. He, G. C. Joshi, H. Lew, and R. R. Volkas, Phys. Rev. D **44**, 2118 (1991).
- [73] A. Crivellin, A. Kokulu, and C. Greub, Phys. Rev. D **87**, 094031 (2013), 1303.5877.
- [74] Ref. [69] considered light NP. However, the interactions are still mediated by heavy particles, such that no peak in the q^2 spectrum occurs.
- [75] A light Z' boson in $b \rightarrow s\ell^+\ell^-$ has been proposed and studied in the literature [44–52], however it cannot explain the $b \rightarrow s\ell^+\ell^-$ anomalies because of di-muon invariant mass distribution in Drell-Yan production close to the Z mass [49] or $e^+e^- \rightarrow \mu^+\mu^-$ + invisible at Belle II [70] and $B \rightarrow K^{(*)}\nu\bar{\nu}$ [57].
- [76] Also a CLEO [71] analysis exists which is however not competitive.
- [77] If one considers specifically the Belle II ITA, which is the most sensitive of the two presented, the significance of the signal is slightly higher, providing $\text{Br}[B^+ \rightarrow K^+\nu\bar{\nu}] = (2.8 \pm 0.7) \times 10^{-5}$, which is in tension with the SM predictions at the 3.3σ level. We assume that the two analyses, ITA and HTA, select two independent populations of events; in other words, we assume no statistical correlation between the two individual ITA and HTA measurements, which is justified by the small sample overlap.
- [78] We checked that the significance is to a good approximation independent of the function used to parameterize the resonance X as long as its width captures the resolution of the detector. However, if e.g. a Crystal-ball function is used instead, a slightly lower mass of X is obtained.
- [79] This requirement excludes the often studied $L_\mu - L_\tau Z'$ boson [67, 72].
- [80] Note that in case the new CP-odd and CP-even Higgses of each doublet are quasi-degenerate, the effect in $B_q - \bar{B}_q$ mixing is suppressed [73].
- [81] This differs from the results found in Ref. [19] where a mass of X of a few hundred GeV was proposed.
- [82] Note that this does not fully capture the systematic uncertainty associated with the Belle-II measurement, which includes other sources attributed to experimental effects that cannot be included in this study. Furthermore, the fit conducted here has larger statistical uncertainties due to the lack of access to full Monte Carlo samples. However, it should provide a reasonable estimate.