QUASI-OPTIMAL DISCONTINUOUS GALERKIN DISCRETISATIONS OF THE p-DIRICHLET PROBLEM[∗]

JAN BLECHTA[†], PABLO ALEXEI GAZCA-OROZCO[‡], ALEX KALTENBACH[§], AND MICHAEL RŮŽIČKA‡

Abstract. The classical arguments employed when obtaining error estimates of Finite Element (FE) discretisations of elliptic problems lead to more restrictive assumptions on the regularity of the exact solution when applied to non-conforming methods. The so-called minimal regularity estimates available in the literature relax some of these assumptions, but are not truly of minimal regularity, since a data oscillation term appears in the error estimate. Employing an approach based on a smoothing operator, we derive for the first time error estimates for Discontinuous Galerkin (DG) type discretisations of non-linear problems with (p, δ) -structure that only assume the natural $W^{1,p}$ -regularity of the exact solution, and which do not contain any oscillation terms.

Key words. p-Dirichlet problem, Discontinuous Galerkin, a priori error estimates, quasioptimality, best-approximation

MSC codes. 35J66, 35J92, 65N12, 65N30

1. Introduction. In this paper, we examine Local Discontinuous Galerkin (LDG) and Incomplete Interior penalty Discontinuous Galerkin (IIDG) discretisations of non-linear problems of p-Dirichlet type, i.e.,

(1.1)
$$
-\operatorname{div} \mathcal{S}(\nabla u) = f \quad \text{in } \Omega, \mathbf{u} = \mathbf{0} \quad \text{on } \partial \Omega.
$$

Here $\Omega \subseteq \mathbb{R}^d$, $d \geq 2$, is a bounded polyhedral domain having a Lipschitz continuous boundary $\partial\Omega, f: \Omega \to \mathbb{R}^n$ is a given vector field, and we seek a vector field $u: \overline{\Omega} \to \mathbb{R}^n$ solving the system [\(1.1\).](#page-0-0) The non-linear operator $\mathcal{S} \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ is assumed to have (p, δ) -structure, cf. [Definition 2.1;](#page-3-0) the prototypical example falling in to this class is

$$
\mathbf{S}(\nabla \mathbf{u}) = (\delta + |\nabla \mathbf{u}|)^{p-2} \nabla \mathbf{u},
$$

where $p \in (1, \infty)$ and $\delta \geq 0$.

The central objective of this work is to establish a quasi-optimal (a priori) error estimate, i.e., a best-approximation result of the form

(1.2)
$$
\|\boldsymbol{F}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h) \n\lesssim \inf_{\boldsymbol{v}_h \in V_h} \left(\|\boldsymbol{F}(\tilde{\nabla}_h \boldsymbol{v}_h) - \boldsymbol{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{v}_h) \right),
$$

where $u_h \in V_h$ is the discrete solution, $\tilde{\nabla}_h \colon W^{1,p}(\mathcal{T}_h)^n \to L^p(\Omega)^{n \times d}$ is a suitable discrete gradient, and $\mathcal F$ and $m_{\varphi_{\beta_k}(\mu_k),h}$ are appropriate measures of the error related

[∗]Submitted to the editors November 27, 2023.

Funding: AK acknowledges support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) — within the Walter–Benjamin-Program (project number: 525389262) and the hospitality of the University of Pisa. JB's contribution to this work has been supported by Charles University Research program No. UNCE/SCI/023.

[†]Faculty of Mathematics and Physics, Charles University, 186 75 Prague, Czech Republic [\(blechta@karlin.mff.cuni.cz\)](mailto:blechta@karlin.mff.cuni.cz).

[‡]Department of Applied Mathematics, University of Freiburg, 79104, Freiburg, Germany [\(alexei.gazca@mathematik.uni-freiburg.de,](mailto:alexei.gazca@mathematik.uni-freiburg.de) [rose@mathematik.uni-freiburg.de\)](mailto:rose@mathematik.uni-freiburg.de).

[§] Institute of Mathematics, Technical University of Berlin, 10623, Berlin, Germany [\(kaltenbach@math.tu-berlin.de\)](mailto:kaltenbach@math.tu-berlin.de).

to the (p, δ) -structure of S; here, V_h is typically a space of broken polynomials, i.e., $V_h \coloneqq \mathbb{P}^k(\mathcal{T}_h)^n$, on a triangulation \mathcal{T}_h of Ω , which is meant to approximate the full space $W^{1,p}(\Omega)^n$. Crucially, when deriving the estimate [\(1.2\)](#page-0-1) we only assume the natural regularity of the continuous problem; namely, $u oldsymbol{\in} W_0^{1,p}(\Omega)^n$ and $\boldsymbol{f} \in W^{-1,p'}(\Omega)^n$. Then, as a corollary we obtain convergence rates, depending on additional regularity conditions.

The main issue with classical approaches based on Strang's Lemma (cf. [\[37\]](#page-32-0); see also [\[7,](#page-30-0) p. 106] and [\[17,](#page-31-0) Sec. 2.3]) or similar tools (cf. [\[14\]](#page-31-1)) when deriving error estimates for non-conforming discretisations, is that they rely either on integratingby-parts (and, e.g., requiring that div $\mathcal{S}(\nabla u)$ is an integrable function) or on assuming that traces of $\mathcal{S}(\nabla u)$ are well-defined on the mesh edges to handle consistency in the jump terms. The first work that got around the assumption of additional unnatural regularity is that of Gudi (cf. [\[20\]](#page-31-2)), where the author proved, using tools from a posteriori error analysis, the following estimate for the Dirichlet problem, assuming just $\boldsymbol{u} \in W_0^{1,2}(\Omega)^n$:

(1.3)
$$
\|\boldsymbol{u}-\boldsymbol{u}_h\|_{1,h}\lesssim \inf_{\boldsymbol{v}_h\in V_h}\|\boldsymbol{u}-\boldsymbol{v}_h\|_{1,h}+\mathrm{osc}_h(\boldsymbol{f}),
$$

where $\|\cdot\|_{1,h}$ is a broken Sobolev norm and $\operatorname{osc}_h(f)$ is a measure for the oscillation of $f \in L^2(\Omega)^n$. Such estimates are in the literature often qualified as being of minimal regularity, and the idea has been extended to various other contexts; see, e.g., [\[10,](#page-30-1) [22,](#page-31-3) [9,](#page-30-2) [21,](#page-31-4) [30,](#page-31-5) [2,](#page-30-3) [31,](#page-31-6) [3\]](#page-30-4) and, in particular, [\[25\]](#page-31-7) for systems of p-Dirichlet type. While this is certainly an improvement over the classical approach, it should be noted that such estimates are not quite minimal in their regularity assumptions, since an additional assumption on the data is needed, namely $-\Delta u = f \in L^2(\Omega)^n$. In contrast, we strive for minimal regularity estimates which entail the equivalence of the error and the distance to V_h .

A whole theory dealing with the characterisation of truly quasi-optimal discretisations of symmetric and elliptic linear problems in $W_0^{1,2}(\Omega)^n$ was developed in [\[38,](#page-32-1) [40,](#page-32-2) [39\]](#page-32-3) (see also [\[41,](#page-32-4) [29,](#page-31-8) [28\]](#page-31-9) for similar results for the Stokes system). In those works, the authors established that quasi-optimality is achieved exactly when the discretisation is fully algebraically consistent (i.e., if the exact solution belongs to the discrete space, then it coincides with the discrete solution), and fully stable (i.e., the map $(f \mapsto u_h)$: $W^{-1,2}(\Omega)^n \to V_h$ is well-defined and bounded). In particular, it is necessary that the scheme is entire, meaning that it is well-defined for general forcing terms $f \in W^{-1,2}(\Omega)^n$. This is achieved by working with modified schemes of the form: Find $u_h \in V_h$ such that for every $v_h \in V_h$, it holds that

(1.4)
$$
(\tilde{\nabla}_h \boldsymbol{u}_h, \nabla \boldsymbol{\mathcal{E}}_h \boldsymbol{v}_h)_{\Omega} = \langle \boldsymbol{f}, \boldsymbol{\mathcal{E}}_h \boldsymbol{v}_h \rangle_{W_0^{1,p}(\Omega)}.
$$

where $\mathcal{E}_h: V_h \to W_0^{1,2}(\Omega)$ is a bounded operator (a so-called *smoothing operator*) leaving $V_h \cap W_0^{1,2}(\Omega)$ invariant; see also [\[18\]](#page-31-10), where $\mathcal{E}_h: V_h \to W_0^{1,2}(\Omega)$ is applied to the unknown $u_h \in V_h$ as well. In order to obtain a quasi-optimal scheme, the smoothing operator is constructed in such a way that it preserves certain moments. More precisely, in the DG setting, for every $z_h \in V_h$, one has that

(1.5a)
$$
(\boldsymbol{q}_F, \boldsymbol{\mathcal{E}}_h \boldsymbol{z}_h)_F = (\boldsymbol{q}_F, \{\!\!\{\boldsymbol{z}_h\}\!\!\})_F \quad \text{for all } F \in \Gamma_h^i, \ \boldsymbol{q}_F \in \mathbb{P}^{k-1}(F),
$$

(1.5b)
$$
(\boldsymbol{q}_K, \boldsymbol{\mathcal{E}}_h \boldsymbol{z}_h)_K = (\boldsymbol{q}_K, \boldsymbol{z}_h)_K \quad \text{for all } K \in \mathcal{T}_h, \ \boldsymbol{q}_K \in \mathbb{P}^{k-2}(K),
$$

where $\{\cdot\}$ stands for facet average, Γ_h^i is the set of internal facets, and we set $\mathbb{P}^{-1}(K) := \emptyset$; see [section 2](#page-2-0) for more details on the DG notation.

An important consequence of the preservation properties [\(1.5\)](#page-1-0) (at least in the linear case) is that in practice one does not need to implement the smoothing operator on the left-hand side of [\(1.4\).](#page-1-1) To see this, take an arbitrary element $\mathbf{T}_h \in \Sigma_h :=$ $\mathbb{P}^{k-1}(\mathcal{T}_h)^{n\times d}$ (which will represent the flux). Then, using the preservation properties (1.5) , for every $z_h \in V_h$, integration-by-parts yields that

$$
(\mathbf{T}_h, \nabla \mathcal{E}_h \mathbf{z}_h)_{\Omega} = -(\text{div}_h \mathbf{T}_h, \mathcal{E}_h \mathbf{z}_h)_{\Omega} + (\llbracket \mathbf{T}_h \mathbf{n} \rrbracket, \mathcal{E}_h \mathbf{z}_h)_{\Gamma_h^i} + (\llbracket \mathbf{T}_h \rrbracket, \llbracket \mathcal{E}_h \mathbf{z}_h \mathbf{n} \rrbracket)_{\Gamma_h}
$$
\n
$$
= -(\text{div}_h \mathbf{T}_h, \mathbf{z}_h)_{\Omega} + (\llbracket \mathbf{T}_h \mathbf{n} \rrbracket, \mathbf{z}_h)_{\Gamma_h^i}
$$
\n
$$
= (\mathbf{T}_h, \mathcal{G}_h \mathbf{z}_h)_{\Omega},
$$

where $\mathcal{G}_h: W^{1,p}(\mathcal{T}_h)^n \to L^p(\Omega)^{n \times d}$ is the usual DG gradient (cf. [subsection 2.2.2\)](#page-6-0). In other words, if the fluxes are broken polynomials of one degree less than polynomial degree of the solution $u_h \in V_h$, the smoothing operator needs to be implemented only in the right-hand side (akin to modifying only the forcing term as $\mathcal{E}_h^* f \in V_h^*$, where ${\mathcal E}_h^*: (W^{1,p}(\mathcal{T}_h)^n)^* \to V_h^*$ is the adjoint to ${\mathcal E}_h: V_h \to W^{1,p}(\mathcal{T}_h)^n$.

In this work, we first prove a best-approximation result of the type [\(1.2\)](#page-0-1) for a non-linear analogue of the problem (1.4) (cf. [Theorem 4.6\)](#page-11-0); since we do not have a Hilbert structure at our disposal (in contrast to the works [\[38,](#page-32-1) [40,](#page-32-2) [39\]](#page-32-3)), our approach is based on working directly with an error equation. This is to our knowledge the first best-approximation result for non-conforming discretisations of non-linear systems of p-Dirichlet type that is genuinely minimal in its regularity assumptions. One disadvantage of the scheme in the non-linear case is that for polynomial degree $k \geq$ 2, the argument [\(1.6\)](#page-2-1) cannot be employed (since non-linear functions of elementwise polynomials are in general not element-wise polynomials) and it is, therefore, necessary to implement the smoothing operator also in the left-hand side. As an alternative to this, we propose also a mixed discretisation in which the flux variable belongs, by construction, to $\Sigma_h = \mathbb{P}^{k-1}(\mathcal{T}_h)^{n \times d}$ and, as a result, [\(1.6\)](#page-2-1) applies. For this mixed discretisation, we prove a minimal regularity error estimate as well (cf. [Theorem 5.5\)](#page-18-0).

This paper is organized as follows: In [section 2,](#page-2-0) we introduce the employed notation, define relevant function spaces, basic assumptions on S , and the used discrete operators. In [section 3,](#page-7-0) we recall the construction of the smoothing operator and prove an interpolation error estimate in terms of N-functions. In [section 4,](#page-9-0) we introduce the continuous and the discrete primal problem, establish their well-posedness and the validity of a quasi-optimal (a priori) error estimate of the form [\(1.2\).](#page-0-1) Then, from the latter, we deduce the convergence of the discrete primal problem under minimal regularity assumptions and derive fractional error decay rates given fractional regularity assumptions expressed in Nikolski˘ı spaces. In addition, aided by the quasi-optimal error estimate, we carry out an ansatz class competition that shows that the approximation capabilities of LDG and IIDG approximations and continuous Lagrange approximations of the problem (1.1) are comparable. In [section 5,](#page-14-0) we introduce the continuous and the discrete mixed problem, establish their well-posedness and the validity of a quasi-optimal (a priori) error estimate in a similar form as [\(1.2\).](#page-0-1) In [section 6,](#page-24-0) we carry out numerical experiments to complement the theoretical findings.

2. Preliminaries. We employ $c, C > 0$ to denote generic constants, that may change from line to line and may depend only on the polynomial degree k , the chunkiness ω_0 , the characteristics of S, and the dimensions n, d.

Moreover, we write $f \leq g$ if there exist a constant $c > 0$ such that $f \leq cg$, and $f \sim g$ if and only if there exists constants $c, C > 0$ such that $cf \leq g \leq Cf$.

Throughout the entire paper, let $\Omega \subseteq \mathbb{R}^d$, $d \in \mathbb{N}$, is a bounded, polyhedral Lipschitz domain and $M \subseteq \mathbb{R}^d$, $d \in \mathbb{N}$, a (Lebesgue) measurable set. Then, for every $k \in \mathbb{N}$ and $p \in [1,\infty]$, we employ the customary Lebesgue spaces $(L^p(M), \|\cdot\|_{p,M})$ and Sobolev spaces $(W^{k,p}(M), \|\cdot\|_{k,p,M})$. The space $W_0^{1,p}(M)$ is defined as the closure of the vector space of smooth and compactly supported functions $C_c^{\infty}(M)$ in $W^{1,p}(M)$. We equip $\overline{W}^{1,p}_0(M)$ with the norm $\|\nabla \cdot \|_{p,M}$.

We always denote vector-valued functions by boldface letters and tensor-valued functions by capital boldface letters. The Euclidean scalar product between two vectors $\mathbf{a} = (a_1, \ldots, a_n)^\top$, $\mathbf{b} = (b_1, \ldots, b_n)^\top \in \mathbb{R}^n$, $n \in \mathbb{N}$, is defined by $\mathbf{a} \cdot \mathbf{b} \coloneqq \sum_{i=1}^n a_i b_i$, while the Frobenius scalar product between two tensors $\mathbf{A} = (A_{ij})_{1 \leq i \leq n; 1 \leq j \leq \ell}$, $\mathbf{B} =$ $(B_{ij})_{1\leq i\leq n;1\leq j\leq \ell} \in \mathbb{R}^{n\times \ell}, n,\ell \in \mathbb{N}$, is defined by $\mathbf{A} : \mathbf{B} \coloneqq \sum_{i=1}^{n} \sum_{j=1}^{\ell} A_{ij} B_{ij}$. Then, the Euclidean norm of a vector $\mathbf{a} \in \mathbb{R}^n$, $n \in \mathbb{N}$, is defined by $|\mathbf{a}| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$, while the Frobenius norm of a tensor $A \in \mathbb{R}^{n \times \ell}$, $n, \ell \in \mathbb{N}$, is defined by $|A| := \sqrt{A : A}$.

The mean value of a locally integrable function f is denoted by $\langle f \rangle_M := \int_M f \, dx$ $\mathcal{L} := \frac{1}{|M|} \int_M f \, dx$. Moreover, we employ the notation $(f, g)_M := \int_M fg \, dx$, whenever the right-hand side is well-defined.

Drawing from the theory of Orlicz spaces $L^{\psi}(M)$ (cf. [\[33\]](#page-31-11)) and generalized Orlicz spaces $L^{\psi(\cdot)}(M)$ (cf. [\[24\]](#page-31-12)), we employ N-functions $\psi: \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ and generalized N-functions $\psi: M \times \mathbb{R}^{\geq 0} \to \mathbb{R}^{20}$, i.e., ψ is a Caratheodory function such that $\psi(x, \cdot)$ is an N-function for a.e. $x \in M$, respectively. The modular is defined via $\rho_{\psi,M}(f) \coloneqq \int_M \psi(|f|) dx$ if ψ is an N-function, and via $\rho_{\psi,M}(f) \coloneqq \int_M \psi(x, |f(x)|) dx$, if ψ is a generalized N-function. An N-function ψ satisfies the Δ_2 -condition (in short, $\psi \in \Delta_2$), if there exists $K > 2$ such that for every $t \geq 0$, it holds that $\psi(2t) \leq K \psi(t)$. We denote the smallest such constant by $\Delta_2(\psi) > 0$. We define the (convex) conjugate (generalized) N-function $\psi^*: M \times \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ via $\psi^*(x, t) \coloneqq \sup_{s \geq 0} ts - \psi(x, s)$ for all $t \geq 0$ and a.e. $x \in M$. If $\psi, \psi^* \in \Delta_2$, then we have that

$$
\psi^* \circ \psi' \sim \psi,
$$

with constants depending only on $\Delta_2(\psi), \Delta_2(\psi^*) > 0$. We will also need the ε -Young inequality: for every $\varepsilon > 0$, there exits a constant $c_{\varepsilon} > 0$, depending only on $\Delta_2(\psi), \Delta_2(\psi^*) < \infty$, such that for every $s, t \geq 0$, it holds that

(2.2)
$$
t s \leq \varepsilon \psi(t) + c_{\varepsilon} \psi^*(s).
$$

2.1. Basic properties of the non-linear operator. Throughout the paper, we always assume that the non-linear operator $\mathcal S$ has (p, δ) -structure. A detailed discussion and proofs can be found, e.g., in [\[12,](#page-30-5) [35\]](#page-32-5).

For $p \in (1,\infty)$ and $\delta \geq 0$, we define a special N-function $\varphi = \varphi_{p,\delta} : \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ via αt

(2.3)
$$
\varphi(t) \coloneqq \int_0^t \varphi'(s) \, ds, \quad \text{where} \quad \varphi'(t) \coloneqq (\delta + t)^{p-2} t, \quad \text{ for all } t \geq 0.
$$

The properties of φ are discussed in detail in [\[12,](#page-30-5) [35,](#page-32-5) [27\]](#page-31-13). An important tool in our analysis play *shifted N-functions* $\{\psi_a\}_{a\geq 0}$, cf. [\[13,](#page-30-6) [35\]](#page-32-5). For a given N-function $\psi: \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ we define the family of shifted N-functions $\psi_a: \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$, $a \geq 0$, via

(2.4)
$$
\psi_a(t) := \int_0^t \psi_a'(s) \, ds, \quad \text{where} \quad \psi_a'(t) := \psi'(a+t) \frac{t}{a+t}, \quad \text{ for all } t \ge 0.
$$

DEFINITION 2.1 $((p, \delta)$ -structure). Let $S \in C^0(\mathbb{R}^{n \times d}, \mathbb{R}^{n \times d})$ satisfy $S(0) = 0$. Then, we say that S has (p, δ) -structure if for some $p \in (1, \infty)$, $\delta \in [0, \infty)$, and the N-function $\varphi = \varphi_{p,\delta}$ (cf. [\(2.3\)](#page-3-1)), there exist constants $C_0, C_1 > 0$ such that

(2.5)
$$
(\mathcal{S}(\mathbf{Q}) - \mathcal{S}(\mathbf{P})) : (\mathbf{Q} - \mathbf{P}) \geq C_0 \varphi_{|\mathbf{Q}|} (|\mathbf{Q} - \mathbf{P}|),
$$

$$
|\mathcal{S}(\mathbf{Q}) - \mathcal{S}(\mathbf{P})| \leq C_1 \varphi'_{|\mathbf{Q}|} (|\mathbf{Q} - \mathbf{P}|).
$$

are satisfied for all $\mathbf{Q}, \mathbf{P} \in \mathbb{R}^{n \times d}$ with $\mathbf{Q} \neq \mathbf{0}$. The constants $C_0, C_1 > 0$ and $p \in (1, \infty)$ are called the characteristics of S .

REMARK 2.2. (i) Assume that S has (p, δ) -structure for some $\delta \in [0, \delta_0]$. Then, if not otherwise stated, the constants in the estimates depend only on the characteristics of S and on $\delta_0 \geq 0$, but are independent of $\delta \geq 0$.

(ii) Let φ be defined in [\(2.3\)](#page-3-1) and $\{\varphi_a\}_{a\geq 0}$ be the corresponding family of the shifted N-functions. Then, the operators $\mathcal{S}_a : \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}_{sym}$, $a \ge 0$, for every $a \ge 0$ and $\mathbf{Q} \in \mathbb{R}^{n \times d}$, defined via

(2.6)
$$
\mathcal{S}_a(\mathbf{Q}) := \frac{\varphi_a'(|\mathbf{Q}|)}{|\mathbf{Q}|} \mathbf{Q} = (\delta + a + |\mathbf{Q}|)^{p-2} \mathbf{Q},
$$

have $(p, \delta + a)$ -structure. In this case, the characteristics of S_a depend only on $p \in (1,\infty)$ and are independent of $\delta \geq 0$ and $a \geq 0$.

(iii) Note that $\varphi_a(t) \sim (\delta + a + t)^{p-2}t^2$ and $(\varphi_a)^*(t) \sim ((\delta + a)^{p-1} + t)^{p'-2}t^2$ uniformly with respect to $t, a \geq 0$. The families ${\{\varphi_a\}}_{a\geq 0}$ and ${\{\varphi_a\}}^*_{a\geq 0}$ satisfy the Δ_2 -condition uniformly with respect to $a \geq 0$, with $\Delta_2(\varphi_a) \leq 2^{\max\{2,p\}}$ and $\Delta_2((\varphi_a)^*) \lesssim 2^{\max\{2,p'\}}$, respectively. Moreover, for all $0 \leq a \leq b$, we have that for every $t \geq 0$, it holds that $(\varphi_a)^*(t) \geq (\varphi_b)^*(t)$, $\varphi_a(t) \leq \varphi_b(t)$ if $p \geq 2$ and $(\varphi_a)^*(t) \leq$ $(\varphi_b)^*(t), \varphi_a(t) \ge \varphi_b(t)$ if $p \le 2$.

Closely related to the non-linear operator S with (p, δ) -structure are the nonlinear operators $\mathcal{F}, \mathcal{F}^* \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$, for every $\mathsf{Q} \in \mathbb{R}^{n \times d}$, defined via

(2.7)
$$
\mathcal{F}(\mathbf{Q}) \coloneqq (\delta + |\mathbf{Q}|)^{\frac{p-2}{2}} \mathbf{Q},
$$

$$
\mathcal{F}^*(\mathbf{Q}) \coloneqq (\delta^{p-1} + |\mathbf{Q}|)^{\frac{p'-2}{2}} \mathbf{Q}.
$$

The connections between $\mathcal{S}, \mathcal{F}, \mathcal{F}^* \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ and $\varphi_a, (\varphi_a)^* \colon \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}, a \geq 0,$ are best explained by the following result (cf. [\[12,](#page-30-5) [35,](#page-32-5) [14\]](#page-31-1)).

PROPOSITION 2.3. Let S have (p, δ) -structure, let φ be defined in [\(2.3\)](#page-3-1), and let $\mathcal{F}, \mathcal{F}^*$ be defined in [\(2.7\)](#page-4-0). Then, uniformly with respect to $\mathbf{Q}, \mathbf{P} \in \mathbb{R}^{n \times d}$, we have that

(2.8)
\n
$$
(\mathcal{S}(\mathbf{Q}) - \mathcal{S}(\mathbf{P})) : (\mathbf{Q} - \mathbf{P}) \sim |\mathcal{F}(\mathbf{Q}) - \mathcal{F}(\mathbf{P})|^2
$$
\n
$$
\sim \varphi_{|\mathbf{Q}|} (|\mathbf{Q} - \mathbf{P}|)
$$
\n
$$
\sim (\varphi_{|\mathbf{Q}|})^* (|\mathcal{S}(\mathbf{Q}) - \mathcal{S}(\mathbf{P})|)
$$
\n
$$
\sim (\varphi^*)_{|\mathcal{S}(\mathbf{Q})|} (|\mathcal{S}(\mathbf{Q}) - \mathcal{S}(\mathbf{P})|),
$$

(2.9)
$$
|\mathcal{S}(\mathbf{Q}) - \mathcal{S}(\mathbf{P})| \sim \varphi'_{|\mathbf{Q}|}(|\mathbf{Q} - \mathbf{P}|),
$$

(2.10)
$$
\left|\mathcal{F}^*(\mathbf{Q})-\mathcal{F}^*(\mathbf{P})\right|^2 \sim (\varphi^*)_{|\mathbf{Q}|}(|\mathbf{Q}-\mathbf{P}|),
$$

(2.11)
$$
|\mathcal{F}^*(\mathcal{S}(\mathsf{P})) - \mathcal{F}^*(\mathcal{S}(\mathsf{Q}))|^2 \sim |\mathcal{F}(\mathsf{P}) - \mathcal{F}(\mathsf{Q})|^2.
$$

The constants in (2.8) – (2.11) depend only on the characteristics of S.

REMARK 2.4. For the operators $\mathcal{S}_a: \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}_{sym}$, $a \ge 0$, defined in [\(2.6\),](#page-4-3) the assertions of [Proposition 2.3](#page-4-4) hold with $\varphi \colon \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ replaced by $\varphi_a \colon \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$, $a \geq 0$.

The following results can be found in [\[13,](#page-30-6) [35\]](#page-32-5).

LEMMA 2.5 (Change of shift). Let φ be defined in [\(2.3\)](#page-3-1) and let $\mathcal F$ be defined in [\(2.7\)](#page-4-0). Then, for each $\varepsilon > 0$, there exists $c_{\varepsilon} \geq 1$ (depending only on $\varepsilon > 0$ and p) such that for every $\mathbf{Q}, \mathbf{P} \in \mathbb{R}^{n \times d}$ and $t \geq 0$, it holds that

$$
\varphi_{|\mathbf{P}|}(t) \le c_{\varepsilon} \varphi_{|\mathbf{Q}|}(t) + \varepsilon |\mathcal{F}(\mathbf{P}) - \mathcal{F}(\mathbf{Q})|^2,
$$

$$
(\varphi_{|\mathbf{P}|})^*(t) \le c_{\varepsilon} (\varphi_{|\mathbf{Q}|})^*(t) + \varepsilon |\mathcal{F}(\mathbf{P}) - \mathcal{F}(\mathbf{Q})|^2.
$$

2.2. Mesh regularity. Throughout the entire paper, $\{\mathcal{T}_h\}_{h>0}$ always denotes a family of conforming triangulations of $\overline{\Omega} \subseteq \mathbb{R}^d$, $d \geq 2$, cf. [\[11\]](#page-30-7), consisting of ddimensional closed simplices K. The parameter $h > 0$, refers to the maximal meshsize of \mathcal{T}_h , i.e., if we define $h_K := \text{diam}(K)$ for all $K \in \mathcal{T}_h$, then $h := \max_{K \in \mathcal{T}_h} h_K$. For simplicity, we assume that $h \leq 1$. For a simplex $K \in \mathcal{T}_h$, we denote by $\rho_K > 0$, the supremum of diameters of inscribed balls. We assume that there exists a constant $\omega_0 > 0$, independent of $h > 0$, such that $h_K \rho_K^{-1} \leq \omega_0$ for every $K \in \mathcal{T}_h$. The smallest such constant is called the chunkiness of $\{\mathcal{T}_h\}_{h>0}$.

We define the sets of $(d-1)$ -dimensional faces Γ_h , interior faces Γ_h^i , and boundary faces Γ_h^{∂} of the partition \mathcal{T}_h via

$$
\Gamma_h^i := \{ K \cap K' \mid K, K' \in \mathcal{T}_h, \dim_{\mathscr{H}} (K \cap K') = d - 1 \},\
$$

\n
$$
\Gamma_h^{\partial} := \{ K \cap \partial \Omega \mid K \in \mathcal{T}_h, \dim_{\mathscr{H}} (K \cap \partial \Omega) = d - 1 \},\
$$

\n
$$
\Gamma_h := \Gamma_h^i \cup \Gamma_h^{\partial},
$$

where for every subset $S \subseteq \mathbb{R}^d$, we denote by $\dim_{\mathscr{H}}(S) := \inf \{d' \geq 0 \mid \mathscr{H}^{d'}(S) =$ 0}, the Hausdorff dimension $(\mathcal{H}^{d'}(S)$ represents here the d'-dimensional Hausdorff measure of S). The (local) mesh-size function $h_{\mathcal{T}}: \overline{\Omega} \to \mathbb{R}$ is defined via $h_{\mathcal{T}}|_K := h_K$ for all $K \in \mathcal{T}_h$. The (local) face-size function $h_{\Gamma} \colon \Gamma_h \to \mathbb{R}$ is defined via $h_{\Gamma}|_F \coloneqq h_F \coloneqq$ diam(F) for all $F \in \Gamma_h$.

2.2.1. Broken function spaces and projectors. For every $k \in \mathbb{N}_0$ and $K \in$ \mathcal{T}_h , we denote by $\mathbb{P}^k(K)$, the space of polynomials of degree at most k on K. Then, for given $k \in \mathbb{N}_0$, we define the space of *broken polynomials of global degree at most* k via

$$
\mathbb{P}^k(\mathcal{T}_h) \coloneqq \left\{ v_h \in L^{\infty}(\Omega) \mid v_h|_K \in \mathbb{P}^k(K) \text{ for all } K \in \mathcal{T}_h \right\},\
$$

and the space of *broken polynomials* via $\mathbb{P}(\mathcal{T}_h) := \bigcup_{k \in \mathbb{N}} \mathbb{P}^k(\mathcal{T}_h)$. In addition, we define the space of element-wise continuous functions via

$$
C^0(\mathcal{T}_h) \coloneqq \{ w_h \in L^\infty(\Omega) \mid w_h|_K \in C^0(K) \text{ for all } K \in \mathcal{T}_h \},
$$

and for given $p \in (1,\infty)$, we define the *broken Sobolev space* via

$$
W^{1,p}(\mathcal{T}_h) \coloneqq \left\{ w_h \in L^p(\Omega) \mid w_h|_K \in W^{1,p}(K) \text{ for all } K \in \mathcal{T}_h \right\}.
$$

For each $w_h \in W^{1,p}(\mathcal{T}_h)^n$, we denote by $\nabla_h w_h \in L^p(\Omega)^{n \times d}$ the local gradient: for every $K \in \mathcal{T}_h$ it is defined via $(\nabla_h \boldsymbol{w}_h)|_K := \nabla(\boldsymbol{w}_h|_K)$ for all $K \in \mathcal{T}_h$. For each $K \in \mathcal{T}_h$, $w_h \in W^{1,p}(\mathcal{T}_h)^n$ admits a trace $\mathrm{tr}^K(w_h) \in L^p(\partial K)^n$. For each face $F \in \Gamma_h$ of a given

element $K \in \mathcal{T}_h$, we define this interior trace via $\operatorname{tr}_F^K(\boldsymbol{w}_h) \in L^p(F)^n$. Then, given some multiplication operator $\odot: \mathbb{R}^n \times \mathbb{R}^d \to \mathbb{R}^\ell$, $\ell \in \mathbb{N}$, for every $w_h \in W^{1,p}(\mathcal{T}_h)^n$ and interior faces $F \in \Gamma_h^i$ shared by adjacent elements $K_F^-, K_F^+ \in \mathcal{T}_h$, we denote by

$$
\begin{aligned} \{\!\!\{\boldsymbol{w}_h\}\!\!\}_F &\coloneqq \tfrac{1}{2} \big(\mathrm{tr}_{F}^{K^+}(\boldsymbol{w}_h) + \mathrm{tr}_{F}^{K^-}(\boldsymbol{w}_h)\big) \in L^p(F)^n \,, \\ \[\![\boldsymbol{w}_h\odot\boldsymbol{n}]\!]_F &\coloneqq \mathrm{tr}_{F}^{K^+}(\boldsymbol{w}_h) \odot\boldsymbol{n}_F^+ + \mathrm{tr}_{F}^{K^-}(\boldsymbol{w}_h) \odot\boldsymbol{n}_F^- \in L^p(F)^{\ell} \,, \end{aligned}
$$

the *average* and *jump*, respectively, of w_h on F. Moreover, for every $w_h \in W^{1,p}(\mathcal{T}_h)^n$ and boundary faces $F \in \Gamma_h^{\partial}$, we define boundary averages and boundary jumps, respectively, via

$$
\{\!\!\{\boldsymbol{w}_h\}\!\!\}_F \coloneqq \operatorname{tr}_F^{\Omega}(\boldsymbol{w}_h) \in L^p(F)^n,
$$

$$
[\![\boldsymbol{w}_h \odot \boldsymbol{n}]\!]_F \coloneqq \operatorname{tr}_F^{\Omega}(\boldsymbol{w}_h) \odot \boldsymbol{n} \in L^p(F)^{\ell}.
$$

If there is no danger of confusion, we will omit the index $F \in \Gamma_h$; in particular, if we interpret jumps and averages as global functions defined on the whole of Γ_h . In addition, for every $w_h \in W^{1,p}(\mathcal{T}_h)^n$, we introduce the DG norm via

$$
\|\boldsymbol{w}_h\|_{h,p}\coloneqq \left(\|\nabla_h \boldsymbol{w}_h\|_{p,\Omega}^p+\big\|h_\Gamma^{-1/p'}\llbracket \boldsymbol{w}_h\otimes \boldsymbol{n}\rrbracket\big\|_{p,\Gamma_h}^p\right)^{1/p}
$$

which turns $W^{1,p}(\mathcal{T}_h)^n$ into a Banach space. We denote by $\Pi_h^k: L^1(\Omega) \to \mathbb{P}^k(\mathcal{T}_h)$, which turns $W \nightharpoonup (h)$ linto a Banach space. We denote by Π_h . $E(u) \nightharpoonup \mathbb{I}^*(h)$,
the (local) L^2 -projection onto $\mathbb{P}^k(\mathcal{T}_h)$, defined for every $v \in L^1(\Omega)$ via $(\Pi_h^k v, v_h)$ $\Omega =$ $(v, v_h)_{\Omega}$ for all $v_h \in \mathbb{P}^k(\mathcal{T}_h)$. Analogously, we define the (local) L^2 -projections into $\mathbb{P}^k(\mathcal{T}_h)^n$, i.e., $\Pi_h^k: L^1(\Omega)^n \to \mathbb{P}^k(\mathcal{T}_h)^n$, and into $\mathbb{P}^k(\mathcal{T}_h)^{n \times d}$, i.e., $\Pi_h^k: L^1(\Omega)^{n \times d} \to$ $\mathbb{P}^k(\mathcal{T}_h)^{n \times d}.$

2.2.2. DG gradient and jump operator. For every $k \in \mathbb{N}$, we define the (global) jump lifting operator $\mathcal{R}_h^k: W^{1,p}(\mathcal{T}_h)^n \to \mathbb{P}^{k-1}(\mathcal{T}_h)^{n \times d}$ (using Riesz representation) for every $w_h \in W^{1,p}(\mathcal{T}_h)^n$ via

$$
(\mathcal{R}_h \mathbf{w}_h, \mathbf{T}_h)_{\Omega} \coloneqq \langle [\![\mathbf{w}_h \otimes \mathbf{n}]\!], \{\!\!\{\mathbf{T}_h\}\!\!\} \rangle_{\Gamma_h} \quad \text{ for all } \mathbf{T}_h \in \mathbb{P}^{k-1}(\mathcal{T}_h)^{n \times d}.
$$

Then, for every $k \in \mathbb{N}_0$, the DG gradient $\mathcal{G}_h: W^{1,p}(\mathcal{T}_h)^n \to L^p(\Omega)^{n \times d}$, for every $w_h \in W^{1,p}(\mathcal{T}_h)^n$, is defined via

$$
\mathbf{G}_h \mathbf{w}_h \coloneqq \nabla_h \mathbf{w}_h - \mathbf{R}_h \mathbf{w}_h \quad \text{ in } L^p(\Omega)^{n \times d}.
$$

Owing to [\[14,](#page-31-1) (A.26)–(A.28)], there exists a constant $c > 0$ such that $w_h \in W^{1,p}(\mathcal{T}_h)^n$, it holds that

$$
(2.12) \t c^{-1} \|w_h\|_{h,p} \leq (||{\mathcal G}_h w_h||_{p,\Omega}^p + ||h_{\Gamma}^{-1/p'}[w_h \otimes n]||_{p,\Gamma_h}^p)^{1/p} \leq c ||w_h||_{h,p}.
$$

For a generalized N-function $\psi \colon \Omega \times \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$, the pseudo-modular^{[1](#page-6-1)} $m_{\psi,h}$: $W^{1,\psi}(\mathcal{T}_h)^n \to \mathbb{R}^{\geq 0}$, for every $\boldsymbol{w}_h \in W^{1,\psi}(\mathcal{T}_h)^n$ is defined via

$$
m_{\psi,h}(\boldsymbol{w}_h) \coloneqq \sum_{F \in \Gamma_h} h_F \int_F \psi(h_F^{-1} [\![\boldsymbol{w}_h \otimes \mathbf{n}]\!]) \, \mathrm{d} s \, .
$$

For $\psi = \varphi_{p,0}$, it holds that $m_{\psi,h}(\boldsymbol{w}_h) = ||h_{\Gamma}^{-1/p'}$ $\int_{\Gamma}^{-1/p'} [\![\boldsymbol{w}_h\otimes \mathbf{n}]\!]\|_{p,\Gamma_h}^p \text{ for all } \boldsymbol{w}_h\in W^{1,\psi}(\mathcal{T}_h)^n.$

,

¹The definition of a pseudo-modular can be found in [\[32\]](#page-31-14). We extend the notion of DG Sobolev spaces to DG Sobolev-Orlicz spaces $W^{1,\psi}(\mathcal{T}_h) := \{w_h \in L^1(\Omega) \mid w_h \in W^{1,1}(K) \}$ with $\psi(\cdot, \nabla(w_h|_K)) \in L^1(K)$ for all $K \in \mathcal{T}_K$.

3. Construction of moment-preserving operators. We employ here the smoothing operator $\mathcal{E}_h \coloneqq \mathcal{E}_{h,k} : C^0(\mathcal{T}_h)^\mathbb{n} \to \mathbb{P}^{k+d}(\mathcal{T}_h)^\mathbb{n} \cap W_0^{1,1}(\Omega)^\mathbb{n}, k \in \mathbb{N}$, introduced in [\[39\]](#page-32-3). The operator, for every $w_h \in C^0(\mathcal{T}_h)^n$, is constructed as:

$$
\boldsymbol{\mathcal{E}}_h \boldsymbol{w}_h \coloneqq \boldsymbol{\mathcal{A}}_h \boldsymbol{w}_h + \boldsymbol{\mathcal{B}}_h (\boldsymbol{w}_h - \boldsymbol{\mathcal{A}}_h \boldsymbol{w}_h) \qquad \text{in } \mathbb{P}^{k+d}(\mathcal{T}_h) \cap W_0^{1,p}(\Omega)^n,
$$

where

$$
\boldsymbol{\mathcal{B}}_h \boldsymbol{w}_h \coloneqq \boldsymbol{\mathcal{B}}_h^{(1)} \boldsymbol{w}_h + \boldsymbol{\mathcal{B}}_h^{(2)} (\boldsymbol{w}_h - \boldsymbol{\mathcal{B}}_h^{(1)} \boldsymbol{w}_h) \qquad \text{in } \mathbb{P}^{k+d}(\mathcal{T}_h) \cap W_0^{1,p}(\Omega)^n,
$$

and

$$
\mathcal{A}_h \coloneqq \mathcal{A}_{h,k} : C^0(\mathcal{T}_h)^n \to \mathbb{P}^k(\mathcal{T}_h)^n \cap W_0^{1,1}(\Omega)^n,
$$

$$
\mathcal{B}_h^{(1)} \coloneqq \mathcal{B}_{h,k}^{(1)} : C^0(\mathcal{T}_h)^n \to \mathbb{P}^{k+d-1}(\mathcal{T}_h)^n \cap W_0^{1,1}(\Omega)^n,
$$

$$
\mathcal{B}_h^{(2)} \coloneqq \mathcal{B}_{h,k}^{(2)} : C^0(\mathcal{T}_h)^n \to \mathbb{P}^{k+d}(\mathcal{T}_h)^n \cap W_0^{1,1}(\Omega)^n
$$

are defined via:

• \mathcal{A}_h (simplified nodal averaging with 1 dof): Denote by $\mathcal{L}_k^{\text{int}}(\mathcal{T}_h)$ the set of interior Lagrange node on \mathcal{T}_h of degree k and for every $y \in \mathcal{L}_k^{\text{int}}(\mathcal{T}_h)$ by $\varphi_y^k \in \mathbb{P}^k(\mathcal{T}_h)$, the corresponding basis function. In addition, for every $y \in \mathcal{L}_{k}^{\text{int}}(\mathcal{T}_{h})$ fix an arbitrary $K_{y} \in \mathcal{T}_{h}$ such that $y \in K_{y}$. Then, the operator $\mathcal{A}_h: C^0(\mathcal{T}_h)^n \to \mathbb{P}^k(\mathcal{T}_h) \cap W_0^{1,1}(\Omega)^n$, for every $w_h \in C^0(\mathcal{T}_h)^n$, is defined via

$$
\boldsymbol{\mathcal{A}}_h \boldsymbol{w}_h \coloneqq \sum_{y\in \mathcal{L}_k^{\mathrm{int}}(\mathcal{T}_h)} (\boldsymbol{w}_h|_{K_y})(y) \varphi_y^k \quad \text{ in } \mathbb{P}^k(\mathcal{T}_h)^n \cap W^{1,1}_0(\Omega)^n\,.
$$

 \bullet $\mathcal{B}_h^{(1)}$ $h_h^{(1)}$ (facet bubble smoother): Denote for $F \in \Gamma_h$ by $\mathcal{L}_{k-1}(F)$ the set of Lagrange nodes on F of degree $k-1$ and by $\varphi_F := \varphi_{y_1}^1 \cdot \ldots \cdot \varphi_{y_d}^1 \in \mathbb{P}^d(F) \cap$ $W_0^{1,1}(F)$, where $y_1, \ldots, y_d \in \mathcal{L}_1(F)$ are such that $F = \text{conv}\{y_1, \ldots, y_d\}$, the corresponding facet bubble function. Then, the operator $\mathcal{B}_h^{(1)}$ $L^{(1)}$: $C^0(\mathcal{T}_h)^n \rightarrow$ $\mathbb{P}^{k+d-1}(\mathcal{T}_h)^n \cap W_0^{1,1}(\Omega)^n$, for every $w_h \in C^0(\mathcal{T}_h)^n$, is defined via

$$
\boldsymbol{\mathcal{B}}_h^{(1)} \boldsymbol{w}_h \coloneqq \sum_{F \in \Gamma_h^i} \sum_{y \in \mathcal{L}_{k-1}(F)} (\boldsymbol{\mathcal{Q}}_F \{\!\!\{ \boldsymbol{w}_h \}\!\!\})(y) \, \varphi_y^{k-1} \, \varphi_F \quad \text{ in } \mathbb{P}^{k+d-1}(\mathcal{T}_h)^n \cap W_0^{1,1}(\Omega)^n \, ,
$$

where $\mathcal{Q}_F: C^0(\mathcal{T}_h)^n \to \mathbb{P}^{k-1}(F)^n$ is the weighted L^2 -projection, for every $w_h \in C^0(\mathcal{T}_h)^n$ and $z_h \in \mathbb{P}^{k-1}(F)^n$, defined via

$$
(\varphi_F\mathcal{Q}_F\bm{w}_h,\bm{z}_h)_F=(\bm{v}_h,\bm{z}_h)_F\,.
$$

 \bullet $\mathcal{B}_h^{(2)}$ ⁽²⁾ (interior bubble smoother): Denote for $K \in \mathcal{T}_h$ by $\mathcal{L}_k(K)$ the set of Lagrange nodes on K of degree k and by $\varphi_K := \varphi_{y_1}^1 \cdot \ldots \cdot \varphi_{y_{d+1}}^1 \in \mathbb{P}^{d+1}(K) \cap$ $W_0^{1,1}(K)$, where $y_1, \ldots, y_{d+1} \in \mathcal{L}_1(K)$ are such that $K = \text{conv}\{y_1, \ldots, y_{d+1}\},$ the corresponding bubble function. Then, the operator $\mathcal{B}_h^{(2)}$ $L^{(2)}$: $C^0(\mathcal{T}_h)^n \rightarrow$ $\mathbb{P}^{k+d}(\mathcal{T}_h)^n \cap W_0^{1,1}(\Omega)^n$, for every $\mathbf{w}_h \in C^0(\mathcal{T}_h)^n$, is defined via

$$
\boldsymbol{\mathcal{B}}_h^{(2)} \boldsymbol{w}_h \coloneqq \sum_{K\in\mathcal{T}_h} (\boldsymbol{\mathcal{Q}}_K \boldsymbol{w}_h)\, \varphi_K\,,
$$

where $\mathcal{Q}_K: C^0(\mathcal{T}_h)^n \to \mathbb{P}^{k-1}(K)^n$ is the weighted L^2 -projection, for every $w_h \in C^0(\mathcal{T}_h)^n$ and $z_h \in \mathbb{P}^{k-1}(K)^n$, defined via

$$
(\varphi_K \mathcal{Q}_K \mathcal{w}_h, \mathcal{z}_h)_K = (\mathcal{w}_h, \mathcal{z}_h)_K.
$$

Note that $\mathcal{B}_h^{(2)}$ $h_h^{(2)}$ is not needed when $k = 1$.

The idea in all these cases is that the bubble smoother helps with the preservation of moments, but as the bubble smoothers are not stable, a nodal averaging operator is added to recover stability. Namely, [\[39,](#page-32-3) (3.17), (3.18)] and $\|\[\mathbf{v}_h\] = \|\[\mathbf{v}_h \otimes \mathbf{n}\]$, for every $\boldsymbol{v}_h \in \mathbb{P}^k(\mathcal{T}_h)^n$, imply that

(3.1)
$$
\|\nabla_h(\boldsymbol{v}_h-\boldsymbol{\mathcal{E}}_h\boldsymbol{v}_h)\|_{2,K}\lesssim \sum_{F\in\Gamma_h(K)}\|h_{\Gamma}^{-1/2}[\![\boldsymbol{v}_h\otimes\boldsymbol{n}]\!]_F\|_{2,F},
$$

where $\Gamma_h(K) \coloneqq \{F \in \Gamma_h \mid K \cap F \neq \emptyset\}$ and the constants depend only on k and ω_0 . We now proceed to generalise the estimate (3.1) to the Orlicz setting.

PROPOSITION 3.1 (Interpolation estimate). Let $\psi \colon \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ be an N-function with $\psi \in \Delta_2$, $\psi^* \in \Delta_2$, and let $k \in \mathbb{N}$. Then, for every $\mathbf{v}_h \in \mathbb{P}^k(K)^n$ and $K \in \mathcal{T}_h$, it holds that

$$
\frac{1}{|K|}\int_K \psi(h_K|\tilde{\nabla}_h(\boldsymbol{v}_h-\boldsymbol{\mathcal{E}}_h\boldsymbol{v}_h)|)\,\mathrm{d} x\lesssim \sum_{F\in\Gamma_h(K)}\frac{1}{|F|}\int_F \psi(\|\boldsymbol{v}_h\otimes\boldsymbol{n}\|_F|\,\mathrm{d} s\,,
$$

where either $\tilde{\nabla}_h = \nabla_h$ or $\tilde{\nabla}_h = \mathcal{G}_h$ and the constants depend only on k, $\Delta_2(\psi), \Delta_2(\psi^*)$ and ω_0 .

Proof. We need to distinguish the cases $\tilde{\nabla}_h = \nabla_h$ and $\tilde{\nabla}_h = \mathcal{G}_h$: Case $\tilde{\nabla}_h = \nabla_h$. Owing to [\(3.1\)](#page-8-0) together with [\[17,](#page-31-0) Lem. 12.1], we have that

(3.2)
$$
h_K \|\nabla (\boldsymbol{v}_h - \boldsymbol{\mathcal{E}}_h \boldsymbol{v}_h)\|_{\infty,K} \lesssim \sum_{F \in \Gamma_h(K)} \frac{1}{|F|} \int_F \|\boldsymbol{v}_h \otimes \boldsymbol{n}\|_F |\,ds\,,
$$

where the constants depends only on k and ω_0 . Using [\(3.2\),](#page-8-1) the Δ_2 -condition and convexity of ψ , in particular, Jensen's inequality, and that $\sup_{h>0} \sup_{K \in \mathcal{T}_h} \text{card}(\Gamma_h(K))$ $\leq c$, where $c > 0$ depends only on ω_0 , we find that

$$
\frac{1}{|K|} \int_{K} \psi(h_{K}|\nabla(\boldsymbol{v}_{h} - \mathcal{E}_{h}\boldsymbol{v}_{h})|) d\boldsymbol{x} \lesssim \psi\bigg(\frac{1}{\text{card}(\Gamma_{h}(K))} \sum_{F \in \Gamma_{h}(K)} \frac{1}{|F|} \int_{F} |\llbracket \boldsymbol{v}_{h} \otimes \boldsymbol{n} \rrbracket_{F}| d\boldsymbol{s}\bigg)
$$
\n
$$
\lesssim \frac{1}{\text{card}(\Gamma_{h}(K))} \sum_{F \in \Gamma_{h}(K)} \frac{1}{|F|} \int_{F} \psi(||\llbracket \boldsymbol{v}_{h} \otimes \boldsymbol{n} \rrbracket_{F}|) d\boldsymbol{s}
$$
\n
$$
\lesssim \sum_{F \in \Gamma_{h}(K)} \frac{1}{|F|} \int_{F} \psi(||\llbracket \boldsymbol{v}_{h} \otimes \boldsymbol{n} \rrbracket_{F}|) d\boldsymbol{s},
$$

where the constants depend only on k, $\Delta_2(\psi), \Delta_2(\psi^*)$ and ω_0 .

Case $\tilde{\nabla}_h = \mathcal{G}_h$. Appealing to [\[14,](#page-31-1) (A.23)], for every $F \in \Gamma_h$ and $K \in \mathcal{T}_h$ with $F \subseteq \partial K$, it holds that

(3.4)
$$
\frac{1}{|K|}\int_K \psi(h_K|\mathcal{R}_h\mathbf{v}_h|)\,\mathrm{d}x \lesssim \frac{1}{|F|}\int_F \psi(\|\mathbf{v}_h\otimes\mathbf{n}\|_F|\,\mathrm{d}s\,,
$$

where the constants depend only on k, $\Delta_2(\psi), \Delta_2(\psi^*)$ and ω_0 . Thus, using [\(3.3\),](#page-8-2) that

 $\mathcal{R}_h \mathcal{E}_h v_h = 0$, and [\(3.4\),](#page-8-3) we obtain

$$
\frac{1}{|K|} \int_{K} \psi(h_K | \tilde{\nabla}_h (\mathbf{v}_h - \mathbf{\mathcal{E}}_h \mathbf{v}_h) |) \, dx \lesssim \frac{1}{|K|} \int_{K} \psi(h_K | \nabla (\mathbf{v}_h - \mathbf{\mathcal{E}}_h \mathbf{v}_h) |) \, dx \n+ \frac{1}{|K|} \int_{K} \psi(h_K | \mathbf{\mathcal{R}}_h \mathbf{v}_h) \, dx \n\lesssim \sum_{F \in \Gamma_h(K)} \frac{1}{|F|} \int_{F} \psi(||[\mathbf{v}_h \otimes \mathbf{n}]_F|) \, ds,
$$

where the constants depend only on k, $\Delta_2(\psi), \Delta_2(\psi^*)$ and ω_0 .

4. Primal formulation. In this section, we examine the primal formulation of the non-linear problem (1.1) and its discretization. In doing so, throughout the entire section, we assume that S has (p, δ) -structure in the sense of [Definition 2.1.](#page-3-0)

4.1. Continuous primal formulation. We abbreviate

$$
V \coloneqq W_0^{1,p}(\Omega)^n.
$$

Then, for given right-hand side $f \in V^*$, the *primal formulation* of [\(1.1\)](#page-0-0) seeks for a function (or vector field, respectively) $u \in V$ such that for every $z \in V$, it holds that

(4.1)
$$
(\mathcal{S}(\nabla u), \nabla z)_{\Omega} = \langle f, z \rangle_V.
$$

Resorting to the celebrated theory of monotone operators, cf. [\[42,](#page-32-6) [34\]](#page-31-15), it is readily seen that the primal formulation admits a unique solution.

4.2. Discrete primal formulation. For given $k \in \mathbb{N}$, we abbreviate

$$
V_h\coloneqq V_h^k\coloneqq \mathbb{P}^k(\mathcal{T}_h)^n
$$

.

 \Box

Then, for given right-hand side $f \in V^*$, the *discrete primal formulation* of [\(1.1\)](#page-0-0) seeks for a discrete function (or vector field, respectively) $u_h \in V_h$ such that for every $z_h \in V_h$, it holds that

$$
(4.2) \quad (\mathcal{S}(\tilde{\nabla}_h \boldsymbol{u}_h), \nabla \mathcal{E}_h \boldsymbol{z}_h)_{\Omega} + \alpha \langle \mathcal{S}_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)}(h_{\Gamma}^{-1}[\![\boldsymbol{u}_h \otimes \boldsymbol{n}]\!]), [\![\boldsymbol{z}_h \otimes \boldsymbol{n}]\!])_{\Gamma_h} = \langle \boldsymbol{f}, \boldsymbol{\mathcal{E}}_h \boldsymbol{z}_h \rangle_V,
$$

where $\alpha > 0$ is the *stabilisation parameter*, $\beta_h : V_h \to \mathbb{R}^{\geq 0}$ is the *max-shift func*tional, for every $v_h \in V_h$, defined via

(4.3)
$$
\beta_h(\boldsymbol{v}_h) \coloneqq \begin{cases} 0 & \text{if } p \leq 2, \\ \|\tilde{\nabla}_h \boldsymbol{v}_h\|_{\infty,\Omega} & \text{if } p > 2, \end{cases}
$$

with $\tilde{\nabla}_h \in {\mathcal{G}_h, \nabla_h}$, and $\mathcal{S}_{\beta_h(u_h)}$ is defined in [\(2.6\).](#page-4-3) More precisely, setting $\tilde{\nabla}_h = \mathcal{G}_h$, results in an LDG-type method and setting $\tilde{\nabla}_h = \nabla_h$, results is an IIDG-type method.

REMARK 4.1. Note that for each choice $\tilde{\nabla}_h \in {\{\mathcal{G}_h, \nabla_h\}}$ and for every $\mathbf{w}_h \in$ $C^0(\mathcal{T}_h)^n$, we have that $\tilde{\nabla}_h \mathcal{E}_h \mathcal{w}_h = \nabla \mathcal{E}_h \mathcal{w}_h$. This enables us to replace the classical gradient in the primal formulation [\(4.2\)](#page-9-1), by $\tilde{\nabla}_h \in {\cal G}_h, \nabla_h$, which is of crucial importance in the proof of a best-approximation type result in [Theorem](#page-11-0) 4.6; see also [Lemma](#page-10-0) 4.4.

REMARK 4.2. It can be seen that if we add to the left-hand side of the LDG discretisation the term $(\mathcal{S}(\mathcal{R}_h\boldsymbol{u}_h), \nabla_h(\mathcal{E}_h\boldsymbol{z}_h - \boldsymbol{z}_h))_{\Omega}$, then we arrive at a SIP-like method (in that it reduces to the usual Symmetric Interior Penalty method in the linear case). Thus, all results proved for the LDG method and the IIDG method can be carried over to this method.

REMARK 4.3. The max-shift functional is constructed in such a way that is has the following two properties. First, for the globally non-constant shift $|\tilde{\nabla}_h\tilde{\bm u}_h|\in L^p(\Omega)^{n\times d}$ and the globally constant shift $\beta_h(\boldsymbol{u}_h) \in \mathbb{R}$, it holds that $\varphi_{|\tilde{\nabla}_h \boldsymbol{u}_h|}(t) \leq \varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)}(t)$ for all $t \geq 0$. To the latter N-function, the approximation properties of the smoothing operator $\mathcal{E}_h : V_h \to V$ apply, cf. [\(4.5\)](#page-10-1) (below). Second, for the conjugate N-functions, it holds that $(\varphi_{\beta_h(u_h)})^*(t) \leq (\varphi_{|\tilde{\nabla}_h u_h|})^*(t)$ for all $t \geq 0$. For the latter N-function, the equivalences (2.8) apply, cf. (4.6) (below).

More precisely, the max-shift functional is precisely constructed in such a way that the following lemma applies.

LEMMA 4.4. For every $\kappa > 0$, there exists constant $c_{\kappa} > 0$, depending on the characteristics of S, such that for every $v_h, z_h \in V_h$ and $v \in V$, it holds that

$$
\begin{aligned} & |(\boldsymbol{\mathcal{S}}(\tilde \nabla_h \boldsymbol{v}_h) - \boldsymbol{\mathcal{S}}(\nabla \boldsymbol{v}), \tilde \nabla_h (\boldsymbol{\mathcal{E}}_h \boldsymbol{z}_h - \boldsymbol{z}_h))_{\Omega}| \\ & \lesssim \kappa \, \| \boldsymbol{\mathcal{F}}(\tilde \nabla_h \boldsymbol{v}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{v}) \|_{2,\Omega}^2 + c_{\kappa} \, m_{\varphi_{\boldsymbol{\beta}_h (\boldsymbol{v}_h)},h}(\boldsymbol{z}_h) \,, \end{aligned}
$$

where the constants depend only on k, ω_0 , and the characteristics of S.

Proof. Using [\(2.9\)](#page-4-5) and the ε -Young inequality [\(2.2\)](#page-3-2) with $\psi = \varphi_{\beta_h(\mathbf{v}_h)}$, we find that

$$
(4.4) \quad |\langle \mathcal{S}(\tilde{\nabla}_h \mathbf{v}_h) - \mathcal{S}(\nabla \mathbf{v}), \tilde{\nabla}_h (\mathcal{E}_h z_h - z_h) \rangle_{\Omega}| \n\lesssim c_{\kappa} \rho_{\varphi_{\beta_h(\mathbf{v}_h)},\Omega}(\tilde{\nabla}_h (\mathcal{E}_h z_h - z_h)) + \kappa \rho_{(\varphi_{\beta_h(\mathbf{v}_h)})^*,\Omega}(\varphi'_{|\tilde{\nabla}_h \mathbf{v}_h}(|\tilde{\nabla}_h \mathbf{v}_h - \nabla \mathbf{v}|)).
$$

Due to $\beta_h(u_h) \in \mathbb{R}$, [Proposition 3.1](#page-8-4) is applicable and yields, also using $h_F \sim h_K$, that

(4.5)
$$
\rho_{\varphi_{\beta_h(\mathbf{v}_h)},\Omega}(\tilde{\nabla}_h(\boldsymbol{\mathcal{E}}_h|\boldsymbol{z}_h-\boldsymbol{z}_h))\lesssim m_{\varphi_{\beta_h(\mathbf{u}_h)},h}(\boldsymbol{z}_h).
$$

By definition of the max-shift functional and [Remark 2.2](#page-4-6) (iii), we obtain $(\varphi_{\beta_h(\mathbf{v}_h)})^* \leq$ $(\varphi_{|\tilde{\nabla}_h v_h|})^*$. Using this, the equivalence (2.1) for $\psi = \varphi_{|\tilde{\nabla}_h v_h|}$, and (2.8) , we obtain

$$
(4.6) \quad \rho_{(\varphi_{\beta_h(v_h)})^*,\Omega}(\varphi'_{|\tilde{\nabla}_h v_h|}(|\tilde{\nabla}_h v_h - \nabla v|)) \leq \rho_{(\varphi_{|\tilde{\nabla}_h v_h|})^*,\Omega}(\varphi'_{|\tilde{\nabla}_h v_h|}(|\tilde{\nabla}_h v_h - \nabla v|)) \\ \lesssim \|\mathcal{F}(\tilde{\nabla}_h v_h) - \mathcal{F}(\nabla v)\|_{2,\Omega}^2.
$$

Eventually, combining [\(4.5\)](#page-10-1) and [\(4.6\)](#page-10-2) in [\(4.4\),](#page-10-3) we conclude the claimed estimate. \Box

In the linear case $\mathcal{S}(\tilde{\nabla}_h \boldsymbol{u}) = \tilde{\nabla}_h \boldsymbol{u}$, non-degeneracy and existence of discrete solutions is a direct consequence of the construction of \mathcal{E}_h , because then the preserva-tion property [\(1.6\)](#page-2-1) means that we can forget the operator \mathcal{E}_h on the left-hand side and so the usual argument for proving coercivity works (note that, by construction, $\tilde{\nabla}_h \mathbf{u}_h \in \mathbb{P}^{k-1}(\mathcal{T}_h)^{n \times d}$. In the non-linear case this argument only works for the lowest order DG ansatz for which the gradients are element-wise constant. Hence, in general, we would need to check that discrete solutions exist.

PROPOSITION 4.5. For $\alpha > 0$ sufficiently large, the primal formulation [\(4.2\)](#page-9-1) admits a solution.

Proof. We equip V_h with the norm $\|\cdot\|_{V_h} := \|\cdot\|_{h,p}$, and consider the operator $\mathcal{T}_h: V_h \to (V_h)^*$, for every $\mathbf{v}_h \in V_h$ and $\mathbf{z}_h \in V_h$, defined via

$$
\langle \boldsymbol{\mathcal{T}}_h \boldsymbol{v}_h, \boldsymbol{z}_h \rangle_{V_h} \coloneqq (\boldsymbol{\mathcal{S}}(\tilde \nabla_h \boldsymbol{v}_h), \nabla \boldsymbol{\mathcal{E}}_h \boldsymbol{z}_h)_{\Omega} + \alpha \langle \boldsymbol{\mathcal{S}}_{\boldsymbol{\beta}_h(\boldsymbol{v}_h)}(h_\Gamma^{-1} [\![\boldsymbol{v}_h \otimes \boldsymbol{n}]\!]), [\![\boldsymbol{z}_h \otimes \boldsymbol{n}]\!])_{\Gamma_h} \,.
$$

Since V_h is finite dimensional, consists of broken polynomials, [\[5,](#page-30-8) Lem. 3.18], the properties of S and \mathcal{S}_a , $a \geq 0$, imply that the operator $\mathcal{T}_h: V_h \to (V_h)^*$, for every fixed $h > 0$, is well-defined, continuous, and monotone. To prove the boundedness of $\mathcal{T}_h: V_h \to (V_h)^*$, we use the properties of $\mathcal{S}, \mathcal{S}_{\beta_h(v_h)}$ (cf. [Proposition 2.3](#page-4-4) and [Re](#page-4-7)[mark 2.4\)](#page-4-7), Young's inequality, [\(2.1\),](#page-3-3) [Proposition 3.1](#page-8-4) with $\psi = \varphi$, the stability of \mathcal{R}_h (cf. [\[14,](#page-31-1) (A.25)]), a shift change (cf. [Lemma 2.5\)](#page-5-0), $\sum_{F \in \Gamma_h} h_F \mathcal{H}^{d-1}(F) \sim |\Omega|$, [\[17,](#page-31-0) Lemma 12.1, Jensen's inequality, and $\varphi(t) + t^p \sim t^p + \delta^p$ for all $t \geq 0$ (cf. [\[5\]](#page-30-8)), to find that

$$
\begin{aligned} |\langle \boldsymbol{\mathcal{T}}_h \boldsymbol{v}_h, \boldsymbol{z}_h \rangle_{V_h}| \lesssim & \rho_{\varphi,\Omega}(\tilde{\nabla}_h \boldsymbol{v}_h) + \rho_{\varphi,\Omega}(\tilde{\nabla}_h (\boldsymbol{\mathcal{E}}_h \boldsymbol{z}_h - \boldsymbol{z}_h)) + \rho_{\varphi,\Omega}(\tilde{\nabla}_h \boldsymbol{z}_h) \\ & + \alpha \, m_{\varphi_{\boldsymbol{\beta}_h (\boldsymbol{v}_h)},h}(\boldsymbol{v}_h) + \alpha \, m_{\varphi_{\boldsymbol{\beta}_h (\boldsymbol{v}_h)},h}(\boldsymbol{z}_h) \\ & \lesssim \rho_{\varphi,\Omega}(\nabla_h \boldsymbol{v}_h) + \rho_{\varphi,\Omega}(\nabla_h \boldsymbol{z}_h) + m_{\varphi,h}(\boldsymbol{v}_h) + m_{\varphi,h}(\boldsymbol{z}_h) + \varphi(\boldsymbol{\beta}_h (\boldsymbol{v}_h)) \\ & \lesssim \| \boldsymbol{v}_h \|_{p,h}^p + \| \boldsymbol{z}_h \|_{p,h}^p + \delta^p \,. \end{aligned}
$$

Thus, $\mathcal{T}_h: V_h \to (V_h)^*$ is pseudo-monotone. If we can prove its coercivity, then, from pseudo-monotone operator theory (cf. [\[42,](#page-32-6) Thm. 27.A]), it follows that $\mathcal{T}_h: V_h \to$ $(V_h)^*$ is surjective. For every $v_h \in V_h$, using first that $\nabla \mathcal{E}_h v_h = \tilde{\nabla}_h \mathcal{E}_h v_h$, then that $\mathcal{S}(\mathsf{Q}) : \mathsf{Q} \sim \varphi(|\mathsf{Q}|) \text{ and } \mathcal{S}_{\beta_h(\bm{v}_h)}(\mathsf{Q}) : \mathsf{Q} \sim \varphi_{\beta_h(\bm{v}_h)}(|\mathsf{Q}|) \text{ for all } \mathsf{Q} \in \mathbb{R}^{n \times d} \text{ (cf. Proposition 1.1)}$ [tion 2.3](#page-4-4) and [Remark 2.4\)](#page-4-7), $\varphi_{\beta_h(\mathbf{v}_h)} \geq \varphi$ (cf. [Remark 2.2](#page-4-6) (iii)), the ε -Young inequality [\(2.2\)](#page-3-2) with $\psi = \varphi$, and [Proposition 3.1](#page-8-4) with $\psi = \varphi$, we find that

$$
\begin{aligned}\n\langle \boldsymbol{\mathcal{T}}_h \boldsymbol{v}_h, \boldsymbol{v}_h \rangle_{V_h} &= (\boldsymbol{\mathcal{S}}(\tilde{\nabla}_h \boldsymbol{v}_h), \tilde{\nabla}_h \boldsymbol{v}_h)_{\Omega} + \alpha \langle \boldsymbol{\mathcal{S}}_{\boldsymbol{\beta}_h(\boldsymbol{v}_h)}(h_\Gamma^{-1}[\![\boldsymbol{v}_h \otimes \boldsymbol{n}]\!]), [\![\boldsymbol{v}_h \otimes \boldsymbol{n}]\!])_{\Gamma_h} \\
&+ (\boldsymbol{\mathcal{S}}(\tilde{\nabla}_h \boldsymbol{v}_h), \tilde{\nabla}_h (\boldsymbol{\mathcal{E}}_h \boldsymbol{v}_h - \boldsymbol{v}_h))_{\Omega} \\
&\geq (c-\varepsilon) \, \rho_{\varphi,\Omega}(\tilde{\nabla}_h \boldsymbol{v}_h) + \alpha \, c \, m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{v}_h)},h}(\boldsymbol{v}_h) - c_\varepsilon \, \rho_{\varphi,\Omega}(\tilde{\nabla}_h (\boldsymbol{\mathcal{E}}_h \boldsymbol{v}_h - \boldsymbol{v}_h)) \\
&\geq (c-\varepsilon) \, \rho_{\varphi,\Omega}(\tilde{\nabla}_h \boldsymbol{v}_h) + (\alpha \, c - c_\varepsilon) \, m_{\varphi,h}(\boldsymbol{v}_h)\, .\n\end{aligned}
$$

Consequently, choosing first $\varepsilon > 0$ sufficiently small and, subsequently, $\alpha > 0$ sufficiently large, also using $\varphi(t) + t^p \sim t^p + \delta^p$ for all $t \geq 0$ (cf. [\[5\]](#page-30-8)), $h \mathcal{H}^{d-1}(\Gamma_h) \sim |\Omega|$, for every $v_h \in V_h$, and the norm equivalence [\(2.12\),](#page-6-2) we arrive at

$$
\begin{aligned}\n\langle \boldsymbol{\mathcal{T}}_h \boldsymbol{v}_h, \boldsymbol{v}_h \rangle_{V_h} &\geq \|\tilde{\nabla}_h \boldsymbol{v}_h\|_{p,\Omega}^p + \alpha \|h_{\Gamma}^{-1/p'}[\![\boldsymbol{v}_h\otimes\boldsymbol{n}]\!]\|_{p,\Gamma_h}^p - \delta^p \, |\Omega|\,(1+\alpha) \\
&\geq \min\{1,\alpha\}\, \|\boldsymbol{v}_h\|_{V_h}^p - \delta^p \, |\Omega| \, \max\{1,\alpha\}\,. \n\end{aligned}
$$

Putting everything together, we proved that the operator $\mathcal{T}_h: V_h \to (V_h)^*$ is welldefined, bounded, pseudo-monotone, and coercive and, thus, surjective (cf. [\[42,](#page-32-6) Thm. 27.A]). П

From [\(4.1\)](#page-9-2) and [\(4.2\),](#page-9-1) it follows that the error equation, for every $z_h \in V_h$, takes the form

$$
(4.7) \quad (\mathcal{S}(\tilde{\nabla}_h \boldsymbol{u}_h) - \mathcal{S}(\nabla \boldsymbol{u}), \nabla \mathcal{E}_h \boldsymbol{z}_h)_{\Omega} + \alpha \langle \mathcal{S}_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)}(h_{\Gamma}^{-1}[\![\boldsymbol{u}_h \otimes \boldsymbol{n}]\!]), [\![\boldsymbol{z}_h \otimes \boldsymbol{n}]\!])_{\Gamma_h} = 0.
$$

THEOREM 4.6. Let $u \in V$ be a solution of (4.1) and $u_h \in V_h$ ba a solution of [\(4.2\)](#page-9-1). If $\alpha > 0$ sufficiently large, we have that

$$
\begin{aligned}&\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\\&\lesssim\inf_{\boldsymbol{v}_h\in V_h}\left(\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{v}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{v}_h)\right)\end{aligned}
$$

,

where the constant depends only on k, ω_0 , and the characteristics of S.

Proof. Adding and substracting $\tilde{\nabla}_h z_h \in \mathbb{P}^{k-1}(\mathcal{T}_h)^{n \times d}$ in [\(4.7\)](#page-11-1) for arbitrary $z_h \in$ V_h , using that $\tilde{\nabla}_h \tilde{\boldsymbol{\mathcal{E}}}_h \tilde{\boldsymbol{\mathcal{z}}}_h = \nabla \boldsymbol{\mathcal{E}}_h \tilde{\boldsymbol{\mathcal{z}}}_h$, we get

(4.8)
\n
$$
0 = (\mathcal{S}(\tilde{\nabla}_h \mathbf{u}_h) - \mathcal{S}(\nabla \mathbf{u}), \tilde{\nabla}_h z_h)_{\Omega}
$$
\n
$$
+ \alpha \langle \mathcal{S}_{\beta_h(\mathbf{u}_h)}(h_{\Gamma}^{-1}[\![\mathbf{u}_h \otimes \mathbf{n}]\!]) , [\![z_h \otimes \mathbf{n}]\!] \rangle_{\Gamma_h}
$$
\n
$$
+ (\mathcal{S}(\tilde{\nabla}_h \mathbf{u}_h) - \mathcal{S}(\nabla \mathbf{u}), \tilde{\nabla}_h (\mathcal{E}_h z_h - z_h))_{\Omega}
$$
\n
$$
=: \mathfrak{I}_1 + \mathfrak{I}_2 + \mathfrak{I}_3.
$$

Next, we choose $z_h = u_h - v_h \in V_h$, where $v_h \in V_h$ is arbitrary, and estimate $\mathfrak{I}_1, \mathfrak{I}_2$, \mathfrak{I}_3 :

ad \mathfrak{I}_1 . Using [\(2.8\),](#page-4-1) and [\(2.9\),](#page-4-5) the ε -Young inequality [\(2.2\)](#page-3-2) with $\psi = \varphi_{|\nabla u|}$, we find that

$$
(4.9) \quad\n\begin{aligned}\n\mathfrak{I}_1 &= (\mathcal{S}(\tilde{\nabla}_h \boldsymbol{u}_h) - \mathcal{S}(\nabla \boldsymbol{u}), \tilde{\nabla}_h \boldsymbol{u}_h - \nabla \boldsymbol{u} + \nabla \boldsymbol{u} - \tilde{\nabla}_h \boldsymbol{v}_h)_{\Omega} \\
&\geq c \, \|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 - |(\mathcal{S}(\tilde{\nabla}_h \boldsymbol{u}_h) - \mathcal{S}(\nabla \boldsymbol{u}), \nabla \boldsymbol{u} - \tilde{\nabla}_h \boldsymbol{v}_h)_{\Omega}| \\
&\geq (c - \varepsilon) \, \|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 - c_{\varepsilon} \, \|\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u}) - \boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{v}_h)\|_{2,\Omega}^2.\n\end{aligned}
$$

ad \mathfrak{I}_2 . Using that $\mathcal{S}_{\beta_h(u_h)}(Q)$: $Q \sim \varphi_{\beta_h(u_h)}(Q)$ uniformly in $Q \in \mathbb{R}^{n \times d}$ (cf. [Proposition 2.3](#page-4-4) and [Remark 2.4\)](#page-4-7) and the ε -Young inequality [\(2.2\)](#page-3-2) with $\psi = \varphi_{|\beta_h(u_h)|}$, we obtain

$$
\mathfrak{I}_2 = \alpha \langle \mathcal{S}_{\beta_h(u_h)}(h_{\Gamma}^{-1}[\![u_h \otimes n]\!]) , [\![u_h \otimes n]\!] - [\![v_h \otimes n]\!] \rangle_{\Gamma_h}
$$
\n
$$
(4.10) \qquad \geq \alpha \, cm_{\varphi_{\beta_h(u_h)},h}(u_h) - \alpha \langle h_{\Gamma}(\varphi_{\beta_h(u_h)})'(h_{\Gamma}^{-1}[\![u_h \otimes n]\!]), h_{\Gamma}^{-1}[\![v_h \otimes n]\!] \rangle_{\Gamma_h}
$$
\n
$$
\geq \alpha \, (c - \varepsilon) \, m_{\varphi_{\beta_h(u_h)},h}(u_h) - \alpha \, c_{\varepsilon} \, m_{\varphi_{\beta_h(u_h)},h}(v_h).
$$

 $ad\,3_3$. Using [Lemma 4.4,](#page-10-0) we find that

$$
(4.11) \quad \begin{aligned} \mathfrak{I}_3 &\leq \kappa \|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + c_{\kappa} \, m_{\varphi_{\beta_h(\boldsymbol{u}_h)},h}(\boldsymbol{z}_h) \\ &\lesssim \kappa \, \|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + c_{\kappa} \, (m_{\varphi_{\beta_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h) + m_{\varphi_{\beta_h(\boldsymbol{u}_h)},h}(\boldsymbol{v}_h)). \end{aligned}
$$

Combining [\(4.9\)–](#page-12-0)[\(4.11\),](#page-12-1) for $\varepsilon, \kappa > 0$ sufficiently small and $\alpha > 0$ sufficiently large, we arrive at the claimed best-approximation result. \Box

As a first immediate consequence of the best-approximation result in [Theorem 4.6,](#page-11-0) we obtain the convergence of the method under minimal regularity assumptions, i.e., merely $u \in V$ and $f \in V^*$.

COROLLARY 4.7 (Convergence). For $\alpha > 0$ sufficiently large, it holds that

$$
\|\boldsymbol{F}(\tilde{\nabla}_h \boldsymbol{u}_h)-\boldsymbol{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\to 0 \quad (h\to 0).
$$

Proof. Let Π_h^{SZ} : $V \to V_h \cap V$ denote the Scott–Zhang quasi-interpolation operator (cf. [\[36\]](#page-32-7)). Then, due to $[\Pi_h^{SZ} \mathbf{u} \otimes \mathbf{n}] = \mathbf{0}$ a.e. on Γ_h , we have that

(4.12)
$$
m_{\varphi_{\beta_h(u_h)},h}(\Pi_h^{SZ}u) = 0.
$$

Therefore, using the stability and convergence properties of Π_{h}^{SZ} : $V \to V_h \cap V$ (cf. [\[36\]](#page-32-7)) and the density of smooth functions in V, we conclude that $\prod_{h}^{SZ} u \to u$ in $V(h \to 0)$. This, together with [\(2.8\),](#page-4-1) [Remark 2.2](#page-4-6) (iii), Hölder's inequality implies that

$$
(4.13) \qquad \|\mathcal{F}(\nabla \mathbf{u}) - \mathcal{F}(\nabla \Pi_h^{SZ} \mathbf{u})\|_{2,\Omega}^2 \lesssim \|\nabla \mathbf{u} - \nabla \Pi_h^{SZ} \mathbf{u}\|_p^{\min\{p,2\}} \to 0 \quad (h \to 0)
$$

with a constant depending possibly on δ , $\|\nabla u\|_p$. Putting everything together, choosing $\mathbf{v}_h = \prod_h^{SZ} \mathbf{u} \in V_h$ in [Theorem 4.6,](#page-11-0) using [\(4.12\)](#page-12-2) and [\(4.13\),](#page-12-3) we conclude that

$$
\|\boldsymbol{F}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h) \n\lesssim \|\boldsymbol{F}(\nabla \boldsymbol{u}) - \boldsymbol{F}(\nabla \Pi_h^{SZ} \boldsymbol{u})\|_{2,\Omega}^2 \to 0 \quad (h \to 0),
$$

which is the claimed convergence under minimal regularity assumptions.

As a second immediate consequence of the best-approximation result in [Theo](#page-11-0)[rem 4.6,](#page-11-0) we obtain fractional convergence rates of the method given fractional regularity assumptions on the solution of the continuous primal problem [\(4.1\).](#page-9-2) In order to express the fractional regularity of the solution of (4.1) , we make use of Nikolski $\check{\mathbf{u}}$ spaces. For given $p \in [1,\infty)$, $\beta \in (0,1]$, and $v \in L^p(\Omega)$, the Nikolski is semi-norm is defined via

$$
[v]_{N^{\beta,p}(\Omega)} \coloneqq \sup_{h \in \mathbb{R}^d \setminus \{0\}} |h|^{-\beta} \bigg(\int_{\Omega \cap (\Omega - h)} |v(x+h) - v(x)|^p \,\mathrm{d}x \bigg)^{\frac{1}{p}} < \infty \,.
$$

Then, for $p \in [1, \infty)$ and $\beta \in (0, 1]$, the *Nikolskiĭ* space is defined via

$$
N^{\beta,p}(\Omega) \coloneqq \left\{ v \in L^p(\Omega) \mid [v]_{N^{\beta,p}(\Omega)} < \infty \right\},\,
$$

and the *Nikolskiĭ* norm $\|\cdot\|_{N^{\beta,p}(\Omega)} := \|\cdot\|_{p,\Omega} + [v]_{N^{\beta,p}(\Omega)}$ turns $N^{\beta,p}(\Omega)$ into a Banach space.

Corollary 4.8 (Fractional convergence rates). Assume that the family of triangulations $\{\mathcal{T}_h\}_h$ is quasi-uniform, and that $\mathcal{F}(\nabla u) \in N^{\beta,2}(\Omega)^{n \times d}$ for some $\beta \in (0,1]$. Then, for $\alpha > 0$ sufficiently large, it holds that

$$
\|\boldsymbol{\mathcal{F}}(\tilde \nabla_h \boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\lesssim h^{2\beta}\left[\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\right]_{N^{\beta,2}(\Omega)}^2,
$$

where the constant depends only on k, ω_0 , and the characteristics of S.

Proof. Thanks to the quasi-uniformity, appealing to [\[8,](#page-30-9) Thm. 4.2], we have that

(4.14)
$$
\|\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})-\boldsymbol{\mathcal{F}}(\nabla \Pi_h^{SZ} \boldsymbol{u})\|_{2,\Omega}^2 \lesssim h^{2\beta} \left[\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\right]_{N^{\beta,2}(\Omega)}^2.
$$

Thus, choosing $\mathbf{v}_h = \prod_h^{SZ} \mathbf{u} \in V_h \cap V$ in [Theorem 4.6,](#page-11-0) due to $[\prod_h^{SZ} \mathbf{u} \otimes \mathbf{n}] = 0$ on Γ_h , we obtain

$$
\begin{aligned}\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\lesssim\|\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})-\boldsymbol{\mathcal{F}}(\nabla\Pi_h^{SZ}\boldsymbol{u})\|_{2,\Omega}^2\\ \lesssim h^{2\beta}\left[\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\right]_{N^{\beta,2}(\Omega)}^2,\end{aligned}
$$

which is the claimed fractional a priori error estimate.

REMARK 4.9. In view of [\[15,](#page-31-16) [Corollary](#page-13-0) 5.8] the assertion of Corollary 4.8 for $\beta = 1$ is also valid if $\mathcal{F}(\nabla u) \in W^{1,2}(\Omega)^{n \times d}$ without the additional assumption that the triangulation $\{\mathcal{T}_h\}_h$ is quasi-uniform.

Eventually, resorting to the approximation properties of the node-averaging quasiinterpolation operator \mathcal{I}_h^{av} : $C^0(\mathcal{T}_h) \to V_{h,c}$, for every $v_h \in \mathbb{P}(\mathcal{T}_h)$, defined via

$$
\boldsymbol{\mathcal{I}}_h^{av} \boldsymbol{v}_h \coloneqq \sum_{y\in\mathcal{L}_k^{\rm int}(\mathcal{T}_h)} \langle \boldsymbol{v}_h\rangle_y \, \varphi_y^k\,, \quad \text{ where }\quad \langle \boldsymbol{v}_h\rangle_y \coloneqq \sum_{T\in\mathcal{T}_h; y\in T} (\boldsymbol{v}_h|_T)(y)\,,
$$

[Theorem 4.6](#page-11-0) allows to carry out an ansatz class competition, which reveals that the approximation capabilities of the LDG and IIDG approximations and continuous Lagrange approximation of the problem [\(1.1\)](#page-0-0) are comparable.

 \Box

 \Box

COROLLARY 4.10 (Ansatz class competition). Let $u_h^c \in V_{h,c} := V_h \cap V$ be the continuous Lagrange solution of [\(1.1\)](#page-0-0), i.e., for every $v_h \in V_{h,c}$, it holds that

(4.15)
$$
(\mathcal{S}(\nabla u_h^c), \nabla v_h)_{\Omega} = (\mathbf{f}, v_h)_{\Omega}.
$$

Then, for $\alpha > 0$ sufficiently large, it holds that

$$
\|\boldsymbol{\mathcal{F}}(\tilde \nabla_h \boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\sim\|\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u}_h^c)-\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2\,,
$$

with constants depending only on k, ω_0 , and the characteristics of S.

Proof. ad \lesssim . Using [Theorem 4.6](#page-11-0) with $\mathbf{v}_h = \mathbf{u}_h^c \in V_{h,c} \subseteq V_h$ and observing that $m_{\varphi_{\beta_h(u_h)},h}(\boldsymbol u_h^c)=0$, we find that

$$
\|\boldsymbol{\mathcal{F}}(\tilde \nabla_h \boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\lesssim \|\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u}_h^c)-\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2\,.
$$

 $ad \gtrsim$. Using [\(2.8\),](#page-4-1) that $\varphi_{|\tilde{\nabla}_h \mathbf{u}_h|} \leq \varphi_{\boldsymbol{\beta}_h(\mathbf{u}_h)}$, and [\[25,](#page-31-7) Prop. A.1], for every $K \in$ \mathcal{T}_h , we find that

$$
\|\boldsymbol{F}(\nabla \boldsymbol{\mathcal{I}}_h^{av} \boldsymbol{u}_h) - \boldsymbol{F}(\tilde{\nabla}_h \boldsymbol{u}_h)\|_{2,K}^2 \lesssim \rho_{\varphi_{|\tilde{\nabla}_h \boldsymbol{u}_h|},K}(\tilde{\nabla}_h (\boldsymbol{u}_h - \boldsymbol{\mathcal{I}}_h^{av} \boldsymbol{u}_h)) \\ \lesssim \rho_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},K}(\tilde{\nabla}_h (\boldsymbol{u}_h - \boldsymbol{\mathcal{I}}_h^{av} \boldsymbol{u}_h)) \\ \lesssim m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h).
$$

Eventually, using the best-approximation properties of $u_h^c \in V_{h,c}$ (cf. [\[15,](#page-31-16) Lem. 5.2]) and [\(4.16\),](#page-14-1) we conclude that

$$
\begin{aligned}\n\|\mathcal{F}(\nabla \boldsymbol{u}_h^c) - \mathcal{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 &\lesssim \|\mathcal{F}(\nabla \mathcal{I}_h^{av} \boldsymbol{u}_h) - \mathcal{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 \\
&\lesssim \|\mathcal{F}(\tilde{\nabla}_h \boldsymbol{u}_h) - \mathcal{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + \|\mathcal{F}(\nabla \mathcal{I}_h^{av} \boldsymbol{u}_h) - \mathcal{F}(\tilde{\nabla}_h \boldsymbol{u}_h)\|_{2,\Omega}^2 \\
&\lesssim \|\mathcal{F}(\tilde{\nabla}_h \boldsymbol{u}_h) - \mathcal{F}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + m_{\varphi_{\boldsymbol{\beta}_h}(\boldsymbol{u}_h),h}(\boldsymbol{u}_h).\n\end{aligned}
$$

Putting everything together, we arrive ar the claimed equivalence.

 \Box

REMARK 4.11 (Application to Crouzeix–Raviart element). If $V_h := \mathcal{CR}^k(\mathcal{T}_h)$, where $\mathcal{CR}^k(\mathcal{T}_h)$ is the Crouzeix-Raviart element and its generalization to higher orders (cf. [\[40,](#page-32-2) Sec. 3.3 (3.17)]), and if we use the same primal formulation with $\tilde{\nabla}_h = \nabla_h$, then it is possible to prove the same results. However, omitting the stabilization terms in the primal formulation as in the linear case is still an open problem, since, in this case, one cannot simply absorb the term $c_{\kappa} m_{\varphi_{\beta_h(u_h)},h}(v_h)$ in [\(4.11\)](#page-12-1) in the term $\alpha c_{\varepsilon} m_{\varphi_{\beta_h(u_h)},h}(v_h)$ in [\(4.10\)](#page-12-4) via choosing $\alpha > 0$ large enough.

5. Mixed formulation. Since $S: \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ is bounded, continuous, and strictly monotone, by the Browder–Minty theorem (cf. [\[42,](#page-32-6) Thm. 26.A]), it is also bijective and its inverse $\mathcal{D} := \mathcal{S}^{-1} \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ is bounded, continuous, and strictly monotone as well. This motivates to consider the following mixed formulation and corresponding discrete mixed formulation. In doing so, throughout the entire section, we assume that $\mathcal{S} \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ has (p, δ) -structure in the sense of [Defini](#page-3-0)[tion 2.1.](#page-3-0) To ensure the validity of the identity (1.6) , in this formulation, it will be crucial to have that $\tilde{\nabla}_h(V_h) \subseteq \Sigma_h$.

5.1. Continuous mixed formulation. We abbreviate

$$
\Sigma := L^{p'}(\Omega)^{n \times d}.
$$

Then, for given $f \in V^*$, the *(continuous) mixed formulation* of [\(1.1\)](#page-0-0) seeks for $(\mathsf{S}, u)^\top \in \Sigma \times V$ such that for every $(\mathsf{T}, z)^\top \in \Sigma \times V$, it holds that

(5.1a)
$$
(\mathcal{D}(\mathsf{S}),\mathsf{T})_{\Omega}-(\mathsf{T},\nabla u)_{\Omega}=0,
$$

(5.1b)
$$
(\mathbf{S}, \nabla z)_{\Omega} = \langle \mathbf{f}, z \rangle_V.
$$

The unique solvability of the mixed formulation (5.1) is a consequence of the unique solvability of the primal formulation (4.1) . In fact, both formulations are equivalent: If $(\mathsf{S},\mathbf{u})^{\top} \in \Sigma \times V$ is a solution of (5.1) , then $(5.1a)$ implies that $\mathcal{D}(\mathsf{S}) = \nabla \mathbf{u}$, which is equivalent to $S = \mathcal{S}(\nabla u)$, and, thus, together with [\(5.1b\)](#page-15-2) shows that $u \in V$ is a solution of [\(4.1\).](#page-9-2) If, in turn, $u \in V$ is a solution of [\(4.1\),](#page-9-2) then, setting $S :=$ $\mathcal{S}(\nabla u) \in \Sigma$, we have that $\mathcal{D}(\mathsf{S}) = \nabla u$ and, thus, $(5.1a)$. In addition, $\mathsf{S} \in \Sigma$ then satisfies [\(5.1b\),](#page-15-2) so that $(\mathbf{S}, \mathbf{u})^{\top} \in \Sigma \times V$ is a solution of [\(5.1\).](#page-15-0) It is possible to prove the weak solvability of [\(5.1\)](#page-15-0) directly using pseudo-monotone operator theory. For this, however, one first needs to ascertain that $\mathcal{D} \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ has similar but dual-growth conditions to $\mathcal{S} \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$.

LEMMA 5.1. Let $\mathcal{S} \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ have (p, δ) -structure in the sense of [Defini](#page-3-0)[tion](#page-3-0) 2.1. Then, its inverse $\mathcal{D} := \mathcal{S}^{-1} : \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ has (p', δ^{p-1}) -structure in the sense of [Definition](#page-3-0) 2.1.

Proof. Due to (2.8) , for every $\mathbf{Q}, \mathbf{P} \in \mathbb{R}^{n \times d}$, it holds that

(5.2)
\n
$$
(\mathcal{D}(\mathbf{Q}) - \mathcal{D}(\mathbf{P})) : (\mathbf{Q} - \mathbf{P}) = (\mathcal{S}(\mathcal{D}(\mathbf{Q})) - \mathcal{S}(\mathcal{D}(\mathbf{P}))) : (\mathcal{D}(\mathbf{Q}) - \mathcal{D}(\mathbf{P}))
$$
\n
$$
\sim (\varphi^*)_{|\mathcal{S}(\mathcal{D}(\mathbf{Q}))|} (|\mathcal{S}(\mathcal{D}(\mathbf{Q})) - \mathcal{S}(\mathcal{D}(\mathbf{P}))|)
$$
\n
$$
= (\varphi^*)_{|\mathbf{Q}|} (|\mathbf{Q} - \mathbf{P}|).
$$

Due to $((\varphi_{|\mathcal{D}(\mathbf{Q})|})^*)' \circ \varphi'_{|\mathcal{D}(\mathbf{Q})|} = id_{\mathbb{R}}, ((\varphi_{|\mathcal{D}(\mathbf{Q})|})^*)' \sim (\varphi^*)'_{|\mathcal{S}(\mathcal{D}(\mathbf{Q}))|}$, and [\(2.9\),](#page-4-5) for every $\mathbf{Q}, \mathbf{P} \in \mathbb{R}^{n \times d}$, it holds that

(5.3)
\n
$$
|\mathcal{D}(\mathbf{Q}) - \mathcal{D}(\mathbf{P})| = (((\varphi_{|\mathcal{D}(\mathbf{Q})|})^*)' \circ \varphi'_{|\mathcal{D}(\mathbf{Q})|})(|\mathcal{D}(\mathbf{Q}) - \mathcal{D}(\mathbf{P})|)
$$
\n
$$
\lesssim (\varphi^*)'_{|\mathcal{S}(\mathcal{D}(\mathbf{Q}))|}(|\mathcal{S}(\mathcal{D}(\mathbf{Q})) - \mathcal{S}(\mathcal{D}(\mathbf{P}))|)
$$
\n
$$
= (\varphi^*)'_{|\mathbf{Q}|}(|\mathbf{Q} - \mathbf{P}|).
$$

In other words, $\mathcal{D} \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ has (p', δ^{p-1}) -structure in the sense of [Defini](#page-3-0)[tion 2.1.](#page-3-0) П

Since, appealing to [Lemma 5.1,](#page-15-3) \mathcal{D} has (p', δ^{p-1}) -structure in the sense of [Def](#page-3-0)[inition 2.1,](#page-3-0) similar to [Proposition 2.3,](#page-4-4) we have the following connections between $\mathcal{D}, \mathcal{F}, \mathcal{F}^* \colon \mathbb{R}^{n \times d} \to \mathbb{R}^{n \times d}$ and $\varphi_a, (\varphi^*)_a, ((\varphi^*)_a)^* \colon \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}, a \geq 0.$

PROPOSITION 5.2. Let S satisfy [Definition](#page-3-0) 2.1, let φ be defined in [\(2.3\)](#page-3-1), and let $\mathcal{F}, \mathcal{F}^*$ be defined in [\(2.7\)](#page-4-0). Then, uniformly with respect to $\mathbf{Q}, \mathbf{P} \in \mathbb{R}^{n \times d}$, we have that

(5.4)
\n
$$
\left(\mathcal{D}(\mathsf{Q})-\mathcal{D}(\mathsf{P})\right) : (\mathsf{Q}-\mathsf{P}) \sim |\mathcal{F}^*(\mathsf{Q})-\mathcal{F}^*(\mathsf{P})|^2
$$
\n
$$
\sim \left((\varphi^*)_{|\mathsf{Q}|}\right)^*(|\mathcal{D}(\mathsf{Q})-\mathcal{D}(\mathsf{P})|)
$$
\n
$$
\sim \varphi_{|\mathcal{D}(\mathsf{Q})|}(|\mathcal{D}(\mathsf{Q})-\mathcal{D}(\mathsf{P})|),
$$

(5.5)
$$
|\mathcal{D}(\mathbf{Q}) - \mathcal{D}(\mathbf{P})| \sim (\varphi^*)'_{|\mathbf{Q}|}(|\mathbf{Q} - \mathbf{P}|),
$$

(5.6)
$$
|\mathcal{F}(\mathcal{D}(\mathsf{P})) - \mathcal{F}(\mathcal{D}(\mathsf{Q}))|^2 \sim |\mathcal{F}^*(\mathsf{P}) - \mathcal{F}^*(\mathsf{Q})|^2.
$$

The constants in (5.4) – (5.6) depend only on the characteristics of S.

5.2. Discrete mixed formulation. For given $k \in \mathbb{N}$, we abbreviate

$$
\Sigma_h \coloneqq \Sigma_h^{k-1} := \mathbb{P}^{k-1}(\mathcal{T}_h)^{n \times d}.
$$

Then, for given $f \in V^*$, the *discrete mixed formulation* of (1.1) seeks for $(\mathbf{S}_h, \mathbf{u}_h) \in$ $\Sigma_h \times V_h$ such that for every $(\mathbf{T}_h, \mathbf{z}_h) \in \Sigma_h \times V_h$, it holds that

(5.7)
$$
(\mathbf{\mathcal{D}}(\mathbf{S}_h) - \tilde{\nabla}_h \mathbf{u}_h, \mathbf{T}_h)_{\Omega} = 0, (\mathbf{S}_h, \mathbf{\mathcal{G}}_h \mathbf{z}_h)_{\Omega} + \alpha \langle \mathbf{\mathcal{S}}_{\mathbf{\mathcal{B}}_h(\mathbf{u}_h)}(h_{\Gamma}^{-1}[\![\mathbf{u}_h \otimes \mathbf{n}]\!]), [\![z_h \otimes \mathbf{n}]\!])_{\Gamma_h} = \langle \mathbf{f}, \mathbf{\mathcal{E}}_h z_h \rangle_V.
$$

PROPOSITION 5.3. The following statements apply:

- (i) If $\tilde{\nabla}_h = \mathcal{G}_h$, then the mixed formulation [\(5.7\)](#page-16-0) admits a solution.
- (ii) If $\tilde{\nabla}_h = \tilde{\nabla}_h$, then for sufficiently large $\alpha > 0$, the mixed formulation [\(5.7\)](#page-16-0) admits a solution.

Proof. We equip $\Sigma_h \times V_h$ with the norm

$$
\|(\mathsf{T}_h,\boldsymbol{z}_h)\|_{\Sigma_h\times V_h}\coloneqq \|\mathsf{T}_h\|_{p',\Omega}+\|\boldsymbol{z}_h\|_{p,h} \,.
$$

For every $\varepsilon > 0$, we consider the regularized operator $\mathcal{T}_h^{\varepsilon} \colon \Sigma_h \times V_h \to (\Sigma_h \times V_h)^*$, defined, for every $(\mathbf{S}_h, \mathbf{u}_h) \in \Sigma_h \times V_h$ and $(\mathbf{T}_h, \mathbf{v}_h) \in \Sigma_h \times V_h$, via

$$
\langle \mathcal{T}_h^{\varepsilon}(\mathsf{S}_h, \boldsymbol{u}_h), (\mathsf{T}_h, \boldsymbol{v}_h) \rangle_{\Sigma_h \times V_h} \coloneqq (\boldsymbol{\mathcal{D}}(\mathsf{S}_h) - \tilde{\nabla}_h \boldsymbol{u}_h, \mathsf{T}_h)_{\Omega} + (\mathsf{S}_h, \boldsymbol{\mathcal{G}}_h \boldsymbol{v}_h)_{\Omega} + \varepsilon (\boldsymbol{\mathcal{S}}(\nabla_h \boldsymbol{u}_h), \nabla_h \boldsymbol{v}_h)_{\Omega} + \alpha \langle \boldsymbol{\mathcal{S}}_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)}(h_{\Gamma}^{-1} [\![\boldsymbol{u}_h \otimes \boldsymbol{n}]\!]), [\![\boldsymbol{v}_h \otimes \boldsymbol{n}]\!])_{\Gamma_h}.
$$

Since V_h is finite dimensional, consists of broken polynomials, [\[5,](#page-30-8) Lem. 3.18], the properties of \mathcal{D}, \mathcal{S} and $\mathcal{S}_a, a \geq 0$, imply that the operator $\mathcal{T}_h^{\varepsilon} \colon \Sigma_h \times V_h \to (\Sigma_h \times V_h)^*,$ for every fixed $h, \varepsilon > 0$, is well-defined, continuous, and monotone. To prove the boundedness of $\mathcal{T}_{h}^{\varepsilon} \colon \Sigma_{h} \times V_{h} \to (\Sigma_{h} \times V_{h})^{*}$, we use the properties of $\mathcal{D}, \mathcal{S}, \mathcal{S}_{\beta_{h}(v_{h})}$ (cf. [Proposition 2.3](#page-4-4) and [Remark 2.4\)](#page-4-7), Young's inequality, (2.1) , the stability of \mathcal{R}_{h} (cf. [\[14,](#page-31-1) (A.25)]), a shift change (cf. [Lemma 2.5\)](#page-5-0), $\sum_{F \in \Gamma_h} h_F \mathcal{H}^{d-1}(F) \sim |\Omega|$, [\[17,](#page-31-0) Lemma 12.1], Jensen's inequality, $\varphi(t) + t^p \sim t^p + \delta^p$, and $\varphi^*(t) + t^{p'} \sim t^{p'} + \delta^{p'}$ for all $t \ge 0$ (cf. [\[5\]](#page-30-8)), to find that

$$
\begin{split}\n&\|\langle\mathcal{T}_{h}^{\varepsilon}(\mathsf{S}_{h},\boldsymbol{u}_{h}),(\mathsf{T}_{h},\boldsymbol{v}_{h})\rangle_{\Sigma_{h}\times V_{h}}| \\
&\lesssim\rho_{\varphi^{*},\Omega}(\mathsf{S}_{h})+\rho_{\varphi^{*},\Omega}(\mathsf{T}_{h})+\rho_{\varphi,\Omega}(\tilde{\nabla}_{h}\boldsymbol{u}_{h})+\rho_{\varphi,\Omega}(\nabla_{h}\boldsymbol{u}_{h})+\rho_{\varphi,\Omega}(\boldsymbol{G}_{h}\boldsymbol{v}_{h}) \\
&\quad+\rho_{\varphi,\Omega}(\tilde{\nabla}_{h}\boldsymbol{v}_{h})+\alpha\,m_{\varphi_{\boldsymbol{\beta}_{h}}(\boldsymbol{v}_{h}),h}(\boldsymbol{u}_{h})+\alpha\,m_{\varphi_{\boldsymbol{\beta}_{h}}(\boldsymbol{v}_{h}),h}(\boldsymbol{v}_{h}) \\
&\lesssim\rho_{\varphi^{*},\Omega}(\mathsf{S}_{h})+\rho_{\varphi^{*},\Omega}(\mathsf{T}_{h})+\rho_{\varphi,\Omega}(\nabla_{h}\boldsymbol{u}_{h})+\rho_{\varphi,\Omega}(\nabla_{h}\boldsymbol{v}_{h})+m_{\varphi,h}(\boldsymbol{u}_{h})+\varphi(\boldsymbol{\beta}_{h}(\boldsymbol{v}_{h})) \\
&\lesssim\|\mathsf{S}_{h}\|_{p'}^{p'}+\|\mathsf{T}_{h}\|_{p'}^{p'}+\|\boldsymbol{u}_{h}\|_{p,h}^{p}+\|\boldsymbol{v}_{h}\|_{p,h}^{p}+\delta^{p}+\delta^{p'}.\n\end{split}
$$

Thus, $\mathcal{T}_h^{\varepsilon} \colon \Sigma_h \times V_h \to (\Sigma_h \times V_h)^*$ is pseudo-monotone. If we can prove its coercivity, then, owing to the pseudo-monotone operator theory (cf. [\[42,](#page-32-6) Thm. 27.A]), it follows that $\mathcal{T}_h^{\varepsilon} \colon \Sigma_h \times V_h \to (\Sigma_h \times V_h)^*$ is surjective. To this end, we distinguish the cases $\tilde{\nabla}_h = \tilde{\boldsymbol{\mathcal{G}}}_h$ and $\tilde{\nabla}_h = \nabla_h$:

Case $\tilde{\nabla}_h = \mathcal{G}_h$. For every $(\mathsf{S}_h, \mathbf{u}_h) \in \Sigma_h \times V_h$, using that $\mathcal{D}(\mathsf{Q}) : \mathsf{Q} \sim \varphi^*(\vert \mathsf{Q} \vert)$ (cf. [\(5.4\)\)](#page-15-4) and $\mathcal{S}_{\beta_h(u_h)}(\mathbf{Q}): \mathbf{Q} \sim \varphi_{\beta_h(u_h)}(|\mathbf{Q}|)$ for all $\mathbf{Q} \in \mathbb{R}^{n \times d}$ (cf. [Proposition 2.3](#page-4-4)) and [Remark 2.4\)](#page-4-7) as well as that $\varphi_{\beta_h(\mathbf{u}_h)} \geq \varphi$, $\delta^{p'} + \varphi^*(t) \sim t^{p'} + \delta^{p'}$, and $\delta^p + \varphi(t) \sim$ $t^p + \delta^p$ for all $t \geq 0$, and $\sum_{F \in \Gamma_h} h_F \mathcal{H}^{d-1}(F) \lesssim |\Omega|$, we find that

$$
\langle \mathcal{T}_{h}^{\varepsilon}(\mathsf{S}_{h}, \boldsymbol{u}_{h}), (\mathsf{S}_{h}, \boldsymbol{u}_{h}) \rangle_{\Sigma_{h} \times V_{h}} = (\mathcal{D}(\mathsf{S}_{h}), \mathsf{S}_{h})_{\Omega} + \varepsilon (\mathcal{S}(\nabla_{h} \boldsymbol{u}_{h}), \nabla_{h} \boldsymbol{u}_{h})_{\Omega} + \alpha \langle \mathcal{S}_{\beta_{h}(\boldsymbol{u}_{h})}(h_{\Gamma}^{-1}[\![\boldsymbol{u}_{h} \otimes \boldsymbol{n}]\!]), [\![\boldsymbol{u}_{h} \otimes \boldsymbol{n}]\!]\rangle_{\Gamma_{h}} \geq \rho_{\varphi^*,\Omega}(\mathsf{S}_{h}) + \varepsilon \rho_{\varphi,\Omega}(\nabla_{h} \boldsymbol{u}_{h}) + \alpha m_{\varphi_{\beta_{h}(\boldsymbol{u}_{h})},h}(\boldsymbol{u}_{h}) \geq ||\mathsf{S}_{h}||_{p',\Omega}^{p'} + \min\{\varepsilon,\alpha\} ||\boldsymbol{u}_{h}||_{p,h}^{p} - \delta^{p} |\Omega| (\varepsilon + \alpha) - \delta^{p'} |\Omega|.
$$

Putting everything together, $\mathcal{T}_h^{\varepsilon} \colon \Sigma_h \times V_h \to (\Sigma_h \times V_h)^*$ is well-defined, continuous, and coercive and, thus, surjective (cf. [\[42,](#page-32-6) Thm. 27.A]).

Case $\tilde{\nabla}_h = \nabla_h$. For every $(\mathbf{S}_h, \mathbf{u}_h) \in \Sigma_h \times V_h$, proceeding as before, also using the *κ*-Young inequality [\(2.2\)](#page-3-2) with $\psi = |\cdot|^{p'}$ and that $||\mathcal{R}_h \mathbf{u}_h||_{p,\Omega} \lesssim ||h_{\Gamma}^{1/p'}$ $_{\Gamma}^{\scriptscriptstyle 1/\,p}\; [\![u_h\otimes n]\!] \|_{p,\Gamma_h}$ (cf. $[14, (A.25)]$ $[14, (A.25)]$), it holds that

(5.8)
\n
$$
\langle \mathcal{T}_{h}^{\varepsilon}(\mathsf{S}_{h}, \mathbf{u}_{h}), (\mathsf{S}_{h}, \mathbf{u}_{h}) \rangle_{\Sigma_{h} \times V_{h}} = (\mathcal{D}(\mathsf{S}_{h}) - \mathcal{R}_{h} \mathbf{u}_{h}, \mathsf{S}_{h})_{\Omega} + \varepsilon (\mathcal{S}(\nabla_{h} \mathbf{u}_{h}), \nabla_{h} \mathbf{u}_{h})_{\Omega} + \alpha \langle \mathcal{S}_{\beta_{h}(\mathbf{u}_{h})}(h_{\Gamma}^{-1}[\![\mathbf{u}_{h} \otimes \mathbf{n}]\!]), [\![\mathbf{u}_{h} \otimes \mathbf{n}]\!] \rangle_{\Gamma_{h}} \n\geq (c - \kappa) \|\mathsf{S}_{h}\|_{p',\Omega}^{p'} + \min\{\varepsilon, \alpha - c_{\kappa}\} \|\mathbf{u}_{h}\|_{p,h}^{p} \n- c \delta^{p} |\Omega| (\varepsilon + \alpha) - c \delta^{p'} |\Omega|.
$$

Therefore, choosing first $\kappa > 0$ sufficiently small and, subsequently, $\alpha > 0$ sufficiently large in [\(5.8\),](#page-17-0) for every $(\mathbf{S}_h, \mathbf{u}_h) \in \Sigma_h \times V_h$, we conclude that

$$
\begin{aligned}\n\langle \mathcal{T}_{h}^{\varepsilon}(\mathsf{S}_{h},\boldsymbol{u}_{h}),(\mathsf{S}_{h},\boldsymbol{u}_{h})\rangle_{\Sigma_{h}\times V_{h}} \\
&\gtrsim \|\mathsf{S}_{h}\|_{p',\Omega}^{p'}+\min\{\varepsilon,\alpha\}\left\|\boldsymbol{u}_{h}\right\|_{p,h}^{p}-\delta^{p}\left|\Omega\right|(\varepsilon+\alpha)-\delta^{p'}\left|\Omega\right|.\n\end{aligned}
$$

Putting everything together, $\mathcal{T}_{h}^{\varepsilon} \colon \Sigma_{h} \times V_{h} \to (\Sigma_{h} \times V_{h})^{*}$ is well-defined, continuous, and coercive and, thus, surjective (cf. [\[42,](#page-32-6) Thm. 27.A]).

Therefore, for every $\varepsilon, h > 0$ and $\tilde{\nabla}_h \in {\mathcal{G}_h, \nabla_h},$ there exists $(\mathbf{S}_h^{\varepsilon}, \mathbf{u}_h^{\varepsilon}) \in \Sigma_h \times V_h$ such that $\mathcal{T}_{h}^{\varepsilon}(\mathbf{S}_{h}^{\varepsilon}, \mathbf{u}_{h}^{\varepsilon}) = (\mathbf{0}, \mathcal{E}_{h}^{*} \mathbf{f})$ in $(\Sigma_{h} \times V_{h})^{*} \cong \Sigma_{h}^{*} \times V_{h}^{*}$, i.e., for every $(\mathbf{T}_{h}, \mathbf{z}_{h}) \in$ $\Sigma_h \times V_h$, it holds that

(5.9a) $(\mathcal{D}(\mathsf{S}_h^\varepsilon), \mathsf{T}_h)_{\Omega} - (\mathsf{T}_h, \tilde{\nabla}_h \mathbf{u}_h^\varepsilon)_{\Omega} = 0,$

(5.9b)
$$
(\mathbf{S}_{h}^{\varepsilon}, \mathbf{G}_{h} \mathbf{z}_{h})_{\Omega} + \varepsilon (\mathbf{S}(\nabla_{h} \mathbf{u}_{h}^{\varepsilon}), \nabla_{h} \mathbf{z}_{h})_{\Omega} + \alpha \langle \mathbf{S}_{\beta_{h}(\mathbf{u}_{h}^{\varepsilon})}(h_{\Gamma}^{-1}[\![\mathbf{u}_{h}^{\varepsilon} \otimes \mathbf{n}]\!]), [\![\mathbf{z}_{h} \otimes \mathbf{n}]\!]\rangle_{\Gamma_{h}} = \langle \mathbf{f}, \mathbf{E}_{h} \mathbf{z}_{h} \rangle_{V}.
$$

Then, choosing $z_h = u_h^{\varepsilon} \in V_h$, $\mathbf{T}_h = \mathbf{S}_h^{\varepsilon} \in \Sigma_h$ in [\(5.9\),](#page-17-1) and using [Remark 4.1,](#page-9-3) [\(1.6\)](#page-2-1) we find that

$$
\langle f, \mathcal{E}_h \mathbf{u}_h^{\varepsilon} \rangle_V = (\mathcal{D}(\mathsf{S}_h^{\varepsilon}), \mathsf{S}_h^{\varepsilon})_{\Omega} + \varepsilon (\mathcal{S}(\nabla_h \mathbf{u}_h^{\varepsilon}), \nabla_h \mathbf{u}_h^{\varepsilon})_{\Omega} + \alpha \langle \mathcal{S}_{\beta_h}(\mathbf{u}_h^{\varepsilon}) (h_{\Gamma}^{-1} [\![\mathbf{u}_h^{\varepsilon} \otimes \mathbf{n}]\!]), [\![\mathbf{u}_h^{\varepsilon} \otimes \mathbf{n}]\!] \rangle_{\Gamma_h} \geq \rho_{\varphi^*,\Omega}(\mathsf{S}_h^{\varepsilon}) + \varepsilon \rho_{\varphi,\Omega}(\nabla_h \mathbf{u}_h^{\varepsilon}) + \alpha m_{\varphi_{\beta_h}(\mathbf{u}_h^{\varepsilon})}, h}(\mathbf{u}_h^{\varepsilon}).
$$

From [\(5.9a\),](#page-17-2) we obtain $\tilde{\nabla}_h \boldsymbol{u}_h^{\varepsilon} = \Pi_h^{k-1} \boldsymbol{\mathcal{D}}(\mathbf{S}_h^{\varepsilon})$. Using this, the Orlicz stability of Π_h^{k-1} (cf. [\[14,](#page-31-1) (A.11)]), that $\mathcal D$ has (p', δ^{p-1}) -structure (cf. [Lemma 5.1\)](#page-15-3), and the equivalence (2.1) , we obtain \overline{a}

(5.11)
$$
\rho_{\varphi,\Omega}(\tilde{\nabla}_h \mathbf{u}_h^{\varepsilon}) = \rho_{\varphi,\Omega}(\Pi_h^{k-1} \mathcal{D}(\mathbf{S}_h^{\varepsilon})) \n\lesssim \rho_{\varphi,\Omega}((\varphi^*)'(\mathbf{S}_h^{\varepsilon})) \n\lesssim \rho_{\varphi^*,\Omega}(\mathbf{S}_h^{\varepsilon}).
$$

Using [\(5.11\)](#page-17-3) in [\(5.10\)](#page-17-4) and $\varphi_{\mathcal{B}_h(u_h)} \geq \varphi, \delta^{p'} + \varphi^*(t) \sim t^{p'} + \delta^{p'},$ and $\delta^p + \varphi(t) \sim t^p + \delta^p$ for all $t \geq 0$, $\sum_{F \in \Gamma_h} h_F \mathcal{H}^{d-1}(F) \lesssim |\Omega|$, and (2.12) if $\tilde{\nabla}_h = \mathcal{G}_h$, yields that

(5.12)
$$
\langle \mathbf{f}, \mathbf{\mathcal{E}}_h \mathbf{u}_h^{\varepsilon} \rangle_V \gtrsim \rho_{\varphi^*,\Omega}(\mathbf{S}_h^{\varepsilon}) + (1+\varepsilon) \rho_{\varphi,\Omega}(\tilde{\nabla}_h \mathbf{u}_h^{\varepsilon}) + \alpha m_{\varphi_{\mathbf{\mathcal{B}}_h}(\mathbf{u}_h^{\varepsilon})}, h(\mathbf{u}_h^{\varepsilon})
$$

$$
\gtrsim \|\mathbf{S}_h^{\varepsilon}\|_{p',\Omega}^{p'} + \min\{1,\alpha\} \| \mathbf{u}_h^{\varepsilon}\|_{p,h}^{p} - \delta^p |\Omega| (1+\alpha) - \delta^{p'} |\Omega|.
$$

Crucially, the constant in the last inequality is independent of ε . On the other hand, using the κ -Young inequality [\(2.2\)](#page-3-2) with $\psi = |\cdot|^p$ and [Proposition 3.1](#page-8-4) with $\psi = |\cdot|^p$, we find that ′

(5.13)
$$
|\langle \mathbf{f}, \mathbf{\mathcal{E}}_h \mathbf{u}_h^{\varepsilon} \rangle_V| \leq c_{\kappa} \| \mathbf{f} \|_{V^*}^{p'} + \kappa \| \mathbf{\mathcal{E}}_h \mathbf{u}_h^{\varepsilon} \|_{V}^{p'}
$$

$$
\lesssim c_{\kappa} \| \mathbf{f} \|_{V^*}^{p'} + \kappa \| \mathbf{u}_h^{\varepsilon} \|_{p,h}^{p}.
$$

Therefore, combining [\(5.12\)](#page-18-1) and [\(5.13\)](#page-18-2) as well as choosing $\kappa > 0$ sufficiently small, we conclude that

$$
\|\mathbf{S}_{h}^{\varepsilon}\|_{p',\Omega}^{p'} + \min\{1,\alpha\} \, \|u_{h}^{\varepsilon}\|_{p,h}^{p} \lesssim 1 \, .
$$

Therefore, due to the finite dimensionality of Σ_h and V_h , the Bolzano–Weierstraß compactness theorem yields the existence of a not re-labeled subsequence as well as limits $\mathbf{S}_h \in \Sigma_h$ and $\mathbf{u}_h \in V_h$ such that

(5.14)
$$
\begin{aligned}\n\mathbf{S}_h^{\varepsilon} \to \mathbf{S}_h & \text{strongly in } \Sigma_h & (\varepsilon \to 0), \\
\mathbf{u}_h^{\varepsilon} \to \mathbf{u}_h & \text{strongly in } V_h & (\varepsilon \to 0).\n\end{aligned}
$$

Using [\(5.14\),](#page-18-3) by passing for $\varepsilon \to 0$ in [\(5.9\),](#page-17-1) we conclude that

$$
(\mathcal{D}(\mathsf{S}_h), \mathsf{T}_h)_{\Omega} - (\mathsf{T}_h, \tilde{\nabla}_h \mathbf{u}_h)_{\Omega} = 0,
$$

$$
(\mathsf{S}_h, \mathcal{G}_h, z_h)_{\Omega} + \alpha \langle \mathcal{S}_{\beta_h(\mathbf{u}_h)}(h_{\Gamma}^{-1}[\![\mathbf{u}_h \otimes \mathbf{n}]\!]), [\![z_h \otimes \mathbf{n}]\!] \rangle_{\Gamma_h} = \langle \mathbf{f}, \mathcal{E}_h z_h \rangle_V,
$$

i.e., $(\mathbf{S}_h, \mathbf{u}_h) \in \Sigma_h \times V_h$ is a solution of the discrete mixed formulation [\(5.7\).](#page-16-0) \Box

Let us next prove a best-approximation result similar to [Theorem 4.6,](#page-11-0) but now for the mixed formulation [\(5.7\).](#page-16-0) To this end, we need to restrict ourselves either to the case of element-wise affine functions or the case $p \leq 2$ together with the assumption that $(\delta + |\overline{\nabla u}|)^{2-p}$, where $\overline{\nabla u} := \nabla u$ in Ω and $\overline{\nabla u} := \mathbf{0}$ in $\mathbb{R}^d \setminus \Omega$, belongs to the Muckenhoupt class $A_2(\mathbb{R}^d)$. More precisely, given $p \in [1, \infty)$, a weight $\sigma \colon \mathbb{R}^d \to (0, +\infty)$, i.e., $\sigma \in L^1_{loc}(\mathbb{R}^d)$ and $0 < \sigma(x) < +\infty$ for a.e. $x \in \mathbb{R}^d$, is said to satisfy the A_p -condition, if

$$
[\sigma]_{A_p(\mathbb{R}^d)} \coloneqq \sup_{B \subseteq \mathbb{R}^d; B \text{ ball}} \langle \sigma \rangle_B (\langle \sigma^{1-p'} \rangle_B)^{p-1} < \infty \, .
$$

We denote by $A_p(\mathbb{R}^d)$ the class of all weights satisfying the A_p -condition and use weighted Lebesgue spaces $L^p(\Omega; \sigma)$ equipped with the norm $||v||_{p,\sigma,\Omega} := (\int_{\Omega} |v|^p \sigma dx)^{\frac{1}{p}}$.

Remark 5.4 (Comments on Muckenhoupt weights).

- (i) If $\sigma \in L^1_{loc}(\mathbb{R}^d)$ is a weight and there exist $c, C > 0$ such that $c \leq \sigma \leq C$ a.e. in \mathbb{R}^d , then $\sigma \in A_p(\mathbb{R}^d)$ for all $p \in [1,\infty)$.
- (ii) If $\sigma \coloneqq |\cdot|^{\eta} \in L^1_{loc}(\mathbb{R}^d)$, then $\sigma \in A_p(\mathbb{R}^d)$, $p \in (1, \infty)$, if $-d < \eta < d(p-1)$.

THEOREM 5.5. Assume that $k = 1$ or that $p \leq 2$ and $(\delta + |\overline{\nabla u}|)^{2-p} \in A_2(\mathbb{R}^d)$, where $\overline{\nabla u} := \nabla u$ in Ω and $\overline{\nabla u} := 0$ in $\mathbb{R}^d \setminus \Omega$. Then, for $\alpha > 0$ sufficiently large, we have that

$$
\begin{aligned}\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_h)-\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\\ \lesssim&\inf_{(\boldsymbol{\mathsf{T}}_h,\boldsymbol{v}_h)}\left(\|\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})-\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{v}_h)\|_{2,\Omega}^2+\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})-\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{T}}_h)\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{v}_h)\right).\end{aligned}
$$

where the constant depends only on k, ω_0 , $[\sigma]_{A_2(\mathbb{R}^d)}$, and the characteristics of S and D.

Key ingredient in the proof of [Theorem 5.5](#page-18-0) is the following convex conjugation type inequality for the natural distance, which applies, if, e.g., $k = 1$ or if $p \leq 2$ and $(\delta + |\overline{\nabla u}|)^{2-p} \in A_2(\mathbb{R}^d)$. In particular, note that if one is capable of establishing these convex conjugation formula for more general assumptions, then [Theorem 5.5](#page-18-0) immediately applies for these assumptions.

Lemma 5.6. The following statements apply: (i) If $k = 1$, then for every $v_h, w_h \in V_h$, it holds that

$$
\begin{split} &\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{v}_h)-\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{w}_h)\|_{2,\Omega}^2\\ &\lesssim \sup_{\boldsymbol{\mathsf{T}}_h\in\Sigma_h}\left[(\tilde{\nabla}_h\boldsymbol{v}_h-\tilde{\nabla}_h\boldsymbol{w}_h,\boldsymbol{\mathsf{T}}_h)_\Omega-\frac{1}{c}\rho_{(\varphi_{|\tilde{\nabla}_h\boldsymbol{v}_h|})^*,\Omega}(\boldsymbol{\mathsf{T}}_h)\right], \end{split}
$$

where the constants depend only on k, ω_0 , $[\sigma]_{A_2(\mathbb{R}^d)}$, and the characteristics of S and D .

(ii) If $p \leq 2$, $\mathcal{F}(\nabla u) \in L^2(\Omega)^{n \times d}$ and $(\delta + |\nabla u|)^{2-p} \in A_2(\mathbb{R}^d)$, then for every $v_h, w_h \in V_h$, it holds that

$$
\begin{aligned} &\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{v}_h)-\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{w}_h)\|_{2,\Omega}^2\\ &\lesssim \sup_{\boldsymbol{\mathsf{T}}_h\in \Sigma_h}\left[(\tilde{\nabla}_h\boldsymbol{v}_h-\tilde{\nabla}_h\boldsymbol{w}_h,\boldsymbol{\mathsf{T}}_h)_\Omega-\frac{1}{c}\rho_{(\varphi_{\mid \tilde{\nabla}_h\boldsymbol{v}_h\mid)^*,\Omega}(\boldsymbol{\mathsf{T}}_h)\right]\\ &+\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{v}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2\,, \end{aligned}
$$

where the constants depend only on k, ω_0 , $[\sigma]_{A_2(\mathbb{R}^d)}$, and the characteristics of S and D .

Lemma 5.6 (ii), in turn, is essentially based on the following (local) stability result for the L^2 -projection in terms of weighted Lebesgue norms.

LEMMA 5.7. If $\sigma \in A_p(\mathbb{R}^d)$, $p \in (1,\infty)$, then for every $\mathbf{T} \in L^p(\Omega;\sigma)^{n \times d}$ and $K \in \mathcal{T}_h$, we have that

$$
\|\Pi_h^{k-1}\mathbf{T}\|_{p,\sigma,K}\lesssim \|\mathbf{T}\|_{p,\sigma,K},
$$

where the constant depends only on k, ω_0 , p, and $[\sigma]_{A_p(\mathbb{R}^d)}$.

Proof. Using Hölder's inequality, a local norm equivalence (cf. [\[17,](#page-31-0) Lem. 12.1]), the L¹-stability of Π_h^{k-1} (cf. [\[14\]](#page-31-1)), $|B_{h_K}(x_K)| \sim |K|$, where x_K is the barycenter of K,

and $\sigma \in A_p(\mathbb{R}^d)$, we observe that

$$
\|\Pi_h^{k-1}\mathbf{T}\|_{p,\sigma,K} \le \|\Pi_h^{k-1}\mathbf{T}\|_{\infty,K} \|\sigma\|_{1,K}^{1/p}
$$

\n
$$
\lesssim |K|^{-1} \|\Pi_h^{k-1}\mathbf{T}\|_{1,K} \|\sigma\|_{1,K}^{1/p}
$$

\n
$$
\lesssim |K|^{-1} \|\mathbf{T}\|_{1,K} \|\sigma\|_{1,K}^{1/p}
$$

\n
$$
\le |K|^{-1} \|\mathbf{T}\sigma^{1/p} \sigma^{-1/p}\|_{1,K} \|\sigma\|_{1,K}^{1/p}
$$

\n
$$
\le \|\mathbf{T}\|_{p,\sigma,K} \||K|^{-1} \sigma^{1-p'}\|_{1,K}^{1/p'} \||K|^{-1} \sigma\|_{1,K}^{1/p}
$$

\n
$$
\lesssim \|\mathbf{T}\|_{p,\sigma,K} \||B_{h_K}(x_K)|^{-1} \sigma^{1-p'}\|_{1,B_{h_K}(x_K)}^{1/p'} \|B_{R_K}(x_K)|^{-1} \sigma\|_{1,B_{h_K}(x_K)}^{1/p}
$$

\n
$$
\lesssim \|\mathbf{T}\|_{p,\sigma,K},
$$

where the constants depend only on k, ω_0 , p, and $[\sigma]_{A_p(\mathbb{R}^d)}$.

Proof (of [Lemma](#page-19-0) 5.6). ad (i). For $\tilde{\nabla} \mathbf{v}_h \in \Sigma_h$, the local Orlicz-stability of Π_h^{k-1} (cf. [\[14,](#page-31-1) (A.11)]), for every $\mathbf{T} \in L^{(\varphi_1 \tilde{\nabla}_h \mathbf{v}_h)^*}(\Omega)^{n \times d}$, implies that

$$
\rho_{(\varphi_\mid\bar{\nabla}_h v_h\mid)^*,\Omega}(\Pi_h^{k-1}\mathbf{T})\leq \rho_{(\varphi_\mid\bar{\nabla}_h v_h\mid)^*,\Omega}(\mathbf{T})\,.
$$

Using this, [Proposition 2.3,](#page-4-4) that $\varphi_{\vert \tilde{\nabla}_h v_h \vert} = (\varphi_{\vert \tilde{\nabla}_h v_h \vert})^{**} = ((\varphi_{\vert \tilde{\nabla}_h v_h \vert})^*)^*$, the convex conjugation formula for integral functionals (cf. [\[16,](#page-31-17) Prop. 1.2]), and $\tilde{\nabla} \mathbf{v}_h, \tilde{\nabla} \mathbf{w}_h \in \Sigma_h$, we find that

$$
\begin{split}\n&\|\mathcal{F}(\tilde{\nabla}_{h}\mathbf{v}_{h})-\mathcal{F}(\tilde{\nabla}_{h}\mathbf{w}_{h})\|_{2,\Omega}^{2} \lesssim \rho_{\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert,\Omega}}(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h}) \\
&= \rho_{(\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert})^{**},\Omega}(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h}) \\
&= \sup_{\mathbf{T}\in L^{(\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert})^{*}}(\Omega)^{n\times d}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\mathbf{T})_{\Omega}-\rho_{(\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert})^{*},\Omega}(\mathbf{T})\right] \\
&= \sup_{\mathbf{T}\in L^{(\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert})^{*}}(\Omega)^{n\times d}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\Pi_{h}^{k-1}\mathbf{T})_{\Omega}-\rho_{(\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert})^{*},\Omega}(\mathbf{T})\right] \\
&\lesssim \sup_{\mathbf{T}\in L^{(\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert})^{*}}(\Omega)^{n\times d}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\Pi_{h}^{k-1}\mathbf{T})_{\Omega}-\frac{1}{c}\rho_{(\varphi_{\lvert \tilde{\nabla}_{h}\mathbf{v}_{h}\rvert})^{*},\Omega}(\Pi_{h}^{k-1}\mathbf{T})\right] \\
&\lesssim \sup_{\mathbf{T}_{h}\in\Sigma_{h}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\mathbf{T}_{h})_{\Omega}-\frac{1}{c}\rho_{(\varphi_{\l
$$

which is the claimed convex conjugation type inequality for the natural distance in the case $k = 1$.

ad (ii). Abbreviate $\sigma := (\delta + |\overline{\nabla u}|)^{2-p}$. Due $p \le 2$, for a.e. $x \in \Omega$ and every $t \ge 0$, it holds that

(5.15)
$$
((\delta + |\nabla \mathbf{u}(x)|)^{p-1} + t)^{p'-2} t^2 \le 2^{p'-2} (\sigma(x) t^2 + t^{p'}) \le 2^{p'-2} ((\delta + |\nabla \mathbf{u}(x)|)^{p-1} + t)^{p'-2} t^2.
$$

Therefore, using [\(5.15\),](#page-20-0) the stability of Π_h^{k-1} in $L^{p'}(\Omega)^{n \times d}$ (cf. [\[26,](#page-31-18) Cor. A.8]), and in $L^2(\Omega,\sigma)^{n\times d}$ (cf. [Lemma 5.7\)](#page-19-1) together with $(\delta + |\nabla u|)^{2-p} \in A_2(\mathbb{R}^d)$, for every $\mathbf{T} \in L^{p'}(\Omega)^{n \times d} \cap L^2(\Omega;\sigma)^{n \times d} \sim L^{(\varphi_{|\nabla u|})^*}(\Omega)^{n \times d}$, it holds that

′

(5.16)
$$
\rho_{(\varphi_{|\nabla u|})^*}(\Pi_h^{k-1}\mathbf{T}) \lesssim \|\Pi_h^{k-1}\mathbf{T}\|_{2,\omega}^2 + \|\Pi_h^{k-1}\mathbf{T}\|_{p'}^{p'}
$$

$$
\lesssim \|\mathbf{T}\|_{2,\omega}^2 + \|\mathbf{T}\|_{p'}^{p'} \lesssim \rho_{(\varphi_{|\nabla u|})^*}(\mathbf{T}),
$$

 \Box

which, using a shift change in [Lemma 2.5,](#page-5-0) implies that

$$
\rho_{(\varphi_{|\tilde{\nabla}_h v_h|})^*,\Omega}(\Pi_h^{k-1}\mathbf{T}) \lesssim \rho_{(\varphi_{|\nabla u|})^*,\Omega}(\Pi_h^{k-1}\mathbf{T}) + \|\mathcal{F}(\tilde{\nabla}_h v_h) - \mathcal{F}(\nabla u)\|_{2,\Omega}^2
$$

\n
$$
\lesssim \rho_{(\varphi_{|\nabla u|})^*,\Omega}(\mathbf{T}) + \|\mathcal{F}(\tilde{\nabla}_h v_h) - \mathcal{F}(\nabla u)\|_{2,\Omega}^2
$$

\n
$$
\lesssim \rho_{(\varphi_{|\tilde{\nabla}_h v_h|})^*,\Omega}(\mathbf{T}) + \|\mathcal{F}(\tilde{\nabla}_h v_h) - \mathcal{F}(\nabla u)\|_{2,\Omega}^2.
$$

Using (5.17) and proceeding as in (i) , we find that

$$
\begin{split}\n&\|\mathcal{F}(\tilde{\nabla}_{h}\mathbf{v}_{h})-\mathcal{F}(\tilde{\nabla}_{h}\mathbf{w}_{h})\|_{2,\Omega}^{2} \lesssim \rho_{\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|},\Omega}(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h}) \\
&= \rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{**},\Omega}(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h}) \\
&= \sup_{\mathbf{T}\in L^{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*}}(\Omega)^{n\times d}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\mathbf{T})_{\Omega}-\rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*},\Omega}(\mathbf{T})\right] \\
&= \sup_{\mathbf{T}\in L^{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*}}(\Omega)^{n\times d}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\Pi_{h}^{k-1}\mathbf{T})_{\Omega}-\rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*},\Omega}(\mathbf{T})\right] \\
&\lesssim \sup_{\mathbf{T}\in L^{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*}}(\Omega)^{n\times d}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\Pi_{h}^{k-1}\mathbf{T})_{\Omega}-\frac{1}{c}\rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*},\Omega}(\Pi_{h}^{k}\mathbf{T})\right] \\
&\quad + \|\mathcal{F}(\tilde{\nabla}_{h}\mathbf{v}_{h})-\mathcal{F}(\nabla\mathbf{u})\|_{2,\Omega}^{2} \\
&\lesssim \sup_{\mathbf{T}_{h}\in\Sigma_{h}}\left[(\tilde{\nabla}_{h}\mathbf{v}_{h}-\tilde{\nabla}_{h}\mathbf{w}_{h},\mathbf{T}_{h})_{\Omega}-\
$$

which is the claimed convex conjugation type inequality for the natural distance in the case $p \leq 2$ and $\sigma \in A_2(\mathbb{R}^d)$. \Box

Proof (of [Theorem](#page-18-0) 5.5). By the continuous mixed formulation (5.1) , the discrete mixed formulation [\(5.7\)](#page-16-0) and the crucial identity [\(1.6\),](#page-2-1) for every $(\mathbf{T}_h, z_h) \in \Sigma_h \times V_h$, we have that

(5.18)
$$
(\mathcal{D}(\mathsf{S}_h) - \mathcal{D}(\mathsf{S}), \mathsf{T}_h)_{\Omega} = (\tilde{\nabla}_h \mathbf{u}_h - \nabla \mathbf{u}, \mathsf{T}_h)_{\Omega},
$$

(5.19)
$$
(\mathsf{S}_h - \mathsf{S}, \nabla_h \mathcal{E}_h z_h)_{\Omega} = -\alpha \langle \mathcal{S}_{\beta_h(\mathbf{u}_h)}(h_{\Gamma}^{-1}[\![\mathbf{u}_h \otimes \mathbf{n}]\!]), [\![z_h \otimes \mathbf{n}]\!]\rangle_{\Gamma_h}.
$$

Therefore, taking an arbitrary $v_h \in V_h$, using [Lemma 5.6,](#page-19-0) [\(5.18\),](#page-21-1) the ε -Young in-equality [\(2.2\)](#page-3-2) with $\psi = \varphi_{|\tilde{\nabla}_h \mathbf{v}_h|}$, [\(2.8\),](#page-4-1) and a shift change in [Lemma 2.5,](#page-5-0) choosing $\varepsilon > 0$ sufficiently small, and using [\(5.4\)](#page-15-4) with $\nabla u = \mathcal{D}(\mathsf{S})$, we find that

$$
\|\mathcal{F}(\tilde{\nabla}_{h}\mathbf{u}_{h}) - \mathcal{F}(\tilde{\nabla}_{h}\mathbf{v}_{h})\|_{2,\Omega}^{2} \n\lesssim \sup_{\mathbf{T}_{h}\in\Sigma_{h}} [(\tilde{\nabla}_{h}\mathbf{u}_{h} - \tilde{\nabla}_{h}\mathbf{v}_{h}, \mathbf{T}_{h})_{\Omega} - \frac{1}{c}\rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h})^{*}},\Omega}(\mathbf{T}_{h})] + \|\mathcal{F}(\nabla\mathbf{u}) - \mathcal{F}(\tilde{\nabla}_{h}\mathbf{v}_{h})\|_{2,\Omega}^{2} \n= \sup_{\mathbf{T}_{h}\in\Sigma_{h}} [(\mathcal{D}(\mathsf{S}_{h}) - \mathcal{D}(\mathsf{S}), \mathbf{T}_{h})_{\Omega} + (\nabla\mathbf{u} - \tilde{\nabla}_{h}\mathbf{v}_{h}, \mathbf{T}_{h})_{\Omega} - \frac{1}{c}\rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h})^{*}},\Omega}(\mathbf{T}_{h})] \n+ \|\mathcal{F}(\nabla\mathbf{u}) - \mathcal{F}(\tilde{\nabla}_{h}\mathbf{v}_{h})\|_{2,\Omega}^{2} \n\leq c_{\varepsilon} \rho_{\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|},\Omega}(\mathcal{D}(\mathsf{S}_{h}) - \mathcal{D}(\mathsf{S})) + (c_{\varepsilon} + 1) \|\mathcal{F}(\nabla\mathbf{u}) - \mathcal{F}(\tilde{\nabla}_{h}\mathbf{v}_{h})\|_{2,\Omega}^{2} \n+ \sup_{\mathbf{T}_{h}\in\Sigma_{h}} [2\varepsilon \rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*}},\Omega}(\mathbf{T}_{h}) - \frac{1}{c}\rho_{(\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|)^{*},\Omega}(\mathbf{T}_{h})] \n\lesssim \rho_{\varphi_{|\tilde{\nabla}_{h}\mathbf{v}_{h}|},\Omega}(\mathcal{D}(\mathsf{S}_{h}) - \mathcal{D}(\mathsf{S})) + \|\mathcal{F}(\nabla\mathbf{u}) - \mathcal{F}(\tilde{\
$$

As a direct consequence of this, we obtain

 $(5.20) \quad \Vert \boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u}) \Vert_{2,\Omega}^2 \lesssim \Vert \boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_h) - \boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}) \Vert_{2,\Omega}^2 + \Vert \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u}) - \boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{v}_h) \Vert_{2,\Omega}^2 \,.$ On the other hand, for every $(T_h, v_h) \in \Sigma_h \times V_h$, using [\(5.4\),](#page-15-4) [\(5.18\),](#page-21-1) and the ε -Young inequality [\(2.2\)](#page-3-2) with $\psi = (\varphi^*)_{|\mathbf{S}|} \sim (\varphi_{|\nabla u|})^*$ and $\psi = \varphi_{|\nabla u|}$, we obtain

$$
\|\mathcal{F}^*(\mathsf{S}_h) - \mathcal{F}^*(\mathsf{S})\|_{2,\Omega}^2 \lesssim (\mathsf{D}(\mathsf{S}_h) - \mathsf{D}(\mathsf{S}), \mathsf{S}_h - \mathsf{S})_{\Omega} \n= (\mathsf{D}(\mathsf{S}_h) - \mathsf{D}(\mathsf{S}), \mathsf{S}_h - \mathsf{T}_h)_{\Omega} + (\mathsf{D}(\mathsf{S}_h) - \mathsf{D}(\mathsf{S}), \mathsf{T}_h - \mathsf{S})_{\Omega} \n= (\tilde{\nabla}_h \mathbf{u}_h - \nabla \mathbf{u}, \mathsf{S}_h - \mathsf{T}_h)_{\Omega} + (\mathsf{D}(\mathsf{S}_h) - \mathsf{D}(\mathsf{S}), \mathsf{T}_h - \mathsf{S})_{\Omega} \n= (\tilde{\nabla}_h \mathbf{u}_h - \nabla \mathbf{u}, \mathsf{S} - \mathsf{T}_h)_{\Omega} + (\tilde{\nabla}_h \mathbf{u}_h - \nabla \mathbf{u}, \mathsf{S}_h - \mathsf{S})_{\Omega} \n+ (\mathsf{D}(\mathsf{S}_h) - \mathsf{D}(\mathsf{S}), \mathsf{T}_h - \mathsf{S})_{\Omega} \n= (\tilde{\nabla}_h \mathbf{u}_h - \nabla \mathbf{u}, \mathsf{S} - \mathsf{T}_h)_{\Omega} + (\tilde{\nabla}_h \mathbf{v}_h - \nabla \mathbf{u}, \mathsf{S}_h - \mathsf{S})_{\Omega} \n+ (\tilde{\nabla}_h \mathbf{u}_h - \tilde{\nabla}_h \mathbf{v}_h, \mathsf{S}_h - \mathsf{S})_{\Omega} + (\mathsf{D}(\mathsf{S}_h) - \mathsf{D}(\mathsf{S}), \mathsf{T}_h - \mathsf{S})_{\Omega} \n\leq \varepsilon \|\mathcal{F}(\tilde{\nabla}_h \mathbf{u}_h) - \mathcal{F}(\nabla \mathbf{u})\|_{2,\Omega}^2 + c_{\varepsilon} \|\mathcal{F}^*(\mathsf{T}_h) - \mathcal{F}^*(\mathsf{S})\|_{2,\Omega}^2 \n+ \varepsilon \|\mathcal{F}^*(\mathsf{S}_h)
$$

Moreover, abbreviating $z_h := u_h - v_h \in V_h$, using [\(5.19\)](#page-21-2) and $\tilde{\nabla}_h \mathcal{E}_h z_h = \nabla \mathcal{E}_h z_h$ (cf. [Remark 4.1\)](#page-9-3), we have that

$$
(5.22) \qquad (\mathbf{S}_h - \mathbf{S}, \tilde{\nabla}_h z_h)_{\Omega}
$$
\n
$$
= (\mathbf{S}_h - \mathbf{S}, \tilde{\nabla}_h (z_h - \mathcal{E}_h z_h))_{\Omega} + (\mathbf{S}_h - \mathbf{S}, \tilde{\nabla}_h \mathcal{E}_h z_h)_{\Omega}
$$
\n
$$
= (\mathbf{S}_h - \mathbf{S}, \tilde{\nabla}_h (z_h - \mathcal{E}_h z_h))_{\Omega} - \alpha \langle \mathbf{S}_{\beta_h (u_h)} (h_{\Gamma}^{-1} [\![u_h \otimes n]\!]), [\![z_h \otimes n]\!] \rangle_{\Gamma_h}.
$$

Using the ε -Young inequality [\(2.2\)](#page-3-2) with $\psi = \varphi_{|\tilde{\nabla}_h \mathbf{u}_h|}$, that $\varphi_{|\tilde{\nabla}_h \mathbf{u}_h|} \leq \varphi_{\boldsymbol{\beta}_h(\mathbf{u}_h)}$, a shift change in [Lemma 2.5,](#page-5-0) [Proposition 3.1,](#page-8-4) $z_h = u_h - v_h \in V_h$, and [\(5.4\)](#page-15-4) with $\nabla u = \mathcal{D}(\mathsf{S})$, we obtain

$$
\begin{aligned}\n &\text{(S}_h - \mathsf{S}, \tilde{\nabla}_h (z_h - \mathcal{E}_h z_h))_{\Omega} \\
 &\leq \varepsilon \, \rho_{(\varphi_{|\tilde{\nabla}_h u_h|})^*, \Omega}(\mathsf{S}_h - \mathsf{S}) + c_{\varepsilon} \, \rho_{\varphi_{|\tilde{\nabla}_h u_h|}, \Omega}(\tilde{\nabla}_h (z_h - \mathcal{E}_h z_h)) \\
 &\leq \varepsilon \, \rho_{(\varphi_{|\tilde{\nabla}_h u_h|})^*, \Omega}(\mathsf{S}_h - \mathsf{S}) + c_{\varepsilon} \, \rho_{\varphi_{\beta_h(u_h)}, \Omega}(\tilde{\nabla}_h (z_h - \mathcal{E}_h z_h)) \\
 &\leq \varepsilon \, \rho_{(\varphi_{|\nabla u|})^*, \Omega}(\mathsf{S}_h - \mathsf{S}) + \varepsilon \, \|\mathcal{F}(\tilde{\nabla}_h u_h) - \mathcal{F}(\nabla u)\|_{2,\Omega}^2 \\
 &\quad + c_{\varepsilon} \, m_{\varphi_{\beta_h(u_h)}, h}(u_h) + c_{\varepsilon} \, m_{\varphi_{\beta_h(u_h)}, h}(v_h) \\
 &\leq \varepsilon \, \|\mathcal{F}^*(\mathsf{S}_h) - \mathcal{F}^*(\mathsf{S})\|_{2,\Omega}^2 + \varepsilon \, \|\mathcal{F}(\tilde{\nabla}_h u_h) - \mathcal{F}(\nabla u)\|_{2,\Omega}^2 \\
 &\quad + c_{\varepsilon} \, m_{\varphi_{\beta_h(u_h)}, h}(u_h) + c_{\varepsilon} \, m_{\varphi_{\beta_h(u_h)}, h}(v_h),\n \end{aligned}
$$

and

(5.24)
$$
\begin{array}{ll}\n(\mathcal{S}_{\beta_h(\boldsymbol{u}_h)}(h_\Gamma^{-1}[\![\boldsymbol{u}_h\otimes\boldsymbol{n}]\!]) , [\![z_h\otimes\boldsymbol{n}]\!])_\Omega \\
\geq (1-\varepsilon) m_{\varphi_{\beta_h(\boldsymbol{u}_h)}}, h(\boldsymbol{u}_h) - c_\varepsilon m_{\varphi_{\beta_h(\boldsymbol{u}_h)}}, h(\boldsymbol{v}_h).\n\end{array}
$$

Combining (5.20) – (5.24) , choosing $\varepsilon > 0$ sufficiently small, we arrive at

$$
\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + \|\boldsymbol{\mathcal{F}}^*(\boldsymbol{S}_h) - \boldsymbol{\mathcal{F}}^*(\boldsymbol{S})\|_{2,\Omega}^2 + (\alpha - c) m_{\varphi_{\beta_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)
$$

\$\lesssim \inf_{(\boldsymbol{\mathsf{T}}_h, \boldsymbol{v}_h) \in \Sigma_h \times V_h} \left(\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{S}) - \boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{T}}_h)\|_{2,\Omega}^2 + \|\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u}) - \boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{v}_h)\|_{2,\Omega}^2 + (\alpha + c) m_{\varphi_{\beta_h(\boldsymbol{u}_h)},h}(\boldsymbol{v}_h) \right).

Eventually, for $\alpha > 0$ sufficiently large, we obtain the desired best-approximation result. \Box

As a first immediate consequence of the best-approximation result in [Theorem 5.5,](#page-18-0) we obtain the convergence of the method under minimal regularity assumptions, i.e., merely $u \in V$, $S \in \Sigma$, and $f \in V^*$.

COROLLARY 5.8 (Convergence). For $\alpha > 0$ sufficiently large, it holds that

$$
\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + \|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_h) - \boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2 + m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h) \to 0 \quad (h \to 0).
$$

Proof. Using the stability and approximation properties of \prod_{h}^{k-1} , and the density of smooth functions, we obtain Π_h^{k-1} **S** \to **S** in $L^{p'}(\Omega)^{n \times d}$ $(h \to 0)$, which implies that

(5.25)
$$
\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{S})-\boldsymbol{\mathcal{F}}^*(\Pi_h^{k-1}\boldsymbol{S})\|_{2,\Omega}^2 \lesssim \|\boldsymbol{S}-\Pi_h^{k-1}\boldsymbol{S}\|_{p'}^{\min\{p',2\}} \to 0 \quad (h \to 0)
$$

with a constant depending possibly on δ , $||\mathbf{S}||_{p'}$. Therefore, choosing $(\mathbf{T}_h, \mathbf{v}_h)$ = $(\prod_{h}^{k-1}S, \Pi_h^{SZ}u) \in \Sigma_h \times V_h$ in [Theorem 5.5,](#page-18-0) using [\(5.25\),](#page-23-0) [\(4.12\),](#page-12-2) and [\(4.13\),](#page-12-3) we conclude that

$$
\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + \|\boldsymbol{\mathcal{F}}^*(\boldsymbol{S}_h) - \boldsymbol{\mathcal{F}}^*(\boldsymbol{S})\|_{2,\Omega}^2 + m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h) \\ \lesssim \|\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u}) - \boldsymbol{\mathcal{F}}(\nabla \Pi_h^{SZ} \boldsymbol{u})\|_{2,\Omega}^2 + \|\boldsymbol{\mathcal{F}}^*(\boldsymbol{S}) - \boldsymbol{\mathcal{F}}^*(\Pi_h^{k-1} \boldsymbol{S})\|_{2,\Omega}^2 \to 0 \quad (h \to 0),
$$

which is the claimed convergence under minimal regularity assumptions.

COROLLARY 5.9 (Fractional convergence rates). Assume that the family of triangulations $\{\mathcal{T}_h\}_h$ is quasi-uniform, and that $\mathcal{F}(\nabla u) \in N^{\beta,2}(\Omega)^{n \times d}$ for some $\beta \in (0,1]$. Then, for $\alpha > 0$ sufficiently large, it holds that

$$
\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h \boldsymbol{u}_h) - \boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\|_{2,\Omega}^2 + \|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_h) - \boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2 + m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h) \\ \lesssim h^{2\beta} \left[\boldsymbol{\mathcal{F}}(\nabla \boldsymbol{u})\right]_{N^{\beta,2}(\Omega)}^2.
$$

Proof. First, we note that $\mathcal{F}(\nabla u) \in N^{\beta,2}(\Omega)^{n \times d}$ for $\beta \in (0,1]$ is equivalent to $\mathcal{F}^*(\mathsf{S}) \in N^{\beta,2}(\Omega)^{n \times d}$ for $\beta \in (0,1]$ and that $[\mathcal{F}(\nabla u)]_{N^{\beta,2}(\Omega)} \sim [\mathcal{F}^*(\mathsf{S})]_{N^{\beta,2}(\Omega)}$. This is an immediate consequence of the fact that, due to (2.11) , for every $h \in \mathbb{R}^d \setminus \{0\}$ and $x \in \Omega \cap (\Omega - h)$, we have that

$$
|\mathcal{F}(\nabla \boldsymbol{u}(x+h)) - \mathcal{F}(\nabla \boldsymbol{u}(x))|^2 \sim |\mathcal{F}^*(\mathsf{S}(x+h)) - \mathcal{F}^*(\mathsf{S}(x))|^2.
$$

Using that Π_h^0 **S** = $\Pi_h^{k-1} \Pi_h^0$ **S**, [\(2.10\),](#page-4-8) the Orlicz-stability of Π_h^{k-1} (cf. [\[26,](#page-31-18) Cor. A.8]), $(2.10), [14, \text{ Lem. } 4.4], \text{ and } [8, (4.6), (4.7)], \text{ we obtain}$ $(2.10), [14, \text{ Lem. } 4.4], \text{ and } [8, (4.6), (4.7)], \text{ we obtain}$ $(2.10), [14, \text{ Lem. } 4.4], \text{ and } [8, (4.6), (4.7)], \text{ we obtain}$ $(2.10), [14, \text{ Lem. } 4.4], \text{ and } [8, (4.6), (4.7)], \text{ we obtain}$ $(2.10), [14, \text{ Lem. } 4.4], \text{ and } [8, (4.6), (4.7)], \text{ we obtain}$ $(2.10), [14, \text{ Lem. } 4.4], \text{ and } [8, (4.6), (4.7)], \text{ we obtain}$

$$
\|\mathcal{F}^*(\mathsf{S}) - \mathcal{F}^*(\Pi_h^{k-1}\mathsf{S})\|_{2,\Omega}^2 \lesssim \|\mathcal{F}^*(\mathsf{S}) - \mathcal{F}^*(\Pi_h^0\mathsf{S})\|_{2,\Omega}^2 + \|\mathcal{F}^*(\Pi_h^0\mathsf{S}) - \mathcal{F}^*(\Pi_h^{k-1}\mathsf{S})\|_{2,\Omega}^2
$$

\n
$$
\lesssim \|\mathcal{F}^*(\mathsf{S}) - \mathcal{F}^*(\Pi_h^0\mathsf{S})\|_{2,\Omega}^2 + \rho_{\varphi_{|\Pi_h^0\mathsf{S}|}^*}(\Pi_h^{k-1}(\Pi_h^0\mathsf{S}-\mathsf{S}))
$$

\n
$$
\lesssim \|\mathcal{F}^*(\mathsf{S}) - \mathcal{F}^*(\Pi_h^0\mathsf{S})\|_{2,\Omega}^2 + \rho_{\varphi_{|\Pi_h^0\mathsf{S}|}^*}(\Pi_h^0\mathsf{S}-\mathsf{S})
$$

\n(5.26)
\n
$$
\lesssim \|\mathcal{F}^*(\mathsf{S}) - \mathcal{F}^*(\Pi_h^0\mathsf{S})\|_{2,\Omega}^2
$$

\n
$$
\lesssim \|\mathcal{F}^*(\mathsf{S}) - \Pi_h^0\mathcal{F}^*(\mathsf{S})\|_{2,\Omega}^2
$$

\n
$$
\lesssim h^{2\beta} \|\mathcal{F}^*(\mathsf{S})\|_{N^{\beta,2}(\Omega)}^2
$$

\n
$$
\sim h^{2\beta} \|\mathcal{F}(\nabla u)\|_{N^{\beta,2}(\Omega)}^2.
$$

 \Box

Therefore, choosing $(\mathbf{T}_h, \mathbf{v}_h) = (\Pi_h^{k-1} \mathbf{S}, \Pi_h^{SZ} \mathbf{u}) \in \Sigma_h \times V_h$ in [Theorem 5.5,](#page-18-0) using [\(5.26\),](#page-23-1) [\(4.12\),](#page-12-2) and [\(4.14\),](#page-13-1) we conclude that

$$
\begin{aligned}&\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_h)-\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\\ &\lesssim \|\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})-\boldsymbol{\mathcal{F}}(\nabla\Pi_h^{SZ}\boldsymbol{u})\|_{2,\Omega}^2+\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})-\boldsymbol{\mathcal{F}}^*(\Pi_h^{k-1}\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2\\ &\lesssim h^{2\beta}\,[\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})]_{N^{\beta,2}(\Omega)}^2\,, \end{aligned}
$$

which is the claimed fractional a priori error estimate.

REMARK 5.10. In view of [\[15,](#page-31-16) [Corollary](#page-23-2) 5.8] the assertion of Corollary 5.9 for $\beta = 1$ is also valid if $\mathcal{F}(\nabla u) \in W^{1,2}(\Omega)^{n \times d}$ without the additional assumption that the triangulation $\{\mathcal{T}_h\}_h$ is quasi-uniform.

COROLLARY 5.11 (Ansatz class competition). Let $k = 1$ and $u_h^c \in V_{h,c} \coloneqq$ $V_h \cap W_0^{1,p}(\Omega)$ the continuous Lagrange solution of [\(1.1\)](#page-0-0), cf. [\(4.15\)](#page-14-2). Then, for $\alpha > 0$ sufficiently large, it holds that

$$
\begin{aligned}\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_h)-\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\\ &\sim\|\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u}_h^c)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2\,,\end{aligned}
$$

i.e., the approximation capabilities of the discrete mixed formulation [\(5.7\)](#page-16-0) and the continuous Lagrange approximation (4.15) of (1.1) are comparable.

Proof. ad \lesssim . Using [Theorem 5.5](#page-18-0) with $(\mathbf{T}_h, \mathbf{v}_h) = (\mathbf{S}(\nabla \mathbf{u}_h^c), \mathbf{u}_h^c) \in \Sigma_h \times V_{h,c} \subseteq$ $\Sigma_h \times V_h$, that $m_{\varphi_{\mathcal{B}_h(u_h)},h}(\boldsymbol{u}_h^c) = 0$, and (2.8) , we find that

$$
\begin{aligned}\|\boldsymbol{\mathcal{F}}(\tilde{\nabla}_h\boldsymbol{u}_h)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_h)-\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2+m_{\varphi_{\boldsymbol{\beta}_h(\boldsymbol{u}_h)},h}(\boldsymbol{u}_h)\\ \lesssim \|\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u}_h^c)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2+\|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}(\nabla\boldsymbol{u}_h^c))-\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}^2\\ \lesssim \|\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u}_h^c)-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}^2\,. \end{aligned}
$$

 $ad \geq$. This is proved in [\(4.17\).](#page-14-3)

Putting everything together, we arrive at the claimed equivalence.

6. Numerical experiments. In this section, we show numerical results that confirm our theoretical findings, in particular, [Corollaries 4.8](#page-13-0) and [5.9.](#page-23-2) In our implementation, the max-shift in the jump penalisation is handled through a fixed point iteration, i.e., when solving the discrete primal formulation [\(4.2\)](#page-9-1) starting from a solution guess $u_h^{(k-1)} \in V_h$, we define the residual $\mathfrak{F}(u_h^{(k-1)})$ $\binom{(k-1)}{h}$; \cdot \in $(V_h)^*$, for every $\boldsymbol{v}_h, \boldsymbol{z}_h \in V_h,$ via

$$
\begin{aligned} &\langle \mathfrak{F}(\bm{u}_h^{k-1};\bm{v}_h),\bm{z}_h\rangle_{V_h}\coloneqq (\bm{\mathcal{S}}(\tilde\nabla_h\bm{v}_h),\nabla \bm{\mathcal{E}}_h\bm{z}_h)_{\Omega} \\ &\quad + \alpha \langle \bm{\mathcal{S}}_{\beta_h(\bm{u}_h^{k-1})}(h_\Gamma^{-1}[\![\bm{v}_h\otimes \bm{n}]\!]), [\![\bm{z}_h\otimes \bm{n}]\!])_{\Gamma_h} - \langle \bm{f},\bm{\mathcal{E}}_h\bm{v}_h\rangle_V\,, \end{aligned}
$$

and, then, apply Newton's method to find the next guess $u_h^{(k)} \in V_h$. At a given Newton step, the linear systems are solved using the sparse direct solver MUMPS [\[1\]](#page-30-10). All the examples were implemented using Firedrake [\[23\]](#page-31-19) and PETSc [\[4\]](#page-30-11). The complete code for reproducing the experiments can be found at $[6]$ with exact version of its dependencies being recorded at [\[43\]](#page-32-8), additionally using Gmsh version 4.8.4 [\[19\]](#page-31-20).

Only polynomial degree $k = 1$ is considered. Note that, in this case, the smoothing operator only needs to be applied on the forcing term, since $\mathcal{S}(\tilde{\nabla}_h\boldsymbol{u}_h) \in \mathbb{P}^0(\mathcal{T}_h)^{n\times d}$. In addition, in all the examples, we restrict to the case $d = 2$ and $n = 1$.

 \Box

 \Box

FIG. 1. Initial mesh \mathcal{T}_{h_0}

6.1. Primal formulation. For the experiments based on the discrete primal formulation [\(4.2\),](#page-9-1) we employ the non-linear term $\mathcal{S} \colon \mathbb{R}^{1 \times 2} \to \mathbb{R}^{1 \times 2}$, for every $\mathsf{D} \in$ $\mathbb{R}^{1\times 2}$, defined via

(6.1)
$$
\mathcal{S}(\mathsf{D}) \coloneqq (\delta + |\mathsf{D}|)^{p-2} \mathsf{D},
$$

which has (p, δ) -structure according to [Definition 2.1.](#page-3-0) We choose $\delta = 0.01$ and various values of $p \in [1.5, 4.5]$. The shifted constitutive relation S_a in the jump penalty term in [\(4.2\)](#page-9-1) is then [\(2.6\)](#page-4-3) and $\alpha = 10$ is chosen. The computational domain is defined via $\Omega := (-1,1)^2$ and, for $\beta > 1 - \frac{2}{p}$, the exact solution $u \in V$ with a point singularity at the origin, for every $x \coloneqq (x_1, x_2)^\top \in \Omega$, is defined via

$$
\mathbf{u}(x) = (1-x_1^2)(1-x_2^2)|x|^\beta.
$$

Note that $\beta > 1 - \frac{2}{p}$ guarantees that at least $u \in V$. We start with an initial unstructured mesh \mathcal{T}_{h_0} with 517 elements, 257 vertices, of which one is at the origin, and $h_0 \approx 0.1668$; see [Figure 1.](#page-25-0) We consider six additional levels of uniform refinement, i.e., \mathcal{T}_{h_l} , $l \in \{1, \ldots, 6\}$, where $h_l = \frac{h_{l-1}}{2}$ for all $l \in \{1, \ldots, 6\}$. The error corresponding to the discrete solution $u_{h_\ell} \in V_{h_\ell}$ of (4.2) , associated to a given refinement level $\ell \in \{0,\ldots,6\}$, is defined via

$$
e_{\ell} \coloneqq \Vert \bm{\mathcal{F}}(\nabla_{h_{\ell}}\bm{u}_{h_{\ell}})-\bm{\mathcal{F}}(\nabla \bm{u})\Vert_{2,\Omega}+\big(\alpha\, m_{\varphi_{\beta_h}(\bm{u}_{h_{\ell}}),h}(\bm{u}_{h_{\ell}})\big)^{\frac{1}{2}}.
$$

The experimental rate of convergence is then set to

(6.2)
$$
\operatorname{EOC}_{\ell} := \frac{\log(e_{\ell}/e_{\ell-1})}{\log(h_{\ell}/h_{\ell-1})}.
$$

An important observation is that $\beta > 0$ determines the regularity of the exact solution and, thus, also the expected rate of convergence. More precisely, to obtain a rate of convergence $\rho \in (0, 1]$, one needs to choose $\beta > 1 - \frac{2(1-\rho)}{n}$ $\frac{(-\rho)}{p}$. [Tables 1,](#page-26-0) [3,](#page-26-1) and [5](#page-27-0) show the results for the IIDG formulation for expected rates of convergence of 1, 0.5, and 0.2, respectively; [Tables 2,](#page-26-2) [4,](#page-27-1) and [6](#page-27-2) show the same for the LDG formulation. It can be observed that the values are in agreement with the theoretical predictions.

 ${\bf Table ~1}$ Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal IIDG formulation, i.e., [\(4.2\)](#page-9-1) with $\tilde{\nabla}_{h_{\ell}} = \nabla_{h_{\ell}}$, with $\beta = 1.01$ and, thus, $\rho = 1.0$.

TABLE 2 Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal LDG formulation, i.e., (4.2) with $\tilde{\nabla}_{h_{\ell}} = \mathcal{G}_{h_{\ell}}^1$, with $\beta = 1.01$ and, thus, $\rho = 1.0$.

\mathcal{p} h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.932	0.928	0.910	0.939	0.970
0.0417	0.951	0.953	0.944	0.963	0.972
0.0208	0.961	0.960	0.956	0.971	0.978
0.0104	0.963	0.963	0.961	0.976	0.985
0.0052	0.965	0.965	0.964	0.979	0.989
0.0026	0.967	0.966	0.967	0.980	0.991
Expected	$1.0\,$	1.0	1.0	1.0	1.0

TABLE 3 Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal IIDG formulation, i.e., (4.2) with $\tilde{\nabla}_{h_{\ell}} = \nabla_{h_{\ell}}$, with $\beta = 1.01 - \frac{1}{p}$ and, thus, $\rho = 0.5$.

$\boldsymbol{\eta}$ h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.672	0.659	0.662	0.758	0.863
0.0417	0.631	0.613	0.616	0.684	0.818
0.0208	0.585	0.569	0.573	0.612	0.762
0.0104	0.553	0.542	0.544	0.566	0.667
0.0052	0.533	0.526	0.528	0.541	0.600
0.0026	0.522	0.518	0.519	0.528	0.562
Expected	0.5	0.5	0.5	0.5	0.5

TABLE $\,4\,$ Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal LDG formulation, i.e., (4.2) with $\tilde{\nabla}_{h_{\ell}} = \mathcal{G}_{h_{\ell}}^1$, with $\beta = 1.01 - \frac{1}{p}$ and, thus, $\rho = 0.5$.

\mathcal{p} h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.671	0.659	0.658	0.744	0.857
0.0417	0.630	0.613	0.614	0.675	0.824
0.0208	0.584	0.569	0.572	0.610	0.747
0.0104	0.553	0.542	0.544	0.566	0.661
0.0052	0.533	0.526	0.528	0.541	0.599
0.0026	0.522	0.518	0.519	0.528	0.562
Expected	0.5	0.5	0.5	0.5	0.5

TABLE 5 Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal IIDG formulation, i.e., (4.2) with $\tilde{\nabla}_{h_{\ell}} = \nabla_{h_{\ell}}$, with $\beta = 1.01 - \frac{8}{5p}$ and, thus, $\rho = 0.2$.

\mathcal{D} h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.475	0.560	0.355	0.330	0.461
0.0417	0.332	0.374	0.267	0.255	0.315
0.0208	0.258	0.273	0.231	0.228	0.253
0.0104	0.227	0.231	0.217	0.219	0.232
0.0052	0.215	0.216	0.212	0.216	0.226
0.0026	0.211	0.211	0.211	0.215	0.224
Expected	0.2	0.2	0.2	0.2	0.2

TABLE 6

Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal LDG formulation, i.e., (4.2) with $\tilde{\nabla}_{h_{\ell}} = \mathcal{G}_{h_{\ell}}^1$, with $\beta = 1.01 - \frac{8}{5p}$ and, thus, $\rho = 0.2$.

h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.475	0.557	0.351	0.323	0.440
0.0417	0.332	0.373	0.266	0.254	0.308
0.0208	0.258	0.273	0.230	0.228	0.252
0.0104	0.227	0.231	0.217	0.219	0.232
0.0052	0.215	0.216	0.212	0.216	0.226
0.0026	0.211	0.211	0.211	0.215	0.224
Expected	0.2	0.2	0.2	0.2	0.2

As mentioned in [Remark 4.11,](#page-14-4) our results do not cover the case of a Crouzeix– Raviart discretisation without jump stabilisation terms. However, as seen in [Tables 7](#page-28-0) to [9,](#page-28-1) the rates are roughly in agreement with the same rates as the DG discretisation, suggesting that there might be a proof strategy that also covers this case.

TABLE $\,$ 7 Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal Crouzeix–Raviart for-
mulation (without jump stabilisation), i.e., $\tilde{\nabla}_{h_{\ell}} = \nabla_{h_{\ell}}$ and $\alpha = 0$, with $\beta = 1.01$ and, thus, $\rho = 1.0$.

\mathcal{p} h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.770	0.817	0.878	0.899	1.061
0.0417	0.783	0.834	0.903	0.903	0.929
0.0208	0.835	0.883	0.944	0.963	0.995
0.0104	0.873	0.913	0.961	0.981	1.006
0.0052	0.900	0.931	0.966	0.986	1.004
0.0026	0.918	0.942	0.968	0.986	1.000
Expected	$1.0\,$	1.0	1.0	1.0	1.0

TABLE 8

Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal Crouzeix–Raviart for-
mulation (without jump stabilisation), i.e., $\overline{\nabla}_{h_{\ell}} = \nabla_{h_{\ell}}$ and $\alpha = 0$, with $\beta = 1.01 - \frac{1}{p}$ and, thus, $\rho = 0.5$.

$\,p\,$ h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.517	0.546	0.599	0.643	0.910
0.0417	0.550	0.574	0.607	0.643	0.675
0.0208	0.554	0.565	0.579	0.643	0.697
0.0104	0.540	0.545	0.550	0.614	0.679
0.0052	0.525	0.528	0.531	0.583	0.651
0.0026	0.515	0.518	0.521	0.560	0.625
Expected	0.5	0.5	0.5	0.5	0.5

 ${\bf Table~9}$ Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the primal Crouzeix–Raviart for-
mulation (without jump stabilisation), i.e., $\hat{\nabla}_{h_{\ell}} = \nabla_{h_{\ell}}$ and $\alpha = 0$, with $\beta = 1.01 - \frac{8}{5p}$ and, thus, $\rho = 0.2$.

6.2. Mixed formulation. For the experiments based on the discrete mixed formulation [\(5.7\),](#page-16-0) for which now the following nonlinear term is employed:

$$
\mathcal{D}(\mathsf{S}) := (\delta^{2(p-1)} + |\mathsf{S}|^2)^{\frac{p'-2}{2}} \mathsf{S},
$$

which has (p', δ^{p-1}) -structure in the sense of [Definition 2.1.](#page-3-0) Note that this relation is the inverse of [\(6.1\)](#page-25-1) when $\delta = 0$. In the experiments, we set $\delta = 0.01$. The jump penalty term in [\(5.7\)](#page-16-0) is again defined using [\(2.6\)](#page-4-3) and $\alpha = 10$. The computational domain, once again, $\Omega = (-1, 1)^2$ and we choose the exact flux $S \in \Sigma$ as

$$
\mathbf{S} \coloneqq (\delta^2 + |\nabla \tilde{\mathbf{u}}|^2)^{\frac{p-2}{2}} \nabla \tilde{\mathbf{u}}, \qquad \tilde{\mathbf{u}}(x) = (1 - x_1^2)(1 - x_2^2)|x|^\beta,
$$

where $p > 1$ is specified beforehand, and $\beta \in \mathbb{R}$. The gradient of the exact solution can, then, be computed as $\nabla u := \mathcal{D}(\mathsf{S})$. In the implementation, we set $\tilde{\nabla}_h = \mathcal{G}_h$ in the discrete formulation [\(5.7\),](#page-16-0) which corresponds to an LDG method. For the mixed formulation, we define the error corresponding to a solution $(\mathbf{S}_{h_\ell}, \boldsymbol{u}_{h_\ell}) \in \Sigma_{h_\ell} \times V_{h_\ell}$ associated to a refinement level $\ell \in \{0, \ldots, 6\}$ as:

$$
e_{\ell} \coloneqq \|\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}}_{h_{\ell}})-\boldsymbol{\mathcal{F}}^*(\boldsymbol{\mathsf{S}})\|_{2,\Omega}+\|\boldsymbol{\mathcal{F}}(\nabla_{h_{\ell}}\boldsymbol{u}_{h_{\ell}})-\boldsymbol{\mathcal{F}}(\nabla\boldsymbol{u})\|_{2,\Omega}+\big(\alpha\,m_{\varphi_{\beta_h}(\boldsymbol{u}_{h_{\ell}}),h}(\boldsymbol{u}_{h_{\ell}})\big)^{\frac{1}{2}}.
$$

The experimental order of convergence is then defined analogously to [\(6.2\).](#page-25-2) The results in this case can be found in [Tables 10](#page-29-0) to [12.](#page-30-13)

TABLE 10

Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the mixed LDG formulation, with $\beta = 1.01$ and, thus, $\rho = 1.0$.

\boldsymbol{p} h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.935	0.932	0.916	0.932	0.935
0.0417	0.949	0.951	0.945	0.965	0.976
0.0208	0.958	0.958	0.956	0.973	0.986
0.0104	0.959	0.961	0.961	0.976	0.991
0.0052	0.952	0.964	0.964	0.978	0.992
0.0026	0.920	0.965	0.966	0.979	0.991
Expected	1.0	1.0	1.0	1.0	1.0

TABLE 11 Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, ..., 6\}$, for the mixed LDG formulation, with $\beta = 1.01 - \frac{1}{p}$ and, thus, $\rho = 0.5$.

\boldsymbol{v} h_{ℓ}	1.5	1.7	2.0	3.0	4.5
0.0834	0.691	0.680	0.673	0.744	0.838
0.0417	0.640	0.622	0.620	0.677	0.826
0.0208	0.591	0.575	0.575	0.612	0.745
0.0104	0.556	0.545	0.546	0.568	0.660
0.0052	0.535	0.528	0.529	0.543	0.600
0.0026	0.522	0.519	0.520	0.529	0.564
Expected	0.5	0.5	0.5	0.5	0.5

Experimental order of convergence EOC_{ℓ} , $\ell \in \{1, \ldots, 6\}$, for the mixed LDG formulation, with $\beta = 1.01 - \frac{8}{5p}$ and, thus, $\rho = 0.2$.

REFERENCES

- [1] P. R. Amestoy, I. S. Duff, J.-Y. L'Excellent, and J. Koster, A fully asynchronous multifrontal solver using distributed dynamic scheduling, SIAM J. Matrix Anal. & amp; Appl., 23 (2001), pp. 15–41, [https://doi.org/10.1137/s0895479899358194.](https://doi.org/10.1137/s0895479899358194)
- [2] S. Badia, R. Codina, T. Gudi, and J. Guzman, Error analysis of discontinuous Galerkin methods for the Stokes problem under minimal regularity, IMA J. Num. Anal., 34 (2013), pp. 800–819, [https://doi.org/10.1093/imanum/drt022.](https://doi.org/10.1093/imanum/drt022)
- [3] A. Baier-Reinio, S. Rhebergen, and G. N. Wells, Analysis of pressure-robust embeddedhybridized discontinuous Galerkin methods for the Stokes problem under minimal regularity, J. Sci. Comput., 92 (2022), p. 51, [https://doi.org/10.1007/s10915-022-01889-6.](https://doi.org/10.1007/s10915-022-01889-6)
- [4] S. Balay, S. Abhyankar, M. Adams, S. Benson, J. Brown, P. Brune, K. Buschelman, E. Constantinescu, L. Dalcin, A. Dener, V. Eijkhout, J. Faibussowitsch, W. Gropp, V. Hapla, T. Isaac, P. Jolivet, D. Karpeev, D. Kaushik, M. Knepley, F. Kong, S. Kruger, D. May, L. McInnes, R. Mills, L. Mitchell, T. Munson, J. Roman, K. Rupp, P. Sanan, J. Sarich, B. Smith, S. Zampini, H. ZHANG, H. ZHANG, AND J. ZHANG, PETSc/TAO users manual (Rev. 3.19), Tech. Report ANL-21/39-Rev.3.19, Office of Scientific and Technical Information (OSTI), Mar. 2023, [https://doi.org/10.2172/1968587.](https://doi.org/10.2172/1968587)
- [5] L. C. Berselli, L. Diening, and M. Růžička, Existence of strong solutions for incompressible fluids with shear dependent viscosities, J. Math. Fluid Mech., 12 (2010), pp. 101–132.
- [6] J. BLECHTA AND P. A. GAZCA-OROZCO, Supporting software for article "Quasi-optimal Discontinuous Galerkin discretisations of the p-Dirichlet problem" v1.2. Zenodo, Nov. 2023, [https://doi.org/10.5281/zenodo.10202168.](https://doi.org/10.5281/zenodo.10202168)
- [7] D. Braess, Finite Elemente, Springer Berlin Heidelberg, 4th revised and extended ed., 2013, [https://doi.org/10.1007/978-3-642-34797-9.](https://doi.org/10.1007/978-3-642-34797-9)
- [8] D. BREIT, L. DIENING, J. STORN, AND J. WICHMANN, The parabolic p-Laplacian with fractional differentiability, IMA J. Num. Anal., 41 (2020), pp. 2110–2138, [https://doi.org/10.](https://doi.org/10.1093/imanum/draa081) [1093/imanum/draa081.](https://doi.org/10.1093/imanum/draa081)
- [9] S. C. BRENNER, S. GU, T. GUDI, AND L.-Y. SUNG, A quadratic C^0 interior penalty method for linear fourth order boundary value problems with boundary conditions of the Cahn–Hilliard
- type, SIAM J. Numer. Anal., 50 (2012), pp. 2088–2110, [https://doi.org/10.1137/110847469.](https://doi.org/10.1137/110847469)
[10] S. C. BRENNER AND M. NEILAN, A \mathcal{C}^0 interior penalty method for a fourth order elliptic singular perturbation problem, SIAM J. Numer. Anal., 49 (2011), pp. 869–892, [https://](https://doi.org/10.1137/100786988) [doi.org/10.1137/100786988.](https://doi.org/10.1137/100786988)
- [11] S. C. BRENNER AND L. R. SCOTT, The Mathematical Theory of Finite Element Methods, vol. 15 of Texts in Applied Mathematics, Springer New York, third ed., 2008, [https://doi.](https://doi.org/10.1007/978-0-387-75934-0) [org/10.1007/978-0-387-75934-0.](https://doi.org/10.1007/978-0-387-75934-0)
- [12] L. Diening and F. Ettwein, Fractional estimates for non-differentiable elliptic systems with general growth, Forum Math., 20 (2008), pp. 523–556, [https://doi.org/10.1515/forum.2008.](https://doi.org/10.1515/forum.2008.027) [027.](https://doi.org/10.1515/forum.2008.027)
- [13] L. Diening and C. Kreuzer, Linear convergence of an adaptive finite element method for the p-Laplacian equation, SIAM J. Numer. Anal., 46 (2008), pp. 614–638, [https://doi.org/](https://doi.org/10.1137/070681508) [10.1137/070681508.](https://doi.org/10.1137/070681508)
- [14] L. Diening, D. Kröner, M. Růžička, and I. Toulopoulos, A Local Discontinuous Galerkin approximation for systems with p-structure, IMA J. Num. Anal., 34 (2013), pp. 1447–1488, [https://doi.org/10.1093/imanum/drt040.](https://doi.org/10.1093/imanum/drt040)
- [15] L. Diening and M. Růžička, Interpolation operators in Orlicz–Sobolev spaces, Numer. Math., 107 (2007), pp. 107–129, [https://doi.org/10.1007/s00211-007-0079-9.](https://doi.org/10.1007/s00211-007-0079-9)
- [16] I. EKELAND AND R. TEMAM, Convex analysis and variational problems, vol. 28 of Classics in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, english ed., 1999, [https://doi.org/10.1137/1.9781611971088.](https://doi.org/10.1137/1.9781611971088) Translated from the French.
- [17] A. ERN AND J.-L. GUERMOND, Finite elements I—Approximation and interpolation, vol. 72 of Texts in Applied Mathematics, Springer, Cham, 2021, [https://doi.org/10.1007/](https://doi.org/10.1007/978-3-030-56341-7) [978-3-030-56341-7.](https://doi.org/10.1007/978-3-030-56341-7)
- [18] E. H. Georgoulis and T. Pryer, Recovered finite element methods, Computer Methods in Applied Mechanics and Engineering, 332 (2018), pp. 303–324, [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.cma.2017.12.026) [cma.2017.12.026.](https://doi.org/10.1016/j.cma.2017.12.026)
- [19] C. Geuzaine and J.-F. Remacle, Gmsh: A 3-D finite element mesh generator with builtin pre- and post-processing facilities, Internat. J. Numer. Methods Engrg., 79 (2009), pp. 1309–1331, [https://doi.org/10.1002/nme.2579.](https://doi.org/10.1002/nme.2579)
- [20] T. GUDI, A new error analysis for discontinuous finite element methods for linear elliptic problems, Math. Comp., 79 (2010), pp. 2169–2189, [https://doi.org/10.1090/](https://doi.org/10.1090/s0025-5718-10-02360-4) [s0025-5718-10-02360-4.](https://doi.org/10.1090/s0025-5718-10-02360-4)
- [21] T. GUDI, H. S. GUPTA, AND N. NATARAJ, Analysis of an interior penalty method for fourth order problems on polygonal domains, J. Sci. Comput., 54 (2012), pp. 177–199, [https:](https://doi.org/10.1007/s10915-012-9612-9) $\frac{1}{\dot{c}}$ /doi.org/10.1007/s10915-012-9612-9.
- [22] T. GUDI AND M. NEILAN, An interior penalty method for a sixth-order elliptic equation, IMA J. Num. Anal., 31 (2011), pp. 1734–1753, [https://doi.org/10.1093/imanum/drq031.](https://doi.org/10.1093/imanum/drq031)
- [23] D. A. Ham, P. H. J. Kelly, L. Mitchell, C. J. Cotter, R. C. Kirby, K. Sagiyama, N. Bouziani, S. Vorderwuelbecke, T. J. Gregory, J. Betteridge, D. R. Shapero, R. W. Nixon-Hill, C. J. Ward, P. E. Farrell, P. D. Brubeck, I. Marsden, T. H. Gibson, M. Homolya, T. Sun, A. T. T. McRae, F. Luporini, A. Gregory, M. Lange, S. W. Funke, F. Rathgeber, G.-T. Bercea, and G. R. Markall, Firedrake User Manual, Imperial College London and University of Oxford and Baylor University and University of Washington, first edition ed., 5 2023, [https://doi.org/10.](https://doi.org/10.25561/104839) [25561/104839.](https://doi.org/10.25561/104839)
- [24] P. HARJULEHTO AND P. HÄSTÖ, Orlicz spaces and generalized Orlicz spaces, vol. 2236 of Lecture Notes in Mathematics, Springer, 2019, [https://doi.org/10.1007/978-3-030-15100-3.](https://doi.org/10.1007/978-3-030-15100-3)
- [25] A. KALTENBACH, Error analysis for a Crouzeix–Raviart approximation of the p-Dirichlet problem, J. Numer. Math., (2023), [https://doi.org/10.1515/jnma-2022-0106.](https://doi.org/10.1515/jnma-2022-0106)
- [26] A. Kaltenbach and M. Růžička, Convergence analysis of a local discontinuous Galerkin approximation for nonlinear systems with balanced Orlicz-structure, ESAIM Math. Model. Numer. Anal., 57 (2023), pp. 1381–1411, [https://doi.org/10.1051/m2an/2023028.](https://doi.org/10.1051/m2an/2023028)
- [27] A. Kaltenbach and M. Růžička, A local discontinuous Galerkin approximation for the p-Navier–Stokes system, Part I: Convergence analysis, SIAM J. Numer. Anal., 61 (2023), pp. 1613–1640, [https://doi.org/10.1137/22M151474X.](https://doi.org/10.1137/22M151474X)
- [28] C. KREUZER, R. VERFÜRTH, AND P. ZANOTTI, Quasi-optimal and pressure robust discretizations of the Stokes equations by moment- and divergence-preserving operators, Comput. Methods Appl. Math., 21 (2020), pp. 423–443, [https://doi.org/10.1515/cmam-2020-0023.](https://doi.org/10.1515/cmam-2020-0023)
- [29] C. KREUZER AND P. ZANOTTI, Quasi-optimal and pressure-robust discretizations of the Stokes equations by new augmented Lagrangian formulations, IMA J. Num. Anal., 40 (2019), pp. 2553–2583, [https://doi.org/10.1093/imanum/drz044.](https://doi.org/10.1093/imanum/drz044)
- [30] M. Li, S. MAO, AND S. ZHANG, New error estimates of nonconforming mixed finite element methods for the Stokes problem, Math. Meth. Appl. Sci., 37 (2013), pp. 937–951, [https:](https://doi.org/10.1002/mma.2849) [//doi.org/10.1002/mma.2849.](https://doi.org/10.1002/mma.2849)
- [31] A. LINKE, C. MERDON, M. NEILAN, AND F. NEUMANN, Quasi-optimality of a pressure-robust nonconforming finite element method for the Stokes-problem, Math. Comp., 87 (2018), pp. 1543–1566, [https://doi.org/10.1090/mcom/3344.](https://doi.org/10.1090/mcom/3344)
- [32] J. Musielak, Orlicz Spaces and Modular Spaces, Springer Berlin Heidelberg, Berlin, 1983, [https://doi.org/10.1007/bfb0072210.](https://doi.org/10.1007/bfb0072210)
- [33] M. M. RAO AND Z. D. REN, Theory of Orlicz spaces, vol. 146 of Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker Inc., New York, 1991.
- [34] M. Růžička, Nonlinear functional analysis. An introduction. (Nichtlineare Funktionalanalysis. Eine Einführung.), Berlin: Springer. xii, 208 p., 2004.
- [35] M. Růžička and L. Diening, Non-Newtonian fluids and function spaces, in Nonlinear Analysis, Function Spaces and Applications, Proceedings of NAFSA 2006 Prague, vol. 8, Institute of Mathematics of the Academy of Sciences of the Czech Republic, 2007, pp. 95–143, [http://eudml.org/doc/221529.](http://eudml.org/doc/221529)
- [36] L. R. SCOTT AND S. ZHANG, Finite element interpolation of nonsmooth functions satisfying boundary conditions, Math. Comp., 54 (1990), pp. 483–493, [https://doi.org/10.1090/](https://doi.org/10.1090/s0025-5718-1990-1011446-7) [s0025-5718-1990-1011446-7.](https://doi.org/10.1090/s0025-5718-1990-1011446-7)
- [37] G. STRANG, Variational crimes in the finite element method, in The mathematical foundations of the finite element method with applications to partial differential equations (Proc. Sympos., Univ. Maryland, Baltimore, Md., 1972), Academic Press, New York-London, 1972, pp. 689–710.
- [38] A. VEESER AND P. ZANOTTI, Quasi-optimal nonconforming methods for symmetric elliptic problems. I—Abstract theory, SIAM J. Numer. Anal., 56 (2018), pp. 1621–1642, [https:](https://doi.org/10.1137/17m1116362) [//doi.org/10.1137/17m1116362.](https://doi.org/10.1137/17m1116362)
- [39] A. VEESER AND P. ZANOTTI, Quasi-optimal nonconforming methods for symmetric elliptic problems. III—Discontinuous Galerkin and other interior penalty methods, SIAM J. Numer. Anal., 56 (2018), pp. 2871–2894, [https://doi.org/10.1137/17m1151675.](https://doi.org/10.1137/17m1151675)
- [40] A. VEESER AND P. ZANOTTI, Quasi-optimal nonconforming methods for symmetric elliptic problems. II—Overconsistency and classical nonconforming elements, SIAM J. Numer. Anal., 57 (2019), pp. 266–292, [https://doi.org/10.1137/17m1151651.](https://doi.org/10.1137/17m1151651)
- [41] R. VERFÜRTH AND P. ZANOTTI, A quasi-optimal Crouzeix–Raviart discretization of the Stokes equations, SIAM J. Numer. Anal., 57 (2019), pp. 1082–1099, [https://doi.org/10.1137/](https://doi.org/10.1137/18m1177688) [18m1177688.](https://doi.org/10.1137/18m1177688)
- [42] E. ZEIDLER, Nonlinear functional analysis and its applications. II/B, Springer New York, New York, 1990, [https://doi.org/10.1007/978-1-4612-0981-2.](https://doi.org/10.1007/978-1-4612-0981-2) Nonlinear monotone operators.
- [43] Software used in "Quasi-optimal Discontinuous Galerkin discretisations of the p-Dirichlet equation", Nov. 2023, [https://doi.org/10.5281/zenodo.10061067.](https://doi.org/10.5281/zenodo.10061067)